

On the Vertex-Degree-Function Indices of Connected (n, m) -Graphs of Maximum Degree at Most Four

Abeer M. Albalahi¹, Igor Ž. Milovanović², Zahid Raza³,
Akbar Ali^{1,*}, Amjad E. Hamza¹

¹*Department of Mathematics, College of Science,
University of Ha'il, Ha'il, Saudi Arabia*

a.albalahi@uoh.edu.sa, akbarali.maths@gmail.com, aboaljod2@hotmail.com

²*Faculty of Electronic Engineering,
University of Niš, Niš, Serbia*
igor.milovanovic@elfak.ni.ac.rs

³*Department of Mathematics, College of Sciences,
University of Sharjah, Sharjah, UAE*
zraza@sharjah.ac.ae

Abstract

Consider a graph G and a real-valued function f defined on the degree set of G . The sum of the outputs $f(d_v)$ over all vertices $v \in V(G)$ of G is usually known as the vertex-degree-function indices and is denoted by $H_f(G)$, where d_v represents the degree of a vertex v of G . This paper gives sharp bounds on the index $H_f(G)$ in terms of order and size of G when G is connected and has the maximum degree at most 4. All the graphs achieving the derived bounds are also determined. Bounds involving several existing indices – including the general zeroth-order Randić index and coindex, the general multiplicative first/second Zagreb index, the variable sum lodeg index, the variable sum exdeg index – are deduced as the special cases of the obtained ones.

Keywords: chemical graph theory; topological index; vertex-degree-function indices; degree of a vertex.

AMS Subject Classification: 05C07, 05C09, 05C92.

*Corresponding author

1 Introduction

This study is concerned with only connected and finite graphs. The (chemical-)graph theoretical concepts used in the this paper without providing their definitions can be found in the related books like [1–4].

A topological index is a function defined on the set of all graphs with the condition that it remains the same under the graph isomorphism. The degree set of a graph G is the set consisting of all distinct elements of the degree sequence of G . Consider a graph G and a real-valued function f defined on the degree set of G . The sum of the outputs $f(d_v)$ over all vertices $v \in V(G)$ of G is usually known as the vertex-degree-function indices and is denoted by $H_f(G)$, where d_v represents the degree of a vertex v of G . Thus,

$$H_f(G) = \sum_{v \in V(G)} f(d_v). \quad (1)$$

Although the terminology and notation of the topological index H_f that is being used by several researchers was coined by Yao et al. [5], to the best of authors' knowledge such indices were studied first by Linial and Rozenman in [6]. The indices H_f have been the subject of several recent papers; see for example the recent articles [7–9], recent review paper [10], and related publications cited therein.

If vertices u and v are adjacent in G , we write $u \sim v$, otherwise we write $u \not\sim v$. Let $TI(G)$ be a vertex–degree–based topological index of the form

$$TI(G) = \sum_{u \sim v} (f(d_u) + f(d_v)) = \sum_{u \in V(G)} d_u f(d_u).$$

Then, the corresponding coindex, $\overline{TI}(G)$ can be defined as [11–13]:

$$\overline{TI}(G) = \sum_{u \not\sim v} (f(d_u) + f(d_v)) = \sum_{u \in V(G)} (n - 1 - d_u) f(d_u).$$

The following identity is valid

$$TI(G) + \overline{TI}(G) = (n - 1) \sum_{u \in V(G)} f(d_u) = (n - 1) H_f(G). \quad (2)$$

In what follows, some existing indices are given that are special cases of Equation (1).

- Equation (1) gives the general zeroth-order Randić index if $f(x) = x^\alpha$ (see for example [14–18]), where α is a real number.

- The general zeroth-order Randić coindex is obtained from equation (1) corresponding to the choice $f(x) = (n - 1 - x)x^{\alpha-1}$, where α is a real number (see e.g. [19,20]). If $\alpha = 2$, then the forgotten topological coindex $\overline{F}(G) = \sum_{u \in V(G)} (n - 1 - d_u)d_u^2$ is obtained (see for example [21,22])
- One gets the general multiplicative first Zagreb index (general multiplicative second Zagreb index, respectively) [23] by taking $f(x) = \ln x^a$ ($f(x) = \ln x^{ax}$, respectively), where $a \in \mathbb{R}$ (that is the set of real numbers).
- The substitution $f(x) = x(\ln x)^a$ in Equation (1) yields the variable sum lodeg index [24], where $a \in \mathbb{R}$
- If $f(x) = xa^x$ then Equation (1) gives the variable sum exdeg index (see for example [24,25]), where $a > 0$ with $a \neq 1$.

A graph with n vertices and m edges is known as an (n, m) -graph. A chemical graph is the one with the maximum degree at most four. This paper gives sharp bounds on the index $H_f(G)$ for chemical (n, m) -graphs in terms of m and n . All the graphs achieving the derived bounds are also identified. Bounds involving the above-mentioned existing indices (that is, the general zeroth-order Randić index and coindex, the general multiplicative first/second Zagreb index, the variable sum lodeg index, the variable sum exdeg index) are deduced as the special cases of the obtained bounds.

2 Main Results

For a graph G , its number of vertices having the degree r is denoted by n_r . If G is a chemical (n, m) -graph, then

$$H_f(G) = \sum_{i=1}^4 n_i f(i), \quad (3)$$

$$\sum_{i=1}^4 n_i = n, \quad (4)$$

$$\sum_{i=1}^4 i n_i = 2m, \quad (5)$$

We solve Equations (4) and (5) for the quantities n_1, n_4 , and then substitute their values in Equation (3):

$$H_f(G) = \frac{1}{3} \left(4f(1) - f(4) \right) n + \frac{2}{3} \left(f(4) - f(1) \right) m$$

$$+ \left(f(2) - \frac{2}{3}f(1) - \frac{1}{3}f(4) \right) n_2 + \left(f(3) - \frac{1}{3}f(1) - \frac{2}{3}f(4) \right) n_3. \quad (6)$$

Let us take

$$\Gamma_f(G) = \left(f(2) - \frac{2}{3}f(1) - \frac{1}{3}f(4) \right) n_2 + \left(f(3) - \frac{1}{3}f(1) - \frac{2}{3}f(4) \right) n_3. \quad (7)$$

Now, Equation (6) yields

$$H_f(G) = \frac{1}{3} \left(4f(1) - f(4) \right) n + \frac{2}{3} \left(f(4) - f(1) \right) m + \Gamma_f(G). \quad (8)$$

From Equation (8), it is evident that if one wants to establish a bound on H_f for chemical (n, m) -graphs in terms of m and n , it is enough to determine the least or greatest Γ_f -value for chemical (n, m) -graphs. Thence, in the next lemma, we derive a bound Γ_f for chemical (n, m) -graphs.

Lemma 1 *Let G be a chemical (n, m) -graph such that $n_2 + n_3 \geq 2$. Take*

$$\xi_1 = f(2) - \frac{2}{3}f(1) - \frac{1}{3}f(4) \quad \text{and} \quad \xi_2 = f(3) - \frac{1}{3}f(1) - \frac{2}{3}f(4).$$

(i). *If both ξ_1 and ξ_2 are negative such that $2\xi_2 < \xi_1 < \xi_2/2$, then*

$$\Gamma_f(G) < \min \{ \xi_1, \xi_2 \}.$$

(ii). *If both ξ_1 and ξ_2 are positive such that $\xi_2/2 < \xi_1 < 2\xi_2$, then*

$$\Gamma_f(G) > \max \{ \xi_1, \xi_2 \}.$$

Proof: (i) Let $\max \{ \xi_1, \xi_2 \} = \xi_{max}$. Note that

$$\Gamma_f(G) = \xi_1 n_2 + \xi_2 n_3 \leq \xi_{max} (n_2 + n_3) \leq 2\xi_{max} < \min \{ \xi_1, \xi_2 \}.$$

(ii) Let $\min \{ \xi_1, \xi_2 \} = \xi_{min}$. Note that

$$\Gamma_f(G) = \xi_1 n_2 + \xi_2 n_3 \geq \xi_{min} (n_2 + n_3) \geq 2\xi_{min} > \max \{ \xi_1, \xi_2 \}.$$

□

Recall that the degree set of a graph G is the set of all unequal degrees of vertices of G .

Theorem 2 Let G be a chemical (n, m) -graph. Let ξ_1 and ξ_2 be the numbers defined in Lemma 1.

(i). If both ξ_1 and ξ_2 are negative such that $2\xi_2 < \xi_1 < \xi_2/2$, then

$$H_f(G) \leq \frac{1}{3} \left(4f(1) - f(4) \right) n + \frac{2}{3} \left(f(4) - f(1) \right) m$$

$$+ \begin{cases} f(2) - \frac{2}{3}f(1) - \frac{1}{3}f(4) & \text{if } n \equiv 0 \pmod{3} \\ f(3) - \frac{1}{3}f(1) - \frac{2}{3}f(4) & \text{if } n \equiv 1 \pmod{3} \\ 0 & \text{if } n \equiv 2 \pmod{3} \end{cases}$$

with equality if and only if the degree set of G is

- $\{1, 2, 4\}$ and G contains only one vertex of degree 2 whenever $n \equiv 0 \pmod{3}$;
- $\{1, 3, 4\}$ and G contains only one vertex of degree 3 whenever $n \equiv 1 \pmod{3}$;
- $\{1, 4\}$ whenever $n \equiv 2 \pmod{3}$.

(ii) If both ξ_1 and ξ_2 are positive such that $\xi_2/2 < \xi_1 < 2\xi_2$, then

$$H_f(G) \geq \frac{1}{3} \left(4f(1) - f(4) \right) n + \frac{2}{3} \left(f(4) - f(1) \right) m$$

$$+ \begin{cases} f(2) - \frac{2}{3}f(1) - \frac{1}{3}f(4) & \text{if } n \equiv 0 \pmod{3} \\ f(3) - \frac{1}{3}f(1) - \frac{2}{3}f(4) & \text{if } n \equiv 1 \pmod{3} \\ 0 & \text{if } n \equiv 2 \pmod{3} \end{cases}$$

with equality if and only if the degree set of G is

- $\{1, 2, 4\}$ and G contains only one vertex of degree 2 whenever $n \equiv 0 \pmod{3}$;
- $\{1, 3, 4\}$ and G contains only one vertex of degree 3 whenever $n \equiv 1 \pmod{3}$;
- $\{1, 4\}$ whenever $n \equiv 2 \pmod{3}$.

Proof: Because the proofs of the both parts are similar to each other, we prove only Part (i). If the inequality $n_2 + n_3 \geq 2$ holds, then by using Lemma 1 and Equation (8), one has

$$H_f(G) < \frac{1}{3} \left(4f(1) - f(4) \right) n + \frac{2}{3} \left(f(4) - f(1) \right) m$$

$$\begin{aligned}
& + \min \left\{ f(2) - \frac{2}{3}f(1) - \frac{1}{3}f(4), f(3) - \frac{1}{3}f(1) - \frac{2}{3}f(4) \right\} \\
& < \frac{1}{3} \left(4f(1) - f(4) \right) n + \frac{2}{3} \left(f(4) - f(1) \right) m
\end{aligned}$$

as desired.

In the remaining proof, assume that $n_2 + n_3 \leq 1$. Then, $(n_2, n_3) \in \{(0, 0), (1, 0), (0, 1)\}$.

From Equations (4) and (5), it follows that $n_2 + 2n_3 \equiv n - 2 \pmod{3}$, which gives

$$(n_2, n_3) = \begin{cases} (1, 0) & \text{if } n \equiv 0 \pmod{3}, \\ (0, 1) & \text{if } n \equiv 1 \pmod{3}, \\ (0, 0) & \text{if } n \equiv 2 \pmod{3}. \end{cases}$$

The required result follows now from Equation (6). \square

In what follows, we consider some well-known topological indices that satisfy the assumptions of Theorem 2 and hence yield different corollaries of Theorem 2.

First, we take $f(x) = x^\alpha$. Then H_f is the general zeroth-order Randić index ${}^0R_\alpha$. Here, we have

$$\xi_1 = f(2) - \frac{2}{3}f(1) - \frac{1}{3}f(4) = \begin{cases} -\frac{(2^\alpha - 2)(2^\alpha - 1)}{3} < 0 & \text{if either } \alpha > 1 \text{ or } \alpha < 0, \\ -\frac{(2^\alpha - 2)(2^\alpha - 1)}{3} > 0 & \text{if } 0 < \alpha < 1, \end{cases}$$

and

$$\xi_2 = f(3) - \frac{1}{3}f(1) - \frac{2}{3}f(4) = \begin{cases} \frac{3^{\alpha+1} - 2^{2\alpha+1} - 1}{3} < 0 & \text{if either } \alpha > 1 \text{ or } \alpha < 0, \\ \frac{3^{\alpha+1} - 2^{2\alpha+1} - 1}{3} > 0 & \text{if } 0 < \alpha < 1. \end{cases}$$

Also,

$$2\xi_2 = \frac{2(3^{\alpha+1} - 2^{2\alpha+1} - 1)}{3} < \xi_1 = -\frac{(2^\alpha - 2)(2^\alpha - 1)}{3} < \frac{\xi_2}{2} = \frac{3^{\alpha+1} - 2^{2\alpha+1} - 1}{6} \quad (9)$$

holds if either $\alpha > 1$ or $\alpha < 0$. If each inequality sign “<” of (9) is replaced with “>” then the resulting inequality holds for $0 < \alpha < 1$. Thus, we have the following known [26] result as a direct consequence of Theorem 2.

Corollary 1 *Let G be a chemical (n, m) -graph. If either $\alpha > 1$ or $\alpha < 0$, then*

$${}^0R_\alpha(G) \leq \frac{4 - 4^\alpha}{3} n + \frac{2(4^\alpha - 1)}{3} m + \begin{cases} -\frac{(2^\alpha - 2)(2^\alpha - 1)}{3} & \text{if } n \equiv 0 \pmod{3} \\ \frac{3^{\alpha+1} - 2^{2\alpha+1} - 1}{3} & \text{if } n \equiv 1 \pmod{3} \\ 0 & \text{if } n \equiv 2 \pmod{3} \end{cases}$$

with equality if and only if the degree set of G is the same as specified in Theorem 2. If $0 < \alpha < 1$ then the above inequality for ${}^0R_\alpha(G)$ is reversed.

Now, we take $f(x) = xa^x$ with $a > 0$ but $a \neq 1$. Then H_f is the variable sum exdeg index SEI_a . Here, we have

$$\xi_1 = f(2) - \frac{2}{3}f(1) - \frac{1}{3}f(4) = \begin{cases} -\frac{2a(a-1)(2a^2+2a-1)}{3} < 0 & \text{if } a > 1 \\ -\frac{2a(a-1)(2a^2+2a-1)}{3} > 0 & \text{if } \frac{\sqrt{33}+1}{16} < a < 1, \end{cases}$$

and

$$\xi_2 = f(3) - \frac{1}{3}f(1) - \frac{2}{3}f(4) = \begin{cases} -\frac{a(a-1)(8a^2-a-1)}{3} < 0 & \text{if } a > 1 \\ -\frac{a(a-1)(8a^2-a-1)}{3} > 0 & \text{if } \frac{\sqrt{33}+1}{16} < a < 1, \end{cases}$$

Also,

$$\begin{aligned} 2\xi_2 = -\frac{2a(a-1)(8a^2-a-1)}{3} < \xi_1 = -\frac{2a(a-1)(2a^2+2a-1)}{3} \\ < \frac{\xi_2}{2} = -\frac{a(a-1)(8a^2-a-1)}{6} \end{aligned} \quad (10)$$

holds if $a > 1$. If each inequality sign “ $<$ ” in (10) is replaced with “ $>$ ” then the resulting inequality holds for $\frac{\sqrt{33}+1}{16} < a < 1$. Thus, we have the next result follows directly from Theorem 2.

Corollary 2 *Let G be a chemical (n, m) -graph. If $a > 1$, then*

$$SEI_a(G) \leq \frac{4a(1-a^3)n}{3} + \frac{2a(4a^3-1)m}{3} + \begin{cases} \frac{2a(1-a)(2a^2+2a-1)}{3} & \text{if } n \equiv 0 \pmod{3} \\ \frac{a(1-a)(8a^2-a-1)}{3} & \text{if } n \equiv 1 \pmod{3} \\ 0 & \text{if } n \equiv 2 \pmod{3} \end{cases}$$

with equality if and only if the degree set of G is the same as specified in Theorem 2. If $\frac{\sqrt{33}+1}{16} < a < 1$ then the above inequality for $SEI_a(G)$ is reversed.

Next, we take $f(x) = x(\ln x)^a$ with $a > 0$. Then H_f is the variable sum loddeg index SLI_a . Here, for $a > \frac{\ln 3 - \ln 4}{\ln(\ln 2) - \ln(\ln 3)}$ (≈ 0.6246), we have

$$\xi_1 = f(2) - \frac{2}{3}f(1) - \frac{1}{3}f(4) = \frac{2(3(\ln 2)^a - 2(\ln 4)^a)}{3} < 0,$$

$$\xi_2 = f(3) - \frac{1}{3}f(1) - \frac{2}{3}f(4) = \frac{9(\ln 3)^a - 8(\ln 4)^a}{3} < 0$$

and

$$2\xi_2 = \frac{2(9(\ln 3)^a - 8(\ln 4)^a)}{3} < \xi_1 = \frac{2(3(\ln 2)^a - 2(\ln 4)^a)}{3} < \frac{\xi_2}{2} = \frac{9(\ln 3)^a - 8(\ln 4)^a}{6}.$$

Hence, the following corollary is another direct consequence of Theorem 2.

Corollary 3 *Let G be a chemical (n, m) -graph. If*

$$a > \frac{\ln 3 - \ln 4}{\ln(\ln 2) - \ln(\ln 3)} \quad (\approx 0.6246),$$

then

$$SLI_a(G) \leq \frac{8(\ln 4)^a}{3} m - \frac{4(\ln 4)^a}{3} n + \begin{cases} \frac{2(3(\ln 2)^a - 2(\ln 4)^a)}{3} & \text{if } n \equiv 0 \pmod{3} \\ \frac{9(\ln 3)^a - 8(\ln 4)^a}{3} & \text{if } n \equiv 1 \pmod{3} \\ 0 & \text{if } n \equiv 2 \pmod{3} \end{cases}$$

with equality if and only if the degree set of G is the same as specified in Theorem 2.

Finally, if we take $f(x) = (n - 1 - x)x^2$, or $f(x) = \ln x^{ax}$, or $f(x) = \ln x^a$, then H_f is the forgotten topological coindex $\overline{F}(G)$ (see [21, 22]), or the natural logarithm of the general multiplicative first Zagreb index $\ln \Pi_{1,a}$, or the natural logarithm of the general multiplicative second Zagreb index $\ln \Pi_{2,a}$, respectively.

- If we take $f(x) = (n - 1 - x)x^2$ with $n \geq 11$, or $f(x) = \ln x^{ax}$ with $a > 0$, or $f(x) = \ln x^a$ with $a < 0$, then f satisfies the conditions of Theorem 2(i).
- If we take $f(x) = \ln x^a$ with $a > 0$, or $f(x) = \ln x^{ax}$ with $a < 0$, then f satisfies the conditions of Theorem 2(ii).

Hence, the next result follows immediately from Theorem 2.

Corollary 4 *Let G be a chemical (n, m) -graph. If $a < 0$ then*

$$\Pi_{1,a}(G) \leq \begin{cases} 2^{\frac{a(4m-2n+1)}{3}} & \text{if } n \equiv 0 \pmod{3} \\ 2^{\frac{2a(2m-n-2)}{3}} 3^a & \text{if } n \equiv 1 \pmod{3} \\ 0 & \text{if } n \equiv 2 \pmod{3} \end{cases}$$

$$\Pi_{2,a}(G) \geq \begin{cases} 2^{\frac{2a(8m-4n-1)}{3}} & \text{if } n \equiv 0 \pmod{3} \\ 2^{\frac{8a(2m-n-2)}{3}} 3^{3a} & \text{if } n \equiv 1 \pmod{3} \\ 0 & \text{if } n \equiv 2 \pmod{3} \end{cases}$$

and if $n \geq 11$ then

$$\overline{F}(G) \leq \begin{cases} 2(m(5n-26) - (2n-11)n + 8) & \text{if } n \equiv 0 \pmod{3} \\ 2(m(5n-26) - (2n-11)n + 9) & \text{if } n \equiv 1 \pmod{3} \\ 0 & \text{if } n \equiv 2 \pmod{3} \end{cases}$$

where the equality sign in any of the above inequalities involving $\Pi_{1,a}(G)$, $\Pi_{2,a}(G)$, $\overline{F}(G)$, holds if and only if the degree set of G is the same as specified in Theorem 2. If $a > 0$ then the above inequalities involving $\Pi_{1,a}(G)$ and $\Pi_{2,a}(G)$ are reversed.

From Theorem 2 and the identity (2), the next result follows.

Theorem 3 Let G be a chemical (n, m) -graph. Let ξ_1 and ξ_2 be the numbers defined in Lemma 1.

(i) If both ξ_1 and ξ_2 are negative such that $2\xi_2 < \xi_1 < \xi_2/2$, then

$$TI(G) + \overline{TI}(G) \leq (n-1) \left(\frac{1}{3} (4f(1) - f(4))n + \frac{2}{3} (f(4) - f(1))m \right) \\ + (n-1) \begin{cases} f(2) - \frac{2}{3}f(1) - \frac{1}{3}f(4) & \text{if } n \equiv 0 \pmod{3} \\ f(3) - \frac{1}{3}f(1) - \frac{2}{3}f(4) & \text{if } n \equiv 1 \pmod{3} \\ 0 & \text{if } n \equiv 2 \pmod{3} \end{cases}$$

with equality if and only if the degree set of G is

- $\{1, 2, 4\}$ and G contains only one vertex of degree 2 whenever $n \equiv 0 \pmod{3}$;
- $\{1, 3, 4\}$ and G contains only one vertex of degree 3 whenever $n \equiv 1 \pmod{3}$;
- $\{1, 4\}$ whenever $n \equiv 2 \pmod{3}$.

(ii) If both ξ_1 and ξ_2 are positive such that $\xi_2/2 < \xi_1 < 2\xi_2$, then

$$TI(G) + \overline{TI}(G) \geq (n-1) \left(\frac{1}{3} (4f(1) - f(4))n + \frac{2}{3} (f(4) - f(1))m \right)$$

$$+ (n - 1) \begin{cases} f(2) - \frac{2}{3}f(1) - \frac{1}{3}f(4) & \text{if } n \equiv 0 \pmod{3} \\ f(3) - \frac{1}{3}f(1) - \frac{2}{3}f(4) & \text{if } n \equiv 1 \pmod{3} \\ 0 & \text{if } n \equiv 2 \pmod{3} \end{cases}$$

with equality if and only if the degree set of G is

- $\{1, 2, 4\}$ and G contains only one vertex of degree 2 whenever $n \equiv 0 \pmod{3}$;
- $\{1, 3, 4\}$ and G contains only one vertex of degree 3 whenever $n \equiv 1 \pmod{3}$;
- $\{1, 4\}$ whenever $n \equiv 2 \pmod{3}$.

3 Funding Statement

This research has been funded by Scientific Research Deanship at University of Ha'il, Saudi Arabia.

4 Data Availability Statement

Data about this study may be requested from the authors.

References

- [1] J. A. Bondy, U. S. R. Murty, Graph Theory, Springer, London, 2008.
- [2] G. Chartrand, L. Lesniak, P. Zhang, Graphs & Digraphs, Sixth Edition, CRC Press, Boca Raton, 2016.
- [3] N. Trinajstić, Chemical Graph Theory, CRC Press, Boca Raton, 1992.
- [4] S. Wagner, H. Wang, Introduction to Chemical Graph Theory, CRC Press, Boca Raton, 2018.
- [5] Y. Yao, M. Liu, F. Belardo, C. Yang, Unified extremal results of topological index and graph spectrum, Discrete Appl. Math. 271 (2019) 218–232.
- [6] N. Linial, E. Rozenman, An extremal problem on degree sequences of graphs, Graphs Combin. 18 (2002) 573–582.

- [7] Z. Hu, L. Li, X. Li, D. Peng, Extremal graphs for topological index defined by a degree-based edge-weight function, *MATCH Commun. Math. Comput. Chem.* 88 (2022) 505–520.
- [8] I. Tomescu, Graphs with given cyclomatic number extremal relatively to vertex degree function index for convex functions, *MATCH Commun. Math. Comput. Chem.* 87 (2022) 109–114.
- [9] I. Tomescu, Extremal vertex-degree function index for trees and unicyclic graphs with given independence number, *Discrete Appl. Math.* 306 (2022) 83–88.
- [10] X. Li, D. Peng, Extremal problems for graphical function-indices and f -weighted adjacency matrix, *Discrete Math. Lett.* 9 (2022) 57–66.
- [11] T. Došlić, T. Réti, D. Vukičević, On the vertex degree indices of connected graphs, *Chem. Phys. Lett.* 512 (2011) 283–286.
- [12] I. Gutman, On coindices of graphs and their complements, *Appl. Math. Comput.* 305 (2017) 161–165.
- [13] M. Matejić, I. Milovanović, E. Milovanović, Some remarks on the general zeroth-order Randić coindex, *Filomat*, 36 (6) (2022) 2083–2090.
- [14] X. Li, J. Zheng, A unified approach of the extremal trees for different indices, *MATCH Commun. Math. Comput. Chem.* 54 (2005) 195–208.
- [15] A. Milićević, S. Nikolić, On variable Zagreb indices, *Croat. Chem. Acta*, 27 (2004) 97–101.
- [16] A. Ali, I. Gutman, E. Milovanović, I. Milovanović, Sum of powers of the degrees of graphs: extremal results and bounds, *MATCH Commun. Math. Comput. Chem.* 80 (2018) 5–84.
- [17] A. Ali, D. Dimitrov, Z. Du, F. Ishfaq, On the extremal graphs for general sum-connectivity index (χ_α) with given cyclomatic number when $\alpha > 1$, *Discrete Appl. Math.* 257 (2019) 19–30.
- [18] P. Milošević, I. Milovanović, E. Milovanović, M. Matejić, Some inequalities for general zeroth-order Randić index, *Filomat*, 33 (16) (2019) 5249–5258.

- [19] T. Monsour, C. Song, The a and (a, b) -analogs of Zagreb indices and coindices of graphs, *Int. J. Comb.* 2012 (2012) art #909285
- [20] I. Milovanović, M. Matejić, E. Milovanović, A note on the general zeroth-order Randić coindex of graphs, *Contrib. Math.* 1 (2020) 17–21.
- [21] A. Ghalavand, A. R. Ashrafi, On forgotten coindex of chemical graphs, *MATCH Commun. Math. Comput. Chem.* 83 (2020) 221–232.
- [22] M. Azari, F. Falahati-Nezhad, Some results on forgotten topological coindex, *Iranian J. Math. Chem.* 10 (2019) 307–318.
- [23] T. Vetrík, S. Balachandran, General multiplicative Zagreb indices of trees, *Discrete Appl. Math.* 247 (2018) 341–351.
- [24] D. Vukičević, Bond additive modeling 5. mathematical properties of the variable sum exdeg index, *Croat. Chem. Acta* 84 (2011) 93–101.
- [25] A. Ali, D. Dimitrov, On the extremal graphs with respect to bond incident degree indices, *Discrete Appl. Math.* 238 (2018) 32–40.
- [26] Y. Hu, X. Li, Y. Shi, T. Xu, I. Gutman, On molecular graphs with smallest and greatest zeroth-order general Randić index, *MATCH Commun. Math. Comput. Chem.* 54 (2005) 425–434.