

CONTINUOUSLY MANY QUASI-ISOMETRY CLASSES OF RESIDUALLY FINITE GROUPS

HIP KUEN CHONG AND DANIEL T. WISE

ABSTRACT. We study a family of finitely generated residually finite small cancellation groups. These groups are quotients of F_2 depending on a subset S of positive integers. Varying S yields continuously many groups up to quasi-isometry.

1. INTRODUCTION

Grigorchuk exhibited continuously many quasiisometry classes of residually finite 3-generator groups by producing continuously many growth types [Gri84, Thm 7.2]. *Continuously many* means having the cardinality of \mathbb{R} . Here, we describe another family of such groups by building upon Bowditch’s method for distinguishing quasi-isometry classes [Bow98], and use consequences of the theory of special cube complexes to obtain residual finiteness [Ago13].

Consider the rank-2 free group $F_2 = \langle a, b \rangle$. Let $w_n = [a, b^{2^{2^n}}][a^2, b^{2^{2^n}}] \cdots [a^{100}, b^{2^{2^n}}]$ for $n \in \mathbb{N}$.

Each subset $S \subseteq \mathbb{N}$ is associated to the following group:

$$G(S) = \langle a, b \mid w_n : n \in S \rangle$$

In Section 3, we show that $G(S)$ is residually finite when $S \subseteq \mathbb{N}_{>100}$. We also observe that $G(S)$ and $G(S')$ are not quasi-isometric when $S \Delta S'$ is infinite.

In fact, our proof of residual finiteness for $G(S)$ works in precisely the same way to prove the residual finiteness for the original examples of Bowditch having torsion. But it appears to fail for Bowditch’s torsion-free examples. We refer to Remark 3.3.

Date: July 4, 2022.

2020 *Mathematics Subject Classification.* 20E26, 20F06.

Key words and phrases. Residually finite, small cancellation groups, quasi-isometry.

Research supported by NSERC.

We also produced an uncountable family of pairwise non-isomorphic residually finite groups in [CW21], and perhaps an appropriate subfamily also yields continuously many quasi-isometry classes.

Our simple approach arranges for certain infinitely presented small-cancellation groups to be residually finitely presented small-cancellation groups. This approach is likely to permit the construction of other interesting families of finitely generated groups.

2. REVIEW OF BOWDITCH'S RESULT

We first recall some small-cancellation background. See [LS01, Ch.V].

Definition 2.1. For a presentation, a *piece* p is a word appearing in more than one way among the relators. Note that for a relator $r = q^n$, subwords that differ by a \mathbb{Z}_n -action are regarded as appearing in the same way. A presentation is $C'(\frac{1}{6})$ if $|p| < \frac{1}{6}|r|$ whenever a piece p occurs in a relator r .

A *major subword* v of a relator r is a subword of a cyclic permutation of r^\pm with $|v| > \frac{|r|}{2}$. Any $g \in G$ is represented by a word u that is *majority-reduced* in the sense that u does not contain a major subword of a relator r . We will use the following proposition for $C'(\frac{1}{6})$ groups, which are groups given by a $C'(\frac{1}{6})$ presentation.

Proposition 2.2. Let G be a $C'(\frac{1}{6})$ group. If u is a nonempty majority-reduced word, then $u \neq 1_G$.

We now recall definitions leading to the main theorem of [Bow98]. Let $\mathbb{N}^+ = \{n \in \mathbb{Z} : n \geq 1\}$.

Definition 2.3. Two subsets $L, L' \subseteq \mathbb{N}^+$ are *related* if for some $k \geq 1$:

- (1) for any $m \in L$ with $m > (k+1)^2$, there is $m' \in L'$ with $m' \in [\frac{m}{k}, km]$; and
- (2) for any $m' \in L'$ with $m' > (k+1)^2$, there is $m \in L$ with $m \in [\frac{m'}{k}, km']$.

We write $L \sim L'$ if L and L' are related, and write $L \not\sim L'$ otherwise.

Lemma 2.4. The relation \sim in Definition 2.3 is an equivalence relation on subsets of \mathbb{N}^+ .

Proof. The relation \sim is reflexive by letting $k = 1$. The relation \sim is symmetric by definition. Hence it suffices to show \sim is transitive.

Let $S, S', S'' \subseteq \mathbb{N}^+$. Suppose $S \sim S'$ via k and $S' \sim S''$ via k' . We claim that $S \sim S''$ via kk' . Let $m \in S$. There is $m' \in S'$ with $m' \in [\frac{m}{k}, km]$ and there is $m'' \in S''$ with $m'' \in [\frac{m'}{k'}, k'm']$. Thus, $m'' \in [\frac{m}{kk'}, kk'm]$. Similarly, $m \in [\frac{m''}{kk'}, kk'm'']$ for $m'' \in S''$. \square

For sets S, S' , their *symmetric difference* is $S\Delta S' = (S - S') \cup (S' - S)$.

Example 2.5. If $S, S' \subseteq \mathbb{N}^+$ with infinite $S\Delta S'$, then $\{2^{2^n}\}_{n \in S} \not\sim \{2^{2^m}\}_{m \in S'}$ [Bow98, Lem 4].

With the notion of \sim , the following is a simplified version of the main theorem in [Bow98].

Theorem 2.6. Let G and G' be the finitely generated $C'(\frac{1}{6})$ groups presentated below.

If G is quasi-isometric to G' , then $\{|w_i|\}_{i \in I} \sim \{|w'_j|\}_{j \in J}$.

$$G = \langle A \mid w_i : i \in I \rangle, \quad G' = \langle A \mid w'_j : j \in J \rangle.$$

3. PROVING THE FAMILY OF GROUPS HAVE DESIRED PROPERTIES

3.1. Small Cancellation. Let $\mathbb{N}_{>100} = \{n \in \mathbb{N} : n > 100\}$.

Proposition 3.1. For any infinite subset $S \subseteq \mathbb{N}_{>100}$, the associated group G is $C'(\frac{1}{6})$.

Furthermore, $G_k = \langle a, b \mid b^{2^k}, w_n : n \in S, n < k \rangle$ is $C'(\frac{1}{6})$ for each $k \in \mathbb{N}$.

Proof. For the first statement, it suffices to show that w_n and w_m have small overlap for $n > m > 100$. The longest piece between w_n and w_m is $b^{-2^{2^m}} a^{100} b^{2^{2^m}}$. Thus, $C'(\frac{1}{6})$ holds since:

$$\left| b^{-2^{2^m}} a^{100} b^{2^{2^m}} \right| = 100 + 2 \cdot 2^{2^m} < \frac{1}{6} (10100 + 200 \cdot 2^{2^m}) = \frac{1}{6} |w_m| < \frac{1}{6} |w_n|$$

For the second statement, we additionally show that w_n and $b^{2^{2^k}}$ satisfy the $C'(\frac{1}{6})$ condition for $100 < n < k$. Their longest piece is $b^{2^{2^n}}$, which is shorter than $\frac{1}{6}$ of the lengths of w_n and $b^{2^{2^k}}$. \square

3.2. Residual Finiteness. Observe that $G_k = G / \langle\langle b^{2^{2^k}} \rangle\rangle$ since $w_m \in \langle\langle b^{2^{2^k}} \rangle\rangle$ for $m \geq k$.

Proposition 3.2. For any infinite subset $S \subseteq \mathbb{N}_{>100}$, the associated group G is residually finite.

Proof. Since G_k is a finitely presented $C'(\frac{1}{6})$ group, the hyperbolic group G_k is cocompactly cubulated by [Wis04]. Thus G_k is residually finite by [Ago13].

By Proposition 2.2, each $g \in G - \{1\}$ is represented by a majority-reduced word v , which does not contain a majority subword of $b^{2^{2^{|v|}}}$ since $|v| < \frac{1}{2} \cdot 2^{2^{|v|}} = \frac{1}{2} \left| b^{2^{2^{|v|}}} \right|$. Hence, $v \neq 1_{G_{|v|}}$. Thus, G is residually residually finite and hence residually finite. \square

Remark 3.3. Bowditch's original examples were $B(S) = \langle a, b \mid (a^{2^{2^n}} b^{2^{2^n}})^7 : n \in S \subseteq \mathbb{N} \rangle$. As in Proposition 3.2, $B(S)$ is residually finite since it is residually finitely presented $C'(\frac{1}{6})$ using the quotients to $B/\langle\langle a^{2^{2^n}}, b^{2^{2^n}} \rangle\rangle$ for $n \geq 3$. However, the analogous argument fails for Bowditch's torsion-free examples $B'(S) = \langle a, b \mid a(a^{2^{2^n}} b^{2^{2^n}})^{12} : n \in S \subseteq \mathbb{N} \rangle$.

3.3. Pairwise Non-quasi-isometric. We first prove a lemma about the relation \sim .

Lemma 3.4. $S \sim nS \sim (S+n)$ for $n \in \mathbb{N}^+$ and $S \subseteq \mathbb{N}^+$.

Proof. First, $S \sim nS$ via n . Indeed, for any $s \in S$, $ns \in [\frac{s}{n}, ns]$; for any $ns \in nS$, $s \in [\frac{ns}{n}, n \cdot ns]$.

Moreover, $S \sim (S+n)$ via $n+1$. For any $s \in S$, $s+n \leq (n+1)s$, so $s+n \in [\frac{s}{n+1}, (n+1)s]$. On the other hand, for $s+n \in S+n$, $(n+1)s \geq s+n$ implies $s \geq \frac{s+n}{n+1}$. Hence, $s \in [\frac{s+n}{n+1}, (n+1)(s+n)]$. \square

Proposition 3.5. Let $S, S' \subseteq \mathbb{N}^+$ have infinite $S\Delta S'$, then $\{|w_n|\}_{n \in S} \not\sim \{|w_m|\}_{m \in S'}$.

Proof. $\{|w_n| : n \in S\} = \{10100 + 200 \cdot 2^{2^n} : n \in S\} = 10100 + 200 \cdot \{2^{2^n} : n \in S\}$. By Lemma 3.4, $\{|w_n| : n \in S\} \sim \{2^{2^n} : n \in S\}$. Similarly, $\{|w_m| : m \in S'\} \sim \{2^{2^m} : m \in S'\}$. By Example 2.5, $\{2^{2^n} : n \in S\} \not\sim \{2^{2^m} : m \in S'\}$, so $\{|w_n|\}_{n \in S} \not\sim \{|w_m|\}_{m \in S'}$. \square

Corollary 3.6. If $S, S' \subseteq \mathbb{N}_{>100}$ have infinite $S\Delta S'$, then $G(S)$ and $G(S')$ are not quasi-isometric.

Proof. $\{|w_n|\}_{n \in S} \not\sim \{|w_m|\}_{m \in S'}$, hence $G(S)$ and $G(S')$ are not quasi-isometric by Theorem 2.6. \square

For $A, B \subseteq \mathbb{N}$, declare $A \sim_{\Delta} B$ if $|A\Delta B| < \infty$. As noted by Bowditch, each \sim_{Δ} equivalence class is countable. Hence, there are continuously many \sim_{Δ} equivalence classes. Our construction thus produces continuously many pairwise non-quasi-isometric groups $G(S)$, which are $C'(\frac{1}{6})$ and residually finite.

REFERENCES

- [Ago13] Ian Agol. The virtual Haken conjecture. *Doc. Math.*, 18:1045–1087, 2013. With an appendix by Agol, Daniel Groves, and Jason Manning.
- [Bow98] B. H. Bowditch. Continuously many quasi-isometry classes of 2-generator groups. *Comment. Math. Helv.*, 73(2):232–236, 1998.
- [CW21] Hip Kuen Chong and Daniel T. Wise. An uncountable family of finitely generated residually finite groups. *Journal of Group Theory*, pages 1–10, 2021. In Press.

- [Gri84] R. I. Grigorchuk. Degrees of growth of finitely generated groups and the theory of invariant means. *Izv. Akad. Nauk SSSR Ser. Mat.*, 48(5):939–985, 1984.
- [LS01] Roger C. Lyndon and Paul E. Schupp. *Combinatorial group theory*. Classics in Mathematics. Springer-Verlag, Berlin, 2001. Reprint of the 1977 edition.
- [Wis04] D. T. Wise. Cubulating small cancellation groups. *Geom. Funct. Anal.*, 14(1):150–214, 2004.

DEPT. OF MATH. & STATS., MCGILL UNIV., MONTREAL, QC, CANADA H3A 0B9

Email address: `chonghk1997@gmail.com`

Email address: `wise@math.mcgill.ca`