

# On Stein’s method for stochastically monotone single-birth chains

Fraser Daly\*

July 4, 2022

**Abstract** We discuss Stein’s method for approximation by the stationary distribution of a single-birth Markov chain, in conjunction with stochastic monotonicity and similar assumptions. We use bounds on the increments of the solution of Poisson’s equation for such a process. Our first applications are to rates of convergence to stationarity. In our second set of applications we bound the total variation distance between the stationary distributions of two Markov chains, including quantifying the effect of truncation of the state space.

**Key words and phrases:** Markov chain; Poisson’s equation; stochastic monotonicity; truncation; total variation distance

**MSC 2020 subject classification:** 60J10; 60E15; 60F05; 62E17

## 1 Introduction

Let  $\{Z_t : t = 0, 1, \dots\}$  be a positive recurrent single-birth Markov chain in discrete time on the non-negative integers, with transition matrix  $P$  whose  $(i, j)$ th entry we denote by  $P_{i,j}$ . Throughout we assume that  $P_{i,i+1} > 0$  for each  $i \in \mathbb{Z}^+ = \{0, 1, 2, \dots\}$ , and that  $P_{i,j} = 0$  for  $j > i + 1$ . We let this chain have stationary distribution denoted by  $(\pi_0, \pi_1, \pi_2 \dots)$ , and  $\pi$  be a random variable with this stationary distribution.

For a given function  $h : \mathbb{Z}^+ \rightarrow \mathbb{R}$ , we let  $f : \mathbb{Z}^+ \rightarrow \mathbb{R}$  denote the solution to Poisson’s equation

$$h(i) - \mathbb{E}h(\pi) = f(i) - \sum_{j=0}^{\infty} P_{i,j}f(j), \quad (1)$$

with  $f(0) = 0$ . This equation can be used in conjunction with Stein’s method to find explicit bounds in approximation by the stationary distribution of our single-birth chain. We will introduce the elements of Stein’s method that we need in our work, but refer the reader to [12] and references therein for an introduction to this technique.

In the remainder of this section we will derive Theorem 1 below, which gives an explicit bound in approximation by our stationary distribution, before exploring applications of this

---

\*Department of Actuarial Mathematics and Statistics and the Maxwell Institute for Mathematical Sciences, Heriot-Watt University, Edinburgh EH14 4AS, UK. E-mail: f.daly@hw.ac.uk

bound in Sections 2 and 3. In many of these applications it will be convenient to exploit stochastic monotonicity of the transition matrix  $P$  or another similar monotonicity property. Recall that  $P$  is defined to be stochastically monotone if the distributions in successive rows of  $P$  are stochastically non-decreasing; in this case stochastic ordering is preserved under transitions taken according to  $P$ . See [3] for further details. For random variables  $X$  and  $Y$ , we say that  $Y$  is stochastically larger than  $X$  if  $\mathbb{P}(Y > j) \geq \mathbb{P}(X > j)$  for all  $j$ , and denote this by  $Y \geq_{st} X$ .

One central aim of our work is to demonstrate how assumptions of stochastic monotonicity can be used in deriving bounds in approximations by the distribution of  $\pi$  by Stein's method, in a similar spirit to [4]. This is somewhat different to earlier work on Stein's method for stationary distributions of Markov processes, including that of Brown and Xia [2], who considered approximation by the stationary distribution of a continuous-time birth-death process without any assumptions of monotonicity, and whose applications are of a quite different flavour to ours. In the discrete-time setting, [1, 11] have recently used Poisson's equation as a starting point for Stein's method, but in the case of a finite state space and again without monotonicity assumptions. The focus of our work is thus somewhat different to these other papers. While there are many other tools available in the literature for tackling examples and applications such as those we consider here (see the discussion and references in the examples below), our purpose here is to show how Stein's method can be added to such a toolkit.

Letting  $\widehat{h}(j) = h(j) - \mathbb{E}h(\pi)$ , Jiang *et al.* [6] have shown that Poisson's equation (1) is solved by a function  $f$  satisfying  $f(j+1) - f(j) = -m_j(h)$  for  $j \geq 0$ , where  $m_0(h) = \widehat{h}(0)/P_{0,1}$  and

$$m_j(h) = \frac{1}{P_{j,j+1}} \left( \widehat{h}(j) + \sum_{k=0}^{j-1} m_k(h) \sum_{l=0}^k P_{j,l} \right),$$

for  $j \geq 1$ ; see their Theorem 2.1. Our approximation results will make use of bounds on  $\sup_{j \in \mathbb{Z}^+} |m_j(h)|$  for  $h \in \mathcal{H} = \{h : \mathbb{Z}^+ \rightarrow \mathbb{R} : |h(j)| \leq 1 \text{ for all } j\}$ . This is the relevant class of functions for us since we derive bounds on the total variation distance between  $\pi$  and the random variable  $X_t$ , where  $\{X_t : t = 0, 1, \dots\}$  is another Markov chain on  $\mathbb{Z}^+$ . This total variation distance is defined by

$$d_{TV}(X_t, \pi) = \sup_{h \in \mathcal{H}} |\mathbb{E}h(X_t) - \mathbb{E}h(\pi)|.$$

Following Stein's method, this will be bounded by rewriting the right-hand side using Poisson's equation (1):

$$\begin{aligned} \mathbb{E}h(X_t) - \mathbb{E}h(\pi) &= \mathbb{E}f(X_t) - \mathbb{E} \sum_{j=0}^{\infty} P_{X_t,j} f(j) \\ &= \sum_{j=0}^{\infty} f(j) \left[ \mathbb{P}(X_t = j) - \sum_{i=0}^{\infty} \mathbb{P}(X_t = i) \mathbb{P}(Z_1 = j | Z_0 = i) \right] \\ &= \sum_{j=0}^{\infty} \Delta f(j) \left[ \mathbb{P}(X_t > j) - \sum_{k=j+1}^{\infty} \sum_{i=0}^{\infty} \mathbb{P}(X_t = i) \mathbb{P}(Z_1 = k | Z_0 = i) \right], \end{aligned}$$

where we recall that  $f(0) = 0$  and write  $\Delta f(j) = f(j+1) - f(j)$ . This immediately yields the following result.

**Theorem 1.** Let  $\{Z_t : t = 0, 1, \dots\}$  be a positive recurrent single-birth Markov chain on  $\mathbb{Z}^+$  with transition matrix  $P$  and stationary distribution  $(\pi_0, \pi_1, \dots)$ . Let  $\{X_t : t = 0, 1, \dots\}$  be a Markov chain on  $\mathbb{Z}^+$ . Then

$$d_{TV}(X_t, \pi) \leq \sup_{h \in \mathcal{H}} \sup_{l \in \mathbb{Z}^+} |m_l(h)| \sum_{j=0}^{\infty} \left| \mathbb{P}(X_t > j) - \sum_{k=j+1}^{\infty} \sum_{i=0}^{\infty} \mathbb{P}(X_t = i) \mathbb{P}(Z_1 = k | Z_0 = i) \right|.$$

Note that this result makes use of bounds on the increments of the solution  $f$  of Poisson's equation, not bounds on  $f$  itself. We conclude this section by noting one case in which a bound on the increments of  $f$  may be easily found. Suppose that  $\{Z_t : t = 0, 1, \dots\}$  is a birth-death chain with  $P_{i,i+1} = b_i > 0$  for each  $i = 0, 1, \dots$  and  $P_{i,i-j} = 0$  for each  $j > 1$  and all  $i$ . In this case, Jiang *et al.* [6] have shown that

$$m_j(h) = \frac{1}{b_j \pi_j} \sum_{k=0}^j \widehat{h}(k) \pi_k = -\frac{1}{b_j \pi_j} \sum_{k=j+1}^{\infty} \widehat{h}(k) \pi_k;$$

see their equation (15), and note that the final equality follows from  $\mathbb{E} \widehat{h}(\pi) = 0$ . This immediately gives the bound

$$\sup_{h \in \mathcal{H}} \sup_{j \in \mathbb{Z}^+} |m_j(h)| \leq \sup_{j \in \mathbb{Z}^+} \left\{ \frac{\mathbb{P}(\pi > j)}{b_j \pi_j} \right\}. \quad (2)$$

In the remainder of this note we will explore some applications of Theorem 1 for a variety of choices of the Markov chain  $\{X_t : t = 0, 1, \dots\}$ , and see how we may make use of stochastic monotonicity in such applications. In Section 2 we will take  $\{X_t : t = 0, 1, \dots\}$  and  $\{Z_t : t = 0, 1, \dots\}$  identical in distribution, so that Theorem 1 gives a bound on the distance of  $Z_t$  to stationarity. This is illustrated with three simple examples. In Section 3 we will let  $\{X_t : t = 0, 1, \dots\}$  be a stationary Markov chain, so that Theorem 1 bounds the distance between two stationary distributions. Applications of this include quantifying the effect of truncation of the state space of our original chain. We emphasise again that there are many tools available to treat applications such as these, we use them here mainly to illustrate the use of Theorem 1 as another such tool.

## 2 Convergence to stationarity

In the setting of Theorem 1, we choose the Markov chain  $\{X_t : t = 0, 1, \dots\}$  to be  $\{Z'_t : t = 0, 1, \dots\}$ , a single-birth process evolving according to the transition matrix  $P$  and such that  $Z'_t$  has the same distribution as  $Z_t$  for each  $t$ . This lets us bound the total variation distance of  $Z_t$  from stationarity. We set  $Z_0 = Z'_0 = 0$ , and note that

$$\sum_{k=j+1}^{\infty} \sum_{i=0}^{\infty} \mathbb{P}(Z'_t = i) \mathbb{P}(Z_1 = k | Z_0 = i) = \sum_{k=j+1}^{\infty} \mathbb{P}(Z'_{t+1} = k) = \mathbb{P}(Z'_{t+1} > j).$$

Under the assumption that  $P$  is stochastically monotone, we can couple our Markov chains in such a way that  $Z'_{t+1} \geq_{st} Z_t$ , i.e., such that  $\mathbb{P}(Z'_{t+1} > j) \geq \mathbb{P}(Z_t > j)$  for all  $j \in \mathbb{Z}^+$ . Hence,

$$\sum_{j=0}^{\infty} |\mathbb{P}(Z_t > j) - \mathbb{P}(Z'_{t+1} > j)| = \mathbb{E} Z'_{t+1} - \mathbb{E} Z_t,$$

and Theorem 1 gives the following, in which we may choose a coupling of  $Z_t$  and  $Z'_{t+1}$  to bound the expectation.

**Corollary 2.** *Let  $\{Z_t : t = 0, 1, \dots\}$  be a positive recurrent and stochastically monotone single-birth Markov chain on  $\mathbb{Z}^+$ , as defined above, with transition matrix  $P$  and stationary distribution  $(\pi_0, \pi_1, \dots)$ . Then*

$$d_{TV}(Z_t, \pi) \leq \sup_{h \in \mathcal{H}} \sup_{l \in \mathbb{Z}^+} |m_l(h)| (\mathbb{E}Z'_{t+1} - \mathbb{E}Z_t).$$

We use the remainder of this section to illustrate the bound of Corollary 2 with three applications.

## 2.1 Example: Simple random walk with reflection

Consider a simple random walk on  $\mathbb{Z}^+$  with reflection at the origin, where  $\mathbb{P}(Z_1 = 0 | Z_0 = 0) = p$ ,  $\mathbb{P}(Z_1 = i - 1 | Z_0 = i) = p$  for all  $i \geq 1$ , and  $\mathbb{P}(Z_1 = i + 1 | Z_0 = i) = 1 - p$  for all  $i \geq 0$ , for some  $p > 1/2$  to ensure positive recurrence. This example was treated by Lund and Tweedie [9], we use it here to confirm that Corollary 2 gives the rate of convergence we would expect. We note that in this case  $P$  is stochastically monotone, and that  $\pi$  is geometrically distributed with parameter  $\alpha = \frac{2p-1}{p}$  and mass function  $\pi_k = \alpha(1 - \alpha)^k$  for  $k = 0, 1, \dots$

We couple our Markov chains as follows: with  $Z_0 = Z'_0 = 0$ , we let  $Z'_1$  be 0 or 1, with probability  $p$  and  $1 - p$  respectively, independently of all else. The processes  $\{Z_t : t = 1, 2, \dots\}$  and  $\{Z'_{t+1} : t = 1, 2, \dots\}$  then evolve using the same underlying sequence of independent Bernoulli trials so that, roughly speaking, one process moves in the positive direction at a given time if and only if the other process does also; the same is true of steps in the negative direction, except that we need to account for the reflection at the origin where a ‘step in the negative direction’ corresponds to remaining at the origin. This continues until the first time  $t$  at which  $Z_t = Z'_{t+1}$ , following which the two processes move together. This happens at a time which is almost surely no greater than  $T = \min\{t \geq 1 : Z'_t = 0\}$ , the first return time to the origin, and we note that

$$\mathbb{E}Z'_{t+1} - \mathbb{E}Z_t \leq \mathbb{P}(T > t + 1) = \mathbb{P}(r^T \geq r^{t+2}) \leq \frac{\mathbb{E}r^T}{r^{t+2}} = \frac{2pr + 1 - [1 - 4p(1 - p)r^2]^{1/2}}{2r^{t+2}},$$

for any  $1 \leq r \leq [4p(1 - p)]^{-1/2}$ ; see Section XIV.4 of [5] for the final equality. With the observation that

$$\frac{\mathbb{P}(\pi > l)}{b_l \pi_l} = \frac{1 - \alpha}{\alpha(1 - p)} = \frac{1}{2p - 1},$$

Corollary 2 and (2) give us that, for any  $1 \leq r \leq [4p(1 - p)]^{-1/2}$ ,

$$d_{TV}(Z_t, \pi) \leq \left( \frac{2pr + 1 - [1 - 4p(1 - p)r^2]^{1/2}}{2r^2(2p - 1)} \right) r^{-t},$$

yielding the expected geometric rate of convergence to stationarity (see also Example 7.1 of [9]) and an explicit bound on the corresponding total variation distance.

## 2.2 Example: M/M/1 queue

For our next example, we look beyond the class of birth-death processes into Markov chains of the type associated with GI/M/1 queues. For simplicity we will restrict our attention to the M/M/1 queue here. Although this may be formulated as a birth-death process, we instead use an alternative representation which makes it clear how this example can be extended to more general GI/M/1-type chains. Consider the M/M/1 queue in which customer interarrival times are exponentially distributed with mean  $1/\lambda$  and service times are exponentially distributed with mean  $1/\mu$ , where we assume  $\rho = \lambda/\mu < 1$ . Let  $\{Z_t : t = 0, 1, \dots\}$  be the Markov chain embedded at customer arrival times, which may be constructed as a stochastically monotone single-birth chain with stationary distribution  $\pi_i = (1 - \rho)\rho^i$  for  $i = 0, 1, \dots$ . We define

$$a_k = \frac{\rho}{1 + \rho} \left( \frac{1}{1 + \rho} \right)^k,$$

the probability that exactly  $k$  customers are served in the time between two consecutive arrivals. We then define our transition matrix  $P$  by writing  $P_{i,j} = a_{i-j+1}$  for  $j = 1, 2, \dots, i + 1$  and each  $i$ . The remaining non-zero entries of  $P$  are those in the left-hand column, which are given by  $P_{i,0} = \sum_{j>i} a_j = (1 + \rho)^{-(i+1)}$ , for  $i \geq 0$ . Note that, as constructed, this is not a birth-death chain, and so we cannot use the bound (2) here. We will need to calculate separately a bound on the increments  $m_i(h)$  of the solution of (1). These are given by  $m_0(h) = (1 + \rho)\rho^{-1}\widehat{h}(0)$  and

$$m_i(h) = \frac{1 + \rho}{\rho} \left( \widehat{h}(i) + \frac{1}{(1 + \rho)^{i+1}} \sum_{k=0}^{i-1} (1 + \rho)^k m_k(h) \right),$$

for  $i = 1, 2, \dots$ . Solving this system of equations gives

$$m_i(h) = \left( \frac{1 + \rho}{\rho} \right) \widehat{h}(i) + \frac{1}{\rho^{i+1}} \sum_{j=0}^{i-1} \widehat{h}(j) \rho^j = \left( \frac{1 + \rho}{\rho} \right) \widehat{h}(i) - \frac{1}{\rho^{i+1}} \sum_{j=i}^{\infty} \widehat{h}(j) \rho^j,$$

where the final equality uses the fact that  $\mathbb{E}\widehat{h}(\pi) = 0$ . For  $h \in \mathcal{H}$  we therefore have

$$|m_i(h)| \leq \frac{1 + \rho}{\rho} + \frac{1}{\rho^{i+1}} \sum_{j=i}^{\infty} \rho^j = \frac{2 - \rho^2}{\rho(1 - \rho)}.$$

To apply Corollary 2 it thus remains only to couple  $Z'_{t+1}$  and  $Z_t$ . We proceed similarly to Section 2.1. With  $Z_0 = Z'_0 = 0$ , let  $Z'_1$  be 0 or 1, with probability  $(1 + \rho)^{-1}$  and  $\rho(1 + \rho)^{-1}$  respectively, independently of all else. We let subsequent arrivals occur at the same times in both processes, and hence  $Z_t \leq Z'_{t+1}$  almost surely for all  $t$ . As before,  $\mathbb{E}Z'_{t+1} - \mathbb{E}Z_t \leq \mathbb{P}(T > t + 1)$ , where  $T$  is as in Section 2.1, since the numbers of customers in the two systems differ by at most one. We may write  $T$  as  $1 + IN$ , where  $N$  is the number of customers served in a busy period of an M/M/1 queue initiated by the arrival of a single customer to an empty system, and  $I$  is a Bernoulli random variable with mean  $\rho(1 + \rho)^{-1}$  independent of all else. We then have

$$\mathbb{P}(T > t + 1) = \frac{\rho}{1 + \rho} \mathbb{P}(N > t) \leq \frac{\rho \mathbb{E}r^N}{(1 + \rho)r^{t+1}} = \frac{1 + \rho - [(1 + \rho)^2 - 4\rho r]^{1/2}}{2(1 + \rho)r^{t+1}},$$

for all  $1 \leq r \leq \frac{(1+\rho)^2}{4\rho}$ ; the expression for  $\mathbb{E}r^N$  is well-known. Corollary 2 thus gives us, for any  $1 \leq r \leq \frac{(1+\rho)^2}{4\rho}$ ,

$$d_{TV}(Z_t, \pi) \leq \left( \frac{(2 - \rho^2)(1 + \rho - [(1 + \rho)^2 - 4\rho r]^{1/2})}{2\rho(1 - \rho)(1 + \rho)r} \right) r^{-t}.$$

### 2.3 Example: Discrete renewal process with decreasing failure rate

Let  $0 < \alpha_0 \leq \alpha_1 \leq \dots < 1$  and define a stochastically monotone transition matrix  $P$  such that  $P_{i,i+1} = \alpha_i$  and  $P_{i,0} = 1 - \alpha_i$  for all  $i = 0, 1, \dots$ , and all other entries are zero. The corresponding Markov chain is the age of a discrete renewal process with decreasing failure rate. The stationary distribution is given by

$$\pi_k = \frac{\alpha_0 \cdots \alpha_{k-1}}{1 + \sum_{j=1}^{\infty} \prod_{i=0}^{j-1} \alpha_i},$$

for  $k = 0, 1, \dots$ , where we assume that the denominator is finite so that our Markov chain is positive recurrent.

With  $f(0) = 0$ , Poisson's equation (1) becomes  $h(i) - \mathbb{E}h(\pi) = f(i) - \alpha_i f(i+1)$ . If the  $\alpha_i$  were all equal, this is an equation which is well known in the setting of geometric approximation; see [10]. The solution is

$$f(j) = -\frac{1}{\pi_j} \sum_{k=0}^{j-1} \pi_k \widehat{h}(k) = \frac{1}{\pi_j} \sum_{k=j}^{\infty} \pi_k \widehat{h}(k),$$

where the final equality follows from  $\mathbb{E}\widehat{h}(\pi) = 0$ . We thus have that, for  $h \in \mathcal{H}$ ,

$$|f(j)| \leq \frac{\sum_{k=j}^{\infty} \alpha_0 \cdots \alpha_{k-1}}{\alpha_0 \cdots \alpha_{j-1}} = 1 + \alpha_j + \alpha_j \alpha_{j+1} + \dots \leq 1 + \beta + \beta^2 + \dots = \frac{1}{1 - \beta},$$

where  $\beta = \lim_{j \rightarrow \infty} \alpha_j$ . This gives  $|\Delta f(j)| \leq 2(1 - \beta)^{-1}$ .

We construct a coupling in a similar way to previous examples. Set  $Z_0 = Z'_0 = 0$ , and let  $Z'_1$  be 0 or 1, with probability  $1 - \alpha_0$  and  $\alpha_0$  respectively, independently of all else. Let  $T$  be as in Section 2.1. Conditional on the event that  $Z'_{t+1} = n$  and  $Z_t = m$  with  $m \leq n$ , choose one of three possible increments for these processes at the next time step, independently of all else: (i) both move one step in the positive direction with probability  $\alpha_m$ , (ii) both move to zero with probability  $1 - \alpha_n$ , or (iii)  $Z'_{t+2} = Z'_{t+1} + 1$  and  $Z_{t+1} = 0$  with probability  $\alpha_n - \alpha_m$ . We thus have  $\mathbb{E}Z'_{t+1} - \mathbb{E}Z_t \leq (t+1)\mathbb{P}(T > t+1)$ . Since  $\mathbb{P}(T > t) = \alpha_0 \cdots \alpha_{t-1}$ , Corollary 2 gives the subgeometric bound

$$d_{TV}(Z_t, \pi) \leq \frac{2(t+1)}{1 - \beta} \alpha_0 \cdots \alpha_t.$$

## 3 Distance between stationary distributions

In this section we let  $\{X_t : t = 0, 1, \dots\}$  be a positive recurrent Markov chain on  $\mathbb{Z}^+$  started according to its stationary distribution, which we denote by  $(\eta_0, \eta_1, \dots)$ . We let  $\eta$  denote a

random variable with this stationary distribution and bound  $d_{TV}(\eta, \pi)$ . We will focus on the case where our underlying Markov chain  $\{Z_t : t = 0, 1, \dots\}$  is a birth-death chain with  $P_{i,i+1} = b_i > 0$  and  $P_{i,i-j} = 0$  for each  $j > 1$  and all  $i$ , since this allows relatively simple approximating distributions and more explicit error bounds.

Let  $X_{(t+1)}$  denote the random variable obtained from  $X_t$  by taking a further step according to  $P$ . That is, conditional on the event that  $X_t = i$ , we have that  $X_{(t+1)} = j$  with probability given by  $P_{i,j}$ . With this notation,

$$\sum_{k=j+1}^{\infty} \sum_{i=0}^{\infty} \mathbb{P}(X_t = i) \mathbb{P}(Z_1 = k | Z_0 = i) = \mathbb{P}(X_{(t+1)} > j).$$

We also note that, since  $X_0$  is distributed as  $\eta$ , we have  $\mathbb{P}(X_t > j) = \mathbb{P}(X_{(t+1)} > j)$ . If we assume that either  $X_{(t+1)} \geq_{st} X_{t+1}$  or  $X_{t+1} \geq_{st} X_{(t+1)}$ , Theorem 1 and (2) then give us

**Corollary 3.** *Let  $\{Z_t : t = 0, 1, \dots\}$  be a positive recurrent birth-death Markov chain on  $\mathbb{Z}^+$  with transition matrix  $P$  and stationary distribution  $(\pi_0, \pi_1, \dots)$ . Let  $\{X_t : t = 0, 1, \dots\}$  be a positive recurrent Markov chain on  $\mathbb{Z}^+$  started according to its stationary distribution  $(\eta_0, \eta_1, \dots)$ , and  $X_{(t+1)}$  be as above. If  $X_{(t+1)} \geq_{st} X_{t+1}$  or  $X_{t+1} \geq_{st} X_{(t+1)}$ ,*

$$d_{TV}(\eta, \pi) \leq \sup_{l \in \mathbb{Z}^+} \left\{ \frac{\mathbb{P}(\pi > l)}{b_l \pi_l} \right\} |\mathbb{E}X_{t+1} - \mathbb{E}X_{(t+1)}|.$$

### 3.1 Example: Single-birth Markov chain

Let  $\{X_t : t = 0, 1, \dots\}$  be a single-birth chain, i.e.,  $\mathbb{P}(X_1 = j | X_0 = i) = 0$  for  $j > i + 1$ . See Corollary 3.4 of [7] for conditions under which such a chain is positive recurrent. We define the transition matrix  $P$  of our birth-death chain by letting  $P_{i,i-1} = \mathbb{P}(X_1 < i | X_0 = i)$  for each  $i \geq 1$ , and  $P_{i,i} = \mathbb{P}(X_1 = i | X_0 = i)$  and  $P_{i,i+1} = \mathbb{P}(X_1 = i + 1 | X_0 = i)$  for each  $i \geq 0$ . Thus, taking a step according to  $P$  results in a stochastically larger distribution than taking a step according to the original transition matrix of  $\{X_t : t = 0, 1, \dots\}$ . That is,  $X_{(t+1)} \geq_{st} X_{t+1}$ .

We now couple  $X_{t+1}$  and  $X_{(t+1)}$ . Conditional on the event that  $X_t = i \in \mathbb{Z}^+$ , we construct  $X_{t+1}$  by taking a step according to the transition matrix for this chain, and then take  $X_{(t+1)} = X_{t+1}$  if  $X_{t+1} \geq i - 1$ , and  $X_{(t+1)} = i - 1$  otherwise. We then have  $(X_{(t+1)} - X_{t+1} | X_t = i) = 0$  with probability  $\mathbb{P}(X_1 \geq i - 1 | X_0 = i)$ , and  $(X_{(t+1)} - X_{t+1} | X_t = i) = k$  with probability  $\mathbb{P}(X_1 = i - k - 1 | X_0 = i)$ , for each  $k = 1, 2, \dots$ . Hence,

$$\mathbb{E}X_{(t+1)} - \mathbb{E}X_{t+1} = \sum_{i=2}^{\infty} \mathbb{P}(X_t = i) \sum_{k=1}^{i-1} k \mathbb{P}(X_1 = i - k - 1 | X_0 = i) \leq \sum_{k=1}^{\infty} k \beta(k),$$

where  $\beta(k) = \sup_{i \geq 2} \{\mathbb{P}(X_1 = i - k - 1 | X_0 = i)\}$ . Corollary 3 thus gives the bound

$$d_{TV}(\eta, \pi) \leq \sup_{l \in \mathbb{Z}^+} \left\{ \frac{\mathbb{P}(\pi > l)}{\pi_l \mathbb{P}(X_1 = l + 1 | X_0 = l)} \right\} \sum_{k=1}^{\infty} k \beta(k).$$

This is useful in the setting of a single-birth chain whose transition matrix is close to tridiagonal, for which the stationary distribution of the approximating birth-death chain is simpler to compute than that of the original chain. An analogous bound can also be found in the setting where  $\{X_t : t = 0, 1, \dots\}$  is a single-death chain.

### 3.2 Example: Truncation

Let  ${}_{(n)}\tilde{P}$  denote the  $(n+1) \times (n+1)$  northwest truncation of  $P$ , augmented by replacing the  $j$ th entry in the final row by  $\mathbb{P}(Z_1 = j|Z_0 = n) + \nu_j\mathbb{P}(Z_1 = n+1|Z_0 = n)$  for each  $j \geq 0$ , where  $(\nu_0, \nu_1, \dots, \nu_n)$  is a probability distribution. We let  $\{X_t : t = 0, 1, \dots\}$  be the stationary Markov chain with transition matrix  ${}_{(n)}\tilde{P}$ ; this is a special case of the truncation problem for discrete-time Markov chains studied by many authors. Tweedie [13] shows that the corresponding stationary probabilities converge to  $\pi_j$  in the case of a geometrically ergodic chain, a stochastically monotone chain or one dominated by a stochastically monotone chain, when the augmentation is in the first or last column only. These results have since been generalised (see, for example, [7, 8] and references therein) and given improved error bounds. Our purpose here is not to compete with these general bounds, but to use the special case of a birth-death chain to further illustrate Corollary 3 and demonstrate that it yields reasonable bounds.

We construct  $X_{(t+1)}$  as follows: If  $X_t < n$ , the transition rules specified by  $P$  and  ${}_{(n)}\tilde{P}$  coincide, and so we set  $X_{(t+1)} = X_{t+1}$ . If  $X_t = n$ , then we set  $X_{(t+1)} = X_{t+1}$  with probability  $1 - b_n$  and we set  $X_{(t+1)} = X_{t+1} + j$  with probability  $\nu_{n+1-j}b_n$ . With this construction, it is clear that  $X_{(t+1)} \geq_{st} X_{t+1}$  and

$$\mathbb{E}X_{(t+1)} - \mathbb{E}X_{t+1} = b_n\mathbb{P}(X_0 = n) \sum_{j=1}^{n+1} j\nu_{n+1-j} = b_n\mathbb{P}(X_0 = n) \left( n+1 - \sum_{j=1}^n j\nu_j \right),$$

where we recall that the chain  $\{X_t : t = 0, 1, \dots\}$  is started according to its stationary distribution. Letting  ${}_{(n)}\pi$  denote a random variable with this stationary distribution, Corollary 3 then gives

$$d_{TV}({}_{(n)}\pi, \pi) \leq \sup_{l \in \mathbb{Z}^+} \left\{ \frac{\mathbb{P}(\pi > l)}{b_l\pi_l} \right\} b_n\mathbb{P}(X_0 = n) \left( n+1 - \sum_{j=1}^n j\nu_j \right). \quad (3)$$

For example, suppose that  $P$  is a stochastically monotone matrix with  $\pi_k = \alpha(1-\alpha)^k$  for some  $\alpha \in (0, 1)$  and all  $k = 0, 1, \dots$ . The simple random walk with reflection of Section 2.1 is an example of such a chain. Since  $\{X_t : t = 0, 1, \dots\}$  and  $\{Z_t : t = 0, 1, \dots\}$  run together until they hit the state  $n$  and  $\{Z_t : t = 0, 1, \dots\}$  subsequently moves to  $n+1$  while  $\{X_t : t = 0, 1, \dots\}$  remains at or below state  $n$ , stochastic monotonicity of  $P$  gives us that  $\mathbb{P}(X_0 = n) \leq \mathbb{P}(\pi \geq n) = (1-\alpha)^n$ , and (3) becomes

$$d_{TV}({}_{(n)}\pi, \pi) \leq \frac{b_n}{\alpha \inf_{l \in \mathbb{Z}^+} \{b_l\}} (1-\alpha)^{n+1},$$

giving the same rate as the lower bound  $d_{TV}({}_{(n)}\pi, \pi) \geq \mathbb{P}(\pi > n) = (1-\alpha)^{n+1}$  if the  $b_l$  are bounded away from zero.

## References

- [1] G. Bresler and D. Nagaraj (2019). Stein’s method for stationary distributions of Markov chains and application to Ising models. *Ann. Appl. Probab.* 29(5): 3230–3265.

- [2] T. C. Brown and A. Xia (2001). Stein’s method and birth-death processes. *Ann. Probab.* 29(3): 1373–1403.
- [3] D. J. Daley (1968). Stochastically monotone Markov chains. *Z. Wahrscheinlichkeitstheorie verw. Geb.* 10: 305–317.
- [4] F. Daly, C. Lefèvre and S. Utev (2012). Stein’s method and stochastic orderings. *Adv. in Appl. Probab.* 44(2): 343–372.
- [5] W. Feller (1960). *An Introduction to Probability Theory and Its Applications* Vol I, 2nd Ed. John Wiley and Sons, New York.
- [6] S. Jiang, Y. Liu and S. Yao (2014). Poisson’s equation for discrete-time single-birth processes. *Statist. Probab. Lett.* 85: 78–83.
- [7] Y. Liu (2010). Augmented truncation approximations of discrete-time Markov chains. *Oper. Res. Lett.* 38: 218–222.
- [8] Y. Liu and W. Li (2018). Error bounds for augmented truncation approximations of Markov chains via the perturbation method. *Adv. in Appl. Probab.* 50: 645–669.
- [9] R. B. Lund and R. L. Tweedie (1996). Geometric convergence rates for stochastically ordered Markov chains. *Math. Oper. Res.* 21(1): 182–194.
- [10] E. Peköz (1996). Stein’s method for geometric approximation. *J. Appl. Probab.* 33: 707–713.
- [11] G. Reinert and N. Ross (2019). Approximating stationary distributions of fast mixing Glauber dynamics, with applications to exponential random graphs. *Ann. Appl. Probab.* 29(5): 3201–3229.
- [12] N. Ross (2011). Fundamentals of Stein’s method. *Probab. Surv.* 8: 2310–293.
- [13] R. L. Tweedie (1998). Truncation approximations of invariant measures for Markov chains. *J. Appl. Probab.* 35: 517–536.