

Designing Tractable Piecewise Affine Policies for Multi-Stage Adjustable Robust Optimization

Simon Thomä*, Grit Walther*, Maximilian Schiffer†
simon.thomae@om.rwth-aachen.de

July 4, 2022

Abstract

We study piecewise affine policies for multi-stage adjustable robust optimization (ARO) problems with non-negative right-hand side uncertainty. First, we construct new dominating uncertainty sets and show how a multi-stage ARO problem can be solved efficiently when uncertainty is replaced by these new sets. We then demonstrate how solutions for this alternative problem can be transformed to solutions for the original problem. By carefully choosing the dominating sets, we prove strong approximation bounds for our policies and extend many previously best known bounds for the two-staged problem variant to its multi-stage counterpart. In two numerical experiments we identify beneficial and disadvantageous properties for our policies and present effective adjustments to tackle the most critical disadvantages. Overall, the experiments show that our piecewise affine policies can be computed by orders of magnitude faster than affine policies, while often yielding comparable or even better results.

1 Introduction

In practice, most decision-making problems have to be solved in view of uncertain parameters. In the operations research domain, two fundamental frameworks exist to inform such decisions. The *stochastic optimization* framework captures uncertainty by using probability distributions and aims to optimize an expected objective. Initially introduced in the seminal work of Dantzig (1955), the framework has been intensively studied and scholars used it to solve a vast variety of problems including production planning (Beebe et al., 1968; Schmidt, 1996), relieve networks (Noyan et al., 2016), expansion planning (Xie and Huang, 2018; Zou et al., 2018), and newsvendor problems (Porteus, 1990). While stochastic optimization performs well on many problem classes, finding tractable formulations is oftentimes challenging. Additionally, the data needed to approximate probability distributions might not always be available and gathering data is often a costly and time consuming process. The *robust optimization* (RO) framework overcomes many of these shortcomings by capturing uncertainty through distribution free uncertainty sets instead of probability distributions. By choosing well representable uncertainty sets,

*Chair of Operations Management, RWTH Aachen, 52072 Aachen, Germany

†TUM School of Management & Munich Data Science Institute, Technical University of Munich, 80333 Munich, Germany

RO often offers computationally tractable formulations that scale well on a variety of optimization problems. In recent years, these favorable properties have led to a steep increase of research interest (see, e.g., Ben-Tal and Nemirovski, 2002; Ben-Tal et al., 2009a; Bertsimas et al., 2011a; Gorissen et al., 2015; Yanikoglu et al., 2019). Flexibility of uncertainty sets and scalability of solution methods also make RO very attractive for applicational purposes and it has been widely applied to many operations management problems (c.f. Lu and Shen, 2021).

In classical RO all decisions have to be made before the uncertainty realization is revealed, which is often not necessary in practice such that RO may yield overly pessimistic solutions. To remedy this drawback, Ben-Tal et al. (2004) introduce the concept of *adjustable robust optimization* (ARO) where some decisions can be delayed until the uncertainty realization is (partly) known. In general ARO, uncertainty realizations are revealed over multiple stages and decisions can be made after each reveal. A decision made in stage t can thus be modeled as a function of all uncertainties associated with previous stages $t' \leq t$.

While this theoretically improves decision making, Ben-Tal et al. (2004) show that already the special case of two-stage ARO is computationally intractable in general. Accordingly, recent works developed many approximation schemes for ARO that often yield good and sometimes even optimal results in practice (see, e.g., Ben-Tal et al., 2004; Bertsimas et al., 2010; Iancu et al., 2013; Simchi-Levi et al., 2019a). In the special case with only two decision stages, the first stage decisions are fixed before any uncertain parameters are known and the second stage decisions use full knowledge of the uncertainty realization. Two-stage ARO already has many applications in practice and has widely been studied in the literature (see, e.g., Ben-Tal et al., 2020; Bertsimas and Goyal, 2012; Bertsimas et al., 2015; Bertsimas and Dunning, 2016; Bertsimas et al., 2021; Hanasusanto et al., 2015; Housni and Goyal, 2021; Simchi-Levi et al., 2019b). For two-stage adjustable robust linear packing problems, Bertsimas et al. (2015) show that even static solutions are optimal.

Still, many real-world problems show inherent multi-stage characteristics and cannot be modeled by two-stage ARO. Examples include variants of inventory management (Ben-Tal et al., 2009b), humanitarian relief (Ben-Tal et al., 2011) and facility location (Goh and Hall, 2013). The two main challenges distinguishing multi-stage ARO from two-stage ARO are *nonanticipativity restrictions* that forbid decisions to use future information, and exponential growth of scenario trees in the number of stages. As a consequence, many results found for two-stage ARO do not readily generalize to multi-stage scenarios. Against this background, we design piecewise affine policies for multi-stage ARO that overcome the previously mentioned challenges and extend, although it is not straight forward, many of the best known approximation bounds for two-stage problems to a multi-stage setting.

In the remainder of this section we formally introduce our problem (Section 1.1), discuss closely related work (Section 1.2), and summarize our main contributions (Section 1.3).

1.1 Problem Description

In this work we study multi-stage adjustable robust optimization with covering constraints and a positive affine uncertain right hand side. Specifically, we consider the following problem:

$$\begin{aligned} Z_{AR}(\mathcal{U}) = \min_{\mathbf{x}(\boldsymbol{\xi})} \max_{\boldsymbol{\xi} \in \mathcal{U}} \quad & \mathbf{c}^\top \mathbf{x}(\boldsymbol{\xi}) \\ & \mathbf{A}\mathbf{x}(\boldsymbol{\xi}) \geq \mathbf{D}\boldsymbol{\xi} + \mathbf{d} \quad \forall \boldsymbol{\xi} \in \mathcal{U} \end{aligned} \quad (1)$$

where $\mathbf{A} \in \mathbb{R}^{l \times n}$, $\mathbf{c} \in \mathbb{R}^n$, $\mathbf{D} \in \mathbb{R}_+^{l \times m}$, $\mathbf{d} \in \mathbb{R}_+^l$ and $\mathcal{U} \subset \mathbb{R}_+^m$. Here, m is the number of uncertainty parameters, n is the number of decisions, and l is the number of constraints. To model the T stages of the problem we split the uncertainty vector $\boldsymbol{\xi}$ into T sub-vectors $\boldsymbol{\xi} = (\boldsymbol{\xi}^1, \dots, \boldsymbol{\xi}^T)$ with $\boldsymbol{\xi}^t$ being the uncertainty vector realized in stage t . In the following, we denote by $\underline{\boldsymbol{\xi}}^t := (\boldsymbol{\xi}^1, \dots, \boldsymbol{\xi}^t)$ the vector of all uncertainties with known realization in stage t . Similarly, the adjustable decision vector $\mathbf{x}(\boldsymbol{\xi})$ divides into $\mathbf{x}(\boldsymbol{\xi}) := (\mathbf{x}^1(\underline{\boldsymbol{\xi}}^1), \dots, \mathbf{x}^T(\underline{\boldsymbol{\xi}}^T))$ where the decision \mathbf{x}^t made in stage t has to preserve *nonanticipativity* and may only depend on those uncertainties $\underline{\boldsymbol{\xi}}^t$ whose realization is known in stage t . Finally, we denote by $\underline{\mathbf{x}}^t(\boldsymbol{\xi}) := (\mathbf{x}^1(\underline{\boldsymbol{\xi}}^1), \dots, \mathbf{x}^t(\underline{\boldsymbol{\xi}}^t))$ the vector of all decisions in the first t stages.

W.l.o.g. we assume in the following that $\mathcal{U} \subseteq [0, 1]^m$ is convex, full-dimensional with $e_i \in \mathcal{U}$ for all $i \in \{1, \dots, m\}$, and down-monotone. Down-monotonicity holds because $\mathbf{D}, \mathbf{d}, \boldsymbol{\xi}$ are all non-negative and thus constraints only simplify for smaller values of $\boldsymbol{\xi}$. Convexity holds due to the linearity of the problem, and $e_i \in \mathcal{U} \subseteq [0, 1]^m$ holds as \mathbf{D} can be rescaled appropriately. We note that the non-negativity assumption of the right hand side does restrict the problem space. As Bertsimas and Goyal (2012) point out, this assumption prevents the introduction of uncertain or constant upper bounds. However, upper bounds in other decision variables are still possible as \mathbf{A} is not restricted, and \mathbf{D}, \mathbf{d} can be zero.

Prominent problem classes covered by Problem (1) include network design (Ordóñez and Zhao, 2007; Wang and Qi, 2020), capacity planning (Laguna, 1998; Mulvey et al., 1995; Paraskevopoulos et al., 1991), as well as versions of inventory management (Ben-Tal et al., 2009b; Solyali et al., 2016) where capacities are unbounded or subject to the decision makers choice. However, as pointed out above, Problem (1) does not cover fixed or uncertain upper bounds including non-expandable production, storage, or shipment capacity, external (uncertain) resource availability and supply uncertainty. This also implies that we cannot model strict equality constraints with non-zero constant, or uncertain parameters. However, oftentimes only one inequality of the equality constraints is crucial for the problem. By relaxing these constraints, we can thus often get an equivalent formulation that fulfills all the assumptions of Problem (1).

1.2 Related Work

Feige et al. (2007) show that already the two-stage version of Problem (1) with $\mathbf{D} = \mathbf{1}$, $\mathbf{d} = \mathbf{0}$ and \mathbf{A} being a 0-1-matrix is hard to approximate with a factor better than $\Omega(\log m)$, even for budgeted uncertainty sets. As it is thereby impossible to find general solutions for \mathbf{x} , a common technique to get tractable formulations is to restrict the function space.

In this context, Ben-Tal et al. (2004) consider \mathbf{x} to be affine in $\boldsymbol{\xi}$. Specifically, they propose \mathbf{x}^t to be of the form $\mathbf{x}^t(\underline{\boldsymbol{\xi}}^t) = \mathbf{P}^t \underline{\boldsymbol{\xi}}^t + \mathbf{q}^t$. Affine policies have been found to deliver good results in practice (Adida and Perakis, 2006; Ben-Tal et al., 2005) and are even optimal for some special problems (Bertsimas et al., 2010; Iancu et al., 2013; Simchi-Levi et al., 2019a). Kuhn et al. (2011) investigate the

optimality of affine policies by using the gap between affine solutions on the primal and the dual of the problem. Further popular decision rules include segregate affine (Chen et al., 2008; Chen and Zhang, 2009), piecewise constant (Bertsimas et al., 2011a), piecewise affine (Ben-Tal et al., 2020; Georghiou et al., 2015), and polynomial (Bampou and Kuhn, 2011; Bertsimas et al., 2011b) policies, as well as combinations of the previous (Rahal et al., 2021). For surveys on adjustable policies we refer to Delage and Iancu (2015) and Yanıkođlu et al. (2019).

Many approximation schemes have been proposed for the two-staged version of Problem (1). For general uncertainty sets on the two-stage version of (1), Bertsimas and Goyal (2012) show that affine policies yield an $O(\sqrt{m})$ approximation if \mathbf{c} and \mathbf{x} are non-negative. They further construct a set of instances where this bound is tight, showing that no better general bounds for affine policies exist. Using geometric properties of the uncertainty sets, Bertsimas and Bidkhori (2015) improve on these bounds for some commonly used sets including budgeted uncertainty, norm balls, and intersections of norm balls. Ben-Tal et al. (2020) propose new piecewise affine decision rules for the two-stage problem that on some sets improve these bounds even further. In addition to strong theoretical bounds, this new approach also yields promising numerical results that can be found by orders of magnitude faster than solutions for affine adjustable policies. For budgeted uncertainty sets and some generalizations thereof, Housni and Goyal (2021) show that affine policies even yield optimal approximations with an asymptotic bound of $O\left(\frac{\log m}{\log \log m}\right)$. This bound was shown to be tight by Feige et al. (2007) for reasonable complexity assumptions, namely 3SAT cannot be solved in time $2^{O(\sqrt{m})}$ on instances of size m .

To the best of our knowledge, Bertsimas et al. (2011a) are the only ones that provide bounds for multi-stage ARO so far. They show these bounds for piecewise constant policies using geometric properties of the uncertainty sets. More specifically, they consider multi-stage uncertainty networks, where the uncertainty realization is taken from one of multiple independent uncertainty sets in each stage. While the choice of the set selected in each stage may depend on the sets selected before, the uncertainty sets are otherwise independent. Although this assumption is fairly general, it still leaves many commonly used sets where uncertainty is dependent over multiple stages uncovered. Among others, uncovered sets include hypersphere and budgeted uncertainty.

As can be seen, previous work has predominantly focused on providing tighter approximation bounds for two-stage ARO, inevitably raising the question whether similar bounds hold for multi-stage ARO as well. This work contributes towards answering this question by extending many of the currently best known bounds on two-stage ARO to its multi-stage setting. By so doing, we are - to the best of our knowledge - the first ones to provide approximation bounds for the multi-stage ARO Problem (1), where uncertainty sets can range over multiple stages.

1.3 Our Contributions

With this work we extend the existing literature in multiple ways, where our main contributions are as follows.

Tractable Piecewise Affine Policies for Multi-Stage ARO: Motivated by piecewise affine policies for two-stage ARO (cf. Ben-Tal et al., 2020) we present a framework to construct policies that can be used to efficiently find good solutions for the multi-stage ARO Problem (1). Instead of solving the problem

directly for uncertainty \mathcal{U} , we first approximate \mathcal{U} by a dominating set $\hat{\mathcal{U}}$. To do so, we define the concept of nonanticipative multi-stage domination and show that this new definition of domination fulfills similar properties to two-stage domination. Based on this new definition, we then construct dominating sets $\hat{\mathcal{U}}$ such that solutions on $\hat{\mathcal{U}}$ can be found efficiently. More specifically, we choose $\hat{\mathcal{U}}$ to be a polytope for which worst case solutions can be computed by a linear program (LP) over its vertices. In order to ensure nonanticipativity, which is the main challenge in this construction, we introduce a new set of constraints on the vertices that guarantee the existence of nonanticipative extensions from the vertex solutions to the full set $\hat{\mathcal{U}}$. Finally, we show how to use the solution on the dominating set $\hat{\mathcal{U}}$ to construct a valid solution for the original uncertainty set \mathcal{U} .

Approximation Bounds: To the best of our knowledge, we provide the first approximation bounds for the multi-stage Problem (1) with general uncertainty sets. More specifically we show that our policies yield $O(\sqrt{m})$ approximations of fully adjustable policies. While this bound is tight for our type of policies in general, we show that better bounds hold for many commonly used uncertainty sets. In fact we prove that all bounds that were previously shown for two-stage ARO by Ben-Tal et al. (2020) hold for our multi-stage policies as well.

While our main contribution is to extend approximation bounds to multi-stage ARO, Problem (1) is further less restrictive than problems previously discussed in the literature. In addition to being restricted to two-stage ARO, previous work often assumed \mathbf{c} and \mathbf{x} to be non-negative (Bertsimas and Goyal, 2012; Bertsimas and Bidkhori, 2015; Ben-Tal et al., 2020). Ben-Tal et al. (2020) additionally restricted parts of \mathbf{A} , i.e. they require the parts of \mathbf{A} associated with the first stage decision to be non-negative. Our policies do not need this assumption. However, we show that Problem (1) is unbounded whenever there is a feasible \mathbf{x} with $\mathbf{c}^\top \mathbf{x} < 0$, due to the non-negativity of the right hand side. As a consequence our policies do not readily extend to general maximization problems.

While mainly asymptotic bounds are of interest from a theoretical perspective, in practice constant factors are important as well. We thus demonstrate how the bound given in Ben-Tal et al. (2020) can be improved by up to a factor of $\frac{1}{\sqrt{2}}$ for hypersphere uncertainty sets and by up to a factor of $\frac{1}{2}$ for budgeted uncertainty sets.

Comparison with Affine Policies: Using the newly found bounds, we show that no approximation bound for affine policies on hypersphere uncertainty exists that is better than the bound we show for our policies. For budgeted uncertainty on the other hand, we show that affine policies strictly dominate our piecewise affine policies. These findings confirm results that have been reported for the two-stage variant, where affine policies do not perform well for hypersphere uncertainty (Bertsimas and Goyal, 2012), while performing very well for budgeted uncertainty (Housni and Goyal, 2021).

Improvement Heuristic: Due to inherent properties of our policy construction, resulting solutions are overly pessimistic on instances where the impact on the objective varies significantly between different uncertainty dimensions. To diminish this effect, we introduce an improvement heuristic that performs at least as good as our policies and that can be integrated into the LP used to construct our policies. While these modifications come at the cost of higher solution times, they allow for significant improvements on some instance classes.

Numerical Evidence: Finally, we present two sets of numerical experiments showing that our policies solve by orders of magnitude faster than the affine adjustable policies presented by Ben-Tal et al. (2004), often yielding comparable or improving results. First, we study a slightly modified version of the tests presented in Ben-Tal et al. (2020), allowing us to demonstrate our policies’ scalability and the impact of our improvement heuristic. Second, we focus on demand covering instances to demonstrate good performances of our policies for a problem that resembles a practical application.

The rest of this paper is structured as follows. In Section 2, we introduce our policies and elaborate on their construction. In Section 3, we present our approximation bounds for the multi-stage ARO Problem (1). We present an improvement heuristic for our policies in Section 4 and finally provide numerical evidence for the performance of our policy compared to affine adjustable policies in Section 5. Section 6 concludes this paper with a brief reflection of our work and avenues for future research. To keep the paper concise, we defer long proofs that could possibly interrupt the reading flow to appendices A-F and keep only short proof sketches for the respective theorems and lemmata in the main body of the paper.

2 Framework For Piecewise Affine Multi-Stage Policies

In this section, we present our piecewise affine framework for the multi-stage ARO Problem (1). Our approach is similar to the two-stage framework from Ben-Tal et al. (2020) with the main difference being the handling of nonanticipativity constraints. The main rationale of our framework is to construct new uncertainty sets $\hat{\mathcal{U}}$ that dominate the original uncertainty sets \mathcal{U} . For a problem $Z_{AR}(\mathcal{U})$ we construct $\hat{\mathcal{U}}$ in such a way that $Z_{AR}(\hat{\mathcal{U}})$ can be efficiently solved, and a solution of $Z_{AR}(\hat{\mathcal{U}})$ can be used to generate solutions for $Z_{AR}(\mathcal{U})$. Intuitively, an uncertainty set $\hat{\mathcal{U}}$ dominates another set \mathcal{U} if for every point in \mathcal{U} there is a point in $\hat{\mathcal{U}}$ that is at least as large in each component.

Definition 1 (Domination). Given an uncertainty set $\mathcal{U} \subseteq \mathbb{R}_+^m$, we say that \mathcal{U} is *dominated* by $\hat{\mathcal{U}} \subseteq \mathbb{R}_+^m$ if there is a *domination function* $\mathbf{h}: \mathcal{U} \rightarrow \hat{\mathcal{U}}$ with $\mathbf{h}(\boldsymbol{\xi}) \geq \boldsymbol{\xi}$, and \mathbf{h} can be expressed as $\mathbf{h}(\boldsymbol{\xi}) = (\mathbf{h}^1(\boldsymbol{\xi}^1), \dots, \mathbf{h}^T(\boldsymbol{\xi}^T))$ where \mathbf{h}^t maps to the uncertainties in stage t and depends on uncertainties up to that stage.

Due to the non-negativity of the problem’s right-hand side, domination only makes the problem harder. As a consequence, each solution for a realization $\hat{\boldsymbol{\xi}} \in \hat{\mathcal{U}}$ is also a valid solution for all realizations $\boldsymbol{\xi} \in \mathcal{U}$ that are dominated by $\hat{\boldsymbol{\xi}}$. Using this property, we can derive piecewise affine policies for $Z_{AR}(\mathcal{U})$ from solutions of $Z_{AR}(\hat{\mathcal{U}})$. Since \mathcal{U} is full-dimensional, there always exists a factor $\beta \geq 0$ such that scaling \mathcal{U} by β contains $\hat{\mathcal{U}}$. Theorem 1 shows that with this factor β , solutions of problem $Z_{AR}(\hat{\mathcal{U}})$ are β -approximations for problem $Z_{AR}(\mathcal{U})$. It also shows that $Z_{AR}(\hat{\mathcal{U}})$ is unbounded exactly when $Z_{AR}(\mathcal{U})$ is unbounded. Thus, we assume for the remainder of this paper w.l.o.g. that both $Z_{AR}(\mathcal{U})$ and $Z_{AR}(\hat{\mathcal{U}})$ are bounded.

Theorem 1. Consider an uncertainty set \mathcal{U} from Problem (1) and a dominating set $\hat{\mathcal{U}}$. Let $\beta \geq 1$ be such that $\forall \hat{\boldsymbol{\xi}} \in \hat{\mathcal{U}}: \frac{1}{\beta} \hat{\boldsymbol{\xi}} \in \mathcal{U}$. Moreover, let $Z_{AR}(\mathcal{U})$ and $Z_{AR}(\hat{\mathcal{U}})$ be optimal values of Problem (1). Then,

either $Z_{AR}(\mathcal{U})$ and $Z_{AR}(\hat{\mathcal{U}})$ are unbounded or

$$0 \leq Z_{AR}(\mathcal{U}) \leq Z_{AR}(\hat{\mathcal{U}}) \leq \beta \cdot Z_{AR}(\mathcal{U}).$$

We prove Theorem 1 in two steps. First, we show that both problems are unbounded when $Z_{AR}(\mathcal{U})$ or $Z_{AR}(\hat{\mathcal{U}})$ has a feasible solution with negative objective value. We do this by using such a feasible negative solution to construct new solutions for $Z_{AR}(\mathcal{U})$ and $Z_{AR}(\hat{\mathcal{U}})$ that are arbitrarily small. Second, we show that $\hat{\mathbf{x}}(\mathbf{h}(\boldsymbol{\xi}))$ is a feasible solution for $Z_{AR}(\mathcal{U})$ whenever $\hat{\mathbf{x}}(\hat{\boldsymbol{\xi}})$ is a feasible solution for $Z_{AR}(\hat{\mathcal{U}})$, which implies $Z_{AR}(\mathcal{U}) \leq Z_{AR}(\hat{\mathcal{U}})$. Conversely, any solution $\beta\mathbf{x}(\frac{1}{\beta}\hat{\boldsymbol{\xi}})$ is a feasible solution for $Z_{AR}(\hat{\mathcal{U}})$ whenever \mathbf{x} is a feasible solution for $Z_{AR}(\mathcal{U})$, which implies the last inequality. For a detailed proof, we refer to Appendix A.

In the remainder of this section, we demonstrate how the results of Theorem 1 can be used to efficiently construct β -approximations for $Z_{AR}(\mathcal{U})$. In this context, we note that one cannot straight forwardly apply the construction scheme used by Ben-Tal et al. (2020) due to nonanticipativity requirements. More specifically, Ben-Tal et al. (2020) construct $\hat{\mathcal{U}}$ as polytopes, where it is well known that worst case solutions always occur on extreme points, as any solution can be represented by convex combinations of extreme point solutions. The construction of these convex combinations, however, is not guaranteed to be nonanticipative in the multi-stage setting. In Section 2.1 we show how to construct dominating polytopes $\hat{\mathcal{U}}$ and efficiently find solutions $Z_{AR}(\hat{\mathcal{U}})$ that comply with nonanticipativity requirements. Then, we construct the dominating function $\mathbf{h}: \mathcal{U} \rightarrow \hat{\mathcal{U}}$ allowing us to extend these solutions to solutions for $Z_{AR}(\mathcal{U})$ in Section 2.2.

2.1 Construction of the Dominating Set

In the following, we construct a dominating set in the form of a polytope for which the worst case solution can be efficiently found by solving a linear program on its vertices. Specifically, for an uncertainty set \mathcal{U} , we consider dominating sets $\hat{\mathcal{U}}$ of the form

$$\hat{\mathcal{U}} := \text{conv}(\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_m) \quad (2)$$

where for all $i \in \{0, \dots, m\} : \frac{1}{\beta}\mathbf{v}_i \in \mathcal{U}$ and for all $i \in \{1, \dots, m\} : \mathbf{v}_i = \mathbf{v}_0 + \rho_i \mathbf{e}_i$ for some $\rho_i \in \mathbb{R}_+$. We postpone the construction of the domination function \mathbf{h} and the coefficients \mathbf{v}_0 and ρ_1, \dots, ρ_m to Section 2.2, and first focus on the construction of solutions for $Z_{AR}(\hat{\mathcal{U}})$. The key component for our construction is LP (3)

$$Z_{LP}(\hat{\mathcal{U}}) = \min_{\mathbf{x}_0, \dots, \mathbf{x}_m} z \quad (3a)$$

$$z \geq \mathbf{c}^\top \mathbf{x}_i \quad \forall i \in \{0, \dots, m\} \quad (3b)$$

$$\mathbf{A}\mathbf{x}_i \geq \mathbf{D}\mathbf{v}_i + \mathbf{d} \quad \forall i \in \{0, \dots, m\} \quad (3c)$$

$$\mathbf{x}_i^t = \mathbf{x}_j^t \quad \forall i, j \in \{0, \dots, m\}, t \in \{1, \dots, T\}, \underline{\mathbf{v}}_i^t = \underline{\mathbf{v}}_j^t. \quad (3d)$$

Here, we extend the notation on \mathbf{x} and $\boldsymbol{\xi}$ introduced in Section 1.1 to \mathbf{x}_i and \mathbf{v}_i . Consequently, \mathbf{x}_i^t is the sub-vector of \mathbf{x}_i corresponding to decisions made in stage t , and $\underline{\mathbf{v}}_i^t$ is the sub-vector of \mathbf{v}_i corresponding

to uncertainties up to stage t .

Intuitively, the Objective (3a) together with Constraints (3b) minimize the maximal cost over all vertex solutions \mathbf{x}_i . Constraints (3c) ensure that each \mathbf{x}_i is a feasible solution for the respective uncertainty vertex \mathbf{v}_i of $\hat{\mathcal{U}}$. Finally, Constraints (3d) ensure nonanticipativity by forcing vertex solutions to be equal unless different uncertainties were observed. With these constraints, we construct LP (3) such that in order to find an optimal solution for $Z_{AR}(\hat{\mathcal{U}})$, it is sufficient to find an optimal solution for $Z_{LP}(\hat{\mathcal{U}})$.

Lemma 1. *Let $\hat{\mathcal{U}}$ be a dominating set as described in (2), $Z_{LP}(\hat{\mathcal{U}})$ be the solution of LP (3), and $Z_{AR}(\hat{\mathcal{U}})$ be the solution of Problem (1). Then the LP solution (\mathbf{x}_i) on the vertices of $\hat{\mathcal{U}}$ can be extended to a solution on the full set $\hat{\mathcal{U}}$ and we find:*

$$Z_{LP}(\hat{\mathcal{U}}) = Z_{AR}(\hat{\mathcal{U}}).$$

To prove Lemma 1, we show that $\mathbf{x}_i := \mathbf{x}(\mathbf{v}_i)$ is a feasible solution for the LP whenever \mathbf{x} is a feasible solution for $Z_{AR}(\hat{\mathcal{U}})$. This implies $Z_{LP}(\hat{\mathcal{U}}) \leq Z_{AR}(\hat{\mathcal{U}})$. To show $Z_{LP}(\hat{\mathcal{U}}) \geq Z_{AR}(\hat{\mathcal{U}})$, we use that each $\boldsymbol{\xi} \in \hat{\mathcal{U}}$ is given by a unique convex combination $\boldsymbol{\xi} = \sum_{i=0}^m \lambda_i \mathbf{v}_i$. This convex combination can then be used to extend an LP solution $\mathbf{x}_0, \dots, \mathbf{x}_m$ to a solution $\mathbf{x}(\boldsymbol{\xi}) := \sum_{i=0}^m \lambda_i \mathbf{x}_i$ for $Z_{AR}(\hat{\mathcal{U}})$. For a detailed proof, we refer to Appendix B.

2.2 Construction of the Domination Function

In the previous section, we showed how to construct dominating sets $\hat{\mathcal{U}}$ such that $Z_{AR}(\hat{\mathcal{U}})$ can be solved efficiently. In order for $\hat{\mathcal{U}}$ to be a valid dominating set for some uncertainty set \mathcal{U} , we additionally have to construct a nonanticipative dominating function $\mathbf{h}: \mathcal{U} \rightarrow \hat{\mathcal{U}}$ according to Definition 1. Specifically, we use

$$\mathbf{h}(\boldsymbol{\xi}) := (\boldsymbol{\xi} - \mathbf{v}_0)_+ + \mathbf{v}_0 \tag{4}$$

where $(\cdot)_+$ is the element-wise maximum with 0. By construction, \mathbf{h} maps each uncertainty realization $\boldsymbol{\xi}$ to its element-wise maximum with \mathbf{v}_0 . It directly follows that \mathbf{h} is nonanticipative, as each element in $\mathbf{h}(\boldsymbol{\xi})$ solely depends on the corresponding element in $\boldsymbol{\xi}$.

Finally, we have to ensure that $\mathbf{h}(\boldsymbol{\xi}) \in \hat{\mathcal{U}}$ for all $\boldsymbol{\xi} \in \mathcal{U}$. We do so by choosing the coefficients \mathbf{v}_0 and ρ_1, \dots, ρ_m during the construction of $\hat{\mathcal{U}}$ appropriately. Using $\lambda_i(\boldsymbol{\xi}) := \frac{((\boldsymbol{\xi} - \mathbf{v}_0)_+)_i}{\rho_i}$ we find

$$\mathbf{h}(\boldsymbol{\xi}) = \sum_{i=1}^m \lambda_i(\boldsymbol{\xi}) \mathbf{v}_i + \left(1 - \sum_{i=1}^m \lambda_i(\boldsymbol{\xi})\right) \mathbf{v}_0.$$

By definition, any convex combination of \mathbf{v}_i is contained in $\hat{\mathcal{U}}$. Thus, \mathbf{h} is a valid domination function whenever

$$\max_{\boldsymbol{\xi} \in \mathcal{U}} \sum_{i=1}^m \lambda_i(\boldsymbol{\xi}) \leq 1. \tag{5}$$

This gives an easy to check criterion to construct valid dominating sets.

Recall that we showed how to extend a solution $(\mathbf{x}_0, \dots, \mathbf{x}_m)$ of LP (3) to the full set $\hat{\mathcal{U}}$ in the proof of Lemma 1. Combining this with \mathbf{h} and using Theorem 1 we get a piecewise affine solution for $Z_{AR}(\mathcal{U})$

No.	Uncertainty Set \mathcal{U}	Bound	Asymptotic Bound
I	$\left\{ \boldsymbol{\xi} \in \mathbb{R}_+^m \mid \ \boldsymbol{\xi}\ _2^2 \leq 1 \right\}$	$\sqrt[4]{m} \frac{\sqrt{m-1}}{\sqrt{2(m-\sqrt{m})}}$	$O(\sqrt[4]{m})$
II	$\left\{ \boldsymbol{\xi} \in [0, 1]^m \mid \ \boldsymbol{\xi}\ _1 \leq k \right\}$	$\frac{k(m-1)}{m+k(k-2)}$	$O\left(\min\left\{k, \frac{m}{k}\right\}\right)$
III	$\left\{ \boldsymbol{\xi} \in \mathbb{R}_+^m \mid \boldsymbol{\xi}^\top \boldsymbol{\Sigma} \boldsymbol{\xi} \leq 1 \right\}$	$\left(\frac{a}{2} + \frac{\sqrt{1-a}}{\sqrt[4]{am^2+(1-a)m}} \right)^{-1}$	$O\left(m^{\frac{2}{5}}\right)$
IV	$\left\{ \boldsymbol{\xi} \in \mathbb{R}_+^m \mid \ \boldsymbol{\xi}\ _p \leq 1 \right\}$	$\frac{2}{p}(p-1) \frac{p-1}{p} m^{\frac{p-1}{p^2}}$	$O\left(m^{\frac{p-1}{p^2}}\right)$
V	$\left\{ \boldsymbol{\xi} \in \mathbb{R}_+^m \mid \ \boldsymbol{\xi}\ _p \leq 1, \ \boldsymbol{\xi}\ _q \leq r \right\}$	$\min\left\{r^{\frac{1-p}{p}} m^{\frac{p-1}{pq}}, r^{\frac{1}{q}} m^{\frac{q-1}{q^2}}\right\}$	$O\left(\min\left\{r^{\frac{1-p}{p}} m^{\frac{p-1}{pq}}, r^{\frac{1}{q}} m^{\frac{q-1}{q^2}}\right\}\right)$
VI	$\mathcal{U} \subset \mathbb{R}_+^m$	$2\sqrt{m} + 1$	$O(\sqrt{m})$

Table 1: Performance bounds of the piecewise affine policy for different uncertainty sets.

We prove specific bounds for uncertainty sets of the forms I hypersphere uncertainty; II budgeted uncertainty; III ellipsoid uncertainty, with $\boldsymbol{\Sigma} := (1-a)\mathbf{1} + a\mathbf{J}$ where $\mathbf{1}$ is the unity matrix and \mathbf{J} the matrix of all ones; IV p -norm ball, with $p \geq 1$; V intersection of two norm balls, with $m^{\frac{1}{q}-\frac{1}{p}} \geq r \geq 1$; VI general uncertainty sets.

by

$$\mathbf{x}(\boldsymbol{\xi}) = \sum_{i=1}^m \lambda_i(\boldsymbol{\xi}) \mathbf{x}_i + \left(1 - \sum_{i=1}^m \lambda_i(\boldsymbol{\xi})\right) \mathbf{x}_0 \quad (6)$$

that has an optimality bound of β .

3 Optimality Bounds for Different Uncertainty Sets

In the previous section we demonstrated how to construct nonanticipative piecewise affine policies for the multi-stage Problem (1). On this basis, proving approximation bounds mainly depends on geometric properties of the uncertainty sets \mathcal{U} . As these properties do not depend on the number of stages, we can extend all bounds shown for the two-stage problem by Ben-Tal et al. (2020) to our multi-stage problem. We formalize this extension in Proposition 1. After that, we show constant factor improvements for the approximation bounds of hypersphere and budgeted uncertainty and demonstrate how the bounds of an uncertainty set \mathcal{U} generalize to transformations of that set.

Proposition 1. *All performance bounds from Ben-Tal et al. (2020) extend to solutions of the form in (6) for Problem (1). This includes cases III to VI from Table 1.*

Proof. First, note that the domination function \hat{h} defined in Equation (2.3) of Ben-Tal et al. (2020) is the same function as we defined in Equation (4). Second, the uncertainty set used for Theorem 2 in Ben-Tal et al. (2020) effectively is our dominating set $\hat{\mathcal{U}}$ from (2) with $\rho_i = \frac{\beta}{2}$ for a β approximation guarantee. All further results solely depend on the uncertainty sets and the performance bounds in the table follow by Propositions 2, 3, 4 and Theorem 5 in Ben-Tal et al. (2020). \square

Note that Ben-Tal et al. (2020) also showed that the asymptotic approximation bounds in Case VI of Table 1 are tight when using pure domination techniques. More precisely, for any polynomial number of vertices the budgeted uncertainty set with $k = \sqrt{m}$ cannot be dominated with some β better than $O(\sqrt{m})$. While this is a strong result for general sets, Table 1 shows that better bounds exist for many commonly used sets. Moreover, not only asymptotic bounds, but also constant factors, are of interest

for application purposes. We demonstrate that factors can still be improved by slightly tightening the bounds of Ben-Tal et al. (2020) for hypersphere and budgeted uncertainty sets.

Both of these uncertainty sets are permutation invariant and thus $\hat{\mathcal{U}}$ is permutation invariant as well. More specifically, \mathbf{v}_i simplifies to $\mathbf{v}_0 = \mu \mathbf{e}$, $\mathbf{v}_i = \mathbf{v}_0 + \rho \mathbf{e}_i$, for some μ, ρ . With these simplifications Condition (5) becomes

$$\frac{1}{\rho} \max_{\boldsymbol{\xi} \in \mathcal{U}} \sum_{i=1}^m (\xi_i - \mu)_+ \leq 1. \quad (7)$$

With the rotational invariance of the problem, Ben-Tal et al. (2020) show that for any μ the maximization problem in (7) has a solution that is constant on the first j components and zero on all others components.

Lemma 2 (Lemma 4 in Ben-Tal et al. (2020)). *Let $\gamma(j)$ be the maximal average value of the first j components of any $\boldsymbol{\xi} \in \mathcal{U}$*

$$\gamma(j) := \frac{1}{j} \max_{\boldsymbol{\xi} \in \mathcal{U}} \sum_{i=1}^j \xi_i.$$

Then for each μ there exists an optimal solution $\boldsymbol{\xi}^$ for the maximization problem in Equation 7 that has the form*

$$\boldsymbol{\xi}^* = \sum_{i=1}^j \gamma(j) \mathbf{e}_i,$$

for some $j \leq m$.

Hypersphere Uncertainty: We first use Lemma 2 to find a new dominating set for hypersphere uncertainty. By doing so, we find a new approximation bound that improves the bound of $\sqrt[4]{m}$ provided in Ben-Tal et al. (2020) by a factor of $\sqrt{\frac{\sqrt{m+1}}{2\sqrt{m}}}$, which for large m converges towards $\frac{1}{\sqrt{2}}$. While this improvement is irrelevant for the asymptotic complexity of the problem, the new formulation of $\hat{\mathcal{U}}$ does make a difference in practice.

Proposition 2 (Hypersphere). *Consider the hypersphere uncertainty set $\mathcal{U} = \{\boldsymbol{\xi} \in \mathbb{R}_+^m \mid \|\boldsymbol{\xi}\|_2^2 \leq 1\}$. Then a solution for $Z_{AR}(\hat{\mathcal{U}})$ where $\hat{\mathcal{U}}$ is constructed using Criterion (7) with*

$$\mu = \frac{1}{2\sqrt[4]{m}}, \quad \rho = \frac{\sqrt[4]{m}}{2},$$

gives a $\beta = \sqrt{\frac{\sqrt{m+1}}{2}}$ approximation for problem $Z_{AR}(\mathcal{U})$.

To prove Proposition 2, our main rationale is to find μ, ρ such that (7) is always fulfilled and β is minimized. To do so, we use Lemma 2 and $\beta = \|\mathbf{v}_{i>0}\| = \sqrt{m\mu^2 + 2\mu\rho + \rho^2}$. For a detailed proof, we refer to Appendix C.

We can also use this improved performance bound to show that affine adjustable policies cannot yield better bounds than piecewise affine adjustable policies for $m \geq 21$. This is because there are instances of Problem (1) with hypersphere uncertainty where affine adjustable policies perform at least $\sqrt[4]{m} - \frac{1}{\sqrt[4]{m}}$ worse than an optimal policy. We formalize these results in Proposition 3. Note, that these better bounds do not imply that piecewise affine adjustable policies always yield better results than affine adjustable policies for hypersphere uncertainty.

Proposition 3. *Affine adjustable policies cannot achieve better performance bounds than $\sqrt[4]{m} - \frac{1}{\sqrt[4]{m}}$ for Problem (1) with hypersphere uncertainty, even for $\mathbf{c}, \mathbf{x}, \mathbf{A}$ being non negative and \mathbf{A} being a 0,1-matrix.*

We refer to Appendix D for a full proof of Proposition 3. In this proof we construct a class of instances as well as piecewise affine solutions for these instances. We then show that these piecewise affine solutions are better than optimal affine policies by a factor of $\sqrt[4]{m} - \frac{1}{\sqrt[4]{m}}$ by carefully bounding the objective values of both policies.

Budgeted Uncertainty: Next, we tighten the bounds for budgeted uncertainty sets. Proposition 4 shows that our new bound is given by $\beta = \frac{k(m-1)}{m+k(k-2)}$. Using

$$\frac{\beta}{k} = \frac{k(m-1)}{k(m+k(k-2))} = \frac{m-1}{m-1+k^2-2k+1} = \frac{m-1}{m-1+(k-1)^2} \leq 1$$

and

$$\frac{\frac{m}{k}}{\frac{m}{k}} = \frac{k^2(m-1)}{m(m+k(k-2))} = \frac{k^2(m-1)}{k^2(m-1)+k^2+m^2-2km} = \frac{k^2(m-1)}{k^2(m-1)+(m-k)^2} \leq 1,$$

we show $\beta \leq \min(k, \frac{m}{k})$, which is the bound in Ben-Tal et al. (2020). As $\frac{\beta}{k}$ is increasing in k and $\frac{\frac{m}{k}}{\frac{m}{k}}$ is decreasing in k , we obtain a maximum improvement for $k = \frac{m}{k} \Leftrightarrow k = \sqrt{m}$. At this point the improvement of the bound reaches a factor of $\frac{1}{2}$.

Proposition 4 (Budget). *Consider the budgeted uncertainty set $\mathcal{U} = \{\boldsymbol{\xi} \in [0, 1]^m \mid \|\boldsymbol{\xi}\|_1 \leq k\}$ for some $k \in \{1, \dots, m\}$. Then a solution for $Z_{AR}(\hat{\mathcal{U}})$ where $\hat{\mathcal{U}}$ is constructed using Criterion 7 with*

$$\mu = \frac{k(k-1)}{m+k(k-2)}, \quad \rho = \frac{k(m-k)}{m+k(k-2)},$$

gives a $\beta = \frac{k(m-1)}{m+k(k-2)}$ approximation for problem $Z_{AR}(\mathcal{U})$.

Similar to the proof of Proposition 2, our rationale is to find μ, ρ such that (7) is always fulfilled and β is minimized. To do so we again use Lemma 2, but this time we find β to be given by $\beta = \min(\frac{m}{k}\mu + \frac{1}{k}\rho, \mu + \rho)$. For the full proof, we refer to Appendix E.

Note, that there is no analogous result to Proposition 3 for budgeted uncertainty as Housni and Goyal (2021) showed that affine policies are in $O(\frac{\log(m)}{\log \log(m)})$ for two-stage problems with non-negative $\mathbf{c}, \mathbf{x}, \mathbf{A}$. Furthermore, our piecewise affine policies are strictly dominated by affine policies for integer budgeted uncertainty.

Proposition 5. *Consider Problem (1) with budgeted uncertainty and an integer budget. Let Z_{PAP} be the optimal value found by our piecewise affine policy and Z_{AFF} the optimal value found by an affine policy. Then*

$$Z_{AFF} \leq Z_{PAP}.$$

To prove Proposition 5 we introduce more general fully piecewise affine policies and show that under the assumptions of the Proposition the robust counterpart of these policies is identical to the robust counterpart of affine policies. For a detailed proof, we refer to Appendix F.

Transformed Uncertainty Sets: We note that by the right hand side $\mathbf{D}\boldsymbol{\xi} + \mathbf{d}$ of Problem (1) any positive affine transformation of uncertainty sets \mathcal{U} can be dominated by the same affine transformation of the dominating set $\hat{\mathcal{U}}$. As the approximation bounds do not depend on \mathbf{D} and \mathbf{d} , the bounds for the transformed set are the same as for the original set. One well known uncertainty type covered by these transformations is scaled ellipsoidal uncertainty $\{\boldsymbol{\xi} \mid \sum_{i=1}^m w_i \xi_i^2 \leq 1\}$, which was first proposed by Ben-Tal and Nemirovski (1999). These sets can be constructed via transformations from hypersphere uncertainty sets with a diagonal matrix \mathbf{D} with $D_{ii} = \frac{1}{w_i}$. Scaled ellipsoidal uncertainty has been applied to many robust optimization problems, including portfolio optimization (Ben-Tal and Nemirovski, 1999), supply chain contracting (Ben-Tal et al., 2005), network design (Mudchanatongsuk et al., 2008), and facility location (Baron et al., 2011).

Another widely used class that is partially covered by these positive affine transformations are factor-based uncertainties given by sets $\mathcal{U} = \{\mathbf{D}\mathbf{z} + \mathbf{d} \mid \mathbf{z} \in \mathcal{U}^z\}$. In these sets, uncertainties affinely depend on a set of factors \mathbf{z} that are drawn from a factor uncertainty set \mathcal{U}^z . Such uncertainty sets were used by Goldfarb and Iyengar (2003) and Ghaoui et al. (2003) to solve portfolio optimization problems and by See and Sim (2010) and Ang et al. (2012) to solve multi-period inventory management problems. In contrast to the general factor sets, where there are no limitations on \mathbf{D} and \mathbf{d} , our approach is restricted to positive factor matrices which allows only for positive correlations between uncertainties. Nevertheless, even this subset of factor-based uncertainties has wide applicational use. As an intuitive example, one could consider component demands where the factors are demands for finished products.

4 Re-scaling Expensive Vertices

On some instances uncertainties do not have a uniform impact on the objective. While accounting for high values in some of the uncertainty dimensions might drastically influence the objective, accounting for high values in others might barely have an impact. This effect can be crucial in our problem setting, because the creation of dominating sets may overemphasize single uncertainty dimensions by up to a factor of β by design. Accordingly, it might thus be beneficial to dominate these uncertainty dimensions more carefully on instances where a few critical uncertainties cause almost all the cost. In this context, we show that it is possible to shrink critical vertices \mathbf{v}_i at the cost of slightly shifting all other vertices towards that direction.

Lemma 3. *Let $\hat{\mathcal{U}} := \text{conv}(\mathbf{v}_0, \dots, \mathbf{v}_m)$ be a dominating set for \mathcal{U} . Let $\mathbf{s} \in [0, 1]^m$ be a vector of scales. Then $\hat{\mathcal{U}}' := \text{conv}(\mathbf{v}'_0, \dots, \mathbf{v}'_m)$ with $\mathbf{v}'_{ij} := s_j(1 - v_{ij}) + v_{ij}$ is also a dominating set for \mathcal{U} .*

Proof. Let $\mathbf{h}: \mathcal{U} \rightarrow \hat{\mathcal{U}}$ be the domination function of $\hat{\mathcal{U}}$. Then define $\mathbf{h}': \mathcal{U} \rightarrow \hat{\mathcal{U}}'$ by

$$h'_j(\boldsymbol{\xi}) := h_j(\boldsymbol{\xi}) + s_j(1 - h_j(\boldsymbol{\xi})). \quad (8)$$

This new dominating function \mathbf{h}' is nonanticipative, as \mathbf{h} is nonanticipative and the j^{th} component of \mathbf{h}' only depends on the j^{th} component of \mathbf{h} . To see that (8) is also a domination function, consider the two cases $h_j(\boldsymbol{\xi}) \geq 1$ and $h_j(\boldsymbol{\xi}) \leq 1$.

1. For $h_j(\boldsymbol{\xi}) \geq 1$ the second summand on the right hand side of (8) is negative for all $s_j \in [0, 1]$. Thus $h'_j(\boldsymbol{\xi})$ is minimal for $s_j = 1$ and we find $h'_j(\boldsymbol{\xi}) \geq h_j(\boldsymbol{\xi}) + 1(1 - h_j(\boldsymbol{\xi})) = 1$.

2. On the other hand, for $h'_j(\boldsymbol{\xi}) \leq 1$ the second summand is positive and $h'_j(\boldsymbol{\xi})$ is minimal for $s_j = 0$. Thus, we have $h'_j(\boldsymbol{\xi}) \geq h_j(\boldsymbol{\xi})$.

In any case, we find $\mathbf{h}'(\boldsymbol{\xi}) \geq \boldsymbol{\xi}$ by $h'_j \geq \min(1, h_j(\boldsymbol{\xi})) \geq \xi_j$ where we used $\mathcal{U} \subseteq [0, 1]^m$.

Next, we show $\mathbf{h}'(\boldsymbol{\xi}) \in \hat{\mathcal{U}}'$. For this let $\mathbf{h}(\boldsymbol{\xi}) = \sum_{i=0}^m \lambda_i \mathbf{v}_i$ be the convex combination of $\mathbf{h}(\boldsymbol{\xi}) \in \hat{\mathcal{U}}$. Then,

$$h'_j(\boldsymbol{\xi}) = h_j(\boldsymbol{\xi}) + s_j(1 - h_j(\boldsymbol{\xi})) \stackrel{(a)}{=} \sum_{i=0}^m (\lambda_i v_{ij} + s_j(1 - v_{ij})) = \sum_{i=0}^m \lambda_i v'_{ij}.$$

Note that in (a) we used $\sum_{i=0}^m \lambda_i = 1$. This shows $\mathbf{h}'(\boldsymbol{\xi}) = \sum_{i=0}^m \lambda_i \mathbf{v}'_i$ and thus $\mathbf{h}'(\boldsymbol{\xi}) \in \hat{\mathcal{U}}'$. \square

The two extreme cases for the modified dominating sets from Lemma 3 are given by $\mathbf{s} = \mathbf{0}$ and $\mathbf{s} = \mathbf{e}$. While the dominating set does not change for $\mathbf{s} = \mathbf{0}$, all vertices become the unit vector consisting of ones in every component for $\mathbf{s} = \mathbf{e}$. Intuitively, increasing s_i leads to shifting the i^{th} component towards one. In the same way as we constructed our dominating sets in (2) this shift of the i^{th} component towards one increases the value for all \mathbf{v}_j with $j \neq i$ and decrease it for $j = i$. Note that this Lemma is not limited to uncertainty sets that are constructed as described in (2), but holds for any dominating set created as a convex combination of vertices.

The transformation used in Lemma 3 is linear, which allows us to add \mathbf{s} as a further decision variable to the second constraint of LP (3) for a given $\hat{\mathcal{U}}$. As $\mathbf{s} = \mathbf{0}$ gives the original dominating set, any optimal solution found with these additional decision variables is at least as good as a non-modified solution. Thus, all performance bounds shown in Section 3 also hold for these re-scaled piecewise affine policies. Note that Proposition 5 also extends to the re-scaled uncertainty set, thus all re-scaled piecewise affine policies are strictly dominated by affine policies for integer budgeted uncertainty.

Adding \mathbf{s} to the LP increases its size, which in practice will often lead to an increase of solution times. To limit the increase of model size, it is possible to add only those s_i where one expects $s_i > 0$, as not adding a variable s_i is equivalent to fixing $s_i = 0$. Those s_i with $s_i > 0$ correspond to the critical uncertainty dimensions, and an experienced decision maker with sufficient knowledge of the problem might be able to identify them a priori.

5 Numerical Experiments

In this section, we present two numerical experiments to compare the performance of our piecewise affine policies for the different constructions of $\hat{\mathcal{U}}$ with the performance of the affine policies described in Ben-Tal et al. (2004). We compare the performance in terms of both objective value and computational time.

We run both of the following tests only with hypersphere uncertainty sets and budgeted uncertainty sets for two reasons: first, in the experiments of Ben-Tal et al. (2020) piecewise affine policies performed particularly good compared to affine adjustable policies for hypersphere uncertainty, and relatively bad for budgeted uncertainty. Accordingly, considering these two uncertainty types gives a good impression of the benefits and limitations of piecewise affine policies. Second, this experimental design allows us to analyze whether or not the new formulations with the tighter bounds presented in Propositions 2 and 4 have a significant impact in practice.

In our studies, we compare the following policies: the affine policies described in Ben-Tal et al. (2004)

(AFF), the constant policies resulting from a dominating set $\hat{\mathcal{U}} = \{\mathbf{e}\}$ with only a single point which by down-monotonicity corresponds to a box (BOX), our piecewise affine policies constructed as described in Propositions 2 and 4 (PAP), the piecewise affine policies constructed as described in Propositions 1 and 5 in Ben-Tal et al. (2020) (PAPBT), and our piecewise affine policies with the vertex re-scaling heuristic described in Section 4 (SPAP). Recall that by the construction of the re-scaling heuristic we always have $Z_{SPAP} \leq Z_{BOX}$ and $Z_{SPAP} \leq Z_{PAP}$.

For all studies, we used Gurobi Version 9.1 on a 6 core 3.70 GHz Processor using a single core per instance.

5.1 Gaussian Instances

We base our first set of benchmark instances on the experiments of Ben-Tal et al. (2020) and Housni and Goyal (2021). Accordingly, we generate instances of Problem (1) by choosing $m = l = n$, $\mathbf{d} = \mathbf{0}$, $\mathbf{D} = \mathbf{1}_m$ and generate \mathbf{c} , \mathbf{A} randomly as

$$\begin{aligned}\mathbf{c} &= \mathbf{e} + \alpha \mathbf{g}, \\ \mathbf{A} &= \mathbf{1}_m + \mathbf{G}.\end{aligned}$$

Here, \mathbf{e} is the vector of all ones, $\mathbf{1}_m$ is the identity matrix, \mathbf{g} and \mathbf{G} are randomly generated by independent and identically distributed (i.i.d.) standard gaussians, and α is a parameter that increases the asymmetry of the problem. More specifically, \mathbf{G} is given by $G_{ij} = |Y_{ij}|/\sqrt{m}$ and \mathbf{g} is given by $g_i = |y_i|$, where Y_{ij} and y_i are i.i.d. standard gaussians. Uncertainties $\boldsymbol{\xi}$ and decision variables \mathbf{x} are split into $\lfloor \sqrt{m} \rfloor$ stages where the i^{th} decision always belongs to the same stage as the i^{th} uncertainty. For the budgeted uncertainty sets we use a budget of $k = \sqrt{m}$. We consider values of m from $m = 10$ to $m = 100$ in increments of 10 and values of α in $\{0, 0.1, 0.5, 1, 5\}$. For each pair of m, α , we consider 100 instances. To make results more comparable, we scale all objective values presented by the constant policies results (i.e., Z/Z_{BOX}) and take averages over all 100 instances. Additionally, we present all results on a logarithmic scale and artificially lower bound the scale for solution times by 0.01s to make the effects on higher solution times more visible.

Figure 1 shows the performances and solution time results on hypersphere uncertainty sets for the different policies. We observe that all three piecewise adjustable policies perform significantly better than affine adjustable policies for small values of α . The improvement increases for larger values of m reaching almost a factor of 2 for $m = 100$ on our policies PAP and SPAP. As expected, the performance of PAP and SPAP for small values of α is almost indistinguishable due to the construction. For larger values of α the improvements of the piecewise affine adjustable policies vanish and the two policies without re-scaling (PAP and PAPBT) perform even worse than AFF for $\alpha = 5$. More severely, PAPBT even performs worse than BOX, which already is a worst case policy. Only our policy with re-scaling (SPAP) achieves better results than affine adjustable policies for all values of α .

While solution times for all policies except BOX grow exponentially in the instance size, piecewise affine adjustable policies are by orders of magnitude faster than classical affine adjustable policies (AFF). These solution time improvements exceed a factor of 100 for piecewise affine policies PAP and SPAP and a factor of 1,000 for PAPBT. Also, solution times of piecewise affine adjustable policies are barely




Figure 1: Relative objective values (a) and solution times (b) for different policies on gaussian instances with hypersphere uncertainty.

influenced by values of α . This is not the case for AFF, which takes longer to solve for increasing α .

Figure 2 shows the performances and solution time results on budgeted uncertainty sets. We observe that for budgeted uncertainty sets piecewise affine policies perform slightly worse than affine policies throughout all instances. This observation nicely demonstrates that our theoretical and experimental results are aligned, as the worse performance is perfectly explained by Proposition 5, which shows that affine policies strictly dominate our piecewise affine policies. Again, we observe that for higher values of α , PAP and PAPBT perform even worse than BOX. However, solution values for SPAP stay within 5% of affine solution values throughout all instances. Solution times behave similar as for hypersphere uncertainty confirming that piecewise affine policies are found by orders of magnitude faster on different instances and uncertainty types.

Solution time and performance results for $\alpha = 0$ align with the results found by Ben-Tal et al. (2020) for the two-stage variant. This demonstrates that our generalized piecewise affine policies do not only extend all theoretical performance bounds, but also achieve comparable numerical results in a multi-stage setting. However, breaking the symmetry by increasing α we show that pure piecewise adjustable policies




Figure 2: Relative objective values (a) and solution times (b) for different policies on gaussian instances with budgeted uncertainty.

perform poorly on highly asymmetric instances and re-scaling as performed by SPAP constitutes a good technique to overcome this shortcoming.

5.2 Demand Covering Instances

For the second set of test instances, we consider the robust demand covering problem with non-consumed resources and uncertain demands. The problem has various applications, among others in the domains of appointment scheduling, production planning, and dispatching, and is especially relevant for the optimization of service levels. Our instances consist of m^l locations, m^p planning periods and m^e execution periods per planning period. In each execution period t an uncertain demand ξ_{lt} has to be covered at each location l . To do so, the decision maker can buy a fixed number of resources R at a unit cost of c^R in a first stage and then distributes these R resources among the locations at the beginning of each planning period. If a demand cannot be met with the resources assigned to a location, the decision maker will either delay the demand to the next period or redirect it to another location. In either case a fraction

$q_{tl}^d \in [0, 1]$ or $q_{tll'}^r \in [0, 1]$ of the demand is lost. Each unit of lost demand causes costs of c^D . For a detailed problem formulation of the robust demand covering problem with non-consumed resources and uncertain demands as well as for the generation of instances, we refer to Appendix G.

In most real-world applications of the demand covering problem, some of the demand will be revealed before the actual demand occurs, e.g. due to already existing contracts, sign-ups, orders, or due to forecasting. To incorporate the increase in knowledge over time we assume an uncertainty vector of the form $\xi = \xi^c + \xi^p + \xi^e$. Here, ξ^c is constant and known before the first stage decision, ξ^p is revealed before each planning period and ξ^e accounts for the short term uncertainties revealed before each execution period.

To analyze practical expected objectives, we also report a simulated average objective that the respective policies achieved on 500 randomly drawn uncertainty realizations (ξ^p, ξ^e) , where the components are drawn i.i.d. from $\mathcal{N}(0, \frac{1}{m})$, in addition to the robust objective value. We again scale the results by the results achieved by constant policies and use logarithmic scales. Note that we set a hard solution time limit of $2h$ and base our analysis only on those instances that could be solved by all policies within this time limit. For larger instance sizes we exclude some policies, as they were not able to find any solutions within the solution time limit (e.g. AFF for $m^p = 3$ and $m^l = 8$).

Figure 3 shows the performance and solution time results for the different policies on demand covering instances with hypersphere uncertainty. Compared to our previous experiment (see Section 5.1) we no longer observe the strong objective improvements of piecewise affine policies over affine adjustable policies. Still, our PAP formulations give similar results as affine policies, and we observe small improvements on instances with a larger number of planning periods. On the simulated realizations, improvements over affine adjustable policies can already be seen for $m^p \geq 3$, which might be of interest to a decision maker with practical interest beyond worst case solutions.

For the solution times we observe similar improvements to the ones observed on the gaussian instances. Still, all piecewise affine policies can be found by orders of magnitude faster than affine policies. Interestingly, PAP is solved similarly fast as PAPBT on these instances, while still achieving up to 15% better objective values on all instances. The largest instance that could be solved by affine adjustable policies within two hours consisted of 320 uncertainty variables and 700 decision variables, while the largest instance solved by PAPs was more than three times larger with 1,120 uncertainty variables and 6,230 decision variables.

Figure 4 shows the results on demand covering instances with budgeted uncertainty sets. For budgeted uncertainty sets, piecewise affine policies perform worse than affine adjustable policies throughout all instances, which again can be explained by Proposition 5. Notably, PAPBT even performs worse than constant policies on most instances. Piecewise affine policies remain better only from a solution time perspective, as they still solve by orders of magnitude faster than affine policies.

While in this set of experiments piecewise affine policies do not show the same improvements in the objective over affine adjustable policies, they still perform slightly better with hypersphere uncertainty on larger instances. Also, they still solve by orders of magnitude faster, which makes them an attractive alternative for large-scale optimization in practice.




Figure 3: Relative objective values (a) and solution times (b) for different policies on demand covering instances with hypersphere uncertainty.

5.3 Discussion

In the experiments presented in Sections 5.1 and 5.2, some results, e.g. the solution time improvements of piecewise affine policies over affine policies, are consistent throughout all instances. However, other results strongly depend on the set of benchmark instances used. In the following, we discuss two of the main deviations that we observe between our results on gaussian instances, and demand covering instances and give intuitions why these differences occur.

Solution Times of PAPBT: First, we see that PAPBT solves by a factor of 10 to 50 faster than PAP on gaussian instances, but both find solutions similarly fast on demand covering instances. This can be explained by the construction of PAPBT and the structure of the instances' constraints. On the gaussian instances, \mathbf{A} and \mathbf{x} are non-negative and \mathbf{D} is the unit matrix. In the construction of dominating sets $\hat{\mathcal{U}}$ for PAPBT most of the vertices are chosen to be scaled unit vectors. As a consequence, most Constraints (3c) in LP (3) have a zero right hand side, such that they trivially hold. Consequently, these




Figure 4: Relative objective values (a) and solution times (b) for different policies on demand covering instances with budgeted uncertainty.

constraints can be eliminated, which reduces the total number of constraints by a factor of $O(m)$. On demand covering instances however, \mathbf{A} contains negative entries. Thus, no constraints trivially hold and no constraints can be removed. For a decision maker who is primarily interested in fast policies, this gives a good criterion on when PAPBT can improve solution times and when no such improvements can be expected.

Performance Differences Between Gaussian and Demand Covering Instances: Second, we observe that the strong performance improvements of piecewise affine policies over affine policies on gaussian instances with hypersphere uncertainty do not transfer to our demand covering instances. This suggests that the relative performance between piecewise affine policies and affine policies significantly depends on the structure of the problem at hand. An intuitive explanation for this lays in the the policies' construction.

Recall that piecewise affine policies derive solutions by finding vertex solutions \mathbf{x}_i that can be extended to a full solution. Thereby, \mathbf{x}_0 focuses on finding a good solution for uncertainty realizations where all

uncertainties take equal values, and each \mathbf{x}_i focuses on finding a good recourse to uncertainty ξ_i . Thus, good results can be expected when there are (a) synergy effects that can be utilized by \mathbf{x}_0 , and (b) good universal recourse decisions for each uncertainty ξ_i that do not depend on the realization of other uncertainty dimensions and can be exploited by \mathbf{x}_i . On the other hand affine policies, directly find solutions on the original uncertainty set. In doing so, they do not depend as strongly on good universal recourse decisions as piecewise affine policies do. However, they also lack the ability to use synergy effects in the way vertex solutions \mathbf{x}_0 do.

The gaussian instances used fulfill both of the properties that are favorable for piecewise affine policies. By being based on a unity matrix, \mathbf{A} has relatively large values along the diagonal, leading to the existence of good universal recourse decisions. Additionally, the relatively small non-negative entries on the non-diagonals lead to synergy effects for uncertainty realizations with many small values. Demand covering instances however, do not fulfill these properties. The question on how to redirect demand optimally, heavily depends on the demand observed at other locations. Also, the only synergistic effects that can be used solely emerge when multiple demands occur at the same location in a single planning period.

On general instances in practice, we would thus not expect to see the same performance improvements that could be observed on our gaussian benchmark instance. Still, piecewise affine policies find solutions by orders of magnitude faster than affine policies and achieve good results throughout all benchmark instances with hypersphere uncertainty. Additionally, Properties (a) and (b) give intuitive criteria on when strong objective improvements over affine policies can be expected.

6 Conclusion

In this work we presented piecewise affine policies for multi-stage adjustable robust optimization. We construct these policies by carefully approximating uncertainty sets with a dominating polytope, which yields a new problem that we efficiently solve with a linear program. By making use of the problem's structure, we then extend solutions for the new problem with approximated uncertainty to solutions for the original problem. We show strong approximation bounds for our policies that extend many previously best known bounds for two-stage ARO to its multi-stage setting. By doing so, we contribute towards closing the gap between the state of the art for two-stage and multi-stage ARO. Moreover, the new bounds are - to the best of our knowledge - the first bounds shown for the general multi-stage ARO Problem.

In two numerical experiments we find that our policies find solutions by a factor of 10 to 1,000 faster than affine adjustable policies, while mostly yielding similar or even better results. Especially for hypersphere uncertainty sets our new policies perform well and sometimes even outperform affine adjustable policies by up to a factor of two. We observe particularly high improvements on instances that exhibit certain synergistic effects and allow for universal recourse decisions. However, on some instances where few uncertainty dimensions have a high impact on the objective, pure piecewise affine policies perform particularly bad by design, sometimes even worse than constant policies. To mitigate this shortcoming, we present an improvement heuristic that significantly improves the solution quality by rescaling the critical uncertainty dimension.

While this work extends most of the best known approximation results for a relatively general class of ARO problems from the two-stage to the multi-stage setting, it remains an open question, whether other strong two-stage ARO results can be generalized to multi-stage ARO in a similar manner. Answering

this question remains an interesting area for further research.

A Proof of Theorem 1

Proof. We split the proof in two parts. First, we handle the cases where at least $Z_{AR}(\mathcal{U})$ or $Z_{AR}(\hat{\mathcal{U}})$ is negative and show that this already implies that both $Z_{AR}(\mathcal{U})$ and $Z_{AR}(\hat{\mathcal{U}})$ are unbounded. Second, we assume $Z_{AR}(\mathcal{U}), Z_{AR}(\hat{\mathcal{U}}) \geq 0$ are bounded and prove that in this case the desired performance bounds hold.

Part 1: Let $\tilde{\mathcal{U}} \in \{\mathcal{U}, \hat{\mathcal{U}}\}$ such that $Z_{AR}(\tilde{\mathcal{U}}) < 0$. Then, there exists a solution $\tilde{\mathbf{x}}$ such that

$$\max_{\xi \in \tilde{\mathcal{U}}} \mathbf{c}^\top \tilde{\mathbf{x}}(\xi) < 0.$$

Now, assume that $Z_{AR}(\mathcal{U})$ or $Z_{AR}(\hat{\mathcal{U}})$ is bounded and let $\bar{\mathcal{U}} \in \{\mathcal{U}, \hat{\mathcal{U}}\}$ such that $Z_{AR}(\bar{\mathcal{U}})$ is bounded with an optimal solution $\bar{\mathbf{x}}$. Arbitrarily fix one $\tilde{\xi} \in \tilde{\mathcal{U}}$ and consider the constant vector $\tilde{\mathbf{x}}(\tilde{\xi})$. Then $\bar{\mathbf{x}}(\bar{\xi}) + \tilde{\mathbf{x}}(\tilde{\xi})$ is a feasible solution for $Z_{AR}(\bar{\mathcal{U}})$ as for all $\bar{\xi} \in \bar{\mathcal{U}}$:

$$\mathbf{A} \left(\bar{\mathbf{x}}(\bar{\xi}) + \tilde{\mathbf{x}}(\tilde{\xi}) \right) \stackrel{(a)}{\geq} \mathbf{D} \left(\bar{\xi} + \tilde{\xi} \right) + 2\mathbf{d} \stackrel{(b)}{\geq} \mathbf{D}\bar{\xi} + \mathbf{d}.$$

Here, (a) holds because $\tilde{\mathbf{x}}, \bar{\mathbf{x}}$ are feasible solutions and (b) holds as $\tilde{\xi}, \mathbf{D}, \mathbf{d}$ are non-negative. For the objective we then find

$$\max_{\xi \in \bar{\mathcal{U}}} \mathbf{c}^\top \left(\bar{\mathbf{x}}(\bar{\xi}) + \tilde{\mathbf{x}}(\tilde{\xi}) \right) \leq \max_{\xi \in \bar{\mathcal{U}}} \mathbf{c}^\top \bar{\mathbf{x}}(\bar{\xi}) + \max_{\xi \in \tilde{\mathcal{U}}} \mathbf{c}^\top \tilde{\mathbf{x}}(\xi) < Z_{AR}(\bar{\mathcal{U}}).$$

This is a contradiction to $\bar{\mathbf{x}}$ being a minimal solution. Thus, $Z_{AR}(\bar{\mathcal{U}})$ cannot be bounded. We have thus shown that if any of $Z_{AR}(\mathcal{U}), Z_{AR}(\hat{\mathcal{U}})$ are negative, both have to be unbounded.

Part 2: Assume $Z_{AR}(\mathcal{U}), Z_{AR}(\hat{\mathcal{U}}) \geq 0$ are bounded. Let $\hat{\mathbf{x}}$ be an optimal solution for $Z_{AR}(\hat{\mathcal{U}})$. Furthermore, let $\mathbf{h}: \mathcal{U} \rightarrow \hat{\mathcal{U}}$ be the domination function from Definition 1. We claim that $\tilde{\mathbf{x}} := \hat{\mathbf{x}} \circ \mathbf{h}$ is a feasible solution for $Z_{AR}(\mathcal{U})$.

First, we see that by the definition of \mathbf{h} the solution $\tilde{\mathbf{x}}$ fulfills the nonanticipativity requirements. Specifically, we have

$$\tilde{\mathbf{x}}(\xi) = \hat{\mathbf{x}}(\mathbf{h}(\xi)) = \left(\hat{\mathbf{x}}^1 \left(\mathbf{h}^1(\underline{\xi}^1) \right), \dots, \hat{\mathbf{x}}^T \left(\mathbf{h}^T(\underline{\xi}^T) \right) \right)$$

where the decisions in stage t depend only on the uncertainty revealed up to that stage.

For the constraints, we find

$$\mathbf{A}\tilde{\mathbf{x}}(\xi) = \hat{\mathbf{x}}(\mathbf{h}(\xi)) \stackrel{(a)}{\geq} \mathbf{D}\mathbf{h}(\xi) + \mathbf{d} \stackrel{(b)}{\geq} \mathbf{D}\xi + \mathbf{d}.$$

Here, (a) follows from the feasibility of $\hat{\mathbf{x}}$ and (b) follows from the definition of \mathbf{h} and \mathbf{D}, \mathbf{d} being non-

negative. Thus, $\tilde{\mathbf{x}}$ is a well-defined feasible solution for $Z_{AR}(\mathcal{U})$ and we have

$$Z_{AR}(\mathcal{U}) \leq \max_{\boldsymbol{\xi} \in \mathcal{U}} \mathbf{c}^\top \tilde{\mathbf{x}}(\boldsymbol{\xi}) \leq \max_{\hat{\boldsymbol{\xi}} \in \hat{\mathcal{U}}} \mathbf{c}^\top \hat{\mathbf{x}}(\hat{\boldsymbol{\xi}}) = Z_{AR}(\hat{\mathcal{U}}).$$

For the other direction let \mathbf{x}^* be an optimal solution for $Z_{AR}(\mathcal{U})$. Then, for all $\hat{\boldsymbol{\xi}} \in \hat{\mathcal{U}}$ we have $\frac{1}{\beta} \hat{\boldsymbol{\xi}} \in \mathcal{U}$ by definition of β . We define $\tilde{\mathbf{x}}(\hat{\boldsymbol{\xi}}) := \beta \mathbf{x}^* \left(\frac{1}{\beta} \hat{\boldsymbol{\xi}} \right)$ and find

$$\mathbf{A} \tilde{\mathbf{x}}(\hat{\boldsymbol{\xi}}) = \beta \mathbf{A} \mathbf{x}^* \left(\frac{1}{\beta} \hat{\boldsymbol{\xi}} \right) \stackrel{(a)}{\geq} \beta \left(\frac{1}{\beta} \mathbf{D} \hat{\boldsymbol{\xi}} + \mathbf{d} \right) \stackrel{(b)}{\geq} \mathbf{D} \hat{\boldsymbol{\xi}} + \mathbf{d}$$

where (a) follows from the feasibility of \mathbf{x}^* and (b) from the non-negativity of \mathbf{d} . Thus, $\tilde{\mathbf{x}}$ is a well-defined feasible solution for $Z_{AR}(\hat{\mathcal{U}})$ and we have

$$Z_{AR}(\hat{\mathcal{U}}) \leq \max_{\hat{\boldsymbol{\xi}} \in \hat{\mathcal{U}}} \mathbf{c}^\top \tilde{\mathbf{x}}(\hat{\boldsymbol{\xi}}) \leq \beta \max_{\boldsymbol{\xi} \in \mathcal{U}} \mathbf{c}^\top \mathbf{x}^*(\boldsymbol{\xi}) = \beta Z_{AR}(\mathcal{U}).$$

Having shown both inequalities this concludes the proof. \square

B Proof of Lemma 1

Proof. We begin by showing $Z_{AR}(\hat{\mathcal{U}}) \geq Z_{LP}(\hat{\mathcal{U}})$. Let \mathbf{x} be an optimal solution of $Z_{AR}(\hat{\mathcal{U}})$. Then, (\mathbf{x}_i, z) is a valid solution for $Z_{LP}(\hat{\mathcal{U}})$, where \mathbf{x}_i, z are defined by

$$\begin{aligned} \mathbf{x}_i &:= \mathbf{x}(\mathbf{v}_i) & \forall i \in \{0, \dots, m\}, \\ z &:= \max_{i \in \{0, \dots, m\}} \mathbf{c}^\top \mathbf{x}_i. \end{aligned}$$

The first set of LP constraints holds by definition of z . The second set of constraints holds as \mathbf{x} is a solution of $Z_{AR}(\hat{\mathcal{U}})$ and $\mathbf{v}_i \in \hat{\mathcal{U}}$, and the last set of constraints holds by nonanticipativity of \mathbf{x} and the definition of \mathbf{v}_i . Furthermore, we find

$$Z_{LP}(\hat{\mathcal{U}}) \leq z = \max_{i \in \{0, \dots, m\}} \mathbf{c}^\top \mathbf{x}_i \leq \max_{\boldsymbol{\xi} \in \hat{\mathcal{U}}} \mathbf{c}^\top \mathbf{x}(\boldsymbol{\xi}) = Z_{AR}(\hat{\mathcal{U}}).$$

For the other direction let (\mathbf{x}_i, z) be an optimal solution for $Z_{LP}(\hat{\mathcal{U}})$. Define $\lambda_i(\boldsymbol{\xi})$ for each $\boldsymbol{\xi} \in \hat{\mathcal{U}}$ by:

$$\lambda_i(\boldsymbol{\xi}) := \frac{(\boldsymbol{\xi} - \mathbf{v}_0)_i}{\rho_i}.$$

For ease of notation we will in the following drop the explicit dependence on $\boldsymbol{\xi}$ and use λ_i . We directly find

$$\sum_{i=1}^m \lambda_i \leq 1$$

for all $\boldsymbol{\xi} \in \hat{\mathcal{U}}$, as by definition of $\hat{\mathcal{U}}$ each $\boldsymbol{\xi}$ is a convex combination of $\mathbf{v}_0, \dots, \mathbf{v}_m$. Also note that λ_i only depends on the i^{th} uncertainty. Using this we can define a nonanticipative decision vector

$\mathbf{x}(\underline{\xi}) = (\mathbf{x}^1(\underline{\xi}^1), \dots, \mathbf{x}^T(\underline{\xi}^T))$ by

$$\mathbf{x}^t(\underline{\xi}^t) := \sum_{i \in \underline{I}^t} \lambda_i \mathbf{x}_i^t + \left(1 - \sum_{i \in \underline{I}^t} \lambda_i\right) \mathbf{x}_0^t$$

where \underline{I}^t is the index set corresponding to the uncertainties up to stage t .

By the nonanticipativity constraints of (3) we have $\mathbf{x}_i^t = \mathbf{x}_0^t$ for all $1 \leq t \leq T$ and $i \in I^T \setminus I^t$. Thus,

$$\mathbf{x}^t(\underline{\xi}^t) = \sum_{i \in \underline{I}^t} \lambda_i \mathbf{x}_i^t + \left(1 - \sum_{i \in \underline{I}^t} \lambda_i\right) \mathbf{x}_0^t = \sum_{i=1}^m \lambda_i \mathbf{x}_i^t + \left(1 - \sum_{i=1}^m \lambda_i\right) \mathbf{x}_0^t,$$

which shows that \mathbf{x} is a convex combination of $\mathbf{x}_0, \dots, \mathbf{x}_m$ by

$$\mathbf{x}(\underline{\xi}) = \sum_{i=1}^m \lambda_i \mathbf{x}_i + \left(1 - \sum_{i=1}^m \lambda_i\right) \mathbf{x}_0.$$

Using this, we find

$$\mathbf{A}\mathbf{x}(\underline{\xi}) = \sum_{i=1}^m \lambda_i \mathbf{A}\mathbf{x}_i + \left(1 - \sum_{i=1}^m \lambda_i\right) \mathbf{A}\mathbf{x}_0 \stackrel{(a)}{\geq} \sum_{i=1}^m \lambda_i \mathbf{D}\mathbf{v}_i + \left(1 - \sum_{i=1}^m \lambda_i\right) \mathbf{D}\mathbf{v}_0 + \mathbf{d} \stackrel{(b)}{=} \mathbf{D}\underline{\xi} + \mathbf{d}.$$

Here, (a) holds because all \mathbf{x}_i are valid solutions for (3), and (b) holds as by definition of λ_i we have $\underline{\xi} = \sum_{i=1}^m \lambda_i \mathbf{v}_i + (1 - \sum_{i=1}^m \lambda_i) \mathbf{v}_0$.

Now we find the missing inequality

$$\begin{aligned} Z_{AR}(\hat{\mathcal{U}}) &\stackrel{(a)}{\leq} \max_{\underline{\xi} \in \hat{\mathcal{U}}} \mathbf{c}^\top \mathbf{x}(\underline{\xi}) = \max_{\underline{\xi} \in \hat{\mathcal{U}}} \sum_{i=1}^m \lambda_i \mathbf{c}^\top \mathbf{x}_i + \left(1 - \sum_{i=1}^m \lambda_i\right) \mathbf{c}^\top \mathbf{x}_0 \\ &\stackrel{(b)}{\leq} \max_{\underline{\xi} \in \hat{\mathcal{U}}} \max_{i \in \{0, \dots, m\}} \mathbf{c}^\top \mathbf{x}_i = \max_{i \in \{0, \dots, m\}} \mathbf{c}^\top \mathbf{x}_i = Z_{LP}(\hat{\mathcal{U}}), \end{aligned}$$

where we use for (a) that \mathbf{x} is a valid solution for $Z_{AR}(\hat{\mathcal{U}})$. For (b), we use that \mathbf{x} is a convex combination of $\mathbf{x}_0, \dots, \mathbf{x}_m$ with convex coefficients $(1 - \sum_{i=1}^m \lambda_i), \lambda_1, \dots, \lambda_m$. The convex sum is upper bounded by its largest summand $\mathbf{c}^\top \mathbf{x}_i$. We can thus take the maximum over all these $\mathbf{c}^\top \mathbf{x}_i$, which equals the LP solution value $Z_{LP}(\hat{\mathcal{U}})$. \square

C Proof of Proposition 2

Proof. The main idea is to find μ, ρ such that (7) is always fulfilled and β is minimized. Recall that β is given by the minimal value such that $\frac{1}{\beta} \mathbf{v}_i \in \mathcal{U}$ for all \mathbf{v}_i . In the case of the hypersphere uncertainty set this is given when $\|\mathbf{v}_i\|_2^2 \leq \beta^2$. As $\mathbf{v}_{i>0}$ is given by $\mathbf{v}_i = \mathbf{v}_0 + \rho \mathbf{e}_i$ and $\mathbf{v}_0 = \mu \mathbf{e}$ we find:

$$\|\mathbf{v}_0\|_2^2 \leq \|\mathbf{v}_{i>0}\|_2^2 = (m-1)\mu^2 + (\mu + \rho)^2 = m\mu^2 + 2\mu\rho + \rho^2. \quad (9)$$

This is the term for β^2 that we want to minimize in the following.

Using Lemma 2 the left hand side in (7) becomes:

$$\frac{1}{\rho} \max_{j \in \{1, \dots, m\}} \sum_{i=1}^j (\gamma(j) - \mu)_+ \stackrel{(a)}{=} \frac{1}{\rho} \max_{j \in \{1, \dots, m\}} j(\gamma(j) - \mu)_+ \stackrel{(b)}{=} \frac{1}{\rho} \max_{j \in \{1, \dots, m\}} j(\gamma(j) - \mu).$$

For (a) we use that all summands are the same and for (b) we see that any negative results will be eliminated by the maximization.

Using the properties of the hypersphere uncertainty set again, we find $\gamma(j) = \frac{1}{\sqrt{j}}$ for all j . Substituting $\gamma(j)$ in the above equation we find Property (7) to become:

$$\max_{j \in \{1, \dots, m\}} \sqrt{j} - j\mu \leq \rho.$$

The maximum of the left hand side is taken for $j = \frac{1}{4\mu^2}$ and for any optimal choice of μ, ρ the inequality will be tight. Thus we find $\rho = \frac{1}{4\mu}$.

Substituting ρ in Equation (9), we get:

$$\beta = \sqrt{m\mu^2 + \frac{1}{2} + \frac{1}{16\mu^2}}.$$

This is minimized by $\mu = \frac{1}{2\sqrt[4]{m}}$, which gives $\rho = \frac{\sqrt[4]{m}}{2}$ and concludes the proof. \square

D Proof of Proposition 3

Proof. Consider the following two-stage instance of Problem (1):

$$\begin{aligned} Z(\mathcal{U}) &= \min_{\alpha, \mathbf{x}(\boldsymbol{\xi})} \max_{\boldsymbol{\xi} \in \mathcal{U}} \sqrt{m}\alpha + \mathbf{e}^\top \mathbf{x}(\boldsymbol{\xi}) \\ \alpha \mathbf{e} + \mathbf{x} &\geq \boldsymbol{\xi} && \forall \boldsymbol{\xi} \in \mathcal{U} \\ \alpha, \mathbf{x} &\geq 0, \end{aligned}$$

where $\mathcal{U} = \{\boldsymbol{\xi} \mid \|\boldsymbol{\xi}\|_2^2 \leq 1\} \subset [0, 1]^m$ is the hypersphere uncertainty set, $\alpha \in \mathbb{R}$ is the first stage decision variable and $\mathbf{x} \in \mathbb{R}^m$ are the second stage decisions that may depend on the uncertainty realization $\boldsymbol{\xi}$.

We begin by finding a lower bound for affine policies on this problem. By symmetry of the problem every affine adjustable solution $\mathbf{x}(\boldsymbol{\xi})$ will be of the form

$$x_i = a\xi_i + b \sum_{j \neq i} \xi_j + c,$$

for some $a, b, c \in \mathbb{R}$. Let $\boldsymbol{\xi} \in \mathcal{U}$ and for some $i \leq m$ define the vector $\boldsymbol{\xi}'$ by $\boldsymbol{\xi}' := \boldsymbol{\xi} - \xi_i \mathbf{e}_i$. Then $\boldsymbol{\xi}'$ is also in \mathcal{U} and by $x_i(\boldsymbol{\xi}') \geq 0$ we find that every feasible solution has to fulfill

$$b \sum_{j \neq i} \xi_j + c \geq 0 \tag{10}$$

for all $\boldsymbol{\xi} \in \mathcal{U}$. Additionally, by $\alpha \mathbf{e} + \mathbf{x}(\mathbf{e}_i) \geq \mathbf{e}_i$ every feasible solution also fulfills

$$\alpha + a + c \geq 1. \quad (11)$$

Using this we obtain a lower bound for the maximum over \mathcal{U} in the objective function as follows

$$\begin{aligned} & \max_{\boldsymbol{\xi} \in \mathcal{U}} \sqrt{m}\alpha + \sum_i \left(a\xi_i + b \sum_{j \neq i} \xi_j + c \right) \\ & \stackrel{(a)}{\geq} \max_{\boldsymbol{\xi} \in \mathcal{U}} \sqrt{m}\alpha + a \sum_i \xi_i = \max_{\boldsymbol{\xi} \in \mathcal{U}} \sqrt{m}\alpha + a \mathbf{e}^\top \boldsymbol{\xi} \\ & \stackrel{(b)}{=} \sqrt{m}\alpha + \sqrt{m}a \stackrel{(c)}{\geq} \sqrt{m}\alpha + \sqrt{m}(1 - \alpha - c) = \sqrt{m}(1 - c). \end{aligned}$$

Here (a) follows from Equation (10), (b) follows from the Cauchy-Schwarz inequality, and (c) follows from Equation (11). On the other hand, using $\mathbf{0} \in \mathcal{U}$ the maximum can also be lower bounded by

$$\sqrt{m}\alpha + \sum_i c = \sqrt{m}\alpha + mc \geq mc.$$

Thus we find

$$\begin{aligned} Z_{AFF}(\mathcal{U}) &= \min_{\alpha, a, b, c} \max_{\boldsymbol{\xi} \in \mathcal{U}} \sqrt{m}\alpha + \sum_i \left(a\xi_i + b \sum_{j \neq i} \xi_j + c \right) \\ &\geq \min_c \max(\sqrt{m}(1 - c), mc) \stackrel{(a)}{=} \sqrt{m} - \frac{m}{m + \sqrt{m}} \geq \sqrt{m} - 1, \end{aligned}$$

where (a) holds as the maximum is taken at equality of the two terms, which is given by $c = \frac{\sqrt{m}}{m + \sqrt{m}}$.

Having found a lower bound for affine policies, we will now give an upper bound for optimal policies. Consider the policy given by $\alpha = \frac{1}{\sqrt{m}}$ and $\mathbf{x} = (\boldsymbol{\xi} - \alpha \mathbf{e})_+$. By construction this is feasible and we find

$$\begin{aligned} Z_{OPT}(\mathcal{U}) &\leq \max_{\boldsymbol{\xi} \in \mathcal{U}} \sqrt{m}\alpha + \mathbf{e}^\top (\boldsymbol{\xi} - \alpha \mathbf{e})_+ \\ &\stackrel{(a)}{=} \max_{0 \leq k \leq m} 1 + k \left(\frac{1}{\sqrt{k}} - \frac{1}{\sqrt{m}} \right) = \max_{0 \leq k \leq m} 1 + \sqrt{k} - \frac{k}{\sqrt{m}} \stackrel{(b)}{=} \sqrt[4]{m}. \end{aligned}$$

Here (a) follows from Lemma 2 and (b) follows as the maximum over k is taken for $k = \sqrt{m}$. To finish the proof we find the optimality ratio which is given by

$$\frac{Z_{AFF}(\mathcal{U})}{Z_{OPT}(\mathcal{U})} \geq \frac{\sqrt{m} - 1}{\sqrt[4]{m}} = \sqrt[4]{m} - \frac{1}{\sqrt[4]{m}}.$$

□

E Proof of Proposition 4

Proof. We proceed analogously as in the proof for Proposition 2. Again using the definition of \mathcal{U} and \mathbf{v}_i we find $\frac{1}{\beta}\mathbf{v}_i \in \mathcal{U}$ to be fulfilled when

$$\beta = \min \left(\frac{m}{k}\mu + \frac{1}{k}\rho, \mu + \rho \right). \quad (12)$$

Analogously, the left hand side in (7) becomes:

$$\frac{1}{\rho} \max_{j \in \{1, \dots, m\}} j(\gamma(j) - \mu).$$

Using the properties of the budgeted uncertainty set we find $\gamma(j) = \frac{k}{j}$ for all $j \geq k$ and $\gamma(j) = 1$ for all $j \leq k$. Inserting $\gamma(j)$ into the above, we find Property (7) to become:

$$\max_{j \in \{1, \dots, m\}} j \left(\min \left(1, \frac{k}{j} \right) - \mu \right) \leq \rho.$$

As for $\mu \geq 1$ the vector \mathbf{v}_0 would already dominate every uncertainty realization in \mathcal{U} , we can assume $\mu \leq 1$. With this the maximum on the left hand side is taken for $j = k$ and we find the minimal possible ρ to be $\rho = k - k\mu$.

Using this ρ together with β from Equation (12), we obtain:

$$\beta = \min \left(\left(\frac{m}{k} - 1 \right) \mu + 1, k - (k-1)\mu \right).$$

As the left term in the minimization is linearly decreasing in μ and the right term is linearly increasing in μ , the minimum is taken at equality. This is the case when $\mu = \frac{k(k-1)}{m+k(k-2)}$, which gives $\rho = \frac{k(m-k)}{m+k(k-2)}$. \square

F Proof of Proposition 5

To prove Proposition 5 we first introduce a more general form of piecewise affine policies. Recall that affine policies are given by $\mathbf{x}^t(\underline{\xi}^t) = \mathbf{P}^t \underline{\xi}^t + \mathbf{q}^t$, with decision variables \mathbf{P}^t and \mathbf{q}^t . By introducing additional decision variables $\underline{\zeta}$ of the same dimension as the uncertainty set, we extend affine policies to *fully piecewise affine policies* (FPAP)

$$\mathbf{x}^i(\underline{\xi}^t) = \mathbf{P}^t (\underline{\xi}^t - \underline{\zeta}^t)_+ + \mathbf{q}^t,$$

where we use the same notation for $\underline{\zeta}$, as for $\underline{\xi}$. By extending the decision matrices \mathbf{P}^t with zeroes and concatenating them to a single decision matrix

$$\mathbf{P} = \begin{pmatrix} \mathbf{P}^1 & \mathbf{0} & \mathbf{0} & \dots & \mathbf{0} \\ & \mathbf{P}^2 & \mathbf{0} & \dots & \mathbf{0} \\ & & \vdots & & \\ & & & & \mathbf{P}^T \end{pmatrix},$$

we get the compact expressions $\mathbf{x}(\boldsymbol{\xi}) = \mathbf{P}\boldsymbol{\xi} + \mathbf{q}$ and $\mathbf{x}(\boldsymbol{\xi}) = \mathbf{P}(\boldsymbol{\xi} - \boldsymbol{\zeta})_+ + \mathbf{q}$ for affine and fully piecewise affine policies, respectively. We can show that these fully piecewise affine adjustable policies are in fact a generalization of the piecewise affine policies we introduce in this work.

Lemma 4. *Consider Problem (1). Let Z_{PAP} be the optimal value found by our piecewise affine policy and let Z_{FPAP} be the optimal value found by a fully piecewise affine policy where the right hand side uncertainty is replaced by $\mathbf{D}((\boldsymbol{\xi} - \boldsymbol{\zeta})_+ + \boldsymbol{\zeta}) + \mathbf{d}$. Then*

$$Z_{FPAP} \leq Z_{PAP}.$$

Proof. We prove the inequality by showing that every feasible piecewise affine policy is a feasible fully piecewise affine policy. Let $\mathbf{v}_0, \dots, \mathbf{v}_m$ be the chosen vertices of the piecewise affine policy and let $\mathbf{x}_0, \dots, \mathbf{x}_m$ be the vertex solutions received from solving LP (3). Then the piecewise affine policy is given by Equation (6) and we find

$$\begin{aligned} \mathbf{x}(\boldsymbol{\xi}) &= \sum_{i=1}^m \lambda_i(\boldsymbol{\xi}) \mathbf{x}_i + \left(1 - \sum_{i=1}^m \lambda_i(\boldsymbol{\xi})\right) \mathbf{x}_0 \\ &= \sum_{i=1}^m \lambda_i(\boldsymbol{\xi}) (\mathbf{x}_i - \mathbf{x}_0) + \mathbf{x}_0 \\ &\stackrel{(a)}{=} \sum_{i=1}^m \frac{((\boldsymbol{\xi} - \mathbf{v}_0)_+)_i}{\rho_i} (\mathbf{x}_i - \mathbf{x}_0) + \mathbf{x}_0 \\ &\stackrel{(b)}{=} \left(\frac{1}{\rho_1} (\mathbf{x}_1 - \mathbf{x}_0), \dots, \frac{1}{\rho_m} (\mathbf{x}_m - \mathbf{x}_0)\right) (\boldsymbol{\xi} - \mathbf{v}_0)_+ + \mathbf{x}_0. \end{aligned}$$

Here (a) follows from the definition of λ_i in Section 2.2 and (b) follows by replacing the sum with a matrix-vector multiplication. By the nonanticipativity Constraint (3d) the vector $\mathbf{x}_i - \mathbf{x}_0$ is zero for all uncertainty dimensions associated to later stages. Thus setting

$$\begin{aligned} \mathbf{P} &:= \left(\frac{1}{\rho_1} (\mathbf{x}_1 - \mathbf{x}_0), \dots, \frac{1}{\rho_m} (\mathbf{x}_m - \mathbf{x}_0)\right) \\ \boldsymbol{\zeta} &:= \mathbf{v}_0 \\ \mathbf{q} &:= \mathbf{x}_0 \end{aligned}$$

yields a valid fully piecewise affine policy that is identical to the initial piecewise affine policy. Even with the modified right hand side, as demanded in the Lemma, the constructed fully piecewise affine policy fulfills all constraints, as by construction of the policy, and definition of the dominating function \mathbf{h} :

$$\mathbf{A}\mathbf{x}(\boldsymbol{\xi}) \geq \mathbf{D}\mathbf{h}(\boldsymbol{\xi}) + \mathbf{d} = \mathbf{D}((\boldsymbol{\xi} - \boldsymbol{\zeta})_+ + \boldsymbol{\zeta}) + \mathbf{d}.$$

□

Even though fully piecewise affine policies are slightly more flexible than affine policies in general, both are equivalent on Problem (1) under some additional assumptions.

Lemma 5. *Consider Problem (1) with budgeted uncertainty and integer budget. Let Z_{FPAP} be the optimal value found by a fully piecewise affine policy with modified right hand side uncertainty as in*

Lemma 4 and let Z_{AFF} be the optimal value found by an affine policy. Then

$$Z_{AFF} = Z_{FPAP}.$$

For the proof of Lemma 5 we first need the following helpful result that allows us to construct a robust counterpart of Problem (1) with fully piecewise affine adjustable policies in a similar manner as robust counterparts for affine policies are constructed.

Lemma 6. Let $\mathbf{c} \in \mathbb{R}^m$, $\zeta \in [0, 1]^m$ and $k \in \mathbb{N}_+$. Then the following two problems yield the same objective.

$$\begin{aligned} \max_{\xi} \quad & \mathbf{c}^\top (\xi - \zeta)_+ \\ & \mathbf{e}^\top \xi \leq k \\ & \mathbf{0} \leq \xi \leq \mathbf{e} \end{aligned} \tag{13}$$

and

$$\begin{aligned} \min_{\alpha, \beta} \quad & \mathbf{e}^\top \beta + k\alpha \\ & \beta + \alpha \mathbf{e} \geq \text{diag}(\mathbf{e} - \zeta) \mathbf{c} \\ & \alpha \geq 0, \quad \beta \geq \mathbf{0}. \end{aligned} \tag{14}$$

Proof. Let I^k be the indices of the largest k values $c_i(1 - \zeta_i)$ and let \mathbf{e}_{I^k} be the indicator vector of I^k . Then for Problem (13) we find the following set of inequalities.

$$\max \{ \mathbf{c}^\top (\xi - \zeta)_+ \mid \mathbf{e}^\top \xi \leq k, \mathbf{0} \leq \xi \leq \mathbf{e} \} \tag{15}$$

$$\stackrel{(a)}{\leq} \max \{ \mathbf{c}^\top \text{diag}(\mathbf{e} - \zeta) \xi \mid \mathbf{e}^\top \xi \leq k, \mathbf{0} \leq \xi \leq \mathbf{e} \} \tag{16}$$

$$\stackrel{(b)}{\leq} \mathbf{c}^\top \text{diag}(\mathbf{e} - \zeta) \mathbf{e}_{I^k}$$

$$\stackrel{(c)}{\leq} \mathbf{c}^\top (\mathbf{e}_{I^k} - \zeta)_+$$

$$\stackrel{(d)}{\leq} \max \{ \mathbf{c}^\top (\xi - \zeta)_+ \mid \mathbf{e}^\top \xi \leq k, \mathbf{0} \leq \xi \leq \mathbf{e} \} \tag{17}$$

Here (a) follows from $(\xi - \zeta)_+ \leq \text{diag}(\mathbf{e} - \zeta) \xi$. Inequality (b) follows as \mathbf{e}_{I^k} is an optimal choice for Problem (16). For (c), observe that $\text{diag}(\mathbf{e} - \zeta) \mathbf{e}_{I^k} = (\mathbf{e}_{I^k} - \zeta)_+$. Finally, (d) follows, as \mathbf{e}_{I^k} is a feasible choice for Problem (17). As (15) and (17) are the same problem all inequalities are in fact equalities.

To conclude the proof note that Problem (16) is a linear optimization problem, where taking the dual yields Problem (14). \square

Having this result, we now show Lemma 5.

Proof of Lemma 5. To prove this result we show that the robust counterparts of the two policies on Problem (1) with integer budget uncertainty are equivalent. Let $k \in \mathbb{N}_+$ be the budget.

Then Problem (1) with affine policies becomes

$$\begin{aligned} Z_{AFF}(\mathcal{U}) = \min_{\mathbf{P}, \mathbf{q}} \quad & \max_{\xi} (\mathbf{c}^\top \mathbf{P} \xi \mid \mathbf{0} \leq \xi \leq \mathbf{e}, \mathbf{e}^\top \xi \leq k) + \mathbf{c}^\top \mathbf{q} \\ \text{s.t.} \quad & \max_{\xi} ((\mathbf{D}_i - \mathbf{A}_i \mathbf{P}) \xi \mid \mathbf{0} \leq \xi \leq \mathbf{e}, \mathbf{e}^\top \xi \leq k) \leq \mathbf{A}_i \mathbf{q} - d_i \quad \forall i \in \{1, \dots, l\}, \end{aligned}$$

where \mathbf{A}_i and \mathbf{D}_i are the i^{th} rows of \mathbf{A} and \mathbf{D} , respectively, and d_i is the i^{th} entry of \mathbf{d} . Dualizing the subproblems we find this to be equivalent to

$$\begin{aligned}
Z_{AFF}(\mathcal{U}) &= \min_{\mathbf{P}, \mathbf{q}, \alpha^0, \dots, \alpha^l, \beta^0, \dots, \beta^l} \mathbf{e}^\top \beta^0 + k\alpha^0 + \mathbf{c}^\top \mathbf{q} \\
\text{s.t. } & \beta^0 + \alpha^0 \mathbf{e} \geq (\mathbf{c}^\top \mathbf{P})^\top \\
& \mathbf{e}^\top \beta^i + k\alpha^i \leq \mathbf{A}_i \mathbf{q} - d_i & \forall i \in \{1, \dots, l\} \\
& \beta^i + \alpha^i \mathbf{e} \geq (\mathbf{D}_i - \mathbf{A}_i \mathbf{P})^\top & \forall i \in \{1, \dots, l\} \\
& \alpha^i \geq 0, \quad \beta^i \geq \mathbf{0} & \forall i \in \{1, \dots, l\}.
\end{aligned} \tag{18}$$

Similarly Problem (1) with fully piecewise affine policies and modified right hand side becomes

$$\begin{aligned}
Z_{AFF}(\mathcal{U}) &= \min_{\mathbf{P}, \mathbf{q}, \zeta} \max_{\xi} (\mathbf{c}^\top \mathbf{P}(\xi - \zeta)_+ | \mathbf{0} \leq \xi \leq \mathbf{e}, \mathbf{e}^\top \xi \leq k) + \mathbf{c}^\top \mathbf{q} \\
\text{s.t. } & \max_{\xi} ((\mathbf{D}_i \mathbf{A}_i \mathbf{P})(\xi - \zeta)_+ | \mathbf{0} \leq \xi \leq \mathbf{e}, \mathbf{e}^\top \xi \leq k) \leq \mathbf{A}_i \mathbf{q} - \mathbf{D}_i \zeta - d_i \quad \forall i \in \{1, \dots, l\} \\
& \mathbf{0} \leq \zeta \leq \mathbf{e}.
\end{aligned}$$

Using Lemma 6 on the subproblems we find this to be equivalent to

$$\begin{aligned}
Z_{AFF}(\mathcal{U}) &= \min_{\mathbf{P}, \mathbf{q}, \zeta, \alpha^0, \dots, \alpha^l, \beta^0, \dots, \beta^l} \mathbf{e}^\top \beta^0 + k\alpha^0 + \mathbf{c}^\top \mathbf{q} & (19a) \\
\text{s.t. } & \beta^0 + \alpha^0 \mathbf{e} \geq (\mathbf{c}^\top \mathbf{P} \text{diag}(\mathbf{e} - \zeta))^\top & (19b) \\
& \mathbf{e}^\top \beta^i + k\alpha^i \leq \mathbf{A}_i \mathbf{q} - \mathbf{D}_i \zeta - d_i & \forall i \in \{1, \dots, l\} & (19c) \\
& \beta^i + \alpha^i \mathbf{e} \geq ((\mathbf{D}_i - \mathbf{A}_i \mathbf{P}) \text{diag}(\mathbf{e} - \zeta))^\top & \forall i \in \{1, \dots, l\} & (19d) \\
& \alpha^i \geq 0, \quad \beta^i \geq \mathbf{0} & \forall i \in \{1, \dots, l\} & (19e) \\
& \mathbf{0} \leq \zeta \leq \mathbf{e}. & (19f)
\end{aligned}$$

Reformulating Constraints (19c) and (19d) yields:

$$\begin{aligned}
\mathbf{e}^\top (\beta^i + \mathbf{D}_i \text{diag}(\zeta)) + k\alpha^i &\leq \mathbf{A}_i \mathbf{q} - d_i & \forall i \in \{1, \dots, l\} \\
\beta^i + \mathbf{D}_i \text{diag}(\zeta) + \alpha^i \mathbf{e} &\geq (\mathbf{D}_i - \mathbf{A}_i \mathbf{P} \text{diag}(\mathbf{e} - \zeta))^\top & \forall i \in \{1, \dots, l\}.
\end{aligned}$$

First note that as \mathbf{D} and ζ are both non-negative we can replace β^i by $\beta'^i := \beta^i + \mathbf{D}_i \text{diag}(\zeta)$ and set ζ to zero without changing the problem. Thus, we can eliminate ζ from the problem. Next we can substitute $\mathbf{P} \text{diag}(\mathbf{e} - \zeta)$ with a new free variable \mathbf{P}' without changing the problem, as \mathbf{P} is a free variable. Doing both of these steps transforms Problem (19) into Problem (18). Thus the robust counterparts of affine policies and fully piecewise affine policies with modified right hand side are equivalent. \square

Having shown these results, Proposition 5 directly follows by combining Lemmas 4 and 5. Note, that Lemma 6, which is crucial for the construction of tractable reformulations for fully piecewise affine policies, heavily depends on the linear structure of budgeted uncertainty and the existence of integer optimal solutions. Thus the results cannot easily be extended to other settings and in general one cannot hope for tractable reformulations of FPAPs.

G Construction of Demand Covering Instances

Mathematically, the demand covering problem can be stated as a robust LP, which we describe in Problem (20), using the notation given in Table 2. Here Objective (20a) minimizes the sum of resource costs and lost demand costs due to delay and relocation. Constraints (20b) ensure that all demands are fulfilled, delayed, or relocated and Constraints (20c) upper bound the allocated resources in each planning period by the total number of available resources R .

Sets & Elements	
$l \in \mathcal{L}$	locations
$p \in \mathcal{P}$	planning periods
$t \in \mathcal{T}$	all $(m^p m^e)$ execution periods
Parameters	
c^R	cost of buying one unit of resources
c^D	cost of loosing one unit of demand
$q_t^d \in [0, 1]$	fraction of demand lost when delayed in period t
$q_{ll'}^r \in [0, 1]$	fraction of demand lost when redirected form location l to l'
Uncertainties	
$p(t)$	planning period containing execution period t
ξ_{pl}	demand at location $l \in \mathcal{L}$ in execution period $p \in \mathcal{T}$
Decision variables	
R	number of resources bought
r_{pl}	number of resources assigned to location $l \in \mathcal{L}$ in planning period $p \in \mathcal{P}$
$s_{ll'}^r$	demand redirected from location l to location l'
s_{tl}^d	demand delayed to the next period

Table 2: Notation for the robust demand covering problem with non-consumed resources and uncertain demands.

$$\min c^R R + c^D \left(\sum_{t \in \mathcal{T}, l \in \mathcal{L}} q_t^d s_{tl}^d + \sum_{t \in \mathcal{T}, l \neq l' \in \mathcal{L}} q_{ll'}^r s_{ll'}^r \right) \quad (20a)$$

$$r_{p(t)l} + s_{tl}^d - (1 - q_{t-1}^d) s_{(t-1)l}^d + \sum_{l' \in \mathcal{L}, l' \neq l} (s_{ll'}^r - (1 - q_{ll'}^r) s_{ll'}^r) \geq \xi_{tl} \quad \forall \xi \in \mathcal{U}, \forall t \in \mathcal{T}, l \in \mathcal{L} \quad (20b)$$

$$\sum_{l \in \mathcal{L}} r_{pl} \leq R \quad \forall p \in \mathcal{P} \quad (20c)$$

$$R, \mathbf{r}, \mathbf{s}^r, \mathbf{s}^d \geq 0 \quad (20d)$$

To construct our instances we draw $m^l \in \{2, 4, 6, 8, 10\}$ locations at uniform random integer positions in the square $[0, 2\lfloor \sqrt{m^l} \rfloor + 1]^2$. In each of $m^p \in \{1, 3, 5, 7\}$ planning periods we consider $m^e = 8$ execution periods corresponding to the hours in a working day. We assume that a fraction $q_{tl}^d = 0.1$ of the demand is lost, when deferred to a later execution period; we consider a doubled loss-rate ($q_{tl}^d = 0.2$) if this results in a deference to another planning period. Similarly, a fraction of the demand is lost, when assigned to another location. We assume this fraction to be negatively correlated to the distance and given by $q_{ll'}^r := \max(0, 1 - 0.02 \cdot \text{dist}(l, l'))$. Demands are given by $\xi_{lt} = d_{lt} \left(1 + \xi_{lt}^p + \frac{1}{2} \xi_{lt}^e\right)$, where d_{lt} is drawn from the normal distribution $\mathcal{N}(10, 4)$. Here the uncertainty vector (ξ^p, ξ^e) is taken from one uncertainty set \mathcal{U} of dimension $m = 2m^l m^p m^e$ resulting in short term uncertainties having a less

severe effect on demands than uncertainties known in advance. Finally we set $c^R = 1$ and choose $c^D \in \{0.1, 0.25, 0.5\}$. For each combination of m^l, m^p we consider 75 instances, where we use each possible values for $c^D \in \{0.1, 0.25, 0.5\}$ in a third of these instances.

References

- E. Adida and G. Perakis. A robust optimization approach to dynamic pricing and inventory control with no backorders. *Mathematical Programming*, 107(1-2):97–129, dec 2006. doi: 10.1007/s10107-005-0681-5.
- M. Ang, Y. F. Lim, and M. Sim. Robust storage assignment in unit-load warehouses. *Management Science*, 58(11):2114–2130, 2012. doi: 10.1287/mnsc.1120.1543.
- D. Bampou and D. Kuhn. Scenario-free stochastic programming with polynomial decision rules. In *IEEE Conference on Decision and Control and European Control Conference*. IEEE, dec 2011. doi: 10.1109/cdc.2011.6161150.
- O. Baron, J. Milner, and H. Naseraldin. Facility location: A robust optimization approach. *Production and Operations Management*, 20(5):772–785, 2011. doi: 10.1111/j.1937-5956.2010.01194.x.
- J. H. Beebe, C. S. Beightler, and J. P. Stark. Stochastic optimization of production planning. *Operations Research*, 16(4):799–818, 1968. doi: 10.1287/opre.16.4.799.
- A. Ben-Tal and A. Nemirovski. Robust solutions of uncertain linear programs. *Operations Research Letters*, 25(1):1–13, 1999. doi: 10.1016/s0167-6377(99)00016-4.
- A. Ben-Tal and A. Nemirovski. Robust optimization - methodology and applications. *Mathematical Programming*, 92(3):453–480, 2002. doi: 10.1007/s101070100286.
- A. Ben-Tal, A. Goryashko, E. Guslitzer, and A. Nemirovski. Adjustable robust solutions of uncertain linear programs. *Mathematical Programming*, 99(2):351–376, 2004. doi: 10.1007/s10107-003-0454-y.
- A. Ben-Tal, B. Golany, A. Nemirovski, and J.-P. Vial. Retailer-supplier flexible commitments contracts: A robust optimization approach. *Manufacturing & Service Operations Management*, 7(3):248–271, 2005. doi: 10.1287/msom.1050.0081.
- A. Ben-Tal, L. E. Ghaoui, and A. Nemirovski. *Robust Optimization*. Princeton University Press, 2009a. ISBN 0691143684.
- A. Ben-Tal, B. Golany, and S. Shtern. Robust multi-echelon multi-period inventory control. *European Journal of Operational Research*, 199(3):922–935, 2009b. doi: 10.1016/j.ejor.2009.01.058.
- A. Ben-Tal, B. D. Chung, S. R. Mandala, and T. Yao. Robust optimization for emergency logistics planning: Risk mitigation in humanitarian relief supply chains. *Transportation Research Part B: Methodological*, 45(8):1177–1189, 2011. doi: 10.1016/j.trb.2010.09.002.

- A. Ben-Tal, O. E. Housni, and V. Goyal. A tractable approach for designing piecewise affine policies in two-stage adjustable robust optimization. *Mathematical Programming*, 182(1-2):57–102, 2020. doi: 10.1007/s10107-019-01385-0.
- D. Bertsimas and H. Bidkhori. On the performance of affine policies for two-stage adaptive optimization: a geometric perspective. *Mathematical Programming*, 153(2):577–594, 2015. doi: 10.1007/s10107-014-0818-5.
- D. Bertsimas and I. Dunning. Multistage robust mixed-integer optimization with adaptive partitions. *Operations Research*, 64(4):980–998, 2016. doi: 10.1287/opre.2016.1515.
- D. Bertsimas and V. Goyal. On the power and limitations of affine policies in two-stage adaptive optimization. *Mathematical Programming*, 134(2):491–531, 2012. doi: 10.1007/s10107-011-0444-4.
- D. Bertsimas, D. A. Iancu, and P. A. Parrilo. Optimality of affine policies in multistage robust optimization. *Mathematics of Operations Research*, 35(2):363–394, 2010. doi: 10.1287/moor.1100.0444.
- D. Bertsimas, V. Goyal, and X. A. Sun. A geometric characterization of the power of finite adaptability in multistage stochastic and adaptive optimization. *Mathematics of Operations Research*, 36(1):24–54, 2011a. doi: 10.1287/moor.1110.0482.
- D. Bertsimas, D. A. Iancu, and P. A. Parrilo. A hierarchy of near-optimal policies for multistage adaptive optimization. *IEEE Transactions on Automatic Control*, 56(12):2809–2824, dec 2011b. doi: 10.1109/tac.2011.2162878.
- D. Bertsimas, V. Goyal, and B. Y. Lu. A tight characterization of the performance of static solutions in two-stage adjustable robust linear optimization. *Mathematical Programming*, 150(2):281–319, 2015. doi: 10.1007/s10107-014-0768-y.
- D. Bertsimas, S. Shtern, and B. Sturt. Technical note—two-stage sample robust optimization. *Operations Research*, 2021. doi: 10.1287/opre.2020.2096.
- X. Chen and Y. Zhang. Uncertain linear programs: Extended affinely adjustable robust counterparts. *Operations Research*, 57(6):1469–1482, dec 2009. doi: 10.1287/opre.1080.0605.
- X. Chen, M. Sim, P. Sun, and J. Zhang. A linear decision-based approximation approach to stochastic programming. *Operations Research*, 56(2):344–357, apr 2008. doi: 10.1287/opre.1070.0457.
- G. B. Dantzig. Linear programming under uncertainty. *Management Science*, 1(3-4):197–206, 1955. doi: 10.1287/mnsc.1.3-4.197.
- E. Delage and D. A. Iancu. Robust multistage decision making. In *The Operations Research Revolution*, pages 20–46. INFORMS, 2015. doi: 10.1287/educ.2015.0139.
- U. Feige, K. Jain, M. Mahdian, and V. Mirrokni. Robust combinatorial optimization with exponential scenarios. In *Integer Programming and Combinatorial Optimization*, pages 439–453. Springer Berlin Heidelberg, 2007. doi: 10.1007/978-3-540-72792-7-33.

- A. Georghiou, W. Wiesemann, and D. Kuhn. Generalized decision rule approximations for stochastic programming via liftings. *Mathematical Programming*, 152(1-2):301–338, may 2015. doi: 10.1007/s10107-014-0789-6.
- L. E. Ghaoui, M. Oks, and F. Oustry. Worst-case value-at-risk and robust portfolio optimization: A conic programming approach. *Operations Research*, 51(4):543–556, 2003. doi: 10.1287/opre.51.4.543.16101.
- J. Goh and N. G. Hall. Total cost control in project management via satisficing. *Management Science*, 59(6):1354–1372, 2013. doi: 10.1287/mnsc.1120.1653.
- D. Goldfarb and G. N. Iyengar. Robust portfolio selection problems. *Mathematics of Operations Research*, 28(1):1–38, 2003. doi: 10.1287/moor.28.1.1.14260.
- B. L. Gorissen, İ. Yanıkoğlu, and D. den Hertog. A practical guide to robust optimization. *Omega*, 53:124–137, 2015. doi: 10.1016/j.omega.2014.12.006.
- G. A. Hanasusanto, D. Kuhn, and W. Wiesemann. K-adaptability in two-stage robust binary programming. *Operations Research*, 63(4):877–891, 2015. doi: 10.1287/opre.2015.1392.
- O. E. Housni and V. Goyal. On the optimality of affine policies for budgeted uncertainty sets. *Mathematics of Operations Research*, 46(2):674–711, 2021. doi: 10.1287/moor.2020.1082.
- D. A. Iancu, M. Sharma, and M. Sviridenko. Supermodularity and affine policies in dynamic robust optimization. *Operations Research*, 61(4):941–956, 2013. doi: 10.1287/opre.2013.1172.
- D. Kuhn, W. Wiesemann, and A. Georghiou. Primal and dual linear decision rules in stochastic and robust optimization. *Mathematical Programming*, 130(1):177–209, dec 2011. doi: 10.1007/s10107-009-0331-4.
- M. Laguna. Applying robust optimization to capacity expansion of one location in telecommunications with demand uncertainty. *Management Science*, 44(11-part-2):S101–S110, nov 1998. doi: 10.1287/mnsc.44.11.s101.
- M. Lu and Z.-J. M. Shen. A review of robust operations management under model uncertainty. *Production and Operations Management*, 30(6):1927–1943, 2021. doi: 10.1111/poms.13239.
- S. Mudchanatongsuk, F. Ordóñez, and J. Liu. Robust solutions for network design under transportation cost and demand uncertainty. *Journal of the Operational Research Society*, 59(5):652–662, 2008. doi: 10.1057/palgrave.jors.2602362.
- J. M. Mulvey, R. J. Vanderbei, and S. A. Zenios. Robust optimization of large-scale systems. *Operations Research*, 43(2):264–281, apr 1995. doi: 10.1287/opre.43.2.264.
- N. Noyan, B. Balcik, and S. Atakan. A stochastic optimization model for designing last mile relief networks. *Transportation Science*, 50(3):1092–1113, 2016. doi: 10.1287/trsc.2015.0621.
- F. Ordóñez and J. Zhao. Robust capacity expansion of network flows. *Networks*, 50(2):136–145, 2007. doi: 10.1002/net.20183.
- D. Paraskevopoulos, E. Karakitsos, and B. Rustem. Robust capacity planning under uncertainty. *Management Science*, 37(7):787–800, jul 1991. doi: 10.1287/mnsc.37.7.787.

- E. L. Porteus. Stochastic inventory theory. In *Handbooks in Operations Research and Management Science*, pages 605–652. Elsevier, 1990. doi: 10.1016/s0927-0507(05)80176-8.
- S. Rahal, D. J. Papageorgiou, and Z. Li. Hybrid strategies using linear and piecewise-linear decision rules for multistage adaptive linear optimization. *European Journal of Operational Research*, 290(3): 1014–1030, may 2021. doi: 10.1016/j.ejor.2020.08.054.
- R. L. Schmidt. A stochastic optimization model to improve production planning and r&d resource allocation in biopharmaceutical production processes. *Management Science*, 42(4):603–617, 1996. doi: 10.1287/mnsc.42.4.603.
- C.-T. See and M. Sim. Robust approximation to multiperiod inventory management. *Operations Research*, 58(3):583–594, 2010. doi: 10.1287/opre.1090.0746.
- D. Simchi-Levi, N. Trichakis, and P. Y. Zhang. Designing response supply chain against bioattacks. *Operations Research*, 67(5):1246–1268, 2019a. doi: 10.1287/opre.2019.1862.
- D. Simchi-Levi, H. Wang, and Y. Wei. Constraint generation for two-stage robust network flow problems. *INFORMS Journal on Optimization*, 1(1):49–70, 2019b. doi: 10.1287/ijoo.2018.0003.
- O. Solyali, J.-F. Cordeau, and G. Laporte. The impact of modeling on robust inventory management under demand uncertainty. *Management Science*, 62(4):1188–1201, 2016. doi: 10.1287/mnsc.2015.2183.
- Z. Wang and M. Qi. Robust service network design under demand uncertainty. *Transportation Science*, 54(3):676–689, may 2020. doi: 10.1287/trsc.2019.0935.
- F. Xie and Y. Huang. A multistage stochastic programming model for a multi-period strategic expansion of biofuel supply chain under evolving uncertainties. *Transportation Research Part E: Logistics and Transportation Review*, 111:130–148, 2018. doi: 10.1016/j.tre.2018.01.015.
- İ. Yamkoğlu, B. L. Gorissen, and D. den Hertog. A survey of adjustable robust optimization. *European Journal of Operational Research*, 277(3):799–813, 2019. doi: 10.1016/j.ejor.2018.08.031.
- J. Zou, S. Ahmed, and X. A. Sun. Partially adaptive stochastic optimization for electric power generation expansion planning. *INFORMS Journal on Computing*, 30(2):388–401, 2018. doi: 10.1287/ijoc.2017.0782.