

THE TWELFTH MOMENT OF HECKE L -FUNCTIONS IN THE WEIGHT ASPECT

PETER HUMPHRIES AND RIZWANUR KHAN

ABSTRACT. We prove an upper bound for the twelfth moment of Hecke L -functions associated to holomorphic Hecke cusp forms of weight k in a dyadic interval $T \leq k \leq 2T$ as T tends to infinity. This bound recovers the Weyl-strength subconvex bound $L(1/2, f) \ll_{\varepsilon} k^{1/3+\varepsilon}$ and shows that for any $\delta > 0$, the sub-Weyl subconvex bound $L(1/2, f) \ll k^{1/3-\delta}$ holds for all but $O_{\varepsilon}(T^{12\delta+\varepsilon})$ Hecke cusp forms f of weight at most T . Our result parallels a related result of Jutila for the twelfth moment of Hecke L -functions associated to Hecke–Maaß cusp forms. The proof uses in a crucial way a spectral reciprocity formula of Kuznetsov that relates the fourth moment of $L(1/2, f)$ weighted by a test function to a dual fourth moment weighted by a different test function.

1. INTRODUCTION

1.1. Main Result. Let \mathcal{B}_{hol} denote the set of holomorphic Hecke cusp forms on the modular surface $\Gamma \backslash \mathbb{H}$, where $\Gamma := \text{SL}_2(\mathbb{Z})$ denotes the modular group. We denote by $k_f \in 2\mathbb{N}$ the weight of a holomorphic Hecke cusp form $f \in \mathcal{B}_{\text{hol}}$. We prove the following result concerning the twelfth moment of the central value $L(1/2, f)$ of the Hecke L -function of f averaged over cusp forms of weight in a given dyadic interval.

Theorem 1.1. *For $T \geq 1$, we have that*

$$\sum_{\substack{f \in \mathcal{B}_{\text{hol}} \\ T \leq k_f \leq 2T}} \frac{L(\frac{1}{2}, f)^{12}}{L(1, \text{ad } f)} \ll_{\varepsilon} T^{4+\varepsilon}.$$

Note that the harmonic weight $1/L(1, \text{ad } f)$ is positive and satisfies the bounds

$$\frac{1}{(\log k_f)^2} \ll \frac{1}{L(1, \text{ad } f)} \ll \log k_f;$$

in particular, the same bound $O_{\varepsilon}(T^{4+\varepsilon})$ holds for the unweighted twelfth moment of $L(1/2, f)$. Upon dropping all but one term and taking twelfth roots, we recover the well-known Weyl-strength subconvex bound¹ $L(1/2, f) \ll_{\varepsilon} k_f^{1/3+\varepsilon}$; thus this should be thought of as a *Weyl-strength* bound for the twelfth moment of $L(1/2, f)$. But in fact our result gives more information, showing for the first time that the central values can only very rarely be as large as the Weyl bound. This is a consequence of the density result given in [Corollary 1.2](#).

1.2. Moments of Hecke L -Functions. [Theorem 1.1](#) fits into an active field of the study of moments of central values of L -functions in various families of L -functions (see in particular [\[Sou21\]](#) for a recent survey on this and related topics). For the family

$$\mathcal{F}_T := \{f \in \mathcal{B}_{\text{hol}} : T \leq k_f \leq 2T\}$$

2020 *Mathematics Subject Classification.* 11F66 (primary); 11F11 (secondary).

The first author was supported by the Simons Foundation (award 965056). The second author was supported by the National Science Foundation Grants DMS-2001183 and DMS-2140604 and the Simons Foundation (award 630985).

¹Note, however, that we *input* this Weyl-strength subconvex bound for $L(1/2, f)$ during the course of the proof of [Theorem 1.1](#); in particular, [Theorem 1.1](#) does not give a new proof of this bound.

of holomorphic cusp forms of weight k_f lying in a dyadic interval $[T, 2T]$, asymptotic formulæ for the k -th moment

$$\sum_{\substack{f \in \mathcal{B}_{\text{hol}} \\ T \leq k_f \leq 2T}} \frac{L\left(\frac{1}{2}, f\right)^k}{L(1, \text{ad } f)}$$

are known for $k \in \{1, 2\}$ (see [BF21, Theorems 3.1, 6.4, 7.4, and 7.5] for state of the art results); indeed, these are even known for the smaller family

$$\mathcal{G}_T := \{f \in \mathcal{B}_{\text{hol}} : k_f = T\}$$

with $T \in 4\mathbb{N}$. Essentially sharp upper bounds have been proven for the third moment for \mathcal{G}_T [Fro20, Theorem 1.3] (see also [Pen01, Theorem 3.1.1]); current technology should also be able to obtain an asymptotic for the third moment for \mathcal{F}_T (cf. [Qi22, Theorem 1.4]). Essentially sharp upper bounds for the fourth moment for \mathcal{F}_T follow directly from the spectral large sieve, while asymptotics for the fourth moment should be possible via current technology (cf. [Ivi02]). Asymptotics for higher moments remain out of reach, though the second author has proven an upper bound for the fifth moment for the family \mathcal{F}_T that is essentially sharp under the assumption of the Selberg eigenvalue conjecture [Kha20, Theorem 1.1].

Tools from random matrix theory have been used to derive a conjectural recipe for *arbitrary* moments: [CFKRS05, Conjecture 1.5.5] posits an asymptotic formula for

$$\sum_{\substack{f \in \mathcal{B}_{\text{hol}} \\ k_f = T}} \frac{L\left(\frac{1}{2}, f\right)^k}{L(1, \text{ad } f)}$$

for each nonnegative integer k . In particular, it is conjectured that there exists a positive arithmetic factor a_k (given explicitly in terms of an Euler product) and a positive geometric factor g_k (given explicitly in terms of an integral arising from random matrix theory) such that this k -th moment is asymptotic to the product of these two factors times $T(\log T)^{k(k-1)/2}$. Thus these random matrix theory heuristics suggest that [Theorem 1.1](#) is far from optimal: instead of an upper bound of size $O_\varepsilon(T^{4+\varepsilon})$, we should expect an *asymptotic* involving a main term of the form $c_{12}T^2(\log T)^{66}$ for some positive constant c_{12} . On the other hand, unconditional improvements to [Theorem 1.1](#) seemingly remain far out of reach, since any power-saving improvement would break the Weyl barrier and yield a sub-Weyl subconvex bound of the form $L(1/2, f) \ll k_f^{1/3-\delta}$ for some $\delta > 0$, a well-known open problem.

[Theorem 1.1](#) parallels an analogous result of Jutila [Jut04, Theorem 2] on the twelfth moment of Hecke L -functions associated to Hecke–Maaß cusp forms f of spectral parameter t_f in a dyadic interval $T \leq t_f \leq 2T$. We also briefly note that there exist related results for twelfth moments of other families of L -functions that similarly fall shy by a power from the optimal asymptotic predicted by random matrix theory yet also yield Weyl-strength subconvex bounds for individual L -functions. The quintessential result in this regard is Heath-Brown’s bound for the twelfth moment of the Riemann zeta function [H-B78, Theorem 1], which has subsequently been reproven by different means by both Iwaniec [Iwa80, Theorem 4] and Jutila [Jut87, Theorem 4.7] and extended more generally by Ivić [Ivi03, Chapter 8]. Finally, bounds for the twelfth moment of Dirichlet L -functions have recently been proven by Milićević and White [MW21, Theorem 1] and by Nunes [Nun21, Theorem 1.1]; the former is of Weyl-strength in the depth aspect, while the latter is of Weyl-strength for smooth moduli. We remark that the twelfth moment bounds for GL_1 L -functions are somewhat easier than for GL_2 L -functions because they can be proven via short interval second moment bounds, while the latter requires short fourth moment bounds, as we discuss below.

1.3. Applications. Upper bounds or asymptotics for various moments of L -functions have many applications, including information on the limiting distribution of L -functions, upper bounds for the size of L -functions, and lower bounds for the proportion of L -functions that are nonvanishing at the central point; see [CS07] for further discussions of such applications. We highlight two applications of [Theorem 1.1](#) below.

The first application of [Theorem 1.1](#) pertains to upper bounds for the proportion of large values of $L(1/2, f)$, recalling the nonnegativity of $L(1/2, f)$ [[KZ81](#), Corollary 1]. Indeed, while [Theorem 1.1](#) does not yield improved *individual* subconvex bounds for $L(1/2, f)$ beyond the Weyl bound $L(1/2, f) \ll_\varepsilon k_f^{1/3+\varepsilon}$, it does show strong *averaged* subconvex bounds, in the form of the following density result (cf. [[H-B78](#), Theorem 2], [[Ivi03](#), Corollary 8.1], [[MW21](#), Theorem 2], and [[Nun21](#), Theorem 1.2]).

Corollary 1.2. *For $T \geq 1$ and $V > 0$, we have that*

$$(1.3) \quad \#\left\{f \in \mathcal{B}_{\text{hol}} : T \leq k_f \leq 2T, L\left(\frac{1}{2}, f\right) \geq V\right\} \ll_\varepsilon \begin{cases} \frac{T^{2+\varepsilon}}{(1+V)^4} & \text{for } 0 \leq V \leq T^{2\vartheta}, \\ \frac{T^{2+2\vartheta+\varepsilon}}{V^5} & \text{for } T^{2\vartheta} \leq V \leq T^{\frac{2(1-\vartheta)}{7}}, \\ \frac{T^{4+\varepsilon}}{V^{12}} & \text{for } T^{\frac{2(1-\vartheta)}{7}} \leq V \leq T^{\frac{1}{3}}(\log T)^{\frac{13}{6}+\varepsilon}, \\ 0 & \text{for } V \geq T^{\frac{1}{3}}(\log T)^{\frac{13}{6}+\varepsilon}, \end{cases}$$

Here $\vartheta = 7/64$ denotes the best known bound towards the Selberg eigenvalue conjecture [[Kim03](#), Proposition 2 of Appendix 2]. The first range of (1.3) follows from the standard fourth moment upper bound

$$\sum_{\substack{f \in \mathcal{B}_{\text{hol}} \\ T \leq k_f \leq 2T}} \frac{L\left(\frac{1}{2}, f\right)^4}{L(1, \text{ad } f)} \ll_\varepsilon T^{2+\varepsilon},$$

which is a straightforward consequence of the spectral large sieve, while the second range is due to the second author's fifth moment bound [[Kha20](#), Theorem 1.1]

$$(1.4) \quad \sum_{\substack{f \in \mathcal{B}_{\text{hol}} \\ T \leq k_f \leq 2T}} \frac{L\left(\frac{1}{2}, f\right)^5}{L(1, \text{ad } f)} \ll_\varepsilon T^{2+2\vartheta+\varepsilon}.$$

[Theorem 1.1](#) yields the bounds (1.3) in the third range. The fourth range is a consequence of Frolenkov's Weyl-strength subconvex bound [[Fro20](#), Corollary 1.4]

$$(1.5) \quad L\left(\frac{1}{2}, f\right) \ll k_f^{\frac{1}{3}}(\log k_f)^{\frac{13}{6}}.$$

In particular, (1.3) shows that for any $\delta > 0$, the number of Hecke cusp forms f of weight $k_f \in [T, 2T]$ that fail to satisfy the sub-Weyl subconvex bound $L(1/2, f) \ll k_f^{1/3-\delta}$ is $O_\varepsilon(T^{12\delta+\varepsilon})$.

[Theorem 1.1](#) also has applications to improved L^4 -norm bounds for Hecke–Maaß cusp forms [[HK22](#)]. Here one must deal with mixed moments of the form

$$\sum_{\substack{f \in \mathcal{B}_{\text{hol}} \\ T \leq k_f \leq 2T}} \frac{L\left(\frac{1}{2}, f\right) L\left(\frac{1}{2}, \text{ad } g \otimes f\right)}{L(1, \text{ad } f)},$$

where g is a Hecke–Maaß cusp form (see in particular [[HK22](#), Proposition 6.11]). While one could bound this mixed moment simply via the individual Weyl-strength subconvex bound (1.5) for $L(1/2, f)$ together with pre-existing bounds for the first moment of $L(1/2, \text{ad } g \otimes f)$ (for which see [[HK22](#), Proposition 6.6]), stronger bounds can be attained by instead using Hölder's inequality and inputting Weyl-strength bounds for the twelfth moment of $L(1/2, f)$, namely [Theorem 1.1](#).

1.4. Sketch of Proof. The analogue of [Theorem 1.1](#) for Hecke–Maaß cusp forms in place of holomorphic Hecke cusp forms is known via work of Jutila [[Jut04](#), Theorem 2]. The proof that we give of [Theorem 1.1](#) follows in broad strokes the same general strategy of that of Jutila, but we need a crucial new idea to handle the holomorphic case, as we now explain.

The approach to estimating the twelfth moment over the family \mathcal{F}_T involves understanding the fourth moment over smaller families such as \mathcal{G}_T . To motivate the approach, define

$$(1.6) \quad \underline{\mathcal{M}}^{\text{hol}}(k) := \sum_{\substack{f \in \mathcal{B}_{\text{hol}} \\ k_f = k}} \frac{L(\frac{1}{2}, f)^4}{L(1, \text{ad } f)}$$

for $k \in 2\mathbb{N}$ and consider the conjectural bound

$$(1.7) \quad \underline{\mathcal{M}}^{\text{hol}}(k) \ll_{\varepsilon} k^{1+\varepsilon},$$

which is an immediate consequence of the generalised Lindelöf hypothesis. This conjectural bound, together with the very strong individual subconvex bound $L(1/2, f) \ll_{\varepsilon} k_f^{1/4+\varepsilon}$ that it implies, immediately yields [Theorem 1.1](#). Although the proof of (1.7) is undoubtedly out of reach of current technology, we are able to prove it in a mean square sense:

$$(1.8) \quad \sum_{\substack{T \leq k \leq 2T \\ k \equiv 0 \pmod{2}}} \underline{\mathcal{M}}^{\text{hol}}(k)^2 \ll_{\varepsilon} T^{3+\varepsilon}.$$

This is a special case of a more refined result, stated in [Proposition 2.1](#), where the mean square is restricted to $\underline{\mathcal{M}}^{\text{hol}}(k)$ of a given size. It turns out that such mean square estimates are enough to substitute for a result like (1.7) as far as obtaining the twelfth moment bound is concerned. This is shown in [Section 2](#).

To establish (1.8), we apply a spectral reciprocity formula due to Kuznetsov and Motohashi (given in [Theorem 3.1](#)) to relate the fourth moment $\underline{\mathcal{M}}^{\text{hol}}(k)$ to a fourth moment of Hecke–Maaß cusp forms. We end up with a relation very roughly of the shape

$$\underline{\mathcal{M}}^{\text{hol}}(k) \rightsquigarrow k^{-1/2} \sum_{\substack{f \in \mathcal{B}_0 \\ t_f \leq k}} \frac{L(\frac{1}{2}, f)^4}{L(1, \text{ad } f)} k^{2it_f},$$

where \mathcal{B}_0 denotes an orthonormal basis of Hecke–Maaß cusp forms f on $\Gamma \backslash \mathbb{H}$, while t_f denotes the spectral parameter of f . When the right hand side is squared and summed over k in a dyadic interval $[T, 2T]$, the oscillatory weight k^{2it_f} (which is an imprecise version of the weight given in [Lemma 4.21](#) with $K = k$ and $L = 2$) facilitates cancellation, as explicated in [Lemma 5.6](#), and leads to the mean square estimate.

In order to apply the spectral reciprocity formula, one must ensure that (certain transforms of) the test functions involved (suppressed in our sketch above) satisfy delicate decay properties. This requirement is well-known to experts by now, but it was overlooked by Kuznetsov, invalidating parts of his paper [[Kuz89](#)], as pointed out by Motohashi [[Mot03](#)]. It is no major challenge to construct test functions satisfying these decay properties for the fourth moment of Hecke–Maaß cusp forms provided that one uses the root number trick (see [[Mot97](#), Section 3.3]). For a purely holomorphic fourth moment such as $\underline{\mathcal{M}}^{\text{hol}}(k)$, however, most natural choices of test function are not valid, and so the spectral reciprocity formula of Kuznetsov has never been successfully applied until now. Our key idea is to introduce a carefully designed and rather non-obvious test function, given in (4.1), that satisfies the required decay properties.

2. PROOF OF THEOREM 1.1

[Theorem 1.1](#) is predicate on the following auxiliary result concerning the second moment in dyadic intervals of large values of the fourth moment of $L(1/2, f)$ in short intervals (cf. [[Jut04](#), Theorem 1]).

Proposition 2.1. For $k \in 2\mathbb{N}$, let $\mathcal{M}^{\text{hol}}(k)$ be as in (1.6). Then for $T \geq 1$ and $V > 0$, we have that

$$(2.2) \quad \mathfrak{S}(T, V) := \sum_{\substack{T \leq k \leq 2T \\ k \equiv 0 \pmod{2} \\ \mathcal{M}^{\text{hol}}(k) \geq V}} \mathcal{M}^{\text{hol}}(k)^2 \ll_{\varepsilon} \begin{cases} T^{3+\varepsilon} & \text{if } 0 < V < T, \\ \frac{T^{4+\varepsilon}}{V} & \text{if } T \leq V \leq T^{\frac{4}{3}}(\log T)^7, \\ 0 & \text{if } V > T^{\frac{4}{3}}(\log T)^7. \end{cases}$$

Note that the bound (2.2) is essentially sharp for $V < T^{1+\varepsilon}$, whereas the generalised Lindelöf hypothesis shows that we should expect that $\mathfrak{S}(T, V) = 0$ for $V \geq T^{1+\varepsilon}$.

We postpone the proof of Proposition 2.1 for the time being and instead proceed directly to the proof of Theorem 1.1.

Proof of Theorem 1.1. We begin by breaking up the twelfth moment based on the size of $L(1/2, f)$. We deal separately with those terms for which $L(1/2, f) < T^{(1-\vartheta)/7}$. By the fifth moment bound (1.4), we have that

$$\sum_{\substack{f \in \mathcal{B}_{\text{hol}} \\ T \leq k_f \leq 2T \\ L(\frac{1}{2}, f) < T^{(1-\vartheta)/7}}} \frac{L(\frac{1}{2}, f)^{12}}{L(1, \text{ad } f)} \ll_{\varepsilon} T^{4+\varepsilon}.$$

We then break up the remaining terms appearing in the twelfth moment via a dyadic subdivision, namely into sums over cusp forms $f \in \mathcal{B}_{\text{hol}}$ for which $T \leq k_f \leq 2T$ and $V \leq L(1/2, f) < 2V$ with $V \geq T^{2(1-\vartheta)/7}$ a dyadic parameter. The Weyl-strength subconvex bound (1.5) ensures that $V \leq k_f^{1/3}(\log k_f)^{7/3}$, say, so that

$$\sum_{\substack{f \in \mathcal{B}_{\text{hol}} \\ T \leq k_f \leq 2T}} \frac{L(\frac{1}{2}, f)^{12}}{L(1, \text{ad } f)} \ll_{\varepsilon} T^{4+\varepsilon} + T^{\varepsilon} \max_{T^{2(1-\vartheta)/7} \leq V \leq T^{1/3}(\log T)^{7/3}} V^8 \sum_{\substack{f \in \mathcal{B}_{\text{hol}} \\ T \leq k_f \leq 2T \\ V \leq L(\frac{1}{2}, f) < 2V}} \frac{L(\frac{1}{2}, f)^4}{L(1, \text{ad } f)}.$$

We introduce a further dyadic subdivision based on the number of positive even integers $k \in [T, 2T]$ for which the cardinality of the set

$$\mathcal{B}(k; V) := \left\{ f \in \mathcal{B}_{\text{hol}} : k_f = k, V \leq L\left(\frac{1}{2}, f\right) < 2V \right\}$$

lies in a dyadic interval. Since $\#\mathcal{B}(k; V) \leq \frac{k}{12} + 1$, we deduce that

$$\sum_{\substack{f \in \mathcal{B}_{\text{hol}} \\ T \leq k_f \leq 2T \\ V \leq L(\frac{1}{2}, f) < 2V}} \frac{L(\frac{1}{2}, f)^4}{L(1, \text{ad } f)} \ll_{\varepsilon} T^{\varepsilon} \max_{1 \leq W \leq \frac{T}{6} + 1} \sum_{\substack{T \leq k \leq 2T \\ k \equiv 0 \pmod{2} \\ W \leq \#\mathcal{B}(k; V) < 2W}} \sum_{f \in \mathcal{B}(k; V)} \frac{L(\frac{1}{2}, f)^4}{L(1, \text{ad } f)}.$$

If $W \leq \#\mathcal{B}(k; V) < 2W$ and $T \leq k \leq 2T$, then

$$\frac{V^4 W}{(\log T)^2} \ll \sum_{f \in \mathcal{B}(k; V)} \frac{L(\frac{1}{2}, f)^4}{L(1, \text{ad } f)} \leq \mathcal{M}^{\text{hol}}(k)$$

as $L(1, \text{ad } f) \ll (\log k_f)^2$, where $\mathcal{M}^{\text{hol}}(k)$ is as in (1.6), and consequently

$$\sum_{\substack{T \leq k \leq 2T \\ k \equiv 0 \pmod{2} \\ W \leq \#\mathcal{B}(k; V) < 2W}} \sum_{f \in \mathcal{B}(k; V)} \frac{L(\frac{1}{2}, f)^4}{L(1, \text{ad } f)} \ll \frac{(\log T)^3}{V^4 W} \mathfrak{S}\left(T, \frac{V^4 W}{(\log T)^3}\right),$$

where $\mathfrak{S}(T, V)$ is as in (2.2). In this way, we see that

$$\sum_{\substack{f \in \mathcal{B}_{\text{hol}} \\ T \leq k_f \leq 2T}} \frac{L(\frac{1}{2}, f)^{12}}{L(1, \text{ad } f)} \ll_{\varepsilon} T^{4+\varepsilon} + T^{\varepsilon} \max_{T^{2(1-\vartheta)/7} \leq V \leq T^{1/3}(\log T)^{7/3}} V^4 \max_{1 \leq W \leq \frac{T}{6}+1} \frac{1}{W} \mathfrak{S}\left(T, \frac{V^4 W}{(\log T)^3}\right),$$

at which point the result follows immediately from the bound (2.2), noting that $\vartheta = 7/64$ implies that $V^4 W / (\log T)^3 \gg T^{57/56} / (\log T)^3 > T$. \square

We return to the proof of Proposition 2.1. The proof relies on the following estimate, which is the technical heart of Theorem 1.1.

Lemma 2.3. *Fix $0 < \delta < 1/100$. For $T \geq 1$, $T^{1+\delta/2} \leq V \leq T^{4/3}(\log T)^7$, and $L := 2 \lfloor \frac{1}{2} V T^{-1-\delta/4} \rfloor$, we have that*

$$(2.4) \quad \sum_{\substack{T \leq K \leq 2T \\ K \equiv 0 \pmod{2}}} \left(\sum_{\substack{K-L \leq k \leq K+L \\ k \equiv 0 \pmod{2} \\ V \leq \mathcal{M}^{\text{hol}}(k) \leq 2V}} \mathcal{M}^{\text{hol}}(k) \right)^2 \ll_{\varepsilon} T^{3+\varepsilon}.$$

We defer the proof of Lemma 2.3, which takes up the bulk of this paper, to Section 5. We use this to now prove Proposition 2.1.

Proof of Proposition 2.1. Via a dyadic subdivision, in order to prove the bounds (2.2), it suffices to prove the related bounds

$$(2.5) \quad \mathfrak{S}^*(T, V) := \sum_{\substack{T \leq k \leq 2T \\ k \equiv 0 \pmod{2} \\ V \leq \mathcal{M}^{\text{hol}}(k) \leq 2V}} \mathcal{M}^{\text{hol}}(k)^2 \ll_{\varepsilon} \begin{cases} T^{3+\varepsilon} & \text{if } 0 < V < T, \\ \frac{T^{4+\varepsilon}}{V} & \text{if } T \leq V \leq T^{4/3}(\log T)^7, \\ 0 & \text{if } V > T^{4/3}(\log T)^7. \end{cases}$$

The bound (2.5) holds for $V > T^{4/3}(\log T)^7$ due to Frolenkov's refinement [Fro20, Theorem 1.3]

$$\sum_{\substack{f \in \mathcal{B}_{\text{hol}} \\ k_f = k}} \frac{L(\frac{1}{2}, f)^3}{L(1, \text{ad } f)} \ll k(\log k)^{\frac{9}{2}}$$

of Peng's third moment bound for $L(1/2, f)$ [Pen01, Theorem 3.1.1] together with the individual Weyl-strength subconvex bound (1.5) for $L(1/2, f)$, since these combine to show that

$$\mathcal{M}^{\text{hol}}(k) \ll k^{\frac{4}{3}}(\log k)^{\frac{20}{3}}.$$

Moreover, the bound (2.5) trivially holds for $0 < V < T^{1+\varepsilon/2}$. It remains to deal with the range $T^{1+\varepsilon/2} \leq V \leq T^{4/3}(\log T)^7$.

We begin by fixing $0 < \delta < 1/100$, taking $T^{1+\delta/2} \leq V \leq T^{4/3}(\log T)^7$, and breaking up the sum over $T \leq k \leq 2T$ in the definition of $\mathfrak{S}^*(T, V)$ to shorter intervals of length $L := 2 \lfloor \frac{1}{2} V T^{-1-\delta/4} \rfloor$, so that

$$\mathfrak{S}^*(T, V) \leq \sum_{\ell=0}^{\lfloor \frac{T}{L} \rfloor - 1} \sum_{\substack{T+\ell L \leq k \leq T+(\ell+1)L \\ k \equiv 0 \pmod{2} \\ V \leq \mathcal{M}^{\text{hol}}(k) \leq 2V}} \mathcal{M}^{\text{hol}}(k)^2 \leq 2V \sum_{\ell=0}^{\lfloor \frac{T}{L} \rfloor - 1} \sum_{\substack{T+\ell L \leq k \leq T+(\ell+1)L \\ k \equiv 0 \pmod{2} \\ V \leq \mathcal{M}^{\text{hol}}(k) \leq 2V}} \mathcal{M}^{\text{hol}}(k).$$

We introduce a dyadic subdivision based on the number of integers ℓ for which the cardinality of

$$\mathcal{P}(\ell; V) := \left\{ T + \ell L \leq k \leq T + (\ell + 1)L : k \equiv 0 \pmod{2}, V \leq \mathcal{M}^{\text{hol}}(k) \leq 2V \right\}$$

lies in a dyadic interval. If $\#\mathcal{P}(\ell; V) \geq 2^j$, then

$$\sum_{\substack{T+\ell L \leq k \leq T+(\ell+1)L \\ k \equiv 0 \pmod{2} \\ V \leq \mathcal{M}^{\text{hol}}(k) \leq 2V}} \underline{\mathcal{M}}^{\text{hol}}(k) \geq 2^j V,$$

and consequently

$$\begin{aligned} \mathfrak{S}^*(T, V) &\leq 2V \sum_{j=0}^{\infty} \sum_{\substack{\ell=0 \\ 2^j \leq \#\mathcal{P}(\ell; V) < 2^{j+1}}}^{\lceil \frac{T}{L} \rceil - 1} \sum_{\substack{T+\ell L \leq k \leq T+(\ell+1)L \\ k \equiv 0 \pmod{2} \\ V \leq \mathcal{M}^{\text{hol}}(k) \leq 2V}} \underline{\mathcal{M}}^{\text{hol}}(k) \\ &\leq 2 \sum_{j=0}^{\infty} 2^{-j} \sum_{\substack{\ell=0 \\ 2^j \leq \#\mathcal{P}(\ell; V) < 2^{j+1}}}^{\lceil \frac{T}{L} \rceil - 1} \left(\sum_{\substack{T+\ell L \leq k \leq T+(\ell+1)L \\ k \equiv 0 \pmod{2} \\ V \leq \mathcal{M}^{\text{hol}}(k) \leq 2V}} \underline{\mathcal{M}}^{\text{hol}}(k) \right)^2 \\ &\leq 4 \sum_{\ell=0}^{\lceil \frac{T}{L} \rceil - 1} \left(\sum_{\substack{T+\ell L \leq k \leq T+(\ell+1)L \\ k \equiv 0 \pmod{2} \\ V \leq \mathcal{M}^{\text{hol}}(k) \leq 2V}} \underline{\mathcal{M}}^{\text{hol}}(k) \right)^2. \end{aligned}$$

We next bound the sum over ℓ . For each even integer $T + \ell L \leq K \leq T + (\ell + 1)L$, we have that

$$\sum_{\substack{T+\ell L \leq k \leq T+(\ell+1)L \\ k \equiv 0 \pmod{2} \\ V \leq \mathcal{M}^{\text{hol}}(k) \leq 2V}} \underline{\mathcal{M}}^{\text{hol}}(k) \leq \sum_{\substack{K-L \leq k \leq K+L \\ k \equiv 0 \pmod{2} \\ V \leq \mathcal{M}^{\text{hol}}(k) \leq 2V}} \underline{\mathcal{M}}^{\text{hol}}(k).$$

Upon squaring, summing over even integers $T + \ell L \leq K \leq T + (\ell + 1)L$, and dividing through by $L/2$, we obtain

$$\begin{aligned} \mathfrak{S}^*(T, V) &\leq \frac{8}{L} \sum_{\ell=0}^{\lceil \frac{T}{L} \rceil - 1} \sum_{\substack{T+\ell L \leq K \leq T+(\ell+1)L \\ K \equiv 0 \pmod{2}}} \left(\sum_{\substack{K-L \leq k \leq K+L \\ k \equiv 0 \pmod{2} \\ V \leq \mathcal{M}^{\text{hol}}(k) \leq 2V}} \underline{\mathcal{M}}^{\text{hol}}(k) \right)^2 \\ &\leq \frac{8}{L} \sum_{\substack{T \leq K \leq 2T \\ K \equiv 0 \pmod{2}}} \left(\sum_{\substack{K-L \leq k \leq K+L \\ k \equiv 0 \pmod{2} \\ V \leq \mathcal{M}^{\text{hol}}(k) \leq 2V}} \underline{\mathcal{M}}^{\text{hol}}(k) \right)^2. \end{aligned}$$

The desired bound $\mathfrak{S}^*(T, V) \ll_{\varepsilon} T^{4+\varepsilon}/V$ now follows upon invoking (2.4). \square

3. $\text{GL}_4 \times \text{GL}_2 \leftrightarrow \text{GL}_4 \times \text{GL}_2$ SPECTRAL RECIPROCITY

We record in this section a particular form of spectral reciprocity: a GL_2 moment of $\text{GL}_4 \times \text{GL}_2$ Rankin–Selberg L -functions is equal to a main term plus a dual moment, which is a GL_2 moment of $\text{GL}_4 \times \text{GL}_2$ Rankin–Selberg L -functions. In the case that we are interested in, these $\text{GL}_4 \times \text{GL}_2$ Rankin–Selberg L -functions factorise as the product of four GL_2 standard L -functions, so that this is an explicit formula for the fourth moment of $L(1/2, f)$ weighted by a test function $h^{\text{hol}}(k_f)$. This formula is due to Kuznetsov [Kuz89, Kuz99], though the initial proof was incomplete in parts and was subsequently completed by Motohashi [Mot03]. We state the following version of Kuznetsov’s formula, which is implicit in the work of Motohashi [Mot03]².

²Motohashi states a formulation of Theorem 3.1 for which the left-hand side of (3.3) only involves a sum over Hecke–Maaß cusp forms instead of including holomorphic Hecke cusp forms, as well as slightly different

Theorem 3.1 (Kuznetsov [Kuz89, Theorem 15], Motohashi [Mot03, Theorem]). *Let $h(t)$ be an even function that is holomorphic in the horizontal strip $|\Im(t)| \leq 1/2 + \delta$ for some $\delta > 0$ and satisfies $h(t) \ll (1 + |t|)^{-5}$, and let $h^{\text{hol}} : 2\mathbb{N} \rightarrow \mathbb{C}$ be a sequence that satisfies $h^{\text{hol}}(k) \ll k^{-5}$. Suppose additionally that the function*

$$(3.2) \quad H(x) := \frac{i}{\pi} \int_{-\infty}^{\infty} \frac{r}{\cosh \pi r} J_{2ir}(4\pi x) h(r) dr + \frac{1}{\pi} \sum_{\substack{k=2 \\ k \equiv 0 \pmod{2}}}^{\infty} (k-1) i^{-k} J_{k-1}(4\pi x) h^{\text{hol}}(k),$$

where $J_{\alpha}(x)$ denotes the Bessel function of the first kind, is such that its Mellin transform $\widehat{H}(s) := \int_0^{\infty} H(x) x^s \frac{dx}{x}$ is holomorphic in the strip $-4 < \Re(s) < 1$, in which it satisfies the bounds $\widehat{H}(s) \ll (1 + |\Im(s)|)^{\Re(s)-4}$. Then

$$(3.3) \quad \sum_{f \in \mathcal{B}_0} \frac{L(\frac{1}{2}, f)^4}{L(1, \text{ad } f)} h(t_f) + \frac{1}{2\pi} \int_{-\infty}^{\infty} \left| \frac{\zeta(\frac{1}{2} + it)^4}{\zeta(1 + 2it)} \right|^2 h(t) dt + \sum_{f \in \mathcal{B}_{\text{hol}}} \frac{L(\frac{1}{2}, f)^4}{L(1, \text{ad } f)} h^{\text{hol}}(k_f) \\ = \sum_{\pm} \sum_{f \in \mathcal{B}_0} \frac{L(\frac{1}{2}, f)^4}{L(1, \text{ad } f)} \widetilde{h}^{\pm}(t_f) + \sum_{\pm} \frac{1}{2\pi} \int_{-\infty}^{\infty} \left| \frac{\zeta(\frac{1}{2} + it)^4}{\zeta(1 + 2it)} \right|^2 \widetilde{h}^{\pm}(t) dt + \sum_{f \in \mathcal{B}_{\text{hol}}} \frac{L(\frac{1}{2}, f)^4}{L(1, \text{ad } f)} \widetilde{h}^{\text{hol}}(k_f) \\ + \lim_{(z_1, z_2, z_3, z_4) \rightarrow (\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2})} \widetilde{h}_{z_1, z_2, z_3, z_4}.$$

Here the transforms \widetilde{h}^{\pm} and $\widetilde{h}^{\text{hol}}$ of h^{hol} are defined by

$$(3.4) \quad \widetilde{h}^{\pm}(t) = \frac{1}{2\pi i} \int_{\sigma - i\infty}^{\sigma + i\infty} \widehat{H}(s) \mathcal{G}_t^{\pm}(s) ds,$$

$$(3.5) \quad \widetilde{h}^{\text{hol}}(k) = \frac{1}{2\pi i} \int_{\sigma - i\infty}^{\sigma + i\infty} \widehat{H}(s) \mathcal{G}_k^{\text{hol}}(s) ds$$

for $0 < \sigma < 1$, where

$$(3.6) \quad \mathcal{G}_t^+(s) := \frac{1}{\pi^2} (2\pi)^s \Gamma\left(\frac{s}{2} + it\right) \Gamma\left(\frac{s}{2} - it\right) \Gamma\left(\frac{1-s}{2}\right)^4 \cos \frac{\pi s}{2} \left(\sin^2 \frac{\pi s}{2} + 1\right),$$

$$(3.7) \quad \mathcal{G}_t^-(s) := \frac{2}{\pi^2} \cosh \pi t (2\pi)^s \Gamma\left(\frac{s}{2} + it\right) \Gamma\left(\frac{s}{2} - it\right) \Gamma\left(\frac{1-s}{2}\right)^4 \sin \frac{\pi s}{2},$$

$$(3.8) \quad \mathcal{G}_k^{\text{hol}}(s) := \frac{i^{-k}}{\pi} (2\pi)^s \frac{\Gamma\left(\frac{s+k-1}{2}\right)}{\Gamma\left(\frac{1-s+k}{2}\right)} \Gamma\left(\frac{1-s}{2}\right)^4 \left(\sin^2 \frac{\pi s}{2} + 1\right).$$

For (z_1, z_2, z_3, z_4) in a sufficiently small neighbourhood of the point $(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2})$ excluding the lines $z_j = 1 - z_{\ell}$ and $z_j = z_{\ell}$ with $1 \leq j < \ell < 4$, we have that

$$(3.9) \quad \widetilde{h}_{z_1, z_2, z_3, z_4} := 2 \sum_{j=1}^4 \widehat{H}(2(1 - z_j)) \frac{\prod_{\substack{\ell=1 \\ \ell \neq j}}^4 \zeta(1 - z_j + z_{\ell}) \prod_{\substack{1 \leq m < n \leq 4 \\ m, n \neq j}} \zeta(z_m + z_n)}{\zeta\left(1 - z_j + \sum_{\substack{\ell=1 \\ \ell \neq j}}^4 z_{\ell}\right)} \\ - 2 \sum_{j=1}^4 h(i(z_j - 1)) \frac{\prod_{\substack{\ell=1 \\ \ell \neq j}}^4 \zeta(1 - z_j + z_{\ell}) \zeta(z_j + z_{\ell} - 1)}{\zeta(3 - 2z_j)} \\ + \sum_{\pm} 2 \sum_{j=1}^4 \widetilde{h}_{z_1, z_2, z_3, z_4; j}^{\pm} \frac{\prod_{\substack{\ell=1 \\ \ell \neq j}}^4 \zeta(1 + z_j - z_{\ell}) \zeta\left(\sum_{\substack{m=1 \\ m \neq j, \ell}}^4 z_m - 1\right)}{\zeta\left(3 + z_j - \sum_{\substack{\ell=1 \\ \ell \neq j}}^4 z_{\ell}\right)},$$

assumptions on the pair of test functions (h, h^{hol}) ; nonetheless, an examination of the proof shows that the result remains valid in the formulation (3.3) (cf. [Mot03, Remark, p. 4]).

where the transforms $\tilde{h}_{z_1, z_2, z_3, z_4; j}^\pm$ are given by

$$\begin{aligned} \tilde{h}_{z_1, z_2, z_3, z_4; j}^+ &:= \frac{1}{2\pi^3 i} \int_{\mathcal{C}_j} \widehat{H}(s) (2\pi)^s \Gamma\left(\frac{s}{2} + \sum_{\ell=1}^4 z_\ell - z_j - 2\right) \Gamma\left(\frac{s}{2} + z_j\right) \\ &\quad \times \prod_{m=1}^4 \Gamma\left(1 - z_m - \frac{s}{2}\right) \sin \frac{\pi}{2} \left(s + \sum_{\ell=1}^4 z_\ell - 1\right) \\ &\quad \times \left(\cos \frac{\pi}{2}(s + z_1 + z_2) \cos \frac{\pi}{2}(s + z_3 + z_4) + \cos \frac{\pi}{2}(z_1 - z_2) \cos \frac{\pi}{2}(z_3 - z_4)\right) ds \end{aligned}$$

and

$$\begin{aligned} \tilde{h}_{z_1, z_2, z_3, z_4; j}^- &:= \frac{1}{2\pi^3 i} \cos \frac{\pi}{2} \left(\sum_{\ell=1}^4 z_\ell - 2z_j\right) \int_{\mathcal{C}_j} \widehat{H}(s) (2\pi)^s \Gamma\left(\frac{s}{2} + \sum_{\ell=1}^4 z_\ell - z_j - 2\right) \Gamma\left(\frac{s}{2} + z_j\right) \\ &\quad \times \prod_{m=1}^4 \Gamma\left(1 - z_m - \frac{s}{2}\right) \left(\cos \frac{\pi}{2}(z_1 - z_2) \cos \frac{\pi}{2}(s + z_3 + z_4) + \cos \frac{\pi}{2}(s + z_1 + z_2) \cos \frac{\pi}{2}(z_3 - z_4)\right) ds, \end{aligned}$$

where \mathcal{C}_j is a contour from $-i\infty$ to $i\infty$ such that $6 + 2z_j - 2\sum_{\ell=1}^4 z_\ell - 2n$ and $2 - 2z_j - 2n$ are to the left of the contour and $-2z_m + 2n$ is to the right of the contour for each $1 \leq m \leq 4$ and each $n \in \mathbb{N}$.

Remark 3.10. There exist cuspidal analogues of Kuznetsov's formula for the fourth moment of $L(1/2, f)$, namely for the first moment of $L(1/2, F \otimes f)$ with F an automorphic form for $\mathrm{SL}_4(\mathbb{Z})$; Kuznetsov's formula corresponds to taking F to be a minimal parabolic Eisenstein series. The authors have proven an analogue of the identity (3.3) with F an Eisenstein series induced from a dihedral Hecke–Maaß cusp form, a quadratic Dirichlet character, and the principal Dirichlet character [HK20, Proposition 7.1]; more recently, the authors have proven an analogue of this identity with F a maximal parabolic Eisenstein series induced from a self-dual Hecke–Maaß cusp form for $\mathrm{SL}_3(\mathbb{Z})$ [HK22, Theorem 4.1]. Blomer, Li, and Miller have proven a *completely* cuspidal version of this identity with F a Hecke–Maaß cusp form for $\mathrm{SL}_4(\mathbb{Z})$ [BLM19, Theorem 1].

4. TEST FUNCTIONS AND TRANSFORMS

Our next step is to use [Theorem 3.1](#) in order to prove [Lemma 2.3](#). Naïvely, we might hope to achieve this by using the identity (3.3) with the choice of pair of test functions (h, h^{hol}) given by

$$\begin{aligned} h(t) &:= 0, \\ h^{\mathrm{hol}}(k) &:= \begin{cases} 1 & \text{if } K - L \leq k \leq K + L, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

This, however, does not meet the requirements of [Theorem 3.1](#), since the Mellin transform $\widehat{H}(s)$ of the transform $H(x)$ of the pair (h, h^{hol}) given by (3.2) does not decay sufficiently rapidly in the vertical strip $-5 < \Re(s) < 1$. One might hope to get around this by instead working with a smooth approximation to the indicator function of the short interval $[K - L, K + L]$, namely $h^{\mathrm{hol}}(k) := \Omega((k - K)/L)$ with Ω an arbitrary smooth function compactly supported on $[-3/2, 3/2]$ that is nonnegative, equal to 1 on $[-1, 1]$, and has bounded derivatives, but this nonetheless runs into the same issue.

We circumvent this obstacle via two tricks. The first is the root number trick: since $L(1/2, f) = 0$ whenever $k_f \equiv 2 \pmod{4}$, as the root number of $L(s, f)$ is i^{k_f} , we can instead choose our test function such that $i^k h^{\mathrm{hol}}(k)$ approximates the indicator function of $[K - L, K + L]$. The second trick is to carefully choose this approximation in such a way that $\widehat{H}(s)$ is holomorphic on a wide vertical strip in which it decays extremely rapidly.

4.1. **Test Functions.** We define the following pair of test functions (h, h^{hol}) :

$$(4.1) \quad \begin{aligned} h(t) &:= 0, \\ h^{\text{hol}}(k) &:= i^{-k} \frac{\Gamma\left(\frac{L^2}{2} + 1\right)^2 \Gamma\left(K + \frac{L^2}{2}\right) \Gamma\left(\frac{K-L^2+k}{2} - 1\right)}{\Gamma\left(K - \frac{L^2}{2} - 1\right) \Gamma\left(\frac{L^2-K+k}{2} + 1\right) \Gamma\left(\frac{K+L^2+k}{2}\right) \Gamma\left(\frac{K+L^2-k}{2} + 1\right)}. \end{aligned}$$

Here $K, L \in \mathbb{N}$ are auxiliary parameters such that $K \equiv L \equiv 0 \pmod{2}$ and $L \leq K^{1/3}$. This particular function h^{hol} is chosen to localise to short intervals: $i^k h^{\text{hol}}(k)$ localises to the interval $[K - L, K + L]$, as we now show.

Lemma 4.2. Fix $0 < \delta < 1/100$. Let (h, h^{hol}) be the pair of test functions (4.1) with $K, L \in 2\mathbb{N}$ satisfying $K^{\delta/8} \leq L \leq K^{1/3-\delta/8}$. Let H be as in (3.2).

- (1) The Mellin transform $\widehat{H}(s)$ of H is such that for $s = \sigma + i\tau$,
- (a) $\widehat{H}(s)$ is holomorphic for $\sigma > L^2 - K - 1$,
 - (b) for σ bounded, $\widehat{H}(s)$ satisfies the bounds

$$(4.3) \quad \widehat{H}(s) \ll \begin{cases} K^\sigma L & \text{if } |\tau| \leq \frac{K}{L}, \\ K^\sigma L e^{-\left(\frac{L|\tau|}{2K}\right)^2} & \text{if } \frac{K}{L} \leq |\tau| \leq K, \\ L|\tau|^\sigma \left(\frac{|\tau|}{K}\right)^{-L^2-1} e^{-\left(\frac{KL}{2|\tau|}\right)^2} & \text{if } K \leq |\tau| \leq KL, \\ L|\tau|^\sigma \left(\frac{|\tau|}{K}\right)^{-L^2-1} & \text{if } |\tau| \geq KL. \end{cases}$$

- (2) The test function $h^{\text{hol}}(k)$ is such that for $k \in 2\mathbb{N}$,
- (a) $i^k h^{\text{hol}}(k) > 0$ for $K - L^2 \leq k \leq K + L^2$ and $h^{\text{hol}}(k) = 0$ otherwise,
 - (b) $i^k h^{\text{hol}}(k) \asymp 1$ for $K - L \leq k \leq K + L$,
 - (c) $h^{\text{hol}}(k) \ll e^{-\frac{(k-K)^2}{4L^2}}$ whenever $L \leq |k - K| \leq L^2$.

Proof. For $w_1, w_2 \in \mathbb{C}$ with $\Re(w_2) > -1$ and $\Re(w_1) > -2\Re(w_2)$, define

$$H_{w_1, w_2}^{\text{hol}}(x) := \frac{1}{\pi} \frac{\Gamma\left(\frac{w_2}{2} + 1\right)^2 \Gamma\left(w_1 + \frac{w_2}{2}\right)}{\Gamma(w_2 + 1) \Gamma\left(w_1 - \frac{w_2}{2} - 1\right)} (2\pi x)^{-w_2} J_{w_1-1}(4\pi x).$$

By [GR15, 6.574.2], we have that

$$\begin{aligned} (\mathcal{L}^{\text{hol}} H_{w_1, w_2}^{\text{hol}})(k) &:= 2\pi i^{-k} \int_0^\infty H_{w_1, w_2}^{\text{hol}}(x) J_{k-1}(4\pi x) \frac{dx}{x} \\ &= i^{-k} \frac{\Gamma\left(\frac{w_2}{2} + 1\right)^2 \Gamma\left(w_1 + \frac{w_2}{2}\right) \Gamma\left(\frac{w_1-w_2+k}{2} - 1\right)}{\Gamma\left(w_1 - \frac{w_2}{2} - 1\right) \Gamma\left(\frac{w_2-w_1+k}{2} + 1\right) \Gamma\left(\frac{w_1+w_2+k}{2}\right) \Gamma\left(\frac{w_1+w_2-k}{2} + 1\right)}. \end{aligned}$$

Choosing $w_1 = K$ and $w_2 = L^2$ with $K, L \in 2\mathbb{N}$, we obtain $h^{\text{hol}}(k)$. On the other hand, by [GR15, 6.574.2],

$$\begin{aligned} (\mathcal{L}^+ H_{w_1, w_2}^{\text{hol}})(t) &:= \frac{\pi i}{\sinh \pi t} \int_0^\infty H_{w_1, w_2}^{\text{hol}}(x) (J_{2it}(4\pi x) - J_{-2it}(4\pi x)) \frac{dx}{x} \\ &= \frac{1}{\pi} \sin \frac{\pi(w_1 - w_2)}{2} \frac{\Gamma\left(\frac{w_2}{2} + 1\right)^2 \Gamma\left(w_1 + \frac{w_2}{2}\right)}{\Gamma\left(w_1 - \frac{w_2}{2} - 1\right)} \prod_{\pm} \frac{\Gamma\left(\frac{w_1-w_2-1}{2} \pm it\right)}{\Gamma\left(\frac{w_1+w_2+1}{2} \pm it\right)}. \end{aligned}$$

This vanishes if $w_1 - w_2 \in 2\mathbb{N}$, and in particular if $w_1 = K$ and $w_2 = L^2$ with $K, L \in 2\mathbb{N}$. It follows that $(\mathcal{L}^{\text{hol}} H_{K, L^2}^{\text{hol}})(k) = h^{\text{hol}}(k)$ and $(\mathcal{L}^+ H_{K, L^2}^{\text{hol}})(t) = h(t)$, so that $H(x) = H_{K, L^2}^{\text{hol}}(x)$ by the Sears–Titchmarsh inversion formula [Iwa02, Appendix B.5].

(1) By [GR15, 6.561.14], we have that

$$(4.4) \quad \widehat{H}(s) = (2\pi)^{-s-1} \frac{\Gamma\left(\frac{L^2}{2} + 1\right)^2 \Gamma\left(K + \frac{L^2}{2}\right) \Gamma\left(\frac{K-L^2+s-1}{2}\right)}{\Gamma(L^2+1)\Gamma\left(K - \frac{L^2}{2} - 1\right) \Gamma\left(\frac{K+L^2-s+1}{2}\right)}.$$

This is holomorphic for $\sigma > L^2 - K - 1$. To prove the bounds (4.3), it suffices to give an asymptotic expansion for $\Re(\log \widehat{H}(s))$. We compute using Stirling's formula the expansions

$$\begin{aligned} \log \frac{\Gamma\left(\frac{L^2}{2} + 1\right)^2}{\Gamma(L^2+1)} &= \log L - (L^2+1) \log 2 + \dots, \\ \log \frac{\Gamma\left(K + \frac{L^2}{2}\right)}{\Gamma\left(K - \frac{L^2}{2} - 1\right)} &= (L^2+1) \log K + \dots, \end{aligned}$$

where the ellipsis indicates lower order terms in L or K . Next, we have that

$$\begin{aligned} \log \frac{\Gamma\left(\frac{K-L^2+s-1}{2}\right)}{\Gamma\left(\frac{K+L^2-s+1}{2}\right)} &= \left(\frac{K-L^2+s-2}{2}\right) \log\left(\frac{K-L^2+s-1}{2e}\right) \\ &\quad - \left(\frac{K+L^2-s}{2}\right) \log\left(\frac{K+L^2-s+1}{2e}\right) + \dots. \end{aligned}$$

The asymptotic expansion of the first two terms depends on the size of $|\tau|$.

For $|\tau| \leq K$, the real part of the first two terms in (4.15) may be written in the form

$$\begin{aligned} -\frac{|\tau|}{2} \arctan\left(\frac{2|\tau|(L^2-\sigma+1)}{K^2\left(1+\frac{\tau^2}{K^2}\right)\left(1-\frac{(L^2-\sigma+1)^2}{K^2+\tau^2}\right)}\right) &+ \frac{K-1}{4} \log \frac{1 - \frac{2(L^2-\sigma+1)}{K\left(1+\frac{\tau^2}{K^2}\right)\left(1+\frac{(L^2-\sigma+1)^2}{K^2+\tau^2}\right)}}{1 + \frac{2(L^2-\sigma+1)}{K\left(1+\frac{\tau^2}{K^2}\right)\left(1+\frac{(L^2-\sigma+1)^2}{K^2+\tau^2}\right)}} \\ &- \frac{L^2-\sigma+1}{2} \log\left(1 + \frac{\tau^2}{K^2}\right) - \frac{L^2-\sigma+1}{2} \log\left(1 + \frac{(L^2-\sigma+1)^2}{K^2+\tau^2}\right) \\ &- \frac{L^2-\sigma+1}{4} \log\left(1 - \frac{4(L^2-\sigma+1)^2}{K^2\left(1+\frac{\tau^2}{K^2}\right)^2\left(1+\frac{(L^2-\sigma+1)^2}{K^2+\tau^2}\right)^2}\right) - (L^2-\sigma+1) \log \frac{K}{2e}. \end{aligned}$$

Via the Taylor series expansions of arctan and log, we deduce that

$$\Re\left(\log \widehat{H}(s)\right) = \log L + \sigma \log K + \begin{cases} O(1) & \text{if } |\tau| \leq \frac{K}{L}, \\ -\frac{\tau^2(L^2-\sigma+1)}{2K^2\left(1+\frac{\tau^2}{K^2}\right)}(1+o(1)) & \text{if } \frac{K}{L} \leq |\tau| \leq K. \end{cases}$$

For $|\tau| \geq K$, we may instead write the real part of the first two terms in (4.15) in the form

$$\begin{aligned} -\frac{|\tau|}{2} \arctan\left(\frac{2(L^2-\sigma+1)}{|\tau|\left(1+\frac{K^2}{\tau^2}\right)\left(1-\frac{(L^2-\sigma+1)^2}{\tau^2+K^2}\right)}\right) &+ \frac{K-1}{4} \log \frac{1 - \frac{2(L^2-\sigma+1)K}{\tau^2\left(1+\frac{K^2}{\tau^2}\right)\left(1+\frac{(L^2-\sigma+1)^2}{\tau^2+K^2}\right)}}{1 + \frac{2(L^2-\sigma+1)K}{\tau^2\left(1+\frac{K^2}{\tau^2}\right)\left(1+\frac{(L^2-\sigma+1)^2}{\tau^2+K^2}\right)}} \\ &- \frac{L^2-\sigma+1}{2} \log\left(1 + \frac{K^2}{\tau^2}\right) - \frac{L^2-\sigma+1}{2} \log\left(1 + \frac{(L^2-\sigma+1)^2}{\tau^2+K^2}\right) \\ &- \frac{L^2-\sigma+1}{4} \log\left(1 - \frac{4(L^2-\sigma+1)^2K^2}{\tau^4\left(1+\frac{K^2}{\tau^2}\right)\left(1+\frac{(L^2-\sigma+1)^2}{\tau^2+K^2}\right)^2}\right) - (L^2-\sigma+1) \log \frac{|\tau|}{2e}. \end{aligned}$$

Via the Taylor series expansions of arctan and log, we deduce that

$$\Re\left(\log \widehat{H}(s)\right) = \log L + \sigma \log |\tau| - (L^2 - \sigma + 1) \log \frac{|\tau|}{K} + \begin{cases} -\frac{K^2(L^2 - \sigma + 1)}{2\tau^2\left(1 + \frac{K^2}{\tau^2}\right)}(1 + o(1)) & \text{if } K \leq |\tau| \leq KL, \\ O(1) & \text{if } |\tau| \geq KL. \end{cases}$$

- (2) As $\Gamma(s)$ has simple poles whenever s is a nonpositive integer, $h^{\text{hol}}(k)$ vanishes unless $K - L^2 \leq k \leq K + L^2$, in which case we have the simpler expression

$$h^{\text{hol}}(k) = i^{-k} \prod_{\ell=0}^{\frac{|K-k|}{2}-1} \frac{L^2 - |K-k| + 1 + \ell}{\frac{L^2}{2} + 1 + \ell} \prod_{j=0}^{L^2} \frac{K - \frac{L^2}{2} - 1 + j}{\frac{K-L^2+k}{2} - 1 + j}$$

by the fact that $\Gamma(n) = (n-1)!$ for $n \in \mathbb{N}$. In particular, $i^k h^{\text{hol}}(k) > 0$ whenever $K - L^2 \leq k \leq K + L^2$. To estimate the size of $i^k h^{\text{hol}}(k)$ in this range, we use Stirling's formula, which shows that

$$\log(i^k h^{\text{hol}}(k)) = -\frac{(K-k)^2}{2L^2}(1 + o(1)) + \frac{(K-k)L^2}{2K} + O(1).$$

for $K - L^2 \leq k \leq K + L^2$. In particular, this is $O(1)$ for $K - L \leq k \leq K + L$, so that $i^k h^{\text{hol}}(k) \asymp 1$ in this range, while in the range $L \leq |k - K| \leq L^2$, this asymptotic implies that $h^{\text{hol}}(k) \ll e^{-\frac{(k-K)^2}{4L^2}}$. \square

4.2. Transforms. We determine the behaviour of the transforms $(\tilde{h}^+, \tilde{h}^-, \tilde{h}^{\text{hol}})$ as in (3.4) and (3.5) with (h, h^{hol}) the pair of test functions (4.1). The latter two transforms are readily estimated using the bounds (4.3).

Lemma 4.5. *Fix $0 < \delta < 1/100$. Let (h, h^{hol}) be the pair of test functions (4.1) with $K, L \in 2\mathbb{N}$ satisfying $K^{\delta/8} \leq L \leq K^{1/3-\delta/8}$. Then for \tilde{h}^- as in (3.4) and \tilde{h}^{hol} as in (3.5), we have that for $t \in \mathbb{R}$ and $k \in 2\mathbb{N}$,*

$$(4.6) \quad \tilde{h}^-(t) \ll K^{-100} L (1 + |t|)^{-100},$$

$$(4.7) \quad \tilde{h}^{\text{hol}}(k) \ll \begin{cases} K^{1-k} L & \text{if } k \leq 101, \\ K^{-100} L & \text{if } 101 \leq k \leq \frac{K}{L}, \\ k^{-100} & \text{if } k \geq \frac{K}{L}. \end{cases}$$

Proof. Let us first consider $\tilde{h}^-(t)$. Provided that $1 + L^2 - K < \sigma < 1$ and that s is a bounded distance away from the poles of $\mathcal{G}_t^-(s)$ at $s = -2(\pm it + \ell)$ with ℓ a nonnegative integer, we have by the bounds (4.3) for $\widehat{H}(s)$ and Stirling's formula applied to the definition (3.7) of $\mathcal{G}_t^-(s)$ that the integrand in (3.4) with $\pm = -$ satisfies the bounds

$$\begin{aligned} & \widehat{H}(s) \mathcal{G}_t^-(s) \\ & \ll_{\sigma} ((1 + |\tau + 2t|)(1 + |\tau - 2t|))^{\frac{\sigma-1}{2}} (1 + |\tau|)^{-2\sigma} e^{-\frac{\pi}{2}|\tau|} \times \begin{cases} 1 & \text{if } |\tau| \leq 2|t|, \\ e^{-\frac{\pi}{2}(|\tau|-2|t|)} & \text{if } |\tau| \geq 2|t|, \end{cases} \\ & \times \begin{cases} K^{\sigma} L & \text{if } |\tau| \leq \frac{K}{L}, \\ K^{\sigma} L e^{-\frac{L^2 \tau^2}{4K^2}} & \text{if } \frac{K}{L} \leq |\tau| \leq K, \\ L |\tau|^{\sigma} \left(\frac{|\tau|}{K}\right)^{-L^2-1} e^{-\frac{L^2 K^2}{4\tau^2}} & \text{if } K \leq |\tau| \leq KL, \\ L |\tau|^{\sigma} \left(\frac{|\tau|}{K}\right)^{-L^2-1} & \text{if } |\tau| \geq KL. \end{cases} \end{aligned}$$

In particular, the integral of this function along the line $\Re(s) = \sigma$ with $\sigma < 0$ is

$$\ll_{\sigma} K^{\sigma} L(1 + |t|)^{\sigma-1}.$$

Additionally, for ℓ a bounded nonnegative integer, we have by Stirling's formula that

$$\operatorname{Res}_{s=2(\pm it - \ell)} \widehat{H}(s) \mathcal{G}_t^-(s) \ll_{\ell} (1 + |t|)^{3\ell - \frac{1}{2}} e^{-\pi|t|} \times \begin{cases} K^{-2\ell} L & \text{if } |t| \leq \frac{K}{2L}, \\ K^{-2\ell} L e^{-\frac{L^2|t|^2}{K^2}} & \text{if } \frac{K}{2L} \leq |t| \leq \frac{K}{2}, \\ L|t|^{-2\ell} \left(\frac{2|t|}{K}\right)^{-L^2-1} e^{-\frac{L^2 K^2}{16|t|^2}} & \text{if } \frac{K}{2} \leq |t| \leq \frac{KL}{2}, \\ L|t|^{-2\ell} \left(\frac{2|t|}{K}\right)^{-L^2-1} & \text{if } |t| \geq \frac{KL}{2}. \end{cases}$$

Thus by shifting the contour of integration in (3.4) to $\sigma = -100$, we deduce the bounds (4.6).

Next, we consider $\widetilde{h}^{\text{hol}}(k)$. Provided that $1 + L^2 - K < \sigma < 1$ and s is a bounded distance away from the poles at $s = 1 - k - 2\ell$ with ℓ a nonnegative integer, we have by the bounds (4.3) for $\widehat{H}(s)$ and Stirling's formula applied to the definition (3.8) of $\mathcal{G}_k^{\text{hol}}(s)$ that the integrand in (3.5) satisfies the bounds

$$\widehat{H}(s) \mathcal{G}_k^{\text{hol}}(s) \ll_{\sigma} (k + |\tau|)^{\sigma-1} (1 + |\tau|)^{-2\sigma} \times \begin{cases} K^{\sigma} L & \text{if } |\tau| \leq \frac{K}{L}, \\ K^{\sigma} L e^{-\frac{L^2 \tau^2}{4K^2}} & \text{if } \frac{K}{L} \leq |\tau| \leq K, \\ L|\tau|^{\sigma} \left(\frac{|\tau|}{K}\right)^{-L^2-1} e^{-\frac{L^2 K^2}{4\tau^2}} & \text{if } K \leq |\tau| \leq KL, \\ L|\tau|^{\sigma} \left(\frac{|\tau|}{K}\right)^{-L^2-1} & \text{if } |\tau| \geq KL. \end{cases}$$

In particular, the integral of this function along the line $\Re(s) = \sigma$ with $\sigma < 0$ is

$$\ll_{\sigma} \begin{cases} L^{1+\sigma} & \text{if } k \leq \frac{K}{L}, \\ K^{1-\sigma} L^{2\sigma} k^{\sigma-1} & \text{if } k \geq \frac{K}{L}. \end{cases}$$

Additionally, for ℓ a bounded nonnegative integer, we have by Stirling's formula that

$$\operatorname{Res}_{s=1-k-2\ell} \widehat{H}(s) \mathcal{G}_k^{\text{hol}}(s) \ll_{\ell} K^{1-k-2\ell} L.$$

Thus by shifting the contour of integration in (3.5) to $\sigma = -\frac{100 \log K}{\log L}$, we deduce the bounds (4.7). \square

Determining the behaviour of the transform \widetilde{h}^+ as in (3.4) takes significantly more effort. Indeed, while \widetilde{h}^- and $\widetilde{h}^{\text{hol}}$ are, in practice, negligibly small, this is *not* the case for \widetilde{h}^+ in certain ranges. Moreover, for the purposes of proving Lemma 2.3, we require not only upper bounds for \widetilde{h}^+ but additionally an *asymptotic* in the range where it is nonnegligible. As we shall presently show, in this nonnegligible range, the function \widetilde{h}^+ is not so small but is highly oscillatory, a property that we will later exploit in Lemma 5.6.

To determine the asymptotic behaviour of \widetilde{h}^+ , we first determine the behaviour of the integrand in the definition (3.4). From hereon, we make use of the ε -convention: ε denotes an arbitrarily small positive constant whose value may change from occurrence to occurrence; we will always assume that ε is sufficiently small with respect to δ . In the proof of the following result, by a "lower order term" in parameters x_1, \dots, x_n , we will mean a function $f(x_1, x_2, \dots, x_n)$ such that

$$\frac{\partial^{j_1}}{\partial x_1^{j_1}} \cdots \frac{\partial^{j_n}}{\partial x_n^{j_n}} f(x_1, \dots, x_n) \ll K^{-\varepsilon} (1 + |x_1|)^{-j_1} \cdots (1 + |x_n|)^{-j_n}$$

for any integers $j_1, \dots, j_n \geq 0$.

Lemma 4.8. Fix $0 < \delta < 1/100$. Let $\widehat{H}(s)$ be given by (4.4) with $K, L \in 2\mathbb{N}$ satisfying $K^{\delta/8} \leq L \leq K^{1/3-\delta/8}$. Let $K^{2/3+\varepsilon} < t < K(\log K)/L$. Let $s = \sigma + i(2t + y)$ with $\sigma > 0$ fixed, where

$$(4.9) \quad \frac{t^3}{K^{2+\varepsilon}} < y < \frac{K \log K}{L}.$$

For $\mathcal{G}_t^+(s)$ as in (3.6), we have that

$$(4.10) \quad \widehat{H}(s)\mathcal{G}_t^+(s) = A(y, t, K)e^{iB(y, t, K)} + O(K^{-500}),$$

where $A(y, t, K)$ and $B(y, t, K)$ are smooth functions (whose dependence on L we suppress in the notation), complex-valued and real-valued respectively, that satisfy

$$(4.11) \quad \frac{\partial^j}{\partial y^j} A(y, t, K) \ll \frac{K^{10}}{y^j}, \quad \frac{\partial}{\partial y} B(y, t, K) = \log \frac{Ky^{\frac{1}{2}}(y+4t)^{\frac{1}{2}}}{(2t+y)^2} + O(1), \quad \frac{\partial^{j+2}}{\partial y^{j+2}} B(y, t, K)(y) \ll \frac{1}{y^{j+1}}$$

for any integer $j \geq 0$. If additionally

$$(4.12) \quad \frac{t^3}{K^{2+\varepsilon}} < y < \frac{t^3}{K^{2-\varepsilon}},$$

then (4.10) holds with

$$(4.13) \quad A(y, t, K) = \frac{L}{(ty)^{\frac{1}{2}}} \left(\frac{a_1 K^2 y}{t^3} \right)^{\frac{\sigma}{2}} \left(\frac{a_2 K}{t} \right)^{2it} b_1(y, t, K),$$

$$(4.14) \quad B(y, t, K) = \frac{a_3 t^3}{K^2} + \frac{y}{2} \left(\log \left(\frac{a_4 K^2 y}{t^3} \right) + b_2(y, t, K) \right),$$

where a_i are some real constants with $a_1, a_2, a_4 > 0$, and $b_1(y, t, K)$, $b_2(y, t, K)$ are smooth functions (whose dependence on L we suppress in the notation), complex-valued and real-valued respectively, that satisfy

$$\frac{\partial^{j_1}}{\partial y^{j_1}} \frac{\partial^{j_2}}{\partial t^{j_2}} \frac{\partial^{j_3}}{\partial K^{j_3}} b_i(y, t, K) \ll y^{-j_1} t^{-j_2} K^{-j_3} \times \begin{cases} 1 & \text{if } i = 1, \\ K^{-\varepsilon} & \text{if } i = 2, \end{cases}$$

for any $j_1, j_2, j_3 \geq 0$.

Proof. By Stirling's formula, we have that

$$(4.15) \quad \log \frac{\Gamma\left(\frac{K-L^2+s-1}{2}\right)}{\Gamma\left(\frac{K+L^2-s+1}{2}\right)} = \left(\frac{K-L^2+s-2}{2}\right) \log\left(\frac{K-L^2+s-1}{2e}\right) - \left(\frac{K+L^2-s}{2}\right) \log\left(\frac{K+L^2-s+1}{2e}\right) + \dots + O(K^{-1000}),$$

where the ellipsis indicates lower order terms in s, K, L . We follow the same convention below, with parameters that will be obvious from the context. Next, we insert the power series expansion

$$\log\left(\frac{K \pm (s-L^2-1)}{2e}\right) = \log\left(\frac{K}{2}\right) - 1 + \sum_{j=1}^{\infty} \frac{(-1)^{j+1}}{j} \left(\pm \frac{s-L^2-1}{K}\right)^j.$$

The contribution to (4.15) of -1 and the term $j = 1$ is

$$\left(\frac{K-L^2+s-2}{2}\right) \left(-1 + \frac{s-L^2-1}{K}\right) - \left(\frac{K+L^2-s}{2}\right) \left(-1 - \frac{s-L^2-1}{K}\right) = -\frac{s-L^2-1}{K}.$$

Similarly computing the contributions of the terms with j even and odd separately, we get that

$$\log \frac{\Gamma\left(\frac{K-L^2+s-1}{2}\right)}{\Gamma\left(\frac{K+L^2-s+1}{2}\right)} = (s-L^2-1) \log\left(\frac{K}{2}\right) - \frac{s-L^2-1}{K}$$

$$- \sum_{j \in 2\mathbb{N}} \frac{K}{j(j+1)} \left(\frac{s-L^2-1}{K} \right)^{j+1} + \dots + O(K^{-1000}).$$

We also compute using Stirling's formula the expansions

$$\log \frac{\Gamma\left(\frac{L^2}{2} + 1\right)^2}{\Gamma(L^2 + 1)} = \log L - (L^2 + 1) \log 2 + \dots + O(K^{-1000})$$

and

$$\log \frac{\Gamma\left(K + \frac{L^2}{2}\right)}{\Gamma\left(K - \frac{L^2}{2} - 1\right)} = (L^2 + 1) \log K + \dots + O(K^{-1000}).$$

Recalling the definition (4.4) of $\widehat{H}(s)$ and the assumption $K^{\frac{2}{3}+\varepsilon} < t < \frac{K^{1+\varepsilon}}{L}$, we deduce that

$$\widehat{H}(s) = (2\pi)^{-s-1} L \exp\left(s \log\left(\frac{K}{2}\right) - \sum_{j \in 2\mathbb{N}} \frac{K}{j(j+1)} \left(\frac{s-K^2-1}{K}\right)^{j+1} + \dots\right) + O(K^{-750}).$$

Consider first contribution of $j = 2$, which is $-\frac{1}{6} \frac{(\sigma+i(2t+y)-L^2-1)^3}{K^2}$. We expand out the cube and put aside the terms $\frac{i4t^3}{3K^2}$ and $\frac{-(2t+y)^2 L^2}{2K^2}$. The rest of the terms are either real and of lower order or imaginary. The imaginary terms divided by iy are real and of lower order. Expanding out the contribution of $j \geq 4$, we see that the real terms are of lower order and the imaginary terms divided by iy are real and of lower order, using (4.9). In this way, we get that $\widehat{H}(s)$ is equal to (4.16)

$$L \left(\frac{K}{4\pi}\right)^{\sigma+2it} \exp\left(\frac{-(2t+y)^2 L^2}{2K^2}\right) \exp\left(\frac{i4t^3}{3K^2}\right) \exp\left(iy \left(\log\left(\frac{K}{4\pi}\right) + \dots\right) + \dots\right) + O(K^{-750}).$$

Next, we turn to the function $\mathcal{G}_t^+(s)$. We write $\mathcal{G}_t^+(s) = \mathcal{H}(s) \mathcal{I}_t(s)$, where

$$\begin{aligned} \mathcal{H}(s) &:= \frac{1}{\pi^2} (2\pi)^{2s} \Gamma\left(\frac{1-s}{2}\right)^4 \left(\sin^2 \frac{\pi s}{2} + 1\right), \\ \mathcal{I}_t(s) &:= (2\pi)^{-s} \Gamma\left(\frac{s}{2} + it\right) \Gamma\left(\frac{s}{2} - it\right) \cos \frac{\pi s}{2}. \end{aligned}$$

By Stirling's formula, we get that

$$(4.17) \quad \mathcal{H}(s) = - \left(\frac{2t+y}{4\pi}\right)^{-2\sigma} \exp\left((-2iy - 4it) \log\left(\frac{2t+y}{4\pi e}\right) + \dots\right) + O(K^{-1000}).$$

In the range (4.12), we can further simplify this to

$$(4.18) \quad \mathcal{H}(s) = -e^{-2\sigma} \left(\frac{t}{2\pi e}\right)^{-2\sigma-4it} \exp\left(-2iy \left(\log\left(\frac{t}{2\pi}\right) + \dots\right) + \dots\right) + O(K^{-1000}).$$

Finally, we also compute using Stirling's formula the expansion

$$(4.19) \quad \mathcal{I}_t(s) = -\frac{i}{2} \left(\frac{y}{4\pi}\right)^{\frac{\sigma-1}{2}} \left(\frac{y+4t}{4\pi}\right)^{\frac{\sigma-1}{2}} \exp\left(\frac{iy}{2} \log\left(\frac{y}{4\pi e}\right) + \frac{i(y+4t)}{2} \log\left(\frac{y+4t}{4\pi e}\right) + \dots\right) + O(K^{-1000}).$$

In the range (4.12), we have that

$$(4.20) \quad \mathcal{I}_t(s) = -\frac{i}{2} e^{\frac{\sigma-1}{2}} \left(\frac{y}{4\pi}\right)^{\frac{\sigma-1}{2}} \left(\frac{t}{\pi e}\right)^{\frac{\sigma-1}{2}+2it} \exp\left(\frac{iy}{2} \left(\log\left(\frac{yt}{4\pi^2 e}\right) + \dots\right) + \dots\right) + O(K^{-1000}).$$

Taking (4.16), (4.17), (4.19) together gives (4.10). Taking (4.16), (4.18), (4.20) together gives (4.13), after collecting the lower order terms into the functions $b_1(y, t, K)$ and $b_2(y, t, K)$. The

factor $\exp\left(\frac{-(2t+y)^2 L^2}{2K^2}\right)$, which decays for $t \gg (2t+y) \gg K/L$, is also absorbed into $b_1(y, t, K)$. \square

Now we are ready to analyse $\tilde{h}^+(t)$. The main tool is stationary phase analysis.

Lemma 4.21. *Fix $0 < \delta < 1/100$. Let (h, h^{hol}) be the pair of test functions (4.1) with $K, L \in 2\mathbb{N}$ satisfying $K^{\delta/8} \leq L \leq K^{1/3-\delta/8}$. Then for \tilde{h}^+ as in (3.4), we have that for $t \in \mathbb{R}$,*

$$(4.22) \quad \tilde{h}^+(t) = \begin{cases} O\left(L^{1+\varepsilon}(1+|t|)^{-\frac{1}{2}}\right) & \text{for } |t| \leq K^{\frac{2}{3}+\varepsilon}, \\ g(t, K, L) + O(K^{-100}) & \text{for } K^{\frac{2}{3}+\varepsilon} < |t| < \frac{K \log K}{L}, \\ O(|t|^{-1000}) & \text{for } |t| \geq \frac{K \log K}{L}, \end{cases}$$

where

$$(4.23) \quad g(t, K, L) := \sum_{\pm} \frac{L}{|t|^{\frac{1}{2}}} \left(\frac{c_1 K}{|t|}\right)^{\pm 2it} \exp\left(\pm i \frac{c_2 t^3}{K^2} (1 + d_1(\pm t, K))\right) d_2(\pm t, K)$$

with c_i some real constants for which $c_1 > 0$ and $d_1(t, K)$ and $d_2(t, K)$ smooth functions (whose dependence on L we suppress in the notation), real-valued and complex-valued respectively, that satisfy

$$(4.24) \quad \frac{\partial^{j_1}}{\partial t^{j_1}} \frac{\partial^{j_2}}{\partial K^{j_2}} d_i(t, K) \ll t^{-j_1} K^{-j_2} \times \begin{cases} K^{-\varepsilon} & \text{if } i = 1, \\ 1 & \text{if } i = 2, \end{cases}$$

for any $j_1, j_2 \geq 0$.

Proof. The integrand in (3.4) with $\pm = +$ is meromorphic in the open half-plane strip $\Re(s) > 1 + L^2 - K$ with simple poles at $s = 2(\pm it - \ell)$ and poles of order 4 at $s = 2\ell + 1$ for each $\ell \in \mathbb{N}_0$. For $s = \sigma + i\tau$ a bounded distance away from such a pole with σ bounded,

$$(4.25) \quad \widehat{H}(s) \mathcal{G}_t^+(s) \ll_{\sigma} ((1 + |\tau + 2t|)(1 + |\tau - 2t|))^{\frac{\sigma-1}{2}} (1 + |\tau|)^{-2\sigma} \times \begin{cases} e^{-\frac{\pi}{2}(2|t| - |\tau|)} & \text{if } |\tau| \leq 2|t|, \\ 1 & \text{if } |\tau| \geq 2|t|, \end{cases} \\ \times \begin{cases} K^{\sigma} L & \text{if } |\tau| \leq \frac{K}{L}, \\ K^{\sigma} L e^{-\frac{L^2 \tau^2}{4K^2}} & \text{if } \frac{K}{L} \leq |\tau| \leq K, \\ L|\tau|^{\sigma} \left(\frac{|\tau|}{K}\right)^{-L^2-1} e^{-\frac{L^2 K^2}{4\tau^2}} & \text{if } K \leq |\tau| \leq KL, \\ L|\tau|^{\sigma} \left(\frac{|\tau|}{K}\right)^{-L^2-1} & \text{if } |\tau| \geq KL. \end{cases}$$

In particular, the integral of this function along the line $\Re(s) = \sigma$ with $\sigma < -1$ is

$$\ll_{\sigma} L^{1+\sigma} + K^{\sigma} L (1 + |t|)^{-\frac{3\sigma}{2} - \frac{1}{2}}.$$

Moreover,

$$\text{Res}_{s=2(\pm it - \ell)} \widehat{H}(s) \mathcal{G}_t^+(s) \ll_{\ell} (1 + |t|)^{3\ell - \frac{1}{2}} \times \begin{cases} K^{-2\ell} L & \text{if } |t| \leq \frac{K}{2L}, \\ K^{-2\ell} L e^{-\frac{L^2 |t|^2}{K^2}} & \text{if } \frac{K}{2L} \leq |t| \leq \frac{K}{2}, \\ L|t|^{-2\ell} \left(\frac{|t|}{K}\right)^{-L^2-1} e^{-\frac{L^2 K^2}{16|t|^2}} & \text{if } \frac{K}{2} \leq |t| \leq \frac{KL}{2}, \\ L|t|^{-2\ell} \left(\frac{|t|}{K}\right)^{-L^2-1} & \text{if } |t| \geq \frac{KL}{2}. \end{cases}$$

Thus for $|t| \leq K^{2/3-\varepsilon}$, we shift the contour of integration to the line $\Re(s) = -\frac{1}{2} \frac{\log K}{\log L}$. The dominant contribution comes from the poles at $s = \pm 2it$, so that $\tilde{h}^+(t) \ll L(1 + |t|)^{-1/2}$.

For $|t| \geq K(\log K)/L$, we keep the contour of integration at $\sigma \in (0, 1)$ and simply estimate the integral via (4.25). This shows that $\tilde{h}^+(t) \ll |t|^{-1000}$ in this range.

Now suppose that $K^{2/3-\varepsilon} < |t| < K(\log K)/L$. We move the line of integration in (3.4) far to the right. In doing so, we cross poles of order 4 of $\mathcal{G}_t^+(s)$ at $s = 2\ell - 1$ for each $\ell \in \mathbb{N}$. The total contribution of the residues of these poles is $O(K^{-1000})$ by the bounds (4.3) for $\hat{H}(s)$ together with Stirling's formula.

On the new line of integration, we write $s = \sigma + i\tau$ for σ a large fixed even integer, say. By the bounds (4.25), the integrand is negligibly small outside the range

$$(4.26) \quad 2|t| - (\log K)^2 < |\tau| < \frac{K \log K}{L}.$$

In this range, we have that

$$(4.27) \quad \hat{H}(s)\mathcal{G}_t^+(s) \ll_{\sigma} \frac{L}{(1 + |\tau + 2t|)^{\frac{1}{2}}(1 + |\tau - 2t|)^{\frac{1}{2}}} \left(\frac{K^2(1 + |\tau + 2t|)(1 + |\tau - 2t|)}{|t|^4} \right)^{\frac{\sigma}{2}}.$$

Thus for σ sufficiently large with respect to δ and ε , we see that the contribution of the ranges $|\tau \pm 2t| \leq |t|^3/K^{2+\varepsilon}$ is $O(K^{-1000})$. Together with (4.26), this implies that if we define $y := \tau - 2t$, then we may restrict to the range

$$(4.28) \quad \frac{t^3}{K^{2+\varepsilon}} < y < \frac{K \log K}{L}$$

in the case that t is positive, and the analogous range with y negative in the case that t is negative. Since both cases are entirely similar (indeed, they correspond to the \pm cases in (4.22)), we henceforth assume that t is positive and that (4.28) holds.

We may write

$$\tilde{h}^+(t) = \frac{1}{2\pi} \int_{\frac{t^3}{K^{2+\varepsilon}}}^{\frac{K \log K}{L}} A(y, t, K) e^{iB(y, t, K)} dy + O(K^{-100}),$$

where $A(y, t, K)$ and $B(y, t, K)$ are as in Lemma 4.8. We then perform a dyadic subdivision on this integral by inserting a collection of weight functions of the form $W(\frac{y}{R})$ for

$$(4.29) \quad \frac{t^3}{K^{2+\varepsilon}} < R < \frac{K \log K}{L},$$

where $W(y)$ is a smooth function compactly supported on the interval $(1, 2)$ with bounded derivatives. After making the substitution $y \mapsto Ry$, we are left with considering integrals of the form

$$(4.30) \quad \frac{R}{2\pi} \int_0^{\infty} W(y) A(yR, t, K) e^{iB(yR, t, K)} dy.$$

By (4.11), we have that

$$\frac{\partial}{\partial y} B(yR, t, K) = R \log \frac{K(yR)^{\frac{1}{2}}(yR + 4t)^{\frac{1}{2}}}{(2t + yR)^2} + O(R).$$

The key point is that this first derivative is sizeable when R is sufficiently large. Specifically, if $R > t^3/K^{2-\varepsilon}$, then by (4.11), we have that

$$\left| \frac{\partial}{\partial y} B(yR, t, K) \right| \gg R, \quad \frac{\partial^{j+2}}{\partial y^{j+2}} B(yR, t, K) \ll R, \quad \frac{\partial^j}{\partial y^j} W(y) A(yR, t, K) \ll K^{10}$$

for any $j \geq 0$. It follows by integration by parts (for example, by [BKY13, Lemma 8.1] with parameters R as in (4.29), $Y = R$, and $U = Q = 1$) that the integral (4.30) is $O(K^{-1000})$ when $R > t^3/K^{2-\varepsilon}$. We have therefore shown that

$$(4.31) \quad \tilde{h}^+(t) = \frac{1}{2\pi} \int_0^{\infty} \Omega\left(y \frac{K^2}{t^3}\right) A(y, t, K) e^{iB(y, t, K)} dy + O(K^{-100}),$$

where $A(y, t, K)$ and $B(y, t, K)$ are as in Lemma 4.8 and Ω is a smooth function compactly supported on the interval

$$(4.32) \quad (K^{-\varepsilon}, K^\varepsilon)$$

with derivatives satisfying $\Omega^{(j)}(y) \ll_j (K^\varepsilon)^j$.

We now address the range $K^{2/3-\varepsilon} < t < K^{2/3+\varepsilon}$. In this case, the integral in (4.31) is restricted to the range $K^{-\varepsilon} < y < K^\varepsilon$, and simply bounding trivially using (4.27) gives the required bound.

We are left to consider the range $K^{2/3+\varepsilon} < t < K(\log K)/L$. Making the substitution $y \mapsto y \frac{t^3}{K^2}$ in (4.31) and inserting the expansions (4.13) and (4.14) for $A(y, t, K)$ and $B(y, t, K)$, we see that up to a negligible error term, $\tilde{h}^+(t)$ is equal to

$$(4.33) \quad \frac{Lt}{K} \left(\frac{a_2 K}{t} \right)^{2it} \exp \left(i \frac{a_3 t^3}{K^2} \right) \times \int_0^\infty \Omega(y) (a_1 y)^{\frac{\sigma-1}{2}} b_1 \left(y \frac{t^3}{K^2}, t, K \right) \exp \left(\frac{iyt^3}{2K^2} \left(\log(a_4 y) + b_2 \left(y \frac{t^3}{K^2}, t, K \right) \right) \right) dy.$$

Letting

$$\phi_{t,K}(y) := \frac{yt^3}{2K^2} \left(\log(a_4 y) + b_2 \left(y \frac{t^3}{K^2}, t, K \right) \right)$$

denote the phase of the integrand, we have that

$$\begin{aligned} \phi'_{t,K}(y) &= \frac{t^3}{2K^2} \left(\log(ea_4 y) + b_2 \left(y \frac{t^3}{K^2}, t, K \right) + y \frac{\partial}{\partial y} b_2 \left(y \frac{t^3}{K^2}, t, K \right) \right) \\ &= \frac{t^3}{2K^2} \left(\log(ea_4 y) + O(K^{-\varepsilon}) \right), \\ \phi_{t,K}^{(j)}(y) &= \frac{1}{y^{j-1}} \frac{t^3}{2K^2} \left(1 + O(K^{-\varepsilon}) \right) \end{aligned}$$

for $j \geq 2$. In particular, for K sufficiently large, $\phi'_{t,K}(y)$ strictly increases from a negative to positive value through the interval (4.32). Thus there is a unique stationary point

$$y_0 = y_0(t, K) := \frac{1}{ea_4} + r(t, K) = \frac{1}{ea_4} + O(K^{-\varepsilon}),$$

say. Moreover, since $r(t, K)$ is defined implicitly by $F(t, K, r(t, K)) = 0$ for

$$F(t, K, x) := \log(1+ea_4 x) + b_2 \left(\left(\frac{1}{ea_4} + x \right) \frac{t^3}{K^2}, t, K \right) + \left(\frac{1}{ea_4} + x \right) \frac{\partial}{\partial x} b_2 \left(\left(\frac{1}{ea_4} + x \right) \frac{t^3}{K^2}, t, K \right),$$

we can use the implicit function theorem to get that $r(t, K)$ is smooth, and to compute

$$\frac{\partial}{\partial t} r(t, K) = - \frac{\frac{\partial}{\partial t} \Big|_{x=r(t,K)} F(t, K, x)}{\frac{\partial}{\partial x} \Big|_{x=r(t,K)} F(t, K, x)}$$

and the partial derivative with respect to K and higher derivatives similarly. From this, we get that

$$\frac{\partial^{j_1}}{\partial t^{j_1}} \frac{\partial^{j_2}}{\partial K^{j_2}} r(t, K) \ll K^{-\varepsilon} t^{-j_1} K^{-j_2}$$

for $j_1, j_2 \geq 0$. The final step is to derive the stationary phase expansion of the integral in (4.33), with leading term of size $\phi''_{t,K}(y_0)^{-\frac{1}{2}} \asymp \left(\frac{t^3}{K^2} \right)^{-\frac{1}{2}}$ and phase

$$\begin{aligned} \phi_{t,K}(y_0) &= \left(\frac{1}{ea_4} + r(t, K) \right) \frac{t^3}{2K^2} \left(-1 - \left(\frac{1}{ea_4} + r(t, K) \right) \frac{\partial}{\partial y} \Big|_{y=\frac{1}{ea_4}+r(t,K)} b_2 \left(y \frac{t^3}{K^2}, t, K \right) \right) \\ &\asymp \frac{t^3}{K^2}. \end{aligned}$$

The full form of the stationary phase expansion can be obtained using [BKY13, Proposition 8.2] with the main parameter being $Y = \frac{t^3}{K^2}$ and the other parameters V, X, Q lying in the interval $(K^{-\varepsilon}, K^\varepsilon)$. This yields the desired result, where all lower order terms have been collected into the functions $d_1(t, K)$ and $d_2(t, K)$. \square

Our final task is to prove upper bounds for the last term on the right-hand side of (3.3).

Lemma 4.34. *Fix $0 < \delta < 1/100$. Let (h, h^{hol}) be the pair of test functions (4.1) with $K, L \in 2\mathbb{N}$ satisfying $K^{\delta/8} \leq L \leq K^{1/3-\delta/8}$. Then for $\tilde{h}_{z_1, z_2, z_3, z_4}$ as in (3.9), we have that*

$$(4.35) \quad \lim_{(z_1, z_2, z_3, z_4) \rightarrow (\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2})} \tilde{h}_{z_1, z_2, z_3, z_4} \ll_{\varepsilon} K^{1+\varepsilon} L.$$

Proof. Let $0 < |z| \leq \varepsilon$ and $z_j = \frac{1}{2} + jz$ for $1 \leq j \leq 4$, a choice that ensures that the point (z_1, z_2, z_3, z_4) does not lie on the polar lines $z_n = z_m$ and $z_n + z_m = 1$ for $1 \leq n < m \leq 4$. We need to find an upper bound for the limit of $\tilde{h}_{z_1, z_2, z_3, z_4}$ as z tends to 0 in terms of K and L .

In the definition (3.9) of $\tilde{h}_{z_1, z_2, z_3, z_4}$, we see that the second sum vanishes because $h(t)$ is identically zero by construction; see (4.1). The third sum involves the integrals $\tilde{h}_{z_1, z_2, z_3, z_4; j}^{\pm}$. We deform, in a rightward direction, the contours \mathcal{C}_j of these integrals to the vertical line $\Re(s) = 1 + 100\varepsilon$. We denote the shifted integrals by $\tilde{h}_{z_1, z_2, z_3, z_4; j}^{\pm, \text{shifted}}$ and note that they are holomorphic for $|z| \leq \varepsilon$. In deforming the contour of integration, we cross simple poles of $\Gamma(1 - z_m - \frac{s}{2})$ at $s = 2 - 2z_m$ for $1 \leq m \leq 4$, and this gives rise to residues, which we denote $\tilde{h}_{z_1, z_2, z_3, z_4; j}^{\pm, \text{residue}}$. These residues are equal to the product of $\widehat{H}(2(1 - z_m))$ and other meromorphic functions of z that do not depend on K and L .

We now have an expression for $\tilde{h}_{z_1, z_2, z_3, z_4}$ as the first sum in (3.9) plus sums like the third one in (3.9) but with $\tilde{h}_{z_1, z_2, z_3, z_4; j}^{\pm}$ replaced by the sum of $\tilde{h}_{z_1, z_2, z_3, z_4; j}^{\pm, \text{shifted}}$ and $\tilde{h}_{z_1, z_2, z_3, z_4; j}^{\pm, \text{residue}}$. Although various individual terms in these sums may have poles at $z = 0$, arising from the presence of $\zeta(1 + nz)$ and $\Gamma(nz)$ values for nonzero integers n , the total expression $\tilde{h}_{z_1, z_2, z_3, z_4}$ cannot have a pole at $z = 0$ because (3.3) shows that the limit $\lim_{z \rightarrow 0} \tilde{h}_{z_1, z_2, z_3, z_4}$ is finite. By Cauchy's integral formula, we can compute this limit as

$$\frac{1}{2\pi i} \oint_{|z|=\varepsilon} \frac{\tilde{h}_{z_1, z_2, z_3, z_4}}{z} dz,$$

and so it suffices to bound $\tilde{h}_{z_1, z_2, z_3, z_4}$ for $|z| = \varepsilon$. To do this, we may bound all functions that do not depend on K and L by a constant. The only functions that depend on K and L and arise from the contribution of the first sum in (3.9) and $\tilde{h}_{z_1, z_2, z_3, z_4; j}^{\pm, \text{residue}}$ are $\widehat{H}(2(1 - z_m))$ for $1 \leq m \leq 4$. These are values of \widehat{H} evaluated close to 1, and by Lemma 4.2 we may bound them by $O_{\varepsilon}(K^{1+\varepsilon} L)$. The only function with K and L dependence that arises from the contribution of $\tilde{h}_{z_1, z_2, z_3, z_4; j}^{\pm, \text{shifted}}$ is $\widehat{H}(s)$, appearing in the integral over the line $\Re(s) = 1 + 100\varepsilon$. By the rapid decay of the gamma function in the integrand, this integral can be truncated to $|\Im(s)| \leq K^{\varepsilon}$ up to a negligible error. Thus $\widehat{H}(s)$ is also a value of \widehat{H} evaluated close to 1, and we may bound it by $O_{\varepsilon}(K^{1+100\varepsilon} L)$. Since $\varepsilon > 0$ was arbitrary, we deduce the requisite bound (4.35). \square

5. PROOF OF LEMMA 2.3

With the estimates and asymptotics for $\tilde{h}^{\pm}(t)$ and $\tilde{h}^{\text{hol}}(k)$ in hand via Lemmata 4.5 and 4.21, we may now proceed towards the proof of Lemma 2.3. Our strategy is to apply Kuznetsov's formula (3.3) with the choice of pair of test functions (h, h^{hol}) given by (4.1). To control the size of the dual fourth moments appearing on the right-hand side of (3.3), we require the following well-known moment bounds.

Lemma 5.1. *For $T \geq 1$, we have that*

$$(5.2) \quad \left. \begin{aligned} & \sum_{\substack{f \in \mathcal{B}_0 \\ T \leq t_f \leq 2T}} \frac{L(\frac{1}{2}, f)^4}{L(1, \text{ad } f)} \\ & \frac{1}{2\pi} \int_{T \leq |t| \leq 2T} \left| \frac{\zeta(\frac{1}{2} + it)^4}{\zeta(1 + 2it)} \right|^2 dt \\ & \sum_{\substack{f \in \mathcal{B}_{\text{hol}} \\ T \leq k_f \leq 2T}} \frac{L(\frac{1}{2}, f)^4}{L(1, \text{ad } f)} \end{aligned} \right\} \ll_{\varepsilon} T^{2+\varepsilon}.$$

Proof. Each of these estimates is a standard application of the spectral large sieve. \square

We now use Kuznetsov's formula (3.3) to give an upper bound for large values of the fourth moment of $L(1/2, f)$ in short intervals.

Lemma 5.3. *Fix $0 < \delta < 1/100$. Let $T \geq 1$, $K \in 2\mathbb{N}$ with $T \leq K \leq 2T$, $T^{1+\delta/2} \leq V \leq T^{4/3}(\log T)^7$, and define $L := 2 \lfloor \frac{1}{2} VT^{-1-\delta/4} \rfloor$. Then*

$$(5.4) \quad \sum_{\substack{K-L \leq k \leq K+L \\ k \equiv 0 \pmod{2} \\ V \leq \mathcal{M}^{\text{hol}}(k) \leq 2V}} \mathcal{M}^{\text{hol}}(k) \ll \left| \sum_{\substack{f \in \mathcal{B}_0 \\ T^{\frac{2}{3}+\varepsilon} < t_f < \frac{T \log T}{L}}} \frac{L(\frac{1}{2}, f)^4}{L(1, \text{ad } f)} g(t_f, K, L) \right| + \left| \int_{T^{\frac{2}{3}+\varepsilon} < |t| < \frac{T \log T}{L}} \left| \frac{\zeta(\frac{1}{2} + it)^4}{\zeta(1 + 2it)} \right|^2 g(t, K, L) dt \right|$$

with $g(t, K, L)$ as in (4.23).

Proof. From Lemma 4.2 (2) (b) and the fact that $L(1/2, f) = 0$ whenever $k_f \equiv 2 \pmod{4}$, we have that

$$\sum_{\substack{K-L \leq k \leq K+L \\ k \equiv 0 \pmod{2}}} \mathcal{M}^{\text{hol}}(k) \ll \sum_{f \in \mathcal{B}_0} \frac{L(\frac{1}{2}, f)^4}{L(1, \text{ad } f)} h(t_f) + \frac{1}{2\pi} \int_{-\infty}^{\infty} \left| \frac{\zeta(\frac{1}{2} + it)^4}{\zeta(1 + 2it)} \right|^2 h(t) dt + \sum_{f \in \mathcal{B}_{\text{hol}}} \frac{L(\frac{1}{2}, f)^4}{L(1, \text{ad } f)} h^{\text{hol}}(k_f)$$

with (h, h^{hol}) the pair of test functions as in (4.1). We apply Theorem 3.1 with this choice of test functions and bound each term on the right-hand side of (3.3) individually. Via the bounds (4.6) and (4.7) for $\tilde{h}^-(t)$ and $\tilde{h}^{\text{hol}}(k)$ and the spectral large sieve bounds (5.2), we have that

$$\left. \begin{aligned} & \sum_{f \in \mathcal{B}_0} \frac{L(\frac{1}{2}, f)^4}{L(1, \text{ad } f)} \tilde{h}^-(t_f) \\ & \frac{1}{2\pi} \int_{-\infty}^{\infty} \left| \frac{\zeta(\frac{1}{2} + it)^4}{\zeta(1 + 2it)} \right|^2 \tilde{h}^-(t) dt \\ & \sum_{f \in \mathcal{B}_{\text{hol}}} \frac{L(\frac{1}{2}, f)^4}{L(1, \text{ad } f)} \tilde{h}^{\text{hol}}(k_f) \end{aligned} \right\} \ll T^{-10}.$$

Next, the bounds and asymptotics (4.22) for $\tilde{h}^+(t)$ together with the spectral large sieve bounds (5.2) yield

$$\begin{aligned}
(5.5) \quad & \sum_{f \in \mathcal{B}_0} \frac{L(\frac{1}{2}, f)^4}{L(1, \text{ad } f)} \tilde{h}^+(t_f) + \frac{1}{2\pi} \int_{-\infty}^{\infty} \left| \frac{\zeta(\frac{1}{2} + it)^4}{\zeta(1 + 2it)} \right|^2 \tilde{h}^+(t) dt \\
&= \sum_{\substack{f \in \mathcal{B}_0 \\ T^{\frac{2}{3} + \varepsilon} < t_f < \frac{T \log T}{L}}} \frac{L(\frac{1}{2}, f)^4}{L(1, \text{ad } f)} g(t_f, K, L) + \frac{1}{2\pi} \int_{T^{\frac{2}{3} + \varepsilon} < |t| < \frac{T \log T}{L}} \left| \frac{\zeta(\frac{1}{2} + it)^4}{\zeta(1 + 2it)} \right|^2 g(t, K, L) dt \\
& \qquad \qquad \qquad + O_\varepsilon(T^{1+\varepsilon} L).
\end{aligned}$$

Finally, the last term on the right-hand side of (3.3) is $O_\varepsilon(T^{1+\varepsilon} L)$ by (4.35). We end by noting that the left-hand side of (5.4) is either 0 or at least as large as V . Since $T^{1+\delta/2} \leq V \leq T^{4/3}(\log T)^7$ and $L = 2 \lfloor \frac{1}{2} V T^{-1-\delta/4} \rfloor$, so that $T^{1+\varepsilon} L = o(V)$, it follows that the only terms that may contribute towards an upper bound for (5.4) are those appearing in (5.5). \square

We are interested in the second moment of the left-hand side of (5.4) averaged over even integers K in the interval $[T, 2T]$. Bearing in mind the upper bound given by the two terms on the right-hand side of (5.4), we are led to studying the average behaviour of $g(t_1, K, L) \overline{g(t_2, K, L)}$. It is here that we take advantage of the oscillatory behaviour of $g(t, K, L)$.

Lemma 5.6. *Fix $0 < \delta < 1/100$. Let $T \geq 1$, $T^{1+\delta/2} \leq V \leq T^{4/3}(\log T)^7$, and define $L := 2 \lfloor \frac{1}{2} V T^{-1-\delta/4} \rfloor$. Let Ω be a smooth function compactly supported on $(1/2, 5/2)$ that is nonnegative, equal to 1 on $[1, 2]$, and has bounded derivatives. Then for $T^{2/3+\varepsilon} < |t_1|, |t_2| < T(\log T)/L$, we have that*

$$(5.7) \quad \sum_{\substack{K=2 \\ K \equiv 0 \pmod{2}}}^{\infty} \Omega\left(\frac{K}{T}\right) g(t_1, K, L) \overline{g(t_2, K, L)} \ll_A \frac{TL^2}{\sqrt{|t_1 t_2|}} \left(1 + \sum_{\pm} |t_1 \pm t_2|\right)^{-A}$$

for any $A \geq 0$, where $g(t, K, L)$ is as in (4.23).

Proof. By the Poisson summation formula, the left-hand side of (5.7) is equal to

$$\sum_{\ell=-\infty}^{\infty} \frac{T}{2} \int_{-\infty}^{\infty} \Omega(x) g(t_1, Tx, L) \overline{g(t_2, Tx, L)} \left(1 + e\left(\frac{Tx}{2}\right)\right) e(-\ell Tx) dx$$

upon making the change of variables $x \mapsto Tx$.

For $\ell \neq 0$, we repeatedly integrate by parts, antidifferentiating $e(-\ell Tx)$ and differentiating the rest. Recalling the definition (4.23) of $g(t, K, L)$ and the derivative bounds (4.24), we see that each summand with $\ell \neq 0$ is

$$\ll_A \frac{TL^2}{\sqrt{|t_1 t_2|}} \left(\frac{|\ell|T}{|t_1|} + \frac{|\ell|T}{|t_2|} \right)^{-A}$$

for any $A \geq 0$. Since $|t_1|, |t_2| < T(\log T)/L$ and $L \geq T^{\delta/8}$, we deduce that

$$\begin{aligned}
& \sum_{\substack{\ell=-\infty \\ \ell \neq 0}}^{\infty} \frac{T}{2} \int_{-\infty}^{\infty} \Omega(x) g(t_1, Tx, L) \overline{g(t_2, Tx, L)} \left(1 + e\left(\frac{Tx}{2}\right)\right) e(-\ell Tx) dx \\
& \qquad \qquad \qquad \ll_A \frac{TL^2}{\sqrt{|t_1 t_2|}} \left(1 + \sum_{\pm} |t_1 \pm t_2|\right)^{-A}.
\end{aligned}$$

For $\ell = 0$ and $|t_1 - t_2| \ll 1$, we simply bound the integral trivially, so that

$$\frac{T}{2} \int_{-\infty}^{\infty} \Omega(x) g(t_1, Tx, L) \overline{g(t_2, Tx, L)} \left(1 + e\left(\frac{Tx}{2}\right)\right) dx \ll \frac{TL^2}{\sqrt{|t_1 t_2|}}.$$

Finally, for $\ell = 0$ and $|t_1 - t_2| \gg 1$, we instead repeatedly integrate by parts, antidifferentiating $x^{\pm 1} 2it_1 \mp 2it_2$ and differentiating the rest. We deduce that

$$\begin{aligned} \frac{T}{2} \int_{-\infty}^{\infty} \Omega(x) g(t_1, Tx, L) \overline{g(t_2, Tx, L)} \left(1 + e\left(\frac{Tx}{2}\right)\right) dx \\ \ll_A \frac{TL^2}{\sqrt{|t_1 t_2|}} \left(\sum_{\pm} |t_1 \pm t_2| + \frac{T^2}{|t_1|^2} + \frac{T^2}{|t_2|^2} \right)^{-A} \\ \ll_A \frac{TL^2}{\sqrt{|t_1 t_2|}} \sum_{\pm} |t_1 \pm t_2|^{-A} \end{aligned}$$

for any $A \geq 0$, having again recalled that $|t_1|, |t_2| < T(\log T)/L$ and $L \geq T^{\delta/8}$. \square

We are now in a position to complete the proof of [Lemma 2.3](#).

Proof of Lemma 2.3. From [Lemmata 5.3](#) and [5.6](#) and Young's inequality, we have that

$$\begin{aligned} \sum_{\substack{T \leq K \leq 2T \\ K \equiv 0 \pmod{2}}} \left(\sum_{\substack{K-L \leq k \leq K+L \\ k \equiv 0 \pmod{2} \\ V \leq \mathcal{M}^{\text{hol}}(k) \leq 2V}} \mathcal{M}^{\text{hol}}(k) \right)^2 \\ \ll \sum_{\substack{K=2 \\ K \equiv 0 \pmod{2}}}^{\infty} \Omega\left(\frac{K}{T}\right) \left| \sum_{\substack{f \in \mathcal{B}_0 \\ T^{\frac{2}{3}+\varepsilon} < t_f < \frac{T \log T}{L}}} \frac{L\left(\frac{1}{2}, f\right)^4}{L(1, \text{ad } f)} g(t_f, K, L) \right|^2 \\ + \sum_{\substack{K=2 \\ K \equiv 0 \pmod{2}}}^{\infty} \Omega\left(\frac{K}{T}\right) \left| \int_{T^{\frac{2}{3}+\varepsilon} < |t| < \frac{T \log T}{L}} \left| \frac{\zeta\left(\frac{1}{2} + it\right)^4}{\zeta(1 + 2it)} \right|^2 g(t, K, L) dt \right|^2 \\ \ll_A TL^2 \sum_{\substack{f_1, f_2 \in \mathcal{B}_0 \\ T^{\frac{2}{3}+\varepsilon} < t_{f_1}, t_{f_2} < \frac{T \log T}{L}}} \frac{L\left(\frac{1}{2}, f_1\right)^4 L\left(\frac{1}{2}, f_2\right)^4}{L(1, \text{ad } f_1) L(1, \text{ad } f_2)} \frac{1}{\sqrt{t_{f_1} t_{f_2}}} (1 + |t_{f_1} - t_{f_2}|)^{-A} \\ + TL^2 \iint_{T^{\frac{2}{3}+\varepsilon} < t_1, t_2 < \frac{T \log T}{L}} \left| \frac{\zeta\left(\frac{1}{2} + it_1\right)^4 \zeta\left(\frac{1}{2} + it_2\right)^4}{\zeta(1 + 2it_1) \zeta(1 + 2it_2)} \right|^2 \frac{1}{\sqrt{t_1 t_2}} (1 + |t_1 - t_2|)^{-A} dt_1 dt_2 \\ \ll_A TL^2 \iint_{T^{\frac{2}{3}+\varepsilon} < x_1, x_2 < \frac{T \log T}{L}} \left(\sum_{\substack{f \in \mathcal{B}_0 \\ x_1 \leq t_f \leq x_1+1}} \frac{L\left(\frac{1}{2}, f\right)^4}{L(1, \text{ad } f)} \right)^2 \frac{1}{\sqrt{x_1 x_2}} (1 + |x_1 - x_2|)^{-A} dx_1 dx_2 \\ + TL^2 \iint_{T^{\frac{2}{3}+\varepsilon} < t_1, t_2 < \frac{T \log T}{L}} \left| \frac{\zeta\left(\frac{1}{2} + it_1\right)^4}{\zeta(1 + 2it_1)} \right|^4 \frac{1}{\sqrt{t_1 t_2}} (1 + |t_1 - t_2|)^{-A} dt_1 dt_2. \end{aligned}$$

The integral over x_2 is $O(x_1^{-1/2})$ and the integral over t_2 is $O(t_1^{-1/2})$, so that

$$\begin{aligned} \sum_{\substack{T \leq K \leq 2T \\ K \equiv 0 \pmod{2}}} \left(\sum_{\substack{K-L \leq k \leq K+L \\ k \equiv 0 \pmod{2} \\ V \leq \mathcal{M}^{\text{hol}}(k) \leq 2V}} \mathcal{M}^{\text{hol}}(k) \right)^2 \\ \ll TL^2 \int_{T^{\frac{2}{3}+\varepsilon}}^{\frac{T \log T}{L}} \left(\sum_{\substack{f \in \mathcal{B}_0 \\ x \leq t_f \leq x+1}} \frac{L\left(\frac{1}{2}, f\right)^4}{L(1, \text{ad } f)} \right)^2 \frac{dx}{x} + TL^2 \int_{T^{\frac{2}{3}+\varepsilon}}^{\frac{T \log T}{L}} \left| \frac{\zeta\left(\frac{1}{2} + it\right)^4}{\zeta(1 + 2it)} \right|^4 \frac{dt}{t}. \end{aligned}$$

To bound these two terms, we dyadically subdivide both integrals, additionally using the classical lower bound $|\zeta(1+it)| \gg 1/\log(3+|t|)$ for $t \in \mathbb{R}$ for the latter. We then appeal to the bounds

$$\int_U^{2U} \left(\sum_{\substack{f \in \mathcal{B}_0 \\ x \leq t_f \leq x+1}} \frac{L(\frac{1}{2}, f)^4}{L(1, \text{ad } f)} \right)^2 dx \ll_\varepsilon U^{3+\varepsilon},$$

$$\int_U^{2U} \left| \zeta\left(\frac{1}{2} + it\right) \right|^{16} dt \ll_\varepsilon U^{\frac{283}{108} + \varepsilon}$$

for any $U \geq 1$, with the former due to Jutila [Jut04, Theorem 1] and the latter due to Ivić [Ivi03, Theorem 8.3]. In this way, we deduce the desired upper bound $O_\varepsilon(T^{3+\varepsilon})$. \square

REFERENCES

- [BF21] Olga Balkanova and Dmitry Frolenkov, “Moments of L -Functions and the Liouville–Green Method”, *Journal of the European Mathematical Society* **23**:4 (2021), 1333–1380.
- [BKY13] Valentin Blomer, Rizwanur Khan, and Matthew Young, “Distribution of Mass of Holomorphic Cusp Forms”, *Duke Mathematical Journal* **162**:14 (2013), 2609–2644.
- [BLM19] Valentin Blomer, Xiaoqing Li, and Stephen D. Miller, “A Spectral Reciprocity Formula and Non-Vanishing for L -Functions on $\text{GL}(4) \times \text{GL}(2)$ ”, *Journal of Number Theory Prime* **205** (2019), 1–43.
- [CFKRS05] J. B. Conrey, D. W. Farmer, J. P. Keating, M. O. Rubinstein, and N. C. Snaith, “Integral Moments of L -Functions”, *Proceedings of the London Mathematical Society* **91**:1 (2005), 33–104.
- [CS07] J. B. Conrey and N. C. Snaith, “Applications of the L -Functions Ratios Conjectures”, *Proceedings of the London Mathematical Society* **94**:3 (2007), 594–646.
- [Fro20] Dmitry Frolenkov, “The Cubic Moment of Automorphic L -Functions in the Weight Aspect”, *Journal of Number Theory* **207** (2020), 247–281.
- [GR15] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products, Eighth Edition*, editors Daniel Zwillinger and Victor Moll, Academic Press, Burlington, 2015.
- [H-B78] D. R. Heath-Brown, “The Twelfth Power Moment of the Riemann-Function”, *The Quarterly Journal of Mathematics* **29**:4 (1978), 443–462.
- [HK20] Peter Humphries and Rizwanur Khan, “On the Random Wave Conjecture for Dihedral Maaß Forms”, *Geometric and Functional Analysis* **30**:1 (2020), 34–125.
- [HK22] Peter Humphries and Rizwanur Khan, “ L^p -Norm Bounds for Automorphic Forms via Spectral Reciprocity”, preprint (2022), 57 pages.
- [Ivi02] Aleksandar Ivić, “On the Moments of Hecke Series at Central Points”, *Functiones et Approximatio Commentarii Mathematici* **30** (2002), 49–82.
- [Ivi03] Aleksandar Ivić, *The Riemann Zeta-Function*, Dover Publications, Inc., Mineola, New York, 2003.
- [Iwa80] Henryk Iwaniec, “Fourier Coefficients of Cusp Forms and the Riemann Zeta-Function”, *Seminaire de Théorie des Nombres de Bordeaux* **18** (1980), 1–36.
- [Iwa02] Henryk Iwaniec, *Spectral Methods of Automorphic Forms, Second Edition*, Graduate Studies in Mathematics **53**, American Mathematical Society, Providence, 2002.
- [Jut87] M. Jutila, *Lectures on a Method in the Theory of Exponential Sums*, Tata Institute of Fundamental Research Lectures on Mathematics and Physics **80**, Springer–Verlag, Berlin, 1987.
- [Jut04] Matti Jutila, “The Twelfth Moment of Central Values of Hecke Series”, *Journal of Number Theory* **108**:1 (2004), 157–168.
- [Kha20] Rizwanur Khan, “The Fifth Moment of Hecke L -Functions in the Weight Aspect”, *Mathematical Proceedings of the Cambridge Philosophical Society* **168**:3 (2020), 543–566.
- [Kim03] Henry H. Kim, “Functoriality for the Exterior Square of GL_4 and the Symmetric Fourth of GL_2 ”, with appendix 1 by Dinakar Ramakrishnan and appendix 2 by Henry H. Kim and Peter Sarnak, *Journal of the American Mathematical Society* **16**:1 (2003), 139–183.
- [KZ81] W. Kohnen and D. Zagier, “Values of L -Series of Modular Forms at the Center of the Critical Strip”, *Inventiones Mathematicae* **64**:2 (1981), 175–198.
- [Kuz89] N. V. Kuznetsov, “Sums of Kloosterman Sums and the Eighth Power Moment of the Riemann Zeta-Function”, in *Number Theory and Related Topics*, Tata Institute of Fundamental Research: Studies in Mathematics **12**, Oxford University Press, 1989, 57–117.
- [Kuz99] N. V. Kuznetsov, “The Hecke Series at the Center of the Critical Strip”, preprint (1999), 27 pages.
- [MW21] Djordje Milićević and Daniel White, “Twelfth Moment of Dirichlet L -Functions to Prime Power Moduli”, *Annali della Scuola Normale Superiore di Pisa, Classe di Scienze* **22**:4 (2021), 1879–1898.
- [Mot97] Yoichi Motohashi, *Spectral Theory of the Riemann Zeta-Function*, Cambridge Tracts in Mathematics **127**, Cambridge University Press, Cambridge, 1997.

- [Mot03] Yoichi Motohashi, “A Functional Equation for the Spectral Fourth Moment of Modular Hecke L -Functions”, in *Proceedings of the Session in Analytic Number Theory and Diophantine Equations*, Bonner mathematische Schriften **360**, Universität Bonn, Bonn, 2003, 19 pages.
- [Nun21] Ramon M. Nunes, “The Twelfth Moment of Dirichlet L -Functions with Smooth Moduli”, *International Mathematics Research Notices* **2021**:12 (2021), 9180–9202.
- [Pen01] Zhuangzhuang Peng, *Zeros and Central Values of Automorphic L -Functions*, Ph.D. Thesis, Princeton University, 2001.
- [Qi22] Zhi Qi, “Proof of the Strong Ivić Conjecture for the Cubic Moment of Maass-Form L -Functions”, preprint (2022), 20 pages.
- [Sou21] Kannan Soundararajan, “The Distribution of Values of Zeta and L -Functions”, preprint (2021), 49 pages.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF VIRGINIA, CHARLOTTESVILLE, VA 22904, USA

Email address: pclhumphries@gmail.com

URL: <https://sites.google.com/view/peterhumphries/>

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSISSIPPI, UNIVERSITY, MS 38677, USA

Email address: rrkhan@olemiss.edu