A Stochastic Contraction Mapping Theorem

Anthony Almudevar Department of Biostatistics and Computational Biology University of Rochester

July 5, 2022

Abstract. In this paper we define contractive and nonexpansive properties for adapted stochastic processes X_1, X_2, \ldots which can be used to deduce limiting properties. In general, nonexpansive processes possess finite limits while contractive processes converge to zero *a.e.* Extensions to multivariate processes are given. These properties may be used to model a number of important processes, including stochastic approximation and least-squares estimation of controlled linear models, with convergence properties derivable from a single theory. The approach has the advantage of not in general requiring analytical regularity properties such as continuity and differentiability.

Keywords. 93E20 Optimal stochastic control; 93E24 Least squares and related methods; 93E35 Stochastic learning and adaptive control

1 Introduction

Let $\mathbf{X} = \{X_n : n \ge 0\}$ be a sequence of random variables adapted to filtration $\tilde{\mathcal{F}} = \{\mathcal{F}_n : n \ge 0\}$. Following the Doob decomposition theorem we may always write

$$X_n = X_{n-1} + A_n + \epsilon_n, \ n \ge 1,$$
(1.1)

where $\epsilon_n = X_n - E[X_n | \mathcal{F}_{n-1}]$ are \mathcal{F}_n -measurable martingale differences, and $A_n = E[X_n | \mathcal{F}_{n-1}] - X_{n-1}$ defines a predictable process, in the sense that A_n is \mathcal{F}_{n-1} -measurable. We let X_0 be any suitable \mathcal{F}_0 -measurable initial value. We can think of A_n as a control effector, and we will sometimes have $A_n = f_n(X_{n-1})$ for some sequence of deterministic mappings f_n , $n \ge 1$. In this case, we can imagine setting $\epsilon_n = 0$ in Equation (1.1), obtaining a deterministic iterative process. To fix ideas, suppose each f_n has common fixed point $0 = f_n(0)$ and that there exists positive constants ρ_n such that $0 \le f_n(x)/x \le \rho_n$ when $x \ne 0$. In a manner similar to the Banach fixed point theorem, it can be shown that if $\prod_{n\ge 1} \rho_n = 0$, then $\lim_n X_n = 0$. We can refer to this as a contraction property, although we need not assume f_n is formally a contraction mapping (or even Lipschitz continuous). If we restore ϵ_n to Equation (1.1), the question becomes how the additional stochastic variation affects the convergence properties.

In Almudevar (2008) (see also Almudevar (2014)) a general theorem for such noisy iterative algorithms was developed, for which X_n is defined on any Banach space $(\mathcal{X}, \|\cdot\|)$ (not necessarily stochastic). For example, suppose operator T_n possesses Lipschitz constant $\rho_n \leq \rho < 1$, and x^* is the fixed point of each T_n . In this case the "exact algorithm" $X_n = T_n(X_{n-1}), n \geq 1$ converges to x^* . The "approximate algorithm" $X_n = T_n(X_{n-1}) + \epsilon_n$, $n \ge 1$, is then a noisy version of the exact algorithm. Then if $\limsup_n \|\epsilon_n\| = d$, we may claim that $\limsup_n \|X_n - x^*\| = O(d)$, and if d = 0 then X_n converges to x^* . In fact, the convergence rate of $\|X_n - x^*\|$ is bounded by $O(\max(\|\epsilon_n\|, (\rho + \delta)^n))$ for any $\delta > 0$ (Almudevar, 2008, 2014).

The theory extends to nonexpansive operators T_n ($\rho_n \leq 1$), especially "weakly contractive" sequences which satisfy $\prod_{n\geq 1} \rho_n = 0$, while, typically, the limit $\lim_n \rho_n = 1$ also holds. In this case, stronger convergence results are obtainable when $(\mathcal{X}, \|\cdot\|)$ is assumed to be additionally a Hilbert space (for example, adapted stochastic processes endowed with the L_2 norm).

In this paper, we extend some of these ideas to filtered stochastic processes, with strong convergence in place of convergence in a Banach space. We will characterize precisely contractive or nonexpansive properties based on the predictable process A_n of Equation (1.1). These force X_n to behave like a supermartingale above 0, and a submartingale below. If $M_{1,n} = \sum_{i=1}^{n} \epsilon_i$ is a martingale of finite variation, it follows naturally that the process will possess a finite limit. However, if we then impose sufficient contractive properties, it can be shown that X_n converges to 0 *a.e.*

The main result is given in Section 2 for $X_n \in \mathbb{R}^1$. We introduce the nonexpansive and contractive properties, then Theorem 2.1 summarizes the important limiting properties.

In Section 3 we consider the "nonuniform contractive" case, by which we mean that the nonexpansive or contractive properties required of Theorem 2.1 are now required to hold only outside a neighborhood of zero (Theorems 3.1 and 3.2).

In Section 4, Theorem 2.1 is extended to $\mathbf{X} = {\mathbf{X}_n \in \mathbb{R}^p : n \ge 0}, p \ge 1$, by replacing the constraint $0 \le E[X_n | \mathcal{F}_{n-1}]/X_{n-1} \le \rho$ with $||E[\mathbf{X}_n | \mathcal{F}_{n-1}]|/||\mathbf{X}_{n-1}|| \le \rho$, $||\mathbf{X}_{n-1}|| \ne 0$. It is then quite straightforward to apply Theorem 2.1 to $||\mathbf{X}_n|| \in \mathbb{R}^1$ (the conditions of which permit the process to be, for example, strictly positive). The argument is given in Theorem 4.1.

We include two important applications which can be modeled using the proposed theory. The Robbins-Monro stochastic approximation algorithm is a filtered stochastic process intended to converge to the solution to $g(\mathbf{x}) = \mathbf{0}$ for a given mapping $g : \mathbb{R}^p \to \mathbb{R}^p$ (Robbins and Monro, 1951). The univariate case will demonstrate the main theorem of Section 2. Extension to the nonuniform contractive model will significantly expand the class of g to which stochastic approximation can be applied. The multivariate extension is then developed. This may be of some interest, since regularity conditions usually imposed on g for $p \geq 1$ are not needed by the methods introduced here (Spall, 1992; Pham *et al.*, 2009; Lai, 2003; Kushner and Yin, 2003; Pham *et al.*, 2009).

In Section 5 we consider the controlled least-squares regression model introduced in Christopeit and Helmes (1980). We can directly derive the same conditions for strong convergence, while also proving that they are also necessary. In general, we can prove that the least-squares estimator always possess a finite limit, hence weak convergence implies strong convergence.

Finally, we bring attention to a machine learning application proposed in Zhou and Hooker (2018), involving regularization of stochastic gradient boosted trees. This makes use of an earlier version of the methodology proposed here that appeared in an unpublished preprint written by this author. It demonstrates how the approach can be used to prove strong consistency where differentiability cannot be assumed.

Throughout the manuscript proofs and a number of technical lemmas are given in the appendix.

2 Main Result

Let $\mathbf{X} = \{X_n : n \ge 0\}$ be an L_1 stochastic process defined on probability space (Ω, \mathcal{F}, P) . Suppose there is a filtration $\tilde{\mathcal{F}}$ defined defined by the σ -algebras $\mathcal{F}_0 \subset \mathcal{F}_1 \subset \mathcal{F}_2 \subset \cdots \subset \mathcal{F}$ to which \mathbf{X} is adapted. Define residuals

$$\epsilon_n = X_n - E[X_n \mid \mathcal{F}_{n-1}], \ n \ge 1, \tag{2.1}$$

and the partial sums $M_{s,t} = \sum_{i=s}^{t} \epsilon_i$ for $t \ge s \ge 1$, with $M_{s,t} = 0$ whenever t < s. Then define the events

$$D_n^+ = \{X_n > 0\}, \ D_n^- = \{X_n < 0\}, \ D_n^0 = \{X_n = 0\}, \ n \ge 0, D_n = D_n^+ (D_{n-1}^+)^c \cup D_n^- (D_{n-1}^-)^c \cup D_n^0 (D_{n-1}^0)^c, \ n \ge 1,$$

and the compound events $D_{s,t}^+ = \bigcap_{i=s}^t D_i^+$, $D_{s,\infty}^+ = \bigcap_{i\geq s} D_i^+$, $0 \leq s \leq t$. Define similarly $D_{s,t}^-$, $D_{s,\infty}^-$, $D_{s,t}^0$ and $D_{s,\infty}^0$. We may then construct the increasing sequence of stopping times T_1, T_2, \ldots

$$T_j \le t \Leftrightarrow \sum_{i=1}^t I\{D_i\} \ge j$$

for $j \geq 1$. Then T_1, T_2, \ldots may be interpreted as the times at which sequence **X** changes sign, changes from zero to a nonzero value or changes from a nonzero value to zero. We can refer to any such event as a crossing. Note that T_j may be infinite, so it will be useful to define $N_T = \sup\{j \geq 1 : T_j < \infty\}$. Then let

$$W_j = \begin{cases} \sup_{T_j \le i < T_{j+1}} |X_i| & ; \quad T_j < \infty \\ 0 & ; \quad T_j = \infty \end{cases}$$

for $j \ge 1$. We will make use throughout the paper of the following sequence:

$$U_n = E[X_n \mid \mathcal{F}_{n-1}]I\{X_{n-1} = 0\}, \ n \ge 1.$$
(2.2)

Our model will be based on the following assumptions:

(A1) There exist nonnegative constants $\alpha_n \ge 0$, $n \ge 1$, such that

$$0 \le \frac{E[X_n \mid \mathcal{F}_{n-1}]}{X_{n-1}} I\{X_{n-1} \ne 0\} \le 1 + \alpha_n, \ a.e$$

and $\sum_{i\geq 1} \alpha_i < \infty$.

(A2) There exist nonnegative constants $0 \le k_n \le 1$, $n \ge 1$, such that

$$0 \le (E[X_n \mid \mathcal{F}_{n-1}]/X_{n-1}) I\{X_{n-1} \ne 0\} \le k_n, \ a.e.$$

and $\sum_{i \ge 1} (1 - k_i) = \infty$ (equivalently, $\prod_{i \ge s} k_i = 0$ for all $s \ge 1$). (A3) $\lim_{n \to \infty} U_n = 0$ a.e.

(A4) The partial sums $M_{1,n}$ possess a finite limit *a.e.* as $n \to \infty$.

Assumption (A1) is analogous to a nonexpansive condition on $E[X_n | \mathcal{F}_{n-1}]$, and is related to the "almost martingale" introduced in Robbins and Siegmund (1971). Assumption (A2) is the stronger, or contractive, version of (A1). The central result of this section is that under assumptions (A3) and (A4) the process defined by (A1) converges to a finite limit, while that defined by (A2) converges to 0.

In some applications, (A3) will be a natural extension of (A1) or (A2) to the case $X_{n-1} = 0$. That is, we may be able to claim $0 \leq E[X_n | \mathcal{F}_{n-1}] \leq X_{n-1}(1 + \alpha_n)$ for $X_{n-1} \geq 0$ and $0 \geq E[X_n | \mathcal{F}_{n-1}] \geq X_{n-1}(1 + \alpha_n)$ for $X_{n-1} \leq 0$. In this case (A3) holds trivially. However, we find that our theory can be extended to a broader class of interesting models by permitting somewhat more flexibility when conditioning on $\{X_{n-1} = 0\}$. Finally, since $M_{1,n}$ $n \geq 1$ is a martingale, conditions under which (A4) holds are well known. For example, if $M_{1,n}$ is an L_2 martingale, by Theorem 4.5.2 of Durrett (2019), we may claim

$$\sum_{i=1}^{\infty} E[\epsilon_i^2 \mid \mathcal{F}_{i-1}] < \infty \ a.e. \ \text{implies} \ \lim_n M_{1,n} \text{ exists and is finite } a.e.$$
(2.3)

We are now give the main results. We first consider the nonexpansive model. Given (A1), (A3) and (A4) we may claim that **X** possesses a finite limit. When (A2) holds as well we have convergence to zero. It will be useful, however, to consider the case for which (A3) need not hold. The results are essentially partitioned into the cases $\{N_T < \infty\}$ and $\{N_T = \infty\}$, since the analysis differs between these cases in some important ways. For example, on $\{N_T = \infty\}$ the nonexpansive property (A1) is sufficient for convergence to zero.

Theorem 2.1 Suppose \mathbf{X} satisfies (A1) and (A4). Then the following statements hold.

- (i) $X_n I\{N_T < \infty\}$ possesses a finite limit *a.e.* as $n \to \infty$.
- (ii) If in addition (A2) holds then $X_n I\{N_T < \infty\}$ converges to zero *a.e.*
- (iii) The inequality $\limsup_n |X_n| I\{N_T = \infty\} \le \limsup_n |U_n|$ holds a.e.
- (iv) If in addition (A3) holds, then $\lim_n X_n I\{N_T = \infty\} = 0$ a.e.

2.1 The Strong Law of Large Numbers as a Contractive Process

We now discuss the relationship between Theorem 2.1 and the strong law of large numbers. We suppose that there is an L_1 sequence of random variables $\mathbf{Y} = \{Y_n : n \ge 0\}$ on a probability space (Ω, \mathcal{F}, P) such that $S_n = \sum_{i=0}^n Y_i$, $n \ge 0$, is a martingale with respect to some filtration $\tilde{\mathcal{F}}$.

Suppose a_0, a_1, a_2, \ldots is a nondecreasing sequence of positive real numbers such that $a_n \to_n \infty$. Then for any $s \ge 1$,

$$\lim_{n \to \infty} \prod_{i=s}^{n} \frac{a_{i-1}}{a_i} = \lim_{n \to \infty} \frac{a_{s-1}}{a_n} = 0.$$
(2.4)

We are interested in conditions under which $S_n/a_n \to_n 0$ a.e. We first note that, following Equation (2.4), the sequence $X_n = S_n/a_n$ satisfies assumption (A2), since

$$\frac{E[X_n \mid \mathcal{F}_{n-1}]}{X_{n-1}} I\{X_{n-1} \neq 0\} = \frac{S_{n-1}/a_n}{S_{n-1}/a_{n-1}} I\{X_{n-1} \neq 0\} \le \frac{a_{n-1}}{a_n}, \ n \ge 1.$$

Then (A3) holds since $E[X_n | \mathcal{F}_{n-1}]I\{X_{n-1} = 0\} = (S_{n-1}/a_n)I\{X_{n-1} = 0\} =$ 0. The residual becomes $\epsilon_n = X_n - E[X_n \mid \mathcal{F}_{n-1}] = S_n/a_n - S_{n-1}/a_n = Y_n/a_n$. Direct application of Theorem 2.1 gives

Theorem 2.2 Let $S_n = \sum_{i=0}^n Y_i$ be an L_1 martingale with respect to filtration $\tilde{\mathcal{F}}$. If $a_n \uparrow_n \infty$, and if $\sum_{i=1}^{\infty} Y_i/a_i$ is a convergent series *a.e.* then $S_n/a_n \to_n 0$ a.e.

But this is simply a restatement of Kronecker's lemma, which states that if $a_n \uparrow_n \infty$ then for any sequence of real numbers x_1, x_2, \ldots the convergence of the series $\sum_{i=1}^{\infty} x_i/a_i$ implies $\lim_{n \to \infty} (1/a_n) \sum_{i=1}^n x_i = 0$.

2.2**Stochastic Approximation**

The stochastic approximation algorithm is a method used to determine the solution $x^* \in E_g \subset \mathbb{R}$ to

$$g(x) = 0 \tag{2.5}$$

where mapping $g: E_q \to \mathbb{R}$ can only be evaluated with noise. In Robbins and Monro (1951) the iterative algorithm

$$X_n = X_{n-1} - \alpha_n U_n, \quad n \ge 1 \tag{2.6}$$

was proposed, giving conditions under which X_n converges in L_2 to x^* . Here, α_n is a sequence of positive constants, and the sequence X_n is adapted to filtration \mathcal{F} and constructed so that $E[U_n \mid \mathcal{F}_{n-1}] = g(X_{n-1}).$

Strong convergence has been since been established under a variety of conditions (Blum, 1954; Robbins and Siegmund, 1971; Ljung, 1978; Lai, 2003; Kushner and Yin, 2003; Benveniste et al., 2012). Our first task with respect to stochastic approximation will be to show how it may be interpreted as a stochastic contraction process.

Given a sequence \mathbf{X} constructed by the iterations of Equation (2.6) we define the following assumptions.

- (B1) The mapping $g: E_q \to \mathbb{R}$, possesses root $x^* \in E_q$.
- (B2) $P(X_n \in E_g) = 1$ for all $n \ge 1$.
- (B3) There exists $0 < m \le M < \infty$ such that $m \le g(x)/(x x^*) \le M$ for all $x \in E_g - \{x^*\}.$
- (B4) $\alpha_n \ge 0$, $\lim_{n \to \infty} \alpha_n = 0$. (B5) The series $\sum_{i=0}^{\infty} \alpha_i \{U_i E[U_i \mid \mathcal{F}_{i-1}]\}$ converges *a.e.* (B6) $\sum_{i=1}^{\infty} \alpha_i = \infty$.
- (B7) $\overline{E[U_n]} \mid \mathcal{F}_{n-1}] = g(X_{n-1}) \ a.e. \text{ for } n \ge 1.$

Assumption (B1) does not require that x^* be in the interior of E_g . Nor does it state that the root is unique, but this is forced by (B3). Then, if finite nonzero left and right derivatives of g exist at x^* , (B3) will hold on some open neighborhood of x^* , if not the entire domain of g.

It should be noted that, following Equation (2.3), if the variance of U_n conditional on \mathcal{F}_{n-1} is bounded, then (B5) may be replaced by $\sum_{i>1} \alpha_i^2 < \infty$, which, with (B6), are the conditions on the sequence α_n commonly given in the literature

We now state the main theorem of this section.

Theorem 2.3 If (B1)-(B7) hold then the process X defined by Equation (2.6) converges to x^* a.e.



Figure 3.1: Figure (a) shows the Lipschitz envelope defined by gradients M and m of condition (B3), which essentially represent sufficient conditions on the function g(x) of Equation (2.5) for convergence of the stochastic approximation algorithm using Theorem 2.1. Note that monotonicity and continuity are not required. Figure (b) gives a comparison of the regularity conditions for objective function g(x) based on Theorem 2.1, and the nonuniform contraction model of Theorem 3.1. The Lipschitz envelope shown in Figure 3.1a is superimposed here, along with a graphical representation of the comparable envelope forced by regularity conditions (C1)-(C3) introduced in Blum (1954). The important feature is that for any $\delta > 0$ we may construct an upper Lipschitz envelope which bounds above the upper envelope of (C1) for all $|x| > \delta$, and for any $0 < \delta_1 < \delta_2 < \infty$, we may construct a lower Lipschitz envelope which bounds below the lower envelope implied by (C3) (see the gray shaded area).

It is interesting to note that the stochastic approximation algorithm will converge if g(x) can be perfectly evaluated (in contrast with simulated annealing, which relies on stochastic variation for convergence). In this case Equation (2.6) becomes

$$x_n = x_{n-1} - \alpha_n g(x_{n-1}) = T_n(x_{n-1}), \ n \ge 1,$$

where T_n is a sequence of operators with Lipschitz constant $\rho_n = (1 - m\alpha_n)$ for all large enough n. Given (B6) we have $\prod_{i\geq 1} \rho_i = 0$, so that although the Lipschitz constants are not bounded away from one, they approach one from below slowly enough to allow contraction to force convergence. Two conditions on α_n are commonly associated with stochastic approximation, in particular, (a) $\sum_{i=1}^{\infty} \alpha_i = \infty$; and (b) $\sum_{i=1}^{\infty} \alpha_i^2 < \infty$. As can be seen, condition (a) forces sufficient contraction to ensure convergence for the noiseless algorithm, while condition (b) ensures that the cumulative effect of noise vanishes in the limit (as a consequence of contraction property). For more on this point of view, see Almudevar (2008, 2014).

3 Nonuniform Contractions

Figure 3.1a demonstrates that the conditions imposed on g(x) by Theorem 2.3 are quite general. Essentially, g(x) must be contained within a Lipschitz

envelope, but otherwise need not be monotone or continuous (for convenience we will assume $x^* = 0$).

However, a variety of conditions on g have been derived in the literature. In the seminal paper Robbins and Monro (1951), alternative conditions for L_2 convergence are given, in particular, $g(x) \leq -\delta$ for x < 0 and $g(x) \geq \delta$ for x > 0, for some $\delta > 0$ (Theorem 1); and g(0) = 0, g(x) is nondecreasing and g'(0) > 0 (Theorem 2). In Blum (1954), the following regularity conditions are offered:

- (C1) $|g(x)| \le c + d|x|$ for some constants $c, d \ge 0$;
- (C2) g(x) < 0 for x < 0 and g(x) > 0 for x > 0;
- (C3) $\inf_{\delta_1 \le |x| \le \delta_2} |g(x)| > 0$ for every pair of numbers δ_1, δ_2 for which $0 < \delta_1 < \delta_2 < \infty$.

While conditions (C1)-(C3) are considerably weaker than the envelope bound of Figure 3.1a, they can be usefully compared. If (C1) holds, then some Lipschitz upper bound (Figure 3.1a) holds for $|x| > \delta$. Then, with (C2), we can show that (A1) will hold *outside* a neighborhood \mathcal{N} of the origin. Then if (C3) holds we may show that (A2) holds for $\delta_1 \leq |x| \leq \delta_2$, for any positive constants $\delta, \delta_1 < \delta_2$. In this case, it will be possible to show, first, that **X** will be bounded in the limit. In this case, (A2) need only hold for a bounded subset of E_g , and we will be able to show that **X** approaches \mathcal{N} a.e., essentially "fattening" the origin. If the neighborhood \mathcal{N} can then be made arbitrarily small, we obtain strong convergence.

Thus, we can say that, in the context of stochastic approximation, (C1)-(C3) define models that are contractive outside any neighborhood of the origin, hence it seems reasonable to refer to such a model as "nonuniformly contractive". Below, we formally define such a model. We again define an adapted sequence \mathbf{X} , on which we may, as an alternative to (A1)-(A4), impose the following assumptions.

(D1) For $\delta > 0$ (A1) holds for $|X_{n-1}| > \delta$ a.e.

- (D2) For $0 < \delta_1 < \delta_2 < \infty$ (A2) holds for $|X_{n-1}| \in (\delta_1, \delta_2)$ a.e.
- (D3) $|X_{n-1}| \in (0, \delta]$ implies $|E[X_n | \mathcal{F}_{n-1}]| \le \kappa < \infty$ a.e.
- (D4) For all $\delta > 0$, (D3) holds for constants κ_{δ} which satisfy the limit $\lim_{\delta \downarrow 0} \kappa_{\delta} = 0$.

The key to the argument is to consider the process derived from **X**:

$$X_n^{\delta+\tau} = X_n I\{ |E[X_n | \mathcal{F}_{n-1}]| \ge \delta + \tau \}, \ n \ge 1.$$
(3.1)

for given $\delta, \tau > 0$ (it will prove to be more intuitive to introduce δ and τ separately in our notation). We denote this process $\mathbf{X}^{\delta+\tau}$. We may define the residuals in the same manner as in Equation (2.1):

$$\epsilon_n^{\delta+\tau} = X_n^{\delta+\tau} - E[X_n^{\delta+\tau} \mid \mathcal{F}_{n-1}], \ n \ge 1,$$
(3.2)

Finally, following Equation (2.2) we may condition the process on $\{X_{n-1}^{\delta+\tau}=0\}$:

$$U_n^{\delta+\tau} = E[X_n^{\delta+\tau} \mid \mathcal{F}_{n-1}]I\{X_{n-1}^{\delta+\tau} = 0\}.$$
 (3.3)

We briefly outline the remaining strategy. Suppose we can assume that $\epsilon_n \rightarrow_n 0$ a.e., where ϵ_n are the residuals of the original process **X**. This will be the case under (A4). Then under (D1), there exists wp1 a finite integer

 N_0 such that $\mathbf{X}^{\delta+\tau}$ satisfies (A1) for all $n \geq N_0$. Accordingly, we define the offset process $\mathbf{X}_{N_0}^{\delta+\tau} = \{X_{N_0+n}^{\delta+\tau} : n \geq 0\}$, so we may claim that $\mathbf{X}_{N_0}^{\delta+\tau}$ satisfies (A1). The remaining argument is of the same style, that is, by assuming the various conditions (D1)-(D4) hold for X, we may verify that the corresponding properties (A1)-(A4) hold for $\mathbf{X}_{N_0}^{\delta+\tau}$. Then the convergence properties of \mathbf{X} follow from those of $\mathbf{X}_{N_0}^{\delta+\tau}$. The first theorem follows.

Theorem 3.1 Suppose $\epsilon_n \rightarrow_n 0$ a.e. Suppose (D1) holds for some $\delta > 0$. Consider the derived process of Equation (3.1). Then

- (i) For all small enough $\tau > 0$, wp1 there exists finite N_0 such that $\mathbf{X}_{N_0}^{\delta+\tau}$ satisfies (A1) for all $n \ge 0$.
- (ii) If in addition, **X** satisfies (D2) for $\delta_1 \geq \delta$, then $\mathbf{X}_{N_0}^{\delta+\tau}$ satisfies (A2) whenever $|X_{N_0+n-1}^{\delta+\tau}| < \delta_2$. (iii) If in addition (D3) holds, then wp1

$$U_{N_0+n}^{\delta+\tau} \le (|U_{N_0+n}| + \delta + 2\tau + \kappa), \ n \ge 0.$$
(3.4)

(iv) If (A4) holds for **X**, then it also holds for $\mathbf{X}_{N_0}^{\delta+\tau}$.

Theorem 3.1 allows us to deduce some convergence properties of $\mathbf{X}^{\delta+\tau}$. If (D1) and (A4) hold for \mathbf{X} then (A1) and (A4) hold for $\mathbf{X}_{N_0}^{\delta+\tau}$. As a consequence, by Theorem 2.1 either $X_n^{\delta+\tau}$ converges to a finite limit, or $\limsup_{n\to\infty} |X_n^{\delta+\tau}| \leq \limsup_{n\to\infty} |U_n^{\delta+\tau}|$. The final step is to deduce the implications of this for the limit of X itself.

Theorem 3.2 Suppose **X** satisfies (D1) for all $\delta > 0$ *a.e.*, and in addition (D4), (A3) and (A4). Then **X** possesses a finite limit *a.e.* If in addition **X** satisfies (D2) for all pairs $0 < \delta_1 < \delta_2 < \infty$ then **X** converges to zero *a.e.*

3.1Stochastic Approximation and Nonuniform Contraction

We continue our discussion of stochastic approximation. Consider assumptions (B1)-(B7) (as before, assuming $x^* = 0$). Then replace assumption (B3), which is placed on g(x), with (C1)-(C3). Fix any $\delta > 0$. By (C1) $|g(x)| \leq c + d|x|$. With (C2) we may conclude

$$0 \le q(x)/x \le (c/\delta) + d$$
, for all $|x| > \delta$.

Since $\alpha_n \to_n 0$ we have, for all large enough n, following Equation (A.9),

$$0 \le \left(1 - \alpha_n \left(\frac{c}{\delta} + d\right)\right) I\{X_{n-1} \ne 0\} \le \frac{E[X_n \mid \mathcal{F}_{n-1}]}{X_{n-1}} I\{X_{n-1} \ne 0\} \le 1,$$

so that assumption (D1) will be satisfied. Thus, (C1)-(C2) force the nonexpansive property.

To verify (D4), note that by (C1)-(C2) $|E[X_n | \mathcal{F}_{n-1}]| \le |X_{n-1}|$ for $|X_{n-1}| \le |X_{n-1}|$ δ for all large enough n. Thus, (D4) holds by setting $\kappa_{\delta} = \delta$.

We next show that (C3) forces the contractive property. Fix $0 < \delta_1 < \delta_2 <$ ∞ . Define the quantity

$$K_{\delta_1,\delta_2} = \inf_{\delta_1 \le |x| \le \delta_2} g(x)/x.$$

Given (C2), under (C3) $K_{\delta_1,\delta_2} > 0$, so that, for all large enough n,

$$0 \le \frac{E[X_n \mid \mathcal{F}_{n-1}]}{X_{n-1}} I\{X_{n-1} \ne 0\} \le (1 - \alpha_n K_{\delta_1, \delta_2}).$$

Following the proof of Theorem 2.3, we may then conclude that (D2) holds for all finite pairs δ_1, δ_2 . Thus, by Theorem 3.2, the stochastic approximation algorithm will converge when (B3) is replaced by (C1)-(C3).

4 Multivariate Processes

We now extend Theorem 2.1 to processes in \mathbb{R}^p , replacing the ratio in (A1) and (A2) with the absolute value of the ratio. Suppose $\mathbf{X} = \{\mathbf{X}_n \in \mathbb{R}^p; n \geq 0\}$ is stochastic process adapted to filtration $\tilde{\mathcal{F}}$. The residual vector is $\boldsymbol{\epsilon}_n = \mathbf{X}_n - E[\mathbf{X}_n \mid \mathcal{F}_{n-1}], n \geq 1$. Then $E[\boldsymbol{\epsilon}_n \mid \mathcal{F}_{n-1}] = 0$, so that as before $\boldsymbol{\epsilon}_n, n \geq 1$ form vectors of martingale differences. Denote the components of \mathbf{X}_n and $\boldsymbol{\epsilon}_t$ by $\mathbf{X}_n(t) \in \mathbb{R}$ and $\boldsymbol{\epsilon}_n(t) \in \mathbb{R}, t \in \mathcal{T} = \{1, \dots, p\}$. Consider the following assumptions, which may be taken as multivariate extensions of (A1)-(A4) (here, $\|\cdot\|$ is the Euclidean norm).

(A5) There exist nonnegative constants $\alpha_n \ge 0$, $n \ge 1$, such that

$$\frac{\|E[\mathbf{X}_n \mid \mathcal{F}_{n-1}]\|}{\|\mathbf{X}_{n-1}\|} I\{\|\mathbf{X}_{n-1}\| \neq 0\} \le 1 + \alpha_n \ a.e.$$

and $\sum_{i>1} \alpha_i < \infty$.

(A6) There exists positive constants $k_n \leq 1, n \geq 1$, such that

$$\frac{\|E[\mathbf{X}_n \mid \mathcal{F}_{n-1}]\|}{\|\mathbf{X}_{n-1}\|} I\{\|\mathbf{X}_{n-1}\| \neq 0\} \le k_n \ a.e.$$

and $\sum_{i>1} (1-k_i) = \infty$ (equivalently, $\prod_{i>s} k_i = 0$ for all $s \ge 1$).

- (A7) $\lim_{n\to\infty} E[\mathbf{X}_n \mid \mathcal{F}_{n-1}]I\{\mathbf{X}_{n-1}=0\} = 0 \ a.e.$
- (A8) There exists a sequence of constants $v_n \ge 0$, $n \ge 1$ such that $\sum_{i=1}^{\infty} v_i < \infty$ and $\max_{t \in \mathcal{T}} \operatorname{var}[\boldsymbol{\epsilon}_n(t) \mid \mathcal{F}_{n-1}] \le v_n$ for all $n \ge 1$.

Assumptions (A5) and (A6) are strict generalizations of (A1) and (A2) even for p = 1, since the contraction implied by $E[\mathbf{X}_n | \mathcal{F}_{n-1}]$ is no longer required to be of the same sign as \mathbf{X}_n . Assumptions (A7)-(A8) are directly comparable to (A3)-(A4), respectively.

The strategy will be to apply Theorem 2.1 directly to $\|\mathbf{X}_n\|$, $n \ge 0$.

Theorem 4.1 Suppose $\mathbf{X} = {\mathbf{X}_n ; n \ge 0}$ is an L_2 process in \mathbb{R}^p , $p \ge 1$ which is adapted to filtration $\tilde{\mathcal{F}}$.

- (i) If (A5), (A7), (A8) hold then $\lim_{n\to\infty} ||\mathbf{X}_n||$ exists and is finite *a.e.*
- (ii) If (A6), (A7), (A8) hold then $\lim_{n\to\infty} \|\mathbf{X}_n\| = 0$ a.e.

4.1 Multivariate Stochastic Approximation

In this section we extend the stochastic approximation algorithm to the multivariate case. In particular, we consider the problem of determining the solution to $\mathbf{x}^* \in E_g \subset \mathbb{R}^p$ to $g(\mathbf{x}) = \mathbf{0}$ where $g : E_g \to \mathbb{R}^p$ is only observable with noise. The extension to $\mathbf{x}^* \in \mathbb{R}^p$ has been widely considered in the literature (Ljung, 1978; Spall, 1992; Lai, 2003; Kushner and Yin, 2003). Regularity conditions usually impose differentiability assumptions on $g(\mathbf{x})$, although in Section 2.2 and Section 3.1 it can be seen that such conditions play little role for the univariate case. We show in this section how Theorem 4.1 can be used to extend this nonsmooth case to \mathbb{R}^p (we do not consider here the nonuniform contractive model, although presumably it would also apply to this application).

We keep more or less intact conditions (B1)-(B7), with (B3) becoming a type of positive definiteness assumption on $q(\mathbf{x})$. Assuming $\mathbf{x}^* = \mathbf{0}$, the algorithm retains form

$$\mathbf{X}_n = \mathbf{X}_{n-1} - \alpha_n \mathbf{U}_n, \ n \ge 1, \tag{4.1}$$

here interpreted as a multivariate process, where $\mathbf{X}_n, \mathbf{U}_n \in \mathbb{R}^p$, and α_i is a positive constant. Again, \mathbf{X}_n , $n \geq 0$, is adapted to some filtration \mathcal{F} . We replace (B1)-(B7) with the following assumptions (some are identical to the corresponding assumption in (B1)-(B7), but are included for convenience).

- (B1a) The mapping $g: E_g \to \mathbb{R}^p$, $E_g \subset \mathbb{R}^p$, possesses root $\mathbf{0} \in E_g$.
- (B2a) $P(\mathbf{X}_n \in E_q) = 1$ for all $n \ge 1$.
- (B3a) There exists $0 < m \leq M < \infty$ such that g satisfies

$$m \leq (g(\mathbf{x}) \circ \mathbf{x}) / \|\mathbf{x}\|^2$$
 and $\|g(\mathbf{x})\| / \|\mathbf{x}\| \leq M$

for all $\mathbf{x} \in E_g - \{\mathbf{0}\}$, where $\mathbf{x} \circ \mathbf{x}'$ is the inner product on \mathbb{R}^p . (B4a) $\alpha_n \ge 0$, $\lim_n \alpha_n = 0$. (B5a) $\sum_{i=1}^{\infty} \alpha_i^2 \operatorname{var}[\mathbf{U}_i(t) \mid \mathcal{F}_{i-1}] < \infty$ a.e. for all $t \in \mathcal{T}$. (B6a) $\sum_{i=1}^{\infty} \alpha_i = \infty$.

- (B7a) $\overline{E}[\mathbf{U}_n \mid \mathcal{F}_{n-1}] = g(\mathbf{X}_{n-1}) \ a.e. \text{ for } n \ge 1.$

Under these conditions the multivariate stochastic approximation algorithm remains strongly convergent.

Theorem 4.2 If (B1a)-(B7a) hold then the process \mathbf{X}_n , $n \geq 1$, defined by Equation (4.1) converges to **0** *a.e.*

Least Squares Regression and Control $\mathbf{5}$

We consider the example of multivariate linear regression, possibly controlled. We are given probability measure space (Ω, \mathcal{F}, P) on which \mathcal{F} is a filtration. Suppose $\{\mathbf{x}_i = (x_{i1}, \ldots, x_{ip}); i \geq 1\}$ is a sequence of random $p \times 1$ vectors, where \mathbf{x}_i is \mathcal{F}_{i-1} -measurable. Let \mathbf{X}_n be the $n \times p$ matrix with *i*th row equal to \mathbf{x}_i^T . Suppose u_1, u_2, \ldots is a sequence of zero mean random variables. Define $n \times 1$ vector $\boldsymbol{u}_n = [u_1 \dots u_n]^T$. We assume \boldsymbol{u}_n is \mathcal{F}_n -measurable, and defines a sequence of martingale differences, that is, $E[u_n \mid \mathcal{F}_{n-1}] = 0, n \geq 1$. Let $\boldsymbol{\beta} = [\beta_1 \dots \beta_p]^T$ be a fixed $p \times 1$ vector. Define $d_{n,t}^2 = \sum_{i=1}^n x_{it}^2, d_{0,t}^2 = 0$ and $\mathbf{A}_n = \mathbf{X}_n^T \mathbf{X}_n$. Then consider the nested sequence of linear models

$$\mathbf{Y}_n = \mathbf{X}_n \boldsymbol{\beta} + \boldsymbol{u}_n, \ n \ge 1.$$
(5.1)

If \mathbf{X}_n contains p linearly independent rows, $\mathbf{X}_{n'}$ does as well for any n' > n. It will therefore be reasonable to assume that wp1 there exists finite n_0 such that \mathbf{A}_n is nonsingular for $n \ge n_0$.

The ordinary least squares estimate of $\boldsymbol{\beta}$ based on the *n*th model is well known to be $\mathbf{b}_n = \mathbf{A}_n^{-1} \mathbf{X}_n^T \mathbf{Y}_n$. We then center the estimates by setting $\bar{\mathbf{b}}_n =$ $\mathbf{b}_n - \boldsymbol{\beta} = \mathbf{A}_n^{-1} \mathbf{X}_n^T \boldsymbol{u}_n$. The consistency problem involves finding conditions on the sequences \mathbf{x}_n and u_n under which $\bar{\mathbf{b}}_n$ converges strongly to zero.

In Drygas (1976) and Anderson and Taylor (1976), conditions are given for deterministic \mathbf{X}_n under which $\mathbf{A}_n^{-1} \to 0$ is necessary and sufficient for weak convergence of \mathbf{b}_n to zero. Here it is assumed that the maximum eigenvalue of the covariance matrix of \boldsymbol{u}_n remains bounded as $n \to \infty$. In Lai *et al.* (1979) it is shown that $\mathbf{A}_n^{-1} \to 0$ is sufficient for strong convergence, assuming that the series $\sum_{i=1}^{\infty} c_i u_i$ converges *a.e.* whenever $\sum_{i=1}^{\infty} c_i^2 < \infty$.

Thus, under quite general conditions, for the deterministic case weak convergence implies strong convergence. Our purpose in this section is to extend the same general rule to the control model. A commonly used approach is to, in essence, turn \mathbf{A}_n into a scalar by provisionally replacing it with a diagonal matrix \mathbf{G}_n , the assumption then being that $\mathbf{A}_n^{-1}\mathbf{G}_n$ is uniformly bounded over n. In Anderson and Taylor (1979), this is accomplished by assuming that the eigenvalue ratio $\lambda_{max}(\mathbf{A}_n)/\lambda_{min}(\mathbf{A}_n)$ is uniformly bounded over n, hence the sequence of matrices \mathbf{A}_n^{-1} trace (\mathbf{A}_n) is similarly bounded. A more general condition is used in Christopeit and Helmes (1980), by assuming $\mathbf{A}_n^{-1}\mathbf{G}_n$ is uniformly bounded over n, where $\mathbf{G}_n = \text{diag}(g(d_{n,1}^2), \dots, g(d_{n,p}^2))$ for some function g(x). Here we will use this model. Under regularity conditions to be given below, it is verified in Christopeit and Helmes (1980) that $\lim_{n \to \infty} \min_{t} d_{n,t}^2 = \infty$ is sufficient for strong consistency. We will also add to this result by allowing $\lim_n d_{n,t}^2 < \infty$ for $t \leq q$ and $\lim_{n \to \infty} d_{n,t}^2 = \infty$ for t > q. Possibly, q = 0, in which case we will have strong consistency. Otherwise we have intermediate cases. We show that \mathbf{b}_n always possesses a finite limit for any q. Indeed, the component $\mathbf{b}_n(t)$ is strongly consistent for β_t when t > q. On the other hand, if $q \ge 1$, and we define subvector $\bar{\mathbf{b}}_n^1 = (\bar{\mathbf{b}}_n(1), \dots, \bar{\mathbf{b}}_n(q))^T$, then $P(\lim_n \bar{\mathbf{b}}_n^1 = \mathbf{0}) < 1$. This means that $\lim_{n \to \infty} \min_{t} d_{n,t}^2 = \infty$ is also necessary for strong consistency.

Interestingly, although Equation (5.1) defines a multivariate process, the main analysis will consider univariate processes, so we make use of Theorem 2.1. Conditions (E1)-(E4) will define our model. Condition (E5) then suffices for strong consistency, but, as already discussed, we will also consider intermediate cases.

- (E1) u_1, u_2, \ldots is a sequence of L_2 martingale differences adapted to \mathcal{F} .
- (E2) $\sup_n \operatorname{var}[u_n \mid \mathcal{F}_{n-1}] \leq \sigma^2 \ a.e.$ for some $\sigma^2 < \infty$.
- (E3) There exists finite n_0 such that \mathbf{A}_{n_0} is nonsingular.
- (E4) $\mathbf{A}_n^{-1} \mathbf{G}_n$ is uniformly bounded over n, where g(x) is a nondecreasing function satisfying $\int_c^{\infty} g^{-2}(x) dx < \infty$ for some c > 0.
- (E5) $\lim_{n \to \infty} d_{n,t}^2 = \infty$ for $t = 1, \dots, p$.

Note that (E4) implies $\lim_{x\to\infty} g(x) = \infty$.

We may write $\bar{\mathbf{b}}_n = \mathbf{A}_n^{-1} \boldsymbol{v}_n$, where

$$\boldsymbol{v}_n = \mathbf{X}_n^T \boldsymbol{u}_n = \mathbf{X}_{n-1}^T \boldsymbol{u}_{n-1} + u_n \mathbf{x}_n = \boldsymbol{v}_{n-1} + u_n \mathbf{x}_n.$$

However, at this point we replace \mathbf{A}_n^{-1} with \mathbf{G}_n^{-1} , and define the process

$$\mathbf{z}_n = \mathbf{G}_n^{-1} \boldsymbol{v}_n, \ n \ge 1$$

If necessary, set $\mathbf{z}_0 = \mathbf{v}_0 = \mathbf{0}$. Since \mathbf{G}_n^{-1} is \mathcal{F}_{n-1} -measurable we have

$$E[\mathbf{G}_n^{-1}\boldsymbol{v}_n \mid \mathcal{F}_{n-1}] = \mathbf{G}_n^{-1}E[\boldsymbol{v}_n \mid \mathcal{F}_{n-1}] = \mathbf{G}_n^{-1}\boldsymbol{v}_{n-1}.$$

Fix t and consider the component process $\mathbf{z}_n(t) = \mathbf{v}_n(t)/g\left(d_{n,t}^2\right)$. If $\mathbf{z}_{n-1}(t) = 0$, then $\mathbf{v}_{n-1}(t) = 0$, and hence $E[\mathbf{z}_n(t) \mid \mathcal{F}_{n-1}] = \mathbf{v}_{n-1}(t)//g\left(d_{n,t}^2\right) = 0$. Therefore (A3) holds for $\mathbf{z}_n(t)$. The ratio of assumption (A1) is given by

$$\frac{E[\mathbf{z}_{n}(t) \mid \mathcal{F}_{n-1}]}{\mathbf{z}_{n-1}(t)} I\{\mathbf{z}_{n-1}(t) \neq 0\} = \frac{g\left(d_{n-1,t}^{2}\right)}{g\left(d_{n,t}^{2}\right)} I\{\mathbf{z}_{n-1}(t) \neq 0\}$$
$$= k_{n,t} I\{\mathbf{z}_{n-1}(t) \neq 0\}.$$

Suppose (E4) holds. Then $d_{n,t}^2$ is nonnegative and nondecreasing in n and g(x) is nondecreasing in x, so we may conclude that $0 \le k_{n,t} \le 1$, and therefore (A1) holds for $\mathbf{z}_n(t)$. If $d_{n,t}^2 \to_n \infty$, then $g(d_{n,t}^2) \to_n \infty$. It follows from Equation (2.4) that (A2) also holds.

Since \mathbf{G}_n^{-1} is \mathcal{F}_{n-1} -measurable, we have residual vector

$$\boldsymbol{\epsilon}_n = \mathbf{G}_n^{-1} \boldsymbol{v}_n - \mathbf{G}_n^{-1} E[\boldsymbol{v}_n \mid \mathcal{F}_{n-1}] = \mathbf{G}_n^{-1} \mathbf{x}_n u_n.$$

This gives

$$E[\boldsymbol{\epsilon}_{n}^{T}\boldsymbol{\epsilon}_{n} \mid \mathcal{F}_{n-1}] \leq \sigma^{2} \mathbf{x}_{n}^{T} \mathbf{G}_{n}^{-2} \mathbf{x}_{n} = \sigma^{2} \sum_{t=1}^{p} \frac{x_{nt}^{2}}{g^{2} \left(d_{n,t}^{2}\right)}.$$
(5.2)

By Lemma D.1 we may conclude that $\sum_{n=1}^{\infty} E[\boldsymbol{\epsilon}_n^T \boldsymbol{\epsilon}_n \mid \mathcal{F}_{n-1}] < \infty$, so that (A4) holds, following Equation (2.3). Thus, by Theorem 2.1, $\mathbf{z}_n(t)$ converges to a finite limit, and if $g(d_{n,t}^2) \to_n \infty$, $\mathbf{z}_n(t)$ converges to zero *a.e.*

The convergence properties of \mathbf{b}_n follow directly, by writing

$$\bar{\mathbf{b}}_n = \mathbf{A}_n^{-1} \boldsymbol{v}_n = \mathbf{A}_n^{-1} \mathbf{G}_n \mathbf{G}_n^{-1} \boldsymbol{v}_n = \mathbf{A}_n^{-1} \mathbf{G}_n \mathbf{z}_n.$$
(5.3)

Then, by (E4), $\mathbf{A}_n^{-1}\mathbf{G}_n$ is bounded uniformly as $n \to \infty$, so we have just proven that (E5) is sufficient for strong consistency, as reported in Christopeit and Helmes (1980).

If we then more precisely characterize properties of \mathbf{A}_n^{-1} implied by (E4) we can deduce convergence properties for intermediate cases, and show that (E5) is also necessary for strong consistency.

Theorem 5.1 Suppose the regression model of Equation (5.1) satisfies (E1)-(E4). Then the following statements hold:

- (i) The least-squares estimator \mathbf{b}_n possesses a finite limit *a.e.*
- (ii) If $\lim_{n \to \infty} d_{n,t}^2 = \infty$, then $\mathbf{b}_n(t)$ is a strongly consistent estimator of β_t .

(iii) \mathbf{b}_n is a strongly consistent estimator of $\boldsymbol{\beta}$ if and only if (E5) holds.

References

- Almudevar, A. (2008). Approximate fixed point iteration with an application to infinite horizon markov decision processes. SIAM Journal on Control and Optimization, 47(5), 2303–2347.
- Almudevar, A. (2014). Approximate Iterative Algorithms. CRC Press.
- Anderson, T. and Taylor, J. B. (1979). Strong consistency of least squares estimates in dynamic models. *The Annals of Statistics*, 7(3), 484–489.

- Anderson, T. W. and Taylor, J. B. (1976). Strong consistency of least squares estimates in normal linear regression. *The Annals of Statistics*, 4(4), 788 – 790.
- Benveniste, A., Métivier, M., and Priouret, P. (2012). Adaptive Algorithms and Stochastic Approximations, volume 22. Springer Science & Business Media.
- Blum, J. R. (1954). Approximation methods which converge with probability one. The Annals of Mathematical Statistics, **25**(2), 382 386.
- Christopeit, N. and Helmes, K. (1980). Strong consistency of least squares estimators in linear regression models. *The Annals of Statistics*, **8**(4), 778 788.
- Drygas, H. (1976). Weak and strong consistency of the least squares estimators in regression models. Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete, **34**(2), 119–127.
- Durrett, R. (2019). *Probability: Theory and Examples*. Cambridge University Press.
- Horn, R. A. and Johnson, C. R. (2012). *Matrix Analysis*. Cambridge University Press.
- Kushner, H. and Yin, G. (2003). Stochastic Approximation and Recursive Algorithms, volume 35. Springer-Verlag NY.
- Lai, T. L. (2003). Stochastic approximation. The Annals of Statistics, 31(2), 391–406.
- Lai, T. L., Robbins, H., and Wei, C. Z. (1979). Strong consistency of least squares estimates in multiple regression II. Journal of Multivariate Analysis, 9(3), 343–361.
- Ljung, L. (1978). Strong convergence of a stochastic approximation algorithm. *The Annals of Statistics*, **6**(3), 680–696.
- Pham, Q.-C., Tabareau, N., and Slotine, J.-J. (2009). A contraction theory approach to stochastic incremental stability. *IEEE Transactions on Automatic Control*, **54**(4), 816–820.
- Robbins, H. and Monro, S. (1951). A stochastic approximation method. The Annals of Mathematical Statistics, **22**(3), 400 407.
- Robbins, H. and Siegmund, D. (1971). A convergence theorem for non negative almost supermartingales and some applications. In J. S. Rustagi, editor, *Optimizing Methods in Statistics*, pages 233– 257. Elsevier.
- Spall, J. C. (1992). Multivariate stochastic approximation using a simultaneous perturbation gradient approximation. *IEEE Trans*actions on Automatic Control, **37**(3), 332–341.

- Taylor, J. B. (1974). Asymptotic properties of multiperiod control rules in the linear regression model. *International Economic Re*view, 15(2), 472–484.
- Zhou, Y. and Hooker, G. (2018). Boulevard: Regularized stochastic gradient boosted trees and their limiting distribution. Preprint https://arxiv.org/abs/1806.09762.

Appendix A Proof of Theorem 2.1, with Technical Lemmas

Lemma A.1 If (A1) holds then for any $i \ge 1$ $X_i I\{D_{i-1}^-\} \le \epsilon_i I\{D_{i-1}^-\}$ a.e.

Proof. We have $(X_i - \epsilon_i) I\{D_{i-1}^-\} = E[X_i | \mathcal{F}_{i-1}] I\{D_{i-1}^-\} \leq 0$ a.e. by (A1), which completes the proof.

The following lemma allows us to bound the process **X** in terms of the partial sums of the form $M_{s,t}$. Define the kernel $\lambda_t^k = \prod_{i=t-k+1}^t (1 + \alpha_i), t \ge 1, k \ge 1$, and $\lambda_t^0 = 1$. Under (A1), $\lambda^{max} = \sup_{t,k} \lambda_t^k < \infty$. Then set $M_{s,t}^{abs} = \sum_{i=s}^t |\epsilon_i|$.

The lemma also relies on the following device. Let $\overline{\mathbb{I}}$ be the extended integers. For any $t \in \overline{\mathbb{I}}$ define $t^- = \sup\{i \in \overline{\mathbb{I}} : i < t\}$.

Lemma A.2 If (A1) holds then for $j \ge 1$

$$W_{j} \leq \lambda^{max} \left(\sup_{T_{j} \leq i < T_{j+1}} M^{abs}_{T_{j}, T_{j+1}-1} + |U_{T_{j}}| \right) I\{T_{j} < \infty\} \ a.e.$$

Proof. We have by Lemma A.1 and the definition of ϵ_i , for $i \geq 1$

$$X_{i}I\{D_{i-1}^{-}\} \leq \epsilon_{i}I\{D_{i-1}^{-}\}$$

$$X_{i}I\{D_{i-1}^{0}\} = (E[X_{i} | \mathcal{F}_{i-1}] + \epsilon_{i})I\{D_{i-1}^{0}\}$$

which when combined give

$$X_i I\{(D_{i-1}^+)^c\} \le U_i + \epsilon_i I\{(D_{i-1}^+)^c\} = (U_i + \epsilon_i) I\{(D_{i-1}^+)^c\},$$
(A.1)

noting that $I\{D_{i-1}^0\} = I\{D_{i-1}^0\}I\{(D_{i-1}^+)^c\}.$

Fix integers $1 \le s < t$ where s is finite but, possibly, $t = \infty$. For now, assume $s < t^-$, and select i, s < i < t. Then

$$X_{i}I\{D_{s,t^{-}}^{+}\} = \left([X_{i} - E[X_{i} | \mathcal{F}_{i-1}]] + E[X_{i} | \mathcal{F}_{i-1}] \right) I\{D_{s,t^{-}}^{+}\}$$

$$\leq \left(\epsilon_{i} + (1 + \alpha_{i})X_{i-1}\right) I\{D_{s,t^{-}}^{+}\}, \qquad (A.2)$$

after applying (A1). Applying Equation (A.2) iteratively gives

$$X_{i}I\{D_{s,t^{-}}^{+}\} \leq \left(\sum_{k=s+1}^{i} \lambda_{i}^{i-k} \epsilon_{k} + \lambda_{i}^{i-s} X_{s}\right) I\{D_{s,t^{-}}^{+}\}.$$
 (A.3)

Note that Equation (A.3) holds also for i = s, using the standard summation convention, so we now permit $s = t^-$. With Lemma A.1 this leads to

$$X_{i}I\{D_{s,t^{-}}^{+}\}I\{(D_{s-1}^{+})^{c}\} \leq \left(\lambda_{i}^{i-s}(U_{s}+\epsilon_{s})+\sum_{k=s+1}^{i}\lambda_{i}^{i-k}\epsilon_{k}\right)I\{D_{s,t^{-}}^{+}\}I\{(D_{s-1}^{+})^{c}\}$$
$$= \left(\lambda_{i}^{i-s}U_{s}+\sum_{k=s}^{i}\lambda_{i}^{i-k}\epsilon_{k}\right)I\{D_{s,t^{-}}^{+}\}I\{(D_{s-1}^{+})^{c}\} \quad (A.4)$$

for $s \leq i < t$. It follows from (A.4) that

$$\sup_{s \le i < t} X_i I\{D_{s,t^-}^+\} I\{(D_{s-1}^+)^c\} \le \lambda^{max} (U_s + M_{s,t^-}^{abs}) I\{D_{s,t^-}^+\} I\{(D_{s-1}^+)^c\}.$$
(A.5)

Then on $\{T_j < \infty\}$ we have $\{T_j = s, T_{j+1} = t\} \cap D_s^+ \subset D_{s,t^-}^+ \cap (D_{s-1}^+)^c$. This gives

$$\sup_{\substack{T_j \le i < T_{j+1} \\ \le \lambda^{max} (U_{T_j} + M_{T_j, T_{j+1}^-}^{abs})} X_i I\{T_j = s, T_{j+1} = t\} I\{D_{T_j}^+\}.$$
(A.6)

Summing Equation (A.6) over s and t yields

$$W_{j}I\{D_{T_{j}}^{+}\} = \sup_{T_{j} \leq i < T_{j+1}} X_{i}I\{D_{T_{j}}^{+}\}I\{T_{j} < \infty\}$$

$$\leq \lambda^{max}(|U_{T_{j}}| + M_{T_{j},T_{j+1}}^{abs})I\{D_{T_{j}}^{+}\}I\{T_{j} < \infty\}.$$

Then note that assumption (A1) also applies to $\{-X_i\}$, hence we may similarly conclude

$$W_{j}I\{D_{T_{j}}^{-}\} \leq \lambda^{max}(|U_{T_{j}}| + M_{T_{j},T_{j+1}}^{abs})I\{D_{T_{j}}^{-}\}I\{T_{j} < \infty\}.$$

Also, by construction we have $X_i I\{D_{T_i}^0\} = 0$ for $T_j \leq i < T_{j+1}$ so that we may conclude

$$W_j \le \lambda^{max} (|U_{T_j}| + M^{abs}_{T_j, T_{j+1}^-}) I\{D^+_{T_j}\} I\{T_j < \infty\}.$$

completing the proof.

We are now in a position to prove the main results. We first consider the nonexpansive model. Given (A1), (A3) and (A4) we may claim that **X** possesses a finite limit. It will be useful, however, to consider the case for which (A3) need not hold.

Appendix A.1 Proof of Theorem 2.1

Proof. We first consider Statement (i), that is, the case $N_T < \infty$. Fix $s \ge 1$, and define stopping time

$$M = \inf\{i \ge s : X_i \le 0\}.$$

Then construct the sequence

$$X'_{s+n} = (X_{s+n} - \epsilon_{s+n})I\{D^+_{s+n-1}\}, \ n \ge 0,$$

then

$$X_{s+n}^* = X'_{(s+n)\wedge M}, \ n \ge 0.$$

Following Lemma A.1 we have $X_{s+n}^* \ge 0$ for all $n \ge 0$. Then $X_{s+n}^* = 0$ for all $s+n \ge M$ and $X_{s+n}^* = X_{s+n}$ when for all s+n < N, noting that we may have $M = \infty$. This leads to

$$E[X_{s+n}^* \mid \mathcal{F}_{s+n-1}] = E[X_{s+n}^*I\{M \ge s+n\} + X_{s+n}^*I\{M < s+n\} \mid \mathcal{F}_{s+n-1}]$$

$$= E[X_{s+n}' \mid \mathcal{F}_{s+n-1}]I\{M \ge s+n\}I\{D_{s+n-1}^+\}$$

$$+ X_M'I\{M < s+n\}I\{D_{s+n-1}^+\}$$

$$= (1 + \alpha_{s+n})(X_{s+n-1}' + \epsilon_{s+n-1})I\{M \ge s+n\}$$

$$+ X_{s+n-1}'I\{M < s+n\}$$

$$\le (1 + \alpha_{s+n})X_{s+n-1}^* + (1 + \alpha_{s+n})\epsilon_{s+n-1}^+.$$

By assumption $\sum_{n\geq 0} \alpha_{s+n} < \infty$ and $\sum_{n\geq 0} (1 + \alpha_{s+n}) < \infty$ a.e., so that the almost martingale conditions of Theorem 1 of Robbins and Siegmund (1971) hold. We may then conclude that X_{s+n}^* possesses a finite limit *a.e.*. This in turn implies that $X_n I\{N_t < \infty\}$ possesses a finite limit.

We next consider Statement (ii). Since (A2) implies (A1), the conclusions of Statement (i) hold. By assumption (A2), for $i \ge s$,

$$(A_{s+1,i} - A_{s+1,i+1})I\{D_{s,\infty}^+\} = (X_i - E[X_{i+1} \mid \mathcal{F}_i])I\{D_{s,\infty}^+\}$$

$$\geq (1 - k_i)X_iI\{D_{s,\infty}^+\} \ a.e.$$
(A.7)

If $X_i I\{D_{s,\infty}^+\}$ does not converge to 0 then there exists c > 0 such that for large enough s' > s, $X_i I\{D_{s,\infty}^+\} \ge c$ a.e. on $D_{s,\infty}^+$ when $i \ge s'$. Applying Equation (A.7) gives

$$\lim_{i \to \infty} A_{s+1,i} I\{D_{s,\infty}^+\} = \lim_{i \to \infty} \left(A_{s+1,s'} - \sum_{i'=s'}^i (A_{s+1,i'} - A_{s+1,i'+1}) \right) I\{D_{s,\infty}^+\}$$

$$\leq \lim_{i \to \infty} \left(A_{s+1,s'} - \sum_{i'=s'}^i (1 - k_{i'}) X_{i'} \right) I\{D_{s,\infty}^+\}$$

$$\leq \lim_{i \to \infty} \left(A_{s,s'} - c \sum_{i'=s'}^i (1 - k_{i'}) \right) I\{D_{s,\infty}^+\}$$

$$= -\infty I\{D_{s,\infty}^+\} \ a.e.$$

But following the proof of Statement (i) $A_{s,i}$ must possess a finite limit, hence $X_i I\{D_{s,\infty}^+\}$ must converge to 0 *a.e.* on $\{N_T < \infty\} \cap \{T_{N_T} = s\} \cap D_s^+ \subset D_{s,\infty}^+$. The remaining proof is essentially the same as for Statement (i).

We next consider Statement (iii), the case $N_T = \infty$. Since $M_{1,i}$ has a limit *a.e.*, by the Cauchy criterion $M_{m,n} \to 0$ as $m, n \to \infty$. which implies wp1

$$\limsup_{j \to \infty} \left(\sup_{T_j \le i < T_{j+1}} |M_{T_j,i}| + |U_{T_j}| \right) I\{N_T = \infty\} = \left(\limsup_{j \to \infty} |U_j| \right) I\{N_T = \infty\}$$

since $\{N_T = \infty\} \subset \{T_j < \infty\}$ for all j, and $T_j \to_j \infty$. Statement (iii) follows by applying Lemma A.2. Given (A3), Statement (iv) follows from a direct application of Statement (ii).

Appendix A.2 Proof of Theorem 2.3

Proof. Assumption (B2) ensures that the process (2.6) is always defined. By (B4), $\alpha_i \rightarrow_i 0$, so we may assume without loss of generality that $M\alpha_i \leq 1$ for all *i*. Subtracting x^* from both sides of (2.6) gives the equivalent process

$$X_i - x^* = X_{i-1} - x^* - \alpha_i U_i, \ i \ge 1,$$
(A.8)

with initial value $X_0 - x^*$. We then apply Theorem 2.1 to the process (A.8). Accordingly, by (B7), we have

$$E[X_i - x^* \mid \mathcal{F}_{i-1}] = X_{i-1} - x^* - \alpha_i g(X_{i-1}), \ i \ge 1.$$

Since $g(x^*) = 0$ we have $E[X_i - x^* | \mathcal{F}_{i-1}]I\{X_{i-1} - x^* = 0\} = 0$ so that (A3) holds. Then by (B3)

$$\frac{E[X_i - x^* \mid \mathcal{F}_{i-1}]}{X_{i-1} - x^*} I\{X_{i-1} - x^* \neq 0\} = \left(1 - \alpha_i \frac{g(X_{i-1})}{X_{i-1} - x^*}\right) I\{X_{i-1} - x^* \neq 0\},$$
(A.9)

which gives

$$(1 - M\alpha_i)I\{X_{i-1} - x^* \neq 0\} \le \frac{E[X_i - x^* \mid \mathcal{F}_{i-1}]}{X_{i-1} - x^*}I\{X_{i-1} - x^* \neq 0\}$$

$$\le (1 - m\alpha_i)I\{X_{i-1} - x^* \neq 0\}.$$
(A.10)

Then note that $1 - M\alpha_i \ge 0$ and $1 - m\alpha_i \le 1$. Also, (B6) implies $\sum_i m\alpha_i = \infty$ so that (A2) holds. Then (B5) is equivalent to (A4), so that $X_i - x^* \to_i 0$ a.e. by Theorem 2.1.

Appendix B Proofs of Theorems 3.1 and 3.2

Appendix B.1 Proof of Theorem 3.1

Proof. Let $\tilde{\mathcal{F}}$ be the filtration for the process X_i . For any $\tau > 0$ there exists a.e. finite N_0 such that $|\epsilon_i| < \tau$ for all $i \ge N_0$. Select $\tau < \delta$, then consider the offset process $X_{N_0+i}^{\delta+\tau}$, $i \ge 0$. Suppose $X_{N_0+i}^{\delta+\tau} > 0$. Since $|\epsilon_i| < \tau$ and $X_{N_0+i} = E[X_{N_0+i} \mid \mathcal{F}_{N_0+i-1}] + \epsilon_i$ we must have $X_{N_0+i} > \delta$. By (D1) we have $0 \le E[X_{N_0+i+1} \mid \mathcal{F}_{N_0+i}] \le X_{N_0+i} = X_{N_0+i}^{\delta+\tau}$. We may then write

$$E[X_{N_0+i+1}^{\delta+\tau} \mid \mathcal{F}_i] = E[X_{N_0+i+1}I\{|E[X_{N_0+i+1} \mid \mathcal{F}_{N_0+i}]| \ge \delta + \tau\} \mid \mathcal{F}_{N_0+i}]$$

= $E[X_{N_0+i+1} \mid \mathcal{F}_{N_0+i}]I\{|E[X_{N_0+i+1} \mid \mathcal{F}_{N_0+i}]| \ge \delta + \tau\}$
 $\in [0, E[X_{N_0+i+1} \mid \mathcal{F}_{N_0+i}]].$

A similar argument holds for the case $X_{N_0+i}^{\delta+\tau} < 0$, hence it follows that (A1) holds for $X_{N_0+i}^{\delta+\tau}$, $i \ge 1$.

The proof of Statement (ii) is similar to that of Statement (i).

To prove Statement (iii), suppose $X_{N_0+i-1}^{\delta+\tau} = 0$. First consider the case $X_{N_0+i-1} = 0$. We may write

$$U_{N_0+i}^{\delta+\tau} I\{X_{N_0+i-1} = 0\}$$

= $E[X_{N_0+i} \mid \mathcal{F}_{N_0+i-1}]I\{X_{N_0+i-1} = 0\}I\{|E[X_{N_0+i} \mid \mathcal{F}_{N_0+i-1}]| \ge \delta + \tau\}$
 $\times I\{X_{N_0+i-1}^{\delta+\tau} = 0\}$
= $U_{N_0+i}W_{N_0+i},$ (B.1)

where $W_{N_0+i} \in \{0,1\}$. Otherwise, suppose $X_{N_0+i-1} \neq 0$ but $|E[X_{N_0+i-1} | \mathcal{F}_{N_0+i-1}]| \in [0, \delta + \tau)$. We must then have $|X_{N_0+i-1}| \in (0, \delta + 2\tau)$. Suppose $|X_{N_0+i-1}| > \delta$. Then by (A1) $|E[X_{N_0+i} | \mathcal{F}_{i-1}]| \leq |X_{N_0+i-1}| < \delta + 2\tau$. Hence, for this case

$$|U_{N_0+i}^{\delta+\tau}|I\{|X_{N_0+i-1}| > \delta\} \le (\delta+2\tau).$$
(B.2)

Finally, suppose $|X_{N_0+i-1}| \leq \delta$. By (D3) $|E[X_{N_0+i} | \mathcal{F}_{N_0+i-1}]| < \kappa$, hence

$$U_{N_0+i}^{\delta+\tau}|I\{|X_{N_0+i-1}|\in(0,\delta]\}<\kappa.$$
(B.3)

Then Equation (3.4) follows from Equations (B.1), (B.2) and (B.3).

We finally consider Statement (iv). Assume (A4) holds for X_i . Then for $X_i^{\delta+\tau}$ the residuals are given by

$$\begin{aligned} \epsilon_i^{\delta+\tau} &= X_i^{\delta+\tau} - E[X_i^{\delta+\tau} \mid \mathcal{F}_{N_0+i-1}] \\ &= (X_i - E[X_i])I\{|E[X_{N_0+i} \mid \mathcal{F}_{N_0+i-1}]| \ge \delta + \tau\} \\ &= \epsilon_i I\{|E[X_{N_0+i} \mid \mathcal{F}_{N_0+i-1}]| \ge \delta + \tau\}, \end{aligned}$$

Thus, $|\epsilon_i^{\delta+\tau}| \leq |\epsilon_i|$, hence (A4) extends to $\epsilon_i^{\delta+\tau}$.

Appendix B.2 Proof of Theorem 3.2

Proof. For the first (nonexpansive) case, we may conclude by Theorem 3.1 that there exists finite N_0 such that $\mathbf{X}_{N_0}^{\delta+\tau}$ satisfies (A1) and (A4) for all small enough $\delta, \tau > 0$. We then note that

$$|X_{N_0+i}| \le \max\{|X_{N_0+i}^{\delta+\tau}|, \delta+2\tau\}.$$
(B.4)

Applying (D4) and (A3) we conclude by Theorem 3.1 (iii) that $\limsup_i |U_{N_0+i}^{\delta+\tau}| \leq \delta + 2\tau + \kappa_{\delta}$. That **X** possesses a finite limit follows by applying Theorem 2.1 to $\mathbf{X}_{N_0}^{\delta+\tau}$, allowing δ, τ to approach zero, and noting Equation (B.4).

For the contractive case, note that under the given assumptions $|\mathbf{X}_{N_0+i}^{\delta+\tau}|$ may be bounded by some finite constant δ_2 . If (D2) holds for all $0 < \delta_1 < \delta_2 < \infty$ pairs then it follows from Theorem 3.1 (ii) that (A2) holds for $\mathbf{X}_{N_0}^{\delta+\tau}$, which therefore satisfies the hypothesis of Theorem 2.1 (ii). The remaining argument follows that of the nonexpansive case.

Appendix C Proofs of Theorems 4.1 and 4.2

Appendix C.1 Proof of Theorem 4.1

Proof. Conditions (A5), (A6) and (A7) directly imply conditions (A1), (A2) and (A3) for $\mathbf{X}^* = \{ \| \mathbf{X}_i \|; i \ge 0 \}$. It remains to verify that (A8) implies (A4) for \mathbf{X}^* . We have residual process

$$\epsilon_i^* = \|\mathbf{X}_i\| - E[\|\mathbf{X}_i\| \mid \mathcal{F}_{i-1}], \ i \ge 1.$$

Then

$$\operatorname{var}[\boldsymbol{\epsilon}_{i}^{*} \mid \mathcal{F}_{i-1}] = E[\|\mathbf{X}_{i}\|^{2} \mid \mathcal{F}_{i-1}] - E[\|\mathbf{X}_{i}\| \mid \mathcal{F}_{i-1}]^{2}$$
$$= E\left[\sum_{t} \mathbf{X}_{i}(t)^{2} \mid \mathcal{F}_{i-1}\right] - E[\|\mathbf{X}_{i}\| \mid \mathcal{F}_{i-1}]^{2}. \quad (C.1)$$

However, Euclidean norm $\|\cdot\|$ is convex, so by Jensen's inequality we have $E[\|\mathbf{X}_i\| \mid \mathcal{F}_{i-1}]^2 \geq \sum_t E[\mathbf{X}_i(t) \mid \mathcal{F}_{i-1}]^2$, which, when combined with Equation (C.1), gives

$$\operatorname{var}[\boldsymbol{\epsilon}_{i}^{*} \mid \mathcal{F}_{i-1}] \leq \sum_{t} \operatorname{var}[\mathbf{X}_{i}(t) \mid \mathcal{F}_{i-1}] \leq \sum_{t} \operatorname{var}[\boldsymbol{\epsilon}_{i}(t) \mid \mathcal{F}_{i-1}], \quad (C.2)$$

which, with (A8), implies that \mathbf{X}^* satisfies (A4). The proof is completed by a direct application of Theorem 2.1.

Appendix C.2 Proof of Theorem 4.2

Proof. The main task is to verify condition (A6) for all large enough *i*. By construction we have $E[\mathbf{X}_i | \mathcal{F}_{i-1}] = \mathbf{X}_{i-1} - \alpha_i g(\mathbf{X}_{i-1})$, so that, by (B3a),

$$\begin{aligned} \|E[\mathbf{X}_{i} \mid \mathcal{F}_{i-1}]\|^{2} &= \|\mathbf{X}_{i-1}\|^{2} - 2\alpha_{i}g(\mathbf{X}_{i-1}) \circ \mathbf{X}_{i-1} + \alpha_{i}^{2}\|g(\mathbf{X}_{i-1})\|^{2} \\ &\leq \|\mathbf{X}_{i-1}\|^{2} \left[1 - 2\alpha_{i}m + \alpha_{i}^{2}M^{2}\right]. \end{aligned}$$

It follows that

$$\frac{\|E[\mathbf{X}_i \mid \mathcal{F}_{i-1}]\|}{\|\mathbf{X}_{i-1}\|} I\{\|\mathbf{X}_{i-1}\| \neq 0\} \le \left[1 - 2\alpha_i m + \alpha_i^2 M^2\right]^{1/2} = k_i.$$

Since $\alpha_i \to 0$, we must have $\left[1 - 2\alpha_i m + \alpha_i^2 M^2\right]^{1/2} < 1$ for all large enough i. Suppose we have function $h(u) = 1 - [1 - u]^{1/2}$. It is easily verified that h'(0) = 1/2, and that h(u) is convex for u < 1. Furthermore, there exists finite K > 0 and finite index i_0 such that for all $i \ge i_0$ we have $2\alpha_i m - \alpha_i^2 M^2 \ge K\alpha_i$. We then have $\sum_{i\ge i_0} (1-k_i) \ge (K/2) \sum_{i\ge i_0} \alpha_i = \infty$ so that (A6) holds. That (A7) and (A8) hold follow from arguments similar to those of Theorem 4.2. The theorem is then proved following an application of Theorem 4.1 (ii).

Appendix D Proof of Theorem 5.1 with Technical Lemmas

The following lemma is a generalization of Lemma 1 of Taylor (1974), which states that for any sequence of real numbers x_1, x_2, \ldots with $x_1 \neq 0$ we must have $S_N = \sum_{n=1}^N x_n^2 / (\sum_{i=1}^n x_i^2)^2 \leq 2/x_1^2$ for all $N \geq 1$. A recursive argument is used in Taylor (1974), but we may also interpret S_N as an approximation of the integral $\int_c^\infty x^{-2} dx$. Doing so will allow us to expand the class of models to be studied.

Lemma D.1 Let a_1, a_2, \ldots be a sequence of nonnegative real numbers, not uniformly equal to zero. Assume $a_1 > 0$ (otherwise delete a_1 from the sequence). Let $A_n = \sum_{i=1}^n a_i$ Suppose we are given a nondecreasing function $f: (0, \infty) \to (0, \infty)$ for which $\int_c^{\infty} f^{-1}(x) dx < \infty$ for any c > 0. Then

$$S_N = \sum_{n=1}^N a_n / f(A_n) \le a_1 / A_1 + \int_{a_1}^\infty f^{-1}(x) dx$$
 (D.1)

for all $N \geq 1$.

Proof. Let $A_{max} = \lim_{n \to \infty} A_n$. Construct a step function h(x) on $x \in (0, A_{max})$ with discontinuities at A_n , $n \ge 1$. Set the left limit of h(x) at A_n to be $f^{-1}(A_n)$. If $A_{max} < \infty$ set h(x) = 0 for $x \ge A_{max}$. Then

$$S_N = \sum_{n=1}^N a_n / f(A_n) = \int_0^{A_N} h(x) dx \le \int_0^\infty h(x) dx = a_1 / f(A_1) + \int_{a_1}^\infty h(x) dx.$$

Equation (D.1) follows from the fact that $0 \le h(x) \le f^{-1}(x)$ for x > 0.

Appendix D.1 Proof of Theorem 5.1

For $\mathbf{y} = (y_1, \ldots, y_p) \in \mathbb{R}^p$ we have ℓ_{∞} norm $|\mathbf{y}|_{\infty} = \max_{t=1,\ldots,p} |y_t|$. Then let \mathbf{C} be a $p \times p$ matrix with elements c_{ij} . We will make use of the matrix norm $\|\mathbf{C}\|_{\infty} = p \max_{i,j} |c_{ij}|$. Note that $\|\mathbf{C}\|_{\infty}$ is a true matrix norm, in particular, $|\mathbf{Cy}|_{\infty} \leq \|\mathbf{C}\|_{\infty} |\mathbf{y}|_{\infty}$ Horn and Johnson (2012).

Proof. Suppose for model (5.1) conditions (E1)-(E4) hold. Then assume for some index $q, 1 \leq q < p$ we have $\lim_{n} d_{n,t}^2 < \infty$ for $t \leq q$ and $\lim_{n} d_{n,t}^2 = \infty$ for

t > q. Set $\mathbf{B}_n = \mathbf{A}_n^{-1}$, and let $b_{n,ij}$ be the elements of \mathbf{B}_n . By (E4) we have $\|\mathbf{A}_n^{-1}\mathbf{G}_n\|_{\infty} \leq \kappa$ for all *n* for some finite κ . It follows that

$$|b_{n,ij}| \leq p^{-1} \kappa / g(d_{n,j}^2)$$
 for all i, j .

Thus, for j > q we must have $\lim_{n \to j} b_{n,ij} = 0$, and, since \mathbf{B}_n is symmetric, for i > q as well. Then consider the matrix partitions

$$\mathbf{A}_n = \begin{bmatrix} \mathbf{A}_n^{11} & \mathbf{A}_n^{12} \\ \mathbf{A}_n^{21} & \mathbf{A}_n^{22} \end{bmatrix}, \ \mathbf{B}_n = \begin{bmatrix} \mathbf{B}_n^{11} & \mathbf{B}_n^{12} \\ \mathbf{B}_n^{21} & \mathbf{B}_n^{22} \end{bmatrix}$$

where \mathbf{A}_n^{11} , \mathbf{B}_n^{11} are $q \times q$ square matrices. We then have

$$\mathbf{A}_n^{11}\mathbf{B}_n^{11} + \mathbf{A}_n^{12}\mathbf{B}_n^{21} = \mathbf{I}_q, \tag{D.2}$$

where \mathbf{I}_q is the $q \times q$ identity matrix. Under our given assumptions, the finite limit $\lim_n \mathbf{A}_n^{11} = \mathbf{A}_\infty^{11}$ exists and is positive definite. Let $\mathbf{C}_n = \mathbf{A}_n^{12} \mathbf{B}_n^{21}$ and let $c_{n,rc}$ be the elements of \mathbf{C}_n . Then (noting that $b_{n,ij} = b_{n,ji}$)

$$c_{n,rc}| \leq \sum_{j=q+1}^{p} |a_{n,rj}| |b_{n,jc}|$$

= $\sum_{j=q+1}^{p} \left| \sum_{i=1}^{n} x_{i,r} x_{n,j} \right| |b_{n,jc}|$
 $\leq p^{-1} \kappa \left(\sum_{i=1}^{\infty} x_{i,r}^2 \right)^{1/2} \sum_{j=q+1}^{p} \frac{(d_{n,j}^2)^{1/2}}{g(d_{n,j}^2)}.$ (D.3)

By assumption $\sum_{i=1}^{\infty} x_{i,r}^2 < \infty$ for $r \leq q$. Furthermore, if (E4) holds we must have $g(x) = x^{1/2}g^*(x)$, where $\lim_{x\to\infty} g^*(x) = \infty$. By assumption $\lim_n d_{n,j}^2 = \infty$ for j > q, so by Equation (D.3) it follows that $\lim_n c_{n,rc} = 0$. After applying Equation (D.2) we have $\lim_n \mathbf{B}_n^{11} = (\mathbf{A}_\infty^{11})^{-1}$. We then have $\mathbf{b}_n - \boldsymbol{\beta} = \mathbf{A}_n^{-1} \boldsymbol{v}_n$. Define the partition $\boldsymbol{v}_n = [\boldsymbol{v}_n^1 \ \boldsymbol{v}_n^2]$, where

We then have $\mathbf{b}_n - \boldsymbol{\beta} = \mathbf{A}_n^{-1} \boldsymbol{v}_n$. Define the partition $\boldsymbol{v}_n = [\boldsymbol{v}_n^1 \ \boldsymbol{v}_n^2]$, where \boldsymbol{v}_n^1 contains the first q elements of \boldsymbol{v}_n . For $t \leq q$ it is easily verified that $\boldsymbol{v}_n(t)$ is a finite variance martingale, and hence possesses limit $\bar{\boldsymbol{v}}(t) = \lim_n \boldsymbol{v}_n(t)$ with $\operatorname{var}[\bar{\boldsymbol{v}}(t)] > 0$. It follows that $P(\lim_n \boldsymbol{v}_n^1 = \mathbf{0}) < 1$. Suppose $\lim_n \boldsymbol{v}_n^1 = \bar{\boldsymbol{v}}^1$. We then have the limit

$$\lim_{n} \mathbf{A}_{n}^{-1} \begin{bmatrix} \boldsymbol{v}_{n}^{1} \\ \mathbf{0} \end{bmatrix} = \lim_{n} \begin{bmatrix} \mathbf{B}_{n}^{11} \boldsymbol{v}_{n}^{1} \\ \mathbf{B}_{n}^{21} \boldsymbol{v}_{n}^{1} \end{bmatrix} = \begin{bmatrix} (\mathbf{A}_{\infty}^{11})^{-1} \bar{\boldsymbol{v}}^{1} \\ \mathbf{0} \end{bmatrix},$$

noting that the elements of \mathbf{B}_n^{21} vanish as $n \to \infty$. Since \mathbf{A}_{∞}^{11} is nonsingular, $\bar{\boldsymbol{v}}^1 \neq 0$ implies $(\mathbf{A}_{\infty}^{11})^{-1} \bar{\boldsymbol{v}}^1 \neq 0$, so that $P((\mathbf{A}_{\infty}^{11})^{-1} \bar{\boldsymbol{v}}^1 = \mathbf{0}) < 1$. Next, we evaluate

$$\mathbf{W}_n = \mathbf{A}_n^{-1} \begin{bmatrix} \mathbf{0} \\ oldsymbol{v}_n^2 \end{bmatrix} = \mathbf{A}_n^{-1} \mathbf{G}_n \mathbf{G}_n^{-1} \begin{bmatrix} \mathbf{0} \\ oldsymbol{v}_n^2 \end{bmatrix}$$

However, in Section 5 it was shown that the components of index t > q of $\mathbf{G}_n^{-1} \boldsymbol{v}_n$ converge to zero *a.e.* Since $\mathbf{A}_n^{-1} \mathbf{G}_n$ is uniformly bounded over *n*, it follows that $\lim_n \mathbf{W}_n = \mathbf{0}$ *a.e.* We then have, wp1, the finite limit

$$\lim_{n} \mathbf{b}_{n} - \boldsymbol{\beta} = \lim_{n} \mathbf{A}_{n}^{-1} \begin{bmatrix} \boldsymbol{v}_{n}^{1} \\ \mathbf{0} \end{bmatrix} + \lim_{n} \mathbf{A}_{n}^{-1} \begin{bmatrix} \mathbf{0} \\ \boldsymbol{v}_{n}^{2} \mathbf{0} \end{bmatrix} = \begin{bmatrix} (\mathbf{A}_{\infty}^{11})^{-1} \bar{\boldsymbol{v}}^{1} \\ \mathbf{0} \end{bmatrix}.$$
(D.4)

Equation (D.4) can clearly be extended to the case q = p, in which case the finite limit $\lim_n \mathbf{b}_n - \boldsymbol{\beta} = \lim_n \mathbf{A}_n^{-1} \bar{\boldsymbol{v}}$ exists *a.e.* This is true also for condition (E5) (that is, q = 0), in which case $\lim_n \mathbf{b}_n - \boldsymbol{\beta} = \mathbf{0}$ a.e. Hence, $\lim_n \mathbf{b}_n - \boldsymbol{\beta}$ always possesses a finite limit, and that limit is zero wp1 if and only if q = 0. Otherwise $\mathbf{b}_n(t)$ converges to $\beta_t wp1$ if $\lim_n d_{n,t}^2 = \infty$.