

On the compactification of bounded edge Morse flow trees

Kenneth Blakey

Abstract

Let $L \subset J^1(M)$ be a closed Legendrian submanifold of the 1-jet space of a closed Riemannian manifold (M, g) and consider the moduli space $\mathcal{L}(N)$ of Morse flow trees in L with at most N edges and vertices. We construct a compactification of $\mathcal{L}(N)$ by considering the moduli space $\overline{\mathcal{L}}(N)$ of broken flow trees in L with at most N edges and vertices, provided that L is front generic, has simple front singularities, and satisfies Ekholm's preliminary transversality condition. In particular, we show that a sequence of Morse flow trees T_n , with a uniform edge bound, always has a subsequence that converges to some broken flow tree T .

Introduction. Let M be a closed smooth n -manifold with a Riemannian metric g , let $J^0(M) = M \times \mathbb{R}$ be M 's 0-jet space, and let $J^1(M) = T^*M \times \mathbb{R}$ be M 's 1-jet space. We endow $J^1(M)$ with the standard contact structure ξ given as the kernel of $dz - \alpha$, where z is the \mathbb{R} -coordinate and α is the tautological 1-form on T^*M . An n -dimensional submanifold $L \subset J^1(M)$ is called Legendrian if it is an integral submanifold of ξ . We also endow T^*M with the standard symplectic structure given as $-d\alpha$. An n -dimensional submanifold $X \subset T^*M$ is called Lagrangian if $-d\alpha$ restricted to X vanishes. This paper concerns the moduli space of Morse flow trees (flow trees for short) determined by L , with at most N edges and vertices, and its compactification.

Flow trees were introduced in [Ekh07] in order to compute the contact homology of Legendrian submanifolds in 1-jet spaces. Specifically, Legendrian contact homology is a part of Symplectic Field Theory (see [EGH00]) and is intended to find isotopy invariants of Legendrian submanifolds. The rigorous construction of Legendrian contact homology in 1-jet spaces was done in [EES07] and involves counting rigid pseudo-holomorphic disks with punctures asymptotic to Reeb chords on the Legendrian. Finding pseudo-holomorphic disks involves solving a non-linear first order partial differential equation which makes explicit computations quite difficult. The main result of [Ekh07] reduces the computation to a finite-dimensional problem in Morse theory by building a 1-1 correspondence between (certain types of) flow trees and (certain types of) pseudo-holomorphic disks, after sufficient perturbations of the involved data. The relation of Morse theory and pseudo-holomorphic disks can be found in the Floer-theoretic case, for example [Flo89] and [FO97], and the power of flow trees can be seen in the computations of Legendrian contact homology in [Riz11], [RG19], and [EN18].

Given Ekhholm’s correspondence of flow trees and pseudo-holomorphic disks, and given the existence of compactness results for pseudo-holomorphic disks (see [BEH⁺03] and [Abb14]), we would expect the existence of compactness results for flow trees. The main result of this paper is Theorem 0.1, which shows that if we have a sequence of flow trees T_n with a uniform bound N on the number of edges and vertices then, perhaps after passing to a subsequence, T_n converges in an appropriate space to a broken flow tree T .

Let M be as before and let $L \subset J^1(M)$ be a closed Legendrian submanifold. We denote by $\Pi_{\mathbb{C}}$ the Lagrangian projection, $\Pi_{\mathbb{C}} : J^1(M) \rightarrow T^*M$. The image of L , under this projection, is an immersed Lagrangian submanifold. After small perturbations of L we may assume that the only self intersections of $\Pi_{\mathbb{C}}(L)$ are self-transverse double points, and we note these double points are in 1-1 correspondence with Reeb chords of L .

For any point p in M , the smooth sheets of L lying over p are 1-jet graphs of a finite number of locally defined functions on M . We denote by Π the base projection, $\Pi : J^1(M) \rightarrow M$. We also denote by Π_F the front projection, $\Pi_F : J^1(M) \rightarrow J^0(M)$. We will assume the restriction $\Pi : L \rightarrow M$ is an immersion outside of a codimension 1 submanifold $\Sigma \subset L$ and at points in Σ , Π has a standard fold singularity, and the corresponding singularity of Π_F is a cusp edge. In this case we say L is front generic with simple front singularities. Note that any Legendrian submanifold in $J^1(M)$ can be perturbed to be front generic, that is to say every point in Σ outside of a codimension 1 subset $\Sigma' \subset \Sigma$ is a fold singularity. However not every Legendrian can be perturbed to have simple front singularities (the case when Σ' is the empty set), there may be topological obstructions (see [AG18]). However, we assume L is front generic with simple front singularities as in [Ekh07] and [Kar17]. Also, after perturbations we assume L satisfies the preliminary transversality condition detailed in Pages 1101-1103 of [Ekh07], and we note this condition is an open condition.

A flow tree determined by L is a continuous map of a source tree T to M whose restriction to any edge is a parameterization of a part of a gradient flow of some local function difference. A broken flow tree determined by L is a flow tree where we relax the condition that edges parameterize parts of gradient flows. In particular, we consider (unparameterized) broken gradient flows of some local function difference.

We define the combinatorial type of a broken flow tree, which characterizes broken flow trees whose associated broken gradient flows are similar. The notion of combinatorial type allows us to formally define what the limit of a sequence of broken flow trees, all of the same combinatorial type, should be. We call this stratum Floer-Gromov convergence. We prove there is a unique topology, the stratum Floer-Gromov topology, on the space of broken flow trees of a fixed combinatorial type (united with all stratum Floer-Gromov limits) where the stratum Floer-Gromov convergence is the topological convergence.

If $\overline{\mathcal{L}}(N)$ denotes the space of broken flow trees with at most N edges and vertices, then combinatorial type gives a finite decomposition of $\overline{\mathcal{L}}(N)$. We may “glue” together the stratum Floer-Gromov topologies to construct the Floer-Gromov topology on $\overline{\mathcal{L}}(N)$.

Theorem 0.1. *The space $\overline{\mathcal{L}}(N)$ is sequentially compact with respect to the Floer-Gromov topology.*

By the natural inclusion of the space of (unparameterized) flow trees with at most N edges and vertices into $\overline{\mathcal{L}}(N)$, we have our desired compactification.

This paper is organized as follows. In Section 1 we recall the the definition and elementary properties of local gradients and flow trees following [Ekh07]. In Section 2 we compactify the various moduli spaces of gradient flows associated to a fixed local function difference following techniques of Section 3.2 of [AD14]. In Section 3 we investigate broken flow trees, construct the (stratum) Floer-Gromov topology, and prove Theorem 0.1.

Acknowledgements. This work was partially conducted while participating at the 2021 DIMACS REU Program at Rutgers University. The author acknowledges support from the Rutgers Department of Mathematics and NSF grant DMS-1711070.

The author would like to thank Christopher Woodward and Soham Chanda for their mentorship and support throughout this project.

1. Flow trees. In this Section we introduce local gradients and flow trees following [Ekh07], and the proofs of the statements of this section may be found there. The main result we are concerned with is Lemma 1.7 (Lemma 3.12 in [Ekh07]) which provides an edge bound for flow trees through geometric data.

1.1. Local gradients. As before, let M be a closed n -manifold equipped with a Riemannian metric g and let $L \subset J^1(M)$ be a closed Legendrian submanifold. We will assume that $\Pi_{\mathbb{C}}(L)$ has only self-transverse double points. Suppose c is a double point of $\Pi_{\mathbb{C}}(L)$, then we write

$$\Pi_{\mathbb{C}}^{-1}(c) \cap L = \{c^+, c^-\},$$

where $z(c^+) > z(c^-)$. Since ∂_z is the Reeb field of the contact form $dz - \alpha$, we note there exists a 1-1 correspondence between self-transverse double points of $\Pi_{\mathbb{C}}(L)$ and Reeb chords on L .

We will assume that L is *front generic* and has *simple front singularities*. In particular, we will assume that L satisfies the following.

- The base projection, $\Pi : L \rightarrow M$, is an immersion outside of a codimension 1 submanifold $\Sigma \subset L$.
- For any point p in Σ , the front projection, $\Pi_F : L \rightarrow J^0(M)$, has a standard cusp edge singularity. Concretely, there exists coordinates $y = (y_1, \dots, y_n)$ around p on L and coordinates $x = (x_1, \dots, x_n)$ around $\Pi(p)$ on M such that, if z is the \mathbb{R} -coordinate in $J^0(M)$, then $\Pi_F(y) = (x(y), z(y))$ where

$$\begin{aligned} x_1(y) &= \frac{1}{2}y_1^2, \\ x_i(y) &= y_i, \quad 2 \leq i \leq n, \\ z(y) &= \frac{1}{3}y_1^3 + b\frac{1}{2}y_1^2 + a_2y_2 + \dots + a_ny_n, \end{aligned} \tag{1}$$

where a_j , $2 \leq j \leq n$, and b are arbitrary real numbers

- The base projection, $\Pi : \Sigma \rightarrow M$, is a self-transverse immersion. We also note that $\Pi(\Sigma) \subset M$ is a closed submanifold of codimension 1.

In dimensions greater than 1, not all Legendrians can be perturbed to have simple front singularities since there may be topological obstructions, however this is a mild genericity condition. We will refer to $\Pi(\Sigma) \subset M$ as the *singular set*. The singular set is naturally stratified as

$$\Pi(\Sigma) = \Sigma^1 \supset \Sigma^2 \supset \cdots \supset \Sigma^k,$$

where Σ^i is the subset of points p in $\Pi(\Sigma)$ such that $\Pi^{-1}(p) \cap \Sigma = \{p_1, \dots, p_i\}$. We also have that $k \leq n$.

Locally, we may describe L as the multi 1-jet graph of *local functions* defined on M . In particular, if p is a point in $M \setminus \Pi(\Sigma)$, there is a maximal open neighborhood $U \subset M \setminus \Pi(\Sigma)$ of p and r disjoint open sets, $U_1, \dots, U_r \subset L \setminus \Sigma$, such that

$$U_i = \{(x, d(f_i)_x, f_i(x)) : x \in U\}$$

for some functions $f_i : U \rightarrow \mathbb{R}$, and each U_i is diffeomorphic to U via the base projection.

Similarly, if p is a point in Σ^j that has r points in $\Pi^{-1}(p) \cap L \setminus \Sigma$, then as above there are r functions defined in a maximal open neighborhood of p in $M \setminus \Pi(\Sigma)$. There is also a maximal open neighborhood $V \subset \Sigma^j$ and j disjoint open sets, $V_1, \dots, V_j \subset \Sigma$, such that any V_i is diffeomorphic to V via the base projection. If we consider a sufficiently small neighborhood U of p in M , then $\Pi(V_i)$ divides U into two components U_i^+ and U_i^- and there are two functions f_i^1 and f_i^2 defined on U_i^+ which extend to $\overline{U_i^+}$, agree on ∂U_i^+ , and the limits of $d(f_i^1)$ and $d(f_i^2)$ agree on ∂U_i^+ when approached from U_i^+ . Since we assume L has only standard cusp edge singularities, these local functions are of the form

$$\begin{aligned} f_i^1(x) &= \frac{1}{3}x_1^{3/2} + bx_1 + a_2x_2 + \cdots + a_nx_n, \\ f_i^2(x) &= -\frac{1}{3}x_1^{3/2} + bx_1 + a_2x_2 + \cdots + a_nx_n, \end{aligned} \tag{2}$$

for $x_1 > 0$, and the common limits of the differentials is

$$bdx_1 + a_2dx_2 + \cdots + a_n dx_n,$$

on $x_1 = 0$, where we use the coordinates of Equation 1 associated to p_i in $\Pi^{-1}(p) \cap \Sigma$. Finally, we note that since L is assumed to be compact, there are only finitely many local functions under consideration.

When specifying a local function f_i we determine a maximal connected component L_i of $L \setminus \Sigma$. If we consider the closure $\overline{L_i} \subset L$, then $\overline{L_i}$ intersected with Σ determines a connected component $\Sigma_{ij} \subset \Sigma$, where the j index refers to $\overline{L_i}$ meeting another such $\overline{L_j}$ determined by some other local function f_j . We call such an L_i a *sheet* of L .

Let f_i and f_j be local functions defined on some common domain $U \subset M$ and consider their *local function difference*

$$F_{ij} = f_i - f_j : U \rightarrow \mathbb{R}.$$

We define the *local gradient of F_{ij}* as the vector field, defined on U , given by

$$\nabla F_{ij} \in \mathfrak{X}(U).$$

Consider a path $\phi : I \rightarrow M$, where $I \subset \mathbb{R}$ is an interval. A *1-jet lift* of ϕ is an unordered pair $\{\tilde{\phi}_1, \tilde{\phi}_2\}$ of continuous lifts $\tilde{\phi}_k : I \rightarrow L$, such that either $\tilde{\phi}_1(t) \neq \tilde{\phi}_2(t)$ or

$$\tilde{\phi}_1(t) = \tilde{\phi}_2(t) \in \Pi^{-1}(\phi(t)) \cap \Sigma.$$

We also define the *cotangent lift* of ϕ as

$$\bar{\phi}_k = \Pi_{\mathbb{C}} \circ \tilde{\phi}_k.$$

Let $\phi : I \rightarrow M$ be a path with 1-jet lift $\{\tilde{\phi}_1, \tilde{\phi}_2\}$ specifying local functions f_i and f_j , and suppose ϕ satisfies the gradient flow equation

$$\dot{\phi}(t) = -\nabla F_{ij}(\phi(t)). \quad (3)$$

Then ϕ is called a *gradient flow of L* associated to F_{ij} . By existence and uniqueness theorems of ODEs, we have that any gradient flow of L has a maximal interval of definition. The asymptotic behavior of gradient flows of L is described as follows.

Lemma 1.1 ([Ekh07]). *Suppose that ϕ is a gradient flow of L that specifies the local functions f_i and f_j . If ϕ is defined on an interval I with an infinite end, (a, ∞) or $(-\infty, b)$, then*

$$\lim_{t \rightarrow \pm\infty} \phi(t) \in \text{Crit}(F_{ij}),$$

where $\text{Crit}(F_{ij})$ is the set of critical points of F_{ij} . If ϕ is defined on an interval I with a compact end, $(a, l]$ or $[l, b)$, then

$$\lim_{t \rightarrow l} \tilde{\phi}_k(t) \in \Sigma, \quad k = 1 \text{ or } k = 2.$$

In particular, $\lim_{t \rightarrow l} \phi(t)$ lies in $\Pi(\Sigma) \cap \bar{U}$.

The preceding Lemma allows us to give a natural categorization of the gradient flows of L associated to some local function difference.

Definition 1.2. Let ϕ be a maximally extended gradient flow of L that satisfies Equation 3 with respect to F_{ij} . Then ϕ is called a

- *Morse flow* if the image of ϕ connects two points of $\text{Crit}(F_{ij})$,
- *fold terminating flow* if the image of ϕ connects a point of $\text{Crit}(F_{ij})$ to a point of $\Pi(\Sigma) \cap \bar{U}$,
- *fold emanating flow* if the image of ϕ connects a point of $\Pi(\Sigma) \cap \bar{U}$ to a point of $\text{Crit}(F_{ij})$,
- *singular flow* if the image of ϕ connects two points of $\Pi(\Sigma) \cap \bar{U}$.

1.2. Flow trees. We now introduce the main objects of study in this paper, flow trees. We will also present some elementary properties of flow trees, culminating in the statement of Lemma 1.7.

A *source tree* is a tree T with finitely many edges. It also has the following extra structure. At each k -valent vertex v of T , $k \geq 2$, the edges adjacent to v are cyclically labelled.

Definition 1.3 ([Ekh07]). A *flow tree* of L is a continuous map $\phi : T \rightarrow M$, where T is a source tree, satisfying the following.

1. If E is an edge of T , then $\phi : E \rightarrow M$ is an injective parametrization of a part of a gradient flow of L .
2. Let v be a k -valent vertex of T with cyclically ordered edges E_1, \dots, E_k . Let $\{\bar{\phi}_j^1, \bar{\phi}_j^2\}$ be the cotangent lift corresponding to E_j . We require that

$$\bar{\phi}_j^2(v) = \bar{\phi}_{j+1}^1(v) = \bar{p} \in \Pi_{\mathbb{C}}(L),$$

such that the flow orientation of $\bar{\phi}_j^2$ at \bar{p} is directed toward \bar{p} if and only if the flow orientation of $\bar{\phi}_{j+1}^1$ at \bar{p} is directed away from \bar{p} .

3. We require that the cotangent lifts of the edges of T yield a loop in $\Pi_{\mathbb{C}}(L)$.

We denote the 1-jet lift of a flow tree T as \tilde{T} . Similarly, we denote the cotangent lift as \bar{T} .

Let T be a flow tree and let v be a k -valent vertex of T with cyclically ordered edges E_1, \dots, E_k . Suppose $\bar{\phi}_j^2, \bar{\phi}_{j+1}^1$ are two paired cotangent lifts, with corresponding 1-jet lifts $\tilde{\phi}_j^2, \tilde{\phi}_{j+1}^1$, at v . If

$$\tilde{\phi}_j^2(v) \neq \tilde{\phi}_{j+1}^1(v),$$

then $\{\tilde{\phi}_j^2(v), \tilde{\phi}_{j+1}^1(v)\}$ are Reeb chord endpoints, and we say that the vertex v contains a *puncture* after the edge E_j .

It may be the case that, at a k -valent vertex v of T , $k \geq 2$, the vertex v may contain a puncture after multiple edges. However, the following Lemma shows that we may assume each vertex has at most 1 puncture.

Lemma 1.4 ([Ekh07]). *Any flow tree T with a vertex v which contains more than one puncture is a union of flow trees*

$$T = T_1 \cup \dots \cup T_n$$

such that every vertex of each T_i contains at most one puncture.

Lemma 1.4 allows us to only consider flow trees T such that, at any vertex v of T , the vertex v contains at most one puncture. For brevity, we will now say that any vertex which contains a puncture is a puncture. Punctures of a flow tree may be one of two types.

Definition 1.5 ([Ekh07]). Let v be a puncture of a flow tree T and let $\bar{\phi}_j^2, \bar{\phi}_{j+1}^1$ be the two paired cotangent lifts, with corresponding 1-jet lifts $\tilde{\phi}_j^2, \tilde{\phi}_{j+1}^1$, such that $\tilde{\phi}_j^2(v) \neq \tilde{\phi}_{j+1}^1(v)$.

Choose notation such that $\tilde{\phi}_{j+1}^1$ is oriented towards $\tilde{\phi}_{j+1}^1(v)$ and $\tilde{\phi}_j^2$ is oriented away from $\tilde{\phi}_j^2(v)$. We say that v is a *positive* puncture if

$$z(\tilde{\phi}_{j+1}^1(v)) < z(\tilde{\phi}_j^2(v)),$$

where we recall that z is the coordinate in the \mathbb{R} -direction of $J^1(M) = T^*M \times \mathbb{R}$. We say v is a *negative* puncture if the opposite inequality is true.

We will now assume that L satisfies the *preliminary transversality condition*, which is found in pages 1101-1103 in [Ekh07]. In short, the preliminary transversality condition affects how the various sheets of the Legendrian meet, and the consequences of the condition ensure flows of L meet the singular set in a locally stable fashion. In particular, any flow line of L may only meet the singular set with order of contact at most n (hence no flow line of L may flow along the singular set). We must assume L satisfies the preliminary transversality condition in order for the remaining Definition and Lemma of this Section to remain true. We also note that the preliminary transversality condition may be obtained after small perturbations of L and is an open condition.

Before the next definition, we shall set up some notation. Let T be a flow tree. We have the follow types of vertices of T .

- $P(T)$ is the set of positive punctures of T .
- $Q(T)$ is the set of negative punctures of T .
- $R(T)$ is the set of vertices of T that are not punctures.

Let v be a puncture of T , then there exists a Reeb chord c corresponding to v . Let c^\pm in L be the endpoints of c such that $z(c^+) > z(c^-)$. The *index* $I(v)$ of v is equal to the Morse index of the critical point $\Pi(c)$ in M of $f^+ - f^-$, where f^\pm are the local functions determined by the sheets of L containing c^\pm .

Let v be a vertex of T that is not a puncture and consider the given map $\phi : T \rightarrow M$. Suppose x in Σ is a point such that $\Pi(x) = \phi(v)$ in M and x lies in \tilde{T} . We define a function $\tilde{\mu} : R(T) \rightarrow \{-1, 0, 1\}$ by

- $\tilde{\mu}(x) = 1$ if the incoming segment of \tilde{T} at x lies in the upper sheet determined by \tilde{T} and the outgoing segment of \tilde{T} at x lies in the lower sheet determined by \tilde{T} ,
- $\tilde{\mu}(x) = -1$ if the incoming segment of \tilde{T} at x lies in the lower sheet determined by \tilde{T} and the outgoing segment of \tilde{T} at x lies in the upper sheet determined by \tilde{T} ,
- $\tilde{\mu}(x) = 0$ otherwise.

Then, for a vertex v of T that is not a puncture, we define the *Maslov content* of v as

$$\mu(v) = \sum_{x \in \Pi^{-1}(\phi(v)) \cap \tilde{T}} \tilde{\mu}(x).$$

With the notation set, we may define the formal dimension of a flow tree.

Definition 1.6 ([Ekh07]). The *formal dimension* of a flow tree T is

$$\dim(T) = \sum_{p \in P(T)} I(p) + \sum_{q \in Q(T)} (n - I(q)) + \sum_{r \in R(T)} \mu(r) - (|P(T)| + |Q(T)| - 1)n + (|P(T)| + |Q(T)| - 3),$$

where we recall that n is the dimension of the base manifold M .

With the background on flow trees complete, we may now state the main Lemma of this Section.

Lemma 1.7 ([Ekh07]). *There exists a constant $N > 0$ such that, for any flow tree T with at most P positive punctures and of formal dimension at most $D > 0$, the flow tree T has no more than N edges and vertices.*

2. Gradient flows of local function differences. If we are given a local function difference F_{ij} , then the assumption that the Lagrangian projection $\Pi_{\mathbb{C}}$ has only self-transverse double points implies F_{ij} is a Morse function on its domain $U \subset M$. Thus, in this Section we compactify the moduli spaces of gradient flows of a local function difference, arising from Definition 1.2, using techniques from Morse theory. In particular, we will follow Section 3.2 of [AD14] and explicitly build our topologies on the moduli spaces, instead of considering the moduli spaces as function spaces as, for example, [FN20] does.

Throughout this section, we fix a local function difference F_{ij} on its domain $U \subset M$.

2.1. Morse theory. For convenience, we restate the Morse Lemma.

Lemma 2.1. *Let c be a critical point of the Morse function F_{ij} . Then there exists a neighborhood $\Omega(c) \subset U$ of c and a diffeomorphism $\varphi : \Omega(c) \rightarrow \mathbb{R}^n$ such that*

$$(F_{ij} \circ \varphi^{-1})(x_1, \dots, x_n) = F_{ij}(c) - \sum_{k=1}^m x_k^2 + \sum_{k=m+1}^n x_k^2.$$

Even though U is not compact, Lemma 2.1 is a local statement that still holds and allows us to construct nice neighborhoods of our critical points.

In the case of an arbitrary closed manifold and a single Morse function defined on the entire manifold, a Corollary of the Morse Lemma is that the Morse function under consideration has finitely many critical points that are isolated. It is fortunate that in our case, the same holds true for F_{ij} . After small perturbation, the self-transverse double points of $\Pi_{\mathbb{C}}(L)$ are isolated (note this equivalent to assuming the Reeb chords on L are isolated, which may always be achieved after perturbation, and L is said to be *chord generic*). Since the critical points of F_{ij} correspond to self-transverse double points of $\Pi_{\mathbb{C}}(L)$ and L is compact, the claim follows. In fact, after small perturbation, we may also assume the following. If p in M is a critical point of F_{ij} , then for any other distinct local function f_k that determines a sheet L_k which meets L_i or L_j , we have

$$p \notin \Pi(\Sigma_{ik}), \quad p \notin \Pi(\Sigma_{kj}).$$

We define the set $W^s(c)$ to be

$$W^s(c) = \{\phi \text{ is a flow of } -\nabla F_{ij} : \lim_{t \rightarrow \infty} \phi(t) = c\},$$

and the set $W^u(c)$ to be

$$W^u(c) = \{\phi \text{ is a flow of } -\nabla F_{ij} : \lim_{t \rightarrow -\infty} \phi(t) = c\},$$

where c is a critical point of F_{ij} . We call $W^s(c)$ and $W^u(c)$ the *stable manifold of c* and the *unstable manifold of c* , respectively. If the critical point c of F_{ij} has index $I(c)$, then an investigation of the Morse neighborhood $\Omega(c)$ of c yields the following Lemma, and we refer to Pages 27-28 of [AD14] for the details.

Lemma 2.2. *The stable manifold of a critical point c intersected with the boundary of a Morse neighborhood of c is diffeomorphic to a sphere of dimension $n - I(c) - 1$, i.e.*

$$W^s(c) \cap \partial\Omega(c) \cong S^{n-I(c)-1}.$$

Similarly, we have

$$W^u(c) \cap \partial\Omega(c) \cong S^{I(c)-1}.$$

We these local results of Morse theory, we are now ready to consider the sets of gradient flows of our local function difference, categorized by Definition 1.2.

2.2. Morse flows. Let c^-, c^+ be critical points of the local function difference F , where c^- and c^+ are not necessarily distinct, and let $\mathcal{M}(F_{ij}, c^-, c^+)$ be the set of Morse flows of F . Note that if c^- is equal to c^+ , then $\mathcal{M}(F_{ij}, c^-, c^+)$ is a set with only one element.

If ϕ is a gradient flow of L in $\mathcal{M}(F_{ij}, c^-, c^+)$, then the domain of ϕ is an infinite interval $(-\infty, \infty)$. There is a natural \mathbb{R} -action on $\mathcal{M}(F_{ij}, c^-, c^+)$, given by

$$(r, \phi) \mapsto r_*\phi(t) = \phi(t + r), \quad r \in \mathbb{R},$$

where the Morse flow $r_*\phi$ is defined on the infinite interval $(-\infty, \infty)$. Given this \mathbb{R} -action, we may define the *set of unparameterized Morse flows connecting c^- to c^+* as

$$\mathcal{L}(F_{ij}, c^-, c^+) = \mathcal{M}(F_{ij}, c^-, c^+)/\mathbb{R}.$$

Again, note that if c^- is equal to c^+ , then $\mathcal{L}(F_{ij}, c^-, c^+)$ is a set with only one element.

Definition 2.3. Let c_0, \dots, c_{q+1} be critical points of F_{ij} such that

$$c_0 = c^- \quad \text{and} \quad c_{q+1} = c^+.$$

A q -times broken Morse flow connecting c^- to c^+ , denoted $\lambda = (\lambda_1, \dots, \lambda_{q+1})$, is a concatenation of paths λ_k such that

$$\lambda_k \in \mathcal{L}(F_{ij}, c_{k-1}, c_k).$$

It should be noted that a 0-times broken Morse flow is simply an unparameterized Morse flow.

We denote by $\overline{\mathcal{L}}(F_{ij}, c^-, c^+)$ the *moduli space of broken Morse flows connecting c^- to c^+* , which has a natural stratification

$$\overline{\mathcal{L}}(F_{ij}, c^-, c^+) = \bigcup_{q=0}^{\infty} \mathcal{L}_q(F_{ij}, c^-, c^+),$$

where $\mathcal{L}_q(F_{ij}, c^-, c^+)$ is the *moduli space of q -times broken Morse flows connecting c^- to c^+* . We will now construct a topology on $\overline{\mathcal{L}}(F_{ij}, c^-, c^+)$.

We assume that c^- and c^+ are distinct, for if they are the same, then $\overline{\mathcal{L}}(F_{ij}, c^-, c^+)$ consists of a single element and is given the trivial topology. Consider an element $\lambda = (\lambda_1, \dots, \lambda_{q+1})$ in $\overline{\mathcal{L}}(F, c^-, c^+)$, let $c^- = c_0$ and $c^+ = c_{q+1}$, and let $\Omega(c_k)$ be a Morse neighborhood of c_k . Consider a neighborhood U_{k-1}^- of the exit point of the image of λ_k from $\Omega(c_{k-1})$ and consider a neighborhood U_k^+ of the entry point of the image of λ_k into $\Omega(c_k)$, such that U_{k-1}^- and U_k^+ are both contained in their respective level sets. See Figure 1.




Figure 1: An arbitrary λ in $\overline{\mathcal{L}}(F_{ij}, c^-, c^+)$ ($c^- \neq c^+$), with λ 's exit and entry neighborhoods.

Denote by U^- and U^+ the collection of the U_k^- 's and U_k^+ 's respectively. We define the set $\mathcal{W}(\lambda, U^-, U^+)$ to be the set of tuples $\mu = (\mu_1, \dots, \mu_r)$, for an arbitrary $r \leq q + 1$, with the property that there exists integers

$$0 = k_0 < k_1 < \dots < k_{r-1} < k_r = q + 1$$

such that

- μ_s is in $\mathcal{L}(F_{ij}, c_{k_{s-1}}, c_{k_s})$ for $1 \leq s \leq r$,

- and the image of μ_s exits $\Omega(c_{k_{s-1}})$ through $U_{k_{s-1}}^-$ and enters $\Omega(c_{k_s})$ through $U_{k_s}^+$ for $1 \leq s \leq r$.

Then the collection of $\mathcal{W}(\lambda, U^-, U^+)$ is a neighborhood basis for a topology on $\overline{\mathcal{L}}(F_{ij}, c^-, c^+)$. And since the topology was constructed from the topology of U , it follows that the constructed topology is first countable and Hausdorff. We will require the following standard Lemma, which is Lemma 3.2.5 in [AD14].

Lemma 2.4. *Let x be in $U \setminus \text{Crit}(F_{ij})$ and let x_n in U a sequence converging to x . Let y_n and y be points lying on the same flows of $-\nabla F_{ij}$ as x_n and x , respectively. Suppose that $F_{ij}(y_n) = F_{ij}(y)$ for all n . Then*

$$y_n \rightarrow y.$$

Before proving $\overline{\mathcal{L}}(F_{ij}, c^-, c^+)$ is a compactification of $\mathcal{L}(F_{ij}, c^-, c^+)$, we will prove the following special case.

Lemma 2.5. *Let ϕ_n be a sequence of unparameterized Morse flows of F_{ij} connecting c^- to c^+ . Then there exists a finite set of critical points c_1, \dots, c_q of F_{ij} , for some q , and a broken Morse flow $\lambda = (\lambda_1, \dots, \lambda_{q+1})$ of F_{ij} connecting c^- to c_1 , c_1 to c_2 , and so on, such that (after passing to a subsequence) ϕ_n converges to λ .*

Proof. Again, we let $c^- = c_0$ and $c^+ = c_{q+1}$. If c_0 and c_{q+1} are equal, then the Lemma is true, since the sequence must be the constant sequence. Hence, we may assume c_0 and c_{q+1} are distinct.

We have that the image of ϕ_n exits the boundary of $\Omega(c_0)$ through a point ϕ_n^- in $\partial\Omega(c_0)$. Since the sequence ϕ_n^- is in $W^u(c_0) \cap \partial\Omega(c_0)$, which is a sphere by Lemma 2.2, we may pass to a subsequence and assume ϕ_n^- converges to a point c_0^- in $W^u(c_0) \cap \partial\Omega(c_0)$. Let λ_1 be the unique unparameterized gradient flow of F_{ij} defined by c_0^- , then it follows

$$\lim_{t \rightarrow -\infty} \lambda_1(t) = c_0.$$

We claim that there exists a critical point c_1 of F_{ij} such that λ_1 is in $\mathcal{L}(F_{ij}, c_0, c_1)$. If this were not the case, then λ_1 would flow into the singular set and end there.

- Suppose λ_1 flows into $\Pi(\Sigma_{ij})$ and ends. In a neighborhood of the ending point of λ_1 , our local functions f_i and f_j are given by the local form in Equation 2. It follows the sequence ϕ_n must flow into $\Pi(\Sigma_{ij})$, which is a contradiction.
- Suppose λ_1 flows into $\Pi(\Sigma_{ik})$ and ends, where the index k refers to some other distinct local function f_k determining a sheet L_k which meets L_i . Then since ∇F_{ij} is non-singular on $\Pi(\Sigma_{ik})$, it follows by dependence of solutions to ODEs on initial conditions that the sequence ϕ_n must flow into $\Pi(\Sigma_{ik})$, which is a contradiction. The same conclusion would hold if λ_1 flows into $\Pi(\Sigma_{kj})$.

Hence there exists a critical point c_1 of F_{ij} such that λ_1 is in $\mathcal{L}(F_{ij}, c_0, c_1)$. Let d^+ in $\partial\Omega(c_1)$ be the entry point of the image of λ_1 into $\Omega(c_1)$. Again, by dependence of solutions to ODEs on initial conditions, we have that the image of ϕ_n must also enter $\Omega(c_1)$ through a point

d_n^+ in $\partial\Omega(c_1)$, for n sufficiently large. We have that d_n^+ converges to d^+ by Lemma 2.4. See Figure 2.




Figure 2: The constructed gradient flow λ_1 .

If c_1 is equal to c_{q+1} then we are done. If c_1 is not equal to c_{q+1} , then d_n^+ does not lie in $W^s(c_1)$, hence the image of ϕ_n exits $\Omega(c_1)$ through a point d_n^- in $\partial\Omega(c_1)$. Since $\partial\Omega(c_1)$ is compact, we may pass to a subsequence and assume d_n^- converges to a point d^- in $\partial\Omega(c_1)$. We claim d^- is in $W^u(c_1)$. Suppose this were not the case. Then d^- would lie on the image of a gradient flow γ of F_{ij} whose image enters $\Omega(c_1)$ through a point d_* in $\partial\Omega(c_1)$, such that $F(d_n^+) = F(d_*)$ and d_* is not in $W^s(c_1)$. By Lemma 2.4, we have that d_n^+ converges to d_* and therefore d_* is equal to d^+ , which is a contradiction to the fact that d^+ was shown to be in $W^s(c_1)$.

We may now induct over the number of critical points between c_0 and c_{q+1} , since $\text{Crit}(F_{ij})$ is a finite set, to construct the limit $\lambda = (\lambda_1, \dots, \lambda_{q+1})$ in $\overline{\mathcal{L}}(F_{ij}, c^-, c^+)$ of the sequence ϕ_n . \square

With Lemma 2.5, we may now show that $\overline{\mathcal{L}}(F_{ij}, c^-, c^+)$ compactifies $\mathcal{L}(F_{ij}, c^-, c^+)$, in the sense that every sequence has a convergent subsequence.

Proposition 2.6. $\overline{\mathcal{L}}(F_{ij}, c^-, c^+)$ is sequentially compact.

Proof. If c^- and c^+ are equal, then the Proposition is true, since the sequence must be the constant sequence. Hence, we may assume c^- and c^+ are distinct.

Let λ_n be a sequence of broken Morse flows connecting c^- to c^+ . By passing to a subsequence we may assume the sequence λ_n has a fixed number of components, i.e.

$$\lambda_n = (\lambda_n^1, \dots, \lambda_n^{q+1}) \in \mathcal{L}_q(F_{ij}, c^-, c^+)$$

for some q . We now apply Lemma 2.5 to each λ_n^k , $1 \leq k \leq q+1$, individually. \square

2.3. Fold emanating flows. Let c^+ be a critical point of the local function difference F_{ij} and let $\mathcal{M}(F_{ij}, \Pi(\Sigma), c^+)$ be the set of fold emanating flows of F_{ij} .

If ϕ is a gradient flow of L in $\mathcal{M}(F_{ij}, \Pi(\Sigma), c^+)$, then the domain of ϕ is a semi-infinite interval of the form $[a, \infty)$. There is a natural \mathbb{R} -action on $\mathcal{M}(F_{ij}, \Pi(\Sigma), c^+)$, given by

$$(r, \phi) \mapsto r_*\phi(t) = \phi(t + r), \quad r \in \mathbb{R},$$

where the fold emanating flow $r_*\phi$ is defined on the semi-infinite interval $[a + r, \infty)$. Given this \mathbb{R} -action, we may define the *set of unparameterized fold emanating flows ending at c^+* as

$$\mathcal{L}(F_{ij}, \Pi(\Sigma), c^+) = \mathcal{M}(F_{ij}, \Pi(\Sigma), c^+)/\mathbb{R}.$$

Definition 2.7. Let $c_1 \dots, c_{q+1}$ be critical points of F_{ij} such that

$$c_{q+1} = c^+.$$

A q -times broken fold emanating flow ending at c^+ , denoted $\lambda = (\lambda_1, \dots, \lambda_{q+1})$, is a concatenation of paths λ_k such that

$$\lambda_1 \in \mathcal{L}(F_{ij}, \Pi(\Sigma), c_1) \quad \text{and} \quad \lambda_k \in \mathcal{L}(F_{ij}, c_{k-1}, c_k)$$

where $2 \leq k \leq q + 1$.

It should be noted that a 0-times broken fold emanating flow is simply an unparameterized fold emanating flow.

We denote by $\overline{\mathcal{L}}(F_{ij}, \Pi(\Sigma), c^+)$ the *moduli space of broken fold emanating flows ending at c^+* , which has a natural stratification

$$\overline{\mathcal{L}}(F_{ij}, \Pi(\Sigma), c^+) = \bigcup_{q=0}^{\infty} \mathcal{L}_q(F_{ij}, \Pi(\Sigma), c^+)$$

where $\mathcal{L}_q(F_{ij}, \Pi(\Sigma), c^+)$ is the *moduli space of q -times broken fold emanating flows ending at c^+* . We will now construct a topology on $\overline{\mathcal{L}}(F_{ij}, \Pi(\Sigma), c^+)$.

Consider an element $\lambda = (\lambda_1, \dots, \lambda_{q+1})$ in $\overline{\mathcal{L}}(F_{ij}, \Pi(\Sigma), c^+)$, let $c_{q+1} = c^+$, and let $\Omega(c_k)$ be a Morse neighborhood of c_k , for $k \geq 1$. Consider a neighborhood U_{k-1}^- of the exit point of the image of λ_k from $\Omega(c_{k-1})$ and consider a neighborhood U_k^+ of the entry point of the image of λ_k into $\Omega(c_k)$, such that U_{k-1}^- and U_k^+ are both contained in their respective level sets, for $k \geq 2$. Let c_0 be the point in $\Pi(\Sigma)$ that the image of λ_1 emanates from. Consider a neighborhood U_0^- of c_0 contained in $\Pi(\Sigma)$ and consider a neighborhood U_1^+ of the entry point of the image of λ_1 into $\Omega(c_1)$. See Figure 3.

Denote by U^- and U^+ the collection of the U_k^- 's and U_k^+ 's respectively. We define the set $\mathcal{W}(\lambda, U^-, U^+)$ to be the set of tuples $\mu = (\mu_1, \dots, \mu_r)$, for an arbitrary $r \leq q + 1$, with the property that there exists integers

$$0 = k_0 < k_1 < \dots < k_{r-1} < k_r = q + 1$$

such that




Figure 3: An arbitrary λ in $\overline{\mathcal{L}}(F_{ij}, \Pi(\Sigma), c^+)$, with λ 's exit and entry neighborhoods.

- μ_s is in $\mathcal{L}(F_{ij}, c_{k_{s-1}}, c_{k_s})$ for $2 \leq s \leq r$,
- μ_1 is in $\mathcal{L}(F_{ij}, \Pi(\Sigma), c_{k_2})$,
- the image of μ_s exits $\Omega(c_{k_{s-1}})$ through $U_{k_{s-1}}^-$ and enters $\Omega(c_{k_s})$ through $U_{k_s}^+$ for $2 \leq s \leq r$,
- and the image of μ_1 emanates from U_0^- and enters $\Omega(c_{i_2})$ through $U_{i_2}^+$.

Then the collection of $\mathcal{W}(\lambda, U^-, U^+)$ is a neighborhood basis for a topology on $\overline{\mathcal{L}}(F_{ij}, \Pi(\Sigma), c^+)$. And since the topology was constructed from the topology of U and $\Pi(\Sigma)$, it follows that the constructed topology is first countable and Hausdorff.

Before proving $\overline{\mathcal{L}}(F_{ij}, \Pi(\Sigma), c^+)$ is a compactification of $\mathcal{L}(F_{ij}, \Pi(\Sigma), c^+)$, we will prove the following special case.

Lemma 2.8. *Let ϕ_n be a sequence of unparameterized fold emanating flows of F_{ij} ending at c^+ . Then there exists a point c_0 in $\Pi(\Sigma)$, a finite set of critical points c_1, \dots, c_q of F_{ij} , for some q , and a broken fold emanating flow $\lambda = (\lambda_1, \dots, \lambda_{q+1})$ of F_{ij} connecting c_0 to c_1 , c_1 to c_2 , and so on, such that (after passing to a subsequence) ϕ_n converges to λ .*

Proof. Again, we let $c^+ = c_{q+1}$, and we recall the sequence of ϕ_n is defined on the domain $[a_n, \infty)$. We have that $\Pi(\Sigma) \cap \overline{U}$ is compact, therefore, we may pass to a subsequence and assume $\phi_n(a_n)$ converges to a point c_0 in $\Pi(\Sigma) \cap \overline{U}$. Let λ_1 be the unique unparameterized flow of F_{ij} emanating from c_0 .

We claim that there exists a critical point c_1 of F_{ij} such that λ_1 is in $\mathcal{L}(F_{ij}, \Pi(\Sigma), c_1)$. If this were not the case, then λ_1 would flow into the singular set and end there.

- Suppose λ_1 flows into $\Pi(\Sigma_{ij})$ and ends. In a neighborhood of the ending point of λ_1 , our local functions f_i and f_j are given by the local form in Equation 2. It follows the

sequence ϕ_n must flow into $\Pi(\Sigma_{ij})$, which is a contradiction.

- Suppose λ_1 flows into $\Pi(\Sigma_{ik})$ and ends, where the index k refers to some other distinct local function f_k determining a sheet L_k which meets L_i . Then since ∇F_{ij} is non-singular on $\Pi(\Sigma_{ik})$, it follows by dependence of solutions to ODEs on initial conditions that the sequence ϕ_n must flow into $\Pi(\Sigma_{ik})$, which is a contradiction. The same conclusion would hold if λ_1 flows into $\Pi(\Sigma_{kj})$.

Hence there exists a critical point c_1 of F such that λ_1 is in $\mathcal{L}(F_{ij}, \Pi(\Sigma), c_1)$. We may now apply the argument of Lemma 2.5.

We may now induct over the number of critical points between c_0 and c_{q+1} using the argument of Lemma 2.5, since $\text{Crit}(F_{ij})$ is a finite set, to construct the limit $\lambda = (\lambda_1, \dots, \lambda_{q+1})$ in $\overline{\mathcal{L}}(F_{ij}, \Pi(\Sigma), c^+)$ of the sequence ϕ_n . \square

With Lemma 2.8, we may now show that $\overline{\mathcal{L}}(F_{ij}, \Pi(\Sigma), c^+)$ compactifies $\mathcal{L}(F_{ij}, \Pi(\Sigma), c^+)$, in the sense that every sequence has a convergent subsequence.

Proposition 2.9. $\overline{\mathcal{L}}(F_{ij}, \Pi(\Sigma), c^+)$ is sequentially compact.

Proof. Let λ_n be a sequence of broken fold emanating flows ending at c^+ . By passing to a subsequence we may assume the sequence λ_n has a fixed number of components, i.e.

$$\lambda_n = (\lambda_n^1, \dots, \lambda_n^{q+1}) \in \mathcal{L}_q(F_{ij}, \Pi(\Sigma), c^+)$$

for some q .

- If $q \geq 1$, we now apply Lemma 2.8 to λ_n^1 and Lemma 2.5 to each λ_n^k , $2 \leq k \leq q+1$, individually.
- If $q = 0$, we now apply Lemma 2.8 to λ_n^1 .

\square

2.4. Fold terminating flows. Let c^- be a critical point of the local function difference F_{ij} and let $\mathcal{M}(F_{ij}, c^-, \Pi(\Sigma))$ be the set of fold terminating flows of F_{ij} .

If ϕ is a gradient flow of L in $\mathcal{M}(F_{ij}, c^-, \Pi(\Sigma))$, then the domain of ϕ is a semi-infinite interval of the form $(-\infty, b]$. There is a natural \mathbb{R} -action on $\mathcal{M}(F_{ij}, c^-, \Pi(\Sigma))$, given by

$$(r, \phi) \mapsto r_*\phi(t) = \phi(t+r), \quad r \in \mathbb{R},$$

where the fold terminating flow $r_*\phi$ is defined on the semi-infinite interval $(-\infty, b+r]$. Given this \mathbb{R} -action, we may define the *set of unparameterized fold terminating flows starting at c^-* as

$$\mathcal{L}(F_{ij}, c^-, \Pi(\Sigma)) = \mathcal{M}(F_{ij}, c^-, \Pi(\Sigma)) / \mathbb{R}.$$

Definition 2.10. Let c_0, \dots, c_q be critical points of F_{ij} such that

$$c_0 = c^-.$$

A q -times broken fold terminating flow starting at c^- , denoted $\lambda = (\lambda_1, \dots, \lambda_{q+1})$, is a concatenation of paths λ_k such that

$$\lambda_k \in \mathcal{L}(F_{ij}, c_{k-1}, c_k) \text{ and } \lambda_{q+1} \in \mathcal{L}(F_{ij}, c_q, \Pi(\Sigma))$$

where $1 \leq k \leq q$.

It should be noted that a 0-times broken fold terminating flow is simply an unparameterized fold terminating flow.

We denote by $\overline{\mathcal{L}}(F_{ij}, c^-, \Pi(\Sigma))$ the *moduli space of broken fold terminating flows starting at c^-* , which has a natural stratification

$$\overline{\mathcal{L}}(F_{ij}, c^-, \Pi(\Sigma)) = \bigcup_{q=0}^{\infty} \mathcal{L}_q(F_{ij}, c^-, \Pi(\Sigma))$$

where $\mathcal{L}_q(F_{ij}, c^-, \Pi(\Sigma))$ is the *moduli space of q -times broken fold terminating flows starting at c^-* . We will now construct a topology on $\overline{\mathcal{L}}(F_{ij}, c^-, \Pi(\Sigma))$.

Consider an element $\lambda = (\lambda_1, \dots, \lambda_{q+1})$ in $\overline{\mathcal{L}}(F_{ij}, c^-, \Pi(\Sigma))$, let $c_0 = c^-$, and let $\Omega(c_k)$ be a Morse neighborhood of c_k , for $k \leq q$. Consider a neighborhood U_{k-1}^- of the exit point of the image of λ_k from $\Omega(c_{k-1})$ and consider a neighborhood U_k^+ of the entry point of the image of λ_k into $\Omega(c_k)$, such that U_{k-1}^- and U_k^+ are both contained in their respective level sets, for $k \leq q$. Let c_{q+1} be the point in $\Pi(\Sigma)$ that the image of λ_{q+1} terminates at. Consider a neighborhood U_{q+1}^+ of c_{q+1} contained in $\Pi(\Sigma)$ and consider a neighborhood U_q^- of the exit point of the image of λ_{q+1} from $\Omega(c_q)$. See Figure 4.




Figure 4: An arbitrary λ in $\overline{\mathcal{L}}(F_{ij}, c^-, \Pi(\Sigma))$, with λ 's exit and entry neighborhoods.

Denote by U^- and U^+ the collection of the U_i^- 's and U_i^+ 's respectively. We define the set $\mathcal{W}(\lambda, U^-, U^+)$ to be the set of tuples $\mu = (\mu_1, \dots, \mu_r)$, for an arbitrary $r \leq q + 1$, with the

property that there exists integers

$$0 = k_0 < k_1 < \cdots < k_{r-1} < k_r = q + 1$$

such that

- μ_s is in $\mathcal{L}(F_{ij}, c_{k_{s-1}}, c_{k_s})$ for $1 \leq s \leq r - 1$,
- μ_r is in $\mathcal{L}(F_{ij}, c_{k_{r-1}}, \Pi(\Sigma))$,
- the image of μ_s exits $\Omega(c_{k_{s-1}})$ through $U_{k_{s-1}}^-$ and enters $\Omega(c_{k_s})$ through $U_{k_s}^+$ for $1 \leq s \leq r - 1$,
- and the image of μ_r exits $\Omega(c_{k_{r-1}})$ through $U_{k_{r-1}}^-$ and terminates in U_{q+1}^+ .

Then the collection of $\mathcal{W}(\lambda, U^-, U^+)$ is a neighborhood basis for a topology on $\overline{\mathcal{L}}(F_{ij}, c^-, \Pi(\Sigma))$. And since the topology was constructed from the topology of U and $\Pi(\Sigma)$, it follows that the constructed topology is first countable and Hausdorff.

Before proving $\overline{\mathcal{L}}(F_{ij}, c^-, \Pi(\Sigma))$ is a compactification of $\mathcal{L}(F_{ij}, c^-, \Pi(\Sigma))$, we will prove the following special case.

Lemma 2.11. *Let ϕ_n be a sequence of unparameterized fold terminating flows of F_{ij} starting at c^- . Then there exists a point c_{q+1} in $\Pi(\Sigma)$, a finite set of critical points c_1, \dots, c_q of F_{ij} , for some q , and a broken fold terminating flow $\lambda = (\lambda_1, \dots, \lambda_{q+1})$ of F_{ij} connecting c^- to c_1 , c_1 to c_2 , and so on, such that (after passing to a subsequence) ϕ_n converges to λ .*

Proof. Again, we let $c^- = c_0$, and we recall that the sequence ϕ_n is defined on the domain $(-\infty, b_n]$. We have that $\Pi(\Sigma) \cap \overline{U}$ is compact, therefore, we may pass to a subsequence and assume $\phi_n(b_n)$ converges to a point c_{q+1} in $\Pi(\Sigma) \cap \overline{U}$.

We have that the image of ϕ_n exits the boundary of $\Omega(c_0)$ through a point ϕ_n^- in $\partial\Omega(c_0)$. Since the sequence ϕ_n^- is in $W^u(c_0) \cap \partial\Omega(c_0)$, which is a sphere by Lemma 2.2, we may pass to a subsequence and assume ϕ_n^- converges to a point c_0^- in $W^u(c_0) \cap \partial\Omega(c_0)$. Let λ_1 be the unique unparameterized gradient flow of F_{ij} defined by c_0^- , then it follows

$$\lim_{t \rightarrow -\infty} \lambda_1(t) = c_0.$$

We now have two cases.

1. It may be the case that λ_1 is in $\mathcal{L}(F_{ij}, c_0, \Pi(\Sigma))$. Hence, λ_1 flows into the singular set and ends there. Let λ_1 be defined on the domain $(-\infty, b]$.
 - Suppose λ_1 flows into $\Pi(\Sigma_{ij})$ and ends. In a neighborhood of the ending point of λ_1 , our local functions f_i and f_j are given by the local form in Equation 2. It follows the sequence ϕ_n satisfies $\lim_{n \rightarrow \infty} \phi_n(b_n) = \lambda_1(b)$. In particular, this shows

$$\lim_{t \rightarrow b} \lambda_1(t) = c_{q+1},$$

hence the sequence ϕ_n converges to $\lambda = (\lambda_1)$, and there is no breaking of the sequence of gradient flows.

- Suppose λ_1 flows into $\Pi(\Sigma_{ik})$ and ends, where the index k refers to some other distinct local function f_k determining a sheet L_k which meets L_i . Then since ∇F_{ij} is non-singular on $\Pi(\Sigma_{ik})$, it follows by dependence of solutions to ODEs on initial conditions that the sequence ϕ_n satisfies $\lim_{n \rightarrow \infty} \phi_n(b_n) = \lambda_1(b)$. In particular, this shows

$$\lim_{t \rightarrow b} \lambda_1(t) = c_{q+1},$$

hence the sequence ϕ_n converges to $\lambda = (\lambda_1)$, and there is no breaking of the sequence of gradient flows. The same conclusion would hold if λ_1 flows into $\Pi(\Sigma_{kj})$.

2. It may also be the case that there exists a critical point c_1 of F_{ij} such that λ_1 is in $\mathcal{L}(F_{ij}, c_0, c_1)$. We then apply the argument of Lemma 2.5.

We may now induct over the number of critical points between c_0 and c_{q+1} using the argument of Lemma 2.5, since $\text{Crit}(F_{ij})$ is a finite set, to construct the limit $\lambda = (\lambda_1, \dots, \lambda_q)$ in $\overline{\mathcal{L}}(F_{ij}, c^-, \Pi(\Sigma))$ of the sequence ϕ_n . We note that in any step of the induction, if the constructed λ_k flows into $\Pi(\Sigma)$, then the induction ends. \square

With Lemma 2.11, we may now show that $\overline{\mathcal{L}}(F_{ij}, c^-, \Pi(\Sigma))$ compactifies $\mathcal{L}(F_{ij}, c^-, \Pi(\Sigma))$, in the sense that every sequence has a convergent subsequence.

Proposition 2.12. $\overline{\mathcal{L}}(F_{ij}, c^-, \Pi(\Sigma))$ is sequentially compact.

Proof. Let λ_n be a sequence of broken fold terminating flows starting at c^- . By passing to a subsequence we may assume the sequence λ_n has a fixed number of components, i.e.

$$\lambda_n = (\lambda_n^1, \dots, \lambda_n^{q+1}) \in \mathcal{L}_q(F_{ij}, c^-, \Pi(\Sigma))$$

for some q .

- If $q \geq 1$, we now apply Lemma 2.11 to λ_n^{q+1} and Lemma 2.5 to each λ_n^k , $1 \leq k \leq q$, individually.
- If $q = 0$, we now apply Lemma 2.11 to λ_n^1 .

\square

2.5. Singular flows. Let $\mathcal{M}(F_{ij}, \Pi(\Sigma))$ be the set of singular flows of F_{ij} .

If ϕ is in $\mathcal{M}(F_{ij}, \Pi(\Sigma))$, then the domain of ϕ is a compact interval of the form $[a, b]$. There is a natural \mathbb{R} -action on $\mathcal{M}(F_{ij}, \Pi(\Sigma))$, given by

$$(r, \phi) \mapsto r_*\phi(t) = \phi(t + r), \quad r \in \mathbb{R},$$

where the singular flow $r_*\phi$ is defined on the compact interval $[a + r, b + r]$. Given this \mathbb{R} -action, we may define the *set of unparameterized singular flows* as

$$\mathcal{L}(F_{ij}, \Pi(\Sigma)) = \mathcal{M}(F_{ij}, \Pi(\Sigma)) / \mathbb{R}.$$

Definition 2.13. Let c_1, \dots, c_q be critical points of F_{ij} . A q -times broken singular flow, denoted $\lambda = (\lambda_1, \dots, \lambda_{q+1})$, is a concatenation of paths λ_i such that

$$\lambda_1 \in \mathcal{L}(F_{ij}, \Pi(\Sigma), c_1), \quad \lambda_k \in \mathcal{L}(F_{ij}, c_{k-1}, c_k), \quad \text{and} \quad \lambda_{q+1} \in \mathcal{L}(F_{ij}, c_q, \Pi(\Sigma))$$

where $2 \leq k \leq q$.

It should be noted that a 0-times broken singular flow is simply an unparameterized singular flow.

We denote by $\overline{\mathcal{L}}(F_{ij}, \Pi(\Sigma))$ the *moduli space of broken singular flows*, which has a natural stratification

$$\overline{\mathcal{L}}(F_{ij}, \Pi(\Sigma)) = \bigcup_{q=0}^{\infty} \mathcal{L}_q(F_{ij}, \Pi(\Sigma)),$$

where $\mathcal{L}_q(F_{ij}, \Pi(\Sigma))$ is the *moduli space of q -times broken singular flows*. We will now construct a topology on $\overline{\mathcal{L}}(F_{ij}, \Pi(\Sigma))$.

Consider an element $\lambda = (\lambda_1, \dots, \lambda_q)$ in $\overline{\mathcal{L}}(F_{ij}, \Pi(\Sigma))$ and let $\Omega(c_k)$ be a Morse neighborhood of c_k , for $1 \leq k \leq q$. Consider a neighborhood U_{k-1}^- of the exit point of the image of λ_k from $\Omega(c_{k-1})$ and consider a neighborhood U_k^+ of the entry point of the image of λ_k into $\Omega(c_k)$, such that U_{k-1}^- and U_k^+ are both contained in their respective level sets, for $2 \leq k \leq q$.

Let c_0 be the point in $\Pi(\Sigma)$ that the image of λ_1 emanates from and c_{q+1} be the point in $\Pi(\Sigma)$ that the image of λ_{q+1} terminates at. Consider neighborhoods U_0^- and U_{q+1}^+ of c_0 and c_{q+1} respectively, contained in $\Pi(\Sigma)$. Finally, consider a neighborhood U_1^+ of the entry point of the image of λ_1 into $\Omega(c_1)$ contained in its level set and consider a neighborhood U_q^- of the exit point of the image of λ_{q+1} from $\Omega(c_q)$ contained in its level set. See Figure 5.




Figure 5: An arbitrary λ in $\overline{\mathcal{L}}(F, \Pi(\Sigma))$ with λ 's exit and entry neighborhoods.

Denote by U^- and U^+ the collection of the U_i^- 's and U_i^+ 's respectively. We define the set $\mathcal{W}(\lambda, U^-, U^+)$ to be the set of tuples $\mu = (\mu_1, \dots, \mu_r)$, for an arbitrary $r \leq q+1$, with the property that there exists integers

$$0 = k_0 < k_2 < \dots < k_{r-1} < k_r = q+1$$

such that

- μ_s is in $\mathcal{L}(F_{ij}, c_{k_{s-1}}, c_{k_s})$ for $2 \leq s \leq r-1$,
- μ_1 is in $\mathcal{L}(F_{ij}, \Pi(\Sigma), c_{i_2})$,
- μ_r is in $\mathcal{L}(F_{ij}, c_{i_{r-1}}, \Pi(\Sigma))$,
- the image of μ_s exits $\Omega(c_{k_{s-1}})$ through $U_{k_{s-1}}^-$ and enters $\Omega(c_{k_s})$ through $U_{k_s}^+$ for $2 \leq s \leq r-1$,
- the image of μ_1 emanates from U_0^- and enters $\Omega(c_{i_2})$ through $U_{i_2}^+$,
- and the image of μ_r exits $\Omega(c_{k_{r-1}})$ through $U_{k_{r-1}}^-$ and terminates in U_{q+1}^+ .

Then the collection of $\mathcal{W}(\lambda, U^-, U^+)$ is a neighborhood basis for a topology on $\overline{\mathcal{L}}(F_{ij}, \Pi(\Sigma))$. And since the topology was constructed from the topology of U and $\Pi(\Sigma)$, it follows that the constructed topology is first countable and Hausdorff.

Before proving $\overline{\mathcal{L}}(F_{ij}, \Pi(\Sigma))$ is a compactification of $\mathcal{L}(F_{ij}, \Pi(\Sigma))$, we will prove the following special case.

Lemma 2.14. *Let ϕ_n be a sequence of unparameterized singular flows of F_{ij} . Then there exists points c_0 and c_{q+1} in $\Pi(\Sigma)$, a finite set of critical points c_1, \dots, c_q of F_{ij} , for some q , and a broken singular flow $\lambda = (\lambda_1, \dots, \lambda_{q+1})$ connecting c_0 to c_1 , c_1 to c_2 , and so on, such that (after passing to a subsequence) ϕ_n converges to λ .*

Proof. We recall that the sequence ϕ_n is defined on the domain $[a_n, b_n]$. We have that $\Pi(\Sigma) \cap \overline{U}$ is compact, therefore, we may pass to a subsequence and assume $\phi_n(a_n)$ and $\phi_n(b_n)$ converge to a points c_0 and c_{q+1} in $\Pi(\Sigma) \cap \overline{U}$, respectively. Let λ_1 be the unique unparameterized flow of F_{ij} emanating from c_0 .

We now have two cases.

1. It may be the case that λ_1 is in $\mathcal{L}(F_{ij}, \Pi(\Sigma))$. Hence, λ_1 flows into the singular set and ends there. Let λ_1 be defined on the domain $[a, b]$. Then as in the proof of Lemma 2.11, it follows that $\lim_{t \rightarrow b} \lambda_1(t) = c_{q+1}$. Hence the sequence ϕ_n converges to $\lambda = (\lambda_1)$, and there is no breaking of the sequence of gradient flows.
2. It may also be the case that there exists a critical point c_1 of F_{ij} such that λ_1 is in $\mathcal{L}(F_{ij}, c_0, c_1)$. We then apply the argument of Lemma 2.5.

We may now induct over the number of critical points between c_0 and c_{q+1} using the argument of Lemma 2.5, since $\text{Crit}(F_{ij})$ is a finite set, to construct the limit $\lambda = (\lambda_1, \dots, \lambda_q)$ in $\overline{\mathcal{L}}(F_{ij}, \Pi(\Sigma))$ of the sequence ϕ_n . We note that in any step of the induction, if the constructed λ_k flows into $\Pi(\Sigma)$, then the induction ends. \square

With Lemma 2.14, we may now show that $\overline{\mathcal{L}}(F_{ij}, \Pi(\Sigma))$ compactifies $\mathcal{L}(F_{ij}, \Pi(\Sigma))$, in the sense that every sequence has a convergent subsequence.

Proposition 2.15. $\overline{\mathcal{L}}(F_{ij}, \Pi(\Sigma))$ is sequentially compact.

Proof. Let λ_n be a sequence of broken singular flows. By passing to a subsequence we may assume the sequence λ_n has a fixed number of components, i.e.

$$\lambda_n = (\lambda_n^1, \dots, \lambda_n^{q+1}) \in \mathcal{L}_q(F_{ij}, \Pi(\Sigma))$$

for some q .

- If $q \geq 2$, we now apply Lemma 2.8 to λ_n^1 , Lemma 2.5 to each λ_n^k , $2 \leq k \leq q$ individually, and Lemma 2.11 to λ_n^{q+1} .
- If $q = 1$, we now apply Lemma 2.8 to λ_n^1 and 2.11 to λ_n^2 .
- If $q = 0$, we now apply Lemma 2.14 to λ_n^1 .

□

3. Sequences of broken flow trees and compactification. In this Section we construct a compactification of the space of flow trees with at most N edges and vertices. As a special case, the space of flow trees with at most P positive punctures and at most formal dimension $D > 0$ is compactified. In Section 2, in order to compactify gradient flows of a local function difference, we had to consider broken gradient flows. Similarly, in this Section, we will consider broken flow trees. We begin by defining broken flow trees and investigating elementary properties, and we will end with the proof of Theorem 0.1

3.1. Broken flow trees. Before we begin our discussion of compactifying the moduli space of flow trees with bounded edges, we need to define a generalization of a flow tree.

Definition 3.1. A *broken flow tree* of L is a continuous map $\lambda : T \rightarrow M$, where T is a source tree, satisfying the following.

1. If E is an edge of T , then $\lambda : E \rightarrow M$ is a part of a broken gradient flow of L .
2. Let v be a k -valent vertex of T with cyclically ordered edges E_1, \dots, E_k . Let $\{\overline{\lambda}_j^1, \overline{\lambda}_j^2\}$ be the cotangent lift corresponding to E_j . We require that

$$\overline{\lambda}_j^2(v) = \overline{\lambda}_{j+1}^1(v) = \overline{p} \in \Pi_{\mathbb{C}}(L),$$

such that the flow orientation of $\overline{\lambda}_j^2$ at \overline{p} is directed toward \overline{p} if and only if the flow orientation of $\overline{\lambda}_{j+1}^1$ at \overline{p} is directed away from \overline{p} .

3. We require that the cotangent lifts of the edges of T yield a loop in $\Pi_{\mathbb{C}}(L)$.

In Definition 3.1, given a broken flow tree $\lambda : T \rightarrow M$, it is said in condition one that if E is an edge of T , then $\lambda : E \rightarrow M$ is a part of a broken gradient flow of L (associated to some local function difference F_{ij}). Implicitly, there are two subtleties to this wording. The first

is that it may be noted we have not used the word “parameterization”, this is because in our definition of a broken gradient flow of F_{ij} , the components are already unparameterized. The second is what it means to be a “part of a broken gradient flow”, and this means the following.

- If $q \geq 1$, then $\lambda = (\lambda_1, \dots, \lambda_{q+1})$, and there exists critical points c_1, \dots, c_q of F_{ij} such that

$$\lambda_k \in \mathcal{L}(F_{ij}, c_{k-1}, c_k), \quad 2 \leq k \leq q.$$

Also, the maximal extension Λ_1 of the unparameterized gradient flow λ_1 of F_{ij} is either a Morse flow or a fold emanating flow with image terminating at c_1 . Similarly, the maximal extension Λ_{q+1} of the unparameterized gradient flow λ_{q+1} of F_{ij} is either a Morse flow or a fold terminating flow with image emanating from c_q .

- If $q = 0$, then $\lambda = (\lambda_1)$, and the maximal extension Λ_1 of the unparameterized gradient flow λ_1 of F_{ij} is any of the four types given in Definition 1.2.

We further remark that if $\lambda : T \rightarrow M$ is a broken flow tree such that for any edge E of T , the map $\lambda : E \rightarrow M$ satisfies $\lambda = (\lambda_1)$, then T could also be called an *unparameterized flow tree*. Another way to reach this conclusion would be to take any flow tree T and consider the natural $\mathbb{R}^{\text{edge}(T)}$ -action on T given by reparametrization of each edge, where $\text{edge}(T)$ is the number of edges of T .

We denote by $\overline{\mathcal{L}}(N)$ the *moduli space of broken flow trees with at most N edges and vertices* and by $\mathcal{L}(N)$ the *moduli space of unparameterized flow trees with at most N edges and vertices*.

Definition 3.2. Let T and T' be two broken flow trees. We say that T and T' have the same *combinatorial type* Γ if the following holds.

- We require that T and T' are isomorphic as (combinatorial) trees.
- We require that for any pair of isomorphic edges E and E' of T and T' , respectively, the broken gradient flows

$$\lambda : E \rightarrow M, \quad \lambda' : E' \rightarrow M$$

are broken gradient flows of the same local function difference F_{ij} . We also require that the maximal extensions Λ and Λ' of λ and λ' , respectively, are elements of the same moduli space from Section 2 with respect to their common local function difference F_{ij} . Concretely, Λ and Λ' are either both broken: Morse flows (emanating and terminating at the same point), fold emanating flows (terminating at the same point), fold terminating flows (emanating from the same point), or singular flows.

Given the fact that there are only finitely many isomorphism classes of trees with a bounded number of edges and vertices, and the fact that there are only finitely many local function differences under consideration, Definition 3.2 gives a finite decomposition

$$\mathcal{L}(N) = \bigcup_{\Gamma} \mathcal{L}(N, \Gamma),$$

where $\mathcal{L}(N, \Gamma)$ is the *moduli space of broken flow trees with at most N edges and vertices of combinatorial type Γ* , and the union is over all combinatorial types Γ .

3.2. Stratum Floer-Gromov convergence. Consider a sequence T_n of broken flow trees of the same combinatorial type Γ with the following labelling. We label the k -th edge of T_n as $E_{n,k}$, the associated broken gradient flow $\lambda_{n,k}$, the maximal extension of $\lambda_{n,k}$ as $\Lambda_{n,k}$, and the adjacent vertices of $E_{n,k}$ as $v_{n,k}^1$ and $v_{n,k}^2$ (where $\lambda_{n,k}$ flows from $v_{n,k}^1$ to $v_{n,k}^2$).

Definition 3.3. Given a sequence T_n of broken flow trees of the same combinatorial type Γ with the above labelling, and a broken flow tree T , we say that T_n *stratum Floer-Gromov* converges to T if there exists an auxiliary broken flow tree T' such that the following holds.

- We require that T_n and T' are isomorphic as (combinatorial) trees.
- For every sequence of isomorphic edges $E_{n,k}$ and every corresponding isomorphic edge E'_k of T' of that sequence, the associated (parts of) broken gradient flows $\lambda_{n,k}$ and λ'_k are (parts of) broken gradient flows of the same local function difference F_{ij} determined by the combinatorial type Γ .
- For every edge E'_k of T' and associated maximally extended broken gradient flow Λ'_k , there exists (parts of) broken gradient flows Ψ_k^{in} and Ψ_k^{out} of the local function difference F_{ij} such that the maximally extended broken gradient flow $\Lambda_{n,k}$ of $E_{n,k}$ and the concatenation

$$\Psi_k = \Psi_k^{\text{in}} \# \Lambda'_k \# \Psi_k^{\text{out}}$$

are in the same moduli space of Section 2 with respect to F_{ij} . We require that $\Lambda_{n,k}$ converges to Ψ_k in this moduli space.

- For every sequence of vertices $v_{n,k}^1$ and $v_{n,k}^2$ associated to the sequence of isomorphic edges $E_{n,k}$ and every corresponding isomorphic vertices $v_k^{1'}$ and $v_k^{2'}$ of T' associated to E'_k , we have

$$\lim_{n \rightarrow \infty} \lambda_{n,k}(v_{n,k}^1) = \lambda'_k(v_k^{1'}) \quad \text{and} \quad \lim_{n \rightarrow \infty} \lambda_{n,k}(v_{n,k}^2) = \lambda'_k(v_k^{2'}).$$

- The broken flow tree $\lambda : T \rightarrow M$ is obtained from the broken flow tree $\lambda' : T' \rightarrow M$ through collapsing of degenerate edges and the natural restriction of the map λ' . Concretely, if E'_k is an edge of T' with adjacent vertices $v_k^{1'}$ and $v_k^{2'}$ and associated broken gradient flow λ'_k that satisfies

$$\lambda'_k(v_k^{1'}) = \lambda'_k(v_k^{2'}),$$

then we delete E'_k from T' and glue together the resulting pieces in the natural way. After collapsing all degenerate edges to obtain T , the resulting map $\lambda : T \rightarrow M$ is simply restriction of λ' .

We note that the Ψ_k^{in} and Ψ_k^{out} in Definition 3.3 may simply be the endpoints of Λ'_k if $\Lambda_{k,n}$ and Λ'_k are already in the same moduli space. In particular, this is true for every edge of T' if T' is in the same combinatorial type Γ as the sequence T_n .

Before the Definition of combinatorial type there was no way to talk about the convergence of a sequence of broken flow trees. Even with an edge bound, the broken flow trees may be unrelated since the local function differences and the corresponding broken gradient flows

are unrelated. However, with this Definition we now have a notion of what it means for certain types of sequences of broken flow trees to converge. And fortunately, we have the following Proposition.

Proposition 3.4. *A sequence of broken flow trees T_n of the same combinatorial type Γ has a stratum Floer-Gromov converging subsequence.*

Proof. Let T' be a tree that is isomorphic to T_n , for all n . It suffices to show that a sequence of isomorphic edges $E_{n,k}$ converges to the corresponding isomorphic edge E'_k of T' . Since the maximal extensions $\Lambda_{n,k}$ of the associated broken gradient flows $\lambda_{n,k} : E_{n,k} \rightarrow M$ (with respect to a common local function difference F_{ij}) must be in the same moduli space (from Section 2) by assumption, by passing to a subsequence we may assume there exists a broken gradient flow Ψ_k of F_{ij} in the appropriate moduli space such that

$$\lim_{n \rightarrow \infty} \Lambda_{n,k} = \Psi_k.$$

It follows that the adjacent vertices $v_{n,k}^1$ and $v_{n,k}^2$ of $E_{n,k}$ satisfy

$$\lim_{n \rightarrow \infty} \lambda_{n,k}(v_{n,k}^1) = p_1, \quad \lim_{n \rightarrow \infty} \lambda_{n,k}(v_{n,k}^2) = p_2,$$

where p_1 and p_2 are in the image of Ψ_k . We now define λ'_k to be the restriction of Ψ_k that connects p_1 to p_2 , and assign λ'_k to the edge E'_k , in particular we have the map

$$\lambda'_k : E'_k \rightarrow M.$$

Continuing in this way, we may build the entire auxiliary broken flow tree $\lambda' : T' \rightarrow M$, and we obtain the limit broken flow tree T of the sequence T_n by collapsing degenerate edges. \square

We denote by $\overline{\mathcal{L}}(N, \Gamma)$ the set $\mathcal{L}(N, \Gamma)$ unioned with the set of all formally defined stratum Floer-Gromov limits of all sequences in $\mathcal{L}(N, \Gamma)$ given by Definition 3.3. In the next part, we shall give the set $\overline{\mathcal{L}}(N, \Gamma)$ a topology.

3.3. The stratum Floer-Gromov topology. We must give the set $\overline{\mathcal{L}}(N, \Gamma)$ a topology. We will do this by using convergence structures, where the details of the following general statements about convergence structures may be found in Chapter 5 of [MS12].

Definition 3.5. Let X be a set. A *convergence structure* on X is a subset

$$\mathcal{C} \subset X \times X^{\mathbb{N}}$$

that satisfies.

- *Constant:* If $x_n = x_0$ for all $n \in \mathbb{N}$ then $(x_0, (x_n)_n) \in \mathcal{C}$.
- *Subsequence:* If $(x_0, (x_n)_n) \in \mathcal{C}$ and $g : \mathbb{N} \rightarrow \mathbb{N}$ is strictly increasing, then $(x_0, (x_{g(n)})_n) \in \mathcal{C}$.
- *Subsubsequence:* If for every strictly increasing function $g : \mathbb{N} \rightarrow \mathbb{N}$, there exists a strictly increasing function $f : \mathbb{N} \rightarrow \mathbb{N}$ such that $(x_0, (x_{(g \circ f)(n)})_n) \in \mathcal{C}$, then $(x_0, (x_n)_n) \in \mathcal{C}$.

- *Diagonal*: If $(x_0, (x_k)_k) \in \mathcal{C}$ and $(x_k, (x_{k,n})_n) \in \mathcal{C}$ for every k , then there exist sequences $k_r, n_r \in \mathbb{N}$ such that $(x_0, (x_{k_r, n_r})_r) \in \mathcal{C}$.
- *Uniqueness*: If $(x_0, (x_n)_n), (y_0, (x_n)_n) \in \mathcal{C}$ then $x_0 = y_0$.

We say that (X, \mathcal{C}) is a *convergence space*.

We note that, if we consider a first countable and Hausdorff topological space Y , then it may be checked that the set of convergent sequences \mathcal{C} of Y satisfies the axioms of Definition 3.5.

Intuitively, one should think of an element $(x_0, (x_n)_n)$ in \mathcal{C} , for a convergence space (X, \mathcal{C}) , as a sequence x_n converging to x_0 . However, this intuition is only valid if X has a topology, which we now construct.

Definition 3.6. Let (X, \mathcal{C}) be a convergence space. We say $U \subset X$ is open if and only if for every $(x_0, (x_n)_n)$ in $\mathcal{C} \cap (U \times X^{\mathbb{N}})$ there exists N such that x_n is in U for every $n \geq N$.

Lemma 3.7. Define the set

$$\mathcal{U}(\mathcal{C}) = \{U \subset X : U \text{ open as in Definition 3.6}\}.$$

Then $\mathcal{U}(\mathcal{C})$ is a topology on X with the property that $(x_0, (x_n)_n) \in \mathcal{C}$ if and only if the sequence x_n converges to x_0 with respect to $\mathcal{U}(\mathcal{C})$. Moreover, this topology is unique with respect to this property.

For the moduli space $\overline{\mathcal{L}}(N, \Gamma)$ we consider the subset

$$\Gamma_{FG} \subset \overline{\mathcal{L}}(N, \Gamma) \times \mathcal{L}(N, \Gamma)^{\mathbb{N}} \subset \overline{\mathcal{L}}(N, \Gamma) \times \overline{\mathcal{L}}(N, \Gamma)^{\mathbb{N}}$$

defined by

$$\Gamma_{FG} = \{(T, (T_n)_n) : T_n \in \mathcal{L}(N, \Gamma) \text{ stratum Floer-Gromov converges to } T \in \overline{\mathcal{L}}(N, \Gamma)\}.$$

Proposition 3.8. The subset Γ_{FG} defines a convergence structure on $\overline{\mathcal{L}}(N, \Gamma)$.

Proof. The *constant*, *subsequence*, *subsubsequence*, and *uniqueness* axioms follow immediately, the only difficulty is the *diagonal* axiom.

So suppose $(T, (T_k)_k)$ is in Γ_{FG} and $(T_k, (T_{k,n})_n)$ is in Γ_{FG} for all k . We will let T' denote the broken flow tree of combinatorial type Γ guaranteed by Definition 3.3 of the stratum Floer-Gromov convergence

$$T_k \rightarrow T.$$

Note we do not require auxiliary broken flow trees T'_k since T_k and $T_{k,n}$ are of combinatorial type Γ for all k and n .

We label the s -th edge of T_k as $E_{k,s}$, the associated broken gradient flow as $\lambda_{k,s}$, the maximal extension of $\lambda_{k,s}$ as $\Lambda_{k,s}$, and the adjacent vertices of $E_{k,s}$ as $v_{k,s}^1$ and $v_{k,s}^2$ (where $\lambda_{k,s}$ flows from $v_{k,s}^1$ to $v_{k,s}^2$).

Similarly, we label the s -th edge of T' as E'_s , the associated broken gradient flow as λ'_s , the maximal extension of λ'_s as Λ'_s , the adjacent vertices of E'_s as $v_s^{1'}$ and $v_s^{2'}$ (where λ'_s flows from $v_s^{1'}$ to $v_s^{2'}$), and Ψ_s the concatenated broken flow guaranteed by Definition 3.3.

And finally, we label the s -th edge of $T_{k,n}$ as $E_{k,s}^n$, the associated broken gradient flow as $\lambda_{k,s}^n$, the maximal extension of $\lambda_{k,s}^n$ as $\Lambda_{k,s}^n$, and the adjacent vertices of $E_{k,s}^n$ as $v_{k,s}^{1,n}$ and $v_{k,s}^{2,n}$ (where $\lambda_{k,s}^n$ flows from $v_{k,s}^{1,n}$ to $v_{k,s}^{2,n}$).

The assumptions result in the diagrams

$$\begin{array}{ccccccc}
T_{1,n} \rightarrow T_1 & \Lambda_{1,s}^n \rightarrow \Lambda_{1,s} & \lambda_{1,s}^n(v_{1,s}^{1,n}) \rightarrow \lambda_{1,s}(v_{1,s}^1) & \lambda_{1,s}^n(v_{1,s}^{2,n}) \rightarrow \lambda_{1,s}(v_{1,s}^2) \\
T_{2,n} \rightarrow T_2 & \Lambda_{2,s}^n \rightarrow \Lambda_{2,s} & \lambda_{2,s}^n(v_{2,s}^{1,n}) \rightarrow \lambda_{2,s}(v_{2,s}^1) & \lambda_{2,s}^n(v_{2,s}^{2,n}) \rightarrow \lambda_{2,s}(v_{2,s}^2) \\
\vdots \longrightarrow \vdots & \vdots \longrightarrow \vdots & \vdots \longrightarrow \vdots & \vdots \longrightarrow \vdots \\
T_{k,n} \rightarrow T_k & \Lambda_{k,s}^n \rightarrow \Lambda_{k,s} & \lambda_{k,s}^n(v_{k,s}^{1,n}) \rightarrow \lambda_{k,s}(v_{k,s}^1) & \lambda_{k,s}^n(v_{k,s}^{2,n}) \rightarrow \lambda_{k,s}(v_{k,s}^2) \\
& \downarrow & \downarrow & \downarrow \\
& T & \Psi_s & \lambda'_s(v_s^{1'}) \\
& & & \lambda'_s(v_s^{2'})
\end{array}$$

where going from left to right in a diagram corresponds to letting n go to infinity, and going top to bottom in a diagram corresponds to letting k go to infinity.

Let U_s^Ψ be an arbitrary neighborhood of Ψ_s in the appropriate moduli space from Section 2, then there exists a number K_s^Ψ such that $\Lambda_{k,s}$ is in U_s^Ψ for $k \geq K_s^\Psi$ by the convergence of $\Lambda_{k,s} \rightarrow \Psi_s$. We have that U_s^Ψ is a neighborhood of $\Lambda_{k,s}$ for $k \geq K_s^\Psi$, thus there exists a number $N_{k,s}^\Lambda$ such that $\Lambda_{k,s}^n$ is in U_s^Ψ for $n \geq N_{k,s}^\Lambda$ and $k \geq K_s^\Psi$ by the convergence of $\Lambda_{k,s}^n \rightarrow \Lambda_{k,s}$. Since the number of edges s is finite, we may define

$$K^\Psi = \max_s K_s^\Psi \quad \text{and} \quad N_k^\Lambda = \max_s N_{k,s}^\Lambda.$$

Let $U_s^{(v^1)}$ be an open neighborhood of $\lambda'_s(v_s^{1'})$ in the base manifold M , then there exists a number $K_s^{(v^1)}$ such that $\lambda_{k,s}(v_{k,s}^1)$ is in $U_s^{(v^1)}$ for $k \geq K_s^{(v^1)}$ by the convergence of $\lambda_{k,s}(v_{k,s}^1) \rightarrow \lambda'_s(v_s^{1'})$. We have that $U_s^{(v^1)}$ is a neighborhood of $\lambda_{k,s}(v_{k,s}^1)$ for $k \geq K_s^{(v^1)}$, thus there exists a number $N_{k,s}^{(v^1)}$ such that $\lambda_{k,s}^n(v_{k,s}^{1,n})$ is in $U_s^{(v^1)}$ for $n \geq N_{k,s}^{(v^1)}$ and $k \geq K_s^{(v^1)}$ by the convergence of $\lambda_{k,s}^n(v_{k,s}^{1,n}) \rightarrow \lambda_{k,s}(v_{k,s}^1)$. Since the number of edges s is finite, we may define

$$K^{(v^1)} = \max_s K_s^{(v^1)} \quad \text{and} \quad N_k^{(v^1)} = \max_s N_{k,s}^{(v^1)}.$$

Similarly, for the fourth diagram, we may construct the numbers $K^{(v^2)}$ and $N_k^{(v^2)}$. We finally define the numbers

$$K = \max \left\{ K^\Psi, K^{(v^1)}, K^{(v^2)} \right\} \quad \text{and} \quad N_k = \max \left\{ N_k^\Lambda, N_k^{(v^1)}, N_k^{(v^2)} \right\}.$$

We note that the numbers K and N_k are dependent on the neighborhoods chosen. We now choose any broken flow tree T_{n_0, k_0} such that

$$k_0 \geq K \text{ and } n_0 \geq N_{k_0}.$$

It now follows, by considering smaller neighborhoods contained in U_s^Ψ , $U_s^{(v^1)}$, and $U_s^{(v^2)}$ (for all edges s), we may construct subsequences of numbers k_r, n_r such that T_{k_r, n_r} stratum Floer-Gromov converges to T , hence $(T, (T_{k_r, n_r})_r)$ is in Γ_{FG} . \square

Proposition 3.8 allows us to define the *stratum Floer-Gromov topology* $\mathcal{U}(\Gamma_{FG})$ on $\overline{\mathcal{L}}(N, \Gamma)$. In Section 3.4, we will “glue” together the stratum Floer-Gromov topologies in order to put a topology on $\overline{\mathcal{L}}(N)$.

3.4. The Floer-Gromov topology. In Section 3.2 we defined the set $\overline{\mathcal{L}}(N, \Gamma)$ for a combinatorial type Γ and in Proposition 3.8 we topologized $\overline{\mathcal{L}}(N, \Gamma)$ with the stratum Floer-Gromov topology. Given the inclusions

$$\mathcal{L}(N, \Gamma) \subset \overline{\mathcal{L}}(N, \Gamma) \subset \overline{\mathcal{L}}(N)$$

and the finite decomposition

$$\overline{\mathcal{L}}(N) = \bigcup_{\Gamma} \overline{\mathcal{L}}(N, \Gamma),$$

we may write $\overline{\mathcal{L}}(N)$ as the finite non-disjoint union

$$\overline{\mathcal{L}}(N) = \bigcup_{\Gamma} \overline{\mathcal{L}}(N, \Gamma).$$

Definition 3.9. We have the set of inclusions

$$\mathcal{I} = \{i_\Gamma : \overline{\mathcal{L}}(N, \Gamma) \rightarrow \overline{\mathcal{L}}(N)\}.$$

We define the *Floer-Gromov topology* on $\overline{\mathcal{L}}(N)$ to be the final topology with respect to \mathcal{I} .

Definition 3.10. Let T_n be a sequence of broken flow trees in $\overline{\mathcal{L}}(N)$. We say that T_n *Floer-Gromov converges* to a broken flow tree T in $\overline{\mathcal{L}}(N)$ if T_n converges to T with respect to the Floer-Gromov topology.

In Definition 3.9, we have that the final topology on $\overline{\mathcal{L}}(N)$ with respect to \mathcal{I} is characterized by the following. A set $U \subset \overline{\mathcal{L}}(N)$ is open if and only if $U \cap \overline{\mathcal{L}}(N, \Gamma)$ is open for every combinatorial type Γ . Because of this characterization, we note that Floer-Gromov convergence includes stratum Floer-Gromov convergence as a special case.

Lemma 3.11. *Let T_n in $\mathcal{L}(N, \Gamma)$ be a sequence that stratum Floer-Gromov converges to T in $\overline{\mathcal{L}}(N, \Gamma)$, then T_n Floer-Gromov converges to T in $\overline{\mathcal{L}}(N)$.*

Proof. Let $U \subset \overline{\mathcal{L}}(N)$ be a neighborhood of T , then $U \cap \overline{\mathcal{L}}(N, \Gamma)$ is a neighborhood of T in $\overline{\mathcal{L}}(N, \Gamma)$ by Definition 3.9. If we refer to Definition 3.6, we have that there exists an $N \geq 0$ such that T_n is in $U \cap \overline{\mathcal{L}}(N, \Gamma)$ for $n \geq N$. It follows that T_n is in U for $n \geq N$, hence, T_n Floer-Gromov converges to T . \square

The main result of this paper now follows.

Proof of Theorem 0.1. Let T_n in $\overline{\mathcal{L}}(N)$ denote an arbitrary sequence of broken flow trees. Due to the finite decomposition

$$\overline{\mathcal{L}}(N) = \bigcup_{\Gamma} \mathcal{L}(N, \Gamma),$$

we may pass to a subsequence and assume T_n is in $\mathcal{L}(N, \Gamma)$, for some combinatorial type Γ . By Proposition 3.4, we may pass to another subsequence and assume there exists a T in $\overline{\mathcal{L}}(N, \Gamma)$ such that T_n stratum Floer-Gromov converges to T . Finally, by Lemma 3.11, we have that T_n Floer-Gromov converges to T in $\overline{\mathcal{L}}(N)$, as desired. \square

There is a natural inclusion

$$\mathcal{L}(N) \rightarrow \overline{\mathcal{L}}(N),$$

where we recall $\mathcal{L}(N)$ is the moduli space of unparameterized flow trees with at most N edges and vertices. By endowing $\mathcal{L}(N)$ with the induced topology of the Floer-Gromov topology, we may consider $\overline{\mathcal{L}}(N)$ a compactification of $\mathcal{L}(N)$. However, a priori, we have no reason to believe a sequence arbitrarily has an edge bound. If we consider the *moduli space of unparameterized flow trees with a bounded number of positive punctures P and at most formal dimension $D > 0$* , denoted $\mathcal{L}(N, P, D)$, then by Lemma 1.7, there exists a natural inclusion

$$\mathcal{L}(N, P, D) \rightarrow \overline{\mathcal{L}}(N),$$

for some N . It now follows that by giving $\mathcal{L}(N, P, D)$ the induced topology of the Floer-Gromov topology, we may consider $\overline{\mathcal{L}}(N)$ a compactification of $\mathcal{L}(N, P, D)$.

References.

- [Abb14] Casim Abbas, *An introduction to compactness results in symplectic field theory*, Springer-Verlag, 2014.
- [AD14] Michèle Audin and Mihai Damian, *Morse theory and Floer homology*, Springer-Verlag, 2014.
- [AG18] Daniel Álvarez-Gavela, *The simplification of singularities of Lagrangian and Legendrian fronts*, *Inventiones mathematicae* **214** (2018), 641–737.
- [BEH⁺03] Frederic Bourgeois, Yakov Eliashberg, Helmut Hofer, Kris Wysocki, and Eduard Zehnder, *Compactness results in symplectic field theory*, *Geometry and Topology* **7** (2003), 799–888.
- [EES07] Tobias Ekholm, John Entyre, and Michael Sullivan, *Legendrian contact homology in $P \times \mathbb{R}$* , *Transactions of the American Mathematical Society* **359** (2007), no. 7, 3301–3335.
- [EGH00] Yakov Eliashberg, Alexander Givental, and Helmut Hofer, *Introduction to symplectic field theory*, *Geometric and Functional Analysis* (2000), 560–673.

- [Ekh07] Tobias Ekholm, *Morse flow trees and Legendrian contact homology in 1-jet spaces*, *Geometry and Topology* **11** (2007), 1083–1224.
- [EN18] Tobias Ekholm and Lenhard Ng, *Higher genus knot contact homology and recursion for colored HOMFLY-PT polynomials*, arXiv:1803.04011, March 2018.
- [Flo89] Andreas Floer, *Witten’s complex and infinite dimensional Morse theory*, *Journal of Differential Geometry* **30** (1989), 207–221.
- [FN20] Urs Frauenfelder and Robert Nicholls, *The moduli space of gradient flow lines and Morse homology*, arXiv:2005.10799, May 2020.
- [FO97] Kenji Fukaya and Yong-Geun Oh, *Zero-loop open strings in the cotangent bundle and Morse homotopy*, *Asian Journal of Mathematics* **1** (1997), no. 1, 96–180.
- [Kar17] Cecilia Karlsson, *To compute orientations of Morse flow trees in Legendrian contact homology*, arXiv:1704.05156, April 2017.
- [MS95] Dusa McDuff and Dietmar Salamon, *Introduction to symplectic topology*, Oxford University Press, 1995.
- [MS12] ———, *J-holomorphic curves and symplectic topology*, American Mathematical Society, 2012.
- [RG19] Georgios Dimitroglou Rizell and Roman Golovko, *Legendrian submanifolds from Bohr-Sommerfeld covers of monotone Lagrangian tori*, arXiv:1901.08415, January 2019.
- [Riz11] Georgios Dimitroglou Rizell, *Knotted Legendrian surfaces with few Reeb chords*, *Algebraic and Geometric Topology* **11** (2011), no. 5, 2903–2936.

THE COLLEGE OF BROWN UNIVERSITY, PROVIDENCE, RI 02912
E-mail address: kenneth_blakey@brown.edu