

LINEAR FAMILIES OF SMOOTH HYPERSURFACES OVER FINITE FIELDS

SHAMIL ASGARLI, DRAGOS GHIOCA, AND ZINOVY REICHSTEIN

ABSTRACT. We construct an n -dimensional linear system \mathcal{L} of hypersurfaces of degree d in \mathbb{P}^n defined over \mathbb{F}_q such that each \mathbb{F}_q -member of \mathcal{L} is smooth, under the hypothesis that the characteristic p does not divide $\gcd(d, n+1)$. Moreover, we exhibit a counterexample when p divides $\gcd(d, n+1)$.

1. INTRODUCTION

The study of hypersurfaces varying in a pencil, or more generally, in a linear system of arbitrary dimension, is an active research area. For instance, determining the number of reducible members in a pencil is already a challenging problem [Ste89], [Vis93], [PY08]. When the base field is a number field, the study of pencils has deep connections to Diophantine geometry; see, for example [DGH21]. We will be interested in the case when the base field is a finite field. In this direction, concrete questions about linear families parameterizing hypersurfaces over finite fields have been studied by Ballico in [Bal07], [Bal09]. There is also a version of Bertini's theorem over finite fields applicable to a pencil of hypersurfaces [AG22].

Let \mathbb{F}_q be a finite field with $q = p^s$ for some prime p . Our primary goal in the present paper is to address the following question on the linear families from a recent paper [AG22b] by the first two authors:

Question 1.1. Let (n, d, r) be any triple of positive integers with $n \geq 2$ and $d \geq 2$. Can we find $r+1$ homogeneous polynomials $F_0, F_1, \dots, F_r \in \mathbb{F}_q[x_0, \dots, x_n]$ of degree d such that the hypersurface

$$X_{[a_0:a_1:\dots:a_r]} = \{a_0F_0 + a_1F_1 + \dots + a_rF_r = 0\} \subset \mathbb{P}^n$$

is smooth for every $[a_0 : a_1 : \dots : a_r] \in \mathbb{P}^r(\mathbb{F}_q)$?

The question can be rephrased in geometric terms as follows. Recall that $\mathcal{L} = \langle F_0, \dots, F_r \rangle$ is called a *linear system* of (projective) dimension r . Thus, Question 1.1 asks for existence of a linear system \mathcal{L} of prescribed dimension over \mathbb{F}_q such that every \mathbb{F}_q -member of \mathcal{L} is a smooth hypersurface of degree d in \mathbb{P}^n .

We show that, under a mild assumption on the characteristic, the maximum value of r for which Question 1.1 has a positive answer is $r = n$.

Theorem 1.2. *Let \mathbb{F}_q be a finite field with $q = p^s$ for some prime p and integer s . The following conclusions hold:*

- (1) *If $r \geq n+1$, then Question 1.1 has a negative answer.*
- (2) *If $r = n$ and $p \nmid \gcd(d, n+1)$, then Question 1.1 has a positive answer.*

2020 *Mathematics Subject Classification.* Primary 14N05; Secondary 14J70, 14G15.
Key words and phrases. linear system, hypersurface, finite fields, smoothness.

Note that (1) holds without any assumption on the characteristic. We will show in Section 3 that the characteristic assumption $p \nmid \gcd(d, p+1)$ *cannot* be removed in general. More precisely, Theorem 3.1 will present a non-existence result involving $q = 2^s$, $d = 2$ and any odd integer n .

It is clear that if Question 1.1 has a positive answer for (n, d, r) over \mathbb{F}_q , then it also has a positive answer for $(n, d, r-1)$ over \mathbb{F}_q . In particular, if $p \nmid \gcd(d, n+1)$, then Question 1.1 has a positive answer for the case $r = 1$, which corresponds to a pencil of hypersurfaces. The hypothesis $p \nmid \gcd(d, n+1)$ is equivalent to asking $p \nmid d$ or $p \nmid (n+1)$. In particular, the latter condition means that for each dimension n , the desired result holds for all but finitely many characteristics. Let us call a pencil of hypersurfaces \mathcal{L} a *smooth pencil* if each \mathbb{F}_q -member of \mathcal{L} is smooth. As an illustration, we have that for every value of d :

- There exists a smooth pencil of degree d in \mathbb{P}^2 if $\text{char}(\mathbb{F}_q) \neq 3$.
- There exists a smooth pencil of degree d in \mathbb{P}^3 if $\text{char}(\mathbb{F}_q) \neq 2$.
- There exists a smooth pencil of degree d in \mathbb{P}^4 if $\text{char}(\mathbb{F}_q) \neq 5$.
- There exists a smooth pencil of degree d in \mathbb{P}^5 if $\text{char}(\mathbb{F}_q) \neq 2, 3$.

As a comparison, the main result [AG22b, Theorem 1.3] proved the existence of a pencil \mathcal{L} of degree d hypersurfaces in \mathbb{P}^n where each \mathbb{F}_q -member of \mathcal{L} is smooth under a different hypothesis:

$$q > \left(\frac{1 + \sqrt{2}}{2} \right)^2 ((n+1)(d-1)^n)^2 ((n+1)(d-1)^n - 1)^2 ((n+1)(d-1)^n - 2)^2.$$

Note that the result is true for any characteristic, but q needs to be sufficiently large. It is reasonable to believe that smooth pencils exist over every finite field, even though the stronger result Theorem 1.2 (2) does not hold when the restriction $p \nmid \gcd(d, n+1)$ on the characteristic is removed; see Section 3.

Acknowledgements. The first author is supported by a postdoctoral research fellowship from the University of British Columbia and the NSERC PDF award. The second and third authors are supported by NSERC Discovery grants.

2. PROOF OF THE MAIN RESULT

In this section, we give a proof of Theorem 1.2.

As a preparation, let us recall a few facts from algebraic geometry. It is a well-known fact that the space of all degree d hypersurfaces in \mathbb{P}^n naturally forms a projective space \mathbb{P}^N where $N = \binom{n+d}{d} - 1$. When we write $F \in \mathbb{P}^N$, we will be referring to the hypersurface X defined by the polynomial F . Inside \mathbb{P}^N , there is a special hypersurface $\mathcal{D} \subset \mathbb{P}^N$ called the *discriminant*, which parameterizes all singular hypersurfaces of degree d .

We begin with a simple lemma on existence of linear subspaces contained in the discriminant.

Lemma 2.1. *Let $P \in \mathbb{P}^N(\mathbb{F}_q)$ be a point. Consider the locus*

$$H_P = \{F \in \mathbb{P}^N \mid X = \{F = 0\} \text{ is singular at the point } P\}$$

Then H_P is a codimension $n+1$ linear subspace defined over \mathbb{F}_q inside \mathbb{P}^N .

Proof. Without loss of generality, we can let $P = [1 : 0 : \dots : 0]$. The condition that $X = \{F = 0\}$ is singular at P is equivalent to $F(P) = \frac{\partial F}{\partial x_1}(P) = \dots = \frac{\partial F}{\partial x_n}(P) = 0$.

Each of these $n + 1$ relations impose a linear condition on the defining coefficients of the polynomial F . It is straightforward to check that these conditions are linearly independent. Thus, H_P has expected codimension $n + 1$ in \mathbb{P}^N . It is also evident that H_P is a subspace defined over the ground field \mathbb{F}_q . \square

We are now ready to establish the first half of our result.

Proof of Theorem 1.2, part (1): Consider a linear system \mathcal{L} of dimension $r \geq n + 1$ parametrizing degree d hypersurfaces in \mathbb{P}^n . Let P be an arbitrary \mathbb{F}_q -point in \mathbb{P}^n . Since $\dim(\mathcal{L}) \geq n + 1$ and $\text{codim}(H_P) = n + 1$, these two linear spaces must meet non-trivially in \mathbb{P}^N . Therefore, the intersection $\mathcal{L} \cap H_P$ must contain an \mathbb{F}_q -point, which corresponds to an \mathbb{F}_q -member of \mathcal{L} which is singular at P . Since \mathcal{L} was arbitrary, this illustrates that Question 1.1 has a negative answer when $r \geq n + 1$. \square

Before we prove the second part, we start with two lemmas on the existence of smooth hypersurfaces of a particular shape.

Lemma 2.2. *Suppose $p \nmid d$. Consider the hypersurface $X = \{c_0x_0^d + c_1x_1^d + \dots + c_nx_n^d = 0\}$ where c_i are arbitrary elements of $\overline{\mathbb{F}_q}$. If $\prod_i c_i \neq 0$, then X is smooth.*

Proof. Let $F = c_0x_0^d + c_1x_1^d + \dots + c_nx_n^d$. We have $\frac{\partial F}{\partial x_i} = dc_ix_i^{d-1}$ for each $0 \leq i \leq n$. Let $P = [u_0 : u_1 : \dots : u_n] \in X = \{F = 0\}$ be an arbitrary point. Then $u_j \neq 0$ for some j , and together with $c_j \neq 0$, we see that $\frac{\partial F}{\partial x_j}(P) = dc_ju_j^{d-1} \neq 0$. Thus, X is smooth at P . \square

Lemma 2.3. *Suppose $p \mid d$ but $p \nmid (n + 1)$. Consider the hypersurface $Y = \{c_0x_0^{d-1}x_1 + c_1x_1^{d-1}x_2 + \dots + c_nx_n^{d-1}x_0 = 0\}$ where c_i are arbitrary elements of $\overline{\mathbb{F}_q}$. If $\prod_i c_i \neq 0$, then Y is smooth.*

Proof. We will divide the proof into two cases, according to whether $d = 2$ or $d \neq 2$.

Case 1. $d \neq 2$.

Let $F = c_0x_0^{d-1}x_1 + c_1x_1^{d-1}x_2 + \dots + c_nx_n^{d-1}x_0$. Assume, to the contrary, that $Y = \{F = 0\}$ is singular at a point $P = [u_0 : u_1 : \dots : u_n] \in \mathbb{P}^n$; without loss of generality, we assume $u_1 \neq 0$. Using the hypothesis $p \mid d$, we obtain:

$$(2.1) \quad \frac{\partial F}{\partial x_i}(P) = c_{i-1}u_{i-1}^{d-1} - c_iu_i^{d-2}u_{i+1} = 0$$

for each $0 \leq i \leq n$, where the indices are taken modulo $n + 1$. Multiplying the relation (2.1) with u_i , we get:

$$(2.2) \quad c_{i-1}u_{i-1}^{d-1}u_i = c_iu_i^{d-1}u_{i+1}$$

Combining $F(P) = 0$ with (2.2) for each $0 \leq i \leq n$, we see that

$$0 = F(P) = \sum_{i=0}^n c_iu_i^{d-1}u_{i+1} = (n + 1)c_0u_0^{d-1}u_1$$

Since $p \nmid (n + 1)$ and $c_0 \neq 0$, we get $u_0^{d-1}u_1 = 0$. Since we know that $u_1 \neq 0$, we obtain that $u_0 = 0$. Applying (2.1) to $\frac{\partial F}{\partial x_0}(P) = 0$, we get $u_n = 0$. This is the step where we use the assumption that $d \neq 2$. Similarly, applying (2.1) to $\frac{\partial F}{\partial x_n}(P) = 0$, we get $u_{n-1} = 0$. Continuing in this way inductively, we obtain $u_0 = u_1 = \dots = u_n = 0$, a contradiction.

Case 2. $d = 2$.

We follow the same reasoning as in Case 1 to get $u_0^{d-1}u_1 = 0$. Again, we get that $u_0 = 0$ (since we assumed that $u_1 \neq 0$). Applying (2.1) to $\frac{\partial F}{\partial x_0}(P) = 0$, we get $u_{n-1} = 0$. Similarly, applying (2.1) to $\frac{\partial F}{\partial x_{n-1}}(P) = 0$, we get $u_{n-3} = 0$. Continuing in this way, we obtain $u_0 = u_{n-1} = u_{n-3} = \dots = 0$. Since $d = 2$, the hypothesis implies that $n = 2k$ is even. Thus, after k steps, we get $u_{n-(2k-1)} = u_1 = 0$, a contradiction. \square

We are now ready to establish the remaining half of our result:

Proof of Theorem 1.2, part (2): Let us first suppose $p \nmid d$. We will explicitly construct a linear system \mathcal{L} of dimension $r = n$ with the desired property. By the normal basis theorem, we can find an element $\alpha \in \mathbb{F}_{q^{n+1}}$ such that $\alpha, \alpha^q, \alpha^{q^2}, \dots, \alpha^{q^n}$ form an \mathbb{F}_q -basis for the $(n+1)$ -dimensional vector space $\mathbb{F}_{q^{n+1}}$. Let

$$\begin{aligned} F_0 &= (\alpha x_0 + \alpha^q x_1 + \alpha^{q^2} x_2 + \dots + \alpha^{q^i} x_i + \dots + \alpha^{q^n} x_n)^d \\ F_1 &= (\alpha^q x_0 + \alpha^{q^2} x_1 + \alpha^{q^3} x_2 + \dots + \alpha^{q^{i+1}} x_i + \dots + \alpha x_n)^d \\ F_2 &= (\alpha^{q^2} x_0 + \alpha^{q^3} x_1 + \alpha^{q^4} x_2 + \dots + \alpha^{q^{i+2}} x_i + \dots + \alpha^q x_n)^d \\ &\vdots \\ F_n &= (\alpha^{q^n} x_0 + \alpha^q x_1 + \alpha^{q^2} x_2 + \dots + \alpha^{q^{i+n}} x_i + \dots + \alpha^{q^{n-1}} x_n)^d \end{aligned}$$

Note that each F_i is not defined over \mathbb{F}_q . However, the set $\{F_0, F_1, \dots, F_n\}$ is invariant under the action of the q -th power Frobenius map. Thus, the linear system $\mathcal{L} = \langle F_0, \dots, F_n \rangle$ is defined over \mathbb{F}_q , that is, one can find a set of new generators G_0, G_1, \dots, G_n for \mathcal{L} where each G_i is defined over \mathbb{F}_q . Furthermore, we claim that F_0, F_1, \dots, F_n are linearly independent. To see this, let

$$(2.3) \quad y_j = \alpha^{q^j} x_0 + \alpha^{q^{j+1}} x_1 + \alpha^{q^{j+2}} x_2 + \dots + \alpha^{q^{j+i}} x_i + \dots + \alpha^{q^{j+n}} x_n$$

for each $0 \leq j \leq n$, and observe that $F_i = y_i^d$. The linear map $x_i \mapsto y_i$ is a linear automorphism of \mathbb{P}^n . Indeed, the matrix of the transformation is known as a *Moore matrix*, which has a non-zero determinant in this case. Thus, y_i are algebraically independent, which in particular implies that $\{F_0, F_1, \dots, F_n\}$ is a linearly independent set. Hence, $\mathcal{L} = \langle F_0, F_1, \dots, F_n \rangle$ has (projective) dimension $r = n$.

We claim that each \mathbb{F}_q -member of \mathcal{L} defines a smooth hypersurface. Suppose that we have a singular hypersurface X which belongs to \mathcal{L} :

$$(2.4) \quad X = \{c_0 F_0 + c_1 F_1 + \dots + c_n F_n = 0\}$$

for some $c_i \in \overline{\mathbb{F}_q}$ where not all c_i are zero. In the new coordinates y_i , we can express (2.4) as:

$$X = \{c_0 y_0^d + c_1 y_1^d + \dots + c_n y_n^d = 0\}$$

Since X is singular, we can apply Lemma 2.2 to deduce that $c_i = 0$ for some i . Without loss of generality, we assume $c_0 = 0$. By applying the Frobenius map, we see that X is sent to:

$$X^\sigma = \{c_1^q F_2 + \dots + c_n^q F_0 = 0\}$$

We claim that X and X^σ are distinct. Indeed, their defining equations are not multiple of one another: otherwise, there would exist a constant b such that $c_i^q = b \cdot c_{i+1}$ for each $0 \leq i \leq n$ taken modulo $n+1$. As $c_0 = 0$, this would force $c_i = 0$

for each $0 \leq i \leq n$, which is a contradiction. Thus, X is not defined over \mathbb{F}_q . We have shown that each singular member of \mathcal{L} is not defined over \mathbb{F}_q giving us the desired conclusion.

When $p \mid d$ but $p \nmid (n+1)$, we have an analogous construction. Using the same notation as in (2.3), we define $F_i = y_i^q y_{i+1}$ for $0 \leq i \leq n-1$ and $F_n = y_n^q y_0$. One can check that $\mathcal{L} = \langle F_0, F_1, \dots, F_n \rangle$ is a linear subspace of (projective) dimension n . By following the same argument as above, except applying Lemma 2.3 instead of Lemma 2.2, it follows that each \mathbb{F}_q -member of \mathcal{L} is smooth. \square

3. QUADRICS IN CHARACTERISTIC 2

In this section, we will show that the hypothesis $p \nmid \gcd(d, n+1)$ in our main theorem cannot be removed in general. We will focus on the case $p = d = 2$ and odd n . Our goal is to prove the following result:

Theorem 3.1. *Suppose n is an odd positive integer, and $q = 2^s$ is an even prime power. There does **not** exist a linear system $\mathcal{L} = \langle F_0, F_1, \dots, F_n \rangle$ of dimension n over \mathbb{F}_q such that each \mathbb{F}_q -member of \mathcal{L} is a smooth quadric hypersurface.*

As a result, Question 1.1 has a negative answer when $q = 2^s$, $d = 2$ and $r = n$ is an odd integer.

We start with a crucial lemma on singular quadric hypersurfaces.

Lemma 3.2. *Suppose $q = 2^s$ (for some positive integer s) and n is odd. Consider a quadric hypersurface $X \subset \mathbb{P}^n$ defined over \mathbb{F}_q by the following homogeneous polynomial:*

$$F = x_0^2 + G(x_1, x_2, \dots, x_n)$$

where G is a homogeneous polynomial of degree 2 in the variables x_1, \dots, x_n . Then X is singular.

Proof. We claim that the homogeneous system of linear equations $\frac{\partial G}{\partial x_i} = 0$ for $i = 1, \dots, n$ has a common nontrivial solution. It suffices to show that the matrix M of this linear system is singular. Note that the matrix M is the Hessian matrix of G ; since G is quadratic, the entries of the Hessian matrix are constant. The Hessian matrix M is symmetric and hence also skew-symmetric as the characteristic is 2. It remains to show that a skew-symmetric square $n \times n$ matrix M over a commutative ring has zero determinant, when n is odd.

It suffices to consider the universal skew-symmetric matrix A over the polynomial ring $R = \mathbb{Z}[x_{ij} \mid 1 \leq i < j \leq n]$. By definition, the (i, j) -th entry of A is x_{ij} if $i < j$, 0 if $i = j$ and $-x_{ij}$ if $i > j$. Taking the determinant on both sides of $A^T = -A$, we obtain $\det(A) = -\det(A)$ in R and hence, $\det(A) = 0$, because R is an integral domain of characteristic 0. A simple specialization argument (specializing x_{ij} to the (i, j) -th entry of M) now shows that $\det(M) = 0$.

Thus, we can find a point $P = [t_1 : t_2 : \dots : t_n] \in \mathbb{P}^{n-1}$ such that $\frac{\partial G}{\partial x_i}(P) = 0$ for each i . Now, we want to add one more coordinate to P so that it belongs to $X = \{F = 0\}$. To achieve this goal, we solve for x_0 from the equation:

$$x_0^2 + G(t_1, t_2, \dots, t_n) = 0$$

to get a solution, say $x_0 = t_0$. This is possible because every element in \mathbb{F}_{2^s} has a square root in \mathbb{F}_{2^s} (for any positive integer s). Then $P' = [t_0 : t_1 : \dots : t_n] \in \mathbb{P}^n$

clearly satisfies $F(P') = 0$ and $\frac{\partial F}{\partial x_i}(P') = 0$ for each $0 \leq i \leq n$. Thus, X is singular at P' . \square

Remark 3.3. The conclusion of Lemma 3.2 is false when $n = 2k$ is even. Indeed, the quadric hypersurface in \mathbb{P}^n defined by the polynomial

$$x_0^2 + x_1x_2 + x_3x_4 + \dots + x_{2k-1}x_{2k} = 0$$

is smooth.

We now proceed with a proof of Theorem 3.1.

Proof of Theorem 3.1. Suppose, to the contrary, that $\mathcal{L} = \langle F_0, \dots, F_n \rangle$ is a linear system of quadric hypersurfaces of (projective) dimension n where each \mathbb{F}_q -member of \mathcal{L} is smooth. Let $\mathcal{L}(\mathbb{F}_q)$ denote the set of \mathbb{F}_q -members of the system.

Let us identify \mathbb{F}_q^{n+1} with the quadratic forms involving x_0 . In other words, $(a_i)_{i=0}^n \in \mathbb{F}_q^{n+1}$ is identified with the polynomial $\sum_{i=0}^n a_i x_0 x_i$. Consider the map

$$\Psi : \mathcal{L}(\mathbb{F}_q) \rightarrow \mathbb{F}_q^{n+1}$$

which sends $F \in \mathcal{L}(\mathbb{F}_q)$ to the part of F that only contains the monomials involving x_0 . Note that $\mathcal{L}(\mathbb{F}_q)$ is an $(n+1)$ -dimensional \mathbb{F}_q -vector space, and Ψ is an \mathbb{F}_q -linear map.

We claim that Ψ is injective. Indeed, if $\Psi(H) = (0, \dots, 0)$ for some nonzero H , then such a polynomial defines a quadric hypersurface which is singular at the point $[1 : 0 : \dots : 0]$, a contradiction.

By dimension count, Ψ must also be surjective. In particular, there exists some $F \in \mathcal{L}(\mathbb{F}_q)$ whose image under Ψ is x_0^2 . In other words, $F = x_0^2 + G(x_1, \dots, x_n)$ for some G . By Lemma 3.2, F defines a singular quadric hypersurface, which again contradicts the assumption that each \mathbb{F}_q -member of \mathcal{L} is smooth. We conclude that such an \mathcal{L} does not exist. \square

We have shown that the hypothesis $p \nmid \gcd(d, n+1)$ cannot be removed in the case $p = 2$. It seems very subtle to determine whether our main result Theorem 1.2 continues to hold even when we assume $p \mid \gcd(d, n+1)$ for some other primes p . We finish the paper by presenting a positive example when $p \mid \gcd(d, n+1)$.

Example 3.4. Let $q = d = 3$ and $n = 2$. Consider the following cubic homogeneous polynomials with coefficients in \mathbb{F}_3 :

$$F_0 = x^3 + x^2y - xy^2 + y^3 + x^2z + xyz + y^2z - xz^2 + z^3$$

$$F_1 = x^3 + x^2y - x^2z - xyz + y^2z + z^3$$

$$F_2 = x^3 - x^2y + xy^2 + y^3 + x^2z + xyz + y^2z - yz^2$$

One can check with a computer that $aF_0 + bF_1 + cF_2 = 0$ defines a smooth plane curve for each of the possible $q^2 + q + 1 = 13$ choices $[a : b : c] \in \mathbb{P}^2(\mathbb{F}_3)$. Thus, we have a linear system of dimension $r = n = 2$ which satisfies the conclusion of Theorem 1.2 (2), even though $p \mid \gcd(d, n+1)$ holds.

REFERENCES

- [AG22a] Shamil Asgarli and Dragos Ghioca, *A Bertini type theorem for pencils over finite fields*, Finite Fields Appl. **77** (2022), Paper No. 101936, 13, DOI 10.1016/j.ffa.2021.101936.
- [AG22b] ———, *Smoothness in pencils of hypersurfaces over finite fields*, Bull. Aust. Math. Soc. (to appear), arXiv e-prints (2022), available at <https://arxiv.org/abs/2203.07169>.

- [Bal07] E. Ballico, *Bertini's theorem over a finite field for linear systems of quadrics*, Int. J. Pure Appl. Math. **35** (2007), no. 4, 453–455.
- [Bal09] ———, *Vanishings and non-vanishings of homogeneous forms over a finite field*, Int. J. Pure Appl. Math. **57** (2009), no. 2, 219–224.
- [DGH21] Vesselin Dimitrov, Ziyang Gao, and Philipp Habegger, *Uniform bound for the number of rational points on a pencil of curves*, Int. Math. Res. Not. IMRN **2** (2021), 1138–1159, DOI 10.1093/imrn/rnz248.
- [PY08] Jorge Vitório Pereira and Sergey Yuzvinsky, *Completely reducible hypersurfaces in a pencil*, Adv. Math. **219** (2008), no. 2, 672–688, DOI 10.1016/j.aim.2008.05.014.
- [Ste89] Yosef Stein, *The total reducibility order of a polynomial in two variables*, Israel J. Math. **68** (1989), no. 1, 109–122, DOI 10.1007/BF02764973.
- [Vis93] Angelo Vistoli, *The number of reducible hypersurfaces in a pencil*, Invent. Math. **112** (1993), no. 2, 247–262, DOI 10.1007/BF01232434.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, BC V6T 1Z2

Email address: sasgarli@math.ubc.ca

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, BC V6T 1Z2

Email address: dghioca@math.ubc.ca

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, BC V6T 1Z2

Email address: reichst@math.ubc.ca