

# On fractional sums of the divisor functions

Wei Zhang

**Abstract** In this paper, we consider the fractional sum of the divisor functions. We can improve previous results considered by Bordellés [2] and Liu-Wu-Yang [6].

**Keywords** Divisor function, Exponential sum, Integral Part, Floor function

**2000 Mathematics Subject Classification** 11N37, 11L07

## 1. INTRODUCTION

Recently, the sum

$$S_f(x) = \sum_{n \leq x} f\left(\left[\frac{x}{n}\right]\right)$$

has attracted many experts special attention (for example, see [1, 2, 6, 10, 11]), where  $f$  is a complex-valued arithmetic function and  $[\cdot]$  denotes the floor function (i.e. the greatest integer function). One can call  $S_f$  the fractional sum of  $f$  (see [8]).

Specially, for some fixed  $\eta \in (0, 1)$  and

$$f(n) \ll n^\eta,$$

independently, Wu [10] and Zhai [11] showed that

$$S_f(x) = C_f x + O(x^{(1+\eta)/2}),$$

where  $f$  is a complex-valued arithmetic function and

$$C_f = \sum_{n=1}^{\infty} \frac{f(n)}{n(n+1)}.$$

This formula improves the recent result obtained by Bordellés, Dai, Heyman, Pan and Shparlinski [1].

In fact, for general function  $f$  being a positive real-valued arithmetic function, then one can obtain some much better results by involving the theory of Fourier series [9] and exponential sums [3]. For example, one can refer to [2, 6, 7, 8].

By using the symmetry of divisor functions, in [7], it is proved that

$$S_\tau(x) = \sum_{n \leq x} \tau\left(\left[\frac{x}{n}\right]\right) = \sum_{n=1}^{\infty} \frac{\tau(n)}{n(n+1)} x + O(x^{11/23+\varepsilon}),$$

where  $\tau(n)$  is the number of representations of  $n$  as product of two natural numbers and

$$11/23 \approx 0.4782.$$

Recently, this result was improved by many experts. In [2], by using the Dirichlet hyperbolic method and more effort, it is proved that

$$S_\tau(x) = \sum_{n \leq x} \tau\left(\left[\frac{x}{n}\right]\right) = \sum_{n=1}^{\infty} \frac{\tau(n)}{n(n+1)} x + O(x^{19/40+\varepsilon}),$$

where

$$19/40 = 0.475.$$

By using a new estimate on 3-dimensional exponential sums, in [6], Liu-Wu-Yang showed that

$$S_\tau(x) = \sum_{n \leq x} \tau \left( \left[ \frac{x}{n} \right] \right) = \sum_{n=1}^{\infty} \frac{\tau(n)}{n(n+1)} x + O(x^{9/19+\varepsilon}),$$

where

$$9/19 \approx 0.47368.$$

This sum has relations to the generalized divisor function  $\tau_k(n)$  ( $k \geq 2$ ), where  $\tau_k(n)$  is the number of representations of  $n$  as product of  $k$  natural numbers. Hence, in [2] and [6], some results for these arithmetic functions are also given. Precisely, in [6], it is proved that

$$S_{\tau_k}(x) = \sum_{n \leq x} \tau_k \left( \left[ \frac{x}{n} \right] \right) = \sum_{n=1}^{\infty} \frac{\tau_k(n)}{n(n+1)} x + O(x^{\theta(k)+\varepsilon}), \quad (1.1)$$

where

$$\theta(k) = \frac{5k-1}{10k-1}.$$

These results improve previous results of [2] and can be seen as a generalization of  $S_\tau(x)$ . More recently, by using some deep results of Jutila [4], Stucky [8] showed that

$$S_\tau(x) = \sum_{n \leq x} \tau \left( \left[ \frac{x}{n} \right] \right) = \sum_{n=1}^{\infty} \frac{\tau(n)}{n(n+1)} x + O(x^{5/11+\varepsilon}), \quad (1.2)$$

where

$$5/11 \approx 0.454545.$$

This improves previous results of  $S_\tau(x)$ . It is reasonable to believe that one can obtain much better results for  $S_{\tau_k}(x)$  by using the method of Stucky [8]. However, the method of Stucky [8] is too special to be generalized to the situation of  $S_{\tau_k}(x)$ .

The purpose of this paper is to study  $S_{\tau_k}(x)$ . We can improve (1.1) by giving the following result, which gives improvement for

$$\theta(k) = \frac{5k-1}{10k-1}.$$

**Theorem 1.1.** *Let  $\tau_k(n)$  be the number of representations of  $n$  as product of  $k$  natural numbers. Then we have*

$$S_{\tau_k}(x) = \sum_{n \leq x} \tau_k \left( \left[ \frac{x}{n} \right] \right) = \sum_{n=1}^{\infty} \frac{\tau_k(n)}{n(n+1)} x + O(x^{9/19+\varepsilon}).$$

*Remark 1.* In fact, a little better result can also be given. One can refer to [5] for details.

## 2. PROOF OF THEOREM 1.1

We will start the proof for Theorem 1.1 with some necessary lemmas. The following lemma is from [6].

**Lemma 2.1.** *Let  $\alpha > 0$ ,  $\beta > 0$ ,  $\gamma > 0$  and  $\delta \in \mathbb{R}$  be some constants. For  $X > 0$ ,  $H \geq 1$ ,  $M \geq 1$ , and  $N \geq 1$ , define*

$$S_\delta = S_\delta(H, M, N) := \sum_{h \sim H} \sum_{m \sim M} \sum_{n \sim N} a_{h,n} b_m e \left( X \frac{M^\beta N^\gamma}{H^\alpha} \frac{h^\alpha}{m^\beta n^\gamma + \delta} \right),$$

where  $e(t) = e^{2\pi i t}$ , the  $a_{h,n}$  and  $b_m$  are complex numbers such that  $a_{h,n} \leq 1$ ,  $b_m \leq 1$  and  $m \sim M$  means that  $M < m \leq 2M$ . For any  $\varepsilon > 0$ , we have

$$S_\delta \ll \left( (X^\kappa H^{2+\kappa} M^{1+\kappa+\lambda} N^{2+\lambda})^{1/(2+2\kappa)} + HM^{1/2}N + H^{1/2}MN^{1/2} + X^{-1/2}HMN \right) X^\varepsilon$$

uniformly for  $M \geq 1$ ,  $N \geq 1$ ,  $H \leq N^{\gamma-1}M^\beta$  and  $|\delta| \leq 1/\varepsilon$ , where  $(\kappa, \lambda)$  is an exponent pair and the implied constant may depend on  $\alpha, \beta, \gamma$ , and  $\varepsilon$ .

Next lemma can be seen in Theorem A.6 in [3] or Theorem 18 in [9].

**Lemma 2.2.** *Suppose that  $H \geq 1$ . There is a function  $\psi^*(x)$  such that*

- $\psi^*(x) = \sum_{1 \leq |h| \leq H} \gamma(h) e(hx)$ ,
- $\gamma(h) \ll \frac{1}{h}$ ,  $(\gamma(h))' \ll \frac{1}{h^2}$ ,
- $|\psi^*(x) - \psi(x)| \leq \frac{1}{2H+2} \sum_{h \leq H} \left(1 - \frac{|h|}{H}\right) e(hx)$ .

Let  $\psi(t) = t - [t] - 1/2$  for  $t \in \mathbb{R}$  and  $\delta \geq 0$ . Define

$$\mathfrak{G}(x, D) := \sum_{D < d \leq 2D} f(d) \psi \left( \frac{x}{d + \delta} \right).$$

We also need the following well-known lemma.

**Lemma 2.3.** *Let  $g^{(m')}(x) \sim YX^{1-m'}$  for  $1 < X \leq x \leq 2X$  and  $m' = 1, 2, \dots$ . Then one has*

$$\sum_{X < n \leq 2X} e(g(n)) \ll Y^\kappa X^\lambda + Y^{-1},$$

where  $(\kappa, \lambda)$  is any exponent pair.

Now we begin the proof of Theorem 1.1. Let

$$N = x^{9/19}.$$

We can write

$$S_{\tau_k}(x) := S_{\tau_k,1} + S_{\tau_k,2},$$

where

$$S_{\tau_k,1} = \sum_{n \leq N} \tau_k \left( \left[ \frac{x}{n} \right] \right) \tag{2.1}$$

and

$$S_{\tau_k,2} = \sum_{N < n \leq x} \tau_k \left( \left[ \frac{x}{n} \right] \right). \tag{2.2}$$

Obviously, by  $\tau_k(n) \ll n^\varepsilon$ , we have

$$\begin{aligned} S_{\tau_k,1} &= \sum_{n \leq N} \tau_k \left( \left[ \frac{x}{n} \right] \right) = \sum_{n \leq N} (x/n)^\varepsilon \\ &\ll N^{1+\varepsilon} \\ &\ll x^{9/19+\varepsilon}. \end{aligned}$$

As to  $S_{\tau_k,2}$ , firstly, by  $\tau_k(n) \ll n^\varepsilon$ , we have

$$\sum_{n \leq x} |\tau_k(n)| \ll x^{1+\varepsilon}.$$

Hence we can get

$$\begin{aligned} S_{\tau_k,2} &= \sum_{N < n \leq x} \tau_k \left( \left[ \frac{x}{n} \right] \right) \\ &= \sum_{d \leq x/N} \tau_k(d) \sum_{x/(d+1) < n \leq x/d} 1 \\ &= \sum_{d \leq x/N} \tau_k(d) \left( \frac{x}{d} - \frac{x}{d+1} - \psi\left(\frac{x}{d}\right) + \psi\left(\frac{x}{d+1}\right) \right) \\ &= x \sum_{d=1}^{\infty} \frac{\tau_k(d)}{d(d+1)} + x \sum_{d > x/N} \frac{\tau_k(d)}{d(d+1)} + (\log x) \sum_{D < d \leq 2D} \tau_k(d) \psi\left(\frac{x}{d+\delta}\right) \\ &= x \sum_{d=1}^{\infty} \frac{\tau_k(d)}{d(d+1)} + O(N^{1+\varepsilon}) + (\log x) \sum_{D < d \leq 2D} \tau_k(d) \psi\left(\frac{x}{d+\delta}\right), \end{aligned} \tag{2.3}$$

where  $N \leq D \leq x/N$ . Then we need to estimate

$$\sum_{D < d \leq 2D} \tau_k(d) \psi\left(\frac{x}{d+\delta}\right).$$

By Lemma 2.2, we have

$$\begin{aligned} &\sum_{D < d \leq 2D} \tau_k(d) \psi\left(\frac{x}{d+\delta}\right) \\ &\ll \sum_{1 \leq h \leq H} \frac{1}{h} \sum_{D < d \leq 2D} \tau_k(d) e\left(\frac{hx}{d+\delta}\right) \\ &\quad + \sum_{1 \leq h \leq H} \frac{1}{H} \sum_{D < d \leq 2D} \tau_k(d) e\left(\frac{hx}{d+\delta}\right) + D/H. \end{aligned} \tag{2.4}$$

Then we will focus on the estimate of

$$S_\delta := \sum_{1 \leq h \leq H} \frac{1}{h} \sum_{D < d \leq 2D} \tau_k(d) e\left(\frac{hx}{d+\delta}\right).$$

By using the relation

$$\sum_{n_1 n_2 \cdots n_k = n} = \tau_k(n),$$

and the dichotomy method, we have

$$S_\delta \ll D^\varepsilon \sum_{1 \leq h \leq H} \frac{1}{h} \sum_{d_i \sim D_i, i=1,2,\dots,k} e\left(\frac{hx}{d_1 d_2 \cdots d_k + \delta}\right),$$

where

$$d_i \leq d_{i+1}, \quad D_i \leq D_{i+1}, \quad \text{for } 1 \leq i \leq k-1 \quad (2.5)$$

and

$$\prod_{i=1}^k D_i \sim D \quad (2.6)$$

### Case I

Suppose that  $D_k \geq D^{2/3}$ .

By Lemma 2.3 and choosing  $(\kappa, \lambda) = (1/2, 1/2)$ , we have

$$\begin{aligned} S_\delta &\ll D^\varepsilon \sum_{1 \leq h \leq H} \frac{1}{h} \sum_{d_i \sim D_i, i=1,2,\dots,k-1} \left( \left( \frac{hx}{d_1 d_2 \cdots d_{k-1} D_k^2} \right)^{1/2} (D_k)^{1/2} \right. \\ &\quad \left. + \frac{(d_1 d_2 \cdots d_{k-1})^2}{hx} \right) \\ &\ll D^{2/9} x^{1/3} + D^2/x, \end{aligned}$$

where we have chosen  $H = D^{7/9} x^{-1/3}$ . We assume that  $D > x^{1/2}$ . Hence we can verify that  $H > 1$ . Then for  $D_k \geq D^{2/3}$ , by choosing  $N = x^{9/19}$ , we have

$$S_\delta \ll D^{2/9} x^{1/3} \ll x^{77/171+\varepsilon} \ll x^{9/19}.$$

### Case II

Suppose that  $D^{1/3} \leq D_k \leq D^{2/3}$ .

By choosing  $N = x^{9/19}$  and  $(\kappa, \lambda) = (1/2, 1/2)$  in Lemma 2.1, we have

$$\begin{aligned} S_\delta &\ll (DH)^\varepsilon (x^{1/6} D^{1/2} D^{1/12} + D^{3/4} + x^{-1/2} D^{3/2}) \\ &\ll x^{9/19+\varepsilon}. \end{aligned}$$

### Case III

Suppose that  $D_k \leq D^{1/3}$ . Then by (2.5) and (2.6), we have  $D_i \leq D^{1/3}$ ,  $i = 1, 2, \dots, k$ . We also suppose that  $t$  is the least integer such that  $D_1 D_2 \cdots D_t > D^{1/3}$ . Then we have

$$D^{1/3} \leq (D_1 D_2 \cdots D_{t-1}) D_t \leq D^{2/3}.$$

Let  $l_1 = d_1 d_2 \cdots d_t$  and let  $l_2 = d_{t+1} d_{t+2} \cdots d_k$ . Then we have

$$S_\delta \ll D^\varepsilon \sum_{1 \leq h \leq H} \frac{1}{h} \sum_{l_1 \sim L_1} \sum_{l_2 \sim L_2} e\left(\frac{hx}{l_1 l_2 + \delta}\right),$$

where  $D^{1/3} \leq L_1 \leq D^{2/3}$  and  $D^{1/3} \leq L_2 \leq D^{2/3}$ . Then similar as the second case, we have

$$\begin{aligned} S_\delta &\ll (DH)^\varepsilon (x^{1/6} D^{1/2} D^{1/12} + D^{3/4} + x^{-1/2} D^{3/2}) \\ &\ll x^{9/19+\varepsilon}. \end{aligned}$$

Then from the above three cases, we have

$$S_\delta := \sum_{1 \leq h \leq H} \frac{1}{h} \sum_{D < d \leq 2D} \tau_k(d) e\left(\frac{hx}{d+\delta}\right) \ll x^{9/19+\varepsilon}.$$

Then by (2.3)-(2.4), we have

$$S_{\tau_k,2} = x \sum_{d=1}^{\infty} \frac{\tau_k(d)}{d(d+1)} + O(x^{9/19+\varepsilon}).$$

Recall (2.1) and (2.2), then we can finally give Theorem 1.1.

**Acknowledgements** The author would like to thank the the referee who gives some detailed corrections and suggestions. This work was supported by NSFC(11871307).

#### REFERENCES

- [1] O. Bordellés, L. Dai, R. Heyman, H. Pan and I. Shparlinski, *On a sum involving the Euler function*. J. Number Theory 202(2019), 278-297.
- [2] O. Bordellés, *On certain sums of number theory*, arXiv:2009.05751v2 [math.NT] 25 Nov 2020.
- [3] W. Graham and G. Kolesnik, *Van der Corput's Method of Exponential Sums*. Cambridge Univ. Press, 1991.
- [4] M. Jutila, *Lectures on a method in the theory of exponential sums*, volume 80 of Tata Institute of Fundamental Research Lectures on Mathematics and Physics, Published for the Tata Institute of Fundamental Research, Bombay; by Springer-Verlag, Berlin, 1987.
- [5] K. Liu, J. Wu and Z.S. Yang *A variant of the prime number theorem*. Indag. Math. (N.S.) 33(2022), no. 2, 388-396.
- [6] K. Liu, J. Wu and Z.S. Yang *On some sums involving the integral part function*. arXiv:2109.01382v1 [math.NT] 3 Sep 2021.
- [7] J. Ma and H.Y. Sun, *On a sum involving the divisor function*. Period. Math. Hung. to appear.
- [8] J. Stucky, *The fractional sum of small arithmetic functions*. J. Number Theory, to appear.
- [9] J.D. Vaaler, *Some extremal functions in Fourier analysis*. Bull. Amer. Math. Soc. (N.S.) 12(1985), 183-216.
- [10] J. Wu, *Note on a paper by Bordellés, Dai, Heyman, Pan and Shparlinski*. Period. Math. Hungar. 80(2020), 95-102.
- [11] W.G. Zhai, *On a sum involving the Euler function*. J. Number Theory 211(2020), 199-219.

WEI ZHANG, SCHOOL OF MATHEMATICS AND STATISTICS, HENAN UNIVERSITY, KAIFENG 475004, HENAN, CHINA

*Email address:* zhangweimath@126.com