
Myopic Quantal Response Policy: Thompson Sampling Meets Behavioral Economics

Jingying Ding

Antai College of Economics and Management
Shanghai Jiao Tong University
jingyding@sjtu.edu.cn

Yifan Feng

Department of Analytics and Operations
National University of Singapore
yifan.feng@nus.edu.sg

Ying Rong

Antai College of Economics and Management
Shanghai Jiao Tong University
yrong@sjtu.edu.cn

Abstract

We study a new family of multi-armed bandit (MAB) algorithms called Myopic Quantal Response (MQR). It prescribes a simple way to randomize over arms according to historical data and a "coefficient of exploitation," which explicitly controls the exploration-exploitation trade-off. We show that MQR partially extends the Thompson Sampling (TS) algorithm. It is also a dynamic version of quantal response models where the expected utilities are directly estimated from historical rewards. Based on theoretical analysis and numerical experiments, we believe the significance of MQR is three-fold. First, it provides a conceptual framework to understand further the exploration-exploitation trade-off. Second, it can be used as a structural estimation tool to learn from realized actions and rewards how much a given policy (either generated by human beings or algorithms) is "under" or "over" exploring. Finally, it can inspire new MAB algorithms that improve well-studied algorithms (e.g., TS) in a nonasymptotic setting.

1 Introduction

In a wide range of sequential decision problems, the decision makers (DMs) need to strike a trade-off between exploration and exploitation. When facing several alternative options with unknown random payoff, DMs have to uncover the expected payoff of various options while also maximizing their profits over time, and thus they are required to adopt a strategy that can balance exploration and exploitation.

The stochastic Multi-Armed Bandit problem (MAB) is a classical model that captures the exploration-exploitation trade-off inherently. Many literature have studied the different versions and generalizations of the MAB problem, and there are some classical algorithms available for the stochastic MAB. Gittins Index was used to generate an optimal strategy by selecting the arm with the highest index in the discounted infinite horizon setting (Gittins, 1979; Gittins et al., 2011). Upper Confidence Bound (UCB) family of algorithms were designed based on the principle of optimism in the face of uncertainty and had good theoretical guarantees (Lai et al., 1985; Auer et al., 2002). Thompson Sampling (TS) was first proposed by Thompson (1933), and the basic idea is to randomly draw each arm according to its probability of being optimal. While it has been well-known for a long time, provable guarantees did not appear until recently. TS has been shown to be asymptotically regret optimal in some settings, e.g., by the metric of Bayesian Regret (Russo and Van Roy, 2014) or by the metric of worst-case regret (Agrawal and Goyal, 2013, 2017).

The current literature primarily focus on the design and performance of heuristics to MAB problems. In this paper, we attempt to understand MAB problems from a behavioral perspective and to investigate the balance of exploration and exploitation explicitly. We apply ideas from the quantal choice framework to develop intuitive policies with good performance. The classical *quantal choice theory* can be traced back to Luce (1959). The core idea is that all possible choice alternatives are candidates for selection, but better options are chosen more often. It models choice behavior as probabilistic and has been well applied in behavioral economics. In game-theoretic settings, McKelvey and Palfrey (1995) developed the quantal response equilibrium framework, in which all players make choices according to the quantal choice model. They also investigated the use of standard econometric models for quantal choice in extensive form games and defined agent quantal response equilibrium (McKelvey and Palfrey, 1998).

Inspired by the application of the quantal choice model in behavioral economics, we develop the *myopic quantal response* framework to characterize choices of DMs/ algorithms in the MAB problems. Since the MAB problem is a sequential decision making problem, we consider a dynamic version of quantal choice models. That is, the decision making at each time can be viewed as following a quantal choice model. Furthermore, we modify the model into a myopic way, where the expected utilities are directly estimated from historical rewards. Thus, there is no anticipation of future rewards in decision making.

In this paper, we focus on a new family of behavioral MAB algorithms. The main contributions of this work can be summarized as follows. First, we leverage the idea of quantal response and propose a family of tractable and intuitive policies to the MAB problem from a behavioral perspective. We provide a conceptual framework to understand further the exploration-exploitation trade-off. Second, we asymptotically obtain the upper bound on the expected regret incurred by our proposed policies. Third, we use MQR as a structural estimation tool to learn from realized actions and rewards how much a given policy (either generated by human beings or algorithms) is “under” or “over” exploring. Fourth, we numerically verify that adjusting the degree of exploitation according to the time horizon T can improve the performance compared with well-studied algorithms (e.g., TS) in a nonasymptotic setting.

2 The Multi-armed Bandit Problem and the MQR Policy

2.1 Problem Definition

We consider a Multi-Armed Bandit problem with N independent arms. In a finite horizon T , at each time $t \in \{1, 2, \dots\}$, the decision maker (DM) chooses one among N arms to pull. Decisions of the DM are denoted as $i(t)$, where $i(t) \in \mathbb{N} = \{1, 2, \dots, N\}$. The rewards of each arm are stochastic and randomly drawn from some fixed distribution, but the distribution is unknown to the DM. We assume that the random rewards obtained from any given arm i are independent and identically distributed across t and independent from other arms. The rewards are with support in $[0, 1]$. After the decision $i(t)$ is made, the reward r_t is observed after playing the arm.

An algorithm for the MAB problem is to decide which arm to pull at each time t based on the history of the previous $t - 1$ periods. The DM aims to select the arm which generates the largest expected reward and to maximize the expected total rewards at time T . Let μ_i denote the (unknown) expected reward of arm i and $\mu_i \in [0, 1]$. Let $\mu^* := \max_i \mu_i$, and $\Delta_i := \mu^* - \mu_i$. We use $k_i(t)$ to denote the number of plays of arm i until time $t - 1$. The empirical mean is defined as $\hat{\mu}_i(t) := \frac{\sum_{\tau=1}^{t-1} r_i(\tau)}{k_i(t)+1}$, where $r_i(t)$ is the reward of arm i at time t . The reward only generates when arm i is pulled, otherwise the reward is zero.

2.2 Thompson Sampling

Thompson Sampling (TS) is a classical heuristic for multi-armed bandit problems. The basic idea is that for each time t and arm i , TS computes the posterior probability that arm i is the best arm and samples i with this probability. Specifically speaking, based on Bayesian ideas, assume for each arm i , there is a parameter ϑ_i associated with the reward distribution, but the parameter ϑ_i is unknown to the DM. The DM assumes a simple prior distribution \mathcal{P}_i on the parameters ϑ_i , that is, $\vartheta_i \sim \mathcal{P}_i(y_i)$, where y_i is belief and is updated after reward observations. At any time step, the DM

samples parameters $\tilde{\vartheta}_i \sim \mathcal{P}_i(y_i)$ for each arm i based on the current belief and plays an arm that is believed to be the best according to $\tilde{\vartheta}_i$.

3 Myopic Quantal Response Policies

We apply the quantal response framework (Luce, 1959; McKelvey and Palfrey, 1995, 1998) to the MAB problem and propose a general class of simple and intuitive policies which we call *myopic quantal response* (MQR) policies. The main idea is that we assume the decision maker (DM) makes choices based on a quantal choice model so that the DM is playing stochastic responses and the intensity of exploration can be modeled, which helps us to understand the the exploration-exploitation trade-off in a explicit way.

The quantal choice/response model is a well-developed statistical model. The simplest Luce model states that the choice probability of selection i from a pool of j selections is given by $P(i) = w_i / \sum_j w_j$, where w indicates the stimulus of a particular selection. In economic models, the stimulus can be measured in terms of expected payoffs. The logit choice model is a special case in which the choice probability is proportional to e^u , where u is the utility. It implies that all possible alternatives are candidates for selection, but more attractive alternatives (yielding higher payoffs) are chosen with larger probabilities.

The quantal choice model has close connections with behavioral economics. The quantal choice model assumes individual choices or responses are rational, but are based on latent variables that are not observed. McKelvey and Palfrey (1995) observed that subjects frequently made errors in choosing Nash equilibrium actions in experiments, and they suggested that these errors may be due to bounded rationality. These errors can be modeled by using latent utility components in the quantal choice model that are not reflected in payoffs. Thus, they introduced the quantal response equilibrium (QRE) to incorporate the bounded rationality into game-theoretic settings. The QRE framework is well established in behavioral economics and has good predictive power in many game-theoretic settings. Under this framework, the best response becomes probabilistic rather than deterministic. From a behavioral perspective, the probabilistic choices can be interpreted that players may make errors or utilize private information in choosing among strategies.

In the MAB problem, we leverage the idea of quantal response that players make choices based on some probabilities and derive a class of behavioral algorithms. The DMs collect random rewards from the selected arm sequentially, and they need to make exploration-exploitation trade-off in the process. As the exploitation of obtained information, the DMs are more likely to choose better arms than worse arms based on the historical rewards. Meanwhile, they reserve the probabilities to choose “suboptimal” arms (which are not performed well at the decision-making time) to explore the distribution of rewards. Also, the extent of exploration is adaptive according to the state of arms (i.e., how much information the DM know).

In what follows, we will first describe the general framework and then propose two specific forms (the distribution of the noise terms) as a start point to analyze.

MQR Policies. The main task of the MQR policies is to decide the choice probability of each arm at each time based on the idea of quantal response. A MQR policy consists of three components: an empirical mean, a noise term and an exploration factor, and next we will elaborate these in sequence.

First, an empirical mean is defined as $\hat{\mu}_i(t)$ before and it serves as the exploitation of historical information. Since the expected reward of each action is not available in the MAB setting, we utilize an adjusted sample mean, which is divided by the number of samples plus one. Note that $\hat{\mu}_i(1) = 0$. Since the DMs use the empirical mean based on the history and have no anticipation of future rewards, we say that it is *myopic*.

Second, the noise term ϵ is similar to the payoff disturbance, which represents the randomness of exploration in the MAB problem. We assume that the noise term ϵ is a continuous random variable and is independently and identically distributed across time t and arm i with cumulative distribution function $F_\epsilon(\cdot)$.

Third, an exploration factor $\beta_i(t)$ is introduced to adjust the intensity of exploration. With more data collected, the variability of the empirical mean of rewards decreases, so the scale of the noise term should also decrease as t increases. Therefore, we introduce a state-dependent factor $\beta_i(t)$, which

depends on the number of playing arm i till time $t - 1$. We define that $\beta_i := 1/(k_i + 1)^\alpha$, where $\alpha > 0$. The parameter α can be interpreted as the degree of exploitation. When $\alpha \rightarrow \infty$, $\beta \rightarrow 0$, then MQR would be a pure exploitation heuristic since it only uses the information of the empirical mean. On the other hand, if $\alpha \rightarrow 0$, then β would approach one. The MQR tends to pay more attention to the exploration as the random term achieves largest variance in this case.

Based on these, we propose the following algorithm. See Algorithm 1 for details.

Algorithm 1: Myopic Quantal Response Policies

Initialization. Set $k_i = 0, \hat{\mu}_i = 0, \beta_i = 1/(k_i + 1)^\alpha$ for each arm $i = 1, \dots, N$.

Main Step.

At time $t \in \{1, 2, \dots, T\}$:

For each arm $i = 1, \dots, N$, sample ϵ_i independently from the distribution $F_\epsilon(\cdot)$. Let

$$\theta_i(t) = \hat{\mu}_i(t) + \epsilon_i \beta_i(t).$$

Play arm $i(t) := \arg \max \theta_i(t)$ and observe reward r_t .

$$\text{Set } \hat{\mu}_{i(t)} := \frac{\hat{\mu}_{i(t)}(k_{i(t)}+1) + r_t}{k_{i(t)}+2}, k_{i(t)} := k_{i(t)} + 1.$$

Quantities $\theta_i(t)$ denotes a sample generated by the decision rule for each arm i , which leverage the idea of the quantal choice model. Note that $\beta_i(t)$ in the algorithm can be adjusted according to the state $k_i(t)$ of arm i . It reveals that DM's extent of exploration decreases as more experience accumulates.

Specification of the noise term. In the remainder of the paper, we focus mainly on two specialized versions of the model specification, MQR using Gumbel noise and MQR using Gaussian noise. MQR using Gumbel noise denotes that ϵ_i is independently drawn from the Gumbel distribution with location parameter 0 and scale parameter 1, and MQR using Gaussian noise represents that ϵ is sampled from Gaussian distribution $\mathcal{N}(0, 1)$. Both of these two cases have an origin in the study of individual choice behavior. Given all β_i are equivalent, if ϵ has a Gumbel distribution or we call an extreme value distribution, then the choice model is defined as a logistic quantal response function (Luce, 1959). If β_i varies between different arms, it is a variant of the logit choice model, which is known as heteroscedastic extreme value model (Bhat, 1995). As for ϵ with Gaussian distribution, the choice behavior corresponds to a heteroscedastic probit choice model (Alvarez and Brehm, 1995).

Degree of Exploitation. Note that $\beta_i = 1/(k_i + 1)^\alpha$ in Algorithm 1. The parameter α is the key in MQR policies and it adds more flexibility to the algorithms. Specifically, we can adjust the extent of exploitation by taking different values of α . As we mention before, larger α represents more exploitation in the learning process. For different time horizon T , the DM should distribute appropriate proportion to exploration and exploitation to achieve largest total rewards, so α plays an important role when facing various T . We will prove that when $\alpha = 0.5$, it is asymptotically optimal in respect of regret. But in a finite time horizon where T does not tend to infinity, the MQR policy with $\alpha = 0.5$ would not be optimal. Therefore, we need to adjust the value of α , namely, the degree of exploitation according to T .

Next, we will investigate the performance of the MQR policies and analyze how the DM make exploration-exploitation trade-off throughout the learning process. The analysis will be taken in two perspectives: one is the theoretic analysis from the asymptotic regime, and the other is the numerical analysis considering a finite time horizon process.

4 Theoretical Analysis of MQR and Asymptotic Characterization

In this section, we evaluate the performance of the MQR policies and provide bounds for the finite time expected regret of the policies. First, we define the expected total *regret* as a measurement, that is the difference between the reward of the optimal arm i^* and the reward of the arm played. Then, the expected regret in time horizon T can be represented as

$$\mathbb{E}[\mathcal{R}(T)] = \mathbb{E} \left[\sum_{t=1}^T (\mu^* - \mu_{i(t)}) \right] = \sum_i \Delta_i \cdot \mathbb{E}[k_i(T+1)], \quad (1)$$

where $i(t)$ is the arm played at time t , and $k_i(T+1)$ is the number of playing arm i till period T . The expectation is taken over the actions $\{i(t)\}$, which are specified by the policies.

Assumption 1. *The cumulative distribution function $F_\epsilon(\cdot)$ of the noise term ϵ satisfies the following conditions:*

$$(1) F_\epsilon(x) \leq 1 - \frac{1}{\sqrt{2\pi}} \frac{x}{x^2+1} e^{-x^2/2} \text{ for any } x \geq 0;$$

$$(2) F_\epsilon(-x) \leq \frac{1}{2} e^{-x^2/2} \text{ for } x \geq 8;$$

(3) *The stochastic term ϵ has mean $m \geq 0$ and standard deviation $\sigma < \infty$.*

Assumption 1 proposes the requirements on the distribution of the noise term ϵ . Condition (1) corresponds to the anti-concentration property, while condition (2) can be interpreted as the requirement of the concentration property. Furthermore, condition (3) is a general assumption on distributions. Lemma 2 shows that the two specifications that we study conform to Assumption 1.

Lemma 2. *Gumbel distribution with location parameter 0 and scale parameter 1 and Gaussian distribution $\mathcal{N}(0, 1^2)$ satisfy Assumption 1.*

Our main finding in this subsection is in Theorem 3. The proof is provided in Appendix.

Theorem 3. *With cumulative density function $F_\epsilon(\cdot)$ of the noise term ϵ satisfying Assumption 1 and $\alpha = 0.5$, the MQR policy has the expected regret*

$$\mathbb{E}[\mathcal{R}(T)] \leq O(\sqrt{T}).$$

The proof follows the similar steps as the proof of Theorem 1.3 in Agrawal and Goyal (2017). We highlight the main difference here. Compared to their regret analysis of TS with Gaussian priors, since we generalize the noise term and design a new form of quantities $\theta_i(t)$, we do not have properties of Gaussian variables in analysis. So, we need to modify the proof by decomposing $\theta_i(t)$ and apply conditions in Assumption 1. To be specific, we use the concentration and anti-concentration properties of the noise term ϵ instead of those of Gaussian r.v. $\theta_i(t)$ in Agrawal and Goyal (2017).

Corollary 4. *When $\alpha = 0.5$, the MQR using Gaussian noise that ϵ follows normal distribution $\mathcal{N}(0, 1^2)$ is equivalent to Thompson sampling using Gaussian priors.*

Proof. Note that Thompson sampling using Gaussian priors draw samples independently from the posterior distribution $\mathcal{N}(\hat{\mu}_i, \frac{1}{k_i+1})$. The MQR using Gaussian noise samples quantities $\theta_i(t) = \hat{\mu}_i(t) + \epsilon_i \beta_i(t)$, where $\epsilon_i \sim \mathcal{N}(0, 1^2)$ and $\beta_i(t) = 1/\sqrt{k_i+1}$. By the property of normal distribution, these two heuristics are equivalent.

Next, we claim that the MQR policies with any value of α are long-run-average optimal (i.e., achieve sublinear regrets).

Theorem 5. *For a multi-armed bandit problem with N arms, the MQR algorithm has the expected regret*

$$\mathbb{E}[\mathcal{R}(T)] = o(T).$$

The proof is in Appendix. The result above says although only $\alpha = 0.5$ matches the optimal order for asymptotic performance, all values of α are “plausible” behavioral MAB policies. This is in contrast to, say, ϵ -greedy algorithms, which appear to be very similar to MQR at first sight.

5 Nonasymptotic Analysis of MQR and Numerical Experiments

In this section, we provide numerical analysis in a finite horizon. Considering the practical constraints, the performance in a finite time horizon T that does not tend to infinity is of important value and is significantly different from the asymptotic analysis in Section 4. The algorithms that perform well in the asymptotic analysis pursue the exploration-exploitation trade-off in a long-run time. However, if T does not go to infinity, the benefit of collecting reward information only works for a finite period of time. Therefore, the degree of exploration should be tuned down appropriately according to length of the time horizon T to help algorithms achieve satisfactory rewards. In the MQR policies, the parameter α , which is known as the degree of exploitation, can be adjusted to change the extent

of exploration. As the parameter α increases, the exploitation becomes more important. However, the classical TS algorithm does not take the length of horizon into account, and thus it results in the unnecessary exploration in a finite horizon. So, we pay attention to the performance in a finite horizon and try to find a optimal α structure in this section.

First, the MQR framework can utilize the characteristic of the parameter α to diagnose existed algorithms (e.g., UCB, TS). The UCB and TS algorithms are proved asymptotically optimal with respect to the expected regret in literature, but the performance in a finite horizon is different. By estimating an empirical α (e.g., $\hat{\alpha}$) from a data set which is generated by a specific algorithm, we are able to obtain some intuitions about how the algorithm make exploration-exploitation trade-off.

Second, inspired by the comparison result of estimated $\hat{\alpha}$, we employ the idea of adjusting the extent of exploration based on the time horizon T . We investigate the optimal α structure along T that corresponds to the optimal performance of the MQR policy. We compare the MQR policy that using the fitted α with the TS algorithm to see how the difference of rewards between two methods change as T increases.

We consider a two-armed bandit problem (i.e., $N = 2$). In our numerical experiments, rewards of arm i are generated by Bernoulli distributions with parameter μ_i , where

$$\mu_i \sim \text{Beta}(1, 1), \quad r_{i,t} \sim \text{Bernoulli}(\mu_i), \quad \forall i \in \{1, 2\}.$$

5.1 Estimating α from Realized Actions/Rewards

First, we introduce an estimation procedure to show how we get an estimated $\hat{\alpha}$ from a data set. The information contained in the data set is the action at each time $i(t)$, the accumulated rewards and accumulated times of playing each arm. We assume the noise term ϵ in the MQR framework is distributed as $\mathcal{N}(0, 1^2)$. The α value is estimated using the maximum likelihood technique. The log likelihood function to be maximized can be written as

$$\sum_t \sum_{i \in N} y_{ti} \log p_{ti},$$

where y_{ti} is the indicator function of whether arm i is played at time t and p_{ti} denotes the probability of pulling arm i at time t . The probability of playing arm i can be represented by information we know in the data set. For example,

$$p_{t1} = \Pr(\hat{\mu}_1(t) + \epsilon_1 \beta_1(t) \geq \hat{\mu}_2(t) + \epsilon_2 \beta_2(t)),$$

where $\beta_i(t) = 1/(k_i(t) + 1)^\alpha$. Then, we verify the accuracy of the estimation by comparing the estimated $\hat{\alpha}$ from the MQR-generated data set and true α . For each time horizon T , we generate $K = 1000$ samples to do estimation, and the log likelihood function is $\sum_{k=1}^K \sum_{t=1}^T \sum_{i \in N} y_{ti} \log p_{ti}$. Table 1 shows that our estimation works well when we test the MQR policies with different α values¹.

Table 1: Comparison of true α and estimated $\hat{\alpha}$

T	true α	estimated $\hat{\alpha}$
100	0.2	0.1935
1000	0.2	0.2013
10000	0.2	0.2013
100	0.5	0.4930
1000	0.5	0.4985
10000	0.5	0.4989
100	1	0.9568
1000	1	0.9848
10000	1	0.9926

¹All computations were carried out on MATLAB2016a on a personal computer with an Intel Core i7-8550U processor, which has four cores and 16.0 GB of RAM.

5.2 Diagnose a Given Policy: Are TS/UCB Over Exploring?

With the tool of estimation of α from a data set, we utilize it to make a diagnosis of the existed algorithms (i.e., UCB and TS) in a finite horizon. We try to investigate and understand the exploration-exploitation trade-off of these two methods in a relative short finite time horizon. Specifically, we generate data sets by the Bayesian optimal policy, TS and UCB, respectively. Then, we compare the estimated $\hat{\alpha}$ from these data sets and the regret incurred by using the estimated $\hat{\alpha}$. Since the Bayesian optimal policy achieves the best possible performance among all non-anticipating policies, the $\hat{\alpha}_{BO}$ can be viewed as an optimal benchmark.

Table 2: Comparison of estimated $\hat{\alpha}$ and optimality gap

T	BO		UCB		TS	
	$\hat{\alpha}_{BO}$	optimality gap	$\hat{\alpha}_{UCB}$	optimality gap	$\hat{\alpha}_{TS}$	
10	1.7237	0.51%	1.4119	5.42%	0.4574	
20	1.4701	0.81%	1.1649	6.30%	0.4620	
30	1.3614	1.04%	1.0261	6.37%	0.4618	
40	1.2968	0.99%	0.9783	6.09%	0.4751	
50	1.2567	0.88%	0.9677	5.79%	0.4822	
60	1.2246	0.91%	0.9204	5.67%	0.4850	
70	1.1893	0.85%	0.9121	5.46%	0.4822	
80	1.1829	0.82%	0.8936	5.23%	0.4881	
90	1.1543	0.83%	0.8734	5.14%	0.4881	
100	1.1543	0.78%	0.8684	4.94%	0.4896	

Table 2 shows the time horizon T and the corresponding estimated $\hat{\alpha}$ according to different kinds of data sets. We consider $T = \{10, 20, \dots, 100\}$ here, and there are 1000 samples of each algorithm for each T . From Table 2, we can see that the value of optimal $\hat{\alpha}$ (i.e., Bayesian optimal $\hat{\alpha}_{BO}$) decreases as T increases, and the decline is slowing down. It is intuitive since the larger T means that the exploration would be more valuable, and the degree of exploitation should reduce accordingly. As for the speed of decreasing, it is more sensitive to the change of time horizon when T is small.

Both of the UCB and TS algorithms are over exploration compared to $\hat{\alpha}_{BO}$. Among these two algorithms, the UCB algorithm performs better than the TS algorithm. It is due to that the UCB algorithm has an exploration term correlated with $\log T$, while the TS algorithm does not incorporate the length of the time horizon into consideration and keeps a constant extent of exploration. The UCB algorithm also has a trend of decreasing the extent of exploitation.

In passing, it is also worth noting from Table 2 that it is reasonable to adjust the value of α according to the length of the time horizon, which can improve the performance of the MQR algorithms significantly. This will be the focus on our next numerical experiment.

5.3 Learning a Better MAB Algorithm for Practically Sized T

Now that we are equipped with a tool to learn the coefficient of coefficient from any policy based on the realized actions/rewards. We proceed further to illustrate in a case study how we can “train” a sequence of MQR policies (each for a different time horizon T) with promising performances for practically sized T .

The training procedure works in the following steps.

1. For smaller values of $T \in \mathcal{T}_{train}$ (In the case study, we used $\mathcal{T}_{train} := \{10, 11, \dots, 150\}$), we compute the Bayesian optimal policies for each T . Note that this step could be notoriously intractable for larger T values but implementable for smaller T .
2. We generate simulated paths from the Bayesian optimal policy for each T , and estimate the MQR-implied α values from the Bayes optimal policies. Following notations from earlier sections, let us denote them by $\{\hat{\alpha}_{BO}(T) : T \in \mathcal{T}_{train}\}$.

3. We fit a curve $\{\alpha^*(T) = e^{aT^b+c} + 0.5 : T \in \mathbb{Z}_+\}$ from $\{\hat{\alpha}_{BO}(T) : T \in \mathcal{T}_{train}\}$ where a, b, c are parameters to be fitted.²

The output of our training procedure is therefore the MQR policy with the coefficient of exploitation $\alpha^*(T)$ for any given time horizon T . We note that although there have been many efforts in improving TS in the non-asymptotic regime (e.g., the information relaxation sampling, Min et al. (2019)), this new algorithm is simple to implement.

In the case study, we obtain a result of $\alpha^*(T) = e^{2.741T^{-0.2215}-1.436} + 0.5$. Figure 1.(a) plots the $\hat{\alpha}_{BO}$ and α given by the function $\alpha^*(T)$. To show the advantage of the MQR algorithm, we compare the regret performance of the MQR using fitted α^* with the TS algorithm. We select $T \in \{t : \log(t) = 4, 5, \dots, 9\}$ and calculate the difference of rewards between the MQR algorithm and the TS algorithm at these time T points. The MQR algorithm uses parameter α by the function $\alpha^*(T)$. To better compare the rewards of two algorithms, we use $\log(T)$ and $\log(\text{Difference of rewards})$. Note that $\text{Difference of rewards} \geq CT^\kappa$ implies that $\log(\text{Difference of rewards})$ should be approximately linear in $\log(T)$ with slope κ . Figure 1.(b) plots the relationship between the difference of rewards and time horizons in a log-log form. From the red linear trend line, it can see that there is a linear tendency in Figure 1.(b). Since the slope is in general close to 0.5, we view it as a promising basis that the new policy can be inspired to improve the *leading coefficient* of TS, even though TS is known to be asymptotically optimal with a regret order of $O(\sqrt{T})$.




Figure 1: Comparison with TS

6 Conclusion

We propose a *myopic quantal response* framework for stochastic multi-armed bandit (MAB) problems and develop a simple and intuitive class of policies. We leverage the framework to connect policies with behavioral economics and Thompson sampling, and it also provide an explicit way to understand the exploration-exploitation trade-off. Our policies can adjust the degree of exploitation in the finite time horizon and thus inspire a better MAB algorithm which brings significant improvement into the problem. In the future, the parameter of degree of exploitation has huge potential to be a useful structural estimation tool to diagnose a given policy whether it is "under" or "over" exploring. There are some limitations in our work, and we can view these as directions to follow-up. First, it would be interesting to apply the framework in behavioral experiments to investigate human beings' behavior. Second, more advanced methods can be adopted to discover the optimal structure of α when face more complex reward settings. Finally, we believe that the MQR framework proposed in this paper will lead to more research and deep understanding on MAB problems.

²In the case study, we used the standard Matlab function $fit(\cdot)$ to find the best values of a, b, c .

References

- Shipra Agrawal and Navin Goyal. 2013. Further optimal regret bounds for thompson sampling. In *Artificial intelligence and statistics*. PMLR, PMLR, Scottsdale, Arizona, USA, 99–107.
- Shipra Agrawal and Navin Goyal. 2017. Near-Optimal Regret Bounds for Thompson Sampling. *J. ACM* 64, 5, Article 30 (sep 2017), 24 pages. <https://doi.org/10.1145/3088510>
- R Michael Alvarez and John Brehm. 1995. American ambivalence towards abortion policy: Development of a heteroskedastic probit model of competing values. *American Journal of Political Science* (1995), 1055–1082.
- Peter Auer, Nicolo Cesa-Bianchi, and Paul Fischer. 2002. Finite-time analysis of the multiarmed bandit problem. *Machine learning* 47, 2 (2002), 235–256.
- Chandra R Bhat. 1995. A heteroscedastic extreme value model of intercity travel mode choice. *Transportation Research Part B: Methodological* 29, 6 (1995), 471–483.
- John Gittins, Kevin Glazebrook, and Richard Weber. 2011. *Multi-armed bandit allocation indices*. John Wiley & Sons, Hoboken, NJ.
- John C Gittins. 1979. Bandit processes and dynamic allocation indices. *Journal of the Royal Statistical Society: Series B (Methodological)* 41, 2 (1979), 148–164.
- Tze Leung Lai, Herbert Robbins, et al. 1985. Asymptotically efficient adaptive allocation rules. *Advances in applied mathematics* 6, 1 (1985), 4–22.
- R Duncan Luce. 1959. *Individual Choice Behavior: A Theoretical Analysis*. John Wiley, New York.
- Richard D McKelvey and Thomas R Palfrey. 1995. Quantal response equilibria for normal form games. *Games and economic behavior* 10, 1 (1995), 6–38.
- Richard D McKelvey and Thomas R Palfrey. 1998. Quantal response equilibria for extensive form games. *Experimental economics* 1, 1 (1998), 9–41.
- Seungki Min, Costis Maglaras, and Ciamac C Moallemi. 2019. Thompson sampling with information relaxation penalties. *Advances in Neural Information Processing Systems* 32 (2019).
- Daniel Russo and Benjamin Van Roy. 2014. Learning to optimize via posterior sampling. *Mathematics of Operations Research* 39, 4 (2014), 1221–1243.
- William R Thompson. 1933. On the likelihood that one unknown probability exceeds another in view of the evidence of two samples. *Biometrika* 25, 3-4 (1933), 285–294.

Appendices

A Proof of Theorem 3

A.1 Overview of the Proof

Before proceeding to the analysis, we first give an overview of the proof. Our proof of Theorem 3 has the similar structure as the proof of Theorem 1.3 of Agrawal and Goyal (2017), which provided the upper bound for TS with Gaussian priors. The main idea is to analyze the upper bound of $\mathbb{E}[k_i(T+1)]$. We follow their framework to decompose the probability of choosing arm i into three parts by using two events $E_i^\theta(t)$ and $E_i^\mu(t)$ (see Equation (A.1)). Then, our Lemma 6-9 correspond to Lemma 2.13-2.16 of Agrawal and Goyal (2017), respectively.

The main difference stems from the use of the noise term. Compared to their analysis of TS with Gaussian priors, we generalize the noise term, which only needs to satisfy Assumption 1, so we do not have properties of Gaussian variables in analysis. Also, we design a new form of quantities $\theta_i(t)$, which is different from the posterior distribution in TS. Therefore, we need to modify the proof by decomposing $\theta_i(t)$ and apply conditions in Assumption 1. To be specific, we use the concentration and anti-concentration properties of the noise term ϵ instead of those of Gaussian random variables in Agrawal and Goyal (2017). In Lemma 6, Equation (A.5) applies the anti-concentration property of ϵ and to derive tighter bound for larger j , the analysis utilizes condition (2) of Assumption 1 (Equation (A.11)). Lemma 9 chooses a different $L_i(T)$ and use Chebyshev's Inequality, so both of the analysis and result are presented in another way.

A.2 Analysis

First, we introduce some notations. We set two thresholds, x_i and y_i , such that $\mu_i < x_i < y_i < \mu_1$. Specifically, we choose $x_i = \mu_i + \frac{\Delta_i}{3}$ and $y_i = \mu^* - \frac{\Delta_i}{3}$. Define $p_{i,t}$ as the probability the $p_{i,t} := \Pr(\theta_1(t) > y_i \mid \mathcal{F}_{t-1})$. In the following analysis, we assume the first arm is the optimal arm. We define the event $E_i^\theta(t)$ as $\theta_i(t) \leq y_i$, and the event $E_i^\mu(t)$ as $\hat{\mu}_i(t) \leq x_i$. We decompose the expected number of plays of an arm $i \neq 1$ as follows.

$$\begin{aligned} \mathbb{E}[k_i(T+1)] &= \sum_{t=1}^T \Pr(i(t) = i) \\ &= \sum_{t=1}^T \Pr(i(t) = i, E_i^\mu(t), E_i^\theta(t)) + \sum_{t=1}^T \Pr(i(t) = i, E_i^\mu(t), \overline{E_i^\theta(t)}) \\ &\quad + \sum_{t=1}^T \Pr(i(t) = i, \overline{E_i^\mu(t)}). \end{aligned} \tag{A.1}$$

Next, we will bound each term of the above equation. The proof is similar to Lemma 2.13-2.16 of Agrawal and Goyal (2017). For the first term, since the derivation of Lemma 2.8 and the derivation of Equation (4) still holds, we have

$$\begin{aligned} \sum_{t=1}^T \Pr(i(t) = i, E_i^\mu(t), E_i^\theta(t)) &\leq \sum_{t=1}^T \mathbb{E} \left[\frac{(1-p_{i,t})}{p_{i,t}} I(i(t) = 1, E_i^\mu(t), E_i^\theta(t)) \right] \\ &\leq \sum_{k=0}^{T-1} \mathbb{E} \left[\frac{(1-p_{i,\tau_k+1})}{p_{i,\tau_k+1}} \right], \end{aligned} \tag{A.2}$$

where τ_k denotes the time step at which arm 1 is played for k^{th} time and let $\tau_0 = 0$. We prove the following lemma corresponding to Lemma 2.13.

Lemma 6. *Let τ_j denote the time step of the j^{th} play of the first arm. Then*

$$\mathbb{E} \left[\frac{1}{p_{i,\tau_j+1}} - 1 \right] \leq \begin{cases} e^{64} + 5, & \forall j \\ \frac{5}{T\Delta_i^2}, & j > L_i(T) \end{cases},$$

where $L_i(T) = \frac{288 \ln(T\Delta_i^2 + e^{32})}{\Delta_i^2}$.

Proof. Recall the definition that $p_{i,t}$ denotes the probability of $\theta_1(t) > y_i$ given \mathcal{F}_{t-1} , and $\theta_i(t) = \hat{\mu}_i(t) + \epsilon\beta_i(t)$, where ϵ is a random variable that satisfies Assumption 1. Given \mathcal{F}_{τ_j} , let Θ_j denote a random variable composed by $\hat{\mu}_1(\tau_j + 1) + \epsilon\beta_1(\tau_j + 1)$, where $\beta_1(\tau_j + 1) = 1/\sqrt{j+1}$. Let G_j be the geometric random variable denoting the number of trials until and including the trial where a sample $\Theta_j > y_i$. Then by the definition $p_{i,\tau_j+1} = \Pr(\Theta_j > y_i | \mathcal{F}_{\tau_j})$ and

$$\mathbb{E} \left[\frac{1}{p_{i,\tau_j+1}} \right] = \mathbb{E} [\mathbb{E} [G_j | \mathcal{F}_{\tau_j}]] = \mathbb{E} [G_j]. \quad (\text{A.3})$$

Next, we will bound the expected value of G_j for all j .

Consider any integer $r \geq 1$ and let $z = \sqrt{\ln r}$. Let \mathbf{MAX}_r be a random variable denoting the maximum of r independent samples of Θ_j . We abbreviate $\hat{\mu}_1(\tau_j + 1)$ to $\hat{\mu}_1$ in the following. Note that $\mathbb{E}[G_j] = \sum_{r=0}^{\infty} \Pr(G_j \geq r)$. Then, we consider $\Pr(G_j \leq r)$,

$$\begin{aligned} \Pr(G_j \leq r) &\geq \Pr(\mathbf{MAX}_r > y_i) \\ &\geq \Pr\left(\mathbf{MAX}_r > \hat{\mu}_1 + \frac{z}{\sqrt{j+1}} \geq y_i\right) \\ &= \mathbb{E} \left[\mathbb{E} \left[I\left(\mathbf{MAX}_r > \hat{\mu}_1 + \frac{z}{\sqrt{j+1}} \geq y_i\right) \mid \mathcal{F}_{\tau_j} \right] \right] \\ &= \mathbb{E} \left[I\left(\hat{\mu}_1 + \frac{z}{\sqrt{j+1}} \geq y_i\right) \Pr\left(\mathbf{MAX}_r > \hat{\mu}_1 + \frac{z}{\sqrt{j+1}} \mid \mathcal{F}_{\tau_j}\right) \right] \\ &= \Pr\left(\hat{\mu}_1 + \frac{z}{\sqrt{j+1}} \geq y_i\right) \Pr\left(\mathbf{MAX}_r > \hat{\mu}_1 + \frac{z}{\sqrt{j+1}}\right). \end{aligned} \quad (\text{A.4})$$

Due to the condition (1) of Assumption 1, we have the following inequation holds. For any $x \geq 0$,

$$\Pr(\epsilon > x) = 1 - F_\epsilon(x) \geq \frac{1}{\sqrt{2\pi}} \frac{x}{x^2 + 1} e^{-x^2/2}. \quad (\text{A.5})$$

Then, since Θ_j is a random variable that composed by $\hat{\mu}_1(\tau_j + 1) + \epsilon\beta_1(\tau_j + 1)$, this gives

$$\begin{aligned} \Pr\left(\mathbf{MAX}_r > \hat{\mu}_1 + \frac{z}{\sqrt{j+1}}\right) &\geq 1 - \left(1 - \frac{1}{\sqrt{2\pi}} \frac{z}{z^2 + 1} e^{-z^2/2}\right)^r \\ &\geq 1 - \left(1 - \frac{1}{\sqrt{2\pi}} \frac{\sqrt{\ln r}}{\ln r + 1} \frac{1}{\sqrt{r}}\right)^r \\ &\geq 1 - e^{-\frac{r}{\sqrt{4\pi r \ln r}}}. \end{aligned} \quad (\text{A.6})$$

Observe that for large r (in particular, for any $r \geq e^{12}$), $e^{-\sqrt{\frac{r}{4\pi \ln(r)}}} \leq \frac{1}{r^2}$. Therefore, for any $r \geq e^{12}$,

$$\Pr\left(\mathbf{MAX}_r > \hat{\mu}_1 + \frac{z}{\sqrt{j+1}} \mid \mathcal{F}_{\tau_j}\right) \geq 1 - \frac{1}{r^2}. \quad (\text{A.7})$$

Next, we bound the probability $\Pr\left(\hat{\mu}_1 + \frac{z}{\sqrt{j+1}} \geq y_i\right)$. Recall the definition that $\hat{\mu}_1(t) = \frac{\sum_{s=1}^{t-1} r_1(s)}{k_1(t)+1}$. Since it is a bit different from the average reward by dividing $k_1(t) + 1$ rather than $k_1(t)$, we add a

term $1/(j+1)$ to $\hat{\mu}_1$. Applying Hoeffding Inequality³ for $t = \tau_j + 1$ and $k_1(t) = j$:

$$\begin{aligned} \Pr\left(\hat{\mu}_1 + \frac{1}{j+1} + \frac{x}{\sqrt{j+1}} \geq \mu_1\right) &= \Pr\left(\hat{\mu}_1 + \frac{1}{j+1} \geq \mu_1 - \frac{x}{\sqrt{j+1}}\right) \\ &\geq \Pr\left(\frac{\sum_{s=1}^{\tau_j+1} r_1(s)}{j+1} \geq \mu_1 - \frac{x}{\sqrt{j+1}}\right) \\ &\geq 1 - e^{-2x^2}. \end{aligned}$$

Now, we use $x := z - \frac{1}{\sqrt{j+1}} \geq z - 1$ to get

$$\Pr\left(\hat{\mu}_1 + \frac{z}{\sqrt{j+1}} \geq \mu_1\right) \geq 1 - e^{-2(z-1)^2} \geq 1 - e^{-2z^2+4z} \geq 1 - \frac{1}{r^2} e^{4\sqrt{\ln(r)}}.$$

Using $y_i \leq \mu_1$, this gives

$$\Pr\left(\hat{\mu}_1 + \frac{z}{\sqrt{j+1}} \geq y_i\right) \geq 1 - \frac{1}{r^2} e^{4\sqrt{\ln(r)}}.$$

Observe that for large r (in particular, for any $r \geq e^{64}$), $\frac{1}{r^2} e^{4\sqrt{\ln(r)}} \leq \frac{1}{r^{1.5}}$. Therefore, for any $r \geq e^{64}$,

$$\Pr(G_j \leq r) \geq 1 - \frac{1}{r^2} - \frac{1}{r^{1.5}}. \quad (\text{A.8})$$

This gives,

$$\begin{aligned} \mathbb{E}[G_j] &= \sum_{r=0}^{\infty} \Pr(G_j \geq r) \\ &= 1 + \sum_{r=1}^{\infty} \Pr(G_j \geq r) \\ &\leq 1 + e^{64} + \sum_{r \geq e^{64}} \left(\frac{1}{r^2} + \frac{1}{r^{1.5}}\right) \\ &\leq 1 + e^{64} + 2 + 2.7. \end{aligned} \quad (\text{A.9})$$

This proves a constant bound on $\mathbb{E}\left[\frac{1}{p_{i,\tau_j+1}}\right] - 1 = \mathbb{E}[G_j] - 1 \leq e^{64} + 5$ for all j .

Next, we derive a tighter bound for large j . Consider $j > L_i(T)$, where $L_i(T) = \frac{288 \ln(T \Delta_i^2 + e^{32})}{\Delta_i^2}$.

For any $r \geq 1$, G_j , \mathbf{MAX}_r , and $z = \sqrt{\ln r}$ are defined as before. Then,

$$\begin{aligned} \Pr(G_j \leq r) &\geq \Pr(\mathbf{MAX}_r > y_i) \\ &\geq \Pr\left(\mathbf{MAX}_r > \hat{\mu}_1 + \frac{z}{\sqrt{j+1}} - \frac{\Delta_i}{6} \geq y_i\right) \\ &= \mathbb{E}\left[\mathbb{E}\left[I\left(\mathbf{MAX}_r > \hat{\mu}_1 + \frac{z}{\sqrt{j+1}} - \frac{\Delta_i}{6} \geq y_i\right) \mid \mathcal{F}_{\tau_j}\right]\right] \\ &= \mathbb{E}\left[I\left(\hat{\mu}_1 + \frac{z}{\sqrt{j+1}} + \frac{\Delta_i}{6} \geq \mu_1\right) \Pr\left(\mathbf{MAX}_r > \hat{\mu}_1 + \frac{z}{\sqrt{j+1}} - \frac{\Delta_i}{6} \mid \mathcal{F}_{\tau_j}\right)\right]. \end{aligned} \quad (\text{A.10})$$

³Hoeffding inequality: Let X_1, X_2, \dots, X_n be independent random variables wherein X_i has a finite support $[a_i, b_i]$. Define $\bar{X} = \frac{1}{n} \sum_{s=1}^n X_s$. Then for any $t > 0$,

$$\begin{aligned} \Pr(\bar{X} - E[\bar{X}] \geq t) &\leq \exp\left(-\frac{2n^2 t^2}{\sum_{s=1}^n (b_s - a_s)^2}\right), \\ \Pr(\bar{X} - E[\bar{X}] \leq -t) &\leq \exp\left(-\frac{2n^2 t^2}{\sum_{s=1}^n (b_s - a_s)^2}\right). \end{aligned}$$

where we used that $y_i = \mu_1 - \frac{\Delta_i}{3}$. Now, since $j + 1 \geq L_i(T) = \frac{288 \ln(T\Delta_i^2 + e^{32})}{\Delta_i^2}$,

$$2 \frac{\sqrt{2 \ln(T\Delta_i^2 + e^{32})}}{\sqrt{j+1}} \leq \frac{\Delta_i}{6}.$$

Therefore, for $r \leq (T\Delta_i^2 + e^{32})^2$, we get

$$\frac{z}{\sqrt{j+1}} - \frac{\Delta_i}{6} = \frac{\sqrt{\ln(r)}}{\sqrt{j+1}} - \frac{\Delta_i}{6} \leq -\frac{\Delta_i}{12}.$$

Let $x = \sqrt{2 \ln(T\Delta_i^2 + e^{32})}$, with condition (2) of Assumption 1 and $x \geq 8$, we have

$$F_\epsilon(-x) = F_\epsilon(-\sqrt{2 \ln(T\Delta_i^2 + e^{32})}) \leq \frac{1}{2(T\Delta_i^2 + e^{32})}.$$

Since $j + 1 \geq L_i(T) = \frac{288 \ln(T\Delta_i^2 + e^{32})}{\Delta_i^2}$, we get

$$\begin{aligned} F_\epsilon\left(-\sqrt{2 \ln(T\Delta_i^2 + e^{32})}\right) &= F_\epsilon\left(-\frac{\Delta_i}{12} \sqrt{L_i(T)}\right) \\ &\geq F_\epsilon\left(-\frac{\Delta_i}{12} \sqrt{j+1}\right). \end{aligned}$$

Combining these two inequations, we get

$$F_\epsilon\left(-\frac{\Delta_i}{12} \sqrt{j+1}\right) \leq F_\epsilon(-\sqrt{2 \ln(T\Delta_i^2 + e^{32})}) \leq \frac{1}{2(T\Delta_i^2 + e^{32})}.$$

Thus,

$$\Pr\left(\epsilon > -\frac{\Delta_i}{12\beta_1(\tau_j + 1)}\right) \geq 1 - \frac{1}{2(T\Delta_i^2 + e^{32})}.$$

By the definition of Θ_j , we have

$$\Pr(\Theta_j > \hat{\mu}_1 - \frac{\Delta_i}{12}) \geq 1 - \frac{1}{2(T\Delta_i^2 + e^{32})}. \quad (\text{A.11})$$

This implies

$$\Pr\left(\mathbf{MAX}_r > \hat{\mu}_1 + \frac{z}{\sqrt{j+1}} - \frac{\Delta_i}{6}\right) \geq 1 - \frac{1}{2^r(T\Delta_i^2 + e^{32})^r}.$$

Also, for any $t \geq \tau_j + 1$, with $k_1(t) \geq j$, we use Hoeffding Inequality,

$$\Pr\left(\hat{\mu}_1(t) \geq \mu_1 - \frac{\Delta_i}{6}\right) \geq 1 - e^{-2k_1(t)\Delta_i^2/36} \geq 1 - \frac{1}{(T\Delta_i^2 + e^{32})^{16}}.$$

Let $T' = (T\Delta_i^2 + e^{32})^2$. Then, for $1 \leq r \leq T'$,

$$\Pr(G_j \leq r) \geq 1 - \frac{1}{2^r(T')^{r/2}} - \frac{1}{(T')^8}.$$

When $r \geq T' \geq e^{64}$, using Equation (A.8), we obtain

$$\Pr(G_j \leq r) \geq 1 - \frac{1}{r^2} - \frac{1}{r^{1.5}}.$$

Combining these bounds,

$$\begin{aligned}
E[G_j] &\leq \sum_{r=0}^{\infty} \Pr(G_j \geq r) \\
&\leq 1 + \sum_{r=1}^{T'} \Pr(G_j \geq r) + \sum_{r=T'}^{\infty} \Pr(G_j \geq r) \\
&\leq 1 + \sum_{r=1}^{T'} \frac{1}{(2\sqrt{T'})^r} + \frac{1}{(T')^7} + \sum_{r=T'}^{\infty} \left(\frac{1}{r^2} + \frac{1}{r^{1.5}}\right) \tag{A.12} \\
&\leq 1 + \frac{1}{\sqrt{T'}} + \frac{1}{(T')^7} + \frac{2}{T'} + \frac{3}{\sqrt{T'}} \\
&\leq 1 + \frac{5}{T\Delta_i^2 + e^{32}}
\end{aligned}$$

Above gives an upper bound of $\mathbb{E}\left[\frac{1}{p^{i,\tau_j+1}}\right] - 1 = \mathbb{E}[G_j] - 1 \leq \frac{5}{T\Delta_i^2}$ for $j > L_i(T)$. \square

Substituting the bound from Lemma 6 into Equation (A.2), we obtain the following bound on the first term on the right side of Equation (A.1).

Lemma 7.

$$\sum_{t=1}^T \Pr(i(t) = i, E_i^\mu(t), E_i^\theta(t)) \leq \frac{288 \ln(T\Delta_i^2 + e^{32})}{\Delta_i^2} (e^{64} + 5) + \frac{5}{\Delta_i^2}.$$

The proof of Lemma 2.11 can be adapted to our sample methods. Here, we choose $x_i = \mu_i + \frac{\Delta_i}{3}$ and have the following lemma to bound the third term on the right side of Equation (A.1).

Lemma 8.

$$\sum_{t=1}^T \Pr(i(t) = i, \overline{E_i^\mu(t)}) \leq \frac{9}{\Delta_i^2} + 1.$$

Proof. Let τ_k denotes the time at which the k^{th} play of arm i . Note that $\tau_T \geq T$. Recall that $E_i^\mu(t)$ is the event that $\hat{\mu}_i(t) \leq x_i$. We have

$$\begin{aligned}
\sum_{t=1}^T \Pr(i(t) = i, \overline{E_i^\mu(t)}) &\leq \sum_{k=0}^{T-1} \Pr(\overline{E_i^\mu(\tau_{k+1})}) \\
&= \sum_{k=0}^{T-1} \Pr(\hat{\mu}_i(\tau_{k+1}) > x_i) \\
&\leq 1 + \sum_{k=1}^{T-1} \exp(-k(x_i - \mu_i)^2) \\
&\leq 1 + \frac{9}{\Delta_i^2}.
\end{aligned}$$

The first inequality holds since at time τ_{k+1} , we always have $i(t) = i$ and $\tau_T \geq T$. Note that $\hat{\mu}_i(\tau_{k+1})$ only depends on k rather than τ_{k+1} . Recall the definition of $\hat{\mu}_i(t)$, that $\hat{\mu}_i(\tau_{k+1}) = \frac{\sum_{s=0}^{k-1} r_i(\tau_{s+1})}{k+1}$. It is a bit different from the average reward by dividing $k+1$ rather than k . Using Hoeffding Inequality, we obtain $\Pr(\hat{\mu}_i(\tau_{k+1}) > x_i) \leq \Pr\left(\frac{\sum_{s=0}^{k-1} r_i(\tau_{s+1})}{k} > x_i\right) \leq e^{-k(x_i - \mu_i)^2}$. Thus, the second last inequality holds. \square

Corresponding to Lemma 2.12, we prove the following lemma obtain a bound on the second term of Equation (A.1).

Lemma 9. Suppose the distribution of ϵ has mean m and standard deviation σ .

$$\sum_{t=1}^T \Pr \left(i(t) = i, E_i^\mu(t), \overline{E_i^\theta(t)} \right) \leq L_i(T) + \frac{\sqrt{T}}{\Delta_i} \sigma^2,$$

where $L_i(T) = \frac{9(\sqrt{2}(T\Delta_i^2)^{1/4} + m)^2}{\Delta_i^2}$.

Proof. We decompose each summand into two parts depending on whether $k_i(t)$ is larger than $L_i(T)$ or not.

$$\begin{aligned} \sum_{t=1}^T \Pr \left(i(t) = i, E_i^\mu(t), \overline{E_i^\theta(t)} \right) &= \sum_{t=1}^T \Pr \left(i(t) = i, k_i(t) \leq L_i(T), E_i^\mu(t), \overline{E_i^\theta(t)} \right) \\ &\quad + \sum_{t=1}^T \Pr \left(i(t) = i, k_i(t) > L_i(T), E_i^\mu(t), \overline{E_i^\theta(t)} \right). \end{aligned} \quad (\text{A.13})$$

The first term of Equation (A.13) is trivially bounded by $L_i(T)$. We then bound the second term. Recall that $E_i^\theta(t)$ is the event that $\theta_i(t) \leq y_i$, and $E_i^\mu(t)$ is the event that $\hat{\mu}_i(t) \leq x_i$. Then,

$$\begin{aligned} \sum_{t=1}^T \Pr \left(i(t) = i, k_i(t) > L_i(T), E_i^\mu(t), \overline{E_i^\theta(t)} \right) &\leq \sum_{t=1}^T \Pr \left(k_i(t) > L_i(T), E_i^\mu(t), \overline{E_i^\theta(t)} \right) \\ &= \sum_{t=1}^T \Pr \left(k_i(t) > L_i(T), \hat{\mu}_i(t) \leq x_i, \theta_i(t) > y_i \right). \end{aligned}$$

And since $\theta_i(t) = \hat{\mu}_i(t) + \epsilon\beta_i(t)$, we have

$$\begin{aligned} \Pr \left(k_i(t) > L_i(T), \hat{\mu}_i(t) \leq x_i, \theta_i(t) > y_i \right) &= \Pr \left(\hat{\mu}_i(t) + \epsilon\beta_i(t) > y_i, k_i(t) > L_i(T), \hat{\mu}_i(t) \leq x_i \right) \\ &\leq \Pr \left(\epsilon\beta_i(t) > y_i - x_i, k_i(t) > L_i(T) \right) \\ &= \Pr \left(\epsilon > (y_i - x_i) \sqrt{k_i(t) + 1}, k_i(t) > L_i(T) \right) \\ &\leq \Pr \left(\epsilon > (y_i - x_i) \sqrt{L_i(t)} \right). \end{aligned}$$

Recall that $x_i = \mu_i + \frac{\Delta_i}{3}$, $y_i = \mu_i - \frac{\Delta_i}{3}$ and $L_i(T) = \frac{9(\sqrt{2}(T\Delta_i^2)^{1/4} + m)^2}{\Delta_i^2}$. Since ϵ is a random variable with mean m and standard deviation σ . Using Chebyshev's Inequality⁴, we have

$$\Pr \left(\epsilon > (y_i - x_i) \sqrt{L_i(t)} \right) \leq \frac{\sigma^2}{2\sqrt{T}\Delta_i} \leq \frac{\sigma^2}{\sqrt{T}\Delta_i}. \quad (\text{A.14})$$

Summing over $t = 1, \dots, T$,

$$\sum_{t=1}^T \Pr \left(i(t) = i, k_i(t) > L_i(T), E_i^\mu(t), \overline{E_i^\theta(t)} \right) \leq \frac{\sqrt{T}}{\Delta_i} \sigma^2.$$

The proof of Lemma 9 is completed. \square

Combining the bounds from Lemma 7-9 with Equation (A.1), we get

$$\mathbb{E}[k_i(t)] \leq \frac{288 \ln(T\Delta_i^2 + e^{32})}{\Delta_i^2} (e^{64} + 5) + \frac{5}{\Delta_i^2} + \frac{9(\sqrt{2}(T\Delta_i^2)^{1/4} + m)^2}{\Delta_i^2} + \frac{\sqrt{T}}{\Delta_i} \sigma^2 + \frac{9}{\Delta_i^2} + 1.$$

⁴Chebyshev's Inequality: Let X be a random variable with finite expected value μ and finite non-zero variance σ^2 . Then for any real number $k > 0$,

$$\Pr(|X - \mu| \geq k\sigma) \leq \frac{1}{k^2}.$$

Thus, the expected regret due to arm i is upper bounded by

$$\Delta_i \mathbb{E}[k_i(t)] \leq \frac{288 \ln(T\Delta_i^2 + e^{32})}{\Delta_i} (e^{64} + 5) + \frac{14}{\Delta_i} + \frac{9(\sqrt{2}(T\Delta_i^2)^{1/4} + m)^2}{\Delta_i} + \sqrt{T}\sigma^2 + \Delta_i.$$

The righthand side of the above inequation is decreasing in Δ_i for $\Delta_i \geq \frac{e}{\sqrt{T}}$ due to $m \geq 0$. Given fixed N , for every arm i with $\Delta_i \geq \frac{e}{\sqrt{T}}$, the expected regret is bounded by $O(\sqrt{T})$. For every arm i with $\Delta_i \leq \frac{e}{\sqrt{T}}$, the total regret is bounded by $O(\sqrt{T})$. This completes the proof of Theorem 3.

B Proof of Theorem 5

Without loss of generality, we assume the arm 1 is the optimal arm.

Lemma 10. *The number of playing the optimal arm $k_1(t)$ converges to infinity almost surely as $t \rightarrow \infty$.*

Proof. Let t_1, t_2, \dots, t_j denote the time at which the j^{th} pull of the arm 1 occurs. Define the process $\{\tau_j\}_{j \geq 1}$, where $\tau_j = t_j - t_{j-1}$ and $\tau_0 = 0$. Then, according to the selection rule of the MQR policy and recall that $\theta_i(t) = \hat{\mu}_i(t) + \epsilon_i^t \beta_i(t)$, we have the probability of pulling the arm 1 that

$$\begin{aligned} \Pr(i(t) = 1) &= \Pr(\theta_1(t) \geq \theta_i(t), \forall i \neq 1) \\ &\geq \Pr(\epsilon_1^t \beta_1(t) - \epsilon_i^t \beta_i(t) \geq 1, \forall i \neq 1). \end{aligned} \quad (\text{B.1})$$

The inequality holds due to $\hat{\mu}_i \in [0, 1]$ for any arm i .

We claim that $\tau_j < \infty$ almost surely for all $j \geq 1$. For $t \geq t_{j-1} + 1$, $\beta_1(t)$ is unchanged before the arm 1 has been pulled by j^{th} time. Recall that $\beta_i(t) = 1/(k_i(t) + 1)^\alpha$, so for each $i \neq 1$, $\beta_i(t)$ is non-increasing in t and $\beta_i(t) \leq 1$. Thus, $\Pr(\epsilon_i^t \beta_i(t) \leq c_j, \forall i \neq 1) \geq \Pr(\epsilon_i^t \leq c_j, \forall i \neq 1)$. By the property of ϵ , we can choose a constant $c_j > 0$ such that $\Pr(\epsilon_i^t \beta_i(t) \leq c_j, \forall i \neq 1) > 0$ for $t \geq t_{j-1} + 1$, where c_j depends on the distribution of ϵ . Then,

$$\begin{aligned} \Pr(i(t) = 1) &\geq \Pr(\epsilon_1^t \beta_1(t) - \epsilon_i^t \beta_i(t) \geq 1, \forall i \neq 1) \\ &\geq \Pr(\epsilon_1^t \beta_1(t) > 1 + c_j \text{ and } \epsilon_i^t \beta_i(t) \leq c_j, \forall i \neq 1) \\ &= \Pr(\epsilon_1^t \beta_1(t) > 1 + c_j) \Pr(\epsilon_i^t \beta_i(t) \leq c_j, \forall i \neq 1) \\ &> 0. \end{aligned} \quad (\text{B.2})$$

Since ϵ_i^t are independent for each arm i , given $\beta_i(t)$, we get the equality by independence. The last inequality holds as $\Pr(\epsilon_1^t \beta_1(t) > 1 + c_j) > 0$ by condition (1) of Assumption 1. Let

$$q_j = \min_{t \geq t_{j-1} + 1} \{\Pr(\epsilon_1^t \beta_1(t) - \epsilon_i^t \beta_i(t) \geq 1, \forall i \neq 1)\},$$

and we define a process $\{\tilde{\tau}_j\}_{j \geq 1}$, where $\tilde{\tau}_j$ is a geometric random variable with parameter q_j . By (B.2), we have $q_j > 0$ almost surely. Then, we get $\tilde{\tau}_j < \infty$ almost surely. We can see that both τ_j and $\tilde{\tau}_j$ represent the interarrival time between playing the arm 1. But the process $\{\tilde{\tau}_j\}$ always has a lower probability of pulling arm 1 than the process $\{\tau_j\}$. Hence, for any $j \geq 1$, we have $\Pr(\tilde{\tau}_j < \infty) \leq \Pr(\tau_j < \infty)$ and $\tilde{\tau}_j < \infty$ almost surely, and then $\tau_j < \infty$ almost surely for all $j \geq 1$.

Next, consider a counting process $N_T := \sup_j \{\tau_1 + \tau_2 + \dots + \tau_j \leq T\}$. We claim that $N_T \rightarrow \infty$ almost surely. Otherwise, with some positive probability, $N_T < \infty$ holds. Then, there exists a constant M such that $N_T \leq M$. But we can get that $\Pr\left(\bigcup_j^{M+1} (\tau_j = \infty)\right) \leq \sum_j^{M+1} \Pr(\tau_j = \infty) = 0$. That is, $\tau_1 + \tau_2 + \dots + \tau_{M+1} \leq T$, it contradicts with $N_T \leq M$. So, there does not exist a constant M , and $N_T \rightarrow \infty$ almost surely. We prove that the arm 1 will be played by infinite times almost surely. \square

Lemma 11. $\Pr(i(t) = 1) \rightarrow 1$ as $t \rightarrow \infty$.

Proof. We claim that for any arm $i \neq 1$, $\Pr(i(t) = i) \rightarrow 0$ as $t \rightarrow \infty$. Otherwise, for any $i \neq 1$, there exists $\delta > 0$ such that $\Pr(i(t) = i) > \delta$ for some t as $t \rightarrow \infty$. Note that the probability of choosing arm $i \neq 1$ only depends on the state $F_t = (k_1(t), k_2(t), \dots, k_N(t), \hat{\mu}_1(t), \hat{\mu}_2(t), \dots, \hat{\mu}_N(t))$. Let \mathcal{F}_t

denote that $\mathcal{F}_t = \{F_t \mid \Pr[i(t) = i \mid F_t] > \delta\}$. Thus, we have $\sum_{F_t \in \mathcal{F}_t} \Pr[i(t) = i \mid F_t] = \infty$. And then, we get $\sum_t \Pr(i(t) = i) = \sum_t \sum_{F_t \in \mathcal{F}_t} \Pr[i(t) = i \mid F_t] \Pr(F_t \in \mathcal{F}_t) = \infty$ almost surely. Also, the events $\{i(t) = i\}_{t=1}^\infty$ are independent conditional on F_t . By the second Borel–Cantelli lemma, it shows that the suboptimal arm i is pulled by infinite times almost surely. If the suboptimal arm i are pulled by infinite times as $t \rightarrow \infty$, by the law of large numbers, when $k_i \rightarrow \infty$, $\hat{\mu}_i \rightarrow \mu_i$. Similarly, by Lemma 10, we have $k_1 \rightarrow \infty$, then $\hat{\mu}_1 \rightarrow \mu_1$. Hence,

$$\begin{aligned} \lim_{t \rightarrow \infty} \Pr(i(t) = 1) &= \lim_{t \rightarrow \infty} \Pr(\hat{\mu}_1(t) + \epsilon_1 \beta_1(t) \geq \hat{\mu}_i(t) + \epsilon_i \beta_i(t), \forall i \neq 1) \\ &= \Pr(\mu_1 \geq \mu_i, \forall i \neq 1) = 1. \end{aligned}$$

It is contradicts with $\Pr(i(t) = i)$ does not converge to zero as $t \rightarrow \infty$, and thus the claim holds that for any $i \neq 1$, $\Pr(i(t) = i) \rightarrow 0$ as $t \rightarrow \infty$. We prove the lemma that $\Pr(i(t) = 1) \rightarrow 1$ as $t \rightarrow \infty$. \square

Therefore, the number of playing the suboptimal arms

$$\begin{aligned} \sum_{i=2}^N \mathbb{E}[k_i(T+1)] &= \sum_{i=2}^N \sum_{t=1}^T \Pr(i(t) = i) \\ &= \sum_{t=1}^T (1 - \Pr(i(t) = 1)) = o(T). \end{aligned}$$

The last equality is from Lemma 11. This implies that the MQR algorithms yield the regret $\mathbb{E}[R(T)] = o(T)$.

C Codes

To facilitate further studies, we provide the codes used in this research, which can be accessed via: https://www.dropbox.com/sh/isk0jzqa2kk6l8m/AAA1o5j8_kBsx4tnWl1KmFTFa?dl=0.