

## Stabilization-free HHO a posteriori error control

Fleurianne Bertrand ·  
Carsten Carstensen · Benedikt Gräßle ·  
Ngoc Tien Tran

**Abstract** The known a posteriori error analysis of hybrid high-order methods (HHO) treats the stabilization contribution as part of the error and as part of the error estimator for an efficient and reliable error control. This paper circumvents the stabilization contribution on simplicial meshes and arrives at a stabilization-free error analysis with an explicit residual-based a posteriori error estimator for adaptive mesh-refining as well as an equilibrium-based guaranteed upper error bound (GUB). Numerical evidence in a Poisson model problem supports that the GUB leads to realistic upper bounds for the displacement error in the piecewise energy norm. The adaptive mesh-refining algorithm associated to the explicit residual-based a posteriori error estimator recovers the optimal convergence rates in computational benchmarks.

**Keywords** hybrid high-order · a posteriori · guaranteed upper error bounds · adaptive mesh refinement · equilibration · stabilization-free · computational comparisons

---

This work has been supported by the Deutsche Forschungsgemeinschaft (DFG) in the Priority Program 1748 *Reliable simulation techniques in solid mechanics. Development of non-standard discretization methods, mechanical and mathematical analysis* under the projects BE 6511/1-1 and CA 151/22-2 as well as the European Union's Horizon 2020 research and innovation programme (Grant agreement No. 891734). The third author is also supported by the Berlin Mathematical School.

---

Fleurianne Bertrand  
University of Twente  
E-mail: f.bertrand@utwente.nl

Carsten Carstensen · Benedikt Gräßle  
Humboldt-Universität zu Berlin, Germany  
E-mail: cc, graesslb@math.hu-berlin.de

Tran Ngoc Tien  
Friedrich-Schiller Universität Jena  
E-mail: ngoc.tien.tran@uni-jena.de

## 1 Introduction

Hybrid high-order methods (HHO) were introduced in [25,26] and are examined in the textbooks [24,28] as a promising class of flexible nonconforming discretization methods for partial differential equations that involve a parameter-free stabilization term for the link between the volume and skeletal variables.

### 1.1 Known a posteriori error estimator

The a priori error analysis of HHO involves the stability terms in extended norms as part of the methodology and motivated a first explicit residual-based a posteriori error estimator in [24] with a reformulation of the stabilization in the upper bound. Let  $s_h(u_h, u_h)$  denote the stabilization at the discrete solution  $u_h \in V_h$  and let the (elliptic) reconstruction  $Ru_h$  of  $u_h$  denote a piecewise polynomial of degree at most  $k+1$  that approximates  $u \in H^1(\Omega)$ , cf. (2) and Section 3 below for further details. Then a possible error term reads

$$\text{total error}^2 := \|\nabla_{\text{pw}}(u - Ru_h)\|_{L^2(\Omega)}^2 + s_h(u_h, u_h). \quad (1)$$

It is disputable if  $s_h(u_h, u_h) \geq 0$  is an error contribution, but if the total error includes  $s_h(u_h, u_h)$  (or an equivalent form), then the error estimator may also include this term (or a computable equivalent) for a reliable and efficient a posteriori error control. Amongst the many skeletal schemes like (nonconforming) virtual elements, hybridized (weak) discontinuous Galerkin schemes et al., the HHO methodology has a clear and efficacious stabilization

$$s_h(v_h, w_h) := \sum_{T \in \mathcal{T}} \sum_{F \in \mathcal{F}(T)} h_F^{-1} \langle S_{TF} v_h, S_{TF} w_h \rangle_{L^2(F)} \quad (2)$$

with the abbreviation  $S_{TF} v_h := \Pi_{F,k}(v_{\mathcal{T}} + (1 - \Pi_{T,k})Rv_h)|_T - v_{\mathcal{F}}|_F$  for  $v_h = (v_{\mathcal{T}}, v_{\mathcal{F}}) \in V_h$  in terms of the  $L^2$  projections  $\Pi_{K,k}$  onto polynomials of degree at most  $k$  on a facet or simplex  $K \in \mathcal{F} \cup \mathcal{T}$  of diameter  $h_K = \text{diam}(K)$ ; cf. Subsection 1.4 for further details. The original residual-based estimator  $\eta_{\text{HHO}}$  from the textbook [24] for the Poisson model problem  $-\Delta u = f$  includes (2) and an interpolation  $\mathcal{A}Ru_h \in V$  of  $Ru_h$  by nodal averaging in

$$\begin{aligned} \eta_{\text{HHO}}^2 &= \|h_{\mathcal{T}}(1 - \Pi_0)(f + \Delta_{\text{pw}}Ru_h)\|_{L^2(\Omega)}^2 + \|\nabla_{\text{pw}}(1 - \mathcal{A})Ru_h\|_{L^2(\Omega)}^2 \\ &\quad + s_h(u_h, u_h). \end{aligned}$$

(Multiplicative constants are undisplayed in this introduction for simplicity.) The results from Theorem 4.3 and 4.7 in [24] show reliability and efficiency for the total error (1) and piecewise polynomial source terms  $f \in P_{k+1}(\mathcal{T})$ ,

$$\text{total error}^2 \approx \eta_{\text{HHO}}^2.$$

## 1.2 Stabilization-free a posteriori error control

There are objections against the double role of  $s_h(u_h, u_h)$  on both sides of the efficiency and reliability estimate. First, the term  $s_h(u, u_h)$  may dominate both sides of the error estimate. In other words, the total error might be equivalent to  $s_h(u_h, u_h)$ , but the quantity of interest may exclusively be

$$\text{error}^2 := \|\nabla_{\text{pw}}(u - Ru_h)\|_{L^2(\Omega)}^2.$$

Second, since the stabilization (2) incorporates a negative power of the mesh-size, a reduction property for local refinements remains unclear but is inevitable in the proofs of optimal convergence of an adaptive algorithm [5, 16]. This paper, therefore, asks a different question about the control of the error without the stabilization term (2) in the upper bound and introduces two stabilization-free error estimators (multiplicative constants are undisplayed)

$$\begin{aligned} \eta_{\text{res}}^2 &= \|h_{\mathcal{T}}(f + \Delta_{\text{pw}}Ru_h)\|_{L^2(\Omega)}^2 + \sum_{F \in \mathcal{F}} h_F \|\llbracket \nabla_{\text{pw}}Ru_h \rrbracket_F\|_{L^2(F)}^2, \\ \eta_{\text{eq},p}^2 &= \text{osc}_{k+p}^2(f, \mathcal{T}) + \|Q_p - \nabla_{\text{pw}}Ru_h\|_{L^2(\Omega)}^2 + \|\nabla_{\text{pw}}(1 - \mathcal{A})Ru_h\|_{L^2(\Omega)}^2 \end{aligned}$$

for some parameter  $p \in \mathbb{N}_0$ . The explicit residual-based a posteriori error estimator  $\eta_{\text{res}}$  follows from the a posteriori methodology in the spirit of [14, 20, 18, 21] with a piecewise volume residual  $f + \Delta_{\text{pw}}Ru_h$  and the jumps  $\llbracket \nabla_{\text{pw}}Ru_h \rrbracket_F$  across a facet  $F$  (on the boundary this is only the tangential component of  $\nabla_{\text{pw}}Ru_h$ ). The equilibrated error estimator  $\eta_{\text{eq},p}$  includes the post-processed quantity  $Q_p \in RT_{k+p}(\mathcal{T})$  in the space  $RT_{k+p}(\mathcal{T})$  of Raviart-Thomas functions of degree  $k + p$  for  $p \in \mathbb{N}_0$  and the nodal average  $\mathcal{A}Ru_h \in S_0^{k+1}(\mathcal{T})$  of  $Ru_h$ . The main results establish reliability and efficiency

$$\text{error}^2 + \text{osc}_{k-1}^2(f, \mathcal{T}) \lesssim \eta_{\text{res}}^2 \approx \eta_{\text{eq},p}^2 \lesssim \text{error}^2 + \text{osc}_q^2(f, \mathcal{T})$$

for any  $p, q \in \mathbb{N}_0$  up to data-oscillations  $\text{osc}_q^2(f, \mathcal{T}) := \|h_{\mathcal{T}}(1 - \Pi_q)f\|_{L^2(\Omega)}^2$  of the source term  $f \in L^2(\Omega)$  and without any stabilization terms. Computational benchmarks with adaptive mesh-refinement driven by any of these estimators provide numerical evidence for optimal convergence rates.

## 1.3 Further contributions and outline

The higher-order Crouzeix-Raviart finite element schemes are complicated at least in 3D [23] and then the HHO methodology is an attractive alternative even for simplicial triangulations with partly unexpected advantages like the computation of higher-order guaranteed eigenvalue bounds [15]. Higher convergence rates rely on an appropriate adaptive mesh-refining algorithm and hence stabilization-free a posteriori error estimators are of particular interest. The recent paper [34] establishes the latter for virtual elements with an over-penalization strategy as an extension of [8] for the discontinuous Galerkin

$\omega_{\max}$	$\pi$	$3\pi/2$	$2\pi$
$M_{\text{bd}}$	4	6	8
$c_{\text{apx}}$	2.9568	6.4642	11.3771
$C_{\text{st}}$	26.0893	55.8498	97.5374
$C_1$	2.9718	6.4710	11.3810
$C_2$	7.0495	15.2341	26.7317

**Table 1** Explicit constants  $C_1, C_{\text{st}}, C_2$  for right-isosceles triangles with respect to the maximum interior angle  $\omega_{\max}$  of the polygonal domain  $\Omega$ .

schemes. A disadvantage is the quantification of the restriction on the stabilization parameter in practise and poor condition for larger parameters. The stabilization-free a posteriori error control in this paper is based on two observations for the HHO schemes on simplicial triangulations. First, the  $P_1$ -conforming finite element functions let the stabilization vanish and, second, the divergence-free lowest-order Raviart-Thomas functions are  $L^2$  perpendicular to the piecewise gradients  $\nabla_{\text{pw}} Ru_h$ . In fact, those two fairly general properties lead in Section 2 to a reliable explicit residual-based a posteriori error estimator. In contrast to the simplified introduction above, the paper also focuses on multiplicative constants that lead to the GUB

$$\text{error} \leq \eta_{\text{res}} \quad \text{and} \quad \text{error} \leq \eta_{\text{eq},p};$$

cf. Table 1 for explicit quantities and Theorem 1 and Theorem 3 for further details. Numerical comparisons of  $\eta_{\text{HHO}}$  with  $\eta_{\text{res}}$  and  $\eta_{\text{eq},p}$  favour the latter. Section 2 identifies general building blocks of the a posteriori error analysis for discontinuous schemes with emphasis on explicit constants. An application to HHO leads to the new stabilization-free residual-based estimator  $\eta_{\text{res}}$  in Section 3. The alternative stabilization-free error estimator  $\eta_{\text{eq},p}$  follows from an equilibration strategy plus post-processing in Section 4. This paper also contributes to the HHO literature a local equivalence of two stabilizations and the efficiency of the stabilization terms up to data-oscillations in extension of [31]. Numerical comparisons of the different error estimators and an error estimator competition for guaranteed error control of the piecewise energy norm in 2D conclude this paper in Section 5. Three computational benchmarks provide striking numerical evidences for the optimality of the associated adaptive algorithms. The appendix provides algorithmic details on the computation of the post-processed contribution  $\|Q_p - \nabla_{\text{pw}} Ru_h\|_{L^2(\Omega)}$  in  $\eta_{\text{eq},p}$ .

#### 1.4 Overall notation

Standard notation for Sobolev and Lebesgue spaces and norms apply with  $\|\bullet\| := \|\bullet\|_{L^2(\Omega)}$  and  $\|\nabla\bullet\| := \|\nabla\bullet\|_{L^2(\Omega)}$ . In particular,  $H(\text{div}, \Omega)$  is the space of Sobolev functions with weak divergence in  $L^2(\Omega)$  and  $H(\text{div} = 0, \Omega)$  contains only divergence-free functions in  $H(\text{div}, \Omega)$ . Throughout this paper,  $\mathcal{T}$  denotes a shape-regular triangulation of the polyhedral bounded Lipschitz domain  $\Omega \subset \mathbb{R}^n$  into  $n$ -simplices with facets  $\mathcal{F}$  (edges for  $n = 2$  and faces for

$n = 3$ ) and vertices  $\mathcal{V}$ . Let  $\mathcal{F}(\Omega)$  (resp.  $\mathcal{V}(\Omega)$ ) denote the set of interior facets (resp. vertices). Given  $v \in \Omega \rightarrow \mathbb{R}^n$  and  $w : \Omega \rightarrow \mathbb{R}^{2n-3}$ , let  $\text{curl } v := \partial_1 v_2 - \partial_2 v_1$  and  $\text{Curl } w := (\partial_2 w, -\partial_1 w)^t$  if  $n = 2$  and  $\text{curl } v := (\partial_2 v_3 - \partial_3 v_2, \partial_3 v_1 - \partial_1 v_3, \partial_1 v_2 - \partial_2 v_1)^t$  and  $\text{Curl } w := \text{curl } w$  if  $n = 3$ . For  $s \in \mathbb{R}$ , let  $H^s(\mathcal{T})$ ,  $H(\text{div}, \mathcal{T})$ , and  $H(\text{curl}, \mathcal{T})$  denote the space of piecewise Sobolev functions with restriction to  $T \in \mathcal{T}$  in  $H^s(T)$ ,  $H(\text{div}, T)$ , and  $H(\text{curl}, T)$ . To simplify notation,  $H^s(K)$  abbreviates  $H^s(\text{int}(K))$  for the open interior  $\text{int}(K)$  of a compact set  $K$ . The  $L^2$ -scalar product reads  $(\bullet, \bullet)_{L^2(\omega)}$  for volumes  $\omega \subseteq \Omega$  and  $\langle \bullet, \bullet \rangle_{L^2(\gamma)}$  for surfaces  $\gamma \subset \overline{\Omega}$  of co-dimension one; the same symbol applies to scalars and to vectors. For  $V := H_0^1(\Omega)$  and  $V^* := H^{-1}(\Omega)$ , let  $\langle \bullet, \bullet \rangle$  denote the duality-brackets in  $V^* \times V$  for the dual space  $V^*$  of  $V$  equipped with the operator norm  $\|F\|_* := \sup_{v \in V \setminus \{0\}} |Fv|/\|v\|$  for  $F \in V^*$ .

Define the energy scalar product  $a(v, w) := (\nabla v, \nabla w)_{L^2(\Omega)}$  for  $v, w \in H^1(\Omega)$  and its piecewise version  $a_{\text{pw}}(v, w) = (\nabla_{\text{pw}} v, \nabla_{\text{pw}} w)_{L^2(\Omega)}$  for  $v, w \in H^1(\mathcal{T})$ . The latter induces the piecewise energy norm  $\|\bullet\|_{\text{pw}} := a_{\text{pw}}(\bullet, \bullet)^{1/2}$  in  $H^1(\mathcal{T})$ . Here and throughout the paper,  $\nabla_{\text{pw}}$ ,  $\text{div}_{\text{pw}}$ ,  $\text{curl}_{\text{pw}}$ ,  $\Delta_{\text{pw}}$ , denote the piecewise evaluation of the differential operators  $\nabla$ ,  $\text{div}$ ,  $\text{curl}_{\text{pw}}$ ,  $\Delta$  without explicit reference to the underlying shape-regular triangulation  $\mathcal{T}$ .

The vector space  $P_k(K)$  of polynomials of degrees at most  $k \in \mathbb{N}_0$  over a facet or simplex  $K \in \mathcal{F} \cup \mathcal{T}$  defines the piecewise polynomial spaces

$$\begin{aligned} P_k(\mathcal{T}) &:= \{p \in L^2(\Omega) : p|_T \in P_k(T) \text{ for all } T \in \mathcal{T}\}, \\ P_k(\mathcal{F}) &:= \{p \in L^2(\mathcal{F}) : p|_F \in P_k(F) \text{ for all } F \in \mathcal{F}\} \end{aligned}$$

and the space of piecewise Raviart-Thomas functions

$$RT_k^{\text{pw}}(\mathcal{T}) := P_k(\mathcal{T}; \mathbb{R}^n) + xP_k(\mathcal{T}).$$

The associated  $L^2$  projections read  $\Pi_{K,k} : L^2(\Omega) \rightarrow P_k(K)$ ,  $\Pi_k : L^2(\Omega) \rightarrow P_k(\mathcal{T})$ , and  $\Pi_{\mathcal{F},k} : L^2(\Omega) \rightarrow P_k(\mathcal{F})$  with the convention  $\Pi_{-1} := 0$ . Abbreviate  $S_0^{k+1}(\mathcal{T}) := P_{k+1}(\mathcal{T}) \cap V$  and  $RT_k(\mathcal{T}) := RT_k^{\text{pw}}(\mathcal{T}) \cap H(\text{div}, \Omega)$  for all  $k \in \mathbb{N}_0$ . The piecewise constant mesh-size function  $h_{\mathcal{T}} \in P_0(\mathcal{T})$  satisfies  $h_{\mathcal{T}|T} := h_T$  for  $T \in \mathcal{T}$  with the diameter  $h_K := \text{diam}(K) \in P_0(K)$  of  $K \in \mathcal{F} \cup \mathcal{T}$ .

If not explicitly stated otherwise, constants are independent of the mesh-size in the triangulation but may depend on the shape-regularity and on the polynomial degree  $k$ . The abbreviation  $A \lesssim B$  hides a generic constant  $C$  (independent of the mesh-size) in  $A \leq C B$ ;  $A \approx B$  abbreviates  $A \lesssim B \lesssim A$ .

## 2 Foundations of the a posteriori error analysis

This section investigates general building blocks of the a posteriori error analysis and revisits arguments from [14, 20, 18, 21] with emphasis on multiply connected domains  $\Omega \subset \mathbb{R}^n$  for  $n = 2, 3$ . The general setting of this section results in reliability for an error estimator that is applicable beyond the HHO methodology. Consider the weak solution  $u \in V = H_0^1(\Omega)$  to the Poisson

model problem  $-\Delta u = f$  a.e. in  $\Omega$  for a given source  $f \in L^2(\Omega)$ ; i.e.,  $u \in V$  satisfies

$$a(u, v) = (f, v)_{L^2(\Omega)} \quad \text{for all } v \in V. \quad (3)$$

An approximation  $G \in L^2(\Omega; \mathbb{R}^n)$  of the gradient  $\nabla u \in H(\text{div}, \Omega)$  gives rise to the residual  $f + \text{div } G \in V^* = H^{-1}(\Omega)$  seen as a linear functional on  $V$ , i.e.,

$$\langle f + \text{div } G, \varphi \rangle := (f, \varphi)_{L^2(\Omega)} - (G, \nabla \varphi)_{L^2(\Omega)} \quad \text{for all } \varphi \in V.$$

Let  $\nu_T$  denote the unit outer normal along the boundary  $\partial T$  of each simplex  $T \in \mathcal{T}$  and fix the orientation of a unit normal  $\nu_F = \pm \nu_T$  for each facet  $F \in \mathcal{F}(T)$  of  $T$  such that it matches the outer unit normal  $\nu$  of  $\partial\Omega$  at the boundary. The jump  $[G]_F$  of a piecewise function in  $m \in \mathbb{N}$  components  $G \in H^1(\mathcal{T}; \mathbb{R}^m)$  reads  $[G]_F := G|_{T_+} - G|_{T_-}$  on interior facets  $F = T_+ \cap T_- \in \mathcal{F}(\Omega)$  (with  $T_\pm$  labelled such that  $\nu_{T_+|F} = \nu_F = -\nu_{T_-|F}$ ) and  $[G]_F := G$  on the boundary  $F \in \mathcal{F}(\partial\Omega)$ . The main result of this section establishes the residual-based error estimator

$$\begin{aligned} \eta^2(\mathcal{T}, G) := & \left( C_1 \|h_{\mathcal{T}}(f + \text{div}_{\text{pw}} G)\| + C_2 \sqrt{\sum_{F \in \mathcal{F}(\Omega)} \ell(F) \|[G]_F \cdot \nu_F\|_{L^2(F)}^2} \right)^2 \\ & + C_H^2 \left( C_1 \|h_{\mathcal{T}} \text{curl}_{\text{pw}} G\| + C_2 \sqrt{\sum_{F \in \mathcal{F}} \ell(F) \|[G]_F \times \nu_F\|_{L^2(F)}^2} \right)^2 \end{aligned} \quad (4)$$

as a GUB  $\|\nabla u - G\| \leq \eta(\mathcal{T}, G)$  under minimal assumptions on the approximation  $G \in H^1(\mathcal{T}; \mathbb{R}^n) \subset L^2(\Omega; \mathbb{R}^n)$ . The constants  $C_1$ ,  $C_2$ , and  $C_H$  (or upper bounds thereof) are computable; cf. Table 1 for an example in 2D with details in Example 1 at the end of Section 2. The first assumption is a weakened discrete solution property

$$(G, \nabla w_C)_{L^2(\Omega)} = (f, w_C)_{L^2(\Omega)} \quad \text{for all } w_C \in S_0^1(\mathcal{T}). \quad (5)$$

The second assumption is the orthogonality to the lowest-order divergence-free Raviart-Thomas functions

$$(G, r)_{L^2(\Omega)} = 0 \quad \text{for all } r \in RT_0(\mathcal{T}) \cap H(\text{div} = 0, \Omega). \quad (6)$$

**Theorem 1 (residual-based GUB)** *Suppose that  $G \in H^1(\mathcal{T}; \mathbb{R}^n)$  and  $f \in L^2(\Omega)$  satisfy (5)–(6). Then the error estimator  $\eta(\mathcal{T}, G)$  from (4) is a GUB*

$$\|\nabla u - G\| \leq \eta(\mathcal{T}, G)$$

*of the error  $\|\nabla u - G\|$  for the solution  $u \in V$  to (3). The constants  $C_1, C_2, C_H$  exclusively depend on  $\Omega$  and the shape-regularity of  $\mathcal{T}$ .*

The remaining parts of this section are devoted to the proof of Theorem 1 and the computation of (upper bounds of) the constants  $C_1$ ,  $C_2$ , and  $C_H$  in (4). The point of departure is the subsequent decomposition that appears necessary in the nonconforming and mixed finite element a posteriori error analysis. It leads to a split of the error  $\|\nabla u - G\|$  into some divergence part and some consistency part.

**Lemma 1 (decomposition)** *Any  $v \in V$  and  $G \in L^2(\Omega; \mathbb{R}^n)$  satisfy the decomposition*

$$\|\nabla v - G\|^2 = \|v - w\|^2 + \|G - \nabla w\|^2 \quad (7)$$

with the (unique) minimizer  $w \in V$  of the distance

$$\delta := \min_{\varphi \in V} \|G - \nabla \varphi\|$$

of  $G$  to the gradients  $\nabla V$  of Sobolev functions. The solution  $u \in V$  to (3) satisfies

$$\begin{aligned} \mu &:= \|f + \operatorname{div} G\|_* = \|u - w\| && \text{and} \\ \|\nabla u - G\|^2 &= \|f + \operatorname{div} G\|_*^2 + \|G - \nabla w\|^2 = \mu^2 + \delta^2. \end{aligned} \quad (8)$$

*Proof* The minimizer  $w \in V$  of  $\|G - \nabla \varphi\|$  among  $\varphi \in V$  satisfies the variational formulation  $a(w, \varphi)_{L^2(\Omega)} = (G, \nabla \varphi)_{L^2(\Omega)}$  for all  $\varphi \in V$ . (Notice that  $w$  is the unique weak solution to the Poisson model problem  $-\Delta w = -\operatorname{div} G \in V^*$ .) In particular,  $G - \nabla w$  is  $L^2$  orthogonal onto  $\nabla V$  and the Pythagoras theorem proves (7). Given  $\varphi \in V$  with  $\|\varphi\| = 1$ , the orthogonality of  $G - \nabla w$  to  $\nabla \varphi$  and (3) show

$$a(u - w, \varphi)_{L^2(\Omega)} = (\nabla u - G, \nabla \varphi)_{L^2(\Omega)} = \langle f + \operatorname{div} G, \varphi \rangle \quad (9)$$

with the duality brackets  $\langle \bullet, \bullet \rangle$  in  $V^* \times V$ . Since the supremum of (9) over all  $\varphi \in V$  with  $\|\varphi\| = 1$  is equal to  $\|u - w\| = \|f + \operatorname{div} G\|_*$ , this and (7) conclude the proof of (8).  $\square$

The split (7) of the error  $\|\nabla u - G\|$  allows for and enforces a separate estimation of the equilibrium and consistency contribution in residual-based a posteriori error estimators.

In order to derive explicit constants, two lemmas are recalled. The first has a long tradition in the a posteriori error control in form of a Helmholtz decomposition on simply connected domains [13, 4] and introduces the constant  $C_H$  from Theorem 1. The following version includes the general case of multiply connected domains as in [32] for  $n = 2$  or  $n = 3$  dimensions and weak assumptions on a divergence-free function  $\varrho \in H(\operatorname{div} = 0, \Omega)$ .

**Lemma 2 (Helmholtz-decomposition)** *Suppose the divergence-free function  $\varrho \in H(\operatorname{div} = 0, \Omega)$  is  $L^2$  orthogonal onto  $RT_0(\mathcal{T}) \cap H(\operatorname{div} = 0, \Omega)$ . Then there exists  $\beta \in H^1(\Omega; \mathbb{R}^N)$ ,  $N = 2n - 3$ , such that any  $\beta_C \in S^1(\mathcal{T})^N$  satisfies*

$$\|\varrho\|^2 = \int_{\Omega} \varrho \cdot \operatorname{Curl}(\beta - \beta_C) \, dx \quad \text{and} \quad \|\beta\| \leq C_H \|\varrho\|. \quad (10)$$

The constant  $C_H > 0$  exclusively depends on  $\Omega$ .

*Proof* The compact polyhedral boundary  $\partial\Omega$  of the bounded Lipschitz domain  $\Omega$  has  $J+1$  connectivity components  $\Gamma_0, \dots, \Gamma_J$  for some finite  $J \in \mathbb{N}_0$ . Those connectivity components have a positive surface measure  $|\Gamma_j|$  and a positive distance of each other. So the integral mean

$$\gamma_j := \int_{\Gamma_j} \varrho \cdot \nu \, ds / |\Gamma_j|$$

is well defined and depends continuously on  $\varrho \in H(\operatorname{div} = 0, \Omega)$  in the sense that  $|\gamma_j| \leq c_1 \|\varrho\|$  (recall  $\operatorname{div} \varrho = 0$ ) for each  $j = 0, \dots, J$  and  $c_1 > 0$ . This constant  $c_1$  and the constants  $c_2, c_3, c_4$  below exclusively depend on the domain  $\Omega$ . The finite real numbers  $\gamma_0, \dots, \gamma_J$  define the Neumann data for the harmonic function  $z \in H^1(\Omega)/\mathbb{R}$  with

$$\Delta z = 0 \text{ in } \Omega \quad \text{and} \quad \partial z / \partial \nu = \gamma_j \text{ on } \Gamma_j \text{ for all } j = 0, \dots, J.$$

The elliptic regularity theory for polyhedral domains lead to  $z \in H^{1+\alpha}(\Omega)$  for some  $\alpha > 1/2$  and  $c_2 > 0$  with  $\|z\|_{H^{1+\alpha}(\Omega)} \leq c_2 (|\gamma_0| + \dots + |\gamma_J|)$ . The Fortin interpolation operator defines a bounded linear operator on  $H(\operatorname{div}, \Omega) \cap L^p(\Omega; \mathbb{R}^n)$  for  $p > 2$ . It is generally accepted that, for  $\alpha > 0$  and  $\nabla z \in H(\operatorname{div} = 0, \Omega) \cap H^\alpha(\Omega; \mathbb{R}^n)$ , the Fortin interpolation  $I_F \nabla z \in RT_0(\mathcal{T}) \cap H(\operatorname{div} = 0, \Omega)$  is well defined and  $\|I_F \nabla z\| \leq c_3 \|\nabla z\|_{H^\alpha(\Omega)}$  follows for some  $c_3 > 0$ . The additional property  $\nabla z \in L^p(\Omega; \mathbb{R}^n)$  for some  $p > 2$  allows the definition of  $\int_F \nabla z \cdot \nu_F \, ds$  as a Lebesgue integral over a facet  $F \in \mathcal{F}$ . One consequence for the boundary facets is the vanishing integral

$$\int_{\Gamma_j} (\varrho - I_F \nabla z) \cdot \nu \, ds = 0 \quad \text{for all } j = 0, \dots, J.$$

Since  $\varrho - I_F \nabla z \in H(\operatorname{div} = 0, \Omega)$  is divergence-free, Theorems 3.1 and 3.4 in [32] prove the existence of  $c_4 > 0$  and  $\beta \in H^1(\Omega; \mathbb{R}^N)$  with

$$\varrho = I_F \nabla z + \operatorname{Curl} \beta \quad \text{and} \quad \|\beta\| \leq c_4 \|\varrho - I_F \nabla z\|.$$

Recall that  $\varrho \perp I_F \nabla z$  and  $\varrho \perp \operatorname{curl} \beta_C \in RT_0(\mathcal{T}) \cap H(\operatorname{div} = 0, \Omega)$ . This concludes the proof of (10) with  $C_H := \sqrt{1 + (c_1 c_2 c_3 (1 + J))^2} c_4$ .  $\square$

The subsequent version of the trace inequality on the facets  $\mathcal{F}$  leads to the piecewise constant  $\ell \in P_0(\mathcal{F})$  defined by

$$\ell(F) := \begin{cases} (n+1) h_T^2 |F| / |T| & \text{for } F \in \mathcal{F}(\partial\Omega) \cap \mathcal{F}(T), \\ (n+1) |F| / (h_{T_+}^{-2} |T_+| + h_{T_-}^{-2} |T_-|) & \text{for } F = \partial T_+ \cap \partial T_- \in \mathcal{F}(\Omega). \end{cases}$$

**Lemma 3 (trace inequality)** *Any  $f \in H^1(\Omega)$  satisfies*

$$\sum_{F \in \mathcal{F}} \ell(F)^{-1} \|f\|_{L^2(F)}^2 \leq \|h_{\mathcal{T}}^{-1} f\|^2 + \frac{2C_{\operatorname{tr}}}{n} \|h_{\mathcal{T}}^{-1} f\| \|f\|$$

with the constant  $C_{\operatorname{tr}} := \max_{T \in \mathcal{T}} \max_{x \in T} |x - \operatorname{mid}(T)| / h_T < n / (n+1)$ .



*Proof* The center of inertia  $\text{mid}(T) = \sum_{j=0}^n P_j / (n+1)$  of the  $n$ -simplex  $T = \text{conv}\{P_0, \dots, P_n\} \in \mathcal{T}$  and the  $n+1$  faces  $F_j = \text{conv}\{P_0, \dots, P_{j-1}, P_{j+1}, \dots, P_n\} \in \mathcal{F}(T)$  for  $j = 0, \dots, n$  give rise to the decomposition of  $T$  into  $n+1$  sub-simplices  $T'_j = \text{conv}(F_j, \text{mid}(T))$  with volume  $|T'_j| = |T|/(n+1)$ .

Standard arguments like the trace identity on  $T'_j \subset T$  [17, Lemma 2.1] for  $|f|^2 \in W^{1,1}(T')$  and a Cauchy inequality show

$$\frac{1}{|F_j|} \|f\|_{L^2(F_j)}^2 \leq \frac{1}{|T'_j|} \|f\|_{L^2(T'_j)}^2 + \frac{2}{n|T'_j|} \max_{x \in T'_j} |x - \text{mid}(T)| \|f\|_{L^2(T'_j)} \|\nabla f\|_{L^2(T')}.$$

The distance  $\max_{x \in T'_j} |x - \text{mid}(T)| = |P_k - \text{mid}(T)| \leq C_{\text{tr}} h_T$  is attained at a vertex  $P_k$  for  $k \in \{0, \dots, j-1, j+1, \dots, n\}$ . Since the centroid  $\text{mid}(T)$  divides each median of  $T$  in the ratio  $n$  to  $1$  and the length of each median is strictly bounded by  $h_T$ , the bound  $C_{\text{tr}} < n/(n+1)$  follows and cannot be improved in the absence of further assumptions on the shape of the simplex  $T$ . Since  $|T'_j| = |T|/(n+1)$ , the previously displayed estimate leads to

$$\frac{|T|}{(n+1)h_T^2|F_j|} \|f\|_{L^2(F_j)}^2 \leq \|h_T^{-1}f\|_{L^2(T'_j)}^2 + \frac{2C_{\text{tr}}}{n} \|h_T^{-1}f\|_{L^2(T'_j)} \|\nabla f\|_{L^2(T'_j)}.$$

Let  $\mathcal{T}'$  be the refinement of  $\mathcal{T}$ , obtained by replacing  $T \in \mathcal{T}$  with  $T'_0, \dots, T'_d$  from above. The triangulation  $\mathcal{T}'$  allows for the facet based decomposition  $\{\omega'(F)\}_{F \in \mathcal{F}}$  of  $\Omega$ , where  $\omega'(F)$  is either the patch  $\omega'(F) = \text{int}(T'_+ \cup T'_-)$  for an interior facet  $F = T'_+ \cap T'_-$  or  $\omega'(F) = \text{int}(T')$  for  $F \in \mathcal{F}(\partial\Omega) \cap \mathcal{F}(T')$ . This establishes, for any  $F \in \mathcal{F}$ , the estimate

$$\ell(F)^{-1} \|f\|_{L^2(F)}^2 \leq \|h_{\mathcal{T}'}^{-1}f\|_{L^2(\omega'(F))}^2 + \frac{2C_{\text{tr}}}{n} \|h_{\mathcal{T}'}^{-1}f\|_{L^2(\omega'(F))} \|\nabla f\|_{L^2(\omega'(F))}.$$

Since the family  $\{\omega'(F) : F \in \mathcal{F}\}$  has no overlap, the sum of the last displayed inequality over all  $F \in \mathcal{F}$  and a Cauchy inequality conclude the proof of Lemma 3.  $\square$

The next lemma utilizes a quasi-interpolation operator  $J : H^1(\Omega) \rightarrow S^1(\mathcal{T})$  with the restriction  $J(V) \subset S_0^1(\mathcal{T})$ , e.g.,  $J = J_1 \circ I_{\text{NC}}$  with explicit constants for  $n = 2$  from [19], and the approximation and stability properties

$$\|h_{\mathcal{T}'}^{-1}(\varphi - J\varphi)\| \leq C_1 \|\varphi\| \quad \text{and} \quad \|\varphi - J\varphi\| \leq C_{\text{st}} \|\varphi\| \quad (11)$$

for constants  $C_1$  and  $C_{\text{st}}$  exclusively depending on the shape-regularity of  $\mathcal{T}$ . Recall the constant  $C_{\text{tr}}$  from Lemma 3 and set  $C_2 := (C_1(C_1 + 2C_{\text{tr}}C_{\text{st}}/n))^{1/2}$ .

**Lemma 4 (equilibrium)** *Suppose that  $G \in H(\text{div}, \mathcal{T})$  and  $f \in L^2(\Omega)$  satisfy (5) and suppose  $(G|_T)|_F \cdot \nu_F \in L^2(F)$  for all  $F \in \mathcal{F}(T)$  and  $T \in \mathcal{T}$ . Then*

$$\|f + \text{div} G\|_* \leq C_1 \|h_{\mathcal{T}}(f + \text{div}_{\text{pw}} G)\| + C_2 \sqrt{\sum_{F \in \mathcal{F}(\Omega)} \ell(F) \|[G]_F \cdot \nu_F\|_{L^2(F)}^2}.$$

*Proof* Given  $\varphi \in V$  with  $\|\varphi\| = 1$ , set  $\psi := \varphi - \varphi_C$  for some quasi-interpolation  $\varphi_C := J\varphi \in S_0^1(\mathcal{T})$  with (11). Since (5) implies  $\langle f + \operatorname{div} G, \varphi \rangle = \langle f + \operatorname{div} G, \psi \rangle$ , a piecewise integration by parts and the collection of jump contributions show

$$\langle f + \operatorname{div} G, \varphi \rangle = (f + \operatorname{div}_{\text{pw}} G, \psi)_{L^2(\Omega)} - \sum_{F \in \mathcal{F}(\Omega)} \langle [G]_F \cdot \nu_F, \psi \rangle_{L^2(F)}. \quad (12)$$

The first bound follows from a Cauchy inequality and (11),

$$\begin{aligned} (f + \operatorname{div}_{\text{pw}} G, \psi)_{L^2(\Omega)} &\leq \|h_{\mathcal{T}}(f + \operatorname{div}_{\text{pw}} G)\| \|h_{\mathcal{T}}^{-1} \psi\| \\ &\leq C_1 \|h_{\mathcal{T}}(f + \operatorname{div}_{\text{pw}} G)\| \|\varphi\|. \end{aligned} \quad (13)$$

The second bound additionally exploits the trace inequality of Lemma 3,

$$\begin{aligned} &\sum_{F \in \mathcal{F}(\Omega)} \langle [G]_F \cdot \nu_F, \psi \rangle_{L^2(F)} \\ &\leq \sqrt{\sum_{F \in \mathcal{F}(\Omega)} \ell(F) \| [G]_F \cdot \nu_F \|_{L^2(F)}^2} \sqrt{\sum_{F \in \mathcal{F}(\Omega)} \ell(F)^{-1} \|\psi\|_{L^2(F)}^2} \\ &\leq \sqrt{C_1^2 + \frac{2C_{\text{tr}}}{n} C_{\text{st}} C_1} \sqrt{\sum_{F \in \mathcal{F}(\Omega)} \ell(F) \| [G]_F \cdot \nu_F \|_{L^2(F)}^2} \end{aligned} \quad (14)$$

with  $\|\varphi\| = 1$  in the last step. Since (13)–(14) hold for all  $\varphi \in V$  with  $\|\varphi\| = 1$ , the supremum in (12) over all such  $\varphi$  concludes the proof.  $\square$

The final ingredient for the proof of Theorem 1 controls the second term  $\delta$  in the decomposition of Lemma 1 for  $\varrho := G - \nabla w$ . Recall  $C_H$  from Lemma 2 and  $C_1, C_1$  from Lemma 4.

**Lemma 5 (conformity)** *Suppose the divergence-free function  $\varrho \in H(\operatorname{div} = 0, \Omega) \cap H(\operatorname{curl}, \mathcal{T})$  is  $L^2$  orthogonal onto  $RT_0(\mathcal{T}) \cap H(\operatorname{div} = 0, \Omega)$  and satisfies  $(\varrho|_T)|_F \times \nu_F \in L^2(F)$  for all  $F \in \mathcal{F}(T)$  and  $T \in \mathcal{T}$ . Then*

$$C_H^{-1} \|\varrho\| \leq C_1 \|h_{\mathcal{T}} \operatorname{curl}_{\text{pw}} \varrho\| + C_2 \sqrt{\sum_{F \in \mathcal{F}} \ell(F) \| [\varrho]_F \times \nu_F \|_{L^2(F)}^2}.$$

*Proof* Lemma 2 provides  $\beta \in H^1(\Omega; \mathbb{R}^N)$  with (10) for a (component-wise) quasi-interpolation  $\beta_C \in S^1(\mathcal{T})^N$  with (11) as in the proof of Lemma 4; set  $\psi := \beta - \beta_C$ . A piecewise integration by parts and the collection of jump contributions shows

$$\|\varrho\|^2 = \int_{\Omega} \varrho \cdot \operatorname{Curl}(\beta - \beta_C) dx = \int_{\Omega} \psi \cdot \operatorname{curl}_{\text{pw}} \varrho dx + \sum_{F \in \mathcal{F}} \int_F \psi [\varrho]_F \times \nu_F ds.$$

Stability and approximation properties of the quasi-interpolation (11) and the trace inequality of Lemma 3 eventually lead to

$$\|\varrho\|^2 \leq C_1 \|h_{\mathcal{T}} \operatorname{curl}_{\text{pw}} \varrho\| \|\beta\| + C_2 \sqrt{\sum_{F \in \mathcal{F}} \ell(F) \| [\varrho]_F \times \nu_F \|_{L^2(F)}^2} \|\beta\|.$$

In fact, the routine estimation with element and jump terms is completely analogous to the proof of Lemma 4 and leads to the same constants  $C_1, C_2$ . This and  $\|\beta\| \leq C_H \|\varrho\|$  conclude the proof.  $\square$

*Proof (Theorem 1)* The trace of  $G|_T \in H^1(T; \mathbb{R}^n)$  is well defined on any facet  $F \in \mathcal{F}(T)$  of the simplex  $T \in \mathcal{T}$ . Lemma 1 provides  $w \in V = H_0^1(\Omega)$  with  $\|\nabla u - G\|^2 = \|f + \operatorname{div} G\|_*^2 + \|G - \nabla w\|^2$ . Lemma 4 establishes

$$\|f + \operatorname{div} G\|_* \leq C_1 \|h_{\mathcal{T}}(f + \operatorname{div}_{\text{pw}} G)\| + C_2 \sqrt{\sum_{F \in \mathcal{F}(\Omega)} \ell(F) \|[G]_F \cdot \nu_F\|_{L^2(F)}^2}.$$

Since  $\operatorname{curl} \nabla w = 0$  and  $\nabla w \times \nu = 0$  vanish, Lemma 5 applies to  $\varrho := G - \nabla w \in H(\operatorname{div} = 0, \Omega) \cap H(\operatorname{curl}, \mathcal{T}; \mathbb{R}^n)$  and reveals

$$C_H^{-1} \|G - \nabla w\| \leq C_1 \|h_{\mathcal{T}} \operatorname{curl}_{\text{pw}} G\| + C_2 \sqrt{\sum_{F \in \mathcal{F}} \ell(F) \|[G]_F \times \nu_F\|_{L^2(F)}^2}.$$

The above estimates together with the decomposition of Lemma 1 establish  $\eta(\mathcal{T}, G)$  as a GUB for the error  $\|\nabla u - G\|$ .  $\square$

*Example 1 (constants for right-isosceles triangles)* In two space dimensions,  $\|\operatorname{Curl} \bullet\| = \|\bullet\|$  and so  $C_H \leq 1$  for a simply connected domain  $\Omega$  in Lemma 2. The choice  $J := J_1 \circ I_{\text{NC}}$  from [19, Section 5] of the quasi-interpolation operator  $J$  in the proof of Lemma 4 allows for the explicit estimates

$$C_1 \leq \sqrt{48^{-1} + j_{1,1}^{-2} + c_{\text{apx}}^2} \quad \text{and} \quad C_{\text{st}} \leq 1 + \sqrt{72} c_{\text{apx}}.$$

For triangulations into right-isosceles triangles, the constant  $c_{\text{apx}} \leq \sqrt{3}/(2 - 2\cos(\pi/\max\{4, M_{\text{bd}}\}))$  from [19, Lemma 4.8] depends on the domain by the maximal number  $M_{\text{bd}} \leq 4 \max\{\pi, \omega_{\text{max}}\}/\pi \leq 8$  of triangles sharing a boundary vertex. Given the maximal interior angle  $\omega_{\text{max}}$  of  $\Omega$ , Table 1 displays those constants for the maximal possible value  $M_{\text{bd}} = 4 \max\{\pi, \omega_{\text{max}}\}/\pi$ . The geometric quantity  $\max_{x \in T} |x - \operatorname{mid}(T)|$  equals two-thirds of the maximum median of  $T$ . Thus,  $C_{\text{tr}} = \sqrt{5}/(3\sqrt{2}) \leq 0.5271$  and  $\ell(F) = 6h_F$  for interior edges  $F \in \mathcal{F}(\Omega)$  and  $\ell(F) = 12h_F$  for boundary edges  $F \in \mathcal{F}(\partial\Omega)$  of triangulations into right-isosceles triangles. Consequently,

$$C_1 \leq \sqrt{48^{-1} + j_{1,1}^{-2} + c_{\text{apx}}^2} =: C_{\mathcal{T}}, \quad (15)$$

$$C_2 \leq \sqrt{C_{\mathcal{T}}(C_{\mathcal{T}} + 0.5271(1 + \sqrt{72}c_{\text{apx}}))} =: C_{\mathcal{F}}. \quad (16)$$

### 3 Explicit residual-based a posteriori HHO error estimator

The arguments from Section 2 apply to the HHO method and result in a stabilization-free reliable a posteriori error control. In combination with the efficiency estimate from this section, this leads to a new explicit residual-based a posteriori error estimator for the HHO method that is equivalent to the error up to data oscillations.

### 3.1 Hybrid high-order methodology

The HHO ansatz space reads  $V_h := P_k(\mathcal{T}) \times P_k(\mathcal{F}(\Omega))$  for  $k \in \mathbb{N}_0$  with the subspace  $P_k(\mathcal{F}(\Omega)) \subset P_k(\mathcal{F})$  of piecewise polynomials  $p \in P_k(\mathcal{F})$  under the convention  $p|_{\partial\Omega} = 0$ . The interpolation  $\mathbb{I} : V \rightarrow V_h$  maps  $v \in V$  onto  $\mathbb{I}v := (\mathbb{I}_k v, \mathbb{I}_{\mathcal{F},k} v) \in V_h$ . Given any  $v_h = (v_{\mathcal{T}}, v_{\mathcal{F}}) \in V_h$ , the reconstruction operator  $R : V_h \rightarrow P_{k+1}(\mathcal{T})$  defines the unique piecewise polynomial  $Rv_h \in P_{k+1}(\mathcal{T})$  with  $\mathbb{I}_0(Rv_h - v_{\mathcal{T}}) = 0$  such that, for all  $w_{k+1} \in P_{k+1}(\mathcal{T})$ ,

$$\begin{aligned} a_{\text{pw}}(Rv_h, w_{k+1}) &= a_{\text{pw}}(v_{\mathcal{T}}, w_{k+1}) - \sum_{T \in \mathcal{T}} \langle v_{\mathcal{T}|T} - v_{\mathcal{F}}, \nabla w_{k+1}|T \cdot \nu_T \rangle_{L^2(\partial T)}. \end{aligned} \quad (17)$$

Let  $u_h \in V_h$  solve the HHO discrete formulation of (3) with

$$a_h(u_h, v_h) = (f, v_{\mathcal{T}})_{L^2(\Omega)} \quad \text{for all } v_h = (v_{\mathcal{T}}, v_{\mathcal{F}}) \in V_h \quad (18)$$

for the HHO bilinear form

$$a_h(u_h, v_h) := a_{\text{pw}}(Ru_h, Rv_h) + s_h(u_h, v_h) \quad (19)$$

and the stabilization term  $s_h(u_h, v_h)$  from (2). Given any  $w_C \in S_0^1(\mathcal{T}) = P_1(\mathcal{T}) \cap H_0^1(\Omega)$ , the definition of the reconstruction operator  $R$  in (17) verifies  $R\mathbb{I}w_C = w_C$  with the interpolation  $\mathbb{I}$  onto  $V_h$ . Hence,  $S_{TF}\mathbb{I}w_C = 0$  vanishes for all  $F \in \mathcal{F}(T)$  and  $T \in \mathcal{T}$ . This and (18) show, for all  $w_C \in S_0^1(\mathcal{T})$ , that

$$a_{\text{pw}}(Ru_h, w_C) = (f, \mathbb{I}_k w_C)_{L^2(\Omega)} = (\mathbb{I}_k f, w_C)_{L^2(\Omega)}. \quad (20)$$

### 3.2 Explicit a posteriori error estimator

As a result of (20),  $\nabla_{\text{pw}} Ru_h$  satisfies the solution property (5) if  $k \geq 1$  and, in the lowest order case  $k = 0$ , (5) holds with  $f$  replaced by  $\mathbb{I}_0 f$ . This allows the application of the theory from Section 2 to the HHO method with minor modifications for the case  $k = 0$ . Define the error estimator contributions

$$\begin{aligned} \eta_{\text{res},1}(\mathcal{T})^2 &:= \begin{cases} \|h_{\mathcal{T}}(f + \Delta_{\text{pw}} Ru_h)\|^2 & \text{for } k \geq 1, \\ \|h_{\mathcal{T}} \mathbb{I}_0 f\|^2 & \text{for } k = 0, \end{cases} \\ \eta_{\text{res},2}(\mathcal{T})^2 &:= \begin{cases} 0 & \text{for } k \geq 1, \\ \text{osc}_0^2(f, \mathcal{T}) & \text{for } k = 0, \end{cases} \\ \eta_{\text{res},3}(\mathcal{T})^2 &:= \sum_{F \in \mathcal{F}(\Omega)} \ell(F) \|[\nabla_{\text{pw}} Ru_h]_F \cdot \nu_F\|_{L^2(F)}^2, \\ \eta_{\text{res},4}(\mathcal{T})^2 &:= \sum_{F \in \mathcal{F}} \ell(F) \|[\nabla_{\text{pw}} Ru_h]_F \times \nu_F\|_{L^2(F)}^2. \end{aligned} \quad (21)$$

Since  $\nabla_{\text{pw}} Ru_h$  is a piecewise gradient, its piecewise curl vanishes. This leads to the explicit residual-based a posteriori error estimator

$$\eta_{\text{res}}(\mathcal{T})^2 := (C_1 \eta_{\text{res},1}(\mathcal{T}) + C_P \eta_{\text{res},2}(\mathcal{T}) + C_2 \eta_{\text{res},3}(\mathcal{T}))^2 + C_H^2 C_2^2 \eta_{\text{res},4}^2(\mathcal{T}). \quad (22)$$

(Recall  $C_1, C_2$  from Lemma 4 and  $C_H$  from Lemma 2 as well as the Poincaré constant  $C_P \leq \pi^{-1}$ .) The main result of this section verifies the assumptions in Theorem 1 and proves reliability and efficiency of  $\eta_{\text{res}}(\mathcal{T})$ .

**Theorem 2 (residual-based GUB for HHO)** *Let  $u \in V$  solve the Poisson equation (3) and let  $u_h \in V_h$  solve the discrete formulation (18). Then*

$$\|u - Ru_h\|_{\text{pw}} \leq \eta_{\text{res}}(\mathcal{T}) \leq C_3 (\|u - Ru_h\|_{\text{pw}} + \text{osc}_q(f, \mathcal{T}))$$

and  $\text{osc}_{k-1}(f, \mathcal{T}) \leq C_4 \eta_{\text{res}}(\mathcal{T})$  hold for any  $q \in \mathbb{N}_0$ . The constants  $C_3$  and  $C_4$  exclusively depend on  $k, q$  and on the shape-regularity of the triangulation  $\mathcal{T}$ .

### 3.3 Proof of Theorem 2

The orthogonality of  $\nabla_{\text{pw}} Ru_h$  to the divergence-free Raviart-Thomas space of lowest degree is an assumption in Theorem 1 and verified below.

**Lemma 6 (orthogonality)** *The piecewise gradients  $\nabla_{\text{pw}} Rv_h$  are  $L^2$  orthogonal to the space  $RT_0(\mathcal{T}) \cap H(\text{div} = 0, \Omega)$ , i.e., any  $v_h \in V_h$  and  $q_{RT} \in RT_0(\mathcal{T}) \cap H(\text{div} = 0, \Omega)$  satisfy*

$$(\nabla_{\text{pw}} Rv_h, q_{RT})_{L^2(\Omega)} = 0. \quad (23)$$

*Proof* Given any  $q_{RT} \in RT_0(\mathcal{T}) \cap H(\text{div} = 0, \Omega)$ ,  $\text{div } q_{RT} = 0$  shows  $q_{RT} \in P_0(\mathcal{T}; \mathbb{R}^n)$  [27, Lemma 14.9]. Since  $P_0(\mathcal{T}; \mathbb{R}^n) = \nabla_{\text{pw}} P_1(\mathcal{T})$ , there exists a piecewise affine function  $\phi_1 \in P_1(\mathcal{T})$  with  $q_{RT} = \nabla_{\text{pw}} \phi_1$  a.e. in  $\Omega$ . This and the definition of  $Rv_h$  from (17) imply, for any  $v_h = (v_{\mathcal{T}}, v_{\mathcal{F}}) \in V_h$ , that

$$\begin{aligned} (\nabla_{\text{pw}} Rv_h, q_{RT})_{L^2(\Omega)} &= a_{\text{pw}}(Rv_h, \phi_1) \\ &= a_{\text{pw}}(v_{\mathcal{T}}, \phi_1) - \sum_{T \in \mathcal{T}} \langle v_{\mathcal{T}}|_T - v_{\mathcal{F}}, \nabla_{\text{pw}} \phi_1 \cdot \nu_T \rangle_{L^2(\partial T)}. \end{aligned}$$

This, a piecewise integration by parts, and  $\Delta_{\text{pw}} \phi_1 \equiv 0$  lead to

$$\begin{aligned} (\nabla_{\text{pw}} Rv_h, q_{RT})_{L^2(\Omega)} &= \sum_{T \in \mathcal{T}} \langle v_{\mathcal{F}}, \nabla_{\text{pw}} \phi_1 \cdot \nu_T \rangle_{L^2(\partial T)} \\ &= \sum_{F \in \mathcal{F}} \langle v_{\mathcal{F}}, [q_{RT} \cdot \nu_F]_F \rangle_{L^2(F)}. \end{aligned} \quad (24)$$

Since  $v_{\mathcal{F}}$  is single-valued on  $F$  and  $q_{RT} \in RT_0(\mathcal{T})$  has continuous normal components, the jump term  $\langle v_{\mathcal{F}}, [q_{RT}]_F \cdot \nu_F \rangle_{L^2(F)} = 0$  vanishes for all  $F \in \mathcal{F}(\Omega)$ . This,  $v_{\mathcal{F}} \equiv 0$  on  $\partial\Omega$ , and (24) conclude  $(\nabla_{\text{pw}} Rv_h, q_{RT})_{L^2(\Omega)} = 0$ .  $\square$

The following lemma concerns the efficiency of the jump contributions. Each facet  $F \in \mathcal{F}$  has at most two adjacent simplices that define a triangulation  $\mathcal{T}(F) := \{T \in \mathcal{T} : F \in \mathcal{F}(T)\}$  of the facet-patch  $\omega(F) := \text{int}(\bigcup_{T \in \mathcal{T}(F)} T)$ .

**Lemma 7 (efficiency of jumps)** *The solution  $u \in V$  to (3) and the discrete solution  $u_h \in V_h$  to (18) satisfy (a) for all  $F \in \mathcal{F}$  and (b) for all  $F \in \mathcal{F}(\Omega)$ .*

$$(a) \quad h_F^{1/2} \|[\nabla_{\text{pw}} R u_h]_F \times \nu_F\|_{L^2(F)} \lesssim \min_{v \in V} \|\nabla v - \nabla_{\text{pw}} R u_h\|_{L^2(\omega(F))},$$

$$(b) \quad h_F^{1/2} \|[\nabla_{\text{pw}} R u_h]_F \cdot \nu_F\|_{L^2(F)} \lesssim \|\nabla u - \nabla_{\text{pw}} R u_h\|_{L^2(\omega(F))} + \text{osc}_k(f, \mathcal{T}(F)).$$

*Proof* The proof is based on the following extension argument. Given a polynomial  $p \in P_k(F)$  of degree at most  $k$  along the side  $F \in \mathcal{F}$ , the coefficients determine a polynomial (also denoted by  $p$ ) along the hyperplane  $H$  that enlarges  $F$ . The intersection  $\widehat{F} := H \cap \text{conv}(\omega(F))$  of the hyperplane  $H$  with the convex hull of the facet-patch  $\omega(F)$  may be strictly larger than  $F$ . The shape-regularity of  $\mathcal{T}$  bounds the size of  $\widehat{F}$  in terms of  $F$  and an inverse estimate leads to a bound  $\|p\|_{L^\infty(\widehat{F})} \leq C(k) \|p\|_{L^\infty(F)}$  with a constant  $C(k)$  that depends on the shape-regularity of  $\mathcal{T}$  and on  $k$ . The extension of  $p$  from  $H$  to  $\mathbb{R}^n$  by constant values along the side normal  $\nu_F$  leads to a polynomial  $\widehat{p} \in P_k(\mathbb{R}^n)$  with

$$\|\widehat{p}\|_{L^\infty(\omega(F))} \leq \|p\|_{L^\infty(\widehat{F})} \leq C(k) \|p\|_{L^\infty(F)}. \quad (25)$$

Proof of (a). The tangential jump  $\varrho_F := [\nabla_{\text{pw}} R v_h]_F \times \nu_F \in P_k(F; \mathbb{R}^N)$  is a polynomial in  $N = 2n - 3$  components on  $F \in \mathcal{F}$  for  $n = 2, 3$ . Let  $p = \varrho_F(j)$  be one of the components of  $\varrho_F \in P_k(F)^N$ , for  $j = 1, \dots, N$ , and extend it as explained above to  $\widehat{p} \in P_k(\mathbb{R}^n)$  and call this  $\widehat{\varrho}(j)$  in the vector  $\widehat{\varrho}_F \in P_k(\mathbb{R}^n; \mathbb{R}^N)$ . The proof involves the piecewise polynomial facet-bubble function  $b_F := n^n \prod_{j=1}^n \varphi_j$  for the  $n$  nodal basis function  $\varphi_1, \dots, \varphi_n \in S^1(\mathcal{T})$  associated with the vertices of  $F$ . An inverse estimate [36, Proposition 3.37] shows

$$\|\varrho_F\|_{L^2(F)}^2 \lesssim \|b_F^{1/2} \varrho_F\|_{L^2(F)}^2 = \langle b_F \varrho_F, \varrho_F \rangle_{L^2(F)}. \quad (26)$$

Since  $\varrho := b_F \widehat{\varrho}_F \in S_0^{k+n}(\mathcal{T}(F); \mathbb{R}^N)$  vanishes on  $\partial\omega(F) \setminus \text{int}(F)$ , (26) and a piecewise integration by parts show

$$\|\varrho_F\|_{L^2(F)}^2 \lesssim \langle \varrho, [\nabla_{\text{pw}} R v_h]_F \times \nu_F \rangle_{L^2(F)} = (\text{curl } \varrho, \nabla_{\text{pw}} R v_h)_{L^2(\omega(F))}. \quad (27)$$

This and  $(\text{curl } \varrho, \nabla v)_{L^2(\Omega)} = 0$  for any  $v \in V$  imply

$$\begin{aligned} \|\varrho_F\|_{L^2(F)}^2 &\lesssim (\text{curl } \varrho, \nabla_{\text{pw}}(R v_h - v))_{L^2(\omega(F))} \\ &\leq \|\text{curl } \varrho\|_{L^2(\omega(F))} \|\nabla_{\text{pw}}(R v_h - v)\|_{L^2(\omega(F))}. \end{aligned} \quad (28)$$

An inverse estimate,  $\|b_F\|_{L^\infty(\omega(F))} = 1$ , and (25) imply

$$\begin{aligned} \|\text{curl } \varrho\|_{L^2(\omega(F))} &\lesssim \|\nabla \varrho\|_{L^2(\omega(F))} \lesssim h_F^{-1+n/2} \|\varrho\|_{L^\infty(\omega(F))} \\ &\lesssim h_F^{-1+n/2} h_F^{-(n-1)/2} \|\varrho_F\|_{L^2(F)} = h_F^{-1/2} \|\varrho_F\|_{L^2(F)}. \end{aligned} \quad (29)$$

In combination with (28), this concludes the proof of (a).

Proof of (b). The efficiency of normal jumps (b) follows from the arguments for conforming FEMs, cf. [36, Section 1.4.5]; further details are omitted.  $\square$

The following lemma reveals that the order  $k \geq \mathbb{N}_0$  of the oscillations  $\text{osc}_k(f, T)$  in Lemma 7 (b) can be any natural number. It is certainly known to the experts but hard to find in the literature. Recall the convention  $\Pi_{-1} := 0$ .

**Lemma 8 (efficiency of lower-order oscillations)** *Given any simplex  $T \in \mathcal{T}$  and parameters  $k, q \in \mathbb{N}_0$ , the solution  $u \in V$  to (3) satisfies*

$$C_5^{-1} \text{osc}_{k-1}^2(f, T) \leq \min_{v_{k+1} \in P_{k+1}(T)} \|\nabla_{\text{pw}}(u - v_{k+1})\|_{L^2(T)}^2 + \text{osc}_q^2(f, T). \quad (30)$$

The constant  $C_5$  exclusively depends on  $q$  and the shape of  $T$ .

*Proof* The assertion (30) is trivial for  $q \leq k - 1$ , so suppose  $k \leq q$ . Any  $v_{k+1} \in P_{k+1}(T)$  and  $\varrho_T := \Pi_q f + \Delta v_{k+1} \in P_q(T)$  satisfy

$$\text{osc}_{k-1}^2(f, T) \leq h_T^2 \|f + \Delta v_{k+1}\|_{L^2(T)}^2 = \text{osc}_q^2(f, T) + h_T^2 \|\varrho_T\|_{L^2(T)}^2. \quad (31)$$

Let  $b_T \in S_0^{n+1}(T)$  with  $0 \leq b_T \leq 1 = \max b_T$  denote the volume bubble-function on  $T \in \mathcal{T}$ . The equivalence of norms in the finite-dimensional space  $P_q(T)$  provides

$$\|b_T^{1/2} \varrho_T\|_{L^2(T)} \leq \|\varrho_T\|_{L^2(T)} \leq C_6 \|b_T^{1/2} \varrho_T\|_{L^2(T)}. \quad (32)$$

A more detailed analysis of the mass matrices reveals that the constant  $C_6$  exclusively depends on the polynomial degree  $q$ . An integration by parts with  $b_T \varrho_T \in S_0^{q+n+1}(T) \subset V$  and the weak formulation (3) result in

$$\begin{aligned} \|b_T^{1/2} \varrho_T\|_{L^2(T)}^2 &= (\Pi_q f + \Delta v_{k+1}, b_T \varrho_T)_{L^2(T)} \\ &= (\nabla(u - v_{k+1}), \nabla(b_T \varrho_T))_{L^2(T)} - (f - \Pi_q f, b_T \varrho_T)_{L^2(T)}. \end{aligned}$$

A Cauchy inequality, the inverse estimate  $h_T \|\nabla(b_T \varrho_T)\|_{L^2(T)} \leq C_7 \|\varrho_T\|_{L^2(T)}$  with a constant  $C_7$  that exclusively depends on  $q + n + 1$  and the shape of  $T$ , and (32) lead to

$$C_6^{-2} h_T \|\varrho_T\|_{L^2(T)} \leq C_7 \|\nabla(u - v_{k+1})\|_{L^2(T)} + \text{osc}_q(f, T) \quad (33)$$

The combination of (31) with (33) and a Cauchy inequality conclude the proof of (30), e.g., with  $C_5 = 1 + C_6^4(1 + C_7^2)$ .  $\square$

*Proof (of Theorem 2)* Recall the definition of  $\eta_{\text{res}}(\mathcal{T})$  for  $k \geq 1$  and  $k = 0$  in (21). Since  $\text{osc}_{k-1}^2(f, \mathcal{T}) \leq \eta_{\text{res},1}^2(\mathcal{T}) + \eta_{\text{res},2}^2(\mathcal{T}) \lesssim \eta_{\text{res}}^2(\mathcal{T})$ , the remaining parts of this proof discuss the reliability and efficiency of  $\eta_{\text{res}}(\mathcal{T})$ .

Lemma 6 provides the orthogonality of  $\nabla_{\text{pw}} R u_h \in H^1(\mathcal{T}; \mathbb{R}^n)$  to the divergence-free Raviart-Thomas function  $RT_0(\mathcal{T}) \cap H(\text{div} = 0, \Omega)$ . This and (20) show that the assumptions in Theorem 1 hold for  $G := \nabla_{\text{pw}} R u_h$  and  $k \geq 1$ , whence the reliability of  $\eta_{\text{res}}(\mathcal{T})$  follows with a reliability constant 1.

Minor modifications to the proof of Theorem 1 lead to reliability in the case  $k = 0$ . In fact, the only modifications required concern the upper bound of  $\|f + \operatorname{div} \nabla_{\text{pw}} Ru_h\|_* \leq \|(1 - \Pi_0)f\|_* + \|\Pi_0 f + \operatorname{div} \nabla_{\text{pw}} Ru_h\|_*$ . A piecewise Poincaré inequality shows  $\|(1 - \Pi_0)f\|_* \leq C_P \operatorname{osc}_0(\mathcal{T}, f)$  with the Poincaré constant  $C_P (\leq 1/\pi$  for simplices). Lemma 4 proves  $\|\Pi_0 f + \operatorname{div} \nabla_{\text{pw}} Ru_h\|_* \leq C_1 \eta_{\text{res},1}(\mathcal{T}) + C_2 \eta_{\text{res},3}(\mathcal{T})$ . Hence, the decomposition of Lemma 1 and Lemma 5 result in  $\|u - Ru_h\|_{\text{pw}} \leq \eta_{\text{res}}(\mathcal{T})$ .

This provides the reliability and it remains to verify the efficiency  $\eta_{\text{res}}(\mathcal{T}) \lesssim \|u - Ru_h\|_{\text{pw}} + \operatorname{osc}_q(f, \mathcal{T})$  for any  $q \in \mathbb{N}_0$ . The Pythagoras theorem and (33) with  $\varrho_T := \Pi_k f + \Delta Ru_h \in P_k(T)$  and  $v_{k+1} := Ru_h \in P_{k+1}(T)$  lead to the local efficiency of the volume contributions

$$\begin{aligned} \|h_T(f + \Delta_{\text{pw}} Ru_h)\|_{L^2(T)}^2 &= \operatorname{osc}_k^2(f, T) + \|h_T \varrho_T\|_{L^2(T)}^2 \\ &\lesssim \|\nabla(u - Ru_h)\|_{L^2(T)}^2 + \operatorname{osc}_k^2(f, T). \end{aligned}$$

Lemma 7 considers the remaining terms in the error estimator and establishes their efficiency namely,

$$\sum_{F \in \mathcal{F}} h_F \|\nabla_{\text{pw}} Ru_h\|_{L^2(F)} \lesssim \|u - Ru_h\|_{\text{pw}} + \operatorname{osc}_k(f, \mathcal{T})$$

with the modified jump  $[\nabla_{\text{pw}} Ru_h]_F = \nabla_{\text{pw}} Ru_h \times \nu_F$  on boundary facets  $F \in \mathcal{F}(\partial\Omega)$ . This and Lemma 8 establish the existence of some mesh-independent constant  $C_3 > 0$  with  $C_3^{-1} \eta_{\text{res}}(\mathcal{T}) \leq \|u - Ru_h\|_{\text{pw}} + \operatorname{osc}_q(f, \mathcal{T})$  for arbitrary  $q \in \mathbb{N}_0$ . This concludes the proof.  $\square$

## 4 Equilibrium-based a posteriori HHO error analysis

The residual-based guaranteed upper bound (GUB) of the error  $\|u - Ru_h\|_{\text{pw}}$  from Subsection 3.2 employs explicit constants that may lead to overestimation in higher dimensions and for different triangular shapes. This section utilizes equilibrated flux reconstructions [1, 2, 29, 6, 5] to establish a constant-free guaranteed upper bounds for a tight error control.

### 4.1 Guaranteed error control

The guaranteed upper bounds of this section involves two post-processings of the potential reconstruction  $Ru_h \in P_{k+1}(\mathcal{T})$  of the discrete solution  $u_h$  to (18). First, the patch-wise design of a flux reconstruction  $Q_p \in RT_{k+p}(\mathcal{T})$  with  $p \in \mathbb{N}_0$  from [3, 10, 30] provides an  $H(\operatorname{div}, \Omega)$ -conforming approximation to  $\nabla_{\text{pw}} Ru_h$  with the equilibrium  $\Pi_r f + \operatorname{div} Q_p = 0$  in  $\Omega$  and  $r$  from (35) below. Second, the nodal average  $\mathcal{A}Ru_h \in S_0^{k+1}(\mathcal{T}) \subset V$  results in an  $V$ -conforming approximation of  $Ru_h$  by averaging all values of the discontinuous



function  $Ru_h$  at each Lagrange point of  $S_0^{k+1}(\mathcal{T})$ . This, the split (7), and the solution property (20) give rise to the guaranteed upper bound (GUB)

$$\eta_{\text{eq},p}^2(\mathcal{T}) := (C_P \text{osc}_r(f, \mathcal{T}) + \|Q_p - \nabla_{\text{pw}} Ru_h\|)^2 + \|(1 - \mathcal{A})Ru_h\|_{\text{pw}}^2 \quad (34)$$

with  $r \in \mathbb{N}_0$  defined by

$$r := 0 \text{ if } k = 0 \quad \text{and} \quad r := k + p \text{ if } k \geq 1. \quad (35)$$

The main result of this section states the reliability and efficiency (up to data oscillations) of  $\eta_{\text{eq},p}$  for all parameters  $p \in \mathbb{N}_0$ .

**Theorem 3 (equilibrium-based GUB for HHO)** *Let  $u \in V$  resp.  $u_h \in V_h$  solve (3) resp. (18). Given a parameter  $p \in \mathbb{N}_0$ , there exists  $Q_p \in RT_{k+p}(\mathcal{T})$  such that the error estimator  $\eta_{\text{eq},p}(\mathcal{T})$  from (34) is an efficient GUB*

$$\|u - Ru_h\|_{\text{pw}} \leq \eta_{\text{eq},p}(\mathcal{T}) \leq C_8 (\|u - Ru_h\|_{\text{pw}} + \text{osc}_q(f, \mathcal{T})). \quad (36)$$

for any  $q \in \mathbb{N}_0$  and  $\text{osc}_{k-1}(f, \mathcal{T}) \leq C_9 \eta_{\text{eq},p}(\mathcal{T})$ . The constants  $C_8$  and  $C_9$  exclusively depend on the polynomial degree  $k \in \mathbb{N}_0$ , the parameter  $q \in \mathbb{N}_0$ , and the shape-regularity of  $\mathcal{T}$ .

At least two technical contributions for the proof of Theorem 3 are of broader interest. A first contribution to the HHO literature is the local equivalence of the original HHO stabilization  $s_h$  from (2) and the alternative stabilization  $\tilde{s}_h(v_h, v_h) := \sum_{T \in \mathcal{T}} \tilde{s}_T(v_h, v_h)$  from [24] defined, for  $v_h = (v_{\mathcal{T}}, v_{\mathcal{F}}) \in V_h$ , by

$$\begin{aligned} \tilde{s}_T(v_h, v_h) &:= h_T^{-2} \|\Pi_{T,k}(v_{\mathcal{T}} - Rv_h)\|_{L^2(T)}^2 \\ &+ \sum_{F \in \mathcal{F}(T)} h_F^{-1} \|\Pi_{F,k}(v_{\mathcal{F}} - Rv_h|_T)\|_{L^2(F)}^2. \end{aligned} \quad (37)$$

A second result of separate interest in the HHO literature (cf. [24] where the efficiency in (38) is left open) is the efficiency of the stabilizations from Theorems 4–5 below,

$$\tilde{s}_h(v_h, v_h)^{1/2} \approx s_h(u_h, u_h)^{1/2} \lesssim \|u - Ru_h\|_{\text{pw}} + \text{osc}_{k+p}(f, \mathcal{T}). \quad (38)$$

The subsequent subsection continues with some explanations on the flux reconstruction  $Q_p \in RT_{k+p}(\mathcal{T})$  that is defined by local minimization problems on each vertex patch. Appendix A complements the discussion with an algorithmic two-step procedure for the computation of  $Q_p - \nabla_{\text{pw}} Ru_h$  in 2D. The efficiency of the averaging  $\|(1 - \mathcal{A})Ru_h\|_{\text{pw}}$  up to data oscillations follows in Subsections 4.3–4.4. Subsection 4.6 concludes with the proof of Theorem 3.

## 4.2 Construction of equilibrated flux

This subsection defines the post-processed  $H(\operatorname{div}, \Omega)$ -conforming equilibrated flux  $Q_p \in RT_{k+p}(\mathcal{T})$  that enters the GUB  $\eta_{\text{eq},p}$  from (34) based on local patch-wise minimization problems in the spirit of [10, 29, 30].

Consider the shape-regular vertex-patch  $\omega(z) := \operatorname{int}(\bigcup\{T \in \mathcal{T}(z)\})$  covered by the neighbouring simplices  $\mathcal{T}(z) := \{T \in \mathcal{T} : z \in T\}$  sharing a given vertex  $z \in \mathcal{V}$  with the facet spider  $\mathcal{F}(z) := \{F \in \mathcal{F} : z \in F\}$ . Recall the space of piecewise Raviart-Thomas functions  $RT_k^{\text{pw}}(\mathcal{T})$  from Subsection 1.4 and define

$$\begin{aligned} L_0^2(\omega(z)) &:= \{f \in L^2(\omega(z)) : (f, 1)_{L^2(\omega(z))} = 0\}, \\ L_*^2(\omega(z)) &:= \begin{cases} L_0^2(\omega(z)) & \text{if } z \in \mathcal{V}(\Omega), \\ L^2(\omega(z)) & \text{else,} \end{cases} \\ H_*^1(\omega(z)) &:= \begin{cases} H^1(\omega(z)) \cap L_0^2(\omega(z)) & \text{if } z \in \mathcal{V}(\Omega), \\ \{v \in H^1(\omega(z)) : v = 0 \text{ on } \partial\Omega \cap \bigcup\mathcal{F}(z)\} & \text{else,} \end{cases} \\ H_0(\operatorname{div}, \omega(z)) &:= \begin{cases} \{r \in H(\operatorname{div}, \omega(z)) : r \cdot \nu = 0 \text{ on } \partial\omega(z)\} & \text{if } z \in \mathcal{V}(\Omega), \\ \{r \in H(\operatorname{div}, \omega(z)) : r \cdot \nu = 0 \text{ on } \partial\omega(z) \setminus \bigcup\mathcal{F}(z)\} & \text{else,} \end{cases} \\ RT_k^0(\mathcal{T}(z)) &:= RT_k^{\text{pw}}(\mathcal{T}(z)) \cap H_0(\operatorname{div}, \omega(z)). \end{aligned}$$

Throughout the remaining parts of this section, abbreviate  $G_h := \nabla_{\text{pw}} Ru_h \in P_k(\mathcal{T}; \mathbb{R}^n)$ . Given a vertex  $z \in \mathcal{V}$  with the  $P_1$ -conforming nodal basis function  $\varphi_z \in S^1(\mathcal{T})$ , the property (20) provides compatible data

$$f_z := \begin{cases} \Pi_p(\varphi_z \Pi_0 f - G_h \cdot \nabla \varphi_z) & \text{if } k = 0, \\ \Pi_{k+p}(\varphi_z f - G_h \cdot \nabla \varphi_z) & \text{if } k \geq 1 \end{cases} \in L_*^2(\omega(z)) \quad (39)$$

such that the discrete affine space

$$\mathcal{Q}_h(z) := \{\tau_z \in RT_{k+p}^0(\mathcal{T}(z)) : \operatorname{div} \tau_z + f_z = 0 \text{ in } \Omega\} \neq \emptyset \quad (40)$$

is not empty. Consequently,

$$Q_{z,h} := \arg \min_{\tau_z \in \mathcal{Q}_h(z)} \|\tau_z - \mathcal{I}_{RT}(\varphi_z G_h)\|_{L^2(\omega(z))} = \Pi_{\mathcal{Q}_h(z)} \mathcal{I}_{RT}(\varphi_z G_h) \quad (41)$$

is well defined as the  $L^2$  projection  $\Pi_{\mathcal{Q}_h(z)} \mathcal{I}_{RT}(\varphi_z G_h)$  of  $\mathcal{I}_{RT}(\varphi_z G_h)$  onto  $\mathcal{Q}_h(z)$  with the piecewise Raviart-Thomas interpolation  $\mathcal{I}_{RT} : H^1(\mathcal{T}; \mathbb{R}^n) \rightarrow RT_{k+p}^{\text{pw}}(\mathcal{T})$  [7, Section III.3.1]. In the case  $p \geq 1$ ,  $\varphi_z G_h \in P_{k+1}(\mathcal{T}(z)) \subset RT_{k+p}^{\text{pw}}(\mathcal{T}(z))$  is a piecewise Raviart-Thomas function of degree  $k+p$ . Hence  $\mathcal{I}_{RT}(\varphi_z G_h) = \varphi_z G_h$  and the interpolation  $\mathcal{I}_{RT}$  could be omitted in the formula (41). The partition of unity  $\sum_{z \in \mathcal{V}} \varphi_z \equiv 1$  shows that the sum  $Q_p =$

$\sum_{z \in \mathcal{V}} Q_{z,h} \in H(\operatorname{div}, \Omega)$  of the patch-wise contributions satisfies

$$\operatorname{div} Q_p = \begin{cases} \sum_{z \in \mathcal{V}} \Pi_p(G_h \cdot \nabla \varphi_z - \varphi_z \Pi_0 f) = -\Pi_0 f & \text{if } k = 0, \\ \sum_{z \in \mathcal{V}} \Pi_{k+p}(G_h \cdot \nabla \varphi_z - \varphi_z f) = -\Pi_{k+p} f & \text{if } k \geq 1, \end{cases} \quad (42)$$

$$\|Q_p - G_h\|_{L^2(\Omega)}^2 = \sum_{z \in \mathcal{V}} \|Q_{z,h} - \mathcal{I}_{RT}(\varphi_z G_h)\|_{L^2(\omega(z))}^2. \quad (43)$$

This establishes the flux reconstruction  $Q_p$ . The efficiency of the flux reconstruction will be based on the following general equivalence.

**Lemma 9 (control of  $H(\operatorname{div})$  minimization by residual [10, 30])** *Given any vertex  $z \in \mathcal{V}$ , a piecewise Raviart-Thomas function  $\sigma_z \in RT_q^{\text{pw}}(\mathcal{T}(z))$  and a piecewise polynomial  $r_z \in P_q(\mathcal{T}(z))$  of degree  $q \in \mathbb{N}_0$ , define the residual*

$$\operatorname{Res}_z(v) := \sum_{T \in \mathcal{T}(z)} \left( (r_z, v)_{L^2(T)} + \langle \sigma_z \cdot \nu_T, v \rangle_{L^2(\partial T)} \right) \quad (44)$$

for all  $v \in H^1(\Omega)$ . If  $z \in \mathcal{V}(\Omega)$  is an interior vertex, then suppose additionally that  $\operatorname{Res}_z(1) = 0$ . Then

$$\min_{\substack{\tau_z \in RT_q^0(\mathcal{T}(z)) \\ \operatorname{div} \tau_z = r_z + \operatorname{div}_{\text{pw}} \sigma_z}} \|\tau_z - \sigma_z\|_{L^2(\omega(z))} \leq C_s \max_{\substack{v \in H_*^1(\omega(z)) \\ \|\nabla v\|_{L^2(\omega(z))} = 1}} \operatorname{Res}_z(v) \quad (45)$$

holds for a constant  $C_s$  that exclusively depends on the shape-regularity (and is in particular independent of the polynomial degree  $q$ ).

*Proof* The assertion follows from [10, Theorem 7] in  $n = 2$  dimensions and [30, Corollaries 3.3, 3.6, and 3.8] in  $n = 3$  dimensions.  $\square$

*Remark 1* The patch-wise construction of  $Q_p$  in Subsection 4.2 typically generates local data oscillation  $\operatorname{osc}_{k+p}(\varphi_z f, \mathcal{T}(z))$  in the error analysis as in the proof of Theorem 3 in Subsection 4.6 below or, e.g., [29, Theorem 3.17]. A straightforward computation  $\operatorname{osc}_{k+p}(\varphi_z f, \mathcal{T}(z)) \leq \operatorname{osc}_{k+p-1}(f, \mathcal{T}(z))$  apparently leads to a loss of one degree in the data oscillation but Lemma 8 verifies

$$\operatorname{osc}_{k+p}^2(\varphi_z f, \mathcal{T}(z)) \lesssim \min_{v_h \in P_{k+1}(\mathcal{T}(z))} \|\nabla_{\text{pw}}(u - v_h)\|_{L^2(\omega(z))}^2 + \operatorname{osc}_q^2(f, \mathcal{T}(z)) \quad (46)$$

for any  $p, q \in \mathbb{N}_0$ . This allows for efficiency of the data oscillations on the right-hand side of the efficiency estimate [29, Formula (3.42)] and leads to a corresponding refinement in [29, Theorem 3.17].

### 4.3 Local equivalence of stabilizations

The first improvement to the current HHO literature is the local equivalence of the two stabilizations  $\tilde{s}_h$  from (37) and  $s_h$  from (2). The authors of this paper could not find any motivation for the alternative stabilization  $\tilde{s}_h$  in the error analysis of [24, Section 4] and suggest to apply Theorem 4 below to [24, Theorem 4.7] to recover the results therein for the original HHO stabilization  $s_h$ . Recall the local stabilization  $\tilde{s}_T$  in  $\tilde{s}_h(v_h, v_h) := \sum_{T \in \mathcal{T}} \tilde{s}_T(v_h, v_h)$  from (37) and  $S_{TF}v_h = \Pi_{F,k}(Rv_{h|T} - v_{\mathcal{F}}|_F - \varphi_k)$  in the definition of  $s_h$  from (2).

**Theorem 4 (local equivalence of stabilizations)** *Any  $v_h = (v_{\mathcal{T}}, v_{\mathcal{F}}) \in V_h$  and  $T \in \mathcal{T}$  satisfy*

$$C_{10}^{-1} \tilde{s}_T(v_h, v_h) \leq \sum_{F \in \mathcal{F}(T)} h_F^{-1} \|S_{TF}v_h\|_{L^2(F)}^2 \leq C_{11} \tilde{s}_T(v_h, v_h). \quad (47)$$

*The constants  $C_{10}$  and  $C_{11}$  exclusively depend on the polynomial degree  $k$  and the shape regularity of  $\mathcal{T}$ .*

*Proof* The second inequality in (47) follows directly from a triangle inequality and an inverse estimate. Therefore, the proof focuses on the first inequality in (47). Given  $v_h = (v_{\mathcal{T}}, v_{\mathcal{F}}) \in V_h$  and  $T \in \mathcal{T}$ , let  $\varphi_k := (\Pi_k Rv_h - v_{\mathcal{T}})|_T \in P_k(T)$ . Since  $S_{TF}v_h = \Pi_{F,k}(Rv_{h|T} - v_{\mathcal{F}}|_F - \varphi_k)$ , the triangle inequality  $\|\Pi_{F,k}(Rv_{h|T} - v_{\mathcal{F}})\|_{L^2(F)} \leq \|S_{TF}v_h\|_{L^2(F)} + \|\varphi_k\|_{L^2(F)}$ , the discrete trace inequality  $\|\varphi_k\|_{L^2(F)} \lesssim h_F^{-1/2} \|\varphi_k\|_{L^2(T)}$ , and the shape-regularity  $h_F \approx h_T$  for all  $F \in \mathcal{F}(T)$  reveal

$$\begin{aligned} & \sum_{F \in \mathcal{F}(T)} h_F^{-1} \|\Pi_{F,k}(Rv_{h|T} - v_{\mathcal{F}})\|_{L^2(F)}^2 \\ & \lesssim \sum_{F \in \mathcal{F}(T)} h_F^{-1} \|S_{TF}v_h\|_{L^2(F)}^2 + h_T^{-2} \|\varphi_k\|_{L^2(T)}^2. \end{aligned} \quad (48)$$

Since  $\Pi_0 \varphi_k = 0$  (from the design of  $Rv_h$ ), a Poincaré inequality shows

$$h_T^{-2} \|\varphi_k\|_{L^2(T)}^2 \leq C_P^2 \|\nabla \varphi_k\|_{L^2(T)}^2. \quad (49)$$

On the one hand, an integration by parts provides

$$\|\nabla \varphi_k\|_{L^2(T)}^2 = -(\Pi_k Rv_h - v_{\mathcal{T}}, \Delta \varphi_k)_{L^2(T)} + \langle \varphi_k, \nabla \varphi_k \cdot \nu_T \rangle_{L^2(\partial T)}. \quad (50)$$

On the other hand, an integration by parts and the definition of  $R$  imply

$$\begin{aligned} - (Rv_h, \Delta \varphi_k)_{L^2(T)} &= (\nabla Rv_h, \nabla \varphi_k)_{L^2(T)} - \langle Rv_{h|T}, \nabla \varphi_k \cdot \nu_T \rangle_{L^2(\partial T)} \\ &= -(v_{\mathcal{T}}, \Delta \varphi_k)_{L^2(T)} + \sum_{F \in \mathcal{F}(T)} \langle v_{\mathcal{F}} - Rv_{h|T}, \nabla \varphi_k \cdot \nu_T \rangle_{L^2(F)}. \end{aligned} \quad (51)$$

Since  $\Delta\varphi_k \in P_k(T)$ , the  $L^2$  projection  $\Pi_k$  on the right-hand side of (50) is redundant. Hence, the combination of (50)–(51) with  $\nabla\varphi_k \cdot \nu_{T|F} \in P_k(F)$  for all  $F \in \mathcal{F}(T)$  results in

$$\|\nabla\varphi_k\|_{L^2(T)}^2 = \sum_{F \in \mathcal{F}(T)} \langle \Pi_{F,k}(v_{\mathcal{F}} - Rv_{h|T} + \varphi_k), \nabla\varphi_k \cdot \nu_{T|F} \rangle_{L^2(F)}. \quad (52)$$

A Cauchy inequality on the right-hand side of (52), a discrete trace inequality, and  $S_{TF}v_h = \Pi_{F,k}(Rv_{h|T} - v_{\mathcal{F}|F} - \varphi_k)$  for all  $F \in \mathcal{F}(T)$  lead to

$$\|\nabla\varphi_k\|_{L^2(T)}^2 \lesssim \sum_{F \in \mathcal{F}(T)} h_F^{-1} \|S_{TF}v_h\|_{L^2(F)}^2. \quad (53)$$

Since  $\tilde{s}_T(v_h, v_h) = \sum_{F \in \mathcal{F}(T)} h_F^{-1} \|\Pi_{F,k}(Rv_{h|T} - v_{\mathcal{F}})\|_{L^2(F)}^2 + h_T^{-2} \|\varphi_k\|_{L^2(T)}^2$ , the combination of (48)–(49) with (53) concludes the proof of (47).  $\square$

#### 4.4 Efficiency of the stabilization

The second improvement to the HHO literature is a quasi-best approximation estimate along the lines of the seminal paper [31, Theorem 4.10]. In combination with Theorem 4, this, in particular, provides the efficiency (38) of the stabilization up to data oscillation.

**Theorem 5 (quasi-best approximation up to data oscillation)** *For any  $p \in \mathbb{N}_0$ , the solution  $u$  to (3) and the discrete solution  $u_h$  to (18) satisfy*

$$\begin{aligned} \|u - Ru_h\|_{pw} + s_h(u_h, u_h)^{1/2} \\ \leq C_{12} \left( \min_{v_{k+1} \in P_{k+1}(\mathcal{T})} \|u - v_{k+1}\|_{pw} + \text{osc}_{k+p}(f, \mathcal{T}) \right). \end{aligned}$$

The constant  $C_{12}$  exclusively depends on  $k$ ,  $p$ , and the shape regularity of  $\mathcal{T}$ .

*Proof* Given  $k, p \in \mathbb{N}_0$ , let  $\tilde{u} \in V$  solve the Poisson model problem  $-\Delta\tilde{u} = \Pi_{k+p}f$  with the right-hand side  $\Pi_{k+p}f$ . Subsection 4.3 in [31] constructs a stable enriching operator  $\mathcal{J} : V_h \rightarrow V$  with local bubble-functions from [36]. A modification, where the polynomial degree  $k-1$  in [31, Eq. (4.16)] is replaced by  $k+p$ , leads to a right-inverse  $\mathcal{J} : V_h \rightarrow V$  of the interpolation  $\mathbb{I} : V_h \rightarrow V$  with the stability  $\|\mathcal{J}v_h\|^2 \lesssim a_h(v_h, v_h)$  and the additional  $L^2$  orthogonality  $\mathcal{J}v_h - v_{\mathcal{T}} \perp P_{k+p}(\mathcal{T})$  for all  $v_h = (v_{\mathcal{T}}, v_{\mathcal{F}}) \in V_h$ . The extra orthogonality allows for

$$(\Pi_{k+p}f, \mathcal{J}v_h)_{L^2(\Omega)} = (\Pi_{k+p}f, v_{\mathcal{T}})_{L^2(\Omega)} = (f, v_{\mathcal{T}})_{L^2(\Omega)} \quad \text{for all } (v_{\mathcal{T}}, v_{\mathcal{F}}) \in V_h.$$

Consequently, the smoother  $\mathcal{J}$  leads to a discrete solution  $u_h = (u_{\mathcal{T}}, u_{\mathcal{F}}) \in V_h$  in the modified HHO discretization of [31] as a quasi-best approximation of the above solution  $\tilde{u}$ . The point is that  $u_h \in V_h$  coincides with the original

HHO solution  $u_h$  from (18). The arguments from the proof of [31, Theorem 4.10] reveal the quasi-best approximation

$$\|\tilde{u} - Ru_h\|_{\text{pw}} + s_h(u_h, u_h)^{1/2} \lesssim \min_{v_{k+1} \in P_{k+1}(\mathcal{T})} \|\tilde{u} - v_{k+1}\|_{\text{pw}}$$

also for the above modified smoother  $\mathcal{J}$ . This, the triangle inequalities  $\|u - Ru_h\|_{\text{pw}} \leq \|u - \tilde{u}\| + \|\tilde{u} - Ru_h\|_{\text{pw}}$  and  $\|\tilde{u} - v_{k+1}\|_{\text{pw}} \leq \|u - \tilde{u}\| + \|u - v_{k+1}\|_{\text{pw}}$  for any  $v_{k+1} \in P_{k+1}(\mathcal{T})$ , and the standard perturbation bound  $\|u - \tilde{u}\| \leq C_P \text{osc}_{k+p}(f, \mathcal{T})$  conclude the proof.  $\square$

#### 4.5 Stabilization-free efficiency of averaging

The main result of this subsection establishes the stabilization-free efficiency of the nodal averaging technique.

**Theorem 6 (averaging is efficient)** *Let  $u \in V$  resp.  $u_h \in V_h$  solve (3) resp. (18). Then  $Ru_h$  and  $\mathcal{A}Ru_h$  satisfy, for any  $p \in \mathbb{N}_0$ , that*

$$C_{13}^{-1} \|(1 - \mathcal{A})Ru_h\|_{\text{pw}} \leq \|u - Ru_h\|_{\text{pw}} + \text{osc}_{k+p}(f, \mathcal{T}). \quad (54)$$

The constant  $C_{13}$  exclusively depends on  $k$ ,  $p$ , and the shape regularity of the triangulation  $\mathcal{T}$ .

The proof can follow the proof of [24, Theorem 4.7] but additionally utilizes the two significant improvements from Subsections 4.3–4.4 that allow a stabilization-free efficiency in (54).

*Proof* Theorem 4.7 in [24] shows  $\|(1 - \mathcal{A})Ru_h\|_{\text{pw}}^2 \lesssim \|u - Ru_h\|_{\text{pw}}^2 + \tilde{s}_h(u_h, u_h)$ . This, the equivalence  $\tilde{s}_h(u_h, u_h) \approx s_h(u_h, u_h)$  of stabilizations (from Theorem 4), and the efficiency  $s_h(u_h, u_h) \lesssim \|u - Ru_h\|_{\text{pw}}^2 + \text{osc}_{k+p}^2(f, \mathcal{T})$  (from Theorem 5) imply the assertion.  $\square$

*Remark 2 ( $p$ -robustness)* The  $H^1(\Omega)$ -conforming post-processing of  $Ru_h$  from [30] provides an efficiency constant independent of the polynomial degree  $k$ . The right-hand side of [30, Corollary 4.2] involves the stabilization-related term  $\sum_{F \in \mathcal{F}} h_F^{-1} \|\Pi_{F,0}[Ru_h]_F\|_{L^2(F)}^2$ . It remains an open question whether this term can be controlled  $p$ -robustly by  $\|u - Ru_h\|_{\text{pw}} + \text{osc}_k(f, \mathcal{T})$  (with a multiplicative constant independent of the polynomial degree  $k$ ).

#### 4.6 Proof of Theorem 3

Let  $p \in \mathbb{N}_0$  and  $r = 0$  if  $k = 0$  and  $r = k + p$  if  $k \geq 1$  as in (35) be given. Recall the abbreviation  $G_h = \nabla_{\text{pw}} Ru_h \in P_k(\mathcal{T})$  with the discrete solution  $u_h \in V_h$  to (18) and let  $Q_p = \sum_{z \in \mathcal{V}} Q_{z,h} \in H(\text{div}, \Omega)$  denote the flux reconstruction from Subsection 4.2. The proof establishes (36) in five steps.

Step 1 provides the GUB  $\|u - Ru_h\|_{\text{pw}} \leq \eta_{\text{eq},p}(\mathcal{T})$ . This can follow from the paradigm of [1, 2, 29] as outlined below. The choice  $G := G_h$  and  $w := \mathcal{A}Ru_h$  in (8) and a triangle inequality lead to

$$\|u - Ru_h\|_{\text{pw}}^2 \leq (\|f + \text{div } Q_p\|_* + \|\text{div}(Q_p - G_h)\|_*)^2 + \|(1 - \mathcal{A})Ru_h\|_{\text{pw}}^2.$$

Since  $\text{div } Q_p + \Pi_r f = 0$  vanishes in  $\Omega$  by (42), a piecewise Poincaré inequality shows  $\|f + \text{div } Q_p\|_* \leq C_P \text{osc}_r(f, \mathcal{T})$ . This, the bound  $\|\text{div}(Q_p - G_h)\|_* \leq \|Q_p - G_h\|$  from the definition of  $\|\bullet\|_*$ , and the previously displayed formula result in the reliability  $\|u - Ru_h\|_{\text{pw}} \leq \eta_{\text{eq},p}(\mathcal{T})$ .

Step 2 establishes  $\text{osc}_{k-1}(f, \mathcal{T}) \lesssim \eta_{\text{eq},p}(\mathcal{T})$ . Lemma 8 provides

$$\text{osc}_{k-1}(f, \mathcal{T}) \lesssim \|u - Ru_h\|_{\text{pw}} + \text{osc}_r(f, \mathcal{T}). \quad (55)$$

This, Step 1 and  $\text{osc}_r(f, \mathcal{T}) \lesssim \eta_{\text{eq},p}(\mathcal{T})$  from (34) conclude the proof of Step 2.

Step 3 reveals, for any  $q \in \mathbb{N}_0$ , the efficiency of the flux reconstruction

$$\|Q_p - G_h\| \lesssim \|u - Ru_h\|_{\text{pw}} + \text{osc}_q(\mathcal{T}, f) \quad (56)$$

for any polynomial degree  $k \geq 1$ . The case  $k = 0$  follows in Step 4 below. Recall  $f_z = \Pi_{k+p}(\varphi_z f - G_h \cdot \nabla \varphi_z)$  from (39) for any vertex  $z \in \mathcal{V}$  in the construction of  $Q_p$  from Subsection 4.2 and set  $\sigma_z := \mathcal{I}_{\text{RT}}(\varphi_z G_h) \in RT_{k+p}^{\text{pw}}(\mathcal{T}(z))$ . The piecewise Raviart-Thomas interpolation  $\mathcal{I}_{\text{RT}} : H^1(\mathcal{T}; \mathbb{R}^n) \rightarrow RT_{k+p}^{\text{pw}}(\mathcal{T})$  satisfies the well-known commuting diagram properties [7, Section 2.5.1]

$$\text{div}_{\text{pw}} \circ \mathcal{I}_{\text{RT}} = \Pi_{k+p} \circ \text{div}_{\text{pw}} \quad \text{and} \quad \gamma_T \circ \mathcal{I}_{\text{RT}}|_F = \Pi_{F,k+p} \circ \gamma_T \quad (57)$$

for any facet  $F \in \mathcal{F}(T)$  of a simplex  $T \in \mathcal{T}$  and the normal trace  $\gamma_T : H^1(T; \mathbb{R}^n) \rightarrow L^2(\partial T)$  with  $\gamma_T \sigma := \sigma \cdot \nu_T$  for  $\sigma \in H^1(T; \mathbb{R}^n)$ . This and elementary algebra with the product rule  $\text{div}_{\text{pw}}(\varphi_z G_h) = \varphi_z \text{div}_{\text{pw}} G_h + \nabla \varphi_z \cdot G_h \in P_k(\mathcal{T}(z))$  imply

$$r_z := -\text{div}_{\text{pw}} \sigma_z - f_z = -\varphi_z \text{div}_{\text{pw}} G_h - \Pi_{k+p}(\varphi_z f) \in P_{k+p}(\mathcal{T}(z)). \quad (58)$$

Recall the residual  $\text{Res}_z(v)$  with  $v \in H^1(\Omega)$  for the vertex  $z \in \mathcal{V}$  from (44). The commuting diagram property (57)–(58) establish the identity

$$\text{Res}_z(1) = -(\text{div}_{\text{pw}} G_h + f, \varphi_z)_{L^2(\omega(z))} + \sum_{T \in \mathcal{T}(z)} \langle G_h \cdot \nu_T, \varphi_z \rangle_{L^2(\partial T)}.$$

This, a piecewise integration by parts, and the property (20) verify  $\text{Res}(1) = (G_h, \nabla \varphi_z)_{L^2(\omega(z))} - (f, \varphi_z)_{L^2(\omega(z))} = 0$  for any interior vertex  $z \in \mathcal{V}(\Omega)$ . Hence, Lemma 9 applies for any vertex  $z \in \mathcal{V}$  and provides

$$\|Q_{z,h} - \mathcal{I}_{\text{RT}}(\varphi_z G_h)\|_{L^2(\omega(z))} \leq C_s \sup_{\substack{v \in H_*^1(\omega(z)) \\ \|\nabla v\|_{L^2(\omega(z))} = 1}} \text{Res}_z(v) \quad (59)$$

for the local contributions  $Q_{z,h}$  of  $Q_p = \sum_{z \in \mathcal{V}} Q_{z,h}$  from (41). The identity  $\mathcal{I}_{RT}(\varphi_z G_h) = \varphi_z G_h$  for  $p \geq 1$  from  $\varphi_z G_h \in P_{k+1}(\mathcal{T}(z)) \subset RT_{k+p}^{\text{pw}}(\mathcal{T}(z))$  allows for a  $k$ - and  $p$ -robust efficiency control of the flux reconstruction error

$$C_{14}^{-1} \|Q_p - G_h\|^2 \leq \|\nabla u - G_h\|^2 + \sum_{z \in \mathcal{V}} \text{osc}_r^2(\varphi_z f, \mathcal{T}(z)) \quad (60)$$

with a constant  $C_{14}$  that solely depends on the shape regularity of  $\mathcal{T}$ . This is deemed noteworthy and motivates two different approaches for the bound of the residual  $\text{Res}_z(v)$  on the right-hand side of (59) for  $p = 0$  and  $p \geq 1$  below.

Step 3.1 provides (56) for  $p = 0$ . Given any normalized  $v \in H_*^1(\omega(z))$  with  $\|\nabla v\|_{L^2(\omega(z))} = 1$ , the product  $\mathcal{I}_{RT}(\varphi_z G_h) \cdot \nu_F v$  vanishes on any boundary facet  $F \in \mathcal{F}(\partial\omega(z))$  of the patch  $\omega(z)$ . This, the commuting diagram property (57), and (58) with  $\varphi_z \text{div}_{\text{pw}} G_h \in P_k(\mathcal{T}(z))$  in the definition of the residual  $\text{Res}_z(v)$  from (44) verify

$$\begin{aligned} \text{Res}_z(v) &= -(\varphi_z f + \varphi_z \text{div}_{\text{pw}} G_h, \Pi_k v)_{L^2(\omega(z))} \\ &\quad + \sum_{F \in \mathcal{F}(z) \cap \mathcal{F}(\Omega)} \langle \varphi_z [G_h]_F \cdot \nu_F, \Pi_{F,k} v \rangle_{L^2(F)}. \end{aligned} \quad (61)$$

The shape regularity of  $\mathcal{T}$ , a Poincaré inequality for interior vertices  $z \in \mathcal{V}(\Omega)$ , and a Friedrichs inequality for boundary vertices  $z \in \mathcal{V}(\partial\Omega)$  provide

$$\|h_{\mathcal{T}}^{-1} v\|_{L^2(\omega(z))} \approx \text{diam}(\omega(z))^{-1} \|v\|_{L^2(\omega(z))} \lesssim \|\nabla v\|_{L^2(\omega(z))} = 1. \quad (62)$$

Given any facet  $F \in \mathcal{F}(z) \cap \mathcal{F}(\Omega)$  in the facet spider  $F \in \mathcal{F}(z)$  with facet patch  $\omega(F)$ , a trace inequality thus shows

$$h_F^{-1/2} \|v\|_{L^2(F)} \lesssim \|h_{\mathcal{T}}^{-1} v\|_{L^2(\omega(F))} + \|\nabla v\|_{L^2(\omega(F))} \lesssim 1. \quad (63)$$

Cauchy inequalities on the right-hand side of (61), the stability of the  $L^2$  projection,  $\|\varphi_z\|_{L^\infty(\omega_z)} = 1$ , and (62)–(63) prove that  $\text{Res}_z(v)$  is controlled by

$$\|h_{\mathcal{T}}(f + \text{div}_{\text{pw}} G_h)\|_{L^2(\omega(z))} + \left( \sum_{F \in \mathcal{F}(z) \cap \mathcal{F}(\Omega)} h_F \|[G_h]_F \cdot \nu_F\|_{L^2(F)}^2 \right)^{1/2}.$$

The efficiency of those residual terms follows from the proof of Theorem 2 in Section 3. In combination with (43) and (59), this results in the global efficiency (56) for  $p = 0$  and concludes the proof of Step 3.1.

Step 3.2 provides (56) for  $p \geq 1$ . Given any normalized  $v \in H_*^1(\omega(z))$  with  $\|\nabla v\|_{L^2(\omega(z))} = 1$ , (58) and the identity  $\mathcal{I}_{RT}(\varphi_z G_h) = \varphi_z G_h$  in the definition of the residual  $\text{Res}_z(v)$  from (44) verify that  $\text{Res}_z(v)$  is equal to

$$-(\varphi_z \text{div}_{\text{pw}} G_h + \Pi_{k+p}(\varphi_z f), v)_{L^2(\omega(z))} + \sum_{T \in \mathcal{T}(z)} \langle G_h \cdot \nu_T, \varphi_z v \rangle_{L^2(\partial T)}.$$



This, a piecewise integration by parts, and the product rule  $\nabla(\varphi_z v) = \varphi_z \nabla v + v \nabla \varphi_z$  reveal

$$\text{Res}_z(v) = -(\Pi_{k+p}(\varphi_z f), v)_{L^2(\omega(z))} + (G_h, \nabla(\varphi_z v))_{L^2(\omega(z))}. \quad (64)$$

The weak formulation (3) with the test function  $\varphi_z v \in H_0^1(\omega(z)) \subset V$  shows  $(\nabla u, \nabla(\varphi_z v))_{L^2(\omega(z))} = (f, \varphi_z v)_{L^2(\omega(z))}$ . Consequently, (64) implies

$$\text{Res}_z(v) = ((1 - \Pi_{k+p})(\varphi_z f), v)_{L^2(\omega(z))} - (\nabla u - G_h, \nabla(\varphi_z v))_{L^2(\omega(z))}.$$

This, a Cauchy-Schwarz, and a piecewise Poincaré inequality with the normalization  $\|\nabla v\|_{L^2(\omega(z))} = 1$  provide

$$\text{Res}_z(v) \leq \|\nabla u - G_h\|_{L^2(\omega(z))} \|\nabla(\varphi_z v)\|_{L^2(\omega(z))} + C_P \text{osc}_{k+p}(\varphi_z f, \mathcal{T}(z)). \quad (65)$$

Since  $\|\nabla \varphi_z\|_{L^\infty(\omega(z))} \approx h^{-1}$ , the Leibniz rule, a triangle inequality, and (62) show that

$$\begin{aligned} \|\nabla(\varphi_z v)\|_{L^2(\omega(z))} &\leq \|\varphi_z\|_{L^\infty(\omega(z))} \|\nabla v\|_{L^2(\omega(z))} + \|\nabla \varphi_z\|_{L^\infty(\omega(z))} \|v\|_{L^2(\omega(z))} \\ &\lesssim \|\nabla v\|_{L^2(\omega(z))} = 1 \end{aligned}$$

can be bounded by a constant independent of  $h$ . Thus, the combination of (43) with (59) and (65) results in the efficiency of  $\|Q_p - G_h\|$  in (60) with a  $k$ - and  $p$ -robust constant  $C_{14}$ . Since  $\varphi_z \Pi_{k-1} f \in P_k(T) \subset P_{k+p}(T)$ , the Pythagoras theorem and  $\|\varphi_z\|_{L^\infty(\omega(z))} = 1$  show  $\|(1 - \Pi_{k+p})(\varphi_z f)\|_{L^2(T)} \leq \|\varphi_z(1 - \Pi_{k-1})f\|_{L^2(T)} \leq \|(1 - \Pi_{k-1})f\|_{L^2(T)}$  for all  $T \in \mathcal{T}(z)$ , whence

$$\text{osc}_{k+p}(\varphi_z f, \mathcal{T}(z)) \leq \text{osc}_{k-1}(f, \mathcal{T}(z)).$$

This and Lemma 8 implies the efficiency

$$\text{osc}_{k+p}(\varphi_z f, \mathcal{T}(z)) \lesssim \|\nabla_{\text{pw}}(u - Ru_h)\|_{L^2(\omega(z))} + \text{osc}_q(f, \mathcal{T}(z))$$

of the local oscillations from (65) for any  $q \in \mathbb{N}_0$  as in Remark 1. This and (60) prove (56) for  $p \geq 1$  and conclude the proof of Step 3.2.

Step 4 affirms the efficiency of the flux reconstruction

$$C_{15}^{-1} \|Q_p - G_h\| \leq \|u - Ru_h\|_{\text{pw}} + \text{osc}_q(f, \mathcal{T}) \quad (66)$$

for the polynomial degree  $k = 0$  and any  $q \in \mathbb{N}_0$ . Let  $\tilde{u} \in V$  solve the Poisson model problem  $-\Delta \tilde{u} = \Pi_0 f$  with piecewise constant right-hand side  $\Pi_0 f \in P_0(\mathcal{T})$ . A careful inspection reveals that all arguments from Step 3 apply to the case  $k = 0$  for  $f$  replaced by  $\Pi_0 f$ . This leads to  $\|Q_p - G_h\| \leq C_{15} \|\tilde{u} - Ru_h\|_{\text{pw}}$  with a constant  $C_{15}$  that solely depends on the shape of  $\mathcal{T}$  (because the data oscillations on the right-hand side of (60) vanish). Therefore, the triangle inequality  $\|\tilde{u} - Ru_h\|_{\text{pw}} \leq \|u - \tilde{u}\| + \|u - Ru_h\|_{\text{pw}}$ , the standard bound  $\|u - \tilde{u}\| \leq C_P \text{osc}_0(f, \mathcal{T})$ , and  $C_P = 1/\pi < 1$  on simplicial domains lead to (66). This and the efficiency of the data oscillations from Lemma 8 conclude the proof of Step 4. Notice that the constant  $C_{15}$  is independent of the parameter  $p$ .

Step 5 finishes the proof. On the one hand, the reliability  $\|u - Ru_h\|_{\text{pw}} + \text{osc}_{k-1}(f, \mathcal{T}) \lesssim \eta_{\text{eq},p}$  is established in Step 1–2. On the other hand, the efficiency of the averaging operator from Theorem 6, Lemma 8, and the efficiency of the flux reconstruction in (56) for  $k \geq 1$  and in (66) for  $k = 0$  imply the efficiency  $\eta_{\text{eq},p} \lesssim \|u - Ru_h\|_{\text{pw}} + \text{osc}_q(f, \mathcal{T})$  for any  $q \in \mathbb{N}_0$ . This concludes the proof of Theorem 3.  $\square$

## 5 Numerical experiments

This section provides numerical evidence for optimal convergence and a comparison of the stabilization-free GUBs  $\eta_{\text{res}}$  and  $\eta_{\text{eq},p}$  from the Sections 3 and 4 with the original error estimator  $\eta_{\text{HHO}}$  from [24, Theorem 4.3] for the HHO method in three 2D benchmarks.

### 5.1 A posteriori error estimation with explicit constants

All triangulations in this section consist of right-isosceles triangles with the Poincaré constant  $C_P = (\sqrt{2}\pi)^{-1}$  [33]. With the estimates  $C_1 \leq C_{\mathcal{T}}$ ,  $C_H \leq 1$ , and  $C_2 \leq C_{\mathcal{F}}$  from Example 1, the residual-based error estimator from (22) reads

$$\eta_{\text{res}}(\mathcal{T})^2 := (C_{\mathcal{T}}\eta_{\text{res},1}(\mathcal{T}) + C_P\eta_{\text{res},2}(\mathcal{T}) + C_{\mathcal{F}}\eta_{\text{res},3}(\mathcal{T}))^2 + C_{\mathcal{F}}^2\eta_{\text{res},4}^2(\mathcal{T}).$$

The equilibrated GUB from Theorem 3,

$$\eta_{\text{eq},p}^2(\mathcal{T}) := (C_P\text{osc}_{k+p}(f, \mathcal{T}) + \|Q_p^\Delta\|_{L^2(\Omega)})^2 + \|(1 - \mathcal{A})Ru_h\|_{\text{pw}}^2,$$

depends on the post-processed quantity  $Q_p^\Delta := Q_p - \nabla_{\text{pw}} Ru_h$  with the Raviart-Thomas function  $Q_p \in RT_{k+p}(\mathcal{T})$  of degree  $k + p$  for  $p \in \mathbb{N}_0$  from Subsection 4.2. An algorithmic description of the computation of  $Q_p^\Delta$  for arbitrary  $p$  follows in Appendix A. Theorem 3 shows that  $\eta_{\text{eq},p}$  is efficient and reliable for all  $p \in \mathbb{N}_0$ . The residual-based error estimator  $\eta_{\text{HHO}}$  from [24, Theorem 4.3] reads

$$\begin{aligned} \eta_{\text{HHO}}^2(\mathcal{T}) := & \sum_{T \in \mathcal{T}} \left( C_P h_T \|(I - \Pi_{T,0})(f + \Delta_{\text{pw}} Ru_h)\|_{L^2(T)} \right. \\ & \left. + \sqrt{\sum_{F \in \mathcal{F}(T)} C_{\partial T} h_T \|R_{T,F}^k u_h\|_{L^2(F)}^2} \right)^2 + \|(I - \mathcal{A})Ru_h\|_{\text{pw}}^2 \end{aligned}$$

with the operator  $R_{T,F}^k$  from [24, Eq. (2.59)] for the original HHO stabilization [24, Eq. (2.22)] that induces the global stabilization  $s_h$ . The constant  $C_{\partial T} = 12C_P(C_P + C_{\text{tr}}) \leq 2.0315$  for right-isosceles triangles bounds the trace of  $f - \Pi_0 f$  for  $f \in H^1(T)$  and improves on the estimate  $C_{\partial T} \leq (C_P(h_T |\partial T|/|T|)(1 + C_P))^2 = 2.524$  from [24, Subs. 4.1.1].

**Lemma 10 (Poincaré-type inequality on trace)** *Given a simplex  $T \subset \mathbb{R}^n$ , any  $f \in H^1(T)$  and  $C_{\partial T} := (n+1)h_T^2|T|^{-1}C_P(C_P + 2C_{\text{tr}}/n)$  satisfy*

$$\|f - \Pi_0 f\|_{L^2(\partial T)}^2 \leq C_{\partial T} h_T \|\nabla f\|_{L^2(T)}^2. \quad (67)$$

*Proof* Abbreviate  $\tilde{\ell}(F) := (n+1)h_F h_T^2 |T|^{-1}$  for the facet  $F \in \mathcal{F}(T)$  of  $T$  and apply Lemma 3 to the singleton triangulation  $\{T\}$  and  $f - \Pi_0 f \in H^1(T)$ . This and the Poincaré inequality  $\|f - \Pi_{0,T} f\|_{L^2(T)} \leq C_P h_T \|\nabla f\|_{L^2(T)}$  show

$$\sum_{F \in \mathcal{F}(T)} \tilde{\ell}(F)^{-1} \|f - \Pi_{0,T} f\|_{L^2(F)}^2 \leq (C_P + 2C_{\text{tr}}/n) C_P \|\nabla f\|_{L^2(T)}^2.$$

The assertion (67) follows from the observation that  $\tilde{\ell}(F)$  is maximized on the facet  $F \in \mathcal{F}(T)$  with  $h_F = h_T$ .  $\square$

## 5.2 Implementation and adaptive algorithm

Our implementation of the HHO method in MATLAB uses nodal bases for the spaces  $P_k(\mathcal{F})$ ,  $P_k(\mathcal{T})$ , and  $P_{k+1}(\mathcal{T})$  and the direct solver *mldivide* (behind the  $\backslash$ -operator) for the discrete system of equations representing (18). For implementation details on the HHO method itself we refer to [24, Appendix B]. For the numerical integration over the reference triangle  $T_{\text{ref}} := \text{conv}\{(0,0), (1,0), (0,1)\}$ , the product quadrature rule

$$Q(g) := \frac{1}{8} \sum_{j=1}^m w_j^0 \sum_{k=1}^m w_k^1 g(F(q_j^0, q_k^1)) \quad \text{for } g \in C(T_{\text{ref}}) \quad (68)$$

with the Duffy transform  $F : (-1,1)^2 \rightarrow T_{\text{ref}}$  from the square to  $T_{\text{ref}}$ , defined by  $F(X, Y) := ((1+X)(1-Y)/4, (1+Y)/2)$ , is applied. Here,  $q_j^\alpha \in (-1,1)$  (resp.  $w_j^\alpha \in \mathbb{R}$ ) for  $j = 1, \dots, m$  denote the  $m \in \mathbb{N}$  points (resp. weights) of the Gauß-Jacobi quadrature over the interval  $(-1,1)$  with respect to the weight function  $w(x) = (1-x)^\alpha$ ,  $\alpha = 0, 1$ . It is well-known that the Gauß-Jacobi quadrature is exact up to degree  $2m-1$  and an integration by substitution verifies

$$\int_{T_{\text{ref}}} g \, dx = \frac{1}{8} \int_{-1}^1 \int_{-1}^1 (1-Y) g(F(X, Y)) \, dY \, dX.$$

This shows that the quadrature rule (68) for the integration of  $g$  over  $T_{\text{ref}}$  is exact for polynomials  $g \in P_{2m-1}(T_{\text{ref}})$  of total degree at most  $2m-1$  and an integration by substitution allows the exact numerical integration of polynomials over  $T \in \mathcal{T}$ . The errors in approximating non-polynomial expressions, such as exact solutions  $u$  and source terms  $f$ , by polynomials of sufficiently high degree are expected to be very small and are neglected for simplicity.

Algorithm 1 displays the standard adaptive algorithm (AFEM) [16,22] driven by the refinement indicators, for any triangle  $T \in \mathcal{T}$ ,

$$\eta_{\text{res}}^2(T) := |T| \|f + \Delta_{\text{pw}} Ru_h\|_{L^2(T)}^2 + |T|^{1/2} \sum_{F \in \mathcal{F}(T)} \|[\nabla_{\text{pw}} Ru_h]_F\|_{L^2(F)}^2 \quad (69)$$

with the modified jump  $[\bullet]_F := \bullet \times n_F$  along a boundary side  $F \in \mathcal{F}(\partial\Omega)$ .

---

**Algorithm 1** AFEM algorithm
 

---

**Input:** Initial regular triangulation  $\mathcal{T}_0$  and polynomial degree  $k \in \mathbb{N}_0$  of the HHO method

**for** levels  $\ell := 0, 1, 2, \dots$  **do**

**Solve** (18) for discrete solution  $u_\ell \in V_\ell$  exactly on  $\mathcal{T}_\ell$  and compute  $Ru_\ell$

**Compute** (refinement indicators)  $\eta_{\text{res}}^2(T)$  for all  $T \in \mathcal{T}_\ell$

**Mark** minimal subset  $\mathcal{M}_\ell \subset \mathcal{T}_\ell$  with  $\frac{1}{2} \sum_{T \in \mathcal{T}_\ell} \eta_{\text{res}}(T)^2 \leq \sum_{T \in \mathcal{M}_\ell} \eta_{\text{res}}(T)^2$

**Refine**  $\mathcal{T}_\ell$  to smallest NVB refinement  $\mathcal{T}_{\ell+1}$  with  $\mathcal{M}_\ell \subseteq \mathcal{T}_\ell \setminus \mathcal{T}_{\ell+1}$

**end for**

**Output:** sequences of triangulations  $\mathcal{T}_\ell$  and  $Ru_\ell$

---

### 5.3 High oscillations on the unit square

This benchmark on the unit square  $\Omega := (0, 1)^2$  considers the Laplace equation  $-\Delta u = f$  with source term  $f$  matching the smooth exact solution

$$u(x, y) = x(x-1)y(y-1)e^{-100((x-1/2)^2 + (y-117/1000)^2)} \in C^\infty(\Omega).$$

Figure 1 displays the energy norm  $\|e\|_{\text{pw}}$  of the error  $e := u - Ru_h$  for uniform and adaptive refinement by Algorithm 1 on the left. The smooth solution allows for optimal convergence rates  $(k+1)/2$  in the number ndof of degrees of freedom, while the adaptive mesh sequence leads to a lower energy error with respect to ndof. The GUBs  $\eta_{\text{res}}$ ,  $\eta_{\text{eq},p}$ , and  $\eta_{\text{HHO}}$  are efficient and therefore equivalent to the energy error  $\|e\|_{\text{pw}}$ , see Figure 1 on the right for  $k = 0, 2$ .


Figure 2 shows the efficiency indices  $EF(\eta) := \eta / \|e\|_{\text{pw}}$  for the residual-based GUBs  $\eta = \eta_{\text{res}}, \eta_{\text{HHO}}$  and the equilibration-based GUB  $\eta = \eta_{\text{eq},p}$  for  $p = 0, 1$ . Higher values of  $p$  for a more expensive postprocessing in  $\eta_{\text{eq},p}$  do not significantly improve on  $\eta_{\text{eq},1}$ .

### 5.4 Analytical solution for the slit domain


The source term  $f \in L^2(\Omega)$  in the second benchmark on the slit domain  $\Omega := (0, 1)^2 \setminus ([0, 1] \times \{0\})$  matches the singular solution (in polar coordinates)

$$u(r, \varphi) = r^{1/2}(r^2 \sin(\varphi)^2 - 1)(r^2 \cos(\varphi)^2 - 1) \sin(\varphi/2).$$

The singularity of  $u$  at the origin  $(0, 0)$  leads to reduced convergence rates  $1/4$  under uniform refinement, regardless of the polynomial degree  $k$ . Figure




**Fig. 1** Convergence history of the energy error  $\|e\|_{pw}$  (left) and the GUBs  $\eta_{res}$ ,  $\eta_{HHO}$ ,  $\eta_{eq,0}$  (right) on the square domain




**Fig. 2** History of the overestimation factor  $EF(\eta) = \eta/\|e\|_{pw}$  for the residual-based error estimators  $\eta_{res}$ ,  $\eta_{HHO}$  (left) and  $\eta_{eq,0}$ ,  $\eta_{eq,1}$  (right) on the unit square

3 shows that the adaptive algorithm recovers optimal rates and verifies the equivalence of the GUBs  $\eta_{res}$ ,  $\eta_{eq,p}$ , and  $\eta_{HHO}$  to the energy error  $\|e\|_{pw}$ .

The efficiency indices in Figure 4 show a strong overestimation by  $\eta_{res}$  in the preasymptotic regime (undisplayed) with values  $EF(\eta_{res}) > 60$ . However, asymptotically the quotients  $EF(\eta) = \eta/\|e\|_{pw}$  for the two residual-based GUBs  $\eta_{res}$ ,  $\eta_{HHO}$  differ only by a factor 10, while the equilibrated GUBs  $\eta_{eq,p}$  provide the closest values to 1.




**Fig. 3** Convergence history of the energy error  $\|e\|_{pw}$  (left) and the GUBs  $\eta_{res}, \eta_{HHO}, \eta_{eq,0}$  (right) on the slit domain



**Fig. 4** History of the overestimation factor  $EF(\eta) = \eta/\|e\|_{pw}$  for the residual-based error estimators  $\eta_{res}, \eta_{HHO}$  (left) and  $\eta_{eq,0}, \eta_{eq,1}$  (right) on the slit domain

### 5.5 Corner singularity in the L-shaped domain

The third benchmark problem is set in the L-shaped domain  $\Omega = (-1, 1)^2 \setminus [0, 1)^2$  with constant right-hand side  $f \equiv 1$ . Figure 5 displays the convergence history of the error  $e := u - Ru_h$  and compares the adaptive scheme, Algorithm



**Fig. 5** Convergence history plot of the energy error  $\|e\|_{\text{pw}}$  on the L-shaped domain with uniform and adaptive refinement with AFEM, driven by  $\eta_{\text{res}}(T)$ , (left) and for AFEM, driven by  $\eta_{\text{res}}(T)$ ,  $\eta_{\text{HHO}}(T)$ , and  $\eta_{\text{eq},0}(T)$ , (right).


1, driven by the refinement indicators  $\eta_{\text{res}}(T)$  from (69) and

$$\begin{aligned} \eta_{\text{HHO}}^2(T) &:= |T| \|(I - \Pi_0)(f + \Delta_{\text{pw}} Ru_h)\|_{L^2(T)}^2 + \|\nabla(1 - \mathcal{A})Ru_h\|_{L^2(T)}^2 \\ &\quad + |T|^{1/2} \sum_{F \in \mathcal{F}(T)} \|R_{T,F}^k u_h\|_{L^2(F)}^2, \\ \eta_{\text{eq},0}^2(T) &:= \text{osc}_k^2(f, T) + \|Q_0^\Delta\|_{L^2(T)}^2 + \|\nabla(1 - \mathcal{A})Ru_h\|_{L^2(T)}^2 \end{aligned}$$

for  $T \in \mathcal{T}$  that are induced from the GUB  $\eta_{\text{res}}$ ,  $\eta_{\text{HHO}}$ , and  $\eta_{\text{eq},0}$ . Here, the norm  $\|e\|_{\text{pw}}$  of the distance  $e$  from the discrete solution  $Ru_h \in P_{k+1}(\mathcal{T})$  over  $\mathcal{T}$  to the unknown solution  $u \in H^1(\Omega)$  is approximated by  $\|\hat{u} - Ru_h\|_{\text{pw}}$ , where  $\hat{u} \in P_{k+1}(\hat{\mathcal{T}})$  is the HHO approximation of  $u$  on an adaptive refinement  $\hat{\mathcal{T}}$  of  $\mathcal{T}$  with at least  $2|\mathcal{T}| \leq |\hat{\mathcal{T}}|$  elements. The three adaptive schemes (Algorithm 1, driven by  $\eta_{\text{res}}(T)$ ,  $\eta_{\text{HHO}}(T)$ , or  $\eta_{\text{eq},0}(T)$ ) recover optimal rates of convergence and lead to similar local refinement of the adaptive mesh sequences as in Figure 6.

## 5.6 Conclusion

The adaptive mesh-refining algorithm recovers optimal convergence rates in all three benchmarks. This holds for AFEM driven by any of the three refinement indicators derived from the GUB  $\eta_{\text{res}}$ ,  $\eta_{\text{HHO}}$ , and  $\eta_{\text{eq},p}$ . The generated mesh sequences from the adaptive schemes, driven by the different estimators, display a very similar concentration of the local mesh-refinement as in Figure 6. All three benchmarks verify that the considered error estimators are GUB with reliability constant 1 while the post-processing in the equilibrated GUB  $\eta_{\text{eq},p}$  produces minimal overestimation.



**Fig. 6** Adaptive triangulations on the L-shaped domain for  $k = 3$  from AFEM, driven by  $\eta_{\text{res}}(T)$  (left,  $|\mathcal{T}| = 882$ ), driven by  $\eta_{\text{HHO}}(T)$  (middle,  $|\mathcal{T}| = 907$ ), and driven by  $\eta_{\text{eq},0}(T)$  (right,  $|\mathcal{T}| = 919$ ).

## References

1. Ainsworth, M.: Robust a posteriori error estimation for nonconforming finite element approximation. *SIAM J. Numer. Anal.* **42**(6), 2320–2341 (2005)
2. Ainsworth, M.: A posteriori error estimation for lowest order Raviart-Thomas mixed finite elements. *SIAM J. Sci. Comput.* **30**, 189–204 (2007)
3. Ainsworth, M., Oden, J.T.: A unified approach to a posteriori error estimation using element residual methods. *Numer. Math.* **65**, 23–50 (1993)
4. Alonso, A.: Error estimators for a mixed method. *Numer. Math.* **74**(4), 385–395 (1996)
5. Bertrand, F., Boffi, D.: The Prager-Synge theorem in reconstruction based a posteriori error estimation. In: 75 years of mathematics of computation, vol. 754, pp. 45–67. Amer. Math. Soc., [Providence], RI (2020)
6. Bertrand, F., Kober, B., Moldenhauer, M., Starke, G.: Weakly symmetric stress equilibration and a posteriori error estimation for linear elasticity. *Numer. Methods Partial Differential Equations* **37**(4), 2783–2802 (2021)
7. Boffi, D., Brezzi, F., Fortin, M.: *Mixed finite element methods and applications*, vol. 44. Springer, Heidelberg (2013)
8. Bonito, A., Nochetto, R.H.: Quasi-optimal convergence rate of an adaptive discontinuous Galerkin method. *SIAM J. Numer. Anal.* **48**(2), 734–771 (2010)
9. Braess, D.: *Finite Elements: theory, fast solvers, and applications in Solid Mechanics*, third edn. Cambridge University Press, Cambridge (2007)
10. Braess, D., Pillwein, V., Schöberl, J.: Equilibrated residual error estimates are  $p$ -robust. *Comput. Methods Appl. Mech. Engrg.* **198**, 1189–1197 (2009)
11. Brezzi, F., Fortin, M.: *Mixed and hybrid finite element methods*, vol. 15. Springer, New York (1991)
12. Cai, Z., Zhang, S.: Robust equilibrated residual error estimator for diffusion problems: conforming elements. *SIAM J. Numer. Anal.* **50**(1), 151–170 (2012)
13. Carstensen, C.: A posteriori error estimate for the mixed finite element method. *Math. Comp.* **66**(218), 465–476 (1997)
14. Carstensen, C.: A unifying theory of a posteriori finite element error control. *Numer. Math.* **100**(4), 617–637 (2005)
15. Carstensen, C., Ern, A., Puttkammer, S.: Guaranteed lower bounds on eigenvalues of elliptic operators with a hybrid high-order method. *Numer. Math.* **149**(2), 273–304 (2021)
16. Carstensen, C., Feischl, M., Page, M., Praetorius, D.: Axioms of adaptivity. *Comput. Math. Appl.* **67**(6), 1195–1253 (2014)
17. Carstensen, C., Gedicke, J., Rim, D.: Explicit error estimates for Courant, Crouzeix-Raviart and Raviart-Thomas finite element methods. *J. Comput. Math.* **30**(4), 337–353 (2012)
18. Carstensen, C., Gudi, T., Jensen, M.: A unifying theory of a posteriori error control for discontinuous Galerkin FEM. *Numer. Math.* **112**(3), 363–379 (2009)



19. Carstensen, C., Hellwig, F.: Constants in discrete Poincaré and Friedrichs inequalities and discrete quasi-interpolation. *Comput. Methods Appl. Math.* **18**(3), 433–450 (2018)
20. Carstensen, C., Hu, J.: A unifying theory of a posteriori error control for nonconforming finite element methods. *Numer. Math.* **107**(3), 473–502 (2007)
21. Carstensen, C., Peterseim, D., Schröder, A.: The norm of a discretized gradient in  $H(\text{div})^*$  for a posteriori finite element error analysis. *Numer. Math.* **132**, 519–539 (2016)
22. Carstensen, C., Rabus, H.: Axioms of adaptivity with separate marking for data resolution. *SIAM J. Numer. Anal.* **55**(6), 2644–2665 (2017)
23. Ciarlet, P., Dunkl, C.F., Sauter, S.A.: A family of Crouzeix-Raviart finite elements in 3D. *Anal. Appl. (Singap.)* **16**(5), 649–691 (2018)
24. Di Pietro, D.A., Droniou, J.: The hybrid high-order method for polytopal meshes, vol. 19. Springer, Cham (2020)
25. Di Pietro, D.A., Ern, A.: A hybrid high-order locking-free method for linear elasticity on general meshes. *Comput. Methods Appl. Mech. Engrg.* **283**, 1–21 (2015)
26. Di Pietro, D.A., Ern, A., Lemaire, S.: An arbitrary-order and compact-stencil discretization of diffusion on general meshes based on local reconstruction operators. *Comput. Methods Appl. Math.* **14**(4), 461–472 (2014)
27. Ern, A., Guermond, J.L.: Finite elements I—Approximation and interpolation, vol. 72. Springer, Cham (2021)
28. Ern, A., Guermond, J.L.: Finite elements II—Galerkin approximation, elliptic and mixed PDEs, vol. 73. Springer, Cham (2021)
29. Ern, A., Vohralík, M.: Polynomial-degree-robust a posteriori estimates in a unified setting for conforming, nonconforming, discontinuous Galerkin, and mixed discretizations. *SIAM J. Numer. Anal.* **53**(2), 1058–1081 (2015)
30. Ern, A., Vohralík, M.: Stable broken  $H^1$  and  $H(\text{div})$  polynomial extensions for polynomial-degree-robust potential and flux reconstruction in three space dimensions. *Math. Comp.* **89**(322), 551–594 (2020)
31. Ern, A., Zanotti, P.: A quasi-optimal variant of the hybrid high-order method for elliptic partial differential equations with  $H^{-1}$  loads. *IMA J. Numer. Anal.* **40**(4), 2163–2188 (2020)
32. Girault, V., Raviart, P.A.: Finite element methods for Navier-Stokes equations, vol. 5. Springer, New York (1986)
33. Kikuchi, F., Liu, X.: Estimation of interpolation error constants for the  $P_0$  and  $P_1$  triangular finite elements. *Comput. Methods Appl. Mech. Engrg.* **196**(37-40), 3750–3758 (2007)
34. da Veiga, L.B., Canuto, C., Nochetto, R.H., Vacca, G., Verani, M.: Adaptive VEM: Stabilization-free a posteriori error analysis (2021)
35. Verfürth, R.: A note on constant-free a posteriori error estimates. *SIAM J. Numer. Anal.* **47**(4), 3180–3194 (2009)
36. Verfürth, R.: A posteriori error estimation techniques for finite element methods. Oxford University Press, Oxford (2013)

## Appendix A: Equilibration algorithm for higher order

The post-processed quantity  $Q_p \in RT_{k+p}(\mathcal{T})$  from Subsection 4.2 enters the equilibrated error estimator  $\eta_{\text{eq},p}(\mathcal{T})$  in Theorem 3 and could be computed by a minimization problem on the vertex patches. The solution property (20) gives rise to the two cases  $r = 0$  if  $k = 0$  and  $r = k + p$  if  $k \geq 1$  for the polynomial degree  $r$  in the equilibrium  $\text{div } Q_p + \Pi_r f = 0$  in  $\Omega$  from (42). This appendix follows [35, 12, 6, 9] to compute the quantity of interest  $Q_p - \nabla_{\text{pw}} R u_h$  directly in an efficient two-step procedure in 2D. Throughout this appendix, fix  $k, p \in \mathbb{N}_0$  and abbreviate  $q := k + p$  and  $G_h := \nabla_{\text{pw}} R u_h \in P_k(\mathcal{T}; \mathbb{R}^2)$ . Let the data  $f \in L^2(\Omega)$  be given and assume, for the sake of brevity, that  $f \in P_0(\mathcal{T})$  if  $k = 0$ .

### A.1 Overview

Recall the definition (41) of the summand  $Q_{z,h}$  in  $Q_p := \sum_{z \in \mathcal{V}} Q_{z,h}$  from Subsection 4.2 with the piecewise Raviart-Thomas interpolation  $\mathcal{I}_{\text{RT}} : H^1(\mathcal{T}; \mathbb{R}^2) \rightarrow RT_q^{\text{pw}}(\mathcal{T})$  [7, Section III.3.1]. The focus is on one vertex  $z \in \mathcal{V}$  with vertex-patch  $\omega(z)$  and its triangulation  $\mathcal{T}(z) = \{T \in \mathcal{T} \mid z \in T\}$ . Consider set of edges  $\mathcal{F}$  and the facet-spider  $\mathcal{F}(z) = \{E \in \mathcal{F} \mid z \in E\}$  as in Figure 7. The nodal basis function  $\varphi_z \in S^1(\mathcal{T}(z))$  gives rise to the discrete spaces

$$RT_q^{\text{pw},0}(\mathcal{T}(z)) := \{\sigma_z \in RT_q^{\text{pw}}(\mathcal{T}(z)) : \sigma_z \cdot \nu_E = 0 \text{ for all } E \in \mathcal{F} \setminus \mathcal{F}(z)\},$$

$$\mathcal{S}(z) := \left\{ \sigma_z \in RT_q^{\text{pw},0}(z) : \begin{array}{ll} \text{div } \sigma_z &= -\Pi_{T,q}(\varphi_z(f + \text{div } G_h)) & \text{for all } T \in \mathcal{T}(z) \\ [\sigma_z \cdot \nu_E]_E &= -\Pi_{E,q}(\varphi_z[G_h \cdot \nu_E]_E) & \text{for all } E \in \mathcal{F}(\omega(z)) \end{array} \right\}.$$

**Proposition 1 (alternative minimization)** *It holds*

$$Q_{z,h}^\Delta := Q_{z,h} - \mathcal{I}_{\text{RT}}(\varphi_z G_h) = \arg \min_{\sigma_z \in \mathcal{S}(z)} \|\sigma_z\|_{L^2(\omega(z))}. \quad (70)$$

*Proof* Recall  $f_z = \Pi_{\tilde{p}}(\varphi_z f - G_h \cdot \nabla \varphi_z)$  from (39). (Notice that this formula coincide with the definition (39) for  $k = 0$  because  $f \in P_0(\mathcal{T})$ .) Given any  $\sigma_z \in \mathcal{S}(z)$ , the commuting diagram property  $\text{div}_{\text{pw}} \circ \mathcal{I}_{\text{RT}} = \Pi_q \circ \text{div}_{\text{pw}}$  [7, Proposition 2.5.2] shows


$$\text{div}_{\text{pw}}(\sigma_z + \mathcal{I}_{\text{RT}}(\varphi_z G_h)) = \text{div}_{\text{pw}} \sigma_z + \Pi_{\tilde{p}} \text{div}_{\text{pw}}(\varphi_z G_h) = -f_z. \quad (71)$$

By design of the interpolation  $\mathcal{I}_{\text{RT}}$ ,  $(\mathcal{I}_{\text{RT}}(\varphi_z G_h)|_T \cdot \nu_E)|_E = \Pi_{E,q}(\varphi_z G_h)|_T \cdot \nu_E$  holds and so  $[\mathcal{I}_{\text{RT}}(\varphi_z G_h) \cdot \nu_E]_E = \Pi_{E,q}(\varphi_z [G_h \cdot \nu_E]_E)$  follows for any  $E \in \mathcal{F}(T)$  and  $T \in \mathcal{T}$ . Therefore, the jump  $[\sigma_z + \mathcal{I}_{\text{RT}}(\varphi_z G_h)]_E \cdot \nu_E \equiv 0$  vanishes on  $E \in \mathcal{F}(\omega(z))$ , whence  $\sigma_z + \mathcal{I}_{\text{RT}}(\varphi_z G_h) \in RT_{\tilde{p}}(\mathcal{T}(z))$ . Since  $RT_{\tilde{p}}^{\text{pw},0}(\mathcal{T}(z)) \cap H(\text{div}, \omega(z)) = RT_{\tilde{p}}^0(\mathcal{T}(z))$ , this and (71) imply  $\sigma_z + \mathcal{I}_{\text{RT}}(\varphi_z G_h) \in \mathcal{Q}_h(z)$  for any  $\sigma_z \in \mathcal{S}(z)$ . In particular,  $\mathcal{S}(z) + \mathcal{I}_{\text{RT}}(\varphi_z G_h) \subseteq \mathcal{Q}_h(z)$ . On the other hand, similar arguments verify the reverse inclusion  $\mathcal{Q}_h(z) \subseteq \mathcal{S}(z) + \mathcal{I}_{\text{RT}}(\varphi_z G_h)$ . The substitution  $\mathcal{S}(z) + \mathcal{I}_{\text{RT}}(\varphi_z G_h) = \mathcal{Q}_h(z)$  in (41) concludes the proof.  $\square$

This establishes that the norm  $\|Q_p^\Delta\|$  of  $Q_p^\Delta := \sum_{z \in \mathcal{V}} Q_{z,h}^\Delta = Q_p - G_h$  contributes to the equilibrated error estimator and the remaining parts of this appendix compute the minimizer  $Q_{z,h}^\Delta$  of (70) in a two-step procedure. First, Algorithm 2 generates the coefficients of a particular solution  $\tilde{\sigma}_z^\Delta \in \mathcal{S}(z)$  in terms of the finite element basis  $\mathcal{B}_{RT}$  of  $RT_q^{\text{pw}}(\mathcal{T}(z))$  from Subsection A.2. The second step computes the correction

$$\sigma_z^\Delta := Q_{z,h}^\Delta - \tilde{\sigma}_z^\Delta = \arg \min_{\sigma_z \in V(z)} \|\sigma_z + \tilde{\sigma}_z^\Delta\|_{L^2(\omega(z))} \quad (72)$$

in terms of the low-dimensional unconstrained minimization problem over the linear space  $V(z) := \mathcal{S}(z) - \tilde{\sigma}_z^\Delta = RT_q^{\text{pw}}(\mathcal{T}(z)) \cap H(\text{div} = 0, \omega(z))$  characterized in Lemma 12. Because [12, Lemma 3.1] is wrong (take, e.g.,  $\tau_K = \text{curl } b_K \neq 0$  for the element bubble function  $b_K$  in their notation to see that uniqueness for general polynomial degrees  $q$  cannot hold) and [12] omits algorithmic details, this appendix focuses on the explicit characterization of the degrees of freedom for the minimization problem (72) over  $V(z)$ .



**Fig. 7** Triangulation  $\mathcal{T}(z)$  and enumeration of the edges  $\mathcal{F}(z)$  of the vertex-patch  $\omega(z)$  for an internal vertex  $z \in \mathcal{V}(\Omega)$  with  $N = |\mathcal{T}(z)| = 5$  (left) or boundary vertex  $z \in \mathcal{V}(\partial\Omega)$  with  $N = |\mathcal{T}(z)| = 4$  (right).

## A.2 Degrees of freedom for $RT_q^{\text{pw}}(T)$

This subsection introduces a basis for the Raviart-Thomas finite element on  $T \in \mathcal{T}(z)$  and starts with the definition of some linear functionals on  $H(\text{div}, \mathcal{T})$ . For any  $\sigma \in H(\text{div}, \mathcal{T})$ , set

$$\begin{aligned} \lambda_{T,\alpha}^{\ell,m}(\sigma) &:= \int_T \sigma \cdot e_\alpha x_1^\ell x_2^m dx & 0 \leq \ell + m \leq q-1, \alpha = 1, 2, \\ \lambda_{T,\text{div}}^{\ell,m}(\sigma) &:= \int_T \text{div} \sigma x_1^\ell x_2^m dx & 0 \leq \ell + m \leq q, \\ \lambda_{T,E}^j(\sigma) &:= \int_E \sigma|_T \cdot \nu_T s^j ds & 0 \leq j \leq q, E \in \mathcal{F}(T). \end{aligned}$$

Here and throughout,  $e_1 = (1, 0)$  and  $e_2 = (0, 1)$  denote the canonical unit vectors in  $\mathbb{R}^2$ . Note that the (classical) degrees of freedom for the Raviart-Thomas finite element  $RT_q(T)$  of degree  $q \in \mathbb{N}_0$  from [11] read

$$\widetilde{\Lambda}_T := \{\lambda_{T,E}^j, \lambda_{T,\alpha}^{\ell,m} : \text{for } 0 \leq j \leq q, 0 \leq \ell + m \leq q-1, \alpha = 1, 2, E \in \mathcal{F}(T)\}.$$

This appendix requires, for the construction of  $\tilde{\sigma}_z^\Delta \in \mathcal{S}(z)$ , a different set of (unisolvant) degrees of freedom  $\Lambda_T$  for  $RT_q(T)$  that includes the edge and divergence moments

$$\Lambda_T^0 := \{\lambda_{T,E}^j, \lambda_{T,\text{div}}^{\ell,m} : \text{for } 0 \leq j \leq q, 1 \leq \ell + m \leq q, E \in \mathcal{F}(T)\} \subseteq \Lambda_T. \quad (73)$$

(The set  $\Lambda_T^0$  itself is linear independent [35, Lemma 3.1].) Given any  $\Lambda_T$  with (73), denote the remaining  $N_q = q(q-1)/2$  degrees of freedom  $\Lambda_T \setminus \Lambda_T^0$  by  $\lambda_T^r$  for  $r = 1, \dots, N_q$ . Let  $\mathcal{B}_{RT,T} = \{\varphi_{T,E}^j, \varphi_{T,\text{div}}^{\ell,m}, \varphi_T^r\}$  be the unique basis of  $RT_q(T)$  dual to  $\Lambda_T$  with inferred indices from  $\Lambda_T$ . Then, the collection  $\mathcal{B}_{RT} := \bigcup_{T \in \mathcal{T}(z)} \mathcal{B}_{RT,T}$  is a basis of  $RT_q^{\text{pw}}(\mathcal{T}(z))$  and any function  $\sigma_z \in RT_q^{\text{pw}}(\mathcal{T}(z))$  has the representation

$$\sigma_z := \sum_{T \in \mathcal{T}(z)} \left( \sum_{E \in \mathcal{F}(T)} \sum_{j=0}^q c_{T,E}^j \varphi_{T,E}^j + \sum_{1 \leq \ell+m \leq q} c_{T,\text{div}}^{\ell,m} \varphi_{T,\text{div}}^{\ell,m} + \sum_{r=1}^{N_q} c_T^r \varphi_T^r \right) \quad (74)$$

with coefficients  $c_{T,E}^j = \lambda_{T,E}^j(\sigma_z)$ ,  $c_{T,\text{div}}^{\ell,m} = \lambda_{T,\text{div}}^{\ell,m}(\sigma_z)$ , and  $c_T^r = \lambda_T^r(\sigma_z)$  for all  $T \in \mathcal{T}(z)$ ,  $E \in \mathcal{F}(T)$ , and  $0 \leq j \leq q$ ,  $1 \leq \ell + m \leq q$ ,  $1 \leq r \leq N_q$ . By duality, the coefficients  $c_{T,E}^j$  with  $0 \leq j \leq q$  uniquely determine the normal trace  $(\sigma_z|_T)|_E \cdot \nu_E \in P_q(E)$  on the edge  $E \in \mathcal{F}(T)$  of  $T \in \mathcal{T}(z)$ . Any set of degrees of freedom  $\Lambda_T$  with (73) works with the equilibration algorithm in A.3.

*Example 2 (Construction of  $\Lambda_T$ )* This example presents a generic procedure to obtain such a set from  $\widehat{\Lambda}_T$ . The integration by parts formula shows that the lowest-order divergence moment  $\lambda_{T,\text{div}}^{0,0} = \sum_{E \in \mathcal{F}(T)} \lambda_{T,E}^0$  depends linearly on the lowest-order edge moments and, similarly, the sums

$$\lambda_{T,\text{div}}^{\ell,m} + \ell \lambda_{T,1}^{\ell-1,m} + m \lambda_{T,2}^{\ell,m-1} \in (P_{\ell+m}(\mathcal{F}(T)))^* \quad (75)$$

are functionals on  $P_{\ell+m}(\mathcal{F}(T))$  (summands with negative indices are understood as zero). This relation allows for the substitution of volume moments in  $\widehat{\Lambda}_T$  for divergence moments  $\lambda_{T,\text{div}}^{\ell,m}$ ,  $1 \leq \ell+m \leq q$ , and leads to  $\Lambda_T$  with (73). The remaining degrees of freedom  $\Lambda_T \setminus \Lambda_T^0$  are volume moments of the form  $\lambda_T^\alpha = \lambda_{T,\alpha}^{\ell,m}$  for a fixed  $\alpha \in \{1, 2\}$ , e.g.,

$$\Lambda_T \setminus \Lambda_T^0 = \{\lambda_{T,2}^{\ell,m} : 1 \leq \ell \leq q-1, 0 \leq m \leq q-1-\ell\}.$$

### A.3 Equilibration algorithm

---

#### Algorithm 2 Particular solution in $\mathcal{S}(z)$

---

**Input:** Data  $f \in L^2(\omega(z))$  and  $G_h \in H^1(\mathcal{T}(z))^2$  for vertex  $z \in \mathcal{V}$ .

- 1: Initialize all coefficients  $c_{T,E}^j, c_{T,\text{div}}^{\ell,m}, c_T^r$  in (74) with zero.
- 2: **for**  $a := 1 : N$  **do**
- 3:  $c_{T_a, E_{a-1}}^0 := \begin{cases} 0 & \text{if } a = 1, \\ ([G_h]_E \cdot \nu_E, \varphi_z)_{L^2(E_{a-1})} - c_{T_{a-1}, E_{a-1}}^0 & \text{else} \end{cases}$
- 4:  $c_{T_a, E_a}^0 := (f + \text{div } G_h, \varphi_z)_{L^2(T_a)} - c_{T_a, E_{a-1}}^0$
- 5: **for**  $1 \leq \ell + m \leq q$  **do**
- 6:  $c_{T_a, \text{div}}^{\ell,m} := (f + \text{div } G_h, \varphi_z x_1^\ell x_2^m)_{L^2(T_a)}$
- 7: **end for**
- 8: **for**  $1 \leq j \leq q$  **do**
- 9:  $c_{T_a, E_{a-1}}^j := 0$
- 10:  $c_{T_a, E_a}^j := ([G_h]_E \cdot \nu_E, \varphi_z s^j)_{L^2(E_a)}$
- 11: **end for**
- 12: **end for**

**Output:**  $\tilde{\sigma}_z^\Delta \in RT_q^{\text{pw}}(\mathcal{T}(z))$  defined by (74) with coefficients  $c_{T,E}^j, c_{T,\text{div}}^{\ell,m}, c_T^r$ .

---

This subsection presents the equilibration procedure, starting with Algorithm 2, that computes an admissible function  $\tilde{\sigma}_z^\Delta \in \mathcal{S}(z)$  in terms of the representation (74). Enumerate the  $N := |\mathcal{T}(z)|$  triangles  $T \in \mathcal{T}(z)$  from 1 to  $N$  as in Figure 7. Any two neighbouring triangles  $T_a, T_{a+1}$  share an edge  $E_a := T_a \cap T_{a+1}$  for  $a = 1, \dots, N-1$ . If  $z \in \mathcal{V}(\Omega)$  is an interior vertex,  $T_1$  and  $T_N$  share an additional edge  $E_0 := E_N := T_1 \cap T_N$ . For a boundary vertex  $z \in \mathcal{V}(\partial\Omega)$ ,  $T_1, T_N$  have the distinct boundary edges  $E_0, E_N \in \mathcal{F}(z) \cap \mathcal{F}(\partial\Omega)$ . The following lemma shows correctness of Algorithm 2 under the compatibility condition (5) and represents step one of the equilibration algorithm. The final step is the local minimization problem in Lemma 12 that provides  $Q_{z,h}^\Delta$  from (70). Both proofs are provided in A.4.

**Lemma 11** *Given  $z \in \mathcal{V}$ , let  $\{\varphi_{T,E}^j, \varphi_{T,\text{div}}^{\ell,m}, \varphi_T^r\}$  be the basis of  $RT_q(T)$  dual to  $\Lambda_T$  with (73) for all  $T \in \mathcal{T}(z)$ . Suppose  $f \in L^2(\omega(z))$  and  $G_h \in P_q(\mathcal{T}(z); \mathbb{R}^2)$  satisfy*

$$(G_h, \nabla \varphi_z)_{L^2(\omega(z))} = (f, \varphi_z)_{L^2(\omega(z))} \quad \text{if } z \in \mathcal{V}(\Omega). \quad (76)$$

*Then the output of Algorithm 2 with input  $f$  and  $G_h$  defines a function  $\tilde{\sigma}_z^\Delta \in \mathcal{S}(z)$ .*

Note that (76) is a local version of (5) and therefore holds for the HHO method with the choice  $G_h := \nabla_{\text{pw}} R u_h$  as proven in (20). This allows for the computation of  $Q_p^\Delta := \sum_{z \in \mathcal{V}} Q_{z,h}^\Delta = Q_p - G_h$  in terms of local and unconstrained minimization problems on the vertex-patches  $\omega(z)$ .

**Lemma 12** *Given  $z \in \mathcal{V}$ , let  $\{\varphi_{T,E}^j, \varphi_{T,\text{div}}^{\ell,m}, \varphi_T^r\}$  be as in Lemma 11 for all  $T \in \mathcal{T}(z)$  and let  $\tilde{\sigma}_z^\Delta \in \mathcal{S}(z)$  be arbitrary. Then  $V(z) := \mathcal{S}(z) - \tilde{\sigma}_z^\Delta$  is a linear vector space and consists of all functions of the form*

$$\sum_{a=1}^N \left( d_0(\varphi_{T_a, E_{a-1}}^0 - \varphi_{T_a, E_a}^0) + \sum_{\ell=1}^q (d_{E_{a-1}}^\ell \varphi_{T_a, E_{a-1}}^\ell - d_{E_a}^\ell \varphi_{T_a, E_a}^\ell) + \sum_{r=1}^{N_q} d_{T_a}^r \varphi_{T_a}^r \right) \quad (77)$$

for arbitrary  $d_0, d_{E_a}^\ell, d_{T_a}^r \in \mathbb{R}$  with  $\ell = 1, \dots, q, r = 1, \dots, N_q, a = 1, \dots, N$  (and  $d_{E_0}^\ell = d_{E_N}^\ell$  for  $z \in \mathcal{V}(\Omega)$ ) and the enumeration of  $\mathcal{T}(z)$  as in Figure 7. Furthermore,  $Q_{z,h}^\Delta = \tilde{\sigma}_z^\Delta + \sigma_z^\Delta$  holds for the solution  $\sigma_z^\Delta \in V(z)$  to the  $1+q|\mathcal{F}(z)|+q(q-1)/2N$ -dimensional minimization problem (72).

## A.4 Proofs

The remaining parts of this appendix are devoted to the verification of Lemmas 11–12.

*Proof (of Lemma 11)* Enumerate  $\mathcal{T}(z)$  as in A.3 and recall the definition of the jump  $[G_h]_E = G_h|_{T_+} - G_h|_{T_-}$  on the interior edge  $E = T_+ \cap T_-$  shared by  $T_+, T_- \in \mathcal{T}$ , and  $[G_h]_E = G_h|_{T_+}$  for the unique triangle  $T_+ \in \mathcal{T}$  with  $E \subset T_+$  for the boundary edge  $E \in \mathcal{F}(\partial\Omega)$ . First, observe that  $\sigma_z \in RT_q^{\text{pw}}(\mathcal{T}(z))$  lies in  $RT_q^{\text{pw},0}(\mathcal{T}(z))$  if and only if the coefficients  $c_{T,E}^j = 0$  in the representation (74) are zero for  $0 \leq j \leq q$  at the edge  $E \in \mathcal{F}(T) \setminus \mathcal{F}(z)$  in  $T \in \mathcal{T}$  opposing  $z$ . By definition,  $\sigma_z \in RT_q^{\text{pw},0}(\mathcal{T}(z))$  belongs to  $\mathcal{S}(z)$  if and only if

$$\lambda_{T,\text{div}}^{\ell,m}(\sigma_z) = (f + \text{div } G_h, \varphi_z x_1^\ell x_2^m)_{L^2(T)} \quad \text{for all } 0 \leq \ell + m \leq q, T \in \mathcal{T}(z), \quad (78)$$

$$(\lambda_{T_+,E}^j + \lambda_{T_-,E}^j)(\sigma_z) = ([G_h]_E \cdot \nu_E, \varphi_z s^j)_{L^2(E)} \quad \text{for all } 0 \leq j \leq q, E \in \mathcal{F}(\omega(z)). \quad (79)$$

This translates into equivalent conditions on the coefficients of  $\sigma_z$  in the representation (74), namely, for all  $a = 1, \dots, N$ ,

$$c_{T_a, E_a}^0 = (f + \text{div } G_h, \varphi_z)_{L^2(T_a)} - c_{T_a, E_{a-1}}^0, \quad (80)$$

$$c_{T_a, \text{div}}^{\ell,m} = (f + \text{div } G_h, \varphi_z x_1^\ell x_2^m)_{L^2(T_a)} \quad \text{for all } 1 \leq \ell + m \leq q, \quad (81)$$

$$c_{T_a, E_{a-1}}^j = d_{E_{a-1}}^j \quad \text{for all } 0 \leq j \leq q, \quad (82)$$

$$c_{T_a, E_a}^j = ([G_h]_E \cdot \nu_E, \varphi_z s^j)_{L^2(E_a)} - d_{E_a}^j \quad \text{for all } 0 \leq j \leq q, \quad (83)$$

where  $d_{E_a}^\ell \in \mathbb{R}$ . Since  $\sigma_z \in RT_q^{\text{pw},0}(\mathcal{T}(z))$  vanishes at the other edges  $E \in \mathcal{F} \setminus \mathcal{F}(z)$ ,  $\lambda_{T_a, \text{div}}^{0,0}(\sigma_z) = c_{T_a, E_{a-1}}^0 + c_{T_a, E_a}^0$  and (80)–(81) are equivalent to (78). The identification  $d_{E_0}^\ell = d_{E_N}^\ell$  for an interior vertex  $z \in \mathcal{V}(\Omega)$  with  $E_0 = E_N \in \mathcal{F}(z)$  shows that (82)–(83) are equivalent to (79). This identification is well defined. Note that, whereas there is no condition on  $d_{E_a}^\ell$  for  $1 \leq \ell \leq q$ , the combination of (80) and (82) with (83) shows the implicit extra condition

$$d_{E_a}^0 = d_{E_0}^0 + \sum_{\alpha=1}^a \left( ([G_h]_E \cdot \nu_E, \varphi_z)_{L^2(E_\alpha)} - (f + \text{div } G_h, \varphi_z)_{L^2(T_\alpha)} \right) \quad \text{for } a = 0, \dots, N.$$

For an interior vertex  $z \in \mathcal{V}(\Omega)$ , an integration by parts and (76) show that the sum on the right-hand side above vanishes for  $a = N$ , whence  $d_{E_N}^0 = d_{E_0}^0$  is indeed well defined. Furthermore, there is no condition on the coefficients  $c_{T_a}^r$  for all  $T_a \in \mathcal{T}(z)$  and  $r = 1, \dots, N_q$  and  $c_{T_a}^r = d_{T_a}^r$  is a further degree of freedom.

Algorithm 2 finds coefficients that satisfy (80)–(83) in a loop over  $a = 1, \dots, N$  and therefore defines  $\tilde{\sigma}_z^\Delta \in \mathcal{S}(z)$  by (74).  $\square$

*Proof (of Lemma 12)* This follows immediately after revisiting the proof of Lemma 11 for an arbitrary function  $\sigma_z \in RT_q^{\text{pw},0}(\mathcal{T}(z))$ . Since  $\sigma_z \in \mathcal{S}(z)$  is equivalent to (80)–(83) for the representation (74) of  $\sigma_z$  in the given basis, all functions  $\sigma_z \in \mathcal{S}(z) - \tilde{\sigma}_z^\Delta$  are of the form (77) for arbitrary  $d_0, d_{E_a}^\ell, d_{T_a}^r \in \mathbb{R}$  with  $\ell = 1, \dots, q$ ,  $r = 1, \dots, N_q$ , and  $a = 1, \dots, N$  (and  $d_{E_0}^\ell = d_{E_N}^\ell$  for  $z \in \mathcal{V}(\Omega)$ ). Hence, the dimension of the linear space  $V(z) = \mathcal{S}(z) - \tilde{\sigma}_z^\Delta$  is  $1 + q|\mathcal{F}(z)| + q(q-1)/2N$ . The claim follows from Proposition 1 by observing

$$Q_{z,h}^\Delta := \arg \min_{\sigma_z \in \mathcal{S}(z)} \|\sigma_z\|_{L^2(\omega(z))} = \tilde{\sigma}_z^\Delta + \arg \min_{\sigma_z \in \mathcal{S}(z) - \tilde{\sigma}_z^\Delta} \|\tilde{\sigma}_z^\Delta + \sigma_z\|_{L^2(\omega(z))}. \quad \square$$