

Generalized Henneberg stable minimal surfaces

David Moya*

Joaquín Pérez*

Abstract

We generalize the classical Henneberg minimal surface by giving an infinite family of complete, finitely branched, non-orientable, stable minimal surfaces in \mathbb{R}^3 . These surfaces can be grouped into subfamilies depending on a positive integer (called the *complexity*), which essentially measures the number of branch points. The classical Henneberg surface H_1 is characterized as the unique example in the subfamily of the simplest complexity $m = 1$, while for $m \geq 2$ multiparameter families are given. The isometry group of the most symmetric example H_m with a given complexity $m \in \mathbb{N}$ is either isomorphic to the dihedral isometry group D_{2m+2} (if m is odd) or to $D_{m+1} \times \mathbb{Z}_2$ (if m is even). Furthermore, for m even H_m is the unique solution to the Björling problem for a hypocycloid of $m + 1$ cusps (if m is even), while for m odd the conjugate minimal surface H_m^* to H_m is the unique solution to the Björling problem for a hypocycloid of $2m + 2$ cusps.

1 Introduction

A celebrated result obtained independently by do Carmo & Peng [1], Fischer-Colbrie & Schoen [2] and Pogorelov [7] establishes that if M is a complete orientable stable minimal surface in \mathbb{R}^3 , then M is a plane. Ros [8] proved that the same characterization holds without assuming orientability. Nevertheless, a plethora of complete stable minimal surfaces in \mathbb{R}^3 appear if we allow these stable minimal surfaces to have branch points, with the simplest example being the classical Henneberg minimal surface [3].

The class of complete, finitely connected and finitely branched minimal surfaces with finite total curvature (among which stable ones are a particular case) appears naturally in the following situation: Given $\varepsilon_0 > 0$, $I \in \mathbb{N} \cup \{0\}$ and $H_0, K_0 \geq 0$, let $\Lambda = \Lambda(I, H_0, \varepsilon_0, K_0)$ be the set of immersions $F: M \looparrowright X$ where X is a complete Riemannian 3-manifold with injectivity radius $\text{Inj}(X) \geq \varepsilon_0$ and absolute sectional curvature bounded from above by K_0 , M is a complete surface, F has constant mean curvature $H \in [0, H_0]$ and Morse index at most I . The second fundamental form $|A_{F_n}|$ of a sequence $\{F_n\}_n \subset \Lambda$ may fail to be uniformly bounded, which leads to lack of compactness of Λ . Nevertheless, the interesting ambient geometry of the immersions F_n can be proven to be well organized locally around at most I points $p_{1,n}, \dots, p_{k,n} \in M_n$ ($k \leq I$) where $|A_{F_n}|$ takes on arbitrarily large local

*This work is supported in part by the IMAG–Maria de Maeztu grant CEX2020-001105-M / AEI / 10.13039/501100011033, MICINN grant PID2020-117868GB-I00 and Junta de Andalucía grants P18-FR-4049 and A-FQM-139-UGR18.

maximum values. Around any of these points $p_{i,n}$, one can perform a blow-up analysis and find a limit of (a subsequence of) expansions $\lambda_n F_n$ of the F_n (that is, we view F_n as an immersion with constant mean curvature H_n/λ_n in the scaled ambient manifold $\lambda_n X_n$ for a sequence $\{\lambda_n\}_n \subset \mathbb{R}^+$ tending to ∞). This limit is a complete immersed minimal surface $f: \Sigma \looparrowright \mathbb{R}^3$ with finite total curvature, passing through the origin $\vec{0} \in \mathbb{R}^3$. Recall that such an f has finitely many ends, each of which is a multi-valued graph of finite multiplicity (spinning) $s \in \mathbb{N}$, over the exterior of a disk in the tangent plane at infinite for f at that end. Thus, arbitrarily small almost perfectly formed copies of large compact portions of $f(\Sigma)$ can be reproduced in $F_n(M_n)$ around $F_n(p_{i,n})$ for n sufficiently large. Complete, finitely connected and finitely branched minimal surfaces with finite total curvature in \mathbb{R}^3 appear naturally when considering clustering phenomena in this framework: It may occur that different blow-up limits of the F_n around $p_{i,n}$ at different scales $\lambda_{1,n} > \lambda_{2,n}$ with $\lambda_{1,n}/\lambda_{2,n} \rightarrow \infty$ as $n \rightarrow \infty$, produce different limits $f_j: \Sigma_j \looparrowright \mathbb{R}^3$, $j = 1, 2$, with $\text{Index}(f_1) + \text{Index}(f_2) \leq I$; in this case, all the geometry of $f_1(\Sigma_1)$ collapses around $\vec{0} \in f_2(\Sigma_2)$, and every end of $f_1(\Sigma_1)$ with multiplicity $m \geq 3$ produces a branch point at the origin for $f_2(\Sigma_2)$ of branching order $s - 1$. For details about this clustering phenomenon and how to organize these blow-up limits in hierarchies appearing around $\{p_{i,n}\}_n$, see the paper [4] by Meeks and the second author.

The main goal of this paper is to generalize the classical Henneberg minimal surface H_1 to an infinite family of connected, 1-sided, complete, finitely branched, stable minimal surfaces in \mathbb{R}^3 . Branch points are unavoidable if we seek for complete, non-flat stable minimal surfaces by the aforementioned results [1, 2, 7, 8]; 1-sidedness is also necessary condition for stability (see Proposition 3 below). Our examples can be grouped into subfamilies depending on the number of branch points (this will be encoded by an integer $m \in \mathbb{N}$ called the *complexity*). The most symmetric examples H_m in each subfamily of complexity m will be studied in depth (Section 5.3). Depending on the parity of m , either H_m or its conjugate minimal surface H_m^* (which does not give rise to a 1-sided surface, see Section 5.4) can be viewed as the unique solution of a Björling problem for a planar hypocycloid (Section 5.7). The isometry group of H_m is isomorphic to the dihedral group D_{2m+2} if m is odd and to the group $D_{m+1} \times \mathbb{Z}_2$ if m is even (Section 5.8). We will also prove that H_1 is the only element in the subfamily with complexity $m = 1$ (Theorem 11), while for $m \geq 2$, H_m can be deformed in multiparameter families: Proposition 14 gives an explicit 1-parameter family of examples with complexity $m = 2$, interpolating between H_2 and a limit which turns out to be H_1 (Section 6.2.1), and the subfamily of examples with complexity $m = 2$ is a two-dimensional real analytic manifold around H_2 (Section 6.2.2).

2 1-sided branched stable minimal surfaces

We start with the Weierstrass data (g, ω) on a Riemann surface Σ , so that (g, ω) solves the period problem and produces a conformal harmonic map $X: \Sigma \looparrowright \mathbb{R}^3$ given by the classical formula

$$X = \text{Re} \int (\phi_1, \phi_2, \phi_3) = \text{Re} \int \left(\frac{1}{2}(1 - g^2)\omega, \frac{i}{2}(1 + g^2)\omega, g\omega \right). \quad (1)$$

We will assume that X is an immersion outside of a locally finite set of points $\mathcal{B} \subset \Sigma$, where X fails to be an immersion (points of \mathcal{B} are called *branch points* of X). Such an X will be

called a *branched minimal immersion*. The induced (possible branched) metric is given by

$$ds^2 = \frac{1}{4}(1 + |g|^2)^2 |\omega|^2. \quad (2)$$

The local structure of X around a branch point in \mathcal{B} is well-known, see e.g. Micallef and White [5, Theorem 1.4] for details. Given $p \in \mathcal{B}$, there exists a conformal coordinate (D, z) for Σ centered at p (here D is the closed unit disk in the plane), a diffeomorphism u of D and a rotation ϕ of \mathbb{R}^3 such that $\phi \circ X \circ u$ has the form

$$z \mapsto (z^q, x(z)) \in \mathbb{C} \times \mathbb{R} \sim \mathbb{R}^3$$

for z near 0, where $q \in \mathbb{N}$, $q \geq 2$, x is of class C^2 , and $x(z) = o(|z|^q)$. In this setting, the *branching order of p* is defined to be $q - 1 \in \mathbb{N}$.

Let us assume that X produces a 1-sided branched minimal surface; this means that there exists an anti-holomorphic involution without fixed points $I : \Sigma \rightarrow \Sigma$ such that $I \circ \phi_j = \overline{\phi_j}$ for $j = 1, 2, 3$. This is equivalent to

$$-1/\overline{g} = g \circ I, \quad I^* \omega = -\overline{g^2 \omega}. \quad (3)$$

In particular, I must preserve the set \mathcal{B} . $\Sigma/\langle I \rangle$ is a non-orientable differentiable surface endowed with a conformal class of metrics, and the harmonic map X induces another harmonic map $\widehat{X} : \Sigma/\langle I \rangle \looparrowright \mathbb{R}^3$ such that $\widehat{X} \circ \pi = X$, where $\pi : \Sigma \rightarrow \Sigma/\langle I \rangle$ is the natural projection (\widehat{X} is a branched minimal immersion). Reciprocally, every 1-sided conformal harmonic map can be constructed in this way.

Remark 1. In the particular case that the compactification of Σ is $\overline{\mathbb{C}}$, we can assume that $I(z) = -1/\overline{z}$ and write $\omega = f dz$ globally. In this setting, the above equations give

$$-1/\overline{g(z)} = g(-1/\overline{z}), \quad f \circ I = -\overline{z^2 g^2 f}. \quad (4)$$

Definition 2. Given a 1-sided conformal harmonic map $\widehat{X} : \Sigma/\langle I \rangle \looparrowright \mathbb{R}^3$, we denote by Δ , $|A|^2$ the Laplacian and squared norm of the second fundamental form of \widehat{X} . The *index* of \widehat{X} is defined as the number of negative eigenvalues of the elliptic, self-adjoint operator $L = \Delta + |A|^2$ (Jacobi operator of X) defined over the space of compactly supported smooth functions $\phi : \Sigma \rightarrow \mathbb{R}$ such that $\phi \circ I = -\phi$. \widehat{X} is said to be *stable* if its index is zero.

In the case \widehat{X} is finitely branched, the eigenvalues and eigenfunctions of the Jacobi operator of X are well defined via a variational approach, since the codimension of the singularity set \mathcal{B} is two (see [9]), and stability also makes sense.

The next result is proven by Meeks and the second author in [4].

Proposition 3. *Let $X : \Sigma \looparrowright \mathbb{R}^3$ be complete, non-flat, finitely branched minimal immersion with branch locus $\mathcal{B} \subset \Sigma$. Then:*

1. [4, Proposition 3] *If X is stable, then Σ is non-orientable and $X(\mathcal{B})$ contains more than 1 point.*
2. [4, Remark 3.6] *Suppose that Σ is non-orientable, X has finite total curvature and its extended unoriented Gauss map $G : \mathbb{P}^2 = \mathbb{S}^2/\{\pm 1\} \rightarrow \mathbb{P}^2$ is a diffeomorphism. Then, X is stable.*

3 The Björling problem

We next recall the basics of the classical Björling problem, to be used later. Let $\gamma: I \subset \mathbb{R} \rightarrow \mathbb{R}^3$ be an analytic regular curve and η an analytic vector field along γ such that $\langle \gamma(t), \eta(t) \rangle = 0$ and $\|\eta(t)\| = 1$ for all $t \in I$. The classical result due to E.G. Björling asserts that the following parametrization generates a minimal surface S which contains γ and has η as unit normal vector along γ :

$$X(u, v) = \operatorname{Re} \left(\tilde{\gamma}(w) - i \int_{w_0}^w \tilde{\eta}(w) \times \tilde{\gamma}'(w) dw \right),$$

where $\tilde{\gamma}, \tilde{\eta}$ are analytic extensions of the corresponding γ, η and $w = u + iv$ is defined in a simply connected domain $\Omega \subset \mathbb{C}$ with $I \subset \Omega$. In particular, the surface S is locally unique around γ with this data (it is called the solution to the Björling problem with data γ, η).

In what follows, we will consider different Björling problems for analytic planar curves $\gamma \subset \{z = 0\}$ that fail to be regular at finitely many points. The above construction can be applied to each of the regular arcs of these curves after removing the zeros of γ' . In all our applications, η will be taken as the (unit) normal vector field to γ as a planar curve.

4 The classical Henneberg surface

The classical Henneberg minimal surface H_1 is the 1-sided, complete, stable minimal surface in \mathbb{R}^3 given by the Weierstrass data:

$$g(z) = z, \quad \omega = z^{-4}(z \pm i)(z \pm 1)dz = z^{-4}(z^4 - 1)dz, \quad z \in \overline{\mathbb{C}} - \{0, \infty\}. \quad (5)$$

H_1 has two branch points¹ at $[1] = \{1, -1\}$, $[i] = \{i, -i\} \in \mathbb{P}^2 = \overline{\mathbb{C}}/\langle A \rangle$, where $A(z) = -1/\bar{z}$ is the antipodal map. By Proposition 3, H_1 is stable.

H_1 can be conformally parameterized (up to translations) by equation (1). After translating X so that $X(e^{i\pi/4}) = \vec{0}$, the branch points of H_1 are mapped by X to $(0, 0, \pm 1)$ and a parametrization of H_1 in polar coordinates $z = re^{i\theta}$ is given by

$$X(re^{i\theta}) = \begin{pmatrix} \frac{\cos \theta}{2} \left(r - \frac{1}{r} \right) - \frac{\cos(3\theta)}{6} \left(r^3 - \frac{1}{r^3} \right) \\ -\frac{\sin \theta}{2} \left(r - \frac{1}{r} \right) - \frac{\sin(3\theta)}{6} \left(r^3 - \frac{1}{r^3} \right) \\ \frac{\cos(2\theta)}{2} \left(r^2 + \frac{1}{r^2} \right) \end{pmatrix}. \quad (6)$$

Since $X(e^{i\theta}) = (0, 0, \cos(2\theta))$, then X maps the unit circle into the vertical segment $\{(0, 0, t) \mid t \in [-1, 1]\}$. In this way, $\theta \in [0, 2\pi] \mapsto X(e^{i\theta})$ bounces between the two branch points of H_1 (observe that the complement of this closed segment in the x_3 -axis is not contained in H_1), see Figure 1.

¹Branch points of H_1 all have order 1 (locally the surface winds twice around the branch point); this follows from direct computation, or from Proposition 21 in White's "Lectures on minimal surfaces theory".




Figure 1: The Henneberg surface H_1 . After a traslation, the branch points of H_1 are contained in the x_3 -axis. H_1 contains two horizontal, orthogonal lines that bisect the x_1 - and x_2 -axis. Left: Intersection of H_1 with a ball of radius 8. Right: top view of H_1 .

4.1 Isometries of H_1

It is straightforward to check that

1. The antipodal map $A: \overline{\mathbb{C}} \rightarrow \overline{\mathbb{C}}$ (in polar coordinates $(r, \theta) \mapsto (1/r, \pi + \theta)$) leaves the surface invariant. This is the deck transformation, which is orientation reversing.
2. The map $z \mapsto -z$ (in polar coordinates $(r, \theta) \mapsto (r, \pi + \theta)$) induces the rotation by angle π about the axis x_3 on the surface.
3. The inversion of the z -plane with respect to the unit circle, $z \mapsto 1/z$, (in polar coordinates $(r, \theta) \mapsto (1/r, \theta)$) is the composition of A with $z \mapsto -z$, and thus, it also induces a rotation of angle π about the x_3 -axis on the surface.
4. The conjugation map $z \mapsto \bar{z}$ (in polar coordinates $(r, \theta) \mapsto (r, -\theta)$) induces the reflection of X about the plane (x_1, x_3) .
5. The reflection about the imaginary axis (in polar coordinates $(r, \theta) \mapsto (r, \pi - \theta)$) induces the reflection of X about the plane (x_2, x_3) .
6. X maps the half-line $\{re^{-i\pi/4} \mid r \in (0, \infty)\}$ (respectively $\{re^{i\pi/4} \mid r \in (0, \infty)\}$) injectively into $l_1 = \text{Span}(1, 1, 0)$ (respectively $l_2 = \text{Span}(1, -1, 0)$). Thus, the rotations R_1, R_2 of angle π about l_1, l_2 are isometries of X (R_1 is induced by $z \mapsto -i\bar{z}$ and R_2 by $z \mapsto i\bar{z}$).
7. The map $z \mapsto iz$ (in polar coordinates $(r, \theta) \mapsto (r, \theta + \pi/2)$) induces the rotation of angle $\pi/2$ about the x_3 -axis composed by a reflection in the (x_1, x_2) -plane.

Together with the identity map, the above isometries form a subgroup of the isometry group $\text{Iso}(H_1)$ of H_1 , isomorphic to the dihedral group D_4 .

Lemma 4. *These are all the (intrinsic) isometries of H_1 .*

Proof. This is a direct consequence of the fact that every intrinsic isometry ϕ of H_1 produces a conformal diffeomorphism of $\mathbb{C} \setminus \{0\}$ into itself that preserves the set of branch points of H_1 . In particular, ϕ is of one of the aforementioned eight cases. \square

4.2 Associated family and the conjugate surface H_1^* .

The flux vector of H_1 around the origin in \mathbb{C} vanishes (in other words, the Weierstrass form $\Phi = (\phi_1, \phi_2, \phi_3)$ associated to H_1 is exact). This implies that all associated surfaces $\{\tilde{H}_1(\varphi) \mid \varphi \in [0, 2\pi)\}$ to the orientable cover $\tilde{H}_1 = \tilde{H}_1(0)$ of H_1 are well-defined as surfaces in \mathbb{R}^3 (the branched minimal immersion $\tilde{H}_1(\varphi)$ has Weierstrass data $g_\varphi = g$, $\omega_\varphi = e^{i\varphi}\omega$ and it is isometric to \tilde{H}_1 , in particular it has the same branch locus as \tilde{H}_1).

None of the surfaces $\tilde{H}_1(\varphi)$ except for $\varphi = 0$ descends to the non-orientable quotient $\mathbb{P}^2 \setminus \{[0]\}$, because the second equation in (3) is not preserved if we exchange ω by $e^{i\varphi}\omega$, $\varphi \in (0, 2\pi)$. In particular, none of these associated surfaces are congruent to H_1 .

The conjugate surface $H_1^* := \tilde{H}_1(\pi/2)$ is symmetric by reflection in the (x_1, x_2) -plane. The intersection between H_1^* and $\{z = 0\}$ consists of the *astroid* γ_4 parameterized by

$$t \mapsto \gamma_4(t) = \begin{pmatrix} -\sin(\theta) + \frac{\sin(3\theta)}{3} \\ -\cos(\theta) - \frac{\cos(3\theta)}{3} \\ 0 \end{pmatrix},$$

together with four rays starting at the cusps of the astroid in the direction of their position vectors, see Figure 2.




Figure 2: The astroid γ_4 (red) and the four rays obtained by intersecting H_1^* with the (x_1, x_2) -plane (blue).

In particular, H_1^* is the solution of the Björling problem for the curve γ_4 and the choice of unit normal field the normal vector to γ_4 as a planar curve, see also Remark 8 below.

5 Generalized Henneberg surfaces

We will next search for a 1-sided, complete, stable minimal surface in $X: \Sigma \looparrowright \mathbb{R}^3$ with $\Sigma = \overline{\mathbb{C}} \setminus \mathcal{E}$, \mathcal{E} finite and $g(z) = z$. Hence, $I(z) = -1/\bar{z}$, $\widehat{X} = X/\langle I \rangle: \Sigma/\langle I \rangle \looparrowright \mathbb{R}^3$ is stable and (4) writes

$$f(-1/\bar{z}) = -\overline{z^4 f(z)}. \quad (7)$$

5.1 General form for f

We take a general rational function

$$f(z) = \frac{c}{z^{m+3}} \frac{\prod_{j=1}^M (z - a_j)}{\prod_{j=1}^N (z - b_j)}, \quad (8)$$

where $c, a_j, b_j \in \mathbb{C}^*$, $m \in \mathbb{N}$, $M, N \in \mathbb{N} \cup \{0\}$ are to be determined.

- Remark 5.**
1. Henneberg's surface H_1 has $f(z) = z^{-4}(z^4 - 1)$, hence $c = 1$, $m = 1$, $N = 0$, $M = 4$, $\{a_j\} = \{\pm 1, \pm i\}$.
 2. The zeros of the induced metric (2) (branch points of the surface) occur precisely at the points a_j ; the ends occur at $0, \infty$ and at the points b_j (in particular, both families $\{a_j\}_j, \{b_j\}_j$ must then come in pairs of antipodal points, see also (12) below).
 3. A consequence of the last observation is that when the above rotations in \mathbb{R}^3 of our surfaces (provided that the Weierstrass data close periods) are not allowed unless the axis of rotation is vertical.

Imposing (7) to (8) we get

$$c(-1)^{m-1+M-N} \bar{z}^{3+m-M+N} \frac{\prod_{j=1}^M (1 + a_j \bar{z})}{\prod_{j=1}^N (1 + b_j \bar{z})} = f(-1/\bar{z}) = -\overline{z^4 f(z)} = -\frac{\bar{c}}{\bar{z}^{m-1}} \frac{\prod_{j=1}^M (\bar{z} - \bar{a}_j)}{\prod_{j=1}^N (\bar{z} - \bar{b}_j)},$$

thus

$$\bar{c}(-1)^{m+M-N} z^{2+2m-M+N} \prod_{j=1}^M (1 + \bar{a}_j z) \prod_{j=1}^N (z - b_j) = c \prod_{j=1}^M (z - a_j) \prod_{j=1}^N (1 - \bar{b}_j z), \quad (9)$$

from where we deduce that

$$2 + 2m - M + N = 0, \quad (10)$$

in particular $M - N$ is even. Substituting $z = 0$ in (9) we get

$$\bar{c}(-1)^m \prod_{j=1}^N b_j = c \prod_{j=1}^M a_j. \quad (11)$$

Using (11), we can rewrite (9) as an equality between monic polynomials in z :

$$\prod_{j=1}^M \left(\frac{1}{\bar{a}_j} + z \right) \prod_{j=1}^N (z - b_j) = \prod_{j=1}^M (z - a_j) \prod_{j=1}^N \left(\frac{1}{\bar{b}_j} + z \right),$$

from where we deduce that

$$\{a_1, \dots, a_M\} = \{-1/\bar{a}_1, \dots, -1/\bar{a}_M\}, \quad \{b_1, \dots, b_N\} = \{-1/\bar{b}_1, \dots, -1/\bar{b}_N\}. \quad (12)$$

that is, M, N are even, the a_j (resp. b_j) are given by $M/2$ (resp. $N/2$) pairs of antipodal points in \mathbb{C}^* . Now (10) and (11) give respectively:

$$1 + m - \widetilde{M} + \widetilde{N} = 0, \quad (13)$$

$$-\bar{c} \prod_{j=1}^{N/2} \frac{b_j}{\bar{b}_j} = c \prod_{j=1}^{M/2} \frac{a_j}{\bar{a}_j}. \quad (14)$$

5.2 Solving the period problem in the one-ended case: complexity

From (3) and (8) we see that the points where ds^2 can blow up are $z = 0, b_1, \dots, b_N$ and its antipodal points. In order to keep the computations simple, we will assume there are no b_j 's, i.e. $N = 0$ (or equivalently $M/2 = m + 1$), which reduces the period problem to imposing

$$\overline{\int_{\gamma} g^2 \omega} = \int_{\gamma} \omega, \quad \operatorname{Re} \int_{\gamma} f \omega = 0,$$

where $\gamma = \{|z| = 1\}$, or equivalently,

$$\overline{\operatorname{Res}_0(g^2 f)} = -\operatorname{Res}_0(f), \quad \operatorname{Im} \operatorname{Res}_0(gf) = 0. \quad (15)$$

We can simplify (8) to

$$f(z) = \frac{c}{z^{m+3}} \prod_{j=1}^{m+1} (z - a_j) \left(z + \frac{1}{a_j} \right), \quad (16)$$

which satisfies (7) (this is the condition to descend to the quotient as a 1-sided surface, provided that the period problem (15) is solved) if and only if (14) holds, which in this case reduces to

$$-\frac{\bar{c}}{c} = \prod_{j=1}^{m+1} \frac{a_j}{\bar{a}_j}. \quad (17)$$

We call

$$P(z) := \prod_{j=1}^{m+1} (z - a_j) \left(z + \frac{1}{a_j} \right) = \sum_{h=0}^{2m+2} A_h z^h. \quad (18)$$

Thus,

$$\operatorname{Res}_0(f) = c \operatorname{Res}_0 \left(\sum_{h=0}^{2m+2} A_h z^{h-m-3} \right) = c A_{m+2},$$

$$\operatorname{Res}_0(g^2 f) = c \operatorname{Res}_0 \left(\sum_{h=0}^{2m+2} A_h z^{h-m-1} \right) = c A_m,$$

$$\operatorname{Res}_0(gf) = c \operatorname{Res}_0\left(\sum_{h=0}^{2m+2} A_h z^{h-m-2}\right) = cA_{m+1}.$$

Thus, (15) reduces to

$$\overline{cA_m} = -cA_{m+2}, \quad \operatorname{Im}(cA_{m+1}) = 0. \quad (19)$$

Remark 6. We can assume $|c| = 1$ due to the fact that multiplying the Weierstrass form by a positive number does not affect to solving the period problem and just multiplies the resulting surface by a homothety. Similarly, exchanging c by $-c$ does not affect to solving the period problem.

We also write $a_j = |a_j|e^{i\theta_j}$, $\theta_j \in \mathbb{R}$. Thus,

$$-a_j + \frac{1}{\overline{a_j}} = (-|a_j| + \frac{1}{|a_j|})e^{i\theta_j}, \quad \frac{a_j}{\overline{a_j}} = e^{2i\theta_j},$$

and so,

$$P(z) = \prod_{j=1}^{m+1} \left(z^2 + (-|a_j| + \frac{1}{|a_j|})e^{i\theta_j} z - e^{2i\theta_j} \right) \quad (20)$$

Definition 7. Given $m \in \mathbb{N}$, a list $(c, a_1, \dots, a_{m+1}) \in \mathbb{S}^1 \times (\mathbb{C}^*)^{m+1}$ solving the equations (17),(19) will be called a *solution of the period problem with complexity m* . Note that geometrically, a_1, \dots, a_{m+1} are the Gaussian images of the branch points of the resulting surface.

5.3 The case when the a_j are the $(2m + 2)$ -roots of unity

For each complexity m , there is a most symmetric configuration that gives rise to a solution of the period problem for that complexity, which we describe next.

Take the a_j as the solutions of the equation $a^{2m+2} = 1$ (i.e. $|a_j| = 1$ and $\theta_j = \frac{\pi}{m+1}(j-1)$, $j = 1, \dots, m+1$). Observe that

$$\prod_{j=1}^{m+1} \frac{a_j}{\overline{a_j}} = \prod_{j=1}^{m+1} e^{2i\theta_j} = e^{2i \sum_{j=1}^{m+1} \theta_j} = e^{\frac{2\pi i}{m+1} \sum_{j=1}^{m+1} (j-1)} = e^{\frac{2\pi i}{m+1} \frac{m(m+1)}{2}} = e^{i\pi m},$$

hence the validity of (17) is equivalent in this case to

$$c = \pm i^{m-1}. \quad (21)$$

As for equation (19), note that (20) can be written as

$$P(z) = \prod_{j=1}^{m+1} (z^2 - e^{2i\theta_j}) = z^{2m+2} - 1,$$

and thus $A_m = 0$ (because $m > 0$), $A_{m+2} = 0$ (because $m+2 < 2m+2$) and $A_{m+1} = 0$. In particular, (19) is trivially satisfied for each value of $c \in \mathbb{C}^*$. Therefore, the Weierstrass data

$$g(z) = z, \quad \omega = i^{m-1} z^{-m-3} (z^{2m+2} - 1) dz, \quad z \in \mathbb{C}^*, \quad (22)$$

give rise to a 1-sided, complete, stable minimal surface H_m . For $m = 1$ we recover the classical Henneberg's surface.

5.4 Associated family and the conjugate surface H_m^* .

Since $A_m = A_{m+1} = A_{m+2} = 0$, the flux vector of H_m around the origin in \mathbb{C} vanishes and the Weierstrass form $\Phi = (\phi_1, \phi_2, \phi_3)$ associated to H_m is exact. Thus all associated surfaces $\{\tilde{H}_m(\varphi) \mid \varphi \in [0, 2\pi)\}$ to the orientable cover $\tilde{H}_m = \tilde{H}_m(0)$ of are well-defined. As in the case $m = 1$ (see Section 4.2), none of these associated surfaces descends to the 1-sided quotient, except for $\pm H_m$. Let $H_m^* := \tilde{H}_m(\pi/2)$ be the conjugate surface to H_m .

The behavior of H_m is very different depending on the parity of m . A naive justification of this dependence on the parity of m comes from the fact that the coefficient for ω changes from ± 1 for m odd to $\pm i$ for m even. A more geometric interpretation of this dependence will be given next.

5.5 The case m odd

If $m \in \mathbb{N}$ is odd, (22) gives $\omega = z^{-m-3}(z^{2m+2} - 1)dz$. Although H_m has $m + 1$ branch points in $\Sigma = \mathbb{P}^2 \setminus \{[0]\}$ (the classes of the $(2m + 2)$ -roots of unity under the antipodal map), they are mapped into just two different points in \mathbb{R}^3 : after translating the surface in \mathbb{R}^3 so that $X(e^{i\frac{\pi}{2(m+1)}}) = \vec{0}$ (we are using the notation in (1)), the branch points of H_m are mapped to $(0, 0, \pm 1)$ and a parameterization of H_m in polar coordinates is (compare with (6))

$$X(re^{i\theta}) = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} \frac{\cos(m\theta)}{2m} \left(r^m - \frac{1}{r^m}\right) - \frac{\cos((m+2)\theta)}{2(m+2)} \left(r^{m+2} - \frac{1}{r^{m+2}}\right) \\ -\frac{\sin(m\theta)}{2m} \left(r^m - \frac{1}{r^m}\right) - \frac{\sin((m+2)\theta)}{2(m+2)} \left(r^{m+2} - \frac{1}{r^{m+2}}\right) \\ \frac{\cos((m+1)\theta)}{m+1} \left(r^{m+1} + \frac{1}{r^{m+1}}\right) \end{pmatrix}. \quad (23)$$

X maps the unit circle into the vertical segment $\{(0, 0, t) \mid t \in [-1, 1]\}$. $\theta \in [0, 2\pi] \mapsto X(e^{i\theta})$ bounces between the two branch points of H_m , and the complement of this closed segment in the x_3 -axis is not contained in H_m . $H_m \cap \{x_3 = 0\}$ consists of an equiangular system of $m + 1$ straight lines passing through the origin (the images by X of the straight lines of arguments $\theta = \frac{\pi/2+k\pi}{m+1}$, $k = 0, \dots, m$ in polar coordinates), see Figure 4 right for H_3 .

5.6 The case m even

If m is even (and non-zero), (22) produces $\omega = iz^{-m-3}(z^{2m+2} - 1)dz$. In this case, a parametrization of H_m in polar coordinates is

$$X(re^{i\theta}) = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} -\frac{\sin(m\theta)}{2m} \left(r^m + \frac{1}{r^m}\right) + \frac{\sin((m+2)\theta)}{2(m+2)} \left(r^{m+2} + \frac{1}{r^{m+2}}\right) \\ -\frac{\cos(m\theta)}{2m} \left(r^m + \frac{1}{r^m}\right) - \frac{\cos((m+2)\theta)}{2(m+2)} \left(r^{m+2} + \frac{1}{r^{m+2}}\right) \\ \frac{\sin((m+1)\theta)}{m+1} \left(\frac{1}{r^{m+1}} - r^{m+1}\right) \end{pmatrix}. \quad (24)$$

X maps the unit circle $\{r = 1\}$ into a certain hypocycloid contained in the plane $\{x_3 = 0\}$, as we will explain next.

A *hypocycloid* of inner radius $r > 0$ and outer radius $R > r$ is the planar curve traced by a point on a circumference of radius r which is rolling along the interior of another circumference (which is fixed) of radius R . It can be parametrized by $\alpha(t) = (x(t), y(t))$,

$t \in \mathbb{R}$, where

$$x(t) = -(R - r) \sin t + r \sin \left(\frac{R - r}{r} t \right), \quad y(t) = -(R - r) \cos t - r \cos \left(\frac{R - r}{r} t \right).$$

Using (24), we deduce that the image by X of the unit circle $\mathbb{S}^1 \subset \mathbb{C}$ has the following parametrization:

$$\theta \in [0, 2\pi) \mapsto X(e^{i\theta}) = \begin{pmatrix} -\frac{\sin(m\theta)}{m} + \frac{\sin((m+2)\theta)}{m+2} \\ -\frac{\cos(m\theta)}{m} - \frac{\cos((m+2)\theta)}{m+2} \\ 0 \end{pmatrix}. \quad (25)$$

From (25) we deduce that, up to the reparametrization $t = m\theta$, $X(\mathbb{S}^1)$ is the hypocycloid of inner radius $r = \frac{1}{m+2}$ and outer radius $R = \frac{2m+2}{m(m+2)}$, which has exactly $m + 1$ cusps. These cusp points are the images by X of the $m + 1$ branch points of H_m . In particular, H_m is the unique minimal surface obtained as solution of the Björling problem for the hypocycloid of $m + 1$ cusps (this number of cusps is any odd positive integer, at least three), inner radius $r = \frac{1}{m+2}$ and outer radius $R = \frac{2m+2}{m(m+2)}$, when we take as normal vector field η (see Section 3 for the notation) the normal vector to the hypocycloid as a planar curve.

We depict this planar curve in the simplest cases $m = 2, 4, 6$ in Figure 3 in red.




Figure 3: The intersection of H_m (with $m > 0$ even) with $\{x_3 = 0\}$ consists of a hypocycloid with $m + 1$ cusps (in red) together with half-lines $\{tp \mid t \geq 1\}$ that start from each of these cusp points p . Left: $H_2 \cap \{x_3 = 0\}$, where the branch points have coordinates $(0, -\frac{3}{4}, 0)$, $(-\frac{3\sqrt{3}}{8}, \frac{3}{8}, 0)$, $(\frac{3\sqrt{3}}{8}, \frac{3}{8}, 0)$. Center: $H_4 \cap \{x_3 = 0\}$, Right: $H_6 \cap \{x_3 = 0\}$.

5.7 Revisiting the case m odd: H_m^* as a solution of a Björling problem for a hypocycloid

Using the Weierstrass formula (1), it can be easily seen that the conjugate surface H_m^* of H_m with odd m can be parameterized in polar coordinates $z = re^{i\theta}$ by $X^*(re^{i\theta})$ given by the same formula as the right-hand-side of (24). $X^*(\mathbb{S}^1)$ parameterizes a hypocycloid γ_{2m+2} with inner radius $r = \frac{1}{m+2}$ and outer radius $R = \frac{2m+2}{m(m+2)}$. Since

$$\frac{R}{r} = \frac{2m+2}{m},$$




Figure 4: Left: H_2 . Right: H_3 .

we deduce that γ_{2m+2} has $2m + 2$ cusps². Observe that $2m + 2$ is a positive multiple of 4 because m is odd; and conversely, every positive multiple of 4 can be written as $2m + 2$ for a unique $m \in \mathbb{N}$ odd. This tells us that for any $m \in \mathbb{N}$ odd, H_m^* is the unique solution to the Björling problem for the hypocycloid γ_{2m+2} , when we take as normal vector field η the normal vector to γ_{2m+2} as a planar curve.

Remark 8. 1. In the particular case of a hypocycloid of 4 cusps (called *astroid*), we recover the conjugate surface H_1^* of the classical Henneberg surface. This result was described by Odehnal [6], who also studied the Björling problem for an hypocycloid γ_3 of three cusps from the viewpoint of algebraic surfaces.

2. We have described the minimal surfaces obtained as the solution of a Björling problem over a hypocycloid if the number of its cusps is either any given odd number or a multiple of four. The case that remains is when the hypocycloid has $4k + 2$ cusps, $k \in \mathbb{N}$. The corresponding solution to this Björling problem can be also explicitly described by the parametrization (24), now with a parameter $m \in \mathbb{Q}$. Namely, if we choose m to be of the form $m = \frac{1}{2k}$, $k \in \mathbb{N}$, inner radius $r = \frac{1}{m+2}$ and outer radius $R = \frac{2m+2}{m(m+2)}$, then

$$\frac{R}{r} = \frac{2m+2}{m} = 4k+2,$$

which ensures that the complete branched minimal surface $H_{\frac{1}{2k}} = X(\mathbb{C} \setminus \{0, \infty\})$ (here X is given by (24)) is symmetric by reflection in the (x_1, x_2) -plane, and $X(\mathbb{S}^1)$ is a

²For a hypocycloid of inner radius $r > 0$ and outer radius $R > r$, the quotient R/r expresses the number of times that the inner circumference rolls along the outer circumference until it completes a loop. If R/r is a rational number and a/b is the irreducible fraction of R/r , then $b \cdot a/b = a$ counts the number of times that the inner circumference rolls until the point that generates the hypocycloid reaches its initial position. This number a coincides with the number of cusps.

hypocycloid with $4k + 2$ cusps. $H_{\frac{1}{2k}}$ does not descend to a 1-sided quotient.

5.8 Isometries of H_m

As expected, the isometry group of H_m depends on whether m is even or odd.

Suppose firstly that m is odd. In this case, (23) gives:

- (O1) The reflection of the z -plane about the imaginary axis, $re^{i\theta} \mapsto re^{i(\pi-\theta)}$, produces via X the reflectional symmetry about the (x_2, x_3) -plane in H_m .
- (O2) The rotation $re^{i\theta} \mapsto re^{i(\theta+\pi+\frac{\pi}{m+1})}$ of angle $\pi + \frac{\pi}{m+1}$ about the origin in the z -plane, gives that H_m is symmetric under the rotation of angle $\frac{\pi}{m+1}$ about the x_3 -axis composed by a reflection in the (x_1, x_2) -plane.

(O1), (O2) generate a subgroup of the extrinsic isometry group $\text{Iso}(H_m)$ of H_m , isomorphic to the dihedral group D_{2m+2} .

Now assume that m is even. Using (24), we obtain:

- (E1) The reflection $re^{i\theta} \mapsto re^{i(\pi-\theta)}$ of the z -plane about the imaginary axis produces via X the reflectional symmetry about the (x_2, x_3) -plane in H_m (this is a common feature of both the odd and even cases).
- (E2) The rotation $re^{i\theta} \mapsto re^{i(\theta+\frac{2\pi}{m+1})}$ of angle $\frac{2\pi}{m+1}$ about the origin in the z -plane, gives that H_m is symmetric under the rotation of angle $\frac{2\pi}{m+1}$.
- (E3) The antipodal map $re^{i\theta} \mapsto re^{i(\theta+\pi)}$ in the z -plane, produces a reflectional symmetry of H_m with respect to the (x_1, x_2) -plane.

(E1), (E2), (E3) generate a subgroup of $\text{Iso}(H_m)$ isomorphic to the group $D_{m+1} \times \mathbb{Z}_2$.

Repeating the argument in the proof of Lemma 4, we now deduce the following.

Lemma 9. *Regardless of the parity of m , these are all the (intrinsic) isometries of H_m .*

6 Moduli spaces of examples with a given complexity

Our next goal is to analyze the structure of the family of solutions of the period problem with a given complexity in the sense of Definition 7. For $m = 1$, we will obtain uniqueness of the Henneberg surface H_1 . This uniqueness is a special feature of the case $m = 1$, since continuous families of examples for complexities $m \geq 2$ can be produced.

We define the function $R: (0, \infty) \rightarrow (0, \infty)$, $R(r) = r - \frac{1}{r}$.

6.1 Solutions with complexity $m = 1$

Since $m = 1$, solving the period problem (19) descending to the 1-sided quotient reduces to solving

$$\overline{cA_1} = -cA_3, \quad \text{Im}(cA_2) = 0, \quad -\frac{\bar{c}}{c} = \frac{a_1 a_2}{a_1 a_2}. \quad (26)$$

Suppose that a list $(c, a_1, a_2) \in \mathbb{S}^1 \times (\mathbb{C}^*)^2$ is a solution of the 1-sided period problem, with associated branched minimal immersion X . Recall that $g(z) = z$ is its Gauss map. The list that gives rise to H_1 (Henneberg) is $(\pm 1, 1, i)$.

Remark 10. Since rotations of our surfaces are not allowed unless the rotation axis is vertical (see Remark 5), we we can assume $a_1 \in \mathbb{R}^+$ from now on, although we cannot assume $a_1 = 1$.

Write a_1, a_2 in polar coordinates as $a_1 = r_1$, $a_2 = r_2 e^{i\theta_2}$, $r_1, r_2 > 0$, $\theta_2 \in [0, 2\pi)$. (18) can be written as

$$\begin{aligned} P(z) &= z^4 - [R(r_1) + R(r_2)e^{i\theta_2}] z^3 - [1 + e^{2i\theta_2} - R(r_1)R(r_2)e^{i\theta_2}] z^2 \\ &\quad + [R(r_1)e^{2i\theta_2} + R(r_2)e^{i\theta_2}] z + e^{2i\theta_2}, \end{aligned}$$

hence

$$A_1 = R(r_1)e^{2i\theta_2} + R(r_2)e^{i\theta_2}, \quad (27)$$

$$A_2 = -[1 + e^{2i\theta_2} - R(r_1)R(r_2)e^{i\theta_2}], \quad (28)$$

$$A_3 = -[R(r_1) + R(r_2)e^{i\theta_2}]. \quad (29)$$

Writing $c = e^{i\beta}$, we have

$$\begin{aligned} \overline{cA_1} + cA_3 &= R(r_1) [e^{-i(\beta+2\theta_2)} - e^{i\beta}] + R(r_2) [e^{-i(\beta+\theta_2)} - e^{i(\beta+\theta_2)}] \\ &= R(r_1)e^{-i\theta_2} [e^{-i(\beta+\theta_2)} - e^{i(\beta+\theta_2)}] - 2R(r_2) \sinh(i(\beta + \theta_2)) \\ &= -2e^{-i\theta_2} R(r_1) \sinh(i(\beta + \theta_2)) - 2iR(r_2) \sin(\beta + \theta_2) \\ &= -2i [R(r_1)e^{-i\theta_2} + R(r_2)] \sin(\beta + \theta_2), \quad (30) \\ cA_2 &= -e^{i\beta} (1 + e^{2i\theta_2}) + R(r_1)R(r_2)e^{i(\beta+\theta_2)} \\ &= -e^{i(\beta+\theta_2)} (e^{-i\theta_2} + e^{i\theta_2}) + R(r_1)R(r_2)e^{i(\beta+\theta_2)} \\ &= -[2 \cosh(i\theta_2) - R(r_1)R(r_2)] e^{i(\beta+\theta_2)} \\ &= -[2 \cos \theta_2 - R(r_1)R(r_2)] e^{i(\beta+\theta_2)}. \quad (31) \end{aligned}$$

A list (c, a_1, a_2) solves the period problem if and only if the right-hand-side of (30) vanishes and the right-hand-side of (31) is real.

The third equation in (26) reduces to

$$e^{2i(\beta+\theta_2)} = -1. \quad (32)$$

Theorem 11. *The Henneberg surface H_1 is the only surface with $m = 1$ that solves the period problem and descends to a 1-sided quotient.*

Proof. By the above arguments, the right-hand-side of (30) vanishes, the right-hand-side of (31) is real and (32) holds.

(32) implies that $\sin(\beta + \theta_2) = \pm 1$. Since the right-hand-side of (30) vanishes, we have

$$R(r_1)e^{-i\theta_2} + R(r_2) = 0. \quad (33)$$

We have two possibilities:

- $r_1 = 1$. Thus (33) implies $r_2 = 1$. From, (32) we have $\beta + \theta_2 \equiv \pi/2 \pmod{\pi}$ and from (31) we have $\cos \theta_2 = 0$, thus $\theta_2 = \pi/2$ or $\theta_2 = 3\pi/2$. This gives the lists $(1, 1, i)$, $(-1, 1, i)$, $(1, 1, -i)$ and $(-1, 1, -i)$. All of them give rise to the Henneberg surface.
- $r_1 \neq 1$. This implies $e^{-i\theta_2} = -\frac{R(r_2)}{R(r_1)}$, which is real. Hence $e^{-i\theta_2} = \pm 1$. As the function $r \mapsto R(r)$ is injective, this implies $r_1 = r_2$ and $\theta_2 = \pi$ or $r_2 = 1/r_1$ and $\theta_2 = 0$. Since the right-hand-side of (31) is real and (32) holds, $2 \cos \theta_2 - R(r_1)R(r_2) = 0$. But in both cases $2 \cos \theta_2 - R(r_1)R(r_2)$ does not vanish. Hence this possibility cannot occur. \square

6.2 Solutions with complexity $m = 2$

Suppose that a list $(c = e^{i\beta}, a_1 = r_1, a_2 = r_2e^{i\theta_2}, a_3 = r_3e^{i\theta_3}) \in \mathbb{S}^1 \times \mathbb{R}^+ \times (\mathbb{C}^*)^2$ is a solution of the period problem with 1-sided quotient and associated branched minimal immersion X . The list that gives rise to H_2 is $(\pm i, 1, e^{i\pi/3}, e^{2i\pi/3})$.

Solving the period problem with 1-sided quotient is equivalent to solving

$$\overline{cA_2} = -cA_4, \quad \text{Im}(cA_3) = 0, \quad -\frac{\bar{c}}{c} = \frac{a_2 a_3}{a_2 \overline{a_3}} \quad (34)$$

The third equation in (34) reduces to

$$e^{2i(\beta+\theta_2+\theta_3)} = -1. \quad (35)$$

(18) can be written as

$$P(z) = z^6 + A_5z^5 + A_4z^4 + A_3z^3 + A_2z^2 + A_1z + A_0,$$

where

$$A_2 = e^{2i(\theta_2+\theta_3)} + e^{2i\theta_2} + e^{2i\theta_3} - R(r_1)R(r_2)e^{i(\theta_2+2\theta_3)} - R(r_1)R(r_3)e^{i(2\theta_2+\theta_3)} - R(r_2)R(r_3)e^{i(\theta_2+\theta_3)}, \quad (36)$$

$$A_3 = 2[R(r_2) \cos \theta_3 + R(r_3) \cos \theta_2 + R(r_1) \cos(\theta_2 - \theta_3) - \frac{1}{2}R(r_1)R(r_2)R(r_3)]e^{i(\theta_2+\theta_3)}, \quad (37)$$

$$A_4 = -(1 + e^{2i\theta_2} + e^{2i\theta_3}) + R(r_1)R(r_2)e^{i\theta_2} + R(r_1)R(r_3)e^{i\theta_3} + R(r_2)R(r_3)e^{i(\theta_2+\theta_3)}. \quad (38)$$

Thus,

$$\overline{cA_2} + cA_4 = 2e^{-i[\beta+2(\theta_2+\theta_3)]}F, \quad (39)$$

$$cA_3 = \pm 2iG \quad (40)$$

where

$$F = e^{2i\theta_3} + [2 \cos \theta_2 - R(r_1)R(r_2)]e^{i\theta_2} - R(r_3) [R(r_1) + R(r_2)e^{i\theta_2}]e^{i\theta_3}, \quad (41)$$

$$G = R(r_2) \cos \theta_3 + R(r_3) \cos \theta_2 + R(r_1) \cos(\theta_2 - \theta_3) - \frac{1}{2}R(r_1)R(r_2)R(r_3). \quad (42)$$

Remark 12. (I) From (42) we deduce that G is real, hence the condition $\text{Im}(cA_3) = 0$ only holds if and only if $G = 0$. We deduce that a list (c, a_1, a_2, a_3) solves the 1-sided period problem if and only if (35) holds and $F = G = 0$.

(II) The expression (41) is symmetric in $(r_2, \theta_2), (r_3, \theta_3)$. This can be deduced from the symmetry of A_2, A_4 , or directly checked by using the equality

$$e^{2i\theta_j} = 2 \cos \theta_j e^{i\theta_j} - 1, \quad (43)$$

which transforms (41) into

$$F = (1 + e^{2i\theta_2} + e^{2i\theta_3}) - R(r_1) \sum_{j=2}^3 R(r_j) e^{i\theta_j} - R(r_2)R(r_3) e^{i(\theta_2+\theta_3)}. \quad (44)$$

Lemma 13. *If $F = 0$, then the coefficient of $R(r_1)$ in (44) is non-zero.*

Proof. Suppose $R(r_2)e^{i\theta_2} + R(r_3)e^{i\theta_3} = 0$. This leads to one of the following two possibilities: (a) $e^{i\theta_2} = e^{i\theta_3}$ and $R(r_2) = -R(r_3)$ or else (b) $e^{i\theta_2} = -e^{i\theta_3}$ and $R(r_2) = R(r_3)$. (a) implies $r_3 = 1/r_2$ and thus, (44) gives $F = 1 + e^{2i\theta_2}(\frac{1}{r_2^2} + r_2^2)$. (b) implies $r_2 = r_3$ and (44) gives the same expression for F . In any case, we deduce from $F = 0$ that $e^{2i\theta_2}$ is real negative, hence $\frac{r_2^2}{r_2^4+1} = -e^{2i\theta_2} = 1$. This is impossible, since the function $x > 0 \mapsto \frac{x}{1+x^2}$ has a unique maximum at $x = 1$ with value $1/2$. \square

The next result describes a one-parameter family of non-trivial examples of complexity $m = 2$ different from H_2 .

Proposition 14. *Suppose that a list (c, a_1, a_2, a_3) solves the 1-sided period problem. Then:*

1. *If $r_1 = 1$, and at least one of r_2 or r_3 equals one, then $(c, a_1, a_2, a_3) = (\pm i, 1, e^{i\pi/3}, e^{2i\pi/3})$ and the example is H_2 .*
2. *If $\theta_2 + \theta_3 = 0 \pmod{\pi}$, then $r_2 = r_3$ or $r_2 = 1/r_3$ and (r_1, r_2) are given by the following functions of $\theta_2 \in (\frac{\pi}{4}, \frac{\pi}{3}] \cup [\frac{2\pi}{3}, \frac{3\pi}{4})$:*

$$R(r_1(\theta_2)) = \frac{1}{8\sqrt{2}} \frac{\sqrt{f(\theta_2) - 3}}{\cos \theta_2 \cos(2\theta_2)} [f(\theta_2) + 3 + 4 \cos(2\theta_2)], \quad (45)$$

$$R(r_2(\theta_2)) = -\frac{\sqrt{f(\theta_2) - 3}}{\sqrt{2}}, \quad (46)$$

or else (r_1, r_2) are given by the opposite expressions for both $R(r_1(\theta_2)), R(r_2(\theta_2))$, which exchange (r_1, r_2) by $(\frac{1}{r_1}, \frac{1}{r_2})$. Here, f is the function

$$f(\theta_2) = \sqrt{1 - 8 \cos(2\theta_2) - 8 \cos(4\theta_2)}. \quad (47)$$

Proof. If $r_1 = 1$, and at least one of r_2 or r_3 equals one, then (44) gives $1 + e^{2i\theta_2} + e^{2i\theta_3} = 0$ and (42) gives $R(r_2) \cos \theta_3 + R(r_3) \cos \theta_2 = 0$. Since at least one of r_2 or r_3 equals one, then at least one of R_2 or R_3 equals zero. In fact, both $R_2 = R_3 = 0$ (because otherwise

we get $\cos \theta_2 = 0$ or $\cos \theta_3 = 0$, which prevents $1 + e^{2i\theta_2} + e^{2i\theta_3}$ from cancelling), and thus, $r_2 = r_3 = 1$. In this setting, $1 + e^{2i\theta_2} + e^{2i\theta_3} = 0$ leads to $(c, a_1, a_2, a_3) = (\pm i, 1, e^{i\pi/3}, e^{2i\pi/3})$, which proves item 1.

Now assume $\theta_2 + \theta_3 = 0$. Then (44),(42) give respectively

$$1 + 2 \cos(2\theta_2) - R(r_2)R(r_3) = R(r_1)[R(r_2)e^{i\theta_2} + R(r_3)e^{-i\theta_2}], \quad (48)$$

$$(R(r_2) + R(r_3)) \cos \theta_2 + R(r_1) \cos(2\theta_2) = \frac{1}{2}R(r_1)R(r_2)R(r_3). \quad (49)$$

Observe that $R(r_1)$ cannot vanish by Lemma 13 (another reason is that otherwise, (49) gives $\cos \theta_2 = 0$, and (48) gives $-1 - R(r_2)R(r_3) = 0$ which is absurd). From (48) we deduce that $R(r_2)e^{i\theta_2} + R(r_3)e^{-i\theta_2}$ is real. This implies that $[R(r_2) - R(r_3)] \sin \theta_2 = 0$. We claim that $\sin \theta_2 \neq 0$; otherwise $\theta_2 \equiv 0 \pmod{\pi}$ and (48),(49) give the system

$$\begin{aligned} 3 - R(r_2)R(r_3) &= \pm R(r_1)[R(r_2) + R(r_3)], \\ R(r_1) \pm (R(r_2) + R(r_3)) &= \frac{1}{2}R(r_1)R(r_2)R(r_3), \end{aligned}$$

(with the same choice for signs), which can be easily seen not to have solutions.

Thus, $\sin \theta_2 \neq 0$ hence $R(r_2) = R(r_3)$ and $r_2 = r_3$. In this setting, (48),(49) reduce to

$$1 + 2 \cos(2\theta_2) - R(r_2)^2 = 2R(r_1)R(r_2) \cos \theta_2, \quad (50)$$

$$2R(r_2) \cos \theta_2 + R(r_1) \cos(2\theta_2) = \frac{1}{2}R(r_1)R(r_2)^2. \quad (51)$$

If we assume $\theta_2 + \theta_3 = \pi$, then (44),(42) give respectively

$$1 + 2 \cos(2\theta_2) + R(r_2)R(r_3) = R(r_1)[R(r_2)e^{i\theta_2} - R(r_3)e^{-i\theta_2}], \quad (52)$$

$$(-R(r_2) + R(r_3)) \cos \theta_2 - R(r_1) \cos(2\theta_2) = \frac{1}{2}R(r_1)R(r_2)R(r_3). \quad (53)$$

Again, $R(r_1)$ can not vanish due to Lemma 13. From (52) we deduce that $R(r_2)e^{i\theta_2} - R(r_3)e^{-i\theta_2}$ is real. This implies that $[R(r_2) + R(r_3)] \sin \theta_2 = 0$. We claim that $\sin \theta_2 \neq 0$; otherwise $\theta_2 \equiv 0 \pmod{\pi}$ and (52),(53) give the system

$$\begin{aligned} 3 + R(r_2)R(r_3) &= \pm R(r_1)[R(r_2) - R(r_3)], \\ -R(r_1) \pm (-R(r_2) + R(r_3)) &= \frac{1}{2}R(r_1)R(r_2)R(r_3), \end{aligned}$$

(with the same choice for signs), which again has no solutions. Thus, $\sin \theta_2 \neq 0$ hence $R(r_2) = -R(r_3)$ and $r_2 = 1/r_3$. In this setting, (48),(49) reduce again to (50) and (51).

The system (50),(51) has two equations and three unknowns r_1, r_2, θ_2 . Next we describe its solutions. Consider the function f given by (47). Then,

$$f(\pi - \theta_2) = f(\theta_2), \text{ for each } \theta_2, \quad f(\theta_{2,0}) = 0 = f(\pi - \theta_{2,0}),$$

where $\theta_{2,0} = \frac{1}{2} \cot^{-1} \left(\frac{9}{\sqrt{32\sqrt{10}+95}} \right) \sim 0.499841$, and the domain of f is $[\theta_{2,0}, \pi - \theta_{2,0}] + \pi\mathbb{Z}$.

The set $\{\theta_2 \in [\theta_{2,0}, \pi - \theta_{2,0}] \mid f(\theta_2) \geq 3\}$ equals $A := [\frac{\pi}{4}, \frac{\pi}{3}] \cup [\frac{2\pi}{3}, \frac{3\pi}{4}]$.

The unique solution (r_1, r_2) to the system (50),(51) as a function of θ_2 is given by (45), (46) and the opposite expressions for both $R(r_1(\theta_2)), R(r_2(\theta_2))$, which exchange (r_1, r_2) by $(\frac{1}{r_1}, \frac{1}{r_2})$. \square

6.2.1 The one-parameter family of examples in item 2 of Proposition 14

Observe that the map $\theta_2 \in (\frac{\pi}{4}, \frac{\pi}{3}] \mapsto \pi - \theta_2 \in [\frac{2\pi}{3}, \frac{3\pi}{4})$ is a diffeomorphism. Using the notation in item 2 of Proposition 14, for each $\theta_2 \in (\frac{\pi}{4}, \frac{\pi}{3}]$, we have

$$R(r_1(\pi - \theta_2)) = -R(r_1(\theta_2)), \quad R(r_2(\pi - \theta_2)) = R(r_2(\theta_2)). \quad (54)$$

Each of these lists with $\theta_2 \in (\frac{\pi}{4}, \frac{\pi}{3}] \cup [\frac{2\pi}{3}, \frac{3\pi}{4})$ solves the 1-sided period problem, hence it defines a non-orientable, branched minimal surface $H(\theta_2)$. Furthermore, (54) implies that

$$r_1(\pi - \theta_2) = \frac{1}{r_1(\theta_2)}, \quad r_2(\pi - \theta_2) = r_2(\theta_2). \quad (55)$$

We claim the surfaces $H(\theta_2)$ and $H(\pi - \theta_2)$ are congruent. To see this, note that the set of points $\{a_j, -1/\bar{a}_j : j = 1, 2, 3\}$ that defines f through (16) and generates the surface $H(\theta_2)$, is:

$$\left\{ r_1, \frac{-1}{r_1}, r_2 e^{i\theta_2}, \frac{1}{r_2} e^{i(\pi+\theta_2)}, r_2 e^{-i\theta_2}, \frac{1}{r_2} e^{i(\pi-\theta_2)} \right\}. \quad (56)$$

The analogous set of points for the surface $H(\pi - \theta_2)$ is given through (55):

$$\left\{ \frac{1}{r_1}, -r_1, -r_2 e^{-i\theta_2}, \frac{1}{r_2} e^{-i\theta_2}, -r_2 e^{i\theta_2}, \frac{1}{r_2} e^{i\theta_2} \right\},$$

which is up to sign the set described in (56). Therefore, the function f defined by equation (16) and the corresponding function \tilde{f} defined by the same formula for the surface $H(\pi - \theta_2)$ are related by $\tilde{f}(-z) = -f(z)$, for each $z \in \mathbb{C}$. Using that $\omega = f dz$ and $\tilde{\omega} = \tilde{f} dz$ define, via the Weierstrass representation (1), related branched minimal immersions $X = (x_1, x_2, x_3)$ for $H(\theta_2)$ and $\tilde{X} = (\tilde{x}_1, \tilde{x}_2, \tilde{x}_3)$ for $H(\pi - \theta_2)$, we get that $H(\theta_2)$ and $H(\pi - \theta_2)$ are congruent.

In the sequel, we will reduce our study to the family $\{H(\theta_2) \mid \theta_2 \in (\frac{\pi}{4}, \frac{\pi}{3}]\}$. From (45), (46) we have

$$\lim_{\theta_2 \rightarrow \pi/3^-} R(r_1(\theta_2)) = \lim_{\theta_2 \rightarrow \pi/3^-} R(r_2(\theta_2)) = 0,$$

which implies that

$$\lim_{\theta_2 \rightarrow \pi/3^-} H(\theta_2) = H_2.$$

We next identify the limit (after rescaling) of the surfaces $H(\theta_2)$ as $\theta_2 \rightarrow \pi/4^+$. We first observe that

$$\lim_{\theta_2 \rightarrow \pi/4^+} R(r_1(\theta_2)) = -\infty, \quad \lim_{\theta_2 \rightarrow \pi/4^+} R(r_2(\theta_2)) = 0. \quad (57)$$

This implies that the branch point $a_1 = a_1(\theta_2)$ is tending to zero, hence the limit of $H(\theta_2)$ when $\theta_2 \rightarrow \pi/4^+$ (if it exists) cannot be an example with complexity $m = 2$. Intuitively, it is clear that the complexity cannot increase when taking limits (even with different scales), hence by Theorem 11 it is natural to think that the limit of suitable re-scalings of $H(\theta_2)$ when $\theta_2 \rightarrow \pi/4^+$ be H_1 . We next formalize this idea.

Another consequence of (57) is that the list $(c, a_1, a_2, a_3) = (i, r_1(\theta_2), r_2(\theta_2)e^{i\theta_2}, r_2(\theta_2)e^{-i\theta_2})$ converges as $\theta_2 \rightarrow \pi/4^+$ to $(c, a_1, a_2, a_3) = (i, 0, e^{i\pi/4}, e^{-i\pi/4})$. After applying to $H(\theta_2)$ a homothety of ratio $r_1(\theta_2) > 0$ (which shrinks to zero), the Weierstrass data of the shrunk

surface $r_1(\theta_2)H(\theta_2)$ is $(g(z) = z, r_1(\theta_2)f(z))$, where $f(z)$ is given by (16). For $z \in \mathbb{C} \setminus \{0\}$ fixed,

$$\begin{aligned} & \lim_{\theta_2 \rightarrow \pi/4^+} r_1(\theta_2)f(z) \stackrel{(16)}{=} \lim_{\theta_2 \rightarrow \pi/4^+} r_1(\theta_2) \frac{i}{z^5} \prod_{j=1}^3 (z - a_j) \left(z + \frac{1}{a_j} \right) \\ &= \frac{i}{z^5} (z - e^{i\pi/4}) (z + e^{i\pi/4}) (z - e^{-i\pi/4}) (z + e^{-i\pi/4}) \lim_{\theta_2 \rightarrow \pi/4^+} (z - r_1(\theta_2)) (r_1(\theta_2)z + 1) \\ &= \frac{i}{z^4} (z - e^{i\pi/4}) (z + e^{i\pi/4}) (z - e^{-i\pi/4}) (z + e^{-i\pi/4}) := \widehat{f}(z). \end{aligned}$$

Plugging the Weierstrass data $(g(z) = z\widehat{f} dz)$ into (1), we obtain a parametrization of the limit surface of $r_1(\theta_2)H(\theta_2)$ as $\theta_2 \rightarrow \pi/4^+$ in polar coordinates $z = re^{i\theta}$:

$$\widehat{X}(re^{i\theta}) = \begin{pmatrix} -\frac{\sin\theta}{2}(r - \frac{1}{r}) + \frac{\sin(3\theta)}{6}(r^3 - \frac{1}{r^3}) \\ -\frac{\cos\theta}{2}(r - \frac{1}{r}) - \frac{\cos(3\theta)}{6}(r^3 - \frac{1}{r^3}) \\ -\cos\theta \sin\theta(r^2 + \frac{1}{r^2}) \end{pmatrix}. \quad (58)$$

We claim that this parametrization generates the Henneberg surface H_1 . To see this, observe that if we first perform the change of variables $\theta = \widetilde{\theta} + \pi/4$ and then rotate the surface an angle of $-\frac{\pi}{4}$ around the x_3 -axis, we get

$$\begin{pmatrix} \cos\left(\frac{\pi}{4}\right) & \sin\left(\frac{\pi}{4}\right) & 0 \\ -\sin\left(\frac{\pi}{4}\right) & \cos\left(\frac{\pi}{4}\right) & 0 \\ 0 & 0 & 1 \end{pmatrix} \cdot \widehat{X}(re^{i(\widetilde{\theta} + \frac{\pi}{4})}) = - \begin{pmatrix} \frac{\cos\widetilde{\theta}}{2}(r - \frac{1}{r}) - \frac{\cos(3\widetilde{\theta})}{6}(r^3 - \frac{1}{r^3}) \\ -\frac{\sin\widetilde{\theta}}{2}(r - \frac{1}{r}) - \frac{\sin(3\widetilde{\theta})}{6}(r^3 - \frac{1}{r^3}) \\ \frac{\cos(2\widetilde{\theta})}{2}(r^2 + \frac{1}{r^2}) \end{pmatrix},$$

which is, up to a sign, the parametrization given in (6) for H_1 .




Figure 5: Surfaces generated by the previous lists $(c, a_1, a_2, a_3) = (i, r_1(\theta_2), r_2(\theta_2)e^{i\theta_2}, r_2(\theta_2)e^{-i\theta_2})$ with $\theta_2 = 1$ (left), $\theta_2 = 0.83$ (center), $\theta_2 = 0.7854$ (right). The limit of $r_1(\theta_2)H(\theta_2)$ as $\theta_2 \rightarrow \pi/4^+ \sim 0.785398$ is the Henneberg surface H_1 .

6.2.2 Around H_2 the space of examples with complexity $m = 2$ is two-dimensional

Item 2 of Proposition 14 defines a non-compact family of non-orientable, branched minimal surfaces $\{H(\theta_2) \mid \theta_2 \in (\frac{\pi}{4}, \frac{\pi}{3}]\}$ inside the moduli space of examples with complexity $m = 2$. Apparently, the space of solutions for this complexity has real dimension 2 (the variables are $r_1, r_2, r_3, \theta_2, \theta_3$, $F = 0$ is a complex condition and $G = 0$ is a real condition). We can ensure this at least around H_2 via the implicit function theorem (this is consistent with item 2 of Proposition 14, since it imposes the extra condition $\theta_2 + \theta_3 = 0 \pmod{\pi}$), as we will show next.

Consider the (smooth) period map given by

$$P : \quad (\mathbb{R}^+)^3 \times \mathbb{R}^2 \quad \longrightarrow \quad \mathbb{R}^3 \equiv \mathbb{C} \times \mathbb{R} \\ ((r_1, r_2), (r_3, \theta_2, \theta_3)) \longmapsto (F(r_1, r_2, r_3, \theta_2, \theta_3), G(r_1, r_2, r_3, \theta_2, \theta_3)),$$

where F, G are given by (44), (42) respectively. Given $(r_1, r_2) \in (\mathbb{R}^+)^2$, let $P^{r_1, r_2} : \mathbb{R}^+ \times \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be the restriction of P to $\{(r_1, r_2)\} \times \mathbb{R}^+ \times \mathbb{R}^2$. Then,

$$d(P^{r_1, r_2})_{(r_3, \theta_2, \theta_3)} \equiv \begin{pmatrix} \frac{\partial \operatorname{Re}(F)}{\partial r_3} & \frac{\partial \operatorname{Re}(F)}{\partial \theta_2} & \frac{\partial \operatorname{Re}(F)}{\partial \theta_3} \\ \frac{\partial \operatorname{Im}(F)}{\partial r_3} & \frac{\partial \operatorname{Im}(F)}{\partial \theta_2} & \frac{\partial \operatorname{Im}(F)}{\partial \theta_3} \\ \frac{\partial G}{\partial r_3} & \frac{\partial G}{\partial \theta_2} & \frac{\partial G}{\partial \theta_3} \end{pmatrix}. \quad (59)$$

Recall that the list associated to H_2 is $(r_1, r_2, r_3, \theta_2, \theta_3) = (1, 1, 1, \pi/3, 2\pi/3)$. Imposing this choice of parameters and computing the determinant of (59) we get

$$d(P^{1,1})_{(1, \pi/3, 2\pi/3)} = 2\sqrt{3} \neq 0.$$

Thus, the implicit function theorem gives an open neighborhood $U \subset (\mathbb{R}^+)^2$ of $(r_1, r_2) = (1, 1)$, an open set $W \subset (\mathbb{R}^+)^3 \times \mathbb{R}^2$ with $(r_1, r_2, r_3, \theta_2, \theta_3) = (1, 1, 1, \pi/3, 2\pi/3) \in W$ and a smooth map $\varphi : U \rightarrow \mathbb{R}^3$ such that all the solutions $(r_1, r_2, \theta_1, \theta_2, \theta_3)$ around $(1, 1, 1, \pi/3, 2\pi/3)$ of the equation $P(r_1, r_2, \theta_1, \theta_2, \theta_3) = 0$ are of the form $(\theta_1, \theta_2, \theta_3) = \varphi(r_1, r_2)$. By Remark 12(I), the list

$$(c = e^{i\beta(r_1, r_2)}, r_1, r_2 e^{i\theta_2}, r_3 e^{i\theta_3})$$

with $\beta = \beta(r_1, r_2)$ given by (35) solves the 1-sided period problem and so, it defines a 1-sided branched minimal surface. This produces a 2-parameter deformation of the surface H_2 in the moduli space of examples with $m = 2$ around H_2 , which in turn describes the whole moduli space around H_2 .

Remark 15. A nice consequence of the classical Leibniz formula for the derivative of a product is a recursive law that gives the coefficients of the polynomial $P(z)$ defined by (18) in terms of the coefficients of the related polynomial for one complexity less. To obtain this

recursive law, we first adapt the notation to the complexity:

$$P_{m+1}(z) := \prod_{j=1}^{m+1} (z - a_j) \left(z + \frac{1}{a_j} \right) = \sum_{h=0}^{2m+2} A_{m+1,h} z^h. \quad (60)$$

(19) can now be written

$$\overline{cA_{m+1,m}} = -cA_{m+1,m+2}, \quad \text{Im}(cA_{m+1,m+1}) = 0. \quad (61)$$

We want to find expressions for the above coefficients $A_{m+1,m}$, $A_{m+1,m+2}$, $A_{m+1,m+1}$, depending only on coefficients of the type $A_{m,h}$ (i.e., for one complexity less). Writing $a_j = r_j e^{i\theta_j}$ in polar coordinates, observe that

$$P_{m+1}(z) := P_m(z)Q_{m+1}(z), \quad \text{where } Q_{m+1}(z) = (z - r_{m+1}e^{i\theta_{m+1}}) \left(z + \frac{e^{i\theta_{m+1}}}{r_{m+1}} \right).$$

Hence for $h \in \{m, m+1, m+2\}$,

$$A_{m+1,h} = \frac{1}{h!} P_{m+1}^{(h)}(0) = \frac{1}{h!} (P_m Q_{m+1})^{(h)}(0) = \frac{1}{h!} \sum_{k=0}^h \binom{h}{k} P_m^{(k)}(0) Q_{m+1}^{(h-k)}(0),$$

where in the last equality we have used Leibniz formula. Since Q_{m+1} is a polynomial of degree two, its derivatives of order three or more vanish. Hence we can reduce the last sum to terms where the index k satisfies $h - k \leq 2$, i.e., $k \in \{h-2, h-1, h\}$ and thus,

$$\begin{aligned} A_{m+1,h} &= \frac{1}{h!} \left[\binom{h}{h-2} P_m^{(h-2)}(0) Q_{m+1}''(0) + \binom{h}{h-1} P_m^{(h-1)}(0) Q_{m+1}'(0) + \binom{h}{h} P_m^{(h)}(0) Q_{m+1}(0) \right] \\ &= \frac{1}{h!} \left[\frac{h!}{(h-2)!2} P_m^{(h-2)}(0) \cdot 2 - h P_m^{(h-1)}(0) R(r_{m+1}) e^{i\theta_{m+1}} - P_m^{(h)}(0) e^{2i\theta_{m+1}} \right] \\ &= \left[\frac{1}{(h-2)!} P_m^{(h-2)}(0) - \frac{1}{(h-1)!} P_m^{(h-1)}(0) R(r_{m+1}) e^{i\theta_{m+1}} - \frac{1}{h!} P_m^{(h)}(0) e^{2i\theta_{m+1}} \right] \\ &= A_{m,h-2} - A_{m,h-1} R(r_{m+1}) e^{i\theta_{m+1}} - A_{m,h} e^{2i\theta_{m+1}}, \end{aligned} \quad (62)$$

which is the desired recurrence law. (62) can be used to find solutions to (61) for complexity $m = 3$ besides the most symmetric example H_3 , but the equations are complicated and we will not give them here.

References

- [1] M. do Carmo and C. K. Peng. Stable complete minimal surfaces in \mathbb{R}^3 are planes. *Bull. Amer. Math. Soc. (N.S.)*, 1:903–906, 1979. MR0546314, Zbl 442.53013.
- [2] D. Fischer-Colbrie and R. Schoen. The structure of complete stable minimal surfaces in 3-manifolds of nonnegative scalar curvature. *Comm. on Pure and Appl. Math.*, 33:199–211, 1980. MR0562550, Zbl 439.53060.

- [3] L. Henneberg. Über solche minimalfläche, welche eine vorgeschriebene ebene curve sur geodätischen line haben. Doctoral Dissertation, Eidgenössisches Polytechnikum, Zurich, 1875.
- [4] W. H. Meeks III and J. Pérez. Hierarchy structures in finite index cmc surfaces. Work in progress.
- [5] M. Micallef and B. White. The structure of branch points in minimal surfaces and in pseudoholomorphic curves. *Ann. of Math.*, 141(1):35–85, 1995. MR1314031, Zbl 0873.53038.
- [6] B. Odehnal. On algebraic minimal surfaces. *KoG*, 20:61–78, 2016.
- [7] A. V. Pogorelov. On the stability of minimal surfaces. *Soviet Math. Dokl.*, 24:274–276, 1981. MR0630142, Zbl 0495.53005.
- [8] A. Ros. One-sided complete stable minimal surfaces. *J. Differential Geom.*, 74:69–92, 2006. MR2260928, Zbl 1110.53009.
- [9] J. Tysk. Eigenvalue estimates with applications to minimal surfaces. *Pacific J. of Math.*, 128:361–366, 1987. MR0888524, Zbl 0594.58018.