

BITANGENTS TO PLANE QUARTICS VIA TROPICAL GEOMETRY: RATIONALITY, \mathbb{A}^1 -ENUMERATION, AND REAL SIGNED COUNT

HANNAH MARKWIG, SAM PAYNE, AND KRIS SHAW

ABSTRACT. We explore extensions of tropical methods to arithmetic enumerative problems such as \mathbb{A}^1 -enumeration with values in the Grothendieck-Witt ring, and rationality over Henselian valued fields, using bitangents to plane quartics as a test case. We consider quartic curves over valued fields whose tropicalizations are smooth and satisfy a mild genericity condition. We then express obstructions to rationality of bitangents and their points of tangency in terms of twisting of edges of the tropicalization; the latter depends only on the tropicalization and the initial coefficients of the defining equation modulo squares. We also show that the GW-multiplicity of a tropical bitangent, i.e., the multiplicity with which its lifts contribute to the \mathbb{A}^1 -enumeration of bitangents as defined by Larson and Vogt [21], can be computed from the tropicalization of the quartic together with the initial coefficients of the defining equation. As an application, we show that the four lifts of most tropical bitangent classes contribute $2\mathbb{H}$, twice the class of the hyperbolic plane, to the \mathbb{A}^1 -enumeration, providing evidence for a conjecture of Larson and Vogt on the \mathbb{A}^1 -enumeration of bitangents over the reals. These results rely on a degeneration theorem relating the Grothendieck-Witt ring of a Henselian valued field to the Grothendieck-Witt ring of its residue field, in residue characteristic not equal to two.

CONTENTS

| | |
|--|----|
| 1. Introduction | 2 |
| 2. Preliminaries | 7 |
| 2.1. Field convention | 7 |
| 2.2. Tropicalizations of plane quartics | 8 |
| 2.3. Bitangents to tropicalized quartics | 10 |
| 3. Lifting tropical bitangents and tropical points of tangencies | 15 |
| 3.1. Local lifting equations and their solutions | 15 |
| 3.2. Lifting conditions and twisted edges | 18 |
| 3.3. Comparing lifting for different fields | 23 |
| 4. The Grothendieck-Witt ring and \mathbb{A}^1 -enumerative invariants | 25 |

2010 *Mathematics Subject Classification.* 14N10, 14T20, 14T25, 14G27.

Key words and phrases. bitangents to plane quartics, real enumerative geometry, tropical bitangent classes, \mathbb{A}^1 -enumerative geometry, Grothendieck-Witt ring, rationality of enumerative solutions.

| | |
|---|----|
| 4.1. The Grothendieck-Witt ring | 25 |
| 4.2. Grothendieck-Witt rings over valued fields | 27 |
| 4.3. The Qtype of a bitangent to a quartic | 28 |
| 4.4. The GW-multiplicity of a tropical bitangent class | 29 |
| 4.5. Comparing the Qtype_{L^∞} for different fields | 38 |
| 5. The real GW-multiplicity in tropical geometry | 39 |
| References | 41 |
| Appendix A. Classification, computations and table | 44 |
| A.1. Classification of generic bitangent shapes | 44 |
| A.2. Details on Theorem 4.23 including case-by-case analysis | 52 |
| A.3. Exceptional GW-multiplicities | 59 |

1. INTRODUCTION

Tropical geometry is well-known for its applications to complex and real enumerative geometry, especially for plane curves. Here, we explore extensions of tropical methods to arithmetic enumerative problems such as \mathbb{A}^1 -enumeration with values in the Grothendieck-Witt ring, and rationality over Henselian valued fields, using bitangents to plane quartics as a test case.

If the tropicalization of a plane quartic is smooth (locally isomorphic to the tropicalization of a linear space) then it has exactly 7 deformation classes of tropical bitangents. These are in natural bijection with the odd tropical theta characteristics and each contains the tropicalization of 4 algebraic bitangents [5, 17, 24]. Cueto and the first author have given an exhaustive classification of the combinatorial types of these bitangent classes [6]. As an application, when the ground field K is real Puiseux series $\mathbb{R}\{\{t\}\}$ and the tropicalization satisfies the genericity constraints from [6, Remark 2.10], they showed that the number of K -rational bitangents tropicalizing into a bitangent class is either 0 or 4. In other words, roughly speaking, the obstruction to rationality over $\mathbb{R}\{\{t\}\}$ is independent of the choice of bitangent in a given bitangent class.

Throughout, let K be a Henselian valued field of residue characteristic not equal to 2, with 2-divisible value group, and with a section of the valuation $\sigma: \text{val}(K^\times) \rightarrow K^\times$. Let k be the residue field of K . Let $Q \in K[x, y, z]$ be a homogeneous polynomial of degree 4 such that the projective plane algebraic curve $C = V(Q)$ is smooth and $\text{Trop}(C)$ is tropically smooth. Assume furthermore that $\text{Trop}(C)$ satisfies the genericity constraints from [6, Remark 2.10], the details of which we recall in Section 2.3.

Note that K could be the Puiseux series field $k\{\{t\}\}$ over any field k of characteristic not equal to 2. Such fields are commonly considered in tropical geometry. Other possibilities include mixed characteristic fields such as $\mathbb{Q}_p(p^{1/2^\infty}) = \bigcup_n \mathbb{Q}_p(p^{1/2^n})$, for $p \neq 2$. Our arguments are not sensitive to distinctions between pure and

mixed characteristic; any reader who prefers to do so may safely assume that $K = k\{\{t\}\}$ is a field of Puiseux series.

Lifting of tropical bitangents to general ground fields. Our work begins with the observation that the algebraic calculations used to prove the lifting result from [6] over the reals are only mildly sensitive to the ground field.

Theorem 1.1 (Theorem 3.3). *Let $C = V(Q)$ be a quartic curve defined over K with $\text{Trop}(C)$ a generic tropicalized quartic and suppose Λ is a liftable tropical bitangent to C . Then whether or not Λ lifts over K is determined by $\text{Trop}(C)$ and the equivalence classes of initials of the coefficients of Q in $k^*/(k^*)^2$.*

When the residue field $k = \mathbb{R}$, then $\mathbb{R}^*/(\mathbb{R}^*)^2 \cong \mathbb{Z}/2\mathbb{Z}$. Knowing the tropicalization of C together with the equivalence classes of initials of the coefficients of Q modulo squares is precisely the information required to carry out Viro's patchworking [34, 14]. Motivated by this connection, we express obstructions to rationality in the case of arbitrary fields in terms of *twisting* of edges of the tropicalization, thus extending to general fields the concept of twisting coming from amoebas over the real numbers, see [3].

Theorem 1.2 (Theorem 3.14). *The number of K -rational bitangents to C that tropicalize into each bitangent class of tropical bitangents is either 0 or 4.*

When K is $\mathbb{R}\{\{t\}\}$, Cueto and the first author showed that the points of tangency of a K -rational bitangent are K -rational [6, Theorem 7.1]. Again, we observe that this is only mildly sensitive to the ground field.

Theorem 1.3 (Corollary 3.15). *Assume that 2 and 3 are in $(k^\times)^2$. Then the points of tangency of a K -rational bitangent to C are K -rational.*

Note that Theorem 1.3 assumes the residue characteristic is also not equal to 3.

The proofs of Theorems 1.2 and 1.3 involve solving systems of equations over the residue field to determine the initials of the coefficients of the defining equation of the bitangents and points of tangency in suitable local coordinates, and then lifting via Hensel's lemma. This is carried out case-by-case, according to the Cueto-Markwig classification from [6]. Details of the case analysis are provided in the Appendix.

As a consequence of the lifting theorems discussed above, we also prove that the liftability of tropical bitangents over different fields are related when the groups of the residue fields modulo squares are isomorphic, see Theorem 3.16 and Corollary 3.17.

The tropical approach to the \mathbb{A}^1 -enumeration of bitangents to a plane quartic, and its foundations. Even when it is possible to count K -rational solutions to a geometric problem, the naive enumeration may not be invariant under deformations. Familiar examples include counts of real rational plane curves of

given degree and genus, passing through an appropriate number of specified points in general position. In such cases, there are often *refined* enumerative counts, such as Welschinger invariants for real rational plane curves, where objects are counted with multiplicities in such a way that the result is invariant under deformation. More recently, techniques from arithmetic geometry and \mathbb{A}^1 -homotopy theory have produced more general \mathbb{A}^1 -enumerative invariants taking values in Grothendieck-Witt rings of quadratic forms that specialize to classical enumerative invariants over \mathbb{C} , to Welschinger invariants and related real enumerative invariants over \mathbb{R} , and produce new invariants of interest over other fields [13, 18, 25]. For the \mathbb{A}^1 -enumeration of the 27 lines on cubic surfaces and the 28 bitangents to plane quartics, see [19] and [21], respectively. We recall the relevant definitions in Section 4.

We write $t^v := \sigma(v)$ for $v \in \text{val}(K^\times)$, where σ denotes the section of our valued field K . For $A \in K$, we let $\text{in}(A)$ be the image of $At^{-\text{val}(A)}$ in the residue field and call it the initial of A . If K is the field of Puiseux series, $\text{in}(A)$ is the leading coefficient of the series A .

Theorem 1.4 (Theorem 4.23). *The \mathbb{A}^1 -enumerative multiplicity of a bitangent L to C depends only on the tropicalization of C and on the initials of its coefficients modulo squares.*

Moreover, we give an effective algorithm for computing the \mathbb{A}^1 -enumerative count of all bitangents to C in each of the 7 deformation classes, and hence the \mathbb{A}^1 -enumerative count of all 28 bitangents, as elements of the Grothendieck-Witt ring $\text{GW}(K)$ (see Remark 4.16).

Theorem 1.4 relies on the following observation relating the Grothendieck-Witt ring of K to that of its residue field. Recall our standing assumption that K is a Henselian valued field of residue characteristic not equal to 2, with 2-divisible value group, and with a section of the valuation $\sigma: \text{val}(K^\times) \rightarrow K^\times$.

Theorem 1.5 (Theorem 4.7). *There is an isomorphism of Grothendieck-Witt rings*

$$\text{GW}(K) \xrightarrow{\sim} \text{GW}(k), \quad \langle A \rangle \mapsto \langle \text{in}(A) \rangle.$$

This isomorphism does not depend on the section σ .

The above theorem is a variant of Springer's theorem on the Witt group of a Henselian discretely valued field, as generalized to arbitrary value groups, e.g., in [9], in the special case where the value group is 2-divisible. Theorems of this form are a natural starting point for the application of tropical methods to \mathbb{A}^1 -enumerative geometry. In forthcoming work, Jaramillo Puentes and Pauli define enriched tropical intersection multiplicities for 0-dimensional intersections of tropical hypersurfaces [33]. As a consequence they prove a tropical version of a quadratically enriched Bézout theorem from [26] and also introduce an enriched Bernstein-Khovanskii-Kushnirenko theorem.

Remark 1.6. Our hypothesis that the value group $\text{val}(K^\times)$ is 2-divisible is a simplifying assumption that is necessary for Theorems 1.4 and 1.5, as stated. Nevertheless, this framework, and all of the computations that we carry out for bitangents to plane quartics, can be extended to the case where K is Henselian of residue characteristic not equal to 2 with arbitrary value group, but then one must also keep track of the valuations of the coefficients, modulo valuations of squares. The calculations needed for our case analysis of bitangents are already sufficiently complicated when the value group is 2-divisible that we have chosen to compromise generality in this way. We hope that this work, along with that of Jaramillo Puentes and Pauli [33], may serve as a proof of concept that tropical methods are relevant and useful to \mathbb{A}^1 -enumerative geometry.

If an \mathbb{A}^1 -enumerative multiplicity equals the hyperbolic plane $\mathbb{H} = \langle 1 \rangle \oplus \langle -1 \rangle$, then it contributes 2 to the corresponding complex count and 0 to the corresponding signed count over the real numbers. We prove that for many types of tropical bitangent classes in the classification, the \mathbb{A}^1 -enumerative multiplicities of the four algebraic bitangents that tropicalize into this class add up to $2\mathbb{H}$. The exceptional cases for which this does not hold true are listed in Appendix A.3. To give an impression which tropical bitangent classes contribute $2\mathbb{H}$ to the \mathbb{A}^1 -enumerative count, we list some sufficient conditions here. For the complete list in terms of the classification of tropical bitangent classes, see Theorem A.2.

Theorem 1.7 (Theorems 4.23 and A.2). *Let S be a tropical bitangent class that is either compact or contains no tropical bitangent line with its vertex on $\text{Trop}(C)$. Then the contribution of S to the \mathbb{A}^1 -enumerative count of bitangents to C is $2\mathbb{H}$.*

We also show that the contributions of tropical bitangents classes to the \mathbb{A}^1 -enumerative count of bitangents over different fields are related if the groups of the residue fields modulo squares are isomorphic, see Theorem 4.24 and Corollary 4.25.

The real signed count of bitangents to plane quartics. In [21], Larson and Vogt investigated the \mathbb{A}^1 -enumeration of bitangents to a plane quartic, relative to a fixed line at infinity, and the resulting signed count over reals. Over the reals, they show that the signed count is equal to 4 if the real locus of the quartic does not meet the line at infinity. They also show that the signed count is nonnegative and conjecture that it is bounded above by 8. The field of real Puiseux series $\mathbb{R}\{\{t\}\}$ also has its Grothendieck-Witt ring generated by the two elements $\langle 1 \rangle$ and $\langle -1 \rangle$. The \mathbb{A}^1 -enumeration of bitangents can thus be expressed as $a \cdot \langle 1 \rangle + b \cdot \langle -1 \rangle$ for some a, b , and the signed count is $a - b$.

Theorem 1.8 (Theorem 5.2). *Let $C = V(Q)$ be a quartic over $\mathbb{R}\{\{t\}\}$ whose tropicalization is smooth and generic. Then the signed count of bitangents of C is either 0, 2 or 4.*

As many, but not all, of the tropical bitangent classes contribute $2\mathbb{H}$ to the \mathbb{A}^1 -enumeration of bitangents, we deduce that many tropical bitangents give a total contribution of 0 to the signed count when working over the reals. The total signed count of 2 and 4 can only be reached for those tropicalized quartics which admit exceptional tropical bitangent classes as listed in Appendix A.3. The strategy to prove Theorem 5.2 is to investigate how the dual motifs (see Definition 2.15) of such exceptional tropical bitangents can fit into the dual Newton subdivision of a tropicalized quartic and show that, at most, there can be either one that contributes 4 or one or two that contribute 2 to the signed count, with the rest contributing 0.

Organization of this paper. This paper is organized as follows. In Section 2 we discuss preliminaries. We fix our convention for the fields we study, introduce tropicalizations of plane quartics and discuss tropical bitangents and their bitangent classes.

In Section 3, we discuss obstructions to rationality. Subsection 3.1 reviews known techniques to solve so-called local lifting equations which provide local obstructions for a tropical bitangent to have a K -rational lift. In Subsection 3.2, we generalize the concept of twisted edges from the reals to more general fields. We also generalize to situations with so-called relative twisting, in which the relative behaviour of a tropical bitangent line and the tropicalized quartic play a role. We express obstructions to K -rationality in terms of twisting of edges. In Subsection 3.3, we sum up our results on K -rationality of tropical bitangent classes, and compare the behaviour over different fields.

Section 4 discusses aspects of \mathbb{A}^1 -enumeration. Subsection 4.1 recalls the definition of the Grothendieck-Witt ring. Subsection 4.2 contains our result (Theorem 1.5, 4.7) relating the Grothendieck-Witt ring of K to the Grothendieck-Witt ring of its residue field. This result lays the foundation for the infusion of tropical methods into the study of \mathbb{A}^1 -enumerative geometry, and we apply it in Subsection 4.4 to investigate the \mathbb{A}^1 -enumerative geometry of tropical bitangent classes to tropicalized quartics. Our main result in this Subsection is Theorem 4.23, which sums up the statements of Theorems 1.4 and 1.7 in the introduction. In this Subsection, we provide the methods to prove that many tropical bitangent classes contribute $2\mathbb{H}$ to the \mathbb{A}^1 -enumeration of bitangents to quartics, and spell out the details for one case of the classification of tropical bitangent classes. The remaining cases are revisited in Appendix A.2. Before we turn to the \mathbb{A}^1 -enumerative geometry of tropical bitangents, in Subsection 4.3, we review the results of [21] on the \mathbb{A}^1 -enumeration of bitangents to quartics. Subsection 4.4 presents the results from the tropical approach to the \mathbb{A}^1 -enumeration of bitangents to quartics and Subsection 4.5 explores relations of the \mathbb{A}^1 types of lifts of tropical bitangents over different fields.

In Section 5, we consider tropical \mathbb{A}^1 -enumeration of bitangents to quartics over real closed fields. We prove Theorem 1.8 (5.2) providing evidence in support

of Larson and Vogt’s conjecture on the possible values for the signed count of bitangents over the reals. We also point out that this result can be obtained without referring to the techniques of \mathbb{A}^1 -enumeration: as the signed count of bitangents to a real quartic has a geometric interpretation (see Subsection 4.3), and one can thus also use the method of Viro’s patchworking to associate the correct sign to a tropical bitangent.

The Appendix contains details of the classification of tropical bitangents. Part A.1 spells out the details about all tropical bitangent classes and their dual motifs (see Definition 2.15) up to \mathbb{S}_2 -symmetry. The original classification in [6] is up to \mathbb{S}_3 -symmetry, but we have to break some of the symmetry to fix the line at infinity. Part A.2 adds the details to Theorem 4.23 (see Theorem A.2) by going through the classification and showing which bitangent classes precisely contribute $2\mathbb{H}$ to the \mathbb{A}^1 -enumeration of bitangents to quartics. Part A.3 finally lists the exceptional tropical bitangent classes and their \mathbb{A}^1 -enumerative multiplicity, which is given in terms of the initials of the coefficients of the defining equation of the quartic, as claimed in Theorem 1.4.

Acknowledgements. The authors would like to thank María Angélica Cueto, Alheydis Geiger, Andrés Jaramillo Puentes, Danny Krashen, Mario Kummer, Hannah Larson, Marta Panizzut, Sabrina Pauli, Eugenii Shustin, Isabel Vogt, and Kirsten Wickelgren for valuable discussions related to this work and helpful comments on earlier drafts. The first author acknowledges support by DFG-grant MA 4797/9-1. The second author acknowledges support from NSF grants DMS-2001502 and DMS-2053261. The third author acknowledges support of the Trond Mohn Foundation project “Algebraic and topological cycles in complex and tropical geometry”. The project was initiated during a stay of the second author in Tübingen funded by the Max-Planck Humboldt-Medal 2018. We thank the Max-Planck society and the Humboldt society for support.

2. PRELIMINARIES

2.1. Field convention. Throughout the paper, let K be a Henselian valued field whose value group $\text{val}(K^\times)$ is 2-divisible, and let σ be a section of the valuation, i.e., a group homomorphism

$$\sigma: \text{val}(K^\times) \rightarrow K^\times$$

such that $\text{val} \circ \sigma$ is the identity. Let $R \subset K$ denote the valuation ring $\mathfrak{m} \subset R$ the maximal ideal, and $k = R/\mathfrak{m}$ the residue field. Throughout, we assume the characteristic of k is not 2.

Example 2.1. Such fields exist with any given residue field k ; for instance, one could take the Puiseux series field $k\{\{t\}\}$, the generalized power series field $k((t^{\mathbb{R}}))$, or the completion of either. The hypotheses are also satisfied by mixed characteristic

fields such as $\mathbb{Q}_p(p^{1/2}, p^{1/4}, p^{1/8}, \dots)$, where $p^{1/2}$ is a square root of p , $p^{1/4}$ is a square root of $p^{1/2}$, and so on; a section is given by $\sigma(v) = p^v$.

Definition 2.2. We write $t^v := \sigma(v)$, for $v \in \text{val}(K^\times)$.

This notation intentionally emphasizes the analogy with generalized power series.

Definition 2.3 (Initials). Let $A \in K^\times$. Then the initial of A , $\text{in}(A)$, is the image of $At^{-\text{val}(A)}$ in k^\times .

In case K is a generalized power series field, $\text{in}(A)$ is the ‘‘initial coefficient’’.

2.2. Tropicalizations of plane quartics. Tropicalization can be viewed as degeneration of algebraic varieties defined over K . Here, we introduce it only for plane curves.

Definition 2.4 (Tropicalization). Let $C \in \mathbb{P}_K^2$ be a plane curve and assume that K is algebraically closed. The *tropicalization* is defined by taking $-\text{val}$ componentwise to the torus points:

$$\text{Trop}(C) := \overline{\{(-\text{val } x, -\text{val } y) \mid (x, y) \in C \cap (K^\times)^2\}},$$

where the bar denotes the Euclidean closure in \mathbb{R}^2 .

If K is not algebraically closed, we apply minus valuation coordinatewise to all points of C in the algebraic closure of K to obtain the tropicalization.

Since we require our field to be Henselian, the valuation extends uniquely to the algebraic closure. Equivalently, $\text{Trop}(C)$ is the image of the Berkovich analytification of $C \cap (\mathbb{G}_m^2)$ under coordinatewise valuation [32].

Remark 2.5. One can also define a compactification of \mathbb{R}^2 that is the tropicalization of \mathbb{P}_K^2 , including its toric boundary, as in Section 3 of [32]. The definition above then extends naturally to take boundary points into account. A smooth quartic is never contained in the toric boundary, and hence meets the torus non-trivially. Consequently, this compact extended tropicalization equals the closure of the tropicalization of its points in the torus as defined above (see Lemma 3.1.1 [30]). We can therefore restrict attention to the tropicalization in the torus as described above.

Example 2.6 (Tropical line). Let $C = V(x + y + t)$. We can parametrize C as

$$\{(x, -x - t) \mid x \neq -t, x \in K^\times\}.$$

By considering the three cases $\text{val}(x) > 1$, $\text{val}(x) < 1$, and $\text{val}(x) = 1$, where in the last case the situation $\text{in}(x) = -1$ plays an important role, we can see that the tropicalization consists of three rays starting at the point $(-1, -1)$, one diagonal, one horizontal and one vertical.




FIGURE 1. On the left, the set D for the polynomial Q in Example 2.9, on the right is the Newton subdivision Δ_Q .

Remark 2.7 (Tropical dual plane). By varying the coefficients in Example 2.6, one can see that the tropicalization of any line whose defining equation has three nonzero coefficients (what we are calling a *tropical line*) consists of three rays centered at a vertex whose coordinates are determined by the valuations of the coefficients. Thus, we can and do identify the *tropical dual plane* parametrizing tropical lines in \mathbb{R}^2 with the tropical plane \mathbb{R}^2 itself, where a tropical line is identified with its vertex. (Note, however, that this does not extend to an identification of the compact tropicalization of \mathbb{P}^2 with that of $(\mathbb{P}^2)^\vee$.)

Definition 2.8 (Newton subdivision). Let $Q \in K[x, y, z]$ be a homogeneous polynomial of degree d , and denote the coefficient of the monomial $x^i y^j z^{d-i-j}$ by A_{ij} . Let $D \subset \mathbb{R}^2 \times \mathbb{R}$ be the convex hull of the set

$$\{((i, j), -\text{val}(A_{ij}) \mid A_{ij} \neq 0\}.$$

Project the faces of D which can be seen from above down to \mathbb{R}^2 . The images of these faces form a convex subdivision Δ_Q of the Newton polygon, called the *Newton subdivision*.

Example 2.9. Let $Q = t^{-1}x^2 + t^{-2}xy + t^{-1}y^2 + t^{-2}x + t^{-2}y + t^{-1}$. Figure 1 shows the set D on the left and the projection of its upper faces, i.e. its Newton subdivision, on the right.

Theorem 2.10 (Duality theorem). *For $C = V(Q)$, the tropical curve $\text{Trop}(C)$ is the 1-skeleton of a subdivision of \mathbb{R}^2 that is dual to the Newton subdivision Δ_Q , i.e., there is a natural inclusion reversing bijection of faces between these two subdivisions.*

A proof can be found, for example, in [27], Proposition 3.11. This duality is illustrated in Figure 2.

The tropicalization of a plane curve is called a *tropical plane curve*. From duality to a Newton subdivision, we conclude that tropical curves satisfy the *balancing condition*: if we weight each edge by the lattice length of its dual edge, then the weighted sum of the primitive edge generators around a vertex is 0.




FIGURE 2. The tropicalization of the curve from Example 2.9 and the Newton subdivision Δ_Q .

A tropical plane curve is *smooth* of degree d if it is dual to a unimodular triangulation of the triangle with vertices $(0, 0)$, $(0, d)$, $(d, 0)$. Here, unimodular means that it is subdivided to d^2 triangles each having area $\frac{1}{2}$.

2.3. Bitangents to tropicalized quartics. Suppose that two tropical plane curves Γ_1, Γ_2 intersect transversally, i.e. the intersection is a finite set of points each of which is contained in the interior of an edge of both Γ_1 and Γ_2 . Let $P \in \Gamma_1 \cap \Gamma_2$. Choose weighted direction vectors u_1, u_2 for edges of Γ_1, Γ_2 emanating from P . Then the *intersection multiplicity* of the two curves at P is $\Gamma_1 \cdot \Gamma_2 |_P = |\det(u_1, u_2)|$, and $\Gamma_1 \cdot \Gamma_2 = \sum_{P \in \Gamma_1 \cap \Gamma_2} \Gamma_1 \cdot \Gamma_2 |_P$. The balancing condition ensures that this definition does not depend on the choice of direction vectors. If the curves do not intersect transversally, we use the *stable* intersection: choose a direction v so that Γ_1 and $\Gamma_2 + \epsilon \cdot v$ intersect transversally whenever ϵ is small enough. Then the stable intersection is

$$\lim_{\epsilon \rightarrow 0} \Gamma_1 \cdot (\Gamma_2 + \epsilon \cdot v).$$

When the intersection is not transverse, the number of intersection points, counted with multiplicity, in the preimage of a connected component of the intersection of the tropicalizations is the sum of the multiplicities of the points of the stable tropical intersection that lie in this component, see [31], Theorem 6.4.

Definition 2.11 (Bitangent). We say that a tropical line Λ is *bitangent* to a tropicalized quartic $\text{Trop}(C)$ in \mathbb{R}^2 if Λ and $\text{Trop}(C)$ intersect in either

- one connected component with total intersection multiplicity 4, or
- two connected components each with total intersection multiplicity 2.

In the first case, the intersection can be a point, or a segment of an edge of Γ containing a vertex (in which case the two points of tangency tropicalize to the vertex and the midpoint of the segment), or three edges adjacent to the same vertex (in which case the two tangency points tropicalize to the midpoints of the two segments we obtain by subtracting the shortest edge length from the others; see Figure 3).

For a bitangent Λ to $\text{Trop}(C)$, we call a connected component of the intersection $\Lambda \cap \text{Trop}(C)$ a *tropical tangency component*. In the second case of Definition 2.11,




FIGURE 3. A local picture of a tropicalized quartic $\text{Trop}(C)$ together with the tropicalization of the two tangency points of a bitangent ℓ to C . Here the line ℓ is a lift of a tropical line Λ whose vertex coincides with the center vertex of $\text{Trop}(C)$. The intersection of Λ and $\text{Trop}(C)$ is a single connected component with tropical intersection multiplicity 4. The tropical line does not appear in the figure.

the intersection can be a point, or a segment of an edge of Γ (in which case the tangency point tropicalizes to the midpoint of the segment).

It is often the case that a tropicalized quartic $\text{Trop}(C)$ in \mathbb{R}^2 admits infinitely many tropical bitangents. A natural question to ask is: which of the infinitely many tropical bitangent lines are tropicalizations of bitangent lines of C ?

Definition 2.12 (Lift of a bitangent). Let C be a quartic defined over K . Let Λ be a tropical line which is tropically bitangent to $\text{Trop}(C)$. A bitangent line L of C is called a *lift* of Λ if $\text{Trop}(L) = \Lambda$. If such a lift exists, we say Λ is *liftable*. If in addition the lift is defined over K , we say Λ is *liftable over K* .

We declare that two tropical bitangent lines are equivalent if we can move one tropical line to the other (connecting their corresponding points in the tropical dual plane) while maintaining bitangency, i.e., the equivalence classes are the connected components of the locus of tropical bitangents in the tropical dual plane. For tropicalized quartics, this is equivalent to saying that the tropical bitangent lines correspond to the same theta characteristic in the tropical Jacobian [2, Definition 3.8].

When studying tropical bitangents and their lifts, we restrict to *generic* tropicalized quartics (see Remark 2.17 and [24], 3.3). This ensures that the tropicalized quartic is smooth, that the tropical tangency components are not contained on the same ray, and that when the intersection $\text{Trop}(C) \cap \Lambda$ consists of a vertex and three ray segments for a liftable tropical bitangent Λ , then the tropicalization of tangency points are not on the vertex.

Definition 2.13 (Bitangent class). A connected component of the set of tropical bitangents lines to a tropicalized quartic $\text{Trop}(C)$ in the tropical dual plane (see Remark 2.7) is called a *bitangent class* of $\text{Trop}(C)$.

By [2], a smooth tropicalized quartic has precisely 7 bitangent classes. The statement can be generalized for non-smooth tropicalized quartics, in which case bitangent classes must be counted with a suitable multiplicity [23].

By [6], a bitangent class is a polyhedral complex. First, we equip it with the coarsest polyhedral complex structure that is needed (for all the shapes in the classification, such a coarsest polyhedral complex structure exists). By duality (see Theorem 2.10), \mathbb{R}^2 obtains a polyhedral complex structure with cells the vertices and edges of $\text{Trop}(C)$, and the connected components of $\mathbb{R}^2 \setminus \text{Trop}(C)$.

Definition 2.14 (Shape of a bitangent class). The *shape* of a bitangent class is the polyhedral complex structure obtained by refining with the polyhedral complex structure of \mathbb{R}^2 given by $\text{Trop}(C)$, where the containment of a polyhedral cell in $\text{Trop}(C)$ is encoded using colors. We identify the tropical dual plane with the original \mathbb{R}^2 to make this refinement.

In [6], shapes of bitangent classes are classified up to \mathbb{S}_3 -symmetry (see Figure 6). There are 41 such shapes, denoted with letters (A), (B), ... or double letters (EE) etc.

Definition 2.15 (Dual motif, [12]). Let $\text{Trop}(C)$ be a smooth tropicalized quartic and B one of its seven bitangent classes. Let Λ be a tropical bitangent in B . The *dual motif* of the bitangent class B is the set of all triangles and edges in the subdivision whose dual vertices resp. edges intersect a bitangent in B .

For an example, see Figure 5. All tropical bitangents in the bitangent class considered there share the same tropical tangency component on the lower right, the intersection of the vertical segment. Dual to this edge, with its two end vertices, are the two triangles on the right in the dual Newton subdivision. The other tropical tangency component varies for the tropical bitangents in B . It is either on an edge, or an end vertex of the same edge. Accordingly, we obtain another pair of triangles on the left in the dual Newton subdivision for the dual motif.

If, as in Figure 5, all tropical bitangents in a bitangent class have two disjoint tropical tangency components, we can divide the dual motif into two parts, one for each tropical tangency component.

Example 2.16. Figure 4 shows a tropicalized quartic and a tropical bitangent line. The tropical line intersects the tropicalized quartic in two tropical tangency components. For the right tropical tangency component, we have to use stable intersection to check that it is a tangency. The tropicalization of a tangency point for any lift is the midpoint of the segment of intersection. We can see that we can move the vertex of the tropical bitangent upwards or downwards, until we hit vertices of the tropicalized quartic, maintaining the bitangency. The bitangent class is therefore a line segment as depicted in Figure 5. As it is disjoint from $\text{Trop}(C)$, its shape is also just a segment. This is a shape of type (E) in the




FIGURE 4. A tropicalized quartic, with its dual Newton subdivision, and a tropical bitangent line. We can move the vertex of the tropical bitangent upwards or downwards maintaining the bitangency.




FIGURE 5. The bitangent class and dual motif of the tropical bitangent to the tropicalized quartic depicted in Figure 4.

classification of [6]. In [24], it is shown that, if C is any algebraic quartic with this tropicalization, exactly 2 of the 28 bitangent lines to C tropicalize to the tropical line with vertex the upper red point, exactly 2 to the one with vertex the lower red point, and none to a point in the interior of the red segment (see Theorem 2.18). Figure 5 also depicts the dual motifs for the two tropical tangency components. Circled in red is the vertex dual to the connected component of \mathbb{R}^2 in which the bitangent class is contained.

Remark 2.17 (Genericity of tropicalized quartics). Compared to [24], we require an additional genericity assumption on our tropicalized quartics: we subdivide the cone in the secondary fan corresponding to the unimodular triangulation according to the types of tropical bitangent classes that can occur and require our tropicalized quartic to correspond to a point in the interior of a cone in this subdivision. This subdivision of the secondary fan is computed in [11]. Put differently, we require the edge lengths of $\text{Trop}(C)$ to be generic in the sense that no unexpected alignment of vertices happens, see Figure 6. This picture shows a local piece of a tropicalized quartic on the left. For a generic element in the corresponding cone of the secondary fan, we expect the two lengths l_1 and l_2 to be different. If these lengths are equal, as depicted on the right, the two lower vertices align.



FIGURE 6. Local pictures of tropicalized quartics. An unexpected alignment of two vertices is depicted on the right. For a generic quartic, as shown on the left, the two lengths l_1 and l_2 are different.

We call a bitangent shape that occurs for such a generic tropicalized quartic a *generic bitangent shape*.

The following statements concern lifts of tropical bitangent classes over $\mathbb{C}\{\{t\}\}$ and $\mathbb{R}\{\{t\}\}$.

Theorem 2.18 ([5, 24]). *Each bitangent class of a generic tropicalized quartic has 4 lifts over $\mathbb{C}\{\{t\}\}$. More precisely, there are either 4 tropical lines in the bitangent class which each lift once, or 2 which each lift twice, or 3 of which one lifts twice, or 1 which lifts four times.*

Theorem 2.19 ([6]). *Each bitangent class of a generic tropicalized quartic has either 0 or 4 lifts over $\mathbb{R}\{\{t\}\}$, i.e. the possible obstruction for real lifting is the same for all representatives of a given tropical bitangent class that have a complex lift.*

These results are based on the classification of shapes of bitangent classes in [6] up to \mathbb{S}_3 -symmetry. When considering GW-multiplicities as in Section 4, we fix the line $\{z = 0\}$ and do not have \mathbb{S}_3 -symmetry for that reason, only \mathbb{S}_2 -symmetry for exchanging the variables x and y .

In Appendix A.1, we classify tuples of shapes of generic bitangent classes together with their dual motifs up to \mathbb{S}_2 -symmetry exchanging x and y . The classification is built on the classification in [6]. It restricts the cases studied there to generic shapes, considers $\mathbb{S}_3/\mathbb{S}_2$ -orbits, and pairs up with dual motifs. In Appendix A.1, we depict all dual motifs together with local pictures of the tropicalized tangency points of the liftable tropical bitangents in the bitangent class are given.

Example 2.20. Consider the plane quartic $C = V(Q)$ for

$$Q(x, y) = t^{36}x^4 + t^{18}x^3y + t^2x^2y^2 + t^{18}xy^3 + t^{36}y^4 + t^{23}x^3 + t^6x^2y + t^6xy^2 \\ + t^{23}y^3 + t^{12}x^2 + xy + t^{12}y^2 + t^2x + t^2y + 1$$

over the field of Puiseux series $\mathbb{C}\{\{t\}\}$.

Its tropicalization $\text{Trop}(C)$ together with the seven bitangent classes (in red) is depicted in Figure 7. The liftable tropical bitangents are depicted as red dots. If




FIGURE 7. A tropicalized quartic with its seven bitangent classes.

a bitangent class has four red dots, each lifts once, if it has two red dots, each lifts twice and a single dot lifts to four bitangents.

3. LIFTING TROPICAL BITANGENTS AND TROPICAL POINTS OF TANGENCIES

3.1. Local lifting equations and their solutions. We review the local lifting techniques for tropical bitangent lines to tropicalized plane quartics from [6, 24].

As before, we let $C = V(Q)$ be a quartic and denote the coefficients of the defining polynomial Q by A_{ij} . For a point $p \in \text{Trop}(C)$, let $\text{in}_p(Q)$ denote the sum of the (dehomogenized) terms $\text{in}(A_{ij})x^i y^j \in k[x, y]$ for which the maximum $\max\{-\text{val}(A_{ij} + ip_x + jp_y)\}$ is attained. We call it the *initial form* of Q at p .

Definition 3.1 (Local equations for tropicalized tangency points). Let T be a connected component of the intersection of $\text{Trop}(C)$ with a tropical bitangent Λ , and assume a lift of Λ is given by the equation $\ell = y + M + Nx$, where M and N are unknowns. Let $A(T) = \bigcup_{p \in T} \text{supp}\{\text{in}_p Q\}$ be the union of the supports of the initial forms of Q for all $p \in T$. We let Q_T be the restriction of Q to the terms appearing in $A(T)$, and call it the *local equation for C at T* .

Analogously, we let $B(T)$ be the union of the supports of the initial forms of ℓ for all $p \in T$, and ℓ_T , the restriction of ℓ to $B(T)$, the local equation of ℓ .

We let $W_S = \frac{\partial Q_T}{\partial x} \cdot \frac{\partial \ell_T}{\partial y} - \frac{\partial Q_T}{\partial y} \cdot \frac{\partial \ell_T}{\partial x}$ be the local version of the Wronskian.

Notice that vanishing of the Wronskian $W = \frac{\partial Q}{\partial x} \cdot \frac{\partial \ell}{\partial y} - \frac{\partial Q}{\partial y} \cdot \frac{\partial \ell}{\partial x}$ implies that the gradients of $V(Q)$ and $V(\ell)$ coincide, i.e. that the line $V(\ell)$ is tangent to $V(Q)$ at (x, y) .

If $p \in T$ is the tropicalization of a tangency point, we solve the *local lifting equations*

$$Q_T(x, y) = 0, \quad \ell_T(x, y) = 0, \quad W_T(x, y) = 0.$$

If $T = \{p\}$ equals just the tropical tangency component, then this amounts to solving the equations of the initial forms of Q , ℓ and W at p . If T is a segment of intersection involving two vertices of an edge E of $\text{Trop}(C)$, then Q_T contains not only the terms of the edge E^\vee dual to E , but also the terms corresponding to the vertices which form triangles with E^\vee in the dual subdivision. If T is a segment

containing one vertex V of $\text{Trop}(C)$ and the vertex of Λ , then $\ell_T = \ell$ and Q_T contains the terms corresponding to E^V and the vertex of the triangle dual to V . If T consists of three segments joined at a vertex V , Q_T contains the terms of the triangle dual to V plus the terms of its neighbouring triangles, and $\ell_T = \ell$.

If T is more than just a point, the equations above are hard to solve. One uses a technical trick which is called a tropical modification or tropical refinement to produce solutions up to the order which appears as biggest valuation of the terms involved. For a discussion on how to treat the case of a segment T containing two vertices of $\text{Trop}(C)$, see [24] Proposition 3.7 and 3.9. For the case of a segment containing the vertex of Λ , see Lemma 5.2, Case (3a) [6] and Proposition 3.12 in [24]. For the case of three segments joined at a vertex, see Proposition 3.12 in [24]. For general background on the technique of tropical modification, see e.g. [4, 14, 28]. Once we solved in such a way for x, y up to terms of higher valuation, it follows from the Henselian property (see also [8], Exercises 7.25, 7.26 and [14], Chapter 2) that these solutions can uniquely be completed to an element in K . For more details, see Section 2.3 in [24]. If the initials are in the residue field k , then the complete solutions are in K .

Example 3.2. We compute the initials of the four lifts of the bitangent class from Example 2.16. Consider first the upper vertex. There are two lifts over the algebraic closure which tropicalize to it. We denote the equations of those two lifts by $M_1 + N_1x + y = 0$ resp. $M_2 + N_2x + y = 0$. Let p_1 be the left tropical tangency component. In $\text{Trop}(C)$, it is at the vertex dual to $A_{01}y + A_{12}xy^2 + A_{22}x^2y^2$. In the tropical bitangent line, it is in the horizontal ray, i.e. dual to $M_i + y$. We denote the initials of the A_{ij} by a_{ij} and analogously for the coefficients of the line equations. Then the initials (x, y) of the lift of the left tropical tangency component satisfy

$$\begin{aligned} a_{01}y_{i1} + a_{12}x_{i1}y_{i1}^2 + a_{22}x_{i1}^2y_{i1}^2 &= 0, \\ m_i + y_{i1} &= 0, \\ 2 \cdot a_{22} \cdot x_{i1} \cdot y_{i1}^2 + a_{12} \cdot y_{i1}^2 &= 0, \end{aligned}$$

for $i = 1, 2$. Solving for m_i , x_{i1} and y_{i1} (for example by computing a Gröbner basis of the ideal defined by the three equations using a computer algebra system such as SINGULAR [7] or OSCAR [29]) we obtain, for $i = 1, 2$,

$$m_i = -\frac{4 \cdot a_{01} \cdot a_{22}}{a_{12}^2}, \quad x_{i1} = -\frac{a_{12}}{2 \cdot a_{22}}, \quad y_{i1} = \frac{4 \cdot a_{01} \cdot a_{22}}{a_{12}^2}.$$

The second tropical tangency component is the same for all liftable tropical bitangents in this bitangent class (i.e. for the tropical line with vertex the upper vertex of the red edge, and for the tropical line with vertex the lower vertex of the red edge in Figure 5). Therefore, we now first study the two lifts tropicalizing to the lower vertex and their tropical tangency component on the left. It is at the

vertex dual to $A_{01}y + A_{11}xy + A_{22}x^2y^2$. The three equations to solve are thus:

$$\begin{aligned} a_{01}y_{i1} + a_{11}x_{i1}y_{i1} + a_{22}x_{i1}^2y_{i1}^2 &= 0, \\ m_i + y_{i1} &= 0, \\ 2 \cdot a_{22} \cdot x_{i1} \cdot y_{i1}^2 + a_{11} \cdot y_{i1} &= 0, \end{aligned}$$

for $i = 3, 4$, and the solutions are

$$m_i = -\frac{a_{11}^2}{4 \cdot a_{01} \cdot a_{22}}, \quad x_{i1} = -\frac{2 \cdot a_{01}}{a_{11}}, \quad y_{i1} = \frac{a_{11}^2}{4 \cdot a_{01} \cdot a_{22}}.$$

Let p_2 be the right tropical tangency component. This tropical tangency component appears not only for the two lifts of the upper vertex of the bitangent class, but also for the lower ones. That is, we can use the corresponding local lifting equations to solve for one tropical tangency component for each of the four lifts. The point p_2 is contained in a connected component of the intersection $\text{Trop}(C)$ and a tropical bitangent which consists of a bounded edge of $\text{Trop}(C)$. Dual to the vertices of this bounded edge are two triangles of the dual subdivision which meet along the edge joining $(2, 1)$ and $(3, 1)$. Therefore, we must take as local equation

$$A_{21}x^2y + A_{31}x^3y + A_{30}x^3 + A_{22}x^2y^2.$$

In the tropical line, the tropical tangency component p_2 is contained in the vertical ray, thus the local equation to consider is $m_i + n_i x$ for $i = 1, \dots, 4$. As we saw earlier, m_i is determined by the left (upper or lower) tropical tangency component. In the local equations, we now solve for the ratio $\frac{m_i}{n_i}$, which is then sufficient to compute all coefficients of the lifts. It turns out that there are two solutions whose initials coincide. In order to be able to differentiate the two solutions whose initials coincide, we also list some contributions of higher valuation.

The solutions we obtain are

$$\begin{aligned} \frac{M_i}{N_i} &= \frac{A_{21}}{A_{31}} \pm \frac{2A_{22}}{A_{31}} \sqrt{-\frac{A_{30}A_{21}}{A_{22}A_{31}}} + h.o.t. \\ X_{i2} &= -\frac{A_{21}}{A_{31}} \mp \frac{2A_{22}}{A_{31}} \sqrt{-\frac{A_{30}A_{21}}{A_{22}A_{31}}} + h.o.t. \\ Y_{i2} &= \pm \sqrt{-\frac{A_{30}A_{21}}{A_{22}A_{31}}} + h.o.t. \end{aligned}$$

for $i = 1, \dots, 4$. Here, we write the solutions not in terms of the initial a_{ij} but in terms of the whole coefficients A_{ij} , to keep track of the contributions of higher valuation.

Combining the results of the local lifting equations, we can now list the initials of the coefficient of the bitangent equation and the tangency point for all four lifts:

| i | m_i | n_i | x_{i1} | y_{i1} | x_{i2} | y_{i2} |
|-----|---|---|----------------------------------|--|--------------------------|--|
| 1 | $-\frac{4 \cdot a_{01} \cdot a_{22}}{a_{12}^2}$ | $-\frac{4 \cdot a_{01} \cdot a_{22} \cdot a_{31}}{a_{12}^2 \cdot a_{21}}$ | $-\frac{a_{12}}{2 \cdot a_{22}}$ | $\frac{4 \cdot a_{01} \cdot a_{22}}{a_{12}^2}$ | $-\frac{a_{21}}{a_{31}}$ | $\sqrt{-\frac{a_{30} a_{21}}{a_{22} a_{31}}}$ |
| 2 | $-\frac{4 \cdot a_{01} \cdot a_{22}}{a_{12}^2}$ | $-\frac{4 \cdot a_{01} \cdot a_{22} \cdot a_{31}}{a_{12}^2 \cdot a_{21}}$ | $-\frac{a_{12}}{2 \cdot a_{22}}$ | $\frac{4 \cdot a_{01} \cdot a_{22}}{a_{12}^2}$ | $-\frac{a_{21}}{a_{31}}$ | $-\sqrt{-\frac{a_{30} a_{21}}{a_{22} a_{31}}}$ |
| 3 | $-\frac{a_{11}^2}{4 \cdot a_{01} \cdot a_{22}}$ | $-\frac{a_{11}^2 \cdot a_{31}}{4 \cdot a_{01} \cdot a_{22} \cdot a_{21}}$ | $-\frac{2 \cdot a_{01}}{a_{11}}$ | $\frac{a_{11}^2}{4 \cdot a_{01} \cdot a_{22}}$ | $-\frac{a_{21}}{a_{31}}$ | $\sqrt{-\frac{a_{30} a_{21}}{a_{22} a_{31}}}$ |
| 4 | $-\frac{a_{11}^2}{4 \cdot a_{01} \cdot a_{22}}$ | $-\frac{a_{11}^2 \cdot a_{31}}{4 \cdot a_{01} \cdot a_{22} \cdot a_{21}}$ | $-\frac{2 \cdot a_{01}}{a_{11}}$ | $\frac{a_{11}^2}{4 \cdot a_{01} \cdot a_{22}}$ | $-\frac{a_{21}}{a_{31}}$ | $-\sqrt{-\frac{a_{30} a_{21}}{a_{22} a_{31}}}$ |

As in the example above, these local lifting equations can be solved for any case appearing in the combinatorial classification in Appendix A.1, and the solutions are always given by Laurent terms in the initials a_{ij} of the coefficients of Q , or by square roots thereof.

The consequence for lifting can be summed up as follows:

Theorem 3.3. *Let $C = V(Q)$ be a quartic curve defined over K with $\text{Trop}(C)$ a generic tropicalized quartic and suppose Λ is a liftable tropical bitangent to C . Then whether or not Λ lifts over K is determined by $\text{Trop}(C)$ and the equivalence classes of initials of the coefficients of Q in $k^*/(k^*)^2$.*

3.2. Lifting conditions and twisted edges. In the case of a tropical tangency component on a segment of intersection, the potential obstruction for lifting can be phrased in terms of twisted edges. We start by stating the lifting solutions, which can be determined with the methods described in Subsection 3.1. In the following, we denote vertices of the dual subdivision corresponding to the term $A_{ij}x^i y^j$ in (the dehomogenized version of) Q by V_{ij} .

Lemma 3.4. *Assume the tropical line Λ intersects the tropicalized quartic $\text{Trop}(C)$ in two connected components of which one is a segment of a horizontal edge dual to an edge connecting V_{1k} and V_{1k+1} . This edge forms triangles with two points V_{0i} and V_{j2} . Then there are two lifts for the constant coefficient M of a line $V(M + Nx + y)$ and for the tangency points (x, y) tropicalizing to this segment.*

If the vertex of Λ is not contained in the segment, we have

$$\begin{aligned}
M &= \frac{A_{1k}}{A_{1k+1}} \pm 2\sqrt{(-1)^{i+j} A_{0i} A_{2j} \left(\frac{A_{1k}}{A_{1k+1}}\right)^{i+j} \frac{(-1)^k A_{1k+1}^{k-1}}{A_{1k}^k} + \dots}, \\
y &= -\frac{A_{1k}}{A_{1k+1}} \mp 2\sqrt{(-1)^{i+j} A_{0i} A_{2j} \left(\frac{A_{1k}}{A_{1k+1}}\right)^{i+j} \frac{(-1)^k A_{1k+1}^{k-1}}{A_{1k}^k} + \dots}, \\
x &= \pm\sqrt{(-1)^{i-j} \frac{A_{0i}}{A_{2j}} \left(\frac{A_{1k}}{A_{1k+1}}\right)^{i-j} + \dots},
\end{aligned}$$

where we assume that the lift of Λ has the equation $y + M + Nx$. Here, the dots stand for terms of higher valuation.



FIGURE 8. The dual subdivision to a bounded edge e and its endpoints in a tropicalized curve, first in the case when $\delta(e) = 0$ on the left, and then when $\delta(e) = 1$ on the right.

If the vertex of Λ is contained in the segment, we have

$$\begin{aligned}
 M &= \frac{A_{1k}}{A_{1k+1}} \pm 2\sqrt{(-1)^{i+k+1} A_{0i} A_{1k+1} N \left(\frac{A_{1k}}{A_{1k+1}}\right)^{i+k} \frac{(-1)^k A_{1k+1}^{k-1}}{A_{1k}^k} + \dots}, \\
 y &= -\frac{A_{1k}}{A_{1k+1}} \mp 2\sqrt{(-1)^{i+k+1} A_{0i} A_{1k+1} N \left(\frac{A_{1k}}{A_{1k+1}}\right)^{i+k} \frac{(-1)^k A_{1k+1}^{k-1}}{A_{1k}^k} + \dots}, \\
 x &= \pm \sqrt{(-1)^{i-k+1} \frac{A_{0i}}{A_{1k+1} N} \left(\frac{A_{1k}}{A_{1k+1}}\right)^{i-k} + \dots}
 \end{aligned}$$

In the following, for a vertex r in the dual Newton subdivision of Q , we let a_r be the initial of the corresponding coefficient A_r of Q .

Definition 3.5 (Twisted edge). Let C be a curve defined over K with smooth tropicalization $\text{Trop}(C)$. Let e be a bounded edge in $\text{Trop}(C)$ so that the dual subdivision corresponding to e is the segment joined by lattice points $q, q' \in \mathbb{Z}^2$ and the two triangles of the subdivision intersecting along this segment have vertices $r, r' \in \mathbb{Z}^2$. We say that the edge e is *twisted over k* if

$$\sqrt{(-1)^{\delta(e)} a_r a_{r'} (a_q a_{q'})^{\delta(e)}} \notin k,$$

where $\delta(e) = 0$ if $r = r' \pmod{2}$ and $\delta(e) = 1$ if $r \neq r' \pmod{2}$.

Example 3.6. To see the two possible cases for $\delta(e)$ consider the lattice polytopes in Figure 8. Notice that on the left hand side the coordinates of the vertices of the two triangles opposite the overlapping edge e have vector difference $(0, 0)$ modulo 2. Therefore, the points r and r' are equal modulo 2 and $\delta(e) = 0$. On the right hand side the vector difference of the coordinates of the two vertices has difference $(1, 1)$ modulo 2. Therefore we have $\delta(e) = 1$ in this case.

Example 3.7. The motivation for the terminology of twisting comes from considering amoebas of curves defined over the real numbers. Let $k = \mathbb{R}$ and $K = \mathbb{R}\{\{t\}\}$. Let C be a curve defined over K with smooth tropicalization $\text{Trop}(C) = \Gamma$. Suppose that the defining polynomial Q of C has convergent coefficients for t sufficiently small. Then we can consider the family of real curves C_t defined by




FIGURE 9. An illustration of the non-twisting and twisting of the real amoeba (in red) from Example 3.7.

the family of polynomials Q_t for t sufficiently small. The tropical curve Γ is also obtained as the limit as s tends to infinity of amoebas of the family of curves \tilde{C}_s defined by the family of polynomials Q_s , where $s = t^{-1}$. The amoeba of a curve \tilde{C}_s is defined to be $\mathcal{A}_s := \text{Log}_s(\tilde{C}_s)$, where Log_s denotes the coordinatewise base s logarithm of absolute values, see [3, Section 2]. Moreover, for s sufficiently large, the real amoeba $\mathbb{R}\mathcal{A}_s := \text{Log}_s(\mathbb{R}\tilde{C}_s)$ either crosses a bounded edge e of Γ or it does not, see Figure 9. Following [3, Section 3], call the edge e *twisted* about the edge e if the amoeba crosses it.

The twisting of the edges of Γ are determined solely by the signs of the leading terms of the coefficients of the defining polynomial of C as described in [3, Remark 3.9]. Again let q, q' and r, r' denote the lattice points dual to the edge e as in Definition 3.5. Letting $\gamma_q, \gamma_{q'}, \gamma_r$ and $\gamma_{r'}$ denote the signs of the leading terms of the corresponding coefficients, then

- (1) if r and r' are distinct modulo 2 then e is twisted if and only if $\gamma_q \gamma_{q'} \gamma_r \gamma_{r'} > 0$;
- (2) if r and r' are equal modulo 2 then e is twisted if and only if $\gamma_r \gamma_{r'} < 0$.

This is equivalent to the notion of twist in Definition 3.5 when $k = \mathbb{R}$. See Figure 8 for two examples of dual subdivisions exhibiting both cases in terms of r and r' above.

Using our local lifting solutions, obstructions to lifting can be phrased in terms of the twisting of an edge:

Proposition 3.8. *Let C be a curve defined over K such that $\text{Trop}(C)$ is a smooth tropicalized curve. Let Λ be a tropical line and suppose a bounded edge e of $\text{Trop}(C)$ is a tropical tangency component of $\Lambda \cap \text{Trop}(C)$ strictly contained in a ray of Λ . If $L = V(\ell)$ is a lift of Λ which is tangent to C at a point tropicalizing to e then the local equation ℓ_e is defined over K if and only if e is not twisted over k .*

Proof. The statement follows by applying Lemma 3.4. □

We now want to consider the situation when $\text{Trop}(C)$ and Λ have a tropical tangency component T which is a segment strictly contained in an edge e of $\text{Trop}(C)$, see Figure 10, i.e. the vertex of Λ is contained in the segment of intersection. Let s, s' be the lattice points dual to the edge of Λ containing T and q, q' be the lattice points dual to e in C . Moreover assume the points s and q are labeled so as to be on corresponding sides with respect to e . By Lemma 3.4, in order for a lift L of Λ to have a tangency point P with C which tropicalizes to T we must have

$$(1) \quad \frac{b_s}{b_{s'}} = \frac{a_q}{a_{q'}}.$$

Definition 3.9 (Relatively twisted edge). Let $\text{Trop}(C)$ be smooth, and let Λ be a tropical line which has a tropical tangency component T which is a segment strictly contained in an edge e of $\text{Trop}(C)$. Assume e is dual to the edge qq' and Λ meets the vertex of e which is dual to the triangle spanned by qq' and r . Let $a_q, a_{q'}, a_r$ denote the corresponding initials. We use the analogous indices for the initials of the coefficients of the polynomial defining L , i.e. b_s and $b_{s'}$ are the initials of the monomials of the endpoints of the edge in the dual subdivision to the ray of Λ which contains e , and $b_{r'}$ is the initial of the remaining coefficient of the defining polynomial of L .

We say that the edge e is relatively twisted over k with respect to C and L if

$$\frac{b_s}{b_{s'}} = \frac{a_q}{a_{q'}} \quad \text{and} \quad \sqrt{(-1)^{\delta(e,L)+1} a_r (a_q a_{q'})^{\delta(e,L)} b_{r'} a_q b_s} \in k,$$

where $\delta(e, L) = 0$ if the edge in the dual subdivision with endpoints a_r, a_q is parallel modulo 2 to the edge in the dual subdivision with endpoints $b_{r'}, b_s$, and otherwise $\delta(e, L) = 1$.

Notice that the first condition $\frac{b_s}{b_{s'}} = \frac{a_q}{a_{q'}}$ implies that the second condition is equivalent to

$$\sqrt{(-1)^{\delta(e,L)+1} a_r (a_q a_{q'})^{\delta(e,L)} b_{r'} a_q b_{s'}} \in k.$$

Also notice that by the balancing condition for tropical curves, the term $\delta(e, L)$ in the above definition can be equivalently be computed by comparing the directions mod 2 of the edges with endpoints $a_r, a_{q'}$ and $b_{r'}, b_{s'}$.

Example 3.10. Once again the above definition of relative twists is motivated by the geometric picture over the real numbers, as was explained in the case of twisted edges in Example 3.7. When considering real tropical intersections of curves, Le Texier defines the notion of relative twists [22]. As in Example 3.7, consider the real amoebas of both the family of lines and the family of curves, as drawn in Figure 10. The coordinatewise logarithm map is the composition of the absolute value map and the logarithm map $\log_t \mathbb{R}_{>0} \rightarrow \mathbb{R}$. Therefore, each branch of the amoeba of a real curve can be labelled by the orthant of $(\mathbb{R}^*)^2$ from




FIGURE 10. The left hand side depicts the parts of the dual subdivision to Γ and Λ which are involved in the definition of relative twists. On the right hand side is the picture of the real amoebas of the two curves. The segment in the overlap of Γ and Λ is relatively twisted if the two branches of the two amoebas which are on the same side of the tropicalized curve Γ are the image under the coordinatewise logarithm map of pieces of the real algebraic curves which live in the same orthants of \mathbb{R}^2 .

which it came. In Figure 10, the branches of the amoebas of the family C along the edge e are labelled $\varepsilon_1^C, \varepsilon_2^C \in \{+, -\}^2$, and the branches of the amoebas of the family L along the edge e are labelled $\varepsilon_1^L, \varepsilon_2^L \in \{+, -\}^2$. The condition that

$$\frac{b_s}{b_{s'}} = \frac{a_q}{a_{q'}}$$

amounts to the equality of the sets

$$\{\varepsilon_1^C, \varepsilon_2^C\} = \{\varepsilon_1^L, \varepsilon_2^L\}.$$

In the real case, the edge e being relatively twisted with respect to C and L amounts to $\varepsilon_1^C = \varepsilon_1^L$ and $\varepsilon_2^C = \varepsilon_2^L$, see [22, Proposition 4.13].

In order for C and L to have a real tangency point tropicalizing to the segment, the edge e must be relatively twisted, see [22, Theorem 1.4]. Otherwise there would be two real points in the intersection. Notice that this appears to be opposite to the case when the tropical tangency component is an entire edge. In that case, in order to have a real tangency we required the edge to be non-twisted.

Proposition 3.11. *Let C be a curve defined over K such that $\text{Trop}(C)$ is a smooth tropicalized curve. Let Λ be a tropical line and suppose the segment T is a tropical tangency component of $\Lambda \cap \text{Trop}(C)$ strictly contained in an edge of Λ and of $\text{Trop}(C)$. Then a lift $L = V(\ell)$ of Λ which is tangent to C at a point tropicalizing to T has local equation ℓ_T defined over K if and only if the edge T is relatively twisted over k .*

Proof. The statement once again follows from Lemma 3.4. □

A condition similar to twisting can be given also for another type of tangency, as follows.

Proposition 3.12. *Let C be a curve defined over K such that $\text{Trop}(C)$ is a smooth tropicalized curve. Let Λ be a tropical line and suppose p is an isolated point of $\Lambda \cap \text{Trop}(C)$ which is a vertex p of Λ with intersection multiplicity 2. Then a lift L of Λ tangent to C at a point tropicalizing to p is defined over K if and only if $b_{r'}, b_{s'} \in k$ and*

$$\sqrt{-a_r a_s b_{r'} b_{s'}} \in k,$$

where $r, s \in \mathbb{Z}^2$ are the lattice endpoints of the edge dual to the edge E of $\text{Trop}(C)$ and $r', s' \in \mathbb{Z}^2$ are lattice endpoints dual to the unique edge of Λ which has intersection multiplicity 2 with E .

Proof. Let Q_p and ℓ_p denote the local equations for the curve C and a lift of the tropical line Λ at p , respectively. Then Q_p is the binomial $A_r x^{r_1} y^{r_2} + A_s x^{s_1} y^{s_2}$ and $\ell_p = \ell$ is a trinomial. We denote the coefficients of the linear form ℓ by $B_{q'}, B_{r'}, B_{s'}$.

If Λ and C have intersection multiplicity 2 at the vertex p and $r', s' \in \mathbb{Z}^2$ are lattice endpoints dual to the unique edge of Λ which has intersection multiplicity 2 with E , then the direction of E is determined. For example, if $r' = (0, 0)$ and $s' = (1, 0)$ then the direction of E must be $(2, 1)$ and hence $Q_p = A_r x^{r_1} y^{r_2} + A_s x^{r_1+1} y^{r_2-2}$ and set $\ell = B_{q'} y + B_{s'} x + B_{r'}$. Solving this system of two equations in the torus reduces to the degree two equation in y given by

$$A_r y^2 - \frac{A_s B_t}{B_{s'}} y - \frac{A_s B_{r'}}{B_{s'}} = 0.$$

In order to have a tangency tropicalizing to p we require the discriminant of this equation to be equal to zero, so that $A_r^2 B_{q'}^2 + 4A_r A_s B_{r'} B_{s'} = 0$. Passing to the initials this implies that $b_{q'} = \pm \sqrt{\frac{-4a_r a_s b_{r'} b_{s'}}{a_r^2}}$. Thus we have $b_{q'} \in k$ if and only if $\sqrt{-a_r a_s b_{r'} b_{s'}} \in k$ and the statement is proven. The other two cases for r' and s' are solved analogously. \square

3.3. Comparing lifting for different fields. In this subsection, we sum up the results about lifting of tropical bitangents, and use our study of lifting to compare lifting over different fields.

Proposition 3.13. *Let $C = V(Q)$ be a quartic defined over K with generic tropicalization $\text{Trop}(C)$ and Λ a tropical bitangent. The initials of the coefficients of the defining equation ℓ of a lift $L = V(\ell)$ of Λ , viewed in the algebraic closure of k , are Laurent terms in the initials a_{ij} of the coefficients of Q or square roots thereof.*

If Λ is not contained in a bitangent class of shape (II) in the classification in [6], the coefficients of these Laurent terms are of the form $\pm 2^k$ for some $k \in \mathbb{Z}$.

For two of the liftable tropical bitangents of a class of shape (II), the terms for the tangency points involve $\sqrt{2}$ and $\sqrt{3}$.

Proof. This is a generalization of Theorem 1.2 in [6]. As the proof of this theorem, the result follows from a case-by-case analysis involving the classification of shapes of bitangent classes from [6], see also Appendix A.1. Unlike in Theorem 1.2 [6], which focused on the case of lifting over \mathbb{R} , we have to pay attention to the coefficients of the solutions of our local lifting equations in order to decide liftability over K . \square

If we check all 4 local lifting computations for a tropical bitangent class, we observe that the obstruction for lifting to K is the same for all 4, i.e. the radicands which appear are all equal up to square (see e.g. Example 3.2). Thus we can conclude:

Theorem 3.14. *Given a generic tropicalization $\text{Trop}(C)$ of a quartic $C = V(Q)$ defined over K , and a bitangent class S of $\text{Trop}(C)$, then either all four lifts of S are defined over K or none of them are.*

As noted above, the obstructions for lifting are equal not only for the coefficients of the equations of the bitangent lines, but also for the tangency points. Taking the special behaviour of shape (II) into account (see Proposition 3.13), we can deduce:

Corollary 3.15. *Assume $\sqrt{2}, \sqrt{3}$ exist in K . Given a generic tropicalization $\text{Trop}(C)$ of a quartic $C = V(Q)$ defined over K , and let S be a bitangent class of $\text{Trop}(C)$ such that the equations of all four lifts are defined over K . Then the tangency points are also all defined over K .*

To compare lifting over different fields K_1 and K_2 with residue fields k_1 and k_2 respectively, we first suppose there exists an isomorphism $\phi : k_1^\times / (k_1^\times)^2 \rightarrow k_2^\times / (k_2^\times)^2$. For an element $a \in k^\times$, let \bar{a} denote its class in $k^\times / (k^\times)^2$. As a main example, consider the case when k is the real numbers or any finite field. Then

$$k^\times / (k^\times)^2 \cong \mathbb{Z}/2\mathbb{Z},$$

and an isomorphism $k^\times / (k^\times)^2 \rightarrow \mathbb{Z}/2\mathbb{Z}$ is given by the Legendre symbol. For a finite field k we let $\left(\frac{a}{k}\right)$ denote the Legendre symbol of $a \in k^*$. Concretely, we have

$$\bar{a} = \left(\frac{a}{k}\right) = \begin{cases} +1 & \text{if } x^2 = a \text{ has a solution in } k \\ -1 & \text{if } x^2 = a \text{ has no solution in } k \end{cases}$$

If we take $k = \mathbb{R}$ then the Legendre symbol is simply remembering whether $a \in k^\times$ is positive or negative. For two fields k_1, k_2 with $k_i^\times / (k_i^\times)^2 \rightarrow \mathbb{Z}/2\mathbb{Z}$, there is a unique isomorphism $\phi : k_1^\times / (k_1^\times)^2 \rightarrow k_2^\times / (k_2^\times)^2$.

Using Theorem 3.3 we can relate the lifts over different fields when $k_1^\times / (k_1^\times)^2$ and $k_2^\times / (k_2^\times)^2$ are isomorphic.

Theorem 3.16. *Let K_1 and K_2 be fields with residue fields k_1 and k_2 , respectively. Suppose there exists an isomorphism of groups $\phi : k_1^\times / (k_1^\times)^2 \rightarrow k_2^\times / (k_2^\times)^2$ such that $\phi(\overline{-1}) = \overline{-1}$.*

For $i = 1, 2$, let $C_i = V(Q_i)$ be a quartic curve defined over K_i such that $\text{Trop}(C_i)$ is generic. Let $Q_1 = \sum A_{ij}x^i y^j z^{4-i-j}$ and $Q_2 = \sum B_{ij}x^i y^j z^{4-i-j}$. We assume $\text{val}(B_{ij}) = \text{val}(A_{ij})$, in particular $\text{Trop}(C_1) = \text{Trop}(C_2)$, and $\phi(\overline{a_{ij}}) = \overline{b_{ij}}$ for all i, j , where a_{ij}, b_{ij} are the initials of A_{ij} and B_{ij} , respectively.

Then a tropical bitangent Λ to the tropicalized quartic $\text{Trop}(C_1)$ lifts to a bitangent of C_1 over K_1 if and only if Λ lifts to a bitangent of C_2 defined over K_2 .

The following corollary deals with the special case of comparing lifting over a finite field k with the real numbers. Here we can make use of the Legendre symbols. This is particularly useful, because lifting over the real numbers can be checked computationally using the POLYMAKE-extension of Geiger and Panizzut [10, 11, 12].

Corollary 3.17. *Let the residue field k be finite and of characteristic p with $p \equiv 3 \pmod{4}$ and order p^{2^l+1} .*

Let $C = V(Q)$ be a quartic curve defined over K with $Q = \sum A_{ij}x^i y^j$. Then a tropical bitangent Λ to the tropicalized curve $\text{Trop}(C)$ lifts to a bitangent over K if and only if L lifts to a bitangent of $C' = V(Q')$ defined over $\mathbb{R}\{\{t\}\}$, where $Q' = \sum B_{ij}x^i y^j$ and $B_{ij} \in \mathbb{R}\{\{t\}\}$ are such that $\text{val}(B_{ij}) = \text{val}(A_{ij})$ and

$$\left(\frac{a_{ij}}{k}\right) = \left(\frac{b_{ij}}{\mathbb{R}}\right).$$

4. THE GROTHENDIECK-WITT RING AND \mathbb{A}^1 -ENUMERATIVE INVARIANTS

Inspired by and building upon a broader program using \mathbb{A}^1 -homotopy theory to introduce arithmetic refinements to enumerative geometry via quadratic forms [13, 18, 19, 25], Larson and Vogt considered an arithmetic count of bitangents for smooth plane quartics with values in the Grothendieck-Witt ring of the ground field [21]. One of our main goals is to demonstrate, using the arithmetic count of bitangents as an illustrative test case, how tropical methods can be useful for computing such arithmetic counts.

4.1. The Grothendieck-Witt ring. We now recall the definition and basic properties of the Grothendieck-Witt rings in which these arithmetic counts take their values. See [20] for a beautiful and comprehensive expository treatment, which includes the proofs that we omit. In this section, K denotes a field of characteristic not equal to two.

A *quadratic space* is a finite-dimensional K -vector space V equipped with a symmetric bilinear form $q : V \times V \rightarrow K$. Two quadratic spaces (V, q) and (V', q') are isomorphic if there is an isomorphism of k -vector spaces $\phi : V \rightarrow V'$ such that $q(v, w) = q'(\phi(v), \phi(w))$ for all v, w in V .

Definition 4.1. For $a \in K^\times$, we write $\langle a \rangle$ for the 1-dimensional quadratic space (k, q) with $q(x, y) = axy$. The *hyperbolic plane* is $\mathbb{H} = \langle 1 \rangle \oplus \langle -1 \rangle$.

Lemma 4.2. For any $a \in K^\times$, the quadratic space $\langle a \rangle \oplus \langle -a \rangle$ is isomorphic to \mathbb{H} , as is the quadratic space K^2 with quadratic form $((x_1, x_2), (y_1, y_2)) \mapsto ax_1y_2 + ax_2y_1$.

Isomorphism classes of quadratic spaces naturally form a semiring with addition and multiplication given by orthogonal direct sum and tensor product, respectively. A theorem of Witt shows that this addition is cancellative, i.e., if V , W , and W' are quadratic spaces such that $V \oplus W$ is isomorphic to $V \oplus W'$ then W is isomorphic to W' . As a consequence, this semiring injects into its associated ring of formal differences.

Definition 4.3. The *Grothendieck-Witt ring* $\mathrm{GW}(K)$ is the ring of formal differences of isomorphism classes of quadratic spaces over K .

In other words, elements of $\mathrm{GW}(K)$ are formal differences $V - W$, where V and W are isomorphism classes of quadratic spaces.

Definition 4.4. The degree map $\mathrm{deg}: \mathrm{GW}(k) \rightarrow \mathbb{Z}$ takes a quadratic space V to its dimension as a K -vector space.

For instance, $\mathrm{deg}\langle a \rangle = 1$ and $\mathrm{deg}\mathbb{H} = 2$. Note that deg is a map of rings. All of the $\mathrm{GW}(K)$ -valued arithmetic counts that we consider specialize to the classical integer valued enumerative invariants when composed with the degree map.

Lemma 4.5. As an additive group, $\mathrm{GW}(K)$ is generated by $\{\langle a \rangle : a \in K^\times\}$, with relations generated by

- (1) $\langle a \rangle = \langle ab^2 \rangle$ for all a, b in K^\times , and
- (2) $\langle a \rangle + \langle b \rangle = \langle a + b \rangle + \langle ab(a + b) \rangle$, for all a, b in K^\times such that $a + b \neq 0$.

When no confusion seems possible, we write $\langle a \rangle$, \mathbb{H} , and so on, not only for a given quadratic space, but also for its class in $\mathrm{GW}(K)$.

We will frequently consider how Grothendieck-Witt rings behave with respect to finite field extensions. Suppose K'/K is a finite extension. Then any finite dimensional K' -vector space V is also a finite-dimensional as a K -vector space, and we write V_K to denote V , viewed as a K -vector space. If (V, q) is a quadratic space over K' , then $(V_K, \mathrm{Tr}_{K'|K} \circ q)$ is a quadratic space over K . One writes

$$\mathrm{Tr}_{K'|K}: \mathrm{GW}(K') \rightarrow \mathrm{GW}(K)$$

for the induced map of Grothendieck-Witt rings. Note that $\mathrm{Tr}_{K'|K}$ multiplies degrees by a factor of $[K' : K]$, since $\dim_K V_K = [K' : K] \dim_{K'} V$.

Example 4.6. Let K'/K be a field extension of degree 2, and let $a \in (K')^\times$. We now explain how to compute $\mathrm{Tr}_{K'|K}\langle a \rangle \in \mathrm{GW}(K)$. Suppose $K' = K(\beta)$, and $b = \beta^2$. Use $\{1, \beta\}$ as a K -basis for K' , and write

$$a = r + s\beta.$$

Then the bilinear form on $\mathrm{Tr}_{K'|K}\langle a \rangle$ may be expressed by the symmetric matrix

$$\begin{pmatrix} 2r & 2bs \\ 2bs & 2br \end{pmatrix}.$$

If $r = 0$, then $\mathrm{Tr}_{K'|K}\langle a \rangle \cong \mathbb{H}$, by Lemma 4.2. If $r \neq 0$, then we can diagonalize the symmetric matrix to get $\mathrm{Tr}_{K'|K}\langle a \rangle \cong \langle 2r \rangle \oplus \langle 2br(r^2 - bs^2) \rangle$.

A case of particular interest is when $K = \mathbb{R}$ and $b = -1$. Then $r^2 - bs^2$ is positive and hence is a square, so the above quadratic space is isomorphic to $\langle 2r \rangle \oplus \langle -2r \rangle \cong \mathbb{H}$. Thus, for any $a \in \mathbb{C}^\times$, we have $\mathrm{Tr}_{\mathbb{C}|\mathbb{R}}\langle a \rangle \cong \mathbb{H}$.

4.2. Grothendieck-Witt rings over valued fields. We now return to the case where K is a Henselian valued field with residue field k . We assume the characteristic of k is not 2 and fix a section of the valuation $\sigma: \mathrm{val}(K^\times) \rightarrow K^\times$. The initial of an element $a \in K^\times$, denoted $\mathrm{in}(a)$, is the image of $\sigma(-\mathrm{val}(a)) \cdot a$ in k^\times . We also make the simplifying assumption that the value group of K is 2-divisible, cf. Remark 1.6.

Theorem 4.7. *There is an isomorphism of Grothendieck-Witt rings*

$$g: \mathrm{GW}(K) \xrightarrow{\sim} \mathrm{GW}(k), \quad \langle A \rangle \mapsto \langle \mathrm{in}(A) \rangle.$$

Moreover, this isomorphism is independent of the section σ .

Proof. We first show that g is well-defined, i.e., that $\langle A \rangle \mapsto \langle \mathrm{in}(A) \rangle$ respects the relations (1) and (2) from Lemma 4.5. Let $a = \mathrm{in}(A)$ and $b = \mathrm{in}(B)$. For (1), we have $\langle \mathrm{in}(AB^2) \rangle = \langle ab^2 \rangle$. For (2), we consider a few subcases. First, suppose $\mathrm{val}(A) \neq \mathrm{val}(B)$. Without loss of generality we may assume $\mathrm{val}(A) > \mathrm{val}(B)$. Then $\mathrm{in}(\langle A \rangle + \langle B \rangle) = \langle a \rangle + \langle b \rangle$, while $\mathrm{in}\langle A + B \rangle = \langle a \rangle$ and $\mathrm{in}\langle AB(A + B) \rangle = \langle a^2b \rangle$. Using 4.5(1), we see that $g(\langle A + B \rangle + \langle AB(A + B) \rangle) = \langle a \rangle + \langle a^2b \rangle = \langle a \rangle + \langle b \rangle$, as required. If $\mathrm{val}(A) = \mathrm{val}(B)$ but $a + b \neq 0$, then (2) is clear. It remains to consider the case where $\mathrm{val}(A) = \mathrm{val}(B)$ and $a + b = 0$. Let $c = \mathrm{in}(A + B)$. Then $g(\langle A + B \rangle) = \langle c \rangle$ and $g(\langle AB(A + B) \rangle) = \langle abc \rangle = \langle -a^2c \rangle$, which is equal to $\langle -c \rangle$ in $\mathrm{GW}(k)$, again by 4.5(1). Finally, $g(\langle A \rangle + \langle B \rangle) = \langle a \rangle + \langle -a \rangle = \langle c \rangle + \langle -c \rangle$, by Lemma 4.2. This proves that g is well-defined.

It is evident that g respects addition and multiplication, so it is a ring map. We now construct its inverse. For each $a \in k^\times$, choose some $A \in R$ with residue a . We claim that $\langle a \rangle \mapsto \langle A \rangle$ gives a well-defined map $\mathrm{GW}(k) \rightarrow \mathrm{GW}(K)$. Indeed, if A and A' are two lifts of a , then A/A' has valuation 0 and initial 1. Since the characteristic of k is not 2 and K is Henselian, it follows that A/A' is a square and hence $\langle A \rangle \cong \langle A' \rangle$. Moreover, since σ is a section and $\mathrm{val}(K^\times)$ is 2-divisible, t^v has a square root for every $v \in \mathrm{val}(K^\times)$. Thus $\mathrm{GW}(K)$ is generated by quadratic spaces $\langle A \rangle$ such that $\mathrm{val}(A) = 0$, and hence the map so defined surjects onto $\mathrm{GW}(K)$ and gives an inverse to g , as required.

Finally, again using the fact that $\mathrm{val}(K^\times)$ is 2-divisible, the image of the section σ is contained in the multiplicative subgroup of squares. Thus, if we choose a

different section, the resulting initials of any $a \in K^\times$ will differ by a square and hence the isomorphism $\mathrm{GW}(K) \rightarrow \mathrm{GW}(k)$ is independent of this choice. \square

4.3. The Qtype of a bitangent to a quartic. A line L in \mathbb{P}_k^2 is a closed point of $(\mathbb{P}_k^2)^\vee$. Let K_L denote the field of definition of L . So K_L/K is a finite extension and L corresponds to a Galois orbit of geometric lines defined over the finite extension K_L/K . Suppose L is a bitangent to our plane quartic C . We assume furthermore that the intersection $L \cap C$ is disjoint from the line at infinity L_∞ given in homogeneous coordinates by $z = 0$. The *GW-multiplicity of the bitangent L* with respect to the fixed reference line L_∞ is an element of degree $[K_L : K]$ in $\mathrm{GW}(K)$ that is defined as follows.

Let $\mathbb{A}^2 = \mathbb{P}^2 \setminus L_\infty$. Let $Q(x, y)$ be a (non-homogeneous) quartic polynomial that vanishes on the affine plane quartic $C \cap \mathbb{A}^2$, and let ∂_L denote the derivation with respect to a linear form (defined over K_L) that vanishes on L . Note that both $Q(x, y)$ and ∂_L are defined only up to nonzero scalars (in K^\times and K_L^\times , respectively).

Definition 4.8. Let $P_1, P_2 \in C(\overline{K})$ be the points where L is tangent to C . Then the Qtype of L with respect to L_∞ is

$$(2) \quad \mathrm{Qtype}_{L_\infty}(L) := \langle \partial_L Q(P_1) \cdot \partial_L Q(P_2) \rangle \in \mathrm{GW}(K_L),$$

and the *GW-multiplicity of L* is


$$\mathrm{mult}_{\mathrm{GW}}(L) := \mathrm{Tr}_{K_L|K}(\mathrm{Qtype}_{L_\infty}(L)) \in \mathrm{GW}(K).$$

To see that $\mathrm{Qtype}_{L_\infty}(L)$ is well-defined, first suppose P_1 and P_2 are rational over K_L . Note that rescaling Q or ∂_L by a nonzero factor a multiplies $\partial_L Q(P_1) \cdot \partial_L Q(P_2)$ by a^2 . Then since $\langle a^2 b \rangle = \langle b \rangle$, the $\mathrm{Qtype}_{L_\infty}$ of L is well-defined.

Otherwise, if P_1 and P_2 are not rational over K_L then they are rational over some quadratic extension K_1/K_L . In this case $\partial_L Q(P_1)$ is conjugate to $\partial_L Q(P_2)$, and so $\partial_L Q(P_1) \cdot \partial_L Q(P_2)$ is in K_L . Once again, rescaling Q or L changes $\partial_L Q(P_1) \cdot \partial_L Q(P_2)$ by a square, and so $\mathrm{Qtype}_{L_\infty}(L)$ is well-defined in $\mathrm{GW}(K_L)$.

Example 4.9. [21] When $K = \mathbb{R}$, the $\mathrm{Qtype}_{L_\infty}$ has a natural topological interpretation, as explained by Larson and Vogt. Suppose a bitangent is defined over \mathbb{R} . If each of the bitangency points of L is defined over \mathbb{R} , then near these two points in the affine plane $\mathbb{A}^2(\mathbb{R})$, the real locus of C is either on the same side of the bitangent line, in which case the $\mathrm{Qtype}_{L_\infty}$ is $\langle 1 \rangle$, or on different sides, in which case the $\mathrm{Qtype}_{L_\infty}$ is $\langle -1 \rangle$. See Figure 11. When these points are not defined over \mathbb{R} , the $\mathrm{Qtype}_{L_\infty}$ is $\langle 1 \rangle$. If the bitangent line is not defined over \mathbb{R} then the $\mathrm{Qtype}_{L_\infty}$ of L is $\langle 1 \rangle \in \mathrm{GW}(\mathbb{C})$ and $\mathrm{mult}_{\mathrm{GW}}(L) = \langle 1 \rangle + \langle -1 \rangle = \mathbb{H}$ by Example 4.6.

Example 4.10. Suppose $K = \mathbb{R}$ (or $\mathbb{R}\{\{t\}\}$). If L is a bitangent line that is not rational over K , (i.e., if L is the point of $(\mathbb{P}_{\mathbb{R}}^2)^\vee$ corresponding to a pair of complex conjugate lines) then, by Example 4.6, we have $\mathrm{mult}_{\mathrm{GW}}(L) = \mathbb{H}$.


 FIGURE 11. Geometric characterization of $Q_{\text{type}_{L_\infty}}$ for the reals.

4.4. The GW-multiplicity of a tropical bitangent class. In the following, we study GW-multiplicities for the four liftable members of a tropical bitangent class.

We start, in the following Lemma, with picking a vector for the computation of $\partial_L Q(P)$ for a bitangent line L . It turns out that our choice is particularly useful for the tropical degeneration.

Lemma 4.11. *Let $L = V(y + M + Nx)$ be a bitangent to a quartic curve defined by Q , and let P be a point of tangency. Then $\partial_L Q(P) = (\frac{\partial}{\partial y} + \frac{1}{N} \cdot \frac{\partial}{\partial x})Q(P)$.*

Proof. Up to rescaling, the direction vector of the line L is $(-1, N)$. To compute ∂_L , we can take any vector which is not parallel to the direction vector and multiply it with the gradient. If $(\frac{1}{N}, 1)$ was parallel to $(-1, N)$, then $-\frac{1}{N} \cdot (-1, N) = (\frac{1}{N}, -1) = (\frac{1}{N}, 1)$ and thus $-1 = 1$, which is not the case as we are not in characteristic 2. \square

Remark 4.12. In the following Theorem, we make use of our choice of gradient and express GW-multiplicity in terms of derivatives of initial forms, which can be determined using the tropicalization of the quartic.

Using the classification in Appendix A.1, one can show that any such derivative of an initial form that can appear equals, after inserting initials of the tangency point, a Laurent monomial in the initials of the coefficients of the quartic times possibly a square root thereof.

Theorem 4.13. *Assume the bitangent L is given by the equation $M + NX + Y$ with $M, N \in K$, and let m, n denote their initials. If the tropicalization p of a tangency point P is contained in the interior of the horizontal ray of $\text{Trop}(L)$, then*

$$\text{in}(\partial_L Q(P)) = 2 \frac{\partial}{\partial y} \text{in}_p(Q)(\text{in}(P)).$$

If it is contained in the interior of the vertical ray of $\text{Trop}(L)$, then

$$\text{in}(\partial_L Q(P)) = 2 \frac{1}{n} \frac{\partial}{\partial x} \text{in}_p(Q)(\text{in}(P)).$$

If it is contained in the interior of the diagonal ray or at the vertex, then

$$\text{in}(\partial_L Q(P)) = \left(\frac{\partial}{\partial y} + \frac{1}{n} \cdot \frac{\partial}{\partial x}\right) \text{in}_p(Q)(\text{in}(P)).$$

In particular, the GW-multiplicity $\langle \partial_L Q(P_1) \cdot \partial_L Q(P_2) \rangle$ can be expressed in terms of derivatives of initial forms.

Proof. Without restriction, we can assume $p = (0, 0)$ and Q contains terms whose coefficients have valuation 0 or higher. The terms of valuation 0 are then precisely the terms contributing to $\text{in}_p(Q)$.

By Lemma 4.11, $\partial_L Q(P) = \left(\frac{\partial}{\partial y} + \frac{1}{N} \cdot \frac{\partial}{\partial x}\right) Q(P)$. The initial equals

$$(3) \quad \text{in}(\partial_L Q(P)) = \text{in}\left(\left(\frac{\partial}{\partial y} + \frac{1}{N} \cdot \frac{\partial}{\partial x}\right) Q(P)\right) = \text{in}\left(\frac{\partial}{\partial y} Q(P) + \frac{1}{N} \cdot \frac{\partial}{\partial x} Q(P)\right).$$

If the two summands above are of the same valuation, then cancellation could happen, which would be an obstruction to expressing $\text{in}(\partial_L Q(P))$ in terms of derivatives of initial forms.

Assume cancellation happens, then

$$\text{in}\left(\frac{\partial}{\partial y} Q(P)\right) = -\frac{1}{n} \text{in}\left(\frac{\partial}{\partial x} Q(P)\right).$$

But the Wronskian $W = N \cdot \frac{\partial}{\partial y} Q(P) - \frac{\partial}{\partial x} Q(P) = 0$ vanishes, and thus also its initial vanishes, so $n \cdot \text{in}\left(\frac{\partial}{\partial y} Q(P)\right) = \text{in}\left(\frac{\partial}{\partial x} Q(P)\right)$. We substitute $n \cdot \text{in}\left(\frac{\partial}{\partial y} Q(P)\right)$ for $\text{in}\left(\frac{\partial}{\partial x} Q(P)\right)$ in the above equation, obtaining $\text{in}\left(\frac{\partial}{\partial y} Q(P)\right) = -\text{in}\left(\frac{\partial}{\partial y} Q(P)\right)$ from which we deduce $1 = -1$ which is a contradiction as we are not in characteristic 2. Thus no cancellation happens.

There are three cases to distinguish:

- Assume p is on the horizontal ray. Then the vertex of $\text{Trop}(L)$ is at $(a, 0)$ for some $a > 0$. The valuation of N is a , the valuation of M is zero.

Consider the Wronskian $W = N \cdot \frac{\partial}{\partial y} Q(P) - \frac{\partial}{\partial x} Q(P) = 0$. Consequently, $\text{in}(N \cdot \frac{\partial}{\partial y} Q(P) - \frac{\partial}{\partial x} Q(P)) = 0$. But the summand $N \cdot \frac{\partial}{\partial y} Q(P)$ has terms of valuation a or higher, and the summand $\frac{\partial}{\partial x} Q(P)$ has terms of valuation 0 or higher. Cancellation can thus only appear if $\frac{\partial}{\partial x} Q(P)$ vanishes up to valuation a .

Thus both summands in the above equation (3) for $\text{in}(\partial_L Q(P))$ have expected valuation 0.

From the computations for GW-multiplicities we perform in Appendix A.2 and A.3 (see Remark 4.12), we conclude that $\text{in}\left(\frac{\partial}{\partial y} Q(P)\right) = \frac{\partial}{\partial y}(\text{in}_p(Q)(\text{in}(P))) \neq 0$ and thus this summand is of the expected valuation. The equality $\text{in}\left(\frac{\partial}{\partial y} Q(P)\right) = \frac{\partial}{\partial y}(\text{in}_p(Q)(\text{in}(P)))$ holds since for a polynomial $F(x, y)$ and a point $P = (P_x, P_y)$ that we insert, we have $\text{in}(F(P_x, P_y)) = \text{in}_{(p_x, p_y)}(F)(\text{in}(P))$, where (p_x, p_y) contains the valuations,

if no cancellation happens in the initial form when inserting $\text{in}(P)$. Furthermore we have $\text{in}_p(\frac{\partial}{\partial y}Q) = \frac{\partial}{\partial y}(\text{in}_p(Q))$. This is true since the p -weight of a term of Q equals the $p + p_y$ -weight of a term of $\frac{\partial}{\partial y}Q$.

Thus $\text{val}(N \cdot \frac{\partial}{\partial y}Q(P)) = a$ and since there is cancellation in the Wronskian, also $\text{val}(\frac{\partial}{\partial x}Q(P)) = a$. Furthermore, since there is cancellation in W , $n \text{in}(\frac{\partial}{\partial y}Q(P)) = \text{in}(\frac{\partial}{\partial x}Q(P))$.

Inserting this into the above equation (3), we obtain

$$\begin{aligned} \text{in}(\partial_L Q(P)) &= \text{in}\left(\frac{\partial}{\partial y}Q(P)\right) + \text{in}\left(\frac{1}{N} \cdot \frac{\partial}{\partial x}Q(P)\right) \\ &= \text{in}\left(\frac{\partial}{\partial y}Q(P)\right) + \frac{1}{n} \text{in}\left(\frac{\partial}{\partial x}Q(P)\right) \\ &= \text{in}\left(\frac{\partial}{\partial y}Q(P)\right) + \frac{1}{n} \cdot n \text{in}\left(\frac{\partial}{\partial y}Q(P)\right) \\ &= \text{in}\left(\frac{\partial}{\partial y}Q(P)\right) + \text{in}\left(\frac{\partial}{\partial y}Q(P)\right) \\ &= 2 \text{in}\left(\frac{\partial}{\partial y}Q(P)\right) = 2 \frac{\partial}{\partial y}(\text{in}_p(Q))(\text{in}(P)). \end{aligned}$$

- Assume p is on the vertical ray. Then the vertex of $\text{Trop}(L)$ is at $(0, a)$ for some $a > 0$. The valuation of N and M is $-a$.

Consider the Wronskian $W = N \cdot \frac{\partial}{\partial y}Q(P) - \frac{\partial}{\partial x}Q(P) = 0$. Consequently, $\text{in}(N \cdot \frac{\partial}{\partial y}Q(P) - \frac{\partial}{\partial x}Q(P)) = 0$. But the summand $N \cdot \frac{\partial}{\partial y}Q(P)$ has terms of valuation $-a$ or higher, and the summand $\frac{\partial}{\partial x}Q(P)$ has terms of valuation 0 or higher. Cancellation can thus only appear if $\frac{\partial}{\partial y}Q(P)$ vanishes up to valuation a .

Thus both summands in the above equation for $\text{in}(\partial_L Q(P))$ (3) have expected valuation 0.

From the computations for GW-multiplicities we perform in Appendix A.2 and A.3 (see Remark 4.12), we conclude that $\text{in}(\frac{\partial}{\partial x}Q(P)) = \frac{\partial}{\partial x}(\text{in}_p(Q))(\text{in}(P)) \neq 0$ and thus this summand is of the expected valuation. Thus $\text{val}(\frac{\partial}{\partial x}Q(P)) = 0$ and since there is cancellation in the Wronskian, also $\text{val}(N \cdot \frac{\partial}{\partial y}Q(P)) = 0$. Furthermore, since there is cancellation in W , $n \text{in}(\frac{\partial}{\partial y}Q(P)) = \text{in}(\frac{\partial}{\partial x}Q(P))$.

Inserting this into the above equation for $\text{in}(\partial_L Q(P))$, we obtain

$$\begin{aligned} \text{in}(\partial_L Q(P)) &= \text{in}\left(\frac{\partial}{\partial y} Q(P)\right) + \text{in}\left(\frac{1}{N} \cdot \frac{\partial}{\partial x} Q(P)\right) \\ &= \text{in}\left(\frac{\partial}{\partial y} Q(P)\right) + \frac{1}{n} \text{in}\left(\frac{\partial}{\partial x} Q(P)\right) \\ &= \frac{1}{n} \text{in}\left(\frac{\partial}{\partial x} Q(P)\right) + \frac{1}{n} \cdot n \text{in}\left(\frac{\partial}{\partial y} Q(P)\right) \\ &= 2 \frac{1}{n} \text{in}\left(\frac{\partial}{\partial x} Q(P)\right) = 2 \frac{1}{n} \frac{\partial}{\partial x} (\text{in}_p(Q)) (\text{in}(P)). \end{aligned}$$

- Assume p is on the vertical ray or the vertex. Then the vertex of $\text{Trop}(L)$ is at $(-a, -a)$ for some $a \geq 0$. The valuation of M is a , the valuation of N is zero. Then both summands in equation (3) are of valuation zero and we can directly express

$$\begin{aligned} \text{in}(\partial_L Q(P)) &= \text{in}\left(\frac{\partial}{\partial y} Q(P)\right) + \text{in}\left(\frac{1}{N} \cdot \frac{\partial}{\partial x} Q(P)\right) \\ &= \text{in}\left(\frac{\partial}{\partial y} Q(P)\right) + \frac{1}{n} \text{in}\left(\frac{\partial}{\partial x} Q(P)\right). \\ &= \frac{\partial}{\partial y} (\text{in}_p(Q)) (\text{in}(P)) + \frac{1}{n} \frac{\partial}{\partial x} (\text{in}_p(Q)) (\text{in}(P)). \end{aligned}$$

For the statement in particular, note that by Theorem 4.7, we have $\langle \partial_L Q(P_1) \cdot \partial_L Q(P_2) \rangle = \langle \text{in}(\partial_L Q(P_1) \cdot \partial_L Q(P_2)) \rangle$. The initial of the product $\text{in}(\partial_L Q(P_1) \cdot \partial_L Q(P_2))$ equals the product of initials $\text{in}(\partial_L Q(P_1)) \cdot \text{in}(\partial_L Q(P_2))$. The factors can be expressed in terms of derivatives of initial forms as described above. \square

Remark 4.14. By Example 4.10, a tropical bitangent with lifting multiplicity 2 which does not lift to K contributes \mathbb{H} to the arithmetic count of bitangents: Such a lift is defined over a field extension of degree 2. The element whose class in $\text{GW}(K)$ we take equals a Laurent term in the initials of Q times a square root of such a term by Remark 4.12 (see also Appendix A.2 and A.3). Thus, in the notation of Example 4.10, $r = 0$ and we get a contribution of \mathbb{H} .

In the following example, we consider tropical bitangents of lifting multiplicity four whose lifts live in a field extension.

Example 4.15. There are tropical bitangent classes for which we need to add two roots to the ground field to lift to 4 bitangent lines. These are precisely the zero-dimensional classes (A), (B) and (C) in the classification in [6]. Assume these two roots are α and β and $K \subset K(\alpha, \beta)$ is of degree 4.

Let $a \in K^\times$. We now compute $\text{Tr}_{K(\alpha, \beta)|L} \langle a \rangle \in \text{GW}(K)$. A basis for $K(\alpha, \beta)$ is given by $1, \alpha, \beta, \alpha \cdot \beta$. We write a in this basis:

$$a = c_1 \cdot 1 + c_2 \cdot \alpha + c_3 \cdot \beta + c_4 \cdot \alpha \cdot \beta.$$

We study the bilinear map

$$\tilde{q} : K(\alpha, \beta) \times K(\alpha, \beta) \rightarrow K(\alpha, \beta) \rightarrow K,$$

where the first arrow is the map $(x, y) \mapsto axy$ and the second is the trace. Let us first compute the trace for an arbitrary element $b = b_1 \cdot 1 + b_2 \cdot \alpha + b_3 \cdot \beta + b_4 \cdot \alpha \cdot \beta$. The matrix we obtain is

$$\begin{pmatrix} b_1 & b_2\alpha^2 & b_3\beta^2 & b_4\alpha^2\beta^2 \\ b_2 & b_1 & b_4\beta^2 & b_3\beta^2 \\ b_3 & b_4\alpha^2 & b_1 & b_2\alpha^2 \\ b_4 & b_3 & b_2 & b_1 \end{pmatrix}.$$

The trace is thus $4b_1$. Now we can insert basis vectors in the above bilinear map to obtain the symmetric matrix

$$\begin{pmatrix} 4c_1 & 4c_2\alpha^2 & 4c_3\beta^2 & 4c_4\alpha^2\beta^2 \\ 4c_2\alpha^2 & 4c_1\alpha^2 & 4c_4\alpha^2\beta^2 & 4c_3\alpha^2\beta^2 \\ 4c_3\beta^2 & 4c_4\alpha^2\beta^2 & 4c_1\beta^2 & 4c_2\alpha^2\beta^2 \\ 4c_4\alpha^2\beta^2 & 4c_3\alpha^2\beta^2 & 4c_2\alpha^2\beta^2 & 4c_1\alpha^2\beta^2 \end{pmatrix}.$$

As we can see in our computations for lifts of tropical bitangents and their Qtype_{L_∞} (see Remark 4.12 and Appendix A.1), the bilinear map we obtain for a lift of a tropical bitangent class of type (A), (B) or (C) is of the form $c_2\alpha$ or $c_4\alpha\beta$. Thus, we can insert $c_1 = c_3 = c_4 = 0$ or $c_1 = c_2 = c_3 = 0$ in the above matrix. In both cases, we obtain a decomposition into two hyperbolic planes.

This computation shows that the GW-multiplicity of a tropical bitangent class whose 4 lifts live in a field extension of degree 4 which we obtain by adjoining two roots is $2\mathbb{H}$.

Remark 4.16 (Computation of GW-multiplicities). Theorem 4.13 together with the theory of tropical bitangents allows to compute GW-multiplicities of quartics as follows:

- Compute the tropicalization $\text{Trop}(C)$, e.g. using the software system POLYMAKE [10] or the library tropical.lib in the Computeralgebrasystem SINGULAR [15, 7].
- Using the POLYMAKE-extension on tropical bitangents of quartics by Geiger-Panizzut [11], compute all its tropical bitangent classes.
- For each liftable tropical bitangent, use Theorem 4.13 to compute its Qtype_{L_∞} .
- For each tropical bitangent class which does not lift, deduce from Remark 4.14 and 4.15 that the GW-multiplicity of the four bitangents equals $2\mathbb{H}$.

In the following, we prepare statements which will be used for the further study of the Qtype_{L_∞} of tropical bitangent lines which lift over K .

Lemma 4.17. *Assume the initial form of Q at tropical tangency component p_1 equals a monomial m times a form q , and ∂_L is as above. Let P_1 be a lift of p_1 . Then $\partial_L(\text{in}_{p_1}(Q))(P_1) = m \cdot \partial_L(q)(P_1)$.*

Proof. This holds true since the second summand we obtain from the derivative of the product disappears since $q(P_1) = 0$. \square

Lemma 4.18. *Let Q be a quartic polynomial over K and assume $\text{Trop}(V(Q))$ is smooth and generic. Assume there is a tropical tangency component p in the interior of an untwisted horizontal bounded edge E of $\text{Trop}(V(Q))$. Then the lifts of the tropical bitangents at p come in pairs L_1, L_2 such that the tangency points $P_i \in L_i \cap V(Q)$ which tropicalize to p satisfy*

$$\text{in}(\partial_{L_1}(Q(P_1))) = -\text{in}(\partial_{L_2}(Q(P_2))).$$

Proof. By Theorem 4.13, $\text{in}(\partial_{L_i}(Q(P_i))) = 2\frac{\partial}{\partial y}\text{in}_p(Q)(\text{in}(P_i))$, as the tropical tangency component p is on the horizontal ray. The initial form is $\text{in}_p(Q) = a_{1k}xy^k + a_{1k+1}xy^{k+1}$. By Lemma 4.17, after inserting the initials of the solutions (x_0, y_0) of the tangency point we have

$$\text{in}(\partial_{L_i}(Q(P_i))) = 2xy^k \frac{\partial}{\partial y}(a_{1k} + a_{1k+1}y)(x_0, y_0) = 2a_{1k+1}x_0y_0^k.$$

Plugging in the solutions for the tangency point P from Lemma 3.4 we obtain the result. \square

The following is the analogue of Lemma 4.18 where the untwisted horizontal bounded edge E is replaced by a diagonal bounded edge. As we broke symmetry by declaring $\{z = 0\}$ to be our infinite line, we cannot expect a similar statement. It is interesting to note that while in Lemma 4.18, the contributions for the two different lifts were negatives of each other (equal up to sign), here we obtain the same contribution.

Lemma 4.19. *Let Q be a quartic polynomial over K and assume $\text{Trop}(V(Q))$ is smooth and generic.*

Assume there is a tropical tangency component p in the interior of an untwisted diagonal bounded edge E of $\text{Trop}(V(Q))$. As in Lemma 4.18, the lifts of the tropical bitangents at p come in pairs L_1, L_2 but now the tangency points $P_i \in L_i \cap V(Q)$ which tropicalize to p satisfy

$$\text{in}(\partial_{L_1}(Q(P_1))) = \text{in}(\partial_{L_2}(Q(P_2))).$$

Proof. Using Theorem 4.13 we have to derive with $\frac{\partial}{\partial y} + \frac{1}{n} \cdot \frac{\partial}{\partial x}$, and by Lemma 4.17, after inserting the initials of the solutions (x_0, y_0) of the tangency point we have




FIGURE 12. Possible duals of bounded edges intersecting a horizontal ray with multiplicity 2.

$$\partial_L(\text{in}_p(Q))(x_0, y_0) = x^k y^{2-k} \frac{\partial}{\partial y} + \frac{1}{n} \cdot \frac{\partial}{\partial x} (a_{k3-k} y + a_{k+12-k} x) = 2a_{k3-k} x_0^k y_0^{2-k},$$

as $n = \frac{a_{k+12-k}}{a_{k3-k}}$ by Lemma 3.4. Also, $\frac{y_0}{x_0} = -n$, so we obtain

$$2a_{k3-k} \left(-\frac{a_{k+12-k}}{a_{k3-k}} \right)^{2-k} \cdot x_0^{2k-2}.$$

Up to squares, this equals

$$(-1)^k \cdot 2 \cdot a_{k3-k}^{k-1} a_{k+12-k}^k,$$

no matter whether we insert the positive or the negative root which is the solution for x_0 . \square

Lemma 4.20. *Assume the horizontal or vertical ray of a tropical bitangent line can move along a bounded edge maintaining tangency, and the tropicalization of the tangency points p_1 and p_2 of the liftable members L_1, L_2 of the corresponding bitangent class are the two end vertices. Then*

$$\text{in}(\partial_{L_1}(Q(P_1))) = -\text{in}(\partial_{L_2}(Q(P_2))) \text{ up to squares.}$$

Proof. By symmetry, we can without restriction assume that the ray is horizontal. The only direction vectors of dual edges that fit into a the Newton polygon of a quartic, i.e. the triangle with vertices $(0, 0)$, $(0, 4)$ and $(4, 0)$, and intersect with multiplicity 2 a horizontal edge are $\binom{2}{1}$, $\binom{2}{-1}$ and $\binom{2}{3}$. Because of smoothness, the adjacent vertices correspond to triangles as depicted in Figure 12.

We perform the computation for the left picture in Figure 12, for the others, it is completely analogous. Here, $\text{in}_{p_1}(Q) = x^i y^j \cdot (a + bxy + cx^2y)$ and $\text{in}_{p_2}(Q) = x^i y^j \cdot (a + dx + cx^2y)$. As the horizontal ray cannot meet any other point of the tropicalized quartic, we can conclude that $i = 0$. Using local lifting equations, we obtain $\text{in}(P_1) = (-\frac{b}{2c}, \frac{4ac}{b^2})$ and $\text{in}(P_2) = (-\frac{2a}{d}, \frac{d^2}{4ac})$. Using Theorem 4.13 we have to derive with $\frac{\partial}{\partial y}$, and by Lemma 4.17, we obtain

$$\begin{aligned} \text{in}(\partial_{L_1}(Q(P_1))) &= \left(\frac{4ac}{b^2}\right)^j \cdot \left(-b\frac{b}{2c} + c\frac{b^2}{4c^2}\right) = \left(\frac{4ac}{b^2}\right)^j \cdot \left(-\frac{b^2}{4c}\right), \\ -\text{in}(\partial_{L_2}(Q(P_2))) &= \left(\frac{d^2}{4ac}\right)^j \cdot \left(c\frac{4a^2}{d^2}\right). \end{aligned}$$

Up to squares, this equals $\pm a^j c^{j+1}$. \square

Remark 4.21. An analogous statement for intersections with the diagonal ray does not hold. (See e.g. Appendix A.3, shape (Ea), where one tropical tangency component contributes a_{21} and the other $-a_{20}a_{31}a_{30}$).

Lemma 4.22. *Assume a liftable tropical bitangent meets a tropicalized quartic at a point p in the interior of an edge of direction $(1, -1)$ dual to $a_{00} + a_{11}xy$ with its vertex. Assume the second tropical tangency component is on the diagonal ray. The lifts come in pairs L_1, L_2 with tangency points P_1 and P_2 tropicalizing to p , and we have*

$$\text{in}(\partial_{L_1}(Q(P_1))) = -\text{in}(\partial_{L_2}(Q(P_2))).$$

Proof. By a local lifting computation, we obtain for the initials of the two lifts P_1 and P_2 the coordinates $(x_0, y_0) = \left(\mp \frac{1}{2n} \sqrt{-\frac{4a_{00}n}{a_{11}}}, \mp \frac{1}{2} \sqrt{-\frac{4a_{00}n}{a_{11}}} \right)$, where n as usual denotes the coefficient of x in the normalized equation defining the bitangent lift. Its initial is imposed by the second tropical tangency component which is on the diagonal ray.

Using Theorem 4.13 we have to derive with $\frac{\partial}{\partial y} + \frac{1}{n} \frac{\partial}{\partial x}$, and by Lemma 4.17, we obtain

$$a_{11} \cdot x_0 + \frac{1}{n} a_{11} y_0 = 2a_{11} x_0.$$

Inserting the two solutions for x_0 , which are negative of each other, the statement follows. \square

Theorem 4.23. *A bitangent class S of a generic tropicalized quartic $\text{Trop}(C)$ contributes either $2\mathbb{H}$ to the \mathbb{A}^1 -enumerative count of bitangents to C , or a sum of four monomials in the initials of the coefficients of the defining polynomial of C .*

In particular, the total \mathbb{A}^1 -enumerative count of bitangents to C is determined by its tropicalization $\text{Trop}(C)$ and the initials of its coefficients, up to square.

The exceptional cases (which do not give $2\mathbb{H}$) and their GW-multiplicities are listed in the Appendix A.2, details are spelled out in Theorem A.2.

Proof. The strategy of the proof is to use Lemmas 4.18, 4.20 and 4.22 to argue that the four lifts come in pairs contributing $\langle \pm a \rangle$ for some a , which sums up to $2\mathbb{H}$. This works if the lifts exist in the field, else we use Examples 4.6 and 4.15 to show that nevertheless, we obtain $2\mathbb{H}$ as total contribution. The proof is a case-by-case analysis relying on the classification of bitangent shapes in Appendix A.1. Here, we show one case and refer to the detailed version, Theorem A.2 in Appendix A.2, for the remaining ones.

We focus on a bitangent shape of type (D) as in Figure 6 of [6]. The bitangent shape is a bounded segment which partly overlaps with an edge of the tropicalized




FIGURE 13. Liftable tropical bitangents and motifs for shapes (Da), (Db) and (Dc).

quartic. The liftable points are the two end vertices of the segment, each such point has lifting multiplicity 2. Thus there are two tropical bitangent lines in the shape which each have two algebraic lifts.

The $\mathbb{S}_3/\mathbb{S}_2$ -orbit of this shape consists of three cases (Da), (Db) and (Dc). In (Da), the bounded segment is horizontal, in (Db) it is diagonal and in (Dc) vertical. Figure 13 shows the dual motifs and local pictures of the tropicalized quartic and the tropical tangency components. The pictures contain the two liftable tropical bitangent lines in blue. In each case, the bitangent class is the segment connecting the two vertices of the two blue tropical lines in each picture. Observe that the segment partially overlaps with an edge of the tropicalized quartic, as mentioned before.

For (Da) and (Dc), each of the two liftable tropical bitangents has a tropical tangency component which is an overlap of a horizontal resp. vertical edge. By Lemma 4.18, we can pair up the two lifts for each such tropical bitangent such that their Qtypes are negative of each other. Altogether, we obtain $2\mathbb{H}$.

For (Db) this argument does not work, but a computation shows that the contribution of the other tropical tangency component, which is not in the interior of the diagonal edge, equals a_{21} for one resp. $-a_{21}$ for the second, up to squares: for the upper tropical tangency component p_1 , we have $\text{in}_{p_1}(Q) = a_{02}y^2 + a_{12}xy^2 + a_{21}x^2y$. Solving for the initials of the coordinates of the tangency point and the coefficient M of the line equation $y + Mx + N$, we obtain $m = \text{in}(M) = -4 \frac{a_{02}a_{21}}{a_{12}^2}$,

$x = -2\frac{a_{02}}{a_{12}}$ and $y = 4\frac{a_{02}a_{21}}{a_{12}^2}$. Using Lemma 4.17 and Theorem 4.13, we obtain

$$2\frac{\partial}{\partial y} \operatorname{in}_{p_1}(Q(\operatorname{in}(P_1))) = 8\frac{a_{02}a_{21}}{a_{12}^2} \cdot (a_{02} + a_{12} \cdot (-2\frac{a_{02}}{a_{12}})) = -8\frac{a_{02}^2}{a_{12}^2} \cdot a_{21}.$$

For the lower tropical tangency component p_2 , solving for the initials of the coordinates of the tangency point and the ratio $\frac{M}{N}$ of the coefficients of the line equation, we obtain $\frac{m}{n} = \operatorname{in} \frac{M}{N} = -\frac{a_{11}^2}{4a_{02}a_{30}}$, $x = \frac{a_{11}^2}{4a_{02}a_{30}}$ and $y = -\frac{a_{11}^3}{8a_{02}^2a_{30}}$. We have $\operatorname{in}_{p_2}(Q) = a_{02}y^2 + a_{11}xy + a_{30}x^3$ and

$$\begin{aligned} 2\frac{1}{n}\frac{\partial}{\partial x} \operatorname{in}_{p_2}(Q(\operatorname{in}(P_2))) &= 2\frac{1}{m} \cdot \frac{m}{n} \frac{\partial}{\partial x} \operatorname{in}_{p_2}(Q) \\ &= 2\frac{1}{m} \cdot \frac{m}{n} \cdot \left(a_{11} \cdot \left(-\frac{a_{11}^3}{8a_{02}^2a_{30}} \right) + 3a_{30} \cdot \left(\frac{a_{11}^2}{4a_{02}a_{30}} \right)^2 \right) \\ &= 2\frac{1}{m} \cdot \frac{m}{n} \cdot \left(\frac{a_{11}^4}{16a_{02}^2a_{30}} \right) \\ &= 2\left(-\frac{a_{12}^2}{4a_{02}a_{21}} \right) \left(-\frac{a_{11}^2}{4a_{02}a_{30}} \right) \left(\frac{a_{11}^4}{16a_{02}^2a_{30}} \right) \\ &= 2\frac{1}{16^2} \frac{a_{12}^2 a_{11}^6}{a_{21}^4 a_{02}^2 a_{30}^2} a_{21}. \end{aligned}$$

Using Lemma 4.19, we can now pair up a lift of one with a lift of the other tropical bitangent and conclude that the total contribution is again $2\mathbb{H}$. \square

4.5. Comparing the $\operatorname{Qtype}_{L_\infty}$ for different fields. Building on our comparison of lifting for different fields in Subsection 3.3, we can also discuss the comparison of the $\operatorname{Qtype}_{L_\infty}$ for different fields, as a consequence of Theorem 4.23.

Theorem 4.24. *Let K_1 and K_2 be fields with residue fields k_1 and k_2 , respectively. Suppose there exists an isomorphism of groups $\phi : k_1^\times / (k_1^\times)^2 \rightarrow k_2^\times / (k_2^\times)^2$ such that $\phi(\overline{-1}) = \overline{-1}$ and $\phi(\overline{2}) = \overline{2}$.*

For $i = 1, 2$, let $C_i = V(Q_i)$ be a quartic curve defined over K_i such that $\operatorname{Trop}(C_i)$ is generic. Let $Q_1 = \sum A_{ij}x^i y^j z^{4-i-j}$ and $Q_2 = \sum B_{ij}x^i y^j z^{4-i-j}$. We assume $\operatorname{val}(B_{ij}) = \operatorname{val}(A_{ij})$, in particular $\operatorname{Trop}(C_1) = \operatorname{Trop}(C_2)$, and $\phi(\overline{a_{ij}}) = \overline{b_{ij}}$ for all i, j , where a_{ij}, b_{ij} are the initials of A_{ij} and B_{ij} , respectively.

Suppose all lifts of Λ are defined over K_1 and K_2 then

$$\sum_{L \text{ lift of } \Lambda} \phi(\operatorname{QType}(L)) = \sum_{L' \text{ lift of } \Lambda} \operatorname{QType}(L')$$

where the lifts L are over K_1 and the lifts L' are over K_2 and if $\operatorname{QType}(L) = \langle a \rangle$ then $\phi(\operatorname{QType}(L)) := \langle \phi(a) \rangle$.

Proof. If k has order p^{2l+1} and $p \equiv 7 \pmod{8}$, then we have that -1 is not a square in k , 2 is a square in k , and $k^\times / (k^\times)^2 = k^\times / (k^\times)^4$. This is also the case

when $k = \mathbb{R}$. The formulas for the Qtype_{L_∞} of tropical bitangents lifting over k are expressed in terms of a Laurent monomials in the a_{ij} 's or the b_{ij} 's with coefficients of -1 , 2 , or squares in the residue fields. Moreover, the Qtypes are defined modulo squares so the statement follows from the assumptions on the a_{ij} 's and b_{ij} 's. \square

We present the special case comparing QTypes over finite fields and the real numbers.

Corollary 4.25. *Let K be a field with finite residue field k of size p^{2l+1} for a prime p satisfying $p \equiv 7 \pmod{8}$. Let $\phi : k^\times / (k^\times)^2 \rightarrow \mathbb{R}^\times / (\mathbb{R}^\times)^2$ denote the unique isomorphism between these two groups.*

Let $C = V(Q)$ be a quartic curve defined over K with $Q = \sum A_{ij}x^i y^j$, and $C' = V(Q')$ be defined over $\mathbb{R}\{\{t\}\}$, where $Q' = \sum B_{ij}x^i y^j$ and $B_{ij} \in \mathbb{R}\{\{t\}\}$. Assume that $\text{val}(B_{ij}) = \text{val}(A_{ij})$ and

$$\left(\frac{a_{ij}}{k}\right) = \left(\frac{b_{ij}}{\mathbb{R}}\right).$$

Suppose all lifts of Λ are defined over K and $\mathbb{R}\{\{t\}\}$, then

$$\sum_{L \text{ lift of } \Lambda} \phi(\text{QType}(L)) = \sum_{L' \text{ lift of } \Lambda} \text{QType}(L')$$

where the lifts L are over K_1 , the lifts L' are over $\mathbb{R}\{\{t\}\}$, and if $\text{QType}(L) = \langle a \rangle$ then $\phi(\text{QType}(L)) := \langle \phi(a) \rangle$.

Proof. The first statement follows immediately from Theorem 4.24, since in a field k of characteristic p with $p \equiv 7 \pmod{8}$ and with order p^{2l+1} we have $\left(\frac{-1}{k}\right) = \left(\frac{-1}{\mathbb{R}}\right)$ and $\left(\frac{2}{k}\right) = \left(\frac{2}{\mathbb{R}}\right)$. \square

5. THE REAL GW-MULTIPLICITY IN TROPICAL GEOMETRY

Let $K = \mathbb{R}\{\{t\}\}$ the field of Puiseux series with real coefficients. By the Tarski principle [16], $\mathbb{R}\{\{t\}\}$ is equivalent to the reals, so we can count real bitangents by lifting tropical bitangents to $\mathbb{R}\{\{t\}\}$. This principle has been applied to other problems in tropical geometry, see e.g. [1].

Recall from Example 4.9, that the Qtype_{L_∞} of a real bitangent is $\langle 1 \rangle$ or $\langle -1 \rangle$, and the two cases can be characterized geometrically (see [21]): Assume first the tangency points are real. In the affine chart of \mathbb{P}^2 we obtain by taking out the line at infinity, the quartic $V(Q)$ can pass on the same side of the bitangent line for the two tangency points, or on opposite sides (see Figure 11). If the tangency points are not real, the Qtype_{L_∞} is $\langle 1 \rangle$.

Larson and Vogt obtained interesting results for counts of real bitangents according to the sign of their Qtype_{L_∞} : If $V(Q)$ does not meet the line at infinity, then the number of real bitangents with Qtype_{L_∞} equal to $\langle 1 \rangle$ minus the number of

real bitangents with Qtype_{L_∞} equal to $\langle -1 \rangle$ equals 4 [21]. Based on a randomized search, they formulate the following conjecture:

Conjecture 5.1 ([21, Conjecture 2]). *Let $V(Q)$ be a real quartic such that for each bitangent L , $L \cap V(Q) \cap L_\infty = \emptyset$, where L_∞ denotes the line at infinity. Then the number of real bitangents with Qtype_{L_∞} equal to $\langle 1 \rangle$ minus the number of real bitangents with Qtype_{L_∞} equal to $\langle -1 \rangle$ is in*

$$\{0, 2, 4, 6, 8\}.$$

Larson and Vogt prove that the numbers are nonnegative.

In the following, we give a partial proof for the conjecture, for the cases of quartics whose tropicalization is smooth and generic. In particular, the quartics we consider all intersect the infinite line in four points. It is interesting to observe that we do not obtain all the possible numbers with this restriction:

Theorem 5.2. *Let $C = V(Q)$ be a real quartic whose tropicalization is smooth and generic. Then the number of real bitangents with Qtype_{L_∞} equal to $\langle 1 \rangle$ minus the number of real bitangents with Qtype_{L_∞} equal to $\langle -1 \rangle$ is in*

$$\{0, 2, 4\}.$$

Our approach to prove this theorem is via GW-multiplicities of bitangent classes of the tropicalization. It builds on our classification of generic bitangent shapes and their dual motifs in Appendix A.1, on our study of GW-multiplicities of tropical bitangents in Theorem 4.23, A.2 and Appendix A.3. In the following lemma, we use the letters for the bitangent shapes which are introduced in the classification in Appendix A.1.

Lemma 5.3. *Let $V(Q)$ be a real quartic whose tropicalization is smooth and generic. Let S be a liftable tropical bitangent class of $\text{Trop}(V(Q))$. Let s denote the number of lifts of S with Qtype_{L_∞} equal to $\langle 1 \rangle$ minus the number of lifts of S with Qtype_{L_∞} equal to $\langle -1 \rangle$.*

If S is of shape (Nb) , (Ob) , (Oc) , (Pb) , (Qb) , (Rb) , (Rc) , (Sb) , (Ub) , (Vb) , (YaI) , (Yb) , (CCb) or (IIc) , then $s = 2$. If S is of shape (BBb) , then $s = 4$.

Otherwise $s = 0$.

Proof. This follows by going through the table in Appendix A.3 and only paying attention to the signs. \square

Proof of Theorem 5.2. By checking the dual motifs of a tropical bitangent class of one of the shapes with non-zero contribution above (and their symmetric cases w.r.t. $x - y$ -symmetry), we can see that the only ones which are not mutually exclusive are (Qb) , (Rb) , (Ub) , (Vb) , resp. $(YbII)$, (BBb) , (CCb) . For each of those, within a symmetry class, it is always the same edges which contain the tropicalization of the tangency points, thus, it is only the lengths of the edges of the tropical curve which decide which of (Qb) , (Rb) , (Ub) or (Vb) (resp. which of

(YbII), (BBb) or (CCb)) shows up, in particular, they do not show up together either. The only possibility is to combine a shape from (Ub) or (Vb) with another such shape after applying $x - y$ -symmetry. Also, the three possibilities for (Nb) do not exclude each other on the level of Newton subdivisions, but the vertex dual to triangle can only align with one of the possible three edges, and so it is precisely one of the types of (Nb) that can occur together. We can see that we can combine at most two shapes that contribute 2, all other shapes with nonzero contribution appear exclusively. As the maximal nonzero contribution from such a shape is 4, we obtain the desired result. \square

One does not need to rely on our results on GW-multiplicities of tropical bitangent classes from Theorem 4.23, A.2 and Appendix A.3 to show this result. An alternative approach is to use Viro's patchworking method, and read off the signs of the GW-multiplicities using Figure 11. We illustrate this approach in an example:

Example 5.4. Introduce Viro's combinatorial patchworking shortly here [34, 14].

Figure 14 shows the dual Newton subdivision of a tropicalized quartic, and the four copies of the tropicalized quartic. We pick the signs of all coefficients except a_{03} to be positive, and $s_{03} < 0$. The signs in the four copies are distributed accordingly. With thick black lines, the patchworked quartic is depicted. By Viro's Theorem, real quartics close to the tropical limit are homeomorphic to this picture. We consider the bitangent class of shape (E) which is also depicted in Figure 5. It has two liftable members. Their patchworked versions are drawn with thick red lines and denoted Λ_1 and Λ_2 in Figure 14. We can see altogether 4 tangency points, p_1, \dots, p_4 , shown in green in Figure 14. The two lifts of the tropical bitangent line for Λ_1 are represented by the same thick red line, but they are tangent at different points: one lift, call it L_{11} , is tangent at p_1 and p_3 , the other, L_{12} , at p_1 and p_4 . Analogously, one lift, L_{21} , of the tropical bitangent for Λ_2 is tangent at p_2 and p_3 , the other, L_{22} , at p_2 and p_4 . If we move along the thick red line for Λ_1 , starting at the top right corner, we meet the first tangency point, p_1 , on the right, and the second, p_3 , also on the right. Thus, the Qtype_{L_∞} of L_{11} is $\langle 1 \rangle$, see Figure 11. For L_{12} , we follow the same path, but now we meet p_1 on the right and p_4 on the left, so the Qtype_{L_∞} of L_{12} is $\langle -1 \rangle$. Analogously, the Qtype_{L_∞} for L_{21} is $\langle -1 \rangle$ and for L_{22} $\langle 1 \rangle$. With this patchworked picture, we thus confirm the result from Lemma 5.3 that the overall contribution of a tropical bitangent class of shape (E) is zero.

REFERENCES

- [1] Xavier Allamigeon, Pascal Benchimol, Stéphane Gaubert, and Michael Joswig. Tropicalizing the simplex algorithm. *SIAM J. Discrete Math.*, 29(2):751–795, 2015.
- [2] Matthew Baker, Yoav Len, Ralph Morrison, Nathan Pflueger, and Qingchun Ren. Bitangents of tropical plane quartic curves. *Mathematische Zeitschrift*, pages 1–15, 2015.




FIGURE 14. A patchworked quartic and the lifts of a tropical bitangent class.

- [3] Erwan Brugallé, Ilia Itenberg, Grigory Mikhalkin, and Kris Shaw. Brief introduction to tropical geometry. In *Proceedings of the Gökova Geometry-Topology Conference 2014*, pages 1–75. Gökova Geometry/Topology Conference (GGT), Gökova, 2015.
- [4] Erwan Brugallé and Lucia M. López de Medrano. Inflection points of real and tropical plane curves. *J. Singul.*, 4:74–103, 2012.
- [5] Melody Chan and Pakawut Jiradilok. Theta characteristics of tropical K_4 -curves. In *Combinatorial algebraic geometry*, volume 80 of *Fields Inst. Commun.*, pages 65–86. Fields Inst. Res. Math. Sci., Toronto, ON, 2017.
- [6] María Angélica Cueto and Hannah Markwig. The combinatorics and real lifting of tropical bitangents to quartics. Preprint, arXiv:2004.10891, 2020.
- [7] Wolfram Decker, Gert-Martin Greuel, Gerhard Pfister, and Hans Schönemann. SINGULAR 3-1-3. A computer algebra system for polynomial computations, Centre for Computer Algebra, University of Kaiserslautern, 2011. <http://www.singular.uni-kl.de>.
- [8] David Eisenbud. *Commutative algebra with a view toward algebraic geometry*. Graduate Texts in Mathematics. Springer, 150 edition, 1995.
- [9] M. A. Elomary and J.-P. Tignol. Springer’s theorem for tame quadratic forms over Henselian fields. *Math. Z.*, 269(1-2):309–323, 2011.
- [10] Ewgenij Gawrilow and Michael Joswig. polymake: a framework for analyzing convex polytopes. In Gil Kalai and Günter M. Ziegler, editors, *Polytopes — Combinatorics and Computation*, pages 43–74. Birkhäuser, 2000.
- [11] Alheydis Geiger and Marta Panizzut. Computing tropical bitangents to smooth quartic curves in polymake. Preprint, arXiv:2112.04447, 2021.
- [12] Alheydis Geiger and Marta Panizzut. A tropical count of real bitangents to plane quartic curves. Preprint, arXiv:2112.04433, 2021.

- [13] Marc Hoyois. A quadratic refinement of the Grothendieck-Lefschetz-Verdier trace formula. *Algebr. Geom. Topol.*, 14(6):3603–3658, 2014.
- [14] Ilya Itenberg, Grigory Mikhalkin, and Eugenii Shustin. *Tropical algebraic geometry*, volume 35 of *Oberwolfach Seminars*. Birkhäuser Verlag, Basel, second edition, 2009.
- [15] Anders N. Jensen, Hannah Markwig, and Thomas Markwig. tropical.lib. A SINGULAR 3.0 library for computations in tropical geometry. <http://www.mathematik.uni-kl.de/~keilen/de/tropical.html>, 2007.
- [16] Christian U. Jensen and Helmut Lenzing. *Model-theoretic algebra with particular emphasis on fields, rings, modules*, volume 2 of *Algebra, Logic and Applications*. Gordon and Breach Science Publishers, New York, 1989.
- [17] David Jensen and Yoav Len. Tropicalization of theta characteristics, double covers, and Prym varieties. *Selecta Math. (N.S.)*, 24(2):1391–1410, 2018.
- [18] Jesse Leo Kass and Kirsten Wickelgren. The class of Eisenbud-Khimshiashvili-Levine is the local \mathbf{A}^1 -Brouwer degree. *Duke Math. J.*, 168(3):429–469, 2019.
- [19] Jesse Leo Kass and Kirsten Wickelgren. An arithmetic count of the lines on a smooth cubic surface. *Compos. Math.*, 157(4):677–709, 2021.
- [20] Tsit Yuen Lam. *Introduction to quadratic forms over fields*, volume 67 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2005.
- [21] Hannah Larson and Isabel Vogt. An enriched count of the bitangents to a smooth plane quartic curve. *Res. Math. Sci.*, 8(2):Paper No. 26, 21, 2021.
- [22] Cédric Le Texier. Topology of real algebraic curves near the non-singular tropical limit. Preprint arXiv:2111.08607, 2021.
- [23] Heejong Lee and Yoav Len. Bitangents of non-smooth tropical quartics. *Port. Math.*, 75(1):67–78, 2018.
- [24] Yoav Len and Hannah Markwig. Lifting tropical bitangents. *J. Symbolic Comput.*, 96:122–152, 2020.
- [25] Marc Levine. Aspects of enumerative geometry with quadratic forms. *Doc. Math.*, 25:2179–2239, 2020.
- [26] Stephen McKean. An arithmetic enrichment of Bézout’s Theorem. *Mathematische Annalen*, 379, 02 2021.
- [27] Grigory Mikhalkin. Enumerative tropical geometry in \mathbb{R}^2 . *J. Amer. Math. Soc.*, 18:313–377, 2005.
- [28] Grigory Mikhalkin. Tropical geometry and its applications. In M. Sanz-Sole et al., editor, *Invited lectures v. II, Proceedings of the ICM Madrid*, pages 827–852, 2006.
- [29] OSCAR – Open Source Computer Algebra Research system, Version 0.8.3-DEV, 2022. <https://oscar.computeralgebra.de>.
- [30] Brian Osserman and Sam Payne. Lifting tropical intersections. *Doc. Math.*, 18:121–175, 2013.
- [31] Brian Osserman and Joseph Rabinoff. Lifting nonproper tropical intersections. In *Tropical and non-Archimedean geometry*, volume 605 of *Contemp. Math.*, pages 15–44. Amer. Math. Soc., Providence, RI, 2013.
- [32] Sam Payne. Analytification is the limit of all tropicalizations. *Math. Res. Lett.*, 16(3):543–556, 2009.
- [33] Andrés Jaramillo Puentes and Sabrina Pauli. Quadratically enriched tropical intersections. In preparation.
- [34] Oleg Viro. Real plane algebraic curves: constructions with controlled topology. *Algebra i Analiz*, 1(5):1–73, 1989.

APPENDIX A. CLASSIFICATION, COMPUTATIONS AND TABLE

A.1. Classification of generic bitangent shapes. For the classification of bitangent shapes up to \mathbb{S}_3 -symmetry, see Figure 6 in [6]. There are 41 shapes up to \mathbb{S}_3 -symmetry, labeled with capital letters such as (A), double capital letters such as (BB) or dashed capital letters such as (T''). For bitangent shapes, we follow the color coding of Figure 6 in [6]: The black cells of each bitangent class miss the tropicalized quartic, whereas the red ones lie on it. The unfilled dots are vertices of the tropicalized quartic.

In our study, we fix the line $\{z = 0\}$ and do not have \mathbb{S}_3 -symmetry for that reason, only \mathbb{S}_2 -symmetry for exchanging the variables x and y . We label shapes in an $\mathbb{S}_3/\mathbb{S}_2$ -orbit by adding lower case letters such as (Da). Sometimes, the shapes allow different cases of dual motifs in the dual Newton subdivision. These different cases play a role for the computation of Qtypes. We label such different cases of dual motifs by adding Roman numbers such as (TaI).

We consider only generic bitangent shapes, i.e. bitangent shapes which appear in the open cones of the subdivided secondary fan, where the subdivision is chosen such that the bitangent shapes are constant in each cell (see Remark 2.17). This subdivision has been computed in [11].

Lemma A.1. *Let S be a tropical bitangent class of a generic tropicalized quartic. Then, from the classification of possible shapes for S in [6], Figure 6, S can have the shapes (A), (B), (C), (D), (E), (F), (G), (H), (N), (O), (P), (T), (S), (R), (Q), (U), (V), (W), (Y), (BB), (CC), (EE), or (II).*

Proof. This follows since the existence of a tropical bitangent class of one of the remaining shapes ((H'), (I), (J), (K), (L'), (L), (M), (T'), (T''), (U'), (Q'), (X), (Z), (AA), (DD), (FF), (GG), (HH)) requires some equalities on the lengths of the edges of the tropicalized quartic to hold, hence they appear only in lower-dimensional cones of our subdivision of the secondary fan. \square

Figure 15 shows the three zero-dimensional bitangent shapes (A), (B) and (C). Next to the shape, we sketch the tropical bitangent in blue and parts of the tropicalized quartic which carry the tropicalization of the tangency points. Next, we draw an exemplary dual motif. The red edges are dual to the edges that overlap with the tropicalized quartic. The red edges could also be shifted for case (A) and (B). In case (A) and (B), the vertices on the vertical and diagonal boundary edge of the Newton triangle could also be shifted. In case (C), all vertices on the boundary could be shifted. We refrain from drawing the $\mathbb{S}_3/\mathbb{S}_2$ -orbits of these bitangent shapes, as the arguments for the computation for the Qtypes do not change. Note that each such shape has a $\mathbb{S}_3/\mathbb{S}_2$ -orbit of size two, as the tropicalization of the tangency points can be on the horizontal and vertical edge, or on the diagonal and horizontal edge.

Figure 13 shows bitangent shape (Da). Shape (Db) and (Dc) are diagonal resp. vertical. Figure 13 depicts the liftable tropical bitangents together with the




FIGURE 15. Bitangent shapes, local pictures and motifs of type (A), (B) and (C).

local parts of the tropicalized quartic carrying the tropicalization of the tangency points, and the dual motifs.

Bitangent shape (E) is a segment which does not meet the tropicalized quartic, i.e. in Figure 6 [6] it is drawn completely in black. The two end points of the segment are the liftable tropical bitangents. Figure 16 shows the behaviour of the tropicalization of the tangency points (i.e. local parts of the tropicalized quartic) together with the two liftable tropical bitangents in blue, and the partial dual motifs for the shapes (Ea), (Eb) and (Ec) which we obtain as the $\mathbb{S}_3/\mathbb{S}_2$ -orbit of (E). In the dual pictures, the red edge could also be shifted (upwards for (Ea) and (Eb), antidiagonally for (Ec)), as could the vertex which forms the left resp. upper triangle with the red edge.

Bitangent shape (F) is a segment which touches the tropicalized quartic, i.e. in the color coding we draw one vertex red. Again, the two end points of the segment are the liftable tropical bitangents. Figure 17 shows the behaviour of the tropicalization of the tangency points together with the two liftable tropical




FIGURE 16. Local pictures of tropicalizations of tangency points and partial motifs for type (Ea), (Eb) and (Ec).

bitangents in blue, and the partial dual motifs for the shapes (Fa), (Fb) and (Fc) which are the $\mathbb{S}_3/\mathbb{S}_2$ -orbit of (F). In the dual pictures, the red edge could also be shifted (upwards for (Fa) and (Fb), antidiagonally for (Fc)), as could the vertex which forms the left resp. upper triangle with the red edge.

Bitangent shape (G) is a segment which is completely contained in the tropicalized quartic. One tropical tangency is an overlap of an edge with the ray spanned by the segment. The two liftable members are the two end points of the segment. As we deform one liftable tropical bitangent to the other, the second tropical tangency component wanders over an edge which can have three possible slopes. We differentiate those three cases by using Roman letters (I), (II) and (III). Figure 18 combines the three possibilities (I), (II), (III) for the three $\mathbb{S}_3/\mathbb{S}_2$ -orbits (Ga), (Gb) and (Gc) in one picture each, both for the local pictures of the tropicalization of the tangency points and for the partial dual motifs.




FIGURE 17. Local pictures of tropicalization of the tangency points and partial dual motifs for type (Fa), (Fb) and (Fc).

Bitangent shape (H) is a ray whose vertex corresponds to the only liftable tropical bitangent in the class. Figure 19 shows the local pictures of the tropicalization of the tangency points and the partial dual motifs for the two cases (Ha) and (Hb). The red edge could be shifted upwards resp. antidiagonally, as could the vertex of its adjacent right resp. upper triangle.

Bitangent shape (N) looks like a reversed tropical line with two rays cut off. The end points of the two cut rays are the two liftable members. As for shape (G) (see Figure 18), one tropical tangency is an overlap with an edge (although the two overlapping edges differ for the two liftable tropical bitangents, different from the situation in (G)), and one tropical tangency component wanders over an edge — which can have three different possible slopes (I), (II) and (III) — as we deform one liftable tropical tangency component to the other. Figure 20 shows local pictures of the tropicalization of the tangency points and partial dual motifs, and combines the three options (I), (II) and (III) in one picture each for the two $\mathbb{S}_3/\mathbb{S}_2$ -orbits (Na) and (Nb).




FIGURE 18. Local pictures of tropical tangency components and partial dual motifs for types (GaI), (GaII), (GaIII), (GbI), (GbII), (GbIII), (GcI), (GcII) and (GcIII).

Shape (O) has two liftable tropical bitangents, the local pictures and dual motifs for the three symmetric cases (Oa), (Ob) and (Oc) are depicted in 21.

Shape (P) has two liftable tropical bitangents. Figure 22 shows local pictures of the tropical tangency components and dual motifs for the two symmetric cases (Pa) and (Pb).

Shape (Q) has two liftable tropical bitangents. Figure 23 shows the local pictures of the tropical tangency components and the dual motifs for the three symmetric cases (Qa), (Qb) and (Qc).

Figure 24 shows the local tropical tangency components and dual motifs for the three symmetric cases (Ra), (Rb) and (Rc) that we have for shape (R). Each has two liftable tropical bitangents.

In shape (S), there are two liftable tropical bitangents, shown in Figure 25 together with the dual motifs for the two symmetric cases (Sa) and (Sb).

Shape (T) is an unbounded 2-dimensional cell show two vertices each lift with multiplicity two. As for shape (G) and (N), there are three possibilities for the




FIGURE 19. Local pictures for the tropical tangency components and partial dual motifs for shapes (Ha), (Hb).




FIGURE 20. Local pictures of tropical tangency components and partial dual motifs for types (NaI), (NaII), (NaIII), (NbI), (NbII), (NbIII).

slope of the edge along which one of the tropical tangency component wanders, which we denote by (I), (II) and (III). Figure 26 combines these possibilities into one picture for (Ta) and one for (Tb), which are the two orbits we have to take into account.




FIGURE 21. Local pictures for tropical tangency components and dual motifs for types (Oa), (Ob) and (Oc).

Similar to shape (G), (N) and (T), shape (U) comes with three different possibilities for slopes for the wandering tropical tangency component, see Figure 27.

The behaviour of the tropicalization of the tangency points for shape (V) is the same as for shape (N), it is only the possible deformations that differ, but this has no effect on Qtypes, see Figure 28.

We now consider the bitangent shapes (W), (Y), (BB), (CC) and (EE) which each have four liftable members. For some of these shapes, tangencies can arise due to edges for which different slopes are possible. We label these different slopes again using Roman numerals (I), (II), (III). Bitangent shape (W) is a parallelogram whose four vertices are the liftable members. The other shapes can be viewed as deformations of the parallelogram, where edges of the tropicalized quartic "cut off" pieces. Figure 29 shows the local pieces of the tropicalized quartic together with the four liftable tropical bitangents in blue for the cases (WaI) and (WbI). Next to it, the partial dual motifs are depicted. For (WaII), (WaIII), (WbIII) and (WcII), the partial dual motifs are depicted in Figure 30. As the local pictures of




FIGURE 22. Local pictures of the tropical tangency components and dual motifs for shapes (Pa) and (Pb).

the tropical tangency components and liftable tropical bitangents are similar to the ones shown in Figure 29, we do not include those.

Concerning shape (Y), the dual motif for (YaI) equals the one for (WbII), (YbI) equals (WcII), (YcI) equals (WaII), (YaII) and (YaIII) equals (WaIII), and (YbII) equals (WbIII). The shape (Y) only differs from (W) by "cutting off" a vertex of the parallelogram — one liftable tropical bitangent now has its vertex on an edge of the tropicalized quartic. Pictures of the partial dual motifs can thus be found in Figure 30. Figure 31 depicts the 4 liftable tropical bitangents in each case together with a local picture of the tropicalized quartic.

Shape (BB) differs from shape (W) by cutting off two vertices of the parallelogram. The dual motif for (BBa) equals the one for (WaIII), the one for (BBb) equals (WbIII). Again, we refer to Figure 30 for these partial dual motifs. The 4 liftable tropical bitangents together with local pictures of the tropicalized quartic are found in Figure 32.

In shape (CC), we also cut off two vertices, but we cut one of those off even further. The dual motif for (CCa) equals (WaIII), the one for (CCb) (WbIII), see Figure 30. The corresponding local pictures of the tangencies are found in Figure 33. (CCaI) and (CCaII) differ by which of the two vertices is cut off further. For (CCb), this is symmetric.

In shape (EE), we only cut off one vertex, but we cut this one off even further again. The dual motif for (EEaI) and (EEaII) equals the one for (WaIII) (those two cases only differ by the choice which vertex of the parallelogram we cut off), the dual motif for (EEaIII) equals the one for (WaII), (EEbI) equals (WbIII), (EEbII)




FIGURE 23. Local pictures of tropical tangency components and dual motifs for shapes (Qa), (Qb) and (Qc).

equals (WbII) and (EEcI) equals (WcII), see Figure 30. The corresponding local pictures of the tangencies appear in Figure 34.

Finally, Figure 35 shows the dual motifs for the bitangent shapes (IIa), (IIb) and (IIc). For (IIa), there is also a local picture of the tropicalized quartic with the liftable tropical bitangent lines. The two left ones have lifting multiplicity one, the right one lifting multiplicity two. The local pictures for (IIb) and (IIc) are similar. For the two left tropical bitangents, the two tangency points tropicalize to the same point, namely a vertex of the tropicalized quartic, where the tropical line and the tropicalized quartic meet with intersection multiplicity four.

A.2. Details on Theorem 4.23 and its proof including the case-by-case analysis of bitangent shapes. The following Theorem spells out the details of the case-by-case analysis of bitangent shapes for Theorem 4.23:

Theorem A.2. *If a tropical bitangent class is of shape (A), (B), (C), (D), (E), (F), (G), (H), (Na), (Oa), (Pa), (Qa), (Qc), (Ra), (Sa), (IIa), (IIb), (T), (Ua), (Uc), (Va), (W), (YcI), (YaII), (CCaII), or (EE), then the total contribution of its Qtype_{L^∞} is $2\mathbb{H}$.*




FIGURE 24. Local tangencies and dual motifs for shapes (Ra), (Rb) and (Rc).

Proof. The strategy of the proof is to use Lemmas 4.18, 4.20 and 4.22 to argue that the four lifts come in pairs contribution $\langle \pm a \rangle$ for some a , which sums up to $2\mathbb{H}$. This works if the lifts exist in the field, else we use Remark 4.14 and Example 4.15 to show that nevertheless, we obtain $2\mathbb{H}$ as total contribution.

If it is of shape (A) or (B), there are two tropical tangency components in the interior of edges, of which at least one has to be vertical or horizontal. Using Lemma 4.18, we can pair up the lifts in such a way that in total we obtain $2\mathbb{H}$. The argument is analogous for shape (C), as locally the tropical tangency components are in the interior of edges here, too. Shape (D) was already discussed in detail in the proof of Theorem 4.23.

For (E),(G), one of the tropical tangency components has to be on the horizontal or vertical ray, and we can use Lemma 4.18 resp. Lemma 4.20, depending on which tropical tangency component it is, to pair up.

For (Fa) and (Fb), we have a tropical tangency component in the interior of a horizontal or vertical edge, and we can use Lemma 4.18 to pair up. For (Fc), we have two liftable tropical bitangents. Both have a segment of intersection with




FIGURE 25. Local tangencies of liftable tropical bitangents and dual motifs for shapes (Sa) and (Sb).




FIGURE 26. Local pictures of tropical tangency components and partial dual motifs for shapes (TaI), (TaII), (TaIII), (TbI), (TbII) and (TbIII).

$\text{Trop}(Q)$ on the diagonal ray, so by Lemma 4.19, we get the same contribution. The second tropical tangency component differs: for the first, it is, as in (Ec), an intersection with the horizontal ray, for the second, it is an intersection with the vertex. A computation shows that the contributions are negative of each other, so we can pair up and obtain $2\mathbb{H}$.




FIGURE 27. Local pictures of tropical tangency components and partial dual motifs for (UaI), (UaII), (UaIII), (UbI), (UbII), (UbIII), (UcI), (UcII) and (UcIII).

For (Ha), one tropical tangency component is in the interior of a horizontal edge. For (Hb), we can pair up using Lemma 4.22 and Lemma 4.19.

For (Na), (Oa), (Pa), (Ra), (Sa) we can pair up using the tropical tangency components on the horizontal resp. vertical ray. For (Qa), for one liftable point we can use the tropical tangency component on the horizontal edge to pair up, for the other, the tropical tangency component at the vertex of the tropical line as in Lemma 4.22. For (Qc), a computation shows that we obtain $\langle 1 \rangle + \langle 1 \rangle + \langle -1 \rangle + \langle -1 \rangle = 2\mathbb{H}$. For (IIa) and (IIb), we can use a computation to show that we obtain $2\mathbb{H}$. For (Ta), Lemma 4.22 shows that we can pair up. For (Tb), we can use Lemma 4.20 to pair up. For (Ua), we can use Lemma 4.22 and Lemma 4.18 to pair up.

For (Uc), a computation shows that the lifts of the right tropical tangency components yield a contribution of $+2$ for the lower liftable tropical bitangent, and -2 for the upper liftable tropical bitangent. Combining with Lemma 4.20, we can pair up.




FIGURE 28. Local pictures of tropical tangency components and partial dual motifs for shapes (VaI), (VaII), (VaIII), (VbI), (VbII) and (VbIII).




FIGURE 29. Local pictures of tropical tangency components, liftable tropical bitangents and partial dual motifs for shapes (WaI) and (WbI).

For (Va), we can use the tropical tangency components on the horizontal and vertical edge to pair up, using Lemma 4.18. For (W), one of the tropical tangency component has to be on a horizontal or vertical ray, and thus we can use Lemma 4.20 to pair up. For (EEaII), (EEbI), (EEbII), (EEcI, and (EEcII) we can use Lemma 4.20 to pair up. For (YcI), (YaII), (CCaII), (EEaI) and (EEaIII), we can pair up two liftable tropical bitangents using Lemma 4.20. For the remaining two, a computation shows that we can again lift up, similar to (Fc).

□




FIGURE 30. Partial dual motifs for the shapes (WaII), (WbII), (WcII), (WaIII), (WbIII).




FIGURE 31. Local pictures of liftable tropical bitangent lines for (YaI), (YbI), (YcI), (YaII), (YaIII), (YbII).




FIGURE 32. Local pictures of liftable tropical bitangent lines for (BBa) and (BBb).




FIGURE 33. Local pictures of liftable tropical bitangent lines for (CCaI), (CCaII), (CCb).




FIGURE 34. Local pictures of liftable tropical bitangent lines for (EEaI), (EEaII), (EEaIII), (EEbI), (EEbII), (EEcI).




FIGURE 35. Dual Motifs for (IIa), (IIb) and (IIc). For (IIa), a local picture involving the three liftable tropical bitangent lines and their tropical tangency components.

A.3. Exceptional GW-multiplicities. Here we discuss the GW-multiplicities of the special bitangent shapes which do not yield $2\mathbb{H}$ by Theorem 4.23 resp. A.2.

The shapes (Nb), (Ob), (Oc), (Pb), (Qb), (Rb), (Rc), (Sb), (Ub), (Vb), (IIc), (YbI), (YbII) and (CCb) all have similar behaviour of the tropical tangency components as described in the classification in Appendix A.1. The GW-multiplicity of a bitangent class of one of these shapes equals

$$\langle 1 \rangle + \langle 1 \rangle + \mathbb{H}.$$

The following table shows the GW-multiplicities of the remaining exceptional cases.

| bitangent | | | | |
|-----------|--|---------------------|---------------------|---------------------|
| (YaI) | $\langle a_{20}a_{31}a_{30}a_{03} \rangle$ | $\langle 2 \rangle$ | \mathbb{H} | |
| (YaIII) | $\langle -a_{01}a_{20}a_{31}a_{30} \rangle$ | $\langle 2 \rangle$ | \mathbb{H} | |
| (BBa) | $\langle -a_{01}a_{20}a_{31}a_{30} \rangle$ | $\langle 2 \rangle$ | \mathbb{H} | |
| (BBb) | $\langle 1 \rangle$ | $\langle 1 \rangle$ | $\langle 1 \rangle$ | $\langle 1 \rangle$ |
| (CCaI) | $\langle -2a_{20}a_{31}a_{30}a_{01} \rangle$ | $\langle 2 \rangle$ | \mathbb{H} | |

HANNAH MARKWIG, EBERHARD KARLS UNIVERSITÄT TÜBINGEN, FACHBEREICH MATHEMATIK, AUF DER MORGENSTELLE 10, 72076 TÜBINGEN, GERMANY

Email address: hannah@math.uni-tuebingen.de

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TEXAS AT AUSTIN, 2515 SPEEDWAY, PMA 8.100, AUSTIN, TX 78712, USA

Email address: sampayne@utexas.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OSLO, POSTBOKS 1053, BLINDERN, 0316 OSLO, NORWAY

Email address: krisshaw@math.uio.no