

ON THE EXISTENCE OF WEAK SOLUTIONS TO STOCHASTIC VOLTERRA EQUATIONS

DAVID J. PRÖMEL AND DAVID SCHEFFELS

ABSTRACT. The existence of weak solutions is established for stochastic Volterra equations with time-inhomogeneous coefficients and with convolutional or bounded kernels. The presented approach is based on a newly formulated local martingale problem associated to stochastic Volterra equations.

Key words: local martingale problem, singular kernel, stochastic Volterra equation, non-Lipschitz coefficients, weak existence.

MSC 2020 Classification: 60H20, 45D05.

1. INTRODUCTION

We investigate the existence of weak solutions to stochastic Volterra equation (SVEs)

$$(1.1) \quad X_t = x_0(t) + \int_0^t K_\mu(s, t)\mu(s, X_s) ds + \int_0^t K_\sigma(s, t)\sigma(s, X_s) dB_s, \quad t \in [0, T],$$

where x_0 is a continuous function, B is a Brownian motion, and the kernels K_μ, K_σ are measurable functions. The time-inhomogeneous coefficients μ, σ are only supposed to be continuous in space uniformly in time. In case of ordinary stochastic differential equations (SDEs), i.e. $K_\sigma = K_\mu = 1$, the existence of weak solutions was first proven by Skorokhod [Sko61] and can, nowadays, be found in different generality in standard textbooks like [SV79, KS91].

A comprehensive study of weak solutions to stochastic Volterra equations was recently initiated by Abi Jaber, Cuchiero, Larsson and Pulido [AJCLP21], see also [MS15]. The extension of the theory of weak solutions from ordinary stochastic differential equations to SVEs constitutes a natural generalization of the classical theory and is motivated by successful applications of SVEs with non-Lipschitz coefficients as volatility models in mathematical finance, see e.g. [EER19, AJEE19]. Assuming that the kernels in the SVE (1.1) are of convolution type, i.e. $K_\mu(s, t) = K_\sigma(s, t) = K(t - s)$ for some function $K: \mathbb{R} \rightarrow \mathbb{R}$, and that the coefficients μ, σ are continuous jointly in space-time, the existence of weak solutions was derived in [AJCLP21], see also [MS15, AJLP19, AJ21]. To that end, Abi Jaber et al. [AJCLP21] introduces a local martingale problem associated to SVEs of convolutional type.

In the present work we establish a local martingale problem associated to general stochastic Volterra equations, see Definition 2.4, and show that its solvability is equivalent to the existence of a weak solution to the associated SVE, see Lemma 2.7. Using this newly formulated Volterra local martingale problem, we obtain the existence of weak solutions to stochastic Volterra equations with convolutional kernels as well as with bounded kernels which are not necessarily of convolutional type, see Theorem 3.3. The presented approach can be considered, roughly speaking, as a generalization of Skorokhod's original construction to the more general case of SVEs, and is developed in a one-dimensional setting to keep the presentation

Date: July 5, 2022.

fairly short without cumbersome notation. However, as for ordinary SDEs, all concepts and results are expected to extend to a multi-dimensional setting in a straightforward manner.

Organization of the paper: In Section 2 we introduce a local martingale problem associated to SVEs. The existence of weak solutions to SVEs is provided in Section 3.

Acknowledgments: D. Scheffels gratefully acknowledges financial support by the Research Training Group “Statistical Modeling of Complex Systems” (RTG 1953) funded by the German Science Foundation (DFG).

2. WEAK SOLUTIONS AND THE VOLTERRA LOCAL MARTINGALE PROBLEM

For $T \in (0, \infty)$ we consider the one-dimensional stochastic Volterra equation

$$(2.1) \quad X_t = x_0(t) + \int_0^t K_\mu(s, t) \mu(s, X_s) ds + \int_0^t K_\sigma(s, t) \sigma(s, X_s) dB_s, \quad t \in [0, T],$$

where $x_0: [0, T] \rightarrow \mathbb{R}$ is a continuous function, $(B_t)_{t \in [0, T]}$ is a Brownian motion on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, and the coefficients $\mu, \sigma: [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ and the kernels $K_\mu, K_\sigma: \Delta_T \rightarrow \mathbb{R}$ are measurable functions, using the notation $\Delta_T := \{(s, t) \in [0, T] \times [0, T]: 0 \leq s \leq t \leq T\}$. The integral $\int_0^t K_\mu(s, t) \mu(s, X_s) ds$ is defined as a Riemann–Stieltjes integral and $\int_0^t K_\sigma(s, t) \sigma(s, X_s) dB_s$ as an Itô integral. Moreover, for $p \in [1, \infty)$ we write $L^p(\Omega \times [0, T])$ and $L^p([0, T])$ for the space of p -integrable functions on $\Omega \times [0, T]$ and on $[0, T]$, respectively.

Analogous to the notion of weak solutions to ordinary stochastic differential equations (see e.g. [KS91, Chapter 5.3, Definition 3.1], we make the following definition.

Definition 2.1. *A weak solution to (2.1) is a triple (X, B) , $(\Omega, \mathcal{F}, \mathbb{P})$, $(\mathcal{F}_t)_{t \in [0, T]}$ such that*

- (i) $(\Omega, \mathcal{F}, \mathbb{P})$ is a probability space, $(\mathcal{F}_t)_{t \in [0, T]}$ is a filtration of sub- σ -algebras of \mathcal{F} satisfying the usual conditions,
- (ii) $X = (X_t)_{t \in [0, T]} \in L^1(\Omega \times [0, T])$ is an (\mathcal{F}_t) -progressively measurable process, $B = (B_t)_{t \in [0, T]}$ is a Brownian motion w.r.t. $(\mathcal{F}_t)_{t \in [0, T]}$,
- (iii) $\int_0^t (|K_\mu(s, t) \mu(s, X_s)| + |K_\sigma(s, t) \sigma(s, X_s)|^2) ds < \infty$ \mathbb{P} -a.s. for any $t \in [0, T]$, and
- (iv) (2.1) holds for (X, B) on $(\Omega, \mathcal{F}, \mathbb{P})$, \mathbb{P} -a.s.

Under suitable assumptions on the coefficients and kernels, the existence of weak solutions to the stochastic Volterra equation (2.1) can be equivalently formulated in terms of solutions to an associated local martingale problem, see Definition 2.4 below. To that end, we make the following assumption.

Assumption 2.2. *Let $K_\mu, K_\sigma: \Delta_T \rightarrow \mathbb{R}$ be measurable functions with $K_\mu(\cdot, t) \in L^1([0, T])$ and $K_\sigma(\cdot, t) \in L^2([0, T])$ for every $t \in [0, T]$, and let $\mu, \sigma: [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ be measurable functions fulfilling the linear growth condition*

$$|\mu(t, x)| + |\sigma(t, x)| \leq C_{\mu, \sigma}(1 + |x|), \quad t \in [0, T], x \in \mathbb{R},$$

for some constant $C_{\mu, \sigma} > 0$.

Let $C^2(\mathbb{R})$ be the space of twice continuously differentiable functions $f: \mathbb{R} \rightarrow \mathbb{R}$ and $C_0^2(\mathbb{R})$ be the space of all $f \in C^2(\mathbb{R})$ with compact support. Given two stochastic processes $X = (X_t)_{t \in [0, T]}$ and $Z = (Z_t)_{t \in [0, T]}$ on a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in [0, T]}, \mathbb{P})$ satisfying

the usual conditions, such that $X \in L^1(\Omega \times [0, T])$ is (\mathcal{F}_t) -progressively measurable and Z is (\mathcal{F}_t) -adapted and continuous, we introduce the process $(\mathcal{M}_t^f)_{t \in [0, T]}$ by

$$(2.2) \quad \mathcal{M}_t^f := f(Z_t) - \int_0^t \mathcal{A}^f(s, X_s, Z_s) ds, \quad t \in [0, T],$$

for $f \in C^2(\mathbb{R})$, where

$$(2.3) \quad \mathcal{A}^f : [0, T] \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R} \quad \text{with} \quad \mathcal{A}^f(t, x, z) := \mu(t, x)f'(z) + \frac{1}{2}\sigma(t, x)^2 f''(z).$$

As we shall see in the next proposition, assuming that $(\mathcal{M}_t^f)_{t \in [0, T]}$ is a local martingale for all $f \in C_0^2(\mathbb{R})$ implies that the stochastic process Z is a semimartingale.

Proposition 2.3. *Suppose Assumption 2.2. Let $(X_t)_{t \in [0, T]}$ be an (\mathcal{F}_t) -progressively measurable process in $L^1(\Omega \times [0, T])$ and $(Z_t)_{t \in [0, T]}$ be an (\mathcal{F}_t) -adapted and continuous process on a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in [0, T]}, \mathbb{P})$ satisfying the usual conditions. If $(\mathcal{M}_t^f)_{t \in [0, T]}$ is a local martingale for every $f \in C_0^2(\mathbb{R})$, then we have:*

- (i) $(Z_t)_{t \in [0, T]}$ is a semimartingale with characteristics $(\int_0^\cdot \mu(s, X_s) ds, \int_0^\cdot \sigma(s, X_s)^2 ds, 0)$.
- (ii) There exists a filtered probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, (\tilde{\mathcal{F}}_t)_{t \in [0, T]}, \tilde{\mathbb{P}})$ satisfying the usual conditions such that $(Z_t)_{t \in [0, T]}$ is a semimartingale on $(\tilde{\Omega}, \tilde{\mathcal{F}}, (\tilde{\mathcal{F}}_t)_{t \in [0, T]}, \tilde{\mathbb{P}})$ and

$$Z_t = \int_0^t \mu(s, X_s) ds + \int_0^t \sigma(s, X_s) dB_s, \quad t \in [0, T],$$

holds $\tilde{\mathbb{P}}$ -a.s., for some Brownian motion $(B_t)_{t \in [0, T]}$ on $(\tilde{\Omega}, \tilde{\mathcal{F}}, (\tilde{\mathcal{F}}_t)_{t \in [0, T]}, \tilde{\mathbb{P}})$.

Proof. (i) By [JS03, Theorem II.2.42], in order to prove the assertion, it is sufficient to show that $(\mathcal{M}_t^f)_{t \in [0, T]}$, defined in (2.2), is a local martingale for every bounded function $f \in C^2(\mathbb{R})$.

Let $f \in C^2(\mathbb{R})$ be bounded and define the hitting times

$$\tau_n := \inf_{t \in [0, T]} \{\max(|X_t|, |Z_t|) \geq n\}, \quad n \in \mathbb{N}.$$

Note that $\tau_n \rightarrow T$ a.s. as $n \rightarrow \infty$ since $X \in L^1(\Omega \times [0, T])$ and Z is continuous. Since the underlying filtered probability space satisfies the usual conditions, by the Début theorem (see [RY99, Chapter I, (4.15) Theorem]), the hitting times $(\tau_n)_{n \in \mathbb{N}}$ are stopping times. It remains to show that $(\tau_n)_{n \in \mathbb{N}}$ is a localizing sequence for $(\mathcal{M}_t^f)_{t \in [0, T]}$. To that end, we approximate f by the functions $(f_n)_{n \in \mathbb{N}} \subset C_0^2(\mathbb{R})$ given by $f_n := \phi_n f$ for some $\phi_n \in C_0^2(\mathbb{R})$ taking values in $[0, 1]$ and being identical to 1 on $[-n, n]$. Hence, $(\mathcal{M}_t^{f_n})_{t \in [0, T]}$ is a local martingale for every $n \in \mathbb{N}$ and, thus, the stopped process $(\mathcal{M}_{t \wedge \tau_n}^{f_n})_{t \in [0, T]}$, given by

$$\mathcal{M}_{t \wedge \tau_n}^{f_n} = (f_n)(Z_{t \wedge \tau_n}) - \int_0^{t \wedge \tau_n} \mathcal{A}^{f_n}(s, X_s, Z_s) ds, \quad t \in [0, T],$$

is a martingale as

$$|\mathcal{M}_{t \wedge \tau_n}^{f_n}| \leq \sup_{x \in \mathbb{R}} |f(x)| + C_{\sigma, \mu, n} n^2 < \infty,$$

for some constant $C_{\sigma, \mu, n} > 0$, using the definition of τ_n and the linear growth condition on μ and σ . Since $\mathcal{M}_{t \wedge \tau_n}^{f_n} = \mathcal{M}_{t \wedge \tau_n}^f$ for $t \in [0, T]$, $(\mathcal{M}_{t \wedge \tau_n}^f)_{t \in [0, T]}$ is a martingale for every $n \in \mathbb{N}$ and, hence, $(\tau_n)_{n \in \mathbb{N}}$ a localizing sequence for $(\mathcal{M}_t^f)_{t \in [0, T]}$.

(ii) Since the process $(Z_t)_{t \in [0, T]}$ is a semimartingale with absolutely continuous characteristics $(\int_0^\cdot \mu(s, X_s) ds, \int_0^\cdot \sigma^2(s, X_s) ds, 0)$, the assertion follows by [JP12, Theorem 2.1.2]. \square

Keeping these preliminary considerations and the classical martingale problem (see e.g. [KS14, Definition 7.1.1]) in mind, we formulate a local martingale problem associated to the stochastic Volterra equation (2.1).

Definition 2.4. A solution to the Volterra local martingale problem given $(x_0, \mu, \sigma, K_\mu, K_\sigma)$ is a triple $(X, Z), (\Omega, \mathcal{F}, \mathbb{P}), (\mathcal{F}_t)_{t \in [0, T]}$ such that

- (i) $(\Omega, \mathcal{F}, \mathbb{P})$ is a probability space, $(\mathcal{F}_t)_{t \in [0, T]}$ is a filtration of sub- σ -algebras of \mathcal{F} satisfying the usual conditions,
- (ii) $X = (X_t)_{t \in [0, T]} \in L^1(\Omega \times [0, T])$ is an (\mathcal{F}_t) -progressively measurable process,
- (iii) $(Z_t)_{t \in [0, T]}$ is a continuous semimartingale with $Z_0 = 0$ and decomposition $Z = A + M$ for some process $(A_t)_{t \in [0, T]}$ of bounded variation and some local martingale $(M_t)_{t \in [0, T]}$,
- (iv) the process $(\mathcal{M}_t^f)_{t \in [0, T]}$, given by

$$(2.4) \quad \mathcal{M}_t^f := f(Z_t) - \int_0^t \mathcal{A}^f(s, X_s, Z_s) ds, \quad t \in [0, T],$$

is a local martingale for every $f \in C_0^2(\mathbb{R})$, where \mathcal{A}^f is defined as in (2.3), and

- (v) the following equality holds:

$$(2.5) \quad X_t = x_0(t) + \int_0^t K_\mu(s, t) dA_s + \int_0^t K_\sigma(s, t) dM_s, \quad t \in [0, T], \quad \mathbb{P}\text{-a.s.}$$

Remark 2.5. The first Volterra local martingale problem was formulated in [AJCLP21] for stochastic Volterra equations of convolution type, that is, the kernels K_μ, K_σ are supposed to be of the form $K(t - s)$ for a deterministic function $K: [0, T] \rightarrow \mathbb{R}$, see [AJCLP21, Definition 3.1]. Compared to [AJCLP21, Definition 3.1], the essential difference is that we reformulated [AJCLP21, (3.3)] to the condition (2.5). While both conditions are equivalent for kernels of convolutional type, the advantage of (2.5) is that it allows for general kernels.

Moreover, notice that the Volterra local martingale problem as presented in Definition 2.4 reduces to the local martingale problem for ordinary stochastic differential equations in the case $K_\mu = K_\sigma = 1$. Indeed, in this case conditions (i) and (iv) imply conditions (iii) and (v) on a possibly extended probability space, see Proposition 2.3.

Remark 2.6. Condition (iii) of Definition 2.4 can be relaxed to the condition “ $(Z_t)_{t \in [0, T]}$ is an (\mathcal{F}_t) -adapted and continuous process” since this together with (iv) of Definition 2.4 already implies the semimartingale property of $(Z_t)_{t \in [0, T]}$, see Proposition 2.3. However, we decided to directly postulate the semimartingale property of $(Z_t)_{t \in [0, T]}$ in the formulation of the Volterra local martingale problem to ensure that condition (v) is obviously well-defined.

As for ordinary stochastic differential equations, the existence of weak solutions to SVEs is equivalent to the solvability of the associated Volterra local martingale problem.

Lemma 2.7. Suppose Assumption 2.2. There exists a weak solution to the SVE (2.1) if and only if there exists a solution to the Volterra local martingale problem given $(x_0, \mu, \sigma, K_\mu, K_\sigma)$.

Proof. Let (X, B) be a (weak) solution to (2.1) on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Setting

$$Z_t := A_t + M_t := \int_0^t \mu(s, X_s) ds + \int_0^t \sigma(s, X_s) dB_s, \quad t \in [0, T],$$

Itô's formula applied to $f(Z_t)$ for $f \in C_0^2(\mathbb{R})$ yields that

$$\begin{aligned} \mathcal{M}_t^f &= f(Z_t) - \int_0^t f'(Z_s)\mu(s, X_s) ds - \frac{1}{2} \int_0^t f''(Z_s)\sigma(s, X_s)^2 ds \\ &= f(Z_0) + \int_0^t f'(Z_s)\sigma(s, X_s) dB_s, \end{aligned}$$

which is a local martingale and, by its definition, Z is a semimartingale satisfying (2.5).

Conversely, if there exists a solution to the Volterra local martingale problem, we obtain a weak solution to the SVE (2.1) by using (2.5) and Proposition 2.3, which yields that $A_t = \int_0^t \mu(s, X_s) ds$ and $M_t = \int_0^t \sigma(s, X_s) dB_s$ for some Brownian motion $(B_t)_{t \in [0, T]}$. \square

3. EXISTENCE OF WEAK SOLUTIONS

In this section we establish the existence of a weak solution to the SVE (2.1) and, equivalently, of a solution to the associated Volterra local martingale problem, under suitable assumptions on the initial condition, coefficients and kernels, which we state in the following.

Assumption 3.1. *There is some $p \in (4, \infty)$ and some $\gamma \in (\frac{2}{p}, \frac{1}{2})$ such that:*

(i) *There is a constant $C_p > 0$ such that, for all $(t, t') \in \Delta_T$,*

$$(3.1) \quad \begin{aligned} \int_0^t |K_\mu(s, t') - K_\mu(s, t)|^{\frac{p}{p-1}} ds + \int_t^{t'} |K_\mu(s, t')|^{\frac{p}{p-1}} ds &\leq C_p |t' - t|^{\frac{\gamma p}{p-1}}, \\ \int_0^t |K_\sigma(s, t') - K_\sigma(s, t)|^{\frac{2p}{p-2}} ds + \int_t^{t'} |K_\sigma(s, t')|^{\frac{2p}{p-2}} ds &\leq C_p |t' - t|^{\frac{2\gamma p}{p-2}}. \end{aligned}$$

(ii) *The coefficients $\mu, \sigma: [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ are measurable functions such that for every compact set $\mathcal{K} \subset \mathbb{R}$ and every $\epsilon > 0$ there exists a $\delta > 0$ such that*

$$|\mu(t, x) - \mu(t, y)| + |\sigma(t, x) - \sigma(t, y)| \leq \epsilon, \quad t \in [0, T], x, y \in \mathcal{K} \text{ with } |x - y| \leq \delta,$$

and μ, σ fulfill the linear growth condition

$$(3.2) \quad |\mu(t, x)| + |\sigma(t, x)| \leq C_{\mu, \sigma}(1 + |x|), \quad t \in [0, T], x \in \mathbb{R},$$

for a constant $C_{\mu, \sigma} > 0$

(iii) *The initial condition $x_0: [0, T] \rightarrow \mathbb{R}$ is β -Hölder continuous for every $\beta \in (0, \gamma - 1/p)$.*

To formulate our second assumption, for $K: \Delta_T \rightarrow \mathbb{R}$, we say $K(\cdot, t)$ is absolutely continuous for every $t \in [0, T]$ if there exists an integrable function $\partial_1 K: \Delta_T \rightarrow \mathbb{R}$ such that $K(s, t) - K(0, t) = \int_0^s \partial_1 K(u, t) du$ for $(s, t) \in \Delta_T$.

Assumption 3.2. *Each kernel $K \in \{K_\mu, K_\sigma\}$ satisfies at least one of the following condition:*

(i) *K is a measurable and bounded function. In the case of $K = K_\sigma$, $K_\sigma(\cdot, t)$ is absolutely continuous for every $t \in [0, T]$ such that $\partial_1 K_\sigma$ fulfills*

$$\sup_{t \in [0, T]} \left| \int_0^t |\partial_1 K_\sigma(s, t)|^p ds \right|^{\frac{1}{p}} \leq C$$

for some $p > 1$ and some constant $C > 0$.

(ii) *$K(s, t) = \tilde{K}(t - s)$ for all $(s, t) \in \Delta_T$ for a function $\tilde{K} \in L^1([0, T])$ if $K = K_\mu$ and $\tilde{K} \in L^2([0, T])$ if $K = K_\sigma$.*

Note, that in case of Assumption 3.2 (i), the bound on the second summand in (3.1) is trivially fulfilled. With these assumptions at hand we are ready to state our main result.

Theorem 3.3. *Suppose Assumptions 3.1 and 3.2. Then, there exists a weak solution (in the sense of Definition 2.1) to the stochastic Volterra equation (2.1).*

Before proving the aforementioned existence result, let us briefly discuss some properties of weak solutions to the SVE (2.1) and some exemplary kernels.

Remark 3.4. *Suppose Assumption 3.1. Due to [PS22, Lemma 3.1 and 3.2], any weak solution $(X)_{t \in [0, T]}$ to the SVE (2.1) satisfies $\sup_{t \in [0, T]} \mathbb{E}[|X_t|^q] < \infty$ for any $q \in [1, \infty)$ and $(X_t)_{t \in [0, T]}$ possesses a β -Hölder continuous modification for any $\beta \in (0, \gamma - 1/p)$.*

Remark 3.5. *Assumptions 3.1 and 3.2 are satisfied, e.g., by the following type of kernels:*

- (i) $K(s, t) := (t - s)^{-\alpha}$ for $\alpha \in (0, \frac{1}{2})$ for any $p \in (\frac{6}{1-2\alpha}, \infty)$ with $\gamma = \frac{1}{2} - \alpha - \frac{1}{p}$,
- (ii) $K(s, t) := \tilde{K}(t - s)$ for a Lipschitz continuous function $\tilde{K}: [0, T] \rightarrow \mathbb{R}$,
- (iii) kernels fulfilling [PS22, Assumption 2.1], and
- (iv) weakly differentiable kernels such that $\partial_1 K_\sigma(s, t) \leq C(t - s)^{-\alpha}$.

The remainder of the paper is devoted to implement the proof of Theorem 3.3 based on several auxiliary lemmas. Note that Lemma 3.9 implies Theorem 3.3 due to Lemma 2.7.

Assuming the coefficients μ, σ satisfy Assumption 3.1, the next lemma provides a way to approximate μ, σ locally uniformly by Lipschitz continuous coefficients.

Lemma 3.6. *Let $f: [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ be a measurable function such that for every compact set $\mathcal{K} \subset \mathbb{R}$ and every $\epsilon > 0$ there exists $\delta > 0$ such that*

$$|f(t, x) - f(t, y)| \leq \epsilon, \quad t \in [0, T], x, y \in \mathcal{K} \text{ with } |x - y| \leq \delta,$$

and such that f fulfills the linear growth condition

$$(3.3) \quad |f(t, x)| \leq C_f(1 + |x|), \quad t \in [0, T], x \in \mathbb{R},$$

for some constant $C_f > 0$. Then, there is a sequence $(f_n)_{n \in \mathbb{N}}$ of measurable functions $f_n: [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$, which satisfies:

- (i) *linear growth:* for $C_f > 0$ as in (3.3), we have

$$|f_n(t, x)| \leq 2C_f(1 + |x|), \quad t \in [0, T], x \in \mathbb{R};$$

- (ii) *Lipschitz continuity:* for each $n \in \mathbb{N}$ there is a $C_n > 0$ such that

$$|f_n(t, x) - f_n(t, y)| \leq C_n|x - y|, \quad t \in [0, T], x, y \in \mathbb{R};$$

- (iii) *locally uniform convergence:* for all $r \in (0, \infty)$ we have

$$\sup_{t \in [0, T], x \in [-r, r]} |f(t, x) - f_n(t, x)| \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

Proof. We explicitly choose the sequence $(f_n)_{n \in \mathbb{N}}$ by

$$f_n(t, x) := \phi_n(x) \int_{\mathbb{R}} f(t, x - y) \delta_n(y) dy, \quad n \in \mathbb{N},$$

for some $\phi_n \in C_0^2(\mathbb{R})$ with support in $[-(n + 1), n + 1]$, taking values in $[0, 1]$ and being identical to 1 on $[-n, n]$, where $\delta_n(y) := \frac{1}{c_n}(1 - y^2)^n \mathbf{1}_{[-1, 1]}(y)$ with $c_n := \int_{[-1, 1]} (1 - y^2)^n dy$.

(i) For $t \in [0, T]$ and $x \in \mathbb{R}$, using the linear growth condition on f , we get

$$|f_n(t, x)| \leq C_f \int_{[-1, 1]} (1 + |x - y|) \delta_n(y) dy \leq C_f \int_{[-1, 1]} (2 + |x|) \delta_n(y) dy \leq 2C_f(1 + |x|).$$

(ii) Let $t \in [0, T]$, $x, y \in \mathbb{R}$ and $n \in \mathbb{N}$. Using the compact support of f_n and the fact, that every δ_n is Lipschitz continuous as a smooth function with compact support, we get

$$|f_n(t, x) - f_n(t, y)| \leq C_f c_n |x - y| \int_{-(n+2)}^{n+2} (1 + |z|) dz \leq C_n |x - y|$$

for some constant C_n .

(iii) Due to the continuity property of f , we can find for every $r > 0$ and for every $\epsilon > 0$ some $\delta > 0$ such that for all $x, y \in [-r, r]$ with $|x - y| \leq \delta$ and all $t \in [0, T]$ holds $|f(t, x) - f(t, y)| \leq \epsilon$. Assuming $n \in \mathbb{N}$ to be large enough that $\phi_n \equiv 1$ on $[-r, r]$, we get for any $x \in [-r, r]$,

$$\begin{aligned} & |f(t, x) - f_n(t, x)| \\ &= \int_{[-\delta, \delta]} \delta_n(y) |f(t, x) - f(t, x - y)| dy + \int_{[-1, 1] \setminus [-\delta, \delta]} \delta_n(y) |f(t, x) - f(t, x - y)| dy. \end{aligned}$$

Let now $N(\epsilon, r) > 0$ be big enough, such that $\int_{[-1, 1] \setminus [-\delta, \delta]} \delta_n(y) dy < \epsilon$ and $\phi_n \equiv 1$ on $[-r, r]$ for all $n \geq N(\epsilon, r)$. Then, setting $\tilde{r} := r + 1$ for all $n \geq N(\epsilon, r)$

$$|f(t, x) - f_n(t, x)| \leq \int_{[-\delta, \delta]} \delta_n(y) \epsilon dy + 2\epsilon \sup_{\substack{s \in [0, T], \\ \tilde{x} \in [-\tilde{r}, \tilde{r}]}} |f(s, \tilde{x})| \leq \epsilon \left(1 + 2 \sup_{\substack{s \in [0, T], \\ \tilde{x} \in [-\tilde{r}, \tilde{r}]}} |f(s, \tilde{x})| \right),$$

which tends to zero as $\epsilon \rightarrow 0$. □

A suitable approximation, like the one provided in Lemma 3.6, ensures the convergence of associated Riemann–Stieltjes integrals. We denote by $C([0, T]; \mathbb{R}^d)$ the space of all continuous functions $g: [0, T] \rightarrow \mathbb{R}^d$, which is equipped with the supremum norm $\|\cdot\|_\infty$.

Lemma 3.7. *Let $f: [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ be a function such that for every compact set $\mathcal{K} \subset \mathbb{R}$ and every $\epsilon > 0$ there exists $\delta > 0$ such that*

$$(3.4) \quad |f(t, x) - f(t, y)| \leq \epsilon, \quad t \in [0, T], \quad x, y \in \mathcal{K} \text{ with } |x - y| \leq \delta,$$

and $(f_k)_{k \in \mathbb{N}}$ be a sequence of functions such that $f_k: [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ and $|f(t, x)| + |f_k(t, x)| \leq C(1 + |x|^2)$, $t \in [0, T]$, $x \in \mathbb{R}$, for all $k \in \mathbb{N}$ and for some $C > 0$, and $f_k \rightarrow f$ locally uniformly. Let $K: \Delta_T \rightarrow \mathbb{R}$ be measurable and bounded. If $(X^k)_{k \in \mathbb{N}}$ is a sequence of continuous stochastic processes such that $X^k \rightarrow X$ in $C([0, T]; \mathbb{R})$ as $k \rightarrow \infty$ \mathbb{P} -a.s., then

$$\left(\int_0^\cdot K(s, \cdot) f_k(s, X_s^k) ds \right)_{t \in [0, T]} \xrightarrow{\mathbb{P}} \left(\int_0^\cdot K(s, \cdot) f(s, X_s) ds \right)_{t \in [0, T]} \quad \text{w.r.t. } \|\cdot\|_\infty, \quad k \rightarrow \infty,$$

where $\xrightarrow{\mathbb{P}}$ denotes convergence in probability.

Proof. First, note that due to the continuity condition (3.4), for every $n \in \mathbb{N}$ there exists some continuous function $g_n: [0, \infty) \rightarrow [0, \infty)$ with $g_n(0) = 0$, such that for all $x, y \in [-n, n]$,

$$|f(t, x) - f(t, y)| \leq g_n(|x - y|), \quad t \in [0, T].$$

Let $\epsilon > 0$ and $\delta > 0$ be fixed but arbitrary. Choose $N \in \mathbb{N}$ and $K \in \mathbb{N}$ big enough such that

$$\mathbb{P}(\|X\|_\infty \geq n/2) \leq \delta/4 \quad \text{and} \quad \mathbb{P}(\|X^k - X\|_\infty \geq n/2) \leq \delta/4,$$

for all $n \geq N$ and $k \geq K$. Then,

$$\begin{aligned} \mathbb{P}(\|X\|_\infty \vee \|X^k\|_\infty \geq n) &\leq \mathbb{P}\left(\{\|X\|_\infty \geq n\} \cup \{\|X^k - X\|_\infty + \|X\|_\infty \geq n\}\right) \\ &\leq \mathbb{P}(\|X\|_\infty \geq n/2) + \mathbb{P}(\|X^k - X\|_\infty \geq n/2) \\ &\leq \delta/4 + \delta/4 = \delta/2. \end{aligned}$$

For every $n, k \in \mathbb{N}$, on $\{\|X\|_\infty \vee \|X^k\|_\infty \leq n\}$ we can bound

$$\begin{aligned} A_t^k - A_t &:= \int_0^t K(s, t) f_k(s, X_s^k) ds - \int_0^t K(s, t) f(s, X_s) ds \\ &\leq K \int_0^t |f_k(s, X_s^k) - f(s, X_s^k)| ds + K \int_0^t |f(s, X_s^k) - f(s, X_s)| ds \\ (3.5) \quad &\leq TK \left(\sup_{t \in [0, T], x \in [-n, n]} |f_k(t, x) - f(t, x)| + g_n(\|X^k - X\|_\infty) \right), \quad t \in [0, T], \end{aligned}$$

where $K := \sup_{(s, t) \in \Delta_T} |K(s, t)|$. For every $n \in \mathbb{N}$ we choose $K_{\epsilon\delta}^n \in \mathbb{N}$ sufficiently large such that

$$\mathbb{P}\left(\sup_{t \in [0, T], x \in [-n, n]} |f_k(t, x) - f(t, x)| + g_n(\|X^k - X\|_\infty) \geq \epsilon/TK \right) \leq \delta/2, \quad k \geq K_{\epsilon\delta}^n.$$

Setting $K_{\epsilon\delta} := \max\{K_{\epsilon\delta}^N, K\}$, we get

$$\begin{aligned} &\mathbb{P}(\|A^k - A\|_\infty \geq \epsilon) \\ &\leq \mathbb{P}\left(\{\|A^k - A\|_\infty \geq \epsilon\} \cup \{\|X\|_\infty \vee \|X^k\|_\infty < N\}\right) + \mathbb{P}(\|X\|_\infty \vee \|X^k\|_\infty \geq N) \\ &\leq \mathbb{P}\left(\sup_{t \in [0, T], x \in [-N, N]} |f_k(t, x) - f(t, x)| + g_N(\|X^k - X\|_\infty) \geq \epsilon/TK \right) + \delta/2 \leq \delta, \end{aligned}$$

for all $k \geq K_{\epsilon\delta}$, which shows the desired convergence. \square

Given coefficients μ, σ satisfying Assumption 3.1, we fix, relying on Lemma 3.6, two sequences $(\mu_n)_{n \in \mathbb{N}}$ and $(\sigma_n)_{n \in \mathbb{N}}$ with

$$\mu_n : [0, T] \times \mathbb{R} \rightarrow \mathbb{R} \quad \text{and} \quad \sigma_n : [0, T] \times \mathbb{R} \rightarrow \mathbb{R},$$

that fulfill properties (i)-(iii) of Lemma 3.6. For every $n \in \mathbb{N}$, we define $(X_t^n)_{t \in [0, T]}$ as the unique (strong) solution (see e.g. [PS22, page 4] for the definition of unique strong solutions to SVEs) to the stochastic Volterra equation

$$(3.6) \quad X_t^n = x_0(t) + \int_0^t K_\mu(s, t) \mu_n(s, X_s^n) ds + \int_0^t K_\sigma(s, t) \sigma_n(s, X_s^n) dB_s, \quad t \in [0, T],$$

given a Brownian motion $(B_t)_{t \in [0, T]}$ on some probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Note that $(X_t^n)_{t \in [0, T]}$ exists by [Wan08, Theorem 1.1] due to the Lipschitz continuity of μ_n and σ_n . Furthermore, we introduce the sequences $(A^n)_{n \in \mathbb{N}}$ and $(M^n)_{n \in \mathbb{N}}$ by

$$(3.7) \quad A_t^n := \int_0^t \mu_n(s, X_s^n) ds \quad \text{and} \quad M_t^n := \int_0^t \sigma_n(s, X_s^n) dB_s, \quad t \in [0, T].$$

In the following, we denote $X \stackrel{\mathcal{L}}{\sim} Y$ for equality in law of stochastic processes X and Y .

Lemma 3.8. *Suppose Assumption 3.1 and let $(X^n)_{n \in \mathbb{N}}$, $(A^n)_{n \in \mathbb{N}}$ and $(M^n)_{n \in \mathbb{N}}$ be given by (3.6) and (3.7). Then, there exist continuous stochastic processes $(\hat{X}^k)_{k \in \mathbb{N}}$, $(\hat{A}^k)_{k \in \mathbb{N}}$, $(\hat{M}^k)_{k \in \mathbb{N}}$, X , A , M and a Brownian motion $(\tilde{B}_t)_{t \in [0, T]}$ on a common probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$ such that $(\hat{X}^k, \hat{A}^k, \hat{M}^k) \rightarrow (X, A, M)$ in $C([0, T]; \mathbb{R}^3)$ as $k \rightarrow \infty$ $\tilde{\mathbb{P}}$ -a.s., $(\hat{X}^k, \hat{A}^k, \hat{M}^k) \stackrel{\mathcal{L}}{\rightarrow} (X^{n_k}, A^{n_k}, M^{n_k})$ and M is a local martingale with the representation*

$$M_t = \int_0^t \sigma(s, X_s) d\tilde{B}_s, \quad t \in [0, T],$$

where $(X^{n_k}, A^{n_k}, M^{n_k})_{k \in \mathbb{N}}$ denotes some subsequence of $(X^n, A^n, M^n)_{n \in \mathbb{N}}$.

Proof. First we want to apply Kolmogorov's tightness criterion (see [KS91, Problem 2.4.11]) to the probability measures $(\mathbb{P}_{(X^n, A^n, M^n, B)})_{n \in \mathbb{N}}$ associated to the four-dimensional stochastic processes $(X^n, A^n, M^n, B)_{n \in \mathbb{N}}$. By Lemma 3.6 (i) we know, that the coefficients μ_n and σ_n fulfill the linear growth condition (3.2) with uniformly bounded constants, i.e. $C_{\mu_n, \sigma_n} \leq 2C_{\mu, \sigma}$ for all $n \in \mathbb{N}$. Hence, using $p \in (4, \infty)$ from Assumption 3.1, we deduce, by [PS22, Lemma 3.1], that

$$\sup_{n \in \mathbb{N}} \sup_{s \in [0, T]} \mathbb{E}[|X_s^n|^p] \leq C \left(1 + \sup_{s \in [0, T]} |x_0(s)| \right)^p < \infty,$$

and, by [PS22, Lemma 3.2 and Remark 3.3], that

$$\mathbb{E}[|X_{t'}^n - X_t^n|^p] \leq C|t' - t|^{\beta p}, \quad n \in \mathbb{N},$$

for every $\beta \in (0, \gamma - 1/p)$, where the constant $C > 0$ depends only on p , T , K_μ , K_σ and $C_{\mu, \sigma}$. Moreover, it is straightforward to show that

$$\mathbb{E}[|A_{t'}^n - A_t^n|^p] \leq C|t' - t|^{\frac{\beta}{2}} \quad \text{and} \quad \mathbb{E}[|M_{t'}^n - M_t^n|^p] \leq C|t' - t|^{\frac{\beta}{2}}$$

for all $0 \leq t \leq t' \leq T$ and some constant $C > 0$, by Hölder's inequality and Burkholder–Davis–Gundy's inequality, respectively. Choosing β sufficiently close to $\gamma - 1/p$ so that $\beta p > 1$, which is possible due to Assumption 3.1, and noting that the initial distributions $(X_0^n, A_0^n, M_0^n, B_0^n)_{n \in \mathbb{N}}$ are independent of n , we can apply Kolmogorov's tightness criterion to obtain the tightness of the sequence $(\mathbb{P}_{(X^n, A^n, M^n, B)})_{n \in \mathbb{N}}$. Hence, by Prohorov's theorem ([KS91, Theorem 2.4.7]) we get relatively compactness ([KS91, Definition 2.4.6]) of the sequence of measures $(\mathbb{P}_{(X^n, A^n, M^n, B)})_{n \in \mathbb{N}}$ in $\mathcal{M}_1(C([0, T]; \mathbb{R}^4))$, which denotes the space of all probability measures on $C([0, T]; \mathbb{R}^4)$. Consequently, there exists a converging subsequence $(\mathbb{P}_{(X^{n_k}, A^{n_k}, M^{n_k}, B)})_{k \in \mathbb{N}}$ such that

$$\mathbb{P}_{(X^{n_k}, A^{n_k}, M^{n_k}, B)} \rightarrow \mathbb{P}_{(X, A, M, B)} \quad \text{weakly as } k \rightarrow \infty,$$

for some measure $\mathbb{P}_{(X, A, M, B)}$ in $\mathcal{M}_1(C([0, T]; \mathbb{R}^4))$.

The Skorokhod representation theorem (see e.g. [Dud02, Theorem 11.7.2]) yields the existence of some probability space $(\hat{\Omega}, \hat{\mathcal{F}}, \hat{\mathbb{P}})$ with continuous stochastic processes $(\hat{X}^k)_{k \in \mathbb{N}}$, $(\hat{A}^k)_{k \in \mathbb{N}}$, $(\hat{M}^k)_{k \in \mathbb{N}}$, $(\hat{B}^k)_{k \in \mathbb{N}}$ and X , A , M , \hat{B} on it such that

$$(X^{n_k}, A^{n_k}, M^{n_k}, \hat{B}) \stackrel{\mathcal{L}}{\rightarrow} (\hat{X}^k, \hat{A}^k, \hat{M}^k, \hat{B}^k), \quad k \in \mathbb{N},$$

and

$$(\hat{X}^k, \hat{A}^k, \hat{M}^k, \hat{B}^k) \rightarrow (X, A, M, \hat{B}) \quad \text{in } C([0, T]; \mathbb{R}^4) \quad \text{as } k \rightarrow \infty, \quad \hat{\mathbb{P}}\text{-a.s.}$$

From a general version of the Yamada–Watanabe result, see [Kur14, Theorem 1.5], we can deduce that $\hat{M}_t^k = \int_0^t \sigma_{n_k}(s, \hat{X}_s^k) d\hat{B}_s^k$, for $t \in [0, T]$ and for all $k \in \mathbb{N}$, and the stochastic

processes $(B^k)_{k \in \mathbb{N}}$ are Brownian motions as $\hat{B}^k \stackrel{\mathcal{D}}{\sim} B$. Thus, \hat{M}^k is a local $\hat{\mathbb{P}}$ -martingale with quadratic variation $\langle \hat{M}^k \rangle_t = \int_0^t \sigma_{n_k}(s, \hat{X}_s^k)^2 ds$.

Due to the $\hat{\mathbb{P}}$ -a.s. convergence of $(\hat{M}^k)_{k \in \mathbb{N}}$ to M , [JS03, Proposition IX.1.17] implies that M is also a local $\hat{\mathbb{P}}$ -martingale, and the convergence of $\int_0^t \sigma_{n_k}(s, X_s^{n_k}) ds$ in probability, see Lemma 3.7, together with [JS03, Corollary VI.6.29] implies that quadratic variation of M is $\langle M \rangle_t = \int_0^t \sigma(s, X_s)^2 ds$. Therefore, the representation theorem for local martingales with absolutely continuous quadratic variations (see e.g. [KS91, Theorem 3.42]) yields the existence of some probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$, which is an extension of $(\hat{\Omega}, \hat{\mathcal{F}}, \hat{\mathbb{P}})$, and a Brownian motion $(\tilde{B}_t)_{t \in [0, T]}$ on it, such that $M_t = \int_0^t \sigma(s, X_s) d\tilde{B}_s$ for $t \in [0, T]$. \square

Using the stochastic processes X , A and M from Lemma 3.8, we can construct a solution to the Volterra local martingale problem in the sense of Definition 2.4.

Lemma 3.9. *Suppose Assumptions 3.1 and 3.2. There exists a solution to the Volterra local martingale problem given $(x_0, \mu, \sigma, K_\mu, K_\sigma)$.*

Proof. Recall, the stochastic processes $(X^n)_{n \in \mathbb{N}}$, $(A^n)_{n \in \mathbb{N}}$ and $(M^n)_{n \in \mathbb{N}}$ on $(\Omega, \mathcal{F}, \mathbb{P})$ are given in (3.7) and $(\hat{X}^k)_{k \in \mathbb{N}}$, $(\hat{A}^k)_{k \in \mathbb{N}}$, $(\hat{M}^k)_{k \in \mathbb{N}}$, X , A and M on $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$ are given by Lemma 3.8. We introduce the stochastic processes $(Z^n)_{n \in \mathbb{N}}$, $(\hat{Z}^k)_{k \in \mathbb{N}}$ and Z by

$$Z_t^n := A_t^n + M_t^n, \quad \hat{Z}_t^k := \hat{A}_t^k + \hat{M}_t^k \quad \text{and} \quad Z_t := A_t + M_t, \quad t \in [0, T].$$

We shall show that the triple (X, Z) , $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$, $(\mathcal{F}_t^X)_{t \in [0, T]}$, where $(\mathcal{F}_t^X)_{t \in [0, T]}$ denotes the augmented natural filtration of X (cf. [KS91, Definition 2.7.2]), solves the Volterra local martingale problem given $(x_0, \mu, \sigma, K_\mu, K_\sigma)$. Since the properties (i)-(iii) of Definition 2.4 are fairly easy to check, we verify here that

- (iv) the process $(\mathcal{M}_t^f)_{t \in [0, T]}$ defined by (2.4) is a local $\tilde{\mathbb{P}}$ -martingale for every $f \in C_0^2(\mathbb{R})$,
- (v) the equality (2.5) holds $\tilde{\mathbb{P}}$ -a.s.

- (iv) For $k \in \mathbb{N}$ and $f \in C_0^2(\mathbb{R})$, the stochastic process $(\mathcal{M}_t^{f,k})_{t \in [0, T]}$ is defined by

$$\mathcal{M}_t^{f,k} := f(\hat{Z}_t^k) - \int_0^t \mathcal{A}^{f,k}(s, \hat{X}_s^k, \hat{Z}_s^k) ds, \quad t \in [0, T],$$

where $\mathcal{A}^{f,k}(t, x, z) := \mu_{n_k}(t, x) f'(z) + \frac{1}{2} \sigma_{n_k}(t, x)^2 f''(z)$. Due to $(\hat{X}^k, \hat{Z}^k) \stackrel{\mathcal{D}}{\sim} (X^{n_k}, Z^{n_k})$ and since (X^{n_k}, Z^{n_k}) solves the Volterra local martingale problem given $(x_0, \mu_{n_k}, \sigma_{n_k}, K_\mu, K_\sigma)$ on $(\Omega, \mathcal{F}, \mathbb{P})$ by construction and Lemma 2.7, it follows that $(\mathcal{M}_t^{f,k})_{t \in [0, T]}$ is a local martingale on $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$ for every $k \in \mathbb{N}$. Moreover, Lemma 3.7 implies that $\mathcal{M}^{f,k} \rightarrow \mathcal{M}^f$ weakly as $k \rightarrow \infty$ and, thus, by [JS03, Proposition IX.1.17], the limiting process $(\mathcal{M}_t^f)_{t \in [0, T]}$ is a local martingale on $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$.

(v) Since $(\hat{X}^k, \hat{M}^k) \stackrel{\mathcal{D}}{\sim} (X^{n_k}, M^{n_k})$ for every $k \in \mathbb{N}$ and pathwise uniqueness holds for SVEs with Lipschitz continuous coefficients (see e.g. [Wan08, Theorem 1.1]), the general version of the Yamada–Watanabe result ([Kur14, Theorem 1.5]) yields

$$(3.8) \quad \hat{X}_t^k = x_0(t) + \int_0^t K_\mu(s, t) \mu_{n_k}(s, \hat{X}_s^k) ds + \int_0^t K_\sigma(s, t) d\hat{M}_s^k, \quad t \in [0, T], \quad \tilde{\mathbb{P}}\text{-a.s.}$$

To continue the proof of (v), we need to distinguish between (a) bounded kernels and (b) kernels of convolutional type.

(a) We start with the bounded kernels as in Assumption 3.2 (i). Due to the absolute continuity of K_σ in the first variable, we can apply the integration by part formula for semimartingales (see [RW00, Theorem (VI).38.3]) to rewrite (3.8) to

$$(3.9) \quad \hat{X}_t^k = x_0(t) + \int_0^t K_\mu(s, t) \mu_{n_k}(s, \hat{X}_s^k) ds + K_\sigma(t, t) \hat{M}_t^k + \int_0^t \hat{M}_s^k \partial_1 K_\sigma(s, t) ds.$$

Since $(\hat{X}^k, \hat{M}^k) \rightarrow (X, M)$ in $C([0, T]; \mathbb{R}^2)$ as $k \rightarrow \infty$, $\tilde{\mathbb{P}}$ -a.s., and K_σ is bounded, we obtain by Lemma 3.7 that $\hat{X}^k \rightarrow X$ and $K_\sigma \hat{M}^k \rightarrow K_\sigma M$ in $C([0, T]; \mathbb{R})$ as $k \rightarrow \infty$, $\tilde{\mathbb{P}}$ -a.s., and $\int_0^\cdot K_\mu(s, \cdot) \mu_{n_k}(s, \hat{X}_s^k) ds \rightarrow \int_0^\cdot K_\mu(s, \cdot) dA_s$ in $C([0, T]; \mathbb{R})$ in probability as $k \rightarrow \infty$. Furthermore, applying Hölder's inequality with $p > 4$ (see Assumption 3.2) and denoting $q = p/(p-1)$, we get by the integrability of $\partial_1 K_\sigma$ that

$$\begin{aligned} \left\| \int_0^\cdot (\hat{M}_s^k - M_s) \partial_1 K_\sigma(s, \cdot) ds \right\|_\infty &\leq \left(\int_0^T |\hat{M}_s^k - M_s|^q ds \right)^{\frac{1}{q}} \left\| \int_0^\cdot |\partial_1 K_\sigma(s, \cdot)|^p ds \right\|_\infty^{\frac{1}{p}} \\ &\leq C \|\hat{M}^k - M\|_\infty. \end{aligned}$$

Hence, the $\tilde{\mathbb{P}}$ -a.s. convergence $(\hat{M}^k)_{k \in \mathbb{N}}$ to M implies $\int_0^\cdot \hat{M}_s^k K_\sigma(s, \cdot) ds \rightarrow \int_0^\cdot M_s K_\sigma(s, \cdot) ds$ as $k \rightarrow \infty$ $\tilde{\mathbb{P}}$ -a.s., and we can take the limit in probability in (3.9) or the $\tilde{\mathbb{P}}$ -a.s. limit for some subsequence, to obtain that (2.5) holds $\tilde{\mathbb{P}}$ -a.s.

(b) For convolution kernels as in Assumption 3.2 (ii), we integrate both sides of (3.8) and use the stochastic Fubini theorem (see e.g. [Ver12, Theorem 2.2]) twice to obtain

$$\begin{aligned} \int_0^t \hat{X}_s^k ds &= \int_0^t x_0(s) ds + \int_0^t \int_0^s K_\mu(s-u) d\hat{A}_u^k ds + \int_0^t \int_0^s K_\sigma(s-u) d\hat{M}_u^k ds \\ &= \int_0^t x_0(s) ds + \int_0^t \int_u^t K_\mu(s-u) ds d\hat{A}_u^k + \int_0^t \int_u^t K_\sigma(s-u) ds d\hat{M}_u^k \\ &= \int_0^t x_0(s) ds + \int_0^t \int_0^{t-u} K_\mu(s) ds d\hat{A}_u^k + \int_0^t \int_0^{t-u} K_\sigma(s) ds d\hat{M}_u^k \\ &= \int_0^t x_0(s) ds + \int_0^t K_\mu(s) \int_0^{t-s} d\hat{A}_u^k ds + \int_0^t K_\sigma(s) \int_0^{t-s} d\hat{M}_u^k ds \\ (3.10) \quad &= \int_0^t x_0(s) ds + \int_0^t K_\mu(t-s) \hat{A}_s^k ds + \int_0^t K_\sigma(t-s) \hat{M}_s^k ds. \end{aligned}$$

Since

$$\left\| \int_0^\cdot K_\sigma(\cdot - s) (\hat{M}_s^k - M_s) ds \right\|_\infty \leq \|\hat{M}^k - M\|_\infty \int_0^T |K_\sigma(T-s)| ds \leq C \|\hat{M}^k - M\|_\infty$$

and $\hat{M}^k \rightarrow M$ as $k \rightarrow \infty$, $\tilde{\mathbb{P}}$ -a.s., we obtain $\int_0^\cdot K_\sigma(t-s) \hat{M}_s^k ds \rightarrow \int_0^\cdot K_\sigma(t-s) M_s ds$ as $k \rightarrow \infty$, $\tilde{\mathbb{P}}$ -a.s. The analogue observation holds for $\int_0^\cdot K_\mu(t-s) \hat{A}_s^k ds$. Thus, taking the $\tilde{\mathbb{P}}$ -a.s. limit of both sides of (3.10) and then taking the derivative yields that (2.5) holds for (X, Z) , $\tilde{\mathbb{P}}$ -a.s.

If one of K_μ and K_σ is bounded and the other one of convolutional form, one can of course combine the calculations from (a) and (b) to obtain the desired result as well. \square

REFERENCES

- [AJ21] Eduardo Abi Jaber, *Weak existence and uniqueness for affine stochastic Volterra equations with L^1 -kernels*, Bernoulli **27** (2021), no. 3, 1583–1615.

- [AJCLP21] Eduardo Abi Jaber, Christa Cuchiero, Martin Larsson, and Sergio Pulido, *A weak solution theory for stochastic Volterra equations of convolution type*, Ann. Appl. Probab. **31** (2021), no. 6, 2924–2952.
- [AJEE19] Eduardo Abi Jaber and Omar El Euch, *Multifactor approximation of rough volatility models*, SIAM J. Financial Math. **10** (2019), no. 2, 309–349.
- [AJLP19] Eduardo Abi Jaber, Martin Larsson, and Sergio Pulido, *Affine Volterra processes*, Ann. Appl. Probab. **29** (2019), no. 5, 3155–3200.
- [Dud02] R. M. Dudley, *Real analysis and probability*, 2 ed., Cambridge Studies in Advanced Mathematics, Cambridge University Press, 2002.
- [EER19] Omar El Euch and Mathieu Rosenbaum, *The characteristic function of rough Heston models*, Math. Finance **29** (2019), no. 1, 3–38.
- [JP12] Jean Jacod and Philip Protter, *Discretization of processes*, Stochastic Modelling and Applied Probability, vol. 67, Springer, Heidelberg, 2012.
- [JS03] Jean Jacod and Albert N. Shiryaev, *Limit theorems for stochastic processes*, 2nd ed., Springer, 2003.
- [KS91] Ioannis Karatzas and Steven E. Shreve, *Brownian motion and stochastic calculus*, second ed., Graduate Texts in Mathematics, vol. 113, Springer-Verlag, New York, 1991.
- [KS14] Gopinath Kallianpur and P. Sundar, *Stochastic analysis and diffusion processes*, Oxford Graduate Texts in Mathematics, vol. 24, Oxford University Press, Oxford, 2014.
- [Kur14] Thomas G. Kurtz, *Weak and strong solutions of general stochastic models*, Electron. Commun. Probab. **19** (2014), no. 58, 16.
- [MS15] Leonid Mytnik and Thomas S. Salisbury, *Uniqueness for Volterra-type stochastic integral equations*, ArXiv preprint arXiv:1502.05513 (2015).
- [PS22] David Prömel and David Scheffels, *Stochastic Volterra equations with Hölder diffusion coefficients*, ArXiv Preprint arXiv:2204.02648 (2022).
- [RW00] L. C. G. Rogers and David Williams, *Diffusions, Markov processes, and martingales. Vol. 2*, Cambridge Mathematical Library, Cambridge University Press, Cambridge, 2000, Itô calculus, Reprint of the second (1994) edition.
- [RY99] Daniel Revuz and Marc Yor, *Continuous martingales and Brownian motion*, third ed., vol. 293, Springer-Verlag, Berlin, 1999.
- [Sko61] A. V. Skorohod, *On the existence and uniqueness of solutions of stochastic differential equations*, Sibirsk. Mat. Ž. **2** (1961), 129–137.
- [SV79] Daniel W. Stroock and S. R. Srinivasa Varadhan, *Multidimensional diffusion processes*, vol. 233, Springer-Verlag, Berlin-New York, 1979.
- [Ver12] Mark Veraar, *The stochastic Fubini theorem revisited*, Stochastics **84** (2012), no. 4, 543–551.
- [Wan08] Zhidong Wang, *Existence and uniqueness of solutions to stochastic Volterra equations with singular kernels and non-Lipschitz coefficients*, Statist. Probab. Lett. **78** (2008), no. 9, 1062–1071.

DAVID J. PRÖMEL, UNIVERSITY OF MANNHEIM, GERMANY
Email address: proemel@uni-mannheim.de

DAVID SCHEFFELS, UNIVERSITY OF MANNHEIM, GERMANY
Email address: dscheffe@mail.uni-mannheim.de