

Cost-Efficient Fixed-Width Confidence Intervals for the Difference of Two Bernoulli Proportions

Ignacio Erazo^a, David Goldsman^a, Yajun Mei^a

^aH. Milton Stewart School of Industrial and Systems Engineering, Georgia Institute of Technology, Atlanta, GA 30332-0205, U.S.A.

ARTICLE HISTORY

Compiled July 5, 2022

Abstract

We study properties of confidence intervals (CIs) for the difference of two Bernoulli distributions' success parameters, $p_x - p_y$, in the case where the goal is to obtain a CI of a given half-width while minimizing sampling costs when the observation costs may be different between the two distributions. Assuming that we are provided with preliminary estimates of the success parameters, we propose three different methods for constructing fixed-width CIs: (i) a two-stage sampling procedure, (ii) a sequential method that carries out sampling in batches, and (iii) an ℓ -stage "look-ahead" procedure. We use Monte Carlo simulation to show that, under diverse success probability and observation cost scenarios, our proposed algorithms obtain significant cost savings versus their baseline counterparts (up to 50% for the two-stage procedure, up to 15% for the sequential methods). Furthermore, for the battery of scenarios under study, our sequential-batches and ℓ -stage "look-ahead" procedures approximately obtain the nominal coverage while also meeting the desired width requirement. Our sequential-batching method turned out to be more efficient than the "look-ahead" method from a computational standpoint, with average running times at least an order-of-magnitude faster over all the scenarios tested.

KEYWORDS

Confidence intervals; Bernoulli success parameters; two-sample differences; cost-optimization; sequential decision-making; Monte Carlo simulation; Markov decision process.

1. Introduction and Motivation

This article proposes various efficient procedures to obtain confidence intervals (CIs) for the difference $p_x - p_y$ of two Bernoulli distributions' success parameters, p_x and p_y . Our goal is to obtain a CI of a pre-specified width while minimizing sampling costs, where the observation costs may be different between the two distributions. This paper complements Erazo and Goldsman (2021), which assumed

that sampling costs are equal and so instead simply sought to minimize the number of observations taken.

Here are some motivational examples illustrating cases involving cost considerations.

- In many biological or biomedical studies, one is often interested in investigating statistical inference on the risk difference, i.e., the difference in the observed risk of events between experimental and control interventions. It stands to reason that the financial burdens and consequences are different in two groups when considering costs and benefits of implementing different interventions.
- Compare the probability that a person will pass an automated online interview (inexpensive to carry out) with the probability that a person will pass a live interview conducted in-person (requiring more-expensive resources to carry out).
- Undertake real-world experiments to estimate the probability that a manufacturing plant can deliver 85% of its orders on time during a given week. Under scenario one, the plant manufactures the orders in a just-in-time manner and maintains minimal inventory (which is relatively inexpensive). But under scenario two, the plant tends to carry additional inventory in the hope of having sufficient stock on hand for upcoming orders (which is relatively more-expensive).

It is well known that taking numerous observations from both Bernoulli populations will eventually whittle down a CI for $p_x - p_y$ to any desired half-width ϵ . Our interest lies in achieving a small half-width ϵ requirement while (approximately) maintaining the nominal coverage probability of $1 - \alpha$, all the while doing so at the lowest cost — which is not necessarily the same as achieving the nominal coverage and desired half-width using the fewest observations. One population’s observations may simply be significantly more expensive than the other’s; and therefore, we ought to consider sampling schemes giving us more “bang for the buck”.

In this paper, we propose a number of sampling schemes, all of which assume that we have in hand preliminary estimates of p_x and p_y (which may be obtained via stage-one samples from the two populations). In particular, we consider the following sampling schemes:

- A two-stage procedure, presented in Algorithm 2;
- A sequential-sampling procedure carried out in batches, presented in Algorithm 4;
- An ℓ -stage “look-ahead” procedure, based loosely on a Markov decision process (MDP) approach, presented in Algorithm 5.

We also present a conservative benchmark procedure (Algorithm 1) to compare against our two-stage sampling scheme, and a “naive” benchmark procedure (Algorithm 3) to compare against our sequential-sampling scheme. Under a diverse battery of practical scenarios for the values of the success probabilities and observation costs, we use Monte Carlo simulation to evaluate procedure performance; and our results indicate that:

- Our procedures (approximately) obtain the nominal coverage $1 - \alpha$;
- The procedures obtain the desired CI width, except in the case of the two-stage procedure (Algorithm 2) which relies on a heuristic that yields a *random* width that is correct “on average”,

yet exhibits substantial variability.

- The procedures do well in terms of reducing costs, especially when compared to the benchmark procedures or to methods that merely minimize the total number of observations used (Erazo and Goldsman, 2021) — none of which take costs into consideration.

The rest of this paper proceeds as follows. §2 provides additional background material, including relevant notation. In §3, we describe a two-stage procedure (Algorithm 2) that incorporates a cost-minimization strategy. §4 presents a sequential procedure (Algorithm 4) in which observations are taken in batches in order to reduce the number of sampling stages. One might think that a natural direction at this point is to attempt a Markov decision process-based approach to establish an optimal sequential sampling procedure. It turns out that this approach is intractable for reasonably sized problems. A meaningful compromise based on an approximate linear programming strategy is considered in §5, where we introduce sequential CIs based on what we call the ℓ -step look-ahead procedure (Algorithm 5). §6 discusses a Monte Carlo evaluation of our various proposed algorithms over an array of parameter configurations. We establish that our proposed procedures perform as advertised, often resulting in substantial savings over their baseline counterparts and their counterparts that attempt to minimize the total number of observations taken without explicitly considering costs. We summarize our work and provide conclusions in §7.

2. Background

§2.1 spells out the general problem at hand and establishes the necessary notation. We also provide a brief summary of relevant work in the area. §2.2 puts forth a simple straw man conservative procedure that we can use for comparison purposes later on.

2.1. Some Notation and Literature

Going forward, we assume that we have at our disposal a stream X_1, X_2, \dots of independent and identically distributed (iid) $\text{Bern}(p_x)$ random variables from population X as well as an iid $\text{Bern}(p_y)$ stream Y_1, Y_2, \dots from population Y ; and moreover, the X 's and Y 's are independent of each other. This independent-samples set-up is common in practice, and is especially useful in the context of computer simulations, where we have the ability to control the random number streams driving the simulations so as to produce independent replications of any experiments.

As a baseline, we consider the following classical Wald CI for $p_x - p_y$, which is familiar from any elementary statistics textbook (Hines et al., 2003),

$$p_x - p_y \in \bar{X} - \bar{Y} \pm H \equiv \bar{X} - \bar{Y} \pm z_{\alpha/2} \sqrt{\frac{\bar{X}(1 - \bar{X})}{m_x} + \frac{\bar{Y}(1 - \bar{Y})}{m_y}}, \quad (1)$$

where the respective sample sizes are m_x and m_y , the corresponding sample means are given by $\bar{X} \equiv \sum_{i=1}^{m_x} X_i / m_x$ and $\bar{Y} \equiv \sum_{i=1}^{m_y} Y_i / m_y$, $z_{\alpha/2}$ is the $(1 - \frac{\alpha}{2})$ quantile of the standard normal distribution, and H denotes the half-width defined by the right-hand-side expression. As described

in Erazo and Goldsman (2021), variants of and alternatives to the one-dimensional Bern(p) version of the Wald CI are surveyed in e.g., Agresti and Coull (1998), Brown et al. (2001, 2002), Frey (2010), and Turner et al. (2013).

Although the Wald CI suffers from poor small-sample properties (e.g., the coverage probability can substantially deviate from the nominal $1 - \alpha$ for small m_x and m_y), the sample sizes we deal with in the current paper will be substantial enough so that our proposed sequential procedures typically obtain the nominal coverage, at least approximately — a claim that we validate via simulation.

In this paper, we take observation costs into consideration — all observations are not created equal. To this end, let $c_x > 0$ and $c_y > 0$ denote the costs of taking a single observation from populations X and Y , respectively. Our job is to deliver a cost-efficient CI on the difference $p_x - p_y$ that satisfies a half-width constraint $H \leq \epsilon$ for some desired value $\epsilon > 0$, while approximately maintaining the nominal coverage. Such a half-width requirement typically requires more than one stage of sampling to be undertaken. With regard to fixed-width sequential methods for the one-dimensional Bern(p) case, there is a rich literature; see, for instance, Armitage (1958), Khan (1998), Robbins and Siegmund (1974), Yaacoub et al. (2018), and Zacks and Mukhopadhyay (2007), just to name a few.

2.2. Conservative Approach

We start with a very simple conservative procedure. At the beginning of the process of constructing a CI for $p_x - p_y$ with the desired half-width $H \leq \epsilon$, the decision maker may have on hand preliminary estimates \hat{p}_x, \hat{p}_y for p_x, p_y (perhaps obtained from historical data, expert opinion, manufacturer specifications, etc.). These estimates may be quite accurate; however, it is not possible for a single stage of sampling to guarantee the desired half-width requirement $H \leq \epsilon$ for all possible sample paths unless a very conservative approach is used. A common way to address this issue is by noting that the function $p(1 - p)$ is maximized at $p = 0.5$, and then assuming $\hat{p}_x = \hat{p}_y = 0.5$ in order to calculate a deterministic “worst-case” sample size that will always produce the desired half-width. Under that worst-case scenario, Expression (1) yields

$$H \leq \frac{z_{\alpha/2}}{2} \sqrt{\frac{1}{m_x} + \frac{1}{m_y}}.$$

Moreover, given a total sample-size budget of $m_x + m_y = m$, we see that the right-hand side of the above expression is minimized by $m_x = m_y$, in which case

$$H \leq z_{\alpha/2} \sqrt{\frac{1}{2m_x}} \leq \epsilon \Leftrightarrow m_x \geq \frac{z_{\alpha/2}^2}{2\epsilon^2},$$

yielding a lower bound on m_x — depending only on ϵ and α — that guarantees $H \leq \epsilon$. This approach is described in Algorithm 1, and the end result is a conservative CI meeting the half-width requirement after taking $m_x = m_y$ observations from both populations X and Y . The procedure is conservative in the sense that the half-width will likely be somewhat smaller than called for (indicating the waste of some unnecessary observations).

Algorithm 1 Conservative Procedure

Notation: m_x, m_y : common number of observations to take from populations X and Y .

Input: Desired coverage probability $(1 - \alpha)$; desired half-width ϵ .

- 1: $m_x, m_y \leftarrow \left\lceil \frac{z_{\alpha/2}^2}{2\epsilon^2} \right\rceil$, where $\lceil \cdot \rceil$ denotes the ‘‘ceiling’’ (integer round-up) function
- 2: Execute the sampling with m_x and m_y
- 3: Compute \bar{X} and \bar{Y}

Output: Confidence Interval $\bar{X} - \bar{Y} \pm z_{\alpha/2} \sqrt{\frac{\bar{X}(1-\bar{X})}{m_x} + \frac{\bar{Y}(1-\bar{Y})}{m_y}}$

3. Two-Stage Confidence Interval Cost-Minimization Procedure

Henceforth, we assume that we have at our disposal preliminary stage-one estimates \hat{p}_x and \hat{p}_y of the respective parameters p_x and p_y . These estimates could be based on, e.g., expert opinion, prior experience, or actual first-stage samples of sizes $m_{x,1}$ and $m_{y,1}$ taken from populations X and Y . We would now like to conduct a second stage of sampling to obtain an ‘‘inexpensive’’ CI that also satisfies the half-width requirement — or at least *aims* at satisfying the requirement. We also assume that each observation from population X [Y] incurs a cost of $c_x > 0$ [$c_y > 0$]. For ease of exposition, we will first assume for now that \hat{p}_x and \hat{p}_y are derived from expert opinion, allowing us to take $m_{x,1} = m_{y,1} = 0$ as the first-stage sample sizes.

We formulate and solve the following optimization problem in order to determine the total sample sizes $m_{x,2}$ and $m_{y,2}$ to be taken by the time stage two is completed.

$$\min m_{x,2}c_x + m_{y,2}c_y \quad (2)$$

$$s.t. \quad z_{\alpha/2} \sqrt{\frac{\hat{p}_x(1-\hat{p}_x)}{m_{x,2}} + \frac{\hat{p}_y(1-\hat{p}_y)}{m_{y,2}}} \leq \epsilon \quad (3)$$

$$m_{x,2}, m_{y,2} \in \mathbb{Z}^+. \quad (4)$$

Constraint (3) arises from Equation (1) with the preliminary estimates \hat{p}_x and \hat{p}_y substituting for \bar{X} and \bar{Y} , respectively. We let $\tau_x \equiv \hat{p}_x(1-\hat{p}_x)$ and $\tau_y \equiv \hat{p}_y(1-\hat{p}_y)$, so that (3) is equivalent to the condition

$$\frac{\tau_x}{m_{x,2}} + \frac{\tau_y}{m_{y,2}} \leq \frac{\epsilon^2}{z_{\alpha/2}^2}.$$

Moreover, by setting $\zeta_x \equiv \frac{1}{m_{x,2}} > 0$ and $\zeta_y \equiv \frac{1}{m_{y,2}} > 0$ and relaxing the integrality of $m_{x,2}$ and $m_{y,2}$,

we obtain the following equivalent problem.

$$\min \frac{c_x}{\zeta_x} + \frac{c_y}{\zeta_y} \quad (5)$$

$$s.t. \quad \zeta_x \tau_x + \zeta_y \tau_y \leq \frac{\epsilon^2}{z_{\alpha/2}^2} \quad (6)$$

$$\zeta_x, \zeta_y > 0. \quad (7)$$

This is a convex problem, and the Karush-Kuhn-Tucker (KKT) first-order sufficient conditions for optimality (Cheridito and Baes, 2018) ensure the existence of a Lagrange multiplier $\lambda \geq 0$ such that

$$\zeta_x = \sqrt{\frac{c_x}{\lambda \tau_x}}, \quad \zeta_y = \sqrt{\frac{c_y}{\lambda \tau_y}}, \quad \text{and} \quad \zeta_x \tau_x + \zeta_y \tau_y = \frac{\epsilon^2}{z_{\alpha/2}^2}.$$

After a little algebra involving the above three equations, we easily find that

$$\lambda = \left(\frac{\sqrt{\tau_x c_x} + \sqrt{\tau_y c_y}}{\epsilon^2 / z_{\alpha/2}^2} \right)^2. \quad (8)$$

Since $\lambda > 0$, all KKT conditions are met, and so we have an optimal solution for the convex problem. In particular, the optimal ζ_x and ζ_y are obtained by substituting the value of λ from (8). Thus, the approximate solution for the two-stage sampling problem is obtained by setting $m_{x,2} = \lceil 1/\zeta_x \rceil$ and $m_{y,2} = \lceil 1/\zeta_y \rceil$.

We now consider the specific case in which \hat{p}_x and \hat{p}_y are sample means arising from recent (first-stage) samples of sizes $m_{x,1}$ and $m_{y,1}$, respectively. How many new observations do we have to take in order to obtain our desired CI? Our strategy is to solve the optimal- $(m_{x,2}, m_{y,2})$ problem just as before, and we see that there are two outcomes of note:

- (i) If we obtain $m_{x,2} \geq m_{x,1}$ and $m_{y,2} \geq m_{y,1}$, then the numbers of new observations to take in stage two will be $b_x = m_{x,2} - m_{x,1}$ and $b_y = m_{y,2} - m_{y,1}$ for each population, respectively.
- (ii) Suppose that either $m_{x,2} < m_{x,1}$ or $m_{y,2} < m_{y,1}$ (but not both, else we would already have the desired CI). Without loss of generality, we assume only that $m_{x,2} < m_{x,1}$; and then all we need to do is to solve a trivial optimization problem that re-sets $m_{x,2} = m_{x,1}$ and optimizes over $m_{y,2}$,

$$\begin{aligned} \min \quad & m_{y,2} \\ s.t. \quad & z_{\alpha/2} \sqrt{\frac{\tau_x}{m_{x,1}} + \frac{\tau_y}{m_{y,2}}} \leq \epsilon. \end{aligned}$$

This immediately leads to the solution in which we re-set

$$m_{y,2} = \left\lceil \frac{\tau_y}{\frac{\epsilon^2}{z_{\alpha/2}^2} - \frac{\tau_x}{m_{x,1}}} \right\rceil,$$

so that it is optimal to take $b_y = m_{y,2} - m_{y,1}$ second-stage observations from population Y .

The two-stage procedure to obtain the CI while minimizing costs is given by Algorithm 2. We note that it is often the case that the two-stage procedure *does not achieve the length requirement* (see the Monte Carlo discussion in §6). This is because the two-stage procedure only *aims* at yielding a CI of the desired length — there is simply no chance after the second (final) stage of sampling to correct for overshooting the specified length. This issue will be addressed in the subsequent discussion on sequential sampling.

Algorithm 2 Two-Stage Procedure that minimizes the total sampling cost

Notation: $m_{x,2}, m_{y,2}$: cumulative numbers of observations taken from populations X and Y , respectively, by the completion of the second stage. \bar{X}_2, \bar{Y}_2 : overall sample means at the end of the second stage of sampling.

Input: initial sample sizes $m_{x,1}, m_{y,1}$; initial estimates \hat{p}_x and \hat{p}_y ; desired coverage probability $(1 - \alpha)$; desired half-width ϵ , and individual observation costs $c_x > 0, c_y > 0$ from populations X and Y .

- 1: $\tau_x \leftarrow \hat{p}_x(1 - \hat{p}_x), \tau_y \leftarrow \hat{p}_y(1 - \hat{p}_y)$
- 2: $\lambda \leftarrow \left(\frac{\sqrt{\tau_x c_x} + \sqrt{\tau_y c_y}}{\epsilon^2 / z_{\alpha/2}^2} \right)^2$
- 3: $\zeta_x = \sqrt{\frac{c_x}{\lambda \tau_x}}, \zeta_y = \sqrt{\frac{c_y}{\lambda \tau_y}}$
- 4: $m_{x,2} \leftarrow \max(\lceil 1/\zeta_x \rceil, m_{x,1}), m_{y,2} \leftarrow \max(\lceil 1/\zeta_y \rceil, m_{y,1})$
- 5: **if** $m_{x,2} - m_{x,1} == 0$ and $m_{y,2} - m_{y,1} == 0$ **then**
- 6: Do not take samples
- 7: **else if** $m_{x,2} - m_{x,1} == 0$ and $m_{y,2} - m_{y,1} > 0$ **then**
- 8: Re-set $m_{y,2} \leftarrow \left\lceil \frac{\tau_y}{\frac{\epsilon^2}{z_{\alpha/2}^2} - \frac{\tau_x}{m_{x,1}}} \right\rceil$
- 9: Sample $m_{y,2} - m_{y,1}$ observations from Population Y
- 10: **else if** $m_{x,2} - m_{x,1} > 0$ and $m_{y,2} - m_{y,1} == 0$ **then**
- 11: Re-set $m_{x,2} \leftarrow \left\lceil \frac{\tau_x}{\frac{\epsilon^2}{z_{\alpha/2}^2} - \frac{\tau_y}{m_{y,1}}} \right\rceil$
- 12: Sample $m_{x,2} - m_{x,1}$ observations from Population X
- 13: **else**
- 14: Sample $m_{x,2} - m_{x,1}$ and $m_{y,2} - m_{y,1}$ observations from populations X and Y , respectively
- 15: **end if**
- 16: Compute \bar{X}_2 and \bar{Y}_2

Output: Confidence Interval $\bar{X}_2 - \bar{Y}_2 \pm z_{\alpha/2} \sqrt{\frac{\bar{X}_2(1 - \bar{X}_2)}{m_{x,2}} + \frac{\bar{Y}_2(1 - \bar{Y}_2)}{m_{y,2}}}$

4. Sequential-Group Sampling Procedure

In this section, we propose sequential-group sampling procedures. Our motivations for this approach are: (i) we must *guarantee* a CI half-width of some desired length, while approximately attaining the nominal coverage; (ii) we would like to do so at minimal cost; (iii) if time is a consideration, we might wish to conduct sequential sampling in a reasonable number of stages; and (iv) there may be a maximum sampling rate per unit time — say, up to a batch of $B \geq 1$ observations per day, where $B = 1$ indicates the fully sequential case. With these considerations in mind, we develop two sequential CI procedures: one employs what we refer to as the *naive* approach, and the other aims to minimize the total sampling cost.

For the remainder of this section, we use the following notation. Let $m_{x,r}$ and $m_{y,r}$ denote the total number of observations taken from populations X and Y , respectively, up to the end of sampling epoch $r \geq 1$; let \bar{X}_r and \bar{Y}_r be the corresponding sample means; and let $\tau_{x,r} \equiv \bar{X}_r(1 - \bar{X}_r)$ and $\tau_{y,r} \equiv \bar{Y}_r(1 - \bar{Y}_r)$. We first present a naive procedure (Algorithm 3) to carry out sequential batch sampling. Although this approach ensures the desired half-width is eventually attained, it is naive in the sense that it does not use information accumulated as sampling progresses to adapt the proportion of observations to be taken from each population during successive stages — when B is even, the procedure simply takes $B/2$ observations from both populations; and if B is odd, the extra observation is assigned arbitrarily.

Algorithm 3 Naive Sequential Batching Procedure

Notation: $m_{x,r}, m_{y,r}$: cumulative numbers of observations taken from populations X and Y , respectively, at the completion of stage r . \bar{X}_r, \bar{Y}_r : sample means at the completion of stage r .

Input: initial sample sizes $m_{x,1}, m_{y,1}$; desired coverage probability $(1 - \alpha)$; sampling capacity per stage B (assume with little loss of generality that B is even); and desired half-width ϵ

```

1:  $r \leftarrow 2$ , Continue  $\leftarrow True$ 
2: while Continue == True do
3:   Sample  $B/2$  observations from population  $X$  and  $B/2$  from population  $Y$ 
4:    $m_{x,r} \leftarrow m_{x,r-1} + B/2$ ,  $m_{y,r} \leftarrow m_{y,r-1} + B/2$ 
5:   Compute  $\bar{X}_r$  and  $\bar{Y}_r$ 
6:    $\tau_{x,r} \leftarrow \bar{X}_r(1 - \bar{X}_r)$ ,  $\tau_{y,r} \leftarrow \bar{Y}_r(1 - \bar{Y}_r)$ 
7:    $H_r \leftarrow z_{\alpha/2} \sqrt{\frac{\tau_{x,r}}{m_{x,r}} + \frac{\tau_{y,r}}{m_{y,r}}}$ 
8:   if  $H_r \leq \epsilon$  then
9:     Continue  $\leftarrow False$ 
10:  else
11:     $r \leftarrow r + 1$ 
12:  end if
13: end while

```

Output: Confidence Interval $\bar{X}_r - \bar{Y}_r \pm H_r$

Next, we propose a slightly more-elaborate sequential procedure (Algorithm 4) that also incorporates batching to save on the number of stages; but in addition makes use of the new information obtained as sampling proceeds, so as to reduce sampling cost. Motivated by the results from §3, and explicitly

detailed in Algorithm 4, the procedure works as follows:

- (i) Compute the intermediate quantities $\lambda_r, \zeta_{x,r}, \zeta_{y,r}$.
- (ii) Compute the sampling proportion (of observations from population X),

$$\gamma_r \equiv \frac{\max(\lceil 1/\zeta_{x,r} \rceil, m_{x,r-1}) - m_{x,r-1}}{(\max(\lceil 1/\zeta_{x,r} \rceil, m_{x,r-1}) - m_{x,r-1}) + (\max(\lceil 1/\zeta_{y,r} \rceil, m_{y,r-1}) - m_{y,r-1})},$$

where the denominator is guaranteed to be positive (otherwise we would have already achieved the desired CI half-width).

- (iii) The numbers of observations to be taken from populations X and Y in the current stage are $b_{x,r} \equiv \lfloor \gamma_r B \rfloor$ and $b_{y,r} \equiv B - b_{x,r}$, respectively, where $\lfloor \cdot \rfloor$ is the rounding (nearest integer) operator.
- (iv) Then the total numbers of observations taken so far are $m_{x,r} = m_{x,r-1} + \lfloor \gamma_r B \rfloor$ and $m_{y,r} = m_{y,r-1} + B - \lfloor \gamma_r B \rfloor$.
- (v) Repeat these computations for subsequent decision epochs until the desired half-width bound ϵ is obtained.

It is useful to point out that so far we have presented “greedy” algorithms that compute the numbers of observations to be taken from each population in each decision epoch — by approximately solving the integer optimization problem described by (2)–(4), using the cumulative results up to the end of the last epoch as input. Algorithms 2 and 4 require initial estimates \hat{p}_x and \hat{p}_y that are in the open interval $(0, 1)$ (else $\tau_{x,r}$ and/or $\tau_{y,r}$ will be zero, which may render the problem trivial). To avoid that issue, both algorithms can instead use the corresponding *minimax* estimators for p_x and p_y , since those estimates are always in $(0, 1)$ and avoid the need for an initial sample or a guess on the parameters (Sheen and Mei, 2021).

5. Markov Decision Process and ℓ -Stage Look-Ahead Sampling Procedures

We now discuss an optimal finite-horizon Markov decision process formulation that minimizes the expected cost of obtaining the CI having the desired half-width. As in §4, we assume that there is a maximum sampling capacity of batch size B per period. We also introduce the following standard MDP notation and relate this in the context of our fixed-width CI problem.

- $\mathcal{T} = \{1, 2, \dots, T\}$ is the *time horizon*, where T is a very large number that is an upper bound on the number of sampling epochs that will be required.
- $\mathcal{S} = \{(w_x, m_x, w_y, m_y) : 0 < w_x < m_x \leq BT, 0 < w_y < m_y \leq BT\}$ is the *state space*, where w_x, w_y represent the numbers of successes, and m_x, m_y represent the numbers of observations taken (so far) from populations X and Y .
- A_s is the *action space* for $s = (w_x, m_x, w_y, m_y) \in \mathcal{S}$, where

Algorithm 4 Sequential Batching Procedure that minimizes the total sampling cost

Notation: $m_{x,r}, m_{y,r}$: cumulative numbers of observations taken from populations X and Y , respectively, at the completion of stage r . \bar{X}_r, \bar{Y}_r : sample means at the completion of stage r .

Input: initial sample sizes $m_{x,1}, m_{y,1}$; initial estimates \hat{p}_x and \hat{p}_y ; desired coverage probability $(1-\alpha)$; sampling capacity (batch size) B ; desired half-width ϵ ; and individual observation costs $c_x > 0, c_y > 0$ from populations X and Y

- 1: $\tau_{x,1} \leftarrow \hat{p}_x(1 - \hat{p}_x), \tau_{y,1} \leftarrow \hat{p}_y(1 - \hat{p}_y)$
- 2: $r \leftarrow 2, \text{Continue} \leftarrow \text{True}$
- 3: **while** Continue == True **do**
- 4: $\lambda_r \leftarrow \left(\frac{\sqrt{\tau_{x,r-1}c_x} + \sqrt{\tau_{y,r-1}c_y}}{\epsilon^2/z_{\alpha/2}^2} \right)^2$
- 5: $\zeta_{x,r} = \sqrt{\frac{c_x}{\lambda_r \tau_{x,r-1}}}, \zeta_{y,r} = \sqrt{\frac{c_y}{\lambda_r \tau_{y,r-1}}}$
- 6: $\gamma_r = \frac{\max(\lceil 1/\zeta_{x,r} \rceil, m_{x,r-1}) - m_{x,r-1}}{(\max(\lceil 1/\zeta_{x,r} \rceil, m_{x,r-1}) - m_{x,r-1}) + (\max(\lceil 1/\zeta_{y,r} \rceil, m_{y,r-1}) - m_{y,r-1})}$
- 7: Sample $\lfloor \gamma_r B \rfloor$ observations from population X and $B - \lfloor \gamma_r B \rfloor$ from population Y
- 8: $m_{x,r} \leftarrow m_{x,r-1} + \lfloor \gamma_r B \rfloor, m_{y,r} \leftarrow m_{y,r-1} + B - \lfloor \gamma_r B \rfloor$
- 9: Compute \bar{X}_r and \bar{Y}_r
- 10: $\tau_{x,r} \leftarrow \bar{X}_r(1 - \bar{X}_r), \tau_{y,r} \leftarrow \bar{Y}_r(1 - \bar{Y}_r)$
- 11: $H_r \leftarrow z_{\alpha/2} \sqrt{\frac{\tau_{x,r}}{m_{x,r}} + \frac{\tau_{y,r}}{m_{y,r}}}$
- 12: **if** $H_r \leq \epsilon$ **then**
- 13: Continue \leftarrow False
- 14: **else**
- 15: $r \leftarrow r + 1$
- 16: **end if**
- 17: **end while**

Output: Confidence Interval $\bar{X}_r - \bar{Y}_r \pm H_r$

$$A_s = \begin{cases} (0, 0), & \text{if } z_{\alpha/2} \sqrt{\frac{\hat{p}_x(1-\hat{p}_x)}{m_x} + \frac{\hat{p}_y(1-\hat{p}_y)}{m_y}} \leq \epsilon, \text{ for } \hat{p}_x = w_x/m_x \text{ and } \hat{p}_y = w_y/m_y; \\ (b_x, b_y) : b_x + b_y = B, & \text{otherwise.} \end{cases}$$

In other words, no actions are taken if the width of the CI has been achieved; otherwise, we take b_x, b_y observations (a total of B) from populations X and Y , respectively, in the actual decision epoch.

- For current state $s = (w_x, m_x, w_y, m_y) \in \mathcal{S}$ and action $a = (b_x, b_y) \in \mathcal{A}_s$, we let $P_{s,a}$ denote a $(b_x + 1) \times (b_y + 1)$ probability matrix having (k_x, k_y) th element

$$\begin{aligned} P_{s,a}(k_x, k_y) &= P(w_x + k_x, m_x + b_x, w_y + k_y, m_y + b_y | s, a) \\ &= \binom{b_x}{k_x} \hat{p}_x^{k_x} (1 - \hat{p}_x)^{b_x - k_x} \binom{b_y}{k_y} \hat{p}_y^{k_y} (1 - \hat{p}_y)^{b_y - k_y}, \end{aligned}$$

for all $0 \leq k_x \leq b_x$, $0 \leq k_y \leq b_y$; this is simply the product of two binomial probabilities, calculated by assuming \hat{p}_x and \hat{p}_y are the true success probabilities and using the fact that all observations from both populations are independent. It follows that, given action a , the transition probabilities from state s into state $s' = (w'_x, m'_x, w'_y, m'_y)$ are given by:

$$P(s'|s, a) = \begin{cases} P_{s,a}(w'_x - w_x, w'_y - w_y), & \text{for } w_x \leq w'_x \leq w_x + b_x, w_y \leq w'_y \leq w_y + b_y \\ & \text{and } m_x + b_x = m'_x, m_y + b_y = m'_y \\ 0, & \text{otherwise.} \end{cases}$$

- The terminal rewards $r_T(w_x, m_x, w_y, m_y) = -(m_x c_x + m_y c_y)$ represent the cost of the sampling procedure at the last possible decision epoch T , whereas all the other (intermediate) rewards are equal to zero.

By maximizing the expected rewards, we minimize the sampling costs and obtain a policy that tells us (i) how to distribute the B observations in each active sampling day/period, and (ii) when to stop sampling. This is a sequential decision/sampling method — at each decision epoch, the corresponding optimal decision minimizing the expected costs can be determined given the actual realization of the system. We note that this method does not necessarily require an initial sample, as the minimax estimators (Sheen and Mei, 2021) can be used for \hat{p}_x and \hat{p}_y to guarantee that both will be in the interval $(0, 1)$ — the use of minimax estimators would change the definition of the probability matrix slightly, where the state space \mathcal{S} would now be $0 \leq w_x \leq m_x \leq BT$, $0 \leq w_y \leq m_y \leq BT$. In any case, the constraint on the action space \mathcal{A}_s can be relaxed to $b_x + b_y \leq B$, which would make sense if only a few observations are needed towards the final decision epochs (perhaps because the actual interval is very close to the desired length).

The issue with this formulation is that it is likely to be intractable, as BT is a potentially large upper bound on the number of observations, and $|\mathcal{S}| = O(B^4 T^4)$. This renders as problematic the computation of the MDP's recursion to obtain the optimal policy; and thus, in practice, we cannot solve this MDP or its dual Linear Programming (LP) version. However, it is possible to implement an algorithm based on the principles of *approximate* LP. To this end, we propose an ℓ -stage *look-ahead* procedure that computes the optimal policy for a state $s = (w_x, m_x, w_y, m_y)$ by considering the next ℓ

decision epochs and yields the smallest expected sampling cost (to obtain the desired CI) along with the policy (for the next ℓ decision epochs) that achieves it. The expected sampling cost is computed over all possible realizations of the observations taken in each stage where a decision is made; and at the final stage (after the ℓ stages, where the sampling may be completed or not), an approximate “cost-to-go” (CTG) function is computed by using Algorithm 2 to determine an approximate number of observations to be taken for each population that is used to compute the expected cost incurred to finish the sampling.

For example, when $\ell = 1$, we compute the expected cost $C(s)$ starting in state s as follows,

$$E_{(b_x, b_y)}[C(s)] = \sum_{s'} P(s'|s, a = (b_x, b_y)) \text{CTG}(s'),$$

where the states s' are all the possible sample states resulting after sampling the new (b_x, b_y) observations in the next stage. The expectation is computed for all (b_x, b_y) such that $b_x + b_y = B$, and $\text{CTG}(s')$ is the cost-to-go of state s' given by the optimal solution obtained from Algorithm 2 applied to state s' .

Similarly, when $\ell = 2$, we have

$$\begin{aligned} E_{(b_x, b_y), (b'_x, b'_y)}[C(s)] &= \sum_{s''} \sum_{s'} P(s'|s, a = (b_x, b_y)) P(s''|s', a = (b'_x, b'_y)) \text{CTG}(s'') \\ &= \sum_{s'} P(s'|s, a = (b_x, b_y)) E_{(b'_x, b'_y)}[C(s')], \end{aligned}$$

where $b_x + b_y = b'_x + b'_y = B$, (b_x, b_y) represents the observations to be taken from populations X , Y respectively, during the next stage, and (b'_x, b'_y) similarly for the subsequent stage. The procedure for $\ell = 1$ is given in Algorithm 5.

Clearly, when the number of look-ahead stages ℓ increases, we will obtain an expected value that becomes closer to the MDP’s optimum expected value because we get increasingly more-accurate CTG functions at the final states of the recursion (as we move to states where the CTG is zero because no more observations are needed, just as the MDP does when looking at complete sample paths). In fact, in the limit (where ℓ is large), this formulation is equivalent to the MDP. Since the number of policies is $O(B)$ at each possible decision point, we see that the total number of expectations to compute is $O(B^\ell)$, which makes it increasingly difficult to find the optimal policy as ℓ increases (as each expectation requires $O(B^{2\ell})$ computations).

6. Experimental Results

This section presents the results of our Monte Carlo experimentation. In particular, the goal of §6.1 is to illustrate the value of the procedures that minimize sampling costs. In order to do so, the procedures proposed in this paper (Algorithms 2 and 4) are benchmarked against their counterpart procedures that minimize the total number of observations taken (see Algorithms 6–8 in Appendix A, reproduced from Erazo and Goldsman (2021)) and the baseline procedures depicted in Algorithms 1 and 3. Table 1 summarizes the purpose of each of the various algorithms discussed in the paper.

Algorithm 5 One-Step Look-Ahead Procedure to minimize total sampling costs

Notation: $m_{x,r}, m_{y,r}$: cumulative numbers of observations taken from populations X and Y , respectively, at the completion of stage r . \bar{X}_r, \bar{Y}_r : sample means at the completion of stage r .

Input: initial sample sizes $m_{x,1}, m_{y,1}$; initial estimates \hat{p}_x and \hat{p}_y ; desired coverage probability $(1-\alpha)$; sampling capacity B ; desired half-width ϵ ; and individual observation costs $c_x > 0, c_y > 0$ from populations X and Y .

```

1:  $\tau_{x,1} \leftarrow \hat{p}_x(1 - \hat{p}_x), \tau_{y,1} \leftarrow \hat{p}_y(1 - \hat{p}_y)$ 
2:  $r \leftarrow 2, \text{Continue} \leftarrow \text{True}$ 
3: while Continue == True do
4:   for all policies  $(b_x, b_y)$  with  $b_x + b_y = B$  do
5:      $E_{(b_x, b_y)} \leftarrow 0$ 
6:     for all pairs  $(i, j)$  with  $0 \leq i \leq b_x$  and  $0 \leq j \leq b_y$  do
7:       Execute Algorithm 2 with inputs:  $m_{x,r-1}, m_{y,r-1}, \bar{X}_{r-1}, \bar{Y}_{r-1}, (1 - \alpha), \epsilon, c_x, c_y$ 
8:       Compute from step 7's output the sampling costs, denoted CTG, of the second stage,
9:        $E_{(b_x, b_y)} \leftarrow E_{(b_x, b_y)} + \text{CTG} \cdot \binom{b_x}{i} \bar{X}_{r-1}^i (1 - \bar{X}_{r-1})^{b_x-i} \binom{b_y}{j} \bar{Y}_{r-1}^j (1 - \bar{Y}_{r-1})^{b_y-j}$ 
10:    end for
11:  end for
12:   $(b_x^*, b_y^*) \leftarrow \text{argmin}_{(b_x, b_y)} E_{(b_x, b_y)}$ 
13:  Sample  $b_x^*, b_y^*$  observations from populations  $X$  and  $Y$ , respectively
14:   $m_{x,r} \leftarrow m_{x,r-1} + b_x^*, m_{y,r} \leftarrow m_{y,r-1} + b_y^*$ 
15:  Compute  $\bar{X}_r$  and  $\bar{Y}_r$ 
16:   $\tau_{x,r} \leftarrow \bar{X}_r(1 - \bar{X}_r), \tau_{y,r} \leftarrow \bar{Y}_r(1 - \bar{Y}_r)$ 
17:   $H_r \leftarrow z_{\alpha/2} \sqrt{\frac{\tau_{x,r}}{m_{x,r}} + \frac{\tau_{y,r}}{m_{y,r}}}$ 
18:  if  $H_r \leq \epsilon$  then
19:    Continue  $\leftarrow \text{False}$ 
20:  else
21:     $r \leftarrow r + 1$ 
22:  end if
23: end while
Output: Confidence Interval  $\bar{X}_r - \bar{Y}_r \pm H_r$ 

```

Algorithm	Description
1	Conservative two-stage procedure
2	Two-stage procedure to minimize cost
3	Naive sequential procedure to minimize cost
4	Batched sequential procedure to minimize cost
5	One-step look-ahead procedure to minimize cost
6	Two-stage procedure to minimize number of observations
7	Fully sequential procedure to minimize number of observations
8	Batched sequential procedure to minimize number of observations

Table 1.: List of algorithms (Algorithms 6–8 are given in the Appendix).

The comparisons are carried out under the three following sampling schemes (within a particular scheme below, the algorithms are arranged in the order: benchmark algorithm vs. min. observations algorithm vs. min. cost algorithm):

- two-stage (Algorithm 1 vs. Algorithm 6 vs Algorithm 2);
- fully sequential (Algorithm 3 vs. Algorithm 7 vs. Algorithm 4); and
- batched sampling with $B = 10$ (Algorithm 3 vs. Algorithm 8 vs. Algorithm 4).

§6.2 discusses the differences between the proposed cost-saving procedures under the different sampling schemes (two-stage, fully sequential, and batched) and the one-step look-ahead procedure defined in §5.

6.1. Cost-Savings Procedures vs. Other Procedures under Different Sampling Schemes

Tables 3, 4, and 5 compare the cost-saving procedures versus the procedures from Erazo and Goldsman (2021) that are presented in Appendix A and the baseline procedures from Algorithms 1 and 3, for each of the sampling schemes (two-stage, fully sequential, and batched). We run nine different experiments/scenarios, where each experiment has $R = 1000$ replications each incorporating initial sample sizes of 50 (used to compute \hat{p}_x and \hat{p}_y) for both populations X and Y . Each experiment is determined by a vector with four values, $(\mathcal{E}, c_x/c_y, p_x, p_y)$, given in column 1 of each table, where \mathcal{E} is the generic designation of the experiments/scenarios that will be used over the discussion. Note that, without loss of generality, we can simply work with the cost ratio c_x/c_y instead of the two individual costs. All experiments use $\epsilon = 0.05$ and $\alpha = 0.05$.

In column 2 of these tables, the label “Coverage Probability (%)” denotes the empirical percentage \hat{c} of CIs that actually covered $p_x - p_y$; this entry has generic standard error $\sqrt{\hat{c}(1 - \hat{c})/R}$. The label “Half-width Achieved (%)” denotes the empirical proportion of replications that actually attained the half-width requirement. Note that Algorithm 1 is a conservative two-stage procedure that *always* attains the desired width, while Algorithms 2 and 6 are two-stage procedures that *do not guarantee* that requirement. All of the other algorithms discussed in these tables are designed to guarantee the half-width requirement. The term “Cost Gap %” is defined/normalized for a generic

procedure as follows. First of all, for replication $i = 1, 2, \dots, R$, define the cost gap for a procedure with respect to the corresponding naive (benchmark) procedure by $\text{Gap}_i(\text{Procedure}) = \text{Cost}_i(\text{Procedure})/\text{Cost}_i(\text{Naive})$. Because Gap_i is a ratio, the overall gap measure reported in the table is the geometric average, i.e., $\text{Cost Gap } \% = (\prod_{i=1}^R \text{Gap}_i)^{1/R}$. Gap calculations are based only on the observations taken outside of the initial estimation stage. We also report the standard deviation of the Gap_i 's taken over all of the R replications, denoted "std.". Column 2 also lists the maximum and minimum gaps from the R replications as performance measures. The performance results for three related algorithms are presented in columns 3–5 of each table. Table 3 has results for three two-stage procedures; Table 4 for three fully sequential procedures; and Table 5 for three procedures employing batched sampling with $B = 10$. Among the three procedures appearing in any table, column 3 represents the baseline algorithm; column 4 the algorithm designed to minimize the number of observations; and column 5 the algorithm designed to minimize costs.

We point out some details about the experiments. Table 2 lists the scenarios on which our Monte Carlo experiments are based.

$(\mathcal{E}, c_x/c_y, p_x, p_y)$	comments
(s1, 1, 0.3, 0.2)	cost-neutral group of scenarios
(s2, 1, 0.5, 0.2)	
(s3, 1, 0.5, 0.5)	
(s4, $\frac{1}{3}$, 0.3, 0.2)	low cost associated with higher p_x
(s5, $\frac{1}{3}$, 0.5, 0.2)	
(s6, $\frac{1}{3}$, 0.5, 0.5)	
(s7, 5, 0.3, 0.2)	high cost associated with higher p_x
(s8, 5, 0.5, 0.2)	
(s9, 5, 0.5, 0.5)	

Table 2.: List of scenarios

- The scenarios can be divided into the following three groups: (s1,s2,s3), (s4,s5,s6), and (s7,s8,s9). The three scenarios within a particular group share the *same cost ratio* c_x/c_y but have different Bernoulli parameters, namely, $(p_x, p_y) = (0.3, 0.2)$, $(0.5, 0.2)$, and $(0.5, 0.5)$. Note that because of the Bernoulli distribution's symmetry (and the definitions of τ_x and τ_y), it is not necessary to study pairs with p_x or p_y greater than 0.5.
- The scenarios can also be divided into a different collection of three groups: (s1,s4,s7), (s2,s5,s8), and (s3,s6,s9). The three scenarios comprising each of these groups share the same (p_x, p_y) -values for the Bernoulli parameters, but sampling costs differ for each of those three scenarios. The first scenario in each group has a cost ratio of 1 (i.e., equal costs, $c_x = c_y$), while the other two scenarios within a group have cost ratios of $\frac{1}{3}$ and 5, respectively.

We dive into the results along with their interpretations:

- When $c_x = c_y$ (scenarios s1, s2, s3), it is obvious that minimizing the number of observations taken is equivalent to minimizing the total cost of the sampling scheme (cf. Equation (2), in which

case we minimize $m_{x,2} + m_{y,2}$). On the other hand, if $p_x = p_y$ (scenarios s3, s6, s9), then sampling from both populations at the same rate actually minimizes the total number of observations taken, which follows from the symmetry of Equation (3) and the fact that minimizing the number of observations taken is equivalent to considering an objective function of $m_{x,2} + m_{y,2}$; thus, for cases (s3, s6, s9) the naive sequential method from Algorithm 3 and the sequential algorithms from Appendix A are equivalent. When both conditions $c_x = c_y$ and $p_x = p_y$ are true (scenario s3), then in each of Tables 3, 4, and 5, the three algorithms that are compared in that respective table are all equivalent. These remarks are corroborated experimentally in Tables 3, 4, and 5.

With respect to the results for scenarios s1 and s2 under the two-stage sampling scheme, the cost is reduced by over 20%. Furthermore, for scenarios s1 and s2 under the sequential sampling schemes, the methods that minimize the number of observations taken (fourth column) and the methods that minimize the total cost (fifth column) have significantly smaller costs than the “naive” sequential method (Algorithm 3), even though the differences in the gap between these two methods and the “naive” method are rather minor ($\leq 2\%$).

- Focusing on Table 3, we note that the Coverage Probability and Half-width Achieved % are equal for Algorithms 1 and 6, when we look at the same value for the Bernoulli parameters (i.e., looking at groups (s1,s4,s7), (s2,s5,s8), and (s3,s6,s9)); but this does not hold true for the procedure that minimizes costs (Algorithm 2). This makes sense as Algorithm 1 and Algorithm 6 do not take into account the cost ratio for decision-making, whereas Algorithm 2 does — so that its performance is not completely determined by the Bernoulli parameters alone.

The previous equality results for the coverage probability and half-width are not repeated in Tables 4 and 5. This stems from the fact that the fully sequential and batching schemes are affected by the inherent randomness of the observations and their ordering (they rely on the sample paths obtained), whereas the two-stage sampling scheme does not have this issue.

- It is easy to see from Tables 4 and 5 that in scenario groups (s1,s2,s3) and (s4,s5,s6), the cost gap for the fourth column (procedures minimizing the number of observations taken) and the fifth column (procedures minimizing total costs) increases when p_x and p_y are farther apart (i.e., the gap of s6 > gap s4 > gap s5), whereas the opposite is true for the group (s7,s8,s9) (where gap s9 < gap s7 < gap s8). This suggests that when the population with the higher sampling cost has a smaller [larger] success probability than the other population, then more discrepancy increases [reduces] the gap. Such behavior with respect to geometric gaps does not occur in Table 3 because the benchmark is versus the conservative approach that does not vary on costs; and so the gap is only reduced when p_x and p_y get closer to 0.5 (and the number of observations taken approaches what happens in the conservative approach).
- With respect to the geometric gaps, we notice that in all the tables of this section, the procedures minimizing costs (column 5) induce a significantly smaller gap compared to the baseline methods (column 3) for all scenarios, except for the one scenario where they are equivalent (s3, under sequential sampling schemes); and a significantly smaller gap than the procedure that minimizes the number of observations taken for all scenarios except for the ones where costs are equivalent (s1,s2,s3). In general, when looking at Tables 4 and 5 the gaps are very similar within equal scenarios (e.g., comparing scenario s5 results in both tables), and an improvement of up to around 15% can be obtained vs. the naive benchmark procedure (Algorithm 3). With respect

to the procedures that minimize observations (column 4), when the population with the higher cost has a smaller [higher] probability parameter, then the cost is significantly smaller [somewhat higher] than that of the baseline; and this means that under those circumstances reducing the number of observations has as a trade-off an increase in the overall cost.

- Furthermore, with respect to maximum and minimum gaps, under the two-stage sampling scheme (Table 3), the two-stage procedures to minimize observations taken and to minimize costs (columns 4 and 5, respectively) never have a cost that exceeds that of the conservative approach (column 3); however, those savings also result due to the already discussed inherent problematic issue underlying two-stage procedures: most of the time the desired half-width is not obtained, as can be seen in the “Half-width Achieved %” rows.
- When looking solely at the two-stage sampling scheme (Table 3), the overall Coverage Probability and Half-width Achieved % (over all 9 scenarios) of the conservative approach (Algorithm 1), the two-stage procedure that minimizes observations (Algorithm 6), and the two-stage procedure minimizing costs (Algorithm 2) are (94.0, 95.6, 94.8) (compared to the desired 95%) and (100.0, 32.7, 33.0), respectively. This shows that our proposed algorithms can produce a significant increase (standard errors from the 1000 replications are below 0.01) in the coverage probability (which is also backed up by the fact that the proposed two-stage algorithms beat the conservative approach in all of the scenarios), while substantially reducing the sampling costs (column 5 has up to 50% of cost reductions, and a minimum of 2%), at the expense of a decrease in the probability of obtaining the desired half-width.
- When looking at Tables 4 and 5, columns 3, 4, and 5, we obtain the following “overall” coverage probabilities by averaging over the nine scenarios: (94.81, 94.7, 95.1) and (95.2, 95.0, 94.6), which tend to be near the nominal level while also satisfying the desired half-width. This suggests that using the cost-saving proposed algorithms (column 5) significantly reduces the sampling costs while preserving similar coverage probabilities and satisfying the half-width criterion.

In terms of an overall conclusion from these comparisons, it is clear that if the goal is to minimize the sampling costs, the cost-saving procedures proposed over the course of this paper are the most-appropriate choices for their respective sampling-scheme (two-stage, fully sequential, or batches). Further, for the sequential sampling schemes, these procedures induce a similar probability of obtaining the desired CI half-width as well as coverage probabilities near the nominal level.

6.2. Comparison of Procedures that Minimize Cost

Now we move on to evaluate the performance of the different cost-minimization procedures and corresponding sampling schemes (Algorithms 2, 4, and 5). Table 6 gives results to enable a comparison for the same nine scenarios previously described, each for 1000 replications, for the two-stage (Algorithm 2), fully sequential (Algorithm 4 with $B = 10$), batched (Algorithm 4 with $B = 10$), and one-step look-ahead (Algorithm 5 with $B = 10$) procedures. It is easy to see that under the initial samples sizes ($m_{x,1} = m_{y,1} = 50$) the two-stage procedure has the smallest cost, and the gap between the sequential methods and the two-stage method is usually between 2% and 3%. Of course, as already mentioned, the two-stage method does not guarantee the desired half-width (in fact, there is no scenario in our study in which the desired half-width is obtained even 50% of the time). The

$(\mathcal{E}, c_x/c_y, p_x, p_y)$	Performance Measures	Algorithm 1 conservative	Algorithm 6 min. obs.	Algorithm 2 min. cost
(s1, 1, 0.3, 0.2)	Coverage Probability (%)	93.9	94.7	93.9
	Half-width Achieved (%)	100	43.3	44.5
	Cost Gap % (std.)	100 (–)	69.1 (9.3)	69.1 (9.3)
	Max Gap % (Min)	—	92.8 (30.2)	92.8 (30.2)
(s2, 1, 0.5, 0.2)	Coverage Probability (%)	94.1	96.9	95.6
	Half-width Achieved (%)	100	44.3	45.5
	Cost Gap % (std.)	100 (–)	77.2 (8.1)	77.2 (8.1)
	Max Gap % (Min)	—	97.1 (36.9)	97.1 (36.9)
(s3, 1, 0.5, 0.5)	Coverage Probability (%)	93.9	95.2	94.6
	Half-width Achieved (%)	100	10.5	10.2
	Cost Gap % (std.)	100 (–)	97.9 (2.0)	97.9 (2.0)
	Max Gap % (Min)	—	100.0 (83.0)	100.0 (83.0)
(s4, $\frac{1}{3}$, 0.3, 0.2)	Coverage Probability (%)	93.9	94.7	94.5
	Half-width Achieved (%)	100	43.3	44.8
	Cost Gap % (std.)	100 (–)	66.3 (10.1)	61.3 (9.5)
	Max Gap % (Min)	—	93.0 (20.4)	86.3 (18.6)
(s5, $\frac{1}{3}$, 0.5, 0.2)	Coverage Probability (%)	94.1	96.9	95.2
	Half-width Achieved (%)	100	44.3	48.0
	Cost Gap % (std.)	100 (–)	72.3 (9.8)	66.7 (9.2)
	Max Gap % (Min)	—	97.4 (24.5)	90.5 (22.6)
(s6, $\frac{1}{3}$, 0.5, 0.5)	Coverage Probability (%)	93.9	95.2	94.5
	Half-width Achieved (%)	100	10.5	9.3
	Cost Gap % (std.)	100 (–)	97.9 (2.1)	90.9 (2.0)
	Max Gap % (Min)	—	100.0 (79.9)	92.8 (73.9)
(s7, 5, 0.3, 0.2)	Coverage Probability (%)	93.9	94.7	94.4
	Half-width Achieved (%)	100	43.3	44.3
	Cost Gap % (std.)	100 (–)	72.6 (8.9)	63.0 (7.7)
	Max Gap % (Min)	—	93.5 (42.2)	80.9 (35.7)
(s8, 5, 0.5, 0.2)	Coverage Probability (%)	94.1	96.9	95.2
	Half-width Achieved (%)	100	44.3	39.5
	Cost Gap % (std.)	100 (–)	83.8 (6.0)	49.4 (4.8)
	Max Gap % (Min)	—	96.8 (53.3)	83.8 (49.4)
(s9, 5, 0.5, 0.5)	Coverage Probability (%)	93.9	95.2	94.7
	Half-width Achieved (%)	100	10.5	10.8
	Cost Gap % (std.)	100 (–)	97.8 (2.1)	84.5 (1.9)
	Max Gap % (Min)	—	100.0 (85.5)	86.3 (73.2)

Table 3.: Results for two-stage sampling scheme

two-stage procedure also has the lowest coverage probability in six of the nine scenarios (including

$(\mathcal{E}, c_x/c_y, p_x, p_y)$	Performance Measures	Algorithm 3 naive seq.	Algorithm 7 min. obs.	Algorithm 4 min. cost
(s1, 1, 0.3, 0.2)	Coverage Probability (%)	94.8	93.7	95.1
	Half-width Achieved (%)	100	100	100
	Cost Gap % (std.)	100 (–)	99.1 (5.4)	99.2 (5.4)
	Max Gap % (Min)	—	118.8 (80.5)	113.8 (83.2)
(s2, 1, 0.5, 0.2)	Coverage Probability (%)	95.2	95.8	96.1
	Half-width Achieved (%)	100	100	100
	Cost Gap % (std.)	100 (–)	98.3 (3.8)	98.3 (3.9)
	Max Gap % (Min)	—	110.2 (85.5)	111.9 (85.4)
(s3, 1, 0.5, 0.5)	Coverage Probability (%)	94.5	94.0	95.4
	Half-width Achieved (%)	100	100	100
	Cost Gap % (std.)	100 (–)	100.0 (0.2)	100.0 (0.2)
	Max Gap % (Min)	—	101.0 (98.9)	100.8 (99.2)
(s4, $\frac{1}{3}$, 0.3, 0.2)	Coverage Probability (%)	94.8	94.4	95.1
	Half-width Achieved (%)	100	100	100
	Cost Gap % (std.)	100 (–)	95.3 (5.7)	88.0 (5.6)
	Max Gap % (Min)	—	114.8 (74.8)	105.8 (69.6)
(s5, $\frac{1}{3}$, 0.5, 0.2)	Coverage Probability (%)	95.2	95.6	95.0
	Half-width Achieved (%)	100	100	100
	Cost Gap % (std.)	100 (–)	92.3 (4.3)	85.1 (4.5)
	Max Gap % (Min)	—	105.1 (78.0)	98.4 (70.8)
(s6, $\frac{1}{3}$, 0.5, 0.5)	Coverage Probability (%)	94.5	94.3	94.2
	Half-width Achieved (%)	100	100	100
	Cost Gap % (std.)	100 (–)	100.0 (0.2)	92.8 (0.2)
	Max Gap % (Min)	—	101.1 (98.7)	93.6 (91.7)
(s7, 5, 0.3, 0.2)	Coverage Probability (%)	94.9	94.0	94.6
	Half-width Achieved (%)	100	100	100
	Cost Gap % (std.)	100 (–)	104.1 (5.3)	90.5 (4.8)
	Max Gap % (Min)	—	124.6 (88.0)	109.1 (75.8)
(s8, 5, 0.5, 0.2)	Coverage Probability (%)	95.4	95.5	95.1
	Half-width Achieved (%)	100	100	100
	Cost Gap % (std.)	100 (–)	106.4 (3.3)	93.0 (2.7)
	Max Gap % (Min)	—	118.1 (95.5)	102.8 (85.7)
(s9, 5, 0.5, 0.5)	Coverage Probability (%)	94.3	94.6	95.4
	Half-width Achieved (%)	100	100	100
	Cost Gap % (std.)	100 (–)	100.0 (0.2)	86.4 (0.2)
	Max Gap % (Min)	—	101.1 (98.8)	87.0 (84.9)

Table 4.: Results for fully sequential sampling scheme

all three with large discrepancies between p_x and p_y) and the lowest overall coverage probability (of

$(\mathcal{E}, c_x/c_y, p_x, p_y)$	Performance Measures	Algorithm 3 naive seq.	Algorithm 8 batch min. obs.	Algorithm 4 batch min. cost
(s1, 1, 0.3, 0.2)	Coverage Probability (%)	96.3	95.4	95.1
	Half-width Achieved (%)	100	100	100
	Cost Gap % (std.)	100 (-)	99.3 (5.2)	99.4 (5.2)
	Max Gap % (Min)	—	114.6 (80.2)	114.9 (82.4)
(s2, 1, 0.5, 0.2)	Coverage Probability (%)	93.0	94.6	95.4
	Half-width Achieved (%)	100	100	100
	Cost Gap % (std.)	100 (-)	98.7 (3.8)	98.6 (3.8)
	Max Gap % (Min)	—	109.9 (85.8)	111.0 (86.3)
(s3, 1, 0.5, 0.5)	Coverage Probability (%)	96.2	94.8	94.9
	Half-width Achieved (%)	100	100	100
	Cost Gap % (std.)	100 (-)	100.0 (0.2)	100.0 (0.2)
	Max Gap % (Min)	—	100.7 (98.6)	100.7 (99.3)
(s4, $\frac{1}{3}$, 0.3, 0.2)	Coverage Probability (%)	94.7	94.6	94.5
	Half-width Achieved (%)	100	100	100
	Cost Gap % (std.)	100 (-)	95.7 (5.7)	88.2 (5.6)
	Max Gap % (Min)	—	116.1 (79.5)	112.0 (71.1)
(s5, $\frac{1}{3}$, 0.5, 0.2)	Coverage Probability (%)	94.9	95.9	93.7
	Half-width Achieved (%)	100	100	100
	Cost Gap % (std.)	100 (-)	92.8 (4.2)	85.5 (4.5)
	Max Gap % (Min)	—	104.9 (78.7)	101.8 (70.9)
(s6, $\frac{1}{3}$, 0.5, 0.5)	Coverage Probability (%)	95.9	95.0	94.3
	Half-width Achieved (%)	100	100	100
	Cost Gap % (std.)	100 (-)	100.0 (0.2)	92.8 (0.2)
	Max Gap % (Min)	—	100.8 (99.1)	93.6 (91.3)
(s7, 5, 0.3, 0.2)	Coverage Probability (%)	94.9	94.3	95.3
	Half-width Achieved (%)	100	100	100
	Cost Gap % (std.)	100 (-)	104.2 (4.9)	90.5 (4.7)
	Max Gap % (Min)	—	118.6 (89.2)	106.2 (77.7)
(s8, 5, 0.5, 0.2)	Coverage Probability (%)	95.4	95.3	94.5
	Half-width Achieved (%)	100	100	100
	Cost Gap % (std.)	100 (-)	106.7 (3.4)	93.1 (2.6)
	Max Gap % (Min)	—	117.6 (94.9)	103.1 (85.1)
(s9, 5, 0.5, 0.5)	Coverage Probability (%)	95.3	94.8	93.8
	Half-width Achieved (%)	100	100	100
	Cost Gap % (std.)	100 (-)	100.0 (0.2)	86.2 (0.2)
	Max Gap % (Min)	—	101.4 (99.0)	87.4 (84.8)

Table 5.: Results for batch sampling scheme ($B = 10$)

93.8 over the nine scenarios, which is well below the specification). It is also easy to see that the

fully sequential algorithm outperforms the other two in all scenarios on average by a small fraction, but at the expense on requiring approximately 10 times as many decision epochs. Both the batching method and the one-step-ahead sampling scheme have the same cost and need approximately the same number of decision epochs. However, the batching method has a more-consistent coverage probability (overall 95.1, slightly greater than nominal) and is considerably easier to compute (at least 500 times faster); and as such should be preferred over the one-step-ahead sampling scheme (the performance of a higher-order-step-ahead procedure could be studied in the future). As a recap, moderate batching almost always results in coverages, half-widths, and costs that are better than or at least more-or-less the equal to any of the methods — but is faster in terms of the number of stages and/or computation time required compared to other competitive methods (e.g., fully sequential or one-step look-ahead).

$(\mathcal{E}, c_x/c_y, p_x, p_y)$	Performance Measures	Algorithm 2 Two-stage	Algorithm 4 Fully seq.	Algorithm 4 Batches	Algorithm 5 One-step ahead
(s1, 1, 0.3, 0.2)	Coverage Probability (%)	93.6	94.5	95.5	94.2
	Half-width Achieved (%)	43.0	100	100	100
	Cost Gap % (std.)	100 (-)	102.7 (15.0)	103.0 (14.7)	103.2 (15.0)
	Observations taken	1002.5	1029.6	1032.5	1034.8
	Running Time (s)	0.00	0.09	0.01	6.42
(s2, 1, 0.5, 0.2)	Coverage Probability (%)	93.8	94.6	94.4	92.6
	Half-width Achieved (%)	44.7	100	100	100
	Cost Gap % (std.)	100 (-)	102.3 (11.4)	102.5 (11.7)	102.5 (11.6)
	Observations taken	1117.0	1142.3	1144.8	1145.1
(s3, 1, 0.5, 0.5)	Coverage Probability (%)	93.7	94.5	96.7	95.4
	Half-width Achieved (%)	11.3	100	100	100
	Cost Gap % (std.)	100 (-)	101.7 (2.2)	102.3 (2.2)	102.3 (2.2)
	Observations taken	1407.7	1435.2	1439.7	1439.8
(s4, $\frac{1}{3}$, 0.3, 0.2)	Coverage Probability (%)	92.3	94.4	94.5	94.6
	Half-width Achieved (%)	45.1	100	100	100
	Cost Gap % (std.)	100 (-)	102.4 (18.2)	102.9 (18.3)	102.2 (18.2)
	Observations taken	1087.0	1114.6	1119.6	1101.4
	Running Time (s)	0.00	0.10	0.01	6.84
(s5, $\frac{1}{3}$, 0.5, 0.2)	Coverage Probability (%)	94.5	95.1	95.6	93.2
	Half-width Achieved (%)	46.3	100	100	100
	Cost Gap % (std.)	100 (-)	102.1 (16.2)	102.3 (16.4)	102.2 (16.2)
	Observations taken	1209.7	1235.7	1237.8	1265.8
(s6, $\frac{1}{3}$, 0.5, 0.5)	Coverage Probability (%)	94.7	94.4	94.5	94.6
	Half-width Achieved (%)	9.8	100	100	100
	Cost Gap % (std.)	100 (-)	101.9 (2.3)	102.3 (2.4)	102.2 (2.4)
	Observations taken	1524.1	1553.5	1559.1	1572.0
(s7, 5, 0.3, 0.2)	Coverage Probability (%)	93.5	95.1	94.5	95.4
	Half-width Achieved (%)	43.3	100	100	100
	Cost Gap % (std.)	100 (-)	102.6 (13.4)	103.0 (13.1)	103.0 (13.5)
	Observations taken	1189.1	1221.0	1223.8	1203.8
	Running Time (s)	0.00	0.11	0.01	7.54
(s8, 5, 0.5, 0.2)	Coverage Probability (%)	93.7	94.5	94.8	95.7
	Half-width Achieved (%)	39.6	100	100	100
	Cost Gap % (std.)	100 (-)	102.3 (7.0)	102.6 (7.0)	102.4 (7.0)
	Observations taken	1321.6	1352.6	1356.3	1333.9
(s9, 5, 0.5, 0.5)	Coverage Probability (%)	94.1	95.9	95.4	94.5
	Half-width Achieved (%)	11.2	100	100	100
	Cost Gap % (std.)	100 (-)	102.0 (2.4)	102.1 (2.4)	102.2 (2.4)
	Observations taken	1664.8	1697.6	1699.7	1734.0

Table 6.: Comparison of results for the proposed cost-minimization procedures

7. Summary and Conclusions

This paper is concerned with the efficient computation of confidence intervals for the difference between parameters from two Bernoulli populations, where efficiency is defined in terms of sampling costs. As such, we proposed four different procedures to minimize the total cost of the sampling, depending on the type of sampling scheme. One procedure is undertaken in two stages, whereas the others carry out several stages and adjust their policies according to the real-time sampling results. We showed that our two-stage cost-minimization procedure (Algorithm 2) can result in savings of up to 50% vs. the conservative approach depicted in Algorithm 1 and up to 14% vs. the counterpart two-stage procedure that minimizes the number of observations taken (Algorithm 6). Furthermore, the proposed sequential methods save up to 15% in costs compared to the baseline “naive” algorithm (Algorithm 3) and up to 14% vs. their counterparts minimizing the number of total observations. In addition to the cost savings, our proposed batching-based sampling scheme (Algorithm 4) always obtains the desired half-width, while approximately attaining the specified coverage. Further, the batching scheme reduces the number of decision epochs (compared to fully sequential methods) at the small price of just an extra 2–3% of costs vs. the two-stage cost-saving procedure (that is *not guaranteed* to actually yield the desired half-width).

Our one-step look-ahead methodology produces similar results, but is much more computationally expensive, so it makes sense to use the batching approach; however, future research will be undertaken in order to study the potential advantage of adding more steps in the relevant expected value computations. Future research may also include testing the performance of our algorithms on borderline scenarios (cases in which p_x and/or p_y are very close to 0 or 1, e.g., rare-event simulation) or considering a guarantee on the coverage probability (for example, worst-case analysis or a weighted average across a large set of scenarios).

References

- Agresti, A. and Coull, B. A. (1998). Approximate is Better than “Exact” for Interval Estimation of Binomial Proportions, *American Statistician* 52: 119–126.
- Armitage, P. (1958). Numerical Studies in the Sequential Estimation of a Binomial Parameter, *Biometrika* 45: 1–15.
- Brown, L. D., Cai, T., and Dasgupta, A. (2001). Interval Estimation for a Binomial Proportion, *Statistical Science* 16: 101–133.
- Brown, L. D., Cai, T., and Dasgupta, A. (2002). Interval Estimation for a Binomial Proportion and Asymptotic Expansions, *Annals of Statistics* 30: 160–201.
- Cheridito, P. and Baes M (2018). KKT Conditions and Applications, accessible at <https://people.math.ethz.ch/~patrickc/co/CO18L06.pdf>.
- Erazo, I., and Goldsman, D. (2021). Efficient Confidence Intervals for the Difference of Two Bernoulli Distributions’ Success Parameters, *Journal of Simulation*, doi: 10.1080/17477778.2021.1955629
- Frey, J. (2010). Fixed-Width Sequential Confidence Intervals for a Proportion, *American Statistician* 64: 242–249.
- Hines, W. W., Montgomery, D. C., Goldsman, D., and Borror, C. M. (2003). *Probability and Statistics in Engineering*, 4th Edition, Hoboken, NJ: John Wiley and Sons.
- Khan, R. A. (1998). Fixed Width Confidence Sequences for the Normal Mean and the Binomial Probability, *Sequential Analysis* 17: 205–217.
- Robbins, H. and Siegmund, D. (1974). Sequential Estimation of p in Bernoulli trials, in *Studies in Probability and Statistics*, E. J. G. Pitman and E. J. Williams, eds., pp. 103–107, Jerusalem: Jerusalem Academic Press.
- Sheen, H. and Mei, Y. (2021). An Elementary Approach for Minimax Estimation of Bernoulli Proportion in the Restricted Parameter Space *ArXiv: Statistics Theory*.
- Turner, A. J., Balestrini-Robinson, S., and Mavris, D. (2013). Heuristics for the Regression of Stochastic Simulations, *Journal of Simulation* 7(4): 229–239.
- Yaacoub, T., Moustakides, G. V., and Mei, Y. (2018). Optimal Stopping for Interval Estimation in Bernoulli Trials, *IEEE Transactions on Information Theory*, doi: 10.1109/TIT.2018.2885405
- Zacks, S. and Mukhopadhyay, N. (2007). Distributions of Sequential and Two-Stage Stopping Times for Fixed-Width Confidence Intervals in Bernoulli Trials: Application in Reliability, *Sequential Analysis* 26: 425–441.

Appendix A. Benchmarking Algorithms from Erazo and Goldsman (2021)

This appendix presents the implementations used by several algorithms from Erazo and Goldsman (2021) to benchmark the algorithms proposed in this paper.

Algorithm 6 Two-Stage Procedure to minimize the number of observations taken

Notation: $m_{x,2}, m_{y,2}$: total numbers of observations to take from populations X and Y , respectively.

Input: initial sample sizes $m_{x,1}, m_{y,1} > 0$, initial sample means \bar{X}_1, \bar{Y}_1 , desired coverage probability $(1 - \alpha)$, and desired half-width ϵ

$$1: \tau_{x,1} \leftarrow \bar{X}_1(1 - \bar{X}_1), \quad \tau_{y,1} \leftarrow \bar{Y}_1(1 - \bar{Y}_1)$$

$$2: \beta^* \leftarrow \frac{1}{1 + \sqrt{\tau_{y,1}/\tau_{x,1}}}$$

$$3: M^* \leftarrow \frac{z_{\alpha/2}^2 (\sqrt{\tau_{x,1}} + \sqrt{\tau_{y,1}})^2}{\epsilon^2}$$

$$4: m_{x,2} \leftarrow \max(\lceil \beta^* M^* \rceil, m_{x,1}), \quad m_{y,2} \leftarrow \max(\lceil (1 - \beta^*) M^* \rceil, m_{y,1})$$

5: Sample $m_{x,2} - m_{x,1}$ observations from population X and $m_{y,2} - m_{y,1}$ from population Y

6: Compute overall sample means \bar{X}_2 and \bar{Y}_2

Output: Confidence Interval $\bar{X}_2 - \bar{Y}_2 \pm z_{\alpha/2} \sqrt{\frac{\bar{X}_2(1-\bar{X}_2)}{m_{x,2}} + \frac{\bar{Y}_2(1-\bar{Y}_2)}{m_{y,2}}}$

Algorithm 7 Fully Sequential Procedure to minimize the number of observations taken

Notation: $m_{x,r}, m_{y,r}$: cumulative numbers of observations taken from populations X and Y by the end of stage r , respectively. \bar{X}_r, \bar{Y}_r : sample means at the completion of stage r for populations X and Y .

Input: initial sample sizes $m_{x,1}, m_{y,1}$, initial sample means \bar{X}_1, \bar{Y}_1 , desired coverage probability $(1 - \alpha)$, and desired half-width ϵ

1: $r \leftarrow 2$, Continue $\leftarrow True$

2: **while** Continue == *True* **do**

$$3: \quad \tau_{x,r-1} = \bar{X}_{r-1}(1 - \bar{X}_{r-1}), \quad \tau_{y,r-1} = \bar{Y}_{r-1}(1 - \bar{Y}_{r-1})$$

4: **if** $\tau_{x,r-1} \left[\frac{1}{m_{x,r-1}} - \frac{1}{m_{x,r-1}+1} \right] > \tau_{y,r-1} \left[\frac{1}{m_{y,r-1}} - \frac{1}{m_{y,r-1}+1} \right]$ **then**

5: Sample from population X

$$6: \quad m_{x,r} \leftarrow m_{x,r-1} + 1, \quad m_{y,r} \leftarrow m_{y,r-1}$$

7: Compute \bar{X}_r and $\bar{Y}_r \leftarrow \bar{Y}_{r-1}$

8: **else**

9: Sample from population Y

$$10: \quad m_{x,r} \leftarrow m_{x,r-1}, \quad m_{y,r} \leftarrow m_{y,r-1} + 1$$

11: $\bar{X}_r \leftarrow \bar{X}_{r-1}$ and compute \bar{Y}_r

12: **end if**

$$13: \quad H_r \leftarrow z_{\alpha/2} \sqrt{\frac{\bar{X}_r(1 - \bar{X}_r)}{m_{x,r}} + \frac{\bar{Y}_r(1 - \bar{Y}_r)}{m_{y,r}}}$$

14: **if** $H_r \leq \epsilon$ **then**

15: Continue $\leftarrow False$

16: **else**

$$17: \quad r \leftarrow r + 1$$

18: **end if**

19: **end while**

Output: Confidence Interval $\bar{X}_r - \bar{Y}_r \pm H_r$

Algorithm 8 Sequential Procedure with Batching to minimize the number of observations taken

Notation: $m_{x,r}, m_{y,r}$: cumulative numbers of observations taken from populations X and Y by the end of stage r , respectively. \bar{X}_r, \bar{Y}_r : sample means at the completion of stage r for populations X and Y .

Input: initial sample sizes $m_{x,1}, m_{y,1}$, initial sample means \bar{X}_1, \bar{Y}_1 , batch size B , desired coverage probability $(1 - \alpha)$, and desired half-width ϵ

```

1:  $\tau_{x,1} \leftarrow \bar{X}_1(1 - \bar{X}_1), \tau_{y,1} \leftarrow \bar{Y}_1(1 - \bar{Y}_1)$ 
2:  $r \leftarrow 2, \text{Continue} \leftarrow \text{True}$ 
3: while Continue == True do
4:    $\gamma_r^* \leftarrow \frac{\sqrt{\tau_{x,r-1}}(m_{y,r-1} + B) - \sqrt{\tau_{y,r-1}}m_{x,r-1}}{B(\sqrt{\tau_{x,r-1}} + \sqrt{\tau_{y,r-1}})}$ 
5:   if  $\gamma_r^* < 0$  then
6:      $\gamma_r^* \leftarrow 0$ 
7:   end if
8:   if  $\gamma_r^* > 0$  then
9:      $\gamma_r^* \leftarrow 1$ 
10:  end if
11:  Sample  $\lfloor \gamma_r^* B \rfloor$  observations from population  $X$  and  $\lfloor (1 - \gamma_r^*) B \rfloor$  observations from  $Y$ 
12:   $m_{x,r} \leftarrow m_{x,r-1} + \lfloor \gamma_r^* B \rfloor, m_{y,r} \leftarrow m_{y,r-1} + \lfloor (1 - \gamma_r^*) B \rfloor$ 
13:  Compute sample means  $\bar{X}_r$  and  $\bar{Y}_r$ 
14:   $\tau_{x,r} \leftarrow \bar{X}_r(1 - \bar{X}_r), \tau_{y,r} \leftarrow \bar{Y}_r(1 - \bar{Y}_r)$ 
15:   $H_r \leftarrow z_{\alpha/2} \sqrt{\frac{\tau_{x,r}}{m_{x,r}} + \frac{\tau_{y,r}}{m_{y,r}}}$ 
16:  if  $H_r \leq \epsilon$  then
17:    Continue  $\leftarrow \text{False}$ 
18:  else
19:     $r \leftarrow r + 1$ 
20:  end if
21: end while

```

Output: Confidence Interval $\bar{X}_r - \bar{Y}_r \pm H_r$
