A NOTE ON IMPROVED DIFFERENTIABILITY FOR THE BANACH-SPACE VALUED FINSLER γ -LAPLACIAN

MAX GOERING AND LUKAS KOCH

MPI for Mathematics in the Sciences, Inselstrasse 22, 04177 Leipzig, Germany

ABSTRACT. We obtain improved fractional differentiability of solutions to the Banach-space valued Finsler γ -Laplacian defined on a σ -convex, τ -smooth Banach space. The operators we consider are non-linear and very degenerately elliptic. Our results are new already in the \mathbb{R} -valued setting.

1. Introduction

Let $\Omega \subset \mathbb{R}^n$ be an open, bounded domain and let (\mathbb{X}, ρ) be a σ -convex, τ -smooth Banach space. Consider $u \colon \Omega \to \mathbb{X}$ and let F be a norm on $\mathbb{X} \otimes \mathbb{R}^n$. We study solutions of the problem

(1.1)
$$\int_{\Omega} \langle F(\mathrm{D}u)^{\gamma-1}(\mathrm{D}F)(\mathrm{D}u), \mathrm{D}\varphi \rangle_{\mathbb{X} \otimes \mathbb{R}^n} = \int_{\Omega} \langle f, \varphi \rangle_{\mathbb{X}} \quad \forall \varphi \in \mathrm{W}_0^{1,\gamma'}(\Omega; \mathbb{X}),$$

for sufficiently regular f. We refer to (1.1) as the Banach-space valued Finsler γ -Laplacian. We refer the reader to Section 2 for precise definitions and notation.

Given a norm ρ on \mathbb{R}^n the (scalar) Finsler Laplacian

(1.2)
$$\int_{\Omega} \langle \rho(\mathrm{D}u)(\mathrm{D}\rho)(\mathrm{D}u), \mathrm{D}\varphi \rangle = 0 \qquad \forall \varphi \in \mathrm{W}_{0}^{1,2}(\Omega; \mathbb{R})$$

is closely related to the anisotropic isoperimetric inequality, [BFK03]. Under additional 2nd order ellipticity conditions, the Finsler Laplacian has also been used to study bubbling phenomena for volume constrained minimizers [DMMN18]. Some qualitative regularity for the Finsler Laplacian was first shown in [FK09], where no quantitative regularity or convexity of ρ was assumed.

Recently, extending results of [BFK03], the Finsler γ -Laplacian

(1.3)
$$\int \langle \rho(\mathrm{D}u)^{\gamma-1}(\mathrm{D}\rho)(\mathrm{D}u), \mathrm{D}\varphi \rangle = 0 \qquad \forall \varphi \in \mathrm{W}_0^{1,\gamma'}(\Omega;\mathbb{R})$$

has been studied in [KN08, Xia19, dBL20, KSYZ22] with a focus on spectral properties, due to applications to the isoperimetric inequality and Finslerian analogs of other classical inequalities in Riemannian geometric analysis. The study of the regularity theory of solutions to (1.3) was initiated in [FK09] and substantially extended and strengthened in [Goe20], where a Harnack inequality and other quantitative regularity results for the Finsler γ -Laplacian are obtained without quantitative regularity or convexity of ρ .

E-mail address: {mlgoering, lkoch}@mis.mpg.de.

Herein, we extend the weak formulation of the Finsler γ -Laplacian to functions taking values in a σ -smooth and τ -convex Banach space (\mathbb{X}, ρ) . We obtain additional differentiability of solutions in the scale of Besov spaces. The gain in differentiability depends upon the convexity and smoothness of the Banach space. In the case that the norm is sufficiently smooth and sufficiently convex, we both recover and extend [BI87] to norms beyond the Euclidean norm.

Our method of proof follows difference quotient arguments well-known in the study of solutions to elliptic systems with p-growth or even so-called non-standard growth, see [Giu03] for an introduction. There are, however, two key difficulties.

On the one hand, the Euclidean-norm is 2-smooth and 2-convex. Weakening these assumptions to τ -smoothness and ρ -convexity fundamentally changes the ellipticity. In particular very degenerate ellipticity behaviour is possible (see Remark 1.4). This also changes the precise form of growth bounds of the system (1.1). In fact, in the special case when (1.1) is orthotropic, improved differentiability results, similar to ours, have been attained in [BCS10] and [BLPV17]. In both cases, the ellipticity and growth bounds that were used, heavily exploited the orthotropic nature of the studied equation. We stress that not only do we deal with a much broader class of problems, but our results also suggest new critical thresholds concerning the regularity of solutions. These thresholds depend on the relationship between the smoothness and convexity of ρ as well as the homogeneity γ of the equation.

On the other hand, the special structure of (1.1) allows us to exploit ideas from convex duality theory in order to obtain better results than a straightforward application of the difference quotient techniques would obtain. Duality ideas in the context of the calculus of variations already appear in [ZKO94] and were used in the context of integrands with linear growth in [Ser93]. More recently, such ideas have been employed in the context of linear growth [KK22], standard growth [CMP05, CP06, CKP13], faster than exponential growth [BCM12, BC13], and (p,q)-growth [dFKK22].

As a further technical point, we prove an inequality concerning the so-called V-functional that is heavily used in the study of systems with p-growth, for general norms. Since this result may be of independent interest in the study of Banach-space valued problems in the calculus of variations, we state it here:

Lemma 1.1. Let (X, ρ) be a Banach space. Then, for any $\tau > -1$,

$$\rho\left(\rho(u)^{\tau}u - \rho(v)^{\tau}v\right) \sim \left(\rho(u) + \rho(v)\right)^{\tau}\rho(u - v).$$

The implicit constants depend only on τ *.*

We are now ready to state our main result.

Theorem 1.2. Fix $1 < \gamma < \infty, \tau \in (1,2]$, and $\sigma \in [2,\infty)$. Let (\mathbb{X}, ρ) be a Banach space, $\Omega \subset \mathbb{R}^n$ a domain, and any σ -convex and τ -smooth norm F on $\mathbb{X} \otimes \mathbb{R}^n$ satisfying

$$F(z \otimes \xi) \le \rho(z)|\xi| \quad \forall z \in \mathbb{X} \ \forall \xi \in \mathbb{R}^n.$$

Suppose $f \in L^{\min(\gamma,\tau)-1}(\Omega; \mathbb{X}^*)$.

If $u \in W^{1,\gamma}(\Omega; \mathbb{X})$ solves (1.1), then

(1.4)
$$F(\mathrm{D}u)^{\gamma-1}(\mathrm{D}F)(\mathrm{D}u) \in \mathrm{B}_{\infty,loc}^{\min(\gamma,\tau)-1,\gamma'}(\Omega;(\mathbb{X}^*\otimes\mathbb{R}^n,F_*))$$

and

(1.5)
$$\mathrm{D}u \in \mathrm{B}^{\frac{\min(\gamma,\tau)}{\max(\gamma,\sigma)},\gamma}_{\infty,loc}(\Omega;(\mathbb{X}\otimes\mathbb{R}^n,F)).$$

Moreover,

(1.6)
$$F(\mathrm{D}u)^{\frac{\gamma-\sigma}{\sigma}}\mathrm{D}u \in \mathrm{B}_{\infty,loc}^{\frac{\min(\gamma,\tau)}{\sigma},\sigma}(\Omega;(\mathbb{X}\otimes\mathbb{R}^n,F))$$

and

(1.7)
$$F(\mathrm{D}u)^{\frac{\gamma}{\tau'}}(\mathrm{D}F)(\mathrm{D}u) \in \mathrm{B}_{\infty,loc}^{\frac{\min(\gamma,\tau)}{\tau'},\tau'}(\Omega;(\mathbb{X}^*\otimes\mathbb{R}^n;F_*))$$

We remark that in the very special case of the spatially homogeneous, scalar-valued Finsler γ -Laplacian with the additional assumption that

(1.8)
$$\langle D^2 \rho(\xi) \zeta, \zeta \rangle \in [\lambda, \Lambda]$$

for some $0 < \lambda < \Lambda < \infty$ for all $\xi, \zeta \in \mathbb{R}^n \setminus \{0\}$, we recover classical results of elliptic divergence form equations and systems [BI87],[Uhl77, Lemma 3.1]. To be precise, Theorem 1.2 in this case reduces to the following statement.

Theorem 1.3. Fix $0 < \lambda \le \Lambda < \infty, \gamma \ge 2$, and suppose $\rho \in C^2(\mathbb{R}^m \otimes \mathbb{R}^n \setminus \{0\}; \mathbb{R})$ satisfies (1.8). If $u \in W^{1,\gamma}(\Omega; \mathbb{R}^m)$ solves (1.2) then $\rho(Du)^{\frac{\gamma}{2}}(D\rho)(Du) \in W^{2,2}_{loc}(\Omega; \mathbb{R})$.

Remark 1.4. Due to the name, Finsler γ -Laplacian, it is easy to assume the PDE has some notion of ellipticity or at least degenerate ellipticity. Here, we emphasize that the ellipticity behaviour can be badly behaved, even in relatively simple cases, and when $\gamma = 2$.

Consider $\rho = \|\cdot\|_{\ell^p}$ for some p > 2. Then, the Finsler Laplacian takes the strong form,

$$-div\left(D|_{\nabla}\|\cdot\|_{p}^{2}\right)=0=-div\left(A(x)Du\right),$$

where

$$(A(x))_{j}^{i} = ||Du(x)||_{p}^{2-p} |\partial_{i}u(x)|^{p-2} \delta_{j}^{i}.$$

Thus when $Du(x_0) = \lambda e_i$ for some standard basis direction, it follows $A(x_0)$ has precisely 1-nonzero entry, no matter the size of $\lambda = |Du(x_0)|$.

We close the introduction by providing some examples of Banach spaces (X, F) to which our results apply.

(1) When $\mathbb{X} = \mathbb{R}$ and $F = \rho$, (1.1) recovers the scalar-valued Finsler γ -Example 1.5.

- (2) When $\mathbb{X} = \mathbb{R}^m$ and $F = \|\cdot\|_{\ell^p(\mathbb{R}^m \otimes \mathbb{R}^n)}$, then F is 2-smooth and p-convex whenever $p \ge 2$ or 2-convex and p-smooth whenever $p \le 2$.
- (3) When (X, ρ) is a σ -convex and τ -smooth Banach space, we can identify $X \otimes \mathbb{R}^n$ with the space of "matrices" with n columns, where each column is an elements in \mathbb{X} after some choice of basis. This is analogous to how $\mathbb{R}^m \otimes \mathbb{R}^n$ is often identified with the space of $m \times n$ matrices. Then, if $F : \mathbb{X} \otimes \mathbb{R}^n \to \mathbb{R}$ is defined by

(1.9)
$$F(x_1, \dots, x_n) = \|(\rho(x_1), \dots, \rho(x_n))\|_{\ell^2(\mathbb{R}^n)} \quad \forall x_i \in \mathbb{X}$$

it follows F is a σ -convex, τ -smooth norm. The details behind this example are contained in Proposition 2.9.

2. Preliminaries

Throughout, given any number $p \in (1, \infty)$ we let p' denote its Hölder conjugate.

2.1. **Vector-valued Sobolev and Besov spaces.** We present the relevant theory of vector-valued Sobolev spaces as found in [HvNVW16] or alternatively [Kre15]. All integrals in this section are to be interpreted in the sense of Bochner integration. Recall that Ω is an open domain in \mathbb{R}^n . We introduce L^p -spaces as in the \mathbb{R}^N -valued case.

Definition 2.1. Let $1 \leq p \leq \infty$. We say a measurable function $f: \Omega \to \mathbb{Y}$ lies in $L^p(\Omega; \mathbb{Y})$ if $|f| \in L^p(\Omega; \mathbb{R})$ and define the L^p -norm as $||f||_{L^p(\Omega; \mathbb{Y})} = |||f||_{L^p(\Omega; \mathbb{R})}$.

We remark that, as in the scalar-valued case, we interpret functions in L^p as equivalence classes of functions equal almost everywhere. It is straightforward to prove that $L^p(\Omega)$ is a Banach space. Further we note that Hölder's inequality holds in this set-up.

Theorem 2.2. Suppose
$$f \in L^p(\Omega, \mathbb{Y})$$
 and $g \in L^{p'}(\Omega, \mathbb{Y}')$ where $\frac{1}{p} + \frac{1}{p'} = 1$. Then

$$\|\langle f,g\rangle\|_{\mathrm{L}^1(\Omega;\mathbb{R})} \leq \|f\|_{\mathrm{L}^p(\Omega,\mathbb{Y})} \|g\|_{\mathrm{L}^{p'}(\Omega,\mathbb{Y}')}.$$

Here $\langle \cdot, \cdot \rangle$ denotes the duality pairing of \mathbb{Y} and \mathbb{Y}' .

We next introduce Sobolev and Besov spaces.

Theorem 2.3. Let $1 \le p \le \infty$ and $m \in \mathbb{N}$. The Sobolev space

$$W^{m,p}(\Omega, \mathbb{Y}) = \{ u \in L^p(\Omega) \colon \nabla^{\alpha} u \in L^p(\Omega, \mathbb{Y}), \ \forall \alpha \in \mathbb{N}^n, \ |\alpha| \le m \}$$

equipped with the norm

(2.1)
$$||u||_{W^{m,p}(\Omega,\mathbb{Y})} = \sum_{|\alpha| \le m} ||\nabla^{\alpha} u||_{L^p(\Omega,\mathbb{Y})}$$

is a Banach space.

Let $0 \le \alpha \le 1$. Then we define

$$B^{\alpha,p}_{\infty}(\Omega) = [L^p(\Omega), W^{1,p}(\Omega)]_{\alpha,\infty}$$

where $[\cdot,\cdot]_{s,p}$ denotes the real-interpolation functor.

We will require a characterisation of Sobolev and Besov spaces in terms of difference quotients. It is useful to denote, for a given function u defined on Ω , $\Delta_h u(x) = u(x) - u(x+h)$. Such a characterisation does not hold in general, but rather requires $\mathbb Y$ to be a Banach space satisfying the Radon-Nikodým property.

Theorem 2.4. Let $1 \le p \le \infty$ and $0 \le \alpha \le 1$. Suppose $\mathbb Y$ is a Banach space satisfying the Radon-Nikodým property. Then $u \in \mathcal{B}^{\alpha,p}_{\infty}(\Omega,\mathbb Y)$ if and only if $u \in \mathcal{L}^p(\Omega,\mathbb Y)$ and there is a constant C > 0 such that, for all subsets $\omega \in \Omega$ and $h \in \mathbb{R}^n$ with $|h| \le \operatorname{dist}(\omega,\partial\Omega)$, we have

$$\|\Delta_h u\|_{L^p(\omega)} \le C|h|^{\alpha}.$$

Moreover we may choose $C = ||u||_{\mathbf{B}^{\alpha,p}(\Omega)}$.

Finally, we remark that we will use the results of this section with the choice $\mathbb{Y} = \mathbb{X} \otimes \mathbb{R}^n$. In particular whenever \mathbb{X} is reflexive, so is $\mathbb{X} \otimes \mathbb{R}^n$, see Proposition 2.9, and consequently both $\mathbb{X} \otimes \mathbb{R}^n$ and its dual have the Radon-Nikodým property.

2.2. Banach space geometry. Let (\mathbb{X}, ρ) be a real Banach space and (\mathbb{X}^*, ρ_*) denote its dual. The modulus of convexity

$$\sigma_{\rho}(\epsilon) = \inf \left\{ 1 - \rho \left(\frac{x+y}{2} \right) : x, y \in \{ \rho = 1 \}, \rho(x-y) \ge \epsilon \right\}$$

and the modulus of smoothness

$$\tau_{\rho}(\epsilon) = \frac{1}{2} \sup \{ \rho(x+y) + \rho(x-y) - 2 : x \in \{ \rho = 1 \}, \rho(y) \le \epsilon \}$$

are fundamental in understanding the geometry of \mathbb{X} . For any $\gamma > 1$, under the assumption that X is reflexive, we define the duality mapping

(2.2)
$$j_{\rho}^{\gamma}(x) = \partial \int_{0}^{\rho(x)} t^{\gamma - 1} dt.$$

We recall the sub-gradient on the right is single valued, and hence j_{ρ}^{γ} is a well-defined function, whenever X is uniformly smooth, that is if $\lim_{\epsilon \downarrow 0} \tau_{\rho}(\epsilon) \epsilon^{-1} = 0$. Moreover, j_{ρ}^{γ} is surjective if and only if X is reflexive, see for example [XR91, Properties (J2,J3)]. Finally, we recall that a Banach space is reflexive if and only if it is uniformly smooth or uniformly convex [LT13, Proposition 1.e.3].

For $\tau \in (1,2]$ and $\sigma \in [2,\infty)$, we define a Banach space \mathbb{X} to be σ -convex if

$$\sigma_{\rho}(\epsilon) \gtrsim \epsilon^{\sigma} \quad \forall \epsilon \in [0, 2]$$

and X is called τ -smooth if

$$\tau_{\rho}(\epsilon) \lesssim \epsilon^{\tau} \quad \forall \epsilon \in (0, \infty).$$

We recall

Proposition 2.5. A Banach space (\mathbb{X}, ρ) is σ -convex and τ -smooth if and only if (\mathbb{X}^*, ρ_*) is τ' -convex and σ' -smooth.

It is also well-known that if $\tau \in (1,2]$ and $\sigma \geq 2$ then

(2.3)
$$j_{\rho}^{\gamma} \circ j_{\rho_*}^{\gamma'} = \operatorname{Id}_{\mathbb{X}_*} \quad \text{and} \quad j_{\rho_*}^{\gamma'} \circ j_{\rho}^{\gamma} = \operatorname{Id}_{\mathbb{X}},$$

see for instance [BM20].

Example 2.6. We note the following examples of σ -smooth and τ -convex spaces.

- (1) $\|\cdot\|_{\ell^p}$ is $\max(p,2)$ -convex and $\min(p,2)$ -smooth for $p \in (1,\infty)$.
- (2) $\|\cdot\|_{L^p}$ is $\max(p,2)$ -convex and $\min(p,2)$ -smooth for $p \in (1,\infty)$.
- (3) Note that for $p \in (1, \infty)$, $W^{m,p}(\Omega, \mathbb{Y}^n)$, equipped with the norm (2.1) is not τ -smooth or σ -convex. However, due to the above examples, endowed with the norm

$$||u||_{W^{m,p}(\Omega,\mathbb{Y})} = \left(\sum_{|\alpha| \le m} ||\nabla^{\alpha} u||_{L^{p}(\Omega,\mathbb{Y})}^{2}\right)^{\frac{1}{2}}$$

it is $\max(p, 2)$ -convex and $\min(p, 2)$ -smooth.

We now recall the main results of [XR91] which are relevant to this article, restated in the special case of σ -convex and τ -smooth Banach spaces.

Theorem 2.7. Let (X, ρ) be a Banach space. The following are equivalent:

- (1) ρ is σ -convex
- (2) For all $\gamma > 1$ and every $x, y \in \mathbb{X}$

(2.4)
$$\langle j_{\rho}^{\gamma}(x) - j_{\rho}^{\gamma}(y), x - y \rangle \gtrsim_{\gamma} (\rho(x) + \rho(y))^{\gamma - \sigma} \rho(x - y)^{\sigma}.$$

(3) There exists a $\gamma > 1$ so that (2.4) holds.

Alternatively, the following are equivalent:

- (a) ρ is τ -smooth
- (b) For all $\gamma > 1$ and every $x, y \in \mathbb{X}$

(c) There exists a $\gamma > 1$ so that (2.5) holds.

Remark 2.8. In light of Proposition 2.5 and Theorem 2.7, we note the following dual characterization of σ -convexity and τ -smoothness of (\mathbb{X}, ρ) . Namely, ρ is τ -smooth if and only if for every $\gamma > 1$

(2.6)

$$(\rho_*(x^*) + \rho_*(y^*))^{\gamma' - \tau'} \rho_* (x^* - y^*)^{\tau'} \lesssim_{\gamma} \left\langle j_{\rho_*}^{\gamma'}(x^*) - j_{\rho_*}^{\gamma'}(y^*), x^* - y^* \right\rangle \qquad \forall x^*, y^* \in \mathbb{X}^*,$$

while ρ is σ -smooth if and only if for every $\gamma > 1$,

(2.7)
$$\rho\left(j_{\rho_*}^{\gamma'}(x^*) - j_{\rho^*}^{\gamma'}(y^*)\right) \lesssim (\rho_*(x^*) + \rho_*(y^*))^{\gamma - \sigma'} \rho_* (x^* - y^*)^{\sigma' - 1}.$$

We gather some information about the relationship between Banach spaces (\mathbb{X}, ρ) and $(\mathbb{X} \otimes \mathbb{R}^n, F)$ where F is as in (1.9).

Proposition 2.9. Let (\mathbb{X}, ρ) be a Banach space and define $F : \mathbb{X} \otimes \mathbb{R}^n \to \mathbb{R}$ by (1.9).

- (i) $(\mathbb{X} \otimes \mathbb{R}^n, F)$ is a Banach space and $(\mathbb{X}^* \otimes \mathbb{R}^n, F_*)$ is its dual. Here we have denoted $F_*(\vec{x^*}) = \left(\sum_i \rho_*(x_i^*)^2\right)^{1/2}$.
- (ii) $(\mathbb{X} \otimes \mathbb{R}^n, F)$ is reflexive if and only if (\mathbb{X}, ρ) is reflexive.
- (iii) If ρ is τ -smooth then so is F.
- (iv) If ρ is σ -convex then so is F.
- (v) $F(x \otimes \xi) = \rho(x)|\xi|$.

Proof. (i) follows from the linearity of the tensor product and (ii) follows from (i).

To check (iii), let $\vec{x} = (x_1, \dots, x_n) \in \mathbb{X} \otimes \mathbb{R}^n$ and $\vec{y} = (y_1, \dots, y_n) \in \mathbb{X} \otimes \mathbb{R}^n$ for some $x_i, y_i \in \mathbb{X}$ we observe

$$(2.8) j_F^{\gamma}(\vec{x}) = (j_{\rho}^{\gamma}(x_1), \dots, j_{\rho}^{\gamma}(x_n)) \in \mathbb{X}^* \otimes \mathbb{R}^n.$$

By Theorem 2.7, (iii) follows given any $\gamma > 1$ so (2.5) holds with F, F_* . To this end, we use (i) and (2.8) to compute

$$F_* \left(j_F^{\gamma}(\vec{x}) - j_F^{\gamma}(\vec{y}) \right) = F_* \left((j_\rho^{\gamma}(x_1) - j_\rho^{\gamma}(y_1), \dots, j_\rho^{\gamma}(x_n) - j_\rho^{\gamma}(y_n)) \right)$$
$$= \left(\sum_{i=1}^n \rho_* \left(j_\rho^{\gamma}(x_i) - j_\rho^{\gamma}(y_i) \right)^2 \right)^{1/2}.$$

Now applying the τ -smoothness of ρ , and choosing some $\gamma > \tau$ the above imply

$$F_* \left(j_F^{\gamma}(\vec{x}) - j_F^{\gamma}(\vec{y}) \right) \lesssim \left(\sum_{i=1}^n \left(\rho(x_i - y_i)^{\tau - 1} (\rho(x_i) + \rho(y_i))^{\gamma - \tau} \right)^2 \right)^{1/2}$$

$$\leq \left(F(\vec{x}) + F(\vec{y}) \right)^{\gamma - \tau} \left(\sum_{i=1}^n \rho(x_i - y_i)^{2(\tau - 1)} \right)^{1/2}$$

$$\lesssim_n \left(F(\vec{x}) + F(\vec{y}) \right)^{\gamma - \tau} \left(\sum_{i=1}^n \rho(x_i - y_i)^2 \right)^{\frac{\tau - 1}{2}}$$

$$= \left(F(\vec{x}) + F(\vec{y}) \right)^{\gamma - \tau} F(\vec{x} - \vec{y})^{\tau - 1},$$

where the last inequality follows by Jensen's since $1 \le \frac{1}{\tau - 1} < \infty$. By Theorem 2.7(c) this confirms the τ -smoothness of F.

To prove (iv), note that by Proposition 2.5, ρ is σ -convex if and only if ρ_* is σ' -smooth. Thus by part (i) and (iii) of this Proposition, F_* is σ' -smooth. Another application of Proposition 2.5 implies $(F_*)_* = F$ is σ -convex as desired.

Finally (v) is a straightforward computation, since
$$x \otimes \xi = (\xi_1 x, \dots, \xi_n x)$$
.

2.3. A vector identity in Banach spaces. We provide an elementary proof of a well-known property of V-functionals in the case when $\mathbb{X} = \mathbb{R}^m$ and ρ is the Euclidean norm. For the convenience of the reader, we restate the lemma here.

Lemma 2.10. Let (X, ρ) be a Banach space. Then, for any $\tau > -1$,

(2.9)
$$\rho\left(\rho(u)^{\tau}u - \rho(v)^{\tau}v\right) \sim \left(\rho(u) + \rho(v)\right)^{\tau}\rho(u - v).$$

The implicit constants depend on τ only.

Proof. Case 1: $\tau \geq 0$.

For any $u, v \in \mathbb{X}$, convexity of ρ implies the difference quotient

$$g(t) = \frac{\rho(u + tv) - \rho(u)}{t}$$

is a non-decreasing function for $t \in [0, \infty)$. We assume without loss of generality $\rho(u) \ge \rho(v)$ and define $\kappa = \frac{\rho(v)}{\rho(u)}$. Since $\tau \geq 0$ ensures $\kappa \leq 1$, it follows

$$\begin{split} \rho\left(\rho(u)^{\tau}u - \rho(v)^{\tau}v\right) &= \rho(u)^{\tau}\kappa^{\tau}\frac{\rho(u - \kappa^{\tau}v) - \rho(u)}{\kappa^{\tau}} + \rho(u)^{\tau+1}\\ &\leq \rho(v)^{\tau}\left[\rho(u - v) - \rho(u)\right] + \rho(u)^{\tau+1}\\ &\leq \rho(v)^{\tau}\rho(u - v) + (\rho(u)^{\tau} - \rho(v)^{\tau})\rho(u)\\ &\lesssim (\rho(u) + \rho(v))^{\tau}\rho(u - v), \end{split}$$

where the last line follows from the mean-value theorem applied to $x \mapsto x^{\tau}$ on $[\rho(v), \rho(u)]$.

We now prove the converse in this case. Without loss of generality, we further assume that $\rho(u) > \rho(v)$, since the identity is trivial when $\rho(u) = \rho(v)$. We have

$$\rho\left(\rho(u)^{\tau}u - \rho(v)^{\tau}v\right) = \rho(u)^{\tau}\rho(u - \kappa^{\tau}v).$$

In particular, it suffices to prove for any $u, v \in \mathbb{X}$ with $\rho(v) < \rho(u)$ and $\mu \leq 1$

$$\rho(u-v) \lesssim \rho(u-\mu v).$$

However, by the triangle inequality,

$$\rho(u - \mu v) + \rho((\mu - 1)v) + \rho(\mu v) + \rho(u - \mu v) \ge \rho(u - v) + \rho(u),$$

from which we deduce, using that $\mu \leq 1$ and that $\rho(u) \geq \rho(v)$,

$$\rho(u - \mu v) \ge \frac{1}{2}(\rho(u - v) + \rho(u) - \rho(v)) \ge \frac{1}{2}\rho(u - v).$$

Case 2: $-1 < \tau < 0$

Note in this case $\kappa^{-\tau} \leq 1$. We estimate

$$\rho(\rho(u)^{\tau}u - \rho(v)^{\tau}v) = \rho(v)^{\tau}\kappa^{-\tau} \frac{\rho(\kappa^{\tau}u - v) - \rho(v)}{\kappa^{-\tau}} + \rho(v)^{\tau+1}$$

$$\leq \rho(v)^{\tau}\kappa^{-\tau}(\rho(u - v) - \rho(v)) + \rho(v)^{\tau+1}$$

$$= \rho(u)^{\tau}\rho(u - v) + \rho(v)^{\tau+1} - \rho(u)^{\tau}\rho(v)$$

$$\lesssim (\rho(u) + \rho(v))^{\tau}\rho(u - v),$$

where the last line follows since

$$\rho(v)^{\tau+1} - \rho(u)^{\tau} \rho(v) \le \rho(u)^{\tau+1} - \rho(u)^{\tau} \rho(v) \le \rho(u)^{\tau} \rho(u-v).$$

We now prove the converse and assume without loss of generality $\rho(u) > \rho(v) > 0$, as the claim is trivial if $\rho(v) = \rho(u)$ or 0. For $\lambda > 0$ to be determined at a later point, we consider

$$(2.10) (2+\lambda)\rho(\rho(u)^{\tau}u - \rho(v)^{\tau}v) = (2+\lambda)\rho(u)^{\tau}\rho(v)^{\tau}\rho(\rho(v)^{-\tau}u - \rho(v)^{-\tau}v)$$

By the triangle inequality

$$2\rho(\rho(v)^{-\tau}u - \rho(u)^{-\tau}v) + \rho(u)^{-\tau}\rho(v) + \left(\rho(u)^{-\tau} - \rho(v)^{-\tau}\right)\rho(v) \ge \rho(v)^{-\tau}\rho(u) + \rho(v)^{-\tau}\rho(u - v).$$
 Further, for any $\lambda > 0$,

$$\lambda \rho(\rho(v)^{-\tau}u - \rho(u)^{-\tau}v) \ge \lambda \rho(v)^{-\tau}\rho(u) - \lambda \rho(u)^{-\tau}\rho(v).$$

Adding these estimates, we deduce

$$(2+\lambda)\rho(\rho(v)^{-\tau}u - \rho(v)^{-\tau}v) \ge (1+\lambda)\rho(v)^{-\tau}\rho(u) - (2+\lambda)\rho(u)^{-\tau}\rho(v) + \rho(v)^{-\tau}\rho(v) + \rho(v)^{-\tau}\rho(u-v).$$
(2.11)

Now consider the function

$$f(x) = (1 + \lambda)\rho(u)x^{-\tau - 1} + x^{-\tau}.$$

Note that as $x \to 0$ and as $x \to \infty$, $f(x) \to \infty$. Moreover, $f(\rho(u)) = (2 + \lambda)\rho(u)^{-\tau}$. Further we note

$$f'(x) = 0 \quad \Leftrightarrow \quad x = \frac{1+\tau}{-\tau}(1+\lambda)\rho(u).$$

Thus, if λ is chosen sufficiently large that $\frac{1+\tau}{-\tau}(1+\lambda) \ge 1$, $f(x) \ge (2+\lambda)\rho(u)^{-\tau}$ in $(0,\rho(u)]$. So, for any such λ , $\rho(v)$ $(f(\rho(v)) - f(\rho(u))) \ge 0$. That is,

$$(2.12) (1+\lambda)\rho(v)^{-\tau}\rho(u) - (2+\lambda)\rho(u)^{-\tau}\rho(v) + \rho(v)^{-\tau}\rho(v) \ge 0.$$

Chaining together (2.10), (2.11), and (2.12) yields the desired result

$$(2+\lambda)\rho(\rho(u)^{\tau}u - \rho(v)^{\tau}v) \ge \rho(u)^{\tau}\rho(u-v) \ge (\rho(u) + \rho(v))^{\tau}\rho(u-v).$$

3. STATEMENT AND PROOF OF MAIN THEOREM

We use the notation from Section 2 to re-write Theorem 1.2 in a way that better elucidates the symmetry between convexity and smoothness.

Theorem 3.1. Fix $1 < \gamma < \infty, \tau \in (1,2]$, and $\sigma \in [2,\infty)$. Let (\mathbb{X}, ρ) be a Banach space, $\Omega \subset \mathbb{R}^n$ a domain, and fix any σ -convex and τ -smooth norm F on $\mathbb{X} \otimes \mathbb{R}^n$ satisfying

(3.1)
$$F(z \otimes \xi) \lesssim \rho(z)|\xi| \quad \forall z \in \mathbb{X} \ \forall \xi \in \mathbb{R}^{n}.$$
If $\alpha_{*} = \min(\gamma, \tau) - 1$, $f \in \mathcal{B}_{loc,\infty}^{\alpha_{*},\gamma'}(\Omega; \mathbb{X}^{*})$, and $u \in \mathcal{W}^{1,\gamma}(\Omega; \mathbb{X})$ solves (1.1), then
$$j_{F}^{\gamma}(\mathcal{D}u) \in \mathcal{B}_{\infty}^{\alpha_{*},\gamma'}(\Omega; ((\mathbb{X} \otimes \mathbb{R}^{n})^{*}, F_{*}))$$

(3.3)
$$Du \in \mathcal{B}_{\infty,loc}^{\alpha,\gamma}(\Omega; (\mathbb{X} \otimes \mathbb{R}^n, F)) \quad \textit{where} \quad \alpha = \frac{\min(\gamma, \tau)}{\max(\gamma, \sigma)}.$$

Moreover,

(3.4)
$$F(\mathrm{D}u)^{\frac{\gamma-\sigma}{\sigma}}\mathrm{D}u \in \mathrm{B}^{\frac{\min(\gamma,\tau)}{\sigma},\sigma}_{\infty,loc}(\Omega;(\mathbb{X}\otimes\mathbb{R}^n,F))$$
 and if $\gamma \leq \sigma$,

and if
$$\int \leq 0$$
,

$$(3.5) F_*\left(j_F^{\gamma}(\mathrm{D}u)\right)^{\frac{\gamma'-\tau'}{\tau'}}j_F^{\gamma}(\mathrm{D}u) \in \mathrm{B}_{\infty,loc}^{\frac{\min(\gamma,\tau)}{\tau'},\tau'}(\Omega;(\mathbb{X}^*\otimes\mathbb{R}^n;F_*))$$

Remark 3.2. A direct comparison of (3.4) and (3.5) follows from the identity

$$F_*\left(j_F^{\gamma}(\mathrm{D} u)\right)^{\frac{\gamma'-\tau'}{\tau'}}j_F^{\gamma}(\mathrm{D} u)=F(\mathrm{D} u)^{\frac{\gamma-\tau'}{\tau'}}j_F^2(\mathrm{D} u).$$

Proof. Suppose $3B \subset \Omega$ and let $\eta \in C_c^\infty(\Omega; \mathbb{R})$ be a smooth non-negative function supported on 2B with $|D\eta| \lesssim 1$ such that $\eta = 1$ on B. Let $h \in \mathbb{R}^n$ and write $u_h = u(x+h)$ for $|h| < \frac{1}{2}d(B,\partial\Omega)$. Recall that we denote $\Delta_h u = u(x) - u(x+h)$. Testing (1.1) against $\Delta_{-h}\eta^{\gamma}\Delta_h$ and using a change of variables, we find

$$I := \int \langle \Delta_h j_F^{\gamma}(\mathrm{D}u), \mathrm{D}(\eta^{\gamma} \Delta_h u) \rangle_{\mathbb{X} \otimes \mathbb{R}^n} = \int \eta^{\gamma} \langle \Delta_h u, \Delta_h f \rangle_{\mathbb{X}} =: II.$$

We write

$$I = \int \langle \Delta_h j_F^{\gamma}(\mathrm{D}u), \eta^{\gamma} \Delta_h \mathrm{D}u \rangle_{\mathbb{X} \otimes \mathbb{R}^n} + \gamma \int \eta^{\gamma - 1} \langle \Delta_h j_F^{\gamma}(\mathrm{D}u), \Delta_h u \otimes \mathrm{D}\eta \rangle =: A_1 + A_2.$$

To prove (3.2) - (3.5) we will repeatedly make use of

$$(3.6) A_1 \le |A_2| + |II|$$

while estimating $|A_2|$ and A_1 above and below by appropriate functions. The main estimate for A_2 follows from Fenchel's inequality and (3.1),

$$(3.7) \quad |A_2| \lesssim \int \eta^{\gamma-1} F(\Delta_h u \otimes \mathrm{D}\eta) F_* \left(\Delta_h j_F^{\gamma}(\mathrm{D}u) \right) \lesssim \int \eta^{\gamma-1} \rho(\Delta_h u) F_* \left(\Delta_h j_F^{\gamma}(\mathrm{D}u) \right).$$

The only estimate on |II| uses Fenchel's inequality,

$$(3.8) |II| \lesssim \int \eta^{\gamma} \rho(\Delta_h u) \rho_*(\Delta_h f) \lesssim \|\rho(\Delta_h u)\|_{L^{\gamma}(2B)} \|\rho_*(\Delta_h f)\|_{L^{\gamma'}(2B)}.$$

Due to (2.4), (2.3), and (2.6), the main estimate for A_1 is

$$(3.9) A_1 \gtrsim \int \eta^{\gamma} \left(F(\mathrm{D}u) + F(\mathrm{D}u_h) \right)^{\gamma - \sigma} F(\Delta_h \mathrm{D}u)^{\sigma}$$

$$+ \int \eta^{\gamma} \left(F_*(j_F^{\gamma}(\mathrm{D}u)) + F_*(j_F^{\gamma}(\mathrm{D}u_h)) \right)^{\gamma' - \tau'} F_*(\Delta_h j_F^{\gamma}(\mathrm{D}u))^{\tau'}.$$

We now split into three cases to verify (3.2) and (3.3).

Case 1: $\gamma \leq \tau$.

Since $\gamma' \geq \tau'$, (3.9) and the triangle inequality further bound A_1 below

(3.10)
$$A_1 \gtrsim \int \eta^{\gamma} F_* \left(\Delta_h j_F^{\gamma}(\mathrm{D}u) \right)^{\gamma'}.$$

Via Young's inequality the bound on $|A_2|$ in (3.7) becomes

$$(3.11) |A_2| \lesssim \int_{\operatorname{spt} D\eta} \rho(\Delta_h u)^{\gamma} + \frac{1}{2} \int \eta^{\gamma} F_* \left(\Delta_h j_F^{\gamma} (\mathrm{D} u)^{\gamma'} \right).$$

Combining (3.6), (3.8), (3.10), and (3.11) before dividing by $|h|^{\gamma}$, we find

$$\int_{B} F_{*} \left(\frac{\Delta_{h} j_{F}^{\gamma}(\mathrm{D}u)}{|h|^{\gamma-1}} \right)^{\gamma'} \lesssim \int_{2B} \rho \left(\frac{\Delta_{h}u}{|h|} \right)^{\gamma} + \left(\int_{2B} \rho \left(\frac{\Delta_{h}u}{|h|} \right)^{\gamma} \right)^{\frac{1}{\gamma}} \left(\int_{2B} \rho_{*} \left(\frac{\Delta_{h}f}{|h|^{\gamma-1}} \right)^{\gamma'} \right)^{\frac{1}{\gamma'}}.$$

Since $f \in \mathcal{B}^{\gamma-1,\gamma'}_{loc,\infty}$ and $u \in W^{1,\gamma}_{loc}$ it follows the RHS is bounded as $h \searrow 0$. Theorem 2.4 and a standard covering argument now verify (3.2) when $\gamma \leq \tau$.

For (3.3), since $\gamma \leq \tau \leq \sigma$, Hölder's inequality provides another bound for A_1

$$\left(\int \eta^{\gamma} F(\Delta_h Du)^{\gamma}\right)^{\frac{\sigma}{\gamma}} \leq \left(\int \eta^{\gamma} \left(F(Du) + F(Du_h)\right)^{\gamma - \sigma} F(\Delta_h Du)^{\sigma}\right) \times \left(\int \eta^{\gamma} \left(F(Du) + F(Du_h)\right)^{\gamma}\right)^{\frac{\sigma - \gamma}{\gamma}}.$$

The term in the second line is bounded above by a constant depending only on $||u||_{W^{1,\gamma}}(3B)$, σ , and γ .

Thus, combining this observation with (3.6), (3.11), (3.2), (3.8) and (3.9) we find

(3.12)
$$\left(\int_{B/2} \left(\frac{F(\Delta_h Du)}{|h|^{\frac{\sigma}{\gamma}}} \right)^{\gamma} \right)^{\frac{\omega}{\gamma}} \lesssim \|F(Du)\|_{L^{\gamma}(3B)}^{\sigma-\gamma} \left(\|u\|_{W^{1,\gamma}(3B)} \|j_F^{\gamma}(Du)\|_{B_{\infty}^{\gamma-1,\gamma'}(3B)} + \|u\|_{W^{1,\gamma}(3B)} \|f\|_{B_{\infty}^{\gamma-1,\gamma'}(3B)} \right),$$

verifying (3.3) in this case.

Case 2:
$$\tau \leq \gamma \leq \sigma$$
.

Similar to the lower bound on A_1 in the proof of (3.3) from Case 1, since $\gamma' \leq \tau'$, we bound A_1 below by

$$\left(\int \eta^{\gamma} F_{*} \left(\Delta_{h} j_{F}^{\gamma}(\mathbf{D}u)\right)^{\gamma'}\right)^{\frac{\tau'}{\gamma'}} \leq \left(\int \eta^{\gamma} F_{*} \left(j_{F}^{\gamma}(\mathbf{D}u) + j_{F}^{\gamma}(\mathbf{D}u_{h})\right)^{\gamma'-\tau'} F_{*} \left(\Delta_{h} j_{F}^{\gamma}(\mathbf{D}u)\right)^{\tau'}\right) \\
\times \left(\int \eta^{\gamma} F_{*} \left(j_{F}^{\gamma}(\mathbf{D}u) + j_{F}^{\gamma}(\mathbf{D}u_{h})\right)^{\gamma'}\right)^{\frac{\tau'-\gamma'}{\gamma'}}.$$
(3.13)

By Hölder's and Young's inequality, we deduce from (3.7)

$$|A_{2}| \lesssim \left(\int_{3B} \rho(\Delta_{h} u)^{\gamma} \right)^{\frac{1}{\gamma}} \left(\int \eta^{\gamma} F_{*} \left(\Delta_{h} j_{F}^{\gamma} (\mathrm{D} u)^{\gamma'} \right) \right)^{\frac{1}{\gamma'}}$$

$$\lesssim \frac{1}{2} \left(\int \eta^{\gamma} F_{*} \left(\Delta_{h} j_{F}^{\gamma} (\mathrm{D} u)^{\gamma'} \right) \right)^{\frac{\tau'}{\gamma'}} + \left(\int_{3B} \rho(\Delta_{h} u)^{\gamma} \right)^{\frac{\tau}{\gamma}}.$$
(3.14)

Combining (3.6), (3.9), (3.13), (3.14), and (3.8) implies

$$\left(\int_{B} F_{*} \left(\Delta_{h} \frac{j_{F}^{\gamma}(\mathrm{D}u)}{|h|^{\tau-1}} \right)^{\gamma'} \right)^{\frac{\tau'}{\gamma'}} \lesssim \|F(\mathrm{D}u)\|_{\mathrm{W}^{1,\gamma}(3B)} \left(\|u\|_{\mathrm{W}^{1,\gamma}(3B)}^{\tau} + \|u\|_{\mathrm{W}^{1,\gamma}(3B)} \|f\|_{B_{\infty(3B)}^{\tau-1,\gamma'}} \right).$$

A standard covering argument gives (3.2). Since $\gamma \leq \sigma$, (3.12) holds by the same argument as in Case 1. Taking into account that now $j_F^{\gamma}(\mathrm{D}u) \in \mathrm{B}^{\tau-1,\gamma'}_{\mathrm{loc},\infty}$, this verifies (3.3).

Case 3:
$$\gamma \geq \sigma \geq \tau$$
.

Since we never used $\gamma \leq \sigma$ in the proof of (3.2) in the previous two cases, we move directly to proving (3.3). Since $\gamma \geq \sigma$, the triangle inequality produces a bound on A_1 analogous to the proof of (3.2) in Case 1. This combined with (3.6), (3.11) and (3.8) imply

$$\int_{B} \frac{F(\Delta_{h} D u)^{\gamma}}{h^{\tau}} \lesssim \|u\|_{W^{1,\gamma}(3B)} \|\Delta_{h}(j_{F}^{\gamma}(D u)\|_{\mathcal{B}_{\infty}^{\tau-1,\gamma'}(3B)} + \|u\|_{Ww^{1,\gamma}(3B)} \|f\|_{\mathcal{B}_{\infty}^{\tau-1,\gamma'}(3B)}.$$

This completes our casework verifying (3.2) and (3.3).

Next we prove (3.4) and (3.5). We return to (3.7) and estimate A_2 by Hölder's inequality

(3.15)
$$|A_2| \lesssim \|\rho(\Delta_h u)\|_{L^{\gamma}(3B)} \|\rho_*(\Delta_h j_F^{\gamma}(Du))\|_{L^{\gamma'}(3B)}.$$

By (3.9) and Lemma 2.10.

$$(3.16) \quad A_1 \gtrsim \int_{\mathcal{B}} F\left(\Delta_h\left(F(\mathrm{D}u)^{\frac{\gamma-\sigma}{\sigma}}\mathrm{D}u\right)\right)^{\sigma} + F_*\left(\Delta_h\left(F_*(j_F^{\gamma}(\mathrm{D}u))^{\frac{\gamma'-\tau'}{\tau'}}J_F^{\gamma}(Du)\right)\right)^{\tau'}.$$

Now (3.4) and (3.5) follow immediately from (3.6) by (3.16), (3.15), (3.8), (our bounds on A_1 , $|A_2|$, and |II| respectively) and the previous part of this theorem (3.2) and (3.3).

REFERENCES

- [BC13] G. Bonfanti and A. Cellina. The nonoccurrence of the Lavrentiev phenomenon for a class of variational functionals. *SIAM J. Control Optim.*, 51:1639–1650, 2013.
- [BCM12] G. Bonfanti, A. Cellina, and M. Mazzola. The higher integrability and the validity of the Euler-Lagrange equation for solutions to variational problems. SIAM J. Control Optim., 50:888–899, 2012.
- [BCS10] Lorenzo Brasco, Guillaume Carlier, and Filippo Santambrogio. Congested traffic dynamics, weak flows and very degenerate elliptic equations. *Journal de mathématiques pures et appliquées*, 93(6):652–671, 2010.
- [BFK03] Marino Belloni, Vincenzo Ferone, and Bernd Kawohl. Isoperimetric inequalities, wulff shape and related questions for strongly nonlinear elliptic operators. *Zeitschrift für angewandte Mathematik und Physik ZAMP*, 54(5):771–783, 2003.
- [BI87] Bogdan Bojarski and Tadeusz Iwaniec. p-harmonic equation and quasiregular mappings. *Banach Center Publications*, 1(19):25–38, 1987.
- [BLPV17] Lorenzo Brasco, Chiara Leone, Giovanni Pisante, and Anna Verde. Sobolev and lipschitz regularity for local minimizers of widely degenerate anisotropic functionals. *Nonlinear Analysis: Theory, Methods & Applications*, 153:169–199, 2017.
- [BM20] Christian Bargetz and Emir Medjic. On the rate of convergence of iterated bregman projections and of the alternating algorithm. *Journal of Mathematical Analysis and Applications*, 481(1):123482, 2020.
- [CKP13] M. Carozza, J. Kristensen, and A. Passarelli di Napoli. Regularity of minimisers of autonomous convex variational integrals. Ann. della Scu. Norm. Sup. di Pisa, 13(4), 2013.
- [CMP05] M. Carozza, G. Moscariello, and A. Passarelli di Napoli. Regularity results via duality for minimizers of degenerate functionals. Asympt. Anal., 44:221–235, 2005.
- [CP06] M. Carozza and A. Passarelli di Napoli. Regularity for minimizers of degenerate elliptic funcitonals. J. Nonlinear Convex Anal., 7:375–383, 2006.
- [dBL20] Giuseppina di Blasio and Pier Domenico Lamberti. Eigenvalues of the finsler p-laplacian on varying domains. *Mathematika*, 66(3):765–776, 2020.
- [dFKK22] Cristiana de Filippis, Lukas Koch, and Jan Kristensen. Regularity in relaxed convex problems. *in Preparation*, 2022.
- [DMMN18] Matias G Delgadino, Francesco Maggi, Cornelia Mihaila, and Robin Neumayer. Bubbling with 12-almost constant mean curvature and an alexandrov-type theorem for crystals. *Archive for Rational Mechanics and Analysis*, 230(3):1131–1177, 2018.
- [FK09] Vincenzo Ferone and Bernd Kawohl. Remarks on a finsler-laplacian. *Proceedings of the American Mathematical Society*, 137(1):247–253, 2009.
- [Giu03] Enrico Giusti. *Direct Methods in the Calculus of Variations*. World Scientific Publishing Co., Inc., 2003.
- [Goe20] Max Goering. A harnack inequality for $W^{1,\gamma}$ -solutions of anisotropic pdes with non-standard growth. arXiv preprint arXiv:2008.08028, 2020.
- [HvNVW16] Tuomas Hytönen, Jan van Neerven, Mark Veraar, and Lutz Weis. *Analysis in Banach spaces. Volume 1: Martingales and Littlewood-Paley theory*. Springer International Publishing, 2016.
- [KK22] Lukas Koch and Jan Kristensen. On the validity of the euler-lagrange system without growth assumptions. *arXiv*:2203.00333, 2022.
- [KN08] B Kawohl and M Novaga. The p-laplace eigenvalue problem as $p \rightarrow 1$ and cheeger sets in a finsler metric. *Journal of Convex Analysis*, 15(3):623, 2008.
- [Kre15] Marcel Kreuter. Sobolev spaces of vector-valued functions. University of Ulm, 2015.
- [KSYZ22] Alexandru Kristály, Zhongmin Shen, Lixia Yuan, and Wei Zhao. Nonlinear spectrums of finsler manifolds. *Mathematische Zeitschrift*, 300(1):81–123, 2022.
- [LT13] Joram Lindenstrauss and Lior Tzafriri. *Classical Banach spaces II: function spaces*, volume 97. Springer Science & Business Media, 2013.
- [Ser93] G. Seregin. Differential perperties of solutions of variational problems for functionals of linear growth. *J. Sov. Math.*, 64:1256–1277, 1993.

- [Uhl77] Karen Uhlenbeck. Regularity for a class of non-linear elliptic systems. *Acta Mathematica*, 138:219–240, 1977.
- [Xia19] Qiaoling Xia. Sharp spectral gap for the finsler p-laplacian. *Science China Mathematics*, 62(8):1615–1644, 2019.
- [XR91] Zong-Ben Xu and Gary F Roach. Characteristic inequalities of uniformly convex and uniformly smooth banach spaces. *Journal of Mathematical Analysis and Applications*, 157(1):189–210, 1991.
- [ZKO94] V. V. Zhikov, S. M. Kozlov, and O. A. Oleinie. *Homogenization of differential operators and integral functionals*. Springer-Verlag, Berlin, 1994.