

SCALING ASYMPTOTICS OF WIGNER DISTRIBUTIONS OF HARMONIC OSCILLATOR ORBITAL COHERENT STATES

NICHOLAS LOHR

ABSTRACT. The main result of this article gives scaling asymptotics of the Wigner distributions $W_{\varphi_N^\gamma, \varphi_N^\gamma}$ of isotropic harmonic oscillator *orbital coherent states* φ_N^γ concentrating along Hamiltonian orbits γ in shrinking tubes around γ in phase space. In particular, these Wigner distributions exhibit a *hybrid* semi-classical scaling. That is, simultaneously, we have an *Airy scaling* when the tube has radius $N^{-2/3}$ normal to the energy surface Σ_E , and a *Gaussian scaling* when the tube has radius $N^{-1/2}$ tangent to Σ_E .

CONTENTS

1. Introduction	1
1.1. Statements of Results	2
1.2. Future Work	4
1.3. Acknowledgments	4
2. Proofs of the Theorems	6
2.1. Preliminaries	6
2.2. Proof of Proposition 1.1	8
2.3. Proof of Theorem 1.2	9
2.4. Proof of Theorem 1.3	12
3. Relation with the Eigenspace Projector	16
4. Generalizations to d dimensions	17
5. Appendix	19
5.1. Airy Function	19
5.2. Stationary Phase Methods	19
5.3. Formula for $\mu_{00}(u)$	21
References	22

1. INTRODUCTION

Coherent states centered at points in phase space were introduced by Schrödinger [S26] in 1926 as minimal uncertainty states. In 1963, Glauber [G63] extended coherent states to quantum electrodynamics. This paper studies *orbital coherent states* centered on periodic classical Hamiltonian orbits. Orbital coherent states were first studied in 1989 for the Hydrogen atom in the papers of Gay et al. [GDB89] and Nauenberg [N89], and more recently in

The author was partially supported by NSF RTG grant DMS-1502632.

Date: July 26, 2022.

papers by Klauder [K96] and Villegas-Blas et al. [VB96, TVB97]. In 1992, De Bièvre [De92] studied orbital coherent states (4) for the two-dimensional isotropic harmonic oscillator:

$$\widehat{H}_\hbar := -\frac{\hbar^2}{2}\Delta_{\mathbb{R}^2} + \frac{|x|^2}{2}. \quad (1)$$

For $\gamma_0(t) := \sqrt{E}(\cos t, \sin t, -\sin t, \cos t)$ in $T^*\mathbb{R}^2$ with fixed energy $E = \hbar(N+1)$, the $L^2(\mathbb{R}^2)$ -normalized orbital coherent state associated to γ_0 is

$$\varphi_N^{\gamma_0}(x_1, x_2) = \frac{(N+1)^{\frac{N+1}{2}}}{E^{\frac{N+1}{2}}\sqrt{\pi N!}}(x_1 + ix_2)^N e^{-\frac{(N+1)}{2E}|x|^2}. \quad (2)$$

From this initial coherent state $\varphi_N^{\gamma_0}$, we define φ_N^γ for a general Hamiltonian orbit γ with the metaplectic representation (see (4) of Section 1.1). This article concerns the Wigner distributions of these orbital coherent states. We recall that in dimension 2, the Wigner distribution $W_{\varphi_N^\gamma, \varphi_N^\gamma} : T^*\mathbb{R}^2 \rightarrow \mathbb{R}$ of φ_N^γ is defined by

$$W_{\varphi_N^\gamma, \varphi_N^\gamma}(x, \xi) := \frac{1}{4\pi^2\hbar^2} \int_{\mathbb{R}^2} \varphi_N^\gamma(x + \frac{v}{2}) \overline{\varphi_N^\gamma(x - \frac{v}{2})} e^{-\frac{i}{\hbar}\langle v, \xi \rangle} dv$$

(see [W32]). It is well-known that for fixed energy, the Wigner distributions tend in the weak sense to a delta function along the orbit γ as $N \rightarrow \infty$, i.e.

$$W_{\varphi_N^\gamma, \varphi_N^\gamma} \rightharpoonup \delta_\gamma \quad (3)$$

(see [De92, De93]). For completeness, we prove (3) in Proposition 1.1. Our purpose is to study the fine structure of this limit in small tubes around γ . The main result of this article gives Airy and Gaussian scaling asymptotics of the Wigner distribution $W_{\varphi_N^\gamma, \varphi_N^\gamma}$ of $\varphi_N^\gamma \in L^2(\mathbb{R}^2)$ (see Theorem 1.3). In this article, we prove all of the results in dimension 2 for notational convenience. These results generalize to dimension d and the same arguments hold, as explained in Section 4.

1.1. Statements of Results. The orbital coherent states (4) are eigenstates of (1),

$$\widehat{H}_\hbar \varphi_N^{\gamma_0} = E \varphi_N^{\gamma_0}$$

where $E = \hbar(N+1)$. The non-constant Hamiltonian orbits of the classical Hamiltonian $H(x, \xi) = \frac{1}{2}|\xi|^2 + \frac{1}{2}|x|^2$ are great circles on the energy surface $\Sigma_E := \{(x, \xi) : H(x, \xi) = E\}$. Modding out by the points on the same orbit, we see that the space of non-constant orbits is isomorphic to $S^3/S^1 = \mathbb{C}P^1$. Since $SU(2)$ acts transitively on $\mathbb{C}P^1$ by rotation, for any non-constant Hamiltonian orbit γ there exists a $U \in SU(2)$ such that $U \cdot \gamma_0 = \gamma$. We define (up to a phase factor) the coherent state centered at the Hamiltonian orbit γ by

$$\varphi_N^\gamma := \mu(U) \varphi_N^{\gamma_0} \quad (4)$$

where μ is the metaplectic representation on $L^2(\mathbb{R}^2)$ (see Chapter 4 of [F89] for details on the metaplectic representation).

The first result of this article is a proof of (3).

PROPOSITION 1.1. *Let $a : T^*\mathbb{R}^2 \rightarrow \mathbb{R}$ be a smooth function with exponential decay. Then in the limit $N \rightarrow \infty$ where $E = \hbar(N+1)$, we have*

$$\int_{T^*\mathbb{R}^2} a(x, \xi) W_{\varphi_N^\gamma, \varphi_N^\gamma}(x, \xi) dx d\xi \xrightarrow{N \rightarrow \infty} \frac{1}{2\pi} \int_0^{2\pi} a(\gamma(t)) dt. \quad (5)$$

It was proved in [De92] that given a smooth function $a : T^*\mathbb{R}^2 \rightarrow \mathbb{R}$ with exponential decay, in the limit $N \rightarrow \infty$ where $E = \hbar(N + 1)$, we have

$$\langle \varphi_N^\gamma, \text{Op}_N^{aw}(a)[\varphi_N^\gamma] \rangle \xrightarrow{N \rightarrow \infty} \frac{1}{2\pi} \int_0^{2\pi} a(\gamma(t)) dt, \quad (6)$$

where $\text{Op}_N^{aw}(a)$ is the anti-Wick quantization of a . Proposition 1.1 proves the analogous result for Weyl quantization, based on the well-known formula

$$\langle \varphi_N^\gamma, \text{Op}_N^w(a)[\varphi_N^\gamma] \rangle = \int_{T^*\mathbb{R}^2} a(x, \xi) W_{\varphi_N^\gamma, \varphi_N^\gamma}(x, \xi) dx d\xi. \quad (7)$$

The next result gives the pointwise asymptotics of $W_{\varphi_N^\gamma, \varphi_N^\gamma}(x, \xi)$ for (x, ξ) on and off the orbit γ (see Figure 1 for a picture).

THEOREM 1.2. *As $N \rightarrow \infty$, we have the following pointwise asymptotics:*

(1) *If (x, ξ) lies on the orbit, then we have*

$$\frac{1}{N^{5/3}} W_{\varphi_N^\gamma, \varphi_N^\gamma}(x, \xi) = \frac{1}{2^{1/3} \pi^2 E^2} \text{Ai}(0) + O(N^{-2/3})$$

where Ai is the Airy function defined in the Section 5.1. In terms of the Gamma function, $\frac{1}{2^{1/3} \pi^2 E^2} \text{Ai}(0) = \frac{\Gamma(1/3) 12^{1/3}}{4\pi^3 E^2 \sqrt{3}}$

(2) *If (x, ξ) lies on the orbit, then for $0 < t < 1$ we have*

$$\frac{1}{N^{5/3}} W_{\varphi_N^\gamma, \varphi_N^\gamma}(t(x, \xi)) = O(N^{-1/6}).$$

(3) *If $(x, \xi) \neq 0$ is not in the above case, we have*

$$W_{\varphi_N^\gamma, \varphi_N^\gamma}(x, \xi) = O(N^{-\infty}).$$

(4) *If $(x, \xi) = 0$, then the Wigner distribution is explicitly*

$$W_{\varphi_N^\gamma, \varphi_N^\gamma}(x, \xi) = \frac{(-1)^N}{\pi^2 E^2} (N + 1)^2.$$

Our main result concerns the scaling asymptotics between the power growth on the orbit and the exponential decay off of it (see Figures 2 and 3 for pictures).

THEOREM 1.3. *Suppose (x, ξ) lies on γ , and let $(x, \xi), v_1, v_2$ be an orthogonal basis (with each vector of norm $\sqrt{2E}$) for the normal space, $N_{(x, \xi)}\gamma$, of γ at (x, ξ) . For $(u, w_1, w_2) \in \mathbb{R}^3$ in a sufficiently small neighborhood of $(0, 0, 0)$, we have the uniform asymptotic expansion*

$$\begin{aligned} & \frac{1}{N^{5/3}} W_{\varphi_N^\gamma, \varphi_N^\gamma} [(1 + u)(x, \xi) + w_1 v_1 + w_2 v_2] \\ &= \frac{e^{-2(N+1)(w_1^2 + w_2^2)}}{4\pi^3 E^2} \mu_{00}(u) \text{Ai}(a(u)N^{2/3}) + \text{Ai}'(a(u)N^{2/3}) \cdot O(N^{-2/3}) \end{aligned}$$

where

$$a(u)^{3/2} = \begin{cases} 3(u + 1)i\sqrt{|u|(u + 2)} - 3i \arccos(1 + u) & \text{if } u < 0 \\ 3(u + 1)\sqrt{u(u + 2)} - 3 \text{arccosh}(1 + u) & \text{if } u \geq 0 \end{cases}$$




FIGURE 1

and $\mu_{00}(u)$ is a smooth function (explicitly given in Section 5.3) with $\mu_{00}(0) = 2^{5/3}\pi$. In particular, if we rescale $u \rightarrow \frac{u}{(2N)^{2/3}}$, $w_1 \rightarrow \frac{w_1}{(2N)^{1/2}}$, and $w_2 \rightarrow \frac{w_2}{(2N)^{1/2}}$, we have the pointwise limit

$$\lim_{N \rightarrow \infty} \frac{1}{N^{5/3}} W_{\varphi_N^{\gamma}, \varphi_N^{\gamma}} \left[\left(1 + \frac{u}{(2N)^{2/3}}\right)(x, \xi) + \frac{w_1}{(2N)^{1/2}} v_1 + \frac{w_2}{(2N)^{1/2}} v_2 \right] = \frac{e^{-(w_1^2 + w_2^2)}}{2^{1/3} \pi^2 E^2} \text{Ai}(2u).$$

Remark: The point (x, ξ) lies on an orbit γ_0 which lies on the energy surface $\Sigma_E := \{(x, \xi) : \frac{1}{2}(|x|^2 + |\xi|^2) = E\}$. Normal to Σ_E at the point (x, ξ) is the vector $(x, \xi)u$ where $u \in \mathbb{R}$, and tangent to Σ_E at the point (x, ξ) are the vectors $w_1 v_1$ and $w_2 v_2$ where $w_1, w_2 \in \mathbb{R}$. The above theorem shows that $W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}$ exhibits Airy scaling normal to the energy surface and Gaussian scaling tangent to the energy surface. This is consistent with the Airy scaling result of the Wigner distributions of the eigenspace projectors in [HZ20], and we explain this in Section 3.

The function $a(u)$ has a geometric interpretation. When $u < 0$, the function $-\frac{2E}{3}ia(u)^{3/2}$ is the area A_u of the circular segment bounded by γ_0 where $|u|\sqrt{2E}$ is the height of the arced portion, as depicted in Figure 4. When $u \geq 0$, we have a similar interpretation but for a hyperbolic segment, also depicted in Figure 4.

1.2. Future Work. The asymptotics of the Wigner distributions in this paper are some of the few explicit results on Wigner distributions of eigenfunctions. In future work, we plan to study the asymptotics of Wigner distributions of orbital coherent states of the hydrogen atom. They are much more difficult to compute, but the isotropic harmonic oscillator of this paper provides a more readily computable model which guides future work.

1.3. Acknowledgments. This article is part of the Ph.D. thesis of the author at Northwestern University under the guidance of Steve Zelditch. The author thanks Stephan De Bièvre




FIGURE 2




FIGURE 3

for beneficial comments. The author also thanks Michael Geis, Erik Hupp, and Abraham Rabinowitz for helpful conversations.




FIGURE 4. This figure shows the geometric interpretation of $a(u)$ in terms of the areas A_u and B_u .

2. PROOFS OF THE THEOREMS

2.1. Preliminaries.

2.1.1. *Formula.* Before we get into the proofs of the theorems, we first write down a formula for the Wigner distribution. Recall from (4) that the coherent state $\varphi_N^{\gamma_0}$ is

$$\varphi_N^{\gamma_0}(x) = \frac{(N+1)^{\frac{N+1}{2}}}{E^{\frac{N+1}{2}} \sqrt{\pi N!}} (x_1 + ix_2)^N e^{-\frac{(N+1)}{2E}|x|^2}.$$

The Wigner distribution for $E = (N+1)\hbar$ is then

$$\begin{aligned} W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}(x, \xi) &= \int_{\mathbb{R}^2} \varphi_N^{\gamma_0}\left(x + \frac{v}{2}\right) \overline{\varphi_N^{\gamma_0}\left(x - \frac{v}{2}\right)} e^{-\frac{i}{\hbar}\langle v, \xi \rangle} \frac{dv}{4\pi^2 \hbar^2} \\ &= \frac{(N+1)^{N+3}}{4\pi^3 E^{N+3} N!} \int_{\mathbb{R}^2} \left(x + \frac{v}{2}\right)^N \overline{\left(x - \frac{v}{2}\right)^N} e^{-\frac{N+1}{2E}|x - \frac{v}{2}|^2} e^{-\frac{N+1}{2E}|x + \frac{v}{2}|^2} e^{-i\frac{(N+1)}{E}\langle v, \xi \rangle} dv \end{aligned} \quad (8)$$

where the multiplication within the integral is complex multiplication. Since

$$\left|x - \frac{v}{2}\right|^2 + \left|x + \frac{v}{2}\right|^2 = 2|x|^2 + \frac{1}{2}|v|^2,$$

we have

$$\begin{aligned} W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}(x, \xi) &= \frac{(N+1)^{N+3}}{4\pi^3 E^{N+3} N!} \int_{\mathbb{R}^2} \left(x + \frac{v}{2}\right)^N \overline{\left(x - \frac{v}{2}\right)^N} e^{-\frac{N+1}{2E}|x - \frac{v}{2}|^2} e^{-\frac{N+1}{2E}|x + \frac{v}{2}|^2} e^{-i\frac{(N+1)}{E}\langle v, \xi \rangle} dv \\ &= \frac{(N+1)^{N+3}}{4\pi^3 E^{N+3} N!} e^{-\frac{(N+1)}{E}|x|^2} \int_{\mathbb{R}^2} \left(x + \frac{v}{2}\right)^N \overline{\left(x - \frac{v}{2}\right)^N} e^{-\frac{N+1}{4E}|v|^2} e^{-i\frac{(N+1)}{E}\langle v, \xi \rangle} dv \\ &= \frac{(N+1)^{N+3}}{4\pi^3 E^{N+2} N!} e^{-\frac{(N+1)}{E}|x|^2} \int_{\mathbb{R}^2} \left(x + \frac{v}{2}\sqrt{E}\right)^N \overline{\left(x - \frac{v}{2}\sqrt{E}\right)^N} e^{-\frac{N+1}{4}|v|^2} e^{-i\frac{(N+1)}{\sqrt{E}}\langle v, \xi \rangle} dv, \end{aligned} \quad (9)$$

where the last line we applied the substitution $v \rightarrow \sqrt{E}v$. To apply the stationary phase method on this formula later on, we write

$$e^{N \log\left(x + \frac{v}{2}\sqrt{E}\right) \overline{\left(x - \frac{v}{2}\sqrt{E}\right)}} = \left(x + \frac{v}{2}\sqrt{E}\right)^N \overline{\left(x - \frac{v}{2}\sqrt{E}\right)^N}$$

where we use the principal branch of log and define the complex phase

$$\Psi(x, \xi, v) := -i \log\left(x + \frac{v}{2}\sqrt{E}\right) \overline{\left(x - \frac{v}{2}\sqrt{E}\right)} + i \log(E) + i\left(\frac{|x|^2}{E} - 1\right) + i\frac{|v|^2}{4} - \frac{1}{\sqrt{E}}\langle v, \xi \rangle$$

$$= -i \log\left(\frac{x}{\sqrt{E}} + \frac{v}{2}\right) \overline{\left(\frac{x}{\sqrt{E}} - \frac{v}{2}\right)} + i\left(\frac{|x|^2}{E} - 1\right) + i\frac{|v|^2}{4} - \frac{1}{\sqrt{E}}\langle v, \xi \rangle \quad (10)$$

The Wigner distribution is then

$$W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}(x, \xi) = \frac{(N+1)^{N+3}}{4\pi^3 E^2 N!} e^{-N - \frac{1}{E}|x|^2} \int_{\mathbb{R}^2} e^{-\frac{1}{4E}|v|^2} e^{-\frac{i}{E}\langle v, \xi \rangle} e^{iN\Psi(x, \xi, v)} dv. \quad (11)$$

We can use Stirling's approximation to write

$$W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}(x, \xi) = \frac{(N+1)^{5/2}}{4\pi^3 \sqrt{2\pi} E^2} e^{1 - \frac{1}{E}|x|^2} \left(1 + O\left(\frac{1}{N}\right)\right) \int_{\mathbb{R}^2} e^{-\frac{1}{4}|v|^2} e^{-\frac{i}{\sqrt{E}}\langle v, \xi \rangle} e^{iN\Psi(x, \xi, v)} dv,$$

which implies

$$\frac{1}{N^{5/3}} W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}(x, \xi) = \frac{N^{5/6}}{4\pi^3 \sqrt{2\pi} E^2} e^{1 - \frac{1}{E}|x|^2} \left(1 + O\left(\frac{1}{N}\right)\right) \int_{\mathbb{R}^2} e^{-\frac{1}{4}|v|^2} e^{-\frac{i}{\sqrt{E}}\langle v, \xi \rangle} e^{iN\Psi(x, \xi, v)} dv. \quad (12)$$

The next lemma will be very useful for our stationary phase approximations.

LEMMA 2.1. *For Ψ defined in (10), we have*

$$\operatorname{Im} \Psi(x, \xi, v) \geq 0$$

with equality if and only if $|v|^2 = 4(1 - \frac{1}{E}|x|^2)$ and $\langle x, v \rangle = 0$.

Proof. Indeed,

$$\begin{aligned} \operatorname{Im} \Psi(x, \xi, v) &\stackrel{(10)}{=} -\log \left| \left(\frac{x}{\sqrt{E}} + \frac{v}{2}\right) \overline{\left(\frac{x}{\sqrt{E}} - \frac{v}{2}\right)} \right| + \frac{|x|^2}{E} - 1 + \frac{|v|^2}{4} \\ &= -\log \left| \frac{|x|^2}{E} - \frac{|v|^2}{4} + i\frac{1}{\sqrt{E}} \operatorname{Im}(v\bar{x}) \right| + \frac{|x|^2}{E} - 1 + \frac{|v|^2}{4} \\ &= -\frac{1}{2} \log \left(\left(\frac{|x|^2}{E} - \frac{|v|^2}{4}\right)^2 + \frac{1}{E} (\operatorname{Im}(v\bar{x}))^2 \right) + \frac{|x|^2}{E} - 1 + \frac{|v|^2}{4} \\ &\geq -\frac{1}{2} \log \left(\left(\frac{|x|^2}{E} - \frac{|v|^2}{4}\right)^2 + \frac{1}{E} |xv|^2 \right) + \frac{|x|^2}{E} - 1 + \frac{|v|^2}{4} \\ &= -\log \left(\frac{|x|^2}{E} + \frac{|v|^2}{4} \right) + \frac{|x|^2}{E} - 1 + \frac{|v|^2}{4} \\ &= -\log \left(1 + \frac{|x|^2}{E} - 1 + \frac{|v|^2}{4} \right) + \frac{|x|^2}{E} - 1 + \frac{|v|^2}{4} \\ &\geq 0, \end{aligned}$$

where the first inequality we use the fact that $|\operatorname{Im} z| \leq |z|$ (with equality if and only if $\operatorname{Re} z = 0$), and the last inequality we use the fact that $-\log(1+x) + x \geq 0$ for all $x \geq -1$ (with equality if and only if $x = 0$), which completes the proof. \square

2.1.2. Symmetries. One important property of Wigner distributions is the so-called metaplectic covariance. In dimension 2, it states that given $f \in L^2(\mathbb{R}^2)$ and μ the metaplectic representation on $L^2(\mathbb{R}^2)$, we have

$$W_{\mu(A)f, \mu(A)f}(x, \xi) = W_{f, f}(A^*(x, \xi)) \quad (13)$$

for all $A \in \operatorname{Sp}(4, \mathbb{R})$ (see page 180 of [F89]). This is useful in proving results for the Wigner distributions of general coherent states φ_N^γ from that of Wigner distributions of $\varphi_N^{\gamma_0}$. In particular, if $U \in \operatorname{SU}(2) \subset \operatorname{Sp}(4, \mathbb{R})$ is such that $U \cdot \gamma_0 = \gamma$, then we have

$$W_{\varphi_N^\gamma, \varphi_N^\gamma}(x, \xi) \stackrel{(13)}{=} W_{\mu(U)\varphi_N^{\gamma_0}, \mu(U)\varphi_N^{\gamma_0}}(x, \xi) \stackrel{(4)}{=} W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}(U^*(x, \xi)). \quad (14)$$

The next lemma will be useful in proving our results for general points in phase space from an initial point.

LEMMA 2.2. For $g_\theta = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \in \text{SO}(2)$, we have

$$W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}(g_\theta \cdot x, g_\theta \cdot \xi) = W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}(x, \xi).$$

In particular, $W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}$ is radial in the plane containing γ .

Proof. One proof is to use the fact that the restricted representation of the metaplectic representation to the subgroup $\text{SO}(2)$ is rotation of functions. The coherent state $\varphi_N^{\gamma_0}$ is invariant under rotation up to a phase factor, which results in equal Wigner distributions. Alternatively, we can use formula (9). We have

$$\begin{aligned} W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}(g_\theta \cdot x, g_\theta \cdot \xi) &= \frac{(N+1)^{N+3}}{4\pi^3 E^{N+3} N!} e^{-\frac{(N+1)}{E}|g_\theta \cdot x|^2} \\ &\quad \int_{\mathbb{R}^2} (g_\theta \cdot x + \frac{v}{2})^N \overline{(g_\theta \cdot x - \frac{v}{2})^N} e^{-\frac{N+1}{4E}|v|^2} e^{-i\frac{(N+1)}{E}\langle v, g_\theta \cdot \xi \rangle} dv \\ &= \frac{(N+1)^{N+3}}{4\pi^3 E^{N+3} N!} e^{-\frac{(N+1)}{E}|x|^2} \int_{\mathbb{R}^2} (g_\theta \cdot x + \frac{v}{2})^N \overline{(g_\theta \cdot x - \frac{v}{2})^N} e^{-\frac{N+1}{4E}|v|^2} e^{-i\frac{(N+1)}{E}\langle g_\theta^* \cdot v, \xi \rangle} dv \\ &= \frac{(N+1)^{N+3}}{4\pi^3 E^{N+3} N!} e^{-\frac{(N+1)}{E}|x|^2} \int_{\mathbb{R}^2} (g_\theta \cdot x + \frac{g_\theta \cdot v}{2})^N \overline{(g_\theta \cdot x - \frac{g_\theta \cdot v}{2})^N} e^{-\frac{N+1}{4E}|g_\theta \cdot v|^2} e^{-i\frac{(N+1)}{E}\langle v, \xi \rangle} dv \\ &= \frac{(N+1)^{N+3}}{4\pi^3 E^{N+3} N!} e^{-\frac{(N+1)}{E}|x|^2} \int_{\mathbb{R}^2} (g_\theta \cdot (x + \frac{v}{2}))^N \overline{(g_\theta \cdot (x - \frac{v}{2}))^N} e^{-\frac{N+1}{4E}|v|^2} e^{-i\frac{(N+1)}{E}\langle v, \xi \rangle} dv \\ &= \frac{(N+1)^{N+3}}{4\pi^3 E^{N+3} N!} e^{-\frac{(N+1)}{E}|x|^2} \int_{\mathbb{R}^2} e^{iN\theta} (x + \frac{v}{2})^N e^{-iN\theta} \overline{(x - \frac{v}{2})^N} e^{-\frac{N+1}{4E}|v|^2} e^{-i\frac{(N+1)}{E}\langle v, \xi \rangle} dv \\ &= W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}(x, \xi), \end{aligned}$$

where the third line we switch variables $v \rightarrow g_\theta \cdot v$. The last statement of the lemma follows from an application of (14). \square

2.2. Proof of Proposition 1.1. We first prove this for $\gamma_0(t) = \sqrt{E}(\cos t, \sin t, -\sin t, \cos t)$. We write

$$I_N := \int_{T^*\mathbb{R}^2} a(x, \xi) W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}(x, \xi) dx d\xi.$$

Using (12), we have

$$I_N = \frac{N^{5/2}}{4\pi^3 \sqrt{2\pi} E^2} (1 + O(\frac{1}{N})) \int_{\mathbb{R}^6} a(x, \xi) e^{1 - \frac{1}{E}|x|^2} e^{-\frac{1}{4}|v|^2} e^{-\frac{i}{\sqrt{E}}\langle v, \xi \rangle} e^{iN\Psi(x, \xi, v)} dv dx d\xi.$$

We now apply the method of stationary phase for Bott-Morse functions (see Theorem 5.4 in Section 5.2). Recall from (10) that Ψ is defined by

$$\Psi(x, \xi, v) := -i \log\left(\frac{x}{\sqrt{E}} + \frac{v}{2}\right) \overline{\left(\frac{x}{\sqrt{E}} - \frac{v}{2}\right)} + i\left(\frac{|x|^2}{E} - 1\right) + i\frac{|v|^2}{4} - \frac{1}{\sqrt{E}}\langle v, \xi \rangle.$$

Using Wirtinger derivatives in ξ and v along with Lemma 2.1, one can deduce

$$C := \{(x, \xi, v) \in \mathbb{R}^6 = \mathbb{C}^3 \mid \text{Im } \Psi = 0, \Psi' = 0\} = \{(\sqrt{E}e^{i\theta}, i\sqrt{E}e^{i\theta}, 0) \in \mathbb{C}^3 \mid \theta \in [0, 2\pi)\}.$$

This is a one-dimensional submanifold where the tangent space $T_\theta C$ is spanned by the vector

$$-\sin(\theta)\partial_{x_1} + \cos(\theta)\partial_{x_2} - \cos(\theta)\partial_{\xi_1} - \sin(\theta)\partial_{\xi_2}.$$

Completing this to an orthonormal basis of \mathbb{R}^6 , we see that normal space, $N_\theta C$, of C at θ is the span of

$$\partial_{v_1}, \partial_{v_2}, \frac{1}{\sqrt{2}}(\partial_{x_2} + \partial_{\xi_1}), \frac{1}{\sqrt{2}}(\partial_{x_1} - \partial_{\xi_2}), \frac{1}{\sqrt{2}}(\cos(\theta)\partial_{x_1} + \sin(\theta)\partial_{x_2} - \sin(\theta)\partial_{\xi_1} + \cos(\theta)\partial_{\xi_2}).$$

Using this basis, one can compute

$$\det(-i\Psi''|_{N_\theta C}) = 8/E^3.$$

It is easy to see that the natural Riemannian measure on C is $d\mu_C = \sqrt{2E}d\theta$. Now applying Theorem 5.4, we see

$$\begin{aligned} I_N &= \frac{N^{5/2}}{4\pi^3\sqrt{2\pi}E^2} (1 + O(\frac{1}{N})) \int_{\mathbb{R}^6} a(x, \xi) e^{1-\frac{1}{E}|x|^2} e^{-\frac{1}{4}|v|^2} e^{-\frac{i}{\sqrt{E}}\langle v, \xi \rangle} e^{iN\Psi(x, \xi, v)} dv dx d\xi \\ &= \frac{N^{5/2}}{4\pi^3\sqrt{2\pi}E^2} (1 + O(\frac{1}{N})) \left[\left(\frac{2\pi}{N}\right)^{\frac{5}{2}} \int_C \frac{a(u)}{\sqrt{8/E^3}} d\mu_C(u) + O(N^{-\frac{7}{2}}) \right] \\ &= \frac{1}{2\pi} \int_0^{2\pi} a(\sqrt{E}e^{i\theta}, i\sqrt{E}e^{i\theta}) d\theta + O(\frac{1}{N}) \\ &= \frac{1}{2\pi} \int_0^{2\pi} a(\gamma_0(\theta)) d\theta + O(\frac{1}{N}), \end{aligned}$$

which is the desired result for γ_0 . Another way to prove this is to initially change x to polar coordinates $re^{i\theta}$, and then use Theorem 5.2 in the variables r, ξ, v for fixed θ .

To prove the proposition for a general orbit γ , let $U \in \text{SU}(2)$ be such that $U \cdot \gamma_0 = \gamma$, and let $a : T^*\mathbb{R}^2 \rightarrow \mathbb{R}$ be a smooth function that decays exponentially. Then $b(x, \xi) := a(U(x, \xi))$ is also a smooth function with exponential decay. Applying what we proved to b , we have

$$\int_{T^*\mathbb{R}^2} b(x, \xi) W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}(x, \xi) dx d\xi \xrightarrow{N \rightarrow \infty} \frac{1}{2\pi} \int_0^{2\pi} b(\gamma_0(\theta)) d\theta. \quad (15)$$

Changing variables and using (14), we have

$$\begin{aligned} \int_{T^*\mathbb{R}^2} b(x, \xi) W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}(x, \xi) dx d\xi &= \int_{T^*\mathbb{R}^2} a(U(x, \xi)) W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}(x, \xi) dx d\xi \\ &= \int_{T^*\mathbb{R}^2} a(x, \xi) W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}(U^*(x, \xi)) dx d\xi \\ &= \int_{T^*\mathbb{R}^2} a(x, \xi) W_{\mu(U)\varphi_N^{\gamma_0}, \mu(U)\varphi_N^{\gamma_0}}(x, \xi) dx d\xi \\ &= \int_{T^*\mathbb{R}^2} a(x, \xi) W_{\varphi_N^\gamma, \varphi_N^\gamma}(x, \xi) dx d\xi. \end{aligned}$$

Substituting this into (15), we see

$$\int_{T^*\mathbb{R}^2} a(x, \xi) W_{\varphi_N^\gamma, \varphi_N^\gamma}(x, \xi) dx d\xi \xrightarrow{N \rightarrow \infty} \frac{1}{2\pi} \int_0^{2\pi} a(\gamma(\theta)) d\theta,$$

as desired.

2.3. Proof of Theorem 1.2. We first prove this theorem for $\gamma_0(t) = \sqrt{E}(\cos t, \sin t, -\sin t, \cos t)$ by applying Theorem 5.1 to (11). Before we do that, we first compute the Wigner distribution explicitly when $(x, \xi) = (0, 0)$

2.3.1. *Proof of (4).* Using (8) and the fact that $\|\varphi_N^{\gamma_0}\|_2 = 1$, we have

$$\begin{aligned} W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}(0, 0) &= \int_{\mathbb{R}^2} \varphi_N^{\gamma_0}\left(\frac{v}{2}\right) \overline{\varphi_N^{\gamma_0}\left(-\frac{v}{2}\right)} \frac{dv}{4\pi^2 \hbar^2} \\ &= \frac{(-1)^N (N+1)}{\pi^2 E^2} \int_{\mathbb{R}^2} |\varphi_N^{\gamma_0}(v)|^2 dv \end{aligned} \quad (16)$$

$$= \frac{(-1)^N (N+1)}{\pi^2 E^2} \quad (17)$$

where $E = (N+1)\hbar$. This shows (4) in the statement of the theorem.

2.3.2. *Proof of (3).* Now we apply the method of stationary phase to (12). Recall the complex phase defined in (10):

$$\Psi(x, \xi, v) := -i \log\left(\frac{x}{\sqrt{E}} + \frac{v}{2}\right) \overline{\left(\frac{x}{\sqrt{E}} - \frac{v}{2}\right)} + i\left(\frac{|x|^2}{E} - 1\right) + i\frac{|v|^2}{4} - \frac{1}{\sqrt{E}} \langle v, \xi \rangle.$$

Computing the Wirtinger derivatives in v , we have the following lemma:

LEMMA 2.3. *The critical point equation $D_v \Psi(x, \xi, v) = 0$ is*

$$\begin{cases} \partial_v \Psi = 0 \\ \bar{\partial}_v \Psi = 0 \end{cases} \iff \begin{cases} iE + \frac{i}{2}v\sqrt{E}\overline{\left(x - \frac{v}{2}\sqrt{E}\right)} - \xi\overline{\left(x - \frac{v}{2}\sqrt{E}\right)} = 0 \\ -iE + \frac{i}{2}\bar{v}\sqrt{E}\left(x + \frac{v}{2}\sqrt{E}\right) - \bar{\xi}\left(x + \frac{v}{2}\sqrt{E}\right) = 0 \end{cases}, \quad (18)$$

which has solutions only if $\langle x, \xi \rangle = 0$.

Proof. Using $\langle v, \xi \rangle = v_1 \xi_1 + v_2 \xi_2 = \operatorname{Re} v \bar{\xi} = \frac{1}{2}(v \bar{\xi} + \bar{v} \xi)$, we have $\bar{\partial}_v(v_1 \xi_1 + v_2 \xi_2) = \frac{1}{2} \bar{\xi}$ and $\partial_v(v_1 \xi_1 + v_2 \xi_2) = \frac{1}{2} \bar{\xi}$. Then have

$$\bar{\partial}_v \Psi = \frac{i}{2} \frac{\frac{x}{\sqrt{E}} + \frac{v}{2}}{\left(\frac{x}{\sqrt{E}} + \frac{v}{2}\right) \overline{\left(\frac{x}{\sqrt{E}} - \frac{v}{2}\right)}} + \frac{i}{4} v - \frac{1}{2\sqrt{E}} \xi = 0,$$

and

$$\partial_v \Psi = -\frac{i}{2} \frac{\overline{\frac{x}{\sqrt{E}} - \frac{v}{2}}}{\left(\frac{x}{\sqrt{E}} + \frac{v}{2}\right) \overline{\left(\frac{x}{\sqrt{E}} - \frac{v}{2}\right)}} + \frac{i}{4} \bar{v} - \frac{1}{2\sqrt{E}} \bar{\xi} = 0.$$

Canceling and multiplying through, we have the desired system. Adding these two equations, we see

$$i\sqrt{E}[\operatorname{Re}(v\bar{x}) - \operatorname{Im}(v\bar{\xi})] = 2 \operatorname{Re}(\xi\bar{x}). \quad (19)$$

In particular, both sides of this equation are 0. We then see $\langle x, \xi \rangle = 0$. \square

We now characterize when the set $C_{x, \xi} := \{v \in \mathbb{R}^2 \mid D_v \Psi(x, \xi, v) = 0, \operatorname{Im} \Psi(x, \xi, v) = 0\}$ is nonempty.

LEMMA 2.4. *The set $C_{x, \xi} := \{v \in \mathbb{R}^2 \mid D_v \Psi(x, \xi, v) = 0, \operatorname{Im} \Psi(x, \xi, v) = 0\}$ is nonempty if and only if $0 \leq |x| \leq \sqrt{E}$ and $\xi = ix$. In this case, when $x \neq 0$,*

$$C_{x, \xi} = \left\{ v_{\pm} := \pm \frac{2}{|x|} ix \sqrt{1 - \frac{|x|^2}{E}} \right\}.$$

Proof. By Lemma 2.3, $D_v\Psi(x, \xi, v) = 0$ has solutions only if $\langle x, \xi \rangle = 0$. This means either $x = 0$ or $\xi = kix$ for some $k \in \mathbb{R}$. We consider each case separately.

$x = \mathbf{0}$: In the case $x = 0$, the top equation of (18) becomes

$$iE - \frac{i}{4}E|v|^2 + \sqrt{E}\xi\bar{v}/2 = 0 \quad (20)$$

By the equality statement of Lemma 2.1, we have $|v| = 2$. With (20), this implies $\xi\bar{v} = 0$ and hence $\xi = 0$.

$\xi = kix$ and $x \neq \mathbf{0}$: In the case $\xi = kix$ and $x \neq 0$, (18) becomes

$$\begin{cases} \frac{\sqrt{E}}{2}(v\bar{x} + kx\bar{v}) = \frac{E}{4}|v|^2 + k|x|^2 - E \\ \frac{\sqrt{E}}{2}(\bar{v}x + k\bar{x}v) = E - k|x|^2 - \frac{E}{4}|v|^2 = -(\frac{E}{4}|v|^2 + k|x|^2 - E). \end{cases} \quad (21)$$

Taking the complex conjugate of the second equation and comparing with the first, we see

$$\frac{1}{4}|v|^2 + \frac{k}{E}|x|^2 - 1 = 0 \quad (22)$$

The equality statement of Lemma 2.1 forces $k = 1$, and we are done. The last statement of the lemma again follows from the equality statement of Lemma 2.1. \square

Note that Lemma 2.1 and Lemma 2.4 along with Theorem 7.7.1 of [H03] implies that $W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}(x, \xi) = O(N^{-\infty})$ when $|x| \notin [0, \sqrt{E}]$ or $\xi \neq ix$, which proves (3) in the statement of the theorem.

2.3.3. *Proof of (2).* We next compute the Hessian of Ψ in v at the critical points v_{\pm} .

LEMMA 2.5. *Suppose $0 < |x| \leq \sqrt{E}$ and $\xi = ix$. Then the determinant of the Hessian $D_v^2\Psi$ evaluated at the critical points $v_{\pm} = \pm \frac{2}{|x|}ix\sqrt{1 - \frac{|x|^2}{E}}$ is*

$$\det D_v^2\Psi(x, ix, v_{\pm}) = 2\frac{|x|^4}{E^2} - 2\frac{|x|^2}{E} \mp i(2\frac{|x|^2}{E} - 1)\frac{|x|}{\sqrt{E}}\sqrt{1 - \frac{|x|^2}{E}}$$

with

$$|\det D_v^2\Psi(x, ix, v_{\pm})| = \frac{|x|}{\sqrt{E}}\sqrt{1 - \frac{|x|^2}{E}}.$$

Proof. Recall from (10) that the phase function is

$$\Psi(x, \xi, v) = -i \log\left(\frac{x}{\sqrt{E}} + \frac{v}{2}\right)\overline{\left(\frac{x}{\sqrt{E}} - \frac{v}{2}\right)} + i\left(\frac{|x|^2}{E} - 1\right) + i\frac{|v|^2}{4} - \frac{1}{\sqrt{E}}\langle v, \xi \rangle$$

with first derivatives

$$\partial_v\Psi = -\frac{i}{2}\frac{1}{\left(\frac{x}{\sqrt{E}} + \frac{v}{2}\right)} + \frac{i}{4}\bar{v} - \frac{1}{2\sqrt{E}}\bar{\xi}, \quad \bar{\partial}_v\Psi = \frac{i}{2}\frac{1}{\left(\frac{x}{\sqrt{E}} - \frac{v}{2}\right)} + \frac{i}{4}v - \frac{1}{2\sqrt{E}}\xi = 0.$$

The second derivatives are then

$$\partial_v^2\Psi = \frac{i}{4}\frac{1}{\left(\frac{x}{\sqrt{E}} + \frac{v}{2}\right)^2}, \quad \bar{\partial}_v\partial_v\Psi = \partial_v\bar{\partial}_v\Psi = \frac{i}{4}, \quad \bar{\partial}_v^2\Psi = \frac{i}{4}\frac{1}{\left(\frac{x}{\sqrt{E}} - \frac{v}{2}\right)^2}.$$

The complex Hessian is

$$H_{\Psi} = \frac{i}{4} \begin{pmatrix} \left(\frac{x}{\sqrt{E}} + \frac{v}{2}\right)^{-2} & \\ & 1 \\ 1 & \left(\frac{x}{\sqrt{E}} - \frac{v}{2}\right)^{-2} \end{pmatrix} \implies \det H_{\Psi}\Psi = \frac{1}{16} \left(1 - \frac{1}{\left[\left(\frac{x}{\sqrt{E}} + \frac{v}{2}\right)\left(\frac{x}{\sqrt{E}} - \frac{v}{2}\right)\right]^2} \right). \quad (23)$$

Note that we have

$$\begin{aligned}
\left(\frac{x}{\sqrt{E}} + \frac{v_{\pm}}{2}\right)\overline{\left(\frac{x}{\sqrt{E}} - \frac{v_{\pm}}{2}\right)} &= \left(\frac{x}{\sqrt{E}} \pm \frac{1}{|x|}ix\sqrt{1 - \frac{|x|^2}{E}}\right)\overline{\left(\frac{x}{\sqrt{E}} \mp \frac{1}{|x|}ix\sqrt{1 - \frac{|x|^2}{E}}\right)} \\
&= \left(\frac{x}{\sqrt{E}} \pm \frac{1}{|x|}ix\sqrt{1 - \frac{|x|^2}{E}}\right)\left(\frac{\bar{x}}{\sqrt{E}} \pm \frac{1}{|\bar{x}|}i\bar{x}\sqrt{1 - \frac{|x|^2}{E}}\right) \\
&= 2\frac{|x|^2}{E} - 1 \pm 2i\frac{|x|}{\sqrt{E}}\sqrt{1 - \frac{|x|^2}{E}},
\end{aligned}$$

which has unit norm. Squaring, we have

$$\begin{aligned}
\left[\left(\frac{x}{\sqrt{E}} + \frac{v_{\pm}}{2}\right)\overline{\left(\frac{x}{\sqrt{E}} - \frac{v_{\pm}}{2}\right)}\right]^2 &= \left[2\frac{|x|^2}{E} - 1 \pm 2i\frac{|x|}{\sqrt{E}}\sqrt{1 - \frac{|x|^2}{E}}\right]^2 \\
&= 8\frac{|x|^4}{E^2} - 8\frac{|x|^2}{E} + 1 \pm i4\left(2\frac{|x|^2}{E} - 1\right)\frac{|x|}{\sqrt{E}}\sqrt{1 - \frac{|x|^2}{E}}
\end{aligned}$$

The determinant of the complex Hessian is

$$\det H_{\Psi}(x, ix, v_{\pm}) = \frac{1}{16}\left(8\frac{|x|^2}{E} - 8\frac{|x|^4}{E^2} \pm i4\left(2\frac{|x|^2}{E} - 1\right)\frac{|x|}{\sqrt{E}}\sqrt{1 - \frac{|x|^2}{E}}\right)$$

with

$$|\det H_{\Psi}(x, ix, v_{\pm})| = \frac{|x|}{4\sqrt{E}}\sqrt{1 - \frac{|x|^2}{E}}.$$

Note that the determinant of the real Hessian $\det D_v^2\Psi$ differs from the determinant of the complex Hessian $\det H_{\Psi}$ by a factor of -4 , and we are done \square

Now we use Theorem 5.1 applied to $W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}(x, \xi)$ when $0 < |x| < \sqrt{E}$ and $\xi = ix$. Indeed, the only critical points of the phase are v_{\pm} (see Lemma 2.4). Theorem 5.1 guarantees

$$\begin{aligned}
\left|\frac{1}{N^{5/3}}W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}(x, \xi)\right| &\stackrel{(12)}{=} \left|\frac{N^{5/6}}{4\pi^3\sqrt{2\pi}E^2}e^{1 - \frac{1}{E}|x|^2}\left(1 + O\left(\frac{1}{N}\right)\right)\int_{\mathbb{R}^2}e^{-\frac{1}{4}|v|^2}e^{-\frac{i}{\sqrt{E}}\langle v, \xi \rangle}e^{iN\Psi(x, \xi, v)}dv\right| \\
&\leq 2\frac{N^{5/6}}{4\pi^3\sqrt{2\pi}E^2}e^{1 - \frac{1}{E}|x|^2}\left(1 + O\left(\frac{1}{N}\right)\right)\left(\frac{2\pi}{N\sqrt{|\det(D_v^2\Psi(x, ix, v_{\pm}))|}}|f(v_{\pm})| + O(N^{-3/2})\right) \\
&= O(N^{-1/6}),
\end{aligned}$$

as desired for (2) in the theorem.

2.3.4. Proof of (1). This follows directly from Theorem 1.3 by setting $u = w_1 = w_2 = 0$. An independent proof is possible with a careful reading of the coefficients of the proof of Proposition 3 on page 334 of [S93].

For general γ , we use (14), and we are done.

2.4. Proof of Theorem 1.3. Again, we start by proving this result for

$\gamma_0(t) = \sqrt{E}(\cos t, \sin t, -\sin t, \cos t)$ and $(x, \xi) = \sqrt{E}((1, 0), (0, 1))$, which in complex notation is $\sqrt{E}(1, i)$. That is, we would like to compute the limit as $N \rightarrow \infty$ of

$$\begin{aligned}
&\frac{1}{N^{5/3}}W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}[\sqrt{E}(1, i)(1 + u) + w_1v_1 + w_2v_2] \\
&= \frac{1}{N^{5/3}}W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}[\sqrt{E} + u\sqrt{E} + iw_1\sqrt{E} + w_2\sqrt{E}, i\sqrt{E} + iu\sqrt{E} + w_1\sqrt{E} - iw_2\sqrt{E}].
\end{aligned}$$

where $v_1 := \sqrt{E}(i, 1)$ and $v_2 := \sqrt{E}(1, -i)$. We define

$$x_{u, w_1, w_2} := \sqrt{E} + u\sqrt{E} + iw_1\sqrt{E} + w_2\sqrt{E}$$

$$\xi_{u,w_1,w_2} := i\sqrt{E} + iu\sqrt{E} + w_1\sqrt{E} - iw_2\sqrt{E}.$$

By (12), we have

$$\begin{aligned} \frac{1}{N^{5/3}} W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}(x_{u,w_1,w_2}, \xi_{u,w_1,w_2}) &= \frac{N^{5/6}}{4\pi^3 \sqrt{2\pi} E^2} e^{1 - \frac{1}{E} |x_{u,w_1,w_2}|^2} \left(1 + O\left(\frac{1}{N}\right)\right) \\ &\int_{\mathbb{R}^2} f(v_1, v_2, u, w_1, w_2) e^{iN\Psi(v_1, v_2, u, w_1, w_2)} dv_1 dv_2. \end{aligned} \quad (24)$$

where

$$\begin{aligned} \Psi(v_1, v_2, u, w_1, w_2) &:= -i \log \left[(1 + u + w_2)^2 + w_1^2 - \frac{v_1^2}{4} - \frac{v_2^2}{4} + i(v_2(1 + u + w_2) - v_1 w_1) \right] \\ &\quad + i \left((1 + u + w_2)^2 + w_1^2 - 1 + \frac{v_1^2}{4} + \frac{v_2^2}{4} \right) - (v_1 w_1 + v_2(1 + u - w_2)) \end{aligned} \quad (25)$$

$$f(v_1, v_2, u, w_1, w_2) := e^{-\frac{1}{4}(v_1^2 + v_2^2) - i[v_1 w_1 + v_2(1 + u - w_2)]}. \quad (26)$$

By Lemma 2.1 and a standard calculation, we have

$$\operatorname{Im} \Psi(0) = 0, \quad \operatorname{Im} \Psi \geq 0, \quad \partial_{v_1} \Psi(0) = 0, \quad \partial_{v_1}^2 \Psi(0) = i.$$

This sets up an application of Theorem 5.2 where we integrate out v_1 . If necessary, we shrink the neighborhood K guaranteed by Theorem 5.2 in order to make the ideal generated by $\partial_{v_1} \Psi$ the same as the ideal generated by $v_1 + 2iw_1$. One can do this because

$$\partial_{v_1} \Psi = \frac{i}{2}(v_1 + 2iw_1) \left(\frac{1}{(1 + u + w_2)^2 + w_1^2 - \frac{v_1^2}{4} - \frac{v_2^2}{4} + i(v_2(1 + u + w_2) - v_1 w_1)} + 1 \right), \quad (27)$$

and the last term in the product vanishes only if

$$|v| = 2\sqrt{(1 + u + w_2)^2 + w_1^2 + 1}.$$

Since they are the same ideal, it is easy to see that $h^0(v_2, u, w_1, w_2) = h^0(-2iw_1, v_2, u, w_1, w_2)$ where h is either f , Ψ or $\partial_{v_1}^2 \Psi$. The amplitude function f in (26) is not in $C_0^\infty(K)$, but it is a Schwartz function in (v_1, v_2) and the theorem still applies by adding and subtracting a bump function whose support lies K . The remainder term is handled by Theorem 7.7.1 because by Lemma 2.1 and (27) we have that $\operatorname{Im} \Psi > 0$ or $\partial_{v_1} \Psi \neq 0$ outside of the support. Now applying Theorem 5.2 to the v_1 integral of (24), we compute

$$\begin{aligned} \frac{1}{N^{5/3}} W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}(x_{u,w_1,w_2}, \xi_{u,w_1,w_2}) &= \frac{N^{5/6}}{4\pi^3 \sqrt{2\pi} E^2} e^{1 - \frac{1}{E} |x_{u,w_1,w_2}|^2} \left(1 + O\left(\frac{1}{N}\right)\right) \\ &\int_{\mathbb{R}^2} f(v_1, v_2, u, w_1, w_2) e^{iN\Psi(v_1, v_2, u, w_1, w_2)} dv \\ &= \frac{N^{5/6}}{4\pi^3 \sqrt{2\pi} E^2} e^{1 - \frac{1}{E} |x_{u,w_1,w_2}|^2} \left(1 + O\left(\frac{1}{N}\right)\right) \\ &\quad \left(\frac{\sqrt{2\pi}}{\sqrt{N}} \int_{\mathbb{R}} \frac{f^0(v_2, u, w_1, w_2)}{\sqrt{-i(\partial_{v_1}^2 \Psi)^0(v_2, u, w_1, w_2)}} e^{iN\Psi^0(v_2, u, w_1, w_2)} dv_2 + O(N^{-3/2}) \right) \\ &= \frac{N^{5/6}}{4\pi^3 \sqrt{2\pi} E^2} e^{1 - \frac{1}{E} |x_{u,w_1,w_2}|^2} \left(1 + O\left(\frac{1}{N}\right)\right) \end{aligned}$$

$$\begin{aligned}
& \left(\frac{\sqrt{2\pi}}{\sqrt{N}} \int_{\mathbb{R}+2iw_2} \frac{f^0(v_2, u, w_1, w_2)}{\sqrt{-i(\partial_{v_1}^2 \Psi)^0(v_2, u, w_1, w_2)}} e^{iN\Psi^0(v_2, u, w_1, w_2)} dv_2 + O(N^{-3/2}) \right) \\
&= \frac{N^{1/3}}{4\pi^3 E^2} e^{(N+1)(-u^2-2u-2w_1^2-2w_2^2)} \left(1 + O\left(\frac{1}{N}\right)\right) \int_{\mathbb{R}} g(v_2, u) e^{iN\Phi(v_2, u)} dv_2 \\
&\quad + e^{1-\frac{1}{E}|x_{u, w_1, w_2}|^2} O(N^{-2/3})
\end{aligned} \tag{28}$$

where the penultimate equality follows from a change of contour, which we can do because of the decay of the integrand at infinity. For the last equality, we define

$$g(v_2, u) := \frac{\sqrt{2}e^{-\frac{1}{4}v_2^2 - iv_2(1+u)}}{\sqrt{\frac{1}{(1+u+i\frac{v_2}{2})^2} + 1}} \tag{29}$$

$$\begin{aligned}
\Phi(v_2, u) &:= -i \log \left[(1+u)^2 + iv_2(1+u) - \frac{v_2^2}{4} \right] + \frac{i}{4}v_2^2 - v_2(1+u) \\
&= -i \log \left[\left(1+u+i\frac{v_2}{2}\right)^2 \right] + \frac{i}{4}v_2^2 - v_2(1+u).
\end{aligned}$$

Note that if we view Φ as a function on \mathbb{C}^2 , we see that it is holomorphic in a neighborhood of $(0, 0)$. Calculating derivatives of Φ , we see

$$\partial_{v_2} \Phi = \frac{1}{1+u+i\frac{v_2}{2}} + \frac{i}{2}v_2 - (1+u) = -\frac{\frac{v_2^2}{4} + u(u+2)}{1+u+i\frac{v_2}{2}} \tag{30}$$

$$\partial_{v_2}^2 \Phi = \frac{i}{2} \left(1 - \frac{1}{(1+u+i\frac{v_2}{2})^2} \right) \tag{31}$$

$$\partial_{v_2}^3 \Phi = -\frac{1}{2} \frac{1}{(1+u+i\frac{v_2}{2})^3}. \tag{32}$$

From (30), we see by the quadratic formula that $\partial_{v_2} \Phi = 0$ if and only if

$$v_2 = v_{\pm}(u) := \pm 2i\sqrt{u(u+2)}.$$

Additionally, note that $\partial_{v_2} \Phi(0, 0) = \partial_{v_2}^2 \Phi(0, 0) = 0$ and $\partial_{v_2}^3 \Phi(0, 0) = -1/2$, so we can apply Theorem 5.3 (which follows from Theorem 2 of [CFU56]) to see

$$\begin{aligned}
e^{-Nb(u)} \int_{\mathbb{R}} g(v_2, u) e^{iN\Phi(v_2, u)} dv_2 &\sim \frac{1}{N^{1/3}} \text{Ai}(a(u)N^{2/3}) \sum_{s=0}^{\infty} \mu_{0s}(u) N^{-s} \\
&\quad + \frac{1}{N^{2/3}} \text{Ai}'(a(u)N^{2/3}) \sum_{s=0}^{\infty} \mu_{1s}(u) N^{-s}
\end{aligned}$$

where a and b are functions satisfying

$$i\Phi(v_2, u) = \frac{1}{3}T(v_2, u)^3 - a(u)T(v_2, u) + b(u) \tag{33}$$

where T is a function holomorphic in v_2 and u . From [CFU56], the functions a and b can be computed explicitly by

$$\begin{aligned}
b(u) &= \frac{i}{2} [\Phi(v_+(u), u) + \Phi(v_-(u), u)] = u(u+2) \\
a(u)^{3/2} &= \frac{3i}{4} [\Phi(v_+(u), u) - \Phi(v_-(u), u)] \\
&= \frac{3}{2} \left(2(u+1)\sqrt{u(u+2)} + \log [1+u-\sqrt{u(u+2)}] - \log [1+u+\sqrt{u(u+2)}] \right)
\end{aligned}$$

$$\begin{aligned}
&= 3(u+1)\sqrt{u(u+2)} - 3\log[1+u+\sqrt{u(u+2)}] \\
&= \begin{cases} 3(u+1)i\sqrt{|u|(u+2)} - 3i\arccos(1+u) & \text{if } u < 0 \\ 3(u+1)\sqrt{u(u+2)} - 3\operatorname{arccosh}(1+u) & \text{if } u \geq 0 \end{cases} \quad (34)
\end{aligned}$$

where the last line follows from the identities

$$\arccos(z) = -i\log(z + i\sqrt{1-z^2}), \quad \operatorname{arccosh}(z) = \log(z + \sqrt{z^2-1})$$

where the principal branches are used for the logarithm and square root functions. The branch in (34) is chosen so as to make

$$a(u) \approx 2^{5/3}u \quad \text{for small } u, \quad (35)$$

that is, we use the principal branch of the cube root (see the top of page 603 in [CFU56]). Now using (28), we have

$$\begin{aligned}
\frac{1}{N^{5/3}} W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}(x_{u,w_1,w_2}, \xi_{u,w_1,w_2}) &\sim \frac{(1 + O(\frac{1}{N}))}{4\pi^3 E^2} e^{-2(N+1)(w_1^2+w_2^2)} \left(\operatorname{Ai}(a(u)N^{2/3}) \sum_{s=0}^{\infty} \mu_{0s}(u)N^{-s} \right. \\
&\quad \left. + \frac{1}{N^{2/3}} \operatorname{Ai}'(a(u)N^{1/3}) \sum_{s=0}^{\infty} \mu_{1s}(u)N^{-s} \right) \\
&\quad + e^{1-\frac{1}{E}|x_{u,w_1,w_2}|^2} O(N^{-2/3}).
\end{aligned}$$

Now we need to compute μ_{00} . At the end of page 601 of [CFU56], we see

$$\frac{\mu_{00}(u)}{2\pi i} = \frac{1}{2} \left(g(v_+, u) (\partial_{v_2} T(v_+, u))^{-1} + g(v_-, u) (\partial_{v_2} T(v_-, u))^{-1} \right), \quad (36)$$

so it suffices to compute $\partial_{v_2} T(v_{\pm}, u)$. For this, by Theorem 2 of [CFU56], we see

$$T(v_{\pm}, u) = \mp a(u)^{1/2}. \quad (37)$$

Note that (37) cannot be $\pm a(u)^{1/2}$ for otherwise this would switch the sign of (39), contradicting (40). Now we differentiate (33) twice in v_2 to obtain

$$i\partial_{v_2}^2 \Phi = \partial_{v_2}^2 T(T^2 - a(u)) + (\partial_{v_2} T)^2 2T. \quad (38)$$

Evaluating at $v_2 = v_{\pm}$, we see

$$\begin{aligned}
i\partial_{v_2}^2 \Phi(v_{\pm}, u) &= \partial_{v_2}^2 T(v_{\pm}, u) (T^2(v_{\pm}, u) - a(u)) + (\partial_{v_2} T(v_{\pm}, u))^2 2T(v_{\pm}, u) \\
&\stackrel{(37)}{=} \mp 2 (\partial_{v_2} T(v_{\pm}, u))^2 a(u)^{1/2}.
\end{aligned}$$

Using (31), we see

$$(\partial_{v_2} T(v_{\pm}, u))^2 = \frac{1}{\mp 4a(u)^{1/2}} - 1 = -\frac{\pm u(u+2) + (1+u)\sqrt{u(u+2)}}{2a(u)^{1/2}}. \quad (39)$$

Since $\partial_{v_2} T$ is holomorphic, it suffices to find the sign of the square root. For this, note $(\partial_{v_2} T(0, 0))^2 = -2^{-4/3}$ by setting $u = 0$ in (39). On the other hand, by setting $u = 0$ in (33), we have

$$2\log(1 + i\frac{v_2}{2}) - \frac{i}{4}v_2^2 + v_2 = \frac{1}{3}T(v_2, 0)^3.$$

Dividing both sides by v_2^3 and letting $v_2 \rightarrow 0$, we see

$$-\frac{i}{4} = (\partial_{v_2} T(0, 0))^3, \quad (40)$$

which implies $\partial_{v_2} T(0, 0) = 2^{-2/3}i$, so the square root in (39) is the principal square root. In particular, we see $\mu_{00}(0) = 2^{5/3}\pi$ (for an explicit formula for $\mu_{00}(u)$, see Section 5.3). Lastly, we use Lemma 2.2 to rotate our initial point $(1, i) = ((1, 0), (0, 1))$ around the orbit giving the desired result for γ_0 . We then use (14) to obtain the result for general γ . Note that the statement of the Theorem is independent of how we extend (x, ξ) to an orthogonal basis (each vector with norm $\sqrt{2E}$) of $N_{(x, \xi)}\gamma$ with the vectors v_1, v_2 since the approximation only depends on the coefficients of these vectors, concluding the proof.

3. RELATION WITH THE EIGENSPACE PROJECTOR

Suppose Π_N is the eigenspace projector to the $\hbar(N+1)$ -eigenspace \mathcal{H}_N of (1). The orbital coherent states have the following reproducing formula.

PROPOSITION 3.1. *If m is the Haar measure on $SU(2)$ and μ is the metaplectic representation on $L^2(\mathbb{R}^2)$, then we have*

$$\Pi_N = (N+1) \int_{SU(2)} (\mu(U)\varphi_N^{\gamma_0}) \otimes (\mu(U)\varphi_N^{\gamma_0})^\dagger dm(U). \quad (41)$$

In particular, if ν is the normalized Fubini-Study measure on $\mathbb{C}P^1$, we have

$$\Pi_N = (N+1) \int_{\mathbb{C}P^1} \varphi_N^\gamma \otimes (\varphi_N^\gamma)^\dagger d\nu(\gamma). \quad (42)$$

Proof. We show (42) as the proof of (41) is the same. We do this by showing that the integral on the right-hand side of (42) is a constant times the identity using Schur's lemma (see Schur's lemma I on page 262 of [F89]). Let $\mu : SU(2) \rightarrow \mathcal{U}(\mathcal{H}_N)$ be the subrepresentation of the metaplectic representation restricted to $SU(2)$. It is a standard fact that this representation is irreducible since it is rotating homogeneous polynomials in two complex variables with constant degree in Fock space. Now for all $g \in SU(2)$ and $f \in \mathcal{H}_N$, we have

$$\begin{aligned} \mu(g) \cdot \int_{\mathbb{C}P^1} \langle \varphi_N^\gamma, f \rangle \varphi_N^\gamma d\nu(\gamma) &= \int_{\mathbb{C}P^1} \langle \varphi_N^\gamma, f \rangle \mu(g)[\varphi_N^\gamma] d\nu(\gamma) \\ &= \int_{\mathbb{C}P^1} \langle \varphi_N^{g^{-1}\cdot\gamma}, f \rangle \mu(g)[\varphi_N^{g^{-1}\cdot\gamma}] d\nu(\gamma) \\ &= \int_{\mathbb{C}P^1} \langle \mu(g^{-1})[\varphi_N^\gamma], f \rangle \varphi_N^\gamma d\nu(\gamma) \\ &= \int_{\mathbb{C}P^1} \langle \varphi_N^\gamma, \mu(g)[f] \rangle \varphi_N^\gamma d\nu(\gamma) \end{aligned}$$

where the first equality we use the Fubini-Tonelli theorem as $\mu(g)$ is given by integral formulas starting on page 191 of [F89]. For the second line, we use the fact that the Fubini-Study measure is $SU(2)$ -invariant. The third line follows because

$$\mu(g^{-1})[\varphi_N^\gamma] = \varphi_N^{g^{-1}\cdot\gamma}$$

up to a phase, which follows from (4). Schur's lemma gives

$$\Pi_N = c_N \int_{\mathbb{CP}^1} \varphi_N^\gamma \otimes (\varphi_N^\gamma)^\dagger d\nu(\gamma)$$

for some constant c_N , which can be computed as $N+1$ by taking the trace of both sides. \square

Taking Wigner distributions of both sides of (42) and using the Fubini-Tonelli theorem, we see that

$$W_{\Pi_N}(x, \xi) := \int_{\mathbb{R}^2} \Pi_N(x + \frac{v}{2}, x - \frac{v}{2}) e^{-\frac{i}{\hbar}\langle v, \xi \rangle} \frac{dv}{4\pi^2 \hbar^2} = (N+1) \int_{\mathbb{CP}^1} W_{\varphi_N^\gamma, \varphi_N^\gamma}(x, \xi) d\nu(\gamma). \quad (43)$$

The Wigner distributions W_{Π_N} were studied extensively in [HZ20]. Theorem 1.4 of [HZ20] proves interface Airy asymptotics for W_{Π_N} , and a consequence of their work is the following:

THEOREM 3.2. *For $u \in \mathbb{R}$ and $(x, \xi) \in \Sigma_E$, we have*

$$\lim_{N \rightarrow \infty} \frac{1}{N^{5/3}} W_{\Pi_N} \left[\left(1 + \frac{u}{(2N)^{2/3}}\right)(x, \xi) \right] = \frac{1}{2^{4/3} \pi^2 E^2} \text{Ai}(2u).$$

This compares nicely with Theorem 1.3. In fact, if we set $w_1 = w_2 = 0$ in Theorem 1.3, the same conclusion is arrived up to a factor of 2. Theorem 1.3 along with (43) shows that the interface Airy asymptotics of $W_{\Pi_N}(x, \xi)$ in [HZ20] are coming from the Airy asymptotics of $W_{\varphi_N^\gamma, \varphi_N^\gamma}(x, \xi)$ in 1.3 for γ in a small neighborhood of the orbit containing (x, ξ) .

4. GENERALIZATIONS TO d DIMENSIONS

The two dimensional coherent state that concentrates on γ_0 in configuration space is

$$\varphi_N^{\gamma_0}(x_1, x_2) = \frac{(N+1)^{\frac{N+1}{2}}}{E^{\frac{N+1}{2}} \sqrt{\pi N!}} (x_1 + ix_2)^N e^{-\frac{(N+1)}{2E}|x|^2} = \frac{1}{\hbar^{\frac{N+1}{2}} \sqrt{\pi N!}} (x_1 + ix_2)^N e^{-\frac{1}{2\hbar}|x|^2}.$$

where $E = \hbar(N+1)$. We can extend this coherent state to a coherent state on dimension d by multiplying by one-dimensional ground states:

$$\begin{aligned} \varphi_{N,d}^{\gamma_0}(x_1, \dots, x_d) &:= \varphi_N^{\gamma_0}(x_1, x_2) \frac{1}{(\hbar\pi)^{\frac{d-2}{4}}} e^{-\frac{1}{2\hbar}x_3^2 - \dots - \frac{1}{2\hbar}x_d^2} \\ &= \frac{1}{(\hbar\pi)^{\frac{d-2}{4}} \hbar^{\frac{N+1}{2}} \sqrt{\pi N!}} (x_1 + ix_2)^N e^{-\frac{1}{2\hbar}|x|^2}. \\ &= \frac{1}{\hbar^{N+\frac{d}{2}} \pi^{\frac{d}{4}} \sqrt{N!}} (x_1 + ix_2)^N e^{-\frac{1}{2\hbar}|x|^2}. \end{aligned}$$

Here we view γ_0 on the left hand side as γ_0 embedded in d -dimensional configuration space \mathbb{R}^d . Recall that the Wigner distribution $W_{\varphi_N^\gamma, \varphi_N^\gamma} : T^*\mathbb{R}^d \rightarrow \mathbb{R}$ of φ_N^γ is defined by

$$W_{\varphi_N^\gamma, \varphi_N^\gamma}(x, \xi) := \frac{1}{(2\pi)^d \hbar^d} \int_{\mathbb{R}^d} \varphi_N^\gamma(x + \frac{v}{2}) \overline{\varphi_N^\gamma(x - \frac{v}{2})} e^{-\frac{i}{\hbar}\langle v, \xi \rangle} dv.$$

Also recall that the one-dimensional ground state $g(x) = (\hbar\pi)^{-\frac{1}{4}} e^{-\frac{1}{2\hbar}x^2}$ has Wigner distribution

$$W_{g,g}(x, \xi) = \frac{1}{\pi \hbar} e^{-\frac{1}{\hbar}(x^2 + \xi^2)}.$$

We then have

$$\begin{aligned} W_{\varphi_{N,d}^{\gamma_0}, \varphi_{N,d}^{\gamma_0}}(x, \xi) &= \frac{1}{(\pi \hbar)^{d-2}} e^{-\frac{1}{\hbar}(x_3^2 + \xi_3^2 + \dots + x_d^2 + \xi_d^2)} W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}((x_1, x_2), (\xi_1, \xi_2)) \\ &= \left(\frac{N + \frac{d}{2}}{\pi E} \right)^{d-2} e^{-\frac{N + \frac{d}{2}}{E}(x_3^2 + \xi_3^2 + \dots + x_d^2 + \xi_d^2)} W_{\varphi_N^{\gamma_0}, \varphi_N^{\gamma_0}}((x_1, x_2), (\xi_1, \xi_2)), \end{aligned} \quad (44)$$

where $E = \hbar(N + \frac{d}{2})$. The non-constant Hamiltonian orbits of the d -dimensional isotropic harmonic oscillator are still great circles lying in two-dimensional planes, and we still have metaplectic covariance

$$W_{\mu(U)f, \mu(U)f}(x, \xi) = W_{f,f}(U^*(x, \xi)) \quad (45)$$

where $f \in L^2(\mathbb{R}^d)$, $U \in \text{SU}(d)$, and μ is the metaplectic representation on $L^2(\mathbb{R}^d)$. The non-constant Hamiltonian orbits can be similarly identified by $\mathbb{C}\mathbb{P}^{d-1}$. Since $\text{SU}(d)$ acts transitively on $\mathbb{C}\mathbb{P}^{d-1}$ by rotation, for any non-constant Hamiltonian orbit γ there exists a $U \in \text{SU}(d)$ such that $U \cdot \gamma_0 = \gamma$. We define (up to a phase) the coherent state centered at the Hamiltonian orbit γ by

$$\varphi_{N,d}^\gamma := \mu(U) \varphi_{N,d}^{\gamma_0}.$$

Using (45), we have

$$W_{\varphi_{N,d}^\gamma, \varphi_{N,d}^\gamma}(x, \xi) = W_{\varphi_{N,d}^{\gamma_0}, \varphi_{N,d}^{\gamma_0}}(U^*(x, \xi)). \quad (46)$$

With (44) along with (46), we can prove all of our results in dimension d . We state them below.

PROPOSITION 4.1. *Let $a : T^*\mathbb{R}^d \rightarrow \mathbb{R}$ be a smooth function with exponential decay. Then in the limit $N \rightarrow \infty$ where $E = \hbar(N + d/2)$, we have*

$$\int_{T^*\mathbb{R}^d} a(x, \xi) W_{\varphi_{N,d}^\gamma, \varphi_{N,d}^\gamma}(x, \xi) dx d\xi \xrightarrow{N \rightarrow \infty} \frac{1}{2\pi} \int_0^{2\pi} a(\gamma(t)) dt. \quad (47)$$

THEOREM 4.2. *As $N \rightarrow \infty$, we have the following pointwise asymptotics:*

(1) *If (x, ξ) lies on the orbit γ , then we have*

$$\frac{1}{N^{d-1/3}} W_{\varphi_{N,d}^\gamma, \varphi_{N,d}^\gamma}(x, \xi) = \frac{1}{2^{1/3} \pi^d E^d} \text{Ai}(0) + O(N^{-2/3}).$$

where $\frac{1}{2^{1/3} \pi^d E^d} \text{Ai}(0) = \frac{\Gamma(1/3) 12^{1/3}}{4\pi^{d+1} E^d \sqrt{3}}$
(2) *If (x, ξ) lies on the orbit γ , then for $0 < t < 1$ we have*

$$\frac{1}{N^{d-1/3}} W_{\varphi_{N,d}^\gamma, \varphi_{N,d}^\gamma}(t(x, \xi)) = O(N^{-1/6}).$$

(3) *If $(x, \xi) \neq 0$ is not in the above case, we have*

$$W_{\varphi_{N,d}^\gamma, \varphi_{N,d}^\gamma}(x, \xi) = O(N^{-\infty}).$$

(4) *If $(x, \xi) = 0$, then the Wigner distribution is explicitly*

$$W_{\varphi_{N,d}^\gamma, \varphi_{N,d}^\gamma}(x, \xi) = \frac{(-1)^N}{\pi^2 E^d} (N + 1)^d.$$

THEOREM 4.3. *Suppose (x, ξ) lies on γ , and let $(x, \xi), v_1, \dots, v_{2(d-1)}$ be an orthogonal basis (with each vector of norm $\sqrt{2E}$) for the normal space, $N_{(x, \xi)}\gamma$, of γ at (x, ξ) . For $(u, w_1, \dots, w_{2(d-1)}) \in \mathbb{R}^{2d-1}$ in a sufficiently small neighborhood of 0, we have the uniform asymptotic expansion*

$$\begin{aligned} & \frac{1}{N^{d-1/3}} W_{\varphi_{N,d}^\gamma, \varphi_{N,d}^\gamma} \left[(1+u)(x, \xi) + w_1 v_1 + \dots + w_{2(d-1)} v_{2(d-1)} \right] \\ &= \frac{e^{-2(N+1)(w_1^2 + \dots + w_{2(d-1)}^2)}}{4\pi^{d+1} E^d} \mu_{00}(u) \text{Ai}(a(u)N^{2/3}) + \text{Ai}'(a(u)N^{2/3}) \cdot O(N^{-2/3}) \end{aligned}$$

where

$$a(u)^{3/2} = \begin{cases} 3(u+1)i\sqrt{|u|(u+2)} - 3i \arccos(1+u) & \text{if } u < 0 \\ 3(u+1)\sqrt{u(u+2)} - 3 \operatorname{arccosh}(1+u) & \text{if } u \geq 0 \end{cases}$$

and $\mu_{00}(u)$ is a smooth function (explicitly given in the appendix) with $\mu_{00}(0) = 2^{5/3}\pi$. In particular, if we rescale $u \rightarrow \frac{u}{(2N)^{2/3}}$, $w_1 \rightarrow \frac{w_1}{(2N)^{1/2}}$, and $w_2 \rightarrow \frac{w_2}{(2N)^{1/2}}$, we have the pointwise limit

$$\lim_{N \rightarrow \infty} \frac{1}{N^{d-1/3}} W_{\varphi_{N,d}^\gamma, \varphi_{N,d}^\gamma} \left[\left(1 + \frac{u}{(2N)^{2/3}}\right)(x, \xi) + \frac{w_1}{(2N)^{1/2}} v_1 + \dots + \frac{w_{2(d-1)}}{(2N)^{1/2}} v_{2(d-1)} \right] = \frac{e^{-(w_1^2 + \dots + w_{2(d-1)}^2)}}{2^{1/3} \pi^d E^d} \text{Ai}(2u).$$

Lastly, the same proof of Proposition 3.1 works for dimension d , and we have similar comparisons to results in [HZ20].

5. APPENDIX

5.1. Airy Function. The Airy function $\text{Ai}(x)$ on \mathbb{R} is defined as the inverse Fourier transform of $\xi \mapsto e^{i\xi^3/3}$. This implies $\text{Ai} \in \mathcal{S}'(\mathbb{R})$, but if we view the integral as an improper Riemann integral on $[-L, L]$ with $L \rightarrow \infty$, we see that $\text{Ai} \in C^\infty(\mathbb{R})$ by an integration by parts argument. So we define

$$\text{Ai}(x) := \lim_{L \rightarrow \infty} \frac{1}{2\pi} \int_{-L}^L \cos(\xi^3/3 + \xi x) d\xi.$$

This function satisfies the differential equation

$$\text{Ai}''(x) - x \text{Ai}(x) = 0,$$

and it can be extended to an entire function on \mathbb{C} using this differential equation.

5.2. Stationary Phase Methods. For the pointwise asymptotics, we use a consequence of Theorem 7.7.5 of [H03], which we state below

THEOREM 5.1. *Suppose f, Ψ are C^∞ complex valued functions on \mathbb{R}^d . Suppose that*

$$\operatorname{Im} \Psi \geq 0, \operatorname{Im} \Psi(x_0) = 0, \Psi'(x_0) = 0, \det \Psi''(x_0) \neq 0,$$

with $\Psi' \neq 0$ in a neighborhood of x_0 . Then

$$\int_{\mathbb{R}^d} f(x) e^{iN\Psi(x)} dx = \left(\frac{2\pi}{N}\right)^{\frac{d}{2}} \det(-i\Psi''(x_0))^{-\frac{1}{2}} f(x_0) e^{iN\Psi(x_0)} + O(N^{-\frac{d}{2}-1})$$

provided that $\operatorname{supp} f$ is in a sufficiently small neighborhood of x_0 .

Also, for the scaling asymptotics we use Theorem 7.7.12 of [H03], which is the above theorem with an extra variable (see below).

THEOREM 5.2. *Let $\Psi(x, y)$ be a complex valued C^∞ function in a neighborhood of $(0, 0)$ in \mathbb{R}^{1+d} , satisfying*

$$\operatorname{Im} \Psi(0, 0) = 0, \quad \operatorname{Im} \Psi \geq 0, \quad \partial_x \Psi(0, 0) = 0, \quad \partial_x^2 \Psi(0, 0) \neq 0,$$

and let $f \in C_0^\infty(K)$ where K is a small neighborhood of $(0, 0)$ in \mathbb{R}^{1+d} . Then

$$\int_{\mathbb{R}} f(x, y) e^{iN\Psi(x, y)} dx = \frac{\sqrt{2\pi}}{\sqrt{N} \sqrt{-i(\partial_x^2 \Psi)^0(y)}} f^0(y) e^{iN\Psi^0(y)} + O(N^{-3/2})$$

where for functions $g(x, y)$ the notation $g^0(y)$ stands for a function of y only which is in the same residue class modulo the ideal generated by $\partial_x \Psi$.

The next theorem we reproduce is a consequence of Theorem 2 of [CFU56], which we use at a critical point in the proof of the scaling asymptotics.

THEOREM 5.3. *Let $\Psi(z, w)$ and $f(z, w) \in \mathcal{S}(\mathbb{C}^2)$ be complex valued functions holomorphic in both variables in a neighborhood of $(0, 0)$ in \mathbb{C}^2 , where Ψ satisfies $\partial_z \Psi(0, 0) = \partial_z^2 \Psi(0, 0) = 0$ and $\partial_z^3 \Psi(0, 0) \neq 0$. Further suppose that Ψ has only two nondegenerate critical points in z , $\pm z(w)$, that depend on w and differ by a sign that coalesce to 0 as w goes to 0 (i.e. $z(0) = 0$). Then uniformly in w , we have the asymptotic formula*

$$\begin{aligned} e^{-iNb(w)} \int_{\mathbb{R}} f(z, w) e^{iN\Psi(z, w)} dz &\sim N^{-1/3} \operatorname{Ai}(a(w)N^{2/3}) \sum_{k=0}^{\infty} u_{0k}(w) N^{-k} \\ &\quad + N^{-2/3} \operatorname{Ai}'(a(w)N^{2/3}) \sum_{k=0}^{\infty} u_{1k}(w) N^{-k}, \end{aligned}$$

provided that w is in a sufficiently small neighborhood of 0. The functions a, b, u_{jk} are holomorphic on this neighborhood.

This theorem is the holomorphic version of Theorem 7.7.18 of [H03]. The functions a and b are related to Ψ by the equation

$$i\Psi(z, w) = \frac{1}{3}T(z, w)^3 - a(w)T(z, w) + b(w)$$

where T is a holomorphic function in z and w . For more information, see [CFU56]. The last theorem is a stationary phase approximation for Bott-Morse functions, and we use it in the weak limit proof.

THEOREM 5.4 (Stationary Phase for Bott-Morse functions). *Let $U \subset \mathbb{R}^d$ be an open set, and let $f, \Psi \in C^\infty(U)$ be such that $\operatorname{Im} \Psi \geq 0$ in U and $\operatorname{supp} f \subset U$. We define*

$$C := \{u \in U \mid \operatorname{Im} \Psi(u) = 0, \Psi'(u) = 0\}.$$

Assume that C is a smooth, compact, and connected manifold of \mathbb{R}^d of dimension k such that for all $u \in C$ the Hessian, $\Psi''(u)$, of Ψ is nondegenerate on the normal space $N_u C$ to C at u . Then we have

$$\int_{\mathbb{R}^d} f(x) e^{iN\Psi(x)} dx = \left(\frac{2\pi}{N}\right)^{\frac{d-k}{2}} e^{i\Psi(u_0)} \int_C \frac{f(u)}{\sqrt{\det(-i\Psi''|_{N_u C})}} d\mu_C(u) + O(N^{-\frac{d-k}{2}-1})$$

where $u_0 \in C$ is arbitrary, μ_C denotes the natural Riemannian metric on C , and the denominator of the integrand is the product of the square roots of the eigenvalues of $-i\Psi''|_{N_u C}$, chosen with positive real part.

For the proof of this theorem, see Theorem 29 on page 134 of [CR12]

5.3. **Formula for $\mu_{00}(u)$.** From (36) in Section 2.4, we have

$$\frac{\mu_{00}(u)}{2\pi i} = \frac{1}{2} \left(g(v_+, u) (\partial_{v_2} T(v_+, u))^{-1} + g(v_-, u) (\partial_{v_2} T(v_-, u))^{-1} \right),$$

where g is defined at (29) by

$$g(v, u) := \frac{\sqrt{2} e^{-\frac{1}{4}v_2^2 - iv_2(1+u)}}{\sqrt{\frac{1}{(1+u+i\frac{v_2}{2})^2} + 1}},$$

and (39) gives

$$\partial_{v_2} T(v_{\pm}, u) = \sqrt{\frac{\frac{1}{(1+u+iv_{\pm}/2)^2} - 1}{\mp 4a(u)^{1/2}}} = i \sqrt{\frac{\pm u(u+2) + (1+u)\sqrt{u(u+2)}}{2a(u)^{1/2}}}.$$

Combing all of these equations, we have

$$\begin{aligned} \mu_{00}(u) &= 2\pi |a(u)|^{1/4} \frac{e^{u(u+2)}}{\sqrt{2u(u+1)(u+2)}} \left| e^{2(u+1)\sqrt{u(u+2)}} \sqrt{\frac{-u(u+2) + (u+1)\sqrt{u(u+2)}}{1+u+\sqrt{u(u+2)}}} \right. \\ &\quad \left. + e^{-2(u+1)\sqrt{u(u+2)}} \sqrt{\frac{u(u+2) + (u+1)\sqrt{u(u+2)}}{1+u-\sqrt{u(u+2)}}} \right|. \end{aligned}$$

REFERENCES

- [CFU56] Chester, C.; Friedman, B.; Ursell, F. An extension of the method of steepest descents. *Proc. Cambridge Philos. Soc.* 53 (1957), 599-611.
- [CR12] Combescure, M.; Robert, D. Coherent states and applications in mathematical physics. Theoretical and Mathematical Physics. *Springer, Dordrecht*, 2012.
- [De92] De Bièvre, S. (1992). Oscillator eigenstates concentrated on classical trajectories. *J. Phys. A* 25(1992), no. 11, 3399-3418.
- [De93] De Bièvre, S.; Houard J.-C.; Irac-Astaud, M. Wave packets localized on closed classical trajectories. *Differential equations with applications to mathematical physics*, 25-32, Math. Sci. Engrg., 192, *Academic Press, Boston, MA*, 1993.
- [F89] Folland, G. B. Harmonic Analysis in Phase Space. Annals of Mathematics Studies, 122. *Princeton University Press, Princeton, NJ*, 1989.
- [GDB89] Gay, J.-C., Delande, D., Bommier, A. Atomic quantum states with maximum localization on classical elliptic orbits. *Phys. Rev. A* 39 (1989), no. 12, 6587-6590.
- [G63] Glauber, Roy J. The quantum theory of optical coherence. *Phys. Rev.*(2) 13 (1963), 2529-2539.
- [HZ20] Hanin, B., Zelditch, S. Interface asymptotics of eigenspace Wigner distributions for the harmonic oscillator. *Comm. Partial Differential Equations* 45 (2020), no. 11, 1589-1620.
- [H03] Hörmander, L. The analysis of linear partial differential operators. I. Distribution theory and Fourier analysis. Reprint of the second (1990) edition [Springer, Berlin; MR1065993]. Classics in Mathematics. *Springer-Verlag, Berlin*, 2003.
- [K96] Klauder, John R. Coherent states for the hydrogen atom. *J. Phys. A* 29 (1996), no. 12, L293-L298.
- [N89] Nauenberg, M. Quantum wave packets on Kepler orbits. *Phys. Rev. A* (3) 40 (1989), no. 2, 1133-1136.
- [S26] Schrödinger, E. Der stetige übergang von der Mikro- zur Makromechanik. *Naturwissenschaften* 14 (1926), 664-666.
- [S93] Stein, E. Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals. With the assistance of Timothy S. Murphy. Princeton Mathematical Series, 43. Monographs in Harmonic Analysis, III. *Princeton University Press, Princeton, NJ*, 1993.
- [TVB97] Thomas, L. E., Villegas-Blas, C. Asymptotics of Rydberg states for the hydrogen atom, *Commun. Math. Phys.* 187 (1997), no. 3, 623-645.
- [VB96] Villegas-Blas, C. The Laplacian on the n -sphere, the hydrogen atom and the Bargmann space representation. Thesis (Ph.D.)—University of Virginia. 1996.
- [W32] Wigner, E. On the quantum correction for thermodynamic equilibrium. *Phys. Rev.* 40 (1932), no. 5, 749-759.