
On A Mallows-type Model For (Ranked) Choices

Yifan Feng

Department of Analytics and Operations
NUS Business School
yifan.feng@nus.edu.sg

Yuxuan Tang

Institute of Operations Research and Analytics
National University of Singapore
yuxuan.tang@u.nus.edu

Abstract

In a preference learning setting, every participant chooses an ordered list of k most preferred items among a displayed set of candidates. (The set can be different for every participant.) We identify a distance-based ranking model for the population's preferences and their (ranked) choice behavior. The ranking model resembles the Mallows model but uses a new distance function called Reverse Major Index (RMJ). We find that despite the need to sum over all permutations, the RMJ-based ranking distribution aggregates into (ranked) choice probabilities with simple closed-form expression. We develop effective methods to estimate the model parameters and showcase their generalization power using real data, especially when there is a limited variety of display sets.

1 Introduction

How to aggregate the population's preferences from their (ranked) choices out of different choice sets? This question is of interest to many communities, such as economics, business, and computer science. A concrete setting is a platform that wishes to learn customer preferences over a universe of n product prototypes. The platform is able to display different subsets of versions to different customers, who then provide feedback in the form of a top- k ranked list of their most preferred items within the subset they see (hereafter referred to as "ranked choices").¹

The population's (ranked) choice behavior could be summarized as a (*ranked*) *choice model* $\Pr(\pi_k|S)$, which specifies the probability that a randomly drawn participant choosing a top- k list π_k from the display set S . An economically rationalizable and yet very general way to model (ranked) choices is to use probabilistic *ranking models*. That is, given a probability distribution λ over preference rankings, $\Pr(\pi_k|S)$ equals the probability that a randomly drawn participant would place π_k as the top- k positions among the items in S .

A popular family of ranking models is *distance-based*, which is the conceptual analog of Gaussian distribution for scalars; see [1]. A distance-based ranking model is specified by a modal (central) ranking π^* and a dispersion parameter $q \in (0, 1)$. Given a ranking π , its probability of being sampled is proportional to $q^{d(\pi^*, \pi)}$. Here $d(\pi_1, \pi_2)$ is a *distance function* that describes the discrepancy of ranking π_1 from π_2 , and different distance functions lead to different models. The most representative distance-based ranking model is the *Mallows model* ([2]), which uses the Kendall-Tau distance as its distance function. It has been studied extensively in the literature regarding topics such as sampling, estimation from sampled (partial) rankings, and learning in a Mallows mixture setting; see [3] and references therein. Due to its popularity and the fact that every distance-based ranking model only differs from the Mallows by distance function, we will also refer to a distance-based ranking model as a *Mallows-type model* interchangeably in the sequel.

¹If $k = 1$, each participant is asked to choose the most preferred candidate. That effectively reduces to a (single) choice, which is a feedback structure extensively studied in the choice modeling literature.

Mallows-type models could be used as “kernels” to “smooth out” the distribution over rankings. As such, they could help mitigate the overfitting issues of generic (i.e., nonparametric) ranking models, which are typically overparameterized. Yet, Mallows models can be highly expressive in a mixture setting: A mixture of Mallows-type models can approximate *any* probability distribution over rankings by an arbitrary precision (as q tends to zero and the number of clusters tends to infinity). Therefore, the Mallows-type model family is a helpful tool to balance capturing (complex) preference heterogeneity across individuals and regularizing the ranking distribution for better out-of-sample predictions. A representative work is by [4], who use the Mallows model to aggregate customer preferences from their choices (i.e., $k = 1$).

Despite the theoretical elegance, the main challenge in applying Mallows-type ranking models to (ranked) choice modeling is analytical and computational tractability. More specifically, $\Pr(\pi_k|S)$ is calculated from summation over all rankings subject to nontrivial conditions. Therefore, even if a Mallows-type ranking model has a simple structure for λ , the resulting choice probabilities $\{\Pr(\pi_k|S)\}$ can be difficult to obtain even when $k = 1$. The state-of-art results are obtained by [4], who develop polynomial-time numerical algorithms to compute $\{\Pr(\pi_k|S)\}$ for $k = 1$ under the Mallows model. The *estimation* problem is even more difficult, which involves finding the central ranking π^* and dispersion parameters q that best explain the (ranked) choice data. To the best of our knowledge, no effective methods to estimate *any* Mallows-type ranking model from (ranked) choice data are known. (Perhaps the best method to date is again by [4], who use a “Mallows smoothing” heuristic to conduct the estimation when $k = 1$, which we will discuss later.)

Summary of results and contributions. This paper identifies and studies a new distance-based (i.e., Mallows-type) ranking model. It is the same as the Mallows model except that it builds on a new distance function (i.e., smoothing kernel), which we call *reverse major index (RMJ)*. Unlike the Mallows model’s Kendall-Tau distance (which weighs all pairwise disagreements equally), RMJ puts more weight on top-position deviations.

The RMJ-based ranking model is a small conceptual deviation from the Mallows model (and, in particular, enjoys the desired properties mentioned above, such as rationalizability, smoothing, and expressive power); see Section 2. However, this twist brings a significant advantage in both analytical and computational simplicity. Specifically, we solve a list of problems under the RMJ-based ranking distribution. That includes:

- *Characterizing (ranked) choice probabilities:* Given $k \geq 1$, calculating $\{\Pr(\pi_k|S)\}$;
- *Sampling:* Given $k \geq 1$, efficiently sampling a top- k list π_k ;
- *Parameter learning:* Estimating the central ranking π^* and dispersion parameter q from the (ranked) choice data through a maximum likelihood estimator (MLE) formulation ;
- *Learning in a mixture setting:* Assuming that there are multiple clusters of participant preferences, learning the central ranking and dispersion parameter for each cluster from choice data.

The solutions to the problems above can be implemented relatively easily. First, we are able to obtain $\{\Pr(\pi_k|S)\}$ in *simple* and *closed-form* expressions for all $k \geq 1$; see Theorems 1 and 4. This is in contrast to, say, its counterpart for the Mallows model ([4]), and we view it as our main theoretical achievement. Second, the sampling can be done in $O(nk)$ time directly; see Lemma 2. Third, the estimation problem can be reduced to a well-studied ranking-aggregation-type formulation; see (5) and Theorem 5. Many off-the-shelf tools are available. For example, it admits a PTAS and can be practically solved via a linear integer programming formulation. The estimation is guaranteed to recover the model parameters asymptotically under mild conditions on the coverage of display sets; see Theorem 2. This stands in contrast to Mallows Smoothing by [4], which cannot recover the Mallows central ranking even under sufficient coverage; see Theorem 3. Finally, the learning problem in a mixture setting can be solved using the standard Expectation-Maximization (EM) algorithm.

We demonstrate the practical effectiveness of our methodology on two data sets on customer preference over different types of sushi. When $k = 1$, we compare it with two representative ranking-based choice models: one based on the Mallows and the other on Plackett-Luce (which leads to the Multinomial Logit choice model). Our tools display superb generalization power, especially when there is a limited variety of display sets in the choice data. When $k > 1$, we demonstrate the robustness of the methodology. (It is difficult to find direct comparisons for prediction power.) Specifically, as long as the underlying population keeps the same, different top- k lists aggregate into the same

central rankings. We also show that our methodology can handle a relatively large number of items ($n = 100$) by achieving high precision solutions ($< 2\%$ optimality gap) in a reasonable time (< 5 minute solving time).

Related literature. Our paper is most closely related to the (extensive) literature on Mallows-type ranking models and their applications to preference and choice modeling (e.g., [1, 5, 3, 6, 7, 8, 9, 4, 10] to name a few.) Besides the new distance function and tractability results explained above, the most notable difference of this paper is the *feedback structure*. In our paper, every participant chooses from an *arbitrary* display set (as opposed to pairwise or the full display set only), and their feedback is in the form of a top- k ranked list (as opposed to single choices or full rankings). To the best of our knowledge, the combination of both generalities makes the setting the first of its kind.² It is also worth mentioning that the analytical and computational simplicity of our model makes it a convenient building block for further the subsequent optimization problems (e.g., new product introduction). It is an advantage that many common methods (e.g., Monte-Carlo algorithms) do not have.

Also related is the literature on *learning to rank* (e.g., [12, 13]). We wish to stress that we do not merely fit a ranking from the data but also the uncertainty quantification of the estimate (and the whole choice model $\{\Pr(\pi_k|S)\}$). Finally, our paper is also closely related to that of [11]. They study a choice model, called the “ordinal attraction model” (OAM), that emerges from an active learning problem of consumer preferences. We rationalize the (surprisingly simple) OAM by showing that it is equivalent to the aggregated choice probabilities $\{\Pr(\pi_k|S)\}$ from the RMJ-based ranking model when $k = 1$. We view it as a nontrivial observation in its own right.

2 A Ranking Model Based on Reverse Major Index

Preliminary. We consider a universe of n candidates (hereafter referred to as “items”), represented by $[n] = \{1, 2, \dots, n\}$. We assume that every participant has a strict preference over them items, represented by a ranking (permutation). We use a bijection $\pi : [n] \rightarrow [n]$ to represent a ranking where $\pi(i)$ is the i th most preferred item. We sometimes find the notation $\sigma := \pi^{-1}$ helpful, where $\sigma(x)$ is the position of item x in the ranking σ . We also use $x \succ_{\pi} y$ if item x is preferred to item y under π , i.e., $\pi^{-1}(x) < \pi^{-1}(y)$. For example, $\pi = (3, 1, 2)$ means $3 \succ_{\pi} 1 \succ_{\pi} 2$ and it corresponds to a “ σ ” notation of $\sigma = (2, 3, 1)$. Finally, we use e to represent the identity ranking, Σ for the set of all rankings over n items, π_k ($k \geq 1$) for a top- k list, and Σ_k for the set of all top- k rankings.

Mallows-type ranking models. Given the distance function $d(\cdot, \cdot)$, the probability mass function of the ranking for a Mallows-type model can be written as

$$\lambda(\pi) = \frac{q^{d(\pi^*, \pi)}}{\sum_{\tilde{\pi}} q^{d(\pi^*, \tilde{\pi})}},$$

where π^* is the central ranking, q is the dispersion parameter, and $\sum_{\tilde{\pi}} q^{d(\pi^*, \tilde{\pi})}$ is the normalization constant. Intuitively, the Mallows-type model defines a population of participants whose preferences are “similar” as they are centered around a common ranking, where the probability for deviations thereof decreases exponentially.

Different distance functions correspond to different models. A common requirement for a valid distance function is that it is invariant to “relabeling.” Formally, that means $d(\cdot, \cdot)$ is left-invariant under the ranking composition, i.e., $d(\pi_1, \pi_2) = d(\pi\pi_1, \pi\pi_2)$ for every π, π_1, π_2 .³ This invariance property enables to make the following conventions without loss of generality: First, we assume that the items have been properly relabeled so that the central ranking $\pi^* = e$ (unless otherwise specified). Second, we may use $d(\pi)$ as shorthand notation for $d(e, \pi)$, which fully represents a distance function with the knowledge that $d(\pi_1, \pi_2) = d(e, \pi_1^{-1}\pi_2)$.

²This setting is also practically relevant. For example, the platform may have capacity constraints on displaying how many items. The platform may also have an incentive to judiciously select the display sets to make feedback collection more efficient; see [11].

³Alternatively, one could use the “ σ ” notation and write the distance function as $\tilde{d}(\sigma_1, \sigma_2) := d(\pi_2, \pi_1)$ and \tilde{d} will be right-invariant under ranking composition.

The Mallows (Kendall’s Tau distance based) ranking model. Commonly studied distance functions include the Spearman’s rank correlation, Spearman’s Footrule, and the Kendall’s Tau distance ([1]). Among those three, Kendall’s Tau corresponds to the Mallows model and is defined as

$$d_K(\pi) = \sum_{i=1}^{n-1} \sum_{j=i+1}^n \mathbb{I}\{\pi(i) > \pi(j)\}. \quad (1)$$

It measures a ranking’s total number of pairwise disagreements (with the identity ranking e). For example, consider the ranking $\pi = (4, 2, 1, 3)$. There are four pairwise disagreements: $\{(4, 2), (4, 1), (4, 3), (2, 1)\}$. Therefore, $d_K((4, 2, 1, 3)) = 4$.

As an appealing property, the Kendall’s tau distance leads to a tractable expression for λ . Other common distances do not have this since the normalization constant involves summing over $n!$ rankings. The constant under the Mallows model can be expressed as (see [14]):

$$\sum_{\pi} q^{d_K(\pi)} = \psi(n, q) := \prod_{i=1}^n \frac{1-q^i}{1-q} = \prod_{i=1}^n (1 + q + \dots + q^{(i-1)}). \quad (2)$$

The RMJ based ranking model. In this paper, we identify a new distance function, which we call *reverse major index (RMJ)*. It is defined as

$$d_R(\pi) := \text{Reverse Major Index}(\pi) = \sum_{i=1}^{n-1} \mathbb{I}\{\pi(i) > \pi(i+1)\} \cdot (n-i) \quad (3)$$

Compared to Kendall’s Tau in (1), RMJ focuses on *adjacent* disagreements and puts more weight on top-position disagreements. For example, consider the ranking $\pi = (4, 2, 1, 3)$ again. The only adjacent disagreements are $\{(4, 2), (2, 1)\}$. Therefore, after including the weights in (3), we have $d_R((4, 2, 1, 3)) = 3 + 2 = 5$. The name of RMJ is inspired by the *major index* from the combinatorics literature ([15]), which is defined as $d_M(\pi) = \sum_{i=1}^{n-1} \mathbb{I}\{\pi(i) > \pi(i+1)\} \cdot i$.

Discussion: Kendall’s Tau vs. RMJ. Both metrics are conceptually similar: they measure a ranking’s deviation from the identity. They coincide in many intuitive cases. For example, in the most extreme ones we have $d_K(e) = d_R(e) = 0$ and $d_K((n, \dots, 1)) = d_R((n, \dots, 1)) = n(n-1)/2$. Therefore, both can be used as “kernels” to smooth out the distribution over rankings. However, they emphasize the deviation in a subtly different way. Therefore, we can also find subtle rankings where their values are different. For example, $d_K((4, 2, 1, 3)) = 4$ but $d_R((4, 2, 1, 3)) = 5$.

It is difficult to tell which kernel is “better” from an axiomatic approach as both distance functions satisfy the basic axioms for ranking distances ([1]). Nevertheless, we will show later that RMJ-based ranking model leads to simple and estimatable (ranked) choice probabilities without compromising the desirable properties of Mallows, such as rationalizability and flexibility for a mixture setting. Moreover, regarding its ability to describe preference distributions that may occur in practice, we will demonstrate its descriptive and predictive power in a case study with real preference data. Therefore, we believe that RMJ produces a promising tool that is more tailed to the application of learning population preferences from (ranked) choice data.

3 Analysis of the (Ranked) Choice Model $\{\text{Pr}(\pi_k|S)\}$

In this section, we will characterize the (ranked) Choice Model $\{\text{Pr}(\pi_k|S)\}$ aggregated from the RMJ-based ranking model, formally defined as

$$\text{Pr}(\pi_k|S) = \sum_{\tilde{\pi}} \lambda(\tilde{\pi}) \mathbf{I}(\pi_k, \tilde{\pi}, S),$$

where $\mathbf{I}(\pi_k, \tilde{\pi}, S)$ means the top- k list π_k is compatible with the ranking $\tilde{\pi}$ in the set S . That is, $\pi_k(i) \in S$ and $\pi_k(i) \succ_{\tilde{\pi}} x$ for all position $i \in \{1, \dots, k\}$ and item $x \in S \setminus \{\pi_k(1), \dots, \pi_k(i)\}$. Note that the summation is over rankings with nontrivial conditions. Therefore it is unclear *in priori* whether *any* distance-based ranking model can aggregate into a tractable $\text{Pr}(\pi_k|S)$ (even for $k = 1$).

We will also discuss how to estimate the ranked choice model parameters from data. In the sequel, we will write $d(\cdot) = d_R(\cdot)$ since RMJ is the distance function of interest.

3.1 The $k = 1$ case

When $k = 1$, a top- k list model reduces to a choice model, which connects to a richer literature (e.g., [4]). Therefore, it is worth a separate discussion, which also helps build intuition for the $k > 1$ case.

Choice probabilities. Our main result for $k = 1$ is summarized below, which characterizes the choice probabilities under the RMJ-based ranking model.

Theorem 1 *Let a display set $S = \{x_1, x_2, \dots, x_M\}$ be such that $x_1 < x_2 < \dots < x_M$. Then*

$$Pr(\{x_i|S\}) = \frac{q^{i-1}}{1 + q + \dots + q^{M-1}}. \quad (4)$$

In other words, for every display set, all the items within the display set are re-ranked so that their choice probabilities decay exponentially fast according to their *relative ranking* within the display set. Noticeably, the choice probabilities in (4) are (much) simpler than that induced by the Mallows model, which even needs a Fast Fourier Transform to evaluate in $O(n^2 \log n)$ time ([14]). Also, (4) rediscovers the ‘‘Ordinal Attraction Model’’ (OAM) in [11], which could also be viewed as a ‘‘multiwise’’ generalization of pairwise noisy comparison models (e.g., [16, 17]). OAM gets its name because the ‘‘attractiveness’’ (i.e., choice probability) of an item within a display set S only depends on its *relative* position in S and therefore is only a function of the ‘‘ordinal’’ information. While OAM emerges from an active learning problem of consumer preferences, we ‘‘rationalize’’ it by showing that it can be aggregated from the RMJ-based ranking model, which we believe is a nontrivial observation in its own right. In the sequel, we will follow its convention and refer to the choice model defined in Theorem 1 as OAM.

Proof outline and key intermediate results for Theorem 1. Let us introduce a few notations for top- k (sub-) rankings. Given a top- k ranking $\pi_k \in \Sigma_k$, we could define RMJ for π_k by truncating the index at position k , i.e., $d(\pi_k) = \sum_{i=1}^{k-1} \mathbb{I}\{\pi_k(i) > \pi_k(i+1)\} \cdot (n - i)$. In addition, let $R(\pi_k) := \{\pi_k(i) : i = 1, \dots, k\}$ be the set of top- k items and $R^c(\pi_k) := [n] \setminus R(\pi_k)$ its complement. Finally, let $L(\pi_k) := |\{x \in R^c(\pi_k) : x < \pi_k(k)\}|$ be the number of items that are (i) not included in π_k and (ii) having smaller indices than (i.e., preferred under $\pi^* = e$) item $\pi_k(k)$. For example, suppose $n = 7$ and $\pi_4 = (7, 4, 6, 2)$. Then $d(\pi_4) = 6 + 4 = 10$, $R(\pi_4) = \{2, 4, 6, 7\}$, $R^c(\pi_4) = \{1, 3, 5\}$, $\pi_4(4) = 2$, and $L(\pi_4) = |\{1\}| = 1$.

Given two subrankings $\pi_k \in \Sigma_k$ and $\pi_{k'} \in \Sigma_{k'}$ with $k \leq k'$, we write $\pi_k \subseteq \pi_{k'}$ if they are compatible, i.e., $\pi_k(i) = \pi_{k'}(i)$ for all $i = 1, \dots, k$. Our first result concerns extending the domain of λ to top- k rankings, formally defined as $\lambda(\pi_k) := \sum_{\tilde{\pi}} \lambda(\tilde{\pi}) \mathbb{I}\{\pi_k \subseteq \tilde{\pi}\}$.

Lemma 1 (Probability distribution of top- k rankings) $\lambda(\pi_k) = q^{d(\pi_k) + L(\pi_k)} \cdot \frac{\psi(n-k, q)}{\psi(n, q)}$.

The significance of the result above is that we could think of a ranking π as a stochastic process on a ‘‘tree’’ with depth n and $n!$ leaves. While $\{\lambda(\pi) : \pi \in \Sigma\}$ describe the probability distribution over the leaves, for every k , $\{\lambda(\pi_k) : \pi_k \in \Sigma_k\}$ describe the probability distribution over the nodes at level k .

As a consequence of Lemma 1, we can write out how to randomly sample a ranking π under the RMJ-based ranking model from the top to the bottom position. Given a top- k ranking π_k and an item z , we write $\pi_k \oplus z$ as the concatenation of π_k and item z . For example, suppose $\pi_3 = (5, 2, 4)$, then $\pi_3 \oplus 3 = (5, 2, 4, 3)$.

Lemma 2 (Random ranking generation) *Given a top- k ranking π_k such that $\pi_k(k) = z$, the conditional probability for the $(k+1)$ -positioned item is*

$$Pr(\pi_{k+1} = \pi_k \oplus y | \pi_k) := \frac{\lambda(\pi_k \oplus y)}{\lambda(\pi_k)} = \frac{q^{h(y|z)-1}}{1 + q + \dots + q^{n-k-1}},$$

$$\text{where } h(y|z) = \begin{cases} \sum_{x \in R^c(\pi_k)} \mathbb{I}\{z < x \leq y\} & \text{if } y > z \\ n - k - \sum_{x \in R^c(\pi_k)} \mathbb{I}\{y < x < z\} & \text{if } y < z \end{cases}.$$

Finally, Lemmas 1 and 2 leads to the following result based on an induction argument.

Lemma 3 *Let a display set $S = \{x_1, x_2, \dots, x_M\}$ be such that $x_1 < x_2 < \dots < x_M$ and a top- k ranking π_k be such that $\pi_k = \pi_{k-1} \oplus z$ and $R(\pi_{k-1}) \cap S = \emptyset$. Then conditional on a participant's top- k preference list is π_k , the probability that (s)he will choose item x_i out of display set S is*

$$Pr(\{x_i|S\}|\pi_k) := \frac{\sum_{\tilde{\pi}} \lambda(\tilde{\pi}) \cdot \mathbf{I}(x_i, \tilde{\pi}, S) \cdot \mathbb{I}\{\pi_k \subseteq \tilde{\pi}\}}{\lambda(\pi_k)} = \begin{cases} \frac{q^{M-p(z|S)+i-1}}{1+q+\dots+q^{M-1}} & \text{if } z > x_i \text{ and } z \notin S, \\ \frac{q^{i-p(z|S)-1}}{1+q+\dots+q^{M-1}} & \text{if } z < x_i \text{ and } z \notin S, \\ 1 & \text{if } z = x_i, \\ 0 & \text{if } z \in S \setminus \{x_i\}. \end{cases}$$

where $p(z|S) := \sum_{x \in S} \mathbb{I}\{x < z\}$.

Note that Lemma 3 could be viewed as a generalization of Theorem 1 by setting $\pi_k = \emptyset$ (which corresponds to $z = 0$).

Parameter learning from choice data. In the parameter learning problem, we are endowed with choice data $H_T = (S_1, x_1, \dots, S_T, x_T)$, where S_t is the display set shown to the t th participant and x_t is his/her choice.

Following [11], the maximum likelihood estimator (MLE) for the central ranking π^* can be obtained from following *choice aggregation* problem

$$\hat{\pi} \in \operatorname{argmin}_{\pi} \sum_{t=1}^T \sum_{x \in S_t} \mathbb{I}\{x \succ_{\pi} x_t\}.$$

It has a further linear integer programming formulation. Let $w_{ij} := \sum_{t=1}^T \mathbb{I}\{\{i, j\} \subseteq S_t \text{ and } x_t = i\}$ be the number of times that item i and item j are displayed together and item i is chosen. Intuitively, a positive $w_{ij} - w_{ji}$ is an indication that item i should be preferred to item j . Invoking Proposition 3 in [11], the integer programming could be written as following:

$$\begin{aligned} \hat{x} \in \operatorname{argmin}_x \quad & \sum_{(i,j):i \neq j} w_{ij} x_{ji} \\ \text{s.t.} \quad & x_{ij} + x_{ji} = 1 \quad \forall 1 \leq i, j \leq n, i \neq j \\ & x_{ij} + x_{jr} + x_{ri} \leq 2 \quad \forall 1 \leq i, j, r \leq n, i \neq j \neq r \\ & x_{ij} \in \{0, 1\} \quad \forall 1 \leq i, j \leq n \end{aligned} \quad (5)$$

In the formulation above, the solution \hat{x} is such that $\hat{x}_{ij} = 1$ if $i \succ_{\hat{\pi}} j$ under the MLE $\hat{\pi}$. Computationally, this integer programming is an instance of the well-studied *feedback arc set problem on tournaments*. Therefore, it admits a polynomial-time approximation scheme (PTAS). From a more practical side, the central ranking π^* can be effectively obtained using off-the-shelf integer programming solvers and with many speeding-up heuristics; see [11] for more discussion. The MLE for dispersion parameter q can be subsequently obtained from an one-dimensional (convex) optimization problem $\hat{\alpha} \in \operatorname{argmin}_{\alpha} \{\alpha \sum_{t=1}^T \sum_{x \in S_t} \mathbb{I}\{x \succ_{\hat{\pi}} x_t\} + \sum_{t=1}^T \log \sum_{j=0}^{|S_t|-1} e^{-j\alpha}\}$ so that $\hat{\alpha} = -\ln \hat{q}$. It is also worth noting that the MLE framework can be easily extended to learning in a mixture setting using the standard EM algorithm, which we refer to the supplementary materials for more details.

Let us conclude this section by providing some theoretical understanding of whether we could recover the ground truth values of π^* and q at least asymptotically. Intuitively, this should depend on the ‘‘coverage’’ of display sets: if only a pair $\{x, y\}$ is repeatedly, there is no hope of recovering the full ranking π^* . It turns out that the parameters can be recovered as long as every pair is ‘‘covered’’ by some display set.

Theorem 2 *The MLE $(\hat{\pi}, \hat{q})$ is an consistent estimator if and only if for every pair of items $\{i, j\}$, there exists a display set S that includes $\{i, j\}$ and is displayed infinite many times.*

We would like to mention that the coverage condition is fairly mild. For example, as long as the full display set is displayed sufficiently many times, the RMJ-implied central ranking can eventually be

recovered. This stands in contrast to its counterpart in the Mallows model. Since the Mallows model does not produce simple choice probabilities, the MLE from choice data is rather difficult to obtain. Perhaps the best method to date is by [4], who use a ‘‘Mallows smoothing’’ heuristic. The following result reveals that the heuristic can be unstable or fail to recover the underlying Mallows parameters even under sufficient coverage of display sets.

Theorem 3 *Even if all display sets with sizes larger than two are displayed infinite times, the estimator from the Mallows Smoothing heuristic is not consistent.*

The intuition behind the result above is that Mallows Smoothing needs a ‘‘choice to ranking’’ step. That is, it needs to first find a distribution over rankings, denoted by $\hat{\lambda} \in \Delta(\Sigma)$, that aggregates into choice probabilities to match the empirical choice probabilities $\{\hat{\Pr}(x|S)\}$. Formally, it corresponds to solving the system of linear equations

$$\sum_{\tilde{\pi}} \hat{\lambda}(\tilde{\pi}) \mathbf{I}(x, \tilde{\pi}, S) = \hat{\Pr}(x|S) \quad (6)$$

for all display set S in the data and $x \in S$. Since there are vastly more variables than equations, the solutions in general form a polytope rather than a singleton. More importantly, we find that there can be solutions $\hat{\lambda}_1$ and $\hat{\lambda}_2$ that aggregate to different Mallows models in the ‘‘ranking aggregation’’ step of the heuristic. Therefore, in general, this heuristic can produce non-unique results and thus be inconsistent.

3.2 The general ($k \geq 1$) case

In the general ($k \geq 1$) case, the ranked choice probabilities and the corresponding estimation remain parsimonious. Roughly speaking, the ranked choice model behaves like a generalized OAM where $\Pr(\pi_k|S)$ depends on the *relative rankings* of the items in π_k among the set S . The tractability comes from the sole dependence on the *relative* ranking. That is, to obtain $\Pr(\pi_k|S)$, one could first treat S as the ‘‘full display set’’ for a subuniverse of items, then re-rank all the items within the display set S , and finally just apply Lemma 1.

Choice probabilities. Given a display set S , let $d_S(\cdot)$, $L_S(\cdot)$ be the originally defined $d(\cdot)$ and $L(\cdot)$ functions, but treating the display set S as the universe. Formally, $d_S(\pi_k) := \sum_{i=1}^{k-1} \mathbb{I}\{\pi_k(i) > \pi_k(i+1)\} \cdot (|S| - i)$ and $L_S(\pi_k) := |\{x \in R^c(\pi_k) \cap S : x < \pi_k(k)\}|$. For example, suppose again that $n = 7$ and $\pi_4 = (7, 4, 6, 2)$, and let $S = \{2, 3, 4, 5, 6, 7\}$. Recall that $d(\pi_4) = 6 + 4 = 10$ and $L(\pi_4) = |\{1\}| = 1$. In comparison, $d_S(\pi_4) = 5 + 3 = 8$ and $L_S(\pi_4) = |\emptyset| = 0$.

Theorem 4 *Given a display set S and a top- k ranking π_k such that $R(\pi_k) \subseteq S$, we have*

$$\Pr(\pi_k|S) = q^{d_S(\pi_k) + L_S(\pi_k)} \cdot \frac{\psi(|S| - k, q)}{\psi(|S|, q)}.$$

Note that Theorem 4 generalizes both Theorem 1 and Lemma 1 (but in different ways). It reduces to Theorem 1 by setting $k = 1$: Given $\pi_1 = (z)$ where $z \in S$, we have $d_S(\pi_1) = 0$, $L_S(\pi_1) = |\{x \in S \setminus \{z\} : x < z\}|$ the relative ranking of z in display set S , and $\frac{\psi(|S| - k, q)}{\psi(|S|, q)} = 1 + q + \dots + q^{|S|-1}$ the appropriate normalizing constant where $k = 1$. In addition, Theorem 4 reduces to Lemma 1 by setting $S = [n]$, as can be easily verified.

Parameter learning from ranked choice data. In the parameter learning problem, we are endowed with choice data $H_T = (S_1, \pi_k^1, \dots, S_T, \pi_k^T)$, where S_t is the display set shown to the t th participant and $\pi_k^t = (x_1^t, \dots, x_k^t)$ are his/her top- k choices ranked from the most preferred to the k th preferred.

It turns out that (somewhat surprisingly) in the $k \geq 1$ case, the central ranking π^* can be estimated from the same integer programming (5) but just with a generalized re-definition of the weight parameters $\{w_{ij}\}$.

Theorem 5 *The MLE for the central ranking π^* , $\hat{\pi}$, can be obtained from the integer program (5) with $\hat{x}_{ij} = 1$ iff $i \succ_{\hat{\pi}} j$ and a generalized definition of $\{w_{ij}\}$ below*

$$w_{ij} = \sum_{t=1}^T \left[\mathbb{I}\{\{i, j\} \in S_t \setminus R(\pi_{k-1}^t) \text{ and } x_k^t = i\} + \sum_{h=1}^{k-1} (|S_t| - h) \cdot \mathbb{I}\{x_h^t = i, x_{h+1}^t = j\} \right].$$

Intuitively, a positive $w_{ij} - w_{ji}$ is still an indication that item i should be preferred to item j , but now taking into the consideration that every participant’s response is actually a *ranked list* rather than a single choice. Practically speaking, the $\{w_{ij}\}$ can be maintained in a relatively simple manner. Suppose a display set S and a ranked choice π_k are given. Then $w_{\pi_k(\ell), \pi_k(\ell+1)}$ will be added by $|S| - \ell$ for all $\ell = 1, \dots, k - 1$. (This captures the ranking information for items included in π_k .) In addition, $w_{\pi_k(k), j}$ will be added by 1 for all $j \in \{R^c(\pi_k) \cap S\}$. (This captures the ranking information between $\pi_k(k)$ and all items that are not included in π_k .) For example, suppose $n = 6$ and $k = 3$. The ranked choice data is such that $S_1 = \{1, 2, 3, 4, 5\}$ and $\pi_k^1 = (3, 1, 2)$. Then we have $w_{3,1} = 4, w_{1,2} = 3, w_{2,4} = w_{2,5} = 1$ and all the other $w_{ij} = 0$.

Discussion: model limitation. A few more words on the ranked choice model $\{Pr(\pi_k|S)\}$. The specific form gives it great simplicity, but as any other parametric model, it may be vulnerable to model misspecification issues in practice. In particular, since the model is calculated from a unimodal ranking distribution, this model (in its “vanilla” form) is best suited when there is an approximately “consensus” ranking in the population. In our numerical studies we will demonstrate how we can effectively mitigate those issues by considering learning the model parameters in a mixture setting to better capture the preference heterogeneity in the population.

4 Numerical Experiments

We investigate the performance of our methodology on two anonymous survey data sets regarding sushi preferences ([18]): The first one consists of 5,000 full preference rankings over 10 kinds of sushi, while the second one consists of 5,000 top-10 rankings over 100 kinds. Throughout the numerical experiments, we use a workstation with dual Intel Xeon Gold 6244 CPU (3.6 GHz and 32 cores in total), no GPUs, and 754 GB of memory.

Experiment 1: Top-1 choice. When $k = 1$, a top- k list corresponds to a choice model, which connects to a richer literature (e.g., [19, 4, 11]). We compare our prediction power with two representative ranking models: Mallows and Plackett-Luce. (The previous is distance-based. The latter is not but leads to the famous MNL choice model.) We use the Mallows Smoothing (MS) heuristic to estimate a Mallows model and MLE to estimate a Plackett-Luce. For all three rankings models (Mallows, PL, and ours), we also use an EM algorithm to perform parameter learning in a mixture setting.


Since the estimation and generalization performances can depend on the display sets, let us define an instance of the comparison by $\{\mathcal{S}_{train}, \mathcal{S}_{test}, \mathcal{C}\}$. Here \mathcal{S}_{train} (resp. \mathcal{S}_{test}) is collection of display sets in the training (resp. test) data, and \mathcal{C} is the number of clusters. In every instance, we split the data into 4000 participants in training and 1000 in testing. We generate empirical training (resp. testing) choice data by first enumerating all display sets in \mathcal{S}_{train} (resp. \mathcal{S}_{test}) and then record every participant’s favorite type of sushi for every display set in the traing (resp. testing) data. Finally, we use the log-likelihood as performance metric. We will use term *explanation power* (resp. *prediction power*) to refer the performance metric evaluated on the training (resp. testing) data.

We use two configurations for \mathcal{S}_{train} and \mathcal{S}_{test} , respectively. For \mathcal{S}_{train} , we take it to be either the full set or a collection of three randomly generated sets, which are realized to be $\{\{1, 3, 4, 7, 8, 10\}, \{2, 4, 5, 6, 8, 9\}, [10]\}$. For \mathcal{S}_{test} , we take it to be either the collection of all pairwise sets or the collection of all display sets with sizes at least two. Note that \mathcal{S}_{train} has low variability and many elements in \mathcal{S}_{test} do not appear in \mathcal{S}_{train} , making the experiment emphasizing more on the *generalization power*. We take $\mathcal{C} \in \{1, 2, 4, 6, 15\}$, therefore running a total of $2 \times 2 \times 5 = 20$ instances.

We summarize our results in Figure 1. We find that our method has supreme performance compared to MS and MNL across the settings. Intuitively, MS and MNL underperform in different ways. MS is vulnerable to the identifiability issue in the “choice-to-rank” step, for which we provide theoretical understanding in Theorem 3. Our numerical experiments confirm the theoretical insight: We randomize over the extreme points of the polytope of solutions to (6), each of which leads to a valid output of the MS heuristic. Every box plot in Figure 1 contains a summary of the performances of those outputs. It is clear that the performances (both in training and testing) span wide ranges. In the meantime, MNL is vulnerable to both the model risk from its specific parametric form (reflected in the limited increase of explanation power when considering more clusters) and the sample risk of

overfitting (reflected in the difference in performance in training vs. testing, especially when trained on the full display set only).

Figure 1: Comparison of Explanation and Prediction Power for Top-1 choice



In each panel, the x-axis represents the number of clusters in the mixture model, and the y-axis represents the log-likelihood metric.

Experiment 2: Top- k choice. In this experiment, we conduct estimation on top-1, top-2 and top-3 ranked choices constructed from the first (i.e., 10-sushi) data set. The collection of display sets is taken to be all display sets with sizes at least k . We find that the estimation results are relatively robust: all three instances of experiments produce the same estimated RMJ-implied central ranking $\hat{\pi} = (8, 5, 6, 3, 2, 1, 4, 9, 7, 10)$.

Experiment 3: 100 sushi types. We apply our method on the large second 100 sushi data set. In this experiment, we offer the full set (100 sushi) to 5000 individuals and record their top-10 choices. Since the problem size is larger, we use the LP-Rand-Pivot heuristic by [11] based on LP relaxation. We bootstrap 10 times (each time drawing 10000 samples) and record the running times and optimality gaps in the Table 1. We find that our method can reasonably handle the 100-sushi-type setup: We can obtain $< 2\%$ optimality gap within 5 minutes (excl. model building time).

Table 1: Computational Time and Optimality Gap on the 100-sushi data

	Model Building Time (min)	Model Solving Time (min)	Optimality Gap
Average	21.10	4.20	1.47%
Max	21.19	4.50	1.79%

5 Conclusions

We identify a novel distance-based (Mallows-type) ranking model. It aggregates into simple probability distributions for top- k subrankings among an arbitrary display set S . In addition, it facilitates effective parameter learning through the MLE formulation. This is the first distance-based ranking model with such properties (even for $k = 1$) to the best of our knowledge.

This ranking model can be used to model population preferences and provide a rationalizable way to model their ranked choices from given display sets. We demonstrate its practical value using real preference data. For example, under a mixture setting with only a few clusters, it shows promising prediction power, especially when there is a limited variety in the display sets.

For future steps, we believe our work can serve as the “infrastructure” for a range of business-related decision problems, such as new product introduction, crowdsourcing, and marketing research, among others.

References

- [1] Michael A Fligner and Joseph S Verducci. Distance based ranking models. *Journal of the Royal Statistical Society: Series B (Methodological)*, 48(3):359–369, 1986.
- [2] Colin L Mallows. Non-null ranking models. i. *Biometrika*, 44(1/2):114–130, 1957.
- [3] Flavio Chierichetti, Anirban Dasgupta, Shahrzad Haddadan, Ravi Kumar, and Silvio Lattanzi. Mallows models for top-k lists. *Advances in Neural Information Processing Systems*, 31, 2018.
- [4] Desir Antoine, Goyal Vineet, Jagabathula Srikanth, and Segev Danny. Mallows-smoothed distribution over rankings approach for modeling choice. *Operations Research*, 2021.
- [5] Tyler Lu and Craig Boutilier. Effective sampling and learning for mallows models with pairwise-preference data. *J. Mach. Learn. Res.*, 15(1):3783–3829, 2014.
- [6] Valeria Vitelli, Øystein Sørensen, Marta Crispino, Arnoldo Frigessi Di Rattalma, and Elja Arjas. Probabilistic preference learning with the mallows rank model. *Journal of Machine Learning Research*, 18(158):1–49, 2018.
- [7] Jagabathula and Rusmevichientong. The limit of rationality in choice modeling: formulation, computation, and implications. *Management Science*, 2018.
- [8] Wenpin Tang. Mallows ranking models: maximum likelihood estimate and regeneration. In *International Conference on Machine Learning*, pages 6125–6134. PMLR, 2019.
- [9] Qinghua Liu, Marta Crispino, Ida Scheel, Valeria Vitelli, and Arnoldo Frigessi. Model-based learning from preference data. *Annual review of statistics and its application*, 6:329–354, 2019.
- [10] Fabien Collas and Ekhine Irurozki. Concentric mixtures of mallows models for top- k rankings: sampling and identifiability. In *International Conference on Machine Learning*, pages 2079–2088. PMLR, 2021.
- [11] Yifan Feng, Rene Caldentey, and Christopher Thomas Ryan. Robust learning of consumer preferences. *Operations Research*, 2021.
- [12] Zhe Cao, Tao Qin, Tie-Yan Liu, Ming-Feng Tsai, and Hang Li. Learning to rank: from pairwise approach to listwise approach. In *Proceedings of the 24th international conference on Machine learning*, pages 129–136, 2007.
- [13] Tie-Yan Liu et al. Learning to rank for information retrieval. *Foundations and Trends® in Information Retrieval*, 3(3):225–331, 2009.
- [14] John I Marden. *Analyzing and modeling rank data*. CRC Press, 1996.
- [15] Thotsaporn Thanatipanonda. Inversions and major index for permutations. *Mathematics Magazine*, 77(2):136–140, 2004.
- [16] Mark Braverman and Elchanan Mossel. Noisy sorting without resampling. In *19th Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 268–276, 2008.
- [17] Mark Braverman and Elchanan Mossel. Sorting from noisy information. *arXiv preprint arXiv:0910.1191*, 2009.
- [18] T Kamishima, H Kazawa, and S Akaho. Supervised ordering—an empirical survey. *Proc. 5th IEEE Internat. Conf. Data Mining*, 2005.
- [19] R Duncan Luce. Individual choice behavior, 1959.

Supplemental Material for:
On A Mallows-type Model For (Ranked) Choices

A Characterizing the Choice Probability (Theorem 1)

A.1 Proofs of Intermediate Results (Lemmas 1, 2, and 3)

Proof of Lemma 1. Let us first assign a weight to each ranking $\pi \in \Sigma$ and subranking $\pi_k \in \Sigma_k$ ($k \in \{0, \dots, n\}$) by

$$w(\pi) := \frac{q^{d(\pi)}}{\psi(n, q)} \quad \text{and} \quad w(\pi_k) := \sum_{\pi \in \Sigma: \pi_k \subseteq \pi} w(\pi). \quad (7)$$

We note that on the one hand, $w \propto \lambda$. On the other hand, it is unclear *in priori* whether $w(\cdot)$ is the probability mass function since one needs to verify that $w(\cdot)$ is equipped with the right normalizing constant. We will first prove that $w(\pi_k)$ can also be written as $w(\pi_k) = q^{d(\pi_k)+L(\pi_k)} \cdot \frac{\psi(n-k, q)}{\psi(n, q)}$. Then we will show that for every $k \in [n]$, $\sum_{\pi_k \in \Sigma_k} w(\pi_k) = 1$. In other words, $w(\cdot) = \lambda(\cdot)$.

Step 1. Recall that for every $\pi_k \in \Sigma_k$, $d(\pi_k) = \sum_{i=1}^{k-1} \mathbb{I}\{\pi_k(i) > \pi_k(i+1)\}(n-i)$ and $L(\pi_k) := |\{x \in R^c(\pi_k) : x < \pi_k(k)\}|$. We prove from (7) that

$$w(\pi_k) = q^{d(\pi_k)+L(\pi_k)} \cdot \frac{\psi(n-k, q)}{\psi(n, q)}$$

by backward induction on k .

Base step. Suppose $k = n$. Pick an arbitrary $\pi_k \in \Sigma_k = \Sigma$. Because π_k is a full ranking,

$$w(\pi_k) = q^{d(\pi_k)} \cdot \frac{1}{\psi(n, q)} \stackrel{(a)}{=} q^{d(\pi_k)+L(\pi_k)} \cdot \frac{\psi(n-k, q)}{\psi(n, q)}.$$

In the derivations above, part (a) is due to the facts that $\psi(n-k, q) = \psi(0, q) = 1$ and $L(\pi_k) = 0$ by definition.

Inductive step. Pick an arbitrary $K \in \{0, 1, \dots, n-1\}$. Suppose our statement holds for every $k = K+1, \dots, n$. We want to show that our statement holds for $k = K$. In other words, pick an arbitrary $\pi_k \in \Sigma_k$. We wish to show that $\sum_k w(\pi_k) = q^{d(\pi_k)+L(\pi_k)} \cdot \frac{\psi(n-k, q)}{\psi(n, q)}$.

We first claim that for every $\pi_{k+1} \in \Sigma_{k+1}$ such that $\pi_k \subseteq \pi_{k+1}$, $\pi_k(k) > \pi_{k+1}(k+1)$ if and only if $L(\pi_{k+1}) \leq L(\pi_k) - 1$. To see why, note that if $\pi_k(k) > \pi_{k+1}(k+1)$, $L(\pi_k) = L(\pi_{k+1}) + 1$. Otherwise, $L(\pi_k) \leq L(\pi_{k+1})$, and hence $L(\pi_k) - 1 < L(\pi_{k+1})$. As a consequence, $d(\pi_{k+1})$ can be expressed in terms of $d(\pi_k)$:

$$\begin{aligned} d(\pi_{k+1}) &= \sum_{i=1}^k \mathbb{I}\{\pi_k(i) > \pi_k(i+1)\} \cdot (n-i) \\ &= \underbrace{\sum_{i=1}^{k-1} \mathbb{I}\{\pi_k(i) > \pi_k(i+1)\} \cdot (n-i)}_{d(\pi_k)} + \underbrace{\mathbb{I}\{\pi_k(k) > \pi_{k+1}(k+1)\} \cdot (n-k)}_{\mathbb{I}\{L(\pi_{k+1}) \leq L(\pi_k) - 1\}} \\ &= d(\pi_k) + \mathbb{I}\{L(\pi_{k+1}) \leq L(\pi_k) - 1\} \cdot (n-k). \end{aligned}$$

Moreover, given π_k , there is a one-to-one correspondence between π_{k+1} and $k+1$ and we use $\pi_k \oplus L^{-1}(i)(k+1)$ to represent the (unique) π_{k+1} such that $\pi_k \subseteq \pi_{k+1}$ and $L(\pi_{k+1}) = i$, ($i \in \{0, 1, \dots, n-k-1\}$).

$$\begin{aligned}
w(\pi_k) &= \sum_{\pi \in \Sigma: \pi_k \subseteq \pi} w(\pi) = \sum_{\pi_{k+1} \in \Sigma_{k+1}: \pi_k \subseteq \pi_{k+1}} w(\pi_{k+1}) \\
&= \sum_{i=0}^{n-k-1} w(\pi_k \oplus L^{-1}(i)(k+1)) \\
&= \sum_{i=0}^{n-k-1} q^{d(\pi_k \oplus L^{-1}(i)(k+1)) + L(\pi_k \oplus L^{-1}(i)(k+1))} \cdot \frac{\psi(n-k-1, q)}{\psi(n, q)} \\
&= \frac{\psi(n-k-1, q)}{\psi(n, q)} \cdot \left(\sum_{i=0}^{L(\pi_k)-1} q^{d(\pi_k \oplus L^{-1}(i)(k+1)) + i} + \sum_{i=L(\pi_k)}^{n-k-1} q^{d(\pi_k \oplus L^{-1}(i)(k+1)) + i} \right) \\
&= \frac{\psi(n-k-1, q)}{\psi(n, q)} \cdot \left(\sum_{i=0}^{L(\pi_k)-1} q^{d(\pi_k) + n - k + i} + \sum_{i=L(\pi_k)}^{n-k-1} q^{d(\pi_k) + i} \right) \\
&= \frac{\psi(n-k-1, q)}{\psi(n, q)} \cdot q^{d(\pi_k)} \cdot q^{L(\pi_k)} \underbrace{\left(\sum_{i=n-k-L(\pi_k)}^{n-k-1} q^i + \sum_{i=0}^{n-k-1-L(\pi_k)} q^i \right)}_{\frac{\psi(n-k, q)}{\psi(n-k-1, q)}} \\
&= q^{d(\pi_k) + L(\pi_k)} \cdot \frac{\psi(n-k, q)}{\psi(n, q)}.
\end{aligned}$$

Hence our statement holds for $k = K$, too, thus $w(\pi_k) = q^{d(\pi_k) + L(\pi_k)} \cdot \frac{\psi(n-k, q)}{\psi(n, q)}$ for any k .

Step 2. We verify that for every $k \in \{0, 1, \dots, n\}$, $\sum_{\pi_k \in \Sigma_k} w(\pi_k) = 1$.

For all $k, k' \in \{0, 1, \dots, n\}$, we invoke (7) and have $\sum_{\pi_k \in \Sigma_k} w(\pi_k) = \sum_{\pi \in \Sigma} w(\pi) = \sum_{\pi_{k'} \in \Sigma_{k'}} w(\pi_{k'})$. Therefore, take an arbitrary $k \in \{0, 1, \dots, n\}$ and $k' = 2$,

$$\begin{aligned}
\sum_{\pi_k \in \Sigma_k} w(\pi_k) &= \sum_{\pi_2 \in \Sigma_2} w(\pi_2) = \sum_{i \in \{0, \dots, n-1\}} w(\pi_1 \oplus i) \\
&= \frac{\psi(n-1, q)}{\psi(n, q)} \cdot (1 + q + \dots + q^{n-1}) \\
&= 1
\end{aligned}$$

Hence, the weight w is the probability mass function λ . \square

Proof of Lemma 2. By Lemma 1, we have

$$\begin{aligned}
\frac{\lambda(\pi_k \oplus y)}{\lambda(\pi_k)} &= \frac{q^{d(\pi_{k+1}) + L(\pi_{k+1})} \cdot \frac{\psi(n-k-1, q)}{\psi(n, q)}}{q^{d(\pi_k) + L(\pi_k)} \cdot \frac{\psi(n-k, q)}{\psi(n, q)}} = \frac{q^{d(\pi_k) + \mathbb{I}\{z > y\} \cdot (n-k) + L(\pi_{k+1})}}{q^{d(\pi_k) + L(\pi_k)}} \frac{1 - q}{1 - q^{n-k}} \\
&= q^{\mathbb{I}\{z > y\} \cdot (n-k) + L(\pi_{k+1}) - L(\pi_k)} \frac{1 - q}{1 - q^{n-k}}.
\end{aligned}$$

By the definition of $L(\cdot)$,

$$\mathbb{I}\{z > y\} \cdot (n-k) + L(\pi_{k+1}) - L(\pi_k) = \begin{cases} \sum_{x \in R^c(\pi_k)} \mathbb{I}\{z < x \leq y\} - 1 & \text{if } y > z \\ n - k - \sum_{x \in R^c(\pi_k)} \mathbb{I}\{y < x < z\} - 1 & \text{if } y < z \end{cases}.$$

The proof is finished. \square

Proof of Lemma 3. We prove this lemma by backward induction on k .

Base step. Since when $k > n - M + 1$, $R(\pi_{k-1}) \cap S = \emptyset$ is not satisfied for all $\pi_{k-1} \in \Sigma_{k-1}$. We suppose $k = n - M + 1$. Pick an arbitrary $\pi_k = \pi_{k-1} \oplus z$ such that $R(\pi_{k-1}) \cap S = \emptyset$. Item z is an item in S , so the choice outcome is deterministic. If $z = x_i$, the conditional choice probability is 1, and if $z \in S \setminus \{x_i\}$, the conditional choice probability is 0.

Inductive step. Pick an arbitrary $K \in \{1, \dots, n - M\}$. Suppose our statement holds for every $k = K + 1, \dots, n - M + 1$. We want to show that our statement holds for $k = K$.

Pick an arbitrary $\pi_k = \pi_{k-1} \oplus z$ such that $R(\pi_{k-1}) \cap S = \emptyset$. Similar to the base step, if $z = x_i$, the conditional choice probability is 1, and if $z \in S \setminus \{x_i\}$, the conditional choice probability is 0. When $z \notin S$, rename the items in $R^c(\pi_k)$ as y_1, y_2, \dots, y_{n-k} so that $y_1 < y_2 < \dots < y_{n-k}$. We have the following decomposition.

$$Q := Pr(\{x_i|S\}|\pi_{k-1} \oplus z) = \sum_{j=1}^{n-k} Pr(\{x_i|S\}|\pi_k \oplus y_j) \cdot Pr(\pi_k \oplus y_j|\pi_{k-1} \oplus z)$$

Since $R(\pi_k) \cap S = \emptyset$, we have $\{x_1, x_2, \dots, x_M\} \subseteq \{y_1, y_2, \dots, y_{n-k}\}$, we use \bar{x}_i to denote the relative position of item x_i in these $n - k$ items, hence item x_i is renamed as $y_{\bar{x}_i}$ under ‘‘y’’ notation. For example, if $\bar{x}_i = 4$, then there are 3 items with smaller indices (i.e., preferred under $\pi^* = e$) than item x_i in $\{y_i\}_{i=1}^{n-k}$. Hence, we can rewrite $\{y_i\}_{i=1}^{n-k}$ as

$$\{y_1, \dots, y_{\bar{x}_1-1}, y_{\bar{x}_1}, y_{\bar{x}_1+1}, \dots, y_{\bar{x}_i-1}, y_{\bar{x}_i}, y_{\bar{x}_i+1}, \dots, y_{n-k}\}.$$

Because different $\pi_{k+1}(k+1)$ could result in the same $p(\pi_{k+1}(k+1)|S)$, we classify $\pi_{k+1}(k+1)$ by its corresponding $p(\pi_{k+1}(k+1)|S)$ and by induction hypothesis, we have

$$Pr(\{x_i|S\}|\pi_k \oplus \pi_{k+1}(k+1)) = \begin{cases} \frac{q^{i-1}}{1+q+\dots+q^{M-1}} & \text{if } \pi_{k+1}(k+1) \in \{y_a\}_{a=\bar{x}_1-1}^{\bar{x}_1-1}, \\ \frac{q^{i-j-1}}{1+q+\dots+q^{M-1}} & \text{if } \pi_{k+1}(k+1) \in \{y_a\}_{a=\bar{x}_j+1}^{\bar{x}_j+1} \text{ for } j \in \{1, \dots, i-1\}, \\ 1 & \text{if } \pi_{k+1}(k+1) = y_{\bar{x}_i}, \\ \frac{q^{M+i-j-1}}{1+q+\dots+q^{M-1}} & \text{if } \pi_{k+1}(k+1) \in \{y_a\}_{a=\bar{x}_j+1}^{\bar{x}_j+1} \text{ for } j \in \{i, \dots, M-1\}, \\ \frac{q^{i-1}}{1+q+\dots+q^{M-1}} & \text{if } \pi_{k+1}(k+1) \in \{y_a\}_{a=\bar{x}_M+1}^{n-k}, \\ 0 & \text{if } \pi_{k+1}(k+1) = y_{\bar{x}_j} \text{ for } x_j \in S \setminus \{x_i\}. \end{cases}$$

The first two cases above correspond to the second case in the induction hypothesis, the 4th, 5th cases above correspond to the first case in the induction hypothesis, the 3rd, 6th cases above correspond to the 3rd, 4th cases in the induction hypothesis, respectively.

By Lemma 2, we know $Pr(\pi_k \oplus y_j|\pi_{k-1} \oplus z)$. Finally, we can get

$$Q = \begin{cases} \frac{q^{M-p(z|S)+i-1}}{1+q+\dots+q^{M-1}} & \text{if } z > x_i \text{ and } z \notin S, \\ \frac{q^{i-p(z|S)-1}}{1+q+\dots+q^{M-1}} & \text{if } z < x_i \text{ and } z \notin S. \end{cases}$$

Hence our statement holds for $k = K$, too, thus finishing the proof. \square

A.2 Putting Things Together for Theorem 1

Proof of Theorem 1. Name all items as $\{y_z\}_{z=1}^n$ such that $y_1 < y_2 < \dots < y_n$. We have

$$Pr(\{x_i|S\}) = \sum_{z=1}^n Pr(\{x_i|S\}|(y_z)) \cdot \lambda((y_z))$$

Use \bar{x}_j to denote the position of item x_j in the universe, hence item x_j is renamed as $y_{\bar{x}_j}$ under ‘‘y’’ notation. By Lemma 3, we have the following equation:

$$Pr(\{x_i|S\}|(y_z)) = \begin{cases} \frac{q^{i-1}}{1+q+\dots+q^{M-1}} & \text{if } y_z \in \{y_a\}_{a=1}^{\bar{x}_1-1}, \\ \frac{q^{i-j-1}}{1+q+\dots+q^{M-1}} & \text{if } y_z \in \{y_a\}_{a=\bar{x}_j+1}^{\bar{x}_{j+1}-1} \text{ for } j \in \{1, \dots, i-1\}, \\ 1 & \text{if } y_z = y_{\bar{x}_i}, \\ \frac{q^{M+i-j-1}}{1+q+\dots+q^{M-1}} & \text{if } y_z \in \{y_a\}_{a=\bar{x}_j+1}^{\bar{x}_{j+1}-1} \text{ for } j \in \{i, \dots, M-1\}, \\ \frac{q^{i-1}}{1+q+\dots+q^{M-1}} & \text{if } y_z \in \{y_a\}_{a=\bar{x}_M+1}^n, \\ 0 & \text{if } y_z = y_{\bar{x}_j} \text{ for } x_j \in S \setminus \{x_i\}. \end{cases}$$

By Lemma 1, we have $\lambda((y_z)) = \frac{q^{z-1}}{1+\dots+q^{n-1}}$. After calculation, we get $Pr(\{x_i|S\}) = \frac{q^{i-1}}{1+q+\dots+q^{M-1}}$. \square

B Consistency of MLE for OAM (Theorem 2)

B.1 Main Body of the Proof

Proof of Theorem 2. First fix the underlying parameters of the RMJ-based ranking model (π^*, q^*) . Let the choice data $H_T = (S_1, x_1, \dots, S_T, x_T)$ be given, where S_t is the display set shown to the t th participant and x_t is his/her choice. Invoking Proposition 3 of [11], the MLE problem for OAM, the choice model induced by the RMJ-based ranking distribution, can be written as

$$(\hat{\pi}, \hat{q}) \in \operatorname{argmax}_{\pi, q} \sum_{t=1}^T \log \left(\frac{1-q}{1-q^{|S_t|}} \right) + \log q \sum_{(i,j):i \neq j} \mathbb{I}\{j \succ_{\pi} i\} w_{ij},$$

where

$$w_{ij} := \sum_{t=1}^T \mathbb{I}\{\{i, j\} \subseteq S_t \text{ and } x_t = i\}$$

is the number of times that both items i and j are displayed and item i is chosen (among the T samples). Furthermore, recall that $\hat{\pi}$ can be obtained from the integer programming formulation (5), which is further equivalent to the following formulation by substituting the relation $x_{ij} + x_{ji} = 1$:

$$\begin{aligned} \{\hat{x}_{ij} : i < j\} \in \operatorname{argmin}_{\{x_{ij}:i < j\}} & \sum_{(i,j):i < j} (w_{ji} - w_{ij}) x_{ij} \\ \text{s.t.} & x_{ij} + x_{jr} - x_{ir} \leq 1 \quad \forall 1 \leq i < j < r \leq n \\ & x_{ij} \in \{0, 1\} \quad \forall 1 \leq i < j \leq n \end{aligned} \quad (8)$$

The ranking $\hat{\pi}$ is obtained by letting $i \succ_{\pi} j$ if and only if $\hat{x}_{ij} = 1$. Given the solution of $\hat{\pi}$, the estimator \hat{q} is obtained by the one-dimensional convex optimization

$$\hat{\alpha} \in \operatorname{argmin}_{\alpha \in (0, +\infty)} L_T(\alpha) := \alpha \sum_{t=1}^T \sum_{x \in S_t} \mathbb{I}\{x \succ_{\hat{\pi}} x_t\} + \sum_{t=1}^T \log \sum_{j=0}^{|S_t|-1} e^{-j\alpha} \quad (9)$$

so that $\hat{\alpha} = -\log \hat{q}$.

We break the rest of the proof into two parts: the ‘‘if’’ part and the ‘‘only if’’ part.

The ‘‘if’’ part. Let \mathcal{S}_{∞} be the collection of display sets that are displayed infinite times. Suppose for every pair of items $\{i, j\}$ is covered infinitely many times. That is, there exists a display set $S \in \mathcal{S}_{\infty}$ such that $\{i, j\} \subseteq S$. We wish to show that $(\hat{\pi}, \hat{q}) \rightarrow (\pi^*, q^*)$ almost surely as the sample size $T \rightarrow \infty$.

We first claim that $\hat{\pi} \rightarrow \pi^*$ almost surely. Without loss of generality, assume $\pi^* = e$, the identity ranking. In other words, we wish to show that with probability one, there exists τ such that for all $T \geq \tau$, the unique solution to (8) is $\{x_{ij} = 1, i < j\}$. Pick an arbitrary pair of items i, j such that $i < j$. Let $N_{ij} := \sum_{t=1}^T \mathbb{I}\{\{i, j\} \subseteq S_t\}$ be the number of times that both items i and j are displayed.

Because both i and j are covered by some $S \in \mathcal{S}_\infty$, we have $N_{ij} \rightarrow \infty$ as $T \rightarrow \infty$. Note that invoking the choice probabilities in (4), OAM is a q^* -separable choice model, i.e., for every display set S such that $\{i, j\} \subseteq S$, $\Pr(\{j|S\}) \leq q \Pr(\{i|S\})$; see [11] for more details. Since the choices are generated independently conditional the display sets, we invoke the law of large numbers and conclude that $w_{ij} \rightarrow \infty$, $w_{ji} \rightarrow \infty$, and $w_{ji}/w_{ij} \rightarrow q'$ for some $q' \leq q$ as the sample size $T \rightarrow \infty$. Furthermore, since there are only a finite number of pairs, with probability one, there exists τ such that for all $T \geq \tau$,

$$w_{ji} - w_{ij} < 0 \quad \text{for all } i < j.$$

In that case, it is straightforward to see that the unique solution to (8) is $\{x_{ij} = 1, i < j\}$. The ‘‘if’’ part is hence completed by noting the result below. Its proof resembles the standard argument for consistency of MLE except for a few technical differences, such as allowing for an arbitrary display set offering process (which leads to no necessarily i.i.d choice data) and non-compactness of the range of α . We provide the proof details in Appendix B.2.

Lemma 4 $\hat{q} \rightarrow q^*$ almost surely.

The ‘‘only if’’ part. Suppose there exists a pair of items $\{i, j\}$ that is not covered infinitely many times. We wish to show that for some underlying parameter (π^*, q^*) , $(\hat{\pi}, \hat{q}) \not\rightarrow (\pi^*, q^*)$ with positive probability as the sample size $T \rightarrow \infty$.

Through a relabeling argument, assume $i = n - 1$ and $j = n$ without loss of generality. Also, for simplicity, assume that $\{n - 1, n\}$ is not covered *at all* without loss of generality. As a consequence, $w_{n-1,n} = w_{n,n-1} = 0$. Therefore, we can verify that the rankings e and $\tilde{\pi} := (1, 2, \dots, n - 2, n, n - 1)$ (i.e., the ranking obtained by swapping the rankings of the bottom two ranked items of e) correspond to the same objective value of (8) for all $T \geq 1$. Therefore, no matter what tie-breaknig rule for (8), one cannot ensure that both e and $\tilde{\pi}$ are almost surely recovered (as ground truth rankings). \square

B.2 Proof of Auxiliary Results (Lemma 4)

Proof of Lemma 4. Let $\alpha^* := -\log q^* \in (0, +\infty)$ and recall that $\hat{\alpha}$ is solution to (9). It suffices to show that $\hat{\alpha} \rightarrow \alpha^*$ almost surely. We employ a pathwise analysis throughout the proof.

Given display set S and item $x \in S$, let $N_S := \sum_{t=1}^T \mathbb{I}\{S_t = S\}$ and $N_S^x := \sum_{t=1}^T \mathbb{I}\{S_t = S \text{ and } x_t = x\}$ be the number of times that S is displayed and x is chosen out of S , respectively. For $i \in \{1, \dots, |S|\}$, let $\pi_S(i)$ be the i^{th} most preferred item within display set under ranking π^* . In other words, if $\pi_S(i) = y$, then $\sum_{x \in S} \mathbb{I}\{x \succ_{\pi^*} y\} = i - 1$. Finally, let us introduce

$$\begin{aligned} L_T^S(\alpha) &:= \frac{1}{N_S} \left(\alpha \sum_{t:S_t=S} \sum_{x \in S} \mathbb{I}\{x \succ_{\pi^*} x_t\} + \sum_{t:S_t=S} \log \sum_{j=0}^{|S|-1} e^{-j\alpha} \right) \\ &= \alpha \sum_{i=1}^{|S|} \frac{N_S^{\pi_S(i)}}{N_S} (i - 1) + \log \sum_{j=0}^{|S|-1} e^{-j\alpha} \end{aligned}$$

to be the (scaled) partial log likelihood loss function when $\hat{\pi} = \pi^*$ and only display set S is considered. Since the choices are generated independently conditional the display sets, we invoke the choice probabilities in (4) as well as the law of large numbers and conclude that with probability one, for all $S \in \mathcal{S}_\infty$ and $\alpha \in (0, +\infty)$,

$$\begin{aligned} L_T^S(\alpha) &\rightarrow L_\infty^S(\alpha) := \alpha \sum_{i=1}^{|S|} \frac{e^{-(i-1)\alpha^*}}{\sum_{j=0}^{|S|-1} e^{-j\alpha^*}} (i - 1) + \log \sum_{j=0}^{|S|-1} e^{-j\alpha} \\ &= \alpha \frac{\sum_{i=1}^{|S|-1} i e^{-i\alpha^*}}{\sum_{j=0}^{|S|-1} e^{-j\alpha^*}} + \log \sum_{j=0}^{|S|-1} e^{-j\alpha}. \end{aligned}$$

In fact, the convergence is also locally uniform: for all $M > 0$,

$$\begin{aligned} \sup_{\alpha \in (0, M]} |L_T^S(\alpha) - L_\infty^S(\alpha)| &= \sup_{\alpha \in (0, M]} \alpha \sum_{i=1}^{|S|} \left| \frac{N_S^{\pi_S(i)}}{N_S} - \frac{e^{-(i-1)\alpha^*}}{\sum_{j=0}^{|S|-1} e^{-j\alpha^*}} \right| (i-1) \\ &\leq M \sum_{i=1}^{|S|} \left| \frac{N_S^{\pi_S(i)}}{N_S} - \frac{e^{-(i-1)\alpha^*}}{\sum_{j=0}^{|S|-1} e^{-j\alpha^*}} \right| (i-1) \rightarrow 0. \end{aligned}$$

as sample size $T \rightarrow \infty$. Therefore, with probability one, $L_T^S(\cdot) \rightarrow L_\infty^S(\cdot)$ uniformly on the set $(0, M]$.

In addition, it is straightforward to verify that for every S , $L_\infty^S(\cdot)$ is strictly convex and attains its unique minimum at α^* . In fact, one can verify that the first order condition is

$$(L_\infty^S)'(\alpha) = 0 \Rightarrow \frac{\sum_{i=1}^{|S|-1} i e^{-i\alpha^*}}{\sum_{j=0}^{|S|-1} e^{-j\alpha^*}} = \frac{\sum_{i=1}^{|S|-1} i e^{-i\alpha}}{\sum_{j=0}^{|S|-1} e^{-j\alpha}},$$

which implies that $\operatorname{argmin}_\alpha L_\infty^S(\alpha) = \alpha^*$. Since $L_T^S(\cdot) \rightarrow L_\infty^S(\cdot)$ uniformly on the set $(0, M]$ for every $M > 0$, we further conclude that with probability one and for an arbitrarily small $\epsilon > 0$, there exists τ_2 such that for all $T \geq \tau_2$,

$$L_T(\alpha) = \sum_{S \in \mathcal{S}_\infty} N_S L_T^S(\alpha) + O(1)$$

attains its minimum in $[\alpha^* - \epsilon, \alpha^* + \epsilon]$. In other words, $\hat{\alpha} \in [\alpha^* - \epsilon, \alpha^* + \epsilon]$. \square

C Inconsistency of the Mallows Smoothing Heuristic (Theorem 3)

C.1 Overview

We restate the Mallows Smoothing heuristic ([4]) in a two-step process.

Step 1: (“Choice to Ranking”) This step takes the empirical choice probabilities $\{\hat{\Pr}(x|S)\}$ as input and produces a distribution over rankings $\hat{\lambda}$ as output. The goal of this step is to find $\hat{\lambda}$, under which the aggregated choice probabilities match the empirical choice probabilities. Formally, let $\hat{\mathcal{S}}$ be the collection of display sets that have appeared in the data. This step corresponds to finding a solution $\hat{\lambda}$ to the following feasibility problem:⁴

$$\begin{aligned} \min_{\hat{\lambda}} \quad & 0 \\ \text{s.t.} \quad & \sum_{\tilde{\pi} \in \Sigma} \hat{\lambda}(\tilde{\pi}) \mathbf{I}(x, \tilde{\pi}, S) = \hat{\Pr}(x|S), \quad \text{for all } S \in \hat{\mathcal{S}} \text{ and } x \in S \\ & \sum_{\tilde{\pi} \in \Sigma} \hat{\lambda}(\tilde{\pi}) = 1 \\ & \hat{\lambda}(\pi) \geq 0, \quad \text{for all } \pi \in \Sigma. \end{aligned} \tag{10}$$

Step 2: (“Smoothing”) This step takes a distribution over rankings $\hat{\lambda}$ as input and produces the Mallows distribution parameters $(\hat{\pi}, \hat{q})$ as output. The goal of this step is to find a Mallows distribution that fits the $\hat{\lambda}$ distribution produced by the previous step.

Formally, recall that the Kendall’s Tau distance between two rankings $\pi, \tilde{\pi}$ is given by $d_K(\pi, \tilde{\pi}) = \sum_{x < y} \mathbb{I}\{(\pi^{-1}(x) - \pi^{-1}(y)) \cdot (\tilde{\pi}^{-1}(x) - \tilde{\pi}^{-1}(y)) < 0\}$. The estimated central ranking $\hat{\pi}$ of the Mallows distribution is obtained from the following ranking aggregation problem:

$$\hat{\pi} \in \operatorname{argmin}_\pi \sum_{\tilde{\pi}} \hat{\lambda}(\tilde{\pi}) \cdot d_K(\pi, \tilde{\pi}). \tag{11}$$

⁴If the system above is not feasible, then solve a “soft” version of it by minimizing a loss function of residues; see [7] for more details.

The dispersion parameter is obtained from solving the following convex problem:

$$\hat{q} = \operatorname{argmin}_q q \cdot \sum_{\tilde{\pi}} \hat{\lambda}(\tilde{\pi}) \cdot d_K(\hat{\pi}, \tilde{\pi}) + T \log \psi(n, q).$$

Note that in system (10), the number of variables is on the order of $\Omega(n!)$. In comparison, the number of constraints is on the order of $O(n2^n)$, which is much smaller. Therefore, whenever feasible, the solution typically forms a high dimensional polytope.

Roughly speaking, we will show that even all display sets with sizes at least three are displayed infinite many times, the Mallows model parameters cannot be identified from the MS heuristic. That is, we suppose that infinite choice data is generated from a Mallows model with the central ranking to be e . However, we can construct a ranking distribution $\tilde{\lambda}$ that solves (10) so that when we use it as input to problem (11), we find a solution $\tilde{\pi}$ that is different from e . That is, a *wrong* ranking parameter can be the output of the MS heuristic.

C.2 Main Body of the Proof

Proof of Theorem 3. Pick $q \in (0, 1)$. Let λ^e be the p.m.f. of the Mallows ranking model with the central ranking to be the identity ranking e and the dispersion parameter to be q . Let $\{\Pr^e(x|S) : |S| \geq 3, x \in S\}$ be the collection of associated choice probabilities for display sets with sizes at least three, formally defined as $\Pr^e(x|S) := \sum_{\tilde{\pi} \in \Sigma} \lambda^e(\tilde{\pi}) \mathbf{I}(x, \tilde{\pi}, S)$ for all S such that $|S| \geq 3$ and $x \in S$.

We claim that we can construct $\tilde{\lambda} \in \Delta(\Sigma)$ that satisfies two properties simultaneously.

1. First, it solves (10) when taking $\{\Pr^e(x|S) : |S| \geq 3, x \in S\}$ as input. That is, $\tilde{\lambda} \in \Delta(\Sigma)$ satisfies

$$\sum_{\tilde{\pi} \in \Sigma} \tilde{\lambda}(\tilde{\pi}) \mathbf{I}(x, \tilde{\pi}, S) = \Pr^e(x|S), \text{ for all } S \text{ such that } |S| \geq 3 \text{ and } x \in S. \quad (12)$$

2. Second, if we take $\tilde{\lambda}$ as input to (11), the (wrong) ranking $\tilde{\pi} := (1, 2, \dots, n-2, n, n-1) \neq e$ can be obtained as solution. That is,

$$(1, 2, \dots, n-2, n, n-1) \in \operatorname{argmin}_{\tilde{\pi}} \sum_{\tilde{\pi}} \tilde{\lambda}(\tilde{\pi}) \cdot d_K(\pi, \tilde{\pi}). \quad (13)$$

Our construction is based on two observations. First, we invoke Theorem 3.7 by [7] and know that under the current display set setup, (10) can only identify a ranking up to its first $n-2$ positions. As a consequence, as long as $\tilde{\lambda}$ satisfies

$$\sum_{\tilde{\pi} \in \Sigma} \tilde{\lambda}(\tilde{\pi}) \mathbb{I}\{\pi_{n-2} \subseteq \tilde{\pi}\} = \sum_{\tilde{\pi} \in \Sigma} \lambda^e(\tilde{\pi}) \mathbb{I}\{\pi_{n-2} \subseteq \tilde{\pi}\}, \quad \text{for all } \pi_{n-2} \in \Sigma_{n-2}, \quad (14)$$

it also satisfies (12).

Second, we observe that problem (11) only depends on the *pairwise* choice probabilities associated with its input ranking distribution. This observation is formalized as the result below, and we present its proof in Appendix C.3.

Lemma 5 Pick $\hat{\lambda} \in \Delta(\Sigma)$ and let $P_{x, \{x, y\}} := \sum_{\tilde{\pi}} \hat{\lambda}(\tilde{\pi}) \mathbb{I}\{\tilde{\pi}^{-1}(x) < \tilde{\pi}^{-1}(y)\}$ be its associated pairwise probability of choosing item x out of $\{x, y\}$. Then

$$\operatorname{argmin}_{\tilde{\pi}} \sum_{\tilde{\pi}} \hat{\lambda}(\tilde{\pi}) \cdot d_K(\pi, \tilde{\pi}) = \operatorname{argmin}_{\tilde{\pi}} \sum_{\pi} \mathbb{I}\{\pi^{-1}(x) < \pi^{-1}(y)\} \cdot (1 - 2P_{x, \{x, y\}}).$$

Let $\tilde{P}_{x, \{x, y\}} := \sum_{\tilde{\pi}} \tilde{\lambda}(\tilde{\pi}) \mathbb{I}\{\tilde{\pi}^{-1}(x) < \tilde{\pi}^{-1}(y)\}$ As a quick consequence of the result above, a sufficient condition of (13) is

$$\tilde{P}_{x, \{x, y\}} > 1/2 \text{ for all } (x, y) \in \{(x, y) : x < y, (x, y) \neq (n-1, n)\} \text{ and } \tilde{P}_{n-1, \{n-1, n\}} < 1/2. \quad (15)$$

In order to construct $\tilde{\lambda}$ that satisfies both (14) and (15) (and therefore, fulfills our claim), let us classify Σ into 3 groups based on its top- $(n-2)$ elements. Specifically, given $\pi \in \Sigma$ and $\pi_{n-2} \in \Sigma_{n-2}$ such that $\pi_{n-2} \subseteq \pi$, we categorize π into three of the groups below:

Group 1: Under π_{n-2} , item $(n-1)$ is preferred to item n . That is, either

- (i) $\{n-1, n\} \subseteq R(\pi_{n-2})$ and $\pi_{n-2}^{-1}(n-1) < \pi_{n-2}^{-1}(n)$; or
- (ii) $(n-1) \in R(\pi_{n-2})$ but $n \in R^c(\pi_{n-2})$.

Group 2: Under π_{n-2} , item n is preferred to item $(n-1)$. That is, either

- (i) $\{n-1, n\} \subseteq R(\pi_{n-2})$ and $\pi_{n-2}^{-1}(n-1) > \pi_{n-2}^{-1}(n)$, or
- (ii) $n \in R(\pi_{n-2})$ but $(n-1) \in R^c(\pi_{n-2})$.

Group 3: Under π_{n-2} , items n and $(n-1)$ are incomparable. That is $\{n-1, n\} = R^c(\pi_{n-2})$.

We explicitly construct $\tilde{\lambda}$ as the following:

$$\tilde{\lambda}(\pi) = \begin{cases} \lambda^e(\pi) & \text{if } \pi \text{ belongs to Group 1 or 2} \\ 0 & \text{if } \pi \in \text{Group 3 and } (n-1) \succ_{\pi} n \\ \lambda^e(\pi_{n-2} \oplus (n-1, n)) + \lambda^e(\pi_{n-2} \oplus (n, n-1)) & \text{if } \pi \in \text{Group 3 and } n \succ_{\pi} (n-1) \end{cases}$$

In other words, $\tilde{\lambda}$ is obtained from λ^e by "transporting" weights in favor of item n over item $(n-1)$ when both of those items are ranked at the bottom two. Since the top- $(n-2)$ rankings are not disturbed, (14) is satisfied by construction.

The rest of proof is devoted to verifying (15). Let us use $\lambda = \lambda^e$ and its associated pairwise choice probability $P_{x, \{x, y\}}$ for shorthand notation. Since only items $(n-1)$ and n are swapped when they are ranked at the bottom, we have for all $(x, y) \in \{(x, y) : x < y, (x, y) \neq (n-1, n)\}$,

$$\tilde{P}_{x, \{x, y\}} = P_{x, \{x, y\}} > 1/2.$$

Denote the sum of probabilities (under the Mallows model) of top- $(n-2)$ rankings in Group 1 as $\lambda(\text{Group 1}) := \sum_{\tilde{\pi}} \lambda(\tilde{\pi}) \mathbb{I}\{\tilde{\pi} \in \text{Group 1}\}$. Given the construction rule, we have $\tilde{P}_{n-1, \{n-1, n\}} = \lambda(\text{Group 1})$. Therefore, to verify (15), it suffices to show

$$\lambda(\text{Group 1}) := \sum_{\pi \in \text{Group 1}} \lambda^e(\pi) < \frac{1}{2}.$$

In order to show that $\lambda(\text{Group 1}) < 1/2$, note that Groups 1, 2, and 3 are defined based on the top- $(n-2)$ items of a ranking. In light of this, let us build on [10], who characterize the probability distribution of top- $(n-2)$ rankings under the Mallows model. Formally speaking, we use $\lambda_{n-2} : \Sigma_{n-2} \rightarrow \mathbb{R}$ to denote the pmf of a top- $(n-2)$ ranking under Mallows model. That is, $\lambda_{n-2}(\pi_{n-2}) := \sum_{\pi} \mathbb{I}\{\pi_{n-2} \subseteq \pi\} \lambda^e(\pi)$. Also, in the proof below, given a collection of top- $(n-2)$ rankings $E \subseteq \Sigma_{n-2}$, we use $\Pr(E) := \sum_{\pi_{n-2} \in E} \lambda_{n-2}(\pi_{n-2})$ as a shorthand notation to represent the total mass of top- $(n-2)$ rankings in E .

We (re-)classify all top- $(n-2)$ rankings depending on the positions of items $(n-1)$ and n . Given a top- $(n-2)$ ranking π_{n-2} , we call an item x a *head item* if it ranks in top- $(n-2)$, i.e., $x \in R(\pi_{n-2})$. Otherwise, we call x a *tail item*. We describe the classification of top- $(n-2)$ rankings below.

1. *Class A: Both items $(n-1)$ and item n are tail items.* That is, π_{n-2} belongs to Class A if $\{n-1, n\} = R^c(\pi_{n-2})$. We can further partition Class A into 2 subclasses depending on the two bottom-ranked items of the top- $(n-2)$ ranking:

- (a) *Class A₁: $\pi_{n-2}(n-3) < \pi_{n-2}(n-2)$.*
- (b) *Class A₂: $\pi_{n-2}(n-3) > \pi_{n-2}(n-2)$.*

There is a one-to-one correspondence (i.e., bijection) between Classes A₁ and A₂. For a top- $(n-2)$ ranking a_1 in Class A₁, we can construct a top- $(n-2)$ ranking a_2 in Class A₂ by swapping two bottom-ranked items ranked at positions $(n-3)$ and $(n-2)$:

$$a_2(i) = \begin{cases} a_1(i) & \text{if } i < n-3; \\ a_1(n-2) & \text{if } i = n-3; \\ a_1(n-3) & \text{if } i = n-2. \end{cases}$$

Before proceeding, let's restate a fact from [10].

Fact (Restated from Lemma 1 of [10]) The probability of a top- k ranking π_k under the Mallows ranking model with the identity central ranking is

$$\lambda_k(\pi_k) = q^{d_K(\pi_k)} \cdot \frac{\psi(n-k, q)}{\psi(n, q)},$$

where $d_K(\pi_k) = \sum_{i=1}^k \left(\sum_{j=i+1}^k \mathbb{I}\{\pi_k(i) > \pi_k(j)\} + \sum_{j \in R^c(\pi_k)} \mathbb{I}\{\pi_k(i) > j\} \right)$ is the Kendall's tau distance between top- k ranking π_k and the identity ranking.

In particular, if we take $k = n - 2$ in the result above, we have

$$\lambda_{n-2}(\pi_{n-2}) = q^{d_K(\pi_{n-2})} \cdot \frac{\psi(2, q)}{\psi(n, q)}, \quad (16)$$

where

$$d_K(\pi_{n-2}) = \sum_{i=1}^{n-2} \left(\sum_{j=i+1}^{n-2} \mathbb{I}\{\pi_{n-2}(i) > \pi_{n-2}(j)\} + \sum_{j \in R^c(\pi_{n-2})} \mathbb{I}\{\pi_{n-2}(i) > j\} \right).$$

For each bijection pair (a_1, a_2) , $d_K(a_2) = d_K(a_1) + 1$. Hence, $\lambda_{n-2}(a_2) = q \cdot \lambda_{n-2}(a_1)$. Let $P_A := \Pr(\text{Class } A)$ and $P_{A_i} := \Pr(\text{Class } A_i)$ for $i \in \{1, 2\}$. In light of the bijection construction above, we have $P_{A_2} = q \cdot P_{A_1}$ and $P_A = P_{A_1} + P_{A_2}$. Therefore, we have

$$P_{A_1} = \frac{1}{1+q} P_A \quad \text{and} \quad P_{A_2} = \frac{q}{1+q} P_A. \quad (17)$$

2. *Class B: Item $(n-1)$ is a head item and item n is a tail item.* That is, π_{n-2} belongs to Class B if item $n-1 \in R(\pi_{n-2})$ but item $n \notin R(\pi_{n-2})$. We can further partition Class B into $n-2$ subclasses by the position of item $(n-1)$. For every $m \in \{1, 2, \dots, n-2\}$, the top- $(n-2)$ ranking π_{n-2} belongs to the subclass B_m if it belongs to Class B and item $(n-1)$ is the m^{th} preferred, i.e., $\pi_{n-2}(m) = n-1$.

For every $m \in \{1, 2, \dots, n-2\}$, there is a bijection between Classes A and B_m . For any top- $(n-2)$ ranking $a \in \text{Class } A$, we can obtain a top- $(n-2)$ ranking $b_m \in \text{Class } B_m$ from the following construction:

$$b_m(i) = \begin{cases} a(i) & \text{if } i < m; \\ n-1 & \text{if } i = m; \\ a(i-1) & \text{if } m+1 \leq i \leq n-2. \end{cases}$$

In other words, b_m is obtained from a by first putting item $(n-1)$ at the m^{th} position and, moving the original $m^{\text{th}}, (m+1)^{\text{th}}, \dots, (n-1)^{\text{th}}$ items back one position, and finally leaving the original $(n-2)^{\text{th}}$ as a tail item. Furthermore, by (16), we have $d_K(b_m) = d_K(a) + n-1-m$ and $\lambda_{n-2}(b_m) = q^{n-1-m} \cdot \lambda_{n-2}(a)$.

Let $P_B := \Pr(\text{Class } B)$ and $P_{B_m} := \Pr(\text{Class } B_m)$ for all $m \in \{1, 2, \dots, n-1\}$. In light of bijection construction above, we have $P_{B_m} = q^{n-1-m} \cdot P_A$ and $P_B = \sum_{m=1}^{n-2} P_{B_m}$. Therefore,

$$P_B = \sum_{m=1}^{n-2} q^{n-1-m} \cdot P_A. \quad (18)$$

3. *Class C: Item n is a head item and item $(n-1)$ is a tail item.* That is, π_{n-2} belongs to Class C if item $n-1 \notin R(\pi_{n-2})$ but item $n \in R(\pi_{n-2})$.

There is a bijection between Classes B and C. For any top- $(n-2)$ ranking $b \in \text{Class } B$, we can obtain a top- $(n-2)$ ranking $c \in \text{Class } C$ from the following construction:

$$c(i) = \begin{cases} b(i) & \text{if } i \neq b^{-1}(n-1); \\ n & \text{if } i = b^{-1}(n-1). \end{cases}$$

In other words, c is obtained from b by replacing item $(n-1)$ by item n . By (16), we have $d_K(c) = d_K(b) + 1$ and $\lambda_{n-2}(c) = q \cdot \lambda_{n-2}(b)$. Let $P_C := \Pr(\text{Class } C)$, and we thus have

$$P_C = q P_B = \sum_{m=1}^{n-2} q^{n-m} \cdot P_A. \quad (19)$$

4. *Class D: Item $(n-1)$ and item n are both head items.* That is, π_{n-2} belongs to Class D if $\{n-1, n\} \subseteq R(\pi_{n-2})$. We can partition Class D into $(n-2)(n-3)$ subclasses based on the preference positions of items $(n-1)$ and n . More specifically, given $j \neq k$ and $j, k \in \{1, \dots, n-2\}$, let the subclass

$$D_{(j,k)} := \{\pi_{n-2} \in \text{Class } D : \pi_{n-2}(j) = n-1 \text{ and } \pi_{n-2}(k) = n\}$$

be the collection of top- $(n-2)$ rankings such that items $(n-1)$ and n are ranked in the j^{th} and k^{th} positions, respectively. Furthermore, let the subclass

$$D_1 := \cup_{j < k} \text{Subclass } D_{(j,k)} \quad (\text{resp.} \quad D_2 := \cup_{j > k} \text{Subclass } D_{(j,k)})$$

be the collection of top- $(n-2)$ rankings that rank item $(n-1)$ higher (resp. lower) than item n .

Given $j < k$, there is a bijection between subclass $D_{(j,k)}$ and Class A_1 . For any ranking $a_1 \in \text{Class } A_1$, we can obtain a top- $(n-2)$ ranking $d_{(j,k)} \in \text{Subclass } D_{(j,k)}$ from the following construction:

$$d_{(j,k)}(i) = \begin{cases} a_1(i) & \text{if } i < j \\ n-1 & \text{if } i = j \\ a_1(i-1) & \text{if } j < i < k \\ n & \text{if } i = k \\ a_1(i-2) & \text{if } i > k \end{cases}$$

In other words, to obtain $d_{(j,k)}$ from a_1 , we can put items $(n-1)$ and n at positions j and k respectively, move the original $j^{\text{th}}, (j+1)^{\text{th}}, \dots, (k-2)^{\text{th}}$ items back one position, move the original $(k-1)^{\text{th}}, \dots, (n-4)^{\text{th}}$ items back two positions, and finally leave $\{a_1(n-3), a_1(n-2)\}$ as tail items. By (16), we have $d_K(d_{(j,k)}) = d_K(a_1) + 2n - 1 - j - k$ and $\lambda_{n-2}(d_{(j,k)}) = q^{2n-1-j-k} \cdot \lambda_{n-2}(a_1)$. Let $P_{D_{(j,k)}} := \Pr(\text{Class } D_{(j,k)})$, and we thus have

$$P_{D_{(j,k)}} = q^{2n-1-j-k} \cdot P_{A_1}, \quad \text{for every } j < k. \quad (20)$$

There is also a bijection between subclass D_1 and subclass D_2 . This bijection is similar to that of Classes A_1 and A_2 . For every $d_1 \in \text{Subclass } D_1$, we can swap the positions of items n and $(n-1)$ to obtain a top- $(n-2)$ ranking $d_2 \in \text{Subclass } D_2$. By (16), we have $d_K(d_2) = d_K(d_1) + 1$ and $\lambda_{n-2}(d_2) = q \cdot \lambda_{n-2}(d_1)$. Let $P_{D_i} := \Pr(\text{Class } D_i)$ for $i = 1, 2$, and we thus have

$$P_{D_2} = q \cdot P_{D_1}. \quad (21)$$

Furthermore, let $P_D := \Pr(\text{Class } D)$. Invoking (17), (20) and (21), we have

$$\begin{aligned} P_D &= P_{D_1} + P_{D_2} = (1+q) \cdot P_{D_1} = (1+q) \cdot \sum_{j=1}^{n-3} \sum_{k=j+1}^{n-2} P_{D_{(j,k)}} \\ &= (1+q) \cdot \sum_{j=1}^{n-3} \sum_{k=j+1}^{n-2} q^{2n-1-j-k} \cdot P_{A_1} \\ &= \sum_{j=1}^{n-3} \sum_{k=j+1}^{n-2} q^{2n-1-j-k} \cdot P_A \end{aligned} \quad (22)$$

Because $P_A + P_B + P_C + P_D = 1$, we align (17) to (19) and (22), and conclude that $P_A = \frac{(1-q)(1-q^2)}{(1-q^{n-1})(1-q^n)}$. Therefore, we can obtain the probability of each aforementioned (sub)class.

Clearly, there is a correspondence between Groups 1-3 and Classes A - D : Class B corresponds Group 1 (ii), Class D_1 corresponds Group 1 (i), Class C corresponds Group 2 (ii), Class D_2 corresponds to Group 2 (i), and finally, Class A corresponds Group 3. Therefore,

$$\lambda(\text{Group 1}) = P_B + P_{D_1} = \left(1 - \frac{(1-q)(1-q^2)}{(1-q^{n-1})(1-q^n)}\right) \frac{1}{1+q}.$$

The last equality is due to the following chain of argument: $P_C = q \cdot P_B$ and $P_{D_2} = q \cdot P_{D_1}$; see (19) and (21). Therefore, $P_A + P_B + P_C + P_D = P_A + P_B + P_C + P_{D_1} + P_{D_2} = P_A + (1+q)(P_B + P_{D_1}) = 1$.

As a further consequence, $\lambda(\text{Group 1}) < 1/2$ when n and q satisfy

$$\left(1 - \frac{(1-q)(1-q^2)}{(1-q^{n-1})(1-q^n)}\right) \frac{1}{1+q} < \frac{1}{2},$$

which simplifies to

$$\mathcal{F}_n(q) := 1 + q^{n-1} + q^n - 2q^2 - q^{2n-1} > 0.$$

Note that $\mathcal{F}_n(0) = 1 > 0$, $\mathcal{F}_n(1) = 0$, and $\mathcal{F}'_n(1) = (n-1) + n - 4 - (2n-1) < 0$. Therefore, we conclude that for all n , $\mathcal{F}_n(q) > 0$ either when q is sufficiently close to 0 or 1. That finishes the proof. \square

C.3 Proof of Auxiliary Results (Lemma 5)

Proof of Lemma 5.

$$\begin{aligned} & \operatorname{argmin}_{\pi} \sum_{\tilde{\pi}} \hat{\lambda}(\tilde{\pi}) \cdot d_K(\pi, \tilde{\pi}) \\ &= \operatorname{argmin}_{\pi} \sum_{\tilde{\pi}} \hat{\lambda}(\tilde{\pi}) \sum_{x < y} \mathbb{I}\{(\pi^{-1}(x) - \pi^{-1}(y)) \cdot (\tilde{\pi}^{-1}(x) - \tilde{\pi}^{-1}(y)) < 0\} \\ &= \operatorname{argmin}_{\pi} \sum_{\tilde{\pi}} \hat{\lambda}(\tilde{\pi}) \sum_{x < y} [\mathbb{I}\{\tilde{\pi}^{-1}(x) < \tilde{\pi}^{-1}(y)\} + \mathbb{I}\{\pi^{-1}(x) < \pi^{-1}(y)\} \cdot (1 - 2\mathbb{I}\{\tilde{\pi}^{-1}(x) < \tilde{\pi}^{-1}(y)\})] \\ &= \operatorname{argmin}_{\pi} \sum_{x < y} \mathbb{I}\{\pi^{-1}(x) < \pi^{-1}(y)\} \cdot \sum_{\tilde{\pi}} \hat{\lambda}(\tilde{\pi}) (1 - 2\mathbb{I}\{\tilde{\pi}^{-1}(x) < \tilde{\pi}^{-1}(y)\}) \\ &= \operatorname{argmin}_{\pi} \sum_{x < y} \mathbb{I}\{\pi^{-1}(x) < \pi^{-1}(y)\} \cdot (1 - 2 \sum_{\tilde{\pi}} \hat{\lambda}(\tilde{\pi}) \mathbb{I}\{\tilde{\pi}^{-1}(x) < \tilde{\pi}^{-1}(y)\}) \\ &= \operatorname{argmin}_{\pi} \sum_{x < y} \mathbb{I}\{\pi^{-1}(x) < \pi^{-1}(y)\} \cdot (1 - 2P_{x, \{x, y\}}) \end{aligned}$$

\square

D On the Ranked Choice Probabilities and Model Estimation for $k \geq 1$ (Theorems 4 and 5)

Proof of Theorem 4. Without loss of generality, suppose the display set is given by $S = \{x_1, \dots, x_M\}$ such that $x_1 < x_2 < \dots < x_M$ for some $M \geq 2$. We prove this theorem by forward induction on k (i.e., the length of the ranked list).

Base step. Suppose $k = 1$. Given $\pi_1 = (z)$ where $z \in S$, we have $d_S(\pi_1) = 0$, $L_S(\pi_1) = |\{x \in S \setminus \{z\} : x < z\}|$ is how z is ranked relatively in the display set S under the identity ranking e , and $\frac{\psi(|S|-1, q)}{\psi(|S|, q)} = \frac{1}{1+q+\dots+q^{|S|-1}}$. Therefore,

$$q^{d_S(\pi_1)+L_S(\pi_1)} \cdot \frac{\psi(|S|-1, q)}{\psi(|S|, q)}$$

equals the choice probability (4) in Theorem 1.

Inductive step. Pick an arbitrary $K \in \{2, \dots, M\}$. Suppose our statement holds for every $k = 1, \dots, K-1$. We want to show that our statement holds for $k = K$.

Pick an arbitrary top- k ranking $\pi_k = (x_1^k, x_2^k, \dots, x_{k-1}^k, x_k^k)$ and display set S such that $R(\pi_k) \subseteq S$. For shorthand notation, let $\pi_{k-2} := (x_1^k, x_2^k, \dots, x_{k-2}^k)$ and $\pi_{k-1} := (x_1^k, x_2^k, \dots, x_{k-1}^k)$ so that $\pi_{k-1} = \pi_{k-2} \oplus x_{k-1}^k$ and $\pi_k = \pi_{k-1} \oplus x_k^k$. Note that $\{\pi_k | S\}$ (resp. $\{\pi_{k-1} | S\}$) is the event that the ranked list π_k (resp. π_{k-1}) belongs to the top k (resp. $k-1$) most preferred items within the display set S . Therefore, we have

$$\Pr(\{\pi_k | S\}) = \Pr(\{\pi_{k-1} | S\}) \cdot \Pr(\{x_k^k | S \setminus R(\pi_{k-1})\} | \{\pi_{k-1} | S\}). \quad (23)$$

There are two parts of RHS of (23). We first evaluate the first part: by the induction hypothesis,

$$\Pr(\{\pi_{k-1}|S\}) = q^{d_S(\pi_{k-1})+L_S(\pi_{k-1})} \cdot \frac{\psi(|S|-k+1, q)}{\psi(|S|, q)}. \quad (24)$$

Now we analyze the second part, which is the probability that x_k^k ranks the first among $S \setminus R(\pi_{k-1})$, given that the ranked list π_{k-1} ranks the top- $(k-1)$ among S . Given any $m \in \{k-1, k, \dots, k-1+n-M\}$, let $\Sigma'_{m-1} \subseteq \Sigma_{m-1}$ be the collection of top- $(m-1)$ ranked lists so that $\pi'_{m-1} \in \Sigma'_{m-1}$ if and only if it satisfies the following two properties:

1. $R(\pi'_{m-1}) \cap S = R(\pi_{k-2})$. That is, items included in π'_{m-1} is either included in π_{k-2} , too, or outside of the display set S .
2. π_{k-2} ranks top- $(k-2)$ among S under π'_{m-1} ;

For some quantity c (to be explained later), we can decompose the second part of RHS of (23) as follows:

$$\begin{aligned} & \Pr\left(\{x_k^k | S \setminus R(\pi_{k-1})\} | \{\pi_{k-1}|S\}\right) \\ &= \sum_{m=k-1}^{n-M+k-1} \sum_{\pi'_{m-1} \in \Sigma'_{m-1}} \Pr\left(\{x_k^k | S \setminus R(\pi_{k-1})\} | \pi'_{m-1} \oplus x_{k-1}^k\right) \cdot \Pr\left(\pi'_{m-1} \oplus x_{k-1}^k | \{\pi_{k-1}|S\}\right) \\ &= c \sum_{m=k-1}^{n-M+k-1} \sum_{\pi'_{m-1} \in \Sigma'_{m-1}} \Pr\left(\pi'_{m-1} \oplus x_{k-1}^k | \{\pi_{k-1}|S\}\right) \\ &= c. \end{aligned}$$

We lay out the reasons for each equality below:

(1st equality:) Rule of total probability.

(2nd equality:) Given $m \in \{k-1, k, \dots, k-1+n-M\}$ and $\pi'_{m-1} \in \Sigma'_{m-1}$, let

$$c := \Pr\left(\{x_k^k | S \setminus R(\pi_{k-1})\} | \pi'_{m-1} \oplus x_{k-1}^k\right).$$

be the probability that a (randomly drawn) participant chooses item x_k^k out of the display set $S \setminus R(\pi_{k-1})$ given that he (she) ranks π'_{m-1} as his (her) top- $(m-1)$ preferred items and item x_{k-1}^k as his (her) m^{th} preferred one. By Lemma 3, we have

$$c = \begin{cases} \frac{q^{|\mathbb{I}\{x_k^k < x_{k-1}^k\}}}{1+q+\dots+q^{|\mathbb{I}\{x_k^k < x_{k-1}^k\}}} & \text{if } x_{k-1}^k > x_k^k, \\ \frac{q^{\sum_{x \in S \setminus R(\pi_{k-1})} \mathbb{I}\{x_{k-1}^k < x < x_k^k\}}}{1+q+\dots+q^{|\mathbb{I}\{x_{k-1}^k < x_k^k\}}} & \text{if } x_{k-1}^k < x_k^k. \end{cases} \quad (25)$$

Note that it is a constant independent of both m and π'_{m-1} (given that π_k and display set S are fixed).

(3rd equality:) Note that

$$\Pr(\pi'_{m-1} \oplus x_{k-1}^k | \{\pi_{k-1}|S\}) = \frac{\Pr(\pi'_{m-1} \oplus x_{k-1}^k \text{ and } \{\pi_{k-1}|S\})}{\Pr(\{\pi_{k-1}|S\})} = \frac{\Pr(\pi'_{m-1} \oplus x_{k-1}^k)}{\Pr(\{\pi_{k-1}|S\})}.$$

Therefore,

$$\begin{aligned}
& \sum_{m=k-1}^{n-M+k-1} \sum_{\pi'_{m-1} \in \Sigma'_{m-1}} \Pr \left(\pi'_{m-1} \oplus x_{k-1}^k \mid \{\pi_{k-1} | S\} \right) \\
&= \sum_{m=k-1}^{n-M+k-1} \sum_{\pi'_{m-1} \in \Sigma'_{m-1}} \frac{\Pr(\pi'_{m-1} \oplus x_{k-1}^k)}{\Pr(\{\pi_{k-1} | S\})} \\
&= \frac{1}{\Pr(\{\pi_{k-1} | S\})} \sum_{m=k-1}^{n-M+k-1} \sum_{\pi'_{m-1} \in \Sigma'_{m-1}} \Pr(\pi'_{m-1} \oplus x_{k-1}^k) \\
&= \frac{\Pr(\{\pi_{k-1} | S\})}{\Pr(\{\pi_{k-1} | S\})} = 1.
\end{aligned}$$

Combining (24) and (25), we have

$$\begin{aligned}
\Pr(\{\pi_k | S\}) &= cq^{d_S(\pi_{k-1}) + L_S(\pi_{k-1})} \cdot \frac{\psi(|S| - k + 1, q)}{\psi(|S|, q)} \\
&= q^{d_S(\pi_k) + L_S(\pi_k)} \cdot \frac{\psi(|S| - k, q)}{\psi(|S|, q)}.
\end{aligned}$$

Hence our statement holds for $k = K$, too, thus finishing the proof. \square

Proof of Theorem 5. Suppose $\pi_k = (x_1, \dots, x_k)$ is an arbitrary top- k ranking. Recall from Theorem 4 that $d_S(\cdot)$ and $L_S(\cdot)$ are discrepancy measures with respect to the identity ranking. Through the relabeling argument explained in Section 2, we may generalize the discrepancy measures $d_S(\cdot)$ and $L_S(\cdot)$ to those with respect to an arbitrary (complete) ranking π . Formally, let

$$d_S^\pi := \sum_{h=1}^{k-1} (|S| - h) \cdot \mathbb{I}\{\pi^{-1}(x_h) > \pi^{-1}(x_{h+1})\}$$

and

$$L_S^\pi := \sum_{j \in S \setminus \{x_1, \dots, x_{k-1}\}} \mathbb{I}\{\pi^{-1}(x_k) > \pi^{-1}(j)\},$$

respectively. Let the voting history be $H_T = (S_1, \pi_k^1, \dots, S_T, \pi_k^T)$, where $\pi_k^t = (x_1^t, \dots, x_k^t)$ and it's the t^{th} participant's top- k preferred items. Invoking Theorem 4, the log likelihood for the RMJ-based ranking model parameter (π, q) is

$$\begin{aligned}
\sum_{t=1}^T \log Pr^\pi(\pi_k^t | S_t) &= \sum_{t=1}^T \left[\log \frac{\psi(|S_t| - k, q)}{\psi(|S_t|, q)} + \log q \cdot (d_{S_t}^\pi(\pi_k^t) + L_{S_t}^\pi(\pi_k^t)) \right] \\
&= \sum_{t=1}^T \sum_{i=|S_t|-k+1}^{|S_t|} \log \frac{1-q}{1-q^i} + \log q \cdot \underbrace{\left(\sum_{t=1}^T d_{S_t}^\pi(\pi_k^t) + L_{S_t}^\pi(\pi_k^t) \right)}_*. \quad (26)
\end{aligned}$$

As seen from above, the log likelihood is an affine transformation of the term (*). In other words, there exist constants a, b , both independent of π , such that the log likelihood value can be written as

$a + b \times (*)$. Now let us take a closer look at the term $(*)$.

$$\begin{aligned}
(*) &= \sum_{t=1}^T d_{S_t}^{\pi}(\pi_k^t) + L_{S_t}^{\pi}(\pi_k^t) \\
&= \sum_{t=1}^T \left(\sum_{h=1}^{k-1} (|S_t| - h) \cdot \mathbb{I}\{\pi^{-1}(x_h^t) > \pi^{-1}(x_{h+1}^t)\} + \sum_{j \in S_t \setminus \{x_1^t, \dots, x_{k-1}^t\}} \mathbb{I}\{\pi^{-1}(x_k^t) > \pi^{-1}(j)\} \right) \\
&= \sum_{t=1}^T \left(\sum_{h=1}^{k-1} (|S_t| - h) \cdot \sum_{(i,j): i \neq j} \mathbb{I}\{\pi^{-1}(i) > \pi^{-1}(j)\} \cdot \mathbb{I}\{x_h^t = i, x_{h+1}^t = j\} + \right. \\
&\quad \left. \sum_{(i,j): i \neq j} \mathbb{I}\{\pi^{-1}(i) > \pi^{-1}(j)\} \cdot \mathbb{I}\{x_k^t = i\} \cdot \mathbb{I}\{j \in S_t \setminus R(\pi_k^t)\} \right) \\
&= \sum_{(i,j): i \neq j} \mathbb{I}\{\pi^{-1}(i) > \pi^{-1}(j)\} \cdot \left(\sum_{t=1}^T \left[\sum_{h=1}^{k-1} (|S_t| - h) \cdot \mathbb{I}\{x_h^t = i, x_{h+1}^t = j\} + \right. \right. \\
&\quad \left. \left. \mathbb{I}\{x_k^t = i\} \cdot \mathbb{I}\{j \in S_t \setminus R(\pi_k^t)\} \right] \right).
\end{aligned}$$

Comparing the form above and Proposition 3 of [11], we conclude that to obtain the MLE for the central ranking π^* , it suffices to solve the integer program (5) with the generalized definition of w_{ij} below:

$$w_{ij} := \sum_{t=1}^T \left[\mathbb{I}\{x_k^t = i\} \cdot \mathbb{I}\{j \in S_t \setminus R(\pi_k^t)\} + \sum_{h=1}^{k-1} (|S_t| - h) \cdot \mathbb{I}\{x_h^t = i, x_{h+1}^t = j\} \right].$$

That finishes the proof. \square

E The Expectation-Maximization (EM) for for (Ranked) Choices under the RMJ-based Ranking Model

In this section, we describe the standard expectation maximization (EM) algorithm used to fit a mixture of M RMJ-based ranking models from (ranked) choices. We break the discussion into two steps: When $k = 1$, the data consists of single choices from (different) display sets. Every cluster follows a choice model described in Theorem 1, which is equivalent to the Ordinal Attraction model (OAM). When $k \geq 1$, the data consists of top- k ranked lists, and every cluster follows a ranked choice model described in Theorem 4.

E.1 Single choices ($k = 1$)

In this subsection, we describe the standard expectation maximization (EM) algorithm used to fit a mixture of M OAMs to a given sample of choice data. This particular version is based on [4] and adapted for the OAM setting.

First, introduce the following notation.

- M : number of clusters.
- $Z_t : (c_{t1}, \dots, c_{tM}), c_{tm} \in \{0, 1\}$ indicates whether sample t comes from cluster m .
- $\Theta = (\{p_m\}, \{\pi_m\}, \{\alpha_m\})$: p_m : mixture probability of cluster m , π_m : central ranking of cluster m , α_m : concentration parameter of cluster m .
- $OAM_m(x|S)$: choice probability of item x given display set S under cluster m .

To fit the mixture distribution, we would like to solve the following maximum likelihood problem:

$$\max_{p, \pi, \alpha} \sum_{m=1}^M \sum_{t=1}^T c_{tm} \log p_m + \sum_{m=1}^M \sum_{t=1}^T c_{tm} \log [OAM_m(x_t|S_t)]$$

The EM algorithm starts with an initial solution and iteratively obtains an improving solution until an appropriate stopping criterion is met. Suppose $\{(p_m, \pi_m, \alpha_m) : 1 \leq m \leq M\}$ is the current solution. Then, an improving solution $\{(\hat{p}_m, \hat{\pi}_m, \hat{\alpha}_m) : 1 \leq m \leq M\}$ is obtained as follows.

In the Expectation step, the algorithm computes "soft counts" c_{tm} , denoting the (posterior) probability that example t was generated from mixture component m . This probability is computed as

$$\hat{c}_{tm} = \frac{\hat{p}_m * O\hat{A}M_m(x_t|S_t)}{\sum_{j=1}^M \hat{p}_j * O\hat{A}M_j(x_t|S_t)}$$

Then, in the Maximization step, we first set $\hat{p}_m = \frac{1}{T} \sum_{t=1}^T \hat{c}_{tm}$, and then solve a separate optimization problem for each segment $1 \leq m \leq M$:

$$\max_{\pi_m, \alpha_m} -\alpha_m \sum_{t=1}^T \hat{c}_{tm} \sum_{x \in S_t} \mathbb{I}\{x \succ_{\pi_m} x_t\} - \sum_{t=1}^T \hat{c}_{tm} \log \sum_{j=0}^{|S_t|-1} e^{-\alpha_m j}.$$

We can obtain an optimal solution of the above problem by solving the following two problems:

$$d_m^* = \min_{\pi_m} \sum_{t=1}^T \hat{c}_{tm} \cdot \sum_{x \in S_t} \mathbb{I}\{x \succ_{\pi_m} x_t\} \quad (27)$$

$$\hat{\alpha}_m = \arg \min_{\alpha_m} \left\{ \alpha_m \cdot d_m^* + \sum_{t=1}^T \hat{c}_{tm} \log \sum_{j=0}^{|S_t|-1} e^{-\alpha_m j} \right\} \quad (28)$$

The problem in (27) can be solved by choice aggregation through the integer program (5) (with properly defined w scores). Also, the optimization problem in (28) is convex.

Once we obtain the new solution, $\{(\hat{p}_m, \hat{\pi}_m, \hat{\alpha}_m) : 1 \leq m \leq M\}$, the above process is repeated until the stopping criterion is met. Our stopping criterion is as follows:

1. $\{\hat{\pi}_m : 1 \leq m \leq M\}$ don't change;
2. $\|\hat{p}_{\text{current round}} - \hat{p}_{\text{previous round}}\|_1 < 1e-3$ or $\|\hat{\alpha}_{\text{current round}} - \hat{\alpha}_{\text{previous round}}\|_1 < 1e-3$, where $\mathbf{p} = (p_1, p_2, \dots, p_M)$ and $\boldsymbol{\alpha} = (\alpha_1, \alpha_2, \dots, \alpha_M)$.

Finally, since the EM algorithm is only guaranteed to converge to local optimality, we run (in parallel) 20 instances the EM algorithm with random initialization and select the parameters with the best log-likelihood value.

E.2 The general setting of ranked choices ($k \geq 1$)

In this subsection, we describe the standard expectation maximization (EM) algorithm used to fit a mixture of M RMJ-based ranking models from a sample of top- k ranked choice data. This particular version is based on [4] and adapted for the ranked choice setting.

First, introduce the following notation.

- k : number of ranked items in each sample
- M : number of clusters.
- $Z_t : (c_{t1}, \dots, c_{tM}), c_{tm} \in \{0, 1\}$ indicates whether sample t comes from cluster m .
- $\Theta = (\{p_m\}, \{\pi_m\}, \{\alpha_m\})$: p_m : mixture probability of cluster m , π_m : central ranking of cluster m , α_m : concentration parameter of cluster m .
- $RCM_m(\pi_k|S)$: ranked choice probability of π_k given display set S under cluster m .

To fit the mixture distribution, we would like to solve the following maximum likelihood problem:

$$\max_{p, \pi, \alpha} \sum_{m=1}^M \sum_{t=1}^T c_{tm} \log p_m + \sum_{m=1}^M \sum_{t=1}^T c_{tm} \log [RCM_m(\pi_k^t|S_t)]$$

The EM algorithm starts with an initial solution and iteratively obtains an improving solution until an appropriate stopping criterion is met. Suppose $\{(p_m, \pi_m, \alpha_m) : 1 \leq m \leq M\}$ is the current solution. Then, an improving solution $\{(\hat{p}_m, \hat{\pi}_m, \hat{\alpha}_m) : 1 \leq m \leq M\}$ is obtained as follows.

In the Expectation step, the algorithm computes "soft counts" c_{tm} , denoting the (posterior) probability that example t was generated from mixture component m . This probability is computed as

$$\hat{c}_{tm} = \frac{\hat{p}_m * RC\hat{M}_m(\pi_k^t | S_t)}{\sum_{j=1}^M \hat{p}_j * RC\hat{M}_j(\pi_k^t | S_t)}$$

Then, in the Maximization step, we first set $\hat{p}_m = \frac{1}{T} \sum_{t=1}^T \hat{c}_{tm}$, and then solve a separate optimization problem for each segment $1 \leq m \leq M$:

$$\max_{\pi_m, \alpha_m} -\alpha_m \sum_{t=1}^T \hat{c}_{tm} (d_{S_t}(\pi_k^t) + L_{S_t}(\pi_k^t)) - \sum_{t=1}^T \hat{c}_{tm} \sum_{i=n-k+1}^n \log \sum_{j=0}^{i-1} e^{-\alpha_m j}$$

We can obtain an optimal solution of the above problem by solving the following two problems:

$$d_m^* = \min_{\pi_m} \sum_{t=1}^T \hat{c}_{tm} (d_{S_t}(\pi_k^t) + L_{S_t}(\pi_k^t)) \quad (29)$$

$$\hat{\alpha}_m = \arg \min_{\alpha_m} \left\{ \alpha_m \cdot d_m^* + \sum_{t=1}^T \hat{c}_{tm} \sum_{i=n-k+1}^n \log \sum_{j=0}^{i-1} e^{-\alpha_m j} \right\} \quad (30)$$

The problem in (29) can be solved by choice aggregation through the integer program (5) (with w defined in Theorem 5). Also, the optimization problem in (30) is convex.

Once we obtain the new solution, $\{(\hat{p}_m, \hat{\pi}_m, \hat{\alpha}_m) : 1 \leq m \leq M\}$, the above process is repeated until until the stopping criterion is met. Our stopping criterion is as follows:

1. $\{\hat{\pi}_m : 1 \leq m \leq M\}$ don't change;
2. $\|\hat{\mathbf{p}}_{\text{current round}} - \hat{\mathbf{p}}_{\text{previous round}}\|_1 < 1e-3$ or $\|\hat{\boldsymbol{\alpha}}_{\text{current round}} - \hat{\boldsymbol{\alpha}}_{\text{last round}}\|_1 < 1e-3$, where $\mathbf{p} = (p_1, p_2, \dots, p_M)$ and $\boldsymbol{\alpha} = (\alpha_1, \alpha_2, \dots, \alpha_M)$.

Finally, since the EM algorithm is only guaranteed to converge to local optimality, we run (in parallel) 20 instances the EM algorithm with random initialization and select the parameters with the best log-likelihood value.