

Dark Energy Black Holes with Intermediate Masses at High Redshifts: an earlier generation of Quasars and observations.

Anupam Singh

Department of Physics, L.N. Mittal I.I.T, Jaipur, India.

(Dated: July 6, 2022)

Dark Energy is the largest fraction of the energy density of our Universe - yet it remains one of the enduring enigmas of our times. Here we show that Dark Energy can be used to solve 2 tantalizing mysteries of the observable universe. We build on existing models of Dark Energy linked to neutrino masses. In these models Dark Energy can undergo Phase Transitions and form Black Holes. Here we look at the implications of the family structure of neutrinos for the phase transitions in dark energy and associated peaks in black hole formation. It has been previously shown that one of these peaks in Black Hole formation is associated with the observed peak in Quasar formation. Here, we predict that there will also be an earlier peak in the Dark Energy Black Holes at high redshifts. These Dark Energy Black Holes formed at high redshifts are Intermediate Mass Black Holes (IMBHs). These Dark Energy Black Holes at large redshift can help explain both the EDGES observations and the observations of large Supermassive Black Holes (SMBHs) at redshifts $z \sim 7$ through the accretion onto the Dark Energy IMBHs at high redshifts. Thus, these Dark Energy Black Holes solve 2 of the most puzzling mysteries of the observable early universe. Not only does the existence of an earlier phase of Dark Energy Black Holes take care of some current challenges to theory implied by existing astronomical data, it also helps us actively look for these Dark Energy Black Holes at these high redshifts as predicted here through targeted searches for these Black Holes at the redshifts $z \sim 18$. There is a slight dependence of the location of the peak on the lightest neutrino mass - so the peak may be located at a slightly lower value of the redshift. This may actually enable a measurement of the lightest neutrino mass - something which has eluded us so far. Finding these Dark Energy Black Holes of Intermediate Mass should be within the reach of upcoming observations - particularly with the James Webb Space Telescope - but perhaps also through the use of other innovative techniques focusing specifically on the redshifts around $z \sim 18$.

I. INTRODUCTION

Cosmology and Astrophysics were once data starved sciences, today they have become data driven sciences. This has pushed Cosmology, Astrophysics and fundamental physics much closer together - benefitting progress in all of these exciting areas. Thus, the measurements of the Hubble constant and the age of the universe forced us to re-visit the simplest cosmological model - a flat universe with a zero cosmological constant[1, 2]. These observations led us[3] to the idea of a non-vanishing vacuum energy due to light fields as playing a significant role in the Universe. We now have a substantial accumulation of research that has bolstered our belief in what we now call dark energy. This work brings together cutting edge observational work in astrophysics and cosmology and theoretical work in fundamental areas of physics including field theory, elementary particle physics and gravitation. For an exhaustive and elaborate discussion on Dark Energy and related issues, please see the review [4] .

From the perspective of field theory and elementary particle physics, Dark Energy is a low energy phenomenon and so a description in terms of fields in curved space-time can be used to understand many aspects of Dark Energy including the current observational implications of Dark Energy. Here, we study Dark Energy and its observational consequences by taking a field theory approach. Field theory coupled to the Standard Model of Elementary Particle Physics has been extremely success-

ful in helping us understand cosmology upto extremely high energies - please see for example[5]. Indeed, at least the expectation was that it would continue to help us understand the low energy universe we currently inhabit. In this context, observations of Dark Energy raised some uncomfortable questions. However, a small extension of the Standard Model may help us understand Dark Energy and its consequences. In the Standard Model of Particle Physics, neutrinos are massless particles. A common extension of the Standard Model is to give the neutrinos a mass. Indeed, massive neutrinos have long been considered to provide the most promising window to the physics beyond the Standard Model. This led to an intense world-wide observational program for the search of neutrino masses resulting in the discovery and measurement of neutrino masses - please see for example[6] for a recent review. It turns out, as we will discuss below, that massive neutrinos may indeed also provide a promising pathway forward to build on top of the standard model of cosmology. In particular, the physics of massive neutrinos may help us understand the phenomena associated with Dark Energy and its observational consequences as discussed below.

Field theory models for dark energy motivated by particle physics were discussed earlier [3]. These fields involved must have very light masses so as to be cosmologically relevant today or in the recent past. Investigations of particle physics models with naturally light masses that can generate interesting cosmological consequences at late times have been done by several authors[7, 8].

Thus, it was noted that the natural models in which we can get interesting effects are models of neutrino masses with Pseudo Nambu Goldstone Bosons (PNGB's). The mass scales in these models can be related to the neutrino masses. Further, any tuning if required can be protected from radiative corrections by the symmetry underlying the Nambu-Goldstone bosons[9].

PNGBs have a celebrated history in the annals of cosmology because at the late times relevant for observational cosmology any dynamical fields must have small masses compared to the usual masses encountered in the standard model and PNGBs provide a natural mechanism from obtaining light fields. Of the PNGBs perhaps the most celebrated is the axion (please see e.g. [10–14] for more details). Of all the particles we currently know and have detected the particles with the smallest non-vanishing masses are the neutrinos and so it should not be surprising that PNGBs linked to neutrino masses can be important for understating the current cosmological dynamics.

Let us now take a closer look at the model and its associated observational implications.

One of the standard ways of generating neutrino masses has been the see-saw mechanism [15]. These models involve a Yukawa interaction between the Higgs fields and leptons. We investigated the finite temperature behaviour of the see-saw model. For this purpose, we selected a very specific and simplified version of the general see-saw model but one which is sufficient to give us some very interesting results.

The relevant Yukawa couplings in the leptonic sector are:

$$-\mathcal{L}_{\text{yuk}} = y_{ai} \bar{l}_L^a N_R^i \Phi + y \overline{N_R^i} N_R^j \sigma_{ij} + \text{h.c.} \quad (1)$$

where a, i, j are family indices and the fields that enter into the see-saw model of neutrino masses are: the standard lepton doublets l_L and the Higgs doublet Φ along with right handed neutrinos N_R^i and a Higgs field σ_{ij} . As has previously been discussed in detail in [3] the PNGB fields in this model can lead to a rapid expansion of the Universe which is consistent with the observed expansion rate of the Universe. This of course is a requirement for any model of what we now call Dark Energy.

Dark Energy is the largest fraction of the energy density of the Universe. Thus, we expect that Dark Energy can play an important role in multiple facets of the Universe.

Thus, we investigated whether this Dark Energy has other important or interesting effects beyond driving the rapid expansion of the Universe on large scales. Since we have a Dark Energy model arising out of fields with a space-time dependence, we can have a rapid expansion of the Universe on large length scales and at the same time have non-trivial and interesting space time dependent dynamics on shorter length scales driven by the dynamical length scale of the Dark Energy field as was shown in [17].

In fact, the physics of these fields is very interesting because it allows for Phase Transitions as the temperature of the Universe drops due to the expansion of the Universe as has previously been shown in [7, 8, 18]. In particular, Black Holes can be formed in these Phase Transitions as previously discussed[17, 18].

In the following sections we will elaborate on these ideas with the goal of arriving at a prediction for the redshift at which an earlier generation of quasars may be observed given the current data on neutrino masses. We will also arrive at an estimate of the BH masses and then compare to existing observations and point out the opportunity to confirm these ideas through future observations which are on our current horizon and feasible in the near future.

II. DARK ENERGY PHASE TRANSITIONS AND BLACK HOLE FORMATION

Extensive discussions of the dynamics of fields and phase transitions in cosmological settings has been done previously (see e.g. Kolb and Turner[5]). Standard discussions also exist for the gravitational dynamics of fields (see e.g. Weinberg[16]). We have built on these techniques to study and understand the physics of phase transitions in a general cosmological setting (please see e.g. [19–24]) and applied it to the phase transitions in dark energy and the dynamics of dark energy fields.

The finite temperature behavior of the see-saw model of neutrino masses has been previously studied[7, 8]. Phase transitions were found which result in the formation of topological defects. The critical temperature in the see-saw model is naturally linked to the neutrino masses.

In earlier papers, it has been shown that Dark Energy can play a significant role in structure formation and the physics of the galaxy [17, 18]. In particular, the dynamical length scale associated with the Dark Energy is comparable to the galactic length scale. Further, Dark Energy collapse can result in the formation of black holes.

In many of our investigations of Dark Energy, we focus mostly on the gravitational interactions of the Dark Energy. It is the dominant interaction of Dark Energy. In this context, the gravitational waves from Dark Energy constitute a new and interesting signal. Thus, we studied the gravitational waves arising Dark Energy dynamics[29]. A result from that paper is useful is that the time period of the gravitational waves emitted is $\sim 10^5$ years.

In a previous paper we have shown that the gravitational waves emitted by Dark Energy can result in the periodicity of the ice ages through the ellipticity variations of the orbit of the earth[30].

It is clear that Dark Energy has many important consequences for our Universe. We now wish to focus on phase transitions in Dark Energy and its implications for our Universe. To do this we will build on the work of Hol-

man and Singh[8]. They investigated the finite temperature see-saw model of neutrino masses and demonstrated the existence of interesting phase transitions. The critical temperature is determined by the neutrino masses. Further the phase transition results in the formation of topological defects. Finally, the gravitational collapse of these field configurations results in black hole formation [17, 18]. It has been shown that these Black Holes are Supermassive Black Holes (SMBHs). In fact, their masses are comparable to the observed masses of Black Holes at the centers of galaxies. Knowing the link between neutrino masses and the critical temperature of the phase transition[8], we could infer when to expect the peak in black hole formation to take place. This was done in [18] and indeed the observations matched the theoretical expectation. It should be noted that at the time of writing of [18] the only data available on neutrino masses was the Solar Neutrino data[31] from which one could arrive at an estimate of m_2 - the mass of the second family of neutrinos. Of course, neutrinos have a family structure like every other known elementary particle and we will label the three families of light neutrinos as 1, 2 and 3 in increasing order of mass, so they will have masses given by m_1 , m_2 and m_3 . We now have an improved understanding of neutrino masses. In particular, atmospheric neutrino data allows us to make some additional inferences and as we will see below implies an additional peak in Black Hole formation at higher redshifts i.e. at an earlier epoch in the history of the Universe. Thus, we expect to see an earlier generation of quasars linked to a more massive generation of neutrinos.

III. NEUTRINO MASSES AND THEIR IMPLICATIONS

A significant amount of progress has been made in determining neutrino masses - particularly using Neutrino Oscillations as a probe of neutrino masses. Please see for example[6] for a recent review. From the neutrino oscillation data using the normal hierarchy of neutrino families, we can extract[6] the neutrino masses - in limit that the lightest neutrino mass is insignificant compared to the masses of the heavier neutrinos (i.e. $m_1 \ll m_2, m_3$) - we get that:

$$m_3 \simeq 5 \times 10^{-2} eV \quad (2)$$

and

$$m_2 \simeq 9 \times 10^{-3} eV. \quad (3)$$

We now wish to use this to relate the epochs at which the Phase Transitions occur resulting in the peaks in the associated Black Hole formation. We have already labeled the three families of light neutrinos as 1, 2 and 3 in increasing order of mass, so they will have masses given

by m_1 , m_2 and m_3 . Let us now label the critical temperatures associated with the phase transitions as T_1 , T_2 , T_3 and scale factors associated with these phase transitions as R_1 , R_2 , R_3 respectively. The scale factors can themselves be associated with the corresponding redshifts z_1 , z_2 , z_3 respectively.

Using the physics of the phase transitions and the evolution of temperature, scale factor and redshift in an expanding universe[5, 8, 16, 18] we note that:

$$\frac{T_3}{T_2} = \frac{m_3}{m_2} \quad (4)$$

$$\frac{T_3}{T_2} = \frac{R_2}{R_3} \quad (5)$$

and

$$\frac{R_2}{R_3} = \frac{1 + z_3}{1 + z_2}. \quad (6)$$

From observations of the Quasar distribution with redshift, see for example[32], we get:

$$z_2 \sim 2.5. \quad (7)$$

The value of z_3 has a dependence on the lightest neutrino mass m_1 and so the result is displayed as a plot of z_3 versus m_1 in the figure below.




FIG. 1. Peak Redshift as a function of m_1 .

Figure 1 displays the location of the Peak in redshift (z) as function of the mass of the lightest neutrino (m_1 in meV).

If the lightest neutrino mass is insignificant compared to the masses of the heavier neutrinos (i.e. $m_1 \ll m_2, m_3$), then we get:

$$z_3 \sim 18. \quad (8)$$

It is not possible to extract the value of m_1 from neutrino oscillation data, but we can actually evaluate the

peak redshift as a function of the mass of lightest neutrino mass this has been done by us. Indeed, the location of the peak at high redshifts may enable us to extract the value of the lightest neutrino mass which has so far eluded us.

At this point it is worthwhile considering the implications of such Black Holes at the relatively high redshifts of $z \sim 18$. It turns out that this might ease some tensions in existing astronomical data.

IV. MASSES OF DARK ENERGY BLACK HOLES FORMED AT HIGH REDSHIFTS

Before we discuss the data, in order to assess the impact of these Dark Energy Black Holes produced at high redshifts, we also need to arrive at the masses of these Dark Energy Black Holes produced due to the phase transition linked to the mass m_3 .

Black Hole (BH) formation as a result of cosmological phase transitions has been extensively studied before by a number of groups. Collapse of false vacuum domains to produce BHs was investigated by Kodama, Sasaki, and Sato[25] and Hsu[27]. BH formation from bubble wall collisions was studied by Hawking, Moss, and Stewart[26]. BH formation from the collapse of closed domain walls was investigated by Ipser and Sikivie [33] and by Widrow[28]. Widrow suggested that the collapse of closed domain walls is likely to produce BHs due to Phase Transitions that occur after the recombination. The mass of such BHs produced has previously been obtained[17, 18] and can be stated as:

$$M_{BH} \simeq 4\pi R^2 \times \sigma. \quad (9)$$

For the case being considered here $\sigma \sim \alpha m_3^2 f$ where f is the heavy neutrino scale in the see-saw model of neutrino masses. Numerical work studying the gravitational collapse of the Dark Energy fields resulting in the formation of Black Holes[17] gives us the numerical value of $\alpha \simeq 10$.

Black Hole formation in Dark Energy Phase Transitions has been studied earlier [17, 18] to arrive at the Black Holes masses. Using the same techniques as earlier but now using the mass of the heaviest neutrino m_3 to arrive at the masses of Black Holes formed at the higher redshifts $z_3 \sim 18$, we get:

$$M_{BH} \simeq 4.3 \times 10^4 \times \left(\frac{R}{25pc} \right)^2 \quad (10)$$

where R is the size of the dark energy field configuration that collapses. It should be noted that the value of $R_c \sim 25pc$ being used here follows from the dynamical length scale f/m_3^2 for the dark energy fields being considered here. Thus, we see that the Dark Energy Black

Holes formed at high redshifts come in the desired Intermediate Mass Black Hole category that can get rid of the current tensions in astronomical data.

V. MEETING THE CHALLENGES IMPLIED BY CURRENT ASTRONOMICAL DATA

In recent years, there has been a recognition that even the existence of quasars at redshifts $z \sim 7$ is at odds with our current theories of supermassive black hole formation - please see for example the recent review[33]. As noted in this review (and also elsewhere), the growth time t_{growth} for an Eddington limited accretion of a black hole growing from an initial mass M_0 to a final mass M_{BH} with radiative efficiency ϵ is given by:

$$t_{growth} \simeq 0.45 \frac{\epsilon}{1-\epsilon} \ln \frac{M_{BH}}{M_0} Gyr. \quad (11)$$

The implication of this is that producing high mass quasars at high redshifts within the available time dictated by the age of the universe becomes a severe challenge for traditional models of black hole formation. This challenge can be overcome if the early phase of black hole formation due to the phase transition linked to neutrinos as described here is taken into account. This is certainly very promising in the context of the ideas discussed in this article. Of course, an explicit observation of such black holes at a redshift $z \sim 18$ would be more compelling and hence searches for black holes at these high redshifts is strongly advocated. This should be within the reach of upcoming observations - particularly with the James Webb telescope - but perhaps also through the use of other innovative techniques focusing specifically on the redshifts around $z \sim 18$.

It turns out that there is already a hint that such Black Holes may have been formed at the high redshifts described here. The EDGES instruments have detected an absorption profile in the 21 cm line centred at redshifts spanning approximately $15 < z < 20$, centered at $z \sim 17$ [34].

Dark Energy Black Holes with the Intermediate Masses as described here can easily explain both the EDGES data as described above and also the existence of large SMBHs at redshifts of $z \sim 7$ which can easily result through accretion from the seed Dark Energy IMBHs produced at the redshifts of $z \sim 18$ as described here.

Thus, the Dark Energy Black Holes described here solve 2 of the most perplexing mysteries in the observed universe today. At this point, it may be worthwhile to look directly for these IMBHs and our current analysis shows that focusing on redshifts $z \sim 18$ in a search targeted in redshift space may help arrive at an early detection of these Black Holes.

VI. SUMMARY AND CONCLUSION

A major takeaway from our discussion here is that Dark Energy can provide a source of Black Holes at high redshift. Further, we note that the masses of the Dark Energy Black Holes at high redshifts that follow from our calculations place these Black Holes in the desired Intermediate Mass Black Hole category. This can help alleviate existing tensions in astronomical data. Further, it provides a target in redshift space to focus on as we look for high redshift black holes. We have predicted here

the redshifts $z \sim 18$ at which these Intermediate Mass Black Holes should be formed. Searches for black holes at these high redshifts is strongly advocated. This should be within the reach of upcoming observations - particularly with the James Webb Space Telescope - but perhaps also through the use of other innovative techniques focusing specifically on the redshifts around $z \sim 18$.

-
- [1] M. J. Pierce, D. L. Welch, R. D. McClure, S. van den Bergh, R. Racine and P. B. Stetson, *Nature*, 371, (September 1994), pp. 385-389.
- [2] Freedman, W.L., et al , *Nature*, **371**, 757-762 (1994).
- [3] Singh, Anupam, *Phys. Rev. D* 52, 6700-6707, (1995).
- [4] A Silvestri, M Trodden, *Reports on Progress in Physics* 72 (9), 096901.
- [5] E. W. Kolb and M. S. Turner : *The Early Universe*, 73. (Addison-Wesley Pub. Co. , 1990).
- [6] Neutrino Masses, Mixing, and Oscillations by M.C. Gonzalez-Garcia and M.Yokoyama in P.A. Zyla et al. (Particle Data Group), *Prog. Theor. Exp. Phys.* 2020, 083C01 (2020).
- [7] A. K. Gupta, C. T. Hill, R. Holman and E. W. Kolb, *Phys. Rev. D* 45, 441 (1992).
- [8] Richard Holman and Anupam Singh, *Phys. Rev. D* 47, 421 (1993).
- [9] see, e. g. , G. 'tHooft "Naturalness, Chiral Symmetry, and Spontaneous Chiral Symmetry Breaking," in *Recent Developments in Gauge Theories*, (Plenum Press, New York, 1980), G. 't Hooft, et al. , eds.
- [10] J. Preskill, M. Wise and F. Wilczek, *Phys. Lett. B* **120**,127 (1983).
- [11] L. Abbott and P. Sikivie, *Phys. Lett. B* **120**,133 (1983).
- [12] M. Dine and W. Fischler, *Phys. Lett. B* **120**,137 (1983).
- [13] R. Peccei and H. Quinn, *Phys. Rev. Lett.* **38**, 1440 (1977) ; S. Weinberg, *Phys. Rev. Lett.* **40**, 223 (1978) ; F. Wilczek, *Phys. Rev. Lett.* **40**, 279 (1978) ; M. Dine, W. Fischler and M. Srednicki, *Phys. Lett. B* **104**, 199 (1981).
- [14] Edward W. Kolb, Anupam Singh, and Mark Srednicki *Phys. Rev. D* 58, 105004 (1998)
- [15] M. Gell-Mann, P. Ramond, R. Slansky, in *Supergravity*, North Holland Pub. (1979); T. Yanagida, in "Proc. of the Workshop on Unified Theories and Baryon Number in the Universe," KEK, Japan (1979).
- [16] Weinberg, S. : *Gravitation and Cosmology*. (John Wiley & Sons, Inc. , 1972).
- [17] Vishal Jhalani, Himanshu Kharkwal and Anupam Singh, Gravitational collapse of dark energy field configurations and supermassive black hole formation, ISSN 1063-7761, *Journal of Experimental and Theoretical Physics*, Vol. 123, No. 5, pp. 827-831, (2016) .
- [18] Quasar production: Topological defect formation due to a phase transition linked with massive neutrinos. A Singh. *Physical Review D* 50 (2), 671, (1994).
- [19] Daniel Boyanovsky, Da-Shin Lee, and Anupam Singh, *Phys. Rev. D* **48** , 800 (1993).
- [20] D Boyanovsky, R Holman, DS Lee, JP Silva, A Singh, *Nuclear Physics B* 441 (3), 595-608 (1995)
- [21] Daniel Boyanovsky, Hector J. de Vega, Richard Holman, Da-Shin Lee, and Anupam Singh,*Phys. Rev. D* **51**, 4419 (1995).
- [22] D Boyanovsky, D Cormier, HJ De Vega, R Holman, A Singh, M Srednicki, *Physical Review D* 56 (4), 1939 (1997).
- [23] A Chodos, F Cooper, W Mao, H Minakata, A Singh, *Physical Review D* 61 (4), 045011 (2000).
- [24] A Chodos, F Cooper, W Mao, A Singh *Physical Review D* 63 (9), 096010 (2001).
- [25] H. Kodama, M. Sasaki, and K. Sato, *Prog. Theor. Phys.* 68, 1979 (1982).
- [26] S. W. Hawking, I. G. Moss, and J. M. Stewart, *Phys. Rev. D* 30, 2681 (1982).
- [27] S. D. H. Hsu, *Phys. Lett. B* 251, 343 (1990)
- [28] L M. Widrow, *Phys. Rev. D* 40, 1002 (1989)
- [29] Vishal Jhalani, Ayushi Mishra, and Anupam Singh, ISSN 1063-7761, *Journal of Experimental and Theoretical Physics*, 2017, Vol. 125, No. 4, pp. 638-643, (2017).
- [30] Anupam Singh, Dark energy gravitational wave observations and ice age periodicity, *Physics Letters B* 802:135226 (2020), DOI: 10.1016/j.physletb.2020.135226.
- [31] X. Shi, D. N. Schramm, and J. N. Bahcall, *Phys. Rev. Lett.* 69, 717 (1992)
- [32] The Sloan Digital Sky Survey Quasar Catalog: Fourteenth data release, I. Paris et.al. *Astronomy and Astrophysics* 613, A51 (2018).
- [33] Titans of the early Universe: The Prato statement on the origin of the first supermassive black holes, *Publications of the Astronomical Society of Australia* , 36, e027,(2019). doi:10.1017/pasa.2019.14
- [34] An absorption profile centred at 78 megahertz in the sky-averaged spectrum, JD Bowman, AEE Rogers, RA Monsalve, TJ Mozdzen, N Mahesh *Nature* 555 , 67-70 (2018).