

Branching Processes in Random Environments with Thresholds

Giacomo Francisci and Anand N. Vidyashankar

George Mason University

July 22, 2022

Abstract

Motivated by applications to COVID dynamics, we describe a branching process in random environments model $\{Z_n\}$ whose characteristics change when crossing upper and lower thresholds. This introduces a cyclical path behavior involving periods of increase and decrease leading to supercritical and subcritical regimes. Even though the process is not Markov, we identify subsequences at random time points $\{(\tau_j, \nu_j)\}$ - specifically the values of the process at crossing times, *viz.*, $\{(Z_{\tau_j}, Z_{\nu_j})\}$ - along which the process retains the Markov structure. Under mild moment and regularity conditions, we establish that the subsequences possess a regenerative structure and prove that the limiting normal distribution of the growth rates of the process in supercritical and subcritical regimes decouple. For this reason, we establish limit theorems concerning the length of supercritical and subcritical regimes and the proportion of time the process spends in these regimes. As a byproduct of our analysis, we explicitly identify the limiting variances in terms of the functionals of the offspring distribution, threshold distribution, and environmental sequences.

Key Words. BPRE, COVID dynamics, Ergodicity of Markov chains, Estimators of growth rate, Length of cycles, Martingales, Random sums, Regenerative structure, Size-dependent branching process, Size-dependent branching process with a threshold, Subcritical regime, Supercritical regime.

Math Subject Classification 2020. *Primary:* 60J80, 60F05, 60J10; *Secondary:* 92D25, 92D30, 60G50, 62F10.

1 Introduction

Branching processes and their variants are used to model various biological, biochemical, and epidemic processes (see Jagers (1975); Haccou et al. (2007); Hanlon and Vidyashankar (2011); Kimmel and Axelrod (2015)). More recently, these methods have been used as a model for spreading COVID cases in a community during the early stages of the pandemic (Yanev et al.; 2020; Atanasov et al.;

2021). As the time progressed, the number of infected members in a community changed due to different containment efforts of the local communities (Falcó and Corral; 2021; Sun et al.; 2022) leading to periods of increase and decrease. In this paper, we describe a stochastic process model built on a branching process model in random environments that explicitly takes into account periods of growth and decrease in the transmission rate of the virus.

Specifically, we consider a branching process model initiated by a random number of ancestors (thought of as initiators of the pandemic within a community). During the first several generations, the process grows uncontrolled, allowing immigration into the system. This initial phase is modeled using a supercritical branching process with immigration in random environments, specifically independent and identically distributed (i.i.d.) environments. When consequences of rapid spread become significant, policymakers introduce restrictions to reduce the rate of growth, hopefully resulting in a reduced number of infected cases. The limitations are modeled using upper thresholds on the number of infected cases, and beyond the threshold the process changes its character to evolve as a subcritical branching process in random environments. During this period - due to strict controls - immigration is also not allowed. In practical terms, this period typically involves a “lockdown” and other social containment efforts, the intensity of which varies across communities.

The period of restrictions is not sustainable for various reasons, including political, social, and economic pressures leading to the easing of controls. The policymakers use multiple metrics to gradually reduce controls, leading to an “opening of communities”, resulting in increased human interactions. As a result or due to changes undergone by the virus, the number of infected cases increases again. We use lower thresholds in the number of “newly infected” to model the period of change and let the process evolve again as a supercritical BPRE in i.i.d. environments after it crosses the lower threshold. The process continues to evolve in this manner alternating between periods of increase and decrease. In this paper, we provide a rigorous probabilistic analysis of this model.

Even though we used the dynamics of COVID spread as a motivation for the proposed model, the aforementioned cyclic behavior is often observed in other biological systems, such as those modeled as a predator-prey model or the SIR model. In some biological populations, the cyclical behavior can be attributed to the decline of fecundity as the population size approaches a threshold (Klebaner; 1993). Deterministic models such as ordinary differential equations, dynamical systems, and chaos theory are used for analysis in the applications mentioned above (Teschl; 2012; Perko; 2013; Iannelli and Pugliese; 2014). While many models described above yield good qualitative descriptions, uncertainty estimates are typically unavailable. It is worthwhile to point out that previously described branching process methods also produce reasonable point estimates for the mean growth during the early stages of the pandemic. However, the above-mentioned point estimates of the growth rate are unreliable during the later stages of the pandemic. In this paper, we address statistical estimation of the mean growth and characterize the variance of the estimates. We end the discussion with a plot, Figure 1, of the total number of confirmed COVID cases per week in Italy from February 23, 2020 to July 20, 2022. The plot also includes the number of cases using the proposed model. Other examples with similar plots include the hare-lynx predator-prey dynamics and measles cases (Tyson et al.; 2010; Iannelli and Pugliese; 2014; Hempel and Earn; 2015).

Weekly COVID cases in Italy




Figure 1: In black weekly COVID cases in Italy from February 23, 2020 to July 20, 2022. In blue a BPRE starting with the same initial value and offspring mean having the negative binomial distribution with predefined number of successful trials $r = 10$ and Gamma-distributed mean with shape parameter equal to the mean of the data and rate parameter 1.

Before we provide a precise description of our model, we begin with a brief description of BPRE with immigration. Let $\zeta = \{\zeta_n\}_{n=0}^\infty$ be a collection of i.i.d. random variables taking values in the space of pairs of probability distributions E . Associated with each ζ_n are two probability generating functions (pgf(s)) $\varphi_{\zeta_n}(\cdot)$ and $\psi_{\zeta_n}(\cdot)$ given by

$$\varphi_{\zeta_n}(s) = \sum_{j=0}^{\infty} p_j(\zeta_n) s^j \text{ and } \psi_{\zeta_n}(s) = \sum_{j=0}^{\infty} p_{I,j}(\zeta_n) s^j.$$

For each realization of the process ζ and the pgfs $\varphi_{\zeta_n}(\cdot)$ and $\psi_{\zeta_n}(\cdot)$, we associate a population process $\{Z_n\}_{n=0}^\infty$ defined recursively as follows: let Z_0 take values on the positive integers and $n \geq 0$

$$Z_{n+1} = \sum_{i=1}^{Z_n} \xi_{n,i} + I_n,$$

where, given ζ_n , $\{\xi_{n,i}\}_{i=1}^\infty$ are i.i.d. with pgf $\varphi_{\zeta_n}(\cdot)$ and I_n is an independent random variable with pgf $\psi_{\zeta_n}(\cdot)$. The random variable $Y = \log(m(\zeta_0))$, where $m(\zeta_0) = \mathbf{E}[\xi_{0,1}|\zeta_0]$, plays an important role in classification of BPRE with immigration. It is well-known that when $\mathbf{E}[Y] > 0$ the process diverges to infinity with probability one and if $\mathbf{E}[Y] \leq 0$ and the immigration is degenerate at zero for all environments then the process becomes extinct with probability one (Athreya and Karlin; 1971). Furthermore, in the subcritical case, that is $\mathbf{E}[Y] < 0$, one can further identify three distinct regimes: (i) weakly subcritical, (ii) moderately subcritical, and (iii) strongly subcritical. (i) corresponds to

when there exists a $0 < \rho < 1$ such that $\mathbf{E}[Ye^{\rho Y}] = 0$, while (ii) corresponds to the case when $\mathbf{E}[Ye^Y] = 0$. Finally, (iii) corresponds to the case when $\mathbf{E}[Ye^Y] < 0$ (Kersting and Vatutin; 2017). In this paper, when working with the subcritical regime, we will assume that the process is strongly subcritical and refer to it as subcritical process in the rest of the manuscript.

We now turn to a description of the model. Let $\zeta^U = \{\zeta_n^U\}_{n=1}^\infty$ denote a collection of environmental sequences yielding supercritical offspring pgf $\varphi_{\zeta_n^U}(\cdot)$ and immigration pgf $\psi_{\zeta_n^U}(\cdot)$. Also, let $\zeta^L = \{\zeta_n^L\}_{n=0}^\infty$ denote a collection of environmental sequences yielding subcritical offspring pgf $\varphi_{\zeta_n^L}(\cdot)$. We note that ζ^U and ζ^L are collections of independent random variables and are independent of each other. We now provide an evolutionary description of the process: at time zero the process starts with a random number of ancestors Z_0 . Each of them live one unit of time and reproduce according to a distribution whose pgf is given by $\varphi_{\zeta_0^U}(\cdot)$. Thus, the size of the first generation population is

$$Z_1 = \sum_{i=1}^{Z_0} \xi_{0,i}^U + I_0^U,$$

where, conditionally on the environment, $\xi_{0,i}^U$ are i.i.d. random variables with offspring pgf $\varphi_{\zeta_0^U}(\cdot)$ and independent of the immigration random variable I_0^U with pgf $\psi_{\zeta_0^U}(\cdot)$. The random variable $\xi_{0,i}^U$ is interpreted the number of children produced by the i^{th} parent in the 0^{th} generation and I_0^U is interpreted as the number of immigrants whose pgf is generated by the same environmental random variable ζ_0^U .

Let U_1 denote the random variable representing the upper threshold. If $Z_1 < U_1$, each of the first generation population live one unit of time and evolve, conditionally on the environment, as the ancestors independent of the population size at time one. That is,

$$Z_2 = \sum_{i=1}^{Z_1} \xi_{1,i}^U + I_1^U.$$

As before, conditionally on the environment, $\xi_{1,i}^U$ are i.i.d. with pgf $\varphi_{\zeta_1^U}(\cdot)$ and I_1^U has a pgf $\psi_{\zeta_1^U}(\cdot)$. The random variables $\xi_{1,i}^U$ are independent of Z_1 , $\xi_{0,i}^U$, and I_0^U, I_1^U . If $Z_1 \geq U_1$, then

$$Z_2 = \sum_{i=1}^{Z_1} \xi_{1,i}^L,$$

where $\xi_{1,i}^L$ are i.i.d. with pgf $\varphi_{\zeta_1^L}(\cdot)$. Thus, the size of the second generation population is

$$Z_2 = \begin{cases} \sum_{i=1}^{Z_1} \xi_{1,i}^U + I_1^U & \text{if } Z_1 < U_1, \\ \sum_{i=1}^{Z_1} \xi_{1,i}^L & \text{if } Z_1 \geq U_1. \end{cases}$$

The process Z_3 is defined recursively as before. As an example, if $Z_1 < U_1$, $Z_2 < U_1$ or $Z_1 \geq U_1$, $Z_2 \leq L_1$, for a random lower threshold L_1 , then the process will evolve like a supercritical BPPE with offspring pgf $\varphi_{\zeta_2^U}(\cdot)$ and immigration pgf $\psi_{\zeta_2^U}(\cdot)$. Otherwise (that is, $Z_1 < U_1$ and $Z_2 \geq U_1$ or $Z_1 \geq U_1$ and $Z_2 > L_1$), the process will evolve like a subcritical BPPE with offspring pgf $\varphi_{\zeta_2^L}(\cdot)$.

This dynamics continues with different thresholds (U_j, L_j) yielding the process $\{Z_n\}_{n=0}^\infty$ which we refer to as branching process in random environments with thresholds (BPRET). The consecutive set of generations where the reproduction is governed by a supercritical BPRE is referred to as the *supercritical regime*, while the other is referred to as the *subcritical regime*. As we will see below, non-trivial immigration in the supercritical regime is required to obtain alternating periods of increase and decrease.

The model described above is related to size dependent branching processes with a threshold as studied by Klebaner (1993) and more recently by Athreya and Schuh (2016). Specifically, in that model the offspring distribution depends on a *fixed threshold* K and the size of the previous generation. As observed in these papers, these Markov processes either explode to infinity or are absorbed at zero. In our model the thresholds are *random and dynamic* resulting in a non-Markov process; however, the offspring distribution does not depend on the size of the previous generation *as long as they belong to the same regime*. Indeed, when $U_j - 1 = L_j = K$ for all $j \geq 1$, the immigration distribution is degenerate at zero, and the environment is fixed, one obtains as a special case the density dependent branching process (see for example Klebaner (1984, 1993); Jagers and Klebaner (2011); Athreya and Schuh (2016)). Additionally, while the model of Klebaner (1993) uses Galton-Watson process as a building block, our model uses branching processes in i.i.d. environments.

Continuing with our discussion on the literature, Athreya and Schuh (2016) show that in the fixed environment case the special case of size-dependent process with a single threshold becomes extinct with probability one. We show that this is also the case for the BPRE when there is no immigration, and the details are in Theorem 2.1. Similar phenomenon have been observed in slightly different contexts in Jagers and Zuyev (2020, 2021). Incorporation of immigration component ensures that the process is not absorbed at zero and hence may be useful for modelling stable populations at equilibrium as done in deterministic models. For an additional discussion see Section 7.

For the ease of further discussion we introduce few notations. Let $Y^U := \log(m(\zeta_0^U))$, $Y^L := \log(m(\zeta_0^L))$, and $Y_I^U := \log(m_I(\zeta_0^U))$, where $m(\zeta_0^U) := \mathbf{E}[\xi_{0,1}^U | \zeta_0^U] = \varphi'_{\zeta_0^U}(1)$, $m(\zeta_0^L) := \mathbf{E}[\xi_{0,1}^L | \zeta_0^L] = \varphi'_{\zeta_0^L}(1)$, and $m_I(\zeta_0^U) = \mathbf{E}[I_0^U | \zeta_0^U] = \psi'_{\zeta_0^U}(1)$; that is, Y^U , Y^L , and Y_I^U represent the logarithm of the conditional offspring means and immigration mean, respectively. Also, let $\sigma^2(\zeta_0^U) = \mathbf{Var}[\xi_{0,1}^U | \zeta_0^U]$ and $\sigma^2(\zeta_0^L) = \mathbf{Var}[\xi_{0,1}^L | \zeta_0^L]$ denote the variance of the offspring distribution, conditional on the environment ζ_0^U and ζ_0^L , respectively.

From the description, it is clear that the crossing times at the thresholds (U_j, L_j) of Z_n , namely τ_j and ν_j will play a significant role in the analysis. It will turn out that $\{Z_{\tau_j}\}$ and $\{Z_{\nu_j}\}$ will form a time homogeneous Markov chain with state space $S_L := \{i \in \mathbb{N}_0 : 0 \leq i \leq L_U\}$ and $S_U := \{i \in \mathbb{N} : i \geq L_U + 1\}$, respectively, where we take $L_j \leq L_U$ and $U_j \geq L_U + 1$ for all $j \geq 1$. Under additional conditions on the offspring distribution and the environment sequence, the processes $\{Z_{\tau_j}\}$ and $\{Z_{\nu_j}\}$ will be uniformly ergodic. These results are established in Section 3.

The amount of time the process spends in the supercritical and subcritical regimes, beyond its mathematical and scientific interest, will also arise when studying the central limit theorem for the estimates of $\mathbf{E}[m(\zeta_0^U)]$ and $\mathbf{E}[m(\zeta_0^L)]$. Using the uniformly ergodicity alluded to above, we will establish that the time averages of $\tau_j - \nu_{j-1}$ and $\nu_j - \tau_j$ converge to finite positive constants, μ^U

and μ^L . Additionally, we establish a central limit theorem (CLT) related to this convergence under a finite second moment hypothesis after an appropriate centering and scaling; that is,

$$\frac{1}{n} \sum_{j=1}^n (\tau_j - \nu_{j-1}) \sim N(\mu^U, n\sigma^{2,U})$$

and we characterize $\sigma^{2,U}$ in terms of the stationary distribution of the Markov chain. A similar result also holds for $\nu_j - \tau_j$. This, in turn, provides qualitative information regarding the proportion of time the process spends in these regimes. That is, if C_n^U is the amount of time the process spends in the supercritical regime up to time $n - 1$ we show that $n^{-1}C_n^U$ converges to $\mu^U(\mu^U + \mu^L)^{-1}$; a related central limit theorem is also established and in the process we also characterize the limiting variance. Interestingly, we show that the CLT prevails even for the joint distribution of the length of time and the proportion of time the process spends in supercritical and subcritical regimes. These results are described in Sections 4 and 5.

An interesting question concerns the rate of growth of the BPRET in the supercritical and subcritical regimes described by the corresponding expectations, namely $\mathbf{E}[m(\zeta_0^T)]$, $T \in \{L, U\}$. Specifically, we establish that the limiting joint distribution of the estimators is bivariate normal with a diagonal covariance matrix yielding asymptotic independence of the mean estimators derived using data from supercritical and subcritical regimes. In the classical setting of a supercritical BPRE *without immigration* this problem has received some attention (see for instance Dion and Esty (1979)). The problem considered here is different in the following four ways: (i) the population size does not converge to infinity, (ii) the lengths of the regimes are random, (iii) in the supercritical regime the population size may be zero, and (iv) there is an additional immigration term. While (iii) and (iv) can be accounted for in the classical settings as well, their effect on the point estimates is minimized due to the exponential growth of the population size. Here, while the exponential growth is ruled out, perhaps as anticipated, the Markov property of the process at crossing times, namely $\{Z_{\tau_j}\}$ and $\{Z_{\nu_j}\}$ and their associated regeneration times play a central role in the proof. It is important to note that, it is possible that both regimes occur between regeneration times. Hence, also the proportion of time the process spends in the supercritical and subcritical regime plays a vital role in deriving the asymptotic limit distribution. The limiting variance of the estimators depend additionally on μ^U and μ^L , beyond $\mathbf{Var}[m(\zeta_0^U)]$, $\mathbf{Var}[m(\zeta_0^L)]$, $\mathbf{E}[\sigma^2(\zeta_0^U)]$, and $\mathbf{E}[\sigma^2(\zeta_0^L)]$. In the special case of fixed environments, the limit behavior of the estimators takes a different form compared to the traditional results as described for example in Heyde (1971). These results are in Section 6.

Finally, in Appendix B we provide some numerical experiments illustrating the behavior of the model. Specifically, we illustrate the effects of different distributions on the path behavior of the process and describe how they change when the thresholds increase. The experiments also suggests that if different regimes are not taken into account the true growth rate of the virus may be underestimated. We now turn to Section 2 where we develop additional notations and provide a precise statement of the main results.

2 Main results

Branching process in random environments with thresholds (BPRET) is a supercritical BPRE with immigration until it reaches an upper threshold after which it transitions to a subcritical BPRE until it crosses a lower threshold. Beyond this time the process reverts to a supercritical BPRE with immigration and the above cycle continues. Specifically, let $\{(U_j, L_j)\}_{j=1}^\infty$ denote a collection of thresholds (assumed to be i.i.d.). Then, the BPRET evolves like a supercritical BPRE with immigration until it reaches the upper threshold U_1 at which time it becomes a subcritical BPRE. The process remains subcritical until it crosses the threshold L_1 ; after that it evolves again as a supercritical BPRE with immigration, and so on. We now provide a precise description of BPRET.

$\{(U_j, L_j)\}_{j=1}^\infty$ are i.i.d. random vectors with support $S_T^U \times S_T^L$, where $S_T^U := \{i \in \mathbb{N} : i \geq L_U + 1\}$, $S_T^L := \{i \in \mathbb{N} : L_0 \leq i \leq L_U\}$, and $1 \leq L_0 \leq L_U$ are fixed integers defined on the probability space $(\Omega_B, \mathcal{F}_B, P_B)$. Also, let $\zeta^L = \{\zeta_n^L\}_{n=0}^\infty$ and $\zeta^U = \{\zeta_n^U\}_{n=0}^\infty$ be subcritical and supercritical independent environmental sequences that are defined on probability spaces $(\Omega_{E_L}, \mathcal{F}_{E_L}, P_{E_L})$ and $(\Omega_{E_U}, \mathcal{F}_{E_U}, P_{E_U})$ and taking values in abstract spaces E_L and E_U , respectively. Associated with each ζ_n^U and ζ_n^L are three probability generating functions (pgf(s)) $\varphi_{\zeta_n^U}^U(\cdot)$, $\varphi_{\zeta_n^L}^L(\cdot)$, and $\psi_{\zeta_n^U}^U(\cdot)$ given by

$$\varphi_{\zeta_n^U}^U(s) = \sum_{j=0}^{\infty} p_j^U(\zeta_n^U) s^j, \quad \varphi_{\zeta_n^L}^L(s) = \sum_{j=0}^{\infty} p_j^L(\zeta_n^L) s^j \quad \text{and} \quad \psi_{\zeta_n^U}^U(s) = \sum_{j=0}^{\infty} p_{I_n^U, j}^U(\zeta_n^U) s^j.$$

Let $(\Omega_U, \mathcal{F}_U, P_U)$ and $(\Omega_L, \mathcal{F}_L, P_L)$ denote probability spaces corresponding to supercritical BPRE with immigration and subcritical BPRE. Hence, the environment sequence $\zeta^U = \{\zeta_n^U\}_{n=0}^\infty$, the offspring sequence $\{\xi_{n,i}^U\}_{i=1}^\infty$, and the immigration sequence $\{I_n^U\}_{n=0}^\infty$ are random variables on $(\Omega_U, \mathcal{F}_U, P_U)$. Similarly, $\zeta^L = \{\zeta_n^L\}_{n=0}^\infty$ and $\{\xi_{n,i}^L\}_{i=1}^\infty$, $n \geq 0$, are random variables on $(\Omega_L, \mathcal{F}_L, P_L)$. We point out here that the probability spaces $(\Omega_U, \mathcal{F}_U, P_U)$ and $(\Omega_{E_U}, \mathcal{F}_{E_U}, P_{E_U})$ are linked; that is, for all integrable functions $H : \Omega_U \rightarrow \mathbb{R}$

$$\int H(z, \zeta^U) dP_U(z, \zeta^U) = \int \int H(z, \zeta^U) dP_{\zeta^U}(z) dP_{E_U}(\zeta^U).$$

Similar comments also holds with U replaced by L in the above. All the above described random variables are defined on the probability space $(\Omega, \mathcal{F}, P) = (\Omega_B \times \Omega_U \times \Omega_L, \mathcal{F}_B \otimes \mathcal{F}_U \otimes \mathcal{F}_L, P_B \times P_U \times P_L)$. Finally, conditionally on the environment, $\xi_{n,i}^U$ are i.i.d. random variables with offspring pgf $\varphi_{\zeta_n^U}^U(\cdot)$ and are independent of the immigration random variable I_n^U which has pgf $\psi_{\zeta_n^U}^U(\cdot)$. Similarly, given ζ_n^L , $\xi_{n,i}^L$ are i.i.d. random variables with offspring pgf $\varphi_{\zeta_n^L}^L(\cdot)$. Finally, let Z_0 be an independent random variable with support included in $\{i \in \mathbb{N} : 1 \leq i \leq L_U\}$. We emphasize that the thresholds are independent of the environmental sequences, offspring random variables, immigration random variables, and Z_0 .

Throughout the manuscript, we make the following assumptions on the environmental sequences, which rule out degenerate behavior for BPRE (see Assumption R of Kersting and Vatutin (2017)), ensure that $\log m(\zeta_n^T)$ and $m(\zeta_n^T)$, where $T \in \{L, U\}$, have well-defined, finite means, and state that ζ_n^U is a supercritical environment whereas ζ_n^L is a (strongly) subcritical environment. Finally,

immigration is positive with positive probability and has finite expectation, and the upper thresholds U_j have finite expectation.

(H1) For $T \in \{L, U\}$ $\{\zeta_n^T\}_{n=0}^\infty$ are i.i.d. with $0 < m(\zeta_0^T) < \infty$ P_{E_T} a.s. and $p_0^T(\zeta_0^T) < 1$ $P_{E_T} < \text{a.s.}$

(H2) $\mathbf{E}[|\log(m(\zeta_0^L))m(\zeta_0^L)|] < \infty$ and $\mathbf{E}[m(\zeta_0^U)] < \infty$.

(H3) $\mathbf{E}[\log(m(\zeta_0^L))m(\zeta_0^L)] < 0$ and $\mathbf{E}[\log m(\zeta_0^U)] > 0$.

(H4) $P(I_0^U > 0) > 0$ and $\mathbf{E}[m_I(\zeta_0^U)] < \infty$.

(H5) $\{(U_j, L_j)\}_{j=1}^\infty$ have support $S_T^U \times S_T^L$, where $L_0 \leq \mathbf{E}[m(\zeta_0^L)]L_U$ and $\mathbf{E}[U_1] < \infty$.

Assumption (H3) and Jensen's inequality yield that $\mathbf{E}[m(\zeta_0^L)] < 1$ and $\mathbf{E}[m(\zeta_0^U)] > 1$. We are now ready to give a precise definition of the BPRET. Let $\nu_0 := 0$. Starting from Z_0 , the BPRET $\{Z_n\}_{n=0}^\infty$ is defined recursively over $j \geq 0$ as follows.

1j. For $n \geq \nu_j$ and until $Z_n < U_{j+1}$

$$Z_{n+1} = \sum_{i=1}^{Z_n} \xi_{n,i}^U + I_n^U. \quad (1)$$

Next, let $\tau_{j+1} := \inf\{n \geq \nu_j : Z_n \geq U_{j+1}\}$.

2j. For $n \geq \tau_{j+1}$ and until $Z_n > L_{j+1}$

$$Z_{n+1} = \sum_{i=1}^{Z_n} \xi_{n,i}^L. \quad (2)$$

Next, let $\nu_{j+1} := \inf\{n \geq \tau_{j+1} : Z_n \leq L_{j+1}\}$.

We will also use the quantities $\Delta_j^U := \tau_j - \nu_{j-1}$, $\Delta_j^L := \nu_j - \tau_j$, $S_n^U := \sum_{j=1}^n \Delta_j^U$, and $S_n^L := \sum_{j=1}^n \Delta_j^L$. We refer to Δ_j^U and Δ_j^L as the length of the j^{th} supercritical and subcritical regime. It is clear from the above definition that ν_j and τ_{j+1} are stopping times with respect to (w.r.t.) the σ -algebra \mathcal{F}_n generated by $\{Z_j\}_{j=0}^n$ and the thresholds $\{(U_j, L_j)\}_{j=1}^\infty$. Thus, if $\nu_j < \infty$ and $\tau_{j+1} < \infty$ with probability one, then Z_{ν_j} , $Z_{\tau_{j+1}}$, $\xi_{\nu_j,i}^U$, $\xi_{\tau_{j+1},i}^L$, and $I_{\nu_j}^U$ are well-defined random variables. Theorem 1 (ii) of Bansaye (2009) shows that a supercritical BPRT with immigration converges in probability to infinity. It follows that, if ν_j is finite almost surely then τ_{j+1} is finite almost surely. Indeed, it holds that

$$P(\Delta_{j+1}^U = \infty | \nu_j < \infty) \leq \lim_{k \rightarrow \infty} P(\Delta_{j+1}^U \geq k + 1 | \nu_j < \infty) \leq \lim_{k \rightarrow \infty} P(Z_{\nu_j+k} < U_{j+1} | \nu_j < \infty) = 0, \quad (3)$$

where $\{Z_n\}_{n=\nu_j}^\infty$ is a supercritical BPRT with immigration. On the other hand, Theorem 2.1 of Kersting and Vatutin (2017) ensures that $\Delta_{j+1}^L < \infty$ whenever $\tau_{j+1} < \infty$. Using $\nu_0 = 0$ and induction over j we see that for all $j \in \mathbb{N}_0$, Δ_{j+1}^U and Δ_{j+1}^L are finite almost surely and so are τ_{j+1} and ν_{j+1} . In the above discussion it is critical that Assumption (H4) holds true, namely $P(I_0^U > 0) > 0$. Indeed, if $I_0^U \equiv 0$ then the process $\{Z_n\}_{n=0}^\infty$ may fail to cross the thresholds

U_j yielding that $\tau_j = \infty$ since some environments may have zero offspring. In this case, $\{Z_n\}_{n=0}^\infty$ extincts almost surely. On the other hand, Theorem 2.1 of Kersting and Vatutin (2017) ensures that

$$P(\nu_{j+1} < \infty | \tau_{j+1} < \infty) = 1. \quad (4)$$

The above equality holds whenever ζ^L is a subcritical or critical (but not strongly critical) environmental sequence (see Definition 2.3 in Kersting and Vatutin (2017)). This is made precise in the next theorem.

Theorem 2.1. *Assume (H1)-(H2), $\mathbf{E}[\log(m(\zeta_0^L))] \leq 0$, $m(\zeta_0^L) \neq 1$, $\mathbf{E}[\log m(\zeta_0^U)] > 0$, and $I_0^U \equiv 0$. Let $T := \inf\{n \geq 1 : Z_n = 0\}$. Then $P(T < \infty) = 1$.*

Theorem 1 of Athreya and Schuh (2016) follows from the above theorem by taking $L_U = K$, $L_j \equiv K$, $U_j \equiv K + 1$, where K is a finite positive integer, and assuming that the environments are fixed in both regimes. We now turn to transience and recurrence of the BPRET $\{Z_n\}_{n=0}^\infty$. Notice that even though $\{Z_n\}_{n=0}^\infty$ is not Markov the concepts of recurrence and transience can be studied using the definition given below (due to Lamperti (1960, 1963)).

Definition 1. *A non-negative stochastic process $\{X_n\}_{n=0}^\infty$ satisfying $P(\limsup_{n \rightarrow \infty} X_n = \infty) = 1$ is said to be recurrent if there exists an $r < \infty$ such that $P(\liminf_{n \rightarrow \infty} X_n \leq r) = 1$ and transient if $P(\lim_{n \rightarrow \infty} X_n = \infty) = 1$.*

Our next result is concerned with the path behavior of $\{Z_n\}_{n=0}^\infty$ and the stopped sequences $\{Z_{\nu_j}\}_{j=0}^\infty$ and $\{Z_{\tau_j}\}_{j=1}^\infty$.

Theorem 2.2. *Assume (H1)-(H5). Then*

- (i) *the process $\{Z_n\}_{n=0}^\infty$ is recurrent;*
- (ii) *$\{Z_{\nu_j}\}_{j=0}^\infty$ and $\{Z_{\tau_j}\}_{j=1}^\infty$ are time homogeneous Markov chains.*

We now turn to the ergodicity properties of $\{Z_{\nu_j}\}_{j=0}^\infty$ and $\{Z_{\tau_j}\}_{j=1}^\infty$. These rely on conditions on the offspring distribution that ensures the Markov chains $\{Z_{\nu_j}\}_{j=0}^\infty$ and $\{Z_{\tau_j}\}_{j=1}^\infty$ are irreducible and aperiodic. While several sufficient conditions are possible, we provide below some possible conditions. For $r \in \mathbb{N}_0$ and $T \in \{L, U\}$, let $p_r^T(\zeta_0^T) := P(\xi_0(\zeta_0^T) = r | \zeta_0^L)$ and $p_{I,r}^U(\zeta_0^U) := P(I_0^U = r | \zeta_0^U)$.

(H6) $P_{E_L}(p_r^L(\zeta_0^L) > 0 \text{ for } r = 0 \text{ and } r = 1) > 0$.

(H7) $P_{E_U}(p_r^U(\zeta_0^U) > 0 \text{ for all } r \in \mathbb{N}_0 \text{ and } p_{I,0}^U(\zeta_0^U) > 0) > 0$ and $P_{E_U}(p_{I,r}^U(\zeta_0^U) > 0 \text{ for some } r \in \mathbb{N} \setminus S_T^U) > 0$.

(H8) $P_{E_U}(p_0^U(\zeta_0^U) > 0 \text{ and } p_{I,r}^U(\zeta_0^U) > 0 \text{ for all } r \in S_T^U) > 0$.

(H6) requires that on a set of positive P_{E_L} probability an individual produces zero or one offspring while (H7) requires that on a set of positive P_{E_U} probability $p_r^U(\zeta_0^U) > 0$ for all $r \in \mathbb{N}_0$ and $p_{I,0}^U(\zeta_0^U) > 0$. A similar interpretation holds for (H8). These are weak conditions on the environment sequences and are part of standard BPRE literature. Recall that $S_T^U = \{i \in \mathbb{N} : i \geq L_U + 1\}$.

Theorem 2.3. *Assume (H1)-(H5). (i) If (H6) also holds, then $\{Z_{\nu_j}\}_{j=0}^{\infty}$ is a uniformly ergodic Markov chain with state space $S_L = \mathbb{N}_0 \setminus S_T^U$. (ii) If (H7) (or (H8)) holds, then $\{Z_{\tau_j}\}_{j=1}^{\infty}$ is a uniformly ergodic Markov chain with state space $S_U = S_T^U$.*

When the assumptions (H1)-(H7) (or (H8)) hold, we denote by $\pi^L = \{\pi_i^L\}_{i \in S_L}$ and $\pi^U = \{\pi_i^U\}_{i \in S_U}$ the stationary distributions of the ergodic Markov chains $\{Z_{\nu_j}\}_{j=0}^{\infty}$ and $\{Z_{\tau_j}\}_{j=1}^{\infty}$, respectively. While π^L has moments of all orders, we show in Proposition 3.1 below, that π^U has a finite first moment. These distributions will play a significant role when studying the length of supercritical and subcritical regime which we now undertake.

Length of supercritical and subcritical regime: We now turn to the law of large numbers and central limit theorem for the differences Δ_j^U and Δ_j^L . To this end, we denote by $\mu^U := \mathbf{E}[\Delta_1^U | Z_{\nu_0} \sim \pi^L]$ the expected value of Δ_1^U when Z_{ν_0} has the stationary distribution π^L (and hence Δ_j^U when $Z_{\nu_{j-1}} \sim \pi^L$ for all $j \in \mathbb{N}$). Similarly, let $\mu^L := \mathbf{E}[\Delta_1^L | Z_{\tau_1} \sim \pi^U]$ be the expected value of Δ_1^L when Z_{τ_1} has the stationary distribution π^U . We also define

$$\sigma^{2,U} := \mathbf{Var}[\Delta_1^U | Z_{\nu_0} \sim \pi^L] + 2 \sum_{j=1}^{\infty} \mathbf{Cov}[\Delta_1^U, \Delta_{j+1}^U | Z_{\nu_0} \sim \pi^L] \text{ and} \quad (5)$$

$$\sigma^{2,L} := \mathbf{Var}[\Delta_1^L | Z_{\tau_1} \sim \pi^U] + 2 \sum_{j=1}^{\infty} \mathbf{Cov}[\Delta_1^L, \Delta_{j+1}^L | Z_{\tau_1} \sim \pi^U]. \quad (6)$$

In our analysis we will encounter the random variables $\Delta_j^U + \Delta_j^L$ and $\Delta_j^L + \Delta_{j+1}^U$ for $j \geq 1$, their variances $\tilde{\sigma}^{2,U}$ and $\tilde{\sigma}^{2,L}$, and their covariances \mathfrak{C}^U and \mathfrak{C}^L , where

$$\begin{aligned} \tilde{\sigma}^{2,U} &:= \mathbf{Var}[\Delta_1^U + \Delta_1^L | Z_{\nu_0} \sim \pi^L] + 2 \sum_{j=1}^{\infty} \mathbf{Cov}[\Delta_1^U + \Delta_1^L, \Delta_{j+1}^U + \Delta_{j+1}^L | Z_{\nu_0} \sim \pi^L], \\ \tilde{\sigma}^{2,L} &:= \mathbf{Var}[\Delta_1^L + \Delta_2^U | Z_{\tau_1} \sim \pi^U] + 2 \sum_{j=1}^{\infty} \mathbf{Cov}[\Delta_1^L + \Delta_2^U, \Delta_{j+1}^L + \Delta_{j+2}^U | Z_{\tau_1} \sim \pi^U], \\ \mathfrak{C}^U &:= \sum_{j=0}^{\infty} \mathbf{Cov}[\Delta_1^U, \Delta_{j+1}^U + \Delta_{j+1}^L | Z_{\nu_0} \sim \pi^L] + \sum_{j=1}^{\infty} \mathbf{Cov}[\Delta_1^U + \Delta_1^L, \Delta_{j+1}^U | Z_{\nu_0} \sim \pi^L], \text{ and} \\ \mathfrak{C}^L &:= \sum_{j=0}^{\infty} \mathbf{Cov}[\Delta_1^L, \Delta_{j+1}^L + \Delta_{j+2}^U | Z_{\tau_1} \sim \pi^U] + \sum_{j=1}^{\infty} \mathbf{Cov}[\Delta_1^L + \Delta_2^U, \Delta_{j+1}^L | Z_{\tau_1} \sim \pi^U]. \end{aligned}$$

In the supercritical regime, we impose an additional assumption (H9) below, so as to not qualify our statements with the phrase ‘‘on the set of non-extinction’’. Assumption (H10) below ensures that the immigration distribution stochastically dominates the upper threshold.

(H9) $p_0^U(\zeta_0^U) = 0$ a.s. $[P_{E_U}]$.

(H10) $\mathbf{E}[\frac{U_1}{P(U_0^U \geq U_1 | U_1)}] < \infty$.

We are now ready to study of the finiteness of the means and variances given above.

Proposition 2.1. *Assume (H1)-(H5). (i) Also, if (H6) and (H9) hold, then $\mu^U < \infty$ and $0 < \sigma^{2,U} < \infty$. Next, (ii) if (H7) (or (H8)) and (H10) hold, then $\mu^L < \infty$ and $0 < \sigma^{2,L} < \infty$. (iii) Additionally, under the assumptions in (i) and (ii) $0 < \tilde{\sigma}^{2,U}, \tilde{\sigma}^{2,L} < \infty$.*

It is easy to see that Proposition 2.1 implies that $|\mathfrak{C}^U|$ and $|\mathfrak{C}^L|$ are finite. We now state the main result of this section.

Theorem 2.4. *Assume (H1)-(H5). (i) If (H6) and (H9) hold, then,*

$$\lim_{n \rightarrow \infty} \frac{1}{n} S_n^U = \mu^U \text{ a.s., and } \frac{1}{\sqrt{n}} (S_n^U - n\mu^U) \xrightarrow[n \rightarrow \infty]{d} N(0, \sigma^{2,U}).$$

(ii) *If (H7) (or (H8)) and (H10) hold, then,*

$$\lim_{n \rightarrow \infty} \frac{1}{n} S_n^L = \mu^L \text{ a.s., and } \frac{1}{\sqrt{n}} (S_n^L - n\mu^L) \xrightarrow[n \rightarrow \infty]{d} N(0, \sigma^{2,L}).$$

Proportion of times spent in supercritical and subcritical regime: We now consider the proportion of time the process spends in the subcritical and supercritical regimes. To this end for $n \geq 0$, let $\chi_n^U := \mathbf{I}_{\cup_{j=1}^{\infty} [\nu_{j-1}, \tau_j)}(n)$ be the indicator function assuming value 1 if at time n the process is in the supercritical regime and 0 otherwise. Similarly, let $\chi_n^L = 1 - \chi_n^U := \mathbf{I}_{\cup_{j=1}^{\infty} [\tau_j, \nu_j)}(n)$ take value 1 if at time n the process is in the subcritical regime and 0 otherwise. Furthermore, let $C_n^U := \sum_{j=1}^n \chi_{j-1}^U$ and $C_n^L = n - C_n^U := \sum_{j=1}^n \chi_{j-1}^L$ be the total time that the process spends in the supercritical and subcritical regime, respectively, up to time $n - 1$. Let

$$\theta_n^U := \frac{C_n^U}{n} \text{ and } \theta_n^L := \frac{C_n^L}{n}$$

denote the proportion of time the process spends in the supercritical and subcritical regimes. Our main result in this section is concerned with the central limit theorem for θ_n^U and θ_n^L . To this end, let

$$\theta^U := \frac{\mu^U}{\mu^U + \mu^L} \text{ and } \theta^L := \frac{\mu^L}{\mu^U + \mu^L},$$

while $\theta^{k,U}$ and $\theta^{k,L}$ denote the k^{th} power of θ^U and θ^L , respectively.

Theorem 2.5. *Assume (H1)-(H7) (or (H8)) and (H9)-(H10). Then, for $T \in \{L, U\}$, θ_n^T converges almost surely to θ^T . Furthermore,*

$$\sqrt{n}(\theta_n^T - \theta^T) \xrightarrow[n \rightarrow \infty]{d} N(0, \eta^{2,T}), \text{ where } \eta^{2,T} := \frac{1}{\mu^T} (\sigma^{2,T} \theta^T - 2\mathfrak{C}^T \theta^{2,T} + \tilde{\sigma}^{2,T} \theta^{3,T}).$$

We use these results to now describe the growth rate of the process, as defined by their expectations, in the supercritical and subcritical regime; that is, $m_o^T := \mathbf{E}[m(\zeta_0^T)]$, $T \in \{L, U\}$.

Offspring mean estimation: We begin by noticing that $Z_{\tau_j} \geq L_U + 1$ and $Z_{\tau_j+1}, \dots, Z_{\nu_j-1} \geq L_0$ are positive for all $j \in \mathbb{N}$. However, there may be instances where $Z_{\nu_j}, \dots, Z_{\nu_j+1}, \dots, Z_{\tau_1-1}$ could be zero. To avoid division by zero in (7) below, we let $\tilde{\chi}_n^U := \chi_n^U \mathbf{I}_{[Z_n \geq 1]}$, $\tilde{C}_n^U := \sum_{j=1}^n \tilde{\chi}_{j-1}^U$, and use

the convention that $0/0 = 0 \cdot \infty = 0$. In the subcritical regime, set $\tilde{\chi}_n^L := \chi_n^L$ and $\tilde{C}_n^L := C_n^L$. The generalized method of moments estimator of m_o^T for $T \in \{L, U\}$ is given by

$$\hat{m}_n^U := \frac{1}{\tilde{C}_n^U} \sum_{j=1}^n \frac{Z_j - I_{j-1}^U}{Z_{j-1}} \tilde{\chi}_{j-1}^U \text{ and } \hat{m}_n^L := \frac{1}{\tilde{C}_n^L} \sum_{j=1}^n \frac{Z_j}{Z_{j-1}} \tilde{\chi}_{j-1}^L, \quad (7)$$

where the last term is non-trivial whenever $\tilde{C}_n^L \geq 1$, that is $n \geq \tau_1 + 1$. The decomposition

$$\hat{m}_n^T = \mathbf{E}[m(\zeta_0^T)] + \frac{1}{\tilde{C}_n^T} (M_{1,n}^T + M_{2,n}^T)$$

will be used in the proof of the main result of this section and involves the martingale structure of $M_{i,n}^T := \sum_{j=1}^n D_{i,j}^T$ where

$$D_{1,j}^T := (m(\zeta_{j-1}^T) - \mathbf{E}[m(\zeta_0^T)]) \tilde{\chi}_{j-1}^T, \text{ and } D_{2,j}^T := \frac{\tilde{\chi}_{j-1}^T}{Z_{j-1}} \sum_{i=1}^{Z_{j-1}} (\xi_{i,j-1}^T - m(\zeta_{j-1}^T))$$

with respect to the σ -fields generated by the process $\{Z_j\}_{j=0}^{n-1}$, the random environments $\{\zeta_j^T\}_{j=0}^{n-1}$, the thresholds $\{(U_j, L_j)_{j=1}^\infty\}$, and the offspring distributions $\{\xi_{i,j}^T\}_{i=0}^\infty$, $j = 0, 1, \dots, n-1$. For $T \in \{L, U\}$ and $s \geq 0$, let $\mathbf{m}_1^s(\zeta_0^T) := |m(\zeta_0^T) - \mathbf{E}[m(\zeta_0^T)]|^s$ and $\mathbf{m}_2^s(\zeta_0^T) := \mathbf{E}[|\xi_{i,0}(\zeta_0^T) - m(\zeta_0^T)|^s | \zeta_0^T]$. Next, let $\mathbf{m}_o := (m_o^U, m_o^L)^\top$ and Σ_o be the 2×2 diagonal matrix with elements

$$\mathbf{Var}[m(\zeta_0^U)] \frac{\mu^U + \mu^L}{\tilde{\mu}^U} + \mathbf{E}[\sigma^2(\zeta_0^U)] \frac{\tilde{a}^U (\mu^U + \mu^L)}{(\tilde{\mu}^U)^2} \text{ and}$$

$$\mathbf{Var}[m(\zeta_0^L)] \frac{\mu^U + \mu^L}{\mu^L} + \mathbf{E}[\sigma^2(\zeta_0^L)] \frac{\tilde{a}^L (\mu^U + \mu^L)}{(\mu^L)^2},$$

where $\tilde{\mu}^U := \mathbf{E}[\sum_{k=1}^{\tau_1} \tilde{\chi}_{k-1}^U | Z_{\nu_0} \sim \boldsymbol{\pi}^L]$, $\tilde{\mu}^L := \mu^L$, $\tilde{a}^U := \mathbf{E}[\sum_{k=1}^{\tau_1} \frac{\tilde{\chi}_{k-1}^U}{Z_{k-1}} | Z_{\nu_0} \sim \boldsymbol{\pi}^L]$, and $\tilde{a}^L := \mathbf{E}[\sum_{k=\tau_1+1}^{\nu_1} \frac{\tilde{\chi}_{k-1}^L}{Z_{k-1}} | Z_{\tau_1} \sim \boldsymbol{\pi}^U]$. Obviously, $0 \leq \tilde{\mu}^U \leq \mu^U$.

Theorem 2.6. *Assume (H1)-(H7) (or (H8)). (i) If $\mu^T < \infty$ and for some $s \in (1, 2]$ $\mathbf{E}[\mathbf{m}_i^s(\zeta_0^T)] < \infty$, where $i = 1, 2$ and $T \in \{L, U\}$, then $(\hat{m}_n^U, \hat{m}_n^L)^\top$ is a strongly consistent estimator of \mathbf{m}_o . (ii) If additionally for some $\delta \in (0, 1]$ $\mathbf{E}[\mathbf{m}_i^{2+\delta}(\zeta_0^T)] < \infty$ for $i = 1, 2$ and $T \in \{L, U\}$, then*

$$\sqrt{n}((\hat{m}_n^U, \hat{m}_n^L)^\top - \mathbf{m}_o) \xrightarrow[n \rightarrow \infty]{d} N(\mathbf{0}, \Sigma_o).$$

Remark 2.1. *In the fixed environment case $m(\zeta_0^T) = m_o^T$ and $\sigma^2(\zeta_0^T) =: \sigma_o^{2,T}$ are deterministic constants. Therefore, $\mathbf{Var}[m(\zeta_0^T)] = 0$ and Σ_o has diagonal elements $\sigma_o^{2,T} \tilde{a}^T (\mu^U + \mu^L) / (\tilde{\mu}^T)^2$, where $T \in \{L, U\}$. In this case,*

$$\sqrt{n}((\hat{m}_n^U, \hat{m}_n^L)_{GW}^\top - \mathbf{m}_o) \xrightarrow[n \rightarrow \infty]{d} N(\mathbf{0}, \Sigma_o),$$

where $(\hat{m}_n^U, \hat{m}_n^L)_{GW}^\top$ is given in (7).

3 Path properties of BPRET

In this section we provide the proofs of Theorems 2.1, 2.2, and 2.3 along with the required probability estimates. We begin with a preparatory lemma regarding time homogeneity of $\{Z_{\nu_j}\}_{j=0}^{\infty}$ and $\{Z_{\tau_j}\}_{j=1}^{\infty}$. Expectedly, this property does not depend on the process being strongly subcritical. When $\tau_{j+1} = \infty$ or $\nu_j = \infty$, then $Z_{\tau_{j+1}} := \infty$ and $Z_{\nu_j} := \infty$ and hence $P(Z_{\tau_{j+1}} = k | \tau_{j+1} = \infty, Z_{\nu_j} = i, \nu_j < \infty) = 0$.

Lemma 3.1. *Assume **(H1)**-**(H2)**, $\mathbf{E}[\log(m(\zeta_0^L))] \leq 0$, $m(\zeta_0^L) \neq 1$, and $\mathbf{E}[\log m(\zeta_0^U)] > 0$. Then, for all $i \in \mathbb{N}_0 \setminus S_T^U$, $k \in \setminus S_T^U$, and $j \geq 0$ (i) $P(Z_{\tau_{j+1}} = k | Z_{\nu_j} = i, \nu_j < \infty) = P(Z_{\tau_1} = k | Z_{\nu_0} = i)$ and $P(Z_{\tau_{j+1}} = k | \tau_{j+1} < \infty, Z_{\nu_j} = i) = P(Z_{\tau_1} = k | \tau_1 < \infty, Z_{\nu_0} = i)$, (ii) $P(Z_{\nu_{j+1}} = i | Z_{\tau_{j+1}} = k, \tau_{j+1} < \infty) = P(Z_{\nu_1} = i | Z_{\tau_1} = k, \tau_1 < \infty)$ and $P(Z_{\nu_{j+1}} = i | \nu_{j+1} < \infty, Z_{\tau_{j+1}} = k) = P(Z_{\nu_1} = i | \nu_1 < \infty, Z_{\tau_1} = k)$. If additionally **(H4)** holds, then (iii) τ_j and ν_j are finite almost surely, $P(Z_{\tau_{j+1}} = k | Z_{\nu_j} = i) = P(Z_{\tau_1} = k | Z_{\nu_0} = i)$ and $P(Z_{\nu_{j+1}} = i | Z_{\tau_{j+1}} = k) = P(Z_{\nu_1} = i | Z_{\tau_1} = k)$.*

Proof of Lemma 3.1. We only prove (i) and (iii). Since $P(Z_{\tau_{j+1}} = k | Z_{\nu_j} = i, \nu_j < \infty)$ is equal to

$$\sum_{s=1}^{\infty} \sum_{u=L_U+1}^{\infty} P(Z_{\nu_{j+s}} = k, Z_{\nu_{j+s-1}} < u, \dots, Z_{\nu_{j+1}} < u | Z_{\nu_j} = i, \nu_j < \infty),$$

it is enough to show that for all $s \geq 1$ and $u \geq L_U + 1$

$$\begin{aligned} & P(Z_{\nu_{j+s}} = k, Z_{\nu_{j+s-1}} < u, \dots, Z_{\nu_{j+1}} < u | Z_{\nu_j} = i, \nu_j < \infty) \\ & = P(Z_s = k, Z_{s-1} < u, \dots, Z_1 < u | Z_0 = i). \end{aligned} \tag{8}$$

To this end, we condition on $\zeta_{\nu_{j+l}}^U = e_l^U$ and $\zeta_l^U = e_l^U$, where $l = 0, 1, \dots, s-1$ and $e_l^U \in E_L$. Since, conditionally on $\zeta_{\nu_{j+l}}^U = e_l^U$ and $\zeta_l^U = e_l^U$, both the sequences $\{\zeta_{\nu_{j+l}, i}^U\}_{i=1}^{\infty}$, $\{\zeta_{l, i}^U\}_{i=1}^{\infty}$ and the random variables $I_{\nu_{j+l}}^U, I_l^U$ are i.i.d., we obtain from (1) that

$$\begin{aligned} & P(Z_{\nu_{j+s}} = k, Z_{\nu_{j+s-1}} < u, \dots, Z_{\nu_{j+1}} < u | Z_{\nu_j} = i, \nu_j < \infty, \zeta_{\nu_{j+l}}^U = e_l^U, l = 0, 1, \dots, s-1) \\ & = P(Z_s = k, Z_{s-1} < u, \dots, Z_1 < u | Z_0 = i, \zeta_l^U = e_l^U, l = 0, 1, \dots, s-1). \end{aligned}$$

By taking expectation w.r.t. $\zeta^U = \{\zeta_n^U\}_{n=0}^{\infty}$ and using that ζ_n^U are i.i.d., we obtain (8). Next, we notice that

$$P(Z_{\tau_{j+1}} = k | \tau_{j+1} < \infty, Z_{\nu_j} = i) = \frac{P(Z_{\tau_{j+1}} = k | Z_{\nu_j} = i, \nu_j < \infty)}{P(\tau_{j+1} < \infty | Z_{\nu_j} = i, \nu_j < \infty)}, \text{ where}$$

$$P(\tau_{j+1} < \infty | Z_{\nu_j} = i, \nu_j < \infty) = \sum_{k=L_U+1}^{\infty} P(Z_{\tau_{j+1}} = k | Z_{\nu_j} = i, \nu_j < \infty)$$

is positive because $\mathbf{E}[m(\zeta_0^U)] > 1$. It follows from part (i) that $P(Z_{\tau_{j+1}} = k | \tau_{j+1} < \infty, Z_{\nu_j} = i) = P(Z_{\tau_1} = k | \tau_1 < \infty, Z_0 = i)$. Finally, (iii) follows from (i) and (ii) using (3) and (4). \square

3.1 Extinction when immigration is zero

In this subsection, we provide the proof of Theorem 2.1, which is an adaptation of Theorem 1 of Athreya and Schuh (2016) for BPRE. Recall that for this theorem there is no immigration in the supercritical regime and hence the extinction time T is finite with probability one.

Proof of Theorem 2.1. Let for simplicity $\tau_0 := -1$. We partition the sample space as

$$\Omega = \{\cup_{j=0}^{\infty} [\tau_{j+1} = \infty, \tau_j < \infty]\} \cup \{\cap_{j=1}^{\infty} [\tau_j < \infty]\}$$

and show that (i) $[\tau_{j+1} = \infty, \tau_j < \infty] \subset [T < \infty]$ for all $j \in \mathbb{N}_0$ and (ii) $P(\cap_{j=1}^{\infty} [\tau_j < \infty]) = 0$. First, we notice that if $\tau_j < \infty$ then, using Theorem 2.1 of Kersting and Vatutin (2017), $\nu_j < \infty$. Thus,

$$[\tau_{j+1} = \infty, \tau_j < \infty] = [Z_n < U_{j+1} \forall n \geq \nu_j, \tau_j < \infty],$$

where $\{Z_n\}_{n=\nu_j}^{\infty}$ is a supercritical BPRE until U_{j+1} is reached. Since $Z_n < U_{j+1}$ for all $n \geq \nu_j$, (2.6) of Kersting and Vatutin (2017) yields that $\lim_{n \rightarrow \infty} Z_n = 0$ a.s. and $[\tau_{j+1} = \infty, \tau_j < \infty] \subset [T < \infty]$. Turning to (ii), since the events $[\tau_j < \infty]$ are nonincreasing, $P(\cap_{j=1}^{\infty} [\tau_j < \infty]) = \lim_{j \rightarrow \infty} P(\tau_{j+1} < \infty)$ and $P(\tau_{j+1} < \infty) = P(\tau_{j+1} < \infty | \tau_j < \infty) P(\tau_j < \infty)$. Since $\tau_j = \infty$, if $Z_{\nu_{j-1}} = 0$, it follows that

$$\begin{aligned} P(\tau_{j+1} < \infty | \tau_j < \infty) &\leq P(\tau_{j+1} < \infty | \tau_j < \infty, Z_{\nu_{j-1}} \in [1, L_U]) \\ &\leq \max_{i \in [1, L_U]} P(\tau_{j+1} < \infty | \tau_j < \infty, Z_{\nu_{j-1}} = i) \\ &\leq 1 - \min_{i \in [1, L_U]} P(Z_{\tau_j+1} = 0 | \tau_j < \infty, Z_{\nu_{j-1}} = i). \end{aligned}$$

Lemma 3.1 yields that for all $k \in S_T^U$

$$P(Z_{\tau_j} = k | \tau_j < \infty, Z_{\nu_{j-1}} = i) = P(Z_{\tau_1} = k | \tau_1 < \infty, Z_{\nu_0} = i). \quad (9)$$

Also, for all $j \geq 1$

$$\begin{aligned} P(Z_{\tau_j+1} = 0 | \tau_j < \infty, Z_{\tau_j} = k) &= P(\cap_{i=1}^k [\xi_{\tau_j, i}^L = 0] | \tau_j < \infty) \\ &= \sum_{n=1}^{\infty} P(\cap_{i=1}^k [\xi_{n, i}^L = 0] | \tau_j = n) P(\tau_j = n) = P(\xi_{0,1}^L = 0)^k. \end{aligned}$$

Multiplying by $P(Z_{\tau_j+1} = 0 | \tau_j < \infty, Z_{\tau_j} = k)$ and $P(Z_{\tau_1+1} = 0 | \tau_1 < \infty, Z_{\tau_1} = k)$ and summing over $k \geq L_U + 1$ in (9), we obtain that

$$\begin{aligned} P(Z_{\tau_j+1} = 0 | \tau_j < \infty, Z_{\nu_{j-1}} = i) &= P(Z_{\tau_1+1} = 0 | \tau_1 < \infty, Z_{\nu_0} = i) \\ &= \sum_{k=L_U+1}^{\infty} P(Z_{\tau_1} = k | \tau_1 < \infty, Z_0 = i) P(\xi_{0,1}^L = 0)^k. \end{aligned}$$

Set $\underline{p} := \min_{i \in [1, L_U]} p_i$, where $p_i := P(Z_{\tau_1+1} = 0 | \tau_1 < \infty, Z_{\nu_0} = i)$. Since $P(\xi_{0,1}^L = 0) > 0$ and $P(Z_{\tau_1} = k | \tau_1 < \infty, Z_0 = i) > 0$ for some k , we have that $p_i > 0$ and $\underline{p} > 0$. Hence, $P(\tau_{j+1} < \infty) \leq (1 - \underline{p}) P(\tau_j < \infty)$. Iterating the above argument it follows that $P(\tau_{j+1} < \infty) \leq (1 - \underline{p})^j P(\tau_1 < \infty)$ yielding $\lim_{j \rightarrow \infty} P(\tau_{j+1} < \infty) = 0$. \square

3.2 Markov property at crossing times

Proof of Theorem 2.2. We begin by proving (i). We first notice that since $\{U_j\}_{j=1}^\infty$ are i.i.d. random variables with unbounded support S_T^U , $\limsup_{j \rightarrow \infty} U_j = \infty$ with probability one. Next, observe that along the subsequence $\{\tau_j\}_{j=1}^\infty$, $Z_{\tau_j} \geq U_j$. Hence, $\limsup_{n \rightarrow \infty} Z_n = \infty$. On the other hand, along the subsequence $\{\nu_j\}_{j=1}^\infty$ $Z_{\nu_j} \leq L_j$. Thus, $0 \leq \liminf_{j \rightarrow \infty} Z_j \leq L_U < \infty$. It follows that $\{Z_n\}_{n=0}^\infty$ is recurrent in the sense of Definition 1. Turning to (ii), we first notice that, since $Z_0 \leq L_U$, $Z_{\nu_j} \leq L_j \leq L_U$ and $Z_{\tau_j} \geq U_j \geq L_U + 1$ for all $j \geq 1$, the state spaces S_L of $\{Z_{\nu_j}\}_{j=0}^\infty$ and S_U of $\{Z_{\tau_j}\}_{j=1}^\infty$ are included in $\mathbb{N}_0 \setminus S_T^U$ and S_T^U , respectively. We now establish the Markov property of $\{Z_{\nu_j}\}_{j=0}^\infty$. For all $j \geq 0$, $k \in S_L$, and $i_0, i_1, \dots, i_j \in S_L$, we consider the probability $P(Z_{\nu_{j+1}} = k | Z_{\nu_0} = i_0, \dots, Z_{\nu_j} = i_j)$. By the law of total expectation this is equal to

$$\mathbf{E}_B[\mathbf{E}_{\zeta^L \times \zeta^U}[P(Z_{\nu_{j+1}} = k | Z_{\nu_0} = i_0, \dots, Z_{\nu_j} = i_j, L_{j+1}, U_{j+1}, \zeta^L, \zeta^U)]],$$

where the expectation \mathbf{E}_B is w.r.t. the thresholds $\{(U_j, L_j)\}_{j=1}^\infty$ and the expectation $\mathbf{E}_{\zeta^L \times \zeta^U}$ is w.r.t. the environmental sequences ζ^L and ζ^U . Now, setting $A_{\nu_j, s} := [Z_{\nu_j+s} \geq u, Z_{\nu_j+s-1} < u, \dots, Z_{\nu_j+1} < u]$, $B_{\nu_j, s, t} := [Z_{\nu_j+t} = k, Z_{\nu_j+t-1} > l, \dots, Z_{\nu_j+s+1} > l]$, we have that

$$\begin{aligned} & P(Z_{\nu_{j+1}} = k | Z_{\nu_0} = i_0, \dots, Z_{\nu_j} = i_j, L_{j+1}, U_{j+1}, \zeta^L, \zeta^U) \\ &= \sum_{s=1}^{\infty} \sum_{t=s+1}^{\infty} P(A_{\nu_j, s} | Z_{\nu_j} = i_j, U_{j+1}, \zeta^U) P(B_{\nu_j, s, t} | A_{\nu_j, s}, L_{j+1}, \zeta^L) \\ &= P(Z_{\nu_{j+1}} = k | Z_{\nu_j} = i_j, L_{j+1}, U_{j+1}, \zeta^L, \zeta^U), \end{aligned}$$

where in the second line we have used that $\{Z_n\}_{n=\nu_j}^\infty$ is a supercritical BPPE with immigration until it crosses the threshold u at time $\tau_{j+1} = \nu_j + s$, and similarly $\{Z_n\}_{n=\tau_{j+1}}^\infty$ is a subcritical BPPE until it crosses the threshold l at time $\nu_{j+1} = \nu_j + t$. Thus,

$$P(Z_{\nu_{j+1}} = k | Z_{\nu_0} = i_0, \dots, Z_{\nu_j} = i_j, L_{j+1}, U_{j+1}, \zeta^L, \zeta^U) = P(Z_{\nu_{j+1}} = k | Z_{\nu_j} = i_j, L_{j+1}, U_{j+1}, \zeta^L, \zeta^U).$$

By taking the expectations \mathbf{E}_B and $\mathbf{E}_{\zeta^L \times \zeta^U}$ on both sides, we obtain that

$$P(Z_{\nu_{j+1}} = k | Z_{\nu_0} = i_0, \dots, Z_{\nu_j} = i_j) = P(Z_{\nu_{j+1}} = k | Z_{\nu_j} = i_j).$$

Turning to the time homogeneity property, we obtain from Lemma 3.1 (iii) that

$$\begin{aligned} P(Z_{\nu_{j+1}} = k | Z_{\nu_j} = i_j) &= \sum_{l=L_U+1}^{\infty} P(Z_{\nu_{j+1}} = k | Z_{\tau_{j+1}} = l) P(Z_{\tau_{j+1}} = l | Z_{\nu_j} = i_j) \\ &= \sum_{l=L_U+1}^{\infty} P(Z_{\nu_1} = k | Z_{\tau_1} = l) P(Z_{\tau_1} = l | Z_{\nu_0} = i_j) \\ &= P(Z_{\nu_1} = k | Z_{\nu_0} = i_j). \end{aligned}$$

The proof for $\{Z_{\tau_j}\}_{j=1}^\infty$ is similar. □

3.3 Uniform ergodicity of $\{Z_{\nu_j}\}_{j=0}^\infty$ and $\{Z_{\tau_j}\}_{j=1}^\infty$

In this subsection we prove Theorem 2.3. The proof relies on the following lemma. We denote by $p_{ik}^L(j) = P(Z_{\nu_j} = k | Z_{\nu_0} = i)$, $i, k \in \mathbb{N}_0 \setminus S_T^U$, and $p_{ik}^U(j) = P(Z_{\tau_{j+1}} = k | Z_{\tau_1} = i)$, $i, k \in S_T^U$, the j -step transition probability of the (time homogeneous) Markov chains $\{Z_{\nu_j}\}_{j=0}^\infty$ and $\{Z_{\tau_j}\}_{j=1}^\infty$. For $j = 1$, we also write $p_{ik}^L = p_{ik}^L(1)$ and $p_{ik}^U = p_{ik}^U(1)$. Finally, let $\mathbf{p}_i^L(j) = \{p_{ik}^L(j)\}_{k \in S_L}$ and $\mathbf{p}_i^U(j) = \{p_{ik}^U(j)\}_{k \in S_U}$ be the j -step transition probability of the Markov chains $\{Z_{\nu_j}\}_{j=0}^\infty$ and $\{Z_{\tau_j}\}_{j=1}^\infty$ from state $i \in S_L$ (resp. $i \in S_U$).

Lemma 3.2. *Assume (H1)-(H5). Then, (i) if (H6) also holds, then $p_{ik}^L \geq \underline{p}^L > 0$ for all $i, k \in \mathbb{N}_0 \setminus S_T^U$. Also, (ii) if (H7) (or (H8)) holds, then $p_{ik}^U \geq \underline{p}_k^U > 0$ for all $i, k \in S_T^U$.*

Proof of Lemma 3.2. The idea of proof is to establish a lower bound on p_{ik}^L and p_{ik}^U using (10) and (11) below, respectively. We begin by proving (i). Using Assumptions (H6), let A be a measurable subset of E_L satisfying $P_{E_L}(A) > 0$ and $p_r^L(\zeta_0^L) > 0$ for all $r = 0, 1$ and $\zeta_0^L \in A$. By the law of total expectation

$$p_{ik}^L = \mathbf{E}_B[\mathbf{E}_{\zeta^L \times \zeta^U}[\mathbf{E}[P(Z_{\nu_1} = k | Z_{\tau_1}, L_1, \zeta^L) | Z_{\nu_0} = i, L_1, \zeta^L]]]. \quad (10)$$

Since $P(Z_{\nu_1} = k | Z_{\tau_1}, L_1, \zeta^L) = 0$ on the event $[L_1 < k]$, it follows that

$$P(Z_{\nu_1} = k | Z_{\tau_1}, L_1, \zeta^L) = P(Z_{\nu_1} = k | Z_{\tau_1}, L_1, \zeta^L) \mathbf{I}_{[L_1 \geq k]}.$$

Now, notice that on the event $[L_1 \geq k]$ the term $P(Z_{\nu_1} = k | Z_{\tau_1}, L_1, \zeta^L)$ is bounded below by the probability of reaching state k from Z_{τ_1} in one step; that is,

$$P(Z_{\nu_1} = k | Z_{\tau_1}, L_1, \zeta^L) \geq P(Z_{\nu_1} = k, \nu_1 = \tau_1 + 1 | Z_{\tau_1}, L_1, \zeta^L) \mathbf{I}_{[L_1 \geq k]}.$$

The RHS of the above inequality is bounded below by the probability that the first k individuals have exactly one offspring and the remaining $Z_{\tau_1} - k$ have no offspring, that is,

$$\mathbf{I}_{[L_1 \geq k]} \prod_{r=1}^k P(\xi_{\tau_1, r}^L = 1 | \zeta_{\tau_1}^L) \prod_{r=k+1}^{Z_{\tau_1}} P(\xi_{\tau_1, r}^L = 0 | \zeta_{\tau_1}^L).$$

Once again, using that conditional on the environment $\zeta_{\tau_1}^L, \xi_{\tau_1, r}^L$ are i.i.d., this is equal to

$$\mathbf{I}_{[L_1 \geq k]} (p_1^L(\zeta_{\tau_1}^L))^k (p_0^L(\zeta_{\tau_1}^L))^{Z_{\tau_1} - k}.$$

Since $\mathbf{I}_{[\zeta_{\tau_1}^L \in A]} \leq 1$ and $[L_1 \geq k] \supset [L_1 = L_U]$ because $k \leq L_U$, the last term is bounded below by

$$\mathbf{I}_{[L_1 = L_U]} (p_1^L(\zeta_{\tau_1}^L))^k (p_0^L(\zeta_{\tau_1}^L))^{Z_{\tau_1} - k} \mathbf{I}_{[\zeta_{\tau_1}^L \in A]}.$$

Finally, again using that ζ_n^L are i.i.d. and taking the expectations $\mathbf{E}_B, \mathbf{E}_{\zeta^L \times \zeta^U}$, and $\mathbf{E}[\cdot | Z_{\nu_0} = i, L_1, \zeta^L]$ as in (10), we obtain that $p_{ik}^L \geq \underline{p}_{ik}^L$, where

$$\underline{p}_{ik}^L := P(L_1 = L_U) \mathbf{E}[(p_1^L(\zeta_0^L))^k (p_0^L(\zeta_0^L))^{Z_{\tau_1} - k} \mathbf{I}_{[\zeta_0^L \in A]} | Z_{\nu_0} = i].$$

Notice that \underline{p}_k^L is positive because $P(L_1 = L_U) > 0$, $P_{E_L}(A) > 0$, and $p_r^L(\zeta_0^L) > 0$ for all $r = 0, 1$ and $\zeta_0^L \in A$. Finally, since $\mathbb{N}_0 \setminus S_T^U$ is finite, $\underline{p}^L := \min_{i,k \in \mathbb{N}_0 \setminus S_T^U} \underline{p}_{ik}^L > 0$.

We now turn to the proof of (ii), which is similar to the proof of (i). Using **(H7)**, let A and B be measurable subsets of E_U satisfying:

- (a) $P_{E_U}(A) > 0$ and $p_{I,0}^U(\zeta_0^U) > 0$, $p_r^U(\zeta_0^U) > 0$ for all $r \in \mathbb{N}_0$ and $\zeta_0^U \in A$; and
- (b) $P_{E_U}(B) > 0$ and $p_{I,s}^U(\zeta_0^U) > 0$ for some $s \in \mathbb{N} \setminus S_T^U$ and $\zeta_0^U \in B$.

Again using the law of total expectation, we obtain

$$p_{ik}^U = \mathbf{E}_B[\mathbf{E}_{\zeta^L \times \zeta^U}[\mathbf{E}[P(Z_{\tau_2} = k | Z_{\nu_1}, U_2, \zeta^U) | Z_{\tau_1} = i, U_2, \zeta^U]]]. \quad (11)$$

Since $P(Z_{\tau_2} = k | Z_{\nu_1}, U_2, \zeta^U) = 0$ on the event $[U_2 > k]$, it follows that

$$P(Z_{\tau_2} = k | Z_{\nu_1}, U_2, \zeta^U) = \sum_{z=0}^{L_U} P(Z_{\tau_2} = k | Z_{\nu_1}, U_2, \zeta^U) \mathbf{I}_{[U_2 \leq k]} \mathbf{I}_{[Z_{\nu_1}=z]}. \quad (12)$$

If $U_2 \leq k$ and $z > 0$, then $P(Z_{\tau_2} = k | Z_{\nu_1}, U_2, \zeta^U)$ is bounded below by the probability that z individuals have a total of exactly k offspring and no immigration occurs; that is,

$$P(Z_{\tau_2} = k | Z_{\nu_1}, U_2, \zeta^U) \mathbf{I}_{[U_2 \leq k]} \mathbf{I}_{[Z_{\nu_1}=z]} \geq P\left(\sum_{r=1}^z \xi_{\nu_1,r}^U = k, I_{\nu_1}^U = 0 | Z_{\nu_1}, U_2, \zeta^U\right) \mathbf{I}_{[U_2 \leq k]} \mathbf{I}_{[Z_{\nu_1}=z]}.$$

The RHS of the above inequality is bounded below by the probability that the first $z_1 := (k_1 + 1)z - k$ individuals have $k_1 := \lfloor \frac{k}{z} \rfloor$ offspring and the last $z_2 := z - z_1$ individuals have $k_2 := k_1 + 1$ offspring (indeed $k_1 z_1 + k_2 z_2 = k$) and no immigration occurs; that is,

$$P(\cap_{r=1}^{z_1} [\xi_{\nu_1,r} = k_1], \cap_{r=z_1+1}^{z_2} [\xi_{\nu_1,r} = k_2], I_{\nu_1}^U = 0 | Z_{\nu_1}, U_2, \zeta^U) \mathbf{I}_{[U_2 \leq k]} \mathbf{I}_{[Z_{\nu_1}=z]}.$$

Using that conditional on the environment $\zeta_{\nu_1}^U$, $\xi_{\nu_1,r}^U$ are i.i.d., the above is equal to

$$(p_{k_1}^U(\zeta_{\nu_1}^U))^{z_1} (p_{k_2}^U(\zeta_{\nu_1}^U))^{z_2} p_{I,0}^U(\zeta_{\nu_1}^U) \mathbf{I}_{[U_2 \leq k]} \mathbf{I}_{[Z_{\nu_1}=z]}. \quad (13)$$

Next, if $U_2 \leq k$ and $z = 0$ then $P(Z_{\tau_2} = k | Z_{\nu_1}, U_2, \zeta^U)$ is bounded below by the probability $P(Z_{\tau_2} = k, \tau_2 = \nu_1 + 2 | Z_{\nu_1}, U_2, \zeta^U)$. Now, this probability is bounded below by the probability that there are s immigrants at time $\nu_1 + 1$, these immigrants have a total of exactly k offspring, and no immigration occurs at time $\nu_1 + 2$; that is,

$$P(I_{\nu_1}^U = s, \sum_{r=1}^s \xi_{\nu_1+1,r}^U = k, I_{\nu_1+1}^U = 0 | Z_{\nu_1}, U_2, \zeta^U) \mathbf{I}_{[U_2 \leq k]} \mathbf{I}_{[Z_{\nu_1}=0]}.$$

As before, this last probability is bounded below by probability that $s_1 := (t_1 + 1)s - k$ individuals have $t_1 := \lfloor \frac{k}{s} \rfloor$ offspring and $s_2 := s - s_1$ individuals have $t_2 := t_1 + 1$ offspring. Thus, the above probability is bounded below by

$$p_{I,s}^U(\zeta_{\nu_1}^U) (p_{t_1}^U(\zeta_{\nu_1+1}^U))^{s_1} (p_{t_2}^U(\zeta_{\nu_1+1}^U))^{s_2} p_{I,0}^U(\zeta_{\nu_1+1}^U) \mathbf{I}_{[U_2 \leq k]} \mathbf{I}_{[Z_{\nu_1}=0]}. \quad (14)$$

Combing (12), (13), and (14) and using that $\mathbf{I}_{[\zeta_{\nu_1}^U \in A]}, \mathbf{I}_{[\zeta_{\nu_1+1}^U \in A]}, \mathbf{I}_{[\zeta_{\nu_1}^U \in B]} \leq 1$, we obtain that $P(Z_{\tau_2} = k | Z_{\nu_1}, U_2, \zeta^U)$ is bounded below by

$$\begin{aligned} & \mathbf{I}_{[U_2 \leq k]} \sum_{z=1}^{L_U} \mathbf{I}_{[Z_{\nu_1}=z]} (p_{k_1}^U(\zeta_{\nu_1}^U))^{z_1} (p_{k_2}^U(\zeta_{\nu_1}^U))^{z_2} p_{I,0}^U(\zeta_{\nu_1}^U) \mathbf{I}_{[\zeta_{\nu_1}^U \in A]} \\ & + \mathbf{I}_{[U_2 \leq k]} \mathbf{I}_{[Z_{\nu_1}=0]} p_{I,s}^U(\zeta_{\nu_1}^U) (p_{t_1}^U(\zeta_{\nu_1+1}^U))^{s_1} (p_{t_2}^U(\zeta_{\nu_1+1}^U))^{s_2} p_{I,0}^U(\zeta_{\nu_1+1}^U) \mathbf{I}_{[\zeta_{\nu_1}^U \in B]} \mathbf{I}_{[\zeta_{\nu_1+1}^U \in A]}. \end{aligned}$$

Using that ζ_n^U are i.i.d. and taking the expectations \mathbf{E}_B , $\mathbf{E}_{\zeta^L \times \zeta^U}$, and $\mathbf{E}[\cdot | Z_{\tau_1} = i, U_2, \zeta^U]$ as in (11), we obtain that

$$p_{ik}^U \geq P(U_1 \geq k) \sum_{z=0}^{L_U} H_k(z) P(Z_{\nu_1} = z | Z_{\tau_1} = i),$$

where $H_k : [0, L_U] \rightarrow \mathbb{R}$ is given by

$$H_k(0) = \mathbf{E}[p_{I,s}^U(\zeta_0^U) (p_{t_1}^U(\zeta_1^U))^{s_1} (p_{t_2}^U(\zeta_1^U))^{s_2} p_{I,0}^U(\zeta_1^U) \mathbf{I}_{[\zeta_0^U \in B]} \mathbf{I}_{[\zeta_1^U \in A]}]$$

and for $z \in \mathbb{N} \setminus S_T^U$

$$H_k(z) = \mathbf{E}[(p_{k_1}^U(\zeta_0^U))^{z_1} (p_{k_2}^U(\zeta_0^U))^{z_2} p_{I,0}^U(\zeta_0^U) \mathbf{I}_{[\zeta_0^U \in A]}].$$

Since $\sum_{z=0}^{L_U} P(Z_{\nu_1} = z | Z_{\tau_1} = i) = 1$, we conclude that $p_{ik}^U \geq \underline{p}_k^U$, where

$$\underline{p}_k^U := P(U_1 \geq k) \min_{z \in [0, L_U]} H_k(z) > 0.$$

This concludes the proof of (ii). If instead of **(H7)**, **(H8)** holds the proof is similar by noticing that for all $z \in [0, L_U]$, on the event $[U_1 \leq k] \cap [Z_{\nu_1} = z]$, there's a positive probability that at time $\nu_1 + 1$ there are k immigrants and the z individuals have no offspring. A detailed proof can be obtained in the same manner. \square

Before we turn to the proof of Theorem 2.3 we introduce few notations. Let $T_{i,0}^L := 0$ and $T_{i,l}^L := \inf\{j > T_{i,l-1}^L : Z_{\nu_j} = i\}$, $l \geq 1$, be the random times in which the Markov chain $\{Z_{\nu_j}\}_{j=0}^\infty$ enters state $i \in S_L$ when the initial state is $Z_{\nu_0} = i$. Similarly, we let $T_{i,0}^U := 1$ and $T_{i,l}^U := \inf\{j > T_{i,l-1}^U : Z_{\tau_j} = i\}$, $l \geq 1$, be the random times in which the Markov chain $\{Z_{\tau_j}\}_{j=1}^\infty$ enters state $i \in S_U$ when the initial state is $Z_{\tau_1} = i$. The expected times to visit state k starting from i is denoted by $f_{ik}^L := \mathbf{E}[T_{k,1}^L | Z_{\nu_0} = i]$ and $f_{ik}^U := \mathbf{E}[T_{k,1}^U | Z_{\tau_1} = i]$, respectively.

Proof of Theorem 2.3. Lemma 3.2 implies that the state spaces of $\{Z_{\nu_j}\}_{j=0}^\infty$ and $\{Z_{\tau_j}\}_{j=1}^\infty$ are $S_L = \mathbb{N}_0 \setminus S_T^U$ and $S_U = S_T^U$. Next, to establish ergodicity of these Markov chains, it is sufficient to verify irreducibility, aperiodicity, and positive recurrence. Irreducibility and aperiodicity follow from Lemma 3.2 in both cases. Now, turning to positive recurrence let $I_n^L(k) := \{(i_1, i_2, \dots, i_{n-1}) : i_j \in S_L \setminus \{k\}\}$ for $k \in S_L$. Then, using Markov property and (i) of Lemma 3.2, it follows that

$$\begin{aligned} f_{kk}^L &= \sum_{n=1}^{\infty} P(T_{k,1}^L \geq n | Z_{\nu_0} = k) = \sum_{n=1}^{\infty} P(\cap_{j=1}^{n-1} [Z_{\nu_j} \neq k] | Z_{\nu_0} = k) \\ &= \sum_{n=1}^{\infty} p_{ki_1} \prod_{I_n^L(k)} p_{i_{j-1}i_j} \leq \sum_{n=1}^{\infty} (1 - \underline{p}^L)^{n-1} < \infty. \end{aligned}$$

Now by the finiteness of S_L it follows that $\{Z_{\nu_j}\}_{j=0}^\infty$ is uniformly ergodic. Next, as above for all $k \in S_U$

$$f_{kk}^U = \sum_{n=1}^{\infty} P(T_{k,1}^U \geq n | Z_{\tau_1} = k) \leq 1 + \sum_{n=2}^{\infty} (1 - \underline{p}_k^U)^{n-1} < \infty.$$

To complete the proof of uniform ergodicity of $\{Z_{\tau_j}\}_{j=1}^\infty$, we will verify the Doeblin's condition for one-step transition: that is, for a probability distribution $\mathbf{q} = \{q_k\}_{k \in S_U}$ and every set $A \subset S_U$ satisfying $\sum_{k \in A} q_k > \epsilon$

$$\inf_{l \in S_U} \left(\sum_{k \in A} p_{lk}^U \right) > \delta.$$

Now, taking $q_k := (\underline{p}_k^U) / (\sum_{k \in S} \underline{p}_k^U)$, it follows from Lemma 3.2 (ii) that

$$\inf_{l \in S_U} \left(\sum_{k \in A} p_{lk}^U \right) \geq \left(\sum_{k \in S} \underline{p}_k^U \right) \sum_{k \in A} q_k.$$

Choosing $\delta = (\sum_{k \in S} \underline{p}_k^U) \epsilon$ uniform ergodicity of $\{Z_{\tau_j}\}_{j=1}^\infty$ follows. \square

Remark 3.1. An immediate consequence of the above theorem is that $\{Z_{\nu_j}\}_{j=0}^\infty$ possesses a proper stationary distribution $\boldsymbol{\pi}^L = \{\pi_k^L\}_{k \in S_L}$, where $\pi_k^L := 1/f_{kk}^L > 0$ and satisfies $\pi_k^L = \sum_{i \in S} \pi_i^L p_{ik}^L$ for all $k \in S_L$. Furthermore, $\lim_{j \rightarrow \infty} \sup_{l \in S_L} \|\mathbf{p}_l^L(j) - \boldsymbol{\pi}^L\| = 0$ where $\|\cdot\|$ denotes the total variation norm. Furthermore, under a finite second moment hypothesis the central limit theorem holds for functions of Z_{ν_j} . Similar comment also holds for $\{Z_{\tau_j}\}_{j=1}^\infty$ with L replaced by U .

Remark 3.2. It is worth noticing that the stationary distributions $\boldsymbol{\pi}^L$ and $\boldsymbol{\pi}^U$ are connected using $\pi_i^L = \sum_{l \in S_U} P(Z_{\nu_1} = i | Z_{\tau_1} = l) \pi_l^U$ for all $i \in S_L$ since by time homogeneity (Lemma 3.1) $P(Z_{\nu_{j+1}} = i | Z_{\tau_{j+1}} = l) = P(Z_{\nu_1} = i | Z_{\tau_1} = l)$. Now, taking the limit as $j \rightarrow \infty$ in

$$P(Z_{\nu_{j+1}} = i) = \sum_{l \in S_U} P(Z_{\nu_{j+1}} = i | Z_{\tau_{j+1}} = l) P(Z_{\tau_{j+1}} = l),$$

the above expression follows. Similarly, $\pi_k^U = \sum_{l \in S_L} P(Z_{\tau_1} = k | Z_{\nu_0} = l) \pi_l^L$ for all $k \in S_U$.

Since the state space S_L is finite $\boldsymbol{\pi}^L$ has moments of all orders. Our next proposition shows that $\boldsymbol{\pi}^U$ has a finite first moment.

Proposition 3.1. Under (H1)-(H5), (H7) (or (H8)), and (H10), $\sum_{k \in S_U} k \pi_k^U < \infty$.

Proof of Proposition 3.1. Using that $\boldsymbol{\pi}^U = \{\pi_k^U\}_{k \in S_U}$ is the stationary distribution of the Markov chain $\{Z_{\tau_j}\}_{j=1}^\infty$, we write for all $k \in S_U$

$$\pi_k^U = P(Z_{\tau_2} = k | Z_{\tau_1} \sim \boldsymbol{\pi}^U) = \mathbf{E}_B[P(Z_{\tau_2} = k | U_2, Z_{\tau_1} \sim \boldsymbol{\pi}^U)].$$

Next, we notice that

$$P(Z_{\tau_2} = k | U_2, Z_{\tau_1} \sim \boldsymbol{\pi}^U) = \sum_{n=3}^{\infty} P(Z_{\tau_2} = k | \tau_2 = n, U_2, Z_{\tau_1} \sim \boldsymbol{\pi}^U) P(\tau_2 = n | U_2, Z_{\tau_1} \sim \boldsymbol{\pi}^U).$$

Now, using that the event $[\tau_2 = n]$ is same as $[Z_n \geq U_2] \cap [Z_k < U_2]$ for all $k = \nu_1, \dots, n-1$, the RHS of the above inequality is bounded above by

$$\max_{i=0,1,\dots,U_2-1} \sum_{n=3}^{\infty} P(Z_n = k | Z_n \geq U_2, U_2, Z_{n-1} = i) P(\tau_2 = n | U_2, Z_{\tau_1} \sim \pi^U).$$

Since BPRE is a time-homogeneous Markov chain and $P(Z_n = k | Z_n \geq U_2, U_2, Z_{n-1} = i)$ is calculated in the supercritical regime, it follows that

$$P(Z_n = k | Z_n \geq U_2, U_2, Z_{n-1} = i) = P(Z_1 = k | Z_1 \geq U_2, U_2, Z_0 = i),$$

where we also use the fact that the process starts in the supercritical regime. Now, using U_1 and U_2 are i.i.d., it follows that

$$\pi_k^U \leq \mathbf{E}_B \left[\max_{i=0,1,\dots,U_1-1} P(Z_1 = k | Z_1 \geq U_1, U_1, Z_0 = i) \right].$$

Since

$$P(Z_1 = k | Z_1 \geq U_1, U_1, Z_0 = i) \leq \frac{P(Z_1 = k | Z_0 = i)}{P(Z_1 \geq U_1 | U_1, Z_0 = i)},$$

using Fubini-Tonelli theorem, we obtain that

$$\begin{aligned} \sum_{k \in S_U} k \pi_k^U &\leq \mathbf{E}_B \left[\max_{i=0,1,\dots,U_1-1} \frac{\sum_{k \in S_U} k P(Z_1 = k | Z_0 = i)}{P(Z_1 \geq U_1 | U_1, Z_0 = i)} \right] \\ &\leq \mathbf{E}_B \left[\max_{i=0,1,\dots,U_1-1} \frac{\mathbf{E}[Z_1 | Z_0 = i]}{P(Z_1 \geq U_1 | U_1, Z_0 = i)} \right]. \end{aligned}$$

Now, for all $i = 0, 1, \dots, U_1 - 1$, we have that

$$P(Z_1 \geq U_1 | U_1, Z_0 = i) \geq P(I_0^U \geq U_1 | U_1).$$

Finally, using the Assumptions **(H3)**, **(H4)**, and **(H10)**, we conclude that

$$\begin{aligned} \sum_{k \in S_U} k \pi_k^U &\leq \mathbf{E}_B \left[\frac{(U_1 - 1) \mathbf{E}[m(\zeta_0^U)] + \mathbf{E}[m_I(\zeta_0^U)]}{P(I_0^U \geq U_1 | U_1)} \right] \\ &\leq \mathbf{E}_B \left[\frac{U_1}{P(I_0^U \geq U_1 | U_1)} \right] \max(\mathbf{E}[m(\zeta_0^U)], \mathbf{E}[m_I(\zeta_0^U)]) < \infty. \end{aligned}$$

□

4 Regenerative property of crossing times

In this section, we establish the law of large numbers and central limit theorem for the length of the supercritical and subcritical regimes $\{\Delta_j^U\}_{j=1}^{\infty}$ and $\{\Delta_j^L\}_{j=1}^{\infty}$. To this end, we will show that $\{\Delta_j^U\}_{j=1}^{\infty}$ and $\{\Delta_j^L\}_{j=1}^{\infty}$ are regenerative over the times $\{T_{i,l}^L\}_{l=0}^{\infty}$ and $\{T_{i,l}^U\}_{l=1}^{\infty}$, respectively. For $l \geq 1$ and $i \in S_L$ let

$$B_{i,l}^L := (T_{i,l}^L - T_{i,l-1}^L, \{\Delta_{j+1}^U : j = T_{i,l-1}^L, \dots, T_{i,l}^L - 1\}, \{\Delta_{j+1}^L + \Delta_{j+1}^U : j = T_{i,l-1}^L, \dots, T_{i,l}^L - 1\})$$

and for $l \geq 1$ and $i \in S_U$ let

$$B_{i,l}^U := (T_{i,l}^U - T_{i,l-1}^U, \{\Delta_j^L : j = T_{i,l-1}^U, \dots, T_{i,l}^U - 1\}, \{\Delta_j^L + \Delta_{j+1}^U : j = T_{i,l-1}^U, \dots, T_{i,l}^U - 1\}).$$

Lemma 4.1. *Assume (H1)-(H5). (i) If (H6) also holds and $Z_{\nu_0} = i \in S_L$, then $\{B_{i,l}^L\}_{l=1}^\infty$ are i.i.d. (ii) If (H7) (or (H8)) holds and $Z_{\tau_1} = i \in S_U$, then $\{B_{i,l}^U\}_{l=1}^\infty$ are i.i.d.*

Proof of Lemma 4.1. We begin by proving (i). It is sufficient to show that for all $n \geq 0$, $k \in \mathbb{N}$, $d_1^U, \dots, d_k^U \in \mathbb{N}$, and $d_1^L, \dots, d_k^L \in \mathbb{N}$

$$P(B_{i,n+1}^L = (k, \{d_j^U\}, \{d_j^U + d_j^L\}) | B_{i,n}^L, \dots, B_{i,1}^L, Z_{\nu_0} = i) = P(B_{i,1}^L = (k, \{d_j^U\}, \{d_j^U + d_j^L\}) | Z_{\nu_0} = i). \quad (15)$$

Indeed, (15) implies that for all $n \geq 1$, $k_1, \dots, k_n \in \mathbb{N}$, $d_{1,1}^U, \dots, d_{1,k_1}^U, \dots, d_{n,1}^U, \dots, d_{n,k_n}^U \in \mathbb{N}$, and $d_{1,1}^L, \dots, d_{1,k_1}^L, \dots, d_{n,1}^L, \dots, d_{n,k_n}^L \in \mathbb{N}$

$$\begin{aligned} & P(B_{i,n}^L = (k_n, \{d_{n,j}^U\}, \{d_{n,j}^U + d_{n,j}^L\}), \dots, B_{i,1}^L = (k_1, \{d_{1,j}^U\}, \{d_{1,j}^U + d_{1,j}^L\}) | Z_{\nu_0} = i) \\ &= \prod_{l=1}^n P(B_{i,l}^L = (k_l, \{d_{l,j}^U\}, \{d_{l,j}^U + d_{l,j}^L\}) | Z_{\nu_0} = i). \end{aligned}$$

We now show (15). For simplicity set $X_j := Z_{\nu_j}$. We notice that

$$\begin{aligned} & P(B_{i,n+1}^L = (k, \{d_j^U\}, \{d_j^U + d_j^L\}) | B_{i,n}^L, \dots, B_{i,1}^L, X_0 = i) \\ &= P(B_{i,n+1}^L = (k, \{d_j^U\}, \{d_j^U + d_j^L\}) | X_{T_{i,n}^L} = i, B_{i,n}^L, \dots, B_{i,1}^L, X_0 = i) \\ &= P(T_{i,n+1}^L - T_{i,n}^L = k, \cap_{j=1}^k [\Delta_{T_{i,n}^L+j}^U = d_j^U, \Delta_{T_{i,n}^L+j}^L = d_j^L] | X_{T_{i,n}^L} = i, B_{i,n}^L, \dots, B_{i,1}^L, X_0 = i) \\ &= P(X_{T_{i,n}^L+k} = i, \cap_{j=1}^{k-1} [X_{T_{i,n}^L+j} \neq i], \cap_{j=1}^k [\Delta_{T_{i,n}^L+j}^U = d_j^U, \Delta_{T_{i,n}^L+j}^L = d_j^L] | X_{T_{i,n}^L} = i, B_{i,n}^L, \dots, B_{i,1}^L, X_0 = i) \end{aligned}$$

We now compute the last term of the above equation. Specifically, by proceeding as in the proof of Lemma 3.1 (involving conditioning on the environments), we obtain that for all $n \geq 1$, $k \geq 1$, $x_0, x_1, \dots, x_k \in S_L$, $d_1^U, \dots, d_k^U \in \mathbb{N}$, and $d_1^L, \dots, d_k^L \in \mathbb{N}$

$$\begin{aligned} & P(\cap_{j=1}^k [X_{T_{i,n}^L+j} = x_j, \Delta_{T_{i,n}^L+j}^U = d_j^U, \Delta_{T_{i,n}^L+j}^L = d_j^L] | X_{T_{i,n}^L} = x_0, B_{i,n}^L, \dots, B_{i,1}^L, X_0 = i) \\ &= \prod_{j=1}^k P(X_{T_{i,n}^L+j} = x_j, \Delta_{T_{i,n}^L+j}^U = d_j^U, \Delta_{T_{i,n}^L+j}^L = d_j^L | X_{T_{i,n}^L+j-1} = x_{j-1}) \\ &= \prod_{j=1}^k P(X_j = x_j, \Delta_j^U = d_j^U, \Delta_j^L = d_j^L | X_{j-1} = x_{j-1}) \\ &= P(\cap_{j=1}^k [X_j = x_j, \Delta_j^U = d_j^U, \Delta_j^L = d_j^L] | X_0 = x_0). \end{aligned}$$

Now, by taking $x_0 = i$ in the above and summing over $x_k \in \{i\}, x_{k-1} \in S_L \setminus \{i\}, \dots, x_1 \in S_L \setminus \{i\}$, we obtain that

$$\begin{aligned} & P(X_{T_{i,n}^L+k} = i, \cap_{j=1}^{k-1} [X_{T_{i,n}^L+j} \neq i], \cap_{j=1}^k [\Delta_{T_{i,n}^L+j}^U = d_j^U, \Delta_{T_{i,n}^L+j}^L = d_j^L] | X_{T_{i,n}^L} = i, B_{i,n}^L, \dots, B_{i,1}^L, X_0 = i) \\ &= P(X_k = i, \cap_{j=1}^{k-1} [X_j \neq i], \cap_{j=1}^k [\Delta_j^U = d_j^U, \Delta_j^L = d_j^L] | X_0 = i). \end{aligned}$$

The last term in the above is

$$P(T_{i,1}^L = k, \cap_{j=1}^k [\Delta_j^U = d_j^U, \Delta_j^L = d_j^L] | X_0 = i) = P(B_{i,1}^L = (k, \{d_j^U\}, \{d_j^U + d_j^L\}) | Z_{\nu_0} = i).$$

We thus obtain (15). The proof of (ii) is similar. \square

We now turn to the proof of Theorem 2.4. As a first step we establish the positivity and finiteness of $\sigma^{2,T}$ and $\tilde{\sigma}^{2,T}$, where $T \in \{L, U\}$. The following lemma is used to control the covariance terms in $\sigma^{2,T}$ and $\tilde{\sigma}^{2,T}$. In the following let $0 < C_T < \infty$ and $\rho_T \in (0, 1)$ be constants.

Lemma 4.2. *Assume (H1)-(H5). The following holds:*

(i) *If (H6) holds and $w_i^L \in \mathbb{R}$, $i \in S_L$, then*

$$\sum_{j=1}^{\infty} \left| \sum_{k \in S_L} \sum_{i \in S_L} w_k^L w_i^L \pi_k^L \pi_i^L p_{ki}(j) - \left(\sum_{k \in S_L} w_k^L \pi_k^L \right) \left(\sum_{i \in S_L} w_i^L \pi_i^L \right) \right| \leq 2C_L^{1/2} \frac{\rho_L^{1/2}}{1 - \rho_L^{1/2}} \left(\sum_{k \in S_L} (w_k^L)^2 \pi_k^L \right).$$

(ii) *If (H7) (or (H8)) holds and $w_k^U \in \mathbb{R}$, $k \in S_U$, then*

$$\sum_{j=1}^{\infty} \left| \sum_{k \in S_U} \sum_{i \in S_U} w_k^U w_i^U \pi_k^U \pi_i^U p_{ki}(j) - \left(\sum_{k \in S_U} w_k^U \pi_k^U \right) \left(\sum_{i \in S_U} w_i^U \pi_i^U \right) \right| \leq 2C_U^{1/2} \frac{\rho_U^{1/2}}{1 - \rho_U^{1/2}} \left(\sum_{k \in S_U} (w_k^U)^2 \pi_k^U \right).$$

The proof of the above lemma can be constructed along the lines of Theorem 17.2.3 of Ibragimov and Linnik (1971) with $p = q = 1/2$ and it involves a repeated use of Cauchy-Schwarz inequality and stationarity in Remark 3.1. The details are included in Appendix A. We now turn to the proof of Proposition 2.1.

Proof of Proposition 2.1. We begin by proving (i). For all $i \in S_L$ it holds that

$$\mathbf{E}[\tau_1 | Z_0 = i] = \sum_{n=1}^{\infty} P(\tau_1 \geq n | Z_0 = i) \leq \sum_{n=0}^{\infty} P(Z_n < U_1 | Z_0 = i) \leq \mathbf{E}[U_1] \sum_{n=0}^{\infty} \mathbf{E} \left[\frac{1}{Z_n} | Z_0 = i \right].$$

By Theorem 2.1 of Grama et al. (2017) there exist $\gamma \in (0, 1)$ and a constant $0 < C_i < \infty$ such that

$$\mathbf{E} \left[\frac{1}{Z_n} | Z_0 = i \right] \leq C_i \gamma^n. \quad (16)$$

Now, using S_L is finite and $\mathbf{E}[U_1] < \infty$ it follows that

$$\mu^U = \sum_{i \in S_L} \mathbf{E}[\tau_1 | Z_0 = i] \pi_i^L \leq \sum_{i \in S_L} C_i \pi_i^L \frac{\mathbf{E}[U_1]}{1 - \gamma} \leq (\max_{i \in S_L} C_i) \frac{\mathbf{E}[U_1]}{1 - \gamma} < \infty.$$

Turning to the finiteness of $\sigma^{2,U}$, replacing n by $\lfloor \sqrt{n} \rfloor$ in (16), one obtains that

$$\mathbf{E}[\tau_1^2 | Z_0 = k] = \sum_{n=1}^{\infty} P(\tau_1 \geq \sqrt{n} | Z_0 = k) \leq \sum_{n=1}^{\infty} P(Z_{\lfloor \sqrt{n} \rfloor} < U_1 | Z_0 = k) < \infty.$$

This together with the finiteness of μ^U yields $\mathbf{Var}[\Delta_1^U | Z_0 \sim \boldsymbol{\pi}^L] < \infty$. Turning to the covariance terms in $\sigma^{2,U}$, we apply Lemma 4.2 (i) with $w_i^L = \mathbf{E}[\tau_1 - \mu^U | Z_0 = i]$ and using $\sum_{i \in S_L} w_i^L \pi_i^L = 0$, we obtain that

$$\begin{aligned} \sum_{j=1}^{\infty} |\mathbf{Cov}[\Delta_1^U, \Delta_{j+1}^U | Z_0 \sim \boldsymbol{\pi}^L]| &= \sum_{j=1}^{\infty} \left| \sum_{k \in S_L} \mathbf{E}[\tau_1 - \mu^U | Z_0 = k] \pi_k^L \sum_{i \in S_L} \mathbf{E}[\tau_1 - \mu^U | Z_0 = i] p_{ki}^L(j) \right| \\ &\leq 2C_L^{1/2} \left(\frac{\rho_L^{1/2}}{1 - \rho_L^{1/2}} \right) \left(\sum_{k \in S_L} (\mathbf{E}[\tau_1 - \mu^U | Z_0 = k])^2 \pi_k^L \right) < \infty. \end{aligned}$$

We conclude that $\sigma^{2,U}$ is finite. For $i \in S_U$, it holds that

$$\mathbf{E}[\nu_1 - \tau_1 | Z_{\tau_1} = i] = \sum_{n=0}^{\infty} P(\nu_1 - \tau_1 > n | Z_{\tau_1} = i) \leq i \sum_{n=0}^{\infty} \mathbf{E}[m(\zeta_0^L)]^n = \frac{i}{1 - \mathbf{E}[m(\zeta_0^L)]},$$

where the inequality follows from the upper bound on the extinction time of the process in the subcritical regime. Hence, using Proposition 3.1 it follows that

$$\mu^L \leq \left(\sum_{i \in S_U} i \pi_i^U \right) \frac{1}{(1 - \mathbf{E}[m(\zeta_0^L)])} < \infty.$$

Next, we show that $\sigma^{2,L}$ is finite. As before, we obtain that for all $i \in S_U$

$$\mathbf{E}[(\nu_1 - \tau_1)^2 | Z_{\tau_1} = i] \leq \sum_{n=0}^{\infty} P(\nu_1 - \tau_1 > \lfloor \sqrt{n} \rfloor | Z_{\tau_1} = i) \leq i \sum_{n=0}^{\infty} \mathbf{E}[m(\zeta_0^L)]^{\lfloor \sqrt{n} \rfloor} \quad (17)$$

yielding that

$$\mathbf{E}[(\nu_1 - \tau_1)^2 | Z_{\tau_1} \sim \boldsymbol{\pi}^U] \leq \left(\sum_{i \in S_U} i \pi_i^U \right) \left(\sum_{n=0}^{\infty} \mathbf{E}[m(\zeta_0^L)]^{\lfloor \sqrt{n} \rfloor} \right) < \infty. \quad (18)$$

This and the finiteness of $\mu^L < \infty$ implies that $\mathbf{Var}[\Delta_1^L | Z_{\tau_1} \sim \boldsymbol{\pi}^U] < \infty$. Turning to covariances we apply Lemma 4.2 (ii) with $w_i^U = \mathbf{E}[\nu_1 - \tau_1 - \mu^L | Z_{\tau_1} = i]$ and using $\sum_{i \in S_U} w_i^U \pi_i^U = 0$, we obtain that

$$\sum_{j=1}^{\infty} |\mathbf{Cov}[\Delta_1^L, \Delta_{j+1}^L | Z_{\tau_1} \sim \boldsymbol{\pi}^U]| \leq 2C_U^{1/2} \left(\frac{\rho_U^{1/2}}{1 - \rho_U^{1/2}} \right) \left(\sum_{k \in S_U} (\mathbf{E}[\nu_1 - \tau_1 - \mu^L | Z_{\tau_1} = k])^2 \pi_k^U \right) < \infty$$

yielding the finiteness of $\sigma^{2,L}$. Turning to (iii), we compute

$$\mathbf{Var}[\Delta_1^U + \Delta_1^L | Z_{\nu_0} \sim \boldsymbol{\pi}^L] \leq 2(\mathbf{Var}[\Delta_1^U | Z_{\nu_0} \sim \boldsymbol{\pi}^L] + \mathbf{Var}[\Delta_1^L | Z_{\nu_0} \sim \boldsymbol{\pi}^L]),$$

where using Part (i) $\mathbf{Var}[\Delta_1^U | Z_{\nu_0} \sim \boldsymbol{\pi}^L] < \infty$ and using Remark 3.2

$$\mathbf{Var}[\Delta_1^L | Z_{\nu_0} \sim \boldsymbol{\pi}^L] = \sum_{k \in S_U} \sum_{i \in S_L} \mathbf{Var}[\Delta_1^L | Z_{\tau_1} = k] P(Z_{\tau_1} = k | Z_0 = i) \pi_i^L = \mathbf{Var}[\Delta_1^L | Z_{\tau_1} \sim \boldsymbol{\pi}^U] < \infty.$$

Turning to the covariance, we again apply Lemma 4.2 (i) with $w_i^L = \mathbf{E}[\Delta_1^U + \Delta_1^L - (\mu^U + \mu^L)|Z_0 = i]$ and conclude that also

$$\sum_{j=1}^{\infty} \mathbf{Cov}[\Delta_1^U + \Delta_1^L, \Delta_{j+1}^U + \Delta_{j+1}^L] \leq 2C_L^{1/2} \left(\frac{\rho_L^{1/2}}{1 - \rho_L^{1/2}} \right) \left(\sum_{k \in S_L} (\mathbf{E}[\Delta_1^U + \Delta_1^L - (\mu^U + \mu^L)|Z_0 = k])^2 \pi_k^L \right)$$

and the finiteness of the RHS yields $\tilde{\sigma}^{2,U} < \infty$. The proof of $\tilde{\sigma}^{2,L} < \infty$ is similar.

We finally establish that $\sigma^{2,U}$, $\sigma^{2,L}$, $\tilde{\sigma}^{2,U}$, and $\tilde{\sigma}^{2,L}$ are positive. We first show that, conditionally on $Z_0 = i \in \mathbb{N}_0 \setminus S_T^U$ and $Z_{\tau_1} = k \in S_T^U$, Δ_1^U and Δ_1^L are non-degenerate. To this end, suppose by contradiction that

$$1 = P(\Delta_1^U = \mu^U | Z_0 = i) = \sum_{u=L_U+1}^{\infty} P(Z_{\mu^U} \geq u, Z_{\mu^U-1} < u, \dots, Z_1 < u | Z_0 = i) P(U_1 = u).$$

Since U_1 has support S_T^U , we obtain that $P(Z_{\mu^U} \geq u, Z_{\mu^U-1} < u, \dots, Z_1 < u | Z_0 = i) = 1$ for all $u \in S_T^U$. In particular,

$$\mathbf{E}[Z_{\mu^U-1} | Z_0 = i] < u \leq \mathbf{E}[Z_{\mu^U} | Z_0 = i] = \mathbf{E}[m(\zeta_0^U)] \mathbf{E}[Z_{\mu^U-1} | Z_0 = i] + \mathbf{E}[m_I(\zeta_0^U)].$$

By taking both $u = L_U + 1$ and $u \geq \mathbf{E}[m(\zeta_0^U)](L_U + 1) + \mathbf{E}[m_I(\zeta_0^U)]$ in the above equation we obtain that both $\mathbf{E}[Z_{\mu^U-1} | Z_0 = i] < L_U + 1$ and $\mathbf{E}[Z_{\mu^U-1} | Z_0 = i] \geq L_U + 1$. Similarly, if

$$1 = P(\Delta_1^L = \mu^L | Z_{\tau_1} = k) = \sum_{l=L_0}^{L_U} P(Z_{\mu^L+\tau_1} \leq l, Z_{\mu^L+\tau_1-1} > l, \dots, Z_{\tau_1+1} > l | Z_{\tau_1} = k) P(L_1 = l),$$

then using that L_1 has support $\{i \in \mathbb{N} : L_0 \leq i \leq L_U\}$, we obtain that $P(Z_{\mu^L+\tau_1} \leq l, Z_{\mu^L+\tau_1-1} > l, \dots, Z_{\tau_1+1} > l | Z_{\tau_1} = k) = 1$ for all $L_0 \leq l \leq L_U$ and

$$\mathbf{E}[m(\zeta_0^L)]^{\mu^L} k \leq l < \mathbf{E}[m(\zeta_0^L)]^{\mu^L-1} k.$$

By taking both $l = L_0$ and $l = L_U$ in the above equation we obtain that $L_0 > \mathbf{E}[m(\zeta_0^L)] L_U$, which contradicts **(H5)**. We deduce that $\sum_{j=0}^{T_{i,1}^L-1} (\Delta_{j+1}^U - \mu^U) = \sum_{n=1}^{\infty} \sum_{j=0}^{n-1} (\Delta_{j+1}^U - \mu^U) \mathbf{I}_{[T_{i,1}^L=n]}$ is non-degenerate and, similarly, $\sum_{j=0}^{T_{i,1}^U-1} (\Delta_{j+1}^L - \mu^L)$, $\sum_{j=0}^{T_{i,1}^L-1} (\Delta_{j+1}^U + \Delta_{j+1}^L - (\mu^U + \mu^L))$, and $\sum_{j=0}^{T_{i,1}^L-1} (\Delta_{j+1}^L + \Delta_{j+2}^U - (\mu^L + \mu^U))$ are non-degenerate. Using Lemma 4.3 below, we conclude that

$$\sigma^{2,U} = \pi_i^L \mathbf{Var}[S_{T_{i,1}^L}^U - \mu^U T_{i,1}^L | Z_0 = i] = \pi_i^L \mathbf{E} \left[\left(\sum_{j=0}^{T_{i,1}^L-1} (\Delta_{j+1}^U - \mu^U) \right)^2 | Z_0 = i \right] > 0$$

and similarly $\sigma^{2,L} > 0$, $\tilde{\sigma}^{2,U} > 0$, and $\tilde{\sigma}^{2,L} > 0$. \square

The proof of the following lemma, which is also required in the proof of the Theorem 2.4, is included in Appendix A. We need the following additional notations: $\tilde{S}_n^U := \sum_{j=1}^n (\nu_j - \nu_{j-1})$ and $\tilde{S}_n^L := \sum_{j=1}^n (\tau_{j+1} - \tau_j)$.

Lemma 4.3. *Under the assumptions of Theorem 2.4, for all $i \in S_L$ the following hold:*

- (i) $\mathbf{E}[S_{T_{i,1}^U}^U | Z_{\nu_0} = i] = (\pi_i^L)^{-1} \mu^U$ and $\mathbf{E}[\tilde{S}_{T_{i,1}^U}^U | Z_{\nu_0} = i] = (\pi_i^L)^{-1} (\mu^U + \mu^L)$,
- (ii) $\mathbf{Var}[S_{T_{i,1}^U}^U | Z_{\nu_0} = i] = (\pi_i^L)^{-1} \sigma^{2,U}$ and $\mathbf{Var}[\tilde{S}_{T_{i,1}^U}^U | Z_{\nu_0} = i] = (\pi_i^L)^{-1} \tilde{\sigma}^{2,U}$; and
- (iii) $\mathbf{Cov}[S_{T_{i,1}^U}^U, \tilde{S}_{T_{i,1}^U}^U | Z_{\nu_0} = i] = (\pi_i^L)^{-1} \mathfrak{C}^U$.

The above statements also hold with U replaced by L .

We are now ready to prove Theorem 2.4. The proof relies on decomposing S_n^U and S_n^L into i.i.d. cycles using Lemma 4.1. Specifically, conditionally on $Z_{\nu_0} = i \in S_L$ (resp. $Z_{\tau_1} = i \in S_U$), the random variables $\{S_{T_{i,l}^U}^U - S_{T_{i,l-1}^U}^U\}_{l=1}^\infty$ (resp. $\{S_{T_{i,l-1}^L}^L - S_{T_{i,l-2}^L}^L\}_{l=1}^\infty$) are i.i.d.

Proof of Theorem 2.4. We begin by proving (i). For $i \in S_L$ and $n \in \mathbb{N}$, let $N_i^L(n) := \sum_{l=1}^\infty \mathbf{1}_{[T_{i,l}^L \leq n]}$ be the number of times $T_{i,l}^L$ is in $\{0, 1, \dots, n\}$. Conditionally on $Z_{\nu_0} = i$, notice that $N_i^L(n)$ is a renewal process (recall that $T_{i,0}^L = 0$). Using the decomposition

$$\frac{1}{n} S_n^U = \frac{N_i^L(n)}{n} \left(\frac{1}{N_i^L(n)} \sum_{l=1}^{N_i^L(n)} (S_{T_{i,l}^U}^U - S_{T_{i,l-1}^U}^U) + \frac{1}{N_i^L(n)} (S_n^U - S_{T_{i,N_i^L(n)}^U}^U) \right), \quad (19)$$

$(S_{T_{i,l}^U}^U - S_{T_{i,l-1}^U}^U)$ are i.i.d. and $\lim_{n \rightarrow \infty} N_i^L(n) = \infty$ a.s., we obtain using the law of large numbers for random sums and Lemma 4.3 (i) that

$$\lim_{n \rightarrow \infty} \frac{1}{N_i^L(n)} \sum_{l=1}^{N_i^L(n)} (S_{T_{i,l}^U}^U - S_{T_{i,l-1}^U}^U) = \mathbf{E}[S_{T_{i,1}^U}^U | Z_{\nu_0} = i] = (\pi_i^L)^{-1} \mu^U \text{ a.s.} \quad (20)$$

Also,

$$\limsup_{n \rightarrow \infty} \frac{1}{N_i^L(n)} (S_n^U - S_{T_{i,N_i^L(n)}^U}^U) \leq \lim_{n \rightarrow \infty} \frac{1}{N_i^L(n)} (S_{T_{i,N_i^L(n)+1}^U}^U - S_{T_{i,N_i^L(n)}^U}^U) = 0 \text{ a.s.}$$

Finally, using the key renewal theorem (Corollary 2.11 of Serfozo (2009)) and Remark 3.1

$$\lim_{n \rightarrow \infty} \frac{N_i^L(n)}{n} = \frac{1}{\mathbf{E}[T_{i,1}^L | Z_{\nu_0} = i]} = \pi_i^L \text{ a.s.} \quad (21)$$

Using (20) and (21) in (19), we obtain the SLLN for S_n^U . Turning to the central limit theorem, conditionally on $Z_{\nu_0} = i$, using the decomposition in (19) and centering, we express $\frac{1}{\sqrt{n}} (S_n^U - n\mu^U)$ as

$$\sqrt{\frac{N_i^L(n)}{n}} \left(\frac{1}{\sqrt{N_i^L(n)}} \sum_{l=1}^{N_i^L(n)} (S_{T_{i,l}^U}^U - S_{T_{i,l-1}^U}^U - \mu^U (T_{i,l}^U - T_{i,l-1}^U)) + \frac{1}{\sqrt{N_i^L(n)}} (S_n^U - S_{T_{i,N_i^L(n)}^U}^U - \mu^U (n - T_{i,N_i^L(n)}^U)) \right),$$

where $S_{T_{i,l}^U}^U - S_{T_{i,l-1}^U}^U - \mu^U (T_{i,l}^U - T_{i,l-1}^U)$ are i.i.d. with mean 0 and variance $\mathbf{Var}[S_{T_{i,1}^U}^U - \mu^U T_{i,1}^U | Z_{\nu_0} = i]$, which using Lemma 4.3 is

$$\mathbf{Var}[S_{T_{i,1}^U}^U - \mu^U T_{i,1}^U | Z_{\nu_0} = i] = (\pi_i^L)^{-1} \sigma^{2,U}. \quad (22)$$

Finally, using the central limit theorem for i.i.d. random sums and (21), it follows that

$$\sqrt{\frac{N_i^L(n)}{n}} \left(\frac{1}{\sqrt{N_i^L(n)}} \sum_{l=1}^{N_i^L(n)} (S_{T_{i,l}^L}^U - S_{T_{i,l-1}^L}^U - \mu^U (T_{i,l}^L - T_{i,l-1}^L)) \right) \xrightarrow[n \rightarrow \infty]{d} N(0, \sigma^{2,U}).$$

To complete the proof notice that

$$\begin{aligned} & \left| \frac{1}{\sqrt{N_i^L(n)}} (S_n^U - S_{T_{i,N_i^L(n)}^L}^U - \mu^U (n - T_{i,N_i^L(n)}^L)) \right| \\ & \leq \frac{1}{\sqrt{N_i^L(n)}} (S_{T_{i,N_i^L(n)+1}^L}^U - S_{T_{i,N_i^L(n)}^L}^U) + \mu^U \frac{1}{\sqrt{N_i^L(n)}} (T_{i,N_i^L(n)+1}^L - T_{i,N_i^L(n)}^L), \end{aligned}$$

where the first term converges to zero almost surely and the second term converges to zero in probability. The proof for S_n^L is similar. \square

When studying the proportion of time the process spends in supercritical and subcritical regimes we will need the above theorem with n replaced by a random time $\tilde{N}(n)$. Our next proposition, whose proof is omitted, addresses this issue.

Proposition 4.1. *Theorem 2.4 holds if n is replaced by a random time $\tilde{N}(n)$, where $\lim_{n \rightarrow \infty} \tilde{N}(n) = \infty$ a.s.*

5 Proportion of times spent in supercritical and subcritical regimes

We recall that $\chi_n^U = \mathbf{1}_{\cup_{j=1}^{\infty} [\nu_{j-1}, \tau_j)}(n)$ is 1 if the process is in the supercritical regime and 0 otherwise and similarly $\chi_n^L = 1 - \chi_n^U$. Also $\theta_n^U = \frac{1}{n} C_n^U$ is the proportion of time the process spends in the supercritical regime up to time $n - 1$. θ_n^L is defined similarly. The limit theorems for θ_n^U and θ_n^L will invoke the i.i.d. blocks developed in Section 4. Let $\mathbf{S}_n^U := (S_n^U, \tilde{S}_n^U)^\top$ and $\mathbf{S}_n^L := (S_n^L, \tilde{S}_n^L)^\top$, where $S_n^U = \sum_{j=1}^n (\tau_j - \nu_{j-1})$, $\tilde{S}_n^U = \sum_{j=1}^n (\nu_j - \nu_{j-1})$, $S_n^L = \sum_{j=1}^n (\nu_j - \tau_j)$, and $\tilde{S}_n^L = \sum_{j=1}^n (\tau_{j+1} - \tau_j)$, and

$$\Sigma^U := \begin{pmatrix} \sigma^{2,U} & \mathfrak{C}^U \\ \mathfrak{C}^U & \tilde{\sigma}^{2,U} \end{pmatrix}, \text{ and } \Sigma^L := \begin{pmatrix} \sigma^{2,L} & \mathfrak{C}^L \\ \mathfrak{C}^L & \tilde{\sigma}^{2,L} \end{pmatrix}.$$

We note that while S_n^U represents the length of the first n supercritical regimes \tilde{S}_n^U is the total time taken for the process to complete the first n cycles.

Lemma 5.1. *Under the conditions of Theorem 2.5, $\frac{1}{\sqrt{n}} (\mathbf{S}_n^U - n\boldsymbol{\mu}^U) \xrightarrow[n \rightarrow \infty]{d} N(\mathbf{0}, \Sigma^U)$, and $\frac{1}{\sqrt{n}} (\mathbf{S}_n^L - n\boldsymbol{\mu}^L) \xrightarrow[n \rightarrow \infty]{d} N(\mathbf{0}, \Sigma^L)$.*

Proof of Lemma 5.1. The proof is similar to that of Theorem 2.4. Conditionally on $Z_{\nu_0} = i$, we write $\frac{1}{\sqrt{n}} (\mathbf{S}_n^U - n\boldsymbol{\mu}^U)$ as

$$\sqrt{\frac{N_i^L(n)}{n}} \left(\frac{1}{\sqrt{N_i^L(n)}} \sum_{l=1}^{N_i^L(n)} (S_{T_{i,l}^L}^U - S_{T_{i,l-1}^L}^U - (T_{i,l}^L - T_{i,l-1}^L) \boldsymbol{\mu}^U) + \frac{1}{\sqrt{N_i^L(n)}} (S_n^U - S_{T_{i,N_i^L(n)}^L}^U - (n - T_{i,N_i^L(n)}^L) \boldsymbol{\mu}^U) \right).$$

Now, by Lemma 4.1 and Lemma 4.3, $\mathbf{S}_{T_{i,l}^L}^U - \mathbf{S}_{T_{i,l-1}^L}^U - \boldsymbol{\mu}^U (T_{i,l}^L - T_{i,l-1}^L)$ are i.i.d. with mean $\mathbf{0} = (0, 0)^\top$ and covariance matrix $(\pi_i^L)^{-1} \Sigma^U$. Using the key renewal theorem we conclude that

$$\sqrt{\frac{N_i^L(n)}{n}} \left(\frac{1}{\sqrt{N_i^L(n)}} \sum_{l=1}^{N_i^L(n)} (\mathbf{S}_{T_{i,l}^L}^U - \mathbf{S}_{T_{i,l-1}^L}^U - \boldsymbol{\mu}^U (T_{i,l}^L - T_{i,l-1}^L)) \right) \xrightarrow[n \rightarrow \infty]{d} N(\mathbf{0}, \Sigma^U)$$

and

$$\frac{1}{\sqrt{N_i^L(n)}} |\mathbf{S}_n^U - \mathbf{S}_{T_{i, N_i^L(n)}^L}^U - \boldsymbol{\mu}^U (n - T_{i, N_i^L(n)}^L)| \xrightarrow[n \rightarrow \infty]{p} 0.$$

The proof of (ii) is similar. \square

Remark 5.1. Lemma 5.1 holds also with n replaced by a random time $\tilde{N}(n)$ such that $\lim_{n \rightarrow \infty} \tilde{N}(n) = \infty$ a.s.

Lemma 5.2. Under the conditions of Theorem 2.5,

(i) $\lim_{n \rightarrow \infty} \frac{\tilde{N}^U(n)}{n+1} = \frac{1}{\mu^U + \mu^L}$ and $\lim_{n \rightarrow \infty} \frac{\tilde{N}^L(n)}{n+1} = \frac{1}{\mu^U + \mu^L}$ a.s.

(ii) $\lim_{n \rightarrow \infty} \frac{C_{\tilde{N}^U(n)}^U}{\tilde{N}^U(n)} = \mu^U$ and $\lim_{n \rightarrow \infty} \frac{C_{\tilde{N}^L(n)}^L}{\tilde{N}^L(n)} = \mu^L$ a.s.

Proof Lemma 5.2. We begin by proving (i). We recall that $\tau_0 = -1$ and $\tau_j < \nu_j < \tau_{j+1}$ a.s. for all $j \geq 0$ yielding that

$$\tilde{N}^L(n) \leq \tilde{N}^U(n) \leq \tilde{N}^L(n) + 1.$$

Since τ_j and ν_j are finite almost surely, we obtain that $\lim_{n \rightarrow \infty} \tilde{N}^U(n) = \infty$ and $\lim_{n \rightarrow \infty} \tilde{N}^L(n) = \infty$ a.s. (i) follows if we show that $\lim_{n \rightarrow \infty} \frac{\tilde{N}^L(n)}{n+1} = \frac{1}{\mu^U + \mu^L}$ a.s. To this end, we notice that $\nu_{\tilde{N}^L(n)} \leq n \leq \nu_{\tilde{N}^L(n)+1}$ and for $n \geq \nu_1$

$$\frac{\nu_{\tilde{N}^L(n)}}{\tilde{N}^L(n)} \leq \frac{n}{\tilde{N}^L(n)} \leq \frac{\nu_{\tilde{N}^L(n)+1}}{\tilde{N}^L(n)+1} \frac{\tilde{N}^L(n)+1}{\tilde{N}^L(n)}$$

Clearly, $\lim_{n \rightarrow \infty} \frac{\tilde{N}^L(n)+1}{\tilde{N}^L(n)} = 1$ a.s. Proposition 4.1 with $\tilde{N}(n) = \tilde{N}^U(n)$ yields that

$$\lim_{n \rightarrow \infty} \frac{\nu_{\tilde{N}^L(n)}}{\tilde{N}^L(n)} = \lim_{n \rightarrow \infty} \frac{1}{\tilde{N}^L(n)} \sum_{j=1}^{\tilde{N}^L(n)} (\nu_j - \nu_{j-1}) = \mu^U + \mu^L \text{ a.s.}$$

Thus, we obtain that

$$\lim_{n \rightarrow \infty} \frac{\nu_{\tilde{N}^L(n)}}{\tilde{N}^L(n)} = \mu^U + \mu^L \text{ a.s.,}$$

which yields

$$\lim_{n \rightarrow \infty} \frac{\tilde{N}^L(n)}{n+1} = \frac{1}{\mu^U + \mu^L} \text{ a.s.} \quad (23)$$

Turning to (ii), we notice that

$$C_{n+1}^U = \sum_{j=0}^n \chi_j^U = \sum_{j=1}^{\tilde{N}^U(n)} (\tau_j - \nu_{j-1}) + \sum_{l=\nu_{\tilde{N}^U(n)}}^n 1,$$

where $\sum_{l=r}^n = 0$ for $r > n$. Proposition 4.1 with $\tilde{N}(n) = \tilde{N}^U(n)$ yield that

$$\lim_{n \rightarrow \infty} \frac{1}{\tilde{N}^U(n)} \sum_{j=1}^{\tilde{N}^U(n)} (\tau_j - \nu_{j-1}) = \mu^U \text{ a.s., and}$$

$$\limsup_{n \rightarrow \infty} \frac{1}{\tilde{N}^U(n)} \sum_{l=\nu_{\tilde{N}^U(n)}}^n 1 \leq \lim_{n \rightarrow \infty} \frac{1}{\tilde{N}^U(n)} (\tau_{\tilde{N}^U(n)+1} - \nu_{\tilde{N}^U(n)}) = 0 \text{ a.s.}$$

Thus, we obtain that $\lim_{n \rightarrow \infty} \frac{C_{n+1}^U}{\tilde{N}^U(n)} = \mu^U$ a.s. Similarly, $\lim_{n \rightarrow \infty} \frac{C_{n+1}^L}{\tilde{N}^L(n)} = \mu^L$ a.s. \square

Our next result is concerned with the joint distribution of the last time the process is in a specific regime and the proportion of time the process spends in that regime under the assumptions of Theorem 2.5. Let

$$\tilde{\Sigma}^U := \begin{pmatrix} \sigma^{2,U}(\mu^U + \mu^L) & -\frac{\mathfrak{e}^U}{\mu^U + \mu^L} \\ -\frac{\mathfrak{e}^U}{\mu^U + \mu^L} & \frac{\sigma^{2,U}}{(\mu^U + \mu^L)^3} \end{pmatrix} \text{ and } \tilde{\Sigma}^L := \begin{pmatrix} \sigma^{2,L}(\mu^U + \mu^L) & -\frac{\mathfrak{e}^L}{\mu^U + \mu^L} \\ -\frac{\mathfrak{e}^L}{\mu^U + \mu^L} & \frac{\sigma^{2,L}}{(\mu^U + \mu^L)^3} \end{pmatrix}.$$

Lemma 5.3. *Under the conditions of Theorem 2.5,*

$$\sqrt{n+1} \begin{pmatrix} \frac{C_{n+1}^U}{\tilde{N}^U(n)} - \mu^U \\ \frac{\tilde{N}^U(n)}{n+1} - \frac{1}{\mu^U + \mu^L} \end{pmatrix} \xrightarrow[n \rightarrow \infty]{d} N(\mathbf{0}, \tilde{\Sigma}^U) \text{ and } \sqrt{n+1} \begin{pmatrix} \frac{C_{n+1}^L}{\tilde{N}^L(n)} - \mu^L \\ \frac{\tilde{N}^L(n)}{n+1} - \frac{1}{\mu^U + \mu^L} \end{pmatrix} \xrightarrow[n \rightarrow \infty]{d} N(\mathbf{0}, \tilde{\Sigma}^L).$$

Proof of Lemma 5.3. We only prove the statement for C_{n+1}^U and $\tilde{N}^U(n)$ since the other case is similar. We write

$$\sqrt{\tilde{N}^U(n)} \begin{pmatrix} \frac{C_{n+1}^U}{\tilde{N}^U(n)} - \mu^U \\ \frac{\tilde{N}^U(n)}{n+1} - (\mu^U + \mu^L) \end{pmatrix} = \frac{1}{\sqrt{\tilde{N}^U(n)}} \sum_{j=1}^{\tilde{N}^U(n)} \begin{pmatrix} \tau_j - \nu_{j-1} - \mu^U \\ \nu_j - \nu_{j-1} - (\mu^U + \mu^L) \end{pmatrix} + \frac{1}{\sqrt{\tilde{N}^U(n)}} \sum_{l=\nu_{\tilde{N}^U(n)}}^n \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

and using Lemma 5.1 and Remark 5.1 with $\tilde{N}(n) = \tilde{N}^U(n)$ we obtain that

$$\sqrt{\tilde{N}^U(n)} \begin{pmatrix} \frac{C_{n+1}^U}{\tilde{N}^U(n)} - \mu^U \\ \frac{\tilde{N}^U(n)}{n+1} - (\mu^U + \mu^L) \end{pmatrix} \xrightarrow[n \rightarrow \infty]{d} N(\mathbf{0}, \Sigma^U).$$

Next, we apply the delta method with $g : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by $g(x, y) = (x, 1/y)$ and obtain that

$$\sqrt{\tilde{N}^U(n)} \begin{pmatrix} \frac{C_{n+1}^U}{\tilde{N}^U(n)} - \mu^U \\ \frac{\tilde{N}^U(n)}{n+1} - \frac{1}{\mu^U + \mu^L} \end{pmatrix} \xrightarrow[n \rightarrow \infty]{d} N(\mathbf{0}, \Sigma_2^U),$$

where

$$\Sigma_2^U = J_g(\boldsymbol{\mu}^U) \Sigma^U J_g(\boldsymbol{\mu}^U)^\top = \begin{pmatrix} \sigma^{2,U} & -\frac{\mathfrak{e}^U}{(\mu^U + \mu^L)^2} \\ -\frac{\mathfrak{e}^U}{(\mu^U + \mu^L)^2} & \frac{\sigma^{2,U}}{(\mu^U + \mu^L)^4} \end{pmatrix}$$

and $J_g(\cdot)$ is the Jacobian matrix of $g(\cdot)$. Using Lemma 5.2 (i), we obtain that

$$\sqrt{n+1} \begin{pmatrix} \frac{C_{n+1}^U}{\tilde{N}^U(n)} - \mu^U \\ \frac{\tilde{N}^U(n)}{n+1} - \frac{1}{\mu^U + \mu^L} \end{pmatrix} \xrightarrow[n \rightarrow \infty]{d} N(\mathbf{0}, \tilde{\Sigma}^U).$$

\square

We are now ready to prove Theorem 2.5. Recall that $\theta_n^U = \frac{C_n^U}{n}$, $\theta_n^L = \frac{C_n^L}{n}$, $\theta^U = \frac{\mu^U}{\mu^U + \mu^L}$, and $\theta^L = \frac{\mu^L}{\mu^L + \mu^U}$.

Proof of Theorem 2.5. Almost sure convergence of θ_n^T follows from Lemma 5.2 upon noticing that $\frac{C_{n+1}^T}{n+1} = (\frac{C_{n+1}^T}{\tilde{N}^T(n)}) (\frac{\tilde{N}^T(n)}{n+1})$. The second statement follows from Lemma 5.2 using the decomposition

$$\sqrt{n+1} \left(\frac{C_{n+1}^T}{n+1} - \frac{\mu^T}{\mu^U + \mu^L} \right) = \frac{\tilde{N}^T(n)}{n+1} \cdot \sqrt{n+1} \left(\frac{C_{n+1}^T}{\tilde{N}^T(n)} - \mu^T \right) + \mu^T \cdot \sqrt{n+1} \left(\frac{\tilde{N}^T(n)}{n+1} - \frac{1}{\mu^U + \mu^L} \right).$$

□

Corollary 5.1. *Under the conditions of Theorem 2.4, for $T \in \{L, U\}$*

$$\sqrt{\tilde{N}^T(n)} \left(\frac{C_{n+1}^T}{\tilde{N}^T(n)} - \mu^T \right) \xrightarrow[n \rightarrow \infty]{d} N(0, \sigma^{2,T}).$$

Proof of Corollary 5.1. We only prove the case $T = U$. We write

$$\sqrt{\tilde{N}^U(n)} \left(\frac{C_{n+1}^U}{\tilde{N}^U(n)} - \mu^U \right) = \frac{1}{\sqrt{\tilde{N}^U(n)}} \sum_{j=1}^{\tilde{N}^U(n)} (\tau_j - \nu_{j-1} - \mu^U) + \frac{1}{\sqrt{\tilde{N}^U(n)}} \sum_{l=\nu_{\tilde{N}^U(n)}}^n 1.$$

Taking the limit in the above equation and using Proposition 4.1 the result follows. □

6 Estimating the mean of the offspring distribution

We recall that $\tilde{\chi}_n^U = \chi_n^U \mathbf{I}_{\{Z_n \geq 1\}}$, $\tilde{C}_n^U = \sum_{j=1}^n \tilde{\chi}_{j-1}^U$, and as before in the subcritical regime $\tilde{\chi}_n^L = \chi_n^L$ and $\tilde{C}_n^L = C_n^L$. We also recall the offspring mean estimate of the BPRET $\{Z_n\}_{n=0}^\infty$ in the supercritical and subcritical regimes are given by

$$\hat{m}_n^U := \frac{1}{\tilde{C}_n^U} \sum_{j=1}^n \frac{Z_j - I_{j-1}^U}{Z_{j-1}} \tilde{\chi}_{j-1}^U \quad \text{and} \quad \hat{m}_n^L := \frac{1}{\tilde{C}_n^L} \sum_{j=1}^n \frac{Z_j}{Z_{j-1}} \tilde{\chi}_{j-1}^L.$$

As explained previously, the decomposition

$$\hat{m}_n^T = \mathbf{E}[m(\zeta_0^T)] + \frac{1}{\tilde{C}_n^T} (M_{1,n}^T + M_{2,n}^T), \tag{24}$$

where for $i = 1, 2$ $M_{i,n}^T := \sum_{j=1}^n D_{i,j}^T$,

$$D_{1,j}^T := (m(\zeta_{j-1}^T) - \mathbf{E}[m(\zeta_0^T)]) \tilde{\chi}_{j-1}^T, \quad \text{and} \quad D_{2,j}^T := \frac{\tilde{\chi}_{j-1}^T}{Z_{j-1}} \sum_{i=1}^{Z_{j-1}} (\xi_{j-1,i}^T - m(\zeta_{j-1}^T))$$

will play a central role in the proof.

Martingale structure of $\mathbf{M}_{1,n}^T$: Let \mathcal{G}_n be the σ -algebra generated by the random environments $\{\zeta_j^T\}_{j=0}^n$, $\mathcal{H}_{1,n}$ the σ -algebra generated by \mathcal{F}_n and \mathcal{G}_{n-1} ; and $\mathcal{H}_{2,n}$ the σ -algebra generated by \mathcal{F}_n ,

\mathcal{G}_{n-1} , and the offspring distributions $\{\xi_{j,i}^T\}_{i=0}^\infty$, $j = 0, 1, \dots, n-1$. Hence, Z_n , $\tilde{\chi}_n^T$, and ζ_{n-1}^T are $\mathcal{H}_{1,n}$ -measurable, whereas ζ_n^T is not $\mathcal{H}_{1,n}$ -measurable. We also denote by $\tilde{\mathcal{H}}_{1,n}$ the σ -algebra generated by \mathcal{F}_{n-1} and \mathcal{G}_{n-1} and the σ -algebra $\tilde{\mathcal{H}}_{2,n}$ generated by \mathcal{F}_{n-1} , \mathcal{G}_{n-1} , and $\{\xi_{j,i}^T\}_{i=0}^\infty$, $j = 0, 1, \dots, n-1$. Hence, Z_{n-1} , $\tilde{\chi}_{n-1}^T$, and ζ_{n-1}^T are all $\tilde{\mathcal{H}}_{1,n}$ -measurable but not $\tilde{\mathcal{H}}_{1,n-1}$ -measurable. Finally, we recall that for $s \geq 0$ $\mathbf{m}_1^s(\zeta_j^T) := |m(\zeta_j^T) - \mathbf{E}[m(\zeta_j^T)]|^s$ and $\mathbf{m}_2^s(\zeta_j^T) := \mathbf{E}[|\xi_{j,i}^T - m(\zeta_j^T)|^s | \zeta_j^T]$.

Proposition 6.1. *The following holds:*

(i) For $i = 1, 2$ $\{(M_{1,n}^T, \mathcal{H}_{i,n})\}_{n=1}^\infty$ is a mean zero martingale sequence and for all $s \geq 0$ $\mathbf{E}[|D_{1,j}^T|^s | \mathcal{H}_{i,j-1}] = \mathbf{E}[\mathbf{m}_1^s(\zeta_0^T)] \tilde{\chi}_{j-1}^T$ a.s. In particular,

$$\mathbf{E}[(D_{1,j}^T)^2 | \mathcal{H}_{i,j-1}] = \mathbf{Var}[m(\zeta_0^T)] \tilde{\chi}_{j-1}^T \text{ a.s., and } \mathbf{E}[(M_{1,n}^T)^2] = \mathbf{Var}[m(\zeta_0^T)] \mathbf{E}[\tilde{C}_n^T].$$

(ii) $\{(M_{2,n}^T, \mathcal{H}_{2,n})\}_{n=1}^\infty$ is a mean zero martingale sequence satisfying

$$\mathbf{E}[(D_{2,j}^T)^2 | \mathcal{H}_{2,j-1}] = \mathbf{E}[\sigma^2(\zeta_0^T)] \frac{\tilde{\chi}_{j-1}^T}{Z_{j-1}} \text{ a.s., and } \mathbf{E}[(M_{2,n}^T)^2] = \mathbf{E}[\sigma^2(\zeta_0^T)] \mathbf{E}\left[\sum_{j=1}^n \frac{\tilde{\chi}_{j-1}^T}{Z_{j-1}}\right].$$

Additionally, for all $s \geq 1$ $\mathbf{E}[|D_{2,j}^T|^s | \mathcal{H}_{2,j-1}] \leq \mathbf{E}[\mathbf{m}_2^s(\zeta_0^T)] \tilde{\chi}_{j-1}^T$ a.s.

(iii) For $i = 1, 2$ $\{(\sum_{j=1}^n ((D_{i,j}^T)^2 - \mathbf{E}[(D_{i,j}^T)^2]), \tilde{\mathcal{H}}_{i,n})\}_{n=1}^\infty$ are mean zero martingale sequences and for all $s \geq 1$

$$\mathbf{E}[|(D_{i,j}^T)^2 - \mathbf{E}[(D_{i,j}^T)^2]|^s | \tilde{\mathcal{H}}_{i,j-1}] \leq 2^s \mathbf{E}[\mathbf{m}_i^{2s}(\zeta_0^T)] \mathbf{E}[\tilde{\chi}_{j-1}^T].$$

(iv) For all $T_1, T_2 \in \{L, U\}$ and $i_1, i_2 \in \{1, 2\}$ such that either $T_1 \neq T_2$ or $i_1 \neq i_2$ it holds that $\mathbf{E}[(D_{i_1,j}^{T_1})(D_{i_2,l}^{T_2})] = 0$ for all $j, l = 1, \dots, n$ and $\mathbf{E}[(M_{i_1,n}^{T_1})(M_{i_2,n}^{T_2})] = 0$. In particular,

$\{(\sum_{j=1}^n (D_{i_1,j}^{T_1})(D_{i_2,j}^{T_2}), \tilde{\mathcal{H}}_{2,n})\}_{n=0}^\infty$ is a mean zero martingale sequence and $\mathbf{E}[|(D_{i_1,j}^{T_1})(D_{i_2,j}^{T_2})|^s | \tilde{\mathcal{H}}_{2,j-1}] \leq \mathbf{E}[\mathbf{m}_{i_1}^s(\zeta_0^{T_1}) \mathbf{m}_{i_2}^s(\zeta_0^{T_2})] \mathbf{E}[\tilde{\chi}_{j-1}^{T_1} \tilde{\chi}_{j-1}^{T_2}]$ for all $s \geq 1$.

Proof of Proposition 6.1. We begin by proving (i) with $i = 1$. We notice that $(M_{1,n}^T, \mathcal{H}_{i,n})$ is a martingale since $M_{1,n}^T$ is $\mathcal{H}_{1,n}$ -measurable and

$$\mathbf{E}[M_{1,n}^T | \mathcal{H}_{1,n-1}] = M_{1,n-1}^T + \mathbf{E}[m(\zeta_{n-1}^T) - \mathbf{E}[m(\zeta_0^T)]] \tilde{\chi}_{n-1}^T = M_{1,n-1}^T.$$

By recursively conditioning on $\mathcal{H}_{1,n-1}, \mathcal{H}_{1,n-2}, \dots, \mathcal{H}_{1,1}$, it also follows that

$$\mathbf{E}[M_{1,n}^T] = \mathbf{E}[\mathbf{E}[M_{1,n}^T | \mathcal{H}_{1,n-1}]] = \mathbf{E}[\mathbf{E}[M_{1,n-1}^T | \mathcal{H}_{1,n-2}]] = \dots = \mathbf{E}[M_{1,1}^T] = 0.$$

Next, notice that for $s \geq 0$

$$\mathbf{E}[|D_{1,j}^T|^s | \mathcal{H}_{1,j-1}] = \mathbf{E}[|m(\zeta_{j-1}^T) - \mathbf{E}[m(\zeta_0^T)]|^s | \tilde{\chi}_{j-1}^T] = \mathbf{E}[\mathbf{m}_1^s(\zeta_0^T)] \tilde{\chi}_{j-1}^T \text{ a.s.}$$

In particular, if $s = 2$ then $\mathbf{E}[(D_{1,j}^T)^2 | \mathcal{H}_{1,j-1}] = \mathbf{Var}[m(\zeta_0^T)] \tilde{\chi}_{j-1}^T$ a.s. and the martingale property yields that

$$\mathbf{E}[(M_{1,n}^T)^2] = \mathbf{E}\left[\sum_{j=1}^n \mathbf{E}[(D_{1,j}^T)^2 | \mathcal{H}_{1,j-1}]\right] = \mathbf{Var}[m(\zeta_0^T)] \mathbf{E}[\tilde{C}_n^T].$$

Finally, we notice that, since $D_{1,j}^T$ do not depend on the offspring distributions $\{\xi_{j,i}^T\}_{i=0}^\infty$, part (i) holds with $\mathcal{H}_{1,n}$ replaced by $\mathcal{H}_{2,n}$. We now turn to the proof of (ii). We notice that $M_{2,n}^T$ is $\mathcal{H}_{2,n}$ -measurable and using that $\mathbf{E}[\xi_{n-1,i}^T - m(\zeta_{n-1}^T) | \zeta_{n-1}^T] = 0$ we obtain that

$$\mathbf{E}[M_{2,n}^T | \mathcal{H}_{2,n-1}] = M_{2,n-1}^T + \frac{\tilde{\chi}_{n-1}^T}{Z_{n-1}} \sum_{i=1}^{Z_{n-1}} \mathbf{E}[\mathbf{E}[\xi_{n-1,i}^T - m(\zeta_{n-1}^T) | \zeta_{n-1}^T]] = M_{2,n-1}^T$$

yielding the martingale property. Next, we recursively condition on $\mathcal{H}_{2,n-1}, \mathcal{H}_{2,n-2}, \dots, \mathcal{H}_{2,0}$ and obtain that

$$\mathbf{E}[M_{2,n}^T] = \mathbf{E}[\mathbf{E}[M_{2,n}^T | \mathcal{H}_{2,n-1}]] = \mathbf{E}[\mathbf{E}[M_{2,n-1}^T | \mathcal{H}_{2,n-2}]] = \dots = \mathbf{E}[\mathbf{E}[M_{2,1}^T | \mathcal{H}_{2,0}]] = 0,$$

since

$$\mathbf{E}[M_{2,1}^T | \mathcal{H}_{2,0}] = \frac{\tilde{\chi}_0^T}{Z_0} \sum_{i=1}^{Z_0} \mathbf{E}[\mathbf{E}[\xi_{0,i}^T - m(\zeta_0^T) | \zeta_0^T]] = 0.$$

We now compute

$$\mathbf{E}[(D_{2,j}^T)^2 | \mathcal{H}_{2,j-1}] = \frac{\tilde{\chi}_{j-1}^T}{Z_{j-1}^2} \mathbf{E} \left[\mathbf{E} \left[\left(\sum_{i=1}^{Z_{j-1}} (\xi_{j-1,i}^T - m(\zeta_{j-1}^T)) \right)^2 | \mathcal{H}_{2,j-1}, \zeta_{j-1}^T \right] | \mathcal{H}_{2,j-1} \right].$$

Using that, conditionally on the environment ζ_{j-1}^T , $\{\xi_{j-1,i}^T\}_{i=1}^\infty$ are i.i.d. with variance $\sigma^2(\zeta_{j-1}^T)$, we obtain that

$$\begin{aligned} \mathbf{E} \left[\left(\sum_{i=1}^{Z_{j-1}} \xi_{j-1,i}^T - m(\zeta_{j-1}^T) \right)^2 | \mathcal{H}_{2,j-1}, \zeta_{j-1}^T \right] &= \sum_{i=1}^{Z_{j-1}} \mathbf{E} \left[(\xi_{j-1,i}^T - m(\zeta_{j-1}^T))^2 | \mathcal{H}_{2,j-1}, \zeta_{j-1}^T \right] \\ &= Z_{j-1} \sigma^2(\zeta_{j-1}^T). \end{aligned}$$

We conclude that

$$\mathbf{E}[(D_{2,j}^T)^2 | \mathcal{H}_{2,j-1}] = \mathbf{E}[\sigma^2(\zeta_0^T)] \frac{\tilde{\chi}_{j-1}^T}{Z_{j-1}} \text{ a.s.}$$

and

$$\mathbf{E}[(M_{2,n}^T)^2] = \mathbf{E} \left[\sum_{j=1}^n \mathbf{E}[(D_{2,j}^T)^2 | \mathcal{H}_{2,j-1}] \right] = \mathbf{E}[\sigma^2(\zeta_0^T)] \mathbf{E} \left[\sum_{j=1}^n \frac{\tilde{\chi}_{j-1}^T}{Z_{j-1}} \right].$$

Additionally, Jensen's inequality yields that for $s \geq 1$

$$\mathbf{E}[|D_{2,j}^T|^s | \mathcal{H}_{2,j-1}] \leq \mathbf{E} \left[\frac{\tilde{\chi}_{j-1}^T}{Z_{j-1}} \sum_{i=1}^{Z_{j-1}} |\xi_{j-1,i}^T - m(\zeta_{j-1}^T)|^s | \mathcal{H}_{2,j-1} \right] = \mathbf{E}[\mathbf{m}_2^s(\zeta_0^T)] \tilde{\chi}_{j-1}^T \text{ a.s.}$$

For (iii) we notice that $\sum_{j=1}^n ((D_{i,j}^T)^2 - \mathbf{E}[(D_{i,j}^T)^2])$ is $\tilde{\mathcal{H}}_{i,n}$ -measurable and since $\tilde{\chi}_{n-1}^T, Z_{n-1}, \zeta_{n-1}^T$, and $\{\xi_{n-1,i}^T\}_{i=0}^\infty$ are not $\tilde{\mathcal{H}}_{i,n-1}$ -measurable we have that $\mathbf{E}[(D_{i,n}^T)^2 | \tilde{\mathcal{H}}_{i,n-1}] = \mathbf{E}[(D_{i,n}^T)^2]$ and

$$\mathbf{E} \left[\sum_{j=1}^n ((D_{i,j}^T)^2 - \mathbf{E}[(D_{i,j}^T)^2]) | \tilde{\mathcal{H}}_{i,n-1} \right] = \sum_{j=1}^{n-1} ((D_{i,j}^T)^2 - \mathbf{E}[(D_{i,j}^T)^2]).$$

Again using the convexity of the function $|\cdot|^s$ for $s \geq 1$, we get that

$$\mathbf{E} \left[\left| (D_{i,j}^T)^2 - \mathbf{E}[(D_{i,j}^T)^2] \right|^s \middle| \mathcal{H}_{i,j-1} \right] \leq 2^{s-1} (\mathbf{E}[(D_{i,j}^T)^{2s}] + (\mathbf{E}[(D_{i,j}^T)^2])^s) \leq 2^s \mathbf{E}[(D_{i,j}^T)^{2s}].$$

If $i = 1$, then by conditioning on $\tilde{\chi}_{j-1}^T$ we have that $\mathbf{E}[(D_{1,j}^T)^{2s}] = \mathbf{E}[\mathbf{m}_1^{2s}(\zeta_0^T)] \mathbf{E}[\tilde{\chi}_{j-1}^T]$. If $i = 2$, then we apply Jensen's inequality and obtain that

$$\mathbf{E}[(D_{2,j}^T)^{2s}] \leq \mathbf{E}[\mathbf{m}_2^{2s}(\zeta_{j-1}^T) \tilde{\chi}_{j-1}^T] = \mathbf{E}[\mathbf{m}_2^{2s}(\zeta_0^T)] \mathbf{E}[\tilde{\chi}_{j-1}^T].$$

Turning to (iv), we show that for all $T_1, T_2 \in \{L, U\}$ and $i_1, i_2 \in \{1, 2\}$ such that either $T_1 \neq T_2$ or $i_1 \neq i_2$ it holds that $\mathbf{E}[(D_{i_1,j}^{T_1})(D_{i_2,l}^{T_2})] = 0$ for all $j, l = 1, \dots, n$. This also yields that

$$\mathbf{E}[(M_{i_1,n}^{T_1})(M_{i_2,n}^{T_2})] = \sum_{j=1}^n \sum_{l=1}^n \mathbf{E}[(D_{i_1,j}^{T_1})(D_{i_2,l}^{T_2})] = 0.$$

First, if $l = j$ and $T_1 \neq T_2$, then $\mathbf{E}[(D_{i_1,j}^{T_1})(D_{i_2,j}^{T_2})] = 0$ because $\tilde{\chi}_{j-1}^{T_1} \tilde{\chi}_{j-1}^{T_2} = 0$. Next, if $l = j$ and $i_1 \neq i_2$ (say $i_1 = 1$ or $i_2 = 2$), then by conditioning on $\mathcal{H}_{1,j-1}$ and $\zeta_{j-1}^{T_2}$ and using that

$$\mathbf{E} \left[\xi_{j-1,i}^{T_2} - m(\zeta_{j-1}^{T_2}) \middle| \mathcal{H}_{1,j-1}, \zeta_{j-1}^{T_2} \right] = 0 \text{ a.s.}$$

and $\zeta_{j-1}^{T_1}, \tilde{\chi}_{j-1}^{T_1}, \tilde{\chi}_{j-1}^{T_2}$, and Z_{j-1} are $\mathcal{H}_{1,j-1}$ -measurable, we obtain that

$$\mathbf{E}[(D_{1,j}^{T_1})(D_{2,j}^{T_2})] = \mathbf{E} \left[(m(\zeta_{j-1}^{T_1}) - \mathbf{E}[m(\zeta_0^{T_1})]) \tilde{\chi}_{j-1}^{T_1} \frac{\tilde{\chi}_{j-1}^{T_2}}{Z_{j-1}} \sum_{i=1}^{Z_{j-1}} \mathbf{E} \left[\xi_{j-1,i}^{T_2} - m(\zeta_{j-1}^{T_2}) \middle| \mathcal{H}_{1,j-1}, \zeta_{j-1}^{T_2} \right] \right] = 0.$$

Finally, if $l \neq j$ (say $l > j$), then by conditioning on $\tilde{H}_{2,l-1}$ and using that $D_{i_1,j}$ is $\tilde{H}_{2,l-1}$ -measurable and $\mathbf{E}[(D_{i_2,l}) | \tilde{H}_{2,l-1}] = \mathbf{E}[D_{i_2,l}] = 0$, we obtain that

$$\mathbf{E}[(D_{i_1,j})(D_{i_2,l})] = \mathbf{E}[D_{i_1,j} \mathbf{E}[D_{i_2,l} | \tilde{H}_{2,l-1}]] = 0.$$

$\{(\sum_{j=1}^n (D_{i_1,j}^{T_1})(D_{i_2,j}^{T_2}), \tilde{H}_{2,n})\}_{n=0}^\infty$ is a mean zero martingale sequence since $\sum_{j=1}^n (D_{i_1,j}^{T_1})(D_{i_2,j}^{T_2})$ is $\tilde{H}_{2,n}$ -measurable and $\mathbf{E}[(D_{i_1,j}^{T_1})(D_{i_2,j}^{T_2}) | \tilde{H}_{2,j-1}] = \mathbf{E}[(D_{i_1,j}^{T_1})(D_{i_2,j}^{T_2})] = 0$ if either $T_1 \neq T_2$ or $i_1 \neq i_2$. If $T_1 \neq T_2$ then both $\mathbf{E}[|(D_{i_1,j}^{T_1})(D_{i_2,j}^{T_2})|^s | \tilde{H}_{2,j-1}] = 0$ and $\mathbf{E}[\tilde{\chi}_{j-1}^{T_1} \tilde{\chi}_{j-1}^{T_2}] = 0$. Finally, if $T_1 = T_2$ and $i_1 \neq i_2$ (say $i_1 = 1$ and $i_2 = 2$) then by Jensen's inequality

$$\mathbf{E}[|(D_{i_1,j}^{T_1})(D_{i_2,j}^{T_2})|^s | \zeta_{j-1}^{T_1}, \mathcal{F}_{j-1}] \leq |(D_{i_1,j}^{T_1})|^s \mathbf{m}_{i_2}^s(\zeta_{j-1}^{T_2}) \tilde{\chi}_{j-1}^{T_2},$$

which yields that

$$\mathbf{E}[|(D_{i_1,j}^{T_1})(D_{i_2,j}^{T_2})|^s | \mathcal{F}_{j-1}] \leq \mathbf{E}[\mathbf{m}_{i_1}^s(\zeta_0^{T_1}) \mathbf{m}_{i_2}^s(\zeta_0^{T_2}) \tilde{\chi}_{j-1}^{T_1} \tilde{\chi}_{j-1}^{T_2}]$$

and

$$\mathbf{E}[|(D_{i_1,j}^{T_1})(D_{i_2,j}^{T_2})|^s | \tilde{H}_{2,j-1}] \leq \mathbf{E}[\mathbf{m}_{i_1}^s(\zeta_0^{T_1}) \mathbf{m}_{i_2}^s(\zeta_0^{T_2}) \mathbf{E}[\tilde{\chi}_{j-1}^{T_1} \tilde{\chi}_{j-1}^{T_2}]].$$

□

Asymptotic variance of martingales: Next, we derive the limit of the variances $\mathbf{E}[(M_{1,n}^T)^2]$ and $\mathbf{E}[(M_{2,n}^T)^2]$ when rescaled by n . By Proposition 6.1, this entails studying the limit behavior of the quantities $\frac{1}{n}\tilde{C}_n^T$ and $\frac{1}{n}\tilde{A}_n^T$, where $\tilde{A}_n^T := \sum_{j=1}^n \frac{\tilde{\chi}_{j-1}^T}{Z_{j-1}}$. For $l \geq 1$ and $i \in S_L$ let

$$\tilde{B}_{i,l}^L := (T_{i,l}^L - T_{i,l-1}^L, \{ \sum_{k=\nu_j+1}^{\tau_{j+1}} \tilde{\chi}_{k-1}^U : j = T_{i,l-1}^L, \dots, T_{i,l}^L - 1 \}, \{ \sum_{k=\nu_j+1}^{\tau_{j+1}} \frac{\tilde{\chi}_{k-1}^U}{Z_{k-1}} : j = T_{i,l-1}^L, \dots, T_{i,l}^L - 1 \})$$

and for $l \geq 1$ and $i \in S_U$ let

$$\tilde{B}_{i,l}^U := (T_{i,l}^U - T_{i,l-1}^U, \{ \nu_j - \tau_j : j = T_{i,l-1}^U, \dots, T_{i,l}^U - 1 \}, \{ \sum_{k=\tau_j+1}^{\nu_j} \frac{\tilde{\chi}_{k-1}^L}{Z_{k-1}} : j = T_{i,l-1}^U, \dots, T_{i,l}^U - 1 \}).$$

Notice that, since $\tilde{\chi}_{k-1}^L = \chi_{k-1}^L$, $\tilde{C}_n^L = C_n^L$, $\sum_{k=\tau_j+1}^{\nu_j} \tilde{\chi}_{k-1}^L = \nu_j - \tau_j$ and Theorem 2.5 already yields that $\lim_{n \rightarrow \infty} \frac{C_n^L}{n} = \frac{\mu^L}{\mu^U + \mu^L}$. We need the following slight modification of Lemma 4.1, whose proof is similar and hence omitted.

Lemma 6.1. *Assume (H1)-(H5). (i) If (H6) holds and $Z_{\nu_0} = i \in S_L$, then $\{\tilde{B}_{i,l}^L\}_{l=1}^\infty$ are i.i.d. (ii) If (H7) (or (H8)) holds and $Z_{\tau_1} = i \in S_U$, then $\{\tilde{B}_{i,l}^U\}_{l=1}^\infty$ are i.i.d.*

Proposition 6.2. *Suppose that (H1)-(H7) (or (H8)) hold and $\mu^U, \mu^L < \infty$. Then*

- (i) $\lim_{n \rightarrow \infty} \frac{\tilde{C}_n^U}{\tilde{N}^L(n)} = \tilde{\mu}^U$ and $\lim_{n \rightarrow \infty} \frac{\tilde{C}_n^U}{n} = \frac{\tilde{\mu}^U}{\mu^U + \mu^L}$ a.s.
- (ii) $\lim_{n \rightarrow \infty} \frac{\tilde{A}_n^U}{\tilde{N}^L(n)} = \tilde{a}^U$ and $\lim_{n \rightarrow \infty} \frac{\tilde{A}_n^U}{n} = \frac{\tilde{a}^U}{\mu^U + \mu^L}$ a.s.
- (iii) $\lim_{n \rightarrow \infty} \frac{\tilde{A}_n^L}{\tilde{N}^U(n)} = \tilde{a}^L$ and $\lim_{n \rightarrow \infty} \frac{\tilde{A}_n^L}{n} = \frac{\tilde{a}^L}{\mu^U + \mu^L}$ a.s.

Since $\frac{\tilde{C}_n^T}{n}$ and $\frac{\tilde{A}_n^T}{n}$ are non-negative and bounded by one, Proposition 6.2 implies convergence in mean of these quantities.

Proof of Proposition 6.2. By Lemma 5.2 (i) it is enough to show the first part of the statements (i)-(iii). Since the proof of the other cases is similar we only prove (i). We recall that for $i \in S_L$ and $j \in \mathbb{N}$, $N_i^L(j) = \sum_{l=1}^j \mathbf{1}_{[T_{i,l}^L \leq j]}$ is the number of times $T_{i,l}^L$ is in $\{0, 1, \dots, j\}$. Conditionally on $Z_{\nu_0} = i$, $T_{i,0}^L = 0$ and we write

$$\frac{\tilde{C}_n^U}{\tilde{N}^L(n)} = \frac{N_i^L(\tilde{N}^L(n))}{\tilde{N}^L(n)} \left(\frac{1}{N_i^L(\tilde{N}^L(n))} \sum_{l=1}^{N_i^L(\tilde{N}^L(n))} (\tilde{C}_{\nu_{T_{i,l}^L}^L}^U - \tilde{C}_{\nu_{T_{i,l-1}^L}^L}^U) + \frac{1}{N_i^L(\tilde{N}^L(n))} (\tilde{C}_n^U - \tilde{C}_{\nu_{T_{i,N_i^L(\tilde{N}^L(n))}^L}^U}) \right).$$

Lemma 6.1 implies that

$$(\tilde{C}_{\nu_{T_{i,l}^L}^L}^U - \tilde{C}_{\nu_{T_{i,l-1}^L}^L}^U) = \sum_{j=T_{i,l-1}^L}^{T_{i,l}^L-1} \sum_{k=\nu_j+1}^{\tau_{j+1}} \tilde{\chi}_{k-1}^U$$

are i.i.d. in l with expectation that using Proposition 1.69 of Serfozo (2009) is given by $\mathbf{E}[\tilde{C}_{\nu_{T_{i,1}^L}^L}^U | Z_{\nu_0} =$

$i] = (\pi_i^L)^{-1} \tilde{\mu}^U$. Since $\lim_{n \rightarrow \infty} N_i^L(\tilde{N}^L(n)) = \infty$ a.s., we obtain that

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{N_i^L(\tilde{N}^L(n))} \sum_{l=1}^{N_i^L(\tilde{N}^L(n))} (\tilde{C}_{\nu_{T_i^L, l}}^U - \tilde{C}_{\nu_{T_i^L, l-1}}^U) &= (\pi_i^L)^{-1} \tilde{\mu}^U \text{ a.s. and} \\ \lim_{n \rightarrow \infty} \frac{1}{N_i^L(\tilde{N}^L(n))} (\tilde{C}_n^U - \tilde{C}_{\nu_{T_i^L, N_i^L(\tilde{N}^L(n))}}^U) &= 0 \text{ a.s.,} \end{aligned}$$

since $(\tilde{C}_n^U - \tilde{C}_{\nu_{T_i^L, N_i^L(\tilde{N}^L(n))}}^U) \leq (\tilde{C}_{\nu_{T_i^L, N_i^L(\tilde{N}^L(n))+1}}^U - \tilde{C}_{\nu_{T_i^L, N_i^L(\tilde{N}^L(n))}}^U)$. Finally, it holds that

$$\lim_{n \rightarrow \infty} \frac{N_i^L(\tilde{N}^L(n))}{\tilde{N}^L(n)} = \pi_i^L \text{ a.s.}$$

□

Asymptotic normality: For $T \in \{L, U\}$, let

$$\mathbf{M}_n^T := \left(\frac{M_{1,n}^T}{\sqrt{\mathbf{E}[(M_{1,n}^T)^2]}}, \frac{M_{2,n}^T}{\sqrt{\mathbf{E}[(M_{2,n}^T)^2]}} \right)^\top \text{ and } \mathbf{M}_n := ((\mathbf{M}_n^U)^\top, (\mathbf{M}_n^L)^\top)^\top.$$

Lemma 6.2. *Under the assumption of Theorem 2.6 (ii), $\mathbf{M}_n \xrightarrow[n \rightarrow \infty]{d} N(\mathbf{0}, \mathbf{I})$.*

Proof of Lemma 6.2. By Cramér–Wold theorem (see Theorem 29.4 of Billingsley (2013)), it is enough to show that for $t_i^T \in \mathbb{R}$, where $i = 1, 2$ and $T \in \{L, U\}$,

$$\sum_{T \in \{L, U\}} \sum_{i=1}^2 t_i^T \frac{M_{i,n}^T}{\sqrt{\mathbf{E}[(M_{i,n}^T)^2]}} \xrightarrow[n \rightarrow \infty]{d} N\left(0, \sum_{T \in \{L, U\}} \sum_{i=1}^2 (t_i^T)^2\right). \quad (25)$$

Using Proposition 6.1, we see that

$$\left\{ \left(\sum_{T \in \{L, U\}} \sum_{i=1}^2 t_i^T M_{i,n}^T, \mathcal{H}_{2,n} \right) \right\}_{n=1}^\infty$$

is a mean zero martingale sequence. In particular,

$$\left\{ \left(\sum_{T \in \{L, U\}} \sum_{i=1}^2 t_i^T \frac{M_{i,j}^T}{\sqrt{\mathbf{E}[(M_{i,n}^T)^2]}}, \mathcal{H}_{2,j} \right) \right\}_{j=1}^n$$

is a mean zero martingale array. We will apply Theorem 3.2 of Hall and Heyde (1980) with $k_n = n$, $X_{nl} = \sum_{T \in \{L, U\}} \sum_{i=1}^2 t_i^T \frac{D_{i,l}^T}{\sqrt{\mathbf{E}[(M_{i,n}^T)^2]}}$, $S_{nj} = \sum_{l=1}^j X_{nl} = \sum_{T \in \{L, U\}} \sum_{i=1}^2 t_i^T \frac{M_{i,j}^T}{\sqrt{\mathbf{E}[(M_{i,n}^T)^2]}}$, $\mathcal{F}_{nj} = \mathcal{H}_{2,j}$, and $B^2 = \sum_{T \in \{L, U\}} \sum_{i=1}^2 (t_i^T)^2$; and obtain (25). To this end, we need to verify the following conditions: (i) $\mathbf{E}[(S_{nj})^2] < \infty$, (ii) $\max_{l=1, \dots, n} |X_{nl}| \xrightarrow[n \rightarrow \infty]{p} 0$, (iii) $\sum_{l=1}^n X_{nl}^2 \xrightarrow[n \rightarrow \infty]{p} B^2$, and (iv)

$\sup_{n \in \mathbb{N}} \mathbf{E}[\max_{l=1, \dots, n} X_{nl}^2] < \infty$. Using Proposition 6.1 (iv) $\mathbf{E}[(M_{i_1, j}^{T_1})(M_{i_2, j}^{T_2})] = 0$ if either $T_1 \neq T_2$ or $i_1 \neq i_2$ and since $\mathbf{E}[(M_{i, j}^T)^2]$ are non-decreasing in j , we obtain that

$$\mathbf{E} \left[\left(\sum_{T \in \{L, U\}} \sum_{i=1}^2 t_i^T \frac{M_{i, j}^T}{\sqrt{\mathbf{E}[(M_{i, n}^T)^2]}} \right)^2 \right] = \sum_{T \in \{L, U\}} \sum_{i=1}^2 (t_i^T)^2 \frac{\mathbf{E}[(M_{i, j}^T)^2]}{\mathbf{E}[(M_{i, n}^T)^2]} \leq \sum_{T \in \{L, U\}} \sum_{i=1}^2 (t_i^T)^2 < \infty$$

yielding Condition (i). Using again that $\mathbf{E}[(D_{i_1, l}^{T_1})(D_{i_2, l}^{T_2})] = 0$ if either $T_1 \neq T_2$ or $i_1 \neq i_2$ and $\mathbf{E}[(M_{i, n}^T)^2] = \sum_{l=1}^n \mathbf{E}[(D_{i, l}^T)^2]$, we obtain that

$$\begin{aligned} \mathbf{E} \left[\max_{l=1, \dots, n} \left(\sum_{T \in \{L, U\}} \sum_{i=1}^2 t_i^T \frac{D_{i, l}^T}{\sqrt{\mathbf{E}[(M_{i, n}^T)^2]}} \right)^2 \right] &\leq \mathbf{E} \left[\sum_{l=1}^n \left(\sum_{T \in \{L, U\}} \sum_{i=1}^2 t_i^T \frac{D_{i, l}^T}{\sqrt{\mathbf{E}[(M_{i, n}^T)^2]}} \right)^2 \right] \\ &= \sum_{T \in \{L, U\}} \sum_{i=1}^2 (t_i^T)^2 \end{aligned}$$

yielding Condition (iv). Turning to Condition (ii), assuming w.l.o.g. that $t_i^T \neq 0$ and using that

$$\left(\sum_{T \in \{L, U\}} \sum_{i=1}^2 t_i^T \frac{D_{i, l}^T}{\sqrt{\mathbf{E}[(M_{i, n}^T)^2]}} \right)^2 \leq 4 \sum_{T \in \{L, U\}} \sum_{i=1}^2 (t_i^T)^2 \frac{(D_{i, l}^T)^2}{\mathbf{E}[(M_{i, n}^T)^2]},$$

we obtain that for all $\epsilon > 0$

$$\begin{aligned} P \left(\max_{l=1, \dots, n} \left| \sum_{T \in \{L, U\}} \sum_{i=1}^2 t_i^T \frac{D_{i, l}^T}{\sqrt{\mathbf{E}[(M_{i, n}^T)^2]}} \right| \geq \epsilon \right) &\leq \sum_{l=1}^n P \left(\left(\sum_{T \in \{L, U\}} \sum_{i=1}^2 t_i^T \frac{D_{i, l}^T}{\sqrt{\mathbf{E}[(M_{i, n}^T)^2]}} \right)^2 \geq \epsilon^2 \right) \\ &\leq \sum_{T \in \{L, U\}} \sum_{i=1}^2 \sum_{l=1}^n P \left((D_{i, l}^T)^2 \geq \left(\frac{\epsilon}{4t_i^T} \right)^2 \mathbf{E}[(M_{i, n}^T)^2] \right). \end{aligned}$$

For $i = 1$, we use that $\tilde{\chi}_{l-1}^T \in \{0, 1\}$ and obtain that

$$\begin{aligned} &\sum_{l=1}^n P \left((D_{1, l}^T)^2 \geq \left(\frac{\epsilon}{4t_1^T} \right)^2 \mathbf{E}[(M_{1, n}^T)^2] \right) \\ &\leq \sum_{l=1}^n P \left((D_{1, l}^T)^2 \geq \left(\frac{\epsilon}{4t_1^T} \right)^2 \mathbf{E}[(M_{1, n}^T)^2] \mid \tilde{\chi}_{l-1}^T = 1 \right) \\ &= \frac{n}{\mathbf{E}[(M_{1, n}^T)^2]} \mathbf{E}[(M_{1, n}^T)^2] P \left((m(\zeta_0^T) - \mathbf{E}[m(\zeta_0^T)])^2 \geq \left(\frac{\epsilon}{4t_1^T} \right)^2 \mathbf{E}[(M_{1, n}^T)^2] \right). \end{aligned}$$

It follows from Proposition 6.1 (i) and Proposition 6.2 (i) that

$$\lim_{n \rightarrow \infty} \frac{n}{\mathbf{E}[(M_{1, n}^T)^2]} = \frac{1}{\mathbf{Var}[m(\zeta_0^T)]} \frac{\mu^U + \mu^L}{\tilde{\mu}^T} < \infty \quad (26)$$

and since $\mathbf{Var}[m(\zeta_0^T)] < \infty$

$$\lim_{n \rightarrow \infty} \mathbf{E}[(M_{1, n}^T)^2] P \left((m(\zeta_0^T) - \mathbf{E}[m(\zeta_0^T)])^2 \geq \left(\frac{\epsilon}{4t_1^T} \right)^2 \mathbf{E}[(M_{1, n}^T)^2] \right) = 0$$

yielding that

$$\lim_{n \rightarrow \infty} \sum_{l=1}^n P \left((D_{1,l}^T)^2 \geq \left(\frac{\epsilon}{4t_1^T} \right)^2 \mathbf{E}[(M_{1,n}^T)^2] \right) = 0.$$

For $i = 2$, we use that if $\tilde{\chi}_{l-1}^T = 1$ then $Z_{l-1} \geq 1$ and obtain that

$$\begin{aligned} & \sum_{l=1}^n P \left((D_{2,l}^T)^2 \geq \left(\frac{\epsilon}{4t_2^T} \right)^2 \mathbf{E}[(M_{2,n}^T)^2] \right) \\ & \leq n \sup_{z \in \mathbb{N}} P \left(\left(\frac{1}{z} \sum_{i=1}^z (\xi_{i,0}(\zeta_0^T) - m(\zeta_0^T)) \right)^2 \geq \left(\frac{\epsilon}{4t_2^T} \right)^2 \mathbf{E}[(M_{2,n}^T)^2] \right). \end{aligned}$$

Next, using Proposition 6.1 (ii) and Proposition 6.2 (ii)-(iii), we have that

$$\lim_{n \rightarrow \infty} \frac{n}{\mathbf{E}[(M_{2,n}^T)^2]} \leq \frac{1}{\mathbf{E}[\sigma^2(\zeta_0^T)]} \frac{\mu^U + \mu^L}{\tilde{a}^T} < \infty. \quad (27)$$

Since by Jensen's inequality

$$\mathbf{E} \left[\left| \frac{1}{z} \sum_{i=1}^z (\xi_{i,0}(\zeta_0^T) - m(\zeta_0^T)) \right|^{2+\delta} \right] \leq \mathbf{E}[\mathfrak{m}_2^{2+\delta}(\zeta_0^T)] < \infty,$$

using Markov inequality, we obtain that

$$\sum_{n=0}^{\infty} \sup_{z \in \mathbb{N}} P \left(\left(\frac{1}{z} \sum_{i=1}^z (\xi_{i,0}(\zeta_0^T) - m(\zeta_0^T)) \right)^2 \geq \left(\frac{\epsilon}{4t_2^T} \right)^2 \mathbf{E}[(M_{2,n}^T)^2] \right) < \infty,$$

which yields that

$$\lim_{n \rightarrow \infty} n \sup_{z \in \mathbb{N}} P \left(\left(\frac{1}{z} \sum_{i=1}^z (\xi_{i,0}(\zeta_0^T) - m(\zeta_0^T)) \right)^2 \geq \left(\frac{\epsilon}{4t_2^T} \right)^2 \mathbf{E}[(M_{2,n}^T)^2] \right) = 0.$$

For (iii), we decompose

$$\sum_{l=1}^n \left(\sum_{T \in \{L,U\}} \sum_{i=1}^2 t_i^T \frac{D_{i,l}^T}{\sqrt{\mathbf{E}[(M_{i,n}^T)^2]}} \right)^2 - \sum_{T \in \{L,U\}} \sum_{i=1}^2 (t_i^T)^2$$

as

$$\sum_{T \in \{L,U\}} \sum_{i=1}^2 (t_i^T)^2 \frac{\sum_{l=1}^n (D_{i,l}^T)^2 - \mathbf{E}[(M_{i,n}^T)^2]}{\mathbf{E}[(M_{i,n}^T)^2]} + \sum_{T_1 \neq T_2 \text{ or } i_1 \neq i_2} t_{i_1}^{T_1} t_{i_2}^{T_2} \frac{\sum_{l=1}^n (D_{i_1,l}^{T_1})(D_{i_2,l}^{T_2})}{\sqrt{\mathbf{E}[(M_{i_1,n}^{T_1})^2] \mathbf{E}[(M_{i_2,n}^{T_2})^2]}} \quad (28)$$

and show that each of the above terms converges to zero with probability one. Since $\mathbf{E}[(M_{i,n}^T)^2] = \sum_{l=1}^n \mathbf{E}[(D_{i,l}^T)^2]$, we use Proposition 6.1 (iii) and obtain that $\{(\sum_{l=1}^n ((D_{i,l}^T)^2 - \mathbf{E}[(D_{i,l}^T)^2]), \tilde{\mathcal{H}}_{i,n})\}_{n=1}^{\infty}$ are mean zero martingale sequences and for $s = 1 + \delta/2$

$$\mathbf{E} \left[\left| (D_{i,l}^T)^2 - \mathbf{E}[(D_{i,l}^T)^2] \right|^s \middle| \tilde{\mathcal{H}}_{i,l-1} \right] \leq 2^s \mathbf{E}[\mathfrak{m}_i^{2s}(\zeta_0^T)] \mathbf{E}[\tilde{\chi}_{l-1}^T] < \infty.$$

We use (26) and (27), and apply Theorem 2.18 of Hall and Heyde (1980) with $S_n = \sum_{l=1}^n ((D_{i,l}^T)^2 - \mathbf{E}[(D_{i,l}^T)^2])$, $X_l = (D_{i,l}^T)^2 - \mathbf{E}[(D_{i,l}^T)^2]$, $\mathcal{F}_n = \tilde{\mathcal{H}}_{i,n}$, $U_n = \mathbf{E}[(M_{i,n}^T)^2]$, and $p = s$, where $i = 1, 2$, and obtain the convergence of the first term in (28). For the second term we proceed similarly. Specifically, using Proposition 6.1 (iv) and Cauchy-Schwartz inequality we obtain that $\{(\sum_{l=1}^n (D_{i_1,l}^{T_1})(D_{i_2,l}^{T_2}), \tilde{\mathcal{H}}_{2,n})\}_{n=0}^\infty$ is a mean zero martingale sequence and for $s = 1 + \delta/2$

$$\mathbf{E}[|(D_{i_1,l}^{T_1})(D_{i_2,l}^{T_2})|^s | \tilde{\mathcal{H}}_{2,l-1}] \leq \mathbf{E}[\mathbf{m}_{i_1}^s(\zeta_0^{T_1}) \mathbf{m}_{i_2}^s(\zeta_0^{T_2})] \leq \sqrt{\mathbf{E}[\mathbf{m}_{i_1}^{2+\delta}(\zeta_0^{T_1})] \mathbf{E}[\mathbf{m}_{i_2}^{2+\delta}(\zeta_0^{T_2})]} < \infty.$$

Finally, we apply Theorem 2.18 of Hall and Heyde (1980) with $S_n = \sum_{l=1}^n (D_{i_1,l}^{T_1})(D_{i_2,l}^{T_2})$, $X_l = (D_{i_1,l}^{T_1})(D_{i_2,l}^{T_2})$, $\mathcal{F}_n = \tilde{\mathcal{H}}_{2,n}$, $U_n = \sqrt{\mathbf{E}[(M_{i_1,n}^{T_1})^2] \mathbf{E}[(M_{i_2,n}^{T_2})^2]}$, and $p = s$; and obtain convergence of the second term in (28). \square

We are now ready to prove the main result of the section.

Proof of Theorem 2.6. Using Proposition 6.1 (i)-(ii) and $\tilde{\chi}_{j-1}^T \leq 1$, we obtain that for $i = 1, 2$ $\{(M_{i,n}^T, \mathcal{H}_{i,n})\}_{n=1}^\infty$ are martingales and

$$\sum_{j=1}^\infty \frac{1}{j^s} \mathbf{E}[|D_{i,j}^T|^s | \mathcal{H}_{i,j-1}] \leq \mathbf{E}[\mathbf{m}_i^s(\zeta_0^T)] \sum_{j=1}^\infty \frac{1}{j^s} < \infty.$$

We apply Theorem 2.18 of Hall and Heyde (1980) with $S_n = M_{i,n}^T$, $X_j = D_{i,j}^T$, where $i = 1, 2$ and $T \in \{L, U\}$, $U_n = n$, $p = s$, and $\mathcal{F}_n = \mathcal{H}_{1,n}$ for $i = 1$ and $\mathcal{F}_n = \mathcal{H}_{2,n}$ for $i = 2$ and obtain that $\lim_{n \rightarrow \infty} \frac{1}{n} M_{i,n}^T = 0$ a.s. From this, Theorem 2.5, and Proposition 6.2 (i), we obtain that $\lim_{n \rightarrow \infty} \frac{1}{\tilde{C}_n^T} M_{i,n}^T = 0$ a.s. Using (24) we conclude that $\lim_{n \rightarrow \infty} \hat{m}_n^T = \mathbf{E}[m(\zeta_0^T)]$ a.s. Turning to the central limit theorem, Lemma 6.2 (iii) yields that

$$\mathbf{M}_n = \left(\frac{M_{1,n}^U}{\sqrt{\mathbf{E}[(M_{1,n}^U)^2]}}, \frac{M_{2,n}^U}{\sqrt{\mathbf{E}[(M_{2,n}^U)^2]}}, \frac{M_{1,n}^L}{\sqrt{\mathbf{E}[(M_{1,n}^L)^2]}}, \frac{M_{2,n}^L}{\sqrt{\mathbf{E}[(M_{2,n}^L)^2]}} \right)^\top$$

is asymptotically normal with mean zero and identity covariance matrix. Let D_n^2 be the 4×4 diagonal matrix

$$D_n^2 := \text{Diag} \left(\frac{n \mathbf{E}[(M_{1,n}^U)^2]}{(\tilde{C}_n^U)^2}, \frac{n \mathbf{E}[(M_{2,n}^U)^2]}{(\tilde{C}_n^U)^2}, \frac{n \mathbf{E}[(M_{1,n}^L)^2]}{(\tilde{C}_n^L)^2}, \frac{n \mathbf{E}[(M_{2,n}^L)^2]}{(\tilde{C}_n^L)^2} \right).$$

By Proposition 6.1 (i)-(ii) and Proposition 6.2, $D_n \mathbf{M}_n$ is asymptotically normal with mean zero and covariance matrix

$$D_o^2 := (\mu^U + \mu^L) \text{Diag} \left(\mathbf{Var}[m(\zeta_0^U)] \frac{1}{\tilde{\mu}^U}, \mathbf{E}[\sigma^2(\zeta_0^U)] \frac{\tilde{a}^U}{(\tilde{\mu}^U)^2}, \mathbf{Var}[m(\zeta_0^L)] \frac{1}{\mu^L}, \mathbf{E}[\sigma^2(\zeta_0^L)] \frac{\tilde{a}^L}{(\mu^L)^2} \right).$$

Using the continuous mapping theorem, it follows that

$$\sqrt{n}((\hat{m}_n^U, \hat{m}_n^L)^\top - \mathbf{m}_o) \xrightarrow[n \rightarrow \infty]{d} N(\mathbf{0}, \Sigma_o).$$

\square

7 Discussion and concluding remarks

In this paper we developed BPRE with Thresholds to describe periods of growth and decrease in the population size arising in several applications including COVID dynamics. Even though the model is non-Markov, we identify Markov subsequences and use them to understand the length of time the process spends in the supercritical and subcritical regimes. Furthermore, using the regeneration technique we also study the rate of growth (or decline) of the process in the supercritical (subcritical) regime. It is possible to start the process using the subcritical BPRE and then move to the supercritical regime; this introduces only minor changes and the qualitative results remain the same. Finally, we note that without incorporating immigration in the supercritical regime the process will become extinct with probability one and hence the cyclical path behavior may not be observed.

An interesting question concerns the choice of strongly subcritical BPRE for the subcritical regime. It is folklore that the generation sizes of moderately and weakly subcritical processes can increase for long periods of time and in that case the time to cross the lower threshold will have a heavier tail. This could lead to lack of identifiability of supercritical and subcritical regimes. Similar issues arise when a subcritical BPRE is replaced by a critical BPRE. The path properties of these alternatives could be useful for modeling other dynamics observed (see Klebaner (1993); Iannelli and Pugliese (2014)). Mathematical issues arising from these alternatives would involve different techniques than used in this paper. Finally, we end this section with a brief discussion concerning the moment conditions in Theorem 2.6. It is possible to reduce the conditions $\mathbf{E}[\mathbf{m}_i^{2+\delta}(\zeta_0^T)] < \infty$ to finite second moment hypothesis. This requires an extension of Lemma 4.1 to joint independence of blocks in $B_{i,l}^L, B_{i,l}^U$, offspring random variables, environments, immigration over cycles. The proof will need the Markov property of the pair $\{(Z_{\nu_{j-1}}, Z_{\tau_j})\}_{j=1}^\infty$ and its uniform ergodicity. Also, the joint Markov property will also yield joint central limit theorem for the length and proportion of time spent in the supercritical and subcritical regimes. The proof is similar to that of Theorem 2.3 and Lemma 4.1 but is more cumbersome with an increased notational burden. Extensions of these ideas under Markovian environments and thresholds is under consideration. The numerical experiments suggest that the estimators of the mean parameters of the supercritical and subcritical regime are not affected by the choice of various distributions. A thorough statistical analysis of the robustness of the estimators and analysis of the datasets are beyond the scope of this paper and will be investigated elsewhere.

References

- Atanasov, D., Stoimenova, V. and Yanev, N. M. (2021). Branching process modelling of COVID-19 pandemic including immunity and vaccination, *Stochastics and Quality Control* **36**: 157–164.
- Athreya, K. B. and Karlin, S. (1971). On branching processes with random environments: I: Extinction probabilities, *The Annals of Mathematical Statistics* **42**: 1499–1520.

- Athreya, K. B. and Schuh, H.-J. (2016). A Galton-Watson process with a threshold, *Journal of Applied Probability* **53**: 614–621.
- Bansaye, V. (2009). Cell contamination and branching processes in a random environment with immigration, *Advances in Applied Probability* **41**: 1059–1081.
- Billingsley, P. (2013). *Convergence of probability measures*, John Wiley & Sons.
- Dion, J. P. and Esty, W. W. (1979). Estimation problems in branching processes with random environments, *The Annals of Statistics* pp. 680–685.
- Falcó, C. and Corral, Á. (2021). Finite-time scaling for epidemic processes with power-law super-spreading events, *arXiv preprint arXiv:2110.10459* .
- Grama, I., Liu, Q. and Miqueu, E. (2017). Harmonic moments and large deviations for a supercritical branching process in a random environment, *Electronic Journal of Probability* **22**: 1–23.
- Haccou, P., Jagers, P. and Vatutin, V. A. (2007). *Branching processes: variation, growth, and extinction of populations*, Cambridge University Press.
- Hall, P. and Heyde, C. C. (1980). *Martingale limit theory and its application*, Academic press.
- Hanlon, B. and Vidyashankar, A. N. (2011). Inference for quantitation parameters in polymerase chain reactions via branching processes with random effects, *Journal of the American Statistical Association* **106**: 525–533.
- Hempel, K. and Earn, D. J. D. (2015). A century of transitions in new york city’s measles dynamics, *Journal of the Royal Society Interface* **12**.
- Heyde, C. C. (1971). Some central limit analogues for supercritical Galton-Watson processes, *Journal of Applied Probability* **8**: 52–59.
- Iannelli, M. and Pugliese, A. (2014). *An introduction to mathematical population dynamics*, Springer.
- Ibragimov, I. A. and Linnik, Y. V. (1971). *Independent and stationary sequences of random variables*, Wolters-Noordhoff Publishing, Groningen.
- Jagers, P. (1975). *Branching processes with biological applications*, Wiley.
- Jagers, P. and Klebaner, F. C. (2011). Population-size-dependent, age-structured branching processes linger around their carrying capacity, *Journal of Applied Probability* **48**: 249–260.
- Jagers, P. and Zuyev, S. (2020). Populations in environments with a soft carrying capacity are eventually extinct, *Journal of Mathematical Biology* **81**: 845–851.
- Jagers, P. and Zuyev, S. (2021). Amendment to: populations in environments with a soft carrying capacity are eventually extinct, *Journal of Mathematical Biology* **83**.

- Kersting, G. and Vatutin, V. A. (2017). *Discrete time branching processes in random environment*, John Wiley & Sons.
- Kimmel, M. and Axelrod, D. E. (2015). *Branching processes in biology*, Springer, New York.
- Klebaner, F. C. (1984). On population-size-dependent branching processes, *Advances in Applied Probability* **16**: 30–55.
- Klebaner, F. C. (1993). Population-dependent branching processes with a threshold, *Stochastic processes and their applications* **46**: 115–127.
- Lamperti, J. (1960). Criteria for the recurrence or transience of stochastic process. I, *Journal of Mathematical Analysis and applications* **1**: 314–330.
- Lamperti, J. (1963). Criteria for stochastic processes II: passage-time moments, *Journal of Mathematical Analysis and Applications* **7**: 127–145.
- Perko, L. (2013). *Differential equations and dynamical systems*, Springer.
- Serfozo, R. (2009). *Basics of applied stochastic processes*, Springer.
- Sun, H., Kryven, I. and Bianconi, G. (2022). Critical time-dependent branching process modelling epidemic spreading with containment measures, *Journal of Physics A: Mathematical and Theoretical* **55**.
- Teschl, G. (2012). *Ordinary differential equations and dynamical systems*, American Mathematical Society.
- Tyson, R., Haines, S. and Hodges, K. E. (2010). Modelling the canada lynx and snowshoe hare population cycle: the role of specialist predators, *Theoretical Ecology* **3**: 97–111.
- Yanev, N. M., Stoimenova, V. K. and Atanasov, D. V. (2020). Stochastic modeling and estimation of COVID-19 population dynamics, *arXiv preprint arXiv:2004.00941* .

A Additional proofs

We recall that uniform ergodicity of the Markov chains $\{Z_{\nu_j}\}_{j=0}^{\infty}$ and $\{Z_{\tau_j}\}_{j=1}^{\infty}$ is equivalent to the existence of constants $C_T \geq 0$ and $\rho_T \in (0, 1)$ such that $\sup_{l \in S_T} \|\mathbf{p}_l^T(j) - \boldsymbol{\pi}^T\| \leq C_T \rho_T^j$.

Proof of Lemma 4.2. Since the proof of (ii) is similar we only prove (i). We proceed along the lines of the proof of Theorem 17.2.3 of Ibragimov and Linnik (1971). Using Cauchy-Schwarz inequality,

we have that

$$\begin{aligned}
& \left| \sum_{k \in S_L} \sum_{i \in S_L} w_k^L w_i^L \pi_k^L p_{ki}^L(j) - \left(\sum_{k \in S_L} w_k^L \pi_k^L \right) \left(\sum_{i \in S_L} w_i^L \pi_i^L \right) \right| \\
&= \left| \sum_{k \in S_L} w_k^L (\pi_k^L)^{1/2} \sum_{i \in S_L} w_i^L (p_{ki}^L(j) - \pi_i^L) (\pi_k^L)^{1/2} \right| \\
&\leq \left(\sum_{k \in S_L} (w_k^L)^2 \pi_k^L \right)^{1/2} \left(\sum_{k \in S_L} \pi_k^L \left(\sum_{i \in S_L} w_i^L (p_{ki}^L(j) - \pi_i^L) \right)^2 \right)^{1/2}.
\end{aligned}$$

Using Cauchy-Schwarz inequality again, we obtain that

$$\begin{aligned}
\left(\sum_{i \in S_L} w_i^L (p_{ki}^L(j) - \pi_i^L) \right)^2 &\leq \left(\sum_{i \in S_L} |w_i^L| (p_{ki}^L(j) + \pi_i^L)^{1/2} |p_{ki}^L(j) - \pi_i^L|^{1/2} \right)^2 \\
&\leq \left(\sum_{i \in S_L} (w_i^L)^2 (p_{ki}^L(j) + \pi_i^L) \right) \left(\sum_{i \in S_L} |p_{ki}^L(j) - \pi_i^L| \right).
\end{aligned}$$

Since $\sum_{k \in S_L} p_{ki}^L(j) \pi_k^L = \pi_i^L$ by Remark 3.1, we deduce that

$$\begin{aligned}
\left(\sum_{k \in S_L} \pi_k^L \left(\sum_{i \in S_L} w_i^L (p_{ki}^L(j) - \pi_i^L) \right)^2 \right)^{1/2} &\leq \left(\sum_{k \in S_L} \pi_k^L \left(\sum_{i \in S_L} (w_i^L)^2 (p_{ki}^L(j) + \pi_i^L) \right) \left(\sum_{i \in S_L} |p_{ki}^L(j) - \pi_i^L| \right) \right)^{1/2} \\
&\leq \left(2 \sum_{i \in S_L} (w_i^L)^2 \pi_i^L \right)^{1/2} \sup_{k \in S_L} \left(\sum_{i \in S_L} |p_{ki}^L(j) - \pi_i^L| \right)^{1/2}.
\end{aligned}$$

Using that $\sup_{i \in S_L} \|p_i^L(j) - \pi^L\| \leq C_L \rho_L^j$, we obtain that

$$\begin{aligned}
\sup_{k \in S_L} \left(\sum_{i \in S_L} |p_{ki}^L(j) - \pi_i^L| \right) &\leq \sup_{k \in S_L} \left(\sum_{i \in S_L: p_{ki}^L(j) - \pi_i^L > 0} (p_{ki}^L(j) - \pi_i^L) \right) + \sup_{k \in S_L} \left(\sum_{i \in S_L: p_{ki}^L(j) - \pi_i^L < 0} (\pi_i^L - p_{ki}^L(j)) \right) \\
&\leq \sup_{k \in S_L} \left(\sum_{i \in S_L: p_{ki}^L(j) - \pi_i^L > 0} p_{ki}^L(j) - \sum_{i \in S_L: p_{ki}^L(j) - \pi_i^L > 0} \pi_i^L \right) \\
&\quad + \sup_{k \in S_L} \left(\sum_{i \in S_L: p_{ki}^L(j) - \pi_i^L < 0} \pi_i^L - \sum_{i \in S_L: p_{ki}^L(j) - \pi_i^L < 0} p_{ki}^L(j) \right) \\
&\leq 2C_L \rho_L^j.
\end{aligned}$$

We have thus shown that

$$\left| \sum_{k \in S_L} \sum_{i \in S_L} w_k^L w_i^L \pi_k^L p_{ki}^L(j) - \left(\sum_{k \in S_L} w_k^L \pi_k^L \right) \left(\sum_{i \in S_L} w_i^L \pi_i^L \right) \right| \leq 2C_L^{1/2} \rho_L^{j/2} \left(\sum_{k \in S_L} (w_k^L)^2 \pi_k^L \right),$$

which yields that

$$\sum_{j=1}^{\infty} \left| \sum_{k \in S_L} \sum_{i \in S_L} w_k^L w_i^L \pi_k^L p_{ki}^L(j) - \left(\sum_{k \in S_L} w_k^L \pi_k^L \right) \left(\sum_{i \in S_L} w_i^L \pi_i^L \right) \right| \leq 2C_L^{1/2} \frac{\rho_L^{1/2}}{1 - \rho_L^{1/2}} \left(\sum_{k \in S_L} (w_k^L)^2 \pi_k^L \right).$$

□

Proof of Lemma 4.3. The first part of (i) follows from Proposition 1.69 of Serfozo (2009) with $X_j = Z_{\nu_j}$, $\pi = \boldsymbol{\pi}^L = \{\pi_i^L\}_{i \in S_L}$, and $V_j = \Delta_{j+1}^U$. For the second part of (i) we use the above proposition with $V_j = \Delta_{j+1}^U + \Delta_{j+1}^L$ and obtain that

$$\mathbf{E}[\tilde{S}_{T_{i,1}^L}^U | Z_{\nu_0} = i] = (\pi_i^L)^{-1} \mathbf{E}[\Delta_1^U + \Delta_1^L | Z_{\nu_0} \sim \boldsymbol{\pi}^L] = (\pi_i^L)^{-1} (\mathbf{E}[\Delta_1^L | Z_{\nu_0} \sim \boldsymbol{\pi}^L] + \mu^U).$$

Remark (3.2) yields that

$$\mathbf{E}[\Delta_1^L | Z_{\nu_0} \sim \boldsymbol{\pi}^L] = \sum_{k \in S_L} \sum_{l \in S_U} \mathbf{E}[\Delta_1^L | Z_{\tau_1} = l] P(Z_{\tau_1} = l | Z_{\nu_0} = k) \pi_k^L = \mu^L.$$

We now prove the first part of (ii). Since, conditionally on $Z_{\nu_0} = i$, Δ_1^U and $\Delta_{T_{i,1}^L+1}^U$ have the same distribution, using (i) we have that

$$\mathbf{Var}[S_{T_{i,1}^L}^U | Z_{\nu_0} = i] = \mathbf{E} \left[\left(\sum_{j=1}^{T_{i,1}^L} (\Delta_{j+1}^U - \mu^U) \right)^2 | Z_{\nu_0} = i \right] = \mathbf{E} \left[\sum_{j=1}^{T_{i,1}^L} (\Delta_{j+1}^U - \mu^U)^2 | Z_{\nu_0} = i \right] + 2C_i^U,$$

$$\text{where } C_i^U := \mathbf{E} \left[\sum_{j=1}^{T_{i,1}^L} (\Delta_{j+1}^U - \mu^U) \sum_{l=j+1}^{T_{i,1}^L} (\Delta_{l+1}^U - \mu^U) | Z_{\nu_0} = i \right].$$

Next, we apply Proposition 1.69 of Serfozo (2009) with $X_j = Z_{\nu_j}$, $\pi = \boldsymbol{\pi}^L = \{\pi_i^L\}_{i \in S_L}$ and $V_j = (\Delta_{j+1}^U - \mu^U)^2$ and obtain that

$$\mathbf{E} \left[\sum_{j=1}^{T_{i,1}^L} (\Delta_{j+1}^U - \mu^U)^2 | Z_{\nu_0} = i \right] = (\pi_i^L)^{-1} \mathbf{E}[(\Delta_1^U - \mu^U)^2 | Z_{\nu_0} \sim \boldsymbol{\pi}^L].$$

Then, we compute

$$\begin{aligned} C_i^U &= \mathbf{E} \left[\sum_{j=1}^{\infty} \mathbf{I}_{[T_{i,1}^L \geq j]} \mathbf{E} \left[(\Delta_{j+1}^U - \mu^U) \sum_{l=j+1}^{T_{i,1}^L} (\Delta_{l+1}^U - \mu^U) | T_{i,1}^L \geq j, Z_{\nu_0} = i \right] | Z_{\nu_0} = i \right] \\ &= \mathbf{E} \left[\sum_{j=1}^{\infty} \mathbf{I}_{[T_{i,1}^L \geq j]} \sum_{k \in S_L} \mathbf{I}_{[Z_{\nu_j} = k]} g^L(k) | Z_{\nu_0} = i \right] \\ &= \sum_{k \in S_L} g^L(k) \mathbf{E} \left[\sum_{j=1}^{T_{i,1}^L} \mathbf{I}_{[Z_{\nu_j} = k]} | Z_{\nu_0} = i \right], \end{aligned}$$

where $g^L : S_L \rightarrow \mathbb{R}$ is given by

$$\begin{aligned} g^L(k) &= \mathbf{E} \left[(\Delta_{j+1}^U - \mu^U) \sum_{l=j+1}^{T_{i,1}^L} (\Delta_{l+1}^U - \mu^U) | T_{i,1}^L \geq j, Z_{\nu_j} = k, Z_{\nu_0} = i \right] \\ &= \sum_{l=j+1}^{\infty} \mathbf{E} \left[(\Delta_{j+1}^U - \mu^U) (\Delta_{l+1}^U - \mu^U) \mathbf{I}_{[T_{i,1}^L \geq l]} | T_{i,1}^L \geq j, Z_{\nu_j} = k, Z_{\nu_0} = i \right]. \end{aligned}$$

Using Theorem 1.54 of Serfozo (2009), we obtain $\mathbf{E}[\sum_{j=1}^{T_{i,1}^L} \mathbf{I}_{[Z_{\nu_j}=k]} | Z_{\nu_0} = i] = (\pi_i^L)^{-1} \pi_k^L$, which yields

$$C_i^U = (\pi_i^L)^{-1} \sum_{k \in S_L} g^L(k) \pi_k^L.$$

Now, using Lemma 4.1, we see that, conditionally on $j \leq T_{i,1}^L < l$, $(\Delta_{l+1}^U - \mu^U)$ is independent of $(\Delta_{j+1}^U - \mu^U)$. If $Z_{\nu_j} \sim \pi^L$, then using stationarity (see Remark 3.1)

$$\mathbf{E}[(\Delta_{l+1}^U - \mu^U) | j \leq T_{i,1}^L < l, Z_{\nu_j} \sim \pi^L] = \mathbf{E}[(\Delta_{l+1}^U - \mu^U) | Z_{\nu_l} \sim \pi^L] = 0.$$

Therefore,

$$\begin{aligned} & \sum_{l=j+1}^{\infty} \sum_{k \in S_L} \pi_k^L \mathbf{E} \left[(\Delta_{j+1}^U - \mu^U) (\Delta_{l+1}^U - \mu^U) \mathbf{I}_{[T_{i,1}^L < l]} | T_{i,1}^L \geq j, Z_{\nu_j} = k, Z_{\nu_0} = i \right] \\ = & \sum_{l=j+1}^{\infty} \mathbf{E} \left[(\Delta_{j+1}^U - \mu^U) \mathbf{E}[(\Delta_{l+1}^U - \mu^U) | j \leq T_{i,1}^L < l, Z_{\nu_j} \sim \pi^L] \mathbf{I}_{[T_{i,1}^L < l]} | T_{i,1}^L \geq j, Z_{\nu_j} \sim \pi^L, Z_{\nu_0} = i \right] = 0. \end{aligned}$$

Adding the above to $\sum_{k \in S_L} g^L(k) \pi_k^L$ we conclude that

$$\begin{aligned} \sum_{k \in S_L} g^L(k) \pi_k^L &= \sum_{l=j+1}^{\infty} \sum_{k \in S_L} \pi_k^L \mathbf{E} \left[(\Delta_{j+1}^U - \mu^U) (\Delta_{l+1}^U - \mu^U) | T_{i,1}^L \geq j, Z_{\nu_j} = k, Z_{\nu_0} = i \right] \\ &= \sum_{l=1}^{\infty} \sum_{k \in S_L} \pi_k^L \mathbf{E} \left[(\Delta_1^U - \mu^U) (\Delta_{l+1}^U - \mu^U) | Z_{\nu_0} = k \right] \\ &= \sum_{l=1}^{\infty} \mathbf{Cov}[\Delta_1^U, \Delta_{l+1}^U | Z_{\nu_0} \sim \pi^L]. \end{aligned}$$

The second part of (ii) and (iii) are obtained similarly. \square

B Numerical experiments

In this section we describe numerical experiments to illustrate the evolution of the process under different distributional assumptions. We also study the empirical distribution of the lengths of supercritical and subcritical regimes and illustrate how the process changes when U_j and L_j exhibit an increasing trend. We emphasize that these experiments illustrate the behavior of the estimates of the parameters of the BPRE T when using a finite number of generations in a single synthetic dataset. In the numerical Experiments 1-4 below, we set $L_0 = 10^2$, $L_U = 10^4$, $n \in \{0, 1, \dots, 10^4\}$, $U_j \sim L_U + \text{Zeta}(3)$, $L_j \sim \text{Unif}_d(L_0, 10L_0)$, and different distributions for Z_0 , I_0^U , $\xi_{1,0}^U$, and $\xi_{1,0}^L$ as follows:

	$Z_0 - L_0$	I_0^U	$\xi_{1,0}^U$	$\xi_{1,0}^L$
Exp. 1	$\text{Pois}(1; L_U - L_0)$	$\text{Pois}(\Lambda_I),$ $\Lambda_I \sim \text{Unif}(0, 10)$	$\text{Pois}(\Lambda_U),$ $\Lambda_U \sim \text{Unif}(0.9, 2.1)$	$\text{Pois}(\Lambda_L),$ $\Lambda_L \sim \text{Unif}(0.5, 1.1)$
Exp. 2	$\text{Pois}(1; L_U - L_0)$	$\text{Pois}(\Lambda_I),$ $\Lambda_I \sim \text{Gamma}(5, 1)$	$\text{Pois}(\Lambda_U),$ $\Lambda_U \sim \text{Gamma}(2, 1)$	$\text{Pois}(\Lambda_L),$ $\Lambda_L \sim \text{Gamma}(0.8, 1)$
Exp. 3	$\text{Nbin}(1, 1; L_U - L_0)$	$\text{Nbin}(R_I, M_I),$ $R_I \sim 1 + \text{Pois}(1),$ $M_I \sim \text{Unif}(0, 10)$	$\text{Nbin}(R_U, M_U),$ $R_U \sim 1 + \text{Pois}(1),$ $M_U \sim \text{Unif}(0.9, 2.1)$	$\text{Nbin}(R_L, M_L),$ $R_L \sim 1 + \text{Pois}(1),$ $M_L \sim \text{Unif}(0.5, 1.1)$
Exp. 4	$\text{Nbin}(1, 1; L_U - L_0)$	$\text{Nbin}(R_I, M_I),$ $R_I \sim 1 + \text{Pois}(1),$ $M_I \sim \text{Gamma}(5, 1)$	$\text{Nbin}(R_U, M_U),$ $R_U \sim 1 + \text{Pois}(1),$ $M_U \sim \text{Gamma}(2, 1)$	$\text{Nbin}(R_L, M_L),$ $R_L \sim 1 + \text{Pois}(1),$ $M_L \sim \text{Gamma}(0.8, 1)$

In the above description, we have used the notation $\text{Unif}(a, b)$ for the uniform distribution over the interval (a, b) and $\text{Unif}_d(a, b)$ for the uniform distribution over integers between a and b . $\text{Zeta}(s)$ is the zeta distribution with exponent $s > 1$. $\text{Pois}(\lambda)$ is the Poisson distribution with parameter λ while $\text{Pois}(\lambda; b)$ is the Poisson distribution truncated to values not larger than b . Similarly, $\text{Nbin}(r, m)$ is the negative binomial distribution with predefined number of successful trials r and mean m while $\text{Nbin}(r, m; b)$ is the negative binomial distribution truncated to values not larger than b . Finally, $\text{Gamma}(\alpha, \beta)$ is the Gamma distribution with shape parameter α and rate parameter β . In these experiments, there were between 400 to 700 crossing of the thresholds depending on the distributional assumptions. The results of the numerical Experiments 1-4 are shown in Figure 2.

We next turn our attention to construction of confidence intervals for the means in the supercritical and subcritical regimes. Values of $\mathbf{E}[m(\zeta_0^T)]$, $\mathbf{Var}[m(\zeta_0^T)]$, and $\mathbf{E}[\sigma^2(\zeta_0^T)]$ in Exp. 1-4 can be deduced from the underlying distributions and are summarized below.

	$\mathbf{E}[m(\zeta_0^U)]$	$\mathbf{Var}[m(\zeta_0^U)]$	$\mathbf{E}[\sigma^2(\zeta_0^U)]$	$\mathbf{E}[m(\zeta_0^L)]$	$\mathbf{Var}[m(\zeta_0^L)]$	$\mathbf{E}[\sigma^2(\zeta_0^L)]$
Exp. 1	1.5	0.12	1.5	0.8	0.03	0.8
Exp. 2	2	2	2	0.8	0.8	0.8
Exp. 3	1.5	0.12	≈ 2.998	0.8	0.03	≈ 1.224
Exp. 4	2	2	≈ 5.793	0.8	0.8	≈ 1.710

In the next table, we provide the estimators \hat{m}_n^U , $C_n^U/\tilde{N}^L(n)$, $\tilde{C}_n^U/\tilde{N}^L(n)$, and $\tilde{A}_n^U/\tilde{N}^L(n)$ of $\mathbf{E}[m(\zeta_0^U)]$, μ^U , $\tilde{\mu}^U$, and \tilde{a}^U , respectively. Notice that

$$\hat{\sigma}_{m,n}^{2,U} := \frac{1}{\tilde{C}_n^U} \sum_{j=1}^n \left(\frac{Z_j - I_{j-1}^U}{Z_{j-1}} - \hat{m}_n^U \right)^2 \tilde{\chi}_{j-1}^U \quad \text{and} \quad \hat{\sigma}_n^{2,U} := \frac{1}{\tilde{C}_n^U} \sum_{j=1}^n \frac{1}{Z_{j-1}} \sum_{i=1}^{Z_{j-1}} \left(\xi_{j-1,i}^U - \frac{Z_j - I_{j-1}^U}{Z_{j-1}} \right)^2 \tilde{\chi}_{j-1}^U$$

are used to estimate $\mathbf{Var}[m(\zeta_0^U)]$ and $\mathbf{E}[\sigma^2(\zeta_0^U)]$. Similar to the proof of Theorem 2.6 it is easy to see that $\hat{\sigma}_{m,n}^{2,U}$ and $\hat{\sigma}_n^{2,U}$ are consistent estimators of $\mathbf{Var}[m(\zeta_0^U)]$ and $\mathbf{E}[\sigma^2(\zeta_0^U)]$. Similar comments hold when U is replaced by L .

	\hat{m}_n^U	$\hat{\sigma}_{m,n}^{2,U}$	$\hat{\sigma}_n^{2,U}$	$C_n^U/\tilde{N}^L(n)$	$\tilde{C}_n^U/\tilde{N}^L(n)$	$\tilde{A}_n^U/\tilde{N}^L(n)$
Exp. 1	1.496	0.122	1.499	9.282	9.282	0.010
Exp. 2	2.009	2.051	2.003	10.807	10.804	0.105
Exp. 3	1.507	0.119	3.016	9.069	9.069	0.011
Exp. 4	2.018	2.157	5.929	10.899	10.893	0.112
	\hat{m}_n^L	$\hat{\sigma}_{m,n}^{2,L}$	$\hat{\sigma}_n^{2,L}$	$C_n^L/\tilde{N}^U(n)$	$\tilde{C}_n^L/\tilde{N}^U(n)$	$\tilde{A}_n^L/\tilde{N}^U(n)$
Exp. 1	0.799	0.031	0.800	14.063	14.063	0.009
Exp. 2	0.809	0.805	0.809	5.118	5.118	0.002
Exp. 3	0.799	0.030	1.214	14.028	14.028	0.010
Exp. 4	0.822	0.945	1.869	5.136	5.136	0.002

Using the above estimators in Theorem 2.6 we obtain the following confidence intervals for \hat{m}_n^U and \hat{m}_n^L . We also provide confidence intervals for the estimator \hat{m}_n defined below, which *does not take into account different regimes*. Specifically,

$$\hat{m}_n := \frac{1}{\sum_{j=1}^n \mathbf{I}_{[Z_{j-1} \geq 1]}} \sum_{j=1}^n \frac{Z_j - I_{j-1}^T}{Z_{j-1}} \mathbf{I}_{[Z_{j-1} \geq 1]},$$

where I_{j-1}^T is equal to I_{j-1}^U if $T = U$ and 0 otherwise.

	95% CI using \hat{m}_n^U	95% CI using \hat{m}_n^L	95% CI using \hat{m}_n
Exp. 1	(1.485, 1.507)	(0.795, 0.804)	(1.068, 1.085)
Exp. 2	(1.975, 2.043)	(0.778, 0.840)	(1.596, 1.651)
Exp. 3	(1.496, 1.518)	(0.794, 0.803)	(1.068, 1.085)
Exp. 4	(1.982, 2.053)	(0.788, 0.855)	(1.607, 1.664)

Next, we investigate the behavior of the process when the thresholds L_j and U_j increase with j . To this end, we let $L_0 = 10^2$, $L_U = 10^4$, and $n \in \{0, 1, \dots, 10^3\}$ and take initial distribution Z_0 , immigration distribution I_0^U , and offspring distributions $\xi_{1,0}^U$ and $\xi_{1,0}^L$ as in the numerical Experiment 1. We consider four different distributions for L_j and U_j as follows:

	$U_j - L_U$	L_j
Exp. 5	Zeta(3)	$\text{Unif}_d(L_0, 10L_0)$
Exp. 6	Zeta(3)	$\text{Unif}_d(L_{j,1}, L_{j,2})$, where $L_{j,1} = \min(L_0 + 100(j-1), L_U)$, $L_{j,2} = \min(10L_0 + 100(j-1), L_U)$
Exp. 7	Zeta(3) + 500(j-1)	$\text{Unif}_d(L_0, 10L_0)$
Exp. 8	Zeta(3) + 500(j-1)	$\text{Unif}_d(L_{j,1}, L_{j,2})$, where $L_{j,1} = \min(L_0 + 100(j-1), L_U)$, $L_{j,2} = \min(10L_0 + 100(j-1), L_U)$

The results of the numerical Experiment 5-8 are shown in Figure 3. From the plots, we see that the number of cases after crossing i.i.d. the upper thresholds are between 10^4 and $2 \cdot 10^4$, whereas




Figure 2: In each column the numerical Experiments 1, 2, 3, and 4. In the first row, the process Z_n for $n = 10^4 - 10^2, \dots, 10^4$. In the second and third row, the empirical probability distribution of $\{\nu_j - \tau_j\}$ and $\{\tau_j - \nu_{j-1}\}$, respectively.




Figure 3: In black the process Z_n for $n \in \{0, 1, \dots, 10^3\}$, in red horizontal lines at L_0 and L_U , and in blue the thresholds U_j and L_j . From left to right, the numerical Experiments 5, 6, 7, and 8.

when the thresholds increase they almost reach the $6 \cdot 10^4$ mark. Also, the number of regimes up to time $n = 10^3$ reduces as it takes more time to reach a larger threshold. As a consequence the overall number of cases also increases.