

Multi-Horseshoe Dense Property and Intermediate Entropy Property of Ergodic Measures with Same Level

Yiwei Dong*, Xiaobo Hou[§] and Xueting Tian^{† ‡}

*School of Mathematical Science, Fudan University
Shanghai 200433, People's Republic of China*

*Email:dongyiwei06@gmail.com; [§]Email:20110180003@fudan.edu.cn; [†]Email:xuetingtian@fudan.edu.cn

Abstract

Katok conjectured that for every C^2 diffeomorphism f on a Riemannian manifold X , the set $\{h_\mu(f) : \mu \text{ is an ergodic measure for } (X, f)\}$ includes $[0, h_{\text{top}}(f))$. In this paper we obtained a refined Katok's conjecture on intermediate metric entropies of ergodic measures with same level that for a homeomorphism of a compact metric space which is expansive, transitive and has the shadowing property, one has

$$\{h_\mu(f) : \mu \in M_{\text{erg}}(f, X) \text{ and } \int \varphi d\mu = a\} = \{h_\mu(f) : \mu \in M(f, X) \text{ and } \int \varphi d\mu = a\}$$

for any $a \in (\inf_{\mu \in M(f, X)} \int \varphi d\mu, \sup_{\mu \in M(f, X)} \int \varphi d\mu)$ and any continuous function φ with bounded variation. In this process, we establish 'multi-horseshoe' entropy-dense property and use it to get the goal combined with L. Barreira and P. Doutor's conditional variational principle.

1 Introduction

Let (X, d) be a nondegenerate (i.e. with at least two points) compact metric space, and $f : X \rightarrow X$ be a continuous map. Such (X, f) is called a dynamical system. For a dynamical system (X, f) , let $M(X)$, $M(f, X)$, $M_{\text{erg}}(f, X)$ denote the space of probability measures, f -invariant, f -ergodic probability measures, respectively. Let ρ be a metric for the weak*-topology on $M(X)$. We denote the sets of natural numbers, integer numbers and nonnegative numbers by \mathbb{N} , \mathbb{Z} , \mathbb{Z}^+ respectively.

It is believed that in most cases, positive topological entropy of (X, f) implies a rich structure of $M(f, X)$. In [24] Katok showed that every $C^{1+\alpha}$ diffeomorphism in dimension 2 has horseshoes of large entropies. This implies that the system has ergodic measures of arbitrary intermediate metric entropies, that is $\{h_\mu(f) : \mu \text{ is an ergodic measure}\}$ includes $[0, h_{\text{top}}(f))$, where $h_{\text{top}}(f)$ is the topological entropy of (X, f) , and $h_\mu(f)$ is the metric entropy of μ . Katok believed that this holds in any dimension.

Conjecture 1. (Katok) For every C^2 diffeomorphism f on a Riemannian manifold X , the set

$$\{h_\mu(f) : \mu \text{ is an ergodic measure for } (X, f)\}$$

includes $[0, h_{\text{top}}(f))$.

In the last decade, a number of partial results have been obtained. See [40, 37, 13, 21, 26, 28, 27, 38]. In the recent works, we find shadowing property is a powerful tool to construct ergodic measures. It is shown in [27] that for a transitive dynamical system with the shadowing property, if the entropy function is upper semi-continuous, then for every $0 \leq c < h_{\text{top}}(f)$ the set of ergodic measures with

[‡] Tian is the corresponding author.

Key words and phrases: Shadowing property; Metric entropy; Horseshoe; Symbolic dynamics.

AMS Review: 37C50; 37A35; 37D20; 37B10.

entropy c is residual in the space of invariant measures with entropy at least c . This implies the set $\{h_\mu(f) : \mu \text{ is an ergodic measure for } (X, f)\}$ includes $[0, h_{\text{top}}(f))$.

In this paper we would like to explore refined Katok's conjecture. Let $\varphi : X \rightarrow \mathbb{R}$ be a continuous function. Denote

$$L_\varphi = \left[\inf_{\mu \in M(f, X)} \int \varphi d\mu, \sup_{\mu \in M(f, X)} \int \varphi d\mu \right]$$

and

$$\text{Int}(L_\varphi) = \left(\inf_{\mu \in M(f, X)} \int \varphi d\mu, \sup_{\mu \in M(f, X)} \int \varphi d\mu \right).$$

Similar as Katok's conjecture, we would like to consider the following refined question.

Question 1.1. For every typical diffeomorphism f on a Riemannian manifold X , every continuous function φ on X , and every $a \in \text{Int}(L_\varphi)$, whether one has

$$\{h_\mu(f) : \mu \in M_{\text{erg}}(f, X) \text{ and } \int \varphi d\mu = a\} = \{h_\mu(f) : \mu \in M(f, X) \text{ and } \int \varphi d\mu = a\}?$$

In the present paper, we give partial answer to this question. We say that a continuous function φ has *bounded variation* if there exists $\varepsilon > 0$ for which $\sup_{n \in \mathbb{N}} \gamma_n(\varphi, \varepsilon) < \infty$ with

$$\gamma_n(\varphi, \varepsilon) = \sup \left\{ \left| \sum_{k=0}^{n-1} \varphi(f^k(x)) - \sum_{k=0}^{n-1} \varphi(f^k(y)) \right| : d(f^k(x), f^k(y)) < \varepsilon \text{ for } k = 0, \dots, n-1 \right\}.$$

For any $a \in L_\varphi$, define

$$M_\varphi(a) = \{\mu \in M(f, X) : \int \varphi d\mu = a\}, \quad M_\varphi^{\text{erg}}(a) = \{\mu \in M_{\text{erg}}(f, X) : \int \varphi d\mu = a\}.$$

Then $M_\varphi(a)$ is closed in $M(f, X)$, and thus $\sup\{h_\mu(f) : \mu \in M_\varphi(a)\} = \max\{h_\mu(f) : \mu \in M_\varphi(a)\}$ for any $a \in L_\varphi$ if the entropy function is upper semi-continuous. We denote the support of a probability measure μ by $S_\mu := \{x \in M : \mu(U) > 0 \text{ for any neighborhood } U \text{ of } x\}$. Now we state our main result as follows.

Theorem A. *Suppose that a homeomorphism $f : X \rightarrow X$ of a compact metric space is transitive, expansive, and has the shadowing property. Let φ be a continuous function on X with bounded variation. Then:*

- (I) *For any $a \in \text{Int}(L_\varphi)$, any $\mu \in M_\varphi(a)$ and any $\eta, \zeta > 0$, there is $\nu \in M_\varphi^{\text{erg}}(a)$ such that $\rho(\nu, \mu) < \zeta$ and $|h_\nu(f) - h_\mu(f)| < \eta$.*
- (II) *For any $a \in \text{Int}(L_\varphi)$, any $\mu \in M_\varphi(a)$, any $0 \leq c \leq h_\mu(f)$ and any $\eta, \zeta > 0$, there is $\nu \in M_\varphi^{\text{erg}}(a)$ such that $\rho(\nu, \mu) < \zeta$ and $|h_\nu(f) - c| < \eta$.*
- (III) *For any $a \in \text{Int}(L_\varphi)$ and $0 \leq c < \max\{h_\mu(f) : \mu \in M_\varphi(a)\}$, the set $\{\mu \in M_\varphi^{\text{erg}}(a) : h_\mu(f) = c, S_\mu = X\}$ is residual in $\{\mu \in M_\varphi(a) : h_\mu(f) \geq c\}$.*
- (IV) $\{(\int \varphi d\mu, h_\mu(f)) : \mu \in M(f, X), \int \varphi d\mu \in \text{Int}(L_\varphi)\} = \{(\int \varphi d\mu, h_\mu(f)) : \mu \in M_{\text{erg}}(f, X), \int \varphi d\mu \in \text{Int}(L_\varphi)\} = \{(a, b) : a \in \text{Int}(L_\varphi), 0 \leq b \leq \max\{h_\mu(f) : \mu \in M_\varphi(a)\}\}.$

The hyperbolic flow and singular hyperbolic flow versions of Theorem A are being prepared [22]. Readers can see the dynamical systems and functions to which Theorem A is applicable in section 6.

Remark 1.2. (1) In [18] Eizenberg, Kifer and Weiss proved for systems with the specification property that any f -invariant probability measure μ is the weak limit of a sequence of ergodic measures $\{\mu_n\}_{n=1}^\infty$, such that the entropy of μ is the limit of the entropies of the $\{\mu_n\}_{n=1}^\infty$. Pfister and Sullivan referred to this property as the entropy-dense property [36] and proved that the entropy-dense property holds for systems with the approximate product property. From Theorem A(I) if a homeomorphism $f : X \rightarrow X$ of


a compact metric space is expansive, transitive, and has the shadowing property, then the entropy-dense property holds in $M_\varphi(a)$ for any $a \in \text{Int}(L_\varphi)$ and any continuous function φ with bounded variation.

(2) Li and Oprocha proved in [27] that for a transitive dynamical system with the shadowing property, for every invariant measure μ and every $0 \leq c \leq h_\mu(f)$, there exists a sequence of ergodic measures $\{\mu_n\}_{n=1}^\infty$ such that $\lim_{n \rightarrow \infty} \mu_n = \mu$ and $\lim_{n \rightarrow \infty} h_{\mu_n}(f) = c$. If further the entropy function is upper semi-continuous, they proved that for every $0 \leq c < h_{\text{top}}(f)$ the set of ergodic measures with entropy c is residual in the space of invariant measures with entropy at least c . From Theorem A(II) and (III) we obtain more refined results for a homeomorphism $f : X \rightarrow X$ of a compact metric space which is expansive, transitive and has the shadowing property, that is for a continuous function φ with bounded variation, for any $a \in \text{Int}(L_\varphi)$, any $\mu \in M_\varphi(a)$, and any $0 \leq c \leq h_\mu(f)$, there exists a sequence of ergodic measures $\{\mu_n\}_{n=1}^\infty \subseteq M_\varphi(a)$ such that $\lim_{n \rightarrow \infty} \mu_n = \mu$ and $\lim_{n \rightarrow \infty} h_{\mu_n}(f) = c$, and for any $a \in \text{Int}(L_\varphi)$ and $0 \leq c < \max\{h_\mu(f) : \mu \in M_\varphi(a)\}$, in $M_\varphi(a)$ the set of ergodic measures with entropy c and full support is residual in the space of invariant measures with entropy at least c .

(3) From Theorem A(IV) we give partial answer to Question 1.1 for a homeomorphism $f : X \rightarrow X$ of a compact metric space which is expansive, transitive and has the shadowing property, that is $\{h_\mu(f) : \mu \in M_\varphi^{erg}(a)\} = \{h_\mu(f) : \mu \in M_\varphi(a)\}$ for any $a \in \text{Int}(L_\varphi)$ and any continuous function φ with bounded variation.

For a continuous function without the assumption of bounded variation, we don't know how to obtain the corresponding results as Theorem A. In the proof of Theorem A, there are two keypoints: 'multi-horseshoe' entropy-dense property (see Theorem 2.6) and conditional variational principle (see Theorem 3.7) proved by L. Barreira and P. Doutor in [5].

Figure 1: Graph of $(\int \varphi d\mu, h_\mu(f))$



Finally, we draw the graph of $(\int \varphi d\mu, h_\mu(f))$. Theorem A shows that every point in the region between the two dotted lines can be attained by ergodic measures.

Organization of this paper. The remainder of this paper is organized as follows. In Section 2 we introduce 'multi-horseshoe' entropy-dense property and prove that it holds for topologically Anosov system which is transitive. In Section 3 and 4, using conditional variational principle (see Theorem 3.7) proved by L. Barreira and P. Doutor in [5], we give abstract conditions on which the results of Theorem A hold in the more general context of almost additive sequences of continuous functions. In Section 5 by 'multi-horseshoe' entropy-dense property we show that the abstract conditions given in Section 3 and 4 are satisfied for topologically Anosov system which is transitive, and thus we obtain Theorem A. In Section 6, we apply the results in the previous sections to transitive locally maximal hyperbolic sets and transitive two-side subshifts of finite type. Finally, we consider homoclinic classes and give corresponding results on refined Katok's conjecture.

2 'Multi-Horseshoe' Entropy-Dense Property

In this section, we introduce 'multi-horseshoe' entropy-dense property which shall serve for our needs in the future. Eizenberg, Kifer and Weiss proved for systems with the specification property that [18] any f -invariant probability measure ν is the weak limit of a sequence of ergodic measures $\{\nu_n\}$, such that the entropy of ν is the limit of the entropies of the $\{\nu_n\}$. This is a central point in large deviations theory, which was first emphasized in [20]. Meanwhile, this also plays an crucial part in the computing of Billingsley dimension [6, 7] on shift spaces [35]. Pfister and Sullivan refered to this property as the *entropy-dense* property [36].

Definition 2.1. We say (X, f) satisfies the *entropy-dense property* (or $M_{erg}(f, X)$ is *entropy-dense* in $M(f, X)$), if for any $\mu \in M(f, X)$, for any neighborhood G of μ in $M(X)$, and for any $\eta > 0$, there exists a $\nu \in M_{erg}(f, X)$ such that $h_\nu(f) > h_\mu(f) - \eta$ and $\nu \in G$.

Definition 2.2. We say (X, f) satisfies the *refined entropy-dense property* (or $M_{erg}(f, X)$ is *refined entropy-dense* in $M(f, X)$), if for any $\mu \in M(f, X)$, for any neighborhood G of μ in $M(X)$, and for any $\eta > 0$, there exists a closed f -invariant set $\Lambda_\mu \subseteq X$ such that $M(f, \Lambda_\mu) \subseteq G$ and $h_{top}(f, \Lambda_\mu) > h_\mu(f) - \eta$. By classical variational principle, it is equivalent that for any neighborhood G of μ in $M(X)$, and for any $\eta > 0$, there exists a $\nu \in M_{erg}(f, X)$ such that $h_\nu(f) > h_\mu(f) - \eta$ and $M(f, S_\nu) \subseteq G$.

Of course, refined entropy-dense \Rightarrow entropy-dense \Rightarrow ergodic measures are dense in the space of invariant measures. For systems with the approximate product property, Pfister and Sullivan in fact had obtained the refined entropy-dense properties by [36, Proposition 2.3]. Note that if a dynamical system is transitive and has the shadowing property, then it has approximate product property by their definitions. Then we have

Proposition 2.3. *Suppose that (X, f) is topologically transitive and satisfies the shadowing property. Then (X, f) has the refined entropy-dense property.*

For C^r diffeomorphisms, a theorem by A. Katok [24] asserts that any ergodic hyperbolic measure can be approximated by a horseshoe. Here for topological dynamical systems we introduce 'multi-horseshoe' entropy-dense property and show that it holds for topologically Anosov system which is transitive.

Definition 2.4. A homeomorphism $f : X \rightarrow X$ of a compact metric space called *topologically hyperbolic* or *topologically Anosov*, if X has infinitely many points, (X, f) is expansive and satisfies the shadowing property.

For any $m \in \mathbb{N}$ and $\{\nu_i\}_{i=1}^m \subseteq M(X)$, we write $\text{cov}\{\nu_i\}_{i=1}^m$ for the convex combination of $\{\nu_i\}_{i=1}^m$, namely,

$$\text{cov}\{\nu_i\}_{i=1}^m = \text{cov}(\nu_1, \dots, \nu_m) := \left\{ \sum_{i=1}^m t_i \nu_i : t_i \in [0, 1], 1 \leq i \leq m \text{ and } \sum_{i=1}^m t_i = 1 \right\}.$$

Definition 2.5. We say (X, f) satisfies the '*multi-horseshoe*' entropy-dense property (abbrev. '*multi-horseshoe*' dense property) if for any $K = \text{cov}\{\mu_i\}_{i=1}^m \subseteq M(f, X)$, any $x \in X$ and any $\eta, \zeta, \varepsilon > 0$, there exist compact invariant subsets $\Lambda_i \subseteq \Lambda \subsetneq X$ such that for each $1 \leq i \leq m$

1. $(\Lambda_i, f|_{\Lambda_i})$ and $(\Lambda, f|_{\Lambda})$ conjugate to transitive subshifts of finite type (and thus they are transitive and topologically Anosov).
2. $h_{top}(f, \Lambda_i) > h_{\mu_i}(f) - \eta$ and consequently, $h_{top}(f, \Lambda) > \sup\{h_\kappa(f) : \kappa \in K\} - \eta$.
3. $d_H(K, M(f, \Lambda)) < \zeta$, $d_H(\mu_i, M(f, \Lambda_i)) < \zeta$.
4. There is a positive integer L such that for any z in Λ_i or Λ one has $f^{j+mL}(z) \in B(x, \varepsilon)$ for some $0 \leq j \leq L - 1$ and any $m \in \mathbb{Z}$.

Theorem 2.6. *Suppose (X, f) is topologically Anosov and transitive. Then (X, f) satisfies the '*multi-horseshoe*' dense property.*

2.1 Some definitions

2.1.1 The space of probability measures

Consider a topological dynamical system (X, f) . The space of Borel probability measures on X is denoted by $M(X)$ and the set of continuous functions on X by $C(X)$. We endow $\varphi \in C(X)$ the norm $\|\varphi\| = \max\{|\varphi(x)| : x \in X\}$. Let $\{\varphi_j\}_{j \in \mathbb{N}}$ be a dense subset of $C(X)$, then

$$\rho(\xi, \tau) = \sum_{j=1}^{\infty} \frac{|\int \varphi_j d\xi - \int \varphi_j d\tau|}{2^j \|\varphi_j\|}$$

defines a metric on $M(X)$ for the *weak** topology [41]. Then we denote the Hausdorff distance between two nonempty subsets of $M(X)$, A and B , by

$$d_H(A, B) := \max \left\{ \sup_{x \in A} \inf_{y \in B} \rho(x, y), \sup_{y \in B} \inf_{x \in A} \rho(y, x) \right\}.$$

For $\nu \in M(X)$ and $r > 0$, we denote a ball in $M(X)$ centered at ν with radius r by

$$\mathcal{B}(\nu, r) := \{\rho(\nu, \mu) < r : \mu \in M(X)\}.$$

One notices that

$$\rho(\xi, \tau) \leq 2 \text{ for any } \xi, \tau \in M(X). \quad (2.1)$$

It is also well known that the natural imbedding $j : x \mapsto \delta_x$ is continuous. Since X is compact and $M(X)$ is Hausdorff, one sees that there is a homeomorphism between X and its image $j(X)$. Therefore, without loss of generality we will assume that

$$d(x, y) = \rho(\delta_x, \delta_y). \quad (2.2)$$

For $x \in X$ and $\varepsilon > 0$, we denote a ball in X centered at x with radius ε by

$$B(x, \varepsilon) := \{d(x, y) < \varepsilon : y \in X\}.$$

A straight calculation using (2.1) and (2.2) gives

Lemma 2.7. *For any $\varepsilon > 0, \delta > 0$, and any two sequences $\{x_i\}_{i=0}^{n-1}, \{y_i\}_{i=0}^{n-1}$ of X , if $d(x_i, y_i) < \varepsilon$ holds for any $i \in [0, n-1]$, then for any $J \subseteq \{0, 1, \dots, n-1\}$ with $\frac{n-|J|}{n} < \delta$, one has:*

- (a) $\rho(\frac{1}{n} \sum_{i=0}^{n-1} \delta_{x_i}, \frac{1}{n} \sum_{i=0}^{n-1} \delta_{y_i}) < \varepsilon$.
- (b) $\rho(\frac{1}{n} \sum_{i=0}^{n-1} \delta_{x_i}, \frac{1}{|J|} \sum_{i \in J} \delta_{y_i}) < \varepsilon + 2\delta$.

Lemma 2.7 is easy to be verified and shows us that if any two orbit of x and y in finite steps are close in the most time, then the two empirical measures induced by x, y are also close.

2.1.2 Topological entropy and metric entropy

Now let us to recall the definition of topological entropy in [11] by Bowen. For $x, y \in X$ and $n \in \mathbb{N}$, the Bowen distance between x, y is defined as

$$d_n(x, y) := \max\{d(f^i(x), f^i(y)) : i = 0, 1, \dots, n-1\}$$

and the Bowen ball centered at x with radius $\varepsilon > 0$ is defined as

$$B_n(x, \varepsilon) := \{y \in X : d_n(x, y) < \varepsilon\}.$$

Let $E \subseteq X$, and $\mathcal{G}_n(E, \sigma)$ be the collection of all finite or countable covers of E by sets of the form $B_u(x, \sigma)$ with $u \geq n$. We set

$$C(E; t, n, \sigma, f) := \inf_{\mathcal{C} \in \mathcal{G}_n(E, \sigma)} \sum_{B_u(x, \sigma) \in \mathcal{C}} e^{-tu} \text{ and } C(E; t, \sigma, f) := \lim_{n \rightarrow \infty} C(E; t, n, \sigma, f).$$

Then we define

$$h_{\text{top}}(E; \sigma, f) := \inf\{t : C(E; t, \sigma, f) = 0\} = \sup\{t : C(E; t, \sigma, f) = \infty\}$$

The *Bowen topological entropy* of E is

$$h_{\text{top}}(f, E) := \lim_{\sigma \rightarrow 0} h_{\text{top}}(E; \sigma, f).$$

We call (X, \mathcal{B}, μ) a probability space if \mathcal{B} is a Borel σ -algebra on X and μ is a probability measure on X . For a finite measurable partition $\xi = \{A_1, \dots, A_n\}$ of a probability space (X, \mathcal{B}, μ) , define

$$H_\mu(\xi) = - \sum_{i=1}^n \mu(A_i) \log \mu(A_i).$$

Let $f : X \rightarrow X$ be a continuous map preserving μ . We denote by $\bigvee_{i=0}^{n-1} f^{-i}\xi$ the partition whose element is the set $\bigcap_{i=0}^{n-1} f^{-i}A_{j_i}, 1 \leq j_i \leq n$. Then the following limit exists:

$$h_\mu(f, \xi) = \lim_{n \rightarrow \infty} \frac{1}{n} H_\mu \left(\bigvee_{i=0}^{n-1} f^{-i}\xi \right)$$

and we define the metric entropy of μ as

$$h_\mu(f) := \sup\{h_\mu(f, \xi) : \xi \text{ is a finite measurable partition of } X\}.$$

2.1.3 Transitive, mixing, expansive and shadowing property

If for every pair of non-empty open sets U, V there is an integer n such that $f^n(U) \cap V \neq \emptyset$ then we call (X, f) *topologically transitive*. Furthermore, if for every pair of non-empty open sets U, V there exists an integer N such that $f^n(U) \cap V \neq \emptyset$ for every $n > N$, then we call (X, f) *topologically mixing*. When $f : X \rightarrow X$ is a homeomorphism of a compact metric space, we say that (X, f) is *expansive* if there exists a constant $c > 0$ such that for any $x, y \in X$, $d(f^i(x), f^i(y)) > c$ for some $i \in \mathbb{Z}$. We call c the expansive constant. When $f : X \rightarrow X$ is a homeomorphism of a compact metric space, we say that a subset Y of X is *f*-invariant if $f(Y) = Y$. If Y is a closed *f*-invariant subset of X , then $(Y, f|_Y)$ also is a dynamical system. We will call it a subsystem of (X, f) .

A finite sequence $\mathfrak{C} = \langle x_1, \dots, x_l \rangle, l \in \mathbb{N}$ is called a *chain*. Furthermore, if $d(f(x_i), x_{i+1}) < \varepsilon, 1 \leq i \leq l-1$, we call \mathfrak{C} an ε -chain with length l . For any $m \in \mathbb{N}$, if there are m ε -chains $\mathfrak{C}_i = \langle x_{i,1}, \dots, x_{i,l_i} \rangle, l_i \in \mathbb{N}, 1 \leq i \leq m$ satisfying that $d(f(x_{i,l_i}), x_{i+1,1}) < \varepsilon, 1 \leq i \leq m-1$, then we can concatenate \mathfrak{C}_i s to constitute a new ε -chain

$$\langle x_{1,1}, \dots, x_{1,l_1}, x_{2,1}, \dots, x_{2,l_2}, \dots, x_{m,1}, \dots, x_{m,l_m} \rangle$$

which we denote by $\mathfrak{C}_1 \mathfrak{C}_2 \dots \mathfrak{C}_m$.

Definition 2.8. Suppose $f : X \rightarrow X$ is a homeomorphism of a compact metric space. For any $\delta > 0$, a sequence $\{x_n\}_{n \in \mathbb{Z}}$ is called a δ -pseudo-orbit if $d(f(x_n), x_{n+1}) < \delta$ for $n \in \mathbb{Z}$. $\{x_n\}_{n \in \mathbb{Z}}$ is ε -shadowed by some $y \in X$ if $d(f^n(y), x_n) < \varepsilon$ for any $n \in \mathbb{Z}$. We say that (X, f) has the *shadowing property* if for any $\varepsilon > 0$, there exists $\delta > 0$ such that any δ -pseudo-orbit is ε -shadowed by some point in X .

2.2 Some lemmas

Proposition 2.9. *Suppose $f : X \rightarrow X$ is a homeomorphism of a compact metric space. Consider $\Delta \subseteq X$ which satisfies $f^n(\Delta) = \Delta$ for some $n \in \mathbb{N}$ and let $\Lambda = \bigcup_{i=0}^{n-1} f^i(\Delta)$. If $f^i(\Delta) \cap f^j(\Delta) = \emptyset$ for any $0 \leq i < j \leq n-1$, then*

(1) *if (Δ, f^n) is expansive, then (Λ, f) is expansive;*

(2) if (Δ, f^n) is topologically transitive, then (Λ, f) is topologically transitive;

(3) if (Δ, f^n) has the shadowing property, then (Λ, f) also has the shadowing property.

Proof. Item (1) and (2) come directly from the uniform continuity of f, \dots, f^{m-1} . Since (Δ, f^n) has the shadowing property and $f : X \rightarrow X$ is a homeomorphism, then $(f^i(\Delta), f^n)$ has the shadowing property for any $0 \leq i \leq n-1$. Combining with $f^i(\Delta) \cap f^j(\Delta) = \emptyset$ for any $0 \leq i < j \leq n-1$, (Λ, f^n) also has the shadowing property. Let $k > 0$ be an integer, then from [2, Theorem 2.3.3] a dynamical system (X, f) has the shadowing property if and only if so does (X, f^k) . So (Λ, f) has the shadowing property. \square

For $\varepsilon > 0$ and $n \in \mathbb{N}$, two points x and y are (n, ε) -separated if $d_n(x, y) > \varepsilon$. A subset E is (n, ε) -separated if any pair of different points of E are (n, ε) -separated. For $x \in X$, we define the empirical measure of x as

$$\mathcal{E}_n(x) := \frac{1}{n} \sum_{j=0}^{n-1} \delta_{f^j(x)},$$

where δ_x is the Dirac mass at x . Let $F \subseteq M(X)$ be a neighbourhood of $\nu \in M(f, X)$. Define $X_{n,F} := \{x \in X : \mathcal{E}_n(x) \in F\}$. For $k \in \mathbb{N}$, let $P_k(f) := \{x \in X : f^k(x) = x\}$.

Lemma 2.10. [17, Corollary 4.13] *Suppose (X, f) is topologically Anosov and transitive. Then for any $\eta > 0$, there exists an $\varepsilon_1^* = \varepsilon_1^*(\eta) > 0$ such that for any $\mu \in M(f, X)$ and its neighborhood F_μ , there exists an $0 < \varepsilon_2^* = \varepsilon_2^*(\eta, \mu, F_\mu) < \varepsilon_1^*$ such that for any $x \in X$, for any $0 < \varepsilon \leq \varepsilon_2^*$, for any $N \in \mathbb{N}$, there exist an $n = n(\eta, \mu, F_\mu, \varepsilon, x) \geq N$ such that for any $p \in \mathbb{N}$, there exists an $(pn, \frac{\varepsilon_1^*}{3})$ -separated set Γ_{pn} so that*

(1) $\Gamma_{pn} \subseteq X_{pn, F_\mu} \cap B(x, \varepsilon) \cap P_{pn}(f)$;

(2) $\frac{\log |\Gamma_{pn}|}{pn} > h_\mu(f) - \eta$;

From Lemma 2.10, the set of periodic points is dense in X , and the set of periodic measures is dense in the space of invariant measures. Then by [39, Proposition 6.5], there is an invariant measure with full support. Since measure supported on periodic points has zero metric entropy, then the set of invariant measures with zero metric entropy is dense in the space of invariant measures. So we have the following.

Corollary 2.11. *Suppose (X, f) is topologically Anosov and transitive. Then we have*

(1) $\{\mu \in M(f, X) : h_\mu(f) = 0\}$ is dense in $M(f, X)$.

(2) there is an invariant measure with full support.

2.3 Proof of Theorem 2.6

Fix $K = \text{cov}\{\nu_i\}_{i=1}^m \subseteq M(f, X)$, $x \in X$ and $\eta_0, \zeta_0, \varepsilon_0 > 0$. Let $\rho_0 = \min\{\rho(\nu_i, \nu_j) : 1 \leq i < j \leq m\}$. Then $K \cap M_{\text{erg}}(f, X)$ is empty or finite (less than m). By Proposition 2.3, (X, f) has the refined entropy-dense property, so there are infinitely many ergodic measures on X . This implies that $K \neq M(f, X)$ and thus $d_H(K, M(f, X)) > 0$. Let $\eta, \zeta > 0$ with $\eta \leq \min\{h_{\text{top}}(f), \eta_0\}$ and $\zeta < \min\{d_H(K, M(f, X)), \rho_0, \zeta_0\}$. By the variational principle of the topological entropy, there exists $\nu_0 \in M(f, X)$ such that $h_{\nu_0}(f) > h_{\text{top}}(f) - \frac{\eta}{8}$.

By Lemma 2.10, there exist $\varepsilon^* > 0$ and $0 < \tilde{\varepsilon}^* < \varepsilon^*$ such that for any $0 < \frac{\delta}{2} < \tilde{\varepsilon}^*$ and each $0 \leq i \leq m$, there exists an $n_i \in \mathbb{N}$ such that for any $p \in \mathbb{N}$, there exists an $(pn_i, \frac{\varepsilon^*}{3})$ -separated set $\Gamma_{pn_i}^{\nu_i}$ with

(a) $\Gamma_{pn_i}^{\nu_i} \subseteq P_{pn_i}(f) \cap X_{pn_i, \mathcal{B}(\nu_i, \frac{\zeta}{4})} \cap B(x, \frac{\delta}{2})$;

(b) $\frac{\log |\Gamma_{pn_i}^{\nu_i}|}{pn_i} > h_{\nu_i} - \frac{\eta}{8}$.

We can assume that $\frac{\varepsilon^*}{3} < \frac{c}{4}$ where $c > 0$ is the expansive constant. Let $s(n, \frac{\varepsilon^*}{3})$ denote the largest cardinality of any $(n, \frac{\varepsilon^*}{3})$ -separated set of X , then by [41, Theorem 7.11] one has

$$h_{\text{top}}(f) = \limsup_{n \rightarrow \infty} \frac{1}{n} \log s(n, \frac{\varepsilon^*}{3}). \quad (2.3)$$

Then there exists $N_1 \in \mathbb{N}$ such that for any $n \geq N$, one has

$$s(n, \frac{\varepsilon^*}{3}) < e^{n(h_{\text{top}}(f) + \frac{\eta}{4})}. \quad (2.4)$$

Set $\varepsilon = \min\{\frac{\zeta}{4}, \frac{\rho_0}{6}, \frac{\varepsilon^*}{27}, \frac{\varepsilon_0}{2}\}$. Then there exists a $0 < \delta < \varepsilon$ such that any δ -pseudo-orbit can be ε -shadowed by some point in X . Set $n = p_0 n_0 n_1 n_2 \cdots n_m$ where p_0 is large enough such that for any $1 \leq i \leq m$

$$n \geq 2N, \quad e^{n(h_{\nu_i} - \frac{\eta}{8})} - n \geq e^{n(h_{\nu_i} - \frac{\eta}{4})} \quad \text{and} \quad e^{n(h_{\text{top}}(f) - \frac{\eta}{4})} > \lceil \frac{n}{2} \rceil e^{\lceil \frac{n}{2} \rceil (h_{\text{top}}(f) + \frac{\eta}{4})} + \sum_{m=1}^{N_1-1} |P_m^*(f)| \quad (2.5)$$

where $P_m^*(f)$ is the set of periodic points with minimal period m . Then for each $0 \leq i \leq m$, $P_{n_i}(f) \subseteq P_n(f)$ by definition and furthermore, we can obtain an $(n, \frac{\varepsilon^*}{3})$ -separated set $\Gamma_n^{\nu_i}$ with

(a) $\Gamma_n^{\nu_i} \subseteq P_n(f) \cap X_{n, \mathcal{B}(\nu_i, \frac{\zeta}{4})} \cap B(x, \frac{\delta}{2});$

(b) $\frac{\log |\Gamma_n^{\nu_i}|}{n} > h_{\nu_i} - \frac{\eta}{8}.$

Since periodic points in $\Gamma_n^{\nu_0}$ with same period l_0 for some $l_0 \in \mathbb{N}$ are $(l_0, \frac{\varepsilon^*}{3})$ -separated, by (2.4) and (2.5) we have

$$\begin{aligned} \sum_{m=1}^{\lceil \frac{n}{2} \rceil} |P_m^*(f) \cap \Gamma_n^{\nu_0}| &\leq \sum_{m=N_1}^{\lceil \frac{n}{2} \rceil} s(m, \frac{\varepsilon^*}{3}) + \sum_{m=1}^{N_1-1} |P_m^*(f)| \\ &< \sum_{m=N_1}^{\lceil \frac{n}{2} \rceil} e^{m(h_{\text{top}}(f) + \frac{\eta}{4})} + \sum_{m=1}^{N_1-1} |P_m^*(f)| \\ &\leq \lceil \frac{n}{2} \rceil e^{\lceil \frac{n}{2} \rceil (h_{\text{top}}(f) + \frac{\eta}{4})} + \sum_{m=1}^{N_1-1} |P_m^*(f)| \\ &< e^{n(h_{\text{top}}(f) - \frac{\eta}{4})} < e^{n(h_{\nu_0} - \frac{\eta}{8})} < |\Gamma_n^{\nu_0}|. \end{aligned}$$

Thus there exists $x_0 \in \Gamma_n^{\nu_0}$ with minimal period n . Together with $\frac{\varepsilon^*}{3} < \frac{c}{4}$, the only sub-intervals of length n of $\langle x_0, f(x_0), \dots, f^{n-1}(x_0), x_0, f(x_0), \dots, f^{n-1}(x_0) \rangle$ that are $\frac{\varepsilon^*}{9}$ -shadowed by $\langle x_0, f(x_0), \dots, f^{n-1}(x_0) \rangle$ are the initial and the final sub-intervals. By the separation assumption, for any $1 \leq i \leq m$ we have

$$|\{y \in \Gamma_n^{\nu_i} : d_n(y, f^j(x_0)) < \frac{\varepsilon^*}{9} \text{ for some } 0 \leq j \leq n-1\}| \leq n.$$

Consequently, by (2.5) one can find a subset $\tilde{\Gamma}_n^{\nu_i} \subset \Gamma_n^{\nu_i}$ with $|\tilde{\Gamma}_n^{\nu_i}| > e^{n(h_{\nu_i} - \frac{\eta}{4})}$ such that $d_n(y, f^j(x_0)) \geq \frac{\varepsilon^*}{9}$ for any $y \in \tilde{\Gamma}_n^{\nu_i}$ and $0 \leq j \leq n-1$. Denote $r_i = |\tilde{\Gamma}_n^{\nu_i}|$ and $r = \sum_{i=1}^m r_i$. Enumerate the elements of each $\tilde{\Gamma}_n^{\nu_i}$ by $\tilde{\Gamma}_n^{\nu_i} = \{p_1^i, \dots, p_{r_i}^i\}$. Let $\tilde{\Gamma}_n = \{p_1^1, \dots, p_{r_1}^1, \dots, p_1^m, \dots, p_{r_m}^m\}$.

Take l large enough such that

$$\frac{1}{l} < \frac{\zeta}{12} \quad \text{and} \quad \frac{(l-2) \log |\tilde{\Gamma}_n^{\nu_i}|}{nl} > h_{\nu_i} - \eta \quad \text{for any } 1 \leq i \leq m. \quad (2.6)$$

Now let $\Gamma^i = \tilde{\Gamma}_n^{\nu_i} \times \tilde{\Gamma}_n^{\nu_i} \times \cdots \times \tilde{\Gamma}_n^{\nu_i}$ whose element is $\underline{y} = (y_1, \dots, y_{l-2})$ with $y_j \in \tilde{\Gamma}_n^{\nu_i}$ for $1 \leq j \leq l-2$, and let $\Gamma = \tilde{\Gamma}_n \times \tilde{\Gamma}_n \times \cdots \times \tilde{\Gamma}_n$ whose element is $\underline{y} = (y_1, \dots, y_{l-2})$ with $y_j \in \tilde{\Gamma}_n$ for $1 \leq j \leq l-2$. For any $y \in X$, let $\mathfrak{C}_y^n = \langle y, fy, \dots, f^{n-1}y \rangle$. Then for $\underline{y} \in \Gamma^i$ or Γ we define the following pseudo-orbit:

$$\mathfrak{C}_{\underline{y}} = \mathfrak{C}_{x_0}^n \mathfrak{C}_{x_0}^n \mathfrak{C}_{y_1}^n \mathfrak{C}_{y_2}^n \cdots \mathfrak{C}_{y_{l-2}}^n.$$

It is clear that $\mathfrak{C}_{\underline{y}}$ is a δ -pseudo-orbit. Moreover, one notes that we can freely concatenate such $\mathfrak{C}_{\underline{y}}$ s to constitute a δ -pseudo-orbit. We write $\mathfrak{C}_{\underline{y}} = \langle \omega_1, \omega_2, \dots, \omega_{ln} \rangle$. If $\underline{y} \in \Gamma^i$ and $d(f^k(z), \omega_{k+1}) \leq \varepsilon$ for any $0 \leq k \leq ln - 1$ then by Lemma 2.7 and (2. 6) one has

$$\begin{aligned} \rho(\mathcal{E}_{ln}(z), \nu_i) &\leq \rho(\mathcal{E}_{ln}(z), \frac{1}{n(l-2)} \sum_{k=1}^{n(l-2)} \delta_{\omega_k}) + \rho(\frac{1}{n(l-2)} \sum_{k=1}^{n(l-2)} \delta_{\omega_k}, \nu_i) \\ &\leq \varepsilon + \frac{4}{l} + \frac{1}{l-2} \sum_{j=1}^{l-2} \rho(\mathcal{E}_n(y_j), \nu_i) \\ &< \varepsilon + \frac{4}{l} + \frac{\zeta}{4} < \frac{3}{4}\zeta. \end{aligned} \quad (2. 7)$$

If $\underline{y} \in \Gamma$ and $d(f^k(z), \omega_{k+1}) \leq \varepsilon$ for any $0 \leq k \leq ln - 1$, we denote $q_i = |\{1 \leq j \leq l-2 : y_j \in \tilde{\Gamma}_n^{\nu_i}\}|$, then $\sum_{i=1}^m q_i = l-2$ and

$$\begin{aligned} \rho(\mathcal{E}_{ln}(z), \frac{\sum_{i=1}^m q_i \nu_i}{\sum_{i=1}^m q_i}) &\leq \rho(\mathcal{E}_{ln}(z), \frac{1}{n(l-2)} \sum_{k=1}^{n(l-2)} \delta_{\omega_k}) + \rho(\frac{1}{n(l-2)} \sum_{k=1}^{n(l-2)} \delta_{\omega_k}, \frac{\sum_{i=1}^m q_i \nu_i}{\sum_{i=1}^m q_i}) \\ &\leq \varepsilon + \frac{4}{l} + \frac{1}{l-2} \sum_{i=1}^m \sum_{y_j \in \tilde{\Gamma}_n^{\nu_i}} \rho(\mathcal{E}_n(y_j), \nu_i) \\ &< \varepsilon + \frac{4}{l} + \frac{\zeta}{4} < \frac{3}{4}\zeta. \end{aligned} \quad (2. 8)$$

by Lemma 2.7 and (2. 6). Now we define

$$\Sigma_{r_i^{l-2}} := \{\theta = \dots \theta_{-2} \theta_{-1} \theta_0 \theta_1 \theta_2 \dots : \theta_j = (\theta_{j,1}, \dots, \theta_{j,l-2}) \in \Gamma^i \text{ for any } j \in \mathbb{Z}\}$$

and

$$\Sigma_{r^{l-2}} := \{\theta = \dots \theta_{-2} \theta_{-1} \theta_0 \theta_1 \theta_2 \dots : \theta_j = (\theta_{j,1}, \dots, \theta_{j,l-2}) \in \Gamma \text{ for any } j \in \mathbb{Z}\}.$$

Then $(\Sigma_{r_i^{l-2}}, \sigma)$ and $(\Sigma_{r^{l-2}}, \sigma)$ are full shifts. This implies $(\Sigma_{r_i^{l-2}}, \sigma)$ and $(\Sigma_{r^{l-2}}, \sigma)$ are mixing and have shadowing property. And for each $\theta = \dots \theta_{-2} \theta_{-1} \theta_0 \theta_1 \theta_2 \dots$ in $\Sigma_{r_i^{l-2}}$ or $\Sigma_{r^{l-2}}$

$$\mathfrak{C}_\theta = \dots \mathfrak{C}_{\theta_{-2}} \mathfrak{C}_{\theta_{-1}} \mathfrak{C}_{\theta_0} \mathfrak{C}_{\theta_1} \mathfrak{C}_{\theta_2} \dots$$

is a δ -pseudo-orbit. We write $\mathfrak{C}_\theta = \dots \omega_{-2} \omega_{-1} \omega_0 \omega_1 \omega_2 \dots$, by the shadowing property,

$$Y_\theta = \{z \in X : d(f^j(z), \omega_j) \leq \varepsilon, j \in \mathbb{Z}\}$$

is nonempty and closed.

We claim that $Y_\theta \cap Y_{\theta'} = \emptyset$ for any $\theta \neq \theta'$ in $\Sigma_{r_i^{l-2}}$ or $\Sigma_{r^{l-2}}$. Next we prove the claim by the following two cases.

Case (1): If $\theta \neq \theta' \in \Sigma_{r_i^{l-2}}$ for some $1 \leq i \leq m$, then there is $t \in \mathbb{Z}$ and $1 \leq s \leq l-2$ such that $\theta_{t,s} \neq \theta'_{t,s}$. Since $\theta_{t,s}$ and $\theta'_{t,s}$ are $(n, \frac{\varepsilon^*}{3})$ -separated, we have $d_n(f^{lnt+sn+n}(z), f^{lnt+sn+n}(z')) > \varepsilon^*/3 - 2\varepsilon > \varepsilon^*/9$ for any $z \in Y_\theta$ and $z' \in Y_{\theta'}$. So $Y_\theta \cap Y_{\theta'} = \emptyset$.

Case (2): For any $\theta \neq \theta' \in \Sigma_{r^{l-2}}$, there is $t \in \mathbb{Z}$ and $1 \leq s \leq l-2$ such that $\theta_{t,s} \neq \theta'_{t,s}$. If $\theta_{t,s}, \theta'_{t,s} \in \tilde{\Gamma}_n^{\nu_i}$ for some $1 \leq i \leq m$, then $Y_\theta \cap Y_{\theta'} = \emptyset$ by Case (1). If there are $1 \leq i \neq i' \leq m$ such that $\theta_{t,s} \in \tilde{\Gamma}_n^{\nu_i}$ and $\theta'_{t,s} \in \tilde{\Gamma}_n^{\nu_{i'}}$, then

$$d_n(f^{lnt+sn+n}(z), f^{lnt+sn+n}(z')) > \varepsilon$$

for any $z \in Y_\theta$ and $z' \in Y_{\theta'}$. Otherwise, we have

$$\rho(\mathcal{E}_n(f^{lnt+sn+n}(z)), \mathcal{E}_n(f^{lnt+sn+n}(z'))) \leq \varepsilon \leq \frac{\rho_0}{6}.$$

Combining with

$$\rho(\mathcal{E}_n(f^{lnt+sn+n}(z)), \mathcal{E}_n(\theta_{t,s})) \leq \varepsilon \leq \frac{\rho_0}{6},$$

$$\rho(\mathcal{E}_n(f^{lnt+sn+n}(z')), \mathcal{E}_n(\theta'_{t,s})) \leq \varepsilon \leq \frac{\rho_0}{6}$$

and

$$\rho(\nu_i, \mathcal{E}_n(\theta_{t,s})) < \frac{\zeta}{4} \leq \frac{\rho_0}{4},$$

$$\rho(\nu_{i'}, \mathcal{E}_n(\theta'_{t,s})) < \frac{\zeta}{4} \leq \frac{\rho_0}{4},$$

we have $\rho(\nu_i, \nu_{i'}) < \rho_0$ which contradicts that $\rho_0 = \min\{\rho(\nu_i, \nu_j) : 1 \leq i < j \leq m\}$. So

$$d_n(f^{lnt+sn+n}(z), f^{lnt+sn+n}(z')) > \varepsilon$$

for any $z \in Y_\theta$ and $z' \in Y_{\theta'}$. This implies $Y_\theta \cap Y_{\theta'} = \emptyset$. Then we can define the following disjoint union:

$$\Delta_i = \bigsqcup_{\theta \in \Sigma_{r_i^{l-2}}} Y_\theta \text{ and } \Delta = \bigsqcup_{\theta \in \Sigma_{r^{l-2}}} Y_\theta.$$

Note that $f^{nl}(Y_\theta) = Y_{\sigma(\theta)}$. Then $f^{nl}(\Delta_i) = \Delta_i$, $1 \leq i \leq m$ and $f^{nl}(\Delta) = \Delta$. Therefore, if we define $\pi : \Delta \rightarrow \Sigma_{r^{l-2}}$ and $\pi_i : \Delta_i \rightarrow \Sigma_{r_i^{l-2}}$ as

$$\pi(x) := \theta \text{ for all } x \in Y_\theta \text{ with } \theta \in \Sigma_{r^{l-2}},$$

$$\pi_i(x) := \theta' \text{ for all } x \in Y_{\theta'} \text{ with } \theta' \in \Sigma_{r_i^{l-2}},$$

then π and π_i are surjective by the shadowing property. Moreover, it is not hard to check that π and π_i are continuous. So Δ and Δ_i are closed. Meanwhile, (X, f) is expansive, so π, π_i are conjugations.

Let $\Lambda_i = \cup_{k=0}^{nl-1} f^k(\Delta_i)$ and $\Lambda = \cup_{k=0}^{nl-1} f^k(\Delta)$. Then $f(\Lambda_i) = \Lambda_i$, $1 \leq i \leq m$ and $f(\Lambda) = \Lambda$. Now let us prove that Λ and Λ_i satisfy the property 1-4.

(1) Since π and π_i are conjugations, the transitivity, expansivity and the shadowing property of $(\Sigma_{r_i^{l-2}}, \sigma)$ and $(\Sigma_{r^{l-2}}, \sigma)$ yield the same properties of (Δ, f^{nl}) and (Δ_i, f^{nl}) . Next we show that $f^k(\Delta_i) \cap f^{k'}(\Delta_i) = \emptyset$ for any $0 \leq k < k' \leq nl - 1$. If $f^k(\Delta_i) \cap f^{k'}(\Delta_i) \neq \emptyset$, then for any $z \in f^k(\Delta_i) \cap f^{k'}(\Delta_i)$, there exist $\theta, \theta' \in \Sigma_{r_i^{l-2}}$ such that

$$d(f^{j-k}(z), \omega_j) \leq \varepsilon \text{ and } d(f^{j-k'}(z), \omega'_j) \leq \varepsilon \quad \forall j \in \mathbb{Z}$$

where $\mathfrak{C}_\theta = \dots \omega_{-2} \omega_{-1} \omega_0 \omega_1 \omega_2 \dots$ and $\mathfrak{C}_{\theta'} = \dots \omega'_{-2} \omega'_{-1} \omega'_0 \omega'_1 \omega'_2 \dots$. Then we have

$$d(\omega_{j+k}, \omega'_{j+k'}) \leq 2\varepsilon \quad \forall j \in \mathbb{Z}. \quad (2.9)$$

Case (1): If $1 \leq k' - k \leq n - 1$, then (2.9) implies $d(\omega_j, \omega'_{j+k'-k}) \leq 2\varepsilon < \frac{\varepsilon^*}{9} \quad \forall 0 \leq j \leq n - 1$. Note that $\omega_0 \omega_1 \omega_2 \dots \omega_{2n-1} = \omega'_0 \omega'_1 \omega'_2 \dots \omega'_{2n-1} = \langle x_0, f x_0, \dots, f^{n-1} x_0, x_0, f x_0, \dots, f^{n-1} x_0 \rangle$, this contradicts that the minimal period of x_0 is n .

Case (2): If $k' - k = n$, then (2.9) implies $d(\omega_{j+n}, \omega'_{j+2n}) \leq 2\varepsilon < \frac{\varepsilon^*}{9} \quad \forall 0 \leq j \leq n - 1$. Note that $\omega_n \omega_{n+1} \dots \omega_{2n-1} = \langle x_0, f x_0, \dots, f^{n-1} x_0 \rangle$ and $\omega'_{2n} \in \tilde{\Gamma}_n^{\nu_i}$, this contradicts that $d_n(y, f^j(x_0)) \geq \frac{\varepsilon^*}{9}$ for any $y \in \tilde{\Gamma}_n^{\nu_i}$ and $0 \leq j \leq n - 1$.

Case (3): If $n < k' - k \leq n(l - 1)$, then $k' - k = tn + s$ for some $1 \leq t \leq l - 2$ and $0 \leq s \leq n - 1$. Thus (2.9) implies $d(\omega_{n-s+j}, \omega'_{(t+1)n+j}) \leq 2\varepsilon \quad \forall 0 \leq j \leq n - 1$. Note that $\omega_0 \omega_1 \omega_2 \dots \omega_{2n-1} = \langle x_0, f x_0, \dots, f^{n-1} x_0, x_0, f x_0, \dots, f^{n-1} x_0 \rangle$, and $\omega'_{(t+1)n} \in \tilde{\Gamma}_n^{\nu_i}$, this contradicts that $d_n(y, f^j(x_0)) \geq \frac{\varepsilon^*}{9}$ for any $y \in \tilde{\Gamma}_n^{\nu_i}$ and $0 \leq j \leq n - 1$.

Case (4): If $n(l - 1) < k' - k \leq nl - 1$, then (2.9) implies $d(\omega_{nl-k'+k+j}, \omega'_{nl+j}) \leq 2\varepsilon \quad \forall 0 \leq j \leq n - 1$. Note that $\omega_0 \omega_1 \omega_2 \dots \omega_{2n-1} = \omega'_{nl} \omega'_{nl+1} \omega'_{nl+2} \dots \omega'_{(l+2)n-1} = \langle x_0, f x_0, \dots, f^{n-1} x_0, x_0, f x_0, \dots, f^{n-1} x_0 \rangle$, and $1 \leq nl - k' + k < n$, this contradicts that the minimal period of x_0 is n .

So we have $f^k(\Delta_i) \cap f^{k'}(\Delta_i) = \emptyset$ for any $0 \leq k < k' \leq nl - 1$. Therefore, Proposition 2.9 ensures that (Λ_i, f) is transitive and topologically Anosov. In fact, if we define

$$\Omega_i = \{\mathfrak{C}_\theta = \dots \omega_{-2} \omega_{-1} \omega_0 \omega_1 \omega_2 \dots : \theta \in \Sigma_{r_i^{l-2}}\},$$

then (Ω_i, σ^{nl}) conjugates to $(\Sigma_{r_i^{l-2}}, \sigma)$. Thus (Ω_i, σ^{nl}) conjugates to (Δ_i, f^{nl}) , and $(\cup_{k=0}^{nl-1} \sigma^k(\Omega_i), \sigma)$ conjugates to (Λ_i, f) . This implies $(\cup_{k=0}^{nl-1} \sigma^k(\Omega_i), \sigma)$ is a subshift which is transitive and topologically Anosov. Recall from [42] a subshift satisfies shadowing property if and only if it is a subshift of finite type. So $(\cup_{k=0}^{nl-1} \sigma^k(\Omega_i), \sigma)$ is a transitive subshift of finite type. By similar method, (Λ, f) is also transitive and topologically Anosov, and (Λ, f) conjugate to a transitive subshift of finite type.

(2) One has $h_{\text{top}}(f, \Lambda_i) = \frac{1}{nl} h_{\text{top}}(f^{nl}, \Delta_i) = \frac{1}{nl} h_{\text{top}}(\sigma, \Sigma_{r_i^{l-2}}) = \frac{(l-2) \log |\tilde{\Gamma}_n^{\nu_i}|}{nl} > h_{\nu_i} - \eta > h_{\nu_i} - \eta_0$ by (2. 6).

(3) For any ergodic measure $\mu_i \in M(f, \Lambda_i)$, pick an arbitrary generic point z_i of μ_i in Δ_i . Then

$$\rho(\mathcal{E}_{ln}(f^{tln}(z_i)), \nu_i) < \frac{3}{4}\zeta \text{ for any } t \in \mathbb{N}$$

by (2. 7). In addition, we have $\mu_i = \lim_{j \rightarrow \infty} \mathcal{E}_j(z_i) = \lim_{t \rightarrow \infty} \mathcal{E}_{tln}(z_i)$. So we have

$$\rho(\mu_i, \nu_i) = \lim_{t \rightarrow \infty} \rho(\mathcal{E}_{tln}(z_i), \nu_i) \leq \frac{3}{4}\zeta.$$

By the ergodic decomposition Theorem, we obtain that $d_H(\nu_i, M(f, \Lambda_i)) \leq \frac{3}{4}\zeta$. Now since K is convex and $\Lambda_i \subseteq \Lambda$, one gets that $K \subseteq \mathcal{B}(M(f, \Lambda), \zeta) \subseteq \mathcal{B}(M(f, \Lambda), \zeta_0)$.

On the other hand, for any ergodic measure $\mu \in M(f, \Lambda)$, pick a generic point z of μ in Δ . Then z ε -shadows some δ -pseudo-orbit \mathfrak{C}_θ with $\theta \in \Sigma_{r^{l-2}}$. Then for any $t \in \mathbb{N}$, there exist nonnegative integers $q_i, 1 \leq i \leq m$ such that

$$\rho\left(\mathcal{E}_{ln}(f^{tln}(z)), \frac{\sum_{i=1}^m q_i \nu_i}{\sum_{i=1}^m q_i}\right) < \frac{3}{4}\zeta$$

by (2. 8). So $\mu \in \mathcal{B}(K, \frac{3}{4}\zeta)$. By the ergodic decomposition Theorem, $M(f, \Lambda) \subseteq \mathcal{B}(K, \zeta) \subseteq \mathcal{B}(K, \zeta_0)$. As a result, $\Lambda \subsetneq X$. For otherwise, $d_H(K, M(f, \Lambda)) = d_H(K, M(f, X)) > \zeta$, a contradiction.

(4) Note that for any θ in $\Sigma_{r_i^{l-2}}$ or $\Sigma_{r^{l-2}}$, one has $\omega_{mnl} = x_0$ for any integer m where $\mathfrak{C}_\theta = \dots \omega_{-2} \omega_{-1} \omega_0 \omega_1 \omega_2 \dots$. Then for any z in Δ_i or Δ ,

$$d(f^{mnl}(z), x) \leq d(f^{mnl}(z), x_0) + d(x_0, x) < \varepsilon + \frac{\delta}{2} < 2\varepsilon < \varepsilon_0.$$

So for any z in Λ_i or Λ one has $f^{j+mnl}(z) \in B(x, \varepsilon)$ for some $0 \leq j \leq nl - 1$ and any $m \in \mathbb{Z}$. \square

3 Almost Additive Sequences and Theorem A(I)

In this section we give abstract conditions on which Theorem A(I) holds in the more general context of almost additive sequences of continuous functions. We first recall some definitions about almost additive sequences from [5]. Consider a topological dynamical system (X, f) . We recall that a sequence of functions $\Phi = (\varphi_n)_{n \in \mathbb{N}}$ is said to be *almost additive* (with respect to (X, f)) if there is a constant $C > 0$ such that for every $n, m \in \mathbb{N}$ we have

$$-C + \varphi_n + \varphi_m \circ f^n \leq \varphi_{n+m} \leq C + \varphi_n + \varphi_m \circ f^n.$$

We denote by $A(f, X)$ the family of almost additive sequences of continuous functions. From [5, Proposition 3] the limit $\lim_{n \rightarrow \infty} \frac{1}{n} \int \varphi_n d\mu$ exists for any $\Phi = (\varphi_n)_{n \in \mathbb{N}} \in A(f, X)$ and any $\mu \in M(f, X)$, and the function,

$$M(f, X) \ni \mu \mapsto \lim_{n \rightarrow \infty} \frac{1}{n} \int \varphi_n d\mu, \quad (3. 10)$$

is continuous with the weak* topology in $M(f, X)$.

Let $d \in \mathbb{N}$ and take $(A, B) \in A(f, X)^d \times A(f, X)^d$. We write

$$A = (\Phi^1, \dots, \Phi^d) \text{ and } B = (\Psi^1, \dots, \Psi^d)$$

and also $\Phi^i = (\varphi_n^i)_{n \in \mathbb{N}}$ and $\Psi^i = (\psi_n^i)_{n \in \mathbb{N}}$. We assume that

$$\liminf_{m \rightarrow \infty} \frac{\psi_m^i(x)}{m} > 0 \quad \text{and} \quad \psi_n^i(x) > 0 \quad (3.11)$$

for every $i = 1, \dots, d, x \in X$, and $n \in \mathbb{N}$. Given $a = (a_1, \dots, a_d) \in \mathbb{R}^d$ we define:

$$R_{A,B}(a) = \bigcap_{i=1}^d \left\{ x \in X : \lim_{n \rightarrow \infty} \frac{\varphi_n^i(x)}{\psi_n^i(x)} = a_i \right\}.$$

We also consider the function $\mathcal{P}_{A,B} : M(f, X) \rightarrow \mathbb{R}$ defined by:

$$\mathcal{P}_{A,B}(\mu) = \left(\lim_{n \rightarrow \infty} \frac{1}{n} \int \varphi_n^1 d\mu, \dots, \lim_{n \rightarrow \infty} \frac{1}{n} \int \varphi_n^d d\mu \right) = \lim_{n \rightarrow \infty} \left(\frac{\int \varphi_n^1 d\mu}{\int \psi_n^1 d\mu}, \dots, \frac{\int \varphi_n^d d\mu}{\int \psi_n^d d\mu} \right). \quad (3.12)$$

(3.10) ensures that the second identity in (3.12) holds, and that the function $\mathcal{P}_{A,B}$ is continuous. Denote

$$L_{A,B} = \{ \mathcal{P}_{A,B}(\mu) : \mu \in M(f, X) \}.$$

For any $a \in L_{A,B}$, define

$$M_{A,B}(a) = \{ \mu \in M(f, X) : \mathcal{P}_{A,B}(\mu) = a \}, \quad M_{A,B}^{erg}(a) = \{ \mu \in M_{erg}(f, X) : \mathcal{P}_{A,B}(\mu) = a \}.$$

Then $M_{A,B}(a)$ is closed in $M(f, X)$ since the function $\mathcal{P}_{A,B}$ is continuous.

Let $\alpha : M(f, X) \rightarrow \mathbb{R}$ be a continuous function. We define the pressure of α with respect to μ by $P(f, \alpha, \mu) = h_\mu(f) + \alpha(\mu)$. Let $E(f, X) \subseteq A(f, X)$ be the family of sequences with a unique equilibrium measure (see definition of equilibrium measures of almost additive sequences in subsection 5.1). Now we give abstract conditions on which Theorem A(I) holds in the more general context of almost additive sequences of continuous functions.

Theorem B. *Suppose (X, f) is a dynamical system whose entropy function is upper semi-continuous. Let $d \in \mathbb{N}$ and $(A, B) \in A(f, X)^d \times A(f, X)^d$ such that B satisfies (3.11). Let $\alpha : M(f, X) \rightarrow \mathbb{R}$ be a continuous function. Assume that the following holds: for any $K = \text{cov}\{\mu_i\}_{i=1}^m \subseteq M(f, X)$, and any $\eta, \zeta > 0$, there exist compact invariant subsets $\Lambda_i \subseteq \Lambda \subsetneq X$ such that for each $i \in \{1, 2, \dots, m\}$*

- (1) $h_{top}(f, \Lambda_i) > h_{\mu_i}(f) - \eta$.
- (2) $d_H(K, M(f, \Lambda)) < \zeta$, $d_H(\mu_i, M(f, \Lambda_i)) < \zeta$.
- (3) $\text{span}\{\Phi^1|_\Lambda, \Psi^1|_\Lambda, \dots, \Phi^d|_\Lambda, \Psi^d|_\Lambda\} \subseteq E(f|_\Lambda, \Lambda)$.

Then for any $a \in \text{Int}(L_{A,B})$, any $\mu_0 \in M_{A,B}(a)$ and any $\eta, \zeta > 0$, there is $\nu \in M_{A,B}^{erg}(a)$ such that $\rho(\nu, \mu_0) < \zeta$ and $|P(f, \alpha, \nu) - P(f, \alpha, \mu_0)| < \eta$.

Example 3.1. The function $\alpha : M(f, X) \rightarrow \mathbb{R}$ can be defined as following:

- (1) $\alpha \equiv 0$. Then $P(f, \alpha, \mu) = h_\mu(f)$ is the metric entropy of μ .
- (2) $\alpha(\mu) = \int \varphi d\mu$ with a continuous function φ . Then from the *weak**-topology on $M(X)$, $\alpha : M(f, X) \rightarrow \mathbb{R}$ is a continuous function. $P(f, \varphi, \mu) = h_\mu(f) + \alpha(\mu)$ is the pressure of φ with respect to μ .
- (3) $\alpha(\mu) = \lim_{n \rightarrow \infty} \frac{1}{n} \int \varphi_n d\mu$ with an almost additive sequences of continuous functions $\Phi = (\varphi_n)_{n \in \mathbb{N}}$. Then $\alpha : M(f, X) \rightarrow \mathbb{R}$ is a continuous function from (3.10). $P(f, \Phi, \mu) = h_\mu(f) + \lim_{n \rightarrow \infty} \frac{1}{n} \int \varphi_n d\mu$ is the pressure of Φ with respect to μ . Readers can refer to [4, 31] for thermodynamic formalism of almost additive sequences.

Remark 3.2. Another type of additivity assumption was introduced by Feng and Huang[19]: a sequence of continuous functions $(\varphi_n)_{n \in \mathbb{N}}$ is said to be asymptotically additive if for each $\varepsilon > 0$, there exists a continuous function $\varphi : X \rightarrow \mathbb{R}$ such that

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \sup_{x \in X} |\varphi_n(x) - S_n \varphi(x)| \leq \varepsilon,$$

where $S_n f(x) = \sum_{i=0}^{n-1} \varphi(f^i(x))$. Every almost additive sequence is asymptotically additive (see, for example, [43, Proposition 2.1]). It's proved in [4, Page 294] that for expansive dynamical system satisfying specification property, every almost additive sequence of continuous functions with bounded variation has a unique equilibrium measure. But in the case of asymptotically additive sequences, the bounded variation condition does not guarantee the uniqueness of the equilibrium measure (see [15, Remark 4.5]). The uniqueness of the equilibrium measure is indispensable on the proof of conditional variational principle (see Theorem 3.7) which is used in our proof. So we don't consider asymptotically additive sequences in this article.

3.1 Some lemmas

To prove Theorem B, we need some lemmas. For any $r \in \mathbb{R}$, denote $r^+ = \{s \in \mathbb{R} : s > r\}$ and $r^- = \{s \in \mathbb{R} : s < r\}$. For any $d \in \mathbb{N}$, $r = (r_1, \dots, r_d) \in \mathbb{R}^d$ and $\xi = (\xi_1, \dots, \xi_d) \in \{+, -\}^d$, we define

$$r^\xi = \{s = (s_1, \dots, s_d) \in \mathbb{R}^d : s_i \in r_i^{\xi_i} \text{ for } i = 1, 2, \dots, d\}.$$

We denote $F^d = \left\{ \left(\frac{p_1}{q_1}, \dots, \frac{p_d}{q_d} \right) : p_i, q_i \in \mathbb{R} \text{ and } q_i > 0 \text{ for any } 1 \leq i \leq d \right\}$. It is easy to check that

Lemma 3.3. *Let $b_i = \frac{p_i}{q_i} \in F^1$ for $i = 1, 2$.*

(1) *If $b_1 = b_2$, then $\frac{\theta p_1^1 + (1-\theta)p_2^2}{\theta q_1^1 + (1-\theta)q_2^2} = b_1 = b_2$ for any $\theta \in [0, 1]$.*

(2) *If $b_1 \neq b_2$, then $\frac{\theta p_1^1 + (1-\theta)p_2^2}{\theta q_1^1 + (1-\theta)q_2^2}$ is strictly monotonic on $\theta \in [0, 1]$.*

Lemma 3.4. *Let $d \in \mathbb{N}$ and $a = \left(\frac{p_1}{q_1}, \dots, \frac{p_d}{q_d} \right) \in F^d$. If $\{b_\xi = \left(\frac{p_1^\xi}{q_1^\xi}, \dots, \frac{p_d^\xi}{q_d^\xi} \right)\}_{\xi \in \{+, -\}^d} \subseteq F^d$ are 2^d numbers satisfies $b_\xi \in a^\xi$ for any $\xi \in \{+, -\}^d$, then there are 2^d numbers $\{\theta_\xi\}_{\xi \in \{+, -\}^d} \subseteq [0, 1]$ such that $\sum_{\xi \in \{+, -\}^d} \theta_\xi = 1$ and*

$$\frac{\sum_{\xi \in \{+, -\}^d} \theta_\xi p_i^\xi}{\sum_{\xi \in \{+, -\}^d} \theta_\xi q_i^\xi} = \frac{p_i}{q_i} \text{ for any } 1 \leq i \leq d.$$

Proof. We prove the lemma inductively. It is clearly true if $d = 1$ from Lemma 3.3.

Now we assume that it is true for $d = k \in \mathbb{N}$. Let $a = (a_1, \dots, a_{k+1}) \in \mathbb{R}^{k+1}$, and $\{b^\xi\}_{\xi \in \{+, -\}^{k+1}}$ is 2^{k+1} numbers satisfies $b^\xi \in a^\xi$ for any $\xi \in \{+, -\}^{k+1}$. Then for the 2^k numbers $\{b^\xi\}_{\xi_{k+1}=+}$, there is 2^k numbers $\{\tau^\xi\}_{\xi_{k+1}=+} \subseteq [0, 1]$ such that $\sum_{\xi_{k+1}=+} \tau^\xi = 1$ and

$$\frac{\sum_{\xi_{k+1}=+} \tau^\xi p_i^\xi}{\sum_{\xi_{k+1}=+} \tau^\xi q_i^\xi} = \frac{p_i}{q_i} \text{ for any } 1 \leq i \leq k. \quad (3.13)$$

Since $\frac{p_{k+1}^\xi}{q_{k+1}^\xi} > \frac{p_{k+1}}{q_{k+1}}$ for any $\xi \in \{+, -\}^{k+1}$ with $\xi_{k+1} = +$, then we have

$$\frac{\sum_{\xi_{k+1}=+} \tau^\xi p_{k+1}^\xi}{\sum_{\xi_{k+1}=+} \tau^\xi q_{k+1}^\xi} > \frac{p_{k+1}}{q_{k+1}}. \quad (3.14)$$

Similarly, the 2^k numbers $\{b^\xi\}_{\xi_{k+1}=-}$, there is 2^k numbers $\{\tau^\xi\}_{\xi_{k+1}=-} \subseteq [0, 1]$ such that $\sum_{\xi_{k+1}=-} \tau^\xi = 1$ and

$$\frac{\sum_{\xi_{k+1}=-} \tau^\xi p_i^\xi}{\sum_{\xi_{k+1}=-} \tau^\xi q_i^\xi} = \frac{p_i}{q_i} \text{ for any } 1 \leq i \leq k. \quad (3.15)$$

Since $\frac{p_{k+1}^\xi}{q_{k+1}^\xi} < \frac{p_{k+1}}{q_{k+1}}$ for any $\xi \in \{+, -\}^{k+1}$ with $\xi_{k+1} = -$, then we have

$$\frac{\sum_{\xi_{k+1}=-} \tau_\xi p_{k+1}^\xi}{\sum_{\xi_{k+1}=-} \tau_\xi q_{k+1}^\xi} < \frac{p_{k+1}}{q_{k+1}}. \quad (3.16)$$

By (3.14), (3.16) and Lemma 3.3(2) there is $\tau_{k+1} \in (0, 1)$ such that

$$\frac{\tau_{k+1} \sum_{\xi_{k+1}=+} \tau_\xi p_{k+1}^\xi + (1 - \tau_{k+1}) \sum_{\xi_{k+1}=-} \tau_\xi p_{k+1}^\xi}{\tau_{k+1} \sum_{\xi_{k+1}=+} \tau_\xi q_{k+1}^\xi + (1 - \tau_{k+1}) \sum_{\xi_{k+1}=-} \tau_\xi q_{k+1}^\xi} = \frac{p_{k+1}}{q_{k+1}}.$$

By (3.13), (3.15) and Lemma 3.3(1), we have

$$\frac{\tau_{k+1} \sum_{\xi_{k+1}=+} \tau_\xi p_i^\xi + (1 - \tau_{k+1}) \sum_{\xi_{k+1}=-} \tau_\xi p_i^\xi}{\tau_{k+1} \sum_{\xi_{k+1}=+} \tau_\xi q_i^\xi + (1 - \tau_{k+1}) \sum_{\xi_{k+1}=-} \tau_\xi q_i^\xi} = \frac{p_i}{q_i} \text{ for any } 1 \leq i \leq k..$$

Let

$$\theta_\xi = \begin{cases} \tau_{k+1} \tau_\xi, & \text{for } \xi_{k+1} = + \\ (1 - \tau_{k+1}) \tau_\xi, & \text{for } \xi_{k+1} = -. \end{cases}$$

Then we have $\sum_{\xi \in \{+, -\}^{k+1}} \theta_\xi = 1$ and

$$\frac{\sum_{\xi \in \{+, -\}^{k+1}} \theta_\xi p_i^\xi}{\sum_{\xi \in \{+, -\}^{k+1}} \theta_\xi q_i^\xi} = \frac{p_i}{q_i} \text{ for any } 1 \leq i \leq k + 1.$$

So we complete the proof of Lemma 3.4. \square

From Lemma 3.4 we have

Corollary 3.5. *Suppose (X, f) is a dynamical system. Let $d \in \mathbb{N}$ and $(A, B) \in A(f, X)^d \times A(f, X)^d$ such that B satisfies (3.11). Then for any $a \in \text{Int}(L_{A,B})$, and 2^d invariant measures $\{\mu_\xi\}_{\xi \in \{+, -\}^d}$ with*

$$\mathcal{P}_{A,B}(\mu_\xi) \in a^\xi \text{ for any } \xi \in \{+, -\}^d,$$

there are 2^d numbers $\{\theta_\xi\}_{\xi \in \{+, -\}^d} \subseteq [0, 1]$ such that $\sum_{\xi \in \{+, -\}^d} \theta_\xi = 1$ and $\mathcal{P}_{A,B}(\sum_{\xi \in \{+, -\}^d} \theta_\xi \mu_\xi) = a$.

Lemma 3.6. *Suppose (X, f) is a dynamical system. Let $d \in \mathbb{N}$ and $(A, B) \in A(f, X)^d \times A(f, X)^d$ such that B satisfies (3.11). Then for any $a \in \text{Int}(L_{A,B})$, any $\mu \in M_{A,B}(a)$ and any $\eta, \zeta > 0$, there exist 2^d invariant measures $\{\mu_\xi\}_{\xi \in \{+, -\}^d}$ such that*

$$\mathcal{P}_{A,B}(\mu_\xi) \in a^\xi, h_{\mu_\xi}(f) > h_\mu(f) - \eta \text{ and } \rho(\mu_\xi, \mu) < \zeta \text{ for any } \xi \in \{+, -\}^d.$$

Proof. By $a \in \text{Int}(L_{A,B})$ there is $\nu_\xi \in M(f, X)$ such that $\mathcal{P}_{A,B}(\nu_\xi) \in a^\xi$ for any $\xi \in \{+, -\}^d$. Then there is $\tau_\xi \in (0, 1)$ close to 1 such that $\mu_\xi = \tau_\xi \mu + (1 - \tau_\xi) \nu_\xi$ satisfies

$$h_{\mu_\xi}(f) > h_\mu(f) - \eta \text{ and } \rho(\mu_\xi, \mu) < \zeta \text{ for any } \xi \in \{+, -\}^d.$$

And we have $\mathcal{P}_{A,B}(\mu_\xi) \in a^\xi$ by $\tau_\xi > 0$ and Lemma 3.3(2). \square

In [5] L. Barreira and P. Doutor give the conditional variational principle as following.

Theorem 3.7. [5, Theorem 3] *Suppose (X, f) is a dynamical system whose entropy function is upper semi-continuous. Let $d \in \mathbb{N}$ and $(A, B) \in A(f, X)^d \times A(f, X)^d$ such that $\text{span}\{\Phi^1, \Psi^1, \dots, \Phi^d, \Psi^d\} \subseteq E(f, X)$ and B satisfies (3.11). If $a \notin L_{A,B}$, then $R_{A,B}(a) = \emptyset$. Otherwise, if $a \in \text{Int}(L_{A,B})$, then $R_{A,B}(a) \neq \emptyset$, and the following properties hold:*

(1) $h_{\text{top}}(f, R_{A,B}(a))$ satisfies the variational principle:

$$h_{\text{top}}(f, R_{A,B}(a)) = \max \{h_\mu(f) : \mu \in M(f, X) \text{ and } \mathcal{P}_{A,B}(\mu) = a\}.$$

(2) *There is an ergodic measure $\mu_a \in M(f, X)$ with $\mathcal{P}_{A,B}(\mu_a) = a$, $\mu_a(R_{A,B}(a)) = 1$, and*

$$h_{\text{top}}(f, R_{A,B}(a)) = h_{\mu_a}(f).$$

3.2 Proof of Theorem B

Fix $a \in \text{Int}(L_{A,B})$, $\mu_0 \in M_{A,B}(a)$ and $\eta, \zeta > 0$. Since the metric entropy is upper semi-continuous and $\alpha : M(f, X) \rightarrow \mathbb{R}$ is continuous, there is $0 < \zeta' < \zeta$ such that

$$h_\omega(f) < h_{\mu_0}(f) + \frac{\eta}{2} \text{ and } |\alpha(\omega) - \alpha(\mu_0)| < \frac{\eta}{2}. \quad (3.17)$$

for any $\omega \in M(f, X)$ with $\rho(\mu_0, \omega) < \zeta'$. By Lemma 3.6 there exist 2^d invariant measures $\{\mu_\xi\}_{\xi \in \{+, -\}^d}$ such that

$$\mathcal{P}_{A,B}(\mu_\xi) \in a^\xi, h_{\mu_\xi}(f) > h_{\mu_0}(f) - \frac{\eta}{6} \text{ and } \rho(\mu_\xi, \mu_0) < \frac{\zeta'}{2} \text{ for any } \xi \in \{+, -\}^d. \quad (3.18)$$

Since the function $\mathcal{P}_{A,B}$ is continuous, there is $0 < \zeta'' < \zeta'$ such that such that

$$\mathcal{P}_{A,B}(\omega_\xi) \in a^\xi \quad (3.19)$$

for any $\omega_\xi \in M(f, X)$ with $\rho(\omega_\xi, \mu_\xi) < \zeta''$. For the 2^d invariant measures $\{\mu_\xi\}_{\xi \in \{+, -\}^d}$, there exist compact invariant subsets $\Lambda_\xi \subseteq \Lambda \subsetneq X$ such that for each $\xi \in \{+, -\}^d$

- (1) $h_{\text{top}}(f, \Lambda_\xi) > h_{\mu_\xi}(f) - \frac{\eta}{6}$.
- (2) $d_H(\text{cov}\{\mu_\xi\}_{\xi \in \{+, -\}^d}, M(f, \Lambda)) < \frac{\zeta''}{2}$, $d_H(\mu_\xi, M(f, \Lambda_\xi)) < \frac{\zeta''}{2}$.
- (3) $\text{span}\{\Phi^1|_\Lambda, \Psi^1|_\Lambda, \dots, \Phi^d|_\Lambda, \Psi^d|_\Lambda\} \subseteq E(f|_\Lambda, \Lambda)$.

By item(1) and the variational principle of the topological entropy, there are $\nu_\xi \in M(f, \Lambda_\xi)$ such that

$$h_{\nu_\xi}(f) > h_{\text{top}}(f, \Lambda_\xi) - \frac{\eta}{6} > h_{\mu_\xi}(f) - \frac{2\eta}{6} > h_{\mu_0}(f) - \frac{\eta}{2}.$$

Then by item(2) and (3.19), we have $\mathcal{P}_{A,B}(\nu_\xi) \in a^\xi$. By Corollary 3.5 there is 2^d numbers $\{\theta_\xi\}_{\xi \in \{+, -\}^d} \subseteq [0, 1]$ such that $\sum_{\xi \in \{+, -\}^d} \theta_\xi = 1$ and $\mathcal{P}_{A,B}(\nu') = a$ where $\nu' = \sum_{\xi \in \{+, -\}^d} \theta_\xi \nu_\xi$. Then on one hand we have

$$\begin{aligned} & \sup\{h_\mu(f) : \mu \in M(f, \Lambda) \text{ and } \mathcal{P}_{A,B}(\mu) = a\} \\ & \geq h_{\nu'}(f) \geq \min\{h_{\nu_\xi}(f) : \xi \in \{+, -\}^d\} \\ & > h_{\mu_0}(f) - \frac{\eta}{2}. \end{aligned} \quad (3.20)$$

On the other hand, by item(2), (3.18) and (3.17) we have

$$\sup\{h_\mu(f) : \mu \in M(f, \Lambda) \text{ and } \mathcal{P}_{A,B}(\mu) = a\} < h_{\mu_0}(f) + \frac{\eta}{2}. \quad (3.21)$$

Now by item(3) and Theorem 3.7, there is an ergodic measure $\nu \in M(f, \Lambda)$ with $\mathcal{P}_{A,B}(\nu) = a$, and

$$h_\nu(f) = h_{\text{top}}(f, R_{A,B}(a)) = \max\{h_\mu(f) : \mu \in M(f, \Lambda) \text{ and } \mathcal{P}_{A,B}(\mu) = a\}.$$

Then $\nu \in M_{A,B}^{\text{erg}}(a)$ and by (3.20), (3.21) we have $|h_\nu(f) - h_{\mu_0}(f)| < \frac{\eta}{2}$. By item(2) and (3.18) we have $\rho(\nu, \mu_0) < \zeta' < \zeta$. Finally, by (3.17) we have $|\alpha(\nu) - \alpha(\mu_0)| < \frac{\eta}{2}$, and thus $|P(f, \alpha, \nu) - P(f, \alpha, \mu_0)| < \eta$. So we complete the proof of Theorem B. \square

4 Almost Additive Sequences and Theorem A(II)-(IV)

In this section we give abstract conditions on which Theorem A(II)-(IV) holds in the more general context of almost additive sequences of continuous functions. Consider a topological dynamical system (X, f) . Let $d \in \mathbb{N}$ and $(A, B) \in A(f, X)^d \times A(f, X)^d$ such that B satisfies (3.11). Let $\alpha : M(f, X) \rightarrow \mathbb{R}$ be a continuous function. Recall that the pressure of α with respect to μ is $P(f, \alpha, \mu) = h_\mu(f) + \alpha(\mu)$. For any $a \in L_{A,B}$, denote

$$H_{A,B}(f, \alpha, a) = \sup\{P(f, \alpha, \mu) : \mu \in M_{A,B}(a)\}.$$

In particular, when $\alpha \equiv 0$, we write

$$H_{A,B}(f, a) = H_{A,B}(f, 0, a) = \sup\{h_\mu(f) : \mu \in M_{A,B}(a)\}.$$

Now we give abstract conditions on which Theorem A(II)-(IV) holds in the more general context of almost additive sequences of continuous functions.

Theorem C. *Assume that, in the conditions of Theorem B, $\{\mu \in M(f, X) : h_\mu(f) = 0\}$ is dense in $M(f, X)$, and α satisfies that for any $\mu_1, \mu_2 \in M(f, X)$ with $\alpha(\mu_1) \neq \alpha(\mu_2)$*

$$\beta(\theta) := \alpha(\theta\mu_1 + (1 - \theta)\mu_2) \text{ is strictly monotonic on } [0, 1], \quad (4. 22)$$

and for any $\mu_1, \mu_2 \in M(f, X)$ with $\alpha(\mu_1) = \alpha(\mu_2)$

$$\beta(\theta) := \alpha(\theta\mu_1 + (1 - \theta)\mu_2) \text{ is constant on } [0, 1]. \quad (4. 23)$$

Then we have

- (II) *For any $a \in \text{Int}(L_{A,B})$, any $\mu_0 \in M_{A,B}(a)$, any $\max_{\mu \in M_{A,B}(a)} \alpha(\mu) \leq c \leq P(f, \alpha, \mu_0)$ and any $\eta, \zeta > 0$, there is $\nu \in M_{A,B}^{erg}(a)$ such that $\rho(\nu, \mu_0) < \zeta$ and $|P(f, \alpha, \nu) - c| < \eta$.*
- (III) *For any $a \in \text{Int}(L_{A,B})$ and $\max_{\mu \in M_{A,B}(a)} \alpha(\mu) \leq c < H_{A,B}(f, \alpha, a)$, the set $\{\mu \in M_{A,B}^{erg}(a) : P(f, \alpha, \mu) = c\}$ is residual in $\{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) \geq c\}$. If further there is an invariant measure with full support, then for any $a \in \text{Int}(L_{A,B})$ and $\max_{\mu \in M_{A,B}(a)} \alpha(\mu) \leq c < H_{A,B}(f, \alpha, a)$, the set $\{\mu \in M_{A,B}^{erg}(a) : P(f, \alpha, \mu) = c, S_\mu = X\}$ is residual in $\{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) \geq c\}$.*
- (IV) *$\{(\mathcal{P}_{A,B}(\mu), P(f, \alpha, \mu)) : \mu \in M(f, X), a = \mathcal{P}_{A,B}(\mu) \in \text{Int}(L_{A,B}), \max_{\mu \in M_{A,B}(a)} \alpha(\mu) \leq P(f, \alpha, \mu) < H_{A,B}(f, \alpha, a)\}$ coincides with $\{(\mathcal{P}_{A,B}(\mu), P(f, \alpha, \mu)) : \mu \in M_{erg}(f, X), a = \mathcal{P}_{A,B}(\mu) \in \text{Int}(L_{A,B}), \max_{\mu \in M_{A,B}(a)} \alpha(\mu) \leq P(f, \alpha, \mu) < H_{A,B}(f, \alpha, a)\}$. If further $\text{span}\{\Phi^1, \Psi^1, \dots, \Phi^d, \Psi^d\} \subseteq E(f, X)$ and $\alpha \equiv 0$, then $\{(\mathcal{P}_{A,B}(\mu), h_\mu(f)) : \mu \in M(f, X), \mathcal{P}_{A,B}(\mu) \in \text{Int}(L_{A,B})\} = \{(\mathcal{P}_{A,B}(\mu), h_\mu(f)) : \mu \in M_{erg}(f, X), \mathcal{P}_{A,B}(\mu) \in \text{Int}(L_{A,B})\}$.*

Example 4.1. The function $\alpha : M(f, X) \rightarrow \mathbb{R}$ can be defined as

- (1) $\alpha \equiv 0$.
- (2) $\alpha(\mu) = \int \varphi d\mu$ with a continuous function φ .
- (3) $\alpha(\mu) = \lim_{n \rightarrow \infty} \frac{1}{n} \int \varphi_n d\mu$ with an almost additive sequences of continuous functions $\Phi = (\varphi_n)_{n \in \mathbb{N}}$.

Then $\alpha : M(f, X) \rightarrow \mathbb{R}$ is a continuous function from Example 3.1. Furthermore, α is affine and thus it satisfies (4. 22) and (4. 23) if it is defined as above.

4.1 Proof of Theorem C

We first establish several auxiliary results.

Lemma 4.2. *Suppose (X, f) is a dynamical system. Let V be a convex subset of $M(f, X)$. If there is an invariant measure $\mu_V \in V$ with $S_{\mu_V} = X$, then $\{\mu \in V : S_\mu = X\}$ is residual in V .*

Proof. Since $\{\mu \in M(f, X) : S_\mu = X\}$ is either empty or residual in $M(f, X)$ from [16, Proposition 21.11]. So if there is an invariant measure $\mu_V \in V$ with $S_{\mu_V} = X$, then $\{\mu \in M(f, X) : S_\mu = X\}$ is residual in $M(f, X)$. Thus $\{\mu \in V : S_\mu = X\}$ is a G_δ subset of V . In addition, for any $\nu \in V$ and $\theta \in (0, 1)$, we have $\nu_\theta = \theta\nu + (1 - \theta)\mu_V \in V$ and $S_{\nu_\theta} = X$. So $\{\mu \in V : S_\mu = X\}$ is dense in V . \square

Lemma 4.3. *Suppose (X, f) is a dynamical system. Let $d \in \mathbb{N}$ and $(A, B) \in A(f, X)^d \times A(f, X)^d$ such that B satisfies (3. 11). If $\alpha : M(f, X) \rightarrow \mathbb{R}$ is a continuous function satisfying (4. 22) and (4. 23), then for any $a \in \text{Int}(L_{A,B})$ and any $\max_{\mu \in M_{A,B}(a)} \alpha(\mu) \leq c < H_{A,B}(f, \alpha, a)$, the following properties hold:*

- (1) If $\{\mu \in M_{A,B}^{erg}(a) : P(f, \alpha, \mu) \geq c\}$ is dense in $\{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) \geq c\}$, then $\{\mu \in M_{A,B}^{erg}(a) : P(f, \alpha, \mu) \geq c\}$ is residual in $\{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) \geq c\}$.
- (2) If there is an invariant measure with full support, then $\{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) \geq c, S_\mu = X\}$ is residual in $\{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) \geq c\}$.
- (3) If $\{\mu \in M(f, X) : h_\mu(f) = 0\}$ is dense in $M(f, X)$, then $\{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) = c\}$ is dense in $\{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) \geq c\}$. If further entropy function is upper semi-continuous, then $\{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) = c\}$ is residual in $\{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) \geq c\}$.

Proof. (1) From [16, Proposition 5.7], $M_{erg}(f, X)$ is a G_δ subset of $M(f, X)$. Then $\{\mu \in M_{A,B}^{erg}(a) : P(f, \alpha, \mu) \geq c\}$ is a G_δ subset of $\{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) \geq c\}$. If $\{\mu \in M_{A,B}^{erg}(a) : P(f, \alpha, \mu) \geq c\}$ is dense in $\{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) \geq c\}$, then $\{\mu \in M_{A,B}^{erg}(a) : P(f, \alpha, \mu) \geq c\}$ is residual in $\{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) \geq c\}$.

(2) Since $\{\mu \in M(f, X) : S_\mu = X\}$ is either empty or residual in $M(f, X)$ from [16, Proposition 21.11]. So if there is an invariant measure with full support, then $\{\mu \in M(f, X) : S_\mu = X\}$ is residual in $M(f, X)$. Now we show that $\{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) \geq c, S_\mu = X\}$ is non-empty. By Lemma 3.6 there exist 2^d invariant measures $\{\mu_\xi\}_{\xi \in \{+, -\}^d}$ such that

$$\mathcal{P}_{A,B}(\mu_\xi) \in a^\xi \text{ for any } \xi \in \{+, -\}^d.$$

Since $\{\mu \in M(f, X) : S_\mu = X\}$ is dense in $M(f, X)$ and the function $\mathcal{P}_{A,B}$ is continuous, then there exists $\omega_\xi \in M(f, X)$ close to μ_ξ such that

$$\mathcal{P}_{A,B}(\omega_\xi) \in a^\xi, S_{\omega_\xi} = X \text{ for any } \xi \in \{+, -\}^d.$$

By Corollary 3.5 there is 2^d numbers $\{\theta_\xi\}_{\xi \in \{+, -\}^d} \subseteq [0, 1]$ such that $\sum_{\xi \in \{+, -\}^d} \theta_\xi = 1$ and $\mathcal{P}_{A,B}(\omega) = a$ where $\omega = \sum_{\xi \in \{+, -\}^d} \theta_\xi \mu_\omega$. Then $S_\omega = X$. Since $c < H_{A,B}(f, \alpha, a)$, there is $\nu \in M_{A,B}(a)$ such that $P(f, \alpha, \nu) > c$. By (4. 22) we can choose $\theta \in (0, 1)$ close to 1 such that $\tilde{\mu} = \theta\nu + (1 - \theta)\omega$ satisfies $P(f, \alpha, \tilde{\mu}) > c$. Then $\tilde{\mu} \in \{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) \geq c, S_\mu = X\}$. Note that $\{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) \geq c\}$ is a convex set by (4. 22), (4. 23) and Lemma 3.3(1). So by Lemma 4.2 we complete the proof of item(2).

(3) Fix $\mu_0 \in M_{A,B}(a)$ with $P(f, \alpha, \mu_0) \geq c$ and $\zeta > 0$. By Lemma 3.6 there exist 2^d invariant measures $\{\mu_\xi\}_{\xi \in \{+, -\}^d}$ such that

$$\mathcal{P}_{A,B}(\mu_\xi) \in a^\xi \text{ and } \rho(\mu_\xi, \mu_0) < \frac{\zeta}{2} \text{ for any } \xi \in \{+, -\}^d. \quad (4. 24)$$

Since $\{\mu \in M(f, X) : h_\mu(f) = 0\}$ is dense in $M(f, X)$ and the function $\mathcal{P}_{A,B}$ is continuous, then there exist $\nu_\xi \in M(f, X)$ close to μ_ξ such that

$$\mathcal{P}_{A,B}(\nu_\xi) \in a^\xi, h_{\nu_\xi}(f) = 0 \text{ and } \rho(\nu_\xi, \mu_\xi) < \frac{\zeta}{2} \text{ for each } \xi \in \{+, -\}^d. \quad (4. 25)$$

By Corollary 3.5 there are 2^d numbers $\{\theta_\xi\}_{\xi \in \{+, -\}^d} \subseteq [0, 1]$ such that $\sum_{\xi \in \{+, -\}^d} \theta_\xi = 1$ and $\mathcal{P}_{A,B}(\nu') = a$ where $\nu' = \sum_{\xi \in \{+, -\}^d} \theta_\xi \nu_\xi$. Then by (4. 25) $h_{\nu'}(f) = 0$. By (4. 24) and (4. 25) we have $\rho(\nu', \mu_0) < \zeta$. Now by (4. 22) we choose $\theta \in [0, 1]$ such that $\nu = \theta\mu_0 + (1 - \theta)\nu'$ satisfies $P(f, \alpha, \nu) = c$. Then by Lemma 3.3(1)

$$\mathcal{P}_{A,B}(\nu) = a \text{ and } \rho(\nu, \mu_0) < \zeta. \quad (4. 26)$$

So $\{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) = c\}$ is dense in $\{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) \geq c\}$. If further entropy function is upper semi-continuous, $\{\mu \in M(f, X) : P(f, \alpha, \mu) \in [c, c + \frac{1}{n}]\}$ is open in $\{\mu \in M(f, X) : P(f, \alpha, \mu) \geq c\}$ for any $n \in \mathbb{N}^+$. Then

$$\{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) \in [c, c + \frac{1}{n}]\} \text{ is open and dense in } \{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) \geq c\},$$

and thus $\{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) = c\}$ is residual in $\{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) \geq c\}$. \square

Now we show that the result of Theorem B is the keystone of the proof of Theorem C.

Lemma 4.4. *Suppose (X, f) is a dynamical system. Let $d \in \mathbb{N}$, $(A, B) \in A(f, X)^d \times A(f, X)^d$ such that B satisfies (3. 11), and $\alpha : M(f, X) \rightarrow \mathbb{R}$ be a continuous function satisfying (4. 22) and (4. 23). Assume that for any $a \in \text{Int}(L_{A,B})$, any $\mu_0 \in M_{A,B}(a)$ and any $\eta, \zeta > 0$, there is $\nu \in M_{A,B}^{erg}(a)$ such that $\rho(\nu, \mu_0) < \zeta$ and $|P(f, \alpha, \nu) - P(f, \alpha, \mu_0)| < \eta$. If $\{\mu \in M(f, X) : h_\mu(f) = 0\}$ is dense in $M(f, X)$, then we have*

- (1) *For any $a \in \text{Int}(L_{A,B})$, any $\mu_0 \in M_{A,B}(a)$, any $\max_{\mu \in M_{A,B}(a)} \alpha(\mu) \leq c \leq P(f, \alpha, \mu_0)$ and any $\eta, \zeta > 0$, there is $\nu \in M_{A,B}^{erg}(a)$ such that $\rho(\nu, \mu_0) < \zeta$ and $|P(f, \alpha, \nu) - c| < \eta$.*

If further entropy function is upper semi-continuous, then

- (2) *For any $a \in \text{Int}(L_{A,B})$ and $\max_{\mu \in M_{A,B}(a)} \alpha(\mu) \leq c < H_{A,B}(f, \alpha, a)$, the set $\{\mu \in M_{A,B}^{erg}(a) : P(f, \alpha, \mu) = c\}$ is residual in $\{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) \geq c\}$. If further there is an invariant measure with full support, then for any $a \in \text{Int}(L_{A,B})$ and $\max_{\mu \in M_{A,B}(a)} \alpha(\mu) \leq c < H_{A,B}(f, \alpha, a)$, the set $\{\mu \in M_{A,B}^{erg}(a) : P(f, \alpha, \mu) = c, S_\mu = X\}$ is residual in $\{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) \geq c\}$.*
- (3) *$\{(\mathcal{P}_{A,B}(\mu), P(f, \alpha, \mu)) : \mu \in M(f, X), a = \mathcal{P}_{A,B}(\mu) \in \text{Int}(L_{A,B}), \max_{\mu \in M_{A,B}(a)} \alpha(\mu) \leq P(f, \alpha, \mu) < H_{A,B}(f, \alpha, a)\}$ coincides with $\{(\mathcal{P}_{A,B}(\mu), P(f, \alpha, \mu)) : \mu \in M_{erg}(f, X), a = \mathcal{P}_{A,B}(\mu) \in \text{Int}(L_{A,B}), \max_{\mu \in M_{A,B}(a)} \alpha(\mu) \leq P(f, \alpha, \mu) < H_{A,B}(f, \alpha, a)\}$.*
- (4) *If further $\text{span}\{\Phi^1, \Psi^1, \dots, \Phi^d, \Psi^d\} \subseteq E(f, X)$ and $\alpha \equiv 0$, then we have $\{(\mathcal{P}_{A,B}(\mu), h_\mu(f)) : \mu \in M(f, X), \mathcal{P}_{A,B}(\mu) \in \text{Int}(L_{A,B})\} = \{(\mathcal{P}_{A,B}(\mu), h_\mu(f)) : \mu \in M_{erg}(f, X), \mathcal{P}_{A,B}(\mu) \in \text{Int}(L_{A,B})\}$.*

Proof. (1) Fix $a \in \text{Int}(L_{A,B})$, $\mu_0 \in M_{A,B}(a)$, $\max_{\mu \in M_{A,B}(a)} \alpha(\mu) \leq c \leq P(f, \alpha, \mu_0)$ and $\eta, \zeta > 0$. By Lemma 4.3(3), there exists $\nu' \in M_{A,B}(a)$ such that $P(f, \alpha, \nu') = c$ and $\rho(\nu', \mu_0) < \frac{\zeta}{2}$. For the $a \in \text{Int}(L_{A,B})$, $\nu' \in M_{A,B}(a)$ and $\eta, \frac{\zeta}{2} > 0$, there is $\nu \in M_{A,B}^{erg}(a)$ such that $\rho(\nu, \nu') < \frac{\zeta}{2}$ and $|P(f, \alpha, \nu) - P(f, \alpha, \nu')| < \eta$. Then we complete the proof of item(1).

(2) Fix $a \in \text{Int}(L_{A,B})$ and $\max_{\mu \in M_{A,B}(a)} \alpha(\mu) \leq c < H_{A,B}(f, \alpha, a)$. First we show that

$$\{\mu \in M_{A,B}^{erg}(a) : P(f, \alpha, \mu) \geq c\} \text{ is dense in } \{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) \geq c\}. \quad (4. 27)$$

Let $\mu_0 \in M_{A,B}(a)$ be an invariant measure with $P(f, \alpha, \mu_0) \geq c$ and $\zeta > 0$. If $P(f, \alpha, \mu_0) > c$, then there is $\eta > 0$ such that $c < c + \eta < P(f, \alpha, \mu_0)$. For the $a \in \text{Int}(L_{A,B})$, $\mu_0 \in M_{A,B}(a)$ and $\eta, \zeta > 0$, there exists an ergodic measure $\nu \in M_{A,B}^{erg}(a)$ such that $\rho(\nu, \mu_0) < \zeta$ and $|P(f, \alpha, \nu) - P(f, \alpha, \mu_0)| < \eta$. If $P(f, \alpha, \mu_0) = c$, then we can pick an invariant measure $\mu' \in M_{A,B}(a)$ such that $c < P(f, \alpha, \mu') \leq H_{A,B}(f, \alpha, a)$, and next pick a sufficiently small number $\theta \in (0, 1)$ such that $\rho(\mu_0, \mu'') < \zeta/2$, where $\mu'' = (1 - \theta)\mu_0 + \theta\mu'$. By (4. 22) we have $P(f, \alpha, \mu'') > c$. By the same argument, there exists an ergodic measure $\nu \in M_{A,B}^{erg}(a)$ such that $\rho(\nu, \mu'') < \zeta/2$ and $P(f, \alpha, \nu) > c$. So $\rho(\nu, \mu_0) < \zeta$.

By (4. 27) and Lemma 4.3(1),

$$\{\mu \in M_{A,B}^{erg}(a) : P(f, \alpha, \mu) \geq c\} \text{ is residual in } \{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) \geq c\} \quad (4. 28)$$

By Lemma 4.3(3)

$$\{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) = c\} \text{ is residual in } \{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) \geq c\}. \quad (4. 29)$$

If there is an invariant measure with full support, then by Lemma 4.3(2) we have

$$\{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) \geq c, S_\mu = X\} \text{ is residual in } \{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) \geq c\} \quad (4. 30)$$

So by (4. 28), (4. 29) and (4. 30), we complete the proof of item(2).

(3) Fix $a \in \text{Int}(L_{A,B})$ and $\mu_0 \in M_{A,B}(a)$ with $\max_{\mu \in M_{A,B}(a)} \alpha(\mu) \leq P(f, \alpha, \mu_0) < H_{A,B}(f, \alpha, a)$. Then by item(2) $\{\mu \in M_{A,B}^{erg}(a) : P(f, \alpha, \mu) = P(f, \alpha, \mu_0)\}$ is residual in $\{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) \geq P(f, \alpha, \mu_0)\}$. In particular, there is $\mu_a \in M_{A,B}^{erg}(a)$ such that $P(f, \alpha, \mu_a) = P(f, \alpha, \mu_0)$.

(4) Fix $a \in \text{Int}(L_{A,B})$ and $\mu_0 \in M_{A,B}(a)$. If $h_{\mu_0}(f) = \sup\{h_\mu(f) : \mu \in M_{A,B}(a)\}$, then by Theorem 3.7, there is an ergodic measure $\mu_a \in M_{A,B}(a)$ such that

$$h_{\mu_a}(f) = h_{\text{top}}(f, R_{A,B}(a)) = h_{\mu_0}(f).$$

If $h_{\mu_0}(f) < \sup\{h_\mu(f) : \mu \in M_{A,B}(a)\}$, then by item(2) $\{\mu \in M_{A,B}^{\text{erg}}(a) : h_\mu(f) = h_{\mu_0}(f)\}$ is residual in $\{\mu \in M_{A,B}(a) : h_\mu(f) \geq h_{\mu_0}(f)\}$. In particular, there is $\mu_a \in M_{A,B}^{\text{erg}}(a)$ such that $h_{\mu_a}(f) = h_{\mu_0}(f)$. So we complete the proof of item(4). \square

Proof of Theorem C: Note that the conditions of Theorem B is contained in Theorem C. Then for any $a \in \text{Int}(L_{A,B})$, any $\mu_0 \in M_{A,B}(a)$ and any $\eta, \zeta > 0$, there is $\nu \in M_{A,B}^{\text{erg}}(a)$ such that $\rho(\nu, \mu_0) < \zeta$ and $|P(f, \alpha, \nu) - P(f, \alpha, \mu_0)| < \eta$. So we obtain Theorem C by Lemma 4.4. \square

5 Proof of Theorem A

In this section, we use 'multi-horseshoe' dense property and the results of almost additive sequences of continuous functions obtained in Section 2-4 to give a more general result than Theorem A.

5.1 Uniqueness of equilibrium measures

We first recall from [3, 4, 31] the notion of nonadditive topological pressure. Consider a dynamical system (X, f) . Let \mathcal{U} be a finite open cover of X . Given $n \in \mathbb{N}$, we denote by $\mathcal{W}_n(\mathcal{U})$ the collection of n -tuples $U = (U_1 \cdots U_n)$ with $U_1, \dots, U_n \in \mathcal{U}$. For each $U \in \mathcal{W}_n(\mathcal{U})$ we write $m(U) = n$, and we define the open set

$$X(U) = \{x \in X : f^{k-1}x \in U_k \text{ for } k = 1, \dots, m(U)\}.$$

We say that a collection $\Gamma \subset \bigcup_{n \in \mathbb{N}} \mathcal{W}_n(\mathcal{U})$ covers the set X if $\bigcup_{U \in \Gamma} X(U) \supset X$. Now let $\Phi = (\varphi_n)_{n \in \mathbb{N}}$ be a sequence of continuous functions $\varphi_n : X \rightarrow \mathbb{R}$. We define the number

$$\gamma_n(\Phi, \mathcal{U}) = \sup\{|\varphi_n(x) - \varphi_n(y)| : x, y \in X(U) \text{ with } U \in \mathcal{W}_n(\mathcal{U})\}$$

We assume that

$$\lim_{\text{diam } \mathcal{U} \rightarrow 0} \limsup_{n \rightarrow \infty} \frac{1}{n} \gamma_n(\Phi, \mathcal{U}) = 0.$$

For each n -tuple $U \in \mathcal{W}_n(\mathcal{U})$ we write $\varphi(U) = \sup_{X(U)} \varphi_n$ when $X(U) \neq \emptyset$, and $\varphi(U) = -\infty$ otherwise.

We also define

$$M(\alpha, \Phi, \mathcal{U}) = \liminf_{n \rightarrow \infty} \inf_{\Gamma} \sum_{U \in \Gamma} \exp(-\alpha m(U) + \varphi(U)) \quad (5.31)$$

where the infimum is taken over all collections $\Gamma \subset \bigcup_{k \geq n} \mathcal{W}_k(\mathcal{U})$ covering X . One can show that the quantity in (5.31) jumps from $+\infty$ to 0 at a unique value of α , and thus we can define

$$P(\Phi, \mathcal{U}) = \inf\{\alpha : M(\alpha, \Phi, \mathcal{U}) = 0\}.$$

Moreover, the limit

$$P(\Phi) = \lim_{\text{diam } \mathcal{U} \rightarrow 0} P(\Phi, \mathcal{U})$$

exists (see [3] for details). The number $P(\Phi)$ is called the nonadditive topological pressure of the sequence of functions Φ (with respect to f on X).

Let $\Phi = (\varphi_n)_{n \in \mathbb{N}}$ be an almost additive sequence of continuous functions. A measure $\mu \in M(f, X)$ is said to be an *equilibrium measure* associated with Φ if

$$P(\Phi) = h_\mu(f) + \lim_{n \rightarrow \infty} \frac{1}{n} \int \varphi_n d\mu.$$

The uniqueness of equilibrium measures was established if Φ has bounded variation. We say that Φ has *bounded variation* if there exists $\varepsilon > 0$ for which $\sup_{n \in \mathbb{N}} \gamma_n(\Phi, \varepsilon) < \infty$ with

$$\gamma_n(\Phi, \varepsilon) = \sup\{|\varphi_n(x) - \varphi_n(y)| : d(f^k(x), f^k(y)) < \varepsilon \text{ for } k = 0, \dots, n\}.$$

Lemma 5.1. [4, Page 294] Suppose that (X, f) is an expansive dynamical system satisfying specification property. Let Φ be an almost additive sequence of continuous functions with bounded variation. Then there is a unique equilibrium measure μ_Φ for Φ .

From [16, Proposition 23.20] it's known that if a homeomorphism of a compact metric space is expansive, mixing and has the shadowing property, then it satisfies specification property. So we have the following.

Corollary 5.2. Suppose that (X, f) is topologically Anosov and mixing. Let Φ be an almost additive sequence of continuous functions with bounded variation. Then there is a unique equilibrium measure μ_Φ for Φ .

Next, using a spectral decomposition theorem due to Bowen, we will show that Corollary 5.2 is still true if (X, f) is just transitive.

Theorem 5.3. Suppose that (X, f) is topologically Anosov and transitive. Let Φ be an almost additive sequence of continuous functions with bounded variation. Then there is a unique equilibrium measure μ_Φ for Φ .

Proof. Since (X, f) is expansive, transitive and has the shadowing property, by [2, Theorem 3.1.11] X admits a decomposition

$$X = \bigsqcup_{i=0}^{m-1} f^i(D)$$

where $m > 0$ is a positive integer, such that $f^i(D), 0 \leq i \leq m-1$, are closed f^m -invariant subsets of X , $f^i(D) \cap f^j(D) = \emptyset$ for any $0 \leq i < j \leq m-1$, and

$$f^m|_{f^i(D)} : f^i(D) \rightarrow f^i(D)$$

is mixing for every $0 \leq i \leq m-1$. Let $k > 0$ be an integer, then from [2, Theorem 2.3.3] a dynamical system (X, f) has the shadowing property if and only if so does (X, f^k) . So $(D, f^m|_D)$ has the shadowing property. Since (X, f) is expansive, from the uniform continuity of f, \dots, f^{m-1} , $(D, f^m|_D)$ is also expansive. So $(D, f^m|_D)$ is topologically Anosov and mixing.

For any $\mu \in M(X)$, define $\sigma(\mu) \in M(D)$ by:

$$\sigma(\mu)(A) = \mu(A \cup f(A) \cup \dots \cup f^{m-1}(A)),$$

where A is a Borel set of D . By [16, Proposition 23.17], σ is a homeomorphism from $M(f, X)$ onto $M(f^m, D)$ and

$$\sigma^{-1}(\nu) = \frac{1}{m}(\nu + f_*\nu + \dots + f_*^{m-1}\nu) \in M(f, X)$$

for any $\nu \in M(f^m, D)$ where $f_*\nu(B) = \nu(f^{-1}(B))$ for any Borel set B . Note that

$$h_{\sigma(\mu)}(f^m) = mh_\mu(f)$$

and

$$\lim_{n \rightarrow \infty} \frac{1}{n} \int \sum_{i=0}^{m-1} \varphi_n \circ f^i d\sigma(\mu) = m \lim_{n \rightarrow \infty} \frac{1}{n} \int \varphi_n d\mu.$$

Thus maximizing

$$h_{\sigma(\mu)}(f^m) + \lim_{n \rightarrow \infty} \frac{1}{n} \int \sum_{i=0}^{m-1} \varphi_n \circ f^i d\sigma(\mu)$$

is equivalent to maximizing

$$h_\mu(f) + \lim_{n \rightarrow \infty} \frac{1}{n} \int \varphi_n d\mu.$$

For $\Phi = (\varphi_n)_{n \in \mathbb{N}}$, define $\Psi = \left(\sum_{i=0}^{m-1} \varphi_n \circ f^i \right)_{n \in \mathbb{N}}$. From the uniform continuity of f, \dots, f^{m-1} , if Φ is an almost additive sequence of continuous functions with bounded variation, then so does Ψ . Since $(D, f^m|_D)$ is topologically Anosov and mixing, there is a unique equilibrium measure ν_Ψ for Ψ by Corollary 5.2. So $\sigma^{-1}(\nu_\Psi)$ is the unique equilibrium measure for Φ . \square

5.2 Proof of Theorem A

Now we show that the results of Theorem B and C hold for topologically Anosov system which is transitive. For a continuous function φ , let $\varphi_n = \varphi + \varphi \circ f + \dots + \varphi \circ f^{n-1}$ then $\{\varphi_n\}_{n \in \mathbb{N}}$ is a additive sequence of continuous functions and has bounded variation if φ has bounded variation. So let $\alpha \equiv 0$ and $d = 1$ in Theorem D, we obtain Theorem A.

Theorem D. *Suppose that (X, f) topologically Anosov and transitive. Let $d \in \mathbb{N}$ and $(A, B) \in A(f, X)^d \times A(f, X)^d$ such that B satisfies (3. 11) and Φ^i, Ψ^i has bounded variation for any $1 \leq i \leq d$. Let $\alpha : M(f, X) \rightarrow \mathbb{R}$ be a continuous function satisfying (4. 22) and (4. 23). Then:*

- (I) *For any $a \in \text{Int}(L_{A,B})$, any $\mu_0 \in M_{A,B}(a)$ and any $\eta, \zeta > 0$, there is $\nu \in M_{A,B}^{erg}(a)$ such that $\rho(\nu, \mu_0) < \zeta$ and $|P(f, \alpha, \nu) - P(f, \alpha, \mu_0)| < \eta$.*
- (II) *For any $a \in \text{Int}(L_{A,B})$, any $\mu_0 \in M_{A,B}(a)$, any $\max_{\mu \in M_{A,B}(a)} \alpha(\mu) \leq c \leq P(f, \alpha, \mu_0)$ and any $\eta, \zeta > 0$, there is $\nu \in M_{A,B}^{erg}(a)$ such that $\rho(\nu, \mu_0) < \zeta$ and $|P(f, \alpha, \nu) - c| < \eta$.*
- (III) *For any $a \in \text{Int}(L_{A,B})$ and $\max_{\mu \in M_{A,B}(a)} \alpha(\mu) \leq c < H_{A,B}(f, \alpha, a)$, $\{\mu \in M_{A,B}^{erg}(a) : P(f, \alpha, \mu) = c, S_\mu = X\}$ is residual in $\{\mu \in M_{A,B}(a) : P(f, \alpha, \mu) \geq c\}$.*
- (IV) *$\{(\mathcal{P}_{A,B}(\mu), P(f, \alpha, \mu)) : \mu \in M(f, X), a = \mathcal{P}_{A,B}(\mu) \in \text{Int}(L_{A,B}), \max_{\mu \in M_{A,B}(a)} \alpha(\mu) \leq P(f, \alpha, \mu) < H_{A,B}(f, \alpha, a)\}$ coincides with $\{(\mathcal{P}_{A,B}(\mu), P(f, \alpha, \mu)) : \mu \in M_{erg}(f, X), a = \mathcal{P}_{A,B}(\mu) \in \text{Int}(L_{A,B}), \max_{\mu \in M_{A,B}(a)} \alpha(\mu) \leq P(f, \alpha, \mu) < H_{A,B}(f, \alpha, a)\}$. If further $\alpha = 0$, then $\{(\mathcal{P}_{A,B}(\mu), h_\mu(f)) : \mu \in M(f, X), \mathcal{P}_{A,B}(\mu) \in \text{Int}(L_{A,B})\} = \{(\mathcal{P}_{A,B}(\mu), h_\mu(f)) : \mu \in M_{erg}(f, X), \mathcal{P}_{A,B}(\mu) \in \text{Int}(L_{A,B})\}$.*

Proof. Since (X, f) is expansive, then the entropy function is upper semi-continuous from [41, Theorem 8.2]. Since Φ^i, Ψ^i has bounded variation for any $1 \leq i \leq d$, then every sequence in $\text{span}\{\Phi^1, \Psi^1, \dots, \Phi^d, \Psi^d\}$ has bounded variation. Thus by Theorem 5.3 we have $\text{span}\{\Phi^1, \Psi^1, \dots, \Phi^d, \Psi^d\} \subseteq E(f, X)$, and $\text{span}\{\Phi^1|_\Lambda, \Psi^1|_\Lambda, \dots, \Phi^d|_\Lambda, \Psi^d|_\Lambda\} \subseteq E(f|_\Lambda, \Lambda)$ if $(\Lambda, f|_\Lambda)$ is transitive and topologically Anosov. Then we complete the proof by Theorem 2.6, Theorem B, Theorem C and Corollary 2.11. \square

6 Applications

In this section, we apply the results in the previous sections to transitive locally maximal hyperbolic sets and transitive two-side subshifts of finite type.

6.1 Hyperbolic diffeomorphisms

We now suppose that $f : M \rightarrow M$ is a diffeomorphism of a compact C^∞ Riemannian manifold M . Then the derivative of f can be considered a map $df : TM \rightarrow TM$ where $TM = \bigcup_{x \in M} T_x M$ is the tangent bundle of M and $df_x : T_x M \rightarrow T_{f(x)} M$. A closed subset $\Lambda \subset M$ is *hyperbolic* if $f(\Lambda) = \Lambda$ and each tangent space $T_x M$ with $x \in \Lambda$ can be written as a direct sum $T_x M = E_x^u \oplus E_x^s$ of subspaces so that

1. $Df(E_x^s) = E_{f(x)}^s, Df(E_x^u) = E_{f(x)}^u$;
2. there exist constants $c > 0$ and $\lambda \in (0, 1)$ so that

$$\|Df^n(v)\| \leq c\lambda^n \|v\| \text{ when } v \in E_x^s, n \geq 1, \text{ and } \|Df^{-n}(v)\| \leq c\lambda^n \|v\| \text{ when } v \in E_x^u, n \geq 1.$$

A hyperbolic set Λ is said to be locally maximal for f if there exists a neighborhood U of Λ in M such that $\Lambda = \bigcap_{n=-\infty}^{+\infty} f^n(U)$. It's known that a locally maximal hyperbolic set is expansive by [25, Corollary 6.4.10] and has shadowing property by [25, Theorem 18.1.2].

Theorem 6.1. *Suppose that (X, f) is a system restricted a transitive locally maximal hyperbolic set. Let $d \in \mathbb{N}$ and $(A, B) \in A(f, X)^d \times A(f, X)^d$ such that B satisfies (3. 11) and Φ^i, Ψ^i has bounded variation for any $1 \leq i \leq d$. Let $\alpha : M(f, X) \rightarrow \mathbb{R}$ be a continuous function satisfying (4. 22) and (4. 23). Then the results of Theorem D hold.*

For a Hölder continuous function φ , let $\varphi_n = \varphi + \varphi \circ f + \dots + \varphi \circ f^{n-1}$ then $\{\varphi_n\}_{n \in \mathbb{N}}$ is a additive sequence of contiuous functions and from [12, Example 2] it has bounded variation. So we have

Theorem 6.2. *Suppose that (X, f) is a system restricted a transitive locally maximal hyperbolic set. Let φ be a Hölder continuous function. Then the results of Theorem A hold.*

6.2 Transitive two-side subshifts of finite type

Let k be a fixed natural number and let $C = \{0, 1, \dots, k-1\}$. Put the discrete topology on C . Consider the two-side full symbolic space $\Sigma = \prod_{-\infty}^{\infty} C$, equipped with the product topology, and the shift homeomorphism $\sigma : \Sigma \rightarrow \Sigma$ defined by $(\sigma(w))_n = w_{n+1}$, where $w = (w_n)_{-\infty}^{\infty}$. A metric on Σ is defined by $d(x, y) = 2^{-m}$ if m is the largest natural number with $x_n = y_n$ for any $|n| < m$, and $d(x, y) = 1$ if $x_0 \neq y_0$. If X is a closed subset of Σ with $\sigma X = X$ then $\sigma|_X : X \rightarrow X$ is called a subshift. We usually write this as $\sigma : X \rightarrow X$. A subshift $\sigma : X \rightarrow X$ is said to be of finite type if there exists some natural number N and a collection of blocks of length $N+1$ with the property that $x = (x_n)_{-\infty}^{\infty} \in X$ if and only if each block (x_i, \dots, x_{i+N}) in x of length $N+1$ is one of the prescribed blocks.

Recall from [42] a subshift satisfies shadowing property if and only if it is a subshift of finite type. As a subsystem of two-side full shift, it is expansive. So we have

Theorem 6.3. *Suppose that (X, f) is a transitive two-side subshit of finite type. Let $d \in \mathbb{N}$ and $(A, B) \in A(f, X)^d \times A(f, X)^d$ such that B satisfies (3. 11) and Φ^i, Ψ^i has bounded variation for any $1 \leq i \leq d$. Let $\alpha : M(f, X) \rightarrow \mathbb{R}$ be a continuous function satisfying (4. 22) and (4. 23). Then the results of Theorem D hold.*

Lemma 6.4. *Let $x, y \in \Sigma_2$, $n \in \mathbb{Z}^+$, and $\varepsilon > 0$. If $d(\sigma^i(x), \sigma^i(y)) < \varepsilon$ for any $0 \leq i \leq n$, then we have $d(\sigma^i(x), \sigma^i(y)) < \varepsilon \cdot 2^{-\min\{i, n-i\}}$ for any $0 \leq i \leq n$.*

Proof. For the $\varepsilon > 0$, there is an integer i such that $\frac{1}{2^{j+1}} < \varepsilon \leq \frac{1}{2^j}$. Then for any $0 \leq i \leq n$, by $d(\sigma^i(x), \sigma^i(y)) < \varepsilon$, we have $(\sigma^i(x))_l = (\sigma^i(y))_l$ for any $-j \leq l \leq j$. Thus we have $x_l = y_l$ for any $-j \leq l \leq n+j$. This implies $(\sigma^i(x))_l = (\sigma^i(y))_l$ for any $-j-i \leq l \leq n+j-i$ and any $0 \leq i \leq n$. So

$$d(\sigma^i(x), \sigma^i(y)) \leq 2^{-\min\{i+j, n+j-i\}-1} = 2^{-j-1} \cdot 2^{-\min\{i, n-i\}} < \varepsilon \cdot 2^{-\min\{i, n-i\}}.$$

for any $0 \leq i \leq n$. □

Let φ a Hölder continuous function on Σ_2 with constant K and exponent α . If $x, y \in \Sigma_2$, $n \in \mathbb{Z}^+$, and $\varepsilon > 0$ satisfy $d(\sigma^i(x), \sigma^i(y)) < \varepsilon$ for any $0 \leq i \leq n$, then we have

$$\begin{aligned} \left| \sum_{k=0}^n \varphi(\sigma^k(x)) - \sum_{k=0}^n \varphi(\sigma^k(y)) \right| &\leq \sum_{k=0}^n |\varphi(\sigma^k(x)) - \varphi(\sigma^k(y))| \\ &\leq \sum_{k=0}^n K(d(\sigma^k(x), \sigma^k(y)))^\alpha \\ &\leq \sum_{k=0}^n K(\varepsilon \cdot 2^{-\min\{i, n-i\}})^\alpha \\ &\leq K(\varepsilon)^\alpha \cdot \frac{1}{1-2^{-\alpha}}. \end{aligned}$$

This implies that every Hölder continuous function on Σ_2 has bounded variation. Same as Theorem 6.2, we have

Theorem 6.5. *Suppose that (X, f) is a transitive two-side subshit of finite type. Let φ be a Hölder continuous function. Then the results of Theorem A hold.*

6.3 Homoclinic classes

In this subsection, we consider homoclinic classes and give corresponding results on refined Katok's conjecture. Let f be a C^1 diffeomorphism on a compact Riemannian manifold M , we recall that the homoclinic class of a hyperbolic saddle p , denoted by $H(p)$, is the closure of the set of hyperbolic saddles q homoclinically related to p (the stable manifold of the orbit of q transversely meets the unstable one of the orbit of p and vice versa).

Let $X = H(p)$ be a homoclinic class. We denote $M^{horse}(H(p))$ the set of measures which can be approximated by horseshoes, that is, an invariant measure $\mu \in M(f, X)$ is in $M^{horse}(H(p))$ if and only if for any $\varepsilon > 0$, there is a f -invariant compact subset $\Lambda_\varepsilon \subseteq H(p)$ such that $\mu_\varepsilon := \mu|_{\Lambda_\varepsilon}$ satisfies the following three properties

- (1) $(\Lambda_\varepsilon, f|_{\Lambda_\varepsilon})$ is a transitive locally maximal hyperbolic set which contains a hyperbolic saddles q homoclinically related to p .
- (2) $\rho(\mu, \mu_\varepsilon) < \varepsilon$.
- (3) $h_{\mu_\varepsilon}(f) > h_\mu(f) - \varepsilon$.

We denote $M_{erg}^{horse}(H(p)) = M^{horse}(H(p)) \cap M_{erg}(f, X)$. Let $d \in \mathbb{N}$ and $(A, B) \in A(f, X)^d \times A(f, X)^d$ such that B satisfies (3. 11). Denote

$$L_{A,B}^{horse} = \{\mathcal{P}_{A,B}(\mu) : \mu \in M^{horse}(H(p))\}.$$

For any $a \in L_{A,B}^{horse}$, define

$$M_{A,B}^{horse}(a) = \{\mu \in M^{horse}(H(p)) : \mathcal{P}_{A,B}(\mu) = a\},$$

$$M_{A,B}^{erg,horse}(a) = \{\mu \in M_{erg}^{horse}(H(p)) : \mathcal{P}_{A,B}(\mu) = a\}.$$

Let $\alpha : M(f, X) \rightarrow \mathbb{R}$ be a continuous function. We define the pressure of α with respect to μ by $P(f, \alpha, \mu) = h_\mu(f) + \alpha(\mu)$. For any $a \in L_{A,B}^{horse}$, denote

$$H_{A,B}^{horse}(f, \alpha, a) = \sup\{P(f, \alpha, \mu) : \mu \in M_{A,B}^{horse}(a)\}.$$

In particular, when $\alpha \equiv 0$, we write

$$H_{A,B}^{horse}(f, a) = H_{A,B}^{horse}(f, 0, a) = \sup\{h_\mu(f) : \mu \in M_{A,B}^{horse}(a)\}.$$

Theorem E. *Let f be a C^1 diffeomorphism on a compact Riemannian manifold M , and $X = H(p)$ be a nontrivial homoclinic class. Assume that the entropy function of (X, f) is upper semi-continuous. Let $d \in \mathbb{N}$ and $(A, B) \in A(f, X)^d \times A(f, X)^d$ such that B satisfies (3. 11) and Φ^i, Ψ^i has bounded variation for any $1 \leq i \leq d$. Let $\alpha : M(f, X) \rightarrow \mathbb{R}$ be a continuous function satisfying (4. 22) and (4. 23). Then:*

- (I) *For any $a \in \text{Int}(L_{A,B}^{horse})$, any $\mu_0 \in M_{A,B}^{horse}(a)$ and any $\eta, \zeta > 0$, there is $\nu \in M_{A,B}^{erg,horse}(a)$ such that $\rho(\nu, \mu_0) < \zeta$ and $|P(f, \alpha, \nu) - P(f, \alpha, \mu_0)| < \eta$.*
- (II) *For any $a \in \text{Int}(L_{A,B}^{horse})$, any $\mu_0 \in M_{A,B}^{horse}(a)$, any $\max_{\mu \in M_{A,B}^{horse}(a)} \alpha(\mu) \leq c \leq P(f, \alpha, \mu_0)$ and any $\eta, \zeta > 0$, there is $\nu \in M_{A,B}^{erg,horse}(a)$ such that $\rho(\nu, \mu_0) < \zeta$ and $|P(f, \alpha, \nu) - c| < \eta$.*
- (III) *For any $a \in \text{Int}(L_{A,B}^{horse})$ and $\max_{\mu \in M_{A,B}^{horse}(a)} \alpha(\mu) \leq c < H_{A,B}^{horse}(f, \alpha, a)$, $\{\mu \in M_{A,B}^{erg,horse}(a) : P(f, \alpha, \mu) = c\}$ is residual in $\{\mu \in M_{A,B}^{horse}(a) : P(f, \alpha, \mu) \geq c\}$.*
- (IV) *$\{(\mathcal{P}_{A,B}(\mu), P(f, \alpha, \mu)) : \mu \in M^{horse}(H(p)), a = \mathcal{P}_{A,B}(\mu) \in \text{Int}(L_{A,B}^{horse}), \max_{\mu \in M_{A,B}^{horse}(a)} \alpha(\mu) \leq P(f, \alpha, \mu) < H_{A,B}^{horse}(f, \alpha, a)\}$ coincides with $\{(\mathcal{P}_{A,B}(\mu), P(f, \alpha, \mu)) : \mu \in M_{erg}^{horse}(H(p)), a = \mathcal{P}_{A,B}(\mu) \in \text{Int}(L_{A,B}^{horse}), \max_{\mu \in M_{A,B}^{horse}(a)} \alpha(\mu) \leq P(f, \alpha, \mu) < H_{A,B}^{horse}(f, \alpha, a)\}$.*

There are many transitive systems for which the whole space is a homoclinic class and the entropy function is upper semi-continuous including

(i) the nonuniformly hyperbolic diffeomorphisms constructed by Katok [23]. For arbitrary compact connected two-dimensional manifold M , A. Katok proved that there exists a C^∞ diffeomorphism f such that the Riemannian volume m is an f -invariant ergodic hyperbolic measure. From [24] (or Theorem S.5.3 on Page 694 of book [25]) we know that the support of any ergodic and non-atomic hyperbolic measure of a $C^{1+\alpha}$ diffeomorphism is contained in a non-trivial homoclinic class, then there is a hyperbolic periodic point p such that $M = S_m = H(p)$. Moreover, J. Buzzi [14] showed that every C^∞ diffeomorphism is asymptotically entropy expansive which implies that the entropy function is upper semi-continuous by [16, Theorem 20.9].

(ii) generic systems in the space of robustly transitive diffeomorphisms $\text{Diff}_{RT}^1(M)$. By the robustly transitive partially hyperbolic diffeomorphisms constructed by Mañé [30] and the robustly transitive nonpartially hyperbolic diffeomorphisms constructed by Bonatti and Viana [10], we know that $\text{Diff}_{RT}^1(M)$ is a non-empty open set in $\text{Diff}^1(M)$. Since any non-trivial isolated transitive set of C^1 generic diffeomorphism is a non-trivial homoclinic class [9], we have that

$$\mathcal{R}_1 = \{f \in \text{Diff}_{RT}^1(M) : \text{there is a hyperbolic periodic point } p \text{ such that } M = H(p)\}$$

is generic in $\text{Diff}_{RT}^1(M)$. Moreover, C^1 generically in any dimension, isolated homoclinic classes are entropy expansive [34]. Since entropy expansive implies upper semi-continuous of the entropy function by [16, Theorem 20.9], then we have that

$$\mathcal{R}_2 = \{f \in \mathcal{R}_1 : \text{the entropy function is upper semi-continuous}\}$$

is generic in $\text{Diff}_{RT}^1(M)$.

(iii) generic systems in the space of volume-preserving diffeomorphisms $\text{Diff}_{vol}^1(M)$. Let M be a compact connected Riemannian manifold. Bonatti and Crovisier proved in [8, Theorem 1.3] that there exists a residual C^1 -subset \mathcal{R}_1 of $\text{Diff}_{vol}^1(M)$ such that if $f \in \mathcal{R}_1$ then f is a transitive diffeomorphism. Moreover, by its proof on page 79 and page 87 of [8], if $f \in \mathcal{R}_1$ then there is a hyperbolic periodic point p such that $M = H(p)$. Since the space of diffeomorphisms away from homoclinic tangencies $\text{Diff}^1(M) \setminus \overline{HT}$ is open in $\text{Diff}^1(M)$, then $\mathcal{R}_2 = \mathcal{R}_1 \cap \text{Diff}^1(M) \setminus \overline{HT}$ is generic in $\text{Diff}_{vol}^1(M) \setminus \overline{HT}$. Moreover, every C^1 diffeomorphism away from homoclinic tangencies is entropy expansive [29]. Note that entropy expansive implies upper semi-continuous of the entropy function by [16, Theorem 20.9], if $f \in \mathcal{R}_2$ then there is a hyperbolic periodic point p such that $M = H(p)$ and the entropy function is upper semi-continuous.

6.3.1 Some lemmas

Now we give some results corresponding to section 2, 3 and 4.

Lemma 6.6. *Let f be a C^1 diffeomorphism on a compact Riemannian manifold M , and $H(p)$ be a nontrivial homoclinic class. Then the set $M^{horse}(H(p))$ is convex.*

Proof. Fix $\mu_1, \mu_2 \in M^{horse}(H(p))$ and $\theta \in [0, 1]$. Then for any $\varepsilon > 0$ and $i \in \{1, 2\}$, there is a f -invariant compact subset $\Lambda_\varepsilon^i \subseteq H(p)$ such that $\mu_\varepsilon^i := \mu^i|_{\Lambda_\varepsilon^i}$ satisfies the following three properties

- (1) $(\Lambda_\varepsilon^i, f|_{\Lambda_\varepsilon^i})$ is a transitive locally maximal hyperbolic set which contains a hyperbolic saddles q_i homoclinically related to p .
- (2) $\rho(\mu_i, \mu_\varepsilon^i) < \varepsilon$.
- (3) $h_{\mu_\varepsilon^i}(f) > h_{\mu_i}(f) - \varepsilon$.

Then q_1 is homoclinically related to q_2 , since homoclinically related is an equivalence relation by [32, Proposition 2.1]. This implies that there is a transitive locally maximal hyperbolic set Λ_ε which contains Λ_ε^1 and Λ_ε^2 (for example, see [33, Lemma 8]). Let $\mu_\varepsilon = \theta\mu_\varepsilon^1 + (1 - \theta)\mu_\varepsilon^2$. Then we have

$$\rho(\theta\mu_1 + (1 - \theta)\mu_2, \mu_\varepsilon) \leq \theta\rho(\mu_1, \mu_\varepsilon^1) + (1 - \theta)\rho(\mu_2, \mu_\varepsilon^2) < \varepsilon.$$

and

$$h_{\mu_\varepsilon}(f) = \theta h_{\mu_\varepsilon^1}(f) + (1 - \theta)h_{\mu_\varepsilon^2}(f) > \theta h_{\mu_1}(f) + (1 - \theta)h_{\mu_2}(f) - \varepsilon = h_{\theta\mu_1 + (1-\theta)\mu_2}(f) - \varepsilon.$$

Note that μ_ε is supported on Λ_ε . So $\theta\mu_1 + (1-\theta)\mu_2 \in M^{\text{horse}}(H(p))$ and thus $M^{\text{horse}}(H(p))$ is convex. \square

Theorem 6.7. *Let f be a C^1 diffeomorphism on a compact Riemannian manifold M , and $X = H(p)$ be a nontrivial homoclinic class. Then (X, f) satisfies the 'multi-horseshoe' entropy-dense property on $M^{\text{horse}}(H(p))$, that is, for any $K = \text{cov}\{\mu_i\}_{i=1}^m \subseteq M^{\text{horse}}(H(p))$, and any $\eta, \zeta > 0$, there exist compact invariant subsets $\Lambda_i \subseteq \Lambda \subsetneq H(p)$ such that for each $1 \leq i \leq m$*

1. $(\Lambda_i, f|_{\Lambda_i})$ and $(\Lambda, f|_\Lambda)$ are transitive locally maximal hyperbolic sets.
2. $h_{\text{top}}(f, \Lambda_i) > h_{\mu_i}(f) - \eta$.
3. $d_H(K, M(f, \Lambda)) < \zeta$, $d_H(\mu_i, M(f, \Lambda_i)) < \zeta$.

Proof. Fix $K = \text{cov}\{\mu_i\}_{i=1}^m \subseteq M^{\text{horse}}(H(p))$, and any $\eta, \zeta > 0$. Denote $\tau = \frac{1}{2} \min\{\eta, \zeta\}$. Then for any $1 \leq i \leq m$ there is a f -invariant compact subset $\Lambda_\tau^i \subseteq H(p)$ such that $\mu_\tau^i := \mu^i|_{\Lambda_\tau}$ satisfies the following three properties

- (1) $(\Lambda_\tau^i, f|_{\Lambda_\tau^i})$ is a transitive locally maximal hyperbolic set which contains a hyperbolic saddles q_i homoclinically related to p .
- (2) $\rho(\mu_i, \mu_\tau^i) < \tau$.
- (3) $h_{\mu_\tau^i}(f) > h_{\mu_i}(f) - \tau$.

Then q_i is homoclinically related to q_j , since homoclinically related is an equivalence relation by [32, Proposition 2.1]. This implies that there is a transitive locally maximal hyperbolic set Λ_τ which contains $\bigcup_{i=1}^m \Lambda_\tau^i$. Note that a transitive locally maximal hyperbolic set is expansive by [25, Corollary 6.4.10] and has shadowing property by [25, Theorem 18.1.2]. Then $(\Lambda_\tau, f|_{\Lambda_\tau})$ has the 'multi-horseshoe' entropy-dense property by Theorem 2.6. Thus for the $K_\tau = \text{cov}\{\mu_\tau^i\}_{i=1}^m \subseteq M(f, \Lambda_\tau)$, and $\tau > 0$, there exist compact invariant subsets $\Lambda_i \subseteq \Lambda \subsetneq X$ such that for each $1 \leq i \leq m$

1. $(\Lambda_i, f|_{\Lambda_i})$ and $(\Lambda, f|_\Lambda)$ conjugate to transitive subshifts of finite type.
2. $h_{\text{top}}(f, \Lambda_i) > h_{\mu_\tau^i}(f) - \tau > h_{\mu_i}(f) - 2\tau > h_{\mu_i}(f) - \eta$.
3. $d_H(K_\tau, M(f, \Lambda)) < \tau$, $d_H(\mu_\tau^i, M(f, \Lambda_i)) < \tau$.

From [1] any hyperbolic set conjugate to a subshift of finite type is a locally maximal. So $(\Lambda_i, f|_{\Lambda_i})$ and $(\Lambda, f|_\Lambda)$ are transitive locally maximal hyperbolic sets. By item 3 we have

$$d_H(K, M(f, \Lambda)) < d_H(K, K_\tau) + d_H(K_\tau, M(f, \Lambda)) < 2\tau < \zeta,$$

$$d_H(\mu_i, M(f, \Lambda_i)) < d_H(\mu_i, \mu_\tau^i) + d_H(\mu_\tau^i, M(f, \Lambda_i)) < 2\tau < \zeta.$$

So we complete the proof. \square

If we replace $M(f, X)$ by $M^{\text{horse}}(H(p))$ in Section 3 and 4, the results of Theorem B and C also hold. In fact, the convexity of $M(f, X)$ is one core property in the proof of Theorem B and C. By Lemma 6.6 the set $M^{\text{horse}}(H(p))$ is convex, then the arguments of Section 3 and 4 are also true. Here, we omit the proof.

Theorem 6.8. *Let f be a C^1 diffeomorphism on a compact Riemannian manifold M , and $X = H(p)$ be a nontrivial homoclinic class. Assume that the entropy function of (X, f) is upper semi-continuous. Let $d \in \mathbb{N}$ and $(A, B) \in A(f, X)^d \times A(f, X)^d$ such that B satisfies (3. 11). Assume that the following holds: for any $K = \text{cov}\{\mu_i\}_{i=1}^m \subseteq M^{\text{horse}}(H(p))$, and any $\eta, \zeta > 0$, there exist compact invariant subsets $\Lambda_i \subseteq \Lambda \subsetneq X$ such that for each $i \in \{1, 2, \dots, m\}$*

- (1) $h_{top}(f, \Lambda_i) > h_{\mu_i}(f) - \eta$.
- (2) $d_H(K, M(f, \Lambda)) < \zeta$, $d_H(\mu_i, M(f, \Lambda_i)) < \zeta$.
- (3) $\text{span}\{\Phi^1|_\Lambda, \Psi^1|_\Lambda, \dots, \Phi^d|_\Lambda, \Psi^d|_\Lambda\} \subseteq E(f|_\Lambda, \Lambda)$.

Let $\alpha : M(f, X) \rightarrow \mathbb{R}$ be a continuous function. Then:

- (I) For any $a \in \text{Int}(L_{A,B}^{horse})$, any $\mu_0 \in M_{A,B}^{horse}(a)$ and any $\eta, \zeta > 0$, there is $\nu \in M_{A,B}^{erg,horse}(a)$ such that $\rho(\nu, \mu_0) < \zeta$ and $|P(f, \alpha, \nu) - P(f, \alpha, \mu_0)| < \eta$.

If further $\{\mu \in M^{horse}(H(p)) : h_\mu(f) = 0\}$ is dense in $M^{horse}(H(p))$, and α satisfies (4. 22) and (4. 23), then we have

- (II) For any $a \in \text{Int}(L_{A,B}^{horse})$, any $\mu_0 \in M_{A,B}^{horse}(a)$, any $\max_{\mu \in M_{A,B}^{horse}(a)} \alpha(\mu) \leq c \leq P(f, \alpha, \mu_0)$ and any $\eta, \zeta > 0$, there is $\nu \in M_{A,B}^{erg,horse}(a)$ such that $\rho(\nu, \mu_0) < \zeta$ and $|P(f, \alpha, \nu) - c| < \eta$.
- (III) For any $a \in \text{Int}(L_{A,B}^{horse})$ and $\max_{\mu \in M_{A,B}^{horse}(a)} \alpha(\mu) \leq c < H_{A,B}^{horse}(f, \alpha, a)$, the set $\{\mu \in M_{A,B}^{erg,horse}(a) : P(f, \alpha, \mu) = c\}$ is residual in $\{\mu \in M_{A,B}^{horse}(a) : P(f, \alpha, \mu) \geq c\}$.
- (IV) $\{(\mathcal{P}_{A,B}(\mu), P(f, \alpha, \mu)) : \mu \in M^{horse}(H(p)), a = \mathcal{P}_{A,B}(\mu) \in \text{Int}(L_{A,B}^{horse}), \max_{\mu \in M_{A,B}^{horse}(a)} \alpha(\mu) \leq P(f, \alpha, \mu) < H_{A,B}^{horse}(f, \alpha, a)\}$ coincides with $\{(\mathcal{P}_{A,B}(\mu), P(f, \alpha, \mu)) : \mu \in M_{A,B}^{erg,horse}(H(p)), a = \mathcal{P}_{A,B}(\mu) \in \text{Int}(L_{A,B}^{horse}), \max_{\mu \in M_{A,B}^{horse}(a)} \alpha(\mu) \leq P(f, \alpha, \mu) < H_{A,B}^{horse}(f, \alpha, a)\}$.

By Theorem 6.7 and Corollary 2.11(1), we have that the set of invariant measures with zero metric entropy is dense in $M^{horse}(H(p))$.

Lemma 6.9. *Let f be a C^1 diffeomorphism on a compact Riemannian manifold M , and $X = H(p)$ be a nontrivial homoclinic class. Then $\{\mu \in M^{horse}(H(p)) : h_\mu(f) = 0\}$ is dense in $M^{horse}(H(p))$.*

6.3.2 Proof of Theorem E

Since Φ^i, Ψ^i has bounded variation for any $1 \leq i \leq d$, then every sequence in $\text{span}\{\Phi^1, \Psi^1, \dots, \Phi^d, \Psi^d\}$ has bounded variation. Thus by Theorem 5.3 we have $\text{span}\{\Phi^1|_\Lambda, \Psi^1|_\Lambda, \dots, \Phi^d|_\Lambda, \Psi^d|_\Lambda\} \subseteq E(f|_\Lambda, \Lambda)$ if $(\Lambda, f|_\Lambda)$ is transitive and topologically Anosov. Then we complete the proof by Theorem 6.7, Theorem 6.8 and Lemma 6.9. \square

Acknowledgements. X. Hou and X. Tian are supported by National Natural Science Foundation of China (grant No. 12071082,11790273) and in part by Shanghai Science and Technology Research Program (grant No. 21JC1400700).

References

- [1] D. Anosov, *Extension of zero-dimensional hyperbolic sets to locally maximal ones*, Sb. Math. 201 (7–8) (2010) 935–946.
- [2] N. Aoki and K. Hiraide, *Topological Theory of Dynamical Systems: Recent Advances*, Vol. 52. Elsevier, Amsterdam, 1994.
- [3] L. Barreira, *A non-additive thermodynamic formalism and application to dimension theory of hyperbolic dynamical systems*, Ergod. Th. & Dyn. Sys. 16 (1996), 871–928.
- [4] L. Barreira, *Nonadditive thermodynamic formalism: equilibrium and Gibbs measures*, Discrete Contin. Dyn. Syst. 16 (2006), 279–305.
- [5] L. Barreira and P. Doutor, *Almost additive multi-fractal analysis*, J. Math. Pures Appl. (9) 92(2009), 1–17.
- [6] P. Billingsley, *Hausdorff dimension in probability theory*, Illinois. J. Math. 4 (1960), 187–209.
- [7] P. Billingsley, *Hausdorff dimension in probability theory II*, Illinois. J. Math. 5 (1961), 291–298.
- [8] C. Bonatti and S. Crovisier, *Réurrence et généricité*, Invent. Math., 158 (2004), 33–104.

- [9] C. Bonatti and L. Díaz, *Connexions hétéroclines et genericité d'une infinité de puits et de sources*, Ann. Sci. École Norm. Sup. 32 (1999) 135–150.
- [10] C. Bonatti and M. Viana, *SRB measures for partially hyperbolic systems whose central direction is mostly contracting*, Israel J. Math. 115(2000), 157–219.
- [11] R. Bowen, *Topological entropy for noncompact sets*, Trans. Amer. Math. Soc. 184 (1973), 125–136.
- [12] R. Bowen, *Some systems with unique equilibrium states*, Math. Syst. Theory 8(3) (1974/75), 193–202.
- [13] D. Burguet, *Topological and almost Borel universality for systems with the weak specification property*, Ergod. Th. & Dynam. Sys. 40 (8) (2020), 2098–2115.
- [14] J. Buzzi, *Intrinsic ergodicity of smooth interval maps*, Israel J. Math., 100 (1997), 125–161.
- [15] N. Cuneo, *Additive, almost additive and asymptotically additive potential sequences are equivalent*, Commun. Math. Phys. 377 (2020), 2579–2595
- [16] M. Denker, C. Grillenberger and K. Sigmund, *Ergodic theory on the compact space*, Lecture Notes in Mathematics 527.
- [17] Y. Dong, X. Hou and X. Tian, *Twelve different statistical future of dynamical orbits: empty syndetic center*, Preprint, 2022, arXiv:1701.01910v4.
- [18] A. Eizenberg, Y. Kifer and B. Weiss, *Large deviations for \mathbb{Z}^d -actions*, Comm. Math. Phys. 164 (1994), 433–454.
- [19] D. Feng and W. Huang, *Lyapunov spectrum of asymptotically sub-additive potentials*, Commun. Math. Phys. 297(1) (2010), 1–43.
- [20] H. Föllmer and S. Orey, *Large deviations for the empirical field of a Gibbs measure*, Ann. Probab. 16 (1988), 961–977.
- [21] L. Guan, P. Sun and W. Wu, *Measures of intermediate entropies and homogeneous dynamics*, Nonlinearity 9 (2017), 3349–3361.
- [22] X. Hou and X. Tian, *Intermediate entropy property of ergodic measures with same level for hyperbolic flows*, in preparation.
- [23] A. Katok, *Bernoulli diffeomorphisms on surfaces*, Ann. of Math. (2) 110(3) (1979), 529–547.
- [24] A. Katok, *Lyapunov exponents, entropy and periodic orbits for diffeomorphisms*, Publ. Math. I. H. E. S. , 51 (1980), 137–174.
- [25] A. Katok and B. Hasselblatt, *Introduction to the Modern Theory of Dynamical Systems (Encyclopedia of Mathematics and Its Applications, 54)*, Cambridge University Press, Cambridge, 1995.
- [26] J. Konieczny, M. Kupsa, and D. Kwietniak, *Arcwise connectedness of the set of ergodic measures of hereditary shifts*, Proc. Amer. Math. Soc. 146 (2018), 3425–3438.
- [27] J. Li and P. Oprocha, *Properties of invariant measures in dynamical systems with the shadowing property*, Ergod. Th. & Dynam. Sys. 38 (2018), 2257–2294.
- [28] M. Li, Y. Shi, S. Wang and X. Wang, *Measures of intermediate entropies for star vector fields*, Israel J. math. 240(2) (2020), 791–819.
- [29] G. Liao, M. Viana, J. Yang, *The entropy conjecture for diffeomorphisms away from tangencies*, J. Eur. Math. Soc. 15 (6) (2013) 2043–2060.
- [30] R. Mañé, *Contributions to the stability conjecture*, Topology 17 (1978) 383–396.
- [31] A. Mummert, *The thermodynamic formalism for almost-additive sequences*, Discrete Contin. Dyn. Syst. 16 (2006), 435–454.
- [32] S. E. Newhouse, *Hyperbolic limit sets*, Trans. Amer. Math. Soc. 167 (1972), 125–150.
- [33] S. E. Newhouse, *The abundance of wild hyperbolic sets and nonsmooth stable sets for diffeomorphisms*, Inst. Hautes Études Sci. Publ. Math. 50 (1979), 101–151.
- [34] M. J. Pacifico and J. L. Vieitez, *Entropy-expansiveness and domination for surface diffeomorphisms*, Rev. Mat. Complut. 21 (2008), 293–317.
- [35] C. E. Pfister and W. G. Sullivan, *Billingsley dimension on shift spaces*, Nonlinearity 16(2) (2003), 661–682.
- [36] C. E. Pfister and W. G. Sullivan, *Large deviations estimates for dynamical systems without the specification property*, Application to the β -shifts, Nonlinearity 18 (2005), 237–261.

- [37] A. Quas, and T. Soo, *Ergodic universality of some topological dynamical systems*, Trans. Amer. Math. Soc. 368 (2016), 4137–4170.
- [38] P. Sun, *Ergodic measures of intermediate entropies for dynamical systems with approximate product property*, Preprint, 2019, arXiv:1906.09862.
- [39] X. Tian, *Different asymptotic behaviour versus same dynamical complexity: recurrence & (ir)regularity.*, Adv. Math. 288 (2016), 464–526.
- [40] R. Ures, *Intrinsic ergodicity of partially hyperbolic diffeomorphisms with a hyperbolic linear part*, Proc. Amer. Math. Soc. 140(6) (2012), 1973–1985.
- [41] P. Walters, *An introduction to ergodic theory*, Springer-Verlag, Berlin, 2001.
- [42] P. Walters, *On the pseudo-orbit tracing property and its relationship to stability*, in: *The Structure of Attractors in Dynamical Systems (Fargo, ND, 1977)*, Lecture Notes in Math. 668, Springer, Berlin, 1978, 231–244.
- [43] Y. Zhao, L. Zhang and Y. Cao, *The asymptotically additive topological pressure on the irregular set for asymptotically additive potentials*, Nonlinear Anal. 74 (2011), 5015–5022.