

EVALUATION BIREPRESENTATIONS OF AFFINE TYPE A SOERTEL BIMODULES

MARCO MACKAAY, VANESSA MIEMIETZ, AND PEDRO VAZ

ABSTRACT. In this paper, we use Soergel calculus to define a monoidal functor, called the evaluation functor, from extended affine type A Soergel bimodules to the homotopy category of bounded complexes in finite type A Soergel bimodules. This functor categorifies the well-known evaluation homomorphism from the extended affine type A Hecke algebra to the finite type A Hecke algebra. Through it, one can pull back the triangulated birepresentation induced by any finitary birepresentation of finite type A Soergel bimodules to obtain a triangulated birepresentation of extended affine type A Soergel bimodules. We show that if the initial finitary birepresentation in finite type A is a cell birepresentation, the evaluation birepresentation in extended affine type A has a finitary cover, which we illustrate by working out the case of cell birepresentations with subregular apex in detail.

CONTENTS

1. Introduction	2
2. The decategorified story	3
2.1. Hecke algebras	4
2.2. Evaluation maps	4
2.3. Graham-Lehrer cell modules	6
2.4. Evaluation modules	8
3. Reminders on Soergel categories	9
3.1. Graded categories and categories with shift	9
3.2. Soergel calculus in finite and non-extended affine type A	10
3.3. Soergel calculus in extended affine type A	14
4. Rouquier complexes	16
4.1. Some diagrammatic shortcuts I: general Rouquier complexes	17
4.2. Some diagrammatic shortcuts II: special Rouquier complexes $T_{\rho}^{\pm 1}$	28
5. Evaluation functors	39
5.1. Definition	39
5.2. Proof of well-definedness	42
6. Evaluation birepresentations and finitary covers	50
6.1. Recollections on birepresentation theory	50
6.2. Finitary covers of evaluation cell birepresentations	51
6.3. The zigzag algebras	51
6.4. The birepresentations	53

1. INTRODUCTION

Finitary birepresentation theory of finite type Soergel bimodules in characteristic zero has been a topic of intensive study, with many interesting results, in the last couple of years [KMMZ2019, MM2017, MMTZ2019, MT2019, Zimm2017]. In this paper, we initiate the study of a class of finitary and triangulated birepresentations of affine type A Soergel bimodules. The bicategories of these Soergel bimodules are no longer finitary and, therefore, new phenomena show up in their birepresentation theory. For example, there are no known interesting triangulated birepresentations in finite type, whereas we do give examples of such birepresentations in affine type A .

To describe these, let us briefly recall the decategorified setting first. In type A , as is well-known, there are evaluation maps from the affine Hecke algebra to the finite type Hecke algebra. These are homomorphisms of algebras, so any representation of the latter algebra can be pulled back to a representation of the former algebra through such a map. These so-called *evaluation representations* form an important and well-studied class of finite-dimensional representations of affine type A Hecke algebras, see e.g. [CP1996, DF2016, LNT2003] and references therein.

Several authors ([MT2017, Introduction] and [E2018, Section 1.6]) have conjectured that these evaluation maps can be categorified by monoidal *evaluation functors* from affine type A Soergel bimodules to the homotopy category of bounded complexes in finite type A Soergel bimodules. In this paper, we indeed define such functors and use them to categorify the aforementioned evaluation representations in the form of triangulated birepresentations, obtained by pulling back the triangulated birepresentations induced by finitary birepresentations of finite type A Soergel bimodules through these functors. Moreover, in case the original finitary birepresentation is simple transitive, we show that the evaluation birepresentation admits a *finitary cover*, i.e., a finitary birepresentation together with an essentially surjective and epimorphic morphism of additive birepresentations from that cover to the evaluation birepresentation. This categorifies the well-known fact that the corresponding evaluation representations are quotients of certain cell representations defined by Graham and Lehrer [GL1998].

Let us finish this introduction with a disclaimer. We do not present a theory of triangulated birepresentations in this paper. Some ingredients for such a theory can already be found in the literature, e.g. [E2018, EH2018, Hog2017, LM2022, Stev2011], but many foundational results are still missing. For a start, it is not clear which parts of finitary birepresentation theory, e.g. the notion of simple transitive birepresentation, the categorical (weak) Jordan-Hölder theorem, the relation with (co)algebra 1-morphisms, the double-centralizer theorem (see [MMMTZ2020] and references therein), generalize to the triangulated setting and/or in which form exactly. These questions need to be answered first, before one can even think of categorifying the induction product of evaluation representations from [LNT2003, Section 2.5]. Finally, all of this is just for affine type A . Hecke algebras of other affine Coxeter types also have interesting finite-dimensional representations, but there are no evaluation morphisms in those cases, so other ideas

will be needed to categorify those representations. In other words, the results in this paper are (hopefully) just the tip of a (tricky) triangulated iceberg.

Plan of the paper. In Section 2, we recall the basics of extended and non-extended affine Hecke algebras of affine type A , the evaluation maps, the Graham-Lehrer cell modules and the evaluation representations. Everything in this section is well-documented in the literature and we only recall the details that are needed in the rest of this paper.

In Section 3, we briefly recall Soergel calculus in finite and affine type A , the latter both in the non-extended and the extended version. Again, nothing new is presented, so the specialists can skip this section and move on to the next one. Of course, in the remainder we often refer to the diagrammatic equations in this section, which is exactly why we recall them.

In Section 4, we first recall some basic results on Rouquier complexes in finite type A and then focus on a special type of Rouquier complex, which is fundamental for the definition of the evaluation functors in the next section. In particular, we develop a mixed diagrammatic calculus for morphisms between products of Bott-Samelson bimodules and these special Rouquier complexes, all in finite type A . To the best of our knowledge, this extension of the usual Soergel calculus is new.

In Section 5, we define the evaluation functors by assigning a bounded complex of finite type A Soergel bimodules (or, more precisely, of finite type A Bott-Samelson bimodules) to each extended affine type A Bott-Samelson bimodule and a map between such complexes to each generating extended affine type A Soergel calculus diagram. The main result of this section, and of this paper, is that this assignment is well-defined up to homotopy equivalence.

In Section 6, we first introduce the notion of a triangulated birepresentation of an additive bicategory and define evaluation birepresentations of Soergel bimodules in extended affine type A , which are important examples. We then prove that each evaluation birepresentation has a (possibly non-unique) finitary cover. Finally, we study in detail the simplest non-trivial evaluation birepresentations, which are the ones induced by cell birepresentations of finite type A with subregular apex. As we show, these admit a simple transitive finitary cover whose underlying algebra is a signed version of the zigzag algebra of affine type A .

Acknowledgments. M.M. was supported in part by Fundação para a Ciência e a Tecnologia (Portugal), projects UID/MAT/04459/2013 (Center for Mathematical Analysis, Geometry and Dynamical Systems - CAMGSD) and PTDC/MAT-PUR/31089/2017 (Higher Structures and Applications). V.M. is partially supported by EPSRC grant EP/S017216/1. P.V. was supported by the Fonds de la Recherche Scientifique - FNRS under Grant no. MIS-F.4536.19.

2. THE DECATEGORIFIED STORY

Fix $d \in \mathbb{N}_{\geq 3}$ and let $\widehat{\mathfrak{S}}_d$ be the affine Weyl group of type \widehat{A}_{d-1} . It is generated by s_0, \dots, s_{d-1} , subject to relations

$$s_i^2 = 1, \quad s_i s_j = s_j s_i \text{ if } |i - j| > 1, \quad s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1},$$

for $i = 0, \dots, d-1$, with indices taken modulo d . The *extended affine Weyl group* $\widehat{\mathfrak{S}}_d^{\text{ext}}$ is the semidirect product

$$\langle \rho \rangle \ltimes \widehat{\mathfrak{S}}_d,$$

where $\langle \rho \rangle$ is an infinite cyclic group generated by ρ and

$$\rho s_i \rho^{-1} = s_{i+1}.$$

The finite Weyl group of type A_{d-1} is the symmetric group on d letters, \mathfrak{S}_d , corresponding to the subgroup of $\widehat{\mathfrak{S}}_d$ generated by s_i for $1 \leq i \leq d-1$.

2.1. Hecke algebras. Let $\mathbb{k} = \mathbb{C}(q)$, where q is a formal parameter. The *extended affine Hecke algebra* $\widehat{H}_d^{\text{ext}}$ is the \mathbb{k} -algebra generated by T_0, \dots, T_{d-1} and $\rho^{\pm 1}$, with relations

$$(1) \quad (T_i + q)(T_i - q^{-1}) = 0, \quad T_i T_j = T_j T_i \text{ if } |i - j| > 1, \quad T_i T_{i+1} T_i = T_{i+1} T_i T_{i+1},$$

$$(2) \quad \rho \rho^{-1} = 1 = \rho^{-1} \rho, \quad \rho T_i \rho^{-1} = T_{i+1},$$

for $i, j = 0, \dots, d-1$ (with indices taken modulo d again). Note that T_i is invertible for every $i = 0, \dots, d-1$ with

$$T_i^{-1} = T_i + q - q^{-1}.$$

As is well-known, $\widehat{H}_d^{\text{ext}}$ is a q -deformation of the group algebra $\mathbb{C}[\widehat{\mathfrak{S}}_d^{\text{ext}}]$ with basis (the *regular basis*) given by $\{\rho^m T_w \mid m \in \mathbb{Z}, w \in \widehat{\mathfrak{S}}_d\}$, where $T_w := T_{i_1} \cdots T_{i_\ell}$ for any *reduced expression* (rex) $s_{i_1} \cdots s_{i_\ell}$ of w .

Another presentation is given in terms of the *Kazhdan–Lusztig generators* $b_i := T_i + q$, for $i = 0, \dots, d-1$, and $\rho^{\pm 1}$, subject to relations

$$(3) \quad b_i^2 = [2]b_i, \quad b_i b_j = b_j b_i \text{ if } |i - j| > 1, \quad b_i b_{i+1} b_i + b_{i+1} = b_{i+1} b_i b_{i+1} + b_i,$$

$$(4) \quad \rho \rho^{-1} = 1 = \rho^{-1} \rho, \quad \rho b_i \rho^{-1} = b_{i+1},$$

for $i = 0, \dots, d-1$, where $[2] := q + q^{-1}$. Note that $T_i = b_i - q$ and $T_i^{-1} = b_i - q^{-1}$, for every $i = 0, \dots, d-1$. The *Kazhdan–Lusztig basis* is given by $\{\rho^m b_w \mid m \in \mathbb{Z}, w \in \widehat{\mathfrak{S}}_d\}$, where b_w is defined for an arbitrary rex of w (and is independent of that choice).

The (non-extended) *affine Hecke algebra* \widehat{H}_d is the subalgebra of $\widehat{H}_d^{\text{ext}}$ generated by either T_0, T_1, \dots, T_{d-1} subject to relations (1), or b_0, b_1, \dots, b_{d-1} subject to relations (3).

The *finite Hecke algebra* H_d is the \mathbb{k} -subalgebra of \widehat{H}_d generated by either T_1, \dots, T_{d-1} subject to relations (1), or b_1, \dots, b_{d-1} subject to relations (3).

2.2. Evaluation maps.

Definition 2.1. For any $a \in \mathbb{k}^\times$, there are two *evaluation maps* $\text{ev}_a, \text{ev}'_a : \widehat{H}_d^{\text{ext}} \rightarrow H_d$. These are defined as the homomorphisms of \mathbb{k} -algebras determined by

$$\text{ev}_a(T_i) = T_i, \quad \text{for } 1 \leq i \leq d-1,$$

$$\text{ev}_a(\rho) = a T_1^{-1} \cdots T_{d-1}^{-1}$$

and

$$\begin{aligned} \text{ev}'_a(T_i) &= T_i, \quad \text{for } 1 \leq i \leq d-1, \\ \text{ev}'_a(\rho) &= aT_1 \cdots T_{d-1}, \end{aligned}$$

respectively.

The definition implies that

$$\text{ev}_a(T_0) = \text{ev}_a(\rho^{-1}T_1\rho) = T_{d-1} \cdots T_2T_1T_2^{-1} \cdots T_{d-1}^{-1}$$

and

$$\text{ev}'_a(T_0) = \text{ev}'_a(\rho^{-1}T_1\rho) = T_{d-1}^{-1} \cdots T_2^{-1}T_1T_2 \cdots T_{d-1},$$

so the restrictions of ev_a and ev'_a to \widehat{H}_d do not depend on a .

In terms of the Kazhdan–Lusztig generators we have

$$\begin{aligned} \text{ev}_a(b_i) &= b_i, \quad \text{for } 1 \leq i \leq d-1, \\ \text{ev}_a(b_0) &= \text{ev}_a(\rho^{-1}b_1\rho) = (b_{d-1} - q) \cdots (b_1 - q)b_1(b_1 - q^{-1}) \cdots (b_{d-1} - q^{-1}) \end{aligned}$$

and

$$\begin{aligned} \text{ev}'_a(b_i) &= b_i, \quad \text{for } 1 \leq i \leq d-1, \\ \text{ev}'_a(b_0) &= \text{ev}'_a(\rho^{-1}b_1\rho) = (b_{d-1} - q^{-1}) \cdots (b_1 - q^{-1})b_1(b_1 - q) \cdots (b_{d-1} - q). \end{aligned}$$

Another way of saying this is that the evaluation maps do not preserve the bar involution, but rather satisfy

$$(5) \quad \overline{\text{ev}_a(x)} = \text{ev}'_a(\bar{x}),$$

for any $x \in \widehat{H}_d^{ext}$ and $a = a(q) \in \mathbb{k}^\times$.

One can also define ev_a and ev'_a using a third presentation of \widehat{H}_d^{ext} , called the *Bernstein presentation*. In that presentation, \widehat{H}_d^{ext} is defined as some sort of semidirect product of H_d and $\mathbb{k}[Y_1^{\pm 1}, \dots, Y_d^{\pm 1}]$. However, there are several possible choices for the algebra of Laurent polynomials. In [E2018], two such choices are given with different variables: y_1, \dots, y_d and y_1^*, \dots, y_d^* respectively. The interaction of H_d and these polynomial algebras is defined by

$$T_i^{-1}y_iT_i^{-1} = y_{i+1}$$

and

$$T_i y_i^* T_i = y_{i+1}^*,$$

respectively, for $i = 1, \dots, d-2$.

The relation between these two Bernstein presentations and our first presentation of \widehat{H}_d^{ext} is given by

$$\begin{aligned} y_1 &= \rho T_{d-1} \cdots T_2 T_1, \\ y_i &= T_{i-1}^{-1} \cdots T_2^{-1} T_1^{-1} \rho T_{d-1} \cdots T_{i+1} T_i, \quad i = 2, \dots, d-1, \end{aligned}$$

resp.


$$\begin{aligned} y_1^* &= \rho T_{d-1}^{-1} \cdots T_2^{-1} T_1^{-1}, \\ y_i^* &= T_{i-1} \cdots T_2 T_1 \rho T_{d-1}^{-1} \cdots T_{i+1}^{-1} T_i^{-1}, \quad i = 2, \dots, d-1. \end{aligned}$$

It follows that the evaluation map $\text{ev}_a: \widehat{H}_d^{\text{ext}} \rightarrow H_d$ is the unique homomorphism of algebras sending T_i to T_i , for $i = 1, \dots, d-1$, and y_1 to a , while $\text{ev}'_a: \widehat{H}_d^{\text{ext}} \rightarrow H_d$ is the unique homomorphism of algebras sending T_i to T_i , for $i = 1, \dots, d-1$, and y_1^* to a . The latter coincides with the flattening map \flat in [E2018, §2.6] for $a = 1$.

We will categorify the evaluation map ev_a in Section 5.1. The categorification of ev'_a is very similar and the relation between the two evaluation maps in (5) also categorifies, since the categorification of the bar-involution is given by flipping diagrams upside-down, inverting the orientation of the differentials in complexes and changing the sign of homological and grading shifts.

Remark 2.2. Some remarks about the various conventions in the literature are in order. We try to follow conventions close to those in [E2018]. Our presentation of the extended affine Hecke algebra in Section 2.1 agrees with [E2018], as does the relation between the standard generators and the Kazhdan–Lusztig generators. Some authors use the inverse of ρ in (2). Our choice of conventions implies the absence of certain powers of q in the definition of the evaluation maps, in comparison with some of the sources in the literature. For more information on evaluation maps, see e.g. [CP1996, §5.1] and [DF2016, (5.0.2)]. There are more possible evaluation maps, but we only consider these two in this paper.

2.3. Graham-Lehrer cell modules. Consider the \widehat{A}_{d-1} Coxeter diagram $\widehat{\Gamma}_{d-1}$ with its vertices ordered as indicated (for e.g. $d = 8$):



For any $z \in \mathbb{k}^\times$, the *Graham-Lehrer cell module* \widehat{M}_z of \widehat{H}_d corresponding to z and the partition $(d-1, 1)$ has underlying vector space

$$\widehat{M}_z := \text{Span}_{\mathbb{k}} \{m_i \mid i = 0, \dots, d-1\},$$

where the indices of the m_i have to be taken modulo d by convention, and the action of \widehat{H}_d on \widehat{M}_z is given by

$$(6) \quad b_i m_j = \begin{cases} [2]m_i, & \text{if } j \equiv i \pmod{d}; \\ zm_1, & \text{if } i-1 \equiv 0 \equiv j \pmod{d}; \\ z^{-1}m_0, & \text{if } i \equiv 0 \equiv j-1 \pmod{d}; \\ m_j, & \text{if } i \equiv j \pm 1 \pmod{d}, \text{ but none of the above}; \\ 0, & \text{else.} \end{cases}$$

It is easy to see that \widehat{M}_z is isomorphic to $W_{d-2, \pm\sqrt{z}}(d)$ in [GL1998, Definition 2.6], where m_i is identified with the cup diagram on a cylinder with $d-2$ endpoints at the bottom, d endpoints

at the top and with only one cup, whose endpoints are i and $i + 1$, and further only straight lines. When $i \neq 0$, the whole diagram corresponding to m_i lives on the front part of the cylinder, but when $i = 0$, the cup of m_0 goes around the back of the cylinder. Note that we have used $\delta = [2]$, rather than $\delta = -[2]$. As remarked in [GL1998, text above Corollary 2.9.1], $W_{d-2, \sqrt{z}}(d)$ and $W_{d-2, -\sqrt{z}}(d)$ are isomorphic, which is clear from the fact that both are isomorphic to \widehat{M}_z .

The Graham-Lehrer cell module \widehat{M}_z can be made into an \widehat{H}_d^{ext} -module, but not in a unique way. As a matter of fact, for each $\lambda \in \mathbb{k}^\times$, we can define

$$(7) \quad \rho m_j = \lambda z^{\delta_{j,0}} m_{j+1},$$

for $j = 0, \dots, d-1$. It is easy to verify that this gives a well-defined action and we denote the corresponding Graham-Lehrer cell module of \widehat{H}_d^{ext} by $\widehat{M}_{z,\lambda}$. Note that the restriction of $\widehat{M}_{z,\lambda}$ to \widehat{H}_d is equal to \widehat{M}_z , for all $\lambda \in \mathbb{k}^\times$, and that the action of ρ^d on $\widehat{M}_{z,\lambda}$ is simply multiplication by $\lambda^d z$.

Graham and Lehrer [GL1998, Theorem 2.8] defined a \mathbb{k} -bilinear form

$$\langle \cdot, \cdot \rangle: \widehat{M}_z \otimes \widehat{M}_{z^{-1}} \rightarrow \mathbb{k},$$

which in our notation is determined by

$$\langle m_i, m_j \rangle = \begin{cases} [2], & \text{if } j \equiv i \pmod{d}; \\ z, & \text{if } i \equiv 0 \equiv j-1 \pmod{d}; \\ z^{-1}, & \text{if } i-1 \equiv 0 \equiv j \pmod{d}; \\ 1, & \text{if } i \equiv j \pm 1 \pmod{d}, \text{ but none of the above}; \\ 0, & \text{else.} \end{cases}$$

This induces a \mathbb{k} -bilinear form on $\widehat{M}_{z,\lambda} \otimes \widehat{M}_{z^{-1},\lambda^{-1}}$, satisfying $\langle \rho^n b_w m_j, m_k \rangle = \langle m_j, b_w^* \rho^{-n} m_k \rangle$, for any $w \in \widehat{W}$, $n \in \mathbb{Z}$ and $j, k = 0, 1, \dots, d-1$, where $b_w^* = b_{w^{-1}}$ is the dual Kazhdan-Lusztig basis element. Therefore, the radical of the bilinear form

$$\text{rad}(\langle \cdot, \cdot \rangle) = \left\{ m \in \widehat{M}_{z,\lambda} \mid \langle m, m' \rangle = 0, \forall m' \in \widehat{M}_{z^{-1},\lambda^{-1}} \right\}$$

is an \widehat{H}_d^{ext} -submodule of $\widehat{M}_{z,\lambda}$. Graham and Lehrer [GL1998, Theorem 2.8] proved that the quotient module $\widehat{M}_z / \text{rad}(\langle \cdot, \cdot \rangle)$ of \widehat{H}_d is simple, and the same holds for the quotient module $\widehat{M}_{z,\lambda} / \text{rad}(\langle \cdot, \cdot \rangle)$ of \widehat{H}_d^{ext} , of course. A straightforward calculation shows that the radical of the bilinear form on $\widehat{M}_{z,\lambda}$ is zero unless $z = (-q)^{\pm d}$ (independently of λ), in which case it has dimension one and is generated by

$$n_\pm := \sum_{k=1}^d (-q)^{\mp k} m_k,$$

where $m_d := m_0$ by convention. Note that, when $z = (-q)^{\pm d}$, we have $\rho n_\pm = \lambda(-q)^{\pm 1} n_\pm$ and $b_i n_\pm = 0$ for all $i = 0, \dots, d-1$.

When $z = (-q)^{\pm d}$, put $\widehat{M}_{d,\lambda}^{\pm} := \widehat{M}_{(-q)^{\pm d}, \lambda^{\pm 1}}$ and let

$$(8) \quad \widehat{L}_{d,\lambda}^{\pm} := \widehat{M}_{d,\lambda}^{\pm} / \langle n_{\pm} \rangle$$

be the simple quotient \widehat{H}_d^{ext} -modules of dimension $d - 1$. Finally, denote the restriction of these simple modules to \widehat{H}_d by

$$(9) \quad \widehat{L}_d^{\pm} := \widehat{M}_d^{\pm} / \langle n_{\pm} \rangle.$$

As explained above, these restrictions do not depend on $\lambda \in \mathbb{k}^{\times}$.

2.4. Evaluation modules. Let M be a finite-dimensional H_d -module (over \mathbb{k}). Recall that, for any $a \in \mathbb{k}^{\times}$, there are two evaluation maps $\text{ev}_a, \text{ev}'_a: \widehat{H}_d^{ext} \rightarrow H_d$ (see Definition 2.1).

Definition 2.3. For any $a \in \mathbb{k}^{\times}$, the *evaluation modules* M^{ev_a} and $M^{\text{ev}'_a}$ of \widehat{H}_d^{ext} are the pull-backs of M through ev_a and ev'_a , respectively.

The actions of \widehat{H}_d^{ext} on M^{ev_a} and $M^{\text{ev}'_a}$ can be computed using the explicit formulas in Definition 2.1 and below. In this paper, we only consider the case when $M := M_d$ is the simple H_d -module corresponding to the partition $(d - 1, 1)$. There are several ways to define M_d explicitly and the definition we choose here is tailor-made for categorification. Take $M_d := \text{span}_{\mathbb{k}}\{m_1, \dots, m_{d-1}\}$, with the action of H_d being given by

$$(10) \quad b_i m_j = \begin{cases} [2]m_i, & \text{if } j = i; \\ m_i, & \text{if } j = i \pm 1; \\ 0, & \text{else,} \end{cases}$$

for $i, j = 1, \dots, d - 1$. It is easy to show that M_d is simple, but this is well-known so we leave it as an exercise to the reader. The action of the $T_i^{\pm 1} = b_i - q^{\pm 1}$ is also easy to give explicitly:

$$T_i^{\pm 1} m_j = \begin{cases} q^{\mp 1} m_i, & \text{if } j = i; \\ m_i - q^{\pm 1} m_j, & \text{if } j = i \pm 1; \\ -q^{\pm 1} m_j, & \text{else.} \end{cases}$$

Note that, as vector spaces, $M_d^{\text{ev}_a} = M_d^{\text{ev}'_a} = M_d$, and the action of $b_i \in \widehat{H}_d^{ext}$, for $i = 1, \dots, d - 1$, is the same as above because $\text{ev}_a(b_i) = b_i$. A simple calculation now shows that

$$(11) \quad \text{ev}_a(\rho) m_j = a T_1^{-1} \cdots T_{d-1}^{-1} m_j = \begin{cases} a(-q)^{2-d} m_{j+1}, & \text{if } j = 1, \dots, d - 2; \\ a q \sum_{k=1}^{d-1} (-q)^{1-k} m_k, & \text{if } j = d - 1, \end{cases}$$

and

$$(12) \quad \text{ev}'_a(\rho) m_j = a T_1 \cdots T_{d-1} m_j = \begin{cases} a(-q)^{d-2} m_{j+1}, & \text{if } j = 1, \dots, d - 2; \\ a q^{-1} \sum_{k=1}^{d-1} (-q)^{k-1} m_k, & \text{if } j = d - 1. \end{cases}$$

The actions of b_0 can then be computed using the equation $b_0 = \rho^{-1} b_1 \rho$, but we omit the calculation because will not need the result.

Recall the simple quotients $\widehat{L}_{d,\lambda}^\pm$ of the Graham-Lehrer cell modules $\widehat{M}_{d,\lambda}^\pm$, defined in (8). We claim that $\widehat{L}_{d,\lambda}^+ \cong M_d^{\text{ev}^a}$ for $a = \lambda(-q)^{d-2}$. To show this, it suffices to compute the action of ρ on $\widehat{L}_{d,\lambda}^+$ and compare it to (11). Let \overline{m}_k be the image of m_k under the projection $\widehat{M}_{d,\lambda}^+ \rightarrow \widehat{L}_{d,\lambda}^+$, for $k = 0, \dots, d-1$. Then $\{\overline{m}_1, \dots, \overline{m}_{d-1}\}$ is a basis of $\widehat{L}_{d,\lambda}^+$, because $\overline{m}_0 = -\sum_{k=1}^{d-1} (-q)^k \overline{m}_{d-k}$. This implies that in $\widehat{L}_{d,\lambda}^+$ we have

$$\rho \overline{m}_j = \begin{cases} \lambda \overline{m}_{j+1}, & \text{if } j = 1, \dots, d-2; \\ -\lambda \sum_{k=1}^{d-1} (-q)^k \overline{m}_{d-k}, & \text{if } j = d-1. \end{cases}$$

This is indeed the same as in (11) because $aq = \lambda(-q)^{d-2}q = -\lambda(-q)^{d-1}$.

Similarly, $\widehat{L}_{d,\lambda}^- \cong M_d^{\text{ev}'^a}$ for $a = \lambda^{-1}(-q)^{2-d}$, as in $\widehat{L}_{d,\lambda}^-$ we have $\overline{m}_0 = -\sum_{k=1}^{d-1} (-q)^{-k} \overline{m}_{d-k}$, so

$$\rho \overline{m}_j = \begin{cases} \lambda^{-1} \overline{m}_{j+1}, & \text{if } j = 1, \dots, d-2; \\ -\lambda^{-1} \sum_{k=1}^{d-1} (-q)^{-k} \overline{m}_{d-k}, & \text{if } j = d-1, \end{cases}$$

which is the same as in (12) because $aq^{-1} = \lambda^{-1}(-q)^{2-d}q^{-1} = -\lambda^{-1}(-q)^{1-d}$.

The two $\widehat{H}_d^{\text{ext}}$ -modules $\widehat{L}_{d,\lambda}^+$ and $\widehat{L}_{d,\lambda}^-$ are actually *dual* to each other. Note that we could also consider the radical defined by

$$\text{rad}'(\langle \cdot, \cdot \rangle) = \left\{ m' \in \widehat{M}_{z^{-1}, \lambda^{-1}} \mid \langle m, m' \rangle = 0, \forall m \in \widehat{M}_{z, \lambda} \right\},$$

which is an $\widehat{H}_d^{\text{ext}}$ -submodule of $\widehat{M}_{z^{-1}, \lambda^{-1}}$. As before, this radical is zero unless $z = (-q)^{\pm d}$. For these two values of z and any value of $\lambda \in \mathbb{k}^\times$, the two simple quotients of $\widehat{M}_{z^{-1}, \lambda^{-1}}$ are isomorphic to $\widehat{L}_{d,\lambda}^\mp$ and the bilinear form descends to a perfect pairing

$$\widehat{L}_{d,\lambda}^+ \otimes \widehat{L}_{d,\lambda}^- \rightarrow \mathbb{k}.$$

By the above, this is equivalent to a perfect pairing

$$M_d^{\text{ev}^a} \otimes M_d^{\text{ev}'^a} \rightarrow \mathbb{k},$$

for $a = \lambda(-q)^{d-2}$.

3. REMINDERS ON SOERTEL CATEGORIES

In this section we briefly recall the definition of the diagrammatic Soergel category of non-extended and extended affine type A and finite type A , but before we do that we start with a brief section on graded categories and categories with shift.

3.1. Graded categories and categories with shift. All categories in this paper are assumed to be essentially small, meaning that they are equivalent to small categories, so set-theoretic questions play no role.

We call a \mathbb{C} -linear category \mathcal{A} *graded* if it is enriched over the category of \mathbb{Z} -graded vector spaces, and we call a \mathbb{C} -linear functor between such graded categories *degree-preserving* if it preserves the degrees of homogeneous morphisms.

We say that a \mathbb{C} -linear category \mathcal{A} has a *shift* (or, alternatively, that it is a *category with shift*) if there is a \mathbb{C} -linear automorphism $\langle 1 \rangle$ of \mathcal{A} . If such a shift exists, we define $\langle r \rangle$ as the composite of r copies of $\langle 1 \rangle$ for any $r \in \mathbb{Z}_{\geq 0}$, and $-r$ copies of the inverse of $\langle 1 \rangle$ for any $r \in \mathbb{Z}_{\leq 0}$. By definition, therefore, we have $\langle r + s \rangle = \langle r \rangle \langle s \rangle = \langle s \rangle \langle r \rangle$, for all $r, s \in \mathbb{Z}$, and $\langle 0 \rangle = \text{Id}_{\mathcal{A}}$.

Given a graded category \mathcal{A} , let \mathcal{A}^{sh} be the associated \mathbb{C} -linear category with shift, whose objects are formal integer shifts of objects in \mathcal{A} and whose hom-spaces are defined by

$$\mathcal{A}^{\text{sh}}(X\langle r \rangle, Y\langle s \rangle) := \mathcal{A}(X, Y)_{s-r}$$

for every $X, Y \in \mathcal{A}$ and $r, s \in \mathbb{Z}$. Note that \mathcal{A}^{sh} is no longer a graded category. If the Hom-spaces of \mathcal{A} are finite-dimensional in every degree, then the hom-spaces of \mathcal{A}^{sh} are finite-dimensional.

Given two graded categories \mathcal{A} and \mathcal{B} , any degree-preserving, \mathbb{C} -linear functor $F: \mathcal{A} \rightarrow \mathcal{B}$ induces a unique \mathbb{C} -linear functor $F: \mathcal{A}^{\text{sh}} \rightarrow \mathcal{B}^{\text{sh}}$, denoted by the same symbol, which commutes with the shifts.

Conversely, given any \mathbb{C} -linear category \mathcal{A} with shift, let \mathcal{A}^{gr} be the associated graded category, whose objects are those of \mathcal{A} and whose graded Hom-spaces are defined by

$$\mathcal{A}^{\text{gr}}(X, Y) := \bigoplus_{s \in \mathbb{Z}} \mathcal{A}(X, Y\langle s \rangle),$$

for any $X, Y \in \mathcal{A}$. Note that \mathcal{A}^{gr} is graded and has a shift, and, moreover, that $X \cong X\langle t \rangle$ for all $X \in \mathcal{A}$ and $t \in \mathbb{Z}$.

Given two \mathbb{C} -linear categories \mathcal{A} and \mathcal{B} with shifts, any \mathbb{C} -linear functor $F: \mathcal{A} \rightarrow \mathcal{B}$ commuting with the shifts induces a unique degree-preserving, \mathbb{C} -linear functor $F: \mathcal{A}^{\text{gr}} \rightarrow \mathcal{B}^{\text{gr}}$, denoted by the same symbol.

Thus $(-)^{\text{sh}}$ and $(-)^{\text{gr}}$ define a pair of 2-functors between the 2-category of graded categories and the 2-category of \mathbb{C} -linear categories with shift. It is not hard to show, see e.g. [EMTW2020, Proposition 11.9], that $(-)^{\text{sh}}$ is left adjoint to $(-)^{\text{gr}}$, i.e., that there is a functorial isomorphism

$$\text{Fun}(\mathcal{A}^{\text{sh}}, \mathcal{B}) \cong \text{Fun}(\mathcal{A}, \mathcal{B}^{\text{gr}})$$

for \mathcal{A} a graded category and \mathcal{B} a \mathbb{C} -linear category with shift. Here the first functor category is between categories with shift and the second between graded categories.


For more details on graded categories and categories with shift, and also on additive closures and idempotent completions (a.k.a. Karoubi closures/envelopes), see e.g. [EMTW2020, Sections 11.2.1-11.24].

3.2. Soergel calculus in finite and non-extended affine type A. The finite type A diagrammatic Soergel calculus was introduced by Elias–Khovanov [EKh2010] and generalized to all Coxeter types by Elias–Williamson [EW2016]. The extended affine Soergel calculus was first defined in [MT2017] and studied more systematically in [E2018]. We refer to the latter two papers for more details. For the specialists, we remark that we use the so-called *root span realization* of the Cartan datum of finite and affine type A below.

Denote by $S = \{s_0, \dots, s_{d-1}\}$ the set of simple reflections of $\widehat{\mathfrak{S}}_d$. The *diagrammatic Bott-Samelson category* of type \widehat{A}_{d-1} , denoted $\widehat{\mathcal{B}}\mathfrak{S}_d$, is the \mathbb{Z} -graded, \mathbb{C} -linear, additive, monoidal category whose objects are formal finite direct sums of finite words in the alphabet S , and whose

graded vector spaces of morphisms are defined below in terms of homogeneous generating diagrams and relations. In general, we can write the objects as vectors of words and morphisms as matrices of equivalence classes of diagrams.

As usual, we will color the strands to facilitate the reading of the diagrams. These colors correspond to the elements of $\mathbb{Z}/d\mathbb{Z}$, so henceforth we will also refer to those elements as colors. When there are too many different colors in a diagram, the colors are sometimes indicated by labels next to the strands. We say that two colors $i, j \in \mathbb{Z}/d\mathbb{Z}$ are *adjacent* if $i = j \pm 1 \pmod d$ and that they are *distant* otherwise. The generating diagrams are



and the diagrams obtained from these by a rotation of 180 degrees (which have the same degrees). The colors of the 4-valent vertices are assumed to be distant, whereas those of the 6-valent vertices are assumed to be adjacent.

Diagrams can be stacked vertically (composition of morphisms) and juxtaposed horizontally (monoidal product of morphisms), while adding the degrees, and are subject to the relations below. We denote by Id_X the identity morphism of X and write fg for the monoidal product of morphisms f and g (or, equivalently, horizontal composition when considering the monoidal category as a one-object bicategory). We also assume isotopy invariance and cyclicity, meaning that closed parts of the diagrams can be moved around freely in the plane as long as they do not cross any other strands and the boundary is fixed, and all diagrams can be bent and rotated and the bent and rotated versions of the relations also hold.

- Relations involving one color:

(13) 

(14) 

(15) 

(16) 

- Relations involving two distant colors:

(17) 

$$(18) \quad \begin{array}{c} \text{blue} \\ \diagdown \\ \text{green} \end{array} \begin{array}{c} \text{green} \\ \diagup \\ \text{blue} \end{array} = \begin{array}{c} \text{blue} \\ \bullet \\ \text{green} \end{array}$$

$$(19) \quad \begin{array}{c} \text{green} \\ \diagup \\ \text{blue} \end{array} = \begin{array}{c} \text{blue} \\ \diagdown \\ \text{green} \end{array}$$

- Relations involving two adjacent colors:

$$(20) \quad \begin{array}{c} \text{red} \\ \diagdown \\ \text{blue} \end{array} \begin{array}{c} \text{blue} \\ \diagup \\ \text{red} \end{array} = \begin{array}{c} \text{red} \\ \bullet \\ \text{blue} \end{array} + \begin{array}{c} \text{blue} \\ \bullet \\ \text{red} \end{array}$$

$$(21) \quad \begin{array}{c} \text{red} \\ | \\ \text{red} \end{array} \begin{array}{c} \text{blue} \\ | \\ \text{blue} \end{array} = \begin{array}{c} \text{red} \\ \diagdown \\ \text{blue} \end{array} \begin{array}{c} \text{blue} \\ \diagup \\ \text{red} \end{array} - \begin{array}{c} \text{blue} \\ \bullet \\ \text{red} \end{array}$$

$$(22) \quad \begin{array}{c} \text{blue} \\ \diagdown \\ \text{red} \end{array} \begin{array}{c} \text{red} \\ \diagup \\ \text{blue} \end{array} = \begin{array}{c} \text{red} \\ \bullet \\ \text{blue} \end{array}$$

$$(23) \quad \begin{array}{c} \text{red} \\ \bullet \\ | \\ \text{red} \end{array} - \begin{array}{c} \text{blue} \\ \bullet \\ | \\ \text{blue} \end{array} = \frac{1}{2} \left(\begin{array}{c} \text{blue} \\ \bullet \\ | \\ \text{blue} \end{array} - \begin{array}{c} \text{red} \\ \bullet \\ | \\ \text{red} \end{array} \right)$$

- Relation involving three distant colors:

$$(24) \quad \begin{array}{c} \text{green} \\ \diagup \\ \text{blue} \end{array} \begin{array}{c} \text{blue} \\ \diagdown \\ \text{green} \end{array} = \begin{array}{c} \text{blue} \\ \diagdown \\ \text{green} \end{array} \begin{array}{c} \text{green} \\ \diagup \\ \text{blue} \end{array}$$

- Relation involving distant dumbbells:

$$(25) \quad \begin{array}{c} \text{green} \\ \bullet \\ | \\ \text{green} \end{array} - \begin{array}{c} \text{blue} \\ \bullet \\ | \\ \text{blue} \end{array} = 0$$

- Relation involving two adjacent colors and one distant from the other two:

$$(26) \quad \begin{array}{c} \text{green} \\ \diagup \\ \text{blue} \end{array} \begin{array}{c} \text{blue} \\ \diagdown \\ \text{green} \end{array} = \begin{array}{c} \text{blue} \\ \diagdown \\ \text{green} \end{array} \begin{array}{c} \text{green} \\ \diagup \\ \text{blue} \end{array}$$

- Relation involving three adjacent colors:

$$(27) \quad \begin{array}{c} \text{green} \\ \diagup \\ \text{blue} \end{array} \begin{array}{c} \text{blue} \\ \diagdown \\ \text{green} \end{array} \begin{array}{c} \text{red} \\ \diagup \\ \text{blue} \end{array} \begin{array}{c} \text{blue} \\ \diagdown \\ \text{green} \end{array} = \begin{array}{c} \text{blue} \\ \diagdown \\ \text{green} \end{array} \begin{array}{c} \text{green} \\ \diagup \\ \text{blue} \end{array} \begin{array}{c} \text{red} \\ \diagup \\ \text{blue} \end{array} \begin{array}{c} \text{blue} \\ \diagdown \\ \text{green} \end{array}$$

Note that the empty word is the identity object in $\widehat{\mathcal{BS}}_d$ and its endomorphisms are the closed diagrams, which by the relations above are equal to polynomials in the colored dumbbells



As each dumbbell has degree 2, the degree of any polynomial in these dumbbells, as a morphism in $\widehat{\mathcal{BS}}$, is twice its polynomial degree. From now on, we denote this polynomial algebra by R .

Note further that, by relations (16), (23) and (25), the morphism

$$(28) \quad \sum_{i=0}^{d-1} \mathbf{i}$$

is central, in the sense that it can be slid through all diagrams (i.e. it commutes horizontally with all morphisms). Note that this morphism is equal to $\boxed{\mathbf{y}}$ (up to sign, depending on conventions) in [MT2017], because it is equal to the sum of all simple roots.

Let $\widehat{\mathcal{BS}}_d^{\text{sh}}$ be the category with shift associated to $\widehat{\mathcal{BS}}_d$, see Section 3.1.

Definition 3.1. The *diagrammatic Soergel category* $\widehat{\mathcal{S}}_d$ is the idempotent completion of the diagrammatic Bott-Samelson category with shift $\widehat{\mathcal{BS}}_d^{\text{sh}}$.

Remark 3.2. In the following sections, we sometimes state and prove diagrammatic equations in $\widehat{\mathcal{BS}}_d$, in which case there are no shifts for the source and target objects, instead of $\widehat{\mathcal{S}}_d$, in which case the source and target objects are carefully shifted. This is just to simplify notation and makes no essential difference in our case. As long as the equations in $\widehat{\mathcal{BS}}_d$ are between homogeneous diagrams of the same degree, they give rise to an equality between morphisms in $\widehat{\mathcal{S}}_d$, which is the key point.

The diagrammatic Bott-Samelson category $\widehat{\mathcal{BS}}_d$ is equivalent to the algebraic category of Bott-Samelson bimodules and bimodule maps and the diagrammatic Soergel category $\widehat{\mathcal{S}}_d$ is equivalent to the algebraic category of Soergel bimodules and degree-preserving bimodule maps, see [EW2016, Theorem 6.28]. For convenience, we will therefore denote the objects of $\widehat{\mathcal{BS}}_d$ by $B_{\underline{w}} = B_{s_{i_1}} \cdots B_{s_{i_\ell}}$, where $\underline{w} = s_{i_1} \cdots s_{i_\ell}$ is a finite word in the alphabet S . In particular, the monoidal product is given by $B_{\underline{u}} B_{\underline{v}} = B_{\underline{uv}}$, where \underline{uv} is the concatenation of the words \underline{u} and \underline{v} .

Let us also recall the so-called *Categorification Theorem*, due to Soergel in finite type A , to Härterich [Har1999] in affine type A and to Elias–Williamson [EW2014, EW2016] in general Coxeter type.

Theorem 3.3. For any $w \in \widehat{\mathcal{S}}_d$ and $\text{rex } \underline{w} = s_{i_1} \cdots s_{i_\ell}$ of w , there is an indecomposable object $B_w \in \widehat{\mathcal{S}}_d$, independent of the choice of rex , such that

$$B_{\underline{w}} \cong B_w \oplus \bigoplus_{u \prec w} B_u^{\oplus h_{w,u}},$$

where \prec is the Bruhat order in $\widehat{\mathcal{S}}_d$ and $h_{w,u} \in \mathbb{N}[q, q^{-1}]$ is the graded multiplicity of B_u in the decomposition of $B_{\underline{w}}$.

Moreover, the $\mathbb{Z}[q, q^{-1}]$ -linear map

$$\begin{aligned} \widehat{H}_d^{\mathbb{Z}[q, q^{-1}]} &\rightarrow [\widehat{\mathcal{S}}_d]_{\oplus} \\ b_w &\mapsto B_w, \quad w \in \widehat{\mathcal{S}}_d \end{aligned}$$

is an isomorphism of algebras, where $\widehat{H}_d^{\mathbb{Z}[q, q^{-1}]}$ is the integral form of \widehat{H}_d .

Let $\widehat{\mathcal{S}}_d^{\text{gr}}$ be the graded monoidal category associated to $\widehat{\mathcal{S}}_d$, see Section 3.1. For every $u, v \in \widehat{\mathcal{S}}$, the graded Hom-space

$$\widehat{\mathcal{S}}_d^{\text{gr}}(B_u, B_v) = \bigoplus_{t \in \mathbb{Z}} \widehat{\mathcal{S}}_d(B_u, B_v \langle t \rangle)$$

is a free left (or right) graded R -module of finite graded rank, given by *Soergel's Hom-formula*:

$$(29) \quad \text{grk}_R \left(\widehat{\mathcal{S}}_d^{\text{gr}}(B_u, B_v) \right) = (b_u, b_v),$$

where $(-, -)$ is the well-known sesquilinear form on \widehat{H}_d , see e.g. [EW2016, Section 2.4 and Theorem 3.15].

Definition 3.4. The diagrammatic Bott-Samelson category and the diagrammatic Soergel category of finite type A_{d-1} , denoted \mathcal{BS}_d and \mathcal{S}_d respectively, are defined as $\widehat{\mathcal{BS}}_d$ and $\widehat{\mathcal{S}}_d$ but only using the colors $1, \dots, d-1$.

Note that \mathcal{BS}_d and \mathcal{S}_d are monoidal subcategories of $\widehat{\mathcal{BS}}_d$ and $\widehat{\mathcal{S}}_d$, respectively, but that the natural embeddings are not full because e.g. the 0-colored dumbbell is not a morphism in \mathcal{BS}_d and \mathcal{S}_d .

3.3. Soergel calculus in extended affine type A . In this subsection we briefly sketch how to enhance $\widehat{\mathcal{BS}}_d$ and $\widehat{\mathcal{S}}_d$ to get the extended diagrammatic Soergel category of type \widehat{A}_{d-1} , denoted $\widehat{\mathcal{BS}}_d^{\text{ext}}$ and $\widehat{\mathcal{S}}_d^{\text{ext}}$, which were introduced in [MT2017] and further studied in [E2018]. We refer to those two papers for more details.

The objects of $\widehat{\mathcal{BS}}_d^{\text{ext}}$ are formal direct sums of words in the alphabet $S \cup \{\rho, \rho^{-1}\}$. Because of the link with algebraic bimodules, we write B_ρ^n for ρ^n , for any $n \in \mathbb{Z}$.

There are also new generating diagrams, all of degree zero, involving oriented strands. The generators involving only oriented strands are

$$(30) \quad \begin{array}{cccccc} \uparrow & \downarrow & \text{cup} & \text{cap} & \text{cup} & \text{cap} \end{array}$$

and the generating diagrams involving oriented strands and adjacent colored strands are

$$(31) \quad \begin{array}{cccc} \begin{array}{c} i \\ \diagup \\ \diagdown \\ i-1 \end{array} & \begin{array}{c} i-1 \\ \diagdown \\ \diagup \\ i \end{array} & \begin{array}{c} i-1 \\ \diagdown \\ \diagup \\ i \end{array} & \begin{array}{c} i \\ \diagdown \\ \diagup \\ i-1 \end{array} \end{array}$$

The new morphisms satisfy the following relations, where we again assume isotopy invariance and cyclicity.

- Relations involving only oriented strands:

$$(32) \quad \begin{array}{c} \circlearrowright \\ = 1 = \\ \circlearrowleft \end{array}$$

(33)

- Relation involving oriented strands and distant colored strands:

(34)

- Relations involving oriented strands and two adjacent colored strands:

(35)

(36)

(37)

- Relations involving oriented strands and three adjacent colored strands:

(38)

By relations (36), the sum of all colored dumbbells in (28) also commutes with oriented strands, so the corresponding morphism is also central in $\widehat{\mathcal{BS}}_d^{\text{ext}}$.

In general, any object in $\widehat{\mathcal{BS}}_d^{\text{ext}}$ is isomorphic to a direct sum of objects of the form $B_\rho^n B_w$, for some $n \in \mathbb{Z}$ and word w in S . By the relations in (32), there is an isomorphism of vector spaces

(and of algebras)

$$\left(\widehat{\mathcal{BS}}_d^{\text{ext}}\right)^0(B_\rho^m, B_\rho^n) \cong \begin{cases} \text{Cid}_{B_\rho^m}, & \text{if } m = n; \\ \{0\}, & \text{else.} \end{cases}$$

Recall that $R = \widehat{\mathcal{BS}}(\emptyset, \emptyset)$ is the polynomial algebra in the colored dumbbells. Then the isomorphism above generalizes to an isomorphism of graded R - R -bimodules

$$\widehat{\mathcal{BS}}_d^{\text{ext}}(B_\rho^m, B_\rho^n) \cong \begin{cases} R^{\tau^m}, & \text{if } m = n; \\ \{0\}, & \text{else,} \end{cases}$$

where τ is the automorphism of R which sends the i -colored dumbbell to the $i + 1$ -colored dumbbell, for any $i \in \mathbb{Z}/d\mathbb{Z}$, and R^{τ^m} is the free rank-one R - R -bimodule with the normal left R -action and the right R -action twisted by τ^m .

Moreover, the black oriented part and the non-oriented colored part of any diagram can be separated by the above relations, resulting in an isomorphism of graded R - R -bimodules

$$\widehat{\mathcal{BS}}_d^{\text{ext}}(B_\rho^m B_{\underline{u}}, B_\rho^m B_{\underline{v}}) \cong \begin{cases} R^{\tau^m} \otimes_R \widehat{\mathcal{BS}}_d(B_{\underline{u}}, B_{\underline{v}}), & \text{if } m = n; \\ \{0\}, & \text{else.} \end{cases}$$

In particular, this implies that the natural embedding $\widehat{\mathcal{BS}}_d \hookrightarrow \widehat{\mathcal{BS}}_d^{\text{ext}}$ is full. For the proofs of these results, see [E2018, Section 3.3].

Definition 3.5. The *extended diagrammatic Soergel category* $\widehat{\mathcal{S}}_d^{\text{ext}}$ is the idempotent completion of $\left(\widehat{\mathcal{BS}}_d^{\text{ext}}\right)^{\text{sh}}$.

The above results on the Hom-spaces in $\widehat{\mathcal{BS}}_d^{\text{ext}}$ and Theorem 3.3 imply the following generalization to the extended case, see [MT2017, Theorem 2.5].

Theorem 3.6. *For any $n \in \mathbb{Z}$ and $w \in \widehat{\mathcal{S}}_d$, the object $B_\rho^n B_w \in \widehat{\mathcal{S}}_d^{\text{ext}}$ is indecomposable. Moreover, the $\mathbb{Z}[q, q^{-1}]$ -linear map*

$$\begin{aligned} \left(\widehat{H}_d^{\text{ext}}\right)^{\mathbb{Z}[q, q^{-1}]} &\rightarrow [\widehat{\mathcal{S}}_d^{\text{ext}}]_{\oplus} \\ \rho^n b_w &\mapsto B_\rho^n B_w, \quad n \in \mathbb{Z}, w \in \widehat{\mathcal{S}}_d \end{aligned}$$

is an isomorphism of algebras.

4. ROUQUIER COMPLEXES

For \mathcal{A} a \mathbb{C} -linear, additive category, we write $\mathcal{K}^b(\mathcal{A})$ for the homotopy category of bounded complexes in \mathcal{A} . If \mathcal{A} is monoidal, then the usual monoidal product of chain complexes equips $\mathcal{K}^b(\mathcal{A})$ with a monoidal structure as well. If \mathcal{A} is graded, then $\mathcal{K}^b(\mathcal{A})$ is bigraded and we denote the shift inherited from \mathcal{A} by $\langle \cdot \rangle$ and the homological shift by $[\cdot]$.

Remark 4.1. Throughout this section, we sometimes state and prove diagrammatic equations in $\mathcal{K}^b(\mathcal{BS}_d)$, instead of $\mathcal{K}^b(\mathcal{S}_d)$. This makes no real difference in our case, as the differentials of the complexes in $\mathcal{K}^b(\mathcal{BS}_d)$ below are always given by matrices of homogeneous diagrams of the same degree, so they always give rise to objects in $\mathcal{K}^b(\mathcal{S}_d)$. See also Remark 3.2.

Let $\mathcal{C} = \mathcal{S}_d$. For the simple reflection $s_i \in W$ the *Rouquier complex* $T_i := T_{s_i} \in \mathcal{K}^b(\mathcal{S}_d)$ is defined by

$$T_i := \underline{B}_i \xrightarrow{\text{⋮}} R\langle 1 \rangle,$$

with B_i placed in homological degree zero (we always underline terms in homological degree zero). This complex is invertible in $\mathcal{K}^b(\mathcal{S}_d)$, with inverse given by

$$T_i^{-1} := R\langle -1 \rangle \xrightarrow{\text{⋮}} \underline{B}_i,$$

as follows from the homotopy equivalences which we recall below. These complexes were introduced in [Rou2006] and categorify the usual generators of the braid group, in particular, they satisfy the braid relations up to homotopy equivalence [Rou2006, Theorem 3.2]. By Matsumoto’s theorem, this implies that, for any $w \in S_n$, the complex T_w can be defined as

$$T_w := T_{i_1} \cdots T_{i_\ell},$$

where $\underline{w} = s_{i_1} \cdots s_{i_\ell}$ is any rex of w (i.e., up to homotopy equivalence, the complex does not depend on the choice of rex).

In subsection 4.1, we briefly recall the results on Rouquier complexes that are relevant for the definition of the evaluation functor. For more details, see [Rou2006], [EKr2010, §3] and [EMTW2020, Chapter 19]. In Subsection 4.2, we introduce a special Rouquier complex, denoted T_ρ , and develop a diagrammatic calculus for morphisms in $\mathcal{K}^b(\mathcal{S}_d)$ whose source and/or target contain tensor powers of T_ρ and T_ρ^{-1} . To the best of our knowledge, this extension of Soergel calculus has not appeared in the literature before.

4.1. Some diagrammatic shortcuts I: general Rouquier complexes. For $i \in \{1, \dots, d - 1\}$, let $\phi_i: T_i^{-1}T_i \rightarrow R$ denote the homotopy equivalence (where 1 stands for the identity map)

(39)

and $\psi_i: \mathbb{T}_i \mathbb{T}_i^{-1} \rightarrow R$ the analogous homotopy equivalence

$$(40) \quad \begin{array}{ccc} & \begin{array}{c} \text{B}_i \mathbb{B}_i \\ \oplus \\ \mathbb{R} \\ \downarrow 1 \\ \mathbb{R} \end{array} & \\ \begin{array}{c} \text{B}_i \langle -1 \rangle \\ \swarrow \\ \text{B}_i \langle 1 \rangle \end{array} & \begin{array}{c} \text{B}_i \mathbb{B}_i \\ \swarrow \\ \mathbb{R} \\ \downarrow 1 \\ \mathbb{R} \end{array} & \begin{array}{c} \text{B}_i \langle 1 \rangle \\ \swarrow \\ \mathbb{R} \\ \downarrow 1 \\ \mathbb{R} \end{array} \end{array}$$

(Note: The diagram above is a schematic representation of the commutative diagram in the image. It shows a central node $\text{B}_i \mathbb{B}_i \oplus \mathbb{R}$ with arrows to $\text{B}_i \langle -1 \rangle$, $\text{B}_i \langle 1 \rangle$, and \mathbb{R} . There are also arrows from $\text{B}_i \langle -1 \rangle$ and $\text{B}_i \langle 1 \rangle$ to \mathbb{R} . A vertical arrow labeled 1 goes from \mathbb{R} to \mathbb{R} . Blue symbols (dots, lines, and brackets) are associated with the nodes and arrows.)

in $\mathcal{K}^b(\mathcal{S}_d)$. These maps are well-known, see e.g. [EKr2010, §3].

Let further $\eta_{i,\pm}: \mathbb{T}_i^{\pm 1} \mathbb{T}_i^{\mp 1} \rightarrow \mathbb{T}_i^{\pm 1} R \mathbb{T}_i^{\mp 1}$ be the canonical isomorphisms $\mathcal{K}^b(\mathcal{S}_d)$, for any $i \in \{1, \dots, d-1\}$, both given by $ab \mapsto a1b$. To simplify notation, we write 1^m for $1 \cdots 1$ (m times) in the sequel.

The following can be checked directly.

Lemma 4.2. *For any $i \in \{1, \dots, d-1\}$, the composite maps*

$$\begin{aligned} \mathbb{T}_i &\xrightarrow{a \mapsto 1a} R \mathbb{T}_i \xrightarrow{\psi_i^{-1} \text{Id}_{\mathbb{T}_i}} \mathbb{T}_i \mathbb{T}_i^{-1} \mathbb{T}_i \xrightarrow{\text{Id}_{\mathbb{T}_i} \phi_i} \mathbb{T}_i R \xrightarrow{ab \mapsto ab} \mathbb{T}_i, \\ \mathbb{T}_i &\xrightarrow{a \mapsto a1} \mathbb{T}_i R \xrightarrow{\text{Id}_{\mathbb{T}_i} \phi_i^{-1}} \mathbb{T}_i \mathbb{T}_i^{-1} \mathbb{T}_i \xrightarrow{\psi_i \text{Id}_{\mathbb{T}_i}} R \mathbb{T}_i \xrightarrow{ba \mapsto ba} \mathbb{T}_i, \end{aligned}$$

are both equal to $\text{Id}_{\mathbb{T}_i}$ in $\mathcal{K}^b(\mathcal{S}_d)$, and the composite maps

$$\begin{aligned} \mathbb{T}_i^{-1} &\xrightarrow{a \mapsto 1a} R \mathbb{T}_i^{-1} \xrightarrow{\phi_i^{-1} \text{Id}_{\mathbb{T}_i^{-1}}} \mathbb{T}_i^{-1} \mathbb{T}_i \mathbb{T}_i^{-1} \xrightarrow{\text{Id}_{\mathbb{T}_i^{-1}} \psi_i} \mathbb{T}_i^{-1} R \xrightarrow{ab \mapsto ab} \mathbb{T}_i^{-1}, \\ \mathbb{T}_i^{-1} &\xrightarrow{a \mapsto a1} \mathbb{T}_i^{-1} R \xrightarrow{\text{Id}_{\mathbb{T}_i^{-1}} \psi_i^{-1}} \mathbb{T}_i^{-1} \mathbb{T}_i \mathbb{T}_i^{-1} \xrightarrow{\phi_i \text{Id}_{\mathbb{T}_i^{-1}}} R \mathbb{T}_i^{-1} \xrightarrow{ba \mapsto ba} \mathbb{T}_i^{-1}, \end{aligned}$$

are both equal to $\text{Id}_{\mathbb{T}_i^{-1}}$ in $\mathcal{K}^b(\mathcal{S}_d)$.

We now introduce the diagrammatics for the maps involving $\mathbb{T}_i^{\pm 1}$ that will be needed in the sequel. For any $i \in \{1, \dots, d-1\}$, we depict the identity morphisms of $\mathbb{T}_i^{\pm 1}$ as

$$\text{Id}_{\mathbb{T}_i} := \begin{array}{c} \uparrow \\ \vdots \\ i \end{array} \quad \text{and} \quad \text{Id}_{\mathbb{T}_i^{-1}} := \begin{array}{c} \downarrow \\ \vdots \\ i \end{array}$$

The degree zero homotopy equivalences in (39) and (40) (which are the units and counits of left and right adjunction of \mathbb{T}_i and \mathbb{T}_i^{-1}) are then depicted as


$$\begin{array}{c} \text{U}_i \\ \text{C}_i \end{array} \quad \text{and} \quad \begin{array}{c} \text{C}_i \\ \text{U}_i \end{array}$$


(Note: The diagram above is a schematic representation of the four homotopy equivalences shown in the image. Each is a blue dashed arc with a dot labeled 'i' at its center.)

and the above remarks translate into the following diagrammatic relations.

Lemma 4.3. For any $i \in \{1, \dots, d - 1\}$, we have the following relations between morphisms of $\mathcal{K}^b(\mathcal{S}_d)$:

(41) 

(42) 


(43) 

Remark 4.4. Just for the record, we give some further results about Rouquier complexes, all well-known to experts.

- For any $i \in \{1, \dots, d - 2\}$, the isomorphism between $T_i T_{i+1} T_i$ and $T_{i+1} T_i T_{i+1}$ in $\mathcal{K}^b(\mathcal{S}_d)$ (see [EKr2010, §3] for the maps) can be represented by the degree zero diagrams



satisfying the relations



There are similar diagrams and relations for braid moves involving the inverses of Rouquier complexes, see e.g. [EW2017, §5]. In Remark 4.12 below, we introduce some new diagrams.

- For any $i \in \{1, \dots, d - 1\}$, the cone of the map $f: T_i \rightarrow T_i^{-1}$, which is the identity on B_i and zero everywhere else, is isomorphic to

$$\underline{R\langle -1 \rangle} \xrightarrow{\bullet_i} R\langle 1 \rangle.$$

in $\mathcal{K}^b(\mathcal{S}_d)$. The distinguished triangle

$$T_i^{-1} \rightarrow \text{Cone}(f) \rightarrow T_i \rightarrow T_i[1]$$

categorifies the quadratic relation in the Hecke algebra H_d .

The remaining lemmas of this subsection are probably all known to experts, but we were not able to find any concrete references for them, so we decided to give all relevant homotopy equivalences explicitly. Further, to keep the notation as simple as possible, we state some equations in $\mathcal{K}^b(\mathcal{BS}_d)$. Being homogeneous, they also give rise to equations between morphisms in $\mathcal{K}^b(\mathcal{S}_d)$, as explained in Remark 4.1.

Lemma 4.5. *For any $i, j \in \{1, \dots, d-1\}$ such that $j = i \pm 1$, the following dumbbell-slide relations hold in $\mathcal{K}^b(\mathcal{BS}_d)$:*

$$\begin{array}{c} \begin{array}{c} \bullet \\ \uparrow \\ \bullet \\ \vdots \\ \bullet \\ \uparrow \\ \bullet \\ i \end{array} = - \begin{array}{c} \bullet \\ \vdots \\ \bullet \\ \uparrow \\ \bullet \\ i \end{array} \\ \\ \begin{array}{c} \bullet \\ \uparrow \\ \bullet \\ \vdots \\ \bullet \\ \uparrow \\ \bullet \\ j \\ i \end{array} = \begin{array}{c} \bullet \\ \vdots \\ \bullet \\ \uparrow \\ \bullet \\ j \\ i \end{array} + \begin{array}{c} \bullet \\ \vdots \\ \bullet \\ \uparrow \\ \bullet \\ \bullet \\ i \end{array} \end{array}$$

Proof. We are actually going to prove the equations in $\mathcal{K}^b(\mathcal{S}_d)$, fixing the shifts of the objects. For the first equation, consider


$$\begin{array}{ccccc} \mathbb{T}_i\langle 2 \rangle : & \mathbb{B}_i\langle 2 \rangle & \xrightarrow{\quad \bullet \quad} & \mathbb{R}\langle 3 \rangle & \\ \uparrow & \uparrow & \swarrow & \uparrow & \\ \mathbb{T}_i : & \mathbb{B}_i & \xrightarrow{\quad \bullet \quad} & \mathbb{R}\langle 1 \rangle & \\ & & & \uparrow & \\ & & & \begin{array}{c} \bullet \\ \vdots \\ \bullet \\ 2 \end{array} & \end{array}$$

The vertical arrows correspond to the map of complexes represented by


$$\begin{array}{c} \bullet \\ \uparrow \\ \bullet \\ \vdots \\ \bullet \\ \uparrow \\ \bullet \\ i \end{array} + \begin{array}{c} \bullet \\ \vdots \\ \bullet \\ \uparrow \\ \bullet \\ i \end{array}$$

and the diagonal arrow is a homotopy. Using (16), we see that the map of complexes is null-homotopic.

For the second equation, consider



The vertical arrows correspond to the map of complexes represented by




and the diagonal arrow is a homotopy. Using (23), we see that the map of complexes is null-homotopic. \square

Lemma 4.6. *There is an isomorphism*

$$T_i^{\pm 1} B_i T_i^{\mp 1} \cong B_i$$

in $\mathcal{K}^b(\mathcal{S}_d)$.

Proof. Recall that $B_i B_i \cong B_i\langle 1 \rangle \oplus B_i\langle -1 \rangle$ in \mathcal{S}_d . Using that isomorphism, it is easy to see that $T_i B_i \cong B_i\langle -1 \rangle$ in $\mathcal{K}^b(\mathcal{S}_d)$, with the homotopy equivalence between the complexes being given by



An analogous homotopy equivalence shows that $B_i T_i \cong B_i\langle -1 \rangle$ in $\mathcal{K}^b(\mathcal{S}_d)$ and thus that $T_i B_i \cong B_i T_i$ in $\mathcal{K}^b(\mathcal{S}_d)$.

Of course, the above also implies that $B_i T_i^{-1} \cong B_i\langle 1 \rangle \cong T_i^{-1} B_i$ in $\mathcal{K}^b(\mathcal{S}_d)$. \square

Lemma 4.7. *For each $1 \leq i \leq d - 2$, there are isomorphisms*

$$f_{i,\pm}: T_{i+1}^{\pm 1} B_i T_{i+1}^{\mp 1} \rightarrow T_i^{\mp 1} B_{i+1} T_i^{\pm 1}$$

in $\mathcal{K}^b(\mathcal{S}_d)$.

Proof. In this case, the complexes are actually isomorphic, not just homotopy equivalent. In the following figure, we exhibit the isomorphism $f_{i,-} : \mathbb{T}_{i+1}^{-1} B_i \mathbb{T}_{i+1} \rightarrow \mathbb{T}_i B_{i+1} \mathbb{T}_i^{-1}$ and its inverse $g_{i,-}$ (to avoid cluttering, we do not write labels in diagrams if they are clear from context):

$$\begin{array}{ccccc}
\mathbb{T}_{i+1}^{-1} B_i \mathbb{T}_{i+1} : & B_i B_{i+1} \langle -1 \rangle & \xrightarrow{\begin{pmatrix} \bullet & | & | \\ | & \bullet & | \\ | & | & \bullet \end{pmatrix}} & \left(\frac{B_{i+1} B_i B_{i+1}}{B_i} \right) & \xrightarrow{\begin{pmatrix} | & | & \bullet \\ | & \bullet & | \\ | & | & \bullet \end{pmatrix}} & B_{i+1} B_i \langle 1 \rangle \\
\updownarrow & \updownarrow -1 & & \updownarrow \bar{g}_{i,-} \quad \downarrow \bar{f}_{i,-} & & \updownarrow 1 \\
\mathbb{T}_i B_{i+1} \mathbb{T}_i^{-1} : & B_i B_{i+1} \langle -1 \rangle & \xrightarrow{\begin{pmatrix} | & | & \bullet \\ | & \bullet & | \\ | & | & \bullet \end{pmatrix}} & \left(\frac{B_i B_{i+1} B_i}{B_{i+1}} \right) & \xrightarrow{\begin{pmatrix} -\bullet & | & | \\ | & \bullet & | \\ | & | & \bullet \end{pmatrix}} & B_{i+1} B_i \langle 1 \rangle
\end{array}$$

Here $\bar{f}_{i,-}$ and $\bar{g}_{i,-}$ are, respectively,

$$\bar{f}_{i,-} = \begin{pmatrix} \begin{array}{c} \text{---} \times \text{---} \\ \text{---} \cup \text{---} \end{array} & \begin{array}{c} \bullet \\ | \\ \cup \\ \bullet \end{array} \\ \begin{array}{c} \cup \\ \bullet \end{array} & 0 \end{pmatrix}, \quad \bar{g}_{i,-} = \begin{pmatrix} \begin{array}{c} \text{---} \times \text{---} \\ \text{---} \cup \text{---} \end{array} & \begin{array}{c} \bullet \\ | \\ \cup \\ \bullet \end{array} \\ \begin{array}{c} \cup \\ \bullet \end{array} & 0 \end{pmatrix}.$$

The maps $f_{i,-} = (1, \bar{f}_{i,-}, 1)$ and $g_{i,-} = (1, \bar{g}_{i,-}, 1)$ are mutual inverses and a pleasant exercise, using the relation in (20), shows that both of them are chain maps. The complexes $\mathbb{T}_{i+1} B_i \mathbb{T}_{i+1}^{-1}$ and $\mathbb{T}_i^{-1} B_{i+1} \mathbb{T}_i$ are isomorphic too, as they are adjoint to $\mathbb{T}_{i+1}^{-1} B_i \mathbb{T}_{i+1}$ and $\mathbb{T}_i B_{i+1} \mathbb{T}_i^{-1}$, respectively. Using the units and counits of adjunction, we obtain the isomorphism $f_{i,+} : \mathbb{T}_{i+1} B_i \mathbb{T}_{i+1}^{-1} \rightarrow \mathbb{T}_i^{-1} B_{i+1} \mathbb{T}_i$ and its inverse $g_{i,+}$. \square

Recall the homotopy equivalences $\phi_i : \mathbb{T}^{-1} \mathbb{T}_i \rightarrow R$ and $\psi_i : \mathbb{T}_i \mathbb{T}_i^{-1} \rightarrow R$ and put $\delta_{i,+} := \phi_i^{-1} \circ \psi_{i+1}$ and $\delta_{i,-} := \psi_i^{-1} \circ \phi_{i+1}$ (we suppress the maps $\eta_{i,\pm}$ whenever we use the diagrams \bullet and \cup). Below, we keep the notation from Lemma 4.7.

Lemma 4.8. *For each $1 \leq i \leq d-2$, the following maps are equal to zero in $\mathcal{K}^b(\mathcal{S}_d)$:*

$$(44) \quad f_{i,\pm} \circ (\text{Id}_{\mathbb{T}_{i+1}^{\pm 1}} \begin{array}{c} \bullet \\ | \\ \cup \\ \bullet \end{array} \text{Id}_{\mathbb{T}_{i+1}^{\mp 1}}) - (\text{Id}_{\mathbb{T}_i^{\mp 1}} \begin{array}{c} \bullet \\ | \\ \cup \\ \bullet \end{array} \text{Id}_{\mathbb{T}_i^{\pm 1}}) \circ \delta_{i,\pm} : \\
\mathbb{T}_{i+1}^{\pm 1} \mathbb{T}_{i+1}^{\mp 1} \rightarrow \mathbb{T}_i^{\mp 1} B_{i+1} \mathbb{T}_i^{\pm 1} \langle 1 \rangle,$$

$$(45) \quad (\text{Id}_{\mathbb{T}_i^{\mp 1}} \begin{array}{c} \bullet \\ | \\ \cup \\ \bullet \end{array} \text{Id}_{\mathbb{T}_i^{\pm 1}}) \circ f_{i,\pm} - \delta_{i,\pm} \circ (\text{Id}_{\mathbb{T}_{i+1}^{\pm 1}} \begin{array}{c} \bullet \\ | \\ \cup \\ \bullet \end{array} \text{Id}_{\mathbb{T}_{i+1}^{\mp 1}}) : \\
\mathbb{T}_{i+1}^{\pm 1} B_i \mathbb{T}_{i+1}^{\mp 1} \langle -1 \rangle \rightarrow \mathbb{T}_i^{\mp 1} \mathbb{T}_i^{\pm 1},$$

$$(46) \quad f_{i,\pm}^{-1} \circ (\text{Id}_{\mathbb{T}_i^{\mp 1}} \begin{array}{c} \bullet \\ | \\ \cup \\ \bullet \end{array} \text{Id}_{\mathbb{T}_i^{\pm 1}}) - (\text{Id}_{\mathbb{T}_{i+1}^{\pm 1}} \begin{array}{c} \bullet \\ | \\ \cup \\ \bullet \end{array} \text{Id}_{\mathbb{T}_{i+1}^{\mp 1}}) \circ \delta_{i,\pm}^{-1} : \\
\mathbb{T}_i^{\mp 1} \mathbb{T}_i^{\pm 1} \rightarrow \mathbb{T}_{i+1}^{\pm 1} B_i \mathbb{T}_{i+1}^{\mp 1} \langle 1 \rangle,$$

$$(47) \quad (\text{Id}_{\mathbb{T}_{i+1}^{\pm 1}} \begin{array}{c} \bullet \\ | \\ \text{Id}_{\mathbb{T}_{i+1}^{\mp 1}} \end{array}) \circ f_{i,\pm}^{-1} - \delta_{i,\pm}^{-1} \circ (\text{Id}_{\mathbb{T}_i^{\mp 1}} \begin{array}{c} \bullet \\ | \\ \text{Id}_{\mathbb{T}_i^{\pm 1}} \end{array}):$$

$$\mathbb{T}_i^{\mp 1} B_{i+1} \mathbb{T}_i^{\pm 1} \langle -1 \rangle \rightarrow \mathbb{T}_{i+1}^{\pm 1} \mathbb{T}_{i+1}^{\mp 1}.$$

Proof. We only need to prove that the maps in (44) and (45) are null-homotopic for $f_{i,-}$, the case of $f_{i,+}$ following by adjunction. Pre- and post-composing those two maps with the appropriate isomorphisms proves the analogous statement for the maps in (46) and (47) as well.

It is not hard to compute that the map of complexes in (44) is given by the vertical arrows in the diagram below:

$$\begin{array}{ccccc} & & \begin{array}{c} \left(\begin{array}{c} | | \bullet \\ \bullet | \end{array} \right) \\ \hline \end{array} & & \\ & & \xrightarrow{\quad} & & \begin{array}{c} \left(\begin{array}{c} \bullet | | \\ | \bullet \end{array} \right) \\ \hline \end{array} \\ & & \xrightarrow{\quad} & & \\ \mathbb{T}_i B_{i+1} \mathbb{T}_i^{-1} \langle 1 \rangle : & \begin{array}{c} B_i B_{i+1} \\ \uparrow \begin{array}{c} - \bullet | \\ | \end{array} \\ B_{i+1} \langle -1 \rangle \end{array} & \xrightarrow{\quad} & \begin{array}{c} \left(\frac{B_i B_{i+1} B_i \langle 1 \rangle}{B_{i+1} \langle 1 \rangle} \right) \\ \uparrow \begin{array}{c} \left(\begin{array}{c} \bullet | \\ | \bullet \end{array} \right) \\ \hline \end{array} \\ \left(\frac{B_{i+1} B_{i+1}}{\underline{R}} \right) \end{array} & \xrightarrow{\quad} & \begin{array}{c} B_{i+1} B_i \langle 2 \rangle \\ \uparrow \begin{array}{c} | \bullet \\ | \end{array} \\ B_{i+1} \langle 1 \rangle \end{array} \\ & & \xrightarrow{H_0} & & \xrightarrow{H_1} \\ & & \xrightarrow{g} & & \\ \mathbb{T}_{i+1}^{-1} \mathbb{T}_{i+1} : & & & & \end{array}$$

where

$$g = \begin{pmatrix} \begin{array}{c} \bullet \swarrow \swarrow \bullet \\ \downarrow \downarrow \end{array} & 0 \\ - \begin{array}{c} \bullet \swarrow \swarrow \\ \downarrow \downarrow \end{array} + \begin{array}{c} \bullet \swarrow \swarrow \\ \downarrow \downarrow \end{array} & - \begin{array}{c} \bullet \\ | \end{array} \end{pmatrix}.$$

It is also easy to check that this map is null-homotopic, with homotopies


$$H_0 = \left(- \begin{array}{c} \bullet \swarrow \swarrow \\ \downarrow \downarrow \end{array}, 0 \right), \quad H_1 = \begin{pmatrix} 0 \\ | \\ \bullet \end{pmatrix}.$$

This establishes (44). The proof of (45) can be obtained by a vertical reflexion of the diagrams above and exchanging the labels i and $i + 1$. \square


Remark 4.9. The isomorphisms in Lemma 4.7 have a diagrammatic interpretation in terms of degree zero generators in $\mathcal{K}^b(\mathcal{BS}_d)$

$$\begin{array}{ccc} \begin{array}{c} \bullet \swarrow \swarrow \bullet \\ \downarrow \downarrow \end{array} & & \begin{array}{c} \bullet \swarrow \swarrow \\ \downarrow \downarrow \end{array} \\ \hline \begin{array}{c} \bullet \swarrow \swarrow \\ \downarrow \downarrow \end{array} & & \begin{array}{c} \bullet \swarrow \swarrow \bullet \\ \downarrow \downarrow \end{array} \\ \hline \begin{array}{c} \bullet \\ | \end{array} & & \begin{array}{c} \bullet \\ | \end{array} \end{array}$$

and relations



Using these diagrams, Lemma 4.8 translates into



There are also analogous diagrams and relations with reversed orientation of the oriented strands.

Lemma 4.10. *There exist the following isomorphisms in $\mathcal{K}^b(\mathcal{S}_d)$:*

$$(48) \quad T_i^{-1} T_{i-1}^{-1} B_i \cong B_{i-1} T_i^{-1} T_{i-1}^{-1}, \quad 1 < i \leq d-1,$$

$$(49) \quad T_{i-1} T_i B_{i-1} \cong B_i T_{i-1} T_i, \quad 1 < i \leq d-1,$$

$$(50) \quad T_i^{\pm 1} T_{i-1}^{\pm 1} B_j \cong B_j T_i^{\pm 1} T_{i-1}^{\pm 1}, \quad j \notin \{i-2, i-1, i, i+1\}.$$

Proof. We start with (48), proving that both chain complexes have retractions that are homotopy equivalent to each other. Here is a homotopy equivalence between the complex $T_i^{-1} T_{i-1}^{-1} B_i$ and

its retraction $(T_i^{-1}T_{i-1}^{-1}B_i)_{\text{retr}}$:

$$\begin{array}{ccccc}
 & & \begin{pmatrix} \text{blue dot} \downarrow \\ \text{red dot} \downarrow \end{pmatrix} & & \\
 & & \downarrow & & \\
 T_i^{-1}T_{i-1}^{-1}B_i: & B_i\langle -2 \rangle & \xrightarrow{\quad} & \begin{pmatrix} B_iB_i\langle -1 \rangle \\ B_{i-1}B_i\langle -1 \rangle \end{pmatrix} & \xrightarrow{\quad} & \underline{B_iB_{i-1}B_i} \\
 & \uparrow 1 & & \uparrow g_l \downarrow f_l & & \uparrow 1 \\
 & & \begin{pmatrix} \frac{1}{2} \downarrow \\ \frac{1}{2} \text{blue dot} \downarrow \\ \text{red dot} \downarrow \end{pmatrix} & & \begin{pmatrix} \text{blue dot} \downarrow, \text{blue dot} \downarrow, - \\ \text{red dot} \downarrow, \text{red dot} \downarrow, \text{red dot} \downarrow \end{pmatrix} & & \\
 & & \downarrow h = (2\cdot, 0, 0) & & \downarrow & & \\
 & B_i\langle -2 \rangle & \xrightarrow{\quad} & \begin{pmatrix} B_i\langle -2 \rangle \\ B_i \\ B_{i-1}B_i\langle -1 \rangle \end{pmatrix} & \xrightarrow{\quad} & \underline{B_iB_{i-1}B_i} \\
 & & & \uparrow g'_l \downarrow f'_l & & \uparrow 1 \\
 & & & \begin{pmatrix} B_i \\ B_{i-1}B_i\langle -1 \rangle \end{pmatrix} & \xrightarrow{\quad} & \underline{B_iB_{i-1}B_i} \\
 & & & \begin{pmatrix} \text{blue dot} \downarrow \\ \text{red dot} \downarrow \end{pmatrix} & & \\
 & & & \downarrow & & \\
 (T_i^{-1}T_{i-1}^{-1}B_i)_{\text{retr}}: & & & & &
 \end{array}$$

The upper vertical arrows correspond to the mutually inverse maps $(1, f_l, 1)$ and $(1, g_l, 1)$ induced by the isomorphism $B_iB_i \cong B_i\langle -1 \rangle \oplus B_i\langle 1 \rangle$. The maps f_l and g_l are

$$f_l = \begin{pmatrix} \begin{pmatrix} \frac{1}{2} \downarrow \\ \text{blue dot} \downarrow \end{pmatrix} & 0 \\ \begin{pmatrix} \frac{1}{2} \downarrow \\ \text{blue dot} \downarrow \\ \text{red dot} \downarrow \end{pmatrix} & 0 \\ 0 & \begin{pmatrix} \text{red dot} \downarrow \\ \text{red dot} \downarrow \end{pmatrix} \end{pmatrix}, \quad g_l = \begin{pmatrix} \begin{pmatrix} \text{blue dot} \downarrow \\ \text{blue dot} \downarrow \end{pmatrix} & \begin{pmatrix} \text{blue dot} \downarrow \\ \text{red dot} \downarrow \end{pmatrix} & 0 \\ 0 & 0 & \begin{pmatrix} \text{red dot} \downarrow \\ \text{red dot} \downarrow \end{pmatrix} \end{pmatrix}.$$

The lower vertical maps are the mutually up-to-homotopy inverse maps $(f'_l, 1)$ and $(g'_l, 1)$ given below.

$$f'_l = \begin{pmatrix} \begin{pmatrix} - \\ \text{blue dot} \downarrow \end{pmatrix} & \begin{pmatrix} \text{red dot} \downarrow \\ \text{red dot} \downarrow \end{pmatrix} & 0 \\ -2 & \begin{pmatrix} \text{red dot} \downarrow \\ \text{red dot} \downarrow \end{pmatrix} & \begin{pmatrix} \text{red dot} \downarrow \\ \text{red dot} \downarrow \end{pmatrix} \end{pmatrix}, \quad g'_l = \begin{pmatrix} 0 & 0 \\ \begin{pmatrix} \text{red dot} \downarrow \\ \text{red dot} \downarrow \end{pmatrix} & 0 \\ 0 & \begin{pmatrix} \text{red dot} \downarrow \\ \text{red dot} \downarrow \end{pmatrix} \end{pmatrix}.$$

The fact that they define a homotopy equivalence uses the homotopy h (whose only non-zero entry is multiplication by 2) in the complex in the middle. We leave the details to the reader.

The following diagram gives a homotopy equivalence between the complex $B_{i-1}T_i^{-1}T_{i-1}^{-1}$ and its retraction $(B_{i-1}T_i^{-1}T_{i-1}^{-1})_{\text{retr}}$:

$$\begin{array}{ccccc}
 B_{i-1}T_i^{-1}T_{i-1}^{-1} : & B_{i-1}\langle -2 \rangle \xrightarrow{\begin{pmatrix} | & | \\ \bullet & \bullet \\ | & | \end{pmatrix}} & \begin{pmatrix} B_{i-1}B_i\langle -1 \rangle \\ B_{i-1}B_{i-1}\langle -1 \rangle \end{pmatrix} & \xrightarrow{\begin{pmatrix} | & | & | \\ \bullet & \bullet & \bullet \\ | & | & | \end{pmatrix}, -\begin{pmatrix} | & | \\ \bullet & \bullet \\ | & | \end{pmatrix}} & \underline{B_{i-1}B_iB_{i-1}} \\
 \updownarrow & \updownarrow 1 & \updownarrow g_r \quad \downarrow f_r & & \updownarrow 1 \\
 & B_{i-1}\langle -2 \rangle \xleftarrow[h=(0, 2\cdot, 0)]{\begin{pmatrix} | & | \\ \frac{1}{2} & | \\ | & \bullet \\ \frac{1}{2} & | \\ | & \bullet \end{pmatrix}} & \begin{pmatrix} B_{i-1}B_i\langle -1 \rangle \\ B_{i-1}\langle -2 \rangle \\ B_{i-1} \end{pmatrix} & \xrightarrow{\begin{pmatrix} | & | & | \\ \bullet & \bullet & \bullet \\ | & | & | \end{pmatrix}, -\begin{pmatrix} | & | \\ \bullet & \bullet \\ | & | \end{pmatrix}, -\begin{pmatrix} | & | \\ \bullet & \bullet \\ | & | \end{pmatrix}} & \underline{B_{i-1}B_iB_{i-1}} \\
 \downarrow & & \updownarrow g'_r \quad \downarrow f'_r & & \updownarrow 1 \\
 (B_{i-1}T_i^{-1}T_{i-1}^{-1})_{\text{retr}} : & & \begin{pmatrix} B_{i-1}B_i\langle -1 \rangle \\ B_{i-1} \end{pmatrix} & \xrightarrow{\begin{pmatrix} | & | & | \\ \bullet & \bullet & \bullet \\ | & | & | \end{pmatrix}, -\begin{pmatrix} | & | \\ \bullet & \bullet \\ | & | \end{pmatrix}} & \underline{B_{i-1}B_iB_{i-1}}
 \end{array}$$

The upper vertical arrows correspond to the mutually inverse maps $(1, f_r, 1)$ and $(1, g_r, 1)$, with f_r and g_r being


$$f_r = \begin{pmatrix} \begin{pmatrix} | & | \\ \bullet & \bullet \\ | & | \end{pmatrix} & 0 \\ 0 & \frac{1}{2} \begin{pmatrix} | & | \\ \cup \\ | & | \end{pmatrix} \\ 0 & \frac{1}{2} \begin{pmatrix} | & | \\ \bullet & \bullet \\ | & | \end{pmatrix} \end{pmatrix}, \quad g_r = \begin{pmatrix} \begin{pmatrix} | & | \\ \bullet & \bullet \\ | & | \end{pmatrix} & 0 & 0 \\ 0 & \begin{pmatrix} | & | \\ \bullet & \bullet \\ | & | \end{pmatrix} & \begin{pmatrix} | & | \\ \cup \\ | & | \end{pmatrix} \end{pmatrix}.$$

The lower vertical arrows correspond to the mutually up-to-homotopy inverse maps $(f'_r, 1)$ and $(g'_r, 1)$ given below.

$$f'_r = \begin{pmatrix} \begin{pmatrix} | & | \\ \bullet & \bullet \\ | & | \end{pmatrix} & -2 \begin{pmatrix} | & | \\ \bullet & \bullet \\ | & | \end{pmatrix} & 0 \\ 0 & -\begin{pmatrix} | & | \\ \bullet & \bullet \\ | & | \end{pmatrix} & | \end{pmatrix}, \quad g'_r = \begin{pmatrix} \begin{pmatrix} | & | \\ \bullet & \bullet \\ | & | \end{pmatrix} & 0 \\ 0 & 0 \\ 0 & | \end{pmatrix}.$$

We leave the details to the reader.

The diagram below shows that the complexes $(T_i^{-1}T_{i-1}^{-1}B_i)_{\text{retr}}$ and $(B_{i-1}T_i^{-1}T_{i-1}^{-1})_{\text{retr}}$ are homotopy equivalent.




This finishes the proof of the existence of the isomorphism in (48).


Tensoring both complexes in (48) with $T_{i-1}T_i$ on the left and on the right yields the isomorphism in (49).

The equivalence in (50) is clear, because the two complexes are canonically isomorphic. \square


Remark 4.11. The isomorphisms in Lemma 4.10 also have a diagrammatic interpretation in terms of degree zero generators in $\mathcal{K}^b(\mathcal{BS}_d)$




and relations



Remark 4.12. The following will not be used in the sequel. The canonical isomorphisms $B_iT_j^{\pm 1} \cong T_j^{\pm 1}B_i$, for distant i and j , translate into the generators




satisfying the relations



There are also maps $B_i \rightarrow T_i^{-1}$, $T_i \rightarrow B_i$, $R \rightarrow T_i$ and $T_i^{-1} \rightarrow R$ of non-zero degree in $\mathcal{K}^b(\mathcal{BS}_d)$, depicted respectively as



and satisfying certain diagrammatic relations, which are easy to deduce. Note also that the composite



is the map $T_i^{-1} \rightarrow T_i$ mentioned in Remark 4.4.


4.2. **Some diagrammatic shortcuts II: special Rouquier complexes $T_\rho^{\pm 1}$.** In this subsection, we introduce and study a special Rouquier complex, denoted T_ρ , which will play an important role in the definition of the evaluation functors.

Definition 4.13. Define

$$T_\rho := T_1 \cdots T_{d-1} \quad \text{and} \quad T_\rho^{-1} := T_{d-1}^{-1} \cdots T_1^{-1}$$

in $\mathcal{K}^b(\mathcal{S}_d)$.

In order to develop a diagrammatic calculus for these special Rouquier complexes, we first picture the *identity morphisms* of T_ρ and T_ρ^{-1} as upward and downward oriented arrows, respectively:



Further, we introduce *oriented cups and caps*





These correspond to the units and counits of left and right adjunction for T_ρ and T_ρ^{-1} in $\mathcal{K}^b(\mathcal{S}_d)$. Algebraically, they can be expressed in terms of the maps given in [Section 4.1](#)

$$\begin{aligned}
 \text{cup} &:= (1^{d-2}\psi_{d-1}^{-1}1^{d-2}) \circ (1^{d-3}\eta_{d-2,+}\psi_{d-2}^{-1}1^{d-3}) \circ \cdots \circ (1\eta_{2,+}\psi_2^{-1}1) \\
 &\quad \circ (\eta_{1,+}\psi_1^{-1}): R \rightarrow T_\rho T_\rho^{-1}, \\
 \text{cap} &:= (1^{d-2}\phi_1^{-1}1^{d-2}) \circ (1^{d-3}\eta_{2,-}\phi_2^{-1}1^{d-3}) \circ \cdots \circ (1\eta_{d-2,-}\phi_{d-2}^{-1}1) \\
 &\quad \circ (\eta_{d-1,-}\phi_{d-1}^{-1}): R \rightarrow T_\rho^{-1} T_\rho, \\
 \text{left cap} &:= (\phi_1\eta_{1,-}^{-1}) \circ (1\phi_2\eta_{2,-}^{-1}1) \circ \cdots \circ (1^{d-3}\phi_2\eta_{2,-}^{-1}1^{d-3}) \\
 &\quad \circ (1^{d-2}\phi_11^{d-2}): T_\rho^{-1} T_\rho \rightarrow R, \\
 \text{right cap} &:= (\psi_{d-1}\eta_{d-1,+}^{-1}) \circ (1\psi_{d-2}\eta_{d-2,+}^{-1}1) \circ \cdots \circ (1^{d-3}\psi_2\eta_{2,+}^{-1}1^{d-3}) \\
 &\quad \circ (1^{d-2}\psi_{d-1}1^{d-2}): T_\rho T_\rho^{-1} \rightarrow R,
 \end{aligned}$$

Lemma 4.14. *The oriented cups and caps satisfy the following relations in $\mathcal{K}^b(\mathcal{S}_d)$*

$$(51) \quad \text{cup} = 1 = \text{cap}$$

$$(52) \quad \begin{array}{c} \downarrow \\ \uparrow \end{array} = \begin{array}{c} \text{cup} \\ \text{cap} \end{array} \quad \begin{array}{c} \uparrow \\ \downarrow \end{array} = \begin{array}{c} \text{cap} \\ \text{cup} \end{array}$$

$$(53) \quad \begin{array}{c} \text{cup} \\ \downarrow \\ \text{cup} \end{array} = \begin{array}{c} \uparrow \\ \text{cap} \\ \uparrow \end{array} = \begin{array}{c} \text{cup} \\ \downarrow \\ \text{cup} \end{array} \quad \begin{array}{c} \text{cap} \\ \downarrow \\ \text{cap} \end{array} = \begin{array}{c} \uparrow \\ \text{cup} \\ \uparrow \end{array} = \begin{array}{c} \text{cap} \\ \downarrow \\ \text{cap} \end{array}$$

Proof. The relations in (53) are a consequence of [Lemma 4.2](#). The other relations are immediate. \square

The next diagrammatic generators involving oriented strands are the *mixed crossings*, which correspond to the following degree-zero isomorphisms in $\mathcal{K}^b(\mathcal{S}_d)$, for $1 < i \leq d-1$:

$$(54) \quad \begin{array}{c} \text{red strand } i-1 \\ \text{blue strand } i \\ \text{crossing} \end{array} = \text{Id}_{T_{d-1}^{-1} \dots T_{i+1}^{-1}} F_{i,r} \text{Id}_{T_{i-2}^{-1} \dots T_1^{-1}}: T_\rho^{-1} B_i \rightarrow B_{i-1} T_\rho^{-1},$$

where in homological degrees -2 , -1 and 0 , respectively, we define

$$(55) \quad F_{i,r} := \left(0, \left(\begin{array}{cc} \begin{array}{c} \text{blue } \uparrow \text{ and } \downarrow \\ \text{red } \uparrow \text{ and } \downarrow \end{array} & \begin{array}{c} \text{red } \uparrow \text{ and } \downarrow \\ \text{blue } \uparrow \text{ and } \downarrow \end{array} \\ \begin{array}{c} \text{red } \uparrow \text{ and } \downarrow \\ \text{blue } \uparrow \text{ and } \downarrow \end{array} & \begin{array}{c} \text{blue } \uparrow \text{ and } \downarrow \\ \text{red } \uparrow \text{ and } \downarrow \end{array} \end{array} \right), \text{blue } \times \text{red} \right).$$

This is the map obtained from the homotopy equivalence in [Lemma 4.10](#) by tensoring on the left with the identity morphism of $T_{d-1}^{-1} \cdots T_{i+1}^{-1}$ and on the right with the identity morphism of $T_{i-2}^{-1} \cdots T_1^{-1}$, and using when necessary the permutation isomorphism between $T_i^{-1}B_j$ and $B_jT_i^{-1}$ if $|i-j| \neq 1$.

Analogously,

$$\begin{array}{c} \text{blue } \uparrow \text{ and } \downarrow \\ \text{red } \uparrow \text{ and } \downarrow \end{array} \begin{array}{c} i \\ i-1 \end{array} = \text{Id}_{T_{d-1}^{-1} \cdots T_{i+1}^{-1}} G_{i,r} \text{Id}_{T_{i-2}^{-1} \cdots T_1^{-1}} : B_{i-1}T_\rho^{-1} \rightarrow T_\rho^{-1}B_i,$$

with

$$(56) \quad G_{i,r} = \left(0, \left(\begin{array}{cc} \begin{array}{c} \text{blue } \uparrow \text{ and } \downarrow \\ \text{red } \uparrow \text{ and } \downarrow \end{array} & \begin{array}{c} \text{red } \uparrow \text{ and } \downarrow \\ \text{blue } \uparrow \text{ and } \downarrow \end{array} \\ \begin{array}{c} \text{red } \uparrow \text{ and } \downarrow \\ \text{blue } \uparrow \text{ and } \downarrow \end{array} & \begin{array}{c} \text{blue } \uparrow \text{ and } \downarrow \\ \text{red } \uparrow \text{ and } \downarrow \end{array} \end{array} \right), \text{blue } \times \text{red} \right).$$

Of course, there are also mixed crossings involving T_ρ , which are depicted as

$$\begin{array}{c} \text{blue } \uparrow \text{ and } \downarrow \\ \text{red } \uparrow \text{ and } \downarrow \end{array} \begin{array}{c} i \\ i-1 \end{array} : T_\rho B_{i-1} \rightarrow B_i T_\rho \quad \text{and} \quad \begin{array}{c} \text{red } \uparrow \text{ and } \downarrow \\ \text{blue } \uparrow \text{ and } \downarrow \end{array} \begin{array}{c} i-1 \\ i \end{array} : B_i T_\rho \rightarrow T_\rho B_{i-1}.$$

Lemma 4.15. *For distant colors $i, j = 1, \dots, d-1$, we have*

$$(57) \quad \begin{array}{c} \text{orange } \uparrow \text{ and } \downarrow \\ \text{purple } \uparrow \text{ and } \downarrow \end{array} \begin{array}{c} j-1 \\ i-1 \end{array} \begin{array}{c} i \\ j \end{array} = \begin{array}{c} \text{orange } \uparrow \text{ and } \downarrow \\ \text{purple } \uparrow \text{ and } \downarrow \end{array} \begin{array}{c} j-1 \\ i-1 \end{array} \begin{array}{c} i \\ j \end{array}$$

in $\mathcal{K}^b(S_d)$.

Proof. It is clear that the map in (54) commutes with the 4-valent crossing for distant colors. \square

The proof of the following lemma is immediate and, therefore, omitted.

Lemma 4.16. *The mixed crossings in $\mathcal{K}^b(\mathcal{S}_d)$ satisfy the relations*

(58)

The diagram shows four equations. The first two are for crossings where the top strand is blue (index i) and the bottom strand is red (index $i-1$). The first equation shows a crossing where the blue strand goes over the red strand, equal to a blue strand on the left and a red strand on the right. The second equation shows a crossing where the red strand goes over the blue strand, equal to a red strand on the left and a blue strand on the right. The last two equations are for crossings where the top strand is red (index $i-1$) and the bottom strand is blue (index i). The third equation shows a crossing where the red strand goes over the blue strand, equal to a red strand on the left and a blue strand on the right. The fourth equation shows a crossing where the blue strand goes over the red strand, equal to a blue strand on the left and a red strand on the right.

Lemma 4.17. *The following diagrammatic relations hold in $\mathcal{K}^b(\mathcal{BS}_d)$:*

(59)

The diagram shows four equations. The first two are for crossings where the top strand is red (index $i-1$) and the bottom strand is blue (index i). The first equation shows a crossing where the red strand goes over the blue strand, equal to a red strand on the left and a blue strand on the right. The second equation shows a crossing where the blue strand goes over the red strand, equal to a blue strand on the left and a red strand on the right. The last two equations are for crossings where the top strand is blue (index i) and the bottom strand is red (index $i-1$). The third equation shows a crossing where the blue strand goes over the red strand, equal to a blue strand on the left and a red strand on the right. The fourth equation shows a crossing where the red strand goes over the blue strand, equal to a red strand on the left and a blue strand on the right.

Proof. We prove the first relation in (59), the proof of the others being similar. By (55), the maps of complexes corresponding to the two sides of (59) are

$$\begin{array}{c} \text{red} \\ \diagdown \\ \text{blue} \\ \diagup \\ \text{red} \end{array} = \text{Id}_{T_{d-1}^{-1} \dots T_{i+1}^{-1}} F \text{Id}_{T_{i-2}^{-1} \dots T_1^{-1}},$$

where in homological degrees $(-2, -1, 0)$, respectively, we have

$$F = \left(0, \left(\begin{array}{cc} \text{red} \downarrow & - \text{blue} \downarrow \\ - \text{red} \downarrow & \text{blue} \downarrow \end{array} \right), \begin{array}{c} \text{red} \diagdown \\ \text{blue} \diagup \\ \text{red} \end{array} \right),$$


and

$$\begin{array}{c} \text{red} \\ \diagdown \\ \text{red} \\ \diagup \\ \text{blue} \end{array} = \text{Id}_{T_{d-1}^{-1} \dots T_{i+1}^{-1}} G \text{Id}_{T_{i-2}^{-1} \dots T_1^{-1}},$$

where in the same homological degrees we have

$$G = \left(\text{red} \downarrow, \left(\begin{array}{cc} \text{red} \downarrow & 0 \\ 0 & \text{red} \downarrow \end{array} \right), \begin{array}{c} \text{red} \downarrow \\ \text{blue} \downarrow \end{array} \right).$$

The diagram below shows that $F - G$ is zero in $\mathcal{K}^b(\mathcal{S}_d)$:




with

$$(F - G)_{-1} = \begin{pmatrix} 0 & -\bullet \\ -\bullet & \text{Y} \end{pmatrix}, \quad H_{-1} = (0, -\bullet), \quad H_0 = \begin{pmatrix} 0 \\ -\bullet \text{Y} \end{pmatrix}.$$

This finishes the proof. □

Lemma 4.18. *The following diagrammatic equalities hold in $\mathcal{K}^b(\mathcal{S}_d)$:*



Proof. We prove the first relation in (60), as the other can be proved in a similar way. The proof is a consequence of the fact that the composites

$$B_{i-1}T_\rho^{-1}B_i \xrightarrow{\text{crossing}} B_{i-1}B_{i-1}T_\rho^{-1} \xrightarrow{\text{cap}} T_\rho^{-1}$$

and

$$B_{i-1}T_\rho^{-1}B_i \xrightarrow{\text{crossing}} T_\rho^{-1}B_iB_i \xrightarrow{\text{cup}} T_\rho^{-1},$$

are both given by

$$\text{Id}_{T_{d-1}^{-1} \dots T_{i+1}^{-1}} \left(0, \left(\begin{array}{cc} \text{red dot} \text{ blue } \downarrow & - \text{red } \downarrow \text{ blue } \\ - \text{red } \downarrow \text{ blue } & \text{red } \downarrow \text{ blue } \end{array} \right), \text{red } \downarrow \text{ blue } \right) \text{Id}_{T_{i-2}^{-1} \dots T_1^{-1}}.$$

This computation is straightforward and uses (55) and (56). □

Remark 4.19. By Lemma 4.18, we can define

$$\begin{array}{c} \text{red } \downarrow \text{ blue } \\ \text{red } \downarrow \text{ blue } \end{array} := \begin{array}{c} \text{red } \downarrow \text{ blue } \\ \text{red } \downarrow \text{ blue } \end{array} = \begin{array}{c} \text{red } \downarrow \text{ blue } \\ \text{red } \downarrow \text{ blue } \end{array}$$

and similarly

$$\begin{array}{c} \text{red } \downarrow \text{ blue } \\ \text{red } \downarrow \text{ blue } \end{array}, \quad \begin{array}{c} \text{red } \downarrow \text{ blue } \\ \text{red } \downarrow \text{ blue } \end{array} \quad \text{and} \quad \begin{array}{c} \text{red } \downarrow \text{ blue } \\ \text{red } \downarrow \text{ blue } \end{array}$$

Lemma 4.20. The following pitchfork relations hold in $\mathcal{K}^b(\mathcal{BS}_d)$:

$$(62) \quad \begin{array}{cc} \begin{array}{c} \text{red } \downarrow \text{ blue } \\ \text{red } \downarrow \text{ blue } \end{array} = \begin{array}{c} \text{red } \downarrow \text{ blue } \\ \text{red } \downarrow \text{ blue } \end{array}, & \begin{array}{c} \text{red } \downarrow \text{ blue } \\ \text{red } \downarrow \text{ blue } \end{array} = \begin{array}{c} \text{red } \downarrow \text{ blue } \\ \text{red } \downarrow \text{ blue } \end{array}, \\ \begin{array}{c} \text{red } \downarrow \text{ blue } \\ \text{red } \downarrow \text{ blue } \end{array} = \begin{array}{c} \text{red } \downarrow \text{ blue } \\ \text{red } \downarrow \text{ blue } \end{array}, & \begin{array}{c} \text{red } \downarrow \text{ blue } \\ \text{red } \downarrow \text{ blue } \end{array} = \begin{array}{c} \text{red } \downarrow \text{ blue } \\ \text{red } \downarrow \text{ blue } \end{array}. \end{array}$$

Proof. We only prove the first relation in (62), as the others can be proved in a similar way. Relations (54) and (55) imply that

$$\begin{array}{c} \text{red } \downarrow \text{ blue } \\ \text{red } \downarrow \text{ blue } \end{array} = \text{Id}_{T_{d-1}^{-1} \dots T_{i+1}^{-1}} F \text{Id}_{T_{i-2}^{-1} \dots T_1^{-1}} : T_{\rho}^{-1} B_i B_i \rightarrow B_{i-1} B_{i-1} T_{\rho}^{-1},$$

where F in homological degrees $-2, -1$ and 0 , respectively, is given by

$$F = \left(0, \left(\begin{array}{cc} \text{red dot} \text{ blue } \downarrow + \text{red } \downarrow \text{ blue } & - \text{red } \downarrow \text{ blue } - \text{red } \downarrow \text{ blue } \\ - \text{red } \downarrow \text{ blue } - \text{red } \downarrow \text{ blue } & \text{red } \downarrow \text{ blue } + \text{red } \downarrow \text{ blue } \end{array} \right), \begin{array}{c} \text{red } \downarrow \text{ blue } \\ \text{red } \downarrow \text{ blue } \end{array} \right).$$

Pre-composing with

$$\begin{array}{c} \text{red } \downarrow \text{ blue } \\ \text{red } \downarrow \text{ blue } \end{array}$$

results in

$$\begin{array}{c} \text{Diagram: a crossing of a red line over a blue line, with a purple arrow pointing down from the crossing.} \end{array} = \text{Id}_{T_{d-1}^{-1} \dots T_{i+1}^{-1}} F_{\text{pitchfork}} \text{Id}_{T_{i-2}^{-1} \dots T_1^{-1}},$$

where

$$F_{\text{pitchfork}} = \left(0, \left(\begin{array}{cc} \text{Diagram: blue Y-shape} & \text{Diagram: red Y-shape with blue dot} \\ \text{Diagram: red Y-shape} & \text{Diagram: blue Y-shape with blue dot} \end{array} \right), \text{Diagram: mixed crossing} \right).$$

In homological degree zero we have used (22) with a blue dot on the leftmost blue endpoint. The proof is now completed by the observation that



is given by exactly the same map, which can be seen immediately by post-composing the mixed crossing in (54) with



and using (55). □

Lemma 4.21. *The following diagrammatic equalities hold in $\mathcal{K}^b(\mathcal{S}_d)$, for any adjacent triple $i-1, i, i+1 \in \{1, \dots, d-1\}$:*

$$(63) \quad \begin{array}{c} \text{Diagram: crossing of red and blue lines with arcs, labeled } i-1, i, i-1 \text{ at top and } i+1, i, i+1 \text{ at bottom.} \end{array} = \begin{array}{c} \text{Diagram: crossing of red and blue lines with arcs, labeled } i-1, i, i-1 \text{ at top and } i+1, i, i+1 \text{ at bottom.} \end{array} \quad \begin{array}{c} \text{Diagram: crossing of red and blue lines with arcs, labeled } i, i-1, i \text{ at top and } i, i+1, i \text{ at bottom.} \end{array} = \begin{array}{c} \text{Diagram: crossing of red and blue lines with arcs, labeled } i, i-1, i \text{ at top and } i, i+1, i \text{ at bottom.} \end{array}$$

Proof. Both diagrams in the first equality represent morphisms between $T_\rho^{-1} B_{i+1} B_i B_{i+1} T_\rho$ and $B_{i-1} B_i B_{i-1}$. By (35), there is an isomorphism $T_\rho^{-1} B_{i+1} B_i B_{i+1} T_\rho \cong B_i B_{i-1} B_i$, so both diagrams correspond to morphisms in


$$\mathcal{S}_d(B_i B_{i-1} B_i, B_{i-1} B_i B_{i-1}).$$

Recall that $B_i B_{i-1} B_i \cong B_{i(i-1)i} \oplus B_i$ and $B_{i-1} B_i B_{i-1} \cong B_{i(i-1)i} \oplus B_{i-1}$, which implies that

$$\mathcal{S}_d(B_i B_{i-1} B_i, B_{i-1} B_i B_{i-1}) \cong \mathcal{S}_d(B_{i(i-1)i}, B_{i(i-1)i}) \cong \mathbb{C}$$

by Soergel's Hom-formula in (29).

In particular, this implies that the two diagrams in the first equality are multiples of each other. To check that they are actually equal, one can attach a dot at an appropriate place. For example, one can easily check that



in $\mathcal{K}^b(\mathcal{BS}_d)$ by using relations (20) and (59), followed by (58) and (62). The second equality of the statement is proved in the same way. □

The *mixed 6-valent vertices* represent the following isomorphisms in $\mathcal{K}^b(\mathcal{S}_d)$, obtained by recursive application of [Lemma 4.7](#):

(64)

$$\begin{array}{c}
 \begin{array}{ccc}
 & d-1 & \\
 \nearrow & & \nwarrow \\
 \text{---} & & \text{---} \\
 \searrow & & \swarrow \\
 & 1 & \\
 \end{array} & : T_\rho^{-1} B_1 T_\rho \rightarrow T_\rho B_{d-1} T_\rho^{-1} \\
 \\
 \begin{array}{ccc}
 & 1 & \\
 \nearrow & & \nwarrow \\
 \text{---} & & \text{---} \\
 \searrow & & \swarrow \\
 & d-1 & \\
 \end{array} & : T_\rho^{-1} B_{d-1} T_\rho \rightarrow T_\rho B_1 T_\rho^{-1}
 \end{array}$$

Remark 4.22. To understand why we have introduced the mixed 6-valent vertices above, recall that the evaluation functors are (yet to be defined) functors from $\widehat{\mathcal{S}}_d^{\text{ext}}$ to $\mathcal{K}^b(\mathcal{S}_d)$, and that in $\widehat{\mathcal{S}}_d^{\text{ext}}$ there are mutually inverse isomorphisms

$$\begin{array}{ccc}
 \begin{array}{ccc}
 & d-1 & \\
 \nearrow & & \nwarrow \\
 \text{---} & & \text{---} \\
 \searrow & & \swarrow \\
 & 1 & \\
 \end{array} & := & \begin{array}{ccc}
 & d-1 & \\
 \curvearrowright & & \curvearrowleft \\
 \text{---} & & \text{---} \\
 \curvearrowleft & & \curvearrowright \\
 & 1 & \\
 \end{array} \\
 \text{and} & & \\
 \begin{array}{ccc}
 & 1 & \\
 \nearrow & & \nwarrow \\
 \text{---} & & \text{---} \\
 \searrow & & \swarrow \\
 & d-1 & \\
 \end{array} & := & \begin{array}{ccc}
 & 1 & \\
 \curvearrowright & & \curvearrowleft \\
 \text{---} & & \text{---} \\
 \curvearrowleft & & \curvearrowright \\
 & d-1 & \\
 \end{array}
 \end{array}$$

Lemma 4.23. *The mixed 6-valent vertices satisfy*

(65)

$$\begin{array}{ccc}
 \begin{array}{ccc}
 & 1 & \\
 \curvearrowright & & \curvearrowleft \\
 \text{---} & & \text{---} \\
 \curvearrowleft & & \curvearrowright \\
 & 1 & \\
 \end{array} = \begin{array}{ccc}
 \text{---} & & \text{---} \\
 \downarrow & & \uparrow \\
 \text{---} & & \text{---} \\
 \downarrow & & \uparrow \\
 \text{---} & & \text{---} \\
 & 1 & \\
 \end{array} \\
 \\
 \begin{array}{ccc}
 & d-1 & \\
 \curvearrowright & & \curvearrowleft \\
 \text{---} & & \text{---} \\
 \curvearrowleft & & \curvearrowright \\
 & d-1 & \\
 \end{array} = \begin{array}{ccc}
 \text{---} & & \text{---} \\
 \uparrow & & \downarrow \\
 \text{---} & & \text{---} \\
 \uparrow & & \downarrow \\
 \text{---} & & \text{---} \\
 & d-1 & \\
 \end{array}
 \end{array}$$

in $\mathcal{K}^b(\mathcal{S}_d)$.

Lemma 4.24. *The mixed 6-valent vertices also satisfy the following dot relations in $\mathcal{K}^b(\mathcal{BS}_d)$:*

(66)

$$\begin{array}{ccc}
 \begin{array}{ccc}
 & d-1 & \\
 \nearrow & & \nwarrow \\
 \text{---} & & \text{---} \\
 \searrow & & \swarrow \\
 & \bullet & \\
 & d-1 & \\
 \end{array} = \begin{array}{ccc}
 & d-1 & \\
 \curvearrowright & & \curvearrowleft \\
 \text{---} & & \text{---} \\
 \curvearrowleft & & \curvearrowright \\
 & \bullet & \\
 & d-1 & \\
 \end{array} \\
 \\
 \begin{array}{ccc}
 & 1 & \\
 \nearrow & & \nwarrow \\
 \text{---} & & \text{---} \\
 \searrow & & \swarrow \\
 & \bullet & \\
 & 1 & \\
 \end{array} = \begin{array}{ccc}
 & 1 & \\
 \curvearrowright & & \curvearrowleft \\
 \text{---} & & \text{---} \\
 \curvearrowleft & & \curvearrowright \\
 & \bullet & \\
 & 1 & \\
 \end{array}
 \end{array}$$

(67)

$$\begin{array}{ccc}
 \begin{array}{ccc}
 & d-1 & \\
 \nearrow & & \nwarrow \\
 \text{---} & & \text{---} \\
 \searrow & & \swarrow \\
 & \bullet & \\
 & d-1 & \\
 \end{array} = \begin{array}{ccc}
 & d-1 & \\
 \curvearrowright & & \curvearrowleft \\
 \text{---} & & \text{---} \\
 \curvearrowleft & & \curvearrowright \\
 & \bullet & \\
 & d-1 & \\
 \end{array} \\
 \\
 \begin{array}{ccc}
 & 1 & \\
 \nearrow & & \nwarrow \\
 \text{---} & & \text{---} \\
 \searrow & & \swarrow \\
 & \bullet & \\
 & 1 & \\
 \end{array} = \begin{array}{ccc}
 & 1 & \\
 \curvearrowright & & \curvearrowleft \\
 \text{---} & & \text{---} \\
 \curvearrowleft & & \curvearrowright \\
 & \bullet & \\
 & 1 & \\
 \end{array}
 \end{array}$$

Proof. Apply [Lemma 4.8](#) recursively. □

Lemma 4.25. *The following mixed dumbbell-slide relation holds in $\mathcal{K}^b(\mathcal{BS}_d)$:*

$$(68) \quad \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \uparrow \\ i \end{array} = \begin{array}{c} \uparrow \\ | \\ \bullet \\ | \\ \bullet \\ | \\ i-1 \end{array} \quad i = 2, \dots, d-1$$

$$(69) \quad \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \uparrow \\ 1 \end{array} = - \begin{array}{c} \uparrow \\ | \\ \sum_{i=1}^{d-1} \bullet \\ | \\ \bullet \\ | \\ i \end{array}$$

$$(70) \quad - \sum_{i=1}^{d-1} \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \uparrow \\ i \end{array} = \begin{array}{c} \uparrow \\ | \\ \bullet \\ | \\ \bullet \\ | \\ d-1 \end{array}$$

Proof. The equality in (68) is an immediate consequence of (59).

For (69) apply the non-oriented dumbbell-slides from Lemma 4.5

$$\begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \uparrow \\ 1 \end{array} = - \begin{array}{c} \uparrow \\ | \\ \bullet \\ | \\ \bullet \\ | \\ 1 \end{array} \quad \text{and} \quad \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \uparrow \\ i+1 \end{array} = \begin{array}{c} \uparrow \\ | \\ \bullet \\ | \\ \bullet \\ | \\ i+1 \end{array} + \begin{array}{c} \uparrow \\ | \\ \bullet \\ | \\ \bullet \\ | \\ i+1 \end{array}$$

recursively.

Finally, for (70) use the same non-oriented dumbbell-slides as above but with the colors i and $i+1$ swapped. \square

To prove Lemmas 4.26 to 4.29 below, we use the same strategy as in the proof of Lemma 4.21: we first check that a certain hom-space is one-dimensional and then conclude that two morphisms in that hom-space are equal by attaching dots to the corresponding diagrams.

Lemma 4.26. *The mixed 6-valent vertices satisfy the following cyclicity relations in $\mathcal{K}^b(\mathcal{S}_d)$:*

$$(71) \quad \begin{array}{c} d-1 \quad 1 \\ \swarrow \quad \searrow \\ \uparrow \quad \downarrow \\ \swarrow \quad \searrow \\ d-1 \quad 1 \end{array} = \begin{array}{c} d-1 \quad 1 \\ \swarrow \quad \searrow \\ \downarrow \quad \uparrow \\ \swarrow \quad \searrow \\ d-1 \quad 1 \end{array} \quad \begin{array}{c} 1 \quad d-1 \\ \swarrow \quad \searrow \\ \uparrow \quad \downarrow \\ \swarrow \quad \searrow \\ 1 \quad d-1 \end{array} = \begin{array}{c} 1 \quad d-1 \\ \swarrow \quad \searrow \\ \downarrow \quad \uparrow \\ \swarrow \quad \searrow \\ 1 \quad d-1 \end{array}$$

$$(72) \quad \begin{array}{c} \uparrow \quad \downarrow \\ \swarrow \quad \searrow \\ \uparrow \quad \downarrow \\ d-1 \quad 1 \end{array} = \begin{array}{c} \uparrow \quad \downarrow \\ \swarrow \quad \searrow \\ \downarrow \quad \uparrow \\ d-1 \quad 1 \end{array} \quad \begin{array}{c} \uparrow \quad \downarrow \\ \swarrow \quad \searrow \\ \downarrow \quad \uparrow \\ 1 \quad d-1 \end{array} = \begin{array}{c} \uparrow \quad \downarrow \\ \swarrow \quad \searrow \\ \downarrow \quad \uparrow \\ 1 \quad d-1 \end{array}$$

Proof. We only prove the first relation in (71), as the remaining ones are proved in the same way. We claim that the two morphisms in (71) are multiples of one another. To see this, note that $T_\rho B_{d-1} T_\rho^{-1} \cong T_\rho^{-1} B_1 T_\rho$ and $B_1 B_1 \cong B_1 \langle -1 \rangle \oplus B_1 \langle 1 \rangle$, whence

$$\begin{aligned} \mathcal{K}^b(\mathcal{S}_d) (R, T_\rho B_{d-1} T_\rho^{-1} T_\rho^{-1} B_1 T_\rho) &\cong \mathcal{K}^b(\mathcal{S}_d) (R, T_\rho B_1 B_1 T_\rho) \\ &\cong \mathcal{S}_d (R, B_1 B_1) \end{aligned}$$

$$\cong \mathcal{S}_d(R, \mathbf{B}_1\langle -1 \rangle \oplus \mathbf{B}_1\langle 1 \rangle),$$

where we have used the biadjointness of \mathbf{T}_ρ and its inverse, and the fullness of the natural embedding of \mathcal{S}_d in $\mathcal{K}^b(\mathcal{S}_d)$, for the second isomorphism. By Soergel's Hom-formula in (29), we know that

$$\dim_{\mathbb{C}}(\mathcal{S}_d(R, \mathbf{B}_1\langle -1 \rangle)) = 0 \quad \text{and} \quad \dim_{\mathbb{C}}(\mathcal{S}_d(R, \mathbf{B}_1\langle 1 \rangle)) = 1$$

and hence

$$\dim_{\mathbb{C}}(\mathcal{K}^b(\mathcal{S}_d)(R, \mathbf{T}_\rho \mathbf{B}_{d-1} \mathbf{T}_\rho^{-1} \mathbf{T}_\rho^{-1} \mathbf{B}_i \mathbf{T}_\rho)) = 1.$$

Attaching a dot to one of the colored strands (say with 1) on both sides of (71) and using the relations in Lemma 4.24 and certain isotopies shows that both morphisms are equal in $\mathcal{K}^b(\mathcal{B}\mathcal{S}_d)$. \square

Lemma 4.27. *For each $j \in \{1, \dots, d-1\}$ distant from 1 and $d-1$, the following equalities hold in $\mathcal{K}^b(\mathcal{S}_d)$:*

$$(73) \quad \begin{array}{c} \begin{array}{ccc} \begin{array}{c} \text{d-1} \\ \nearrow \quad \searrow \\ \text{j} \quad \text{1} \end{array} & = & \begin{array}{c} \text{1} \\ \nearrow \quad \searrow \\ \text{j} \quad \text{d-1} \end{array} \\ \begin{array}{c} \text{d-1} \\ \nearrow \quad \searrow \\ \text{j} \quad \text{1} \end{array} & = & \begin{array}{c} \text{1} \\ \nearrow \quad \searrow \\ \text{j} \quad \text{d-1} \end{array} \end{array}$$

Proof. We only prove the first equality, as the other can be proved in the same way. By adjointness, proving the first equality in (73) is equivalent to proving the equality

$$(74) \quad \begin{array}{c} \begin{array}{ccc} \begin{array}{c} \text{d-1} \quad \text{1} \\ \nearrow \quad \searrow \\ \text{j} \end{array} & = & \begin{array}{c} \text{d-1} \quad \text{1} \\ \nearrow \quad \searrow \\ \text{j} \end{array} \end{array}$$

in $\mathcal{K}^b(\mathcal{S}_d)$.

For any $j \in \{1, \dots, d-1\}$ distant from 1 and $d-1$, the same arguments as before (and the fact that \mathbf{B}_1 and \mathbf{B}_{j+1} commute) prove the following isomorphisms of hom-spaces:

$$\begin{aligned} \mathcal{K}^b(\mathcal{S}_d)(\mathbf{B}_j, \mathbf{T}_\rho \mathbf{B}_{d-1} \mathbf{T}_\rho^{-1} \mathbf{B}_j \mathbf{T}_\rho^{-1} \mathbf{B}_1 \mathbf{T}_\rho) &\cong \mathcal{K}^b(\mathcal{S}_d)(\mathbf{B}_j, \mathbf{T}_\rho^{-1} \mathbf{B}_1 \mathbf{T}_\rho \mathbf{B}_j \mathbf{T}_\rho^{-1} \mathbf{B}_1 \mathbf{T}_\rho) \\ &\cong \mathcal{K}^b(\mathcal{S}_d)(\mathbf{T}_\rho \mathbf{B}_j \mathbf{T}_\rho^{-1}, \mathbf{B}_1 \mathbf{T}_\rho \mathbf{B}_j \mathbf{T}_\rho^{-1} \mathbf{B}_1) \\ &\cong \mathcal{S}_d(\mathbf{B}_{j+1}, \mathbf{B}_1 \mathbf{B}_{j+1} \mathbf{B}_1) \\ &\cong \mathcal{S}_d(\mathbf{B}_{j+1}, \mathbf{B}_{j+1} \mathbf{B}_1 \mathbf{B}_1) \\ &\cong \mathcal{S}_d(\mathbf{B}_{j+1}, \mathbf{B}_{j+1} \mathbf{B}_1 \langle -1 \rangle \oplus \mathbf{B}_{j+1} \mathbf{B}_1 \langle 1 \rangle) \end{aligned}$$

By Soergel's Hom-formula in (29), we know that


$$\dim_{\mathbb{C}}(\mathcal{S}_d(\mathbf{B}_{j+1}, \mathbf{B}_{j+1} \mathbf{B}_1 \langle -1 \rangle)) = 0 \quad \text{and} \quad \dim_{\mathbb{C}}(\mathcal{S}_d(\mathbf{B}_{j+1}, \mathbf{B}_{j+1} \mathbf{B}_1 \langle 1 \rangle)) = 1$$

whence

$$\dim_{\mathbb{C}}(\mathcal{K}^b(\mathcal{S}_d)(\mathbf{B}_j, \mathbf{T}_\rho \mathbf{B}_{d-1} \mathbf{T}_\rho^{-1} \mathbf{B}_j \mathbf{T}_\rho^{-1} \mathbf{B}_1 \mathbf{T}_\rho)) = 1$$

and the equality in (74) can be proved by attaching dots to these diagrams at appropriate places. \square

Lemma 4.28. *The following equalities are true in $\mathcal{K}^b(\mathcal{S}_d)$:*

(75) 

Proof. We first note that

$$\begin{aligned} T_\rho^{-1}B_1T_\rho B_{d-1}T_\rho^{-1}B_1T_\rho &\cong T_\rho B_{d-1}T_\rho^{-1}B_{d-1}T_\rho B_{d-1}T_\rho^{-1} \\ &\cong T_\rho B_{d-1}B_{d-2}B_{d-1}T_\rho^{-1}, \end{aligned}$$

and

$$\begin{aligned} B_{d-1}T_\rho^{-1}B_1T_\rho B_{d-1} &\cong B_{d-1}T_\rho B_{d-1}T_\rho^{-1}B_{d-1} \\ &\cong T_\rho B_{d-2}B_{d-1}B_{d-2}T_\rho^{-1}, \end{aligned}$$

and therefore

$$\begin{aligned} \mathcal{K}^b(\mathcal{S}_d) (B_{d-1}T_\rho^{-1}B_1T_\rho B_{d-1}, T_\rho^{-1}B_1T_\rho B_{d-1}T_\rho^{-1}B_1T_\rho) \\ \cong \mathcal{S}_d (B_{d-1}B_{d-2}B_{d-1}, B_{d-2}B_{d-1}B_{d-2}) \end{aligned}$$

By the decompositions

$$B_{d-1}B_{d-2}B_{d-1} \cong B_{(d-1)(d-2)(d-1)} \oplus B_{d-1} \quad \text{and} \quad B_{d-2}B_{d-1}B_{d-2} \cong B_{(d-1)(d-2)(d-1)} \oplus B_{d-2}$$


and Soergel's Hom-formula in (29), we conclude that

$$\dim_{\mathbb{C}} (\mathcal{K}^b(\mathcal{S}_d) (B_{d-1}T_\rho^{-1}B_1T_\rho B_{d-1}, T_\rho^{-1}B_1T_\rho B_{d-1}T_\rho^{-1}B_1T_\rho)) = 1.$$

Thus the two diagrams in (75) are scalar multiples of each other and the equality now follows by attaching dots to these diagrams at appropriate places. \square

The proof of the following lemma uses exactly the same arguments as above and is left as an exercise to the reader.

Lemma 4.29. *The following equalities hold in $\mathcal{K}^b(\mathcal{S}_d)$:*

(76) 

$$(77) \quad \begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \end{array} = \begin{array}{c} \text{Diagram 3} \\ \text{Diagram 4} \end{array}$$

Diagram 1: A crossing of a blue strand (1) and a green strand (d-1). A purple strand crosses both. An orange strand (3) crosses the blue strand.

Diagram 2: Similar to Diagram 1, but the orange strand (3) crosses the green strand.

Diagram 3: Similar to Diagram 1, but the orange strand (3) crosses the purple strand.

Diagram 4: Similar to Diagram 1, but the orange strand (3) crosses the purple strand.

$$(78) \quad \begin{array}{c} \text{Diagram 5} \\ \text{Diagram 6} \end{array} = \begin{array}{c} \text{Diagram 7} \\ \text{Diagram 8} \end{array}$$

Diagram 5: A crossing of a blue strand (1) and a green strand (d-1). A purple strand crosses both.

Diagram 6: Similar to Diagram 5, but the green strand (d-1) crosses the blue strand.

Diagram 7: A crossing of a blue strand (1) and a green strand (d-1). A purple strand crosses both.

Diagram 8: Similar to Diagram 7, but the green strand (d-1) crosses the blue strand.

5. EVALUATION FUNCTORS

In this section, we finally define the evaluation functors $\mathcal{E}v_{r,s}: \widehat{\mathcal{S}}_d^{\text{ext}} \rightarrow \mathcal{K}^b(\mathcal{S}_d)$, for $r, s \in \mathbb{Z}$, which categorify the evaluation maps ev_a from Definition 2.1, for $a = (-1)^s q^r$ with $r, s \in \mathbb{Z}$. The other evaluation maps in that definition, denoted ev'_a , can be categorified likewise, but we don't work out the details here.

Remark 5.1. To be really precise, we actually define a degree-preserving functor from $\widehat{\mathcal{B}\mathcal{S}}_d^{\text{ext}}$ to $\mathcal{K}^b((\mathcal{B}\mathcal{S}_d^{\text{sh}})^{\text{gr}})$ which uniquely determines $\mathcal{E}v_{r,s}$, see Remarks 3.2 and 4.1. Note that $(\mathcal{B}\mathcal{S}_d^{\text{sh}})^{\text{gr}}$ is a graded category with shift, and that $X\langle t \rangle \cong X$ for every $X \in (\mathcal{B}\mathcal{S}_d^{\text{sh}})^{\text{gr}}$ and $t \in \mathbb{Z}$. The natural, degree-preserving embedding of $\mathcal{B}\mathcal{S}_d$ into $(\mathcal{B}\mathcal{S}_d^{\text{sh}})^{\text{gr}}$ is therefore fully faithful and essentially surjective, although it is not an equivalence of graded categories because its inverse is not degree-preserving. However, for our purposes all that matters is that the monoidal subcategory of degree-zero morphisms $((\mathcal{B}\mathcal{S}_d^{\text{sh}})^{\text{gr}})^0$ is isomorphic with $\mathcal{B}\mathcal{S}_d^{\text{sh}}$, which implies that the idempotent completion of both is \mathcal{S}_d . This might sound a bit complicated, but we can not simply define a functor from $\widehat{\mathcal{B}\mathcal{S}}_d^{\text{ext}}$ to $\mathcal{K}^b(\mathcal{B}\mathcal{S}_d)$ because the image of B_ρ requires non-trivial internal shifts when $r \neq 0$.

5.1. **Definition.** Let $r, s \in \mathbb{Z}$ and $d \in \mathbb{N}_{\geq 3}$ be arbitrary but fixed for the remainder of this section.

The *evaluation functor* is the monoidal, \mathbb{C} -linear functor

$$\mathcal{E}v_{r,s}: \widehat{\mathcal{S}}_d^{\text{ext}} \rightarrow \mathcal{K}^b(\mathcal{S}_d)$$

commuting with shifts which is uniquely determined (see Remark 5.1) by the monoidal, degree-preserving, \mathbb{C} -linear functor

$$\mathcal{E}v_{r,s}: \widehat{\mathcal{B}\mathcal{S}}_d^{\text{ext}} \rightarrow \mathcal{K}^b((\mathcal{B}\mathcal{S}_d^{\text{sh}})^{\text{gr}})$$

defined below. Note that we use the same notation for both functors.

- On the (non-full) subcategory $\mathcal{B}\mathcal{S}_d$ of $\widehat{\mathcal{B}\mathcal{S}}_d^{\text{ext}}$, the evaluation functor $\mathcal{E}v_{r,s}$ is the identity. More specifically, this means that $\mathcal{E}v_{r,s}(B_i) := B_i$ for every $i \in \{1, \dots, d-1\}$ and that $\mathcal{E}v_{r,s}$ sends any diagram without unoriented 0-colored strands and oriented strands to itself.

On other objects of $\widehat{\mathcal{BS}}_d^{\text{ext}}$, it is defined as

$$\begin{aligned}\mathcal{E}v_{r,s}(\mathbf{B}_0) &:= \mathbf{T}_\rho^{-1} \mathbf{B}_1 \mathbf{T}_\rho, \\ \mathcal{E}v_{r,s}(\mathbf{B}_\rho^{\pm 1}) &:= \mathbf{T}_\rho^{\pm 1} \langle \pm r \rangle [\pm s].\end{aligned}$$

On other morphisms it is defined as follows.

- On oriented and 0-colored generators:

$$\begin{aligned}\mathcal{E}v_{r,s}\left(\begin{array}{c} \uparrow \\ \uparrow \end{array}\right) &= \begin{array}{c} \uparrow \\ \uparrow \end{array} & \mathcal{E}v_{r,s}\left(\begin{array}{c} \downarrow \\ \downarrow \end{array}\right) &= \begin{array}{c} \downarrow \\ \downarrow \end{array} \\ \mathcal{E}v_{r,s}\left(\begin{array}{c} \curvearrowright \end{array}\right) &= \begin{array}{c} \curvearrowright \end{array} & \mathcal{E}v_{r,s}\left(\begin{array}{c} \curvearrowleft \end{array}\right) &= \begin{array}{c} \curvearrowleft \end{array} \\ \mathcal{E}v_{r,s}\left(\begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array}\right) &= \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} & \mathcal{E}v_{r,s}\left(\begin{array}{c} \curvearrowleft \\ \curvearrowright \end{array}\right) &= \begin{array}{c} \curvearrowleft \\ \curvearrowright \end{array} \\ \mathcal{E}v_{r,s}\left(\begin{array}{c} 0 \\ \downarrow \end{array}\right) &= \begin{array}{c} 0 \\ \downarrow \end{array} & \mathcal{E}v_{r,s}\left(\begin{array}{c} 1 \\ \downarrow \end{array}\right) &= \begin{array}{c} 1 \\ \downarrow \end{array} \\ \mathcal{E}v_{r,s}\left(\begin{array}{c} 0 \\ \bullet \end{array}\right) &= \begin{array}{c} 1 \\ \bullet \end{array} & \mathcal{E}v_{r,s}\left(\begin{array}{c} 1 \\ \bullet \\ 0 \end{array}\right) &= \begin{array}{c} 1 \\ \bullet \\ 1 \end{array} \\ \mathcal{E}v_{r,s}\left(\begin{array}{c} \text{Y-junction} \\ 0 \end{array}\right) &= \begin{array}{c} \text{Y-junction} \\ 1 \end{array} & \mathcal{E}v_{r,s}\left(\begin{array}{c} \text{Y-junction} \\ 0 \end{array}\right) &= \begin{array}{c} \text{Y-junction} \\ 1 \end{array}\end{aligned}$$

- On generators including strands with *distant colors*:

$$\begin{aligned}\mathcal{E}v_{r,s}\left(\begin{array}{c} \text{Cross} \\ 0 \quad i \end{array}\right) &= \begin{array}{c} \text{Cross} \\ 1 \quad i \end{array} & \text{for } i \neq 1, d-1 \\ \mathcal{E}v_{r,s}\left(\begin{array}{c} \text{Cross} \\ i \quad 0 \end{array}\right) &= \begin{array}{c} \text{Cross} \\ i \quad 1 \end{array} & \text{for } i \neq 1, d-1\end{aligned}$$

- On generators including strands with *adjacent colors*:

$$(79) \quad \mathcal{E}v_{r,s}\left(\begin{array}{c} \text{Cross} \\ i-1 \quad i \end{array}\right) = \begin{array}{c} \text{Cross} \\ i-1 \quad i \end{array} \quad \mathcal{E}v_{r,s}\left(\begin{array}{c} \text{Cross} \\ i-1 \quad i \end{array}\right) = \begin{array}{c} \text{Cross} \\ i-1 \quad i \end{array}$$

$$(80) \quad \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{red } i-1 \\ \text{blue } i \end{array} \right) = \begin{array}{c} \text{red } i-1 \\ \text{blue } i \end{array} \quad \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{blue } i \\ \text{red } i-1 \end{array} \right) = \begin{array}{c} \text{blue } i \\ \text{red } i-1 \end{array}$$

if $i \neq 0, 1$, while

$$(81) \quad \mathcal{E}v_{r,s} \left(\begin{array}{c} 0 \\ \text{blue } 1 \end{array} \right) = \begin{array}{c} \text{blue } 1 \\ \text{purple } 0 \end{array} \quad \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{blue } 1 \\ 0 \end{array} \right) = \begin{array}{c} \text{blue } 1 \\ \text{purple } 0 \end{array}$$

$$(82) \quad \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{purple } 0 \\ \text{blue } 1 \end{array} \right) = \begin{array}{c} \text{purple } 0 \\ \text{blue } 1 \end{array} \quad \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{blue } 1 \\ \text{purple } 0 \end{array} \right) = \begin{array}{c} \text{blue } 1 \\ \text{purple } 0 \end{array}$$

$$(83) \quad \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{green } d-1 \\ \text{purple } 0 \end{array} \right) = \begin{array}{c} \text{green } d-1 \\ \text{purple } 1 \end{array} \quad \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{purple } 0 \\ \text{green } d-1 \end{array} \right) = \begin{array}{c} \text{purple } 1 \\ \text{green } d-1 \end{array}$$

$$(84) \quad \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{purple } d-1 \\ \text{green } 0 \end{array} \right) = \begin{array}{c} \text{purple } d-1 \\ \text{green } 1 \end{array} \quad \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{green } 0 \\ \text{purple } d-1 \end{array} \right) = \begin{array}{c} \text{green } 1 \\ \text{purple } d-1 \end{array}$$

and

$$(85) \quad \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{blue } 1 \\ \text{purple } 0 \\ \text{blue } 1 \end{array} \right) = \begin{array}{c} \text{blue } 1 \\ \text{purple } 1 \\ \text{blue } 1 \end{array}$$

$$(86) \quad \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{purple } 0 \\ \text{blue } 1 \\ \text{purple } 0 \end{array} \right) = \begin{array}{c} \text{purple } 0 \\ \text{blue } 1 \\ \text{purple } 0 \end{array}$$

$$(87) \quad \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram} \\ d-1 \quad 0 \quad d-1 \end{array} \right) = \text{Diagram}$$

$$(88) \quad \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram} \\ 0 \quad d-1 \quad 0 \end{array} \right) = \text{Diagram}$$

This ends the definition of $\mathcal{E}v_{r,s}$.

Remark 5.2. Since $T_\rho^{-1}B_1T_\rho \cong T_\rho B_{d-1}T_\rho^{-1}$ in $\mathcal{K}^b((\mathcal{BS}_d^{\text{sh}})^{\text{gr}})$, we could have defined $\mathcal{E}v_{r,s}(B_0)$ as $T_\rho B_{d-1}T_\rho^{-1}$. These two choices result in naturally isomorphic evaluation functors, the isomorphism being induced by the 6-valent vertices (64), as can be checked by straightforward diagrammatic calculations.

Remark 5.3. The apparent lack of symmetry between the image via $\mathcal{E}v_{r,s}$ of the mixed 4-vertices involving strands colored 0 and 1, and the corresponding image of the mixed 4-vertices involving colored 0 and $d-1$ ((81) to (84)) is explained by Remark 5.2. Note also that

$$\mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram} \\ d-1 \\ 0 \\ 1 \end{array} \right) = \begin{array}{c} \text{Diagram} \\ d-1 \\ 1 \end{array} \quad \text{and} \quad \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram} \\ 1 \\ 0 \\ d-1 \end{array} \right) = \begin{array}{c} \text{Diagram} \\ 1 \\ d-1 \end{array}$$

5.2. Proof of well-definedness.

Theorem 5.4. *The monoidal functor $\mathcal{E}v_{r,s}$ is well-defined.*

Proof. The fact that $\mathcal{E}v_{r,s}$ preserves isotopy invariance follows from Lemma 4.14, Lemma 4.18 and Lemma 4.26, together with isotopy invariance of the usual (non-oriented) Soergel calculus.

- Relations involving only *one color*. We only need to check for color 0. Relations (13) and (14) are clear. For the remaining one-color relations we have

$$\mathcal{E}v_{r,s} \left(\begin{array}{c} \text{circle} \\ 0 \end{array} \right) = \begin{array}{c} \text{circle with dot} \\ 1 \end{array} \stackrel{(51)}{=} \begin{array}{c} \text{circle with dot} \\ 1 \end{array} \stackrel{(15)}{=} 0,$$

and

$$\mathcal{E}v_{r,s} \left(\begin{array}{c} | \\ 0 \end{array} + \begin{array}{c} | \\ 0 \end{array} \right) = \begin{array}{c} \text{circle with dot} \\ 1 \end{array} \downarrow \uparrow + \downarrow \uparrow \begin{array}{c} \text{circle with dot} \\ 1 \end{array} \stackrel{(52)}{=} \begin{array}{c} \text{circle with dot} \\ 1 \end{array} \downarrow \uparrow + \downarrow \uparrow \begin{array}{c} \text{circle with dot} \\ 1 \end{array} \stackrel{(16)}{=} 2 \begin{array}{c} \downarrow \uparrow \\ 1 \end{array} = \mathcal{E}v_{r,s} \left(\begin{array}{c} 2 \\ | \\ 0 \end{array} \right).$$

- Relations involving *two distant colors*. Here $j \neq 1, d - 1$.

$$\mathcal{E}v_{r,s} \left(\begin{array}{c} \text{cross} \\ 0 \quad j \end{array} \right) = \begin{array}{c} \text{cross with dot} \\ 1 \quad j \end{array} \stackrel{(17),(58)}{=} \begin{array}{c} \downarrow \uparrow \\ 1 \quad j \end{array} = \mathcal{E}v_{r,s} \left(\begin{array}{c} | \\ | \end{array} \right),$$

$$\mathcal{E}v_{r,s} \left(\begin{array}{c} \text{dot} \\ \text{cross} \\ j \end{array} \right) = \begin{array}{c} \text{dot with dot} \\ \text{cross} \\ j \end{array} \stackrel{(18),(58)}{=} \begin{array}{c} \text{dot with dot} \\ \text{cross} \\ j \end{array} = \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{dot} \\ \text{cross} \\ j \end{array} \right),$$

$$\mathcal{E}v_{r,s} \left(\begin{array}{c} \text{triple} \\ j \\ 0 \end{array} \right) = \begin{array}{c} \text{triple with dot} \\ 1 \end{array} \stackrel{(19),(58)}{=} \begin{array}{c} \text{triple with dot} \\ 1 \end{array} = \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{triple} \\ j \end{array} \right).$$

The corresponding relations with the colors 0 and j switched are proved in the same way.

- Relations involving *two adjacent colors*. We have to check the cases involving either the pair $(0, 1)$ or the pair $(0, d - 1)$. For the pair $(0, 1)$ we compute:

$$\begin{aligned} \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{cross} \\ 0 \end{array} \right) &= \begin{array}{c} \text{cross with dot} \\ 1 \end{array} \stackrel{(59)}{=} \begin{array}{c} \text{cross with dot} \\ 1 \end{array} \stackrel{(20)}{=} \begin{array}{c} \text{cross with dot} \\ 1 \end{array} + \begin{array}{c} \text{cross with dot} \\ 1 \end{array} \\ &\stackrel{(52),(58),(59)}{=} \begin{array}{c} \text{cross with dot} \\ 1 \end{array} + \begin{array}{c} \text{cross with dot} \\ 1 \end{array} = \mathcal{E}v_{r,s} \left(\begin{array}{c} | \\ | \\ | \end{array} + \begin{array}{c} \text{cross} \\ 1 \end{array} \right), \end{aligned}$$

$$\begin{aligned}
\mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram} \\ 0 \end{array} \right) &= \text{Diagram} \stackrel{(58)}{=} \text{Diagram} \stackrel{(21)}{=} \text{Diagram} + \text{Diagram} \\
&\stackrel{(52),(58),(59)}{=} \text{Diagram} + \text{Diagram} = \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram} \\ \text{Diagram} \end{array} \right), \\
\mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram} \\ 0 \end{array} \right) &= \text{Diagram} \stackrel{(51),(58)}{=} \text{Diagram} \stackrel{(22)}{=} \text{Diagram} \\
&\stackrel{(51),(58)}{=} \text{Diagram} = \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram} \end{array} \right), \\
\mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram} \\ 0 \end{array} \right) - \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram} \\ 0 \end{array} \right) &= \text{Diagram} - \text{Diagram} \stackrel{(68)}{=} \text{Diagram} - \text{Diagram} \stackrel{(23)}{=} \frac{1}{2} \left(\text{Diagram} - \text{Diagram} \right) \\
&\stackrel{(52),(53)}{=} \frac{1}{2} \left(\text{Diagram} - \text{Diagram} \right) = \frac{1}{2} \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram} \\ \text{Diagram} \end{array} \right).
\end{aligned}$$

The relations with the colors 0 and 1 switched are proved in the same way. The relations for the pair $(0, d-1)$ can be proved similarly, using the image of the corresponding mixed 6-valent vertex, of course.

- The relation involving *three distant colors* is straightforward and follows from the observation that the case involving colors 0, i and j , with $1 < i, j < d-1$ and distant implies checking a relation involving the colors 1, $i+1$ and $j+1$, which are still distant.

- The relation involving a *distant dumbbell* colored $i \in \{2, \dots, d - 2\}$ and a straight line colored 0 is straightforward, because (68) implies that it reduces to the same relation involving a distant dumbbell with color $i + 1$ and a straight line colored 1. Similarly, the relation involving a *distant dumbbell* colored 0 and a straight line colored $i \in \{2, \dots, d - 2\}$ reduces to the relation involving a distant dumbbell colored 1 and a straight line colored $i + 1$, thanks to (58).
- Relation involving *two adjacent colors and one distant from the other two*. If the distant color in (26) is 0, the proof is straightforward. Otherwise, we compute

$$\begin{aligned} \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram} \\ 1 \quad 0 \end{array} \right) &= \begin{array}{c} \text{Diagram} \\ 1 \end{array} \stackrel{(26), (58), (57)}{=} \begin{array}{c} \text{Diagram} \\ 1 \end{array} \\ &= \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram} \end{array} \right), \end{aligned}$$

and

$$\begin{aligned} \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram} \\ d-1 \quad 0 \end{array} \right) &= \begin{array}{c} \text{Diagram} \\ d-1 \quad 1 \quad d-1 \end{array} \stackrel{(26), (57), (73)}{=} \begin{array}{c} \text{Diagram} \\ d-1 \quad 1 \quad d-1 \end{array} = \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram} \end{array} \right). \end{aligned}$$

The relations with the adjacent colors exchanged are proved in the same way.


- Relation involving *three adjacent colors*. We need to check the cases of three adjacent colors belonging to $\{d - 2, d - 1, 0, 1, 2\}$. Starting with the case of $(0, 1, d - 1)$, we have

$$\mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram} \\ 1 \quad 0 \quad d-1 \end{array} \right) = \begin{array}{c} \text{Diagram} \end{array}$$

and

$$\mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram 1} \end{array} \right) = \text{Diagram 2}$$

To prove that these are equal, first use the relations in [Lemma 4.28](#) and [Lemma 4.29](#) to write them in the form



Then observe that the parts of the diagrams inside the dashed circle are exactly as the two sides of (27) with colors (1, 2, 3), which completes the proof of this case.

The remaining cases can be proved in similar ways, but they are actually a bit easier. For example, for the colors (0, 1, 2) we have

$$\mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram 3} \end{array} \right) = \text{Diagram 4}$$

and

$$\mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram with 4 strands (red, blue, purple, red) crossing in a grid pattern} \end{array} \right) = \begin{array}{c} \text{Diagram with 4 strands (red, blue, purple, red) with various crossings and arcs} \end{array}$$

Proceeding as in the previous case, but using the relations in [Lemma 4.21](#) and [Lemma 4.16](#), results in two diagrams which differ only by parts that are equal to the two sides of (27) with colors (1, 2, 3) again.

- Relations involving *oriented strands*. Relations (32) and (33) translate under $\mathcal{E}v_{r,s}$ into relations (51) and (52), respectively. The remaining relations (34) to (38) translate into relations (57), (58), (59), (62) and (63) (together with some obvious relations in the usual (non-oriented) Soergel calculus), respectively, if they don't involve the color 0.

However, if one of the strands is colored 0, then there is something to check. For each relation, we prove one case involving the colors 0 and 1 and one case involving the colors 0 and $d - 1$, the other cases being similar.

– For relation (34), we have

$$\mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram with strands colored 0, 1, and 1} \end{array} \right) = \begin{array}{c} \text{Diagram with strands colored 0, 1, and 1} \end{array} \stackrel{(58)}{=} \begin{array}{c} \text{Diagram with strands colored 0, 1, and 1} \end{array} = \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram with strands colored 0, 1, and 1} \end{array} \right)$$

and

$$\mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram with strands colored 0, 1, and } d-1 \end{array} \right) = \begin{array}{c} \text{Diagram with strands colored 0, 1, and } d-1 \end{array} \stackrel{(58),(73)}{=} \begin{array}{c} \text{Diagram with strands colored 0, 1, and } d-1 \end{array} = \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram with strands colored 0, 1, and } d-1 \end{array} \right)$$

– For relation (35), we have

$$\mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram with two strands colored 1} \end{array} \right) = \begin{array}{c} \text{Diagram with two strands colored 1} \end{array} \stackrel{(51)}{=} \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram with two strands colored 1} \end{array} \right)$$

and

$$\mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram 1} \\ 0 \end{array} \right) = \begin{array}{c} \text{Diagram 2} \\ 1 \end{array} \stackrel{(52)}{=} \begin{array}{c} \text{Diagram 3} \\ 1 \end{array} \stackrel{(65)}{=} \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram 4} \\ \text{Diagram 5} \end{array} \right)$$

– for relation (36), we have

$$\mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram 1} \\ 1 \end{array} \right) = \begin{array}{c} \text{Diagram 2} \\ \text{Diagram 3} \end{array} \stackrel{(52)}{=} \begin{array}{c} \text{Diagram 4} \\ \text{Diagram 5} \end{array} = \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram 6} \\ \text{Diagram 7} \end{array} \right)$$

and

$$\mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram 1} \\ 0 \end{array} \right) = \begin{array}{c} \text{Diagram 2} \\ 1 \end{array} \stackrel{(51),(66)}{=} \begin{array}{c} \text{Diagram 3} \\ \text{Diagram 4} \end{array} = \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram 5} \\ \text{Diagram 6} \end{array} \right)$$

– For relation (37), we have

$$\mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram 1} \\ 0 \end{array} \right) = \begin{array}{c} \text{Diagram 2} \\ 1 \end{array} = \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram 3} \\ \text{Diagram 4} \end{array} \right)$$

and

$$\mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram 1} \\ 0 \end{array} \right) = \begin{array}{c} \text{Diagram 2} \\ 1 \end{array} \stackrel{(78)}{=} \begin{array}{c} \text{Diagram 3} \\ 1 \end{array} = \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram 4} \\ \text{Diagram 5} \end{array} \right)$$

– Relation (38) actually consists of two (similar) relations. For the first of them, we have

$$\mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram 1} \\ 0 \\ 2 \quad 1 \end{array} \right) = \begin{array}{c} \text{Diagram 2} \\ \text{Diagram 3} \end{array} \stackrel{(58)}{=} \begin{array}{c} \text{Diagram 4} \\ \text{Diagram 5} \end{array} = \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram 6} \\ \text{Diagram 7} \end{array} \right)$$

and

$$\mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram 1} \\ 1 \quad 0 \\ d-1 \end{array} \right) = \begin{array}{c} \text{Diagram 2} \\ \text{Diagram 3} \end{array} \stackrel{(65)}{=} \begin{array}{c} \text{Diagram 4} \\ \text{Diagram 5} \end{array} \stackrel{(75)}{=} \begin{array}{c} \text{Diagram 6} \\ \text{Diagram 7} \end{array} = \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{Diagram 8} \\ \text{Diagram 9} \end{array} \right)$$

To check this relation with colors $(d-2, d-1, 0)$, use (58) and (65).

For the second relation in (38), we have

$$\mathcal{E}v_{r,s} \left(\begin{array}{c} 0 \\ \text{---} \\ \text{---} \\ \text{---} \\ 1 \quad 2 \end{array} \right) = \text{---} \stackrel{(51),(58)}{=} \text{---} = \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \right)$$

and

$$\mathcal{E}v_{r,s} \left(\begin{array}{c} d-1 \\ \text{---} \\ \text{---} \\ \text{---} \\ 0 \quad 1 \end{array} \right) = \text{---} \stackrel{(52),(65)}{=} \text{---} \stackrel{(75)}{=} \mathcal{E}v_{r,s} \left(\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \right)$$

Checking the relation with colors $(d-2, d-1, 0)$ uses (65), (58) and (52).

This ends the proof of Theorem 5.4. \square

Remark 5.5. (1) The functor $\mathcal{E}v_{r,s}$ is not full: For example, the special Rouquier complex $\mathcal{E}v_{r,s}(\mathbb{B}_\rho^{-1}) = \mathbb{T}_\rho^{-1}\langle -r \rangle[-s]$ has the form

$$\cdots \rightarrow \mathbb{B}_{d-1} \cdots \mathbb{B}_1 \langle -r \rangle[-s] \rightarrow 0.$$

Therefore, there is an obvious (non-null-homotopic) map from $\mathbb{B}_{d-1} \cdots \mathbb{B}_1 \langle -r \rangle[-s]$ to $\mathbb{T}_\rho^{-1}\langle -r \rangle[-s]$, which is the identity on $\mathbb{B}_{d-1} \cdots \mathbb{B}_1 \langle -r \rangle[-s]$ and zero elsewhere, but this map is not in the image of $\mathcal{E}v_{r,s}$.

Note also that $\mathcal{E}v_{r,s}$ induces a functor $\mathcal{E}v_a: \mathcal{K}^b(\widehat{\mathcal{S}}_d^{\text{ext}}) \rightarrow \mathcal{K}^b(\mathcal{S}_d)$ and that

$$\mathcal{K}^b(\widehat{\mathcal{S}}_d^{\text{ext}}) (\mathbb{T}_\rho^{-1}, \mathbb{B}_\rho^{-1}) = 0,$$

whereas

$$\mathcal{K}^b(\mathcal{S}_d) (\mathcal{E}v_{r,s}(\mathbb{T}_\rho^{-1}), \mathcal{E}v_{r,s}(\mathbb{B}_\rho^{-1})) = \mathcal{K}^b(\mathcal{S}_d) (\mathbb{T}_\rho^{-1}, \mathbb{T}_\rho^{-1}) \cong R',$$

where R' is the polynomial ring in the i -colored dumbbells with $i \in \{1, \dots, d\}$, i.e. the endomorphism ring of the identity object for finite type A_{d-1} .

(2) By (30) and (69), the evaluation functor $\mathcal{E}v_{r,s}$ maps the central morphism

$$\sum_{i=0}^{d-1} \text{---} i$$

to zero. We could have defined $\widehat{\mathcal{B}}\mathcal{S}_d^{\text{ext}}$ over the polynomial ring $\mathbb{C}[y, x_1, \dots, x_{d-1}]$ as in [MT2017] and extended $\mathcal{E}v_{r,s}$ to that "base ring". In that case, the central morphism \boxed{y} (which is equal to the above dumbbell sum, as already remarked) would be sent to zero by the evaluation functor, which makes perfect sense as the extended base ring of $\mathcal{B}\mathcal{S}_d$ would be $\mathbb{C}[x_1, \dots, x_d]$.

6. EVALUATION BIREPRESENTATIONS AND FINITARY COVERS

6.1. Recollections on birepresentation theory. In the following, we will work with graded (finitary or triangulated) birepresentations of graded, additive bicategories. The particular bicategory we are interested in is, of course, $\widehat{\mathcal{S}}_d^{\text{ext}}$, which we view as a bicategory with one object in the usual way.

We call a graded, \mathbb{C} -linear, additive category \mathcal{A} *graded-finitary* if \mathcal{A}^{sh} is idempotent complete, morphism spaces between indecomposables are finite-dimensional and there are only finitely many isomorphism classes of indecomposables up to isomorphism and grading shift. Note that \mathcal{A} need not be finitary, because the Hom-spaces might be infinite-dimensional, although they are finite-dimensional in each degree. This is why we write *graded-finitary* and not *graded, finitary*. We denote the 2-category of graded, resp. graded-finitary, \mathbb{C} -linear, additive categories, degree-preserving \mathbb{C} -linear functors and natural transformations by $\mathfrak{A}_{\mathbb{C}}^g$, resp. $\mathfrak{A}_{\mathbb{C}}^{g,f}$. A (locally) *graded, additive bicategory* \mathcal{C} is one whose morphism categories are enriched over $\mathfrak{A}_{\mathbb{C}}^g$ and a (locally) *graded-finitary bicategory* \mathcal{C} is one whose morphism categories are enriched over $\mathfrak{A}_{\mathbb{C}}^{g,f}$ and whose identity 1-morphisms are indecomposable. Note that, to shorten the string of adjectives, we drop the adjective \mathbb{C} -linear, even though it is implicit in the enrichment. A *graded, additive* (resp. *graded-finitary*) *birepresentation* is a degree-preserving pseudofunctor from \mathcal{C} to $\mathfrak{A}_{\mathbb{C}}^g$ (resp. $\mathfrak{A}_{\mathbb{C}}^{g,f}$).

Since we are mainly interested in $\widehat{\mathcal{S}}_d$, we will also abuse notation and call additive (bi)categories of the form \mathcal{A}^{sh} graded-finitary provided \mathcal{A} is. Similarly, given a graded-finitary birepresentation \mathbf{M} of a graded, additive bicategory \mathcal{C} , we will also call the birepresentation \mathbf{M}^{sh} of \mathcal{C}^{sh} (which acts on categories $\mathbf{M}(i)^{\text{sh}}$, for objects i , via functors which commute with shifts) graded-finitary. For more detail on these constructions, we refer to [MMMTZ2019, Section 2.6].

We will also be considering triangulated birepresentations of graded, additive bicategories. Denote by $\mathfrak{T}_{\mathbb{C}}$ the bicategory of triangulated, \mathbb{C} -linear categories, (\mathbb{C} -linear) triangle functors and natural transformations. A *triangulated birepresentation* of a \mathbb{C} -linear, additive bicategory \mathcal{C} is a (\mathbb{C} -linear) pseudofunctor from \mathcal{C} to $\mathfrak{T}_{\mathbb{C}}^{g,f}$. In order to consider graded versions, we restrict ourselves to the 2-full subcategory $\mathfrak{T}_{\mathbb{C}}^g$ of $\mathfrak{T}_{\mathbb{C}}$ whose objects are triangulated categories of the form $\mathcal{K}^b(\mathcal{A}^{\text{sh}})$ for a graded, \mathbb{C} -linear, additive category \mathcal{A} , and whose functors are degree-preserving triangle functors. A *graded-triangulated birepresentation* of an additive, graded bicategory \mathcal{C} is then a degree-preserving (\mathbb{C} -linear) pseudofunctor from \mathcal{C} to $\mathfrak{T}_{\mathbb{C}}^g$.

Similarly to the finitary case above, we will call a birepresentation via which a bicategory of the form \mathcal{C}^{sh} acts on triangulated categories of the form \mathcal{T}^{sh} via triangle functors commuting with shifts (i.e. one obtained by taking a graded birepresentation of \mathcal{C} acting on \mathcal{T} , closing under shifts, and then restricting to morphisms of degree zero) a graded, triangulated birepresentation.

In some cases, graded-finitary birepresentations will have an additional shift functor (coming from the homological shift in a triangulated birepresentation), with respect to which morphisms in the underlying categories will have degree zero. We call such birepresentations bigraded-finitary.

Given a (locally) additive, graded bicategory, the set of isomorphism classes of indecomposable 1-morphisms up to grading shift can be given three natural partial preorders: the *left* preorder ($[F] \leq_L [G]$ if and only if $[G]$ appears as a direct summand of $[HF]$ for some 1-morphism H),

the *right* preorder ($[F] \leq_R [G]$ if and only if $[G]$ appears as a direct summand of $[FH]$ for some 1-morphism H) and the *two-sided* preorder ($[F] \leq_J [G]$ if and only if $[G]$ appears as a direct summand of $[H_1FH_2]$ for some 1-morphisms H_1, H_2), and the corresponding equivalence classes are called *left*, *right* and *two-sided cells*, respectively.

If \mathcal{C} is graded-finitary, we can associate to any left cell a so-called *graded cell 2-representation*, which is the quotient of the left 2-ideal in \mathcal{C} generated by the identities on the 1-morphisms in the cell, by the unique maximal ideal of the resulting birepresentation (i.e. the unique maximal ideal of the underlying categories which is stable under the action of \mathcal{C}). For more details (in the ungraded case, but the graded one is analogous), see e.g. [MM2016, Section 3.3].

6.2. Finitary covers of evaluation cell birepresentations. Let \mathbf{M} be a graded-finitary birepresentation of \mathcal{S}_d , for any $d \in \mathbb{N}_{\geq 2}$. Then $\mathcal{K}^b(\mathbf{M})$, as a graded, triangulated birepresentation of $\mathcal{K}^b(\mathcal{S}_d)$, induces a graded, triangulated birepresentation of $\widehat{\mathcal{S}}_d^{\text{ext}}$, the *evaluation birepresentation* $\mathbf{M}^{\mathcal{E}v_{r,s}}$, resp. $\mathbf{M}^{\mathcal{E}v'_{r,s}}$, by pull-back through the evaluation functors $\mathcal{E}v_{r,s}$, resp. $\mathcal{E}v'_{r,s}$, for any $r, s \in \mathbb{Z}$.

In this subsection we show that, if \mathbf{M} is a cell birepresentation of \mathcal{S}_d , then $\mathbf{M}^{\mathcal{E}v_{r,s}}$ has a bigraded-finitary cover in the following sense.

Definition 6.1. A *bigraded-finitary cover* of a graded, triangulated birepresentation \mathbf{N} of a graded, additive bicategory \mathcal{C} is a bigraded-finitary birepresentation \mathbf{L} of \mathcal{C} together with an epimorphic, faithful and essentially surjective morphism $\Phi: \mathcal{K}^b(\mathbf{L}) \rightarrow \mathcal{K}^b(\mathbf{N})$ of graded, triangulated birepresentations. Here *epimorphic* means that every morphism in $\mathcal{K}^b(\mathbf{N})$ is a composite of morphisms in the image of Φ , possibly with additional isomorphisms.

Proposition 6.2. *Let \mathbf{M} be the graded cell birepresentation associated to some left cell \mathcal{L} of \mathcal{S}_d . Then $\mathbf{M}^{\mathcal{E}v_{r,s}}$ has a bigraded-finitary cover.*

Proof. By [EH2018, Proposition 4.31], T_ρ^d acts as $\text{Id}\langle x \rangle[y]$ on $\mathbf{M}^{\mathcal{E}v_{r,s}}$, for some $x, y \in \mathbb{Z}$. Let \mathbf{L} be the closure under isomorphisms, direct sums, direct summands, grading and homological shifts of the $\mathcal{E}v_{r,s}(T_\rho^i)B_w$, for $0 \leq i \leq d-1$ and $w \in \mathcal{L}$. Relation (35) implies that \mathbf{L} is a bigraded-finitary birepresentation of $\widehat{\mathcal{S}}_d^{\text{ext}}$.

The inclusion functor $\mathbf{L} \hookrightarrow \mathbf{M}^{\mathcal{E}v_{r,s}}$ extends to a morphism of graded triangulated birepresentations $\Phi: \mathcal{K}^b(\mathbf{L}) \hookrightarrow \mathcal{K}^b(\mathbf{M}^{\mathcal{E}v_{r,s}})$ which is epimorphic, faithful and essentially surjective by construction. \square

We refer to Corollary 6.5 for an example demonstrating that Φ is not full in general.

Remark 6.3. It is easy to see that \mathbf{L} is transitive, and it looks likely that calculations, using the explicit descriptions of the representing bimodules for the B_w given in [MMMTZ2019, Section 4.3], one can verify that it is indeed simple transitive.

6.3. The zigzag algebras. Let us first recall the *affine zigzag algebra* \widehat{Z}_d over \mathbb{C} associated to the \widehat{A}_{d-1} Dynkin diagram. As is well-known, there are two isomorphism classes of affine zigzag algebras with invertible integer coefficients, and we use a specific representative of either one or the other depending on the parity of d .

Let e_0, \dots, e_{d-1} denote the orthogonal idempotents associated to the vertices of the zigzag quiver and $i_1|i_2| \dots |i_k$ the path in the quiver from i_k to i_1 via i_{k-1}, \dots, i_2 . As usual, all indices are to be taken modulo d . The relations in \widehat{Z}_d are

$$\begin{aligned} i|i+1|i+2 = 0 = i|i-1|i-2, \quad i = 0, \dots, d-1; \\ i|i+1|i = i|i-1|i, \quad i = 1, \dots, d-1; \\ 0|1|0 = (-1)^d(0|d-1|0). \end{aligned}$$

For convenience, we also use the notation

$$\ell_i := i|i+1|i,$$

for any $i = 0, \dots, d-1$. This algebra has dimension $4d$, it is positively graded by putting the degree of every path equal to its length, and it is a graded Frobenius algebra with non-degenerate trace defined by

$$\mathrm{tr}(\ell_i) = 1 \text{ for every } i = 0, \dots, d-1; \quad \mathrm{tr}(a) = 0 \text{ when } \deg(a) \neq 2.$$

This means that $\widehat{Z}_d^* \cong \widehat{Z}_d\langle 2 \rangle$ as graded left, resp. right, \widehat{Z}_d -modules. Define the non-degenerate bilinear pairing $\langle \cdot, \cdot \rangle: \widehat{Z}_d \otimes \widehat{Z}_d \rightarrow \mathbb{C}$ as usual

$$\langle a, b \rangle := \mathrm{tr}(ab), \quad a, b \in \widehat{Z}_d,$$

and recall that two bases of \widehat{Z}_d , say $\{a_i \mid i = 1, \dots, 4d\}$ and $\{a_i^* \mid 1, \dots, 4d\}$, are called *dual* to each other if they satisfy

$$\langle a_i, a_j^* \rangle = \delta_{i,j}, \quad i, j = 1, \dots, 4d,$$

where $\delta_{i,j}$ is the Kronecker delta. With respect to the bilinear form on \widehat{Z}_d , there is a natural pair of dual bases $\{e_i, \ell_i, i|i \pm 1 \mid i = 0, \dots, d-1\}$ and $\{e_i^*, \ell_i^*, (i|i \pm 1)^* \mid i = 0, \dots, d-1\}$, such that

$$e_i^* = \ell_i; \quad \ell_i^* = e_i; \quad (0|(d-1))^* = (-1)^d((d-1)|0); \quad (i|(i \pm 1))^* = (i \pm 1|i), \quad i = 1, \dots, d-1.$$

for $i = 0, \dots, d-1$. Note that \widehat{Z}_d is symmetric when d is even and only weakly symmetric when d is odd.

Let $\widehat{Z}_d\text{-fgproj}$, resp. $\mathrm{fgproj}\text{-}\widehat{Z}_d$, be the category of finite-dimensional, graded, projective left, resp. right, \widehat{Z}_d -modules and degree-preserving module maps. The indecomposable objects in these categories are isomorphic to $\widehat{Z}_d e_i \langle t \rangle$, resp. $e_i \widehat{Z}_d \langle t \rangle$, for some $i = 0, \dots, d-1$ and $t \in \mathbb{Z}$.

Finally, let $\widehat{Z}_d\text{-fgbiproj}\text{-}\widehat{Z}_d$ be the monoidal category of all finite-dimensional, graded, biprojective $\widehat{Z}_d\text{-}\widehat{Z}_d$ -bimodules and degree-preserving bimodule maps. A bimodule is called biprojective if it is projective as a graded left module and as a graded right module, but not necessarily as a graded bimodule. Every indecomposable projective object in this category is isomorphic to

$$\widehat{Z}_d e_i \otimes e_j \widehat{Z}_d \langle t \rangle,$$

for some $i, j = 0, \dots, d-1$ and $t \in \mathbb{Z}$. The monoidal structure of $\widehat{Z}_d\text{-fgbiproj}\text{-}\widehat{Z}_d$ is given by tensoring over \widehat{Z}_d and the unit object is \widehat{Z}_d , which is biprojective but not projective as a bimodule over itself. Recall that any exact, graded endofunctor of $\widehat{Z}_d\text{-fgproj}$ is naturally isomorphic to

$B \otimes_{\widehat{Z}_d} -$, for some $B \in \widehat{Z}_d\text{-fgbiproj-}\widehat{Z}_d$. Natural transformations between exact, graded endofunctors correspond to $\widehat{Z}_d\text{-}\widehat{Z}_d$ -bimodule maps and the composition of endofunctors corresponds to the tensor product of the corresponding bimodules over \widehat{Z}_d .

Let τ be the degree-preserving algebra automorphism of \widehat{Z}_d induced by the counterclockwise rotation of the Dynkin diagram defined by

$$e_i \mapsto e_{i+1}, \quad 0|(d-1) \mapsto (-1)^d(1|0), \quad i|j \mapsto (i+1)|(j+1),$$

for $i, j = 0, \dots, d-1$, such that $j = i \pm 1$ but $(i, j) \neq (0, d-1)$. Note that $\tau^d = \text{id}$ when d is even, and $(\tau)^{2d} = \text{id}$ when d is odd. By definition, the *twisted bimodule*

$$\widehat{Z}_d^\tau \in \widehat{Z}_d\text{-fgbiproj-}\widehat{Z}_d$$

has underlying vector space \widehat{Z}_d , while the left and right \widehat{Z}_d -actions are defined by

$$a \cdot_L b \cdot_R c := ab\tau(c),$$

for $a, b, c \in \widehat{Z}_d$. It is clear that $\widehat{Z}_d^\tau \cong \widehat{Z}_d$ as left and as right \widehat{Z}_d -modules, but not as $\widehat{Z}_d\text{-}\widehat{Z}_d$ -bimodules. In other words, \widehat{Z}_d^τ is biprojective, but not projective as a $\widehat{Z}_d\text{-}\widehat{Z}_d$ -bimodule. We record the existence of an isomorphism

$$(89) \quad \widehat{Z}_d^{\tau^k} \otimes_{\widehat{Z}_d} \widehat{Z}_d^{\tau^m} \cong \widehat{Z}_d^{\tau^{k+m}}$$

in $\widehat{Z}_d\text{-fgbiproj-}\widehat{Z}_d$, for every pair $k, m \in \mathbb{Z}$.

Note further that there exist isomorphisms of left, resp. right, \widehat{Z}_d -modules

$$(90) \quad \widehat{Z}_d^\tau \otimes_{\widehat{Z}_d} \widehat{Z}_d e_i \cong \widehat{Z}_d e_{i+1} \quad \text{and} \quad e_i \widehat{Z}_d \otimes_{\widehat{Z}_d} \widehat{Z}_d^\tau \cong e_{i-1} \widehat{Z}_d$$

and, therefore, an isomorphism of $\widehat{Z}_d\text{-}\widehat{Z}_d$ -bimodules

$$(91) \quad \widehat{Z}_d^\tau \otimes_{\widehat{Z}_d} \widehat{Z}_d e_i \otimes e_i \widehat{Z}_d \cong \widehat{Z}_d e_{i+1} \otimes e_{i+1} \widehat{Z}_d \otimes_{\widehat{Z}_d} \widehat{Z}_d^\tau$$

for every $i = 0, \dots, d-1$.

The *zigzag algebra* Z_d of finite type A_{d-1} is by definition the idempotent subalgebra

$$(e_1 + \dots + e_{d-1}) \widehat{Z}_d (e_1 + \dots + e_{d-1}).$$

6.4. The birepresentations. Let $Z = Z_d$ denote the zigzag algebra of finite type A_{d-1} for some fixed $d \geq 3$. Recall the finitary birepresentation \mathbf{M}_d of \mathcal{S}_d acting on $Z\text{-gproj}$, the finitary category of finite-dimensional, graded projective Z -modules, by graded, biprojective $Z\text{-}Z$ -bimodules. Under this birepresentation, $\mathbb{1} = R$ acts by tensoring (over Z) with Z and each B_i acts by tensoring (over Z) with $Z e_i \otimes e_i Z \langle 1 \rangle$, for $i = 1, \dots, d-1$. The image of the generating

Soergel diagrams is given by

$$\begin{aligned}
\mathbf{M}_d \left(\begin{array}{c} \bullet \\ | \\ i \end{array} \right) & : & Z e_i \otimes e_i Z \langle 1 \rangle & \rightarrow Z \\
& & a e_i \otimes e_i b & \mapsto a e_i b, \\
\mathbf{M}_d \left(\begin{array}{c} i \\ | \\ \bullet \end{array} \right) & : & Z & \rightarrow Z e_i \otimes e_i Z \langle 1 \rangle \\
& & e_j & \mapsto \begin{cases} (-1)^i (\ell_i \otimes e_i + e_i \otimes \ell_i), & j = i; \\ (-1)^i (j|i \otimes i|j), & j \pm 1 = i, \end{cases} \\
\mathbf{M}_d \left(\begin{array}{c} i \\ / \quad \backslash \\ i \quad i \end{array} \right) & : & Z e_i \otimes e_i Z e_i \otimes e_i Z \langle 2 \rangle & \rightarrow Z e_i \otimes e_i Z \langle 1 \rangle \\
& & e_i \otimes e_i a e_i \otimes e_i & \mapsto (-1)^i \text{tr}(e_i a e_i) e_i \otimes e_i \\
\mathbf{M}_d \left(\begin{array}{c} i \quad i \\ \backslash \quad / \\ i \end{array} \right) & : & Z e_i \otimes e_i Z \langle 1 \rangle & \rightarrow Z e_i \otimes e_i Z e_i \otimes e_i Z \langle 2 \rangle \\
& & e_i \otimes e_i & \mapsto e_i \otimes e_i \otimes e_i,
\end{aligned}$$

while all other generating Soergel diagrams are sent to zero. The proof that this is well-defined is a straightforward computation and similar to the proof of [MT2019, Theorem I]. It is easy to see that this birepresentation decategorifies to the representation M_d of H_d , given in (10).

Now, consider the triangulated birepresentations $\mathbf{M}^{\mathcal{E}v_{r,s}}$ and $\mathbf{M}^{\mathcal{E}v'_{-r,-s}}$ of $\widehat{\mathfrak{S}}_d^{\text{ext}}$, for $r, s \in \mathbb{Z}$, obtained by pulling $\mathcal{K}^b(\mathbf{M})$ back through the evaluation functors $\mathcal{E}v_{r,s}$ and $\mathcal{E}v'_{-r,-s}$. These decategorify to M^{ev_a} and $M^{\text{ev}_{a-1}}$ defined in (11) and (12), respectively, where $a = (-1)^s q^r$. The case $(r, s) = (d-2, 2-d)$ is somewhat special, as it corresponds to the so-called *Tate twist*, but the general case can easily be derived from this one by shifting the bigrading in all arguments below. To keep the notation simple, we therefore consider $\mathbf{M}^{\mathcal{E}v_{r,s}}$ for the fixed choice $(r, s) = (d-2, 2-d)$ first.

Define the complex

$$X_0 := \underline{Z e_{d-1} \langle 1 \rangle} \rightarrow Z e_{d-2} \langle 2 \rangle \rightarrow \cdots \rightarrow Z e_1 \langle d \rangle$$

where the term $Z e_{d-1} \langle 1 \rangle$ is in homological degree 0 and the differential in position i is given by right multiplication by $d-i-1|d-i-2$. We further set $X_i := Z e_i$, for $i = 1, \dots, d-1$.

In Proposition 6.2, the rank of the bigraded-finitary cover \mathbf{L} of an evaluation cell birepresentation is not necessarily minimal. In the following proposition, we give a minimal finitary cover for $\mathbf{M}^{\mathcal{E}v_{r,s}}$.

Proposition 6.4. *The bigraded-finitary subcategory*

$$\widehat{\mathbf{M}}_{d-2,2-d} := \text{add} \{ (X_0 \oplus X_1 \oplus \cdots \oplus X_{d-1}) \langle i \rangle [j] \mid i, j \in \mathbb{Z} \}$$

is stable under the action of $\widehat{\mathcal{S}}_d^{\text{ext}}$, and hence carries the structure of a finitary birepresentation of $\widehat{\mathcal{S}}_d^{\text{ext}}$, which we denote by the same symbol.

Proof. We need to check stability under $B_1 \dots, B_{d-1}$ and T_ρ . The action of $B_1 \dots, B_{d-1}$ stabilises $\text{add} \{X_1 \oplus \dots \oplus X_{d-1} \langle i \rangle [j] \mid i, j \in \mathbb{Z}\}$ since this is just the finitary birepresentation of \mathcal{S}_d described above. We therefore first compute $B_i(X_0)$ for $i \in \{1, \dots, d-1\}$ and then verify stability of $\text{add} \{X_1 \oplus \dots \oplus X_{d-1} \langle i \rangle [j] \mid i, j \in \mathbb{Z}\}$ under T_ρ .

Notice that, for $i \in \{2, \dots, d-2\}$, $B_i(X_0)$ is given by the complex

$$Ze_i \otimes (e_i Ze_{i+1} \langle d-i \rangle \rightarrow e_i Ze_i \langle d-i+1 \rangle \rightarrow e_i Ze_{i-1} \langle d-i+2 \rangle)$$

and the complex of vector spaces in the right tensor factor is null-homotopic, since the first map embeds a one-dimensional space into a two-dimensional space, and the second map is a surjection onto another one-dimensional space. Hence the whole complex is null-homotopic.

Further, $B_1(X_0)$ is given by

$$Ze_1 \otimes (e_1 Ze_2 \langle d-1 \rangle \rightarrow e_1 Ze_1 \langle d \rangle)$$

with map $1|2 \mapsto \ell_1$, which is injective, hence the summand surviving Gaussian elimination is $Ze_1 \otimes e_1 \langle d \rangle$ in homological degree $d-2$. Thus the result is homotopy equivalent to $Ze_1 \langle d \rangle [2-d]$.

On the other extreme, $B_{d-1}(X_0)$ is given by

$$Ze_{d-1} \otimes (\underline{e_{d-1} Ze_{d-1} \langle 2 \rangle} \rightarrow e_{d-1} Ze_{d-2} \langle 3 \rangle)$$

where the map is right multiplication by $d-1|d-2$, which is surjective. The kernel is thus $Ze_{d-1} \otimes \ell_{d-1} \langle 2 \rangle$ and the result is homotopy equivalent to Ze_{d-1} without any shifts.


Thus $\text{add} \{X_1 \oplus \dots \oplus X_{d-1} \langle i \rangle [j] \mid i, j \in \mathbb{Z}\}$ is stable under the action of B_1, \dots, B_{d-1} .

It remains to show that $\widehat{\mathbf{M}}$ is stable under the action of T_ρ . Recall from Section 5.1 that $\mathcal{E}v_{d-2,2-d}(T_\rho) = T_1^{-1} \dots T_{d-1}^{-1} \langle d-2 \rangle [2-d]$ and

$$T_i^{-1} = R \langle -1 \rangle \xrightarrow{\quad \downarrow \quad} \underline{B}_i.$$

Using the definition of \mathbf{M}_d above, it is easy to see that the complex representing $\mathcal{E}v_{d-2,2-d}(T_\rho)$ is

(92)



Here $d^{-1} = (-d_1^{-1}, d_2^{-1}, \dots, (-1)^{d-1} d_{d-1}^{-1})$, where each $d_i^{-1}: Z\langle -1 \rangle \rightarrow Ze_i \otimes e_i Z\langle 1 \rangle$ is given by

$$e_j \mapsto \begin{cases} \ell_i \otimes e_i + e_i \otimes \ell_i, & \text{if } i = j; \\ j|i \otimes i|j, & \text{if } i \neq j. \end{cases}$$

The other differentials are all vectors of Z - Z -bimodule maps which are equal to the tensor product of $\pm \text{id}$ on one tensor factor and $i|i+1$, for some $i = 1, \dots, d-2$, on the other tensor factor. For our arguments below, the signs of these maps are not important.

We are first going to prove that $\mathcal{E}v_{d-2,2-d}(\mathbb{T}_\rho)(X_i) \simeq X_{i+1}$, for any $i = 1, \dots, d-2$. Since $e_j Z e_i = \{0\}$ when $|i-j| > 1$, the non-zero part of the complex corresponding to $\mathcal{E}v_{d-2,2-d}(\mathbb{T}_\rho)(X_i)$ is

$$\begin{array}{ccccccc} & & \underline{Ze_{i-1} \otimes e_{i-1} Ze_i \langle 1 \rangle} & \longrightarrow & Ze_{i-2} \otimes e_{i-1} Ze_i \langle 2 \rangle & \longrightarrow & \dots & Ze_1 \otimes e_{i-1} Ze_i \langle i-1 \rangle \\ & \nearrow & & \searrow & & & & \searrow \\ Ze_i \langle -1 \rangle & \longrightarrow & \underline{Ze_i \otimes e_i Ze_i \langle 1 \rangle} & \longrightarrow & Ze_{i-1} \otimes e_i Ze_i \langle 2 \rangle & \longrightarrow & \dots & Ze_2 \otimes e_i Ze_i \langle i-1 \rangle & \longrightarrow & Ze_1 \otimes e_i Ze_i \langle i \rangle \\ & \searrow & & \nearrow & & & & \nearrow & & \searrow \\ & & \underline{Ze_{i+1} \otimes e_{i+1} Ze_i \langle 1 \rangle} & \longrightarrow & Ze_i \otimes e_{i+1} Ze_i \langle 2 \rangle & \longrightarrow & \dots & Ze_3 \otimes e_{i+1} Ze_i \langle i-1 \rangle & \longrightarrow & Ze_2 \otimes e_{i+1} Ze_i \langle i \rangle & \longrightarrow & Ze_1 \otimes e_{i+1} Ze_i \langle i+1 \rangle \end{array}$$

By Gaussian elimination, one can then see that this is homotopy equivalent to the purple $Ze_{i+1} \otimes e_{i+1} Ze_i \langle 1 \rangle$ in homological degree zero, which is isomorphic to X_{i+1} . To explain this, we identify each vertex of the diagram above by its pair of coordinates (row number, column number), where we number the rows of the complex by 1,2,3 from top to bottom and the columns by their homological degree. As in the diagram above, we omit the signs of all maps below, since they are not important for our argument. Using these conventions, first note that the the part of the complex $(2, -1) \rightarrow (2, 0) \rightarrow (3, 1)$ is given by

$$Ze_i \otimes (\mathbb{k} \langle -1 \rangle \rightarrow \underline{e_i Ze_i \langle 1 \rangle} \rightarrow e_{i+1} Ze_i \langle 2 \rangle)$$

where the complex of vector spaces is split by the same arguments as above and hence null-homotopic. Thus these three terms cancel in the Gaussian elimination procedure. Similarly, every part of the complex of the form $(1, j) \rightarrow (2, j + 1) \rightarrow (3, j + 2)$, for $j = 0, \dots, i - 2$, is given by

$$Ze_{i-j-1} \otimes (e_{i-1}Ze_i\langle j + 1 \rangle \rightarrow e_iZe_i\langle j + 2 \rangle \rightarrow e_{i+1}Ze_i\langle j + 3 \rangle)$$

is split and hence null-homotopic. Hence all these triples of terms cancel in the Gaussian elimination procedure, which in the end only leaves the purple one, proving the desired homotopy equivalence.

The next homotopy equivalence we are going to prove is $\mathcal{E}v_{d-2,2-d}(\mathbb{T}_\rho)(X_{d-1}) \simeq X_0$. The non-zero part of the complex $\mathcal{E}v_{d-2,2-d}(\mathbb{T}_\rho)(X_{d-1})$ is

$$\begin{array}{ccccccc}
 & \underline{Ze_{d-2} \otimes e_{d-2}Ze_{d-1}\langle 1 \rangle} & \longrightarrow & Ze_{d-3} \otimes e_{d-2}Ze_{d-1}\langle 2 \rangle & \longrightarrow & \dots & \longrightarrow & Ze_{e_1} \otimes e_{d-2}Ze_{d-1}\langle d-1 \rangle \\
 & \nearrow & & \searrow & & & & \searrow \\
 Ze_{d-1}\langle -1 \rangle & & & & & & & \\
 & \searrow & & \nearrow & & & & \nearrow \\
 & \underline{Ze_{d-1} \otimes e_{d-1}Ze_{d-1}\langle 1 \rangle} & \longrightarrow & Ze_{d-2} \otimes e_{d-1}Ze_{d-1}\langle 2 \rangle & \longrightarrow & \dots & \longrightarrow & Ze_{e_2} \otimes e_{d-1}Ze_{d-1}\langle d-1 \rangle \\
 & & & & & & & \nearrow \\
 & & & & & & & Ze_{e_1} \otimes e_{d-1}Ze_{d-1}\langle d \rangle
 \end{array}$$

The differentials are as above and by Gaussian elimination this complex is homotopy equivalent to the direct summand of the purple subcomplex for which the right tensor factor is restricted to multiples of e_{d-1} . This direct summand is indeed isomorphic to X_0 . (row number, column number). Note that again all descending maps in the complex are given by the tensor product the identity of some Ze_{d-j} with an injective map of vector spaces hence split. This implies that all black terms in the complex are killed and only one direct summand of each purple term (the one given by $Ze_{d-1-j} \otimes e_{d-1}\langle j + 1 \rangle$) survives in the Gaussian elimination procedure. Thus the complex is homotopy equivalent to X_0 , as claimed..

The remaining case of the action of $\mathcal{E}v_{d-2,2-d}(\mathbb{T}_\rho)$ on X_0 can be replaced by considering the action of $\mathcal{E}v_{d-2,2-d}(\mathbb{T}_\rho)^{-1}$ on X_1 , which is analogous to the action of $\mathcal{E}v_{d-2,2-d}(\mathbb{T}_\rho)$ on X_{d-2} . \square

Similarly, we can define an additive birepresentation $\widehat{\mathbf{M}}_{r,s}$ of $\widehat{\mathcal{S}}_d^{\text{ext}}$, for any $r, s \in \mathbb{Z}$.

Corollary 6.5. *For any $r, s \in \mathbb{Z}$, there is a morphism of additive $\widehat{\mathcal{S}}_d^{\text{ext}}$ -birepresentations $\Phi: \widehat{\mathbf{M}}_{r,s} \rightarrow \mathbf{M}^{\mathcal{E}v_{r,s}}$, induced by the embedding from Proposition 6.4 with a suitable bigrading shift. Moreover, Φ extends to a morphism of triangulated $\widehat{\mathcal{S}}_d^{\text{ext}}$ -birepresentations $\Phi: \mathcal{K}^b(\widehat{\mathbf{M}}_{r,s}) \rightarrow \mathbf{M}^{\mathcal{E}v_{r,s}}$, which is essentially surjective and faithful, but not full.*

Proof. All assertions follow immediately from Proposition 6.4, except the lack of fullness. Without loss of generality, assume that $(r, s) = (d - 2, 2 - d)$ again. Note that in $\widehat{\mathbf{M}}$ the only non-radical morphisms between X_0 and X_i , for $i = 1, \dots, d - 1$, are from X_0 to X_{d-1} and from $X_1[2 - d]$ to X_0 . This implies that $X_0 \not\cong \underline{(X_{d-1}\langle 1 \rangle \rightarrow X_{d-2}\langle 2 \rangle \rightarrow \dots \rightarrow X_1\langle d - 1 \rangle)}$ in $\mathcal{K}^b(\widehat{\mathbf{M}})$. \square


Remark 6.6. Note that $\widehat{\mathbf{M}}_{r,s}$ decategorifies to the Graham–Lehrer cell module $\widehat{M}_{d,\lambda}$ with $\lambda = (-1)^{s-(2-d)}q^{r-(d-2)}$, as can be easily seen by comparing the action of the generators on the X_i

with the decategorified action in (6) and (7). Moreover, Φ decategorifies to the projection of $\widehat{M}_{r,s}$ onto $L_{d,(-1)^{s-(2-d)}q^{r-(d-2)}}^+$.

Proposition 6.7. *For any $r, s \in \mathbb{Z}$, there is an isomorphism of ungraded algebras*

$$\text{End}_{\mathbf{M}^{\varepsilon_{vr,s}}}(X_0 \oplus \cdots \oplus X_{d-1}) \cong \widehat{Z}.$$

Proof. Without loss of generality, we assume that $(r, s) = (d-2, 2-d)$, as before. Denote by $p_{d-1}: X_0 \rightarrow X_{d-1}$ the projection onto the component in homological degree 0 and by $j_d: X_{d-1} \rightarrow X_0$ the map induced by multiplication with ℓ_{d-1} . Similarly, denote by $j_1: X_1[2-d] \rightarrow X_0$ the inclusion of the component in homological degree $d-2$ and by $p_1: X_0 \rightarrow X_1[2-d]$ the map induced by multiplication with ℓ_1 . We remark that $p_{d-1}, j_{d-1}, j_1, p_1$ have degrees $1, 1, 1-d, d+1$, respectively. Moreover, we denote the maps $Ze_i \rightarrow Ze_{i\pm 1}$ given by right multiplication by $i|i \pm 1$ by $r_{i|i\pm 1}$. Then it is a straightforward calculation to verify that $\text{End}_{\mathbf{M}^{\varepsilon_{va}}}(X_0 \oplus \cdots \oplus X_{d-1})$ is given by the path algebra of the quiver



modulo the relations defining \widehat{Z} under the isomorphism sending $r_{i|i\pm 1}$ to $i \pm 1|i$, p_i to $i|0$ and j_i to $0|i$ for $i \in \{1, d-1\}$. To verify the sign in the relation involving 0 we observe that the endomorphism of X_0 given by $j_1 p_1 + (-1)^{d-1} j_{d-1} p_{d-1}$ is (omitting shifts for readability) given by the solid arrows in the diagram

$$\begin{array}{ccccccc} Ze_{d-1} & \longrightarrow & Ze_{d-2} & \longrightarrow & \cdots & \longrightarrow & Ze_2 & \longrightarrow & Ze_1 \\ \ell_{d-1} \downarrow & & \swarrow r_{d-2|d-1} & \downarrow 0 & & \swarrow r_{2|3} & \downarrow 0 & \swarrow r_{1|2} & \downarrow \ell_1 \\ Ze_{d-1} & \longrightarrow & Ze_{d-2} & \longrightarrow & \cdots & \longrightarrow & Ze_2 & \longrightarrow & Ze_1 \end{array}$$

and is null-homotopic via the homotopy indicated by the dashed arrows. \square

Remark 6.8. The natural bigrading of $\text{End}_{\mathbf{M}^{\varepsilon_{vr,s}}}(X_0 \oplus \cdots \oplus X_{d-1})$ induces a bigrading on \widehat{Z} via the isomorphism in Proposition 6.7. The first entry of this bigrading is compatible with the above grading of \widehat{Z} except for the degrees of the arrows between 0 and 1.

The explicit 2-action of $\widehat{S}_d^{\text{ext}}$ on $\widehat{M}_{r,s}$ is given

- on 1-morphisms by

$$\begin{aligned} F(i) &:= \widehat{Z}_d e_i \otimes e_i \widehat{Z}_d \langle 1 \rangle, \quad i = 0, \dots, n; \\ F(\pm) &:= \widehat{Z}_d^{r \pm 1} \langle r \rangle [s], \end{aligned}$$

- on 2-morphisms by

$$\begin{aligned}
 F \left(\begin{array}{c} \bullet \\ | \\ i \end{array} \right) &: \widehat{Z}_d e_i \otimes e_i \widehat{Z}_d \langle 1 \rangle \rightarrow \widehat{Z}_d \\
 &ae_i \otimes e_i b \mapsto ae_i b, \\
 F \left(\begin{array}{c} i \\ | \\ \bullet \end{array} \right) &: \widehat{Z}_d \rightarrow \widehat{Z}_d e_i \otimes e_i \widehat{Z}_d \langle 1 \rangle \\
 e_j &\mapsto \begin{cases} (-1)^i (\ell_i \otimes e_i + e_i \otimes \ell_i), & j = i; \\ (-1)^i (j|i \otimes i|j), & j \pm 1 = i \neq 0; \\ 1|0 \otimes 0|1, & j = 1, i = 0; \\ (-1)^d (d-1|0 \otimes 0|d-1), & j = d-1, i = 0, \end{cases} \\
 F \left(\begin{array}{c} i \\ / \quad \backslash \\ i \quad i \end{array} \right) &: \widehat{Z}_d e_i \otimes e_i \widehat{Z}_d e_i \otimes e_i \widehat{Z}_d \langle 2 \rangle \rightarrow \widehat{Z}_d e_i \otimes e_i \widehat{Z}_d \langle 1 \rangle \\
 &e_i \otimes e_i a e_i \otimes e_i \mapsto (-1)^i \text{tr}(e_i a e_i) e_i \otimes e_i \\
 F \left(\begin{array}{c} i \quad i \\ \backslash \quad / \\ \quad i \end{array} \right) &: \widehat{Z}_d e_i \otimes e_i \widehat{Z}_d \langle 1 \rangle \rightarrow \widehat{Z}_d e_i \otimes e_i \widehat{Z}_d e_i \otimes e_i \widehat{Z}_d \langle 2 \rangle \\
 &e_i \otimes e_i \mapsto e_i \otimes e_i \otimes e_i.
 \end{aligned}$$

The generating 2-morphisms involving an oriented black strand in (30) and (31) are sent to the isomorphisms in (89) and (91), respectively, and all other generating 2-morphisms are sent to zero.

Remark 6.9. We could alternatively have used the evaluation functor $\mathcal{E}v'_{-r,-s}$ to obtain another evaluation birepresentation and its finitary cover.

REFERENCES

- [CP1996] V. Chari and A. Pressley, Quantum affine algebras and affine Hecke algebras. *Pacific J. Math.* 174, 295-326 (1996).
- [DF2016] J. Du and Q. Fu, Small representations for affine q -Schur algebras. *Algebr. Represent. Theory* 19 (2016), no. 2, 355-376.
- [E2018] B. Elias, Gaitsgory's central sheaves via the diagrammatic Hecke category (2018). [arXiv: 1811.06188v1](https://arxiv.org/abs/1811.06188v1).
- [EH2018] B. Elias and M. Hogancamp, Categorical diagonalization of full twists. [arXiv: 1801.00191v1](https://arxiv.org/abs/1801.00191v1)
- [EKh2010] B. Elias and M. Khovanov, Diagrammatics for Soergel categories. *Int. J. Math. and Math. Sci.* 2010.
- [EKr2010] B. Elias and D. Krasner, Rouquier complexes are functorial over braid cobordisms. *Homology Homotopy Appl.* 12 (2010), no. 2, 109-146.

- [EMTW2020] B. Elias, S. Makisumi, U. Thiel, G. Williamson, *Introduction to Soergel bimodules* RSME Springer Series, volume 5, Springer, 2020.
- [EW2014] B. Elias and G. Williamson, Hodge theory for Soergel bimodules. *Ann. of Math. (2)* 180 (2014), no. 3, 1089–1136.
- [EW2016] B. Elias and G. Williamson, Soergel calculus. *Represent. Theory* 20 (2016), 295–374.
- [EW2017] B. Elias and G. Williamson, Diagrammatics for Coxeter groups and their braid groups. *Quantum Topol.* 8 (2017), no. 3, 413–457.
- [GL1998] J. J. Graham and G. I. Lehrer, The representation theory of affine Temperley-Lieb algebras. *L'Enseignement Mathématique* 44 (1998), 173–218.
- [Har1999] M. Härterich, Kazhdan-Lusztig-Basen, unzerlegbare Bimoduln, und die Topologie der Fahnenmannigfaltigkeit einer Kac-Moody-Gruppe. Dissertation in Freiburg, 1999
- [Hog2017] M. Hogancamp, Idempotents in triangulated monoidal categories, Preprint, arXiv:1703.01001.
- [KMMZ2019] T. Kildetoft, M. Mackaay, V. Mazorchuk, J. Zimmermann, Simple transitive 2-representations of small quotients of Soergel bimodules. *Trans. Amer. Math. Soc.* 371 (2019), 5551–5590.
- [LM2022] R. Laugwitz and V. Miemietz, Pretriangulated 2-representations via dg algebra 1-morphisms. Preprint, arXiv:2205.09999.
- [LNT2003] B. Leclerc, M. Nazarov, J.-Y. Thibon, Induced representations of affine Hecke algebras and canonical bases of quantum groups. *Studies in Memory of Issai Schur*, A. Joseph et al, Birkhäuser, Boston, 2003, 115–153.
- [LW2014] N. Libedinsky and G. Williamson, Standard objects in 2-braid groups. *Proc. Lond. Math. Soc. (3)* 109 (2014), no. 5, 1264–1280.
- [MM2017] M. Mackaay and V. Mazorchuk, Simple transitive 2-representations of some 2-subcategories of Soergel bimodules. *Journal of Pure and Applied Algebra* 221(3) (2017), 565–587.
- [MMMTZ2019] M. Mackaay, V. Mazorchuk, V. Miemietz, D. Tubbenhauer, X. Zhang, Simple transitive 2-representations of Soergel bimodules for finite Coxeter types. Preprint, arXiv:1906.11468.
- [MMMTZ2020] M. Mackaay, V. Mazorchuk, V. Miemietz, D. Tubbenhauer, X. Zhang, Finitary birepresentations of finitary bicategories. *Forum Mathematicum* 33(5) (2021), 1261–1320.
- [MT2017] M. Mackaay and A.-L. Thiel, Categorifications of the extended affine Hecke algebra and the affine q -Schur algebra $\widehat{S}(n, r)$ for $3 \leq r < n$. *Quantum Topol.*, 8(1):113–203, 2017.
- [MT2019] M. Mackaay and D. Tubbenhauer, Two-color Soergel calculus and simple transitive 2-representations. *Canad. J. Math.* 71 (2019), no. 6, 1523–1566.
- [MM2016] V. Mazorchuk, V. Miemietz, Transitive 2-representations of finitary 2-categories. *Trans. Amer. Math. Soc.* 368 (2016), no. 11, 7623–7644.
- [Rou2006] R. Rouquier, Categorification of \mathfrak{sl}_2 and braid groups, *Trends in representation theory of algebras and related topics* (American Mathematical Society, Providence, RI, 2006) 137–167.
- [Soe2007] W. Soergel, Kazhdan-Lusztig-Polynome und unzerlegbare Bimoduln über Polynomringen. *J. Inst. Math. Jussieu* 6 (2007), no. 3, 501–525.
- [Stev2011] G. Stevenson, Tensor actions on locally complete intersections. PhD-thesis, Australian National University, 2011.
- [Zimm2017] J. Zimmermann, Simple transitive 2-representations of Soergel bimodules in type B_2 . *Journal of Pure and Applied Algebra* 221(3) (2017), 666–690.

CENTER FOR MATHEMATICAL ANALYSIS, GEOMETRY, AND DYNAMICAL SYSTEMS, DEPARTAMENTO DE MATEMÁTICA, INSTITUTO SUPERIOR TÉCNICO, 1049-001 LISBOA, PORTUGAL & DEPARTAMENTO DE MATEMÁTICA, FCT, UNIVERSIDADE DO ALGARVE, CAMPUS DE GAMBELAS, 8005-139 FARO, PORTUGAL
Email address: mmackaay@ualg.pt

SCHOOL OF MATHEMATICS, UNIVERSITY OF EAST ANGLIA, NORWICH NR4 7TJ, UNITED KINGDOM
Email address: v.miemietz@uea.ac.uk

INSTITUT DE RECHERCHE EN MATHÉMATIQUE ET PHYSIQUE, UNIVERSITÉ CATHOLIQUE DE LOUVAIN, CHEMIN DU CYCLOTRON 2, 1348 LOUVAIN-LA-NEUVE, BELGIUM
Email address: pedro.vaz@uclouvain.be