

PRIMITIVE IRREDUCIBLE REPRESENTATIONS OF FINITELY GENERATED NILPOTENT GROUPS

ANATOLII. V. TUSHEV

Department of Mathematics, Dnipro National University,
Gagarin Avenue 72, Dnipro, 49065, Ukraine
anavlatu@gmail.com

ABSTRACT. In the paper we show that any irreducible representation of a finitely generated nilpotent group G over a finitely generated field F of characteristic zero is induced from a primitive representation of some subgroup of G .

1. INTRODUCTION

Let G be a group and let R be a ring. Let H be a subgroup of the group G and let U be a right RH -module. Since the group ring RG can be considered as a left RH -module, we can define the tensor product $U \otimes_{RH} RG$, which is a right RG -module named the RG -module induced from the RH -module U . If M is an RG -module and $U \leq M$, then it follows from [7, Chap. 2, Lemma 1.1] that $M = U \otimes_{RH} RG$ if and only if $M = \bigoplus_{t \in T} Ut$, where T is a right transversal to the subgroup H in G .

If a group G has a finite series each of whose factor is either cyclic or quasi-cyclic then G is said to be minimax. If in such a series all infinite factors are cyclic then the group G is said to be polycyclic; the number $h(G)$ of infinite factors in such a series is the Hirsch number of G . By [19, Theorem 2.13], any finitely generated nilpotent group is polycyclic.

Let G be a group, let F be a field, let φ be a representation of G over F and let M be an FG -module of the representation φ . The representation φ is said to be faithful if $\text{Ker} \varphi = 1$. If M is induced from some FH -module U , where H is a subgroup of the group G , then we say that the representation φ is induced from a representation ϕ of the subgroup H , where U is the module of the representation ϕ . The module M and the representation φ are said to be primitive if there are no subgroups $H < G$ such that M is induced from an FH -submodule. If the group G is polycyclic then the module M and the representation

φ are said to be semi-primitive if there are no subgroups $H \leq G$ such that $h(H) < h(G)$ and M is induced from an FH -submodule.

There are many results which show that the existence of a faithful irreducible representation of a group G over a field F may have essential influence on the structure of the group G (see for instance [11–17]). Certainly, primitive irreducible modules are a basic subject for investigations when we are dealing with irreducible representations.

In [4] Harper solved a problem raised by Zaleskii and proved that any not abelian-by-finite finitely generated nilpotent group has an irreducible primitive representation over a not locally finite field. In [16] we proved that if a minimax nilpotent group G of class 2 has a faithful irreducible primitive representation over a finitely generated field of characteristic zero then the group G is finitely generated. In [5] Harper studied polycyclic groups which have faithful irreducible semi-primitive representations. It is well known that any polycyclic group is finitely generated soluble of finite rank and meets the maximal condition for subgroups (in particular, for normal subgroups). In [15] we showed that in the class of soluble groups of finite rank with the maximal condition for normal subgroups only polycyclic groups may have faithful irreducible primitive representations over a field of characteristic zero.

It is well known that any finitely dimensional irreducible representation of a group G is induced from a primitive representation of some subgroup of G . It is evident that any representation of a polycyclic group G over a field F is induced from a semi-primitive representation of some subgroup H of G . We say that a semi-primitive representation of G is purely semi-primitive if it is not induced from a primitive representation of some subgroup H of G and it is still unknown whether polycyclic groups may have purely semi-primitive representations. In the paper we show that under some additional conditions on the group G and the field F semi-primitive irreducible representations do not occur. Our main result (Theorem 1) states that any irreducible representation of a finitely generated nilpotent group G over a finitely generated field F of characteristic zero is induced from a primitive representation of some subgroup of G . So, the following questions naturally arise.

Question 1 *Is it true that any irreducible representation of a finitely generated nilpotent group G over a field F is induced from a primitive representation of some subgroup of G ?*

Question 2 *Is it true that any irreducible representation of a minimax nilpotent group G over a field F is induced from a primitive representation of some subgroup of G ?*

Our methods of investigations are based on the following approach introduced by Brookes in [3]. Let N be a group and let K be a normal subgroup of N such that the quotient group N/K is finitely generated free abelian. Let R be a ring and let W be a finitely generated RN -module. Let I be an N -invariant ideal of RK such that $|K/I^\dagger| < \infty$ and $k = R/(R \cap I)$ is a field, where $I^\dagger = K \cap (1 + I)$. Then N/I^\dagger has a central subgroup A of finite index. So, the quotient module W/WI may be considered as a finitely generated kA -module and we can apply powerful techniques of commutative algebra for studying properties of W/WI . The results of the paper show that properties of W and W/WI are deeply related.

The main results of the paper were announced in [18]. The author is deeply grateful to the referee for helpful notes.

2. AUXILIARY RESULTS

Lemma 1. *Let G be a group and let D be a finite central subgroup of G . Let $\{H_i | i \in \mathbb{N}\}$ be a descending chain of subgroups of G . Then:*

- (i) *if there is $n \in \mathbb{N}$ such that $DH_n = DH_i$ for all $i \geq n$ then there is $m \in \mathbb{N}$ such that $H_m = H_i$ for all $i \geq m$;*
- (ii) *if $\bigcap_{i \in \mathbb{N}} H_i = 1$ then $\bigcap_{i \in \mathbb{N}} DH_i = D$.*

Proof. (i) There is no harm in assuming that $n = 1$. Since $\{D \cap H_i | i \in \mathbb{N}\}$ is a descending chain of subgroups of D and $|D| < \infty$, we can conclude that there is $m \in \mathbb{N}$ such that $D \cap H_m = D \cap H_i$ for all $i \geq m$. We also may assume that $m = 1$. Thus, we can assume that $K = D \cap H_1 = D \cap H_i$ for all $i \in \mathbb{N}$. As the subgroup D is central, passing to the quotient group G/K we can assume that $1 = D \cap H_1 = D \cap H_i$ for all $i \in \mathbb{N}$. Then $DH_i = D \times H_i$ for all $i \in \mathbb{N}$ and, as $H_i \leq H_1$, we can conclude that $DH_i = DH_1$ if and only if $H_i = H_1$.

(ii). Since $\bigcap_{i \in \mathbb{N}} H_i = 1$, there is $m \in \mathbb{N}$ such that $D \cap H_i = 1$ for all $i \geq m$. Evidently, there is no harm in assuming that $m = 1$. Then any element of DH_1 can be unequally written in the form dh , where $d \in D$, $h \in H_1$, and hence $dh \in DH_i$ if and only if $h \in H_i$. As $\bigcap_{i \in \mathbb{N}} H_i = 1$, it implies that $\bigcap_{i \in \mathbb{N}} DH_i = D$. \square

Lemma 2. *Let G be a finitely generated nilpotent group and let D be the derived subgroup of G . Let $\{H_i | i \in \mathbb{N}\}$ be a descending chain of subgroups of finite index in G . Suppose that there is $n \in \mathbb{N}$ such that $DH_n = DH_i$ for all $i \geq n$. Then there is $m \in \mathbb{N}$ such that $H_m = H_i$ for all $i \geq m$.*

Proof. Proof is by induction on the length of the lower central series $1 = G_0 \leq G_1 \leq \dots \leq G_k = G$ of the group G . Put $K = G_1$ then by the induction hypothesis we can assume that there is $n \in \mathbb{N}$ such that $KH_n = KH_i$ for all $i \geq n$. Evidently, we can assume that $n = 1$.

It is well known that $[ab, c] = [a, c]^b[b, c]$ and $[c, ab] = [c, b][c, a]^b$ for any elements $a, b, c \in G$ (see [6, Section 3.2]). As $[G_2, G] = G_1 = K \leq Z(G)$, we see that if some of elements a, c belongs to G_2 then $[ab, c] = [a, c][b, c]$ and $[c, ab] = [c, b][c, a]$. It easily implies that if some of elements a, b belongs to G_2 then

$$(1) \quad [a^u, b^v] = [a, b]^{uv}$$

for any $u, v \in \mathbb{N}$. Since $|G : H_1| < \infty$, there are $u, v \in \mathbb{N}$ such that $a^u, b^v \in H_1$ for any $a \in G_2$ and any $b \in G$. Then it follows from (1) that $[a, b]^{uv} \in [H_1, H_1]$. Since the subgroup $K = G_1 = [G_2, G]$ is abelian and generated by commutators $[a, b]$, where $a \in G_2$ and $b \in G$, it implies that $K^{uv} \leq [H_1, H_1] \leq H_1$. As $KH_1 = KH_i$ for all $i \geq n$, and K is a central subgroup of G , we can conclude that $K^{uv} \leq [H_1, H_1] = [KH_1, KH_1] = [KH_i, KH_i] = [H_i, H_i] \leq H_i$ for all $i \in \mathbb{N}$.

Since the subgroup K^{uv} is central, we can pass to the quotient group G/K^{uv} and hence, as $|K/K^{uv}| < \infty$, we can assume that the subgroup K is finite. Then the assertion follows from Lemma 1.1(i). \square

Let p be a prime integer then $n \in \mathbb{N}$ is said to be a p' -number if p does not divide n . A finite group is called a p' -group if its order is a p' -number.

Proposition 1. *Let G be a finitely generated nilpotent group and let D be the derived subgroup of G . Let $\{H_i | i \in \mathbb{N}\}$ be a descending chain of subgroups of finite index in G such that $|G : \bigcap_{i \in \mathbb{N}} H_i| = \infty$. Then there is a descending chain $\{G_i | i \in \mathbb{N}\}$ of subgroups of finite index in G such that $DH_i \leq G_i$, the quotient group $G/(\bigcap_{i \in \mathbb{N}} G_i)$ is free abelian and:*

- (i) either for any prime integer p there is $n \in \mathbb{N}$ such that $p \notin \pi(G_n/G_i)$ for all $i \geq n$;
- (ii) or G/G_i is a p -group for all $i \in \mathbb{N}$, where p is a prime integer.

Proof. It is easy to note that $|G : \bigcap_{i \in \mathbb{N}} H_i| < \infty$ if and only if there is $m \in \mathbb{N}$ such that $H_m = H_i$ for all $i \geq m$. Then it follows from Lemma 2, that $|G/(\bigcap_{i \in \mathbb{N}} DH_i)| = \infty$ and hence there is no harm in assuming that $D \leq H_i$ for all $i \geq m$. Then passing to the quotient group G/D we can assume that the group G is abelian.

(i) If for any prime integer p there is $n \in \mathbb{N}$ such that $p \notin \pi(H_n/H_i)$ for all $i \geq n$ then we can put $G_i = H_i$ for all $i \in \mathbb{N}$.

(ii) Suppose that the situation (i) does not hold for $\{H_i | i \in \mathbb{N}\}$. Then there is a prime integer p such that for any $i \in \mathbb{N}$ there is $j \in \mathbb{N}$ such that $j > i$ and $p \in \pi(H_i/H_j)$. Let G_i be the set of all elements $g \in G$ such that $g^n \in H_i$ for some p' -number $n \in \mathbb{N}$. It is easy to check that G_i is a subgroup of G such that $H_i \leq G_i$, $G/H_i = (G_i/H_i) \times (S_i/H_i)$, where G_i/H_i is a p' -group and S_i/H_i is a p -group. It implies that G/G_i is a p -group for all $i \in \mathbb{N}$. As $H_i \geq H_{i+1}$, we can conclude that $G_i \geq G_{i+1}$ and hence $\{G_i | i \in \mathbb{N}\}$ is a descending chain of subgroups of finite index in G . Since for any $i \in \mathbb{N}$ there is $j \in \mathbb{N}$ such that $p \in \pi(H_i/H_j)$, we can conclude that for any $i \in \mathbb{N}$ there is $j \in \mathbb{N}$ such that $j > i$ and $|G/G_j| > |G/G_i|$. It implies that $|G/(\bigcap_{i \in \mathbb{N}} G_i)| = \infty$. Passing to the quotient group $G/(\bigcap_{i \in \mathbb{N}} G_i)$, we can assume that $\bigcap_{i \in \mathbb{N}} G_i = 1$. As the group G is finitely generated abelian, the torsion subgroup T of G is finite. Then it follows from Lemma 1(ii) that $\bigcap_{i \in \mathbb{N}} TG_i = T$. Therefore, changing G_i by TG_i and passing to the quotient group G/T we can assume that G is torsion-free and hence G is free abelian. \square

Lemma 3. *Let G be a group of finite rank which has a finite subgroup D such that the quotient group G/D is torsion-free abelian then G has a characteristic central torsion-free subgroup A of finite index.*

Proof. As G' is a characteristic subgroup of G , it is not difficult to show that $C = C_G(G')$ is a characteristic subgroup of G . Then $B = C^m$ is a characteristic subgroup of G , where $m = |D|$. Since $|G'| < \infty$ and $C = C_G(G')$, we can conclude that $|G : C| < \infty$. As the group G has finite rank, it is easy to note that $|C : C^m| < \infty$ and hence we can conclude that $|G : B| < \infty$. For any $a, b \in C$ and any $g \in G$ we have $[ab, g] = [a, g]^b [b, g]$ and, as $[a, g] \in G'$ and $b \in C = C_G(G')$, we can conclude that $[ab, g] = [a, g][b, g]$.

Any element $a \in B = C^m$ may be presented in the form $a = a_1^m \dots a_t^m$, where $a_i \in C$ and $1 \leq i \leq t$. Since $[ab, g] = [a, g][b, g]$, for any $g \in G$ we have $[a, g] = [a_1, g]^m \dots [a_t, g]^m$. Then, as $m = |D|$ and $[a_i, g] \in G' \leq D$, where $1 \leq i \leq t$, we can conclude that $[a, g] = 1$. It implies that B is a characteristic central subgroup of finite index in G . Since the quotient group G/D is torsion-free, so is $B/(B \cap D)$ and it implies that $A = B^m$ is a torsion-free characteristic central subgroup of finite index in G . \square

3. CULLING IDEALS OF GROUP RINGS OF FINITELY GENERATED NILPOTENT GROUPS

A commutative ring R is Hilbert if every prime ideal in R is an intersection of some maximal ideals of R . A ring R is absolutely Hilbert if it is commutative, Noetherian, Hilbert and all field images of R are locally finite. Let G be a group and let K be a normal subgroup of G . Let R be a ring and let I be a G -invariant ideal of the group ring RK then $I^\dagger = G \cap (I + 1)$ is a normal subgroup of G . We say that a G -invariant ideal I is G -large if $R/(R \cap I) = k$ is a field and $|K/I^\dagger| < \infty$. Let R be an absolutely Hilbert ring, let V be an RK -module and U be a submodule of V . According to [3, Section 1] a G -invariant ideal I of RK culls U in V if

- (a) $VI < V$,
- (b) $VI + U = V$,
- (c) $R/(R \cap I) = k$ is a field,
- (d) $(I \cap RC)RK = I$, where $C = cch_p(K)$ is the centralizer of chief p -factors of K and $p = \text{char} K$.

However, (c) holds for any G -large ideal of RK and (d) holds for any nilpotent group because in this case $K = cch_p(K)$. So, in the case where the group K is nilpotent, a G -invariant ideal I of RK culls a submodule U of V if I satisfies conditions (a), (b) and (c), and if the ideal I is G -large then I culls U in V if I satisfies conditions (a) and (b).

An RK -module V is G -ideal critical if for any non-zero submodule U of V there is a G -invariant ideal I of RK which culls the submodule U in V .

Lemma 4. *Let G be a finitely generated nilpotent group and let K be a normal subgroup of G . Let R be an absolutely Hilbert ring. Let V be an RK -module, let U be a non-zero submodule of V and let J be a G -invariant ideal of RK which culls U in V . Then there is a G -large ideal I of RK which culls U in V and such that $J \leq I$.*

Proof. It is sufficient to show that there is a G -invariant ideal I of RK which culls U in V such that $J \leq I$, $R/(R \cap I) = k$ is a field and $|K/I^\dagger| < \infty$. This assertion was proved in the second paragraph of section 5 of [3]. \square

Let $\pi \subseteq \mathbb{N}$ be a set of prime integers, we will say that a G -large ideal I is (π, G) -large if $\text{char}(R/(R \cap I)) \notin \pi$. We will say that an RK -module V is (π, G) -large critical if for any non-zero submodule U of V there is a (π, G) -large ideal I of RK which culls the submodule U in V . Evidently, any (π, G) -large critical RK -module is G -ideal critical.

Lemma 5. *Let G be a finitely generated nilpotent group and let K be a normal subgroup of G . Let R be an absolutely Hilbert domain of characteristic zero. Let V be a G -ideal critical uniform RK -module which is R -torsion-free. Then V is (π, G) -large critical for any finite set of prime integers $\pi \subseteq \mathbb{N}$.*

Proof. Let $\pi = \{p_1, \dots, p_k\}$ be a finite set of prime integers and let U be a non-zero submodule of V . Put $X = (\bigcap_{i=1}^n Vp_i) \cap U$, as $\text{char} R = 0$ and V is R -torsion-free, we see that $Vp_i \neq 0$ for all $1 \leq i \leq n$ and hence, as the module V is uniform, we can conclude that $X \neq 0$. Since V is a G -ideal critical RK -module, there is a G -invariant ideal I of RK which culls X in V and, by Lemma 4, we can assume that the ideal I is G -large. By [3, Lemma 6], the ideal I culls U and each Vp_i , where, $1 \leq i \leq n$. If $\text{char}(R/(R \cap I)) = p_i$ for some $p_i \in \pi$ then $1p_i \in I$, where 1 is the unite of R . It implies that $Vp_i \leq VI$ but it is impossible because I culls Vp_i . Thus, $\text{char}(R/(R \cap I)) \notin \pi$. \square

We should make some notes to apply Theorems E and D of [3] in the next Proposition. A free abelian group A of finite rank acted on by a group G is a G -plinth if all subgroups of finite index in G act rationally irreducibly on A . A polycyclic normal subgroup K of a group G has a G -plinth series if it has G -invariant subgroups

$$(2) \quad 1 = K_1 \leq K_2 \leq \dots \leq K_n = K$$

with each factor K_{i+1}/K_i either finite or a G -plinth. It is easy to note that if the group G is nilpotent then any central series (2) with cyclic factors is an X -plinth series for any subgroup X of G .

Let R be an absolutely Hilbert ring. According to [3, Section 2] an RK -module V is \mathcal{S} -calibrated (with respect to the series (2)) if there are uniform RK_i -submodules such that

- (1) $V_1 \cong R/Q$ for some prime ideal Q of R ,
- (2) $V_i = V_1 RK_i$ for $1 \leq i \leq n$ and $V_n = V$,
- (3) either
 - (i) K_{i+1}/K_i is infinite cyclic and $V_{i+1} = V_i \otimes_{RK_i} RK_{i+1}$, or
 - (ii) $u - \dim_{RK_i}(V_{i+1}) < \infty$ and V_i is K_{i+1} -essential in V_{i+1} , i.e. $V_i \cap Y \neq 0$ for any non-zero RK_{i+1} -submodule $Y \leq V_{i+1}$.

Two RK -modules V and V_1 are similar if their injective hulls are isomorphic. If V and V_1 are similar we denote $V \approx V_1$. By [2, Lemma 3.2(i)], $S_G(V) = \{g \in G | Vg \approx V\}$ is a subgroup of G which is called the the similiser of the RK -module V in G .

Proposition 2. *Let K be a normal subgroup of a finitely generated nilpotent group G . Let R be an absolutely Hilbert domain of characteristic zero. Let W be a finitely generated RK -module which is R -torsion-free. Then there is a cyclic uniform RK -submodule $aRK = V \leq W$ which is (π, G) -large critical for any finite set of prime integers $\pi \subseteq \mathbb{N}$.*

Proof. By [19, Theorem 3.7], W is a Noetherian RK -module and hence, by [8, Chap.6, Lemma 2.3], W has Krull dimension. Then, by [8, Chap.6, Lemma 2.10], W has a critical submodule U which, by [8, Chap.6, Lemma 2.13], is uniform. Let $S = S_G(U)$ then $S = S_S(U)$. As the group G is finitely generated nilpotent, it has a central series with cyclic factors and hence the subgroup K has a finite series (2) with cyclic factors which are centralized by the group G . Then, as it was mentioned above, series (2) may be considered as an X -plinth series for any subgroup $X \leq G$. So, we can assume that (2) is an S -plinth series. Therefore, by [3, Theorem E], U has an \mathcal{S} -calibrated submodule V with respect to the S -plinth series (2). However, (2) is also a G -plinth series. Then, by [3, Theorem D], the module V is G -ideal critical. By the definition of an \mathcal{S} -calibrated submodule, $V = V_1RK$, where $V_1 \cong R/Q$. It implies that $V_1 = aR$, where $a \in V_1$ and $\text{Ann}_R(a) = Q$, and hence $V = aRK$, i.e. the RK -module V is cyclic. Then, as the module V is G -ideal critical, the assertion follows from Lemma 5. \square

4. COMMUTATIVE INVARIANTS FOR MODULES OVER FINITELY GENERATED NILPOTENT GROUPS

Let P be a prime ideal of a commutative ring S and let S_P be the localization of S at the ideal P . Let M be an S -module, the support $\text{Supp}_S M$ of the module M consists of all prime ideals P of S such that $M_P = M \otimes_S S_P \neq 0$ (see. [1, Chap. II, §4.4]).

By [1, Chap. IV, §1.4, Theorem 2], if S and M are Noetherian then the set $\mu_S(M)$ of minimal elements of $\text{Supp}_S M$ is finite and coincides with $\mu_S(\text{Ann}_S M)$, where $\mu_S(\text{Ann}_S M)$ is the set of prime ideals of S which are minimal over $\text{Ann}_S M$, i.e. we have

$$(3) \quad \mu_S(M) = \mu_S(\text{Ann}_S M)$$

Further in the paper, if J is an ideal of S then $\mu_S(J)$ will denote the set of minimal prime ideals over J .

Let G be a nilpotent group and let K be a normal subgroup of G such that the quotient group G/K is finitely generated free abelian. Let R be an absolutely Hilbert ring, if I is a G -large ideal of the group ring RK then the derived subgroup of the quotient group G/I^\dagger is finite and

it follows from Lemma 3 that G/I^\dagger has a characteristic central subgroup A of finite index. Since the quotient group G/K is finitely generated free abelian, the subgroup A may be chosen finitely generated free abelian. Further in the paper, A will denote a finitely generated free abelian central subgroup of G/I^\dagger . Put $k = R/(R \cap I)$, by [19, Theorem 3.7], the group ring kA is Noetherian and, by [10, Chap.1, Lemma 1.1], kA is an integral domain.

Let W be a finitely generated RG -module then W/WI is a finitely generated kA -module and hence W/WI is a Noetherian kA -module. So, the finite set $\mu_{kA}(W/WI) = \mu_{kA}(\text{Ann}_{kA}(W/WI))$ is defined.

Further in the paper, for any RG -module W we consider W/WI as a kA -module, where I is a G -large ideal of RK , A is a central free abelian subgroup of finite index in G/I^\dagger and $k = R/(R \cap I)$.

Lemma 6. *Let G be a finitely generated nilpotent group and let K be a normal subgroup of G such that the quotient group G/K is free abelian. Let R be an absolutely Hilbert domain and let W be a finitely generated RG -module. Let I be a G -large ideal of RK , let A be a central free abelian subgroup of finite index in G/I^\dagger and let $k = R/(R \cap I)$. Then for any RG -submodule $W_1 \leq W$:*

- (i) W_1/W_1I has a finite series each of whose quotient is isomorphic to some section of W/WI ;
- (ii) $\text{Supp}_{kA}W_1/W_1I \subseteq \text{Supp}_{kA}W/WI$;

Proof. (i). By [10, Chap.11, Theorem 2.9], $WI^n \cap W_1 \leq W_1I$ for some $n \in \mathbb{N}$. It implies that W_1/W_1I has a finite series each of whose factor is isomorphic to some section of WI^{k-1}/WI^k , where $1 \leq k \leq n$. Then it is sufficient to show that each section WI^{k-1}/WI^k has a finite series each of whose factor is isomorphic to some section of W/WI . Applying induction on k , it is sufficient to show that WI/WI^2 has a finite series each of whose factor is isomorphic to some section of W/WI . As the ideal I is G -invariant, $J = IRG$ is an ideal of RG . Evidently, $WI^{k-1}/WI^k = WJ^{k-1}/WJ^k$ for all $1 \leq k \leq n$. By [10, Chap.11, Theorem 3.12], the ideal $J = IRG$ is polycentral, i.e. J has generating elements $\{a_1, a_2, \dots, a_r\}$ such that $a_i + J_{i-1}$ is a central element of RG/J_{i-1} , where J_{i-1} is the ideal of J generated by $\{a_1, \dots, a_{i-1}\}$. As it was mentioned above, it is sufficient to show that WJ/WJ^2 has a finite series each of whose factor is isomorphic to some section of W/WJ . We apply induction on the number of the generating elements $\{a_1, a_2, \dots, a_r\}$. Passing to the quotient module W/WJ^2 we can assume that $WJ^2 = 0$. The mapping $\varphi : W \rightarrow W$ given by $\varphi : x \mapsto xa_1$ is an endomorphism such that $WJ \leq \text{Ker}\varphi$. Therefore, the submodule $\varphi(W) = Wa_1$ of WJ is isomorphic to the section $W/\text{Ker}\varphi$ of W/WJ .

Thus, passing to the quotient module W/Wa_1 we can assume that $Wa_1 = 0$. Then we can pass to the ideal J/J_1 of RG/J_1 and the assertion follows from the induction hypothesis.

(ii) The assertion follows from (i) and [1, Chap.II, §4, Proposition 16]. □

Proposition 3. *Let G be a finitely generated nilpotent group and let K be a normal subgroup of G such that the quotient group G/K is free abelian. Let R be an absolutely Hilbert domain, let F be the field of fractions of R and let M be an irreducible FG -module. Let I be a G -large ideal of RK , let A be a central free abelian subgroup of finite index in G/I^\dagger and let $k = R/(R \cap I)$. Then for any non-zero elements $a, b \in M$:*

- (i) aRG contains a submodule which is isomorphic to bRG ;
- (ii) $\mu_{kA}((akG)/(akG)I) = \mu_{kA}((bkG)/(bkG)I)$;
- (iii) if $(akG)/(akG)I \neq 0$ then $(bkG)/(bkG)I \neq 0$;
- (iv) if $(akG)/(akG)I$ is kA -torsion then $(bkG)/(bkG)I$ is kA -torsion.

Proof. (i) As the module M is irreducible, there is $c \in FG$ such that $ac = b$. As F is the field of fractions of the domain R , there is $d \in R$ such that $cd \in RG$. Therefore, $bd = acd \in aRG$. As $d \in R$ we see that RG -modules $bdRG$ and bRG are isomorphic and the assertion follows.

(ii) By (i), we can assume that $bRG \leq aRG$. Then it follows from Lemma 6(ii) that $Supp_{kA}(bRG)/(bRG)I \subseteq Supp_{kA}(aRG)/(aRG)I$. However, the arguments of (i) show that we can assume that $aRG \leq bRG$. Therefore, $Supp_{kA}(aRG)/(aRG)I \subseteq Supp_{kA}(bRG)/(bRG)I$ and hence $Supp_{kA}(aRG)/(aRG)I = Supp_{kA}(bRG)/(bRG)I$. It implies that $\mu_{kA}((akG)/(akG)I) = \mu_{kA}((bkG)/(bkG)I)$.

(iii) By (i), we can assume that $aRG \leq bRG$. By [10, Chap.11, Theorem 2.9], $(bRG)I^n \cap (aRG) \leq (aRG)I$ for some $n \in \mathbb{N}$ and, as $(akG)/(akG)I \neq 0$, it implies that $(bRG)I \neq (bRG)$, i.e. $(bkG)/(bkG)I$.

(iv) By (i), we can assume that $bRG \leq aRG$. Then it follows from Lemma 6(i) that $(bRG)/(bRG)I$ has a finite series each of whose quotient is isomorphic to some section of $(aRG)/(aRG)I$. Since the kA -module $(aRG)/(aRG)I$ is kA -torsion, it implies that $(bRG)/(bRG)I$ has a finite series each of whose quotient is kA -torsion. Then, as kA is a domain, we can conclude that $(bRG)/(bRG)I$ is kA -torsion. □

Proposition 4. *Let G be a finitely generated nilpotent group and let $H \leq G$. Let R be an absolutely Hilbert domain and let $0 \neq W$ be an RG -module. Let K be a normal subgroup of G such that $K \leq H$. Let I be a G -large ideal of RK , let $k = R/(I \cap R)$, let A be a free abelian central subgroup of finite index in G/I^\dagger and $B = (H/I^\dagger) \cap A$.*

Let $0 \neq d \in W$ such that $(dRG)/(dRG)I \neq 0$, let w be the image of d in $(dRG)/(dRG)I$ and let $J = \text{Ann}_{kA}(w)$. Suppose that $dRG = dRH \otimes_{RH} RG$. Then:

- (i) $J = (J \cap kB)kA$;
- (ii) $\mu_{kA}((dRG)/(dRG)I) = \mu_{kA}(J)$;
- (iii) if $(dRG)/(dRG)I \neq 0$ and $(dRG)/(dRG)I$ is kA -torsion then $\mu_{kA}(J)$ consists of proper ideals of kA .

Proof. (i) Since $dRG = dRH \otimes_{RH} RG$, by [7, Chap. 2, Lemma 1.1], we have $dRG = \bigoplus_{t \in T} (dRH)t$, where T is a right transversal to the subgroup H in G . As $K \leq H$, we see that $dRGI = \bigoplus_{t \in T} (dRHI)t$ and hence $dRG/dRGI = \bigoplus_{t \in T} (dRH/dRHI)t$. It implies that $wRG = \bigoplus_{t \in T} (wRH)t$, where w is the image of d in $dRG/dRGI$. Then, as I^\dagger centralizes $dRG/dRGI$ and $R \cap I$ annihilates $dRG/dRGI$, we have $wk(G/I^\dagger) = \bigoplus_{t \in T} (wk(H/I^\dagger))t$, where T is a right transversal to H/I^\dagger in G/I^\dagger . Therefore, $wkA = \bigoplus_{s \in S} (wkB)s$, where S is a right transversal to B in A , and it implies that $\text{Ann}_{kA}(w) = J = (J \cap kB)kA$.

(ii) As I^\dagger centralizes $dRG/dRGI$ and $R \cap I$ annihilates $dRG/dRGI$, we see that $(dRG)/(dRG)I \cong wk(G/I^\dagger)$ and therefore $\mu_{kA}((dRG)/(dRG)I) = \mu_{kA}(wk(G/I^\dagger))$. Since A is a central subgroup of finite index in G/I^\dagger it is not difficult to show that $\text{Ann}_{kA}(wk(G/I^\dagger)) = \text{Ann}_{kA}(w)$ and it implies that $\mu_{kA}(\text{Ann}_{kA}(wk(G/I^\dagger))) = \mu_{kA}(\text{Ann}_{kA}(w)) = \mu_{kA}(J)$. According to Equation (3) we have $\mu_{kA}(wk(G/I^\dagger)) = \mu_{kA}(\text{Ann}_{kA}(wk(G/I^\dagger)))$ and therefore, as $\mu_{kA}((dRG)/(dRG)I) = \mu_{kA}(wk(G/I^\dagger))$, we can conclude that $\mu_{kA}((dRG)/(dRG)I) = \mu_{kA}(J)$.

(iii) Put $U = (dRG)/(dRG)I \neq 0$. According to (ii) and Equation (3) it is sufficient to show that $\mu_{kA}(\text{Ann}_{kA}(U)) = \mu_{kA}(U)$ consists of proper ideals of kA . But, as the kA -module U is finitely generated and kA is a domain, $0 \in \mu_{kA}(\text{Ann}_{kA}(U))$ if and only if U is not kA -torsion, and $kA \in \mu_{kA}(\text{Ann}_{kA}(U))$ if and only if $U = 0$. So, the assertion follows. \square

Lemma 7. *Let A be a finitely generated abelian group, let k be a field and let I be an ideal of kA . Then for any subgroup B of finite index in A the following assertions hold:*

- (i) for any $Q \in \mu_{kB}(I \cap kB)$ there is a prime ideal P of kA such that $I \leq P$, $Q = P \cap kB$ and any such prime P is necessarily in $\mu_{kA}(I)$;
- (ii) if $P \in \mu_{kA}(I)$ then $P \cap kB \notin \mu_{kB}(I \cap kB)$ if and only if there is $S \in \mu_{kA}(I)$ such that $S \cap kB < P \cap kB$;
- (iii) if J is an ideal of kA such that $\mu_{kA}(I) = \mu_{kA}(J)$ then $\mu_{kB}(I \cap kB) = \mu_{kB}(J \cap kB)$;
- (iv) if $\mu_{kA}(I)$ consists of proper ideals of kA then $\mu_{kB}(I \cap kB)$ consists of proper ideals of kB .

Proof. (i) Evidently, the quotient ring kA/I is integer over $(kB+I)/I \cong kB/(kB \cap I)$. Then it follows from [20, Ch.V, §2, Theorem 3] that there is a prime ideal $P \leq kA$ such that $I \leq P$ and $P \cap kB = Q$. If $P \notin \mu_{kA}(I)$ then there is an ideal $P' \in \mu_{kA}(I)$ such that $I \leq P' < P$ and hence $I \cap kB \leq P' \cap kB \leq P \cap kB = Q$ and, as $Q \in \mu_{kB}(I \cap kB)$, we can conclude that $I \cap kB \leq P' \cap kB = P \cap kB = Q$. Since $I \leq P' < P$, it contradicts Complement 1 to [20, Ch.V, §2, Theorem 3]. Thus, $P \in \mu_{kA}(I)$.

(ii) Let $P \in \mu_{kA}(I)$. Suppose that $P \cap kB \notin \mu_{kB}(I \cap kB)$ then there is a prime ideal $Q \in \mu_{kB}(I \cap kB)$ such that $Q < P \cap kB$. By (i), there is $S \in \mu_{kA}(I)$ such that $Q = S \cap kB$ and hence $S \cap kB < P \cap kB$. Suppose now that there is $S \in \mu_{kA}(I)$ such that $S \cap kB < P \cap kB$. Then $I \cap kB \leq S \cap kB < P \cap kB$ and hence $P \cap kB \notin \mu_{kB}(I \cap kB)$.

(iii) Let $Q \in \mu_{kB}(I \cap kB)$. By (i), there is $P \in \mu_{kA}(I)$ such that $Q = P \cap kB$. Then, by (ii), there are no $S \in \mu_{kA}(I)$ such that $S \cap kB < P \cap kB$. Since $\mu_{kA}(I) = \mu_{kA}(J)$, we can conclude that there is $P \in \mu_{kA}(J)$ such that $Q = P \cap kB$ and there are no $S \in \mu_{kA}(J)$ such that $S \cap kB < P \cap kB$. Then it follows from (ii) that $Q \in \mu_{kB}(J \cap kB)$. The same arguments show that if $Q \in \mu_{kB}(J \cap kB)$ then $Q \in \mu_{kB}(I \cap kB)$ and the assertion follows.

(iv) It follows from [15, Lemma 2.2.3(ii)] that if P is a proper ideal of kA then $P \cap kB \neq 0$ and the assertion follows. \square

Lemma 8. *Let A be a free abelian group of finite rank and let B be a subgroup of finite index in A . Let k be a field such that $\text{char } k \notin \pi(A/B)$ and let I be an ideal of kA such that $I = (I \cap kB)kA$. Suppose that there is a prime ideal $Q \in \mu_{kB}(I \cap kB)$ such that there is only one $P \in \mu_{kA}(I)$ such that $Q = kB \cap P$. Then $P = QkA$.*

Proof. As $I = (I \cap kB)kA$ and $(I \cap kB) \leq Q$, we see that $QkA \geq I$. If $P \in \mu_{kA}(I)$ and $Q = kB \cap P$ then $P \geq QkA$ and hence there is only one prime minimal over $QkA \geq I$. Therefore, it is sufficient to show that the ideal QkA is prime because then, by Lemma 7(i), $QkA \in \mu_{kA}(I)$. So, it is sufficient to show that if the quotient ring kA/QkA has only one minimal prime then kA/QkA is a domain. Since the quotient ring kA/QkA is isomorphic to the cross product $(kB/Q) * (A/B)$ and $\text{char } k \notin \pi(A/B)$, it follows from [9, Theorem 2.1] that the quotient ring kA/QkA is semiprime and hence kA/QkA has no non-zero nilpotent elements. Therefore, if kA/QkA is not a domain then there are zero divisors $a, b \in kA/QkA$ such that $a \neq b$ and hence there are at least two different minimal prime ideals of kA over QkA : minimal prime P_1 such that $a \notin P_1$ and minimal prime Q_2 such that $b \notin Q_2$, but it leads to a contradiction. \square

5. PRIMITIVE REPRESENTATIONS OF FINITELY GENERATED
NILPOTENT GROUPS

Let F be a field and let G be a finitely generated nilpotent group. We say that a semi-primitive FG -module M is purely semi-primitive if it is not induced from a primitive FX -module for any subgroup $X \leq G$.

Lemma 9. *Let G be a finitely generated nilpotent group and let H be a subgroup of G . Let F be a field and let M be an FG -module such that $M = V \otimes_{FH} FG$, where V is an FH -module. Then:*

- (i) *if M is an irreducible FG -module then V is an irreducible FH -module;*
- (ii) *if M is a purely semi-primitive FG -module then V is a purely semi-primitive FH -module.*

Proof. (i) It follows from [7, Lemma 2.8(i, ii)] that if U is a proper submodule of V then $N = U \otimes_{FH} FG$ is a proper submodule of M . As the FG -module M is irreducible, it implies that the FH -module V is irreducible.

(ii) If the module V is not semi-primitive then there is a subgroup K of H such that $V = U \otimes_{FK} FH$, where U is an FK -submodule of V and $h(K) < h(H)$. Then, as $M = V \otimes_{FH} FG$, it follows from [7, Lemma 2.8(i, ii)] that $M = U \otimes_{FK} FG$ but, as $h(K) < h(H) \leq h(G)$, it is impossible because M is semi-primitive. Thus, V is semi-primitive.

If V is not purely semi-primitive then there is a subgroup K of H such that $V = U \otimes_{FK} FH$, where U is a primitive FK -module. Then, as $M = V \otimes_{FH} FG$, it follows from [7, Lemma 2.1] that $M = U \otimes_{FK} FG$ but it is impossible because M is purely semi-primitive. □

Lemma 10. *Let F be a field and let G be a finitely generated nilpotent group. Suppose that there exists a purely semi-primitive irreducible FG -module M . Then there are a descending chain $\{G_i | i \in \mathbb{N}\}$ of subgroups of finite index in G and a set $\{a_i | i \in \mathbb{N}\} \subseteq M$ such that :*

- (i) *G/K is a free abelian group, where $K = \bigcap_{i \in \mathbb{N}} G_i$ and either for any prime integer p there is $n \in \mathbb{N}$ such that $p \notin \pi(G_n/G_i)$ for all $i \geq n$ or G/G_i is a p -group for all $i \in \mathbb{N}$, where p is a prime integer;*
- (ii) *$a_i FG = a_i FG_i \otimes_{FG_i} FG$ for each $i \in \mathbb{N}$.*

Proof. As the module M is irreducible and purely semi-primitive, there are a proper subgroup H_1 of finite index in G and an FH_1 -submodule V_1 of M such that $M = V_1 \otimes_{FH_1} FG$. It follows from Lemma 9 that the FH_1 -submodule V_1 is irreducible and purely semi-primitive. Then the above arguments show that there are a proper subgroup H_2 of finite index in H_1 and an FH_2 -submodule V_2 of V_1 such that $V_1 =$

$V_2 \otimes_{FH_2} FH_1$. As $M = V_1 \otimes_{FH_1} FG$ and $V_1 = V_2 \otimes_{FH_2} FH_1$, it follows from [7, Lemma 2.1] that $M = V_2 \otimes_{FH_2} FG$. It follows from Lemma 9 that the FH_2 -submodule V_2 is irreducible and purely semi-primitive. Continuing this process we obtain an infinite strongly descending chain $\{H_i | i \in \mathbb{N}\}$ of subgroups of finite index in G and a strongly descending chain $\{V_i | i \in \mathbb{N}\}$, where V_i is an irreducible and purely semi-primitive FH_i -module such that $M = V_i \otimes_{FH_i} FG$ for each $i \in \mathbb{N}$. As M is an irreducible FG -submodule and V_i is an irreducible FH_i -module, we can conclude that for any $a_i \in V_i \setminus V_{i+1}$ we have $a_i FG = M$ and $a_i FH_i = V_i$ for each $i \in \mathbb{N}$. Therefore, $a_i FG = a_i FH_i \otimes_{FH_i} FG$ for each $i \in \mathbb{N}$.

Thus, we have an infinite strongly descending chain $\{H_i | i \in \mathbb{N}\}$ of subgroups of finite index in G and a set $\{a_i | i \in \mathbb{N}\} \subseteq M$ such that $a_i FG = a_i FH_i \otimes_{FH_i} FG$ for each $i \in \mathbb{N}$. As the chain $\{H_i | i \in \mathbb{N}\}$ is strongly descending, we see that $|G : \bigcap_{i \in \mathbb{N}} H_i| = \infty$. Then, by Proposition 1, there is a descending chain $\{G_i | i \in \mathbb{N}\}$ of subgroups of finite index in G such that $H_i \leq G_i$ and the quotient group G/K is free abelian, where $K = \bigcap_{i \in \mathbb{N}} G_i$.

(i) By Proposition 1, either for any prime integer p there is $n \in \mathbb{N}$ such that $p \notin \pi(G_n/G_i)$ for all $i \geq n$ or G/G_i is a p -group for all $i \in \mathbb{N}$, where p is a prime integer.

(ii) As $H_i \leq G_i$ and $a_i FG = a_i FH_i \otimes_{FH_i} FG$, it follows from [7, Lemma 2.1] that $a_i FG = a_i FG_i \otimes_{FG_i} FG$ for all $i \in \mathbb{N}$. □

Lemma 11. *Let G be a finitely generated nilpotent group and let K be a normal subgroup of G such that the quotient group G/K is finitely generated free abelian. Let R be an absolutely Hilbert ring. Let W be an RG -module such that $W = VRG$ and $W \neq V \otimes_{RK} RG$, where V is a uniform G -ideal critical RK -module. Then there exists a submodule U of V with the property that when a G -large ideal I of RK culls U in V then $W/WI \neq 0$ and W/WI is kA -torsion, where $k = R/(R \cap I)$ and A is a free abelian central subgroup of finite index in G/I^\dagger .*

Proof. It follows from [3, lemma 8] that there exists a non-zero submodule V_1 of V with the property that when a G -large ideal I of RK culls V_1 in V then $W \neq V \otimes_{RK} RG$ if and only if $W/WI \otimes_{kA} Q = 0$, where Q is the field of fractions of kA . As $W \neq V \otimes_{RK} RG$, if I culls V_1 in V then $W/WI \otimes_{kA} Q = 0$. However, it is well known that $W/WI \otimes_{kA} Q = 0$ if and only if W/WI is kA -torsion.

It follows from [3, lemma 14] that there is a non-zero submodule V_2 of V with the property that when a G -large ideal I of RK culls V_2 in V then $W/WI \neq 0$. Let $U = V_1 \cap V_2$ then, as V is uniform, $U \neq 0$

and, by [3, lemma 6], if a G -invariant ideal I of RK culls U in V then I culls V_1 and V_2 . So, the assertion follows. \square

Lemma 12. *Let F be a finitely generated field of characteristic zero and let G be a finitely generated nilpotent group. Suppose that there exists a purely semi-primitive irreducible FG -module M . Then there are an absolutely Hilbert domain $R \leq F$, a descending chain $\{G_i | i \in \mathbb{N}\}$ of subgroups of finite index in G and a set $\{a_i | i \in \mathbb{N}\} \subseteq M$ such that the quotient group G/K is free abelian, where $K = \bigcap_{i \in \mathbb{N}} G_i$, and $a_i RG = a_i RG_i \otimes_{RG_i} RG$ for all $i \in \mathbb{N}$. Further, there is a G -large ideal I of RK with a free abelian central subgroup A of finite index in G/I^\dagger and $k = R/(R \cap I)$ such that:*

- (i) $(a_i RG)/(a_i RG)I \neq 0$ and $(a_i RG)/(a_i RG)I$ is kA -torsion for all $i \in \mathbb{N}$;
- (ii) $\mu_{kA}((a_i RG)/(a_i RG)I) = \mu_{kA}((a_j RG)/(a_j RG)I)$ for all $i, j \in \mathbb{N}$;
- (iii) $\text{chark} \notin \pi(G_1/G_i)$ for all $i \in \mathbb{N}$.

Proof. As the field F is finitely generated, there is a finitely generated domain R such that F is the field of fractions of R . It follows from [1, Chap.V, §3.4, Corollary 1] that R is an absolutely Hilbert domain.

Let $\{G_i | i \in \mathbb{N}\}$ be a descending chain of subgroups of finite index in G and $\{a_i | i \in \mathbb{N}\} \subseteq M$ be a set which exist by Lemma 10. By Lemma 10(i), the quotient group G/K is free abelian, where $K = \bigcap_{i \in \mathbb{N}} G_i$. By Lemma 10(ii), $a_i FG = a_i FG_i \otimes_{FG_i} FG$ and it implies that $RG = a_i RG_i \otimes_{RG_i} RG$ for all $i \in \mathbb{N}$. By Proposition 2 there is a cyclic uniform RK -submodule $a_0 RK = V \leq M$ which is (π, G) -large critical for any finite set of prime integers $\pi \subseteq \mathbb{N}$.

As the module M is irreducible purely semi-primitive and $h(K) < h(G)$, we see that $M = a_0 FG \neq a_0 FK \otimes_{FK} FG$ and hence $a_0 RG \neq a_0 RK \otimes_{RK} RG$. Then it follows from Lemma 11 that there is a submodule $U \leq a_0 RK$ with the property that when a G -large ideal I of RK culls U in $a_0 RK$ then $(a_0 RG)/(a_0 RG)I \neq 0$ and the kA -module $(a_0 RG)/(a_0 RG)I \neq 0$ is kA -torsion, where $k = R/(R \cap I)$ and A is a free abelian central subgroup of finite index in G/I^\dagger . As the RK -module $a_0 RK$ is (π, G) -large critical, such an ideal I exists.

(i) As $(a_0 RG)/(a_0 RG)I \neq 0$ and $(a_0 RG)/(a_0 RG)I$ is kA -torsion, it follows from Proposition 3(iii, iv) that $(a_i RG)/(a_i RG)I \neq 0$ and $(a_i RG)/(a_i RG)I$ is kA -torsion for all $i \in \mathbb{N}$.

(ii) By Proposition 3(ii), $\mu_{kA}((a_i RG)/(a_i RG)I) = \mu_{kA}((a_j RG)/(a_j RG)I)$ for all $i, j \in \mathbb{N}$.

(iii) By Lemma 10(i), either for any prime integer p there is $n \in \mathbb{N}$ such that $p \notin \pi(G_n/G_i)$ for all $i \geq n$ or G/G_i is a p -group for all $i \in \mathbb{N}$, where p is a prime integer. Since the submodule $a_0 RK$ is (π, G) -large

critical for any finite set of prime integers $\pi \subseteq \mathbb{N}$, we can chose the ideal I such that $\text{char}k \neq p$ if G/G_i is a p -group for all $i \in \mathbb{N}$. Therefore, there is $n \in \mathbb{N}$ such that $\text{char}k \notin \pi(G_n/G_i)$ for all $i \geq n$. Starting the chain $\{G_i | i \in \mathbb{N}\}$ and the set $\{a_i | i \in \mathbb{N}\}$ from the n -th members we obtain the assertion. \square

Lemma 13. *Let F be a finitely generated field of characteristic zero and let G be a finitely generated nilpotent group. Suppose that there exists a purely semi-primitive irreducible FG -module M . Then there exist a finitely generated free abelian group A , a locally finite field k , a descending chain $\{A_i | i \in \mathbb{N}\}$ of subgroups of finite index in A and a set $\{J_i | i \in \mathbb{N}\}$ of ideals of the group algebra kA such that:*

- (i) $\bigcap_{i \in \mathbb{N}} A_i = 1$ and $\text{char}k \notin \pi(A_1/A_i)$ for all $i \in \mathbb{N}$;
- (ii) $J_i = (J_i \cap kA_i)kA$ for all $i \in \mathbb{N}$;
- (iii) $\mu_{kA}(J_i) = \mu_{kA}(J_j)$ and $\mu_{kA}(J_i)$ consists of proper ideals of kA for all $i, j \in \mathbb{N}$.

Proof. Let $R \leq F$ be an absolutely Hilbert domain, $\{G_i | i \in \mathbb{N}\}$ be a descending chain of subgroups of finite index in G and $\{a_i | i \in \mathbb{N}\}$ be a subset of M which exist by Lemma 12. Then the quotient group G/K is free abelian, where $K = \bigcap_{i \in \mathbb{N}} G_i$, and $a_i RG = a_i RG_i \otimes_{RG_i} RG$ for all $i \in \mathbb{N}$.

Further, let I be a G -large ideal I of RK which exists by Lemma 12. Let A be a free abelian central subgroup of finite index in G/I^\dagger and let $k = R/(R \cap I)$. Put $A_i = (G_i/I^\dagger) \cap A$ then A_i is a free abelian group for each $i \in \mathbb{N}$. As $\{G_i | i \in \mathbb{N}\}$ is a descending chain of subgroups of finite index in G , it is easy to note that $\{A_i | i \in \mathbb{N}\}$ is a descending chain of subgroups of finite index in A . Let w_i be the image of a_i in $(a_i RG)/(a_i RG)I$ and put $J_i = \text{Ann}_{kA}(w_i)$.

(i) As A_i is free abelian and K/I^\dagger is finite, we have $A_i \cap (K/I^\dagger) = 1$. Then, as $K/I^\dagger = \bigcap_{i \in \mathbb{N}} (G_i/I^\dagger)$, we can conclude that $\bigcap_{i \in \mathbb{N}} A_i = 1$. By Lemma 12(iii), $\text{char}k \notin \pi(G_1/G_i)$ for all $i \in \mathbb{N}$ and hence $\text{char}k \notin \pi(A_1/A_i)$ for all $i \in \mathbb{N}$.

(ii) As $a_i RG = a_i RG_i \otimes_{RG_i} RG$, it follows from Proposition 4(i) that $J_i = (J_i \cap kA_i)kA$ for all $i \in \mathbb{N}$.

(iii) By Lemma 12(ii), $\mu_{kA}((d_i RG)/(d_i RG)I) = \mu_{kA}((d_j RG)/(d_j RG)I)$ for all $i, j \in \mathbb{N}$. Then it follows from Proposition 4(ii) that $\mu_{kA}(J_i) = \mu_{kA}(J_j)$ for all $i, j \in \mathbb{N}$. By Lemma 12(i), $(d_i RG)/(d_i RG)I \neq 0$ and $(d_i RG)/(d_i RG)I$ is kA -torsion. Therefore, by Proposition 4(iii), $\mu_{kA}(J_i)$ consists of proper ideals of kA for all $i \in \mathbb{N}$. \square

Theorem 1. *Let F be a finitely generated field of characteristic zero and let G be a finitely generated nilpotent group. Let M be an irreducible FG -module. Then there are an element $0 \neq a \in M$ and a subgroup $L \leq G$ such that $aFG = aFL \otimes_{FL} FG$ and aFL is a primitive FL -module.*

Proof. We apply induction on the Hirsch number $h(G)$ of G . If $h(G) = 0$ then G is finite. Therefore, $\dim_F M$ is finite and the assertion follows. Then by the induction hypothesis we can assume that the module M is semi-primitive. Suppose that the assertion does not hold then the module M is purely semi-primitive.

By Lemma 13, there exist a finitely generated free abelian group A , a locally finite field k , a descending chain $\{A_i | i \in \mathbb{N}\}$ of subgroups of finite index in A and a set $\{J_i | i \in \mathbb{N}\}$ of ideals of the group algebra kA such that:

- (i) $\cap_{i \in \mathbb{N}} A_i = 1$ and $\text{char} k \notin \pi(A_1/A_i)$ for all $i \in \mathbb{N}$;
- (ii) $J_i = (J_i \cap kA_i)kA$ for all $i \in \mathbb{N}$;
- (iii) $\mu_{kA}(J_i) = \mu_{kA}(J_j)$ and $\mu_{kA}(J_i)$ consists of proper ideals of kA for all $i, j \in \mathbb{N}$.

As $\mu_{kA}(J_i) = \mu_{kA}(J_j)$ for all $i, j \in \mathbb{N}$, we can put $\mu = \mu_{kA}(J_i)$ for all $i \in \mathbb{N}$.

It follows from (iii) and Lemma 7(iii) that $\mu_{kA_t}(J_i \cap kA_t) = \mu_{kA_t}(J_j \cap kA_t)$ for all $i, j, t \in \mathbb{N}$. Then we can put $\mu_i = \mu_{kA_i}(J_j \cap kA_i)$ for all $i, j \in \mathbb{N}$.

It follows from Lemma 7(i) that for any J_j and any $Q \in \mu_{kA_{i+1}}(J_j \cap kA_{i+1})$ there is $P \in \mu_{kA_i}(J_j \cap kA_i)$ such that $P \cap kA_{i+1} = Q$. It easily implies that $|\mu_i| \geq |\mu_{i+1}|$ for all $i \in \mathbb{N}$. Therefore, there is $n \in \mathbb{N}$ such that $|\mu_n| = |\mu_i|$ for all $i \geq n$.

Then for any $i \geq n$ and any $Q \in \mu_{kA_i}(J_i \cap kA_i) = \mu_i$ there is only one prime $P \in \mu_{kA_n}(J_i \cap kA_n) = \mu_n$ such that $P \cap kA_i = Q$. It follows from (i) that $\text{char} k \notin \pi(A_n/A_i)$ and it follows from (ii) that $J_i \cap kA_n = (J_i \cap kA_i)kA_n$. Then it follows from Lemma 8 that $P = QkA_n = (P \cap kA_i)kA_n$. On the other hand, as $|\mu_n| = |\mu_i|$, for any $P \in \mu_{kA_n}(J_i \cap kA_n) = \mu_n$ we have $P \cap kA_i = Q \in \mu_{kA_i}(J_i \cap kA_i) = \mu_i$. Therefore, for any $i \geq n$ and any $P \in \mu_n$ we have $P = QkA_n = (P \cap kA_i)kA_n$ and, as $\cap_{i \in \mathbb{N}} A_i = 1$, it implies that $P = kA_n$. But, as μ consists of proper ideals of kA , it follows from Lemma 7(iv) that μ_n consists of proper ideals of kA_n and a contradiction is obtained. □

REFERENCES

- [1] Bourbaki, N.: Elements of Mathematics: Commutative Algebra, Chapters 1-7. Springer, London (1998).

- [2] Brookes, C.J.B., Brown, K. A.: Primitive group rings and Noetherian rings of quotients. *Trans. Amer. Math. Soc.* **288**, 605-623 (1985).
- [3] Brookes, C.J.B.: Modules over polycyclic groups. *Proc. London Math. Soc.* **57**, 88-108 (1988)
- [4] Harper, D.L.: Primitive irreducible representations of nilpotent groups. *Math. Proc. Camb. Phil. Soc.* **82**, 241-247 (1977)
- [5] Harper, D.L.: Primitivity in representations of polycyclic groups. *Math. Proc. Camb. Phil. Soc.* **88**, 15-31 (1980)
- [6] Kargapolov, M.I., Merzljakov, Yu.I.: *Fundamentals of the Theory of Groups*, Graduate Texts in Mathematics, Springer, Berlin (1979).
- [7] Karpilovsky, G.: *Induced modules over group algebras*, North-Holland, Amsterdam (1990).
- [8] McConnell, J.C., Robson, J.C.: *Noncommutative Noetherian rings*. Graduate Studies in Mathematics, AMS, Providence (2001).
- [9] Passman, D.S.: *Algebraic crossed products*. *Contemporary Mathematics*, **43**, 209-225 (1985)
- [10] Passman, D.S.: *The algebraic structure of group rings*. Wiley-Interscience, New York (1977).
- [11] Szechtman, F.: Groups having a faithful irreducible representation. *J. Algebra*. **454**, 292-307 (2016)
- [12] Szechtman, F., Tushev, A.: Infinite groups admitting a faithful irreducible representation. *J. Algebra Appl.* **17**, 1850005-1 - 1850005-7 (2017)
- [13] Tushev, A.V.: Irreducible representations of locally polycyclic groups over an absolute field, *Ukrainian Math. J.* **42**, 1233-1238 (1990)
- [14] Tushev, A.V.: On exact irreducible representations of locally normal groups, *Ukrainian Mat. J.* **45**, 1900-1906 (1993)
- [15] Tushev, A.V.: On the primitive representations of soluble groups of finite rank, *Sb. Math.* **191**, 117-159 (2000)
- [16] Tushev, A.V.: On primitive representations of minimax nilpotent groups, *Mathematical Notes* **72**, 117-128 (2002)
- [17] Tushev, A.V.: On the irreducible representations of soluble groups of finite rank, *Asian-European J. Math.* **5**, 1250061-1-1250061-12 (2012)
- [18] Tushev, A.V.: On the primitive irreducible representations of finitely generated nilpotent groups, *Dopov. Nac. akad. nauk Ukr.* **4**, 24-27 (2021)
- [19] Wehrfritz, B.A.F.: *Group and Ring Theoretic Properties of Polycyclic Groups*. Algebra and Applications, vol. 10, Springer, London (2009)
- [20] Zariski, O., Samuel, P.: *Commutative Algebra*. Vol.1. Van Nostrand, Princeton (1958)