

Sprays on Hamel-Funk Functions Model

Guojun Yang

Abstract

Hamel functions of a spray play an important role in the study of the projective metrization of the concerned spray, and Funk functions are special Hamel functions. A Finsler metric is a special Hamel function of the spray induced by the metric itself and a Funk metric is a special Funk function of a Minkowski spray. In this paper, we study sprays on a Hamel or Funk function model. Firstly, we give some basic properties of a Hamel or Funk function of a spray and some curvature properties of a Hamel or Funk function in projective relations. We use the Funk metric to construct a family of sprays and obtain some of their curvature properties and their metrization conditions. Secondly, we consider the existence of Funk functions on certain spray manifold. We prove that there exist local Funk functions on a R-flat spray manifold, and on certain projectively flat Berwald spray manifolds, we construct a multitude of nonzero Funk functions. Finally, we introduce a new class of sprays called Hamel or Funk sprays associated to given sprays and Hamel or Funk functions. We obtain some special properties of a Hamel or Funk spray of scalar curvature, especially on its metrization and a special form of its Riemann curvature.

Keywords: Spray, Finsler Metric, Metrization, Scalar (Isotropic, Constant) Curvature, Hamel-Funk Function, Hamel-Funk Spray

2010 Mathematics Subject Classification: 53C60, 53B40

1 Introduction

Spray geometry studies the properties of sprays on a manifold, and it is closely related to Finsler geometry. Every Finsler metric induces a natural spray but there are a lot of sprays which cannot be induced by any Finsler metric ([4, 7, 8, 16, 17]). A spray \mathbf{G} on a manifold M defines a special vector field on a conical region \mathcal{C} of $TM \setminus \{0\}$, and it naturally defines the integral curves and the projections of the integral curves onto the manifold M are called geodesics. For deep understanding of spray geometry, it is important to develop more basic curvatures, to study some classes of sprays with special geometric properties and to investigate the (projective) metrization of a spray under certain curvature conditions.

Many basic curvatures, such as Riemann curvature, Ricci curvature, Weyl curvature, Berwald curvature and Douglas curvature, appearing in Finsler geometry, are actually defined in spray geometry via the spray coefficients. With a fixed volume form, the S-curvature, originally introduced in Finsler geometry by Z. Shen in [12], can also be generalized to spray geometry. The notions for a Finsler metric of scalar flag curvature, isotropic flag curvature and constant flag curvature, already have the corresponding generalization in spray geometry, and they are respectively called a spray of scalar curvature, isotropic curvature and constant curvature. A spray of scalar curvature first appears in Shen's book [14] (therein, it is originally called an isotropic spray). In [9], B. Li and Z. Shen introduce a spray of isotropic curvature. The present author introduces a spray of constant curvature in [17]. Further studies show that these curvatures or curvature characters play an important role in spray geometry.

Among all sprays, some special classes of sprays are helpful in understanding the geometric structures of a general spray ([4, 8, 17]). A Berwald spray means that its spray coefficients are

polynomials and every Berwald metric induces a Berwald spray. A (locally) projectively flat spray means its geodesics are (local) straight lines, and it's the Hilbert's Fourth Problem to study the structure of a (locally) projectively flat spray, especially when the spray is induced by a Finsler metric. These sprays and sprays of scalar curvature (including isotropic, constant curvature) have some special good properties and deserve further studies.

The metrizable problem for a spray \mathbf{G} seeks for a Finsler metric whose spray is just \mathbf{G} , and the projective metrizable of a spray \mathbf{G} aims to look for a Finsler metric projectively related to \mathbf{G} . So a natural question is to determine whether a given spray is (projectively) Finsler-metrizable or not under certain curvature conditions. Quite a few papers concentrate on this problem ([3, 4, 6, 7, 8, 10, 11, 16, 17]). For instance, any spray of scalar curvature is locally projectively metrizable ([10, 3]); the metrizable for a spray of scalar curvature with nonzero Ricci curvature is solved ([4]); the local structure is determined for a projectively flat Berwald spray of isotropic curvature to be metrizable ([17]).

In this paper, we are going to study some geometric properties of Hamel-Funk functions and Hamel-Funk sprays. The notion of Funk function first appears in Shen's book [14]. Hamel-Funk sprays are a new class of sprays we introduce in this paper.

Let \mathbf{G} be a spray on a manifold M . We can use the Berwald connection of \mathbf{G} to define a Hamel function or a Funk function of \mathbf{G} (Definition 3.1). When a Finsler metric F is projectively flat, F is a Hamel function of the spray $\mathbf{G} = 0$, which satisfies the well-known Euler-Lagrange equations. When F is a Funk metric on a convex domain in R^n , F is a Funk function of the spray $\mathbf{G} = 0$. In Sec. 3, we introduce some basic properties of a Hamel or Funk function of a Minkowski spray or a general spray, and generalize some known equations in [13] for projectively flat Finsler metrics of constant flag curvature to Hamel or Funk functions of a spray (Lemmas 3.7–3.10). We see that a Funk function is a Hamel function and the collection of all Hamel functions forms a linear space (Lemma 3.5).

A Hamel or Funk function plays a special role in projective relations. A simple fact shows that a Hamel function is projectively invariant, but a Funk function is not (Lemmas 4.1, 4.2). Z. Shen proves that The Riemann curvatures of two projectively related sprays keep invariant if the projective factor is a Funk function ([14]). Further, we prove that if one of two projectively related sprays is of isotropic curvature, then the other is also of isotropic curvature if and only if the projective factor is a Hamel function (Proposition 4.4(iii)). Starting from a Funk metric, we obtain the following result.

Theorem 1.1 *Let \mathbf{G} be the spray of the Funk metric F on a convex domain $\Omega \subset R^n$. For a constant c , define a spray*

$$\bar{G}^i(y) := G^i(y) + cF(-y)y^i. \quad (1)$$

- (i) $\bar{\mathbf{G}}$ is of isotropic curvature. Further, $\bar{\mathbf{G}}$ is of constant curvature if and only if $c = 0$ or $c = -1/2$.
- (ii) $\bar{\mathbf{G}}$ is (locally) metrizable if and only if $c = 0$ or $c = -1/2$. In this case, $\bar{\mathbf{G}}$ is induced by F if $c = 0$; and if $c = -1/2$, $\bar{\mathbf{G}}$ is induced by the Klein metric

$$\bar{F}(y) := [F(y) + F(-y)]/2. \quad (2)$$

Theorem 1.1(i) gives a new class of examples to show that a spray of isotropic curvature is not necessarily of constant curvature even in dimension greater than two (cf. [17]). Theorem 1.1(ii) also gives a class of non-metrizable sprays. For $\bar{\mathbf{G}}$ in Theorem 1.1, if $F(-y)$ is replaced by $F(y)$, the result refers to [16].

By some properties of a Hamel function, it is relatively free to construct Hamel functions of certain sprays. For instance, a Finsler metric is a Hamle function of its induced spray, and Propositions 5.2 and 5.3 also give some Hamel functions of certain sprays. However, the

construction of a Funk function of a spray is not easy. On a Minkowski spray $\mathbf{G} = 0$, we can get many Funk functions (a Funk metric itself is). In [14], Z. Shen proves that no nonzero Funk function exists on a compact regular spray manifold (Theorem 5.4). In [2], I. Bucataru and Z. Muzsnay show that no nonzero Funk function exists on a Finsler manifold of nonzero scalar curvature (Theorem 5.6). To construct nontrivial Funk functions of a spray, we first prove Proposition 5.5 on some properties satisfied by a Funk function of a spray of scalar or isotropic or constant curvature, and then we give many nontrivial closed 1-form Funk functions of projectively flat Berwald sprays in Examples 5.7–5.9. Further we have the following local existence theorem.

Theorem 1.2 *There exist non-trivial (local) Funk functions on a R-flat spray manifold.*

A locally Minkowski spray is R-flat and there are many Funk functions of a Minkowski spray (see Sec. 3). If a spray is just R-flat (not Berwaldian), we generally don't have the explicit expression of the Funk function in Theorem 1.2.

Using a spray \mathbf{G} and a nonzero homogeneous function Q of degree one, we define a spray $\bar{\mathbf{G}}$ by

$$\bar{G}^i := G^i + \frac{Q_{;0}}{2Q} y^i. \quad (3)$$

Every spray $\bar{\mathbf{G}}$ in the projective class of \mathbf{G} can be locally written in the form (3) for some Q (Lemma 6.1), and it is projectively invariant for a fixed Q (Lemma 6.2). When Q is a Hamel or Funk function, we obtain a Hamel or Funk spray $\bar{\mathbf{G}}$ associated to (\mathbf{G}, Q) , with \mathbf{G}, Q being the adjoint spray and the adjoint Hamel function respectively (Definition 6.3). Hamel functions and Hamel sprays play an important role in studying the projective metrizability of a given spray. We have the following theorem.

Theorem 1.3 *A spray \mathbf{G} is projectively metrizable if and only if it has a Finsler-Hamel function \bar{F} . In this case, the spray \mathbf{G} is projectively induced by the Finsler metric \bar{F} whose spray is the Hamel spray associated to (\mathbf{G}, \bar{F}) .*

For the projective metrizability of a spray, a different description from Theorem 1.3 is given in [3]. In Theorem 1.3, if the Hamel function is a Finsler metric, then the Hamel spray is a Finsler spray. As we know, for a Finsler spray of scalar curvature, its Riemann curvature has a special form (7). This form is inherited well by the Riemann curvature of a Hamel spray of scalar curvature in a similar way.

Theorem 1.4 (i) *A Hamel spray $\bar{\mathbf{G}}$ is of scalar curvature if and only if its Riemann curvature has the form*

$$\bar{R}^i_{\ k} = \lambda(Q^2 \delta_k^i - QQ_{.k} y^i), \quad (4)$$

where $\lambda = \lambda(x, y)$ is a scalar function and Q is the adjoint Hamel function of $\bar{\mathbf{G}}$.

(ii) *A Funk spray $\bar{\mathbf{G}}$ is of isotropic curvature if and only if its Riemann curvature has the form (4), where $\lambda = \lambda(x)$ is a scalar function and Q is the adjoint Funk function of $\bar{\mathbf{G}}$.*

The scalar function λ in Theorem 1.4 is determined by (55) and (56) respectively, and λ can be a constant for some special cases (cf. Examples 6.10 and 6.11).

A given Hamel spray is possibly associated to (\mathbf{G}, Q_1) and (\mathbf{G}, Q_2) with different Hamel functions Q_1, Q_2 of \mathbf{G} . But under certain condition, a given Hamel spray corresponds a unique Hamel function (up to a constant scaling) (cf. Proposition 6.13). On the metrizability of a Hamel spray, we have the following theorem.

Theorem 1.5 *A Hamel spray of nonzero scalar curvature is metrizable if and only if its adjoint Hamel function is a Finsler metric. In this case, the Hamel spray is uniquely induced by its adjoint Hamel function (the metric is unique up to a constant scaling).*

Theorem 1.5 is not true if the Hamel spray is of zero scalar curvature or not of scalar curvature. A Minkowski spray $\bar{\mathbf{G}}$ is a natural Hamel spray of zero scalar curvature, but any two Minkowski metrics induce $\bar{\mathbf{G}}$. Any Finsler spray $\bar{\mathbf{G}}$ (induced by a Finsler metric \bar{F}) is also a natural Hamel spray associated to $(\bar{\mathbf{G}}, \bar{F})$. If $\bar{\mathbf{G}}$ is not of scalar curvature, we can find many Finsler metrics inducing the same $\bar{\mathbf{G}}$ but not having constant ratio (see [5]: Chapter 9).

Starting from a spray \mathbf{G} of scalar curvature and a closed 1-form, we can obtain many Hamel sprays which are not metrizable (cf. Examples 6.10 and 6.11, Corollary 6.15).

2 Preliminaries

Let M be an n -dimensional manifold. A conical region $\mathcal{C} = \mathcal{C}(M)$ of $TM \setminus \{0\}$ means $\mathcal{C}_x := \mathcal{C} \cap T_x M \setminus \{0\}$ is conical region for $x \in M$ ($\lambda y \in \mathcal{C}_x$ if $\lambda > 0, y \in \mathcal{C}_x$). A *spray* on M is a smooth vector field \mathbf{G} on a conical region \mathcal{C} of $TM \setminus \{0\}$ expressed in a local coordinate system (x^i, y^i) in TM as follows

$$\mathbf{G} = y^i \frac{\partial}{\partial x^i} - 2G^i \frac{\partial}{\partial y^i},$$

where $G^i = G^i(x, y)$ are local functions which are positively homogeneous with degree two. If \mathbf{G} is defined on the whole $TM \setminus \{0\}$, we call \mathbf{G} a regular spray.

The Riemann curvature tensor R^i_k of a given spray G^i is defined by

$$R^i_k := 2\partial_k G^i - y^j (\partial_j \dot{\partial}_k G^i) + 2G^j (\dot{\partial}_j \dot{\partial}_k G^i) - (\dot{\partial}_j G^i) (\dot{\partial}_k G^j), \quad (5)$$

where we define $\partial_k := \partial/\partial x^k, \dot{\partial}_k := \partial/\partial y^k$. The trace of R^i_k is called the Ricci curvature, $Ric := R^i_i$. A spray \mathbf{G} is said to be *R-flat* if $R^i_k = 0$. A spray \mathbf{G} is said to be of *scalar curvature* if its Riemann curvature R^i_k satisfies

$$R^i_k = R\delta_k^i - \tau_k y^i, \quad (6)$$

where $R = R(x, y)$ and $\tau_k = \tau_k(x, y)$ are some homogeneous functions ([14]). If in (6) there holds $R_{,i} = 2\tau_i$, then \mathbf{G} is said to be of *isotropic curvature* ([9]). A spray \mathbf{G} is said to be of *constant curvature* if \mathbf{G} satisfies (6) with ([17])

$$\tau_{i;k} = 0 \quad (\Leftrightarrow R = \tau_k = 0, \text{ or } R_{,i} = 0).$$

The three notions are natural generalizations for a Finsler metric of scalar/isotropic/constant flag curvature in Finsler geometry.

A spray \mathbf{G} is called a Berwald spray if its Berwald curvature vanishes $G^i_{hjk} := \dot{\partial}_h \dot{\partial}_j \dot{\partial}_k G^i = 0$. If G^i is locally given by $G^i = 0$, we call \mathbf{G} a locally Minkowski spray. Two sprays \mathbf{G} and $\bar{\mathbf{G}}$ on M are said to be projectively related if their geodesics coincide as set points, or equivalently, $\bar{G}^i = G^i + P y^i$ for some positively homogeneous function P of degree one. A spray \mathbf{G} is said to be locally projectively flat if locally G^i can be expressed as $G^i = P y^i$. A spray \mathbf{G} is called *reversible* if the spray $G^i(-y)$ is projectively related to $G^i(y)$, and *strictly reversible* if $G^i(y) = G^i(-y)$. We use $Proj(G)$ to denote the projective class of \mathbf{G} .

In the calculation of some geometric quantities of a spray, it is very convenient to use Berwald connection as a tool. For a spray manifold (\mathbf{G}, M) , Berwald connection is usually defined as a linear connection on the pull-back $\pi^* TM$ ($\pi : TM \rightarrow M$ the natural projection) over the base manifold M . The Berwald connection is defined by

$$D(\partial_i) = (G^k_{ir} dx^r) \partial_k, \quad (G^k_{ir} := \dot{\partial}_r \dot{\partial}_i G^k),$$

For a spray tensor $T = T_i dx^i$ as an example, the horizontal and vertical derivatives of T with respect to Berwald connection are given by

$$T_{i;j} = \delta_j T_i - T_r G^r_{ij}, \quad T_{i \cdot j} = \dot{\partial}_j T_i, \quad (\delta_i := \partial_i - G^r_i \dot{\partial}_r).$$

For the Ricci identities and Bianchi identities of Berwald connection, readers can refer to [1].

In this paper, we define a Finsler metric $L(\neq 0)$ on a manifold M as follows (cf. [14]): (i) for any $x \in M$, L_x is defined on a conical region of $T_x M \setminus \{0\}$ and L is C^∞ ; (ii) L is positively homogeneous of degree two; (iii) the fundamental metric tensor $g_{ij} := (\frac{1}{2}L)_{y^i y^j}$ is non-degenerate. A Finsler metric L is said to be regular if additionally L is defined on the whole $TM \setminus \{0\}$ and (g_{ij}) is positively definite. Otherwise, L is called singular. In general case, we don't require that L be regular. If a Finsler metric $L > 0$, we put $L = F^2$, and in this case, F is also called a Finsler metric being positively homogeneous of degree one.

Any Finsler metric L induces a natural spray whose coefficients G^i are given by

$$G^i := \frac{1}{4}g^{il}\{L_{x^k y^l} y^k - L_{x^l}\},$$

where (g^{ij}) is the inverse of (g_{ij}) . L is said to be of *scalar flag curvature* $K = K(x, y)$ if

$$R^i_k = K(L\delta_k^i - y^i y_k) = K(F^2\delta_k^i - FF_{.k}y^i), \quad (L = F^2), \quad (7)$$

where $y_k := (L/2)_{.k} = g_{km}y^m$. If $K_{.i} = 0$, then L is said to be of *isotropic flag curvature*, and in this case, K is a constant if the dimension $n \geq 3$ (Schur's Theorem). L is said to be of *constant flag curvature* if K is constant.

A spray \mathbf{G} is (globally) Finsler-metrizable on M (or on $\mathcal{C}(M)$) if there is a Finsler metric L defined on a conical region $\mathcal{C}(M)$ and L induces \mathbf{G} . A spray \mathbf{G} is locally Finsler-metrizable on M if for each $x \in M$, there is a neighborhood U of x such that \mathbf{G} is Finsler-metrizable on U .

Lemma 2.1 *A Finsler spray $\bar{\mathbf{G}}$ and a spray \mathbf{G} are related by*

$$\bar{G}^i = G^i + \frac{\bar{F}_{;0}}{2\bar{F}}y^i - \frac{1}{2}\bar{F}\bar{g}^{ir}(\bar{F}_{.r} - \bar{F}_{.r;0}), \quad (8)$$

where \bar{F} is a Finsler metric inducing $\bar{\mathbf{G}}$.

Lemma 2.2 *A Finsler spray $\bar{\mathbf{G}}$ (induced by \bar{F}) is projective to a spray \mathbf{G} if and only if \bar{F} satisfies one of the following conditions:*

$$(i) \bar{F}_{.i} = (P\bar{F})_{.i}, \quad (ii) \bar{F}_{.i} = \bar{F}_{.i;0}, \quad (iii) \bar{F}_{.i} = \frac{1}{2}\bar{F}_{;0.i}, \quad (9)$$

where in case (i), P is a homogeneous function of degree one. In this case, P is the projective factor which is given by

$$P = \frac{\bar{F}_{;0}}{2\bar{F}}. \quad (10)$$

Lemma 2.3 *Let $\bar{\mathbf{G}}$ and \mathbf{G} be projectively related with $\bar{G}^i = G^i + Py^i$. Then their Riemann curvatures and Ricci curvatures are related by*

$$\bar{R}^i_k = R^i_k + (3Q_k - Q_{0.k})y^i - Q_0\delta_k^i, \quad (11)$$

$$\bar{Ric} = Ric - (n-1)Q_0 = Ric - (n-1)(P_{;0} - P^2), \quad (12)$$

$$(Q_j := P_{.j} - PP_j, \quad Q_{jk} := P_{j;k} - P_{k;j} = Q_{k.j} - Q_{j.k}).$$

The above three lemmas can be found in some textbooks. For the proofs, readers can refer to Chapter 3 of [1], for instance.

Lemma 2.4 ([17]) *Let T be a positively homogeneous function of degree zero on a spray manifold of scalar curvature with nonzero Ricci curvature. If $T_{.i} = 0$, then T is constant.*

Lemma 2.5 *On a pray manifold (M, \mathbf{G}) , put*

$$T_i := Q_{;r.i}y^r - Q_{;i} (= Q_{.i;0} - Q_{;i}), \quad (13)$$

where Q is an arbitrary positively homogeneous function of degree one. Then T_i is a projective invariant.

Proof : For two projectively related $\bar{\mathbf{G}}$ and \mathbf{G} , let $\bar{G}^i = G^i + Py^i$. A direct computation gives

$$Q_{\bar{;i}} = Q_{;i} - (PQ)_{.i}, \quad Q_{\bar{;r.i}} = Q_{;r.i} - (PQ)_{.i.r}.$$

Therefore, it is easy to get

$$Q_{\bar{;r.i}}y^r - Q_{\bar{;i}} = Q_{;r.i}y^r - Q_{;i},$$

which shows that T_i is projectively invariant. Q.E.D.

3 Hamel-Funk Functions and Basic Properties

In this section, we introduce the definitions of a Hamel function and a Funk function, and then study some of their basic properties on a spray manifold.

Definition 3.1 *Let (M, \mathbf{G}) be a spray manifold. A positively homogeneous function P of degree one is called a Hamel Function (resp. Funk function, weak Funk function) of \mathbf{G} if P satisfies*

$$P_{;k} = P_{k;0} \text{ (resp. } P_{;k} = PP_k, P_{;0} = P^2), \quad (P_k := P_k).$$

Z. Shen introduces Funk function and weak Funk function of a spray in [14].

3.1 Hamel-Funk Functions of a Minkowski Spray

In this subsection, we show some properties of a Hamel-Funk function of the Minkowski spray $\mathbf{G} = 0$ on R^n (cf. [5, 9, 13]).

Lemma 3.2 *Let P be a Funk function of the spray $\mathbf{G} = 0$ on R^n . Then we have*

$$P_{x^{i_1} \dots x^{i_m}} = \frac{1}{m+1} (P^{m+1})_{y^{i_1} \dots y^{i_m}}, \quad (14)$$

$$Q_{x^k} = (PQ)_{y^k} \implies Q_{x^{i_1} \dots x^{i_m}} = (QP^m)_{y^{i_1} \dots y^{i_m}}, \quad (15)$$

where in (15), Q is some function on TR^n .

Lemma 3.3 *Let P be a Funk function of the spray $\mathbf{G} = 0$ on R^n and $k \geq 0$ be an integer. Define*

$$Q := (P^{k+1})_{y^{i_1} \dots y^{i_k}} x^{i_1} \dots x^{i_k} \quad \left(= (k+1)P_{x^{i_1} \dots x^{i_k}} x^{i_1} \dots x^{i_k} \right).$$

Then Q is a Hamel function of \mathbf{G} .

Proof : This proof refers to [9]. By $P_{x^k} = PP_{y^k}$, we have

$$\begin{aligned} Q_{x^j} &= (k+1)(P^k P_{x^j})_{y^{i_1} \dots y^{i_k}} x^{i_1} \dots x^{i_k} + k(P^{k+1})_{y^{i_1} \dots y^{i_{k-1}} y^j} x^{i_1} \dots x^{i_{k-1}} \\ &= \frac{k+1}{k+2} (P^{k+2})_{y^{i_1} \dots y^{i_k} y^j} x^{i_1} \dots x^{i_k} + k(P^{k+1})_{y^{i_1} \dots y^{i_{k-1}} y^j} x^{i_1} \dots x^{i_{k-1}}. \end{aligned}$$

Then we easily obtain

$$Q_{x^j} = Q_{x^m y^j} y^m,$$

which shows that Q is a Hamel function of \mathbf{G} . Q.E.D.

Lemma 3.4 ([5]) *Let $\varphi = \varphi(y)$ be a positively homogeneous function of degree one on R^n and be C^∞ on $R^n \setminus \{0\}$. Then there is a neighborhood U of $0 \in R^n$, and a unique Funk function $P = P(x, y)$ of the spray $\mathbf{G} = 0$ such that for $x \in U$, P satisfies*

$$P(x, y) = \varphi(y + P(x, y)x). \quad (16)$$

In this case, we have $P(0, y) = \varphi(y)$.

By Lemma 3.4, we can construct many Funk functions of the spray $\mathbf{G} = 0$ by solving P from (16) for a given φ . For example, let $\varphi(y) = |y|$ and $\varphi(y) = -\langle a, y \rangle$ respectively, where a is a constant vector. Then solving P from (16) gives respectively

$$P = \frac{\sqrt{(1 - |x|^2)|y|^2 + \langle x, y \rangle^2} + \langle x, y \rangle}{1 - |x|^2}, \quad P = -\frac{\langle a, y \rangle}{1 + \langle a, x \rangle}. \quad (17)$$

3.2 Hamel-Funk Functions of a General Spray

Lemma 3.5 *A Hamel function or a Funk function has the following basic properties:*

- (i) *For a spray, any Funk function is a Hamel function.*
- (ii) *The collection of Hamel functions on a spray manifold is a vector space under the natural addition and number-product for functions. So any linear combination of any two Funk functions is a Hamel function.*

Proof : For case (i), if P is a Funk function of a spray, then we have $P_{;k} = PP_k$, which gives $P_{;0} = P^2$. So we have $P_{;0,k} = 2PP_k$, that is, $P_{;k} + P_{k;0} = 2PP_k$. Thus we obtain $P_{;k} = P_{k;0}$ ($= PP_k$). This means that P is a Hamel function. For case (ii), the former is obvious by the definition of a Hamel function and the latter follows from case (i). Q.E.D.

Lemma 3.6 *Let \mathbf{G} be a spray and Q is a Hamel function of \mathbf{G} . Put $P(y) := Q(-y)$. Then P is a Hamel function of \mathbf{G} if and only if*

$$[G^r(y) - G^r(-y)]Q_{.r.i}(-y) = 0. \quad (18)$$

In particular, (18) holds if \mathbf{G} is locally projectively flat or reversible.

Proof : We first have

$$\begin{aligned} P_{;i}(y) &= (\partial_i P - G_i^r P_{.r})(y) = \partial_i Q(-y) + G_i^r(y) Q_{.r}(-y) \\ &= Q_{;i}(-y) + [G_i^r(y) + G_i^r(-y)] Q_{.r}(-y). \end{aligned} \quad (19)$$

Next we get

$$P_{;m.i}(y) = -Q_{.i;m}(-y) + [G_{mi}^r(y) - G_{mi}^r(-y)] Q_{.r}(-y) - [G_m^r(y) + G_m^r(-y)] Q_{.r.i}(-y).$$

Therefore, we obtain

$$P_{;i,0}(y) = Q_{.i;0}(-y) + [G_i^r(y) + G_i^r(-y)] Q_{.r}(-y) - 2[G^r(y) - G^r(-y)] Q_{.r.i}(-y).$$

Since Q is a Hamel function, we have

$$(P_{;i} - P_{;i,0})(y) = 2[G^r(y) - G^r(-y)] Q_{.r.i}(-y),$$

which shows that P is a Hamel function of \mathbf{G} if (18) holds. Q.E.D.

Lemma 3.6 shows that the Klein metric in (2) is a Hamel function and it is projectively flat.

In [13], Z. Shen gives some equations satisfied by a projectively flat Finsler metric of constant flag curvature and its corresponding projective factor. These properties can be generalized to a general spray space satisfied by a Hamel or Funk function (see Lemmas 3.7–3.10).

Lemma 3.7 Let Q be a Funk function of a spray \mathbf{G} and put $\tilde{Q}(y) := [Q(y) + Q(-y)]/2$. Then Q satisfies

$$P := \frac{Q_{;0}}{2Q} = \frac{1}{2}Q, \quad \frac{P^2 - P_{;0}}{Q^2} = -\frac{1}{4}. \quad (20)$$

If \mathbf{G} is strictly reversible, then \tilde{Q} is a Hamel function of \mathbf{G} and satisfies

$$\tilde{P} := \frac{\tilde{Q}_{;0}}{2\tilde{Q}} = \frac{1}{2}[Q(y) - Q(-y)], \quad \frac{\tilde{P}^2 - \tilde{P}_{;0}}{\tilde{Q}^2} = -1. \quad (21)$$

Proof : We only prove (21). Since \mathbf{G} is strictly reversible, we have $G^i(y) = G^i(-y)$. Then we get $G_i^T(-y) = -G_i^T(y)$. By the calculation of (19) and the definition of a Funk function, we have

$$\begin{aligned} \tilde{P} &= \frac{1}{2} \frac{Q_{;0}(y) - Q_{;0}(-y)}{Q(y) + Q(-y)} = \frac{1}{2} \frac{Q^2(y) - Q^2(-y)}{Q(y) + Q(-y)} = \frac{1}{2} [Q(y) - Q(-y)], \\ \tilde{P}^2 - \tilde{P}_{;0} &= \frac{1}{4} [Q(y) - Q(-y)]^2 - \frac{1}{2} [Q^2(y) + Q^2(-y)] = -\tilde{Q}^2. \end{aligned}$$

By Lemma 3.6, $Q(-y)$ is a Hamel function of \mathbf{G} . Thus \tilde{Q} is a Hamel function of \mathbf{G} by Lemma 3.5. This gives the proof. Q.E.D.

Lemma 3.8 Let Q and \bar{Q} be two Funk functions of a spray \mathbf{G} and put $\tilde{Q} := (Q - \bar{Q})/2$. Then \tilde{Q} is a Hamel function of \mathbf{G} and satisfies

$$\tilde{P} := \frac{\tilde{Q}_{;0}}{2\tilde{Q}} = \frac{1}{2}(Q + \bar{Q}), \quad \frac{\tilde{P}^2 - \tilde{P}_{;0}}{\tilde{Q}^2} = -1. \quad (22)$$

It is easy to verify that a Funk function Q of a spray \mathbf{G} satisfies

$$Q_{;i,j} = Q_{;j,i}. \quad (23)$$

Lemma 3.9 Let P, Q be two positively homogeneous functions of degree one satisfying

$$Q_{;i} = (PQ)_{;i} \quad (24)$$

with respect to a spray \mathbf{G} . Then Q is a Hamel function of \mathbf{G} . Further, if $\mathbf{G} = 0$ and P is a Funk function of \mathbf{G} , then in a neighborhood of $0 \in R^n$, Q can be expressed as

$$Q(x, y) = \psi \left(y + P(x, y)x \right) \left(1 + P_{y^m}(x, y)x^m \right), \quad (25)$$

where ψ is defined by $\psi(y) := Q(0, y)$.

Proof : It is easy to prove that Q is a Hamel function of \mathbf{G} by (24). Now let $\mathbf{G} = 0$ and P be a Funk function of \mathbf{G} satisfying (24), that is, $Q_{x^k} = (PQ)_{y^k}$. Then by (15) we have

$$Q_{x^{i_1} \dots x^{i_m}}(0, y) = (\psi \varphi^m)_{y^{i_1} \dots y^{i_m}}(y), \quad (\varphi(y) := P(0, y)). \quad (26)$$

On the other hand, if Q is given by (25), then using $P_{x^k} = PP_{y^k}$ and $P_{x^k y^m} = P_{x^m y^k}$ (see (23)), we can directly verify $Q_{x^k} = (PQ)_{y^k}$, which gives (26) again. So by the uniqueness of expansion, Q is determined by (25) in a neighborhood of $0 \in R^n$. Q.E.D.

Lemma 3.10 Let $H = P + iQ$ be a complex-valued Funk function of a spray \mathbf{G} . Then P, Q are related by

$$Q_{;k} = (PQ)_{;k}, \quad P_{;k} = PP_{;k} - QQ_{;k}. \quad (27)$$

So Q is a Hamel function.

Let $\phi = \phi(y)$ be a complex-valued homogeneous function of degree one on R^n . If a complex-valued function $H = H(x, y)$ satisfies

$$H(x, y) = \phi(y + H(x, y)x), \quad (28)$$

then H is a Funk function, $H_{x^k} = HH_{y^k}$, of $\mathbf{G} = 0$ on R^n (see [13]).

Example 3.11 Let $\phi(y) = i|y|$ in (28), that is,

$$H = i\sqrt{|y|^2 + 2\langle x, y \rangle H + |x|^2 H^2}.$$

Solving this equation yields $H = P + iQ$ satisfying (27) with

$$P = -\frac{\langle x, y \rangle}{1 + |x|^2}, \quad Q = \frac{\sqrt{(1 + |x|^2)|y|^2 - \langle x, y \rangle^2}}{1 + |x|^2}.$$

4 Hamel-Funk Functions in Projective Relations

Now we investigate some projective properties related to a Hamel or Funk function.

Lemma 4.1 *A Hamel function of a spray is projectively invariant.*

Lemma 4.1 directly follows from Lemma 2.5. A Funk function is not projectively invariant, as shown in the following lemma.

Lemma 4.2 *If two projectively related sprays have a same nonzero Funk function, then the two sprays are equal.*

Proof : Let $\bar{\mathbf{G}}$ and \mathbf{G} be related by $\bar{G}^i = G^i + Py^i$. For a positively homogeneous function $Q \neq 0$ of degree one, we have

$$Q_{\bar{y}^k} = Q_{y^k} - (PQ)_{,k}, \quad (29)$$

which implies $PQ = 0$ if Q is a Funk function of \mathbf{G} and also of $\bar{\mathbf{G}}$.

Q.E.D.

Lemma 4.3 *For $\bar{\mathbf{G}} \in Proj(\mathbf{G})$ with $\bar{G}^i = G^i + Py^i$, P is a Funk function of \mathbf{G} if and only if $P_{\bar{y}^k} = -PP_{,k}$, and P is a weak Funk function of \mathbf{G} if and only if $P_{\bar{y}^0} = -P^2$.*

Lemma 4.3 is a direct result of (29) by taking $Q = P$. Now we consider some curvature properties when the projective factor is a Hamel or Funk function.

Proposition 4.4 *Let $\bar{\mathbf{G}}$ and \mathbf{G} be projectively related with the projective factor P .*

- (i) ([14]) $\bar{R}^i_{\bar{k}} = R^i_k$ if and only if P is a Funk function of \mathbf{G} .
- (ii) ([14]) $\bar{Ric} = Ric$ if and only if P is a weak Funk function of \mathbf{G} .
- (iii) If \mathbf{G} is of isotropic curvature, then $\bar{\mathbf{G}}$ is of isotropic curvature if and only if P is a Hamel function of \mathbf{G} .

Proof : Cases (i) and (ii) directly follow from (11) and (12) respectively. For case (iii), it follows from (11) that the Riemann curvature tensor $\bar{R}^i_{\bar{k}}$ of $\bar{\mathbf{G}}$ is given by

$$\begin{aligned} \bar{R}^i_{\bar{k}} &= \bar{R}\delta_k^i - \bar{\tau}_k y^i, \\ (\bar{R} := R + P^2 - P_{,0}, \quad \bar{\tau}_k := \tau_k + PP_{,k} + P_{k;0} - 2P_{;k}), \end{aligned} \quad (30)$$

where we have put

$$\bar{G}^i = G^i + Py^i, \quad R^i_k = R\delta_k^i - \tau_k y^i.$$

Now by $R_{;i} = 2\tau_i$, it can be directly verified that

$$\bar{R}_{;k} = 2\bar{\tau}_k \iff P_{;k;0} = P_{;k}.$$

This completes the proof of Case (iii).

Q.E.D.

Proposition 4.5 *Let \mathbf{G} be a spray of isotropic curvature, and $\bar{\mathbf{G}}$ be projectively related to \mathbf{G} with the projective factor P defined by $P(y) := Q(-y)$ for a Hamel function Q of \mathbf{G} . Then $\bar{\mathbf{G}}$ is of isotropic curvature if and only if (18) holds.*

Proof : It directly follows from Proposition 4.4 (iii) and Lemma 3.6.

Q.E.D.

A Funk metric on a convex domain of R^n has a lot of special curvature properties, and a Klein metric is produce by a Funk metric. Using the Finsler spray induced by a Funk metric, we define a family of sprays given by (1), and we obtain Theorem 1.1 on some properties of the spray (1). Now we give its proof.

Proof of Theorem 1.1: Since

$$\bar{G}^i = G^i + Py^i, \quad G^i = \frac{1}{2}Fy^i, \quad (P(y) := cF(-y)),$$

by (19) and $F_{;i} = 0$, we have

$$\begin{aligned} P_{;i}(y) &= c[G^r_i(y) + G^r_i(-y)]F_{;r}(-y) \\ &= -\frac{1}{2}cF(-y)F_{;i}(y) + c\left[\frac{1}{2}F(y) + F(-y)\right]F_{;i}(-y). \end{aligned} \quad (31)$$

(i) Since \mathbf{G} is of isotropic (actually constant) curvature and F is a Hamel function of \mathbf{G} , it follows from Proposition 4.5 that $\bar{\mathbf{G}}$ is of isotropic curvature with $\bar{R}^i_k = \bar{R}\delta_k^i - \frac{1}{2}\bar{R}_{;k}y^i$, where, by (30) and (31), \bar{R} is written as

$$\bar{R}(y) = -\frac{1}{4}F^2(y) + c^2F^2(-y) + c[F(y)F(-y) + F^2(-y)]. \quad (32)$$

By (32) and (31), a direct computation gives

$$\begin{aligned} \bar{R}_{;i} &= \bar{R}_{;i} - 2\bar{R}P_i - P\bar{R}_{;i} = c(1 + 2c)F(-y)T_i, \\ (T_i &:= 2(1 + c)F(-y)F_{;i}(-y) + 2F(y)F_{;i}(-y) - F(-y)F_{;i}(y)). \end{aligned} \quad (33)$$

If $\bar{R} = 0$ or

$$y^i T_i = -F(-y)[(2(1 + c)F(-y) + 3F(y))] = 0,$$

we have $F(-y) = \sigma F(y)$ for some constant σ . By the geometric meaning, $F(-y) = \sigma F(y)$ holds only on a nowhere dense set \mathcal{A} of $T\Omega \setminus \{0\}$. If $\bar{\mathbf{G}}$ is of constant curvature, we have $\bar{R}_{;i} = 0$ on the open set $\tilde{\mathcal{A}} := T\Omega \setminus \{0\} - \mathcal{A}$ since $\bar{R} \neq 0$ on $\tilde{\mathcal{A}}$. There is some T_i which is not zero on an $\tilde{\mathcal{A}}$. Then by (33), we have $c = 0$ or $c = -1/2$. Conversely, if $c = 0$ or $c = -1/2$, then $\bar{R}_{;i} = 0$ by (33). So $\bar{\mathbf{G}}$ is of constant curvature.

(ii) Using the facts that \mathbf{G} is projectively flat and $\bar{\mathbf{G}}$ is of isotropic curvature by (i), if $\bar{\mathbf{G}}$ is metrizable (on a conical region $\mathcal{C}(\Omega)$), we have $\bar{R}_{;i} = 0$ (on $\mathcal{C}(\Omega)$) according to a result in [17]. Then the above proof shows $c = 0$ or $c = -1/2$.

Conversely, if $c = 0$, \mathbf{G} is induced by F . If $c = -1/2$, then by (32), we obtain

$$-\bar{R} = \frac{1}{4}[F(y) + F(-y)]^2,$$

which is a Finsler metric (called the Klein metric). Further by (33) we have $\bar{R}_{\bar{i}} = 0$. Therefore, \mathbf{G} is induced by the Klein metric $-\bar{R}$ according to a result in [17]. Q.E.D.

5 Constructions of Hamel-Funk Functions

In this section, we consider the existence of Hamel functions or Funk functions on some special spray manifolds.

For a Finsler spray \mathbf{G} induced by a Finsler metric F , it is clear that F is a Hamel function of \mathbf{G} , but it is not a Funk function of \mathbf{G} . For a Minkowski spray $\mathbf{G} = 0$, we can construct many Funk functions \mathbf{G} by solving P from (16) for a given φ . Then by Lemma 3.3, we can obtain a lot of Hamel Function of $\mathbf{G} = 0$. By Lemmas 3.6–3.10, we have different ways of constructing Hamel functions of a spray \mathbf{G} , especially when $\mathbf{G} = 0$. Theses Hamel functions include projectively flat Finsler metrics of constant flag curvature. We know that a Hamel function is projectively invariant (Lemma 4.1); and all Hamel functions of a spray form a vector space (Lemma 3.5), but it might be difficult to know its dimension.

A Hamel function or a Funk function might be also a Finsler metric. A Hamel (resp. Funk) function is called a Finsler-Hamel (resp. Finsler-Funk) function, if it is also a Finsler metric. The following example gives a Finsler-Hamel function of a spray.

Example 5.1 *For an (α, β) -metric $F = \alpha\phi(s)$, $s = \beta/\alpha$, the spray coefficients \bar{G}^i of F is given by (cf. [15])*

$$\begin{aligned}\bar{G}^i &= G^i + \alpha^{-1}\Theta(-2\alpha Qs_0 + r_{00})y^i, \\ G^i &:= G_\alpha^i + \alpha Qs_0^i + \Psi(-2\alpha Qs_0 + r_{00})b^i,\end{aligned}$$

where G_α^i denote the spray coefficients of α and

$$\begin{aligned}Q &:= \frac{\phi'}{\phi - s\phi'}, \quad \Theta := \frac{Q - sQ'}{2\Delta}, \quad \Psi := \frac{Q'}{2\Delta}, \quad \Delta := 1 + sQ + (b^2 - s^2)Q', \\ r_{ij} &:= \frac{1}{2}(b_{i|j} + b_{j|i}), \quad s_{ij} := \frac{1}{2}(b_{i|j} - b_{j|i}), \quad s_j := b^i s_{ij}, \quad s^i_j := a^{im} s_{mj},\end{aligned}$$

where $b_{i|j}$ denotes the covariant derivative of the Levi-Civita connection of α .

It is clear that F is a Finsler-Hamel function of the spray \bar{G}^i , since F is naturally a Hamel function of \bar{G}^i and a Hamel function is projectively invariant (Lemma 4.1).

Proposition 5.2 *A 1-form is a Hamel function of a spray if and only if it is closed. So for a spray, a Hamel function plus a closed 1-form is a Hamel function.*

Proposition 5.3 *If \mathbf{G} is of isotropic curvature, then the S -curvature of \mathbf{G} with respect to any volume form is a Hamel function of \mathbf{G} .*

Proposition 5.2 follows directly from the definition of a Hamel function. For Proposition 5.3, the S -curvature satisfies $S_{;i} - S_{;i;0} = 0$ if \mathbf{G} is of isotropic curvature ([?]). So the S -curvature of \mathbf{G} with respect to any volume form is a Hamel function of \mathbf{G} .

Now we turn to study Funk functions of certain sprays. We have shown the construction of Funk functions of a Minkowski spray $\mathbf{G} = 0$. Actually, we can prove that there exist local Funk functions on a R-flat spray manifold, as shown in Theorem 1.2.

Proof of Theorem 1.2 : Consider the following PDE system

$$Q_{;i} = QQ_{;i}, \quad YP = P, \quad (Y := y^i \partial_i). \quad (34)$$

We only need to prove that (34) is integrable if \mathbf{G} is R-flat. The integrable conditions of (34) are

$$\delta_i YP - Y\delta_i P = [\delta_i, Y]P, \quad \delta_i \delta_j P - \delta_j \delta_i P = [\delta_i, \delta_j]P. \quad (35)$$

The first condition in (35) automatically holds, since by (34), we have

$$[\delta_i, Y] = 0, \quad \delta_i YP = P_{;i}, \quad Y\delta_i P = Y(PP_{;i}) = P_{;i}.$$

The second condition in (35) is equivalent to

$$Q_{;i;j} - Q_{;j;i} = -Q_{;r}R^r_{ij}. \quad (36)$$

Using (34), we have

$$Q_{;i;j} = Q_{;j}Q_{;i} + QQ_{;i;j}.$$

Then using $Q_{;i;j} = Q_{;j;i}$ and (34), we get

$$Q_{;i;j} - Q_{;j;i} = 0.$$

Thus (36) holds if \mathbf{G} is R-flat ($R^r_{ij} = 0$).

Q.E.D.

A similar system of PDEs to (34) is concerned in [6]. Next we consider the construction of Funk functions of sprays with non-zero Riemann curvature, and prove some results on the non-existence of non-zero Funk function on some spray manifolds under certain conditions.

Theorem 5.4 ([14]) *A weak Funk function on a complete regular spray manifold must be zero.*

Proof : Let (\mathbf{G}, M) be a complete regular spray manifold, and Q be a weak Funk function of (\mathbf{G}, M) . Then Q satisfies $Q_{;0} = Q^2$. Along a geodesic $c = c(t)$ of \mathbf{G} with $c(0) = x \in M, \dot{c}(0) = y \in T_x M$, we have

$$Q'(t) - Q^2(t) = 0, \quad (Q(t) := Q(c(t), \dot{c}(t))),$$

whose solution is given by

$$Q(t) = \frac{Q(0)}{1 - Q(0)t}.$$

Since (\mathbf{G}, M) is complete, $Q(t)$ should be defined on $(-\infty, +\infty)$. It is easy to conclude $Q(0) = 0$ since $Q(t)$ is unbounded at the point $t = 1/Q(0)$ if $Q(0) \neq 0$. Thus we get $Q = 0$ since the geodesic c is arbitrarily given. Q.E.D.

In Theorem 5.4, if (\mathbf{G}, M) is not complete, then possibly there exist nontrivial Funk functions. For example, consider the Minkowski spray $\mathbf{G} = 0$ on a convex domain Ω of R^n , and the Funk metric on Ω is a Funk function of \mathbf{G} .

A Funk functions on a spray manifold of scalar curvature has some special properties. First we introduce a basic proposition as follows.

Proposition 5.5 *Let \mathbf{G} be a spray of scalar curvature $R^i_k = R\delta^i_k - \tau_k y^i$ and Q be a Funk function of \mathbf{G} . Then R, τ_k and Q are related by*

$$RQ_{;k} = Q\tau_k. \quad (37)$$

(i) If \mathbf{G} is of isotropic curvature with $R \neq 0$, then (37) gives

$$Q^2 = cR, \quad (38)$$

where $c = c(x)$ is a scalar function.

(ii) If \mathbf{G} is of constant curvature with $R \neq 0$, then Q is a closed 1-form. In this case, if $Q \neq 0$, then for some scalar function $\sigma = \sigma(x)$, we have

$$Q = \sigma_0, \quad R = \pm[(e^{-\sigma})_0]^2, \quad Q^2 = \pm e^{2\sigma}R. \quad (39)$$

Proof: Since Q is a Funk function of \mathbf{G} , we have $Q_{,r}R^r_k = 0$ by (36). Using $R^i_k = R\delta_k^i - \tau_k y^i$ we get (37) immediately.

If \mathbf{G} is of isotropic curvature with $R \neq 0$, then plugging $R_{,k} = 2\tau_k$ into (37) yields

$$\frac{2Q_{,i}}{Q} = \frac{R_{,i}}{R}, \quad (\text{letting } Q \neq 0),$$

which gives (38) by an integration.

If \mathbf{G} is of constant curvature with $R \neq 0$, then we have $R_{,i} = 0$. Thus by (38), we have

$$2Q^2Q_{,i} = 2QQ_{,i} = Rc_i + cR_{,i} = Rc_i, \quad (c_i := c_{x^i}).$$

Contracting the above by y^i and using (38), we have

$$2cQ = c_0.$$

If $c = 0$, then $Q = 0$ by (38). If $c \neq 0$, then we have

$$Q = (\ln \sqrt{|c|})_0.$$

So Q is a closed 1-form. Putting $Q = \sigma_0$ with $c = \pm e^{2\sigma}$, we obtain (39). Q.E.D.

Theorem 5.6 ([2]) *A Funk function on a Finsler manifold of non-zero scalar flag curvature must be zero. So a spray \mathbf{G} of scalar curvature with non-zero Riemann curvature is not metrizable if there is a non-zero Funk function of \mathbf{G} .*

Proof: Let L be a Finsler metric of scalar flag curvature $\lambda \neq 0$, and Q be a Funk metric of \mathbf{G} . Then the spray \mathbf{G} of L is given by

$$R^i_k = R\delta_k^i - \tau_k y^i, \quad (R := \lambda L, \quad \tau_k := \lambda y_k).$$

Suppose $Q \neq 0$. Then by (37) we have

$$\frac{2Q_{,i}}{Q} = \frac{L_{,i}}{L},$$

which gives $Q^2 = cL$ for a scalar function $c = c(x) \neq 0$. By $Q^2 = cL$, we have

$$2Q^2Q_{,i} = 2QQ_{,i} = Lc_i, \quad (c_i := c_{x^i}).$$

Contracting the above by y^i and using $Q^2 = cL$, we see that $Q = c_0/(2c)$ is a closed 1-form. Thus $L = Q^2/c$ is not a Finsler metric, which gives a contradiction. Q.E.D.

In the following examples, we consider the construction of Funk functions of a projectively flat Berwald spray of isotropic curvature. It should be the first time that we obtain non-trivial Funk functions on a spray manifold with non-zero Riemann curvature.

Example 5.7 Let \mathbf{G} be a projectively flat Berwald spray with

$$G^i := \tau_0 y^i, \quad (\tau_i := \tau_{x^i}, \quad \tau := f(\sigma)), \quad (40)$$

where $\sigma = \sigma(x)$ is a scalar function. Suppose $Q := \sigma_0$ is a Funk function of \mathbf{G} . Then by $Q_{;j} = QQ_{,j}$, we have

$$\sigma_{0;j} - \sigma_0 \sigma_j = 0.$$

which can be rewritten as

$$\sigma_{ij} - [1 + 2f'(\sigma)]\sigma_i \sigma_j = 0, \quad \text{or} \quad \left[\int e^{-\sigma - 2f(\sigma)} d\sigma \right]_{ij} = 0.$$

Thus σ is given by

$$\int e^{-\sigma - 2f(\sigma)} d\sigma = \langle a, x \rangle + b, \quad (41)$$

where a is an n -vector and b is a constant.

(i) By the different choices of the function f in (41), we obtain many Funk functions $Q = \sigma_0$ of the spray \mathbf{G} in (40).

(ii) \mathbf{G} in (40) is of isotropic curvature $R^i_k = R\delta_k^i - \frac{1}{2}R_{,k}y^i$ with

$$R := -(\tau_0)^2 - \tau_{0;0} = -\{f''(\sigma) + f'(\sigma)[1 + f'(\sigma)]\}(\sigma_0)^2. \quad (42)$$

It follows from Proposition 5.5(i) that (38) holds, and then by (42), c in (38) is given by

$$c = -\{f''(\sigma) + f'(\sigma)[1 + f'(\sigma)]\}^{-1} \quad (43)$$

(iii) By Theorem 5.6, the spray in (40) with σ given by (41) is non-metrizable.

In Example 5.7, taking $f(t) := t$ as a special case, by (40), (41) and (43), we obtain

$$G^i = -\frac{1}{3} \frac{\langle a, y \rangle}{\langle a, x \rangle + b} y^i, \quad Q = -\frac{1}{3} \frac{\langle a, y \rangle}{\langle a, x \rangle + b}, \quad c = -\frac{1}{2}.$$

Example 5.8 If the spray \mathbf{G} given by (40) is of constant curvature, we can also obtain non-trivial Funk function of \mathbf{G} . Let \mathbf{G} be given by (40) and σ is determined by (41). Then $Q = \sigma_0$ is a Funk function of \mathbf{G} by Example 5.7. Now we can choose f such that \mathbf{G} is of constant curvature. To do so, we need to solve $R_{,i} = 0$ for R given by (42). It is easy to see that $R_{,i} = 0$ if and only if

$$f'''(t) + 3f''(t) + 2f'(t)f''(t) + 2f'(t) + 2(f'(t))^2 = 0,$$

whose solution is given by

$$f(t) = -c_1 e^{-t} + \ln |1 + c_2 e^{2c_1 e^{-t}}| + c_3,$$

where c_1, c_2, c_3 are constant.

Example 5.9 Consider an n -dimensional projectively flat Berwald spray \mathbf{G} :

$$G^i = -\frac{\langle x, y \rangle}{|x|^2} y^i.$$

In [17], we have shown that \mathbf{G} is of constant curvature and it is not metrizable, and further, the Riemann curvature of \mathbf{G} satisfies

$$R^r_{jk} = \frac{1}{2} R_{,j} \delta_k^r - (jk), \quad R := \frac{|x|^2 |y|^2 - \langle x, y \rangle^2}{|x|^4}. \quad (44)$$

We will prove that there is a Funk function $Q \neq 0$ of \mathbf{G} if and only if $n = 2$ and Q is given by

$$Q = \sigma_0, \quad \sigma := \ln \left| \frac{c_1}{c_1 \arctan \frac{x^2}{x^1} + c_2} \right|, \quad (45)$$

where c_1, c_2 are constant. In this case, Q, R satisfy (39), and c in (38) is given by

$$c = \left(\frac{c_1}{c_1 \arctan \frac{x^2}{x^1} + c_2} \right)^2. \quad (46)$$

Actually, by Proposition 5.5(ii), we first have $Q = \sigma_0$ for some scalar function $\sigma = \sigma(x)$. By (36), the integral condition for Q is given by $Q_{.r}R^r_{jk} = 0$. Then by (44), the integral condition becomes

$$(|x|^2 \delta_{ij} - x^i x^j) \sigma_k - (jk) = 0. \quad (47)$$

Taking a sum over i, j in (47) we have

$$(n-2)|x|^2 \sigma_k + \sigma_r x^r x^k = 0. \quad (48)$$

Then a further contraction in (48) by x^k gives $\sigma_r x^r = 0$, which implies that σ is constant by (48) again when $n > 2$. When $n = 2$, the integral condition (47) becomes $x^1 \sigma_1 + x^2 \sigma_2 = 0$, and in this case, solving the following system of PDEs

$$Q_{;i} = QQ_{.i} \quad \left(\iff \sigma_{ij} + \frac{1}{|x|^2} (\sigma_i x^j + \sigma_j x^i) - \sigma_i \sigma_j = 0 \right),$$

we obtain (45). By the way, we also have $n = 2$ by the R in (44) and (39).

6 Hamel-Funk Sprays

In this section we introduce a new spray associated to a spray and a Hamel function, and then study some properties of this new spray.

For a given spray \mathbf{G} and a nonzero homogeneous function Q of degree one, we consider a spray $\bar{\mathbf{G}}$ defined by

$$\bar{G}^i := G^i + \frac{Q_{;0}}{2Q} y^i. \quad (49)$$

When \mathbf{G}, Q satisfies certain good conditions, $\bar{\mathbf{G}}$ is metrizable (Theorem 1.3 below). So in general case, $\bar{\mathbf{G}}$ is pre-metrizable. For short, $\bar{\mathbf{G}}$ is called a PM-spray, and \mathbf{G}, Q are respectively called the adjoint spray and adjoint function of $\bar{\mathbf{G}}$.

Lemma 6.1 *Each spray in the projective class $Proj(\mathbf{G})$ of a spray \mathbf{G} is a local PM-spray $\bar{\mathbf{G}}$ defined by (49) for some Q .*

Proof : For an arbitrary spray $\bar{\mathbf{G}} \in Proj(\mathbf{G})$, we have $\bar{G}^i = G^i + P y^i$ for some function P . Consider the following system of PDEs with the unknown function Q :

$$\bar{Y}(Q) = 2PQ, \quad Y(Q) = Q, \quad (\bar{Y} := y^i \delta_i, \quad Y = y^i \dot{\partial} y^i). \quad (50)$$

The integral condition of (50) is given by

$$[\bar{Y}, Y](Q) = \bar{Y}Y(Q) - Y\bar{Y}(Q).$$

Applying $[\bar{Y}, Y] = -\bar{Y}$ and (50), it is easy to verify that the above integral condition is satisfied. So (50) has a local solution Q . Q.E.D.

In Lemma 6.1, Q is generally not unique. For instance, letting

$$G^i := 0, \quad P := -\frac{x^1 y^1 + x^2 y^2}{1 + (x^1)^2 + (x^2)^2}, \quad Q := \frac{f\left(\frac{y^2}{y^1}, \frac{x^2 y^1 - x^1 y^2}{y^1}\right)}{1 + (x^1)^2 + (x^2)^2} y^1, \quad (51)$$

where $f = f(s, t)$ is some function. Then Q satisfies (50).

Lemma 6.2 *A PM-spray is projectively invariant with a fixed adjoint function.*

Proof : Let $\bar{\mathbf{G}}_1, \bar{\mathbf{G}}_2$ be two PM-sprays associated to $(\mathbf{G}_1, Q), (\mathbf{G}_2, Q)$ respectively, where \mathbf{G}_1 and \mathbf{G}_2 are projectively related with $G_1^i = G_2^i + P y^i$. By a direct computation, we have (see (29))

$$Q_{|i} = Q_{;i} - (PQ)_{.i}, \quad Q_{|0} = Q_{;0} - 2PQ,$$

where $|$ and $;$ denote the horizontal derivatives of \mathbf{G}_1 and \mathbf{G}_2 respectively. Therefore, we obtain

$$\begin{aligned} \bar{G}_1^i &= G_1^i + \frac{Q_{|0}}{2Q} y^i = G_2^i + P y^i + \frac{Q_{;0} - 2PQ}{2Q} y^i \\ &= G_2^i + \frac{Q_{;0}}{2Q} y^i = \bar{G}_2^i, \end{aligned}$$

which completes the proof.

Q.E.D.

Lemma 6.2 is similar to the result of the spray defined by the S-curvature ([?]). For a PM-spray associated to (\mathbf{G}, Q) , one important case is when Q is a Hamel or Funk function of \mathbf{G} . We have the following definition.

Definition 6.3 *Let \mathbf{G} be a spray and Q be a Hamel function of \mathbf{G} . The spray $\bar{\mathbf{G}}$ defined by (49) is called a Hamel spray associated to (\mathbf{G}, Q) , and \mathbf{G}, Q are respectively called the adjoint spray and adjoint Hamel function of $\bar{\mathbf{G}}$. As a special case, if Q is a Funk function of \mathbf{G} , then $\bar{\mathbf{G}}$ is called a Funk spray associated to (\mathbf{G}, Q) .*

It is clear that the Funk spray $\bar{\mathbf{G}}$ associated to (\mathbf{G}, Q) has the form $\bar{G}^i = G^i + (Q/2)y^i$.

Lemma 6.4 *A Hamel spray is projectively invariant with a fixed adjoint Hamel function.*

Proof : It directly follows from Lemma 6.2 and the fact that a Hamel function is projectively invariant, and so Q is a Hamel function of \mathbf{G}_1 if and only if it is a Hamel function of \mathbf{G}_2 , where \mathbf{G}_1 and \mathbf{G}_2 are projectively related. Q.E.D.

It is clear that a Finsler spray $\bar{\mathbf{G}}$ induced by a Finsler metric \bar{F} is a Hamel spray associated to $(\bar{\mathbf{G}}, \bar{F})$, and then by Lemma 6.4, $\bar{\mathbf{G}}$ is a Hamel spray associated to (\mathbf{G}, \bar{F}) for any spray \mathbf{G} projectively related to $\bar{\mathbf{G}}$.

Proof of Theorem 1.3 : If a spray \mathbf{G} is projectively metrizable, then there is a Finsler metric \bar{F} such that $\bar{G}^i := G^i + P y^i$ for some P is the Finsler spray of \bar{F} . By Lemma 2.2(ii), \bar{F} is a Hamel function of \mathbf{G} . This fact can also be proved by that \bar{F} is a natural Hamel function of $\bar{\mathbf{G}}$ and a Hamel function is projectively invariant. In this case, P is given by (10), and so the spray $\bar{\mathbf{G}}$ of \bar{F} is a Hamel spray associated to (\mathbf{G}, \bar{F}) .

Conversely, if a spray \mathbf{G} has a Finsler-Hamel function \bar{F} , it follows from Lemma 2.2 that \mathbf{G} is projectively related to the Finsler spray $\bar{\mathbf{G}}$ of \bar{F} which is the Hamel spray associated to $(\bar{\mathbf{G}}, \bar{F})$. This finishes the proof. Q.E.D.

By Theorem 1.3 we have the following corollary.

Corollary 6.5 *If a spray \mathbf{G} has a Finsler-Funk function \bar{F} , then \mathbf{G} is projectively induced by \bar{F} whose spray is the Funk spray associated to (\mathbf{G}, \bar{F}) .*

Example 6.6 *In Example 5.1, \bar{F} is a Finsler-Hamel function of $\bar{\mathbf{G}}$. So by Theorem 1.3, $\bar{\mathbf{G}}$ is a Hamel spray associated to (\mathbf{G}, \bar{F}) .*

For a Hamel or Funk spray, we give some special properties as follows.

Lemma 6.7 *Let $\bar{\mathbf{G}}$ and \mathbf{G} be two sprays projectively related and $Q \neq 0$ be a Hamel function. Then $\bar{\mathbf{G}}$ is a Hamel spray associated to (\mathbf{G}, Q) if and only if $Q_{;k} = 0$.*

Proof : Put $\bar{G}^i = G^i + Py^i$. A direct computation gives

$$Q_{;\bar{i}} = Q_{;i} - Q_{;r}(P_{.i}y^r + P\delta_i^r) = Q_{;i} - (PQ)_{.i}. \quad (52)$$

If $\bar{\mathbf{G}}$ is a Hamel spray associated to (\mathbf{G}, Q) , then by definition we have

$$P = \frac{Q_{;0}}{2Q}, \quad (53)$$

where Q is a Hamel function of the spray \mathbf{G} . It follows from (52) that

$$Q_{;\bar{i}} = Q_{;i} - \frac{1}{2}Q_{;0.i} = \frac{1}{2}(Q_{;i} - Q_{.i;0}) = 0.$$

Conversely, if $Q_{;\bar{k}} = 0$, then since Q is a Hamel function, it follows from (52) that

$$0 = Q_{;\bar{i}} = Q_{.i;0} - (PQ)_{.i} = \frac{1}{2}Q_{;0.i} - (PQ)_{.i} = \left(\frac{1}{2}Q_{;0} - PQ\right)_{.i},$$

which implies (53). Thus $\bar{\mathbf{G}}$ is a Hamel spray associated to (\mathbf{G}, Q) . Q.E.D.

Theorem 6.8 *(i) An n -dimensional Hamel spray $\bar{\mathbf{G}}$ associated to (\mathbf{G}, Q) is of scalar curvature if and only if its Riemann curvature has the form*

$$\bar{R}^i_k = \lambda(Q^2\delta_k^i - QQ_{.k}y^i), \quad (54)$$

where $\lambda = \lambda(x, y)$ is a scalar function given by

$$\lambda := (R + P^2 - P_{;0})/Q^2, \quad (P := Q_{;0}/(2Q), \quad R := Ric/(n-1)). \quad (55)$$

(ii) A Funk spray $\bar{\mathbf{G}}$ associated to (\mathbf{G}, Q) is of isotropic curvature if and only if its Riemann curvature has the form (54) with $\lambda = \lambda(x)$ being given by

$$\lambda = \eta - \frac{1}{4}, \quad (\eta := R/Q^2). \quad (56)$$

Proof : Put

$$\bar{G}^i = G^i + Py^i, \quad (P := Q_{;0}/(2Q)).$$

Let $\bar{\mathbf{G}}$ be of scalar curvature. Then \mathbf{G} is also of scalar curvature and we put $R^i_k = R\delta_k^i - \tau_k y^i$. Then we have (see (30))

$$\begin{aligned} \bar{R}^i_k &= \bar{R}\delta_k^i - \bar{\tau}_k y^i, \\ (\bar{R} := R + P^2 - P_{;0}, \quad \bar{\tau}_k := \tau_k + PP_{.k} + P_{.k;0} - 2P_{;k}). \end{aligned} \quad (57)$$

It is easy to obtain

$$\begin{aligned} P_{;k} &= \frac{QQ_{;0;k} - Q_{;0}Q_{;k}}{2Q^2}, & P_{;0} &= \frac{QQ_{;0;0} - (Q_{;0})^2}{2Q^2}, \\ P_{.k} &= \frac{2QQ_{;k} - Q_{;0}Q_{.k}}{2Q^2}, & P_{.k;0} &= \frac{Q_{;k;0}}{Q} - \frac{3Q_{;0}Q_{;k} + Q_{;0;0}Q_{.k}}{2Q^2} + \frac{(Q_{;0})^2Q_{.k}}{Q^3}, \end{aligned}$$

where in the third formula, $Q_{;0.k} = 2Q_{;k}$ is used, and in the fourth formula, $Q_{.k;0} = Q_{;k}$ is concerned. Plugging the above formulas into (57) and applying

$$Q_{;0;k} - Q_{;k;0} = -Q_{.r}R^r_k, \quad R^r_k = R\delta^r_k - \tau_k y^r,$$

we obtain

$$\bar{R} = \lambda Q^2, \quad \bar{\tau}_k = \lambda QQ_{.k},$$

where λ is given by (55). This gives the proof of case (i).

If $\bar{\mathbf{G}}$ is of isotropic curvature and Q is a Funk function of \mathbf{G} , then by Proposition 4.4(iii), we see that \mathbf{G} is of isotropic curvature, since $P = Q/2$ is a Hamel function. By (20) we get

$$\frac{P^2 - P_{;0}}{Q^2} = -\frac{1}{4}.$$

By (38), $\eta := R/Q^2$ is just a function of x . So (56) follows from (55). Q.E.D.

Based on Theorem 6.8, we make the following definition.

Definition 6.9 *A Hamel spray $\bar{\mathbf{G}}$ associated to (\mathbf{G}, Q) is called of scalar curvature λ if its Riemann curvature has the form (54). $\bar{\mathbf{G}}$ is called of isotropic (resp. constant) curvature λ if $\lambda = \lambda(x)$ (resp. $\lambda = \text{constant}$).*

Write (54) in the form

$$\bar{R}^i_k = \bar{R}\delta^i_k - \bar{\tau}_k y^i, \quad \bar{R} := \lambda Q^2, \quad \bar{\tau}_k := \lambda QQ_{.k}.$$

Then it is easily to show that for $Q \neq 0$,

$$\lambda = \lambda(x) \iff \bar{R}_{.k} = 2\bar{\tau}_k; \quad \lambda = \text{constant} \iff \bar{R}_{;k} = 0,$$

where we have applied Lemma 6.7 and Lemma 2.4 for the second case in the above.

Example 6.10 *Define a spray \mathbf{G} and 1-form Q by*

$$G^i := f'(\sigma)\sigma_0 y^i, \quad Q := \sigma_0,$$

where f is a function on R^1 and $\sigma = \sigma(x)$ is given by (41). Then by Example 5.7, \mathbf{G} is of isotropic curvature and Q is a Funk function of \mathbf{G} . By Theorem 6.8(ii), the Funk spray $\bar{\mathbf{G}}$ associated to (\mathbf{G}, Q) satisfies

$$\bar{R}^i_k = \lambda(Q^2\delta^i_k - QQ_{.k}y^i), \quad \lambda := -f''(\sigma) - f'(\sigma)[1 + f'(\sigma)] - \frac{1}{4},$$

where we have applied (56) and (43). So $\bar{\mathbf{G}}$ is of isotropic curvature λ . For suitable choice of f , λ in the above can be a constant (cf. Example 5.7). For instance, taking $f(t) = t$ we get $\lambda = -9/4$.

Example 6.11 Define a two-dimensional spray \mathbf{G} and 1-form Q by

$$G^i = -\frac{x^1 y^1 + x^2 y^2}{(x^1)^2 + (x^2)^2} y^i, \quad Q := \sigma_0, \quad \sigma := -\ln \left| \kappa + \arctan \frac{x^2}{x^1} \right|,$$

where κ is a constant. By Example 5.9, \mathbf{G} is of constant curvature and Q is a Funk function of \mathbf{G} . By Theorem 6.8(ii), the Funk spray $\bar{\mathbf{G}}$ associated to (\mathbf{G}, Q) satisfies

$$\bar{R}^i_k = \lambda(Q^2 \delta_k^i - Q Q_{.k} y^i), \quad \lambda := \left(\kappa + \arctan \frac{x^2}{x^1} \right)^2 - \frac{1}{4},$$

where we have applied (56) and (46). So $\bar{\mathbf{G}}$ is of isotropic curvature λ .

By Example 6.11, the Funk spray $\bar{\mathbf{G}}$ associated to (\mathbf{G}, Q) is not necessarily of constant curvature even if \mathbf{G} is of constant curvature.

For a spray \mathbf{G} , define a map

$$\mathcal{H} : [Q] \rightarrow \frac{Q_{;0}}{2Q}, \quad (58)$$

where $Q \neq 0$ is a homogeneous function of degree one and $[Q]$ denotes the equivalent class consisting of all the functions cQ for arbitrary nonzero constant c .

Let \mathcal{A} be the set of all nonzero Hamel functions and set $\mathcal{H}_1 := \mathcal{H}|_{[\mathcal{A}]}$. By definition, we have $Q \in \mathcal{H}_1^{-1}(P)$ if and only if

$$Q_{;i} = Q_{;i;0}, \quad Q_{;0} = 2PQ. \quad (59)$$

It is clear that for a given spray \mathbf{G} and a function P , the spray $\bar{G}^i := G^i + P y^i$ is metrizable if and only if there is a Finsler metric Q satisfying (59).

Generally, \mathcal{H} is not injective (see the example given by (51)). However, for certain function P , there will have a unique element in $\mathcal{H}_1^{-1}(P)$. We first show the following lemma.

Lemma 6.12 Let Q be a Hamel function of a spray \mathbf{G} and $\lambda = \lambda(x, y)$ be a homogeneous function of degree zero. Then λ satisfies $\lambda_{;0} = 0$ and $\bar{Q} := \lambda Q$ is a Hamel function of \mathbf{G} if and only if

$$2Q\lambda_{;i} = Q_{;0}\lambda_{.i}, \quad \text{or} \quad \lambda_{;i} = 0 \quad (60)$$

where in the second case, the derivative is take with respect to the spray

$$\bar{G}^i := G^i + \frac{Q_{;0}}{2Q} y^i.$$

Proof : Let $\bar{Q} = \lambda Q$ with $\lambda_{;0} = 0$. We have

$$\begin{aligned} \bar{Q}_{;i} &= Q\lambda_{;i} + \lambda Q_{;i}, & \bar{Q}_{.i} &= Q\lambda_{.i} + \lambda Q_{.i}, \\ \bar{Q}_{;i;0} &= Q\lambda_{;i;0} + Q_{;0}\lambda_{.i} + \lambda Q_{;i;0} = Q\lambda_{.i;0} + Q_{;0}\lambda_{.i} + \lambda Q_{;i}, \end{aligned}$$

where in the third formula above we have used the assumptions $\lambda_{;0} = 0$ and $Q_{.i;0} = Q_{;i}$. Therefore, \bar{Q} is a Hamel function if and only if

$$Q\lambda_{;i} + \lambda Q_{;i} = Q\lambda_{.i;0} + Q_{;0}\lambda_{.i} + \lambda Q_{;i},$$

which reduces to the first formula in (60) since $\lambda_{.i;0} = \lambda_{;0.i} - \lambda_{;i} = -\lambda_{;i}$. Conversely, if the first formula in (60) holds, it is easy to get $\lambda_{;0} = 0$ by contracting this formula by y^i . Further, the above proof has shown that \bar{Q} is Hamel function.

Now under the spray $\bar{\mathbf{G}}$, a direct computation gives

$$\lambda_{;i} = \lambda_{.i} - \frac{Q_{;0}}{2Q} \lambda_{.i},$$

which implies $\lambda_{;i} = 0 \iff 2Q\lambda_{;i} = Q_{;0}\lambda_{.i}$.

Q.E.D.

Proposition 6.13 For a given spray \mathbf{G} and a function P , let Q_o be a solution of (59). Suppose that the spray $\bar{G}^i := G^i + Py^i$ is of nonzero scalar curvature. Then any solution Q of (59) is given by $Q = cQ_o$ for some constant c , or $\mathcal{H}_1^{-1}(P)$ has a unique element.

Proof : Let $Q = cQ_o$ for a homogeneous function $c = c(x, y)$ of degree zero. Then we have

$$\frac{Q_{o;0}}{2Q_o} = \frac{Q_{;0}}{2Q} (= P),$$

which gives $c_{;0} = 0$. Since Q, Q_o are Hamel functions of \mathbf{G} , by Lemma 6.12, we have $c_{;i} = 0$. By assumption, $\bar{\mathbf{G}}$ is of nonzero scalar curvature. Then c is a constant due to Lemma 2.4, which also shows that $\mathcal{H}_1^{-1}(P)$ has a unique element. Q.E.D.

Example 6.14 In the system (59), put

$$G^i := 0, \quad P := -\frac{\mu\langle x, y \rangle}{1 + \mu|x|^2}, \quad (\mu = \text{constant} \neq 0).$$

A direct verification shows that

$$Q_o := \frac{\sqrt{(1 + \mu|x|^2)|y|^2 - \mu\langle x, y \rangle^2}}{1 + \mu|x|^2}$$

is a solution of (59). The spray $\bar{G}^i := G^i + Py^i$ is a Hamel spray associated to (\mathbf{G}, Q_o) and it is of constant curvature μ . By Proposition 6.13, any solution Q of (59) is given by $Q = cQ_o$ for some constant c . Actually, $\bar{\mathbf{G}}$ is metrizable induced by the metric Q_o . This can be an example of Theorem 1.5.

Proof of Theorem 1.5 : Let $\bar{\mathbf{G}}$ be a Hamel spray associated to (\mathbf{G}, Q) and $\bar{\mathbf{G}}$ be of nonzero scalar curvature (cf. (54) with $\lambda \neq 0$). If $\bar{\mathbf{G}}$ is metrizable, then there is a Finsler metric \bar{F} inducing $\bar{\mathbf{G}}$, and \bar{F} is a Hamel function of \mathbf{G} . By Lemma 2.2, $\bar{\mathbf{G}}$ can be expressed as

$$\bar{G}^i = G^i + \frac{\bar{F}_{;0}}{2\bar{F}}y^i,$$

which implies

$$\bar{F}_{;0} = 2P\bar{F}, \quad (P := Q_{;0}/(2Q)).$$

Thus \bar{F} is a solution of (59). By assumption, Q is also a solution of (59). Then by Proposition 6.13, $Q = c\bar{F}$ for some constant $c \neq 0$ is a Finsler metric. The converse is clear by Theorem 1.3. Q.E.D.

The Funk sprays in Examples 6.10 and 6.11 are not metrizable by Theorem 1.5, since the scalar curvature is not zero, where in Example 6.10, $f(t) \neq -t/2 + \ln|a + bt|$ for any constant a, b . Since any closed 1-form is a Hamel function of any spray, we have the following corollary by Theorem 1.5.

Corollary 6.15 Let $\sigma = \sigma(x) \neq 0$ be a scalar function, and define a spray $\bar{\mathbf{G}}$ by

$$\bar{G}^i = \frac{\sigma_{00}}{2\sigma_0}y^i, \quad (\sigma_i := \sigma_{x^i}, \quad \sigma_{ij} := \sigma_{x^i x^j}). \quad (61)$$

Then $\bar{\mathbf{G}}$ is metrizable if and only if $\bar{\mathbf{G}}$ is of zero scalar curvature, or equivalently,

$$3(\sigma_{00})^2 = 2\sigma_0\sigma_{000}, \quad (\sigma_{ijk} := \sigma_{x^i x^j x^k}). \quad (62)$$

Proof : $\bar{\mathbf{G}}$ is a Hamel spray associated to (\mathbf{G}, σ_0) with $\mathbf{G} = 0$, and $\bar{\mathbf{G}}$ is of scalar curvature. By (55), $\bar{\mathbf{G}}$ is of zero scalar curvature if and only if (62) holds. If $\bar{\mathbf{G}}$ is of zero scalar curvature, then $\bar{\mathbf{G}}$ is (locally) metrizable due to [17]. If $\bar{\mathbf{G}}$ is of nonzero scalar curvature, then $\bar{\mathbf{G}}$ is not metrizable by Theorem 1.5, since σ_0 is not a Finsler metric. Q.E.D.

References

- [1] P. L. Antonelli, R. S. Ingarden and M. Matsumoto, *The theory of sprays and Finsler spaces with applications in physics and biology*, Kluwer Academic Publishers, 1993.
- [2] I. Bucataru and Z. Muzsnay, *Non-existence of Funk functions for Finsler spaces of non-vanishing scalar flag curvature*, C. R. Acad. Sci. Paris, Ser. I **354** (2016), 619-622.
- [3] I. Bucataru and Z. Muzsnay, *Projective metrizable and formal integrability*, SIGMA 7 (2011), 114-125.
- [4] I. Bucataru and Z. Muzsnay, *Finsler metrizable isotropic sprays and Hilbert's Fourth Problem*, J. Aust. Math. Soc. **97** (2014), 27-47.
- [5] S. S. Chern and Z. Shen, *Riemann-Finsler geometry*, World Scientific, Singapore, 2005.
- [6] M. Crampin, *Isotropic and R-flat sprays*, Houston J. Math. **33** (2007), 451-459.
- [7] S. G. Elgendi and Z. Muzsnay, *Metrizability of holonomy invariant projective deformation of sprays*, Cana. Math. Bulletin, 2020.
- [8] Y. Li, X. Mo and Y. Yu, *Inverse problem of sprays with scalar curvature*, Intern. J. Math. **30**(6), 2019.
- [9] B. Li and Z. Shen, *Sprays of isotropic curvature*, Intern. J. math. 2019.
- [10] M. Matsumoto, *Every path space of dimension two is projectively related to a Finsler Space*, Open Systems 1, Information Dynamics, **3**(3) (1995), 291-303.
- [11] Z. Muzsnay, *The Euler-Lagrange PDE and Finsler metrizable*, Houston Journal of Mathematics, **32** (1) (2006), 79-98.
- [12] Z. Shen, *Volume comparison and its applications in Riemann-Finsler geometry*, Adv. in Math. **128**(1997), 306-328.
- [13] Z. Shen, *Projectively flat Finsler metrics of constant flag curvature*, Trans. AMS., **355**(4)(2002), 1713-1728.
- [14] Z. Shen, *Differential geometry of spray and Finsler spaces*, Kluwer Academic Publishers, Dordrecht, 2001.
- [15] Z. Shen, *On a class of Landsberg metrics in Finsler geometry*, Canadian J. of Math., **61**(6) (2009), 1357-1374.
- [16] G. Yang, *Some classes of sprays in projective spray geometry*, Diff. Geom. Appl., **29** (2011), 606-614.
- [17] G. Yang, *On sprays of scalar curvature and metrizable*, preprint.

Guojun Yang
Department of Mathematics
Sichuan University
Chengdu 610064, P. R. China
yangguojun@scu.edu.cn