

Closed embedded self-shrinkers of mean curvature flow

Oskar Riedler*

July 11, 2022

Abstract

In this article we show the existence of closed embedded self-shrinkers in \mathbb{R}^{n+1} that are topologically of type $S^1 \times M$, where $M \subset S^n$ is any isoparametric hypersurface in S^n for which the multiplicities of the principle curvatures agree. If the number of distinct principle curvatures of M is one the resulting self-shrinker is topologically $S^1 \times S^{n-1}$ and the construction recovers Angenent's shrinking doughnut [2].

1 Introduction

Consider a compact n -dimensional manifold Σ that is smoothly immersed in \mathbb{R}^{n+1} via a map $F_0 : \Sigma \rightarrow \mathbb{R}^{n+1}$. A *mean curvature flow* of $F_0(\Sigma)$ is a family of smooth immersions $F_t : \Sigma \rightarrow \mathbb{R}^{n+1}$ where $t \in \mathbb{R}$ varies over some interval and for which

$$\partial_t F_t(x) = \mathbf{H}_t(x)$$

holds for all t . Here $\mathbf{H}_t(x)$ is the mean curvature of $F_t(\Sigma)$ at $F_t(x)$. In other words $F_t(\Sigma)$ flows along its mean curvature vector in \mathbb{R}^{n+1} . Due to compactness of Σ such a flow necessarily becomes singular in finite time, see e.g. [11].

By the work of Huisken [11], Ilmanen [12] and White [18] rescaling $F_t(M)$ near the singular time in an appropriate way leads to weak limits that are so called *self-shrinkers*, that is immersed manifolds whose mean curvature flow is given by dilations. These self-shrinkers then take a special role in the singularity theory of the mean curvature flow.

*Westfälische Wilhelms-Universität Münster, Mathematisches Institut, oskar.riedler@uni-muenster.de

In this paper we use the theory of isoparametric foliations of the sphere S^n to construct new examples of closed embedded self-shrinkers. Concretely we show:

Theorem A. For any isoparametric hypersurface M in S^n , $n \geq 2$, for which the multiplicities m_1 and m_2 of the principal curvatures agree there is a closed embedded self-shrinker of topological type $S^1 \times M$ in \mathbb{R}^{n+1} . This hypersurface is a union of homothetic copies of the leaves of the isoparametric foliation of S^n associated to M .

The terminology of isoparametric foliations will be recalled in Section 2.1. The proof works via a reduction of the shrinker condition to a geodesic equation in a two-dimensional manifold. Simple periodic solutions of this ordinary differential equation are then established by shooting methods very similar to [2], although the equation itself is quite different.

Denoting with g the number of principal curvatures of a regular leaf of the isoparametric foliation one has in the case $g = 1$ that the leaves become the latitudes of a sphere. The hypersurfaces found by Theorem A are then rotationally invariant under the $O(n)$ action on \mathbb{R}^{n+1} and topologically of type $S^1 \times S^{n-1}$, the same topological type as the “shrinking donut” found by Angenent [2]. It is currently an open question whether there exist embedded rotationally invariant self-shrinkers of type $S^1 \times S^{n-1}$ in \mathbb{R}^{n+1} other than Angenent’s example, so we also remark:

Proposition B. In the case $g = 1$ the construction of Theorem A gives Angenent’s shrinking doughnut [2].

The structure of this article is as follows: In section 2 we recall the necessary facts about isoparametric foliations, explain the reduction of the self-shrinker problem to an ordinary differential equation, and remark on some elementary properties of the resulting equation. In section 3 the shooting argument is presented and Theorem A is shown with the exception of a technical proposition. Proposition B is also proved in section 3. The aforementioned technical proposition is shown in section 4.

The author would like to thank Peter McGrath for interesting and helpful discussions. The author gratefully acknowledges the support of Germany’s Excellence Strategy EXC 2044 390685587, Mathematics Münster: Dynamics-Geometry-Structure.

2 Reduction and geodesic equation

In subsection 2.1 we recall some basic definitions and facts about isoparametric foliations. Subsection 2.2 explains the reduction procedure: a result due to Angenent [2] gives that a hypersurface is a self-shrinker if and only if it is minimal in some metric g_{Ang} . The reduction theorem of Palais and Terng [15] is then applied in order to reduce the shrinker property to a geodesic equation on an open subset of \mathbb{R}^2 equipped with a special metric. In subsection 2.3 we present this geodesic equation and simplify its form.

2.1 Isoparametric foliations on spheres

Definition 2.1. Let M be a smooth Riemannian manifold. A smooth function $f : M \rightarrow \mathbb{R}$ is called *isoparametric* if there are smooth functions $a, b : f(M) \rightarrow \mathbb{R}$ so that

$$\|\nabla f\|^2 = a \circ f, \quad (1)$$

$$\Delta f = b \circ f. \quad (2)$$

The geometric meaning of condition (1) is that the fibres of f form a (singular) transnormal system¹, in particular they are all equidistant to each other. Condition (2) implies that the regular fibres of the foliation are of constant mean curvature in M (cf. [6]). If M is a space-form one even has that the individual principal curvatures of such a fibre are constant along the fibre. Foliation that arise from the fibres of an isoparametric function are called *isoparametric foliations*.

The classification of isoparametric foliations in spheres was initiated by Cartan in [3–5]. This has proven to be a difficult problem and, despite a long and active history of research, it is in part still open. A significant part of the structure theory of these foliations was developed by Münzner in two seminal papers [13, 14].

We review now some structural facts of isoparametric foliations of S^n , cf. [6, 10, 13, 14, 17] for proofs and further information:

- (i) The principal curvatures of any regular fibre are constant along the fibre.
- (ii) The number of distinct principal curvatures of a regular fibre is the same for any two regular fibres. Denoting this number by g one has that $g \in \{1, 2, 3, 4, 6\}$.

¹Meaning if a geodesic is perpendicular to a leaf at any time that the geodesic then remains perpendicular to all leaves it intersects, cf. [16] for more details.

- (iii) There are precisely two singular fibres V_1 and V_2 . One has $\text{dist}(V_1, V_2) = \frac{\pi}{g}$. These singular fibres are closed and minimal submanifolds of S^n .
- (iv) Any regular fibre is of the form $M_\varphi := \{x \in S^n \mid \text{dist}(x, V_1) = \varphi\}$, where $\varphi \in (0, \frac{\pi}{g})$. These fibres are all diffeomorphic to one another.
- (v) The principal curvatures of a regular fibre M_φ are of the form $\cot(\varphi), \cot(\varphi + \frac{\pi}{g}), \dots, \cot(\varphi + \frac{(g-1)\pi}{g})$. Denoting with $m_1 := \text{codim}(V_1) - 1$, $m_2 := \text{codim}(V_2) - 1$ the multiplicities of these principle curvatures are m_1, m_2, m_1, \dots . In particular $n - 1 = \frac{g}{2}(m_1 + m_2)$ and $m_1 = m_2$ if g is odd.
- (vi) For $\varphi^* = \frac{2}{g} \arctan(\sqrt{m_1/m_2})$ the hypersurface M_{φ^*} is minimal in S^n .
- (vii) The volume of a regular fibre is given by

$$\text{Vol}(M_\varphi) = c \sin(\frac{g}{2}\varphi)^{m_1} \cos(\frac{g}{2}\varphi)^{m_2} \quad (3)$$

where c a positive constant that does not vary in φ (but will be different for different foliations).

- (viii) There is a homogenous polynomial $F : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$ (called the *Cartan-Münzner polynomial*) of degree g so that $F|_{S^n} = \cos(g\varphi)$ and for which one has:

$$\|\nabla F(x)\|^2 = g^2 \|x\|^{2g-2}, \quad \Delta F(x) = \frac{g^2}{2}(m_1 - m_2) \|x\|^{g-2}.$$

Example. The cases $g \in \{1, 2, 3\}$ were classified by Cartan, see [6] for more detailed remarks also for the other cases. Examples where the fibres of the foliation arise as the orbits of an isometric action on S^n are called homogenous.

- (i) When $g = 1$ the isoparametric foliation is congruent (that is equal up to an isometric transformation) to the latitudes of the sphere S^n . One has $V_1 = \{(1, 0, \dots, 0)\}$, $V_2 = \{(-1, 0, \dots, 0)\}$ and $M_\varphi = \{\cos(\varphi)\} \times \sin(\varphi)S^{n-1}$ for $\varphi \in (0, \pi)$. These examples are homogenous and $m_1 = m_2 = n - 1$.
- (ii) When $g = 2$ the isoparametric foliation is congruent to the foliation by Clifford tori $S^{m_1} \times S^{m_2}$. One has $V_1 = \{(0, \dots, 0)\} \times S^{m_2}$, $V_2 = S^{m_1} \times \{(0, \dots, 0)\}$ and $M_\varphi = \sin(\varphi)S^{m_1} \times \cos(\varphi)S^{m_2}$ for $\varphi \in (0, \frac{\pi}{2})$. The integers m_1, m_2 are arbitrary so long as $m_1 + m_2 = n - 1$, in particular $m_1 \neq m_2$ is possible. The fibres are the orbits of an isometric $O(m_1) \times O(m_2)$ action on S^n .
- (iii) When $g = 3$ one has $m_1 = m_2 \in \{1, 2, 4, 8\}$. The fibres of the foliation arise as the

distance tubes of certain embeddings of the projective planes $\mathbb{F}\mathbb{P}^2 (= V_1)$ in S^{3m_1+1} where $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}, \mathbb{H}, \mathbb{O}\}$ is one of the real division algebras or the octonions. These examples are homogenous.

- (iv) For $g = 4$ there is an infinite family, introduced by Ferrus, Karcher, and Münzner in [10], which contains both homogenous as well as inhomogenous examples. Two additional homogenous cases beyond this family exist, else all examples belong to this family. Here $m_1 \neq m_2$ is possible.
- (v) For $g = 6$ one has $m_1 = m_2 \in \{1, 2\}$, as was shown by Abresch [1]. For both cases there exist homogenous examples. If $m_1 = m_2 = 1$ it was shown by Dorfmeister and Neher [7] that the homogenous example is the only one.

2.2 Reduction for self-shrinkers

Definition 2.2. Let F be the Cartan-Münzner polynomial of an isoparametric foliation of S^n . Define:

$$\mathbf{f} : \mathbb{R}^{n+1} - \mathbb{R}_{\geq 0} \cdot (V_+ \cup V_-) \rightarrow (0, \infty) \times (0, \frac{\pi}{g}), \quad x \mapsto \left(\|x\|, \frac{\arccos(F(x/\|x\|))}{g} \right).$$

A set $X \subset \mathbb{R}^{n+1}$ is called *\mathbf{f} -invariant* if there is a set $N \subset (0, \infty) \times (0, \frac{\pi}{g})$ so that $X = \mathbf{f}^{-1}(N)$. Compare with the notion of F -invariant in [17].

Note that the \mathbf{f} -invariant sets are precisely those sets that are unions of homothetic copies of the regular fibres of the isoparametric foliation - that is unions of sets of the form $r \cdot M_\varphi$ for $(r, \varphi) \in (0, \infty) \times (0, \frac{\pi}{g})$.

Recall (cf. [2, 8]) that a closed submanifold $X \subset \mathbb{R}^{n+1}$ is a *self-shrinker under mean curvature flow* (short: *self-shrinker*) if and only if there is an $\tau > 0$ such that X is a minimal hypersurface in \mathbb{R}^{n+1} equipped with the metric:

$$g_{\text{Ang}} = e^{-\frac{\tau\|x\|^2}{n}} \sum_{i=1}^n dx_i^2 = e^{-\frac{\tau\|x\|^2}{n}} g_{\text{Euc}}.$$

The parameter τ is related to the extinction time of X . By rescaling X if necessary we take $\tau = 1$ in what follows.

Proposition 2.3. *Equipping $\mathbb{R}^{n+1} - \mathbb{R}_{\geq 0} \cdot (V_+ \cup V_-)$ with the metric g_{Ang} (with $\tau = 1$) and $(0, \infty) \times (0, \frac{\pi}{g})$ with the metric $g_{\text{Subm}} := e^{-\frac{r^2}{n}} (dr^2 + r^2 d\varphi^2)$ one has:*

(i) \mathbf{f} is a surjective and proper Riemannian submersion.

(ii) The mean curvature vector of a fiber $\mathbf{f}^{-1}(r, \varphi)$ is given by

$$e^{\frac{r^2}{2n}} \left(\left(\frac{1}{r} - \frac{r}{n} \right) \nu_r + \frac{H(\varphi)}{r} \nu_\varphi \right),$$

where $H(\varphi)$ is the mean curvature of $M_\varphi \subset S^n$, ν_φ is the unit normal of rM_φ in rS^n equipped with g_{Ang} , and ν_r is the unit normal of rS^n in \mathbb{R}^{n+1} equipped with g_{Ang} .

The proof is a standard calculation and from (ii) one sees that the mean curvature of the fibres of \mathbf{f} form a basic field of the Riemannian submersion. Riemannian submersions with this property are the key ingredient in the reduction theory developed by Palais and Terng in [15], recall:

Theorem (Palais-Terng, cf. Theorem 4 in [15]). *Let $\pi : (E, g_E) \rightarrow (B, g_B)$ be a Riemannian submersion for which the mean curvatures of the fibres form a basic field, then for a k -dimensional submanifold $X \subset B$ one has that $\pi^{-1}(X)$ is minimal in E if and only if X is minimal in $(B, V^{2/k}g_B)$. Here $V^{2/k}g_B$ is the metric given by*

$$(V^{2/k}g_B)(b) = \text{Vol}_{g_E}(\pi^{-1}(b))^{2/k} g_B(b).$$

Using (3) one gets (up to a constant factor):

$$\begin{aligned} \text{Vol}_{g_{Ang}}(\mathbf{f}^{-1}(r, \varphi))^2 &= e^{-\frac{r^2(n-1)}{n}} \text{Vol}_{g_{\text{Euc}}}(r \cdot M_\varphi)^2 = r^{2(n-1)} e^{-r^2(1-1/n)} \text{Vol}_{S^n}(M_\varphi)^2 \\ &= r^{2(n-1)} e^{-r^2(1-1/n)} \sin^{2m_1}\left(\frac{g}{2}\varphi\right) \cos^{2m_2}\left(\frac{g}{2}\varphi\right). \end{aligned}$$

The problem of finding \mathbf{f} -invariant hypersurfaces that are minimal with respect to g_{Ang} is then reduced to finding geodesic segments in $(0, \infty) \times (0, \frac{\pi}{g})$ equipped with the metric

$$r^{2n-2} e^{-r^2} \sin\left(\frac{g}{2}\varphi\right)^{2m_1} \cos\left(\frac{g}{2}\varphi\right)^{2m_2} (dr^2 + r^2 d\varphi^2). \quad (4)$$

We conclude:

Proposition 2.4. *A \mathbf{f} -invariant hypersurface $X \subset \mathbb{R}^{n+1}$ is a closed self-shrinker in \mathbb{R}^{n+1} if and only if $N := \mathbf{f}(X)$ is a closed geodesic in $(0, \infty) \times (0, \frac{\pi}{g})$ with respect to the metric (4).*

2.3 Geodesic equation

For the metric (4) one gets the following geodesic equation, where α denotes the angle between $\frac{dr}{dt}$ and $\frac{d\varphi}{dt}$:

$$\begin{aligned}\frac{dr}{dt} &= \cos \alpha G(r, \varphi), \\ \frac{d\varphi}{dt} &= \sin \alpha \frac{G(r, \varphi)}{r}, \\ \frac{d\alpha}{dt} &= \sin \alpha r \partial_r \frac{G(r, \varphi)}{r} - \cos \alpha \frac{1}{r} \partial_\varphi G(r, \varphi).\end{aligned}$$

Here

$$G(r, \varphi) = r^{-n+1} e^{r^2/2} \sin\left(\frac{g}{2}\varphi\right)^{-m_1} \cos\left(\frac{g}{2}\varphi\right)^{-m_2}.$$

Since we are not directly interested in the parametrisation of the geodesic but rather in its orbit we perform a substitution $\frac{dt^{\text{new}}}{dt^{\text{old}}} = \frac{1}{r} G(r, \varphi)$ to simplify the equation:

$$\begin{aligned}\frac{dr}{dt} &= r \cos \alpha, \\ \frac{d\varphi}{dt} &= \sin \alpha, \\ \frac{d\alpha}{dt} &= \sin \alpha (r^2 - n) + \frac{g}{2} \cos \alpha (m_1 \cot\left(\frac{g}{2}\varphi\right) - m_2 \tan\left(\frac{g}{2}\varphi\right)).\end{aligned}$$

We simplify once more by letting $\theta(t) := \frac{g}{2}\varphi(t)$, substituting $\frac{dt^{\text{new}}}{dt^{\text{old}}} = \frac{g}{2} \frac{1}{\sin(2\theta)}$, letting $\xi(t) := \frac{g}{2} \ln\left(\sqrt{\frac{2}{g}} r(t)\right)$ and $m := \frac{2}{g}n = m_1 + m_2 + \frac{2}{g}$ to get:

$$\begin{aligned}\frac{d\xi}{dt} &= \cos \alpha \sin(2\theta), \\ \frac{d\theta}{dt} &= \sin \alpha \sin(2\theta), \\ \frac{d\alpha}{dt} &= \sin \alpha \sin(2\theta) \left(e^{\frac{4}{g}\xi} - m \right) + 2 \cos \alpha l(\theta).\end{aligned} \tag{*}$$

Here $l(\theta) = m_1 \cos^2(\theta) - m_2 \sin^2(\theta)$.

If $\theta'(t) \neq 0$ then ξ may be (locally) given the form of a graph over θ . This graph obeys the following ODE:

$$\frac{d^2\xi}{d\theta^2}(\theta) = \frac{\xi''(t)}{\theta'(t)^2} - \frac{\xi'(t)}{\theta'(t)} \frac{\theta''(t)}{\theta'(t)^2} = - \left(1 + \left(\frac{d\xi}{d\theta} \right)^2 \right) \left(e^{\frac{4}{g}\xi} - m + 2H(\theta) \frac{d\xi}{d\theta} \right). \tag{**}$$

Here

$$H(\theta) = \frac{l(\theta)}{\sin(2\theta)} = \frac{m_1}{2} \cot(\theta) - \frac{m_2}{2} \tan(\theta).$$

The ODE (*) can be formulated for all initial conditions $(\xi, \theta, \alpha) \in \mathbb{R}^3$. But in coordinates (ξ, θ) the domain $(0, \infty) \times (0, \frac{\pi}{g})$ has been transformed to $\mathbb{R} \times (0, \frac{\pi}{2})$; so we are only interested in solutions where the ξ and θ components remain in that domain. We set $\mathcal{D} := \mathbb{R} \times (0, \frac{\pi}{2}) \times \mathbb{R}$.

The ODE (*) admits two trivial families of solutions in \mathcal{D} , namely for any $k \in \mathbb{Z}$:

$$\begin{aligned} (\xi, \theta, \alpha)(t) &= (\xi(0) + 2(-1)^k \frac{\sqrt{m_1 m_2}}{m_1 + m_2} t, \arctan\left(\sqrt{\frac{m_1}{m_2}}\right), \pi k), \\ (\xi, \theta, \alpha)(t) &= \left(\frac{g}{4} \ln m, \operatorname{arccot}\left(e^{(-1)^k 2t} \cot(\theta(0))\right), \frac{\pi}{2} + \pi k\right). \end{aligned}$$

The first of these solutions lifts to the cone $\mathbb{R}_{>0} \cdot M_{\varphi^*}$ over the minimal hypersurface of the isoparametric foliation, which is a minimal submanifold of \mathbb{R}^{n+1} . The second lifts to the round sphere, albeit with the singular fibres V_1, V_2 removed.

2.4 Elementary properties of (*) and symmetry

We briefly note some elementary properties of solutions of (*), proofs are standard and are thus omitted.

Proposition 2.5. (i) For any $(\xi_0, \theta_0, \alpha_0) \in \mathbb{R}^3$ there is a unique solution γ of (*) with initial condition $\gamma(0) = (\xi_0, \theta_0, \alpha_0)$. This solution is smooth and has domain of definition all of \mathbb{R} , i.e. solutions exist for all times.

(ii) Suppose $(\xi_s, \theta_s, \alpha_s)$ converges in \mathbb{R}^3 to a point $(\xi_\infty, \theta_\infty, \alpha_\infty)$. Denote by γ_s the solution of (*) with initial condition $(\xi_s, \theta_s, \alpha_s)$ and γ_∞ the solution of (*) with initial condition $(\xi_\infty, \theta_\infty, \alpha_\infty)$. Then γ_s converges uniformly on compacta to γ_∞ .

(iii) Solutions of (*) with initial condition in \mathcal{D} remain in \mathcal{D} for all times.

For our investigation we are interested in periodic solutions of (*). These will be found with the help of a discrete symmetry of the ODE (*).

Definition 2.6. Define $\theta^* := \arctan(\sqrt{m_1/m_2})$ and let

$$S : \mathbb{R}^3 \rightarrow \mathbb{R}^3, \quad (\xi, \theta, \alpha) \mapsto (\xi, 2\theta^* - \theta, \pi - \alpha).$$

Remark. Note that θ^* is the solution in $(0, \frac{\pi}{2})$ of $l(\theta) = 0$. Additionally the map S is an involution that reflects θ at θ^* while sending $\cos \alpha \rightarrow -\cos \alpha$ and $\sin \alpha \rightarrow \sin \alpha$. In the event that $m_1 = m_2$ one has $\theta^* = \frac{\pi}{4}$ and $l(2\theta^* - \theta) = -l(\theta)$ as well as $\sin(2(2\theta^* - \theta)) = \sin(2\theta)$. For $m_1 = m_2$ one has $S(\mathcal{D}) = \mathcal{D}$.

Proposition 2.7. *If $m_1 = m_2$ then for any $x \in \mathcal{D}$ one has $S(\gamma_1(-t)) = \gamma_2(t)$ for all $t \in \mathbb{R}$, where γ_1, γ_2 are the solutions to $(*)$ with initial conditions x and $S(x)$, respectively.*

If $x = S(x)$ then $\gamma_1 = \gamma_2$ in the above proposition and one gets $S(\gamma_1(t)) = \gamma_1(-t)$. Noting that $(*)$ is further invariant under transformations of the form $\alpha \mapsto \alpha + 2\pi k$ for $k \in \mathbb{Z}$ then immediately gives a criterium for finding periodic solutions:

Corollary 2.8. *Let $m_1 = m_2$ and $x \in \mathcal{D}$ with $S(x) = x$, let γ be solution of $(*)$ with initial condition x . If there are $T \neq 0$ and $k \in \mathbb{Z}$ so that*

$$S(\gamma(T)) = \gamma(T) + \begin{pmatrix} 0 \\ 0 \\ 2\pi k \end{pmatrix}$$

then the ξ and θ components of γ are periodic and $2T$ is a period.

Note that one has $S(\xi, \theta, \alpha) = (\xi, \theta, \alpha + 2\pi k)$ for some $k \in \mathbb{Z}$ if and only if $\theta = \theta^*$ and $\alpha = \frac{\pi}{2} + \pi j$ for some $j \in \mathbb{Z}$.

3 Existence of periodic curves

In light of Corollary 2.8 we wish to find geodesic segments that begin and end on the line $\theta = \theta^*$, with both intersections being orthogonal. We begin with the following definition:

Definition 3.1. For $\xi_0 \in \mathbb{R}$ let $(\theta_{\xi_0}(t), \xi_{\xi_0}(t), \alpha_{\xi_0}(t))$ denote the solution of $(*)$ with initial condition $\xi(0) = \xi_0, \theta(0) = \theta^*$ and $\alpha(0) = \frac{\pi}{2}$. Then:

- (i) ξ_0 is said to be of *type 1* if there is a $T > 0$ so that $\theta_{\xi_0}(T) = \theta^*$ and $\xi'_{\xi_0}(t) \neq 0$ for all $t \in (0, T)$.
- (ii) ξ_0 is said to be of *type 2* if there is a $T > 0$ so that $\xi'_{\xi_0}(T) = 0$ and $\theta_{\xi_0}(t) \neq \theta^*$ for all $t \in (0, T)$.
- (iii) ξ_0 is said to be of *type 3* if $\xi'_{\xi_0}(t) \neq 0$ and $\theta_{\xi_0}(t) \neq \theta^*$ for all $t > 0$.

Note that type 1 and type 2 are not exclusive, whereas a point is type 3 precisely if it is not type 1 or type 2. In fact a point that is both of type 1 and and type 2 corresponds

to a curve segment that orthogonally meets the $\theta = \theta^*$ line at its start and its end. If $m_1 = m_2$ this leads to a solution for which the ξ and θ components are periodic, which corresponds a closed embedded self-shrinker in \mathbb{R}^{n+1} of topological type $S^1 \times M$, here M is diffeomorphic to the leaves of the isoparametric foliation. The following argument then finds a value ξ_0^* that is both type 1 and type 2.




Figure 1: Examples of curves of different type. The blue curve is type 1 but not type 2, the green curve is both type 1 and type 2, the orange curve is type 2 but not type 1, the red curve is type 3.

Define:

$$\xi_0^* = \inf\{r \in \mathbb{R} \mid \xi_0 \text{ is type 1 for all } \xi_0 > r \}.$$

Proposition 3.2. *We have:*

- (i) $\xi_0^* < \infty$.
- (ii) $\xi_0^* > \frac{g}{4} \ln(m)$.
- (iii) ξ_0^* is not type 3.

This proposition will be proven in Section 4. For now we make use of the following elementary lemma:

Lemma 3.3. *For $(\xi, \theta)(t) \in \mathcal{D}$ one has the following characterisation of extrema:*

- (i) If $\xi'(t) = 0$, then $\text{sign}(\xi''(t)) = \text{sign}(\frac{g}{4} \ln(m) - \xi(t))$.
- (ii) If $\theta'(t) = 0$, then $\text{sign}(\theta''(t)) = \text{sign}(\theta^* - \theta)$.

Proof. $\xi'(t) = 0$ if and only if $\cos(\alpha(t)) = 0$, so

$$\xi''(t) = -\sin \alpha \sin(2\theta) \alpha'(t) = -\sin^2 \alpha \sin^2(2\theta) (e^{\frac{4}{g}\xi} - m).$$

In the same way $\theta'(t) = 0$ if and only if $\sin(\alpha(t)) = 0$, so

$$\theta''(t) = \cos \alpha \sin(2\theta) \alpha'(t) = 2 \cos^2 \alpha \sin(2\theta) l(\theta).$$

Now $l(\theta) > 0$ for $\theta < \theta^*$ and $l(\theta) < 0$ for $\theta > \theta^*$. □

This now gives:

Proposition 3.4. ξ_0^* is type 1 and type 2.

Proof. Since ξ_0^* is not type 3 by Proposition 3.2 (iii) it must be at least one of type 1 or type 2. We first assume that ξ_0^* is type 1 but not type 2, then we have a $T > 0$ so that for all $\epsilon > 0$ small enough one gets:

$$\cos(\alpha_{\xi_0^*}(t)) \neq 0 \quad \forall t \in [\epsilon, T + \epsilon], \quad \theta_{\xi_0^*}(T + \epsilon) < \theta^*.$$

Since solutions of the ODE (*) vary continuously in the initial conditions (with respect to the topology of uniform convergence on compacta) one finds a neighbourhood $U(\epsilon)$ of ξ_0^* so that for all $\xi_0 \in U(\epsilon)$ one has

$$\cos(\alpha_{\xi_0}(t)) \neq 0 \quad \forall t \in [\epsilon, T + \epsilon], \quad \theta_{\xi_0}(T + \epsilon) < \theta^*.$$

Additionally one has $|\frac{d}{dt} \xi_{\xi_0}(t)| \leq 1$, whence if one chooses ϵ small enough there is a neighbourhood V of $\xi_0^* > \frac{g}{4} \ln m$ with $\xi_{\xi_0}(t) > \frac{g}{4} \ln m$ for all $t \in [0, \epsilon]$ and $\xi_0 \in V$. By Lemma 3.3 $\xi(t)$ can only have maxima when $\xi(t) > \frac{g}{4} \ln m$, which implies that $\cos(\alpha_{\xi_0}(t)) \neq 0$ for all $t \in (0, \epsilon]$ and $\xi_0 \in V$.

The above shows that for $\xi_0 \in V \cap U(\epsilon)$ one has that ξ_0 is type 1, contradicting the definition of ξ_0^* .

An analogous argument shows that the assumption that ξ_0^* being type 2 but not type 1 leads to a neighbourhood of ξ_0^* consisting entirely of points that are type 2 but not type 1, which again contradicts the definition of ξ_0^* . □

As discussed, this yields a periodic geodesic via Corollary 2.8 in the case $m_1 = m_2$.

Lemma 3.5. *When $m_1 = m_2$ one has that the periodic geodesic $t \mapsto (\xi_{\xi_0^*}, \theta_{\xi_0^*})(t)$ is simple.*

Proof. Let $2T > 0$ denote the period of the geodesic, then $\theta(0) = \theta(T) = \theta^*$. For $t \in (0, T)$ this gives $\xi'_{\xi_0^*}(t) < 0$ and $\theta_{\xi_0^*}(t) > \theta^*$, similarly if $t \in (T, 2T)$ then $\xi'_{\xi_0^*}(t) > 0$ and $\theta_{\xi_0^*}(t) < \theta^*$. This immediately implies that the geodesic is simple. \square

Together with Proposition 2.4 this proves the main theorem of the paper:

Theorem A. For any isoparametric hypersurface M in S^n , $n \geq 2$, for which the multiplicities m_1 and m_2 of the principal curvatures agree there is a closed embedded self-shrinker of topological type $S^1 \times M$ in \mathbb{R}^{n+1} . This hypersurface is a union of homothetic copies of the leaves of the isoparametric foliation of S^n associated to M .

In the case $m_1 \neq m_2$ Proposition 3.4 remains true and yields a simple geodesic segment that starts and ends on the $\theta = \theta^*$, meeting this line orthogonally in both places. In between the two ends one has $\theta > \theta^*$ and the same arguments give another geodesic arc with the same properties, except now $\theta < \theta^*$.

It may be useful to connect the end-points of these two arcs by line segments $\theta = \theta^*$ (which are geodesics). Doing so gives a simple closed curve consisting piecewise geodesic segments and having external angle sum equal to 0 . By the Theorem of Gauß-Bonnet this curve then encloses a total Gauß curvature of 2π . Such a curve is an essential ingredient in [9], where an adapted curve shortening flow is used to generate closed geodesics.

3.1 The case studied by Angenent

The case $g = 1$ yields an embedded self-shrinker in \mathbb{R}^{n+1} of topological type $S^1 \times S^{n-1}$, which is invariant under an isometric $O(n)$ action on \mathbb{R}^{n+1} . This is the case investigated by Angenent in [2]. In this subsection we relate Angenent's construction to ours and show that they give the same self-shrinker.

Let $\omega := \{(x_0, \dots, x_n) \in \mathbb{R}^{n+1} \mid \sum_{i=0}^n x_i^2 = 1, x_0 = 0\}$, $e_0 := (1, 0, \dots, 0) \in \mathbb{R}^{n+1}$. Then the construction of [2] yields a self-shrinker of the form

$$\{x(t)e_0 + \tilde{r}(t)\omega \mid t \in (a, b)\}, \quad (5)$$

where $x : (a, b) \rightarrow \mathbb{R}$, $\tilde{r} : (a, b) \rightarrow \mathbb{R}_{>0}$ are smooth functions and $a, b \in \mathbb{R}$. Note that sets the form (5) are precisely the \mathbf{f} -invariant sets from Definition 2.2, where \mathbf{f} arises from

the isoparametric foliation of S^n with $V_1 = e_0$. To be more precise, if $r : (a, b) \rightarrow \mathbb{R}_{>0}$, $\varphi : (a, b) \rightarrow (0, \pi)$ one has:

$$\mathbf{f}^{-1}(\{(r(t), \varphi(t)) \mid t \in (a, b)\}) = \{r(t) \cos(\varphi(t)) e_0 + r(t) \sin(\varphi(t)) \omega \mid t \in (a, b)\}. \quad (6)$$

In [2] the variational condition that a set of the form (5) is a self-shrinker is reduced to $\{(x(t), r(t)) \mid t \in (a, b)\}$ being a geodesic segment in

$$(\mathbb{R} \times \mathbb{R}_{>0}, \tilde{r}^{2n-2} e^{-\tau(x^2 + \tilde{r}^2)}(dx^2 + d\tilde{r}^2)). \quad (7)$$

Here $\tau > 0$ is related to the extinction time and in [2] one has $\tau = \frac{1}{4}$. For ease of comparison we take $\tau = 1$ and then up to a constant conformal factor the transformation implicit in (6) gives an isometry to the metric (4) on $\mathbb{R}_{>0} \times (0, \pi)$, as is easy to see (recall $g = 1$). It follows that any geodesic of (7) is a reparametrisation of a geodesic in (4). Carrying out the additional coordinate changes of subsection 2.3 one sees that the following map sends the orbits of solutions of (*) to the orbits of geodesics of (7):

$$\Psi : \mathbb{R} \times (0, \frac{\pi}{2}) \rightarrow \mathbb{R} \times \mathbb{R}_{>0}, \quad (\xi, \theta) \mapsto (e^{\frac{2}{g}\xi} \cos(2\theta), e^{\frac{2}{g}\xi} \sin(2\theta)).$$

In particular $x(t) = 0$ if and only if $\theta = \frac{\pi}{4} = \theta^*$.

Denote with $(\tilde{r}_R, x_R)(t)$ the evolution of a geodesic in (7) with initial conditions $(\tilde{r}(0), x(0)) = (R, 0)$ and $(\tilde{r}'(0), x'(0)) = (0, 1)$. Define:

$$R_* := \inf\{\tilde{R} > 0 \mid \forall R > \tilde{R} : \exists t_1 > 0 \text{ so that } x_R(t_1) = 0 \text{ and } \tilde{r}'_R(t) < 0 \forall t \in (0, t_1)\}.$$

Angenent then shows that the geodesic $(\tilde{r}_{R_*}, x_{R_*})$ meets the line $x = 0$ orthogonally after a finite time. A symmetry argument as in Corollary 2.8 then shows that this yields a simple periodic geodesic.

In order to show Proposition B we start with the following lemma. It follows from elementary arguments using continuity of solutions of the relevant ODEs in initial conditions, similar to Proposition 3.4.

Lemma 3.6. *For any neighbourhood U of R_* there are $R \in U$ and $t_1(R) > 0$ so that*

$$x_R(t_1(R)) = 0, \quad \tilde{r}'_R(t_1(R)) > 0, \quad \text{and } x(t) > 0 \text{ for all } t \in (0, t_1(R)).$$

Similarly for any neighbourhood V of ξ_0^* there are $\xi_0 \in V$ and $T(\xi_0) > 0$ so that

$$\theta_{\xi_0}(T(\xi_0)) = \theta^*, \quad \xi'_{\xi_0}(T(\xi_0)) > 0, \quad \text{and } \theta_{\xi_0}(t) > \theta^* \text{ for all } t \in (0, T(\xi_0)).$$

Proposition B. In the case $g = 1$ the construction of Theorem A gives Angenent's shrinking doughnut [2].

Proof. Let $L : \mathbb{R} \rightarrow \mathbb{R}$ be the reparametrisation of a geodesic so that $\Psi((\xi, \theta)(L(t))) = (x, \tilde{r})(t)$. We take $L'(t) > 0$ for all t . Then if $x(t) = 0$ one has by an elementary calculation:

$$\text{sign}(\xi'(L(t))) = \text{sign}(\tilde{r}'(t)) \tag{8}$$

Assume first $R_* > e^{\frac{2}{g}\xi_0^*}$. Then apply the first part of Lemma 3.6 to R_* to get initial conditions R arbitrarily close to R_* and times $t_1(R)$ for which

$$x_R(t_1(R)) = 0, \quad \tilde{r}'_R(t_1) > 0, \quad \text{and } x_R(t) > 0 \text{ for all } t \in (0, t_1(R)).$$

This then transforms under Ψ^{-1} to values ξ_0 close to $\frac{g}{2} \ln(R_*)$ (which is larger than ξ_0^*), by (8) one then gets $\xi'_{\xi_0}(L(t_1)) > 0$, while $\theta_{\xi_0}(L(t)) > \theta^*$ for all $t \in (0, t_1)$.

These points are not type 1, contradicting the definition of ξ_0^* . The contradiction for $R_* < e^{\frac{2}{g}\xi_0^*}$ is similar. \square

4 Proof of Proposition 3.2

4.1 Crossings in finite time

In this section we prove two useful lemmas that expand on the analysis of extrema in Lemma 3.3. The lemmas state that if $\xi'(t)$ points towards the $\frac{g}{4} \ln m$ line then we reach this line in finite time, the same holding true for θ if $\theta'(t)$ points toward θ^* .

Lemma 4.1. *If for some $t_0 \in \mathbb{R}$ one has $\xi(t_0) > \frac{g}{4} \ln m$ ($\xi(t_0) < \frac{g}{4} \ln m$) and $\xi'(t_0) < 0$ ($\xi'(t_0) > 0$) then there exists a time $T \in (0, \infty)$ so that $\xi(t_0 + T) = \frac{g}{4} \ln m$.*

Proof. If this were not true then $\xi(t) > \frac{g}{4} \ln(m)$ for all $t > t_0$. By Lemma 3.3 we would then have that any extremum of ξ is a maximum when $t > t_0$. Since $\xi'(t_0) < 0$ it follows that ξ has no extrema for times $> t_0$ and then $\xi(t)$ is monotonically decreasing in t , hence it must converge.

Since $\xi(t)$ is bounded we find that $\alpha'(t)$ is bounded, whence $\xi'(t) = \cos(\alpha) \sin(2\theta)$ must converge to 0 (since $\theta'(t)$ is also automatically bounded). So either $\lim_{t \rightarrow \infty} \alpha(t) \in \frac{\pi}{2} + \pi\mathbb{Z}$ or $\lim_{t \rightarrow \infty} \theta(t) \in \{0, \frac{\pi}{2}\}$. We briefly show that the first condition implies the second and then continue working only with the second.

The condition $\alpha(t) \rightarrow \frac{\pi}{2} + k\pi$ for some $k \in \mathbb{Z}$ gives $\sin \alpha \rightarrow (-1)^k$. For large times the dynamics of $\theta(t)$ are then given by

$$\theta'(t) = (-1)^k \sin(2\theta) + O(\cos \alpha).$$

This gives that $\theta(t)$ converges to either 0 or $\frac{\pi}{2}$ as $t \rightarrow \infty$, depending on whether k is even or odd.

Assuming now $\theta(t) \rightarrow 0$ as $t \rightarrow \infty$ one gets from Lemma 3.3 that $\theta(t)$ admits no extrema for t large enough, so $\sin \alpha < 0$ for t large enough. This gives $\alpha(t) \in 2\pi\mathbb{Z} + (\pi, \frac{3\pi}{2})$ for t large enough. However for such t

$$\alpha'(t) = \sin \alpha \sin(2\theta)(e^{\frac{4}{g}\xi} - m) + 2 \cos \alpha l(\theta)$$

is a sum of two strictly negative terms. α then decreases for large times, so there is an $\epsilon > 0$ with $|\cos \alpha| > \epsilon$ for large enough t . In particular $\alpha'(t) < -2\epsilon l(\theta)$ for large t , where $l(\theta)$ converges to m_1 . This means that in finite time α exits the interval $2\pi k + (\pi, \frac{3\pi}{2})$ from the bottom, contradicting that $\sin \alpha < 0$ for all t large enough.

The case $\theta(t) \rightarrow \frac{\pi}{2}$ can be treated in the same way. This contradiction then implies the statement for $\xi(t_0) > \frac{g}{4} \ln m$ and $\xi'(t_0) < 0$. The case $\xi(t_0) < \frac{g}{4} \ln m$ and $\xi'(t_0) > 0$ is also completely analogous. \square

Lemma 4.2. *If for some t_0 one has $\theta(t_0) < \theta^*$ ($\theta(t_0) > \theta^*$) and $\theta'(t_0) > 0$ ($\theta'(t_0) < 0$) then there exists a time $T \in (0, \infty)$ so that $\theta(t_0 + T) = \theta^*$.*

Proof. As before assuming that $\theta(t) < \theta^*$ for all $t > t_0$ and $\theta(t_0) > 0$ implies that $\theta(t)$ converges.

We first assume that $e^{\frac{4}{g}\xi(t)}$ remains bounded as $t \rightarrow \infty$, this implies that $\alpha'(t)$ remains bounded and then from convergence of θ one gets that $\theta'(t) = \sin(\alpha) \sin(2\theta)$ converges to 0. Since $\sin(2\theta)$ remains bounded away from 0 one gets that α converges to some element of $\pi\mathbb{Z}$. Since $\xi(t)$ is not allowed to go to $+\infty$ one gets that $\lim_{t \rightarrow \infty} \alpha(t) \in \pi + 2\pi\mathbb{Z}$.

Performing a coordinate transform $R = \ln \xi$ the system of ODEs (*) becomes:

$$\begin{aligned}\theta'(t) &= \sin \alpha \sin(2\theta), \\ R'(t) &= \cos \alpha \sin(2\theta)R, \\ \alpha'(t) &= \sin \alpha \sin(2\theta)(R^{\frac{4}{g}} - m) + 2 \cos \alpha l(\theta).\end{aligned}$$

From $\alpha(t) \rightarrow \pi + 2\pi k$ for some $k \in \mathbb{Z}$ one gets $R(t) \rightarrow 0$ and $\theta(t) \rightarrow \theta^*$. But the fixpoint $(\theta, R, \alpha) = (\theta^*, 0, \pi + 2\pi k)$ is hyperbolic and at this point the above ODE has as linearisation:

$$\frac{d}{dt} \begin{pmatrix} \theta \\ R \\ \alpha \end{pmatrix} \approx \begin{pmatrix} 0 & 0 & -\sin(2\theta^*) \\ 0 & -\sin(2\theta^*) & 0 \\ -2l'(\theta^*) & 0 & \sin(2\theta^*)m \end{pmatrix} \begin{pmatrix} \theta - \theta^* \\ R \\ \alpha - (\pi + 2\pi k) \end{pmatrix}.$$

The system then has a one dimensional stable manifold - this is the line

$$(\theta(t), R(t), \alpha(t)) = (\theta^*, R(0) \exp(-\sin(2\theta^*)t), \pi + 2\pi k).$$

Since we are assuming $\theta(t) < \theta^*$ for all $t > t_0$ the solution cannot lie on the stable manifold, yielding a contradiction.

To complete the proof of the lemma we must show that $e^{\frac{4}{g}\xi(t)}$ cannot be unbounded under the hypothesis $\theta(t) < \theta^*$ for all $t > t_0$ and $\theta'(t_0) > 0$. First recall the graph form (**):

$$\frac{d^2\xi}{d\theta^2}(\theta) = - \left(1 + \left(\frac{d\xi}{d\theta} \right)^2 \right) \left(e^{\frac{4}{g}\xi} - m + 2H(\theta) \frac{d\xi}{d\theta} \right).$$

Whence if $\theta < \theta^*$, $\xi > \frac{g}{4} \ln m$ and $\frac{d\xi}{d\theta} > 0$ one has $\frac{d^2\xi}{d\theta^2} < 0$, even becoming arbitrarily negative if ξ becomes arbitrarily large. So if $\xi(t)$ is unbounded from above it cannot eventually be monotonic in θ (and hence in t by $\theta'(t) > 0$) and must admit maxima, in fact infinitely many such maxima. Between two maxima there must be a minimum, which can only happen for values of $\xi(t)$ less than $\frac{g}{4} \ln m$.

However at each minimum one has $\theta'(t) = \sin(2\theta)$, which may be bounded from below since θ stays away from $\{0, \frac{\pi}{2}\}$. Since the system (*) admits a Lipschitz constant on $\{(\theta, \xi, \alpha) \mid \xi < \frac{g}{4} \ln(m) + 1\}$ one finds that at each minimum of ξ the parameter θ increases by some positive number admitting a bound from below. This contradicts the assumption that $\theta(t) < \theta^*$ for all $t > t_0$.

The case $\theta(t_0) > \theta^*$, $\theta'(t_0) < 0$ can be treated analogously. \square

4.2 Proof of Proposition 3.2 (i)

The proof of Proposition 3.2 (i) is divided into two parts. First we show that for ξ_0 large enough there is a $T_2 > 0$ so that the solution to (*) with initial value $(\xi, \theta, \alpha)(t=0) = (\xi_0, \theta^*, \frac{\pi}{2})$ has the property:

$$\theta'(T_2) = 0, \quad \theta(T_2) - \theta^* < \frac{1}{\xi_0}, \quad \xi_0 - \xi(T_2) < \frac{1}{\xi_0}, \quad \text{while } \xi'(t) < 0 \text{ for all } t \in (0, T_2).$$

After such a time T_2 has been established one has by Lemma 4.2 that θ reaches θ^* in finite time and so ξ_0 cannot be of type 3. The proof then continues by contradiction, assuming that ξ_0 is not of type 1 means it must be of type 2. Being of type 2 means that ξ must travel all the way to some value $< \frac{g}{4} \ln(m)$ where we have an extremum of ξ - all the while θ is not allowed to cross the line θ^* .

The proof by contradiction is carried out in Lemma 4.6, here one assumes that conditions of this scenario have been set: there is some time T_3 at which $\xi(T_3) = \frac{4}{g} \ln(m)$ all the while $\theta(t) > \theta^*$ and $\xi'(t) < 0$ for $t \in (0, T_3]$. Using bounds for the value of $\theta(T_3)$ one is however able to show that even in this worst-case-scenario θ crosses the value θ^* before any extremum of ξ is possible, contradicting the assumption that ξ_0 is type 2. Hence, since it cannot be type 3, it must have been type 1.

In what follows ξ, θ and α will denote the components of the solution of (*) with initial condition $(\xi, \theta, \alpha)(0) = (\xi_0, \theta^*, \frac{\pi}{2})$. The proof begins by establishing an auxilliary time T_1 , at which $\frac{\xi'(T_1)}{\theta'(T_1)} = -1$.

Lemma 4.3. *If ξ_0 is large enough then there is a time $T_1 > 0$ so that $\frac{\xi'(T_1)}{\theta'(T_1)} = -1$ while $\theta'(t) > 0, \xi'(t) < 0$ for all $t \in (0, T_1]$.*

Proof. Note that one initially has $\frac{d}{dt} \cos \alpha|_{t=0} < 0$ whence one gets $\xi'(t) < 0$ for small t . By Lemma 4.1 $\xi(t)$ then descends to $\frac{g}{4} \ln m$ and does not have any extrema until after this value is reached, meaning $\cos \alpha(t) < 0$ for all $t \in (0, t_m]$ and some $t_m \in \mathbb{R}$ at which $\xi(t_m) = \frac{g}{4} \ln m$. For ξ_0 large enough there will then be some intermediate time $T_1 < t_m$ for which $\frac{d}{d\theta} \xi(\theta) = \frac{\xi'(t)}{\theta'(t)} = -1$ holds, since either $\theta'(t) = 0$ for some $t \in [0, t_m]$ or the graph $\xi(\theta)$ must descend from ξ_0 at θ^* to $\frac{g}{4} \ln m$ at some value $\theta < \frac{\pi}{2}$. In the second case the mean-value theorem implies that the graph achieves slope -1 at some point. \square




Figure 2: A sketch of the argument for Proposition 3.2(i). The black curve gives the evolution of (ξ, θ) up until the extremum of θ . The dashed blue line describes the worst-case scenario for the evolution of $(\xi, \theta)(t)$ after this extremum. The red line, which crosses the line $\theta = \theta^*$ without any extrema of ξ , is an estimate of actual evolution starting on a certain point of the worst-case scenario.

Lemma 4.4. *There are constants $c_1, c_2, c_3 \in \mathbb{R}_{>0}$ so that if ξ_0 is large enough one has*

$$c_1 e^{-\frac{4}{g}\xi_0} \leq \theta(T_1) - \theta^* \leq c_2 e^{-\frac{4}{g}\xi_0}, \quad \xi(T_1) \geq \xi_0 + c_3 e^{-\frac{4}{g}\xi_0}$$

Proof. Before beginning with the proper analysis one notes that by well definedness one has $\theta(T_1) \leq \frac{\pi}{2}$, whence by the mean value theorem $\xi_0 - \xi(T_1) \leq \frac{\pi}{2} - \theta^*$, some finite value bounded from above.

For the proof of this lemma it is more convenient to work with the following system of ODEs:

$$\begin{aligned} \xi'(t) &= \cos \alpha, \\ \theta'(t) &= \sin \alpha, \\ \alpha'(t) &= \sin \alpha (e^{\frac{4}{g}\xi} - m) + 2 \cos \alpha H(\theta). \end{aligned}$$

which one can get from (*) by rescaling time by the law $\frac{dt_{\text{new}}}{dt_{\text{old}}} = \sin(2\theta)$. To keep the number of superfluous parameters at a minimum we still use T_1 to denote the time at

which $\frac{\xi'(T_1)}{\theta'(T_1)} = -1$ in this new ODE. One then gets for all $t \in [0, T_1]$:

$$\cos \alpha(t) \in [-\frac{1}{\sqrt{2}}, 0], \quad \sin \alpha(t) \in [\frac{1}{\sqrt{2}}, 1].$$

Which implies:

$$\xi_0 - \xi(T_1) \in [0, \frac{1}{\sqrt{2}}T_1], \quad \theta(T_1) - \theta^* \in [\frac{1}{\sqrt{2}}T_1, T_1].$$

We then proceed by bounding T_1 from above and below. Noting that for $t \in [0, T_1]$:

$$\frac{1}{\sqrt{2}}(e^{\frac{4}{g}\xi(T_1)} - m) \leq \alpha'(t) \leq e^{\frac{4}{g}\xi_0} - \sqrt{2}H(\theta(T_1)).$$

Integrating the left inequality from 0 to T_1 yields:

$$\frac{T_1}{\sqrt{2}}(e^{\frac{4}{g}\xi(T_1)} - m) \leq \alpha(T_1) - \alpha(0) = \frac{\pi}{4}.$$

Recalling that $\xi_0 - \xi(T_1) \leq \frac{\pi}{2} - \theta^*$ one gets $T_1 \leq d_1 e^{-\frac{4}{g}\xi_0}$ for an appropriate constant d_1 . This implies $\theta(T_1) - \theta^* \leq c_2 e^{-\frac{4}{g}\xi_0}$ and $\xi_0 - \xi(T_1) \leq c_3 e^{-\frac{4}{g}\xi_0}$ for appropriate c_2, c_3 .

Combining $\theta(T_1) - \theta^* \leq c_2 e^{-\frac{4}{g}\xi_0}$ with $H(\theta^*) = 0$ gives for ξ_0 large enough that $-\sqrt{2}H(\theta(T_1)) \leq d_2 e^{-\frac{4}{g}\xi_0}$ for some other constant d_2 . Integrating the other inequality for $\alpha'(t)$ from 0 to T_1 then gives:

$$\frac{\pi}{4} \leq T_1(e^{\frac{4}{g}\xi_0} + d_2 e^{-\frac{4}{g}\xi_0}) \implies T_1 \geq d_3 e^{-\frac{4}{g}\xi_0}$$

for another constant d_3 , provided ξ_0 is large enough. This yields the final bound of the lemma, namely: $c_1 e^{-\frac{4}{g}\xi_0} \leq \theta(T_1) - \theta^*$. \square

Lemma 4.5. *For ξ_0 large enough there is a time $T_2 > T_1$ so that $\theta'(T_2) = 0$, $\xi(T_2) > \xi_0 - \frac{1}{\xi_0}$ while $\theta(t) \in \theta^* + (0, \frac{1}{\xi_0})$ and $\xi'(t) < 0$ for all $t \in (0, T_2]$.*

Proof. For all $t \in (0, \frac{1}{\xi_0})$ one has from $|\xi'(t)| \leq 1$ that $\xi(t) > \xi_0 - \frac{1}{\xi_0}$, so for ξ_0 large enough $\xi'(t) < 0$ for such t . Further as long as $\theta > \theta^*$ and $\alpha \in [\frac{3\pi}{4}, \pi)$ one has for such t that

$$\alpha'(t) = \sin \alpha \sin(2\theta)(e^{\frac{4}{g}\xi} - m) + 2 \cos \alpha l(\theta)$$

is a sum of two positive terms and so α is increasing. Recalling that by definition

$\alpha(T_1) = \frac{3\pi}{4}$ and assuming that $\alpha(t) < \pi$ (i.e. $\theta'(t) > 0$) for all $t \in (T_1, \frac{1}{\xi_0})$ yields:

$$\alpha'(t) > Ae^{\frac{4}{g}(\xi_0 - \frac{1}{\xi_0})}(\pi - \alpha).$$

Here $A > 0$ is some constant. For this estimate we used that $\theta(t)$ is bounded away from $\{0, \frac{\pi}{2}\}$ for $t \in (0, \frac{1}{\xi_0})$, which follows from $|\theta'(t)| \leq 1$. From the intermediate value theorem one then gets a $\tilde{t} \in (T_1, \frac{1}{2\xi_0})$ so that

$$\alpha(\tilde{t}) = \pi - A \exp\left(-e^{\frac{4}{g}(\xi_0 - \frac{1}{\xi_0})} \frac{1}{2\xi_0}\right).$$

For ξ_0 large enough one then finds $\alpha(\tilde{t}) > \pi - e^{-\xi_0^2}$. Now $\theta(T_1) > \theta^* + c_1 e^{-\frac{4}{g}\xi_0}$ from Lemma 4.4, so $l(\theta(T_1)) \leq c_1 l'(\theta^*) e^{-\frac{4}{g}\xi_0}$ for ξ_0 large enough. By assumption $\theta(\tilde{t}) > \theta(T_1)$, so another estimate yields for $t \in (\tilde{t}, \frac{1}{\xi_0})$:

$$\alpha'(t) > B e^{-\frac{4}{g}\xi_0}$$

where $B > 0$ is some constant incorporating the $2 \cos(\alpha)$ term (which is close to -2) and $c_1 l'(\theta^*)$ (which is bounded away from 0). This then yields

$$\alpha\left(\tilde{t} + \frac{1}{2\xi_0}\right) > \pi - e^{-\xi_0^2} + B \frac{e^{-\frac{4}{g}\xi_0}}{2\xi_0}$$

which is larger than π , contradicting our assumption that $\alpha(t) < \pi$ for all $t \in (T_1, \frac{1}{\xi_0})$. The lemma then follows. \square

We will now prove that ξ_0 is of type 1 by contradiction.

Lemma 4.6. *If ξ_0 is large enough then it is of type 1.*

Proof. If ξ_0 is large enough by Lemma 4.5 there is a $T_2 > 0$ for which $\theta'(T_2) = 0$, which corresponds to a local maximum of θ . This means θ reaches θ^* in finite time by Lemma 4.2 and so ξ_0 is not type 3. We now assume it is not type 1, so it must be type 2.

Hence $\xi(t)$ has an extremum before $\theta(t)$ reaches θ^* . Since this must extremum must be a minimum one has a time $T_3 > T_2$ for which $\xi(T_3) = \frac{g}{4} \ln m$, while $\theta(t) > \theta^*$ for all $t \in (0, T_3]$. We first show that if ξ_0 is large enough one has $\cos \alpha(T_3) < -\frac{1}{2}$:

If this were not the case then $\sin \alpha(T_3) < -\frac{\sqrt{3}}{2}$. Note that $\alpha'(t)$ is bounded on $\{(\xi, \theta, \alpha) \in \mathbb{R}^3 \mid \xi < \frac{g}{4} \ln m + 1\}$ and that $|\xi'(t)| \leq 1$. From $\sin \alpha(T_3) < -\frac{\sqrt{3}}{2}$ there is then some $b < 1$

independent of ξ_0 so that

$$\sin \alpha(t) \leq -\frac{\sqrt{3}}{4}$$

for all $t \in [T_3 - b, T_3]$. Further $\sin(2\theta(t)) > \frac{1}{2} \sin(2\theta^*)$ for $t \in [T_3 - b, T_3]$ provided ξ_0 is large enough, since θ is confined to $[\theta^*, \theta(T_2)]$. Then from the mean value theorem one gets a $\tilde{t} \in [T_3 - b, T_3]$ for which:

$$(\theta(T_3) - \theta^*) - (\theta(T_2) - \theta^*) \leq \theta(T_3) - \theta(T_3 - b) = \theta'(\tilde{t}) \leq -\frac{\sqrt{3} \sin(2\theta^*)}{8}.$$

Since $\theta(T_2) - \theta^*$ becomes arbitrarily small one sees that $\theta(T_3) - \theta^*$ has to become negative, contradicting the assumption that ξ_0 was not type 1.

Then $\cos \alpha(T_3) < -\frac{1}{2}$ and there is a $T_4 > T_3$ for which $\cos \alpha(T_4) = -\frac{1}{2}$, all the while $\theta^* < \theta(t) < \theta(T_2)$ and $\xi'(t) \neq 0$ for $t \in (T_2, T_4]$. Repeating the previous argument gives also here the contradiction $\theta(T_4) \leq \theta^*$, finishing the proof. \square

4.3 Proof of Proposition 3.2 (ii)

We consider the solution curve with initial condition $\xi_0 = \frac{g}{4} \ln m + \epsilon$ and show that this is not of type 1 for ϵ sufficiently small. To do this we assume that it is of type 1 - only to later arrive at a contradiction. If it were of type 1 then there is a $T > 0$ with $\theta(T) = \theta^*$ and $\xi'(t) < 0$ for all $t \in (0, T)$. Since θ can only have maxima when $\theta > \theta^*$, we find that the trajectory $\{(\theta, \xi)(t) \mid t \in [0, T]\}$ must be the union of two graphs of ξ over θ . The maximum of θ occurs at the point denoted by (ξ_2, θ_2) in Figure 3.

In the upper graph one has that the slope $\frac{d\xi}{d\theta}$ starts at 0 and must go to $-\infty$ (which occurs when $\theta'(t) = 0$). Along the way ξ' has been negative and one verifies that ξ_2 has decreased to a value far enough below $\frac{g}{4} \ln m$ (c.f. Lemma 4.10). When we then switch to the lower graph the $e^{\frac{4}{g}\xi} - m$ term in the ODE for $\frac{d^2\xi}{d\theta^2}(\theta)$ will be large enough to push $\frac{d\xi}{d\theta}$ over the value 0 before θ reaches θ^* , contradicting the assumption that ξ_0 was type 1.

Note that this does not prove that $\frac{g}{4} \ln m + \epsilon$ is of type 2, because the proof by contradiction assumes that $\theta(t)$ has a maximum.

In what follows ξ, θ and α will denote the components of the solution of (*) with initial condition $(\xi, \theta, \alpha)(0) = (\frac{4}{g} \ln m + \epsilon, \theta^*, \frac{\pi}{2})$. The proof begins by investigating an auxilliary value θ_1 , which is defined to be the least (and for small ϵ only) value of θ for which one has $\frac{d\xi}{d\theta}(\theta_1) = -1$ in the upper graph.




Figure 3: The figure sketches the argument for Proposition 3.2(ii). The dashed blue line denotes the form that $(\xi, \theta)(t)$ must be if the initial condition were type 1. The red line, which has an extremum of ξ , is an estimate of the actual evolution starting at (ξ_2, θ_2) .

Lemma 4.7. *If $\xi_0 = \frac{g}{4} \ln m + \epsilon$ is type 1, then there are $T_2(\epsilon) > T_1(\epsilon) > 0$ so that $\theta'(T_2) = 0$ and $\frac{\xi'(T_1)}{\theta'(T_1)} = -1$, while $\xi'(t) < 0$ and $\theta'(t) > 0$ for all $t \in (0, T_2)$.*

Proof. Assuming that ξ_0 is of type 1 means that there is a time $T > 0$ for which $\theta(T) = \theta^*$ and $\xi'(t) \neq 0, \theta(t) \neq \theta^*$ for all $t \in (0, T)$. Since $\theta'(0) = \sin(2\theta^*) > 0$ one finds that $\theta'(t) > 0$ for small t , whence $\theta(t)$ must go through an extremum before it can go back to θ^* and there is a $T_2 < T$ so that $\theta'(T_2) = 0$. On the other hand one has $\xi''(0) = -\sin(2\theta^*)m(e^{\frac{4}{g}\epsilon} - 1) < 0$, whence $\xi'(t) < 0$ for all $t \in (0, T]$, in particular for all $t \in (0, T_2)$.

This means that $\frac{\xi'(t)}{\theta'(t)}$ is 0 at $t = 0$ and diverges to $-\infty$ at $t = T_2$. There must then be a T_1 so that $\frac{\xi'(T_1)}{\theta'(T_1)} = -1$. \square

Lemma 4.8. *For ϵ small enough there is only one pair (T_1, T_2) satisfying the conditions of Lemma 4.7 and $\theta(T_1) \rightarrow \frac{\pi}{2}$ as $\epsilon \rightarrow 0$.*

Proof. The time T_2 is obviously unique.

On the other hand the initial condition $\epsilon = 0$ has as solution the line $(\theta, \xi, \alpha) = (\arctan(\tan(\theta^*)e^{2t}), \frac{g}{4} \ln m, \frac{\pi}{2})$. So one finds that as $\epsilon \rightarrow 0$ the solution and its derivatives converge uniformly on compacta to the above curve, in particular for on any finite interval $[0, T]$ one can make $\frac{\xi'(t)}{\theta'(t)}$ arbitrarily small for all $t \in [0, T]$ (by taking ϵ small), while $\theta(T)$ is arbitrarily close to $\frac{\pi}{2}$ (by taking T large and then ϵ small). This means that as $\epsilon \rightarrow 0$ one must have $\frac{\pi}{2} - \theta(T_1(\epsilon)) \rightarrow 0$, where T_1 is any of the times satisfying Lemma 4.7.

Looking however at the graph ODE (**)

$$\frac{d^2\xi}{d\theta^2} = -\left(1 + \left(\frac{d\xi}{d\theta}\right)^2\right)\left(e^{\frac{4}{g}\xi} - m + 2H(\theta)\frac{d\xi}{d\theta}\right)$$

it is clear that if $\frac{d\xi}{d\theta} \leq -1$ and θ is close enough to $\frac{\pi}{2}$ while ξ is not too large that then $\frac{d^2\xi}{d\theta^2} < 0$. For ϵ small enough one gets that for any pair (T_1, T_2) satisfying Lemma 4.7 there is no pair (T'_1, T_2) satisfying the lemma with $T'_1 > T_1$. \square

This allows us to introduce the following notation:

$$\theta_1 := \theta(T_1), \quad \theta_2 := \theta(T_2), \quad \xi_2 := \xi(T_2).$$

Lemma 4.9. *For ϵ small enough one has that $\xi(T_1) \leq \frac{g}{4} \ln m$.*

Proof. We assume that $\xi(T_1) > \frac{g}{4} \ln m$ and get a contradiction. The first step is to show that this assumption leads to constants $c_1, c_2 > 0$ (independent of ϵ) so that for ϵ small enough one has:

$$c_1\epsilon < \left(\frac{\pi}{2} - \theta_1\right)^{m_2} < c_2\epsilon. \quad (9)$$

Since $\xi'(t) \leq 0$ for all $t \in (0, T_2)$ we may assume $\xi(t) > \frac{g}{4} \ln m$ for such t . Then from the graph ODE

$$\frac{d^2\xi}{d\theta^2} = -\left(1 + \left(\frac{d\xi}{d\theta}\right)^2\right)\left(e^{\frac{4}{g}\xi} - m + 2H(\theta)\frac{d\xi}{d\theta}\right)$$

one gets that $\frac{d^2\xi}{d\theta^2} < 0$ for all $\theta \in (\theta^*, \theta_1)$ and then $\frac{d\xi}{d\theta}$ is strictly decreasing in this interval. The mean value theorem then implies both

$$\frac{d\xi}{d\theta} \left(\frac{\theta_1 - \theta^*}{2}\right) > -\epsilon \frac{\theta_1 - \theta^*}{2}, \quad \xi \left(\frac{\theta_1 - \theta^*}{2}\right) > \frac{g}{4} \ln m + \frac{\epsilon}{2}. \quad (10)$$

Combining the second inequality with the graph ODE yields (recall that $H(\theta) < 0$ if $\theta > \theta^*$) additionally the bound $\frac{d\xi}{d\theta} \left(\frac{\theta_1 - \theta^*}{2}\right) < -\frac{m\epsilon}{g}(\theta_1 - \theta^*) + O(\epsilon^2)$. Using that $\theta_1 \rightarrow \frac{\pi}{2}$ as $\epsilon \rightarrow 0$ then gives constants \tilde{c}_1, \tilde{c}_2 so that

$$\tilde{c}_1\epsilon < \left| \frac{d\xi}{d\theta} \left(\frac{\theta_1 - \theta^*}{2}\right) \right| < \tilde{c}_2\epsilon. \quad (11)$$

Rewriting the graph ODE as

$$\frac{\frac{d^2\xi}{d\theta^2}}{\frac{d\xi}{d\theta}(1 + (\frac{d\xi}{d\theta})^2)} = -\frac{e^{\frac{4}{g}\xi} - m}{\frac{d\xi}{d\theta}} - 2H(\theta), \quad (12)$$

one notes that the first term on the right-hand side is $O(1)$ in the interval $(\frac{\theta_1 - \theta^*}{2}, \theta_1)$ by (11) and monotonicity of $\frac{d\xi}{d\theta}$. Integrating (12) over this interval then gives

$$\ln\left(\left|\frac{d\xi}{d\theta}\left(\frac{\theta_1 - \theta^*}{2}\right)\right|\right) + O(1) = m_2 \ln(\cos(\theta_1)) + O(1).$$

The approximation $\cos(\theta) = \frac{\pi}{2} - \theta + O((\frac{\pi}{2} - \theta)^3)$ then gives the bounds (9) from (11).

For the next step note that $H(\theta) - (\frac{d\xi}{d\theta}(\theta))^{-1}$ becomes $+\infty$ as $\theta \rightarrow \theta^*$ and $H(\theta_1) + 1$ as $\theta \rightarrow \theta_1$, which for ϵ small enough will be negative. Hence for small enough ϵ there is a $\theta_0 \in (\theta^*, \theta_1)$ so that $H(\theta_0) = (\frac{d\xi}{d\theta}(\theta_0))^{-1}$. In the same way as Lemma 4.8 one shows that $\theta_0 \rightarrow \frac{\pi}{2}$ as $\epsilon \rightarrow 0$.

We let

$$q := \frac{\pi/2 - \theta_0}{\pi/2 - \theta_1}$$

and show next that there is a constant c_3 so that $q^{m_2} \geq \frac{c_3}{\pi/2 - \theta_0}$, and hence that q grows unboundedly as $\epsilon \rightarrow 0$.

To show this we again integrate (12), this time from θ_0 to $\theta \leq \theta_1$. The result is:

$$\ln\left(\left|\frac{d\xi}{d\theta}(\theta_0)\right|\right) - \ln\left(\left|\frac{d\xi}{d\theta}(\theta)\right|\right) + O(1) = m_2 \ln\left(\frac{\cos(\theta)}{\cos(\theta_0)}\right) + O(1).$$

Noting that $\frac{d\xi}{d\theta}(\theta_0) = \frac{1}{H(\theta_0)} \approx -m_2(\frac{\pi}{2} - \theta_0) + O((\frac{\pi}{2} - \theta_0)^2)$ and again using the expansion of cosine close to $\frac{\pi}{2}$ implies the existence of $c_3, c_4 > 0$ so that:

$$c_4 \left|\frac{d\xi}{d\theta}(\theta)\right| \geq \left(\frac{\pi/2 - \theta_0}{\pi/2 - \theta}\right)^{m_2} (\frac{\pi}{2} - \theta) \geq c_3 \left|\frac{d\xi}{d\theta}(\theta)\right|. \quad (13)$$

Taking $\theta = \theta_1$ shows $q^{m_2} \geq \frac{c_3}{\pi/2 - \theta_0}$, in particular $q \rightarrow \infty$ as $\epsilon \rightarrow 0$. On the other hand, if one integrates $\frac{d\xi}{d\theta}(\theta)$ over (θ_0, θ_1) one finds by (13) that

$$c_4 |\xi(\theta_1) - \xi(\theta_0)| \geq \begin{cases} (\frac{\pi}{2} - \theta_0)^2 \ln q & m_2 = 1 \\ \frac{1}{m_2 - 1} (\frac{\pi}{2} - \theta_0)^2 (q^{m_2 - 1} - 1) & m_2 > 1 \end{cases}.$$

Noting that $(\frac{\pi}{2} - \theta_0)^2 = \frac{(\frac{\pi}{2} - \theta_1)^{m_2}}{(\frac{\pi}{2} - \theta_0)^{m_2}} q^{m_2} (\frac{\pi}{2} - \theta_0)^2 \geq c_1 c_4 \epsilon (\frac{\pi}{2} - \theta_0)^{1-m_2}$ and using unbound-
edness of q one sees that $|\xi(\theta_1) - \xi(\theta_0)|$ will be larger than ϵ , contradicting our assumption
that $\xi(\theta_1) \geq \frac{g}{4} \ln(m)$. \square

Lemma 4.10. *For θ_1 sufficiently close to $\frac{\pi}{2}$ one has that*

$$\xi(T_2) \leq \frac{g}{4} \ln(m) - \left(2^{\frac{1}{2m_2}} - 1\right) \left(\frac{\pi}{2} - \theta_2\right) + O\left(\left(\frac{\pi}{2} - \theta_2\right)^2\right).$$

Proof. By Lemma 4.9 one has that $\xi(T_1) \leq \frac{g}{4} \ln(m)$. Additionally for ϵ small enough
 $\frac{d^2\xi}{d\theta^2}(\theta) < 0$ for all $\theta \in (\theta_1, \theta_2)$, whence $\xi(T_2) \leq \xi(T_1) - (\theta_2 - \theta_1)$ and the statement reduces
to checking $\theta_2 - \theta_1 = \left(2^{\frac{1}{2m_2}} - 1\right) \left(\frac{\pi}{2} - \theta_2\right) + O\left(\left(\frac{\pi}{2} - \theta_2\right)^2\right)$.

From the graph ODE (**) one recovers:

$$\frac{\frac{d^2\xi}{d\theta^2}}{\frac{d\xi}{d\theta} \left(1 + \left(\frac{d\xi}{d\theta}\right)^2\right)} = -\frac{e^{\frac{4}{g}\xi} - m}{\frac{d\xi}{d\theta}} - 2H(\theta).$$

The absolute value of the first summand on the right-hand side is bounded by m over the
interval (θ_1, θ_2) , so integrating the equation from θ_1 to θ_2 gives:

$$\frac{1}{2} \ln(2) = O\left(\frac{\pi}{2} - \theta_1\right) - m_2 \ln\left(\frac{\cos(\theta_2)}{\cos(\theta_1)}\right) - m_1 \ln\left(\frac{\sin(\theta_2)}{\sin(\theta_1)}\right).$$

As θ becomes arbitrarily close to $\frac{\pi}{2}$ we use that $\cos(\theta) = \frac{\pi}{2} - \theta + O\left(\left(\frac{\pi}{2} - \theta\right)^3\right)$, $\sin(\theta) =$
 $1 + O\left(\left(\frac{\pi}{2} - \theta\right)^2\right)$ to get:

$$\frac{1}{2} \ln(2) = -m_2 \ln\left(\frac{\frac{\pi}{2} - \theta_2}{\frac{\pi}{2} - \theta_1}\right) + O\left(\frac{\pi}{2} - \theta_1\right).$$

Together with some arithmetic this implies the statement about $\theta_2 - \theta_1$. \square

Lemma 4.11. *If ϵ is small enough then $\xi_0 = \frac{g}{4} \ln m + \epsilon$ is not type 1.*

Proof. As noted before the assumption that $\xi_0 = \frac{g}{4} \ln(m) + \epsilon$ is type 1 leads to (ξ, θ)
being the union of two graphs of ξ over θ . In the previous lemmas we investigated the
upper graph and found that it ends at the turning point (ξ_2, θ_2) .

The lower graph is then determined by the graph ODE and the initial conditions $\xi(\theta_2) = \theta_2$,
 $\lim_{\theta \rightarrow \theta_2^-} \frac{d\xi}{d\theta}(\theta) = +\infty$. The assumption that ξ_0 is type 1 necessitates that for all $\theta \in (\theta^*, \theta_2)$

one has $\frac{d\xi}{d\theta}(\theta) > 0$ for the lower graph. In particular if we integrate

$$\frac{\frac{d^2\xi}{d\theta^2}}{\frac{d\xi}{d\theta}(1 + (\frac{d\xi}{d\theta})^2)} = -\frac{e^{\frac{4}{g}\xi} - m}{\frac{d\xi}{d\theta}} - 2H(\theta) \quad (14)$$

from $\frac{\theta^* + \theta_2}{2}$ to θ_2 one gets

$$-\ln\left(\frac{\frac{d\xi}{d\theta}(\frac{\theta^* + \theta_2}{2})}{\sqrt{1 + (\frac{d\xi}{d\theta})^2}}\right) \geq -m_2 \ln\left(\frac{\pi}{2} - \theta_2\right) + O(1),$$

where we used $e^{\frac{4}{g}\xi(\theta)} - m \leq e^{\frac{4}{g}\xi_2} - m \leq 0$. Inverting the above inequality gives the existence of a $c_1 > 0$ so that:

$$\frac{d\xi}{d\theta}\left(\frac{\theta^* + \theta_2}{2}\right) \leq \frac{c_1(\frac{\pi}{2} - \theta_2)^{m_2}}{\sqrt{1 - c_1^2(\frac{\pi}{2} - \theta_2)^{2m_2}}} \leq 2c_1(\frac{\pi}{2} - \theta_2)^{m_2}. \quad (15)$$

For $m_2 > 1$ this implies the lemma, since for all $\theta \in (\theta^*, \theta_2)$ one has

$$\frac{d^2\xi}{d\theta^2}(\theta) > -(e^{\frac{4}{g}\xi(\theta)} - m) \geq c_2\left(\frac{\pi}{2} - \theta_2\right)$$

with $c_2 > 0$ some constant by Lemma 4.10. For $m_2 = 1$ one gets first from (15) that $\frac{d\xi}{d\theta}$ takes on all values in $(\frac{c_1}{2}(\frac{\pi}{2} - \theta_2), \infty)$ as θ varies from $\frac{\theta^* + \theta_2}{2}$ to θ_2 . In particular for θ_2 close enough to $\frac{\pi}{2}$ there exist θ_3, θ_4 so that

$$\frac{d\xi}{d\theta}(\theta_3) = 1, \quad \frac{d\xi}{d\theta}(\theta_4) = -\frac{1}{H(\theta_4)}.$$

By integrating (14) one shows that θ_3, θ_4 get arbitrarily close to $\frac{\pi}{2}$ if $\frac{\pi}{2} - \theta_2$ is small enough. Letting $q := \frac{\pi/2 - \theta_4}{\pi/2 - \theta_2}$ one gets again that q grows unboundedly as θ_2 approaches $\frac{\pi}{2}$ by integrating (14) from θ_4 to θ_2 as in Lemma 4.9.

Integrating (14) from θ_3 to θ_2 gives a constant c_3 so that $c_3(\frac{\pi}{2} - \theta_2) \leq (\frac{\pi}{2} - \theta_3)$, just as in Lemma 4.10. Finally if we integrate (14) from $\theta \geq \theta_4$ to θ_3 one gets:

$$-\ln\left(\frac{d\xi}{d\theta}(\theta)\right) + O(1) = -\ln\left(\frac{\pi/2 - \theta_3}{\pi/2 - \theta}\right) + O\left(\frac{\pi}{2} - \theta\right).$$

Integrating the resulting bound for $\frac{d\xi}{d\theta}$ from θ_3 to θ_4 yields a constant c_4 so that

$$\xi(\theta_4) - \xi(\theta_3) \geq c_3\left(\frac{\pi}{2} - \theta_2\right) \ln q.$$

In particular this gives for any $\theta \in (\theta^*, \frac{\theta^* + \theta_2}{2})$ that

$$\frac{d^2\xi}{d\theta^2}(\theta) \geq m - e^{\frac{4}{g}\xi(\theta_3)} \geq m(1 - e^{-\frac{4}{g}c_3(\frac{\pi}{2} - \theta_2) \ln q}).$$

Together with the bound (15) for $\frac{d\xi}{d\theta}(\frac{\xi^* + \theta_2}{2})$ this yields $\frac{d\xi}{d\theta}(\theta) = 0$ for some $\theta \in (\theta^*, \frac{\theta^* + \theta_2}{2})$, provided $\frac{\pi}{2} - \theta_2$ is small enough. This contradicts our assumption that ξ_0 was type 1. \square

4.4 Proof of Proposition 3.2 (iii)

Lemma 4.12. *If ξ_0^* is of type 3 then:*

- (i) *There is a $\delta > 0$ and a $T > 0$ so that $\xi_{\xi_0^*}(t) < \frac{g}{4} \ln(m) - \delta$ for all $t > T$.*
- (ii) *$\theta'_{\xi_0^*}(t) > 0$ for all $t > 0$.*
- (iii) *$\lim_{t \rightarrow \infty} \theta_{\xi_0^*}(t) = \frac{\pi}{2}$.*

Proof. Since $\xi_0^* > \frac{g}{4}$ and $\xi''_{\xi_0^*}(0) = -\sin(2\theta^*)(e^{\frac{4}{g}\xi_0^*} - m) < 0$, $\xi'_{\xi_0^*}(0) = 0$ one has that $\xi'_{\xi_0^*}(t)$ becomes negative for small times t . Then by Lemma 4.1 it must reach $\frac{g}{4} \ln(m)$ in finite time. Since $\xi'_{\xi_0^*}(t) < 0$ for all $t > 0$ this yields part (i).

One also has that $\theta'_{\xi_0^*}(t) \neq 0$ for all t , as otherwise Lemma 4.2 implies that $\theta_{\xi_0^*} = \theta^*$ in finite time. This gives part (ii).

$\theta_{\xi_0^*}$ is then monotonous in t and bounded by $\frac{\pi}{2}$, whence it converges. Since ξ_0^* is bounded above one has that all derivatives of the parameters are bounded and hence $\theta'_{\xi_0^*}(t) = \sin \alpha \sin(2\theta) \rightarrow 0$. It is clear that $\sin \alpha$ cannot converge to 0 as $\alpha'(t)$ would then be asymptotically equal to $-2l(\theta)$, which does not converge to 0. Hence $\theta \rightarrow \frac{\pi}{2}$, giving part (iii). \square

Lemma 4.13. *ξ_0^* is not of type 3.*

Proof. Let $(\xi_\epsilon(t), \theta_\epsilon(t))$ denote the solution to (*) with initial condition $(\xi, \theta, \alpha)(t=0) = (\xi_0^* + \epsilon, \theta^*, \frac{\pi}{2})$ where $\epsilon > 0$. Note that $\xi_0^* + \epsilon$ is type 1 by the definition of ξ_0^* , in particular there is a $T_1(\epsilon)$ so that $\theta_\epsilon(t)$ has a local maximum and $\xi'_\epsilon(T_1) < 0$ for all $t \in (0, T_1]$.

Assuming that ξ_0^* is of type 3 and using that ξ_ϵ and θ_ϵ converge uniformly on compacta to $\xi_{\epsilon=0}$ and $\theta_{\epsilon=0}$ as $\epsilon \rightarrow 0$ one finds that $\theta_\epsilon(T_1) \rightarrow \frac{\pi}{2}$ as $\epsilon \rightarrow 0$. This also implies $T_1(\epsilon) \rightarrow \infty$ for $\epsilon \rightarrow 0$, giving for ϵ small enough that one has $\xi_\epsilon(T_1) < \frac{g}{4} \ln(m) - \delta$.

After the extremum at $\theta_\epsilon(T_1)$ one has that ξ_ϵ becomes a graph over θ because no more extrema of θ are possible until after $\theta = \theta^*$. As in Lemma 4.9 the graph ODE for ξ yields:

$$\frac{\frac{d^2 \xi_\epsilon}{d\theta^2}}{\frac{d\xi_\epsilon}{d\theta} \left(1 + \left(\frac{d\xi_\epsilon}{d\theta}\right)^2\right)} = -\frac{e^{\frac{4}{g}\xi_\epsilon} - m}{\frac{d\xi_\epsilon}{d\theta}} - 2H(\theta).$$

Integrating this for the lower graph from $x := \frac{\theta^* + \pi/2}{2}$ to $\theta_\epsilon(T_1)$ one gets

$$-\ln \left(\frac{d\xi_\epsilon/d\theta}{\sqrt{1 + \left(\frac{d\xi_\epsilon}{d\theta}\right)^2}}(x) \right) \geq -m_2 \ln\left(\frac{\pi}{2} - \theta_\epsilon(T_1)\right) + O(1)$$

where $-\frac{e^{\frac{4}{g}\xi} - m}{\frac{d\xi}{d\theta}} \geq 0$ and $\frac{d\xi}{d\theta}(\theta(T_1)) = +\infty$ were used.

As $\epsilon \rightarrow 0$ this implies that $\frac{d\xi_\epsilon}{d\theta}(x)$ becomes arbitrarily small, in particular we may assume it to be smaller than $\frac{4m}{g} \frac{1}{2}(x - \theta^*)\delta$. Remarking however that if $\xi < \frac{g}{4} \ln(m) - \delta$ then $-(e^{\frac{4}{g}\xi} - m) > \frac{4}{g}m\delta$, one gets:

$$\frac{d^2 \xi_\epsilon}{d^2 \theta}(\theta) > \frac{4}{g}m\delta$$

for all $\theta \in (\theta^*, x)$. With $\frac{d\xi_\epsilon}{d\theta}(x) < \frac{4m}{g} \frac{1}{2}(x - \theta^*)\delta$ one immediately gets $\frac{d\xi_\epsilon}{d\theta}\left(\frac{\theta^* + x}{2}\right) < 0$, contradicting that ξ_ϵ is type 1. \square

References

- [1] Uwe Abresch, *Isoparametric hypersurfaces with four or six distinct principal curvatures. Necessary conditions on the multiplicities*, Math. Ann. **264** (1983), no. 3, 283–302.
- [2] Sigurd B. Angenent, *Shrinking doughnuts*, Nonlinear diffusion equations and their equilibrium states, 3 (Gregynog, 1989), Progr. Nonlinear Differential Equations Appl., vol. 7, Birkhäuser Boston, Boston, MA, 1992, pp. 21–38.
- [3] Elie Cartan, *Sur des familles remarquables d'hypersurfaces isoparamétriques dans les espaces sphériques*, Math. Z. **45** (1939), 335–367.

- [4] ———, *Sur quelques familles remarquables d'hypersurfaces*, C.R. Congres Math. Liège (1939), 30–41.
- [5] ———, *Sur des familles d'hypersurfaces isoparamétriques des espaces sphériques à 5 et à 9 dimensions*, Univ. Nac. Tucumán. Revista A. **1** (1940), 5–22.
- [6] Miguel Domínguez-Vázquez, *An introduction to isoparametric foliations*, 2018.
- [7] Josef Dorfmeister and Erhard Neher, *Isoparametric hypersurfaces, case $g = 6$, $m = 1$* , Comm. Algebra **13** (1985), no. 11, 2299–2368.
- [8] Gregory Drugan, Hojoo Lee, and Xuan Hien Nguyen, *A survey of closed self-shrinkers with symmetry*, Results Math. **73** (2018), no. 1, Paper No. 32, 32.
- [9] Gregory Drugan and Xuan Hien Nguyen, *Shrinking doughnuts via variational methods*, J. Geom. Anal. **28** (2018), no. 4, 3725–3746.
- [10] Dirk Ferus, Hermann Karcher, and Hans Friedrich Münzner, *Cliffordalgebren und neue isoparametrische Hyperflächen*, Math. Z. **177** (1981), no. 4, 479–502.
- [11] Gerhard Huisken, *Asymptotic behavior for singularities of the mean curvature flow*, J. Differential Geom. **31** (1990), no. 1, 285–299.
- [12] Tom Ilmanen, *Singularities of mean curvature flow of surfaces*, (1995).
- [13] Hans Friedrich Münzner, *Isoparametrische Hyperflächen in Sphären*, Math. Ann. **251** (1980), no. 1, 57–71.
- [14] ———, *Isoparametrische Hyperflächen in Sphären. II. Über die Zerlegung der Sphäre in Ballbündel*, Math. Ann. **256** (1981), no. 2, 215–232.
- [15] Richard S. Palais and Chuu-Lian Terng, *Reduction of variables for minimal submanifolds*, Proc. Amer. Math. Soc. **98** (1986), no. 3, 480–484.
- [16] Gudlaugur Thorbergsson, *Singular Riemannian foliations and isoparametric submanifolds*, Milan J. Math. **78** (2010), no. 1, 355–370.
- [17] Qi Ming Wang, *On a class of minimal hypersurfaces in \mathbb{R}^n* , Math. Ann. **298** (1994), no. 2, 207–251.
- [18] Brian White, *Partial regularity of mean-convex hypersurfaces flowing by mean curvature*, Internat. Math. Res. Notices (1994), no. 4, 186 ff., approx. 8 pp.