

# Compositional Sparsity, Approximation Classes, and Parametric Transport Equations\*

*Dedicated to Ronald DeVore on the occasion of his 80th birthday*

Wolfgang Dahmen<sup>†</sup>

July 14, 2022

## Abstract

Approximating functions of a large number of variables poses particular challenges often subsumed under the term “Curse of Dimensionality”. Unless the approximated function exhibits a very high level of smoothness the Curse can be avoided only by exploiting some typically hidden *structural sparsity*. In this paper we propose a general framework for model classes of functions in high dimensions based on suitable notions of *compositional sparsity* quantifying approximability by highly nonlinear expressions such as deep neural networks. The relevance of these concepts are demonstrated for *solution manifolds* of parametric transport equations which are known not to enjoy the type of high order regularity of parameter-to-solution maps that help to avoid the Curse of Dimensionality in other model scenarios. Compositional sparsity is shown to serve as the key mechanism for proving that sparsity of problem data is inherited in a quantifiable way by the solution manifold. In particular, one obtains convergence rates for deep neural network realizations showing that the Curse of Dimensionality is indeed avoided.

**AMS Subject Classification:** 41A25, 35A35, 41A63, 35B30, 35L04, 41A46

**Keywords:** Tamed compositions, compositional approximation, approximation classes, deep neural networks, parametric transport equations, solution manifolds, nonlinear widths.

## 1 What this is about

**Background:** Like hardly any other topic have Deep Neural Networks (DNNs) been recently influencing numerous vibrant research activities constantly broadening the scope of

---

\*This work has been supported in part by National Science Foundation grant NSF-DMS-2012469, and the SFB 1481, funded by the German Research Foundation

<sup>†</sup>Mathematics Department, University of South Carolina, Columbia, SC, USA

applications of machine learning. In a nutshell, DNNs are compositions of simple finitely parametrized mappings, each one being in turn a composition of an affine map and a componentwise acting nonlinear function. Thus, DNNs are highly nonlinear mappings differing in many respects from classical approximation systems, see [5, 6] for DNNs in the general context of nonlinear approximation.

The empirically observed stunning success of deep networks in unsupervised and supervised learning, covering, in particular, “Big Data” application scenarios, has raised high expectations of a similar impact in other, more science related areas. Again, numerous studies appear to support such hopes there as well. Particular attention has been paid to understanding the *expressive power* of neural networks (see e.g. [5, 13, 10, 27] as well as the excellent exposition in [6]), especially the approximability of functions of a large number of variables which usually suffers from the *Curse of Dimensionality*. This latter aspect is the focus of this work as well. The term “Curse” expresses roughly that the cost of recovering such a function within some target accuracy increases exponentially in the spatial dimension - viz. the number variables. Such *high-dimensional* problems arise, for instance, in Uncertainty Quantification, state estimation, data assimilation and alike.

A first important approach to understanding the *expressive power* of DNNs is “emulation”, see e.g. [17, 27, 25, 6, 24, 13, 10]. In this program it is shown that the approximation power of nearly all known “classical” methods are (perhaps up to some log-factors) matched by deep neural networks (DNNs), see e.g. [22]. The prospect of having a single approximation format that can address essentially any approximation problem with near-optimal expressive power is indeed very tempting. There are perhaps two points suggesting that things are after all not that simple. First, classical methods come typically with well-understood algorithmic realizations while training DNNs is haunted by a significant remaining uncertainty of optimization success. In fact, it is shown in [14] that theoretically possible high approximation orders cannot be realized based on point samples as training input. Thus, celebrated superconvergence results may lose their punch. Second, regarding high-dimensionality, by its very nature emulation can show that DNNs can avoid the *Curse of Dimensionality* if some existing classical method can do so as well. Less seems to be known instead about the question:

*can DNNs help avoiding the Curse of Dimensionality even in scenarios where state of the art methods don't?*

In this regard, interesting rigorous results have indeed been obtained for DNN approximations of solutions of certain *spatially high-dimensional* partial differential equations (PDEs) like Kolmogorov's or Hamilton-Jacobi-Bellman equations where the use of DNNs is shown to break the Curse of Dimensionality, see e.g. in [1, 12, 11]. In this context for instance *Barron classes* have been attracting significant interest where sparsity is tantamount to “dimension-dependent” smoothness rates, see e.g. [8]. Thus, the model classes get slimmer with increasing dimension. The type of results in [11] is somewhat different in that it is roughly shown that sparse problem data give rise to sparse approximate representations of solutions avoiding the Curse of Dimensionality. This is also the general theme in the present

paper.

Comparable results seem to be lacking for a second important class of high-dimensional problems arising, for instance, in Uncertainty Quantifications. Here the objects of interest are so called *solution manifolds* of *parameter dependent* families of PDEs, written in abstract residual form as

$$\mathcal{D}(u, y; f) = 0, \quad y \in \mathcal{Y}, \quad (1.1)$$

where  $\mathcal{D}$  stands for a differential operator involving coefficients depending on parameters  $y \in \mathcal{Y}$ . The parameter domain  $\mathcal{Y}$  is typically a compact subset of  $\mathbb{R}^{d_y}$ , e.g. a hypercube, where  $d_y$  could be large. Accordingly, the parameter-dependent solutions  $u = u(y)$ ,  $y \in \mathcal{Y}$ , of (1.1) become functions of  $m$  spatial and  $d_y$  parametric variables. The collection of viable states covered by the PDE model obtained when  $y$  traverses  $\mathcal{Y}$ , forms the aforementioned solution manifold

$$\mathcal{M} = \{u(y) : \mathcal{D}(u(y), y; f) = 0, \quad y \in \mathcal{Y}\}. \quad (1.2)$$

An efficient numerical exploration of  $\mathcal{M}$  is in one way or the other connected with efficiently approximating the *parameter-to-solution map*  $y \mapsto u(y)$ , preferably without suffering from the Curse of Dimensionality. This depends strongly on the type of the PDE family. For elliptic problems (and their close relatives), under reasonable conditions on parameter representations,  $y \mapsto u(y)$  is then known to be holomorphic. Reduced Bases concepts as well as sparse polynomial approximations are in this case very successful because associated Kolmogorov widths

$$d_n(\mathcal{M})_{\mathbb{U}} := \inf_{\dim \mathbb{W} = n} \sup_{u \in \mathcal{M}} \inf_{w \in \mathbb{W}} \|u - w\|_{\mathbb{U}}$$

decay rather robustly with respect to  $d_y$  where  $\mathbb{U}$  is the space accommodating the solutions of (1.1). So here sparsity means effective approximability from linear spaces of moderate dimension. Via emulation DNNs can also be shown to avoid the Curse of Dimensionality although their preference is perhaps hard to argue, see e.g. [17, 22].

For other PDE models, in particular those dominated by transport rather than diffusion, the situation is grossly different. Pure transport equations could be viewed as extreme examples where parameter-to-solution maps are no longer holomorphic and since one cannot exploit dissipative effects different sparsity notions are needed. In this regard, the approximation of solutions to parameter dependent transport equations by deep neural networks (DNNs) has been investigated in [23]. It is important to note though that the convergence rates established there still reflect the full Curse of Dimensionality. Roughly, achieving accuracy  $\varepsilon$  requires a network complexity of the order  $\varepsilon^{-\frac{m+1+d_y}{\alpha}}$  where  $m+1, d_y$  denote the number of space-time and parametric variables and  $\alpha$  represents the smoothness the solutions resulting from suitable smoothness assumptions on the problem data like initial conditions, right hand sides, convection coefficients. So, the use of DNNs per se is certainly not a guarantee to avoid the Curse. Instead, one can only expect results that quantify the extent to what *structural sparsity of problem data* is inherited by solutions, see [3, 11] for such results on high-dimensional PDEs.

**Objectives and Layout:** This paper addresses two closely intertwined but different foci.

First, the expressive power of DNNs needs to refer to a specific *model class*  $\mathcal{K}$  of target functions. Since according to [2], stable approximation schemes that avoid the Curse of Dimensionality are only possible if the *entropy numbers* of the underlying model class do not exhibit an exponential dependence on the number of variables, balls in classical smoothness spaces are ruled out. The identification of relevant model classes is therefore an issue by itself and typically problem dependent. Solution manifolds (1.2) are certainly relevant but rather than focusing solely on  $\mathcal{M}$  for each specific PDE type we propose and analyze in § 2a more general type of model classes, based on what we refer to as *compositional dimension-sparsity*. This builds to some extent on ideas and concepts discussed in [25] in the context of nonlinear regression. The relevance of compositions for neural approximation is by no means new, see e.g. [8, 20, 21, 25]. However, although similar in spirit to [25], the angle taken here is still different and studies first compositional approximation on a continuous functional level, somewhat in the spirit of [3] for tensors. This gives then rise to model classes whose members are not confined to specific compositional formats but allows one to quantify *compositional approximability*. The proposed framework is neither tied to the discretization of dynamical systems nor to classical smoothness conditions but applies whenever the objects of interest can be quantifiably approximated by iterative processes (here primarily in spaces of continuous functions but with conceivable extensions to more general scenarios).

As a second focus, we study parametric transport equations and corresponding solution manifolds in the context of these compositional model classes. This class of PDEs is, among many other things, of interest because, as mentioned earlier, the classical model reduction concepts fail. In § 3 we describe the parametric model problem and collect a few elementary prerequisites that will be frequently used in what follows.

With these two preparatory sections in place we formulate in § 4 the main results of this paper, concerning the DNN approximability of solution manifolds. In particular, we show that the parameter-to-solution map admits DNN approximations that do not suffer from the Curse of Dimensionality. More precisely, sparse approximability of problem data implies sparse approximability of solutions. For instance, when the parameter dependence of the convection field is affine, the characteristics, as functions of time, space, and parametric variables, can be approximated by DNNs as follows

$$\|z - \mathcal{N}_\varepsilon\|_{L_\infty([0,T] \times D \times \mathcal{Y}; \mathbb{R}^m)} \leq \varepsilon, \quad \#\mathcal{N}_\varepsilon \lesssim e^{LT(m+1)} \left( \frac{Td_y}{\varepsilon} \right)^{m+1} \left| \log_2 \frac{Td_y}{\varepsilon} \right|^2, \quad (1.3)$$

where  $m, d_y$  denote again the spatial, parametric dimension, respectively. Our results yield, in particular, upper bounds for *nonlinear manifold widths* of such solution manifolds. All proofs are given in §6. They are at times technical and tedious which is the price for keeping track of constants in the estimates. This in turn is relevant for the discussion of sharpness of convergence rates, see Theorem 4.5 improving on (1.3).

While linear transport offers a convenient case study for compositional sparsity models, we finally indicate in §5 possible extensions to a wider scope of PDE models.

**Notational Conventions:** In what follows we often write  $a \lesssim b$  to indicate that  $a$  is bounded by a constant multiple of  $b$  where the constant is independent of any parameters  $a$  and  $b$  may depend on, unless specified otherwise. Accordingly  $a \sim b$  means  $a \lesssim b$  and  $b \lesssim a$ .

For notational brevity and convenience we will use, for any pair of finite dimensional metric spaces  $X, Y$  and any continuous function  $g : X \subset \mathbb{R}^{d_0} \rightarrow Y \subset \mathbb{R}^{d_1}$ , the shorthand notation

$$\|g\|_\infty = \|g\|_{L_\infty(X;Y)} = \sup_{x \in X} \sup_{i=1, \dots, d_1} |g_i(x)| = \sup_{x \in X} |g(x)|_\infty,$$

unless we wish to emphasize particular domains or there is a risk of confusion. Likewise we use the domain- and dimension unspecific notation  $\|g\|_{\text{Lip}}, |\cdot|_{\text{Lip}}$  to denote the full Lipschitz norm, respectively semi-norm

$$|g|_{\text{Lip}} := \sup_{x, z \in X} \left| \frac{g(x) - g(z)}{|x - z|_\infty} \right|_\infty, \quad \|g\|_{\text{Lip}} := \max\{\|g\|_\infty, |g|_{\text{Lip}}\}.$$

## 2 Compositions

### 2.1 Rationale

Trying to understand the efficacy of DNNs through compositions is most natural and the common view point of several earlier works, see e.g., [8, 21, 20, 25]. From a rather different angle the aim here is to put forward a general framework that allows one to quantify *compositional approximability*, still on a continuous infinite-dimensional functional level. Here approximants are not yet determined by a finite number of degrees of freedom but prepare the ground for constructing finitely parametrized approximations in a second step.

First, in § 2.2, we fix some notation and collect a few useful structural properties of compositions revolving on the notion of *dimension-sparsity*, inspired to some extent by [25]. In § 2.3 we introduce metric properties of (finite) compositions, that allow us to introduce in § 2.4 *compositional approximation classes* that are to serve later as model classes for characterizing approximability by DNNs.

### 2.2 Basic and Notation and Structural Properties

#### 2.2.1 Compositional Representations

We will interpret “compositions” - denoted in what follows (at times in slight abuse of notation) by the symbol “ $\circ$ ” - in a broad sense, covering iterated applications of global operators. A prominent albeit not exclusive role is, however, played by compositions in a *pointwise sense*, defined for continuous functions. Specifically, we consider compositions of mappings

$$g^1 : D_0 \subset \mathbb{R}^{d_0} \rightarrow \mathbb{R}^{d_1}, \quad g^\ell : \mathbb{R}^{d_{\ell-1}} \rightarrow \mathbb{R}^{d_\ell}, \quad \ell = 2, \dots, n,$$

where we always require that the last factor is linear, i.e., for some  $\alpha_i \in \mathbb{R}$ ,  $i = 1, \dots, d_{n-1}$

$$g^n = \sum_{i=1}^{d_{n-1}} \alpha_i g_i^{n-1}.$$

It will be convenient to abbreviate the ordered array of mappings by  $\mathbf{g} = (g^j)_{j=1}^n$  to denote the *pointwise composition*

$$G(z) = (g^n \circ \dots \circ g^1)(z), \quad \text{where} \quad (g \circ h)(z) := g(h(z)) =: G_{\mathbf{g}}(z), \quad (2.1)$$

as a mapping from  $D_0 \subset \mathbb{R}^{d_0} \rightarrow \mathbb{R}^{d_n}$ . When sometimes writing  $h \circ g$ , say, it will always be assumed that the two factors  $g, h$  are “dimensionally compatible” in the above sense.

We refer to  $G_{\mathbf{g}}$  or simply to  $\mathbf{g}$  in (2.1) as a *compositional representation* of  $G$ .

prominent examples are DNNs whose formal definition can be found in numerous texts, see e.g. [6, 15, 11, 22, 27]. Here we are content with mentioning that DNN realizations (denoted by  $\mathcal{N}$ ) are of the form (2.1) with factors  $g^j(\cdot) = \sigma(A^j \cdot + b^j)$  where  $A^j \in \mathbb{R}^{d_j \times d_{j-1}}$ ,  $b^j \in \mathbb{R}^{d_j}$ , and the *activation function* or *rectifier*  $\sigma$  acts componentwise. An important examples is the ReLU rectifier  $\sigma(t) = \max\{0, t\} =: t_+$ . The entries in  $A^j, b^j$  or in the linear output layer are called *weights* and their number  $\#\mathcal{N}$  is the *size* or *complexity* of the DNN  $\mathcal{N}$ .

## 2.2.2 Lipschitz-Stability

When dealing with pointwise compositions, the first important constraint on (2.1) in what follows is that we require all factors  $g^j$  (and hence  $G$  itself) in  $\mathbf{g}$  to be Lipschitz continuous with Lipschitz constants  $L_j$ . The reason lies in the following simple folklore perturbation bound. Consider dimensionally compatible Lipschitz functions  $g, h$  with constants  $L_g, L_h$  and  $\varepsilon_g, \varepsilon_h$  accurate approximations  $\tilde{g}, \tilde{h}$ . Then

$$\|g \circ h - \tilde{g} \circ \tilde{h}\|_{\mathbb{X}} \leq \|g \circ h - g \circ \tilde{h}\|_{\mathbb{X}} + \|g \circ \tilde{h} - \tilde{g} \circ \tilde{h}\|_{\mathbb{X}} \leq L_g \varepsilon_h + \varepsilon_g. \quad (2.2)$$

Given a compositional representation  $G_{\mathbf{g}}$  of the form (2.1), we denote for  $k \leq n$  by  $L_{[n,k]} = L_{[n,k]}(\mathbf{g})$  the Lipschitz constant of partial compositions  $g^n \circ g^{n-1} \circ \dots \circ g^k$ . It will be convenient to set  $L_{[n,n+1]} = 1$ . Then, using the above argument inductively yields the following convenient folklore bounds to be recorded for later frequent use, see e.g. [25].

**Remark 2.1** *Assume that we have mappings  $\tilde{g}^j : \mathbb{R}^{d_{j-1}} \rightarrow \mathbb{R}^{d_j}$ ,  $j = 1, \dots, n$ , such that*

$$\|g^j - \tilde{g}^j\|_{\mathbb{X}} \leq \varepsilon_j, \quad j = 1, \dots, n.$$

*Then*

$$\|g^n \circ \dots \circ g^1 - \tilde{g}^n \circ \dots \circ \tilde{g}^1\|_{\mathbb{X}} \leq \varepsilon_n + \sum_{j=1}^{n-1} \varepsilon_j L_{[n,j+1]} = \sum_{j=1}^n \varepsilon_j L_{[n,j+1]}.$$

*where in the last term we have used the convention in the definition of  $L_{[n,k]}$ . By a symmetric argument we can replace  $L_{[n,j+1]}(\mathbf{g})$  by  $L_{[n,j+1]}(\tilde{\mathbf{g}})$ . ns.*

*In terms of the individual Lipschitz constants, one has, of course,  $L_{[n,k]} \leq \prod_{j=k}^n L_j$ .*

*Finally, the estimates remain valid for more general iterated applications of operators as long as an estimate like (2.2) holds.*

An important context where this will be used later is the following result.

**Proposition 2.2** *For any  $\delta > 0$ , and any  $g \in \text{Lip}_1((0, 1)^s)$ , there exist a ReLU network  $\mathcal{N}_\delta$  such that*

$$\|g - \mathcal{N}_\delta\|_\infty \leq \delta, \quad \|\mathcal{N}_\delta\|_{\text{Lip}_1} \leq c_3(1 + \|g\|_\infty)\|g\|_{\text{Lip}_1}, \quad (2.3)$$

and

$$\#\mathcal{N}_\delta \leq c_1 \|g\|_{\text{Lip}_1}^s \delta^{-s} \log_2 \frac{1}{\delta}, \quad \text{depth of } \mathcal{N}_\delta \leq c_2 \log_2 \frac{1}{\delta}, \quad (2.4)$$

where the constants  $c_1, c_2, c_3$  depend only on  $s$ . Using if necessary Lipschitz-stable continuation from bounded domains to hypercubes, analogous results hold for more general domains under mild geometric constraints (see e.g. [15]).

There are numerous results of this flavor, perhaps up to the  $\text{Lip}_1$ -stability claim in (2.3). In this latter regard, [15] establishes the existence of ReLU networks that approximate functions of higher Sobolev regularity in weaker Sobolev norms without compromising the standard complexity bounds. This would imply (2.3) if we impose more regularity on  $g$  than just to belong to  $\text{Lip}_1$  (or  $W^1(L_\infty((0, 1)^s))$ ). In the present context we prefer to avoid assuming such “excess regularity” and sketch corresponding arguments in § 6.1.1 for completeness. However, we do build on some of the concepts developed in [15].

### 2.2.3 Compositional Complexity

Obviously, while each  $\mathbf{g}$  determines  $G = G_{\mathbf{g}}$ , a mapping  $G$  may have infinitely many compositional representations. In fact, we can always “reshape” compositional factors by writing

$$f \circ g = (f \circ h) \circ (h^{-1} \circ g) =: \tilde{f} \circ \tilde{g}, \quad (2.5)$$

where the factors in the new representation could have unfavorable regularity or stability properties, or in fact vice versa. In slight abuse of terminology we sometimes write  $\mathbf{g}_G$  to express that  $\mathbf{g}$  is a representation of the mapping  $G$ .

Specifically, we are interested in *large* input-dimension  $d_0$  and ask whether specific compositional structures  $\mathbf{g}_G$  may help avoiding the Curse of Dimensionality. In this regard, the role of sparsely connected neural networks has been already observed in [25] in the context of nonlinear regression and has been a source of motivation for what follows.

To that end, let for a scalar function  $g : \mathbb{R}^{d'} \rightarrow \mathbb{R}$  denote by  $s(g) \leq d'$  the number of variables,  $g$  depends on. Then, for each interior factor  $g^j$ ,  $j = 1, \dots, n-1$ , in a representation  $\mathbf{g}$  define the integer  $s(g^j) \in \mathbb{N}_0$  by

$$s(g^j) := \begin{cases} 0, & \text{if } g^j \text{ is the identity mapping,} \\ 1, & \text{when } g^j \text{ is at most affine or bilinear,} \\ \max_{i=1}^{d_j} s(g_i^j), & \text{else.} \end{cases} \quad (2.6)$$

**Definition 2.3** For a representation  $\mathbf{g}$  of a mapping  $G$  let

$$s(\mathbf{g}) := \max_{j=1, \dots, n(\mathbf{g})-1} s(g^j).$$

A representation  $\mathbf{g}_G$  of the form (2.1) is called  $s$ -dimension sparse if  $s(\mathbf{g}) \leq s$ .

We sometimes encode the compositional structure of  $\mathbf{g}$  by its ‘dimensionality vector’

$$\mathbf{D} = \mathbf{D}(\mathbf{g}) = (d_0, (d_1, \mathbf{s}^1), \dots, (d_{n-1}, \mathbf{s}^{n-1}), d_n), \quad \mathbf{s}^j = (s(g_i^j))_{i=1}^{d_j}, \quad j = 1, \dots, n-1. \quad (2.7)$$

Moreover, we use this notion for *any* array  $\mathbf{D}$  which is dimensionally compatible in the sense that  $0 \leq |\mathbf{s}^j|_\infty \leq d_{j-1}$ .

We introduce next a measure for the *compositional complexity* of a representation  $\mathbf{g}$

$$\mathfrak{N}(\mathbf{g}) := \sum_{j=1}^n \sum_{i=1}^{d_j} s(g_i^j) \leq \sum_{j=1}^n d_{j-1} d_j.$$

When writing  $\mathfrak{N}(\mathbf{D})$  without reference to a specific  $\mathbf{g}$  this is understood by default as  $\mathfrak{N}(\mathbf{D}) := \sum_{j=1}^n \sum_{i=1}^{d_{j-1}} \mathbf{s}_i^j d_j$ .

**Remark 2.4** The rationale behind (2.6) is that in the end, we wish to approximate a mapping  $G$  by a finitely parametrized mapping, taking a certain ‘functional’ compositional structure  $\mathbf{g}_G$  as point of departure. Factors  $g^j$  which are already given as a linear or bilinear, hence are directly computable mappings, do not need to be approximated e.g. by a DNN (perhaps in the bilinear case only at a logarithmic cost). Therefore, such a factor should contribute only weakly to the ‘complexity’ of  $\mathbf{g}$  encoded by  $\mathfrak{N}(\mathbf{g})$ .

We say two representations  $\mathbf{g}, \mathbf{g}'$  are dimensionally compatible (in this order) if the output dimension  $d_{n(\mathbf{g})}$  agrees with the input-dimension  $d_0(\mathbf{g}')$ . One easily checks the following useful facts.

**Remark 2.5** For any two dimensionally compatible representations  $\mathbf{g}, \mathbf{g}'$ , the composition  $\hat{G} := G_{\mathbf{g}'} \circ G_{\mathbf{g}}$  has a representation  $\hat{\mathbf{g}} = (\mathbf{g}' | \mathbf{g})$  satisfying

$$\mathfrak{N}(\hat{\mathbf{g}}) = \mathfrak{N}(\mathbf{g}) + \mathfrak{N}(\mathbf{g}'). \quad (2.8)$$

Likewise, when  $\mathbf{g}, \mathbf{g}'$  have equal in- and output dimension, i.e.,  $d_0(\mathbf{g}) = d_0(\mathbf{g}')$  and  $d_{n(\mathbf{g})} = d_{n(\mathbf{g}')}$ , the sum  $\hat{G} := G_{\mathbf{g}} + G_{\mathbf{g}'}$  has a representation  $\hat{\mathbf{g}} = \begin{pmatrix} \mathbf{g} \\ \mathbf{g}' \end{pmatrix}$  satisfying

$$\mathfrak{N}(\hat{\mathbf{g}}) = \mathfrak{N}(\mathbf{g}) + \mathfrak{N}(\mathbf{g}'). \quad (2.9)$$

This follows easily by (what in the DNN context is called) parallelization (as hinted at by the notation  $\begin{pmatrix} \mathbf{g} \\ \mathbf{g}' \end{pmatrix}$ ), upon possibly inserting identity factors in the representation of smaller depth.



Given any integer  $s$ ,

$$\mathfrak{C}_{N,s}(D, d') := \left\{ G_{\mathbf{g}} : D \rightarrow \mathbb{R}^{d'} : \mathfrak{N}(\mathbf{g}) \leq N, s(\mathbf{g}) \leq s, g^j \in \text{Lip}_1, 1 \leq j \leq n(\mathbf{g}) \right\}$$

denotes the collection of mappings with  $s$ -dimension-sparse representations of complexity at most  $N$ . When  $D, d'$  is clear from the context we write  $\mathfrak{C}_{N,s}$ .

$\mathfrak{C}_{N,s}$  is of course not a linear set but, by Remark 2.5

$$\mathfrak{C}_{N,s} + \mathfrak{C}_{N,s} \subset \mathfrak{C}_{2N}.$$

Likewise for mappings  $G \in \mathfrak{C}_{N,s}(D, d')$ ,  $\tilde{G} \in \mathfrak{C}_{N',s}(\mathbb{R}^{d'}, d'')$  we infer from (2.8) that

$$\tilde{G} \circ G \in \mathfrak{C}_{N+N',s}, \quad (2.10)$$

We'll use at times the notation  $\mathfrak{C}(\mathbf{D}) := \{G_{\mathbf{g}} : \mathbf{D}(\mathbf{g}) = \mathbf{D}\}$  to collect all mappings with representations that have the dimensionality vector  $\mathbf{D}$  and use that

$$\mathfrak{C}_{N,s} = \bigcup \left\{ \mathfrak{C}(\mathbf{D}) : \mathbf{s}_i^j(\mathbf{D}) \leq s, 1 \leq j \leq n(\mathbf{D}), \mathfrak{N}(\mathbf{D}) \leq N \right\}. \quad (2.11)$$

#### 2.2.4 Implanting finitely-parametrized components:

The following comments, taking up on Remark 2.1, show how compositions on a continuous functional level can be used to construct finitely parametrized approximations, see also [25].

First, if the components of the factors  $\tilde{g}^j$  in a representation  $\tilde{\mathbf{g}}$  of  $\tilde{G}$  are neural networks  $\tilde{g}_i^j = \mathcal{N}_i^j$ ,  $i = 1, \dots, d_j$ , and  $\#\mathcal{N}_i^j$  denotes the number of weights in  $\mathcal{N}_i^j$ , one readily sees that

$$\#\mathcal{N}_i^j \approx \mathfrak{N}(\mathcal{N}_i^j), \quad (2.12)$$

uniformly in  $i, j$ . We form approximations to a mapping  $G$  with representation  $\mathbf{g}$  as in (2.1), by approximating each component  $g_i^j$ ,  $i = 1, \dots, d_j$ ,  $j = 1, \dots, n(\mathbf{g}) - 1$ , by a neural network of corresponding in- and output-dimension. We refer to such an operation as ‘‘implanting’’ neural networks. As a consequence,  $\tilde{G}$  has a representation in terms of neural networks.

A simple and frequently used consequence of Remark 2.1 and (2.8) can then be stated as follows.

**Lemma 2.1** *Assume that for some  $s \leq d_0$ , the mapping  $G$  belongs to  $\mathfrak{C}_{N,s}$ , i.e., is  $s$ -dimension-sparse (see Definition 2.3). Then, forming a DNN  $\mathcal{N}$  by replacing each (non-trivial) component  $g_i^j$  by a  $\delta_j$  accurate Lipschitz stable network  $\mathcal{N}_i^j$ , i.e.,*

$$\|g_i^j - \mathcal{N}_i^j\|_{\mathbb{X}_i} \leq \delta_j, \quad j = 1, \dots, n(\mathbf{g}) - 1,$$

one has

$$\|G - \mathcal{N}\|_{L_\infty} \leq \delta_n + \sum_{j=1}^{n(\mathbf{D})-1} \delta_j L_{[n(\mathbf{D}), j+1]}, \quad \#\mathcal{N} \leq \sum_{j=1}^{n(\mathbf{D})-1} \|g^j\|_{\text{Lip}}^s d_j \delta_j^{-s} |\log_2 \delta_j|. \quad (2.13)$$

In absence of any knowledge about  $\mathbf{D}(\mathbf{g})$ , one has

$$\#\mathcal{N} \leq \|G\|_{N, \text{Lip}}^s \mathfrak{N}(G) \max_{j=1, \dots, n(\mathbf{D})} \delta_j^{-s} |\log_2 \delta_j|. \quad (2.14)$$

**Proof:** The first relation in (2.13) follows from Remark 2.1 while the second one is a consequence of Proposition 2.2. Since  $\sum_{j=1}^{n(\mathbf{g})} d_j \leq \mathfrak{N}(\mathbf{g})$  (2.14) follows.  $\square$

### 2.2.5 Comments

The following remarks motivate the discussion in the next section.

**Remark 2.6** *When  $s \geq d_0$ ,  $D \subset \mathbb{R}^{d_0}$ , the constraint of s-dimension sparsity is, of course void. In this case one simply has that  $\mathfrak{C}_{N,s}(D, d') = \text{Lip}_1(D; \mathbb{R}^{d'})$ . In fact, any  $G \in \text{Lip}_1(D, d')$  has a trivial representation*

$$G(x) = (g^2 \circ g^1)(x), \quad g_i^1 = G_i, \quad i = 1, \dots, d' \quad g^2 = \text{id}_{d'}. \quad (2.15)$$

*So, s-dimension-sparsity with  $s < d = d_0$  is essential for such a framework to offer interesting information.*

On the other hand, dimension-sparsity per se is also not yet a sufficient instrument. The famous Arnold-Kolmogorov Superposition Theorem, in a variant established by G.G. Lorentz ([19]), states that every continuous function  $G$  on  $D_0 = [0, 1]^d$  has a representation

$$G(x_1, \dots, x_d) = \sum_{q=0}^{2d} \Phi \left( \sum_{p=1}^d \phi_{q,p}(x_p) \right), \quad (2.16)$$

where  $\Phi, \phi_{q,p}$  are continuous functions. Thus,  $G = g^4 \circ g^3 \circ g^2 \circ g^1$ , where  $g^1(x) = (\phi_{q,p}(x_p))_{q,p=0,1}^{2d,d}$ , i.e.,  $d_1 = (2d + 1)d$ ,  $g^2(z) = \left( \sum_{p=1}^d z_{q,p} \right)_{q=0}^{2d}$ , i.e.,  $d_2 = 2d + 1$ ,  $g^3(z) = (\Phi(z_q))_{q=0}^{2d}$ , i.e.,  $d_3 = 2d + 1$ ,  $g^4(z) = \sum_{q=0}^{2d} z_q$ . Hence, every  $G \in \text{Lip}_1([0, 1]^d)$  belongs to  $\mathfrak{C}_{N_d,1}([0, 1]^d, 1)$  where  $N_d = d(2d + 1) + (2d + 1) + (2d + 1) + 1$ . We refer to [9, 16] for somewhat controversial views on the relevance of the Arnold-Kolmogorov Superposition Theorem for computational aspects, in particular, in neural computation.

Thus, mere s-dimension sparsity just by itself is still not very informative. However, it is well known that the *regularity* of component functions  $\phi_{q,p}$  in (2.16) *cannot* be controlled. Hence, one needs to “*tame*” s-dimension-sparse representations to arrive at a meaningful concept.

## 2.3 Tamed Compositions and Compactness

In view of the preceding comments, we need to *regularize* representations of elements in  $\mathfrak{C}_{N,s}$ . We discuss next one (among many possible) such version(s). Let for any  $G \in \mathfrak{C}_{N,s}$

$$\|G\|_{N,s} := \inf_{\mathbf{g}} \left\{ \max_{\ell \leq n(\mathbf{g})} \{ \|g^\ell\|_{\text{Lip}_1} \} : G_{\mathbf{g}} = G, s(\mathbf{g}) \leq s, \mathfrak{N}(\mathbf{g}) \leq N \right\}. \quad (2.17)$$

This does not quite induce a distance yet but is close to one. In fact, one has

$$\|G + \tilde{G}\|_{2N,s} \leq \|G\|_{N,s} + \|\tilde{G}\|_{N,s}. \quad (2.18)$$

To see this, let  $\mathbf{g}, \tilde{\mathbf{g}}$  be near-infimizing representations of  $G \in \mathfrak{C}_{N,s}$ ,  $\tilde{G} \in \mathfrak{C}_{\tilde{N},s}$ , respectively. Then, by (2.9) there exists a representation  $\hat{\mathbf{g}}$  of  $G + \tilde{G}$  with

$$\mathfrak{N}(\hat{\mathbf{g}}) \leq \mathfrak{N}(\tilde{\mathbf{g}}) + \mathfrak{N}(\mathbf{g}) \leq N + \tilde{N}.$$

Clearly  $s(\hat{\mathbf{g}}) \leq \max\{s(\mathbf{g}), s(\tilde{\mathbf{g}})\}$  and  $\|\hat{g}^j\|_{\text{Lip}_1} \leq \max\{\|g^j\|_{\text{Lip}_1}, \|\tilde{g}^j\|_{\text{Lip}_1}\}$ . Thus

$$\|G + \tilde{G}\|_{2N,s} \leq \max_{\ell \leq n(\hat{\mathbf{g}})} \{\|\tilde{g}^\ell\|_{\text{Lip}_1}, \|g^\ell\|_{\text{Lip}_1}\} \leq \max\{\|G\|_{N,s}, \|\tilde{G}\|_{N,s}\}. \quad (2.19)$$

With some abuse of terminology we refer to  $\|\cdot\|_{N,s}$  as “composition-norm”.

Note that a uniform control of the Lipschitz norms of the factors does not imply a uniform control of  $\|G\|_{\text{Lip}_1}$  when  $N$  increases. Therefore, we will also make use of a stronger regularization:

$$\|G\|_{N,s,\text{Lip}} := \inf_{\substack{\mathbf{g}: G_{\mathbf{g}}=G \\ s(\mathbf{g}) \leq s, \mathfrak{N}(\mathbf{g}) \leq N}} \left\{ \max \left\{ \|g^\ell\|_{\text{Lip}_1}, L_{[n(\mathbf{g}), \ell+1]}(\mathbf{g}) \right\} : \ell = 1, \dots, n(\mathbf{g}) \right\}, \quad (2.20)$$

where  $L_{[n(\mathbf{g}), \ell+1]}(\mathbf{g})$  denotes again the Lipschitz constant of the *partial compositions*  $g^{n(\mathbf{g})} \circ \dots \circ g^{\ell+1}$ . Controlling  $\|\cdot\|_{N,s,\text{Lip}}$ , obviously constrains representations of elements further implying

$$\|G\|_{\text{Lip}_1} \leq \|G\|_{N,s,\text{Lip}}, \quad G \in \mathfrak{C}_{N,s}. \quad (2.21)$$

Since in parallization  $\begin{pmatrix} g^j \\ \tilde{g}^j \end{pmatrix} \begin{pmatrix} z \\ \tilde{z} \end{pmatrix} = \begin{pmatrix} g^j(z) \\ \tilde{g}^j(\tilde{z}) \end{pmatrix}$ , the same reasoning as above shows that (2.19) and hence (2.18) persists to hold for this notion as well.

Similarly, for dimensionally compatible mappings  $G_i \in \mathfrak{C}_{N_i,s}$ ,  $i = 1, 2$ , we know already that  $G := G_2 \circ G_1 \in \mathfrak{C}_{N_1+N_2,s}$ , see (2.10). So, in addition one has

$$\|G\|_{N_1+N_2,s} \leq \max \left\{ \|G_1\|_{N_1,s}, \|G_2\|_{N_2,s} \right\}, \quad (2.22)$$

and for the stronger regularization

$$\|G\|_{N_1+N_2,s,\text{Lip}} \leq \max \left\{ \|G_1\|_{N_1,s}, \|G_2\|_{N_2,s}, \|G_1\|_{N_1,s} \cdot \|G_2\|_{N_2,s} \right\} \quad (2.23)$$

Our next observation concerns compactness of bounded subsets of  $\mathfrak{C}_{N,s}$ .

**Proposition 2.7** *For any fixed constant  $B < \infty$  and any  $s \in \mathbb{N}$  the set*

$$\mathfrak{C}_{N,s}(B) := \{G \in \mathfrak{C}_{N,s} : \|G\|_{N,s} \leq B\} \quad (2.24)$$

*is compact in  $\mathbb{X}_0 = L_\infty(D)$  so that minimizing representation in (2.17) exist. Hence, the analog  $\mathfrak{C}_{N,s,\text{Lip}}(B) := \{G \in \mathfrak{C}_{N,s} : \|G\|_{N,s,\text{Lip}} \leq B\}$  is so as well.*

The proof of Proposition 2.7 is given in § 6.1.2.

Again, in view of Remark 2.6 and (2.21), compactness of  $\mathfrak{C}_{N,s}(B)$  and  $\mathfrak{C}_{N,s,\text{Lip}}(B)$  is trivial when  $s \geq d_0$

## 2.4 Approximation Classes

We are now prepared to quantify *compositional approximability*. In the spirit of [3], consider the “K-functional”

$$K_s(v, N, \delta) := \inf_{G \in \mathfrak{C}_{N,s}} \|v - G\|_\infty + \delta \|G\|_{N,s}. \quad (2.25)$$

In the same way we define  $K_{s,\text{Lip}}(v, N, \delta)$ , replacing  $\|\cdot\|_{N,s}$  by  $\|\cdot\|_{N,s,\text{Lip}}$ .

In what follows putting the subscript Lip in parentheses, like  $K_{s,(\text{Lip})}$  means that respective statements hold for either version.

Obviously,  $K_{s,(\text{Lip})}(v, N, \delta) \leq C\delta$  means that for some  $\bar{G} \in \mathfrak{C}_{N,s,(\text{Lip})}$  one has  $\|v - \bar{G}\|_\infty \leq C\delta$  and  $\|\bar{G}\|_{N,s,(\text{Lip})} \leq C$ , i.e., accuracy  $\delta$  is achieved with a controlled “composition-norm”.

Interrelating  $N$  and  $\delta$  is then a way to define collections of functions with a certain quantifiable *compositional approximability*. A (smooth) strictly increasing function  $\gamma : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  with  $\lim_{s \rightarrow \infty} \gamma(s) = \infty$ , is called a *growth function*. Its inverse  $\gamma^{-1}$  exists and is also a growth function. This is to be distinguished from  $\gamma(\cdot)^{-1} = 1/\gamma(\cdot)$ . Given such a growth function  $\gamma$ , we consider the class

$$\mathcal{A}_{(\text{Lip})}^{\gamma,s} := \{v \in \infty : \|v\|_{\mathcal{A}_{(\text{Lip})}^{\gamma,s}} := \|v\|_\infty + |v|_{\mathcal{A}_{(\text{Lip})}^{\gamma,s}} < \infty\}, \quad (2.26)$$

where

$$|v|_{\mathcal{A}_{(\text{Lip})}^{\gamma,s}} := \sup_{N \in \mathbb{N}} \gamma(N) K_{s,(\text{Lip})}(v, N, \gamma(N)^{-1}). \quad (2.27)$$

Since trivially

$$\|v - G_N\|_\infty \leq \gamma(N)^{-1} \gamma(N) \{ \|v - G_N\|_\infty + \gamma(N)^{-1} \|G_N\|_{N,s,(\text{Lip})} \} \leq \gamma(N)^{-1} |v|_{\mathcal{A}_{(\text{Lip})}^{\gamma,s}},$$

we will use the information  $v \in \mathcal{A}_{(\text{Lip})}^{\gamma,s}$  often in the form that for each  $N \in \mathbb{N}$  there exists  $G_N \in \mathfrak{C}_{N,s}$  such that

$$\|v - G_N\|_\infty \leq \gamma(N)^{-1} |v|_{\mathcal{A}_{(\text{Lip})}^{\gamma,s}}, \quad \|G_N\|_{N,s,(\text{Lip})} \leq \|v\|_{\mathcal{A}_{(\text{Lip})}^{\gamma,s}}, \quad N \in \mathbb{N}. \quad (2.28)$$

To put it differently, realizing target accuracy  $\varepsilon$  by a composition  $G_{\mathbf{g}_\varepsilon}$  is achievable within a complexity  $N = N_\varepsilon$  of the order

$$N_\varepsilon = \left\lceil \gamma^{-1} \left( \frac{|v|_{\mathcal{A}_{(\text{Lip})}^{\gamma,s}}}{\varepsilon} \right) \right\rceil. \quad (2.29)$$

A natural candidate for a *model class* can now be defined as follows. Let for any fixed  $B < \infty$

$$\mathcal{K}_{\gamma,s,(\text{Lip})}(B) := \{v \in \mathcal{A}_{(\text{Lip})}^{\gamma,s} : \|v\|_{\mathcal{A}_{(\text{Lip})}^{\gamma,s}} \leq B\}. \quad (2.30)$$

Again, on account of Remark 2.6 the classes  $\mathcal{A}^{\gamma,s}$ ,  $\mathcal{A}_{\text{Lip}}^{\gamma,s}$  agree with  $\text{Lip}_1$  when  $s \geq d_0$  and hence do not provide any useful information. In particular, by (2.21) and (2.28), balls in these classes are compact when  $s \geq d_0$ . Since  $\|v\|_{\text{Lip}_1} \leq \|v\|_{\mathcal{A}_{\text{Lip}}^{\gamma,s}}$  for all  $s \in \mathbb{N}$ , precompactness of  $\mathcal{K}_{\gamma,s,\text{Lip}}(B)$  is also immediate. Since in general  $\mathcal{A}^{\gamma,s}$  need not be contained in  $\text{Lip}_1$  for  $s < d$  the following claim requires an argument.

**Remark 2.8** (a) Obviously one has

$$\gamma(\cdot) \lesssim \tilde{\gamma}(\cdot) \Rightarrow \mathcal{A}_{(\text{Lip})}^{\tilde{\gamma},s} \subseteq \mathcal{A}_{(\text{Lip})}^{\gamma,s}. \quad (2.31)$$

(b) Whenever  $\gamma$  satisfies  $\gamma(N) \geq c_\gamma \gamma(2N)$  for  $N \in \mathbb{N}$ , the class  $\mathcal{A}_{(\text{Lip})}^{\gamma,s}$  is a linear space.

**Remark 2.9** The sets  $\mathcal{K}_{\gamma,s}(B)$ ,  $\mathcal{K}_{\gamma,s,\text{Lip}}(B)$  (see (2.30)) are for  $B < \infty$  and  $s \in \mathbb{N}$  compact.

**Proof:** Fix  $B < \infty$  and let  $(f_j)_{j \in \mathbb{N}}$  be a sequence in  $\mathcal{K}_{\gamma,s}(B)$ . Take a sequence  $(\varepsilon_k)_{k \in \mathbb{N}}$  of numbers decreasing monotonically to zero. For each  $f_j$  let  $\mathbf{g}_{j,k}$  denote a compositional representation of some function in  $\mathfrak{C}_{N_{\varepsilon_k},s}$  such that (see (2.28))  $\|G_{\mathbf{g}_{j,k}}\|_{N_{\varepsilon_k},s} \leq B$  and  $\|f_j - G_{\mathbf{g}_{j,k}}\|_\infty \leq \varepsilon_k$ . The complexity function  $\mathfrak{N}(\mathbf{g}_{j,k})$  is controlled uniformly in  $j$  by  $N_{\varepsilon_k}$ , defined by (2.29). For fixed  $k$  the class  $\mathfrak{C}_{N_{\varepsilon_k},s}$  is compact (Proposition 2.7). Therefore, for fixed  $k$ ,  $(G_{\mathbf{g}_{j,k}})_{j \in \mathbb{N}}$  contains a subsequence (again denoted by  $(G_{\mathbf{g}_{j,k}})_{j \in \mathbb{N}}$ ), converging uniformly to some  $G_k \in \mathfrak{C}_{N_{\varepsilon_k},s}$ . Now one can take a diagonalization argument, letting  $k$  tend to infinity, extracting a convergent subsequence from  $(f_j)_{j \in \mathbb{N}}$ .  $\square$

**Remark 2.10** (a) If  $\gamma$  satisfies  $\gamma(N) \geq c_\gamma(N_0)\gamma(N + N_0)$  for some  $N_0 \in \mathbb{N}$  and all  $N \in \mathbb{N}$ , then  $\mathcal{A}^{\gamma,s}$  is closed under composition with elements from  $\mathfrak{C}_{N_0,s}$  for all  $s \in \mathbb{N}$ . This covers a range of growth functions covering logarithmic up to exponential growth

(b) For equal input- and output-dimension  $\mathcal{A}_{\text{Lip}}^{\gamma,s}$  is closed under composition whenever  $\gamma(N) \geq c_\gamma \gamma(2N)$ . An analogous statement holds when  $v \in \mathcal{A}_{\text{Lip}}^{\gamma,s}(D, d')$ ,  $w \in \mathcal{A}_{\text{Lip}}^{\gamma,s}(D', d'')$ ,  $\dim D' = d''$ . The growth range covers any polynomial growth.

**Proof:** Let  $G_0 \in \mathfrak{C}_{N_0,s}$  with infimizing representation  $\mathbf{g}_0$ , and  $v \in \mathcal{A}^{\gamma,s}$ . For  $N \in \mathbb{N}$  let  $G_N \in \mathfrak{C}_{N,s}$  realize  $K_s(v, \gamma(N)^{-1})$  with minimizing representation  $\mathbf{g}_N$ . Then, on the one hand,

$$\begin{aligned} \|G_0 \circ v - G_0 \circ G_N\|_\infty &\leq |G_0|_{\text{Lip}_1} \|v - G_N\|_\infty \leq \|G_0\|_{N,s}^{N_0} \|v\|_{\mathcal{A}^{\gamma,s}} \gamma(N)^{-1} \\ &\leq c_\gamma(N_0)^{-1} \|G_0\|_{N,s}^{N_0} \|v\|_{\mathcal{A}^{\gamma,s}} \gamma(N + N_0)^{-1}. \end{aligned}$$

On the other hand  $\|G_0 \circ G_N\|_{N+N_0,s} \leq \max\{\|G_0\|_{N_0,s}, \|G_N\|_{N,s}\}$ . Hence,

$$\gamma(N + N_0) K_s(G_0 \circ v, \gamma(N + N_0)^{-1}) \leq c_\gamma(N_0)^{-1} \|G_0\|_{N,s}^{N_0} \|v\|_{\mathcal{A}^{\gamma,s}} + \max\{\|G_0\|_{N_0,s}, \|v\|_{\mathcal{A}^{\gamma,s}}\},$$

which shows (a).

Regarding (b), in a similar vein for  $v, w \in \mathcal{A}_{\text{Lip}}^{\gamma,s}$ ,  $N \in \mathbb{N}$ , there exist  $G_w, G_v \in \mathfrak{C}_{N,s}$  such that  $\|z - G_z\|_\infty \leq \|z\|_{\mathcal{A}_{\text{Lip}}^{\gamma,s}} \gamma(N)^{-1}$ ,  $z \in \{v, w\}$ . Thus,

$$\begin{aligned} \|w \circ v - G_w \circ G_v\|_\infty &\leq \|(w - G_w) \circ v\|_\infty + \|G_w(v) - G_w(G_v)\|_\infty \\ &\leq \left\{ \|w\|_{\mathcal{A}^{\gamma,s}} + |G_w|_{\text{Lip}_1} \|v\|_{\mathcal{A}^{\gamma,s}} \right\} \gamma(N)^{-1} \\ &\leq c_\gamma^{-1} \|w\|_{\mathcal{A}_{\text{Lip}}^{\gamma,s}} \left\{ 1 + \|v\|_{\mathcal{A}_{\text{Lip}}^{\gamma,s}} \right\} \gamma(2N)^{-1}, \end{aligned} \quad (2.32)$$

which confirms (b). □

Various further modifications of the above setting are conceivable and, depending on the context, useful. For instance,  $\|\cdot\|_{N,s,\text{Lip}}$  could be weakened by replacing the compositional Lipschitz constants  $L_{[n(\mathbf{g}),\ell+1]}(\mathbf{g})$  in (2.20) by  $\zeta(N)L_{[n(\mathbf{g}),\ell+1]}(\mathbf{g})$ , where  $\zeta(N) \rightarrow 0$ , which would permit some growth of the  $L_{[n(\mathbf{g}),\ell+1]}(\mathbf{g})$ , thereby no longer confining compositions to stay in  $\text{Lip}_1$ .

A particular scenario, that will be heavily used later, are classes of the form  $L_\infty(I; \mathcal{A}_{(\text{Lip})}^{\gamma,s})$ ,  $\text{Lip}_1(I; \mathcal{A}_{(\text{Lip})}^{\gamma,s})$  for some “time-interval”  $I = [0, T]$ , where we have used standard notation for Bochner spaces. A useful observation, whose proof is given in § 6.3.2, reads as follows.

**Proposition 2.11** *Let  $f \in \text{Lip}_1(I; \mathcal{A}_{(\text{Lip})}^{\gamma,s})$ . Then  $F(t, w) := \int_0^t f(s, w(s))ds$ ,  $t \in I$ , belongs to  $L_\infty(I; \mathcal{A}_{(\text{Lip})}^{\tilde{\gamma},s})$ , where for  $\phi(r) := r\gamma(r)$  we let  $\tilde{\gamma}(r) := \gamma(\phi^{-1}(r))$ .*

**Remark 2.12** *Typical scenarios where compositional approximations are expected to be relevant concern, of course, the range of dynamical systems, which is the primary context discussed in the literature. In addition, the approximation of solution manifolds (see (1.2)) is an interesting context as well and will be discussed in the following sections. States of interest are then given only implicitly as solutions of operator equations and often approximated through a wider range of iterative schemes. Below we exemplify this for a scenario where, among other things, fixed point iterations involving global operators apply.*

## 3 Linear Parametric Transport Equations

### 3.1 A Model Problem

As the lead example for (1.1), we consider the Cauchy problem for a linear (scalar) transport equation in  $m$  spatial dimensions ( $m \in \{1, 2, 3\}$ , say) with parameter dependent convection field

$$\partial_t u(t, x) + a(t, x, y) \cdot \nabla_x u(t, x) - f(t, x, y) = 0, \quad x \in \mathbb{R}^m, t \in I := [0, T), y \in \mathcal{Y} \quad (3.1)$$

subject to initial conditions

$$u(0, x, y) = u_0(x, y), \quad x \in \mathbb{R}^m, y \in \mathcal{Y}. \quad (3.2)$$

We assume for convenience that  $\text{supp } u_0 = \overline{D} \times \mathcal{Y}$  where  $D$  is a bounded domain. Hence, for a fixed time horizon  $T$  the solution, as a function of  $t, x, y$  can take values different from zero only in a bounded subset  $\Omega$  of  $I \times \mathbb{R}^m$ .

Since solutions depend on temporal, spatial variables and on parametric variables, possibly for large  $d_y$ , one faces approximation problems in high dimensions. We wish to understand under which circumstances one can avoid the Curse of Dimensionality without

requiring excessively high smoothness. Therefore, we impose only low or moderate smoothness conditions on the problem data. In general, we assume throughout:

$$\|a\|_{L_\infty(\Omega \times \mathcal{Y}; \mathbb{R}^m)} = A < \infty, \quad \text{for some } A \geq 1. \quad (3.3)$$

Furthermore we assume throughout that the convection field is uniformly Lipschitz continuous in space, time, and parametric variables

$$|a(t, z; y) - a(t', z', y')| \leq L \max\{|z - z'|, |t - t'|, |y - y'|\}, \quad (t, z, y), (t', z', y') \in \Omega \times \mathcal{Y}. \quad (3.4)$$

Here and later  $|\cdot|$  is understood as the  $\ell_\infty$ -norm when applied to vectors.

The very special case, where  $a$  is independent of  $t, x$ , is instructive. E.g. when  $f = 0$  we have

$$u(t, x, y) = u_0(x - ta(y), y). \quad (3.5)$$

This is a simple *composition* of the initial values with a linear function in  $(t, x)$  involving, however, a  $y$ -dependent coefficient, typically a function of a large number of variables. Using for a fixed time  $t = T$ , the solution field  $u(T, x; y) = u_0(x - Ta(y), y)$  as initial values for the corresponding backward problem, shows that the  $u(T, \cdot; \cdot)$  traverse essentially every element in a Lipschitz ball in  $L_\infty([0, T] \times D \times \mathcal{Y})$  so that stable approximations will suffer from the Curse of Dimensionality. This is in agreement with the findings in [23] where the only conditions on the convection field are given in terms of classical smoothness properties. So, more specific structural constraints on the convection field are needed to ensure that solutions  $u$  exhibit some *structural sparsity* that allows one to avoid the Curse of Dimensionality.

## 3.2 Dimension-Sparse Convection fields

In the light of the preceding comments we consider convection fields  $a$  of the type

$$a \in \text{Lip}_1(I; \mathcal{A}_{\text{Lip}}^{\gamma, s}(D \times \mathcal{Y}; \mathbb{R}^m)) \quad (3.6)$$

the Bochner spaces of functions that are Lipschitz in time with values in  $\mathcal{A}_{\text{Lip}}^{\gamma, s}$ . Here and below we set  $I = [0, T]$  for a fixed time horizon  $T$  and some dimension-sparsity  $s \leq m + 1 + d_y$  where  $d_y := \dim \mathcal{Y}$ .

**Remark 3.1** *We could also consider  $a \in \mathcal{A}_{\text{Lip}}^{\gamma, s}(I \times D \times \mathcal{Y}; \mathbb{R}^m)$  at the expense of more technical elaborations on finitely parametrized approximations.*

A particular case of interest concerns *affine parametric* expansions for the convection field (arising e.g. as Karhunen-Loève expansions of random convection fields)

$$a(t, x; y) = \sum_{j=1}^{d_y} y_j a_j(t, x) \in C(I \times \mathbb{R}^m \times \mathcal{Y}), \quad (3.7)$$

where  $\mathcal{Y} = [-1, 1]^{d_y}$ , i.e.,  $a : \mathbb{R}^{m+1+d_y} \rightarrow \mathbb{R}^m$ . In this case we will assume that

$$\sum_{j=1}^{d_y} \|a_j\|_{L_\infty(\Omega; \mathbb{R}^m)} \leq \Lambda < \infty. \quad (3.8)$$

holds for some constant  $\Lambda$  independent of  $d_y$ .

In fact, (3.7) is a special case of (3.6) because in this case  $a$  is easily shown to belong to  $\text{Lip}_1(I; \mathcal{A}_{\text{Lip}}^{\gamma, m+1})$  for *every* growth function  $\gamma$ .

### 3.3 Characteristics

The field of characteristics, given by the family of ODEs

$$\dot{z}(t) = a(t, z(t); y), \quad z(0) = x, \quad (3.9)$$

plays a pivotal role in what follows. Note then that the characteristics have a natural semi-group property, namely that they can be obtained by composing individual characteristic segments. More precisely, suppressing the dependence on  $y$  for a moment, we consider the solution of the more general initial value problem

$$\dot{z}(t, \tau; \bar{z}) = a(t, z(t)), \quad z(\tau) = \bar{z}, \quad (3.10)$$

often abbreviating  $z(t, 0; \bar{z}) = z(t; \bar{z})$  when  $\tau = 0$ . Thus, one has for any  $\tau$

$$z(t, x) = z(t, \tau; z(\tau; x)), \quad (3.11)$$

i.e., a point on a characteristic at time  $t$  can be reached by following the characteristic curve emanating from a point at an earlier time, in brief by *composing* characteristic segments.

There is a second angle regarding compositional approximations to characteristics, namely that (3.9) is equivalent to

$$z(t, x; y) = x + \int_0^t a(s, z(s; x; y); y) ds. \quad (3.12)$$

Since fixed point iterations are prototypes of compositional approximations that apply in much wider context we will actually stress this aspect in what follows and indicate some ramifications beyond the specific transport equation model later in § 5.

Under the above assumptions characteristics don't cross, i.e., the value of the solution to (3.7) can be determined by tracing back along characteristics. In fact, in view of (3.9), one has for the solution  $u$  of (3.1) (suppressing again the dependence on  $y$  for a moment)  $\frac{d}{dt}u(t, z(t)) = f(t, z(t))$ . Hence, recalling that  $u(0, z(0; x; y)) = u_0(x; y)$ ,

$$u(t, z(t; x; y), y) = u_0(x; y) + \int_0^t f(s, z(s; x; y)) ds, \quad (3.13)$$



or equivalently, using (3.11) and noting that when  $x = z(t, 0; \bar{x})$  one has  $z(s, 0; \bar{x}) = z(-(t - s), t; x) = z(s - t, t; x)$ ,

$$u(t, x, y) = u_0(z(-t, t; x, y), y) - \int_0^t f(s, z(s - t, t, x; y)) ds. \quad (3.14)$$

In summary, if the characteristics have “good (pointwise) compositional approximability”, for  $f = 0$ , the solution results from one additional composition, when  $f \neq 0$  there is another composition plus integration.

The central objective in what follows is to construct finitely parametrized surrogates  $\mathcal{N}(t, z, y)$  for the map

$$(t, z, y) \in I \times D \times \mathcal{Y} \mapsto u(t, z, y), \quad (3.15)$$

that are determined by possibly few degrees of freedom. The strategy rests on a *two-stage process*. First, we identify quantified *compositional approximations* that are not *finitely parametrized* yet. The issue is to identify, among all possible such representations, particular structural properties of compositions that lend themselves, in a second step, to *finitely parametrized* approximations so as to avoid the Curse of Dimensionality. The map in (3.15) results from this second step.

In other words, fixing  $T$ , we seek to construct uniform approximations to the solution manifolds

$$\mathcal{M}_z = z_{\leq T}(\mathcal{Y}) := \{z(\cdot, \cdot; y) : z(\cdot, \cdot; y) \text{ solves (3.1) on } \Omega \text{ for } y \in \mathcal{Y}\} \subset L_\infty(I \times D; \mathbb{R}^m), \quad (3.16)$$

and

$$\mathcal{M}_u = \mathcal{M}_u(u_0, f) := \{u(\cdot, \cdot, y) : u(\cdot, \cdot, y) \text{ solves (3.9) on } \Omega \text{ for } y \in \mathcal{Y}\} \subset L_\infty(I \times D), \quad (3.17)$$

from which we can then draw conclusions on their complexity in terms of widths and entropy numbers.

## 4 Main Results

The general flavor of the following results is: membership of the problem data (convection field, initial conditions, right hand side) to an approximation class (see § 2.4) imply membership of characteristics and solution to a certain approximation class. In view of (3.14), we begin with discussing the field of parameter dependent characteristics focusing on two exemplary types of growth functions

$$(\text{alg}): \quad \gamma(r) \sim C_\alpha r^\alpha, \quad \text{or} \quad (\text{exp}): \quad \gamma(r) \sim C_\alpha e^{\alpha r}, \quad (4.1)$$

for some  $\alpha > 0$ . (With a bit more technical effort the arguments extend to more refined scales like  $\gamma(r) \sim e^{\alpha r^\beta}$ , for some  $0 < \beta \leq 1$ .)

The first result, whose proof is given in § 6.3, asserts that dimension-sparsity of the convection field is inherited by the characteristic field and that the characteristic manifold belongs to a compositional approximation class approaching order-one growth when the order of the convection field increases.

**Theorem 4.1** *Assume that the convection field  $a$  belongs to  $\text{Lip}_1(I; \mathcal{A}_{\text{Lip}}^{\gamma, s})$  (see (3.6)) for some growth function  $\gamma$  of either type in (4.1). Then, the field of characteristics  $z$  belongs to  $L_\infty(I; \mathcal{A}_{\text{Lip}}^{\hat{\gamma}, s})$ , i.e.,  $\mathcal{M}_z \subset L_\infty(I; \mathcal{A}_{\text{Lip}}^{\hat{\gamma}, s})$ , where*

$$\hat{\gamma}(r) \sim e^{-LT} \begin{cases} Cr^{\frac{\alpha}{1+\alpha}} (\log_2 r)^{-1}, & \text{in case (alg), } C = \|a\|^{-\frac{1}{1+\alpha}} (Q_{A,L,T})^{-\frac{\alpha}{1+\alpha}}, \\ Cr(\log_2 r)^{-2}, & \text{in case (exp), } C = \alpha(Q_{A,L,T})^{-1}, \end{cases} \quad (4.2)$$

where  $Q_{A,L,T} = AL^2T^2 \max\{1, T\}$ .

More precisely, one has

$$\|z\|_{L_\infty(I; \mathcal{A}^{\hat{\gamma}, s})} \lesssim \left(1 + \left(\frac{\|a\|}{T}\right)^{\frac{1}{2}} e^{\|a\|T}\right), \quad (4.3)$$

where we abbreviate here and in what follows.

$$\|a\| := \|a\|_{L_\infty(I; \mathcal{A}_{\text{Lip}}^{\gamma, s})}. \quad (4.4)$$

As a next step we establish approximability by *finitely parametrized* functions under the assumptions of Theorem 4.1. For later use we keep tracking the dependence of proportionality factors on problem parameters.

**Theorem 4.2** *Assume that  $a \in \text{Lip}_1(I; \mathcal{A}_{\text{Lip}}^{\gamma, s})$  for growth functions according to (4.1). Let  $T \geq T_0 > 0$ . Then, for each  $\varepsilon > 0$  there exists a DNN  $\mathcal{N}_\varepsilon$  such that*

$$\|z - \mathcal{N}_\varepsilon\|_{L_\infty(\Omega \times \mathcal{Y}; \mathbb{R}^m)} \leq \varepsilon, \quad (4.5)$$

and

$$\#\mathcal{N}_\varepsilon \leq \begin{cases} B \left(\frac{\|a\|}{\varepsilon}\right)^{\frac{(1+\alpha)(1+s)}{\alpha}} \left|\log_2 \frac{e^{T\|a\|}}{\varepsilon}\right|^2, & \gamma(r) \sim r^\alpha, \\ C \left(\frac{\|a\|}{\alpha\varepsilon}\right)^{(1+s)} \left|\log_2 \frac{e^{T\|a\|}}{\varepsilon}\right|^{3+s}, & \gamma(r) \sim e^{\alpha r}. \end{cases} \quad (4.6)$$

where  $B \leq \bar{C}LT^{\frac{3\alpha+s+1}{\alpha}} e^{\frac{\|a\|T(1+\alpha)(1+s)}{\alpha}} \|a\|^s$ ,  $C \leq \bar{C}LT^3 e^{\|a\|T(1+s)} \|a\|^{s-1}$ , and  $\bar{C}$  depends only on  $s, A$  and  $T_0$ .

It is instructive to reformulate the above findings in terms of *convergence rates*, see § 6.3.

**Corollary 4.3** For each  $N \in \mathbb{N}$ , sufficiently large, there exists a DNN  $\mathcal{N}_N$  of complexity  $\#\mathcal{N}_N \leq N$ , such that

$$\|z - \mathcal{N}_N\|_{L_\infty(I \times D \times \mathcal{Y}; \mathbb{R}^m)} \leq e^{T\|a\|} \tilde{\gamma}(N)^{-1}, \quad (4.7)$$

where

$$\tilde{\gamma}(r) \sim \begin{cases} Br^{\frac{\alpha}{(1+\alpha)(1+s)}} |\log_2 r|^{-\frac{2\alpha}{(1+\alpha)(1+s)}}, & \text{when } \gamma(r) \sim r^\alpha, \\ C\alpha r^{\frac{1}{1+s}} |\log_2 r|^{-\frac{3+s}{1+s}}, & \text{when } \gamma(r) \sim e^{\alpha r}, \end{cases} \quad (4.8)$$

with

$$B \sim L^{-\frac{\alpha}{(1+\alpha)(1+s)}} T^{-\frac{3\alpha+s+1}{(1+\alpha)(1+s)}} \|a\|^{-(1+\frac{s\alpha}{(1+\alpha)(1+s)})}, \quad C \sim \alpha (LT^3 \|a\|^{2s})^{-\frac{1}{1+s}}. \quad (4.9)$$

In both theorems the exponential case can be seen as a “limit  $\alpha \rightarrow \infty$ ” of algebraic rates. For  $s = m + d_y$  the obtained rate would reflect the full Curse of Dimensionality. So a non-trivial dimension sparsity of the convection field appears to be one key to avoiding the Curse in the present context.

Nevertheless, some adverse dependence on (the potentially large) parametric dimension  $d_y$  may still be hidden in  $\|a\| = \|a\|_{L_\infty(I; \mathcal{A}_{\text{Lip}}^{\gamma, s})}$ . This will be illustrated by the following more specialized scenario (see §6.4 for a proof).

**Theorem 4.4** Assume that the convection field  $a$  has an affine representation (3.7), satisfying (3.4) and (3.8). Then, for any  $\varepsilon > 0$  there exists a DNN  $\mathcal{N}_\varepsilon$  such that

$$\|z - \mathcal{N}_\varepsilon\|_{L_\infty(I \times D \times \mathcal{Y}; \mathbb{R}^m)} \leq \varepsilon, \quad \#\mathcal{N}_\varepsilon \lesssim B e^{LT(m+1)} \left(\frac{Td_y}{\varepsilon}\right)^{m+1} \left|\log_2 \frac{Td_y}{\varepsilon}\right|^2, \quad (4.10)$$

where  $B \sim \max\{1, T^2\}$  with a uniform constant depending only on  $L, A, m, \Lambda$ .

Moreover, for each  $N \in \mathbb{N}$  (sufficiently large) there exists a DNN  $\mathcal{N}_N$  with complexity  $\#\mathcal{N}_N \leq N$  such that

$$\|z - \mathcal{N}_N\|_{L_\infty(I \times D \times \mathcal{Y})} \leq e^{LT} \gamma(N)^{-1}, \quad \gamma(r) \sim d_y^{-1} r^{\frac{1}{m+1}} |\log_2 r|^{-\frac{2}{m+1}} \quad (4.11)$$

where the proportionality constant depends only on  $m, L, A, \Lambda$ . In fact,  $z \in L_\infty(I; \mathcal{A}_{\text{Lip}}^{\gamma, m})$  and

$$\mathcal{M}_z \subset \left\{v \in L_\infty(I; \mathcal{A}_{\text{Lip}}^{\gamma, m}) : \|v\|_{L_\infty(I; \mathcal{A}_{\text{Lip}}^{\gamma, m})} \leq 2e^{KT}\right\} \quad (4.12)$$

where,  $K$  depends on  $L, \Lambda, A$ .

The network  $\mathcal{N}_\varepsilon$  in Theorem 4.4 is a function of spatial and parametric variables and also, as a piecewise constant, of time. At the expense of further technical stipulations one could arrive at a piecewise affine in time representation (piecewise linear interpolation of time snapshots in time) that inherits the Lipschitz-in-time property (see Remark 6.2 later below).

While the dependence on the number of variables  $1 + m + d_y$  is only algebraic, so that, as in Theorem 4.2 when  $s \ll d_y$ , the Curse of Dimensionality is indeed avoided, one may ask

whether the factor  $d_y^{m+1}$  in (4.10) has the lowest possible order. In this regard, the proofs are so far solely based on compositional approximations derived from *fixed point iterations* for the field of characteristics. They have not made use of the fact that characteristics have a natural compositional structure reflected by (3.11). Combining both angles turns out to derive quantitative improvements regarding the dependence of constants on  $T$  and  $d_y$  (which is the reason for having kept track of the dependence of constants on various problem parameters). We keep both versions nevertheless separately for two reasons. First, the subsequent improvement builds on Theorem 4.4, applied to judiciously chosen intermediate time slabs. Second, in the form presented below, at least formally, it provides an approximation to the characteristic field at a single time snapshot  $T$ .

**Theorem 4.5** *Under the same hypothesis on the convection field as in Theorem 4.4 there exists for every  $\varepsilon > 0$  a DNN  $\mathcal{N}_\varepsilon$  such that*

$$\|z(T, \cdot; \cdot) - \mathcal{N}_\varepsilon\|_{L_\infty(D \times \mathcal{Y}; \mathbb{R}^m)} \leq \varepsilon, \quad (4.13)$$

and

$$\#\mathcal{N}_\varepsilon \lesssim T d_y L^{-(m+1)} e^{L(T+d_y^{-1})(m+1)} \varepsilon^{-(m+1)} \left( \log_2 \frac{e^{LT}}{\varepsilon} \right)^2. \quad (4.14)$$

with an absolute constant, depending only on  $\Lambda, A, m$ . This can be restated in terms of convergence rates with growth function

$$\tilde{\gamma}(r) \sim (T d_y)^{\frac{1}{m+1}} e^{-L(T+d_y^{-1})} r^{\frac{1}{m+1}} \left| \log_2 \frac{r}{T d_y} \right|^{-\frac{2}{m+1}},$$

see (6.110).

As one can see from the proof of Theorem 4.5 in § 6.4, the construction of  $\mathcal{N}_\varepsilon$  in (4.13) does contain the information on a fine time mesh in  $[0, T]$ , i.e., similar bounds can be derived for the full time range. A finitely parametrized representation is simply obtained by piecewise linear in time interpolation of the piecewise constants in time. Again we dispense with stating this formally because of additional technical efforts which are not essential at this point.

**Remark 4.6** *It should be emphasized though that neither construction is based directly on a conventional time-stepping scheme for systems of ODEs and it seems that this latter approach seems to lead to much weaker results.*

We discuss next approximability of the parameter dependent solutions themselves. A preliminary observation is stated without proof and refer to § 6.3.2 for some comments on the arguments.

**Remark 4.7** *We mention in passing that, using (2.31) and Remark 2.10, combined with Proposition 2.11, one can show that, when  $u_0, f$  belong to approximation classes with stronger growth than  $\gamma$  from (4.11), the solution manifold  $\mathcal{M}_u$  is contained in an approximation class  $L_\infty(I; \mathcal{A}_{\text{Lip}}^{\tilde{\gamma}, m})$  where (up to factors with low order algebraic and logarithmic dependence on  $d_y$ )  $\tilde{\gamma}(r) \gtrsim r^{\frac{1}{m+2}} |\log_2 r|^{-\omega}$ ,  $\omega \approx 2m^{-1}$ .*

Instead we turn to more precise statements about DNN approximations and focus on the dependence of constants on the problem data  $u_0, f$ . To simplify the presentation we dispense with tracking the dependence on  $T$ . In addition, aside from the dependencies stated explicitly in the following result, whose proof is given in §6.5, the constants are understood to depend on  $T, m, L, A, \Lambda$  (see §3.1, 3.2).

**Theorem 4.8** *Under the same hypotheses on the convection field  $a$  ((3.7), (3.4), (3.8)) assume that for given growth functions*

$$\gamma_0(r) \sim r^{\alpha_0}, \quad \gamma_f(r) \sim r^{\alpha_f}, \quad r \geq r_0, \quad (4.15)$$

the data  $u_0, f$  satisfy  $u_0 \in \mathcal{A}_{\text{Lip}}^{\gamma_0, m}$ ,  $f \in \text{Lip}_1(I; \mathcal{A}_{\text{Lip}}^{\gamma_f, m})$ . Without loss of generality we assume that  $d_y \geq \|u_0\|_{\mathcal{A}_{\text{Lip}}^{\gamma_0, m}}^{\alpha_0}, \|f\|_{L_\infty(I; \mathcal{A}_{\text{Lip}}^{\gamma_f, m})}$ . Then, for any  $\varepsilon > 0$  there exists a DNN  $\mathcal{N}_{u, \varepsilon}$  such that for the exact solution  $u$  of the transport equation (3.1) satisfies

$$\|u - \mathcal{N}_{u, \varepsilon}\|_{L_\infty(I \times D \times \mathcal{Y})} \leq \varepsilon, \quad \#\mathcal{N}_{u, \varepsilon} \lesssim B d_y^{m+1} \varepsilon^{-\frac{(m+1)(1+\alpha)}{\alpha}} \left| \log_2 \frac{d_y}{\varepsilon} \right|^2, \quad (4.16)$$

where

$$\alpha := \min\{\alpha_0, \alpha_f\}, \quad B \lesssim \left( 1 + \|u_0\|_{\mathcal{A}_{\text{Lip}}^{\gamma_0, m}}^{\frac{\alpha+1}{\alpha}} + \|f\|_{L_\infty(I; \mathcal{A}_{\text{Lip}}^{\gamma_f, m})}^{\frac{\alpha+1}{\alpha}} \right)^{\frac{(m+1)(\alpha+1)}{\alpha}}, \quad (4.17)$$

with a constant depending on  $L_0, L_f, L, A, T, \|a\|$ .

Moreover, for  $N \geq N_0$  there exists a DNN  $\mathcal{N}_N$  with  $\#\mathcal{N}_N \leq N$  such that the approximation rate

$$\|u - \mathcal{N}_N\|_{L_\infty(I \times D \times \mathcal{Y})} \leq \left( 1 + \|u_0\|_{\mathcal{A}_{\text{Lip}}^{\gamma_0, m}}^{\frac{\alpha+1}{\alpha}} + \|f\|_{L_\infty(I; \mathcal{A}_{\text{Lip}}^{\gamma_f, m})}^{\frac{\alpha+1}{\alpha}} \right) \hat{\gamma}(N)^{-1}, \quad N \geq N_0, \quad (4.18)$$

holds where

$$\hat{\gamma}(r) := C d_y^{-\frac{\alpha}{\alpha+1}} r^{\frac{\alpha}{(m+1)(\alpha+1)}} \left| \log_2 r d_y^{\frac{1}{\alpha}} \right|^{-\frac{2\alpha}{(m+1)(\alpha+1)}}, \quad (4.19)$$

with a constant  $C$  depending only on the problem parameters from § 3.

The common trait of all network size bounds is a factor  $\varepsilon^{-(m+1)\beta}$  for some  $\beta \geq 1$ . If  $\beta > 1$  it tends to 1 with increasing compositional approximation orders of the problem data. This corresponds to the entropy numbers of a Lipschitz ball of functions of  $m+1$  variables. According to [2] this marks the best achievable rate by a stable algorithm of any type for approximands ranging over such a Lipschitz ball. Note that even without parameter dependence, when the convection field  $a(t, z)$  is allowed to be any element of such a Lipschitz ball, it is easy to see that the spatial smoothness of the characteristics is at most of order one as well (while one can gain an order of smoothness in time). This suggests that the size order  $\varepsilon^{-(m+1)}$  should be (near-)optimal.

We leave a rigorous detailed discussion to future work but close this section with a brief indication of the direction of thought. Recall from [7] that for a compact set  $\mathcal{K}$  in a Banach space  $\mathbb{X}$  the notion of *nonlinear widths* is based on considering pairs of mappings

$$a_N : \mathcal{K} \rightarrow \mathbb{R}^N, \quad M_N : \mathbb{R}^N \rightarrow \mathbb{X}, \quad (4.20)$$

forming for  $f \in \mathcal{K}$  approximations  $A_N(f) := (M_N \circ a_N)(f)$ , described by  $N$  parameters  $a_N(f) \in \mathbb{R}^N$ . The so called *manifold widths*  $\delta_N(\mathcal{K})_{\mathbb{X}}$  are defined by minimizing  $E_{a_N, M_N}(f) := \|f - A_N(f)\|_{\mathbb{X}}$  over such pairs, subject to some *stability constraints* on  $a_N, M_N$ . Such constraints are necessary to avoid trivial approximations based on space-filling curves, see e.g. [2, 6]. Minimal such constraints require just continuity of the mappings  $a_N, M_N$  leading to the notion of manifold widths, see [7]. Specializing this to  $\mathcal{K} = \mathcal{M}_z$ , a natural realization of  $a_N$  is the collection of weights in the networks with the above convergence properties, viewed as a vector  $\theta \in \mathbb{R}^N$ . Then  $M_N$  is the resulting network realization. It follows from the proofs in § 6 that these mappings are indeed continuous, establishing manifold-widths rates that avoid the Curse of Dimensionality (see Theorems 4.4, 4.5) in contrast to what one expects for Kolmogorov widths.

However, from a practical point of view, continuity is still a rather weak requirement. To warrant *stability* of approximations, somewhat stronger constraints on the mappings  $a_N, M_N$  are needed which leads to the notion of *stable widths* suggested in [2]. Specifically, they still satisfy a version of Carl's inequality asserting (up to logarithmic factors) that such widths are bounded from below by the *entropy numbers* of  $\mathcal{K}$  (here we focus on  $\mathcal{K} = \mathcal{M}_z$ ). Recall that the entropy numbers are defined by

$$\varepsilon_n(\mathcal{K})_{\mathbb{X}} := \min\{\varepsilon > 0 : \mathcal{K} \text{ can be covered by } 2^n \text{ balls of radius } \varepsilon\}$$

specifying the number of bits needed to encode an object in  $\mathcal{K}$  within accuracy  $\varepsilon$ , say. Among these stronger requirements on  $a_N, M_N$ , Lipschitz continuity of  $M_N \circ a_N$  is confirmed later in § 6 (see (6.28), Lemma 6.9). Lipschitz continuity of  $M_N$  follows, in principle, along the lines in [6, § 5]. At least for affine parameter dependence (3.7) Proposition 2.2 helps controlling Lipschitz constants to an extent that one can indeed derive from Theorem 4.4 bounds of the type

$$\varepsilon_n(\mathcal{M}_z)_{\infty} \lesssim d_y n^{-\frac{1}{m+1}} (\log_2 n)^{\frac{3}{m+1}}, \quad n \in \mathbb{N}, \quad (4.21)$$

following in essence the reasoning in [2], [6, § 5]. If these estimates were sharp, optimality of the bounds for the stable manifold widths would indeed follow (up to a logarithmic factor). Sharpness of (4.21) is however not clear from our preliminary arguments so that further work is needed.

## 5 Comments, Consequences, and Outlook

Under stronger smoothness conditions on the convection field, right hand side, and initial values, one could derive better rates along the same lines. In particular, one could use

versions of Proposition 2.2 from [15] that exploit higher Sobolev regularity of approximands. However, from an application viewpoint further weakening smoothness assumptions on the data would perhaps be more interesting.

Specifically, it would be desirable to include less regular right hand sides and initial (boundary) values (e.g. including discontinuities) and even convection fields. For instance, further modifying the  $K$ -functionals could permit dimension sparse approximations to the convection field with gradually increasing Lipschitz constants. Discontinuous right hand sides or initial conditions suggest using  $L_p$  (especially  $L_2$ ) norms for solutions while pointwise accuracy estimates for characteristics would still be reasonable and desirable. In fact, once accurate surrogates on the characteristics are available that can be evaluated efficiently, (e.g. in terms of DNNs), one can apply supervised learning techniques to approximate solutions, again using sampled versions of (3.14) as training data. For deterministic as well as probabilistic estimates sharp bounds on the entropy numbers would be important. These issues will be addressed in forthcoming work.

In a different direction, we plan to apply the above concepts to a wider scope of transport problems. In particular, the accuracy controlled solvers for *Radiative Transfer* models developed in [4] build on fixed point iterations where each step requires an approximate solve of a parametric linear transport problem. Furthermore, nonlinear scalar conservation laws are known to be equivalent to a parametric family of linear transport equations [18] which in turn can be approached via fixed point iterations [26].

## 6 Proofs

### 6.1 Back to Compositions

#### 6.1.1 Proof of Proposition 2.2

In this section we build mainly on findings from [15, 27]. Consider the “hat-function”  $\phi(x) := (1 - |x|)_+ = \max\{0, 1 - |x|\}$ ,  $x \in \mathbb{R}$  as well as the scaled and shifted versions  $\phi_{i,h}(x) := \phi(h^{-1}x - i)$ ,  $i \in \mathbb{Z}$ , with support  $S_i = [(i-1)h, (i+1)h]$ . We let  $h = 1/q$  for some integer  $q \in \mathbb{N}$ , so that the restrictions of the  $\phi_{i,h}$  to  $[0, 1]$  form a stable basis for all piecewise linears on  $(0, 1)$  subordinate to the partition induced by the nodes  $\{ih = i/q : i = 0, \dots, q\}$ . Since each  $\phi_{i,h}$  is a second order divided difference of the ReLU rectifier  $\sigma(x) := x_+$  with respect to the nodes  $ih$  it has an exact representation as a univariate neural network of fixed finite depth and a fixed finite number of weights. The Lipschitz constant of  $\phi_{i,h}$  and hence of this network is clearly  $h^{-1}$ . Abbreviating  $\mathbf{i} := (i_1, \dots, i_s) \in \{0, \dots, q\}^s$  we consider next for  $x = (x_1, \dots, x_s) \in \mathbb{R}^s$  the tensor products  $\phi_{\mathbf{i},h} := \phi_{i_1,h}(x_1) \cdots \phi_{i_s,h}(x_s)$ , which obviously satisfy  $\phi_{\mathbf{i},h}(\mathbf{i}') = \delta_{\mathbf{i},\mathbf{i}'}$  for any  $\mathbf{i}, \mathbf{i}' \in \mathcal{I}_h := \{0, h, \dots, hq\}^s$  while we still have  $\|\partial_j \phi_{\mathbf{i},h}\|_\infty \leq h^{-1}$ . The next step consists in approximating each  $\phi_{\mathbf{i},h}$ , viz. a product of univariate ReLU networks of fixed depth and number of weights by a ReLU network of input dimension  $s$ . This is where one uses that the function  $M : \nu = (\nu_1, \dots, \nu_s) \mapsto \prod_{j=1}^s \nu_j$  can be approximated by a ReLU

network  $\mathcal{N}_{M,\delta}$  according to

$$\|M - \mathcal{N}_{M,\delta}\|_{W^k(L_\infty((0,1)^s))} \leq \delta, \quad k \in \{0, 1\}. \quad (6.1)$$

where the depth of  $\mathcal{N}_{M,\delta}$  as well as  $\#\mathcal{N}_{M,\delta}$  is bounded by a constant multiple of  $\log_2 \delta^{-1}$ , with constants depending only on  $s$ . Moreover,  $\mathcal{N}_{M,\delta}(0) = 0$ . The case  $k = 0$  in (6.1) appears already in [27]. A key observation in [15, § C] is that  $k = 1$  still holds under the same complexity bounds. This is then used to show that for each  $\mathbf{i} \in \mathcal{I}_h$  there exists a ReLU network  $\mathcal{N}_{\mathbf{i},\delta}$  such that

$$\begin{aligned} \|\phi_{\mathbf{i},h} - \mathcal{N}_{\mathbf{i},\delta}\|_{W^k(L_\infty((0,1)^s))} &\leq c^k \delta h^{-1}, \quad k \in \{0, 1\}, \\ \|\mathcal{N}_{\mathbf{i},\delta}\|_{\text{Lip}_1} &\leq c \delta^{-1}, \quad \text{supp } \mathcal{N}_{\mathbf{i},\delta} \subseteq \text{supp } \phi_{\mathbf{i},h}, \end{aligned} \quad (6.2)$$

(with a constant  $c$ , depending on  $s$ ) and

$$\#\mathcal{N}_{\mathbf{i},\delta}, \text{ depth of } \mathcal{N}_{\mathbf{i},\delta} \lesssim \log_2 \frac{1}{\delta}, \quad (6.3)$$

with constants depending only on  $s$ .

Now, given  $g$ , consider the interpolant

$$g_h := \sum_{\mathbf{i} \in \mathcal{I}_h} g(\mathbf{i}h) \phi_{\mathbf{i},h}. \quad (6.4)$$

Obviously  $\|g_h\|_\infty \leq \|g\|_\infty$ . We claim that  $g_h$  is also Lipschitz continuous. To see this, let  $\mathcal{I}_h(s) := \{\mathbf{i} \in \mathcal{I}_h : \phi_{\mathbf{i},h}(x) \neq 0\}$  denote the collection of those nodes whose basis functions contain  $x \in (0,1)^s$  in the interior of their support. Then for  $x \in (0,1)^s$  let  $\mathbf{i}(x) \in \mathcal{I}_h(x)$  denote the node closest to the Chebyshev center of the convex hull  $[\mathcal{I}_h(x)]$  of  $\mathcal{I}_h(x)$ . Then, since  $g(\mathbf{i}(x)h) \sum_{\mathbf{i} \in \mathcal{I}_c(x)} \phi_{\mathbf{i},h}(x)$  is constant in a neighborhood of  $x$ , one has

$$|\partial_j g_h(x)| = \left| \sum_{\mathbf{i} \in \mathcal{I}_h(x)} (g(\mathbf{i}h) - g(\mathbf{i}(x)h)) \partial_j \phi_{\mathbf{i},h}(x) \right| \leq \max_{\mathbf{i} \in \mathcal{I}_h(x)} |g(\mathbf{i}h) - g(\mathbf{i}(x)h)| \frac{1}{h} \#(\mathcal{I}_h(x)).$$

Since  $\text{diam } \mathcal{I}_h(x) \leq ch$  for a constant depending on  $s$ , this yields

$$|\partial_j g_h(x)| \leq c^{-1} \|g\|_{\text{Lip}_1}, \quad x \in (0,1)^s, \quad (6.5)$$

from which it follows that (weakly)

$$|g_h(x) - g_h(x')| = \left| \int_0^1 \nabla g_h(x + t(x' - x)) \cdot (x' - x) dt \right| \leq |x - x'| \sqrt{s} c^{-1} \|g\|_{\text{Lip}_1}. \quad (6.6)$$

Moreover,

$$|g(x) - g_h(x)| = \left| \sum_{\mathbf{i} \in \mathcal{I}_h(x)} (g(x) - g(\mathbf{i}h)) \phi_{\mathbf{i},h} \right| \leq Ch \|g\|_{\text{Lip}_1}, \quad (6.7)$$



since  $\max\{|x - \mathbf{i}h| : \mathbf{i} \in \mathcal{I}_h(x)\} \leq Ch$  with  $C$  depending only on  $s$ . Given  $\delta > 0$ , the choice  $h = h(\delta) \leq \frac{\delta}{2C\|g\|_{\text{Lip}_1}}$  ensures  $|g(x) - g_h(x)| \leq \frac{\delta}{2}$ . Now approximate each  $\phi_{\mathbf{i},h(\delta)}$  by a ReLU network  $\mathcal{N}_{\mathbf{i},\delta}$  with accuracy  $\|\phi_{\mathbf{i},h(\delta)} - \mathcal{N}_{\mathbf{i},\delta}\|_\infty \leq c^*\delta$ , with  $c^*$  to be determined in a moment. We obtain, by (6.2)

$$\begin{aligned} \left\| g - \sum_{\mathbf{i} \in \mathcal{I}_h(\delta)} g(\mathbf{i}h) \mathcal{N}_{\mathbf{i},\delta} \right\|_\infty &\leq \|g - g_h\|_\infty + \sup_{x \in (0,1)^s} \sum_{\mathbf{i} \in \mathcal{I}_h(\delta)(x)} |g(\mathbf{i}h)| |\phi_{\mathbf{i},h(\delta)}(x) - \mathcal{N}_{\mathbf{i},\delta}(x)| \\ &\leq \frac{\delta}{2} + \sup_{x \in (0,1)^s} \#\mathcal{I}_h(\delta)(x) c^* \delta. \end{aligned}$$

Thus, choosing  $c^* = (2 \sup_{x \in (0,1)^s} \#\mathcal{I}_h(\delta)(x))^{-1}$ , we have confirmed

$$\left\| g - \sum_{\mathbf{i} \in \mathcal{I}_h(\delta)} g(\mathbf{i}h) \mathcal{N}_{\mathbf{i},\delta} \right\|_\infty \leq \delta. \quad (6.8)$$

Moreover, setting

$$\mathcal{N}_\delta := \sum_{\mathbf{i} \in \mathcal{I}_h} g(\mathbf{i}h) \mathcal{N}_{\mathbf{i},\delta},$$

we obtain, again by (6.2) and (6.5),

$$\begin{aligned} |\partial_j \mathcal{N}_\delta(x)| &\leq |\partial_j (\mathcal{N}_\delta(x) - g_h(x))| + c^{-1} \|g\|_{\text{Lip}_1} = \left| \sum_{\mathbf{i} \in \mathcal{I}_h(\delta)} g(\mathbf{i}h) (\partial_j \mathcal{N}_{\mathbf{i},\delta}(x) - \partial_j \phi_{\mathbf{i},h}) \right| + c^{-1} \|g\|_{\text{Lip}_1} \\ &\leq c' \|g\|_\infty \#(\mathcal{I}_h(\delta)) h^{-1} \delta + c^{-1} \|g\|_{\text{Lip}_1} \leq c'' (\|g\|_\infty + 1) \|g\|_{\text{Lip}_1}, \end{aligned}$$

where  $c''$  depends only on  $s$  and where we have used that  $\frac{\delta}{h} \leq 4C\|g\|_{\text{Lip}_1}$ . This confirms (2.3). Regarding the complexity (2.4) of  $\mathcal{N}_\delta$ , we have

$$\#(\mathcal{I}_h(\delta)) = h(\delta)^{-s} \leq c\delta^{-s} \|g\|_{\text{Lip}_1}^s,$$

which completes the proof because  $\#\mathcal{N}_\delta \lesssim \#(\mathcal{I}_h(\delta)) \log_2 \frac{1}{\delta}$ .  $\square$

### 6.1.2 Proof of Proposition 2.7

For each finite  $N$  there is only a finite number of feasible dimensionality vectors  $\mathbf{D}$  with  $\mathfrak{N}(\mathbf{D}) \leq N$  which representations of  $G \in \mathfrak{C}_N$  may have. For each such  $\mathbf{D}$  consider first the following auxiliary classes. Let  $\mathbb{F}_\ell \subset C(\mathbb{R}^{d_\ell-1}; \mathbb{R}^{d_\ell})$  be compact and let

$$\mathfrak{C}(\mathbf{D}, \mathbb{F}) := \{G \in \infty_0, G = G_{\mathbf{g}}, \mathbf{D}(\mathbf{g}) = \mathbf{D} : \text{with } g^j \in \mathbb{F}_j, j = 1, \dots, n(\mathbf{D})\}, \quad (6.9)$$

and

$$\mathfrak{C}_N(\mathbb{F}) := \bigcup_{\mathfrak{N}(\mathbf{D}) \leq N} \mathfrak{C}(\mathbf{D}, \mathbb{F}). \quad (6.10)$$

**Lemma 6.1** *The collection  $\mathfrak{C}_N(\mathbb{F})$  is compact in  $C(D_0; \mathbb{R}^{d_n(D)})$  (and so are the subsets  $\mathfrak{C}(D, \mathbb{F})$ ).*

To establish first this Lemma, it is enough to confirm compactness of  $\mathfrak{C}(D, \mathbb{F})$  for each of the eligible  $D$ . To this end, let  $\mathbb{F}, \mathbb{F}'$ , be dimensionally compatible compact subclasses so that for  $g \in \mathbb{F}, h \in \mathbb{F}'$  compositions  $h \circ g$  are defined. Then  $\{h \circ g : h \in \mathbb{F}', g \in \mathbb{F}\}$  is compact in  $\infty$ . In fact, let  $(g_n)_{n \in \mathbb{N}}, (h_n)_{n \in \mathbb{N}}$  be uniformly bounded sequences in  $\mathbb{F}, \mathbb{F}'$ , respectively. By Arzela-Ascoli's Theorem they are equicontinuous and have a convergent subsequence with continuous limits  $g, h$ , say. Denote these subsequences again by  $(g_n)_{n \in \mathbb{N}}, (h_n)_{n \in \mathbb{N}}$ . Then,

$$\|h_n \circ g_n - h \circ g\|_{\infty(D_g)} \leq \|h_n \circ g_n - h_n \circ g\|_{\infty(D_g)} + \|h_n \circ g - h \circ g\|_{\infty(D_g)}.$$

By uniform convergence of the  $g_n$  and equicontinuity of the  $h_n$  the first summand becomes arbitrarily small for  $n$  large enough. By uniform convergence of the  $h_n$  the second summand gets small as well. Iterating this argument, shows that  $\mathfrak{C}(D, \mathbb{F})$  is compact. Since  $\mathfrak{C}_N(\mathbb{F})$  is a finite union of such sets the assertion of Lemma 6.1 follows.

The proof of Proposition 2.7 follows now from noticing that membership to  $\mathfrak{C}_N(B)$  requires all composition factors to have uniformly bounded Lipschitz norm and hence belong to compact classes.  $\square$

## 6.2 Further Prerequisites

### 6.2.1 "Inverting" Growth Functions

As the above theorems indicate we are particularly interested in growth functions  $\gamma$  of algebraic growth whose inverses determine, in view of (2.29), the complexity.

**Lemma 6.2** *Assume that for positive  $b_1, b_2, \zeta$  and real  $\beta$*

$$\gamma(s) = b_1 s^\zeta |\log_2 b_2 s|^\beta, \quad s \geq s_0 \geq 1. \quad (6.11)$$

*Then*

$$\gamma^{-1}(r) \sim b_1^{-1/\zeta} \zeta^{\beta/\zeta} r^{\frac{1}{\zeta}} |\log_2(b_2^\zeta r / b_1)|^{-\frac{\beta}{\zeta}}, \quad r \geq r_0 \geq 1. \quad (6.12)$$

**Proof:** The proof is elementary but tedious, here a sketch. Making the ansatz  $\gamma^{-1}(r) \sim F r^{\frac{1}{\zeta}} |\log_2(Qr)|^\theta$ , we have

$$\begin{aligned} s &= \gamma^{-1}(\gamma(s)) \sim F (b_1 s^\zeta |\log_2(b_2 s)|^\beta)^{\frac{1}{\zeta}} \left| \log_2 (Q b_1 s^\zeta |\log_2(b_2 s)|^\beta) \right|^\theta \\ &= F (b_1 s^\zeta |\log_2(b_2 s)|^\beta)^{\frac{1}{\zeta}} \zeta^\theta \left| \log_2 ((Q b_1)^{\frac{1}{\zeta}} s) \right|^\theta \left| 1 + \frac{|\log_2 |\log_2(b_2 s)|^\beta|}{|\log_2(Q b_1 s^\zeta)|} \right|^\theta. \end{aligned} \quad (6.13)$$

Equating coefficients yields

$$F = b_1^{-\frac{1}{\zeta}} \zeta^{\frac{\beta}{\zeta}}, \quad Q = \frac{b_2^\zeta}{b_1}, \quad \theta = -\frac{\beta}{\zeta},$$

which confirms the claim.  $\square$

Lemma 6.2 will be perpetually used in the following way. Suppose we have established an error bound  $\varepsilon$  at the expense of  $\omega(\varepsilon^{-1})$  degrees of freedom where  $\omega$  is a growth function. Then, one concludes a convergence rate  $O(\hat{\gamma}(N)^{-1})$ , when the number  $N$  of invested degrees of freedom tends to infinity, where  $\hat{\gamma} \sim \omega^{-1}$ .

### 6.2.2 Fixed-Point Iterations and Lipschitz Bounds

For the convenience of the reader we recall the following frequently used classical facts.

**Remark 6.1** *Under the above assumptions on the convection field  $a$ , it follows from a classical Gronwall inequality that one has indeed*

$$\|z(\cdot, x) - z(\cdot, \bar{x})\|_{L_\infty(I)} \leq e^{L|I|}|x - \bar{x}|. \quad (6.14)$$

It can be also seen from (3.12) that  $z(t, \cdot)$  is Lipschitz with respect to  $t$

$$|z(s, x) - z(\xi_i; x)| = \left| \int_{\xi_i}^s a(s, z(s; x)) ds \right| \leq A|s - \xi_i|. \quad (6.15)$$

To find approximate compositions we recall the fixed-point relation (3.12) and consider the corresponding mapping  $\Phi_x = \Phi(\cdot; t, y) : L_\infty(I \times D \times \mathcal{Y}; \mathbb{R}^m) \rightarrow L_\infty(I \times D \times \mathcal{Y}; \mathbb{R}^m)$ , defined by,

$$\Phi_x(t, \bar{z}; y) := x + \int_0^t a(s, \bar{z}(s), y) ds, \quad t \in I = [0, T]. \quad (6.16)$$

A natural strategy is to approximate the fixed point of (3.12) by iterates of the mapping  $\Phi_x(\cdot; t, y)$ . In this case the arguments  $x, \bar{z}$  are not independent of each other. In fact, a natural choice would be the constant-in-time function

$$\bar{z}(x) = x, \quad s \in I. \quad (6.17)$$

Then, we always use the notational convention

$$\Phi_x^k(t, \bar{z}(x); y) := \Phi_x(\Phi_x^{k-1}(t, \bar{z}(x); y)). \quad (6.18)$$

In order to facilitate later applications of Remark 2.1 we do not tie  $\bar{z}$  to the particular choice (6.17), and view  $\bar{z}$  for now as a separate argument. Accordingly, iterates of the form (6.18) are understood with different initializations  $\bar{z}$ .

We collect first some simple facts concerning Lipschitz properties of iterates of these mappings.

**Lemma 6.3** *Under the above assumptions one has*

$$|\Phi_x^k(t, \bar{z}; y) - \Phi_{x'}^k(t, \bar{z}'; y')| \leq e^{LT} |x - x'| + \frac{(LT)^k}{k!} \max\{\|\bar{z} - \bar{z}'\|_{L_\infty(I; \mathbb{R}^m)}, |y - y'|\}, \quad k \in \mathbb{N}, \quad (6.19)$$

for all  $\bar{z}, \bar{z}' \in \Omega$ ,  $y, y' \in \mathcal{Y}$ . In particular, one has for all  $(t, x, y), (t, x', y') \in I \times D \times \mathcal{Y}$

$$|\Phi_x^k(t, \bar{z}(x); y) - \Phi_{x'}^k(t, \bar{z}(x'); y')| \leq \max\{|x - x'|, |y - y'|\} \sum_{\ell=0}^k \frac{(LT)^\ell}{\ell!}, \quad k \in \mathbb{N}, \quad (6.20)$$

**Proof:** By our assumptions (3.4) on the convection field we conclude that

$$|\Phi_x(t, \bar{z}; y) - \Phi_x(t, \bar{z}'; y')| \leq tL \max\{\|\bar{z} - \bar{z}'\|_{L_\infty(I; \mathbb{R}^m)}, |y - y'|\}, \quad (6.21)$$

so that  $|\Phi_x(t, \bar{z}; y) - \Phi_{x'}(t, \bar{z}'; y')| \leq |x - x'| + tL \max\{\|\bar{z} - \bar{z}'\|_{L_\infty(I; \mathbb{R}^m)}, |y - y'|\}$ . One easily verifies inductively that

$$|\Phi_x^k(t, \bar{z}; y) - \Phi_{x'}^k(t, \bar{z}'; y')| \leq \sum_{\nu=0}^k \frac{(t\nu)^k}{\nu!} |x - x'| + \frac{(Lt)^k}{k!} \max\{\|\bar{z} - \bar{z}'\|_{L_\infty(I; \mathbb{R}^m)}, |y - y'|\}. \quad (6.22)$$

Specifically, when  $\bar{z}(s) = \bar{z}(x) = x$  for  $s \in I$ , (6.21) becomes  $|\Phi_x(t, \bar{z}(x); y) - \Phi_x(t, \bar{z}(x'); y')| \leq tL \max\{|x - x'|, |y - y'|\}$  from which the assertion follows.  $\square$

By Stirling's formula one must have  $k_T \sim eLT$  with uniform absolute constants to ensure that

$$\frac{(LT)^{k_T}}{k_T!} \leq \frac{1}{2}, \quad (6.23)$$

say, i.e., one has

$$|\Phi_x^{k_T}(t, z; y) - \Phi_x^{k_T}(t, \bar{z}; y)| \leq \frac{1}{2} \|z - \bar{z}\|_{L_\infty(I; \mathbb{R}^m)}, \quad (6.24)$$

are contractions, as mappings on  $L_\infty(I \times D; \mathbb{R}^m)$ , uniformly in  $(t, x, y)$  with Lipschitz constant  $\frac{1}{2}$ . In what follows we assume that  $k_T = k_T(L, T)$  is minimal with property (6.24).

Moreover, Stirling's formula reveals that

$$\max_{k \geq 1} \frac{(LT)^k}{k!} \sim (LT)^{-1/2} e^{LT}. \quad (6.25)$$

Since (for  $t' < t$ ), keeping (3.3) in mind  $|\Phi_x(t, \bar{z}; y) - \Phi_x(t', \bar{z}; y)| \leq \int_{t'}^t |a(s, \bar{z}(s); y)| ds \leq A|t - t'|$ , we have

$$|\Phi_x^k(t, \bar{z}; y) - \Phi_x^k(t', \bar{z}; y)| \leq \int_{t'}^t |a(s, \Phi_x^{k-1}(s, \bar{z}; y); y)| ds \leq A|t - t'|.$$

These observations can be summarized as follows.

**Remark 6.2** Defining  $k_\varepsilon := |\log_2 \varepsilon| k_T$  and with  $\bar{z}(x)$  from (6.17)

$$Z_\varepsilon(x, t; y) := \Phi_x^{k_\varepsilon}(t, \bar{z}(x); y), \quad (6.26)$$

on has

$$|z(x, t; y) - Z_\varepsilon(x, t; y)| \leq \varepsilon, \quad (x, t, y) \in D \times I \times \mathcal{Y}. \quad (6.27)$$

Moreover, there exists a constant  $\hat{L} < \infty$ , depending only on  $A, L, T$ , such that for all  $(t, x, y), (t', x', y') \in I \times D \times \mathcal{Y}$

$$|Z_\varepsilon(x, t; y) - Z_\varepsilon(x', t'; y')| \leq \hat{L} \max\{|t - t'|, |x - x'|, |y - y'|\}. \quad (6.28)$$

The approximation  $Z_\varepsilon$  to the characteristic field results from applying a *composition of operators*. In a next step operator compositions will be approximated by pointwise compositions of the form (2.1).

## 6.3 Proof of Theorems 4.1 and 4.2

### 6.3.1 Compositional Approximation of $\Phi_x$ by Pointwise Mappings

An obvious way of approximating compositions of integral operators by pointwise compositions is to use quadrature. Since our primary interest is robust approximability in high dimensions we are not interested in high orders, [14]. We are therefore content with the simple midpoint rule which automatically realizes second order when the integrands are  $C^3$ .

We will consistently use the following notation. Consider an equidistant partition of  $I = [0, T]$  with breakpoints  $\tau_i = iT/q$ , for some  $q \in \mathbb{N}$  which will be used to control accuracy. Let  $\xi_i$  denote the respective midpoints of the intervals  $[\tau_{i-1}, \tau_i] =: I_i$ ,  $i = 1, \dots, q$ . The following simple facts will be used frequently.

**Lemma 6.4** Assume that  $g \in \text{Lip}_1(I)$  with Lipschitz constant  $L'$ . Let  $I_i$ ,  $i = 1, \dots, q$  form a uniform partition of  $I$  and let  $\xi_i$  denote the midpoint of  $I_i$ . Denoting further

$$\rho_i(t) := \int_0^t \chi_{I_i}(s) ds, \quad i(t) := \underset{i=1, \dots, q}{\operatorname{argmin}} |t - \xi_i|, \quad (6.29)$$

one has  $\sum_{i=1}^q \rho_i(\xi_k) = \sum_{i=1}^k \rho_i(\xi_k) \leq \frac{|I|k}{q}$ ,  $k \leq q$ , and

$$\left| \int_0^{\xi_{i(t)}} g(s) ds - \sum_{i=1}^q \rho_i(\xi_{i(t)}) g(\xi_i) \right| \leq \frac{T^2 L'}{2q}. \quad (6.30)$$

We approximate now  $\Phi_x$  in a first step by

$$P_{x,q}(t, \bar{z}; y) := x + \sum_{i=1}^{i(t)} \rho_i(\xi_{i(t)}) a(\xi_i, \bar{z}(\xi_i); y), \quad (6.31)$$

which we keep for simplicity piecewise constant in time.

**Lemma 6.5** *Under the assumptions (3.3) and (3.4), one has*

$$\left| \Phi_x(t, \bar{z}; y) - P_{x,q}(t, \bar{z}; y) \right| \leq \frac{AT(LT+1)}{2q}, \quad (6.32)$$

**Proof:** Recall from (3.4) that  $|a(s, \bar{z}(s); y) - a(\xi_i, \bar{z}(\xi_i); y)| \leq L \max\{|s - \xi_i|, |\bar{z}(s) - \bar{z}(\xi_i)|\}$ ,  $s \in I_i$ . If  $\bar{z}$  results from applying  $\Phi_x$ , i.e.,  $\bar{z}(t) = \Phi_x(t, w; y)$  for some  $w$ , we have in view of (3.3)

$$|\bar{z}(t) - \bar{z}(\xi_{i(t)})| = \left| \int_{[t, \xi_{i(t)}]} a(s, w(s), y) ds \right| \leq \frac{TA}{2q}.$$

Hence, on account of Lemma 6.4,

$$\begin{aligned} \left| \Phi_x(t, \bar{z}; y) - P_{x,q}(t, \bar{z}; y) \right| &\leq \sum_{i=1}^{i(t)} \int_{I_i} |a(s, \bar{z}(s), y) - a(\xi_i, \bar{z}(\xi_i); y)| ds + \left| \Phi_x(t, \bar{z}; y) - \Phi_x(\xi_{i(t)}, \bar{z}; y) \right| \\ &\leq \sum_{i=1}^{i(t)} \frac{LT|I_i|}{2q} \max\{1, A\} + \frac{TA}{2q} \leq \frac{LT^2}{2q} \max\{1, A\} + \frac{TA}{2q}, \end{aligned}$$

which, in view of (3.3), confirms the claim.  $\square$

Let  $\bar{\xi} = \bar{\xi}(q) = (\xi_1, \dots, \xi_q)$ ,  $\xi_i = \frac{(i-1/2)T}{q}$ . In what follows we often identify a piecewise constant in time function with the corresponding vector of values

$$\bar{z}(s) = \sum_{i=1}^q \chi_{I_i}(s) z(\xi_i) \quad \leftrightarrow \quad \bar{z}(\bar{\xi}).$$

Note that  $P_{x,q}$  can be written as a composition

$$P_{x,q}(\xi_k, \bar{z}; y) = (g^{2,q} \circ g^{1,q})(x, \bar{z}, \xi_k; y) \quad (6.33)$$

where for  $k = 1, \dots, q$ ,

$$g^{1,q} : (\xi_k, x, \bar{z}; y) \mapsto (k, x, y, a(\xi_1, \bar{z}(\xi_1); y), \dots, a(\xi_q, \bar{z}(\xi_q); y)) \in \mathbb{R}^{(q+1)m+d_y+1}, \quad (6.34)$$

and

$$g^{2,q} : (k, x, w^1, \dots, w^q) \mapsto x + \sum_{i=1}^k \rho_i(\xi_k) w^i \in \mathbb{R}^m. \quad (6.35)$$

We will next exploit that the convection field satisfies  $a \in \text{Lip}_1(I; \mathcal{A}_{\text{Lip}}^{\gamma, s})$ . We know that for each  $t \in I$ ,  $N \in \mathbb{N}$ , there is a composition  $\tilde{A}_{N,t} \in \mathfrak{C}_{N,s}$  such that for each  $i = 1, \dots, q$ ,

$$\begin{aligned} \max_{i=1, \dots, q} |a(\xi_i, \bar{z}, y) - \tilde{A}_{N,i}(\bar{z}, y)| &\leq \gamma(N)^{-1} \|a\|_{L_\infty(I; \mathcal{A}_{\text{Lip}}^{\gamma, s})} \\ \|\tilde{A}_{N,i}\|_{N,s, \text{Lip}} &\leq \|a\|_{L_\infty(I; \mathcal{A}_{\text{Lip}}^{\gamma, s})} =: \|a\|. \end{aligned} \quad (6.36)$$

In a second step we approximate now  $P_{x,q}$  from (6.84) by

$$A_{x,q,N}(t, \bar{z}; y) := x + \sum_{i=1}^{i(t)} \rho_i(\xi_{i(t)}) \tilde{A}_{N,i}(\bar{z}, y), \quad (\tilde{A}_{N,i} := \tilde{A}_{N,\xi_i}) \quad (6.37)$$

to obtain from (6.32) (see also (3.4))

$$\begin{aligned} \left| \Phi_x(t, \bar{z}; y) - A_{x,q,N}(t, \bar{z}; y) \right| &\leq \frac{LT^2}{2q} + \sum_{i=1}^{i(t)} \rho_i(\xi_{i(t)}) |a(\xi_i, \bar{z}, y) - \tilde{A}_{N,i}(\bar{z}, y)| \\ &\leq \frac{(LT+1)AT}{2q} + \frac{T\|a\|}{\gamma(N)}, \end{aligned} \quad (6.38)$$

where we recall the abbreviation  $\|a\| := \|a\|_{L^\infty(I; \mathcal{A}_{\text{Lip}}^{\gamma,s})}$ .

Hence, for any target accuracy  $\eta > 0$ , whenever

$$q = q(\eta) := \left\lceil \frac{(LT+1)AT}{\eta} \right\rceil, \quad N = N(\eta) = \lceil \gamma^{-1}(2T\|a\|/\eta) \rceil, \quad (6.39)$$

we have

$$|\Phi_x(t, \bar{z}; y) - A_{x,q(\eta),N(\eta)}(t, \bar{z}; y)| \leq \eta, \quad (6.40)$$

In what follows we write briefly  $A_{x,\eta} := A_{x,q(\eta),N(\eta)}$  with dimensionality vector  $\mathbf{D}_{x,\eta}$ .

**Remark 6.3** *Suppose that  $\mathbf{D}_{i,q,N}$  is the dimensionality vector of  $\tilde{A}_{N,i}$ . Then, a corresponding realization of  $A_{x,q,N}$  results from parallelization of the  $\tilde{A}_{N,i}$ . Then, one easily checks that one has for the resulting dimensionality vector  $\mathbf{D}_{x,q,N}$*

$$\mathfrak{N}(\mathbf{D}_{x,q,N}) \leq q \max_{i=1,\dots,q} \mathfrak{N}(\mathbf{D}_{i,q,N}) \leq qN. \quad (6.41)$$

Thus, by (6.39), one has for an absolute constant close to one

$$\mathfrak{N}(\mathbf{D}_{x,\eta}) \lesssim q(\eta)\gamma^{-1}(\|a\|2T/\eta) \leq \frac{(LT+1)AT}{\eta}\gamma^{-1}(2T\|a\|/\eta). \quad (6.42)$$

### 6.3.2 Proof of Proposition 2.11

The above considerations have more general ramifications regarding the approximation of primitives which is the subject of Proposition 2.11. Specifically, for  $f \in \text{Lip}_1(I; \mathcal{A}_{\text{Lip}}^{\gamma,s})$  there exists  $L' \leq \|f\|_{L_{\text{Lip}}(I; \mathcal{A}_{\text{Lip}}^{\gamma,s})}$  such that  $|f(t, w) - f(t', w)| \leq L'|t - t'|$ ,  $w \in D_0$ . Moreover, by assumption, there exists for  $N \in \mathbb{N}$  a composition  $f_N(t, \cdot) \in \mathfrak{C}_{N,s}$ ,  $t \in I$ , such that  $\|f(t, \cdot) - f_N(t, \cdot)\|_\infty \leq \gamma(N)^{-1}\|f\|_{L^\infty(I; \mathcal{A}_{\text{Lip}}^{\gamma,s})}$ . Then, for  $q = \gamma(N)$  and  $F_N(t, \cdot) :=$

$\sum_{i=1}^{\gamma(N)} \rho_i(\xi_i(t)) f_N(\xi_i, \cdot)$ , essentially the same arguments as in Lemma 6.4 one can show that for  $t \in I$

$$\begin{aligned} \left| \int_0^t f(s, w) ds - F_N(t, w) \right| &\leq \left| \int_t^{\xi_i(t)} f(s, w) ds \right| + \sum_{i=1}^{i(t)} \int_{I_i \cap [0, \xi_i(t)]} |f(s, w) - f(\xi_i, w)| \\ &\quad + \sum_{i=1}^{i(t)} \int_{I_i \cap [0, \xi_i(t)]} |f(\xi_i, w) - f_N(\xi_i, w)| \lesssim \gamma(N)^{-1} \|f\|_{\text{Lip}_1(I; \mathcal{A}_{\text{Lip}}^{\gamma, s})} \end{aligned}$$

with a constant depending on  $T$ . Since, by parallelization,  $\max_i \|F_N(\xi_i, \cdot)\|_{N\gamma(N), s, \text{Lip}} \leq \max_{t \in I} \|f(t, \cdot)\|_{N, s, \text{Lip}}$  the assertion of Proposition 2.11 follows with  $\phi(N) := N\gamma(N)$ .  $\square$

### 6.3.3 Lipschitz Continuity of Pointwise Mappings

As a final prerequisite we need bounds for the Lipschitz constants of compositions of  $A_{x, q, N}$  which will later be needed when approximating  $A_{x, q, N}$  by DNNs. To that end, suppose that

$$\tilde{A}_{N, i} = (\tilde{A}_{N, i})^{n_i} \circ \cdots \circ (\tilde{A}_{N, i})^1, \quad (6.43)$$

of depth  $n_i$ , where, by definition of  $\mathfrak{C}_{N, s}$ , each component  $(\tilde{A}_{N, i})_\nu^j$ ,  $1 \leq \nu \leq d_j$  depends for  $j < n_i$  only on at most  $s$  variables. Of course, the underlying dimensionality vector  $\mathbf{D}$  depends on  $N$ . We do not know though any exact information on  $\mathbf{D}$  such as widths and depth  $n_i = n_i(N)$ .

To proceed, recall also that the Lipschitz constants of the factors in  $\tilde{A}_{N, i}$  as well as the Lipschitz constants  $L_{[n_i, j]}(\tilde{A}_{N, i})$  of the partial compositions  $(\tilde{A}_{N, i})^{n_i} \circ \cdots \circ (\tilde{A}_{N, i})^j$  are controlled by

$$\max_{1 \leq i \leq q} \|\tilde{A}_{N, i}\|_{N, \mathbb{Y}} \leq \|a\| := \|a\|_{L_\infty(I; \mathcal{A}_{\text{Lip}}^{\gamma, s})}. \quad (6.44)$$

**Lemma 6.6** *For any  $q \in \mathbb{N}$ ,  $k \in \mathbb{N}$ , and  $t = \xi_\ell$  for some  $\ell \leq q$ , one has*

$$|A_{x, q, N}^k(t, \bar{z}; y) - A_{x, q, N}^k(t, \bar{z}'; y')| \leq \frac{(\|a\|T)^k}{k!} \{|y - y'|, \|\bar{z} - \bar{z}'\|_{L_\infty(I; \mathbb{R}^m)}\}, \quad x \in D, \bar{z}, \bar{z}' \in \mathbb{R}^q, \quad (6.45)$$

where in analogy to (6.25)

$$\max_{k \geq 1} \frac{(\|a\|T)^k}{k!} \sim (\|a\|T)^{-1/2} e^{\|a\|T}. \quad (6.46)$$

Similarly, when  $\bar{z} = \bar{z}(x)$ ,  $\bar{z}' = \bar{z}(x')$ , one has for all  $x, x' \in D$ ,  $t = \xi_\ell = (\ell - \frac{1}{2})T/q$

$$|A_{x, q, N}^k(t, \bar{z}(x); y) - A_{x', q, N}^k(t, \bar{z}(x'); y')| \leq \{|y - y'| + |x - x'|\} e^{\|a\|T}. \quad (6.47)$$



**Proof:** The reasoning is analogous to Lemma 6.3. We consider first the case of fixed  $x$ . Using (6.37) and (6.44), one finds for  $t = \xi_\ell$

$$|A_{x,q,N}(t, \bar{z}; y) - A_{x,q,N}(t, \bar{z}'; y')| \leq \sum_{i=1}^{\ell} \rho_i(t) \|a\| \max\{|y - y'|, |\bar{z}(\xi_i) - \bar{z}'(\xi_i)|\}, \quad (6.48)$$

and hence, to see the pattern,

$$\begin{aligned} |A_{x,q,N}^2(t, \bar{z}; y) - A_{x,q,N}^2(t, \bar{z}'; y')| &\leq \sum_{i_1=1}^{\ell} \rho_{i_1}(t) \|a\| |A_{x,q,N}(\bar{z}(\xi_{i_1}), \xi_{i_1}; y) - A_{x,q,N}(\bar{z}(\xi_{i_1}), \xi_{i_1}; y')| \\ &\leq \sum_{i_1=1}^{\ell} \rho_{i_1}(t) \|a\| \left\{ \sum_{i_2=1}^{i_1} \rho_{i_2}(t) \|a\| \max\{|y - y'|, |\bar{z}(\xi_{i_2}) - \bar{z}'(\xi_{i_2})|\} \right\}. \end{aligned}$$

Inductively it follows that for  $t = \xi_\ell$  and  $k \in \mathbb{N}$

$$|A_{x,q,N}^k(t, \bar{z}; y) - A_{x,q,N}^k(t, \bar{z}'; y')| \leq \|a\|^k \sum_{\ell \geq i_1 \geq i_2 \geq \dots \geq i_k \geq 1} \rho_{i_1}(t) \cdots \rho_{i_k}(t) \max\{|y - y'|, |\bar{z}(\xi_{i_k}) - \bar{z}'(\xi_{i_k})|\}.$$

Invoking (6.30), yields

$$\sum_{\ell \geq i_1 \geq i_2 \geq \dots \geq i_k \geq 1} \rho_{i_1}(t) \cdots \rho_{i_k}(t) \leq T^k \sum_{\ell \geq i_1 \geq i_2 \geq \dots \geq i_k \geq 1} \frac{i_1 \cdots i_k}{q^k} \leq \frac{T^k}{k!},$$

providing

$$|A_{x,q,N}^k(t, \bar{z}; y) - A_{x,q,N}^k(t, \bar{z}'; y')| \leq \frac{(\|a\|T)^k}{k!} \max\{|y - y'|, \|\bar{z} - \bar{z}'\|_{L_\infty(I; \mathbb{R}^m)}\}. \quad (6.49)$$

Similarly, for  $\bar{z} = \bar{z}(x)$ ,  $\bar{z}' = \bar{z}(x')$  (see (6.17)), we obtain

$$\begin{aligned} |A_{x,q,N}(t, \bar{z}; y) - A_{x',q,N}(t, \bar{z}'; y')| &\leq |x - x'| + \sum_{i=1}^{\ell} \rho_i(t) \|a\| \max\{|y - y'|, \|\bar{z} - \bar{z}'\|_{L_\infty(I; \mathbb{R}^m)}\} \\ &\leq |x - x'| + \|a\| \sum_{i=1}^{\ell} \rho_i(t) \{|y - y'| + |x - x'|\}, \end{aligned}$$

and hence inductively

$$|A_{x,q,N}^k(t, \bar{z}; y) - A_{x',q,N}^k(t, \bar{z}'; y')| \leq \{|y - y'| + |x - x'|\} \sum_{\nu=0}^k \frac{(\|a\|T)^\nu}{\nu!}, \quad (6.50)$$

which completes the proof.  $\square$

**Remark 6.4** *The same reasoning as in § 6.2.2 applies, see (6.23). From now on we let*

$$k_T := \frac{(\max\{L, \|a\|\}T)^k}{k!} \leq \frac{1}{2}. \quad (6.51)$$

*One still has  $k_T \sim e \max\{L, \|a\|\}T$  and consequently*

$$\max_{k \geq 1} \frac{(\max\{L, \|a\|\}T)^k}{k!} \sim (\max\{L, \|a\|\}T)^{-1/2} e^{\max\{L, \|a\|\}T}. \quad (6.52)$$

*From now on let us assume for simplicity (the natural case) that  $\|a\| \geq L$ .*

### 6.3.4 Approximating Compositions of $\Phi_x$ by Pointwise Compositions

Our approximation of the characteristic field will now proceed in two stages. First we approximate compositions of the operators  $\Phi_x$  by pointwise compositions. The latter ones will be approximated in a second step by DNNs.

To that end, recall that the  $\varepsilon$ -accurate approximation  $Z_\varepsilon$  from (6.27) can be written as

$$\Psi_x^\mu(t, \bar{z}(x); y) = \Phi_x^{\mu k_T}(t, \bar{z}(x); y), \quad \mu = \mu(\varepsilon) = \lceil \log_2 \varepsilon \rceil, \quad (6.53)$$

i.e., we can view this as a  $\mu$ -fold composition of the block-operators  $\Psi_x$  which are  $k_T$  fold compositions of the operator  $\Phi_x$ , where in analogy to (6.18)

$$\Psi_x^\ell(t, \bar{z}(x); y) = \Psi_x(t, \Psi_x^{\ell-1}(t, \bar{z}(x); y); y).$$

Accordingly, we construct a compositional approximation to  $Z_\varepsilon$  by approximating first the blocks  $\Psi_x$  within some target accuracy that will be stipulated later.

The following lemma exploits the favorable Lipschitz constant of the blocks  $\Psi_x$ .

**Lemma 6.7** *In the above terms we have*

$$|\Psi_x(t, \bar{z}; y) - A_{x,\eta}^{k_T}(t, \bar{z}; y)| \leq \eta \sum_{j=1}^{k_T} L_{[k_T, j+1]} \leq \eta e^{LT}, \quad y \in \mathcal{Y}, \quad (6.54)$$

where

$$L_{[k_T, k_T+1]} := 1, \quad L_{[k_T, j+1]} \leq \frac{(LT)^{k_T-j}}{(k_T - j)!}. \quad (6.55)$$

**Proof:** The assertion follows from Remark 2.1 combined with (6.22) and (6.40).  $\square$

We are now ready to approximate  $Z_\varepsilon$  from (6.26) by a pointwise composition. To that end, let

$$C_{\mu,\eta}(t, x; y) := A_{x,\eta}^{\mu k_T}(t, \bar{z}_x; y) = (A_{x,\eta}^{k_T} \circ \dots \circ A_{x,\eta}^{k_T})(t, \bar{z}_x; y), \quad (6.56)$$

where we recall the definition (6.17) of  $\bar{z}(x)$ . The above representation of  $C_{\mu,\eta}$  is obviously  $s$ -dimension sparse.

Next, we need to choose  $\eta$  in (6.54) so as to warrant a given target accuracy  $\varepsilon > 0$ . Remark 2.1 yields on account of (6.23)

$$|Z_\varepsilon(t, x, ; y) - C_{\mu,\eta}(t, x; y)|_\infty \leq \eta e^{LT} \left\{ 1 + \sum_{j=1}^{\mu-1} 2^{-(\mu-j)} \right\} \leq 2\eta e^{LT}, \quad (6.57)$$

where we have exploited the favorable Lipschitz constants  $1/2$  of the factors  $\Psi_x$ , (6.23). Taking now

$$\eta(\varepsilon) := \frac{\varepsilon}{2} e^{-LT}, \quad (6.58)$$

and setting  $C_{\mu(\varepsilon),\eta(\varepsilon)} =: C_\varepsilon$ , yields

$$|z(t, x; y) - C_\varepsilon(t, x; y)| \leq 2\varepsilon, \quad (6.59)$$

while (identifying  $C_\varepsilon$  with its representation), we infer from Remark 6.3, (6.39), and (2.8)

$$\begin{aligned} \mathfrak{N}(C_\varepsilon) &\lesssim LT |\log_2 \varepsilon| (LT + 1) A T e^{LT} \varepsilon^{-1} \gamma^{-1} (4T e^{LT} \|a\| / \varepsilon) \\ &= LT (LT + 1) A T e^{LT} \frac{|\log_2 \varepsilon|}{\varepsilon} \gamma^{-1} \left( \frac{4T e^{LT} \|a\|}{\varepsilon} \right) =: N_\varepsilon, \end{aligned} \quad (6.60)$$

with an absolute proportionality constant. Invoking (6.46) and (6.44), we obtain

$$\|C_\varepsilon\|_{N_\varepsilon, s, \text{Lip}} \lesssim \max_{1 \leq i \leq q(\eta_\varepsilon)} \|A_{N_\varepsilon, i}\|_{N_\varepsilon, s} (\|a\| T)^{-1/2} e^{\|a\| T} \leq \left( \frac{\|a\|}{T} \right)^{\frac{1}{2}} e^{\|a\| T}. \quad (6.61)$$

To finish the proof of Theorem 4.1, it remains to estimate  $N_\varepsilon$  for the two growth function types in (4.1). In the case (alg) one readily checks that  $\gamma^{-1}(\zeta) = \zeta^{\frac{1}{\alpha}}$  with constants depending only on  $\alpha$ . Thus

$$N_\varepsilon \sim \|a\|^{\frac{1}{\alpha}} L T (L T + A) T e^{\frac{L T (1 + \alpha)}{\alpha}} \varepsilon^{-\frac{1 + \alpha}{\alpha}} |\log_2 \varepsilon|^2, \quad (6.62)$$

while in the case (exp), one has

$$N_\varepsilon \sim L T (L T + A) T e^{L T} \frac{|\log_2(4T \|a\| e^{L T} / \varepsilon)|^2}{\varepsilon}. \quad (6.63)$$

It will be convenient to abbreviate

$$Q_{\Lambda, L, T} := L T (L T + 1) A T \lesssim A \begin{cases} L^2 T^3, & T > 1, \\ (L T)^2, & 0 < T \leq 1, \end{cases} \quad (6.64)$$

with a absolute constant depending on  $L$ .

To determine the corresponding convergence rates, we view  $N_\varepsilon$  as a growth function  $\phi(\varepsilon^{-1})$  and wish to determine  $\hat{\gamma}(r) = \phi^{-1}(r)$ . Applying Lemma 6.2 to  $\phi(\zeta) = \|a\|^\frac{1}{\alpha} Q_{A,L,T} e^{\frac{LT(1+\alpha)}{\alpha} \zeta^\frac{1+\alpha}{\alpha}} |\log_2 \zeta|$ ,  $\phi(\zeta) = \alpha^{-1} Q_{A,L,T} e^{LT} \zeta |\log_2 \zeta|^2$ , for the two cases (alg), (exp) in (4.1), respectively, yields

$$\hat{\gamma}(r) \sim \begin{cases} (\|a\|^\frac{1}{\alpha} Q_{A,L,T})^{-\frac{\alpha}{1+\alpha}} e^{-LT} r^\frac{\alpha}{1+\alpha} (\log_2 r)^{-1}, & (r > Q_{A,L,T} e^{LT(1+\alpha)/\alpha}), \quad \text{for (alg),} \\ \alpha (Q_{A,L,T})^{-1} e^{-LT} r \left| \log_2 r^\frac{4\|a\|}{L^2 T^2} \right|^{-2}, & \text{for (exp).} \end{cases} \quad (6.65)$$

Thus, upon collecting the above estimates, we have by (6.59) and (6.61),

$$K_{s,\text{Lip}}(z, \hat{\gamma}(N_\varepsilon)^{-1}) \leq 2\varepsilon + \left( \frac{\|a\|}{T} \right)^\frac{1}{2} e^{\|a\|T} \hat{\gamma}(N_\varepsilon)^{-1} \lesssim \hat{\gamma}(N_\varepsilon)^{-1} \left( 1 + \left( \frac{\|a\|}{T} \right)^\frac{1}{2} e^{\|a\|T} \right).$$

This completes the proof of Theorem 4.1.  $\square$

### 6.3.5 Proof of Theorem 4.2, Implanting DNNs

The final step is to invoke Lemma 2.1 and Lemma 2.1 to approximate each  $\tilde{A}_{N,i}(\bar{z}, y) \in \mathfrak{C}_{N,s}$  in (6.37) by a DNN. Now the role of  $\Psi_x$  is played by  $A_{x,\eta}^{k_T}$  with  $k_T$  from (6.51) so that all previous estimates remain valid and Lemma 6.7 applies, on account of Remark 6.4 in the same way.

For a given tolerance  $\delta > 0$ , let  $\mathcal{N}_{i,\delta,N}(z, y)$  denoted the expanded composition resulting from implanting  $\delta_j$ - accurate neural networks in the  $j$ th composition factor  $(\tilde{A}_{N,i})^j$  of  $\tilde{A}_{N,i}(z, y)$ ,  $j < n_i$  (recalling that the output factor  $(\tilde{A}_{N,i})^{n_i}$  is affine). Invoking Proposition 2.2, let  $(\mathcal{N})_v^j$  denote the neural network implanted at the position of the component  $(\tilde{A}_{N,i})_v^j$ . Moreover, one infers from Lemma 2.1, (2.13), (6.44), that

$$\|\tilde{A}_{N,i} - \mathcal{N}_{i,\delta,N}\|_{L_\infty(D \times I \times \mathcal{Y}; \mathbb{R}^m)} \leq \delta_{n_i} + \sum_{r=1}^{n_i-1} \delta_r L_{[n_i, r+1]}(\tilde{A}_{N,i}) \leq \delta_{n_i} + \|a\| \sum_{r=1}^{n_i-1} \delta_r. \quad (6.66)$$

Taking  $\delta_r := \frac{\delta}{n_i}$ ,  $r = 1, \dots, n_i$ , we have (assuming  $\|a\| \geq 1$ )

$$\|\tilde{A}_{N,i} - \mathcal{N}_{i,\delta,N}\|_{L_\infty(D \times I \times \mathcal{Y}; \mathbb{R}^m)} \leq \|a\| \delta. \quad (6.67)$$

On the other hand, we invoke (2.14) in Lemma 2.1 to conclude

$$\#\mathcal{N}_{i,\delta,N} \leq C_s \|a\|^s N n_i^s \delta^{-s} \left| \log_2 \delta \right| + \log_2 n_i \quad (6.68)$$

where the constant  $C_s$  depends only on  $s$ .

Now we define in analogy to (6.37)

$$\mathcal{N}_{x,q,N,\delta}(z, t; y) := x + \sum_{i=1}^q \rho_i(t) \mathcal{N}_{i,\delta,N}(z, y), \quad t \in \bar{\xi}, \quad (6.69)$$

We need to stipulate the choice of  $\delta, q, N$  so as to ensure that  $\mathcal{N}_{x,\eta} := \mathcal{N}_{x,q(\eta/2),N(\eta/2),\delta(\eta)}$ , with  $q(\eta/2), N(\eta/2)$ , defined according to (6.39). This provides

$$\left| \Phi_x(z, t; y) - \mathcal{N}_{x,\eta}(z, t; y) \right| \leq \eta, \quad (6.70)$$

when

$$\delta = \delta(\eta) := \frac{\eta}{2\|a\|}. \quad (6.71)$$

To estimate the complexity of  $\mathcal{N}_{x,\eta}$  we use (6.68) and bound the depths  $n_i$  by  $N \sim \gamma^{-1}(4T\|a\|/\eta)$ . Arguing as in (6.41) and recalling (6.39), we then have

$$\begin{aligned} \#\mathcal{N}_{x,\eta} &\leq q(\eta/2) \max_{i=1,\dots,q(\eta/2)} \#\mathcal{N}_{i,\delta(\eta/2),N(\eta/2)} \\ &\leq q(\eta/2) \|a\|^s \gamma^{-1}(4T\|a\|/\eta)^{1+s} \delta(\eta)^{-s} |\log_2 \delta(\eta) + \log_2 N(\eta/2)| \\ &\leq C_2 C_s \|a\|^{2s} T(LT + A) \eta^{-(1+s)} \gamma^{-1}(4T\|a\|/\eta)^{1+s} \\ &\quad \times \left\{ \left| \log_2 \frac{\eta}{2\|a\|} \right| + \left| \log_2 \left( \gamma^{-1}(4T\|a\|/\eta) \right) \right| \right\}. \end{aligned} \quad (6.72)$$

We consider first the case of algebraic growth  $\gamma(r) = C_\alpha r^\alpha$  for which

$$\gamma^{-1}(4T\|a\|/\eta) = C_\alpha^{-1/\alpha} (4T\|a\|)^{1/\alpha} \eta^{-1/\alpha}, \quad \log_2 \gamma^{-1}(4T\|a\|/\eta) \sim \log_2(4T\|a\|/\eta).$$

Absorbing for convenience  $C_\alpha$  and the logarithmic dependence on  $T$  and  $\|a\|$  in  $C_2$ , we obtain

$$\begin{aligned} \#\mathcal{N}_{x,\eta} &\leq C_2 C_s \|a\|^{\frac{(2\alpha+1)s+1}{\alpha}} AT(LT + 1) T^{\frac{1+s}{\alpha}} \eta^{-\frac{(1+\alpha)(1+s)}{\alpha}} |\log_2(T\|a\|/\eta)| \\ &= b_1 \|a\|^{\frac{(2\alpha+1)s+1}{\alpha}} \eta^{-\zeta} |\log_2(b_2/\eta)|, \end{aligned} \quad (6.73)$$

where

$$b_1 := C_2 C_s AT(LT + 1) T^{\frac{1+s}{\alpha}}, \quad b_2 := T\|a\|, \quad \zeta := \frac{(1+\alpha)(1+s)}{\alpha}. \quad (6.74)$$

We can now define  $\mathcal{N}_{\mu,\eta}$  in analogy to (6.56) but with  $A_{x,\eta}^{k_T}$  replaced by  $\mathcal{N}_{x,\eta}^{k_T}$ . The choice of  $\eta$  is based on the following lemma.

**Lemma 6.8** *Taking*

$$\eta = \eta(\varepsilon) := \frac{\varepsilon}{2} e^{-\|a\|T}, \quad \mu = \mu(\varepsilon) = \lceil \log_2 \varepsilon \rceil, \quad (6.75)$$

and abbreviating  $\mathcal{N}_\varepsilon := \mathcal{N}_{\mu(\varepsilon),\eta(\varepsilon)}$ , one has

$$|Z_\varepsilon(x, t, y) - \mathcal{N}_\varepsilon(x, t, y)| \leq \varepsilon, \quad t \in I, y \in \mathcal{Y}, \quad (6.76)$$

and  $q(\eta(\varepsilon)) \sim LT^2 e^{\|a\|T} \varepsilon^{-1}$ .

**Proof:** By Remark 6.4, (6.77) remains valid when  $\Psi_x$  is replaced by  $A_{x,\eta}^{k_T}$ . Remark 2.1 yields on account of (6.23)

$$|Z_\varepsilon(x, t, y) - \mathcal{N}_{\mu,\eta}(x, t, y)|_\infty \leq \eta e^{\|a\|T} \left\{ 1 + \sum_{j=1}^{\mu-1} 2^{-(\mu-j)} \right\} \leq 2\eta e^{\|a\|T}, \quad (6.77)$$

where we have exploited again the favorable Lipschitz constants  $1/2$  of the factors  $A_{x,\eta}^{k_T}$ , (6.51). The estimate (6.76) follows now from (6.75).  $\square$

Thus, it remains to bound next  $\#\mathcal{N}_\varepsilon$ . Arguing as before, since  $T \geq T_0$ , and  $k_T \gtrsim T$ , one obtains from (6.73)

$$\begin{aligned} \#\mathcal{N}_\varepsilon &\sim \mu(\varepsilon)k_T \#\mathcal{N}_{x,\eta(\varepsilon)} \sim |\log_2 \varepsilon| \|a\|T b_1 \|a\|^{\frac{(2\alpha+1)s+1}{\alpha}} \eta(\varepsilon)^{-\zeta} |\log_2 T \|a\|/\eta(\varepsilon)| \\ &\sim |\log_2 \varepsilon| T b_1 \|a\|^{\frac{(2\alpha+1)s+1+\alpha}{\alpha}} \varepsilon^{-\zeta} e^{\|a\|T\zeta} \left| \log_2 2 \|a\|T e^{\|a\|T}/\varepsilon \right|, \quad \zeta = \frac{(1+\alpha)(1+s)}{\alpha}, \end{aligned}$$

In summary we obtain

$$\#\mathcal{N}_\varepsilon \sim \tilde{b}_1 \|a\|^\tau \varepsilon^{-\zeta} |\log_2 \tilde{b}_2/\varepsilon|^2, \quad \tilde{b}_1 := C_s L T^{\frac{3\alpha+s+1}{\alpha}} e^{\|a\|T\zeta}, \quad \tilde{b}_2 := 2T \|a\| e^{\|a\|T}, \quad (6.78)$$

where

$$(\zeta, \tau) = \left( \frac{(1+\alpha)(1+s)}{\alpha}, s + \frac{(\alpha+1)(s+1)}{\alpha} \right), \quad (6.79)$$

which confirms the first relation in (4.6).

When  $\gamma(r) \sim e^{\alpha r}$ , we have  $\gamma^{-1}(s) = \frac{1}{\alpha} \ln s$ , so that by (6.72),

$$\begin{aligned} \#\mathcal{N}_{x,\eta} &\lesssim \alpha^{-(1+s)} \|a\|^{2s} L T^2 \eta^{-(1+s)} (\log_2(2T \|a\|/\eta))^{1+s} \left| \log_2 \frac{\eta}{2 \|a\|} \right| \\ &\sim \alpha^{-(1+s)} \|a\|^{2s} L T^2 \eta^{-(1+s)} (\log_2(\|a\|/\eta))^{2+s}, \end{aligned} \quad (6.80)$$

Thus, by (6.75)

$$\begin{aligned} \#\mathcal{N}_\varepsilon &\sim \alpha^{-(1+s)} \|a\|^{2s} L T^3 |\log_2 \varepsilon| e^{\|a\|T(1+s)} \varepsilon^{-(1+s)} \log_2(4T e^{\|a\|T} \|a\|/\varepsilon)^{2+s} \\ &\sim \alpha^{-(1+s)} \|a\|^{2s} L T^3 e^{\|a\|T(1+s)} \varepsilon^{-(1+s)} \log_2(4T e^{\|a\|T} \|a\|/\varepsilon)^{3+s}, \quad \varepsilon \leq \varepsilon_0. \end{aligned} \quad (6.81)$$

This confirms the second relation in (4.6) and finishes the proof of Theorem 4.2.  $\square$

We now turn to establishing the convergence rates stated in Corollary 4.3. To begin with the case of algebraic growth, we need to “invert” the relation (6.78). Arguing exactly as in the previous section, we consider the growth function  $\phi(s) = \tilde{b}_1 s^\zeta |\log_2 \tilde{b}_2 s|^2$  with  $\tilde{b}_1, \tilde{b}_2$  from (6.78), and need to determine  $\tilde{\gamma}(r) = \phi^{-1}(r)$ . Invoking Lemma 6.2, yields

$$\tilde{\gamma}(r) \sim \phi^{-1}(r) \sim \tilde{b}_1^{-1/\zeta} \|a\|^{-(s+\zeta)/\zeta} \zeta^{\zeta(1+s)/\zeta} r^{1/\zeta} \left| \log_2 \left( r \tilde{b}_2^\zeta / \tilde{b}_1 \right) \right|^{-\frac{2}{\zeta}}.$$

For  $\zeta = \frac{(1+\alpha)(1+s)}{\alpha}$ , this remains uniformly proportional we obtain

$$\tilde{\gamma}(r) \sim B e^{-\|a\|T} \|a\|^{-\left(1 + \frac{s\alpha}{(1+s)(1+\alpha)}\right)} r^{\frac{\alpha}{(1+\alpha)(1+s)}} \left| \log_2 r \right|^{-\frac{2\alpha}{(1+\alpha)(1+s)}}, \quad (6.82)$$

with a constant scaling like

$$B \sim C_s L^{-\frac{\alpha}{(1+\alpha)(1+s)}} T^{-\frac{3\alpha+1+s}{(1+\alpha)(1+s)}}. \quad (6.83)$$

Regarding exponential growth, applying the same reasoning to (6.81), Lemma 6.2 yields with

$$\phi(s) = b_1 s^\zeta |\log_2 s|^{3+s}, \quad b_1 := \alpha^{-\zeta} \|a\|^{s+\zeta} L T^3 e^{\|a\|T\zeta}, \quad b_2 = 4T e^{\|a\|T} \|a\|, \quad \zeta = 1 + s$$

the searched for growth function reads

$$\begin{aligned} \tilde{\gamma}(r) = \phi^{-1}(r) &\sim \alpha \|a\|^{-\left(1 + \frac{s}{s+1}\right)} (L T^3)^{-\frac{1}{1+s}} e^{\|a\|T} r^{\frac{1}{1+s}} \left| \log_2 r \right|^{-\frac{3+s}{1+s}} \\ &\sim \alpha (L T^3 \|a\|^{2s+1})^{-\frac{1}{1+s}} e^{-\|a\|T} r^{\frac{1}{1+s}} \left| \log_2 r \right|^{-\frac{3+s}{1+s}}. \end{aligned}$$

This finishes the proof of Corollary 4.3.  $\square$

## 6.4 Proof of Theorems 4.4, 4.4

Under the assumption (3.7), the fixed-point map  $\Phi_x$  is now approximated in a first step by

$$P_{x,q}(t, \bar{z}; y) := x + \sum_{j=1}^{d_y} y_j \sum_{i=1}^{i(t)} \rho_i(\xi_{i(t)}) a_j(\xi_i, \bar{z}(\xi_i)), \quad (6.84)$$

whose accuracy is given by (6.32). To that end, we approximate the low-dimensional functions  $a_j(t, \cdot)$  by finitely parametrized functions such as neural networks invoking Proposition 2.2. Specifically, there exist networks  $\mathcal{N}_{j,i,\delta}$  of depth  $\lesssim \log_2 \delta^{-1}$  such that

$$\|a_j(\xi_i, \cdot) - \mathcal{N}_{j,i,\delta}\|_{L^\infty(D)} \leq \delta, \quad \#\mathcal{N}_{j,i,\delta} \lesssim \delta^{-m} |\log_2 \delta|, \quad j = 1, \dots, d_y, \quad i = 1, \dots, q, \quad (6.85)$$

Setting

$$\mathcal{N}_{x,q,\delta}(t, \bar{z}; y) := x + \sum_{j=1}^{d_y} y_j \sum_{i=1}^{i(t)} \rho_i(\xi_{i(t)}) \mathcal{N}_{j,i,\delta}(\bar{z}(\xi_i)), \quad (6.86)$$

this yields, in view of (6.32) and (6.85),

$$|\Phi_x(t, \bar{z}; y) - \mathcal{N}_{x,q,\delta}(t, \bar{z}; y)| \leq \frac{AT(LT + 1)}{2q} + d_y T \delta. \quad (6.87)$$

Thus, given any target tolerance  $\eta > 0$ , choosing

$$q(\eta) = \left\lceil \frac{TA(LT + 1)}{\eta} \right\rceil, \quad \delta(\eta) = \frac{\eta}{2T d_y}, \quad (6.88)$$

and abbreviating  $\mathcal{N}_{x,\eta} := \mathcal{N}_{x,q(\eta),\delta(\eta)}$ , we indeed obtain

$$|\Phi_x(t, \bar{z}, y) - \mathcal{N}_{x,\eta}(t, \bar{z}; y)| \leq \eta, \quad x \in D, \bar{z} \in \Omega, t \leq T. \quad (6.89)$$

Lets see next that  $\mathcal{N}_{x,\eta}$  is indeed an  $m$ -dimension-sparse neural network. To that end, the input-variables  $t, x, y$  need to be carried across layers, formally in a ‘‘skip-connection’’ format. Thus, formally we have

$$\mathcal{N}_{x,\eta}(t, \bar{z}, y) = (G_2 \circ G_1)(t, x, \bar{z}, y) \quad (6.90)$$

where for clarity the following representation groups variables in a formally incorrect way and should be viewed as a mapping into  $\mathbb{R}^{1+m+d_y+m q(\eta)d_y}$

$$G_1 : (t, x, y) \mapsto \begin{pmatrix} t, x, y, \\ \mathcal{N}_{1,1,\delta(\eta)}, \dots, \mathcal{N}_{d_y,1,\delta(\eta)}, \\ \vdots \\ \mathcal{N}_{1,q(\eta),\delta(\eta)}, \dots, \mathcal{N}_{d_y,q(\eta),\delta(\eta)} \end{pmatrix} \in \mathbb{R}^{1+m+d_y+m q(\eta)d_y}. \quad (6.91)$$

Hence  $G_1$  itself is a neural network whose depth is bounded by  $\log_2 \delta^{-1}$ . The components  $t, x, y$  are treated as skip-connections carried to the next layer. More precisely  $t$  is treated as a parameter.

The factor  $G_2$

$$G_2 : (t, x, y, \zeta^{1,1}, \dots, \zeta^{d_y,q(\eta)}) \mapsto \left( t, x, y, x + \sum_{i=1}^{i(t)} \rho(\xi_{i(t)}) \sum_{j=1}^{d_y} y_j \zeta^{j,i} \right) \in \mathbb{R}^{1+2m+d_y}, \quad (6.92)$$

just represents the affine output layer.

Hence, for each  $t \in I$

$$\begin{aligned} \#\mathcal{N}_{x,\eta}(t, \cdot, \cdot) &\lesssim m d_y q(\eta) \delta(\eta)^{-m} |\log_2 \delta(\eta)| \lesssim m d_y A T (L T + 1) \eta^{-(1+m)} (2 T d_y)^m \left| \log_2 \left( \frac{2 T d_y}{\eta} \right) \right| \\ &\sim m A (L T + 1) T^{m+1} \left( \frac{d_y}{\eta} \right)^{1+m} \left| \log_2 \left( \frac{d_y}{\eta} \right) \right|, \end{aligned} \quad (6.93)$$

with an absolute proportionality factor.

**Remark 6.5** *The mappings  $\mathcal{N}_{x,\eta}(t, \cdot, \cdot)$  should really be viewed as piecewise constants in time  $t \in I$  and the above relations represent the values  $\mathcal{N}_{x,\eta}(\xi_k, \cdot, \cdot)$ ,  $k = 0, \dots, q(\eta)$ , where formally  $\mathcal{N}_{x,\eta}(\xi_0) := x$ .*

Recall from (6.25) that  $Z_\varepsilon(x, t; y) = \Phi_x^{k_T \mu(\varepsilon)}(\bar{z}_x, t; y)$ , with  $\bar{z}_x(s) = x$ ,  $s \in I$ , is an  $\varepsilon$ -accurate approximation to the characteristics  $z(x, t; y)$ ,  $t \in I$ , where  $\mu = \mu(\varepsilon)$  as in (6.53). We define in analogy to (6.56)

$$\mathcal{N}_{\mu,\eta}(t, x; y) := \mathcal{N}_{x,\eta}^{\mu k_T}(t, \bar{z}_x; y). \quad (6.94)$$



In view of (6.89), the same reasoning as earlier (just using Remark 2.1 and the fact that the Lipschitz constants of  $\Phi_x^{kT}$  are at most  $1/2$ ) ensures the validity of (6.77) in the present situation as well, so that

$$\|Z_\varepsilon - \mathcal{N}_{\mu,\eta}\|_{L_\infty(I \times D \times \mathcal{Y}; \mathbb{R}^m)} \leq 2\eta e^{LT}. \quad (6.95)$$

Stipulating again

$$\eta = \eta(\varepsilon) := \frac{\varepsilon}{2} e^{-LT}, \quad (6.96)$$

since these bounds are uniform in  $x \in D$ , we obtain indeed for  $\mathcal{N}_\varepsilon := \mathcal{N}_{\mu(\varepsilon),\eta(\varepsilon)}$

$$\|z - \mathcal{N}_\varepsilon\|_{L_\infty(I \times D \times \mathcal{Y}; \mathbb{R}^m)} \leq 2\varepsilon. \quad (6.97)$$

Recall from Remark 6.2 and (6.23) that the total number of iterations of  $\Phi_x$  is  $k_T |\log_2 \varepsilon| \sim \max\{1, T\} |\log_2 \varepsilon|$  with a constant depending on  $L$ . On account of (6.93) and (6.96), one obtains

$$\begin{aligned} \#\mathcal{N}_\varepsilon &\sim \max\{1, T\} |\log_2 \varepsilon| m A (LT + 1) T^{m+1} \left( \frac{d_y}{\eta(\varepsilon)} \right)^{1+m} \left| \log_2 \left( \frac{d_y}{\eta(\varepsilon)} \right) \right| \\ &\lesssim (\max\{1, T\})^2 T^{m+1} e^{LT(m+1)} \left( \frac{d_y}{\varepsilon} \right)^{m+1} \left| \log_2 \frac{d_y}{\varepsilon} \right|^2, \end{aligned} \quad (6.98)$$

with a constant depending on  $L, A, m$ . This confirms (4.10) and the proof of Theorem 4.4.  $\square$

To express this in terms of a convergence, we apply as before Lemma 6.2, we find that the corresponding growth function  $\hat{\gamma}$  scales like

$$\hat{\gamma}(s) \sim d_y^{-1} e^{-LT} (LT^{m+2})^{-\frac{1}{m+1}} s^{\frac{1}{m+1}} \left| \log_2 s \right|^{-\frac{2}{m+1}} \quad (6.99)$$

which can be restated as (4.11).

Note that in agreement with the previous section  $d_y \sim \|a\|$ . To prove the remaining claim in Theorem 4.4, we need to estimate  $\|\mathcal{N}_\varepsilon\|_{N_\varepsilon, m, \text{Lip}}$  where  $N_\varepsilon := \#\mathcal{N}_\varepsilon$ .

**Lemma 6.9** *The network approximations  $\mathcal{N}_{\cdot, q, \delta}(\xi_i, \cdot, \cdot; \cdot)$  from (6.86) have the following Lipschitz continuity properties: for  $(\xi_i, x, y), (\xi_{i'}, x', y') \in I \times D \times \mathcal{Y}$ ,*

$$|\mathcal{N}_{x, q, \delta}^\ell(\xi_k, \bar{z}; y) - \mathcal{N}_{x, q, \delta}^\ell(\xi_k, \bar{z}'; y')| \leq \frac{(KT)^\ell}{\ell!} \{|y - y'|, \|\bar{z} - \bar{z}'\|_{L_\infty(I; \mathbb{R}^m)}\}, \quad x \in D, \bar{z}, \bar{z}' \in \mathbb{R}^q. \quad (6.100)$$

Similarly, when  $\bar{z} = \bar{z}(x), \bar{z}' = \bar{z}(x')$ , one has for all  $x, x' \in D, t = \xi_\ell = (\ell - \frac{1}{2})T/q$

$$|\mathcal{N}_{x, q, \delta}^\ell(\bar{z}(x), \xi_k; y) - \mathcal{N}_{x', q, \delta}^\ell(\bar{z}(x'), \xi_k; y')| \leq \{|y - y'| + |x - x'|\} e^{KT}, \quad (6.101)$$

where  $K$  depends on  $A, L, T$ .

**Proof:** Recall from (6.86) that  $\mathcal{N}_{x,q,\delta}(\xi_k, \bar{z}; y) := x + \sum_{i=1}^k \rho_i(\xi_k) \sum_{j=1}^{d_y} y_j \mathcal{N}_{j,i,\delta}(\bar{z}(\xi_i))$ . Here it matters that we have employed Lipschitz stable DNNs to approximate the  $a_j(\cdot, \cdot)$  (Proposition 2.2), so that

$$\sum_{j=1}^{d_y} \|\mathcal{N}_{j,i,\delta}\|_\infty \leq \bar{\Lambda}, \quad |\mathcal{N}_{j,i,\delta}(\bar{z}) - \mathcal{N}_{j,i,\delta}(\bar{z}')| \leq \bar{L} \|\bar{z} - \bar{z}'\|_{L_\infty(I; \mathbb{R}^m)}, \quad \bar{z}, \bar{z}' \in L_\infty(I; \mathbb{R}^m), \quad (6.102)$$

holds, by Proposition 2.2, for some  $\bar{L} \lesssim L$ , uniformly in  $i = 1, \dots, d_y$ ,  $j = 1, \dots, d_y$ . Then

$$\begin{aligned} \left| \mathcal{N}_{x,q,\delta}(\xi_k, \bar{z}; y) - \mathcal{N}_{x,q,\delta}(\xi_k, \bar{z}'; y') \right| &\leq \sum_{i=1}^k \rho_{I_i}(\xi_k) \sum_{j=1}^{d_y} \left| y_j \mathcal{N}_{j,i,\delta}(\bar{z}(\xi_i)) - y'_j \mathcal{N}_{j,i,\delta}(\bar{z}'(\xi_i)) \right| \\ &\leq \sum_{i=1}^k \rho_i(\xi_k) \left\{ \sum_{j=1}^{d_y} |y_j - y'_j| |\mathcal{N}_{j,i,\delta}(\bar{z}(\xi_i))| + \sum_{j=1}^{d_y} |\mathcal{N}_{j,i,\delta}(\bar{z}(\xi_i)) - \mathcal{N}_{j,i,\delta}(\bar{z}'(\xi_i))| \right\} \\ &\leq \sum_{i=1}^k \rho_i(\xi_k) \left\{ \bar{\Lambda} |y - y'| + \bar{L} \|\bar{z} - \bar{z}'\|_{L_\infty(I; \mathbb{R}^m)} \right\} \\ &\leq \sum_{i=1}^k \rho_i(\xi_k) K \max\{|y - y'|, \|\bar{z} - \bar{z}'\|_{L_\infty(I; \mathbb{R}^m)}\}, \end{aligned} \quad (6.103)$$

where  $K := \bar{\Lambda} + \bar{L}$ . Hence, we are in the same situation as in (6.48). Therefore, the claim follows by the same arguments as used in the proof of Lemma 6.6.  $\square$

From this result and (6.52) we conclude that indeed  $\|\mathcal{N}_\varepsilon\|_{N_\varepsilon, m, \text{Lip}} \leq e^{KT}$ . For  $\gamma(r) = d_y^{-1} r^1 m + 1 |\log_2 r|^{-\frac{2}{m+1}}$ , we have

$$\gamma(N_\varepsilon) K_{m, \text{Lip}}(z, N, \gamma(N)^{-1}) \leq \gamma(N_\varepsilon) \|z - \mathcal{N}_\varepsilon\|_\infty + \|\mathcal{N}_\varepsilon\|_{N_\varepsilon, m, \text{Lip}} \leq e^{LT} + e^{KT},$$

which confirms (4.12).  $\square$

We turn next to the proof of Theorem 4.5. All the preceding results on parameter dependent characteristics hinge on the same strategy of generating first on a continuous level compositional approximations with the aid of *fixed point iterations*. We have emphasized this because it applies to a larger scope of problems. We now exploit that the characteristics have a natural semi-group compositional structure (3.11). We briefly exemplify next a simple strategy how to combine both structural elements for the case of *affine* parameter-representations of convection fields. The arguments carry over to the more general scenarios as well.

To that end, the role of the time horizon  $T$  is now played by an “intermediate” fixed time-horizon  $T = T^\circ$  which we let depend on  $d_y$ , namely

$$T^\circ d_y = 1. \quad (6.104)$$

We wish to construct a DNN approximation to  $z(T, x; y)$  on  $D \times \mathcal{Y}$  for a fixed time horizon  $T = pT^\circ$  and consider the intermediate time-levels  $T_j := jT^\circ$ ,  $j = 0, \dots, p$ . We will now introduce a compositional notion for characteristics, viewing them as *propagation operators*.

$$z_{j+1,k}(w; y) = z(T_{j+1}, T_k; w; y), \quad k \leq j, \quad (6.105)$$

where we have used (3.11) so that

$$z_{j+1,0}(x; y) = (z_{j+1,j} \circ z_{j,0})(x; y) = (z_{j+1,j} \circ \dots \circ z_{1,0})(x; y) = z(T_{j+1}, x; y). \quad (6.106)$$

Analogously, we denote by

$$\mathcal{N}_{j+1,j,\eta}(w; y) = \mathcal{N}_\eta(T_{j+1}, w; y), \quad \mathcal{N}_\eta(T_j, w; y) = w,$$

the neural network constructed in the proof but now for the fixed small time-interval  $[T_j, T_{j+1}]$  of length  $T^\circ$  with target accuracy  $\eta > 0$ . By Theorem 4.4, we have, on account of (6.104)

$$\|z_{j+1,j}(\cdot; \cdot) - \mathcal{N}_{j+1,j}(\cdot; \cdot)\|_{L_\infty(\mathbb{R}^m \times \mathcal{Y})} \leq \eta, \quad \#\mathcal{N}_{j+1,j} \sim e^{LT^\circ(m+1)} \eta^{-(m+1)} \log_2 \left(\frac{1}{\eta}\right)^2, \quad (6.107)$$

with no dependence on  $d_y$ . Defining in analogy to (6.105)

$$(\mathcal{N}_{J+1,j,\eta} \circ \mathcal{N}_{j,j-1,\eta})(w; y) = \mathcal{N}_\eta(T_{j+1}, \mathcal{N}_{j,j-1,\eta}(w; y); y) = \mathcal{N}_\eta(T_{j+1}; \mathcal{N}_{j,j-1}(w; y); y),$$

one has

$$\begin{aligned} & \|z_{j+1,j} \circ z_{j,j-1} - \mathcal{N}_{J+1,j,\eta} \circ \mathcal{N}_{j,j-1,\eta}\|_{L_\infty(\mathbb{R}^m \times \mathcal{Y})} \leq \|z_{j+1,j} \circ z_{j,j-1} - z_{J+1,j} \circ \mathcal{N}_{j,j-1,\eta}\|_{L_\infty(\mathbb{R}^m \times \mathcal{Y})} \\ & \quad \|z_{J+1,j} \circ \mathcal{N}_{j,j-1,\eta} - \mathcal{N}_{J+1,j,\eta} \circ \mathcal{N}_{j,j-1,\eta}\|_{L_\infty(\mathbb{R}^m \times \mathcal{Y})} \\ & \leq L_{j+1,j} \|z_{j,j-1} - \mathcal{N}_{j,j-1,\eta}\|_{L_\infty(\mathbb{R}^m \times \mathcal{Y})} + \|z_{j+1,j} - \mathcal{N}_{J+1,j,\eta}\|_{L_\infty(\mathbb{R}^m \times \mathcal{Y})}, \end{aligned}$$

where, in view of, (6.14),  $L_{j+1,j} \leq e^{L|T_{j+1}-T_j|} = e^{LT^\circ}$ . Hence, Remark 2.1 applies and yields for  $\mathcal{N}_\eta(T, x; y) := (\mathcal{N}_{p,p-1,\eta} \circ \dots \circ \mathcal{N}_{1,0,\eta})(x; y)$

$$\begin{aligned} |z(T, x; y) - \mathcal{N}_\eta(T, x; y)| & \leq \eta \sum_{j=1}^p e^{L(p-j)T} = \eta \sum_{j=0}^{p-1} e^{LTj} = \eta \frac{e^{pLd_y^{-1}} - 1}{e^{Ld_y^{-1}} - 1} \\ & \leq \eta \frac{d_y}{L} e^{pLd_y^{-1}} = \eta L^{-1} e^{LT}. \end{aligned} \quad (6.108)$$

while

$$\#\mathcal{N}_\eta(T, \cdot; \cdot) \lesssim p e^{Ld_y^{-1}(m+1)} \eta^{-(m+1)} |\log_2 n\eta|^2 = T d_y e^{Ld_y^{-1}(m+1)} \eta^{-(m+1)} |\log_2 \eta|^2. \quad (6.109)$$

Thus, taking  $\eta(\varepsilon) = L e^{-LT} \varepsilon$ , we have

$$\#\mathcal{N}_\eta(\varepsilon)(T, \cdot; \cdot) \lesssim T d_y L^{-(m+1)} e^{L(T+d_y^{-1})(m+1)} \varepsilon^{-(m+1)} \left(\log_2 \frac{e^{LT}}{\varepsilon}\right)^2.$$

We deduce from Lemma 6.2 that this can be restated in terms of a convergence rate with growth function

$$\tilde{\gamma}(r) \sim (T d_y)^{\frac{1}{m+1}} L e^{-L(T+d_y^{-1})} r^{\frac{1}{m+1}} \left| \log_2 \frac{r L^{m+1}}{T d_y e^{Ld_y^{-1}(m+1)}} \right|^{-\frac{2}{m+1}}. \quad (6.110)$$

This finishes the proof of Theorem 4.5.  $\square$

## 6.5 Proof of Theorem 4.8

By assumption,  $f_\varepsilon \in \text{Lip}_1(I; \mathfrak{C}_{N_\varepsilon(f), m})$ , so that given  $\varepsilon > 0$ , we can find a composition  $u_{0,\varepsilon} \in \mathfrak{C}_{N_\varepsilon(u_0), m}$ , (identifying for notational convenience in what follows mappings and representations) with

$$\|u_0 - u_{0,\varepsilon}\|_{L_\infty(D)} \leq \varepsilon, \quad \mathfrak{N}(u_{0,\varepsilon}) \leq \gamma_0^{-1} (\|u_0\|_{\mathcal{A}_{\text{Lip}}^{\gamma_0, m}} / \varepsilon) \sim \|u_0\|_{\mathcal{A}_{\text{Lip}}^{\gamma_0, m}}^{1/\alpha_0} \varepsilon^{-1/\alpha_0}, \quad \varepsilon > 0, \quad (6.111)$$

and likewise

$$\|f - f_\varepsilon\|_{L_\infty(I; L_\infty(D))} \leq \varepsilon, \quad \mathfrak{N}(f_\varepsilon) \leq \gamma_f^{-1} (\|f\|_{\mathcal{A}_{\text{Lip}}^{\gamma_f, m}} / \varepsilon) \sim \|f\|_{\mathcal{A}_{\text{Lip}}^{\gamma_f, m}}^{1/\alpha_f} \varepsilon^{-\frac{1}{\alpha_f}}, \quad \varepsilon > 0. \quad (6.112)$$

Next, we use that the compositional factors in  $u_{0,\varepsilon}, f_\varepsilon$  are Lipschitz continuous with constants controlled by

$$\|u_0\| := \|u_0\|_{\mathcal{A}_{\text{Lip}}^{\gamma_0, m}}, \quad \|f\| := \|f\|_{L_\infty(I; \mathcal{A}_{\text{Lip}}^{\gamma_f, m})},$$

respectively. Implanting  $\eta$ -accurate Lipschitz controlled DNNs, on account of Lemma 2.1, to obtain

$$\|u_{0,\varepsilon} - \mathcal{N}_{u_{0,\varepsilon}, \eta}\|_{L_\infty(D)} \leq \eta \left\{ 1 + \sum_{j=1}^{n(\mathbf{D}(u_{0,\varepsilon})) - 1} \|u_0\| \right\} \leq \eta \mathfrak{N}(u_{0,\varepsilon}) \|u_0\|, \quad (6.113)$$

as well as

$$\|f_\varepsilon(t, \cdot) - \mathcal{N}_{t, f_\varepsilon, \eta}\|_{L_\infty(D \times \mathcal{Y})} \leq \eta \left\{ 1 + \sum_{j=1}^{n(\mathbf{D}(f_\varepsilon)) - 1} \|f_0\| \right\} \leq \eta \mathfrak{N}(f_\varepsilon) \|f_0\|, \quad t \in I, \quad (6.114)$$

while, by Proposition 2.2, (6.111), and (6.112),

$$\#\mathcal{N}_{u_{0,\varepsilon}, \eta} \leq \|u_{0,\varepsilon}\|_{\mathfrak{N}(u_{0,\varepsilon}), m, \text{Lip}}^m \mathfrak{N}(u_{0,\varepsilon}) \eta^{-m} |\log_2 \eta| \leq \|u_0\|^{m + \frac{1}{\alpha_0}} \varepsilon^{-\frac{1}{\alpha_0}} \eta^{-m} |\log_2 \eta|, \quad (6.115)$$

and, uniformly in  $t \in I$ ,

$$\#\mathcal{N}_{t, f_\varepsilon, \eta} \leq \|f_\varepsilon(t, \cdot)\|_{\mathfrak{N}(f_\varepsilon), m, \text{Lip}}^m \mathfrak{N}(f_\varepsilon) \eta^{-m} |\log_2 \eta| \leq \|f\|^{m + \frac{1}{\alpha_f}} \varepsilon^{-\frac{1}{\alpha_f}} \eta^{-m} |\log_2 \eta|. \quad (6.116)$$

Employing again a time-discretization of size  $q$ , (3.14) suggests the following DNN approximation to  $u$  which yields, on account of Lemma 6.4,

$$\left| \int_0^t f_\varepsilon(s, w, y) ds - \sum_{i=1}^q \rho_i(\xi_{i(t)}) f_\varepsilon(\xi_i, w, y) \right| \leq \frac{\|f\| T^2}{2q}, \quad (6.117)$$

where  $i(t) = \max\{i : [\xi_i] \leq t\}$ . Finally, we know from Theorem 4.4 that there exists a DNN  $\mathcal{N}_{z, \varepsilon}$  that approximates the characteristic field  $z$  within accuracy  $\varepsilon$ , i.e., in view of (4.10),

$$\|z - \mathcal{N}_{z, \varepsilon}\|_{L_\infty(I \times D \times \mathcal{Y})} \leq \varepsilon, \quad \#\mathcal{N}_{z, \varepsilon} \lesssim e^{LT(m+1)} \left( \frac{T d_y}{\varepsilon} \right)^{m+1} |\log_2(d_y/\varepsilon)|^2, \quad (6.118)$$

with a constant depending on  $L, A, \Lambda, m$ . Hence the network  $\mathcal{N}_{u,\varepsilon,\eta,q}$  formed by composing the DNNs  $\mathcal{N}_{u_0,\varepsilon,\eta}, \mathcal{N}_{t,f_\varepsilon,\eta}, i = 1, \dots, q$ , with the approximate characteristics  $\mathcal{N}_{z,\varepsilon}$  satisfies, on account of (2.9), (2.8), (6.115), and (6.116),

$$\begin{aligned}
\#\mathcal{N}_{u,\varepsilon,\eta,q} &\leq \#(\mathcal{N}_{u_0,\varepsilon,\eta} \circ \mathcal{N}_{z,\varepsilon}) + q \max_{i=1,\dots,q} \#(\mathcal{N}_{\xi_i,f_\varepsilon,\eta} \circ \mathcal{N}_{z,\varepsilon}) \\
&= \#\mathcal{N}_{u_0,\varepsilon,\eta} + \#\mathcal{N}_{z,\varepsilon} + q \left( \max_{i=1,\dots,q} \#\mathcal{N}_{\xi_i,f_\varepsilon,\eta} + \#\mathcal{N}_{z,\varepsilon} \right) \\
&\lesssim \left( \|u_0\|^{m+\frac{1}{\alpha_0}} \varepsilon^{-\frac{1}{\alpha_0}} + q \|f\|^{m+\frac{1}{\alpha_f}} \varepsilon^{-\frac{1}{\alpha_f}} \right) \eta^{-m} |\log_2 \eta| \\
&\quad + (1+q) C e^{LT(m+1)} \left( \frac{Td_y}{\varepsilon} \right)^{m+1} |\log_2(d_y/\varepsilon)|^2. \tag{6.119}
\end{aligned}$$

To determine  $\eta$ , we have by (3.14),

$$\begin{aligned}
\|u(t, \cdot) - \mathcal{N}_{u,\varepsilon,\eta,q}(t, \cdot)\|_{L_\infty(\Omega \times \mathcal{Y})} &\leq \|u_0(z(-t, t, \cdot; \cdot)) - \mathcal{N}_{u_0,\varepsilon,\eta} \circ \mathcal{N}_{z,\varepsilon}(-t, t, \cdot; \cdot)\|_{L_2(\Omega \times \mathcal{Y})} \\
&\quad + \sup_{x,y} \left| \int_0^t f(s, z(t-s, t, x, y)) ds - \sum_{i=1}^{i(t)} \rho_i(\xi_i) \mathcal{N}_{\xi_i,f_\varepsilon,\eta} \circ \mathcal{N}_{z,\varepsilon}(\xi_i(t) - \xi_i, x, y) \right| \\
&=: Q_1 + \sup_{x,y} Q_2(x, y).
\end{aligned}$$

Regarding  $Q_1$ , because of (6.113),

$$\begin{aligned}
Q_1 &\leq \|u_0(z(-t, t, \cdot; \cdot)) - u_0(\mathcal{N}_{z,\varepsilon}(-t, t, \cdot; \cdot))\|_{L_\infty(D \times \mathcal{Y})} \\
&\quad + \|u_0(\mathcal{N}_{z,\varepsilon}(-t, t, \cdot; \cdot)) - u_{\varepsilon,0}(\mathcal{N}_{z,\varepsilon}(-t, t, \cdot; \cdot))\|_{L_\infty(D \times \mathcal{Y})} \\
&\quad + \|u_{\varepsilon,0}(\mathcal{N}_{z,\varepsilon}(-t, t, \cdot; \cdot)) - \mathcal{N}_{u_0,\varepsilon,\eta}(\mathcal{N}_{z,\varepsilon}(-t, t, \cdot; \cdot))\|_{L_\infty(D \times \mathcal{Y})} \\
&\stackrel{(6.111)}{\leq} (1+L_0)\varepsilon + \|u_0\| \eta \mathfrak{N}(u_0, \varepsilon) \lesssim (1+L_0)\varepsilon + \|u_0\|^{1+\frac{1}{\alpha_0}} \eta \varepsilon^{-\frac{1}{\alpha_0}}, \tag{6.120}
\end{aligned}$$

where we have used (4.15) in the last step. Similarly, by (6.112) (6.114), and (6.117),

$$\begin{aligned}
Q_2(x, y) &\leq TL_f \varepsilon + \int_0^t |f(s, \mathcal{N}_{z,\varepsilon}(t-s, t, x, y)) - f_\varepsilon(s, \mathcal{N}_{z,\varepsilon}(t-s, t, x, y))| ds \\
&\quad + \int_0^t \left| f_\varepsilon(s, \mathcal{N}_{z,\varepsilon}(t-s, t, x, y)) - \sum_{i=1}^{i(t)} \rho_i(\xi_i) f_\varepsilon(\xi_i, \mathcal{N}_{z,\varepsilon}(\xi_i(t) - \xi_i, x, y)) \right| \\
&\quad + \sum_{i=1}^{i(t)} \rho_i(\xi_i) \left| f_\varepsilon(\xi_i, \mathcal{N}_{z,\varepsilon}(\xi_i(t) - \xi_i, x, y)) - \mathcal{N}_{\xi_i,f_\varepsilon,\eta}(\mathcal{N}_{z,\varepsilon}(\xi_i(t) - \xi_i, x, y)) \right|, \tag{6.121}
\end{aligned}$$

which gives

$$\begin{aligned} Q_2(x, y) &\leq (1 + L_f)T\varepsilon + \frac{\|f\|T^2}{2q} + T\eta\mathfrak{R}(f_\varepsilon)\|f\| \\ &\leq (1 + L_f)T\varepsilon + \frac{\|f\|T^2}{2q} + T\eta\|f\|^{1+\frac{1}{\alpha_f}}\varepsilon^{-\frac{1}{\alpha_f}}. \end{aligned} \quad (6.122)$$

Now let

$$\alpha := \min\{\alpha_0, \alpha_f\}, \quad q(\varepsilon) = \frac{T}{2\varepsilon\|f\|^{1/\alpha_f}}, \quad \eta(\varepsilon) = \varepsilon^{1+\max\{\frac{1}{\alpha_f}, \frac{1}{\alpha_0}\}} \quad (6.123)$$

to conclude from (6.120) and (6.122) that the network  $\mathcal{N}_{u,\varepsilon} := \mathcal{N}_{u,\varepsilon,\eta(\varepsilon),q(\varepsilon)}$  satisfies

$$\begin{aligned} \|u - \mathcal{N}_{u,\varepsilon}\|_\infty &\leq \left\{1 + L_0 + \|u_0\|^{1+\frac{1}{\alpha_0}} + T(1 + L_f + 2\|f\|^{1+\frac{1}{\alpha_f}})\right\}\varepsilon \\ &\leq C_{L_0,L_f,T}(1 + \|u_0\|^{1+\frac{1}{\alpha}} + \|f\|^{1+\frac{1}{\alpha}})\varepsilon =: C_*\varepsilon =: \varepsilon_*. \end{aligned} \quad (6.124)$$

This confirms the first part of (4.16).

Now we infer from (6.119), (6.111), (6.112) that (supressing now explicit dependence of constants on  $T$ )

$$\begin{aligned} \#\mathcal{N}_{u,\varepsilon} &\lesssim \left\{\|u_0\|^{m+\frac{1}{\alpha_0}}\varepsilon^{-\frac{1}{\alpha_0}} + \frac{1}{2\varepsilon\|f\|^{1/\alpha_f}}\|f\|^{m+\frac{1}{\alpha_f}}\varepsilon^{-\frac{1}{\alpha_f}}\right\}\varepsilon^{-m(1+\frac{1}{\alpha})}|\log_2 \varepsilon| + \left(\frac{d_y}{\varepsilon}\right)^{m+1}\left|\log_2 \frac{d_y}{\varepsilon}\right|^2 \\ &\leq \left\{\|u_0\|^{m+\frac{1}{\alpha_0}}\varepsilon^{-\frac{1}{\alpha}} + \frac{1}{2}\|f\|^m\varepsilon^{-(1+\frac{1}{\alpha})}\right\}\varepsilon^{-m(1+\frac{1}{\alpha})}|\log_2 \varepsilon| + \left(\frac{d_y}{\varepsilon}\right)^{m+1}\left|\log_2 \frac{d_y}{\varepsilon}\right|^2 \\ &\lesssim (\|u_0\|^{m+\frac{1}{\alpha_0}} + \|f\|^m + d_y^{m+1})\varepsilon^{-(m+1)(1+\frac{1}{\alpha})}\left|\log_3 \frac{d_y}{\varepsilon}\right|^2 \end{aligned}$$

with a constant depending on  $L, A, T$ . Substituting  $\varepsilon = C_*^{-1}\varepsilon_*$ , and using that  $d_y \geq \|u_0\|^{\alpha_0}, \|f\|$ , yields that achieving accuracy  $\varepsilon_*$  requires network size of the order

$$\mathcal{N}_{u,\varepsilon} \lesssim d_y^{m+1}C_*^{\frac{(m+1)(\alpha+1)}{\alpha}}\varepsilon_*^{-(m+1)(1+\frac{1}{\alpha})}\left|\log_3 \frac{d_y}{\varepsilon_*}\right|^2,$$

where  $C_*$  is the constant from (6.124). This confirms the second part of (4.16) as well as (4.17).

Applying Lemma 6.2, assuming that  $\|u_0\|, \|f\|$  are dominated by  $d_y$ , straightforward calculations confirm that the error rate (4.18) holds for

$$\hat{\gamma}(r) \lesssim C_*^{-1}d_y^{-\frac{\alpha}{(\alpha+1)}}r^{\frac{\alpha}{(m+1)(\alpha+1)}}\left|\log_2 r d_y^{\frac{1}{\alpha}}\right|^{-\frac{2\alpha}{(m+1)(\alpha+1)}}.$$

This concludes the proof of Theorem 4.8. □

## References

- [1] C. Beck, S. Becker, P. Grohs, N. Jaafari, A. Jentzen, Solving the Kolmogorov PDE by means of deep learning, *Journal of Scientific Computing* 88 (3), 1–28
- [2] A. Cohen, R. DeVore, G. Petrova, P. Wojtaszczyk, Optimal stable nonlinear approximation, (2020), arXiv:2009.09907.
- [3] W. Dahmen, R. DeVore, L. Grasedyck, E. Süli, Tensor Sparsity of Solutions to High-Dimensional Elliptic Partial Differential Equations, *Foundation of Computational Mathematics*, 16 (No 4) (2016), 813–874. (DOI) 10.1007/s10208-015-9265-9.
- [4] W. Dahmen, F. Gruber, O. Mula, An Adaptive Nested Source Term Iteration for Radiative Transfer Equations, *Mathematics of Computation*, 89 (No 324) (2020), 1605–1646, <https://doi.org/10.1090/mcom/3505>.
- [5] I. Daubechies, R. DeVore, S. Foucart, B. Hanin, G. Petrova, Nonlinear approximation and (deep) ReLU networks, *Constructive Approximation*, 55 (2022), 127–172.
- [6] R. DeVore, B. Hanin, G. Petrova, Neural Network Approximation, *Acta Numerica*, Volume 30 , May 2021, pp. 327 – 444. DOI: <https://doi.org/10.1017/S0962492921000052>
- [7] R. DeVore, R. Howard, C. Micchelli, Optimal non-linear approximation, *Manuscripta Math.* 4 (1989), 469–478.
- [8] W. E, C. Ma, L. Wu, The Barron Space and the Flow-Induced Function Spaces for Neural Network Models, *Constructive Approximation*, 55 (2022), 369–406.
- [9] F. Girosi, T. Poggio, Representation properties of networks: Kolmogorov’s Theorem is irrelevant, *iKun neural computation*, (1), (no. 4) (1989), 465–469, doi: 10.1162/neco.1989.1.4.465.
- [10] R. Gribonval, G. Kutyniok, M. Nielsen, F. Voigtlaender, Approximation spaces of deep neural networks, *Constructive Approximation*, 55(2022), 259–367.
- [11] P. Grohs, L. Herrmann, Deep neural network approximation for high-dimensional parabolic Hamilton-Jacobi-Bellman equations, arXiv:2103.05744v1 [math.NA], March 2021.
- [12] P. Grohs, F. Hornung, A. Jentzen, P. Von Wurstemberger, A proof that artificial neural networks overcome the curse of dimensionality in the numerical approximation of Black-Scholes partial differential equations, *Memoirs of the American Mathematical Society*, <https://doi.org/10.48550/arXiv.1809.02362>
- [13] P. Grohs, D. Perekrstenko, D. Elbrächter, H. Bölcskei, Deep neural network approximation theory, *IEEE Transactions on Information Theory*, (2019), arXiv:1901.02220.

- [14] P. Grohs, F. Voigtlaender, Proof of the Theory-to-Practice Gap in Deep Learning via Sampling Complexity bounds for Neural Network Approximation Spaces, arXiv:2104.02746v1 [cs.LG], 04/06/2021.
- [15] I. Gühring, G. Kutyniok, P. Petersen, Error bounds for approximations with deep ReLU neural networks in  $W^{s,p}$ -norms, Analysis and Applications, 18 (5)(2020), 803–859, <https://doi.org/10.1142/S0219530519410021>, <http://arxiv:1902.07896v1> [math.FA], Feb. 21, 2019.
- [16] V. Kurková, Kolmogorov’s Theorem is relevant, <https://doi.org/10.1162/neco.1991.3.4.617>
- [17] G. Kutyniok, P. Petersen, M. Raslan, R. Schneider, A Theoretical Analysis of Deep Neural Networks and Parametric PDEs, Constructive Approximation, 55 (2022), 73–125. <https://doi.org/10.48550/arXiv.1904.00377>
- [18] J.P. Lions, B. Perthame, E. Tadmor, A kinetic formulation of multidimensional scalar conservation laws and related equations, Journal of the American Mathematical Society, 7 (no. 1)(1994), 169–191.
- [19] G.G. Lorentz, Metric entropy, widths, and superpositions of functions. American Mathematical Monthly. 69 (6)(1962), 469–485. doi:10.1080/00029890.1962.11989915.
- [20] J. Lu, Z. Shen, H. Yang, S. Zhang, Deep network approximation for smooth functions, SIAM J. Math. Anal., 53 (5) (2020), arXiv:2001.03040.
- [21] H. N. Mhaskar, T. Poggio, Function approximation by deep networks, Communications on Pure and Applied Analysis, 19(8)(2020), 4085–4095. doi: 10.3934/cpaa.2020181 (2019). arXiv:1905.12882.
- [22] J.A.A. Opschoor, C. Schwab, J. Zech, Exponential ReLU DNN expression of holomorphic maps in high dimension, Constructive Approximation, 55 (2022), 537–582.
- [23] F. Laakmann, P. Petersen, Efficient Approximation of Solutions of Parametric Linear Transport Equations by ReLU DNNs, Advances in Computational Mathematics, 47 (11) (2021), <https://doi.org/10.1007/s10444-020-09834-7>.
- [24] P. Petersen, F. Voigtlaender, Optimal approximation of piecewise smooth functions using deep relu neural networks, Neural Networks 108, (2018), 296–330.
- [25] J. Schmidt-Hieber, Nonparametric regression using deep neural networks with ReLU activation, Ann. Statist. 48(4)(2020), 1875–1897. DOI: 10.1214/19-AOS1875(2020),
- [26] A. Vasseur, Kinetic semi-discretization of scalar conservation laws and convergence by using averaging lemmas, SIAM J. Numer.Anal., 36 (No.2)(1999), 465–474.
- [27] D. Yarotsky, Error bounds for approximations with deep ReLU networks, Neural Networks 94, (2017), 103–114.