

Systemic-risk and evolutionary stable strategies in a financial network

¹Indrajit Saha and ¹Veeraruna Kavitha

¹ IIT Bombay, India

Abstract

We consider a financial network represented at any time instance by a random liability graph which evolves over time. The agents connect through credit instruments borrowed from each other or through direct lending, and these create the liability edges. These random edges are modified (locally) by the agents over time, as they learn from their experiences and (possibly imperfect) observations. The settlement of the liabilities of various agents at the end of the contract period (at any time instance) can be expressed as solutions of random fixed point equations. Our first step is to derive the solutions of these equations (asymptotically and one for each time instance), using a recent result on random fixed point equations. The agents, at any time instance, adopt one of the two available strategies, risky or risk-free investments, with an aim to maximize their returns. We aim to study the emerging strategies of such replicator dynamics that drives the financial network. We theoretically reduce the analysis of the complex system to that of an appropriate ordinary differential equation (ODE). Using the attractors of the resulting ODE we showed that the replicator dynamics converges to one of the two pure evolutionary stable strategies (all risky or all risk-free agents); one can have mixed limit only when the observations are imperfect. We verified our theoretical findings using exhaustive Monte Carlo simulations. We established that the dynamics avoid the emergence of the systemic-risk regime (where majority default).

1 Introduction

We consider a financial network, where the agents are interconnected to each other through financial commitments (e.g., borrowing, lending, etc). In addition they invest in either risk-free (bonds, fixed deposit, government projects etc) or risky securities (derivative markets, stocks etc). In such a system the agents not only face random economic shocks (received via significantly smaller returns of their risky investments), they are also affected by the percolation of the shocks faced by their neighbours (creditors/borrowers), neighbours of their neighbours and so on. In the recent years from 2007-2008 onwards, there is a surge

of activity to study the financial and systemic level risks caused by such a percolation of shocks ([1, 2, 3, 4]). Systemic risk is the study of the risks related to financial networks, when individual or entity level shocks can trigger severe instability at system level that can collapse the entire economy (e.g., [1, 2, 3]). In this set of papers, the authors study the kind of topology (or graph structure) that is more stable towards the percolation of shocks in a financial network, where stability is measured in terms of number/fraction of defaults.

In contrast to many existing studies related to systemic risk, we consider heterogeneous agents and an evolutionary framework. There are two groups of agents existing simultaneously in the network; one group of agents invest in risk-free assets, while the rest consider risky investments. The second group borrows money from other agents of the network to gather more funds towards the risky investments (with much higher expected returns). These investments are subjected to large (but rare) economic shocks, which can potentially percolate throughout the network and can also affect the ‘risk-free’ agents. The extent of percolation depends upon the relative sizes of the two groups.

We further consider that the system evolves over time, some strategic agents change their choices (investment types) and we also have agents randomly joining the network at various time instances. The new agents choose their investment type (risky or risk-free) based on their observations of the returns of a random sample of agents that invested in the previous round. The existing agents may change their strategy based on their own returns and their observations of a random sample. The relative sizes of the two groups changes, the network structure changes, which influences the (economic shock-affected) returns of the agents in the next round, which in turn influences the decision of the agents for the round after. Thus the system evolves after each round. We study this process using the well known evolutionary game theoretic tools.

From the perspective of a financial network, this type of work is new to the best of our knowledge. We found few papers that consider evolutionary approach in other aspects related to finance; in [5], the authors study the financial safety net (a series of the arrangement of the firms to maintain financial stability), and analyze the evolution of the bank strategies (to take insurance or not); recently in [6] authors consider an evolutionary game theoretic model with three types of players, i) momentum traders ii) contrarian traders iii) fundamentalists and studied the evolution of the relative populations. As already mentioned, these papers relate to very different aspects in comparison with our work.

In [7] we have an initial analysis of this flavor related to systemic risk analysis; the new agents choose their strategy once for all, when they join the network; we showed that all the agents eventually revert to either risky or risk-free portfolios unless they choose their strategies based on a large sample of observations. In the former scenario, the dynamics converge to a pure (only one of the strategies, risky/risk-free, survive) evolutionary stable strategy (ESS), while the latter scenario converges to a mixed ESS. In [7], we derive the analysis, only for the cases in which the dynamics is smooth and always converges to one solution (ESS) for a given set of parameters.

The current paper considers a significant generalization. We consider rational agents

that modify their strategies at various time instances, in addition to the agents that randomly join the network and choose their investment type. These type of processes are often analyzed using ordinary differential equations (ODEs). The resulting ODEs in this context are non-smooth and we completed the analysis of such a complex ODE, by using the approximate solutions of a relevant set of random fixed point equations (derived using the recent results of [4, 8]). The analysis of the simplified representations of the fixed points helped us in deriving the attractors and the corresponding regions of attraction of the otherwise complex (non-smooth) ODE. We finally showed that the network can converge to one of the two pure ESSs based on network parameters. More importantly, this convergence also depends upon the initial configuration. Further the dynamics avert the systemic risk regime by converging to the pure ESS of all risk-free agents, in scenarios that have potential to lead to the systemic risk event (when majority of the agents fail to clear their obligations). One may have mixed ESS, when the observations are partially erroneous. Interestingly even these imperfect observations are (majorly) sufficient to drive the system away from systemic risk regime.

The analysis of these complex networks (in each round) necessitated the study of random fixed point equations (defined sample path-wise in large dimensional spaces), which represent the clearing vectors of all the agents ([1, 3] etc). As already mentioned, the study is made possible because of the recent result in [4, 8], which provided an asymptotically accurate one dimensional equivalent solution.

We conducted exhaustive Monte Carlo (MC) simulation based numerical study with the following details: a) the dynamics (economic shocks, random returns, random set of agents modifying their choices etc.) are randomly generated; and b) the random clearing vector is obtained by numerically solving the relevant fixed point equations, in each round; however, we did not generate random liability connections in each round. The limits of this dynamics well match with the attractors of the corresponding ODE.

Using the random fixed point approximation, we also derive the closed-form expressions for the solution of the ODEs approximating the financial dynamics. Once again using exhaustive MC simulations we showed that the ODE trajectories well approximate the random financial trajectories, even for finite (initial) number of rounds. The approximation is good even for few hundreds of entities and even for few initial (≈ 100) rounds.

We conclude this section with a brief discussion on the evolutionary game theory. The finance network and random fixed point solutions for clearing vectors are discussed respectively in Sections 2 and 3. The ODE approximation is derived in Section 4, while (mixed) evolutionary stable strategies are considered in Section 5.

Evolutionary game theory: Traditionally evolutionary game models have been used in the literature to study animal behavior (e.g., [9]). The main purpose of such a study is to derive asymptotic analysis, in particular to identify the predominant behavioural traits that persist (emerge and remain forever hence after) because of their stability against mutations. Several important questions answered in this context are: will the dynamics converge, i.e., would the relative fractions of various populations (each type has a specific characteristic

trait) settle as the number of rounds/generations increase? will some of the strategies/traits disappear eventually? if more than one population type survives what would be the asymptotic fractions? etc. This kind of analysis is common in various other types of networks (e.g., wireless networks (e.g., [10]), biological networks (e.g., [11, 12])), but is relatively less studied in the context of financial networks (e.g., [6]).

The key ingredients of evolutionary game theory are: a) a large number of players, b) the dynamics, and c) the pay-off function (e.g., see the pioneering work [9]). Replicator dynamics deals with the evolution of reward-based learning strategies. Typically these dynamics converge to a stable equilibrium point called Evolutionary Stable Strategy (ESS), which can be seen as a refinement of a strict Nash equilibrium ([9]). Formally, in a two-player symmetric game, a pure strategy \hat{s} is said to be evolutionary stable if: i) (\hat{s}, \hat{s}) is a Nash equilibrium, i.e., $u(\hat{s}, \hat{s}) \geq u(s', \hat{s}) \forall s'$ and, ii) If (\hat{s}, \hat{s}) is not a strict Nash equilibrium (the above inequality replaced by strict inequality), i.e., \exists some $s' \neq \hat{s}$ such that $u(\hat{s}, \hat{s}) = u(s', \hat{s})$, then $u(\hat{s}, s') > u(s', s')$.

2 Finance Network

We consider random graphs, where the edges represent the financial connection between the nodes. Any two nodes are connected with probability $p_{ss} > 0$ independent of the others. This graph represents a financial network where borrowing and lending are represented by edges and the weights over them. The modeller may not have access to exact connections of the network, but a random graph is a good model to analyse such complex systems. In particular we consider graphs that satisfy the assumptions of [4, 8].

The agents are repeatedly investing in some financial projects. In each round of investment, the agents borrow/lend from/to some random subset of the agents of the network. Some of them may invest the remaining in a risk-free investment (which has a constant rate of interest r_s). While the others invest the rest of their money in risky investments which have random returns; we consider a binomial model in which returns are high (rate u) with high probability δ and can have large shocks (rate d), but with small probability $(1 - \delta)$; it is clear that $d < r_s < u$. We thus have two types of agents, we call the group that invests in risk-free projects as ‘risk-free’ group (G_1), the rest are referred to as ‘risky’ group (G_2).

New agents may join the network in each round of investment. They choose their investment type, either risk-free or risky, for the first time based on the previous experience of the network. The new agents learn from the returns of the agents of the previous round of investments before choosing a suitable investment type. To be precise, they learn from the experience of a random sample (returns of two random agents) of the network corresponding to the previous round.

In any round, a random agent of the network may modify its strategy, based on its own experience and that of another random agent of the network.

Two strategies: As mentioned before, there are two strategies available in the financial market. Risk-free agents of G_1 use strategy 1; these agents lend some amount of their

initial wealth to other agents (of G_2) that are willing to borrow, while the rest is invested in a government security, for example, bonds, government project etc. Risky agents of G_2 are adapting strategy 2, wherein they borrow funds from the other agents and invest in risky security, for example, derivative markets, stocks, corporate loans etc. These agents also lend to other agents of G_2 . Let ϵ_t be the fraction of the agents in G_1 group and let $n(t)$ be the total number of agents in round t . Thus the total number of agents (during round t) in group 1 equals $n_1(t) := |G_1| = n(t)\epsilon_t$, similarly $n_2(t) := |G_2| = n(t)(1 - \epsilon_t)$.

We consider that some new agents may join in each round and a random number of the existing agents may modify their choice. Thus the size of the network is increasing. The agents are homogeneous, i.e., they reserve the same wealth $w > 0$ for investments (at the initial investment period) of each round. *Each round is composed of two time periods, the agents invest during the initial investment period and they obtain their returns after some given time gap.* The two time period model is borrowed from [1, 3, 4] etc. Recall the agents make their choice for the next round, based on their observations of these returns.

Initial investment phases: During the initial investment phases (of any round t), any agent $i \in G_1$ lends to any agent $j \in G_2$ with probability p_{ss} and it lends equal amount $w/n(t)p_{ss}$ to each of the approachers that approached it for loan; let I_{ij} be the indicator of this lending event. Thus any agent of G_1 lends approximately $w(1 - \epsilon_t)$ fraction to agents of G_2 ; note that for large $n(t)$, the number of approachers of G_2 approximately equals $n(t)(1 - \epsilon_t)p_{ss}$. The agents of G_1 invest the rest $w\epsilon_t$ in risk-free investment (returns at fixed interest-rate r_s). We now provide complete details of the network in one round, when $\epsilon_t = \epsilon$.

Let \tilde{w} be amount accumulated by any agent of G_2 out of which a positive fraction α is invested towards the other banks of G_2 and $(1 - \alpha)$ portion is invested in risky security. These amounts could be random and different from agent to agent, but with large networks (by law of large numbers) one can approximate these to be constants. Thus the amount accumulated by a typical G_2 agent is given by the following:

$$\tilde{w} = \underbrace{w + w\epsilon}_{\text{Initial wealth + Borrowed from } G_1} + \underbrace{\tilde{w}\alpha}_{\text{Borrow from } G_2}, \quad \text{and hence} \quad (1)$$

$\tilde{w} = \frac{w(1+\epsilon)}{(1-\alpha)}$; this amount is used for investments/lending in the initial phase. Once again this equation is accurate when $n(t)$ is large and is used for asymptotic analysis. Thus the total investment towards the risky venture equals $\tilde{w}(1 - \alpha) = w(1 + \epsilon)$. The G_2 agents have to settle their liabilities at the end of the return/contract period (in each round) and this would depend upon their returns from the risky investments. Thus the total liability of any agent of G_2 is $y = (w\epsilon + \tilde{w}\alpha)(1 + r_b)$, where r_b is the borrowing rate; by simplifying

$$y = \frac{w(\epsilon + \alpha)(1 + r_b)}{(1 - \alpha)}. \quad (2)$$

For simplicity of explanation, we are considering constant terms to represent all these quantities, in reality they would be i.i.d. quantities which are further independent of other rounds and the asymptotic analysis would go through as in [4, 8]. Similarly, any agent of

G_2 lends the following amount to each of its approachers (of G_2):

$$\frac{\alpha\tilde{w}}{n(t)(1-\epsilon_t)p_{ss}} = \frac{\alpha w(1+\epsilon)}{n(t)(1-\epsilon_t)p_{ss}(1-\alpha)}. \quad (3)$$

Returns and Clearing Vectors: We fix round t and avoid notation t for simpler representation. The agents of G_2 have to clear their liabilities during this phase in every round. Recall the agents of G_2 invested $w(1+\epsilon)$ amount in risky-investments and the corresponding random returns (after economic shocks) are:

$$K_i = \begin{cases} w(1+\epsilon)(1+u) =: k_u, & \text{w.p. (with probability) } \delta \\ w(1+\epsilon)(1+d) =: k_d, & \text{otherwise.} \end{cases}$$

This is the well known *binomial model* ([1, 3]), in which the upward moment occurs with probability δ and downward moment with $(1-\delta)$. The agents have to return y (after the interest rate r_b) amount to their creditors, however may not be able to manage the same because of the above economic shocks. In case of default, the agents return the maximum possible; let X_i be the amount cleared by the i -th agent of group G_2 . Here we consider a standard bankruptcy rule, limited liability and pro-rata basis repayment of the debt contract (see [1, 3]), where the amounts returned are proportional to their liability ratios. Thus node j of G_2 pays back $X_j L_{ji}/y$ towards node i , where L_{ji} the liability/the amount borrowed during initial investment phases equals (see the details of previous sub-section and equation (3)):

$$L_{ji} = \begin{cases} I_{ji} \frac{w}{np_{ss}}, & \text{if } i \in G_1 \\ I_{ji} \frac{\alpha w(1+\epsilon)}{np_{ss}(1-\alpha)(1-\epsilon)}, & \text{if } i \in G_2. \end{cases} \quad (4)$$

Thus the maximum amount cleared by any agent $i \in G_2$, X_i , is given by the following fixed point equation in terms of the clearing vector $\{X_j\}_{j \in G_2}$ composed of clearing values of all the agents (see [1, 3] etc):

$$X_i = \min \left\{ \left(K_i + \sum_{j \in G_2} X_j \frac{L_{ji}}{y} - v \right)^+, y \right\}, \quad (5)$$

with the following details: the term K_i is returns of risky investment, the term $\sum_{j \in G_2} X_j L_{ji}/y$ equals the claims from the other agents (those borrowed from agent i) and v denotes the taxes/operational costs/deposits/senior liabilities (e.g., as in [1, 13, 8, 4]). In other words, agent $i \in G_2$ will pay back the (maximum possible) amount, $(K_i + \sum_{j \in G_2} X_j \frac{L_{ji}}{y} - v)^+$, in case of a default, and otherwise, will exactly pay back the liability amount y .

Surplus of any agent is defined as the amount left behind, after clearing all the liabilities. This represents the utility of the agent in the given round. The surplus/return of the agent $i \in G_2$ and $i \in G_1$ respectively are given by,

$$R_i^2 = \left(K_i + \sum_{j \in G_2} X_j \frac{L_{ji}}{y} - v - y \right)^+ \text{ and } R_i^1 = \left(w\epsilon(1+r_s) + \sum_{j \in G_2} X_j \frac{L_{ji}}{y} - v \right)^+. \quad (6)$$

3 Asymptotic approximation

We thus have dynamic graphs whose size increases, with some probability, in each round. In this section, we obtain appropriate asymptotic analysis of these graphs, with an aim to derive the pay-off of each group after each round. Towards this, we derive the (approximate) closed form expression of the equations (6) - (6) which are the returns of the various agents after the settlement of the liabilities. The returns of the agents depend upon how other agents settle their liabilities to their creditors. Thus our first step is to derive the solution of the clearing vector fixed point equations (5). Observe that the clearing vector $\{X_j\}_{j \in G_2}$ is the solution of the vector-valued random fixed point equations (5) in n -dimensional space (where n is the size of the network), defined sample-path wise.

Clearing vectors using results of [8, 4]: Our financial framework can be analysed using the results of [8], as the details of the model match the assumptions of the paper. Observe that $\alpha(1 + \epsilon)/(\alpha + \epsilon) < 1$. By results of [8, Section 4] (like [4, Theorem 1]), the aggregate claims converge almost surely to constant values, as the network size increases to infinity (see subsection 8.1 of Appendix for more details):

$$\begin{aligned} \text{(claims of } G_1), \quad \sum_{j \in G_2} X_j \frac{L_{ji}}{y} &\rightarrow \frac{(1 - \alpha)(1 - \epsilon)}{\alpha + \epsilon} \bar{x}^\infty \text{ a.s.,} \\ \text{(claims of } G_2), \quad \sum_{j \in G_2} X_j \frac{L_{ji}}{y} &\rightarrow \frac{\alpha(1 + \epsilon)}{(\alpha + \epsilon)} \bar{x}^\infty \text{ a.s.,} \end{aligned} \quad (7)$$

where the common expected clearing value \bar{x}^∞ satisfies the following fixed point equation¹ in one-dimension (see [8, Section 4], [4, Theorem 1]):

$$\bar{x}^\infty = E \left[\min \left\{ \left(K_i + \frac{\alpha(1 + \epsilon)}{\alpha + \epsilon} \bar{x}^\infty - v \right)^+, y \right\} \right]. \quad (8)$$

Also by the same theorem, the clearing vectors converge almost surely to (asymptotically independent) random vectors:

$$X_i \rightarrow \min \left\{ \left(K_i + \frac{\alpha(1 + \epsilon)}{\alpha + \epsilon} \bar{x}^\infty - v \right)^+, y \right\}, \text{ for } i \in G_2. \quad (9)$$

By virtue of the above results, the random returns given by (6) converge almost surely:

$$\begin{aligned} R_i^1 &\rightarrow \left(w\epsilon(1 + r_s) + \frac{(1 - \alpha)(1 - \epsilon)}{(\alpha + \epsilon)} \bar{x}^\infty - v \right)^+, \text{ for } i \in G_1 \\ R_i^2 &\rightarrow \left(K_i + \frac{\alpha(1 + \epsilon)}{\alpha + \epsilon} \bar{x}^\infty - v - y \right)^+, \text{ for } i \in G_2. \end{aligned} \quad (10)$$

Probability of default: It is defined as the fraction of agents of G_2 that failed to pay back their full liability, i.e., $P_d := P(X_i < y)$. For large networks (when the initial network size n_0 itself is sufficiently large), one can use the above approximate expressions and using the same we obtain the default probabilities and the aggregate clearing vectors in the following (proof is in Appendix):

¹This approximation is valid for $\epsilon \in (0, 1)$. In Appendix we also provide the details for $\epsilon \in \{0, 1\}$.

Lemma 1 Consider $w(1+d)(1+\epsilon) \geq v$. There is a unique solution to (8) and the asymptotic average clearing vector and the default probability of G_2 is given by:

$$(\bar{x}^\infty, P_d) = \begin{cases} (y, 0) & \text{if } c_\epsilon \geq a_1 \\ \left(\frac{\delta y + (1-\delta)\underline{w}}{1-(1-\delta)c_\epsilon}, 1-\delta \right) & \text{if } a_2 \leq c_\epsilon < a_1 \\ \left(\frac{E[W]}{1-c_\epsilon}, 1 \right) & \text{if } c_\epsilon < a_2, \end{cases} \quad (11)$$

where, $c_\epsilon = \frac{\alpha + \alpha\epsilon}{\alpha + \epsilon}$, $E[W] = \delta\bar{w} + (1-\delta)\underline{w}$, $\underline{w} = k_d - v$, $\bar{w} = k_u - v$, $a_1 = \frac{y-\underline{w}}{y}$ and $a_2 = \frac{y-\bar{w}}{y-(1-\delta)(\bar{w}-\underline{w})}$. ■

We hence forth, replace (6) with its approximation (10). In [8] we have performed Monte Carlo (MC) simulations to demonstrate that the asymptotic approximation well matches the MC estimates with just few hundred financial entities. The approximation is better for more regular graphs.

4 Financial Replicator Dynamics

In every round of investments, we have a new network, that represents the liability structure of all the agents of that round, formed by their investment choices. New agents may join the network and choose their strategy, as well as the existing agents may modify their choices in each round.

The main purpose of the study of such replicator dynamics is to derive asymptotic analysis and answer questions such as: will the dynamics converge, i.e., would the relative fractions of various populations settle as the number of rounds increase? will some of the strategies disappear eventually? if more than one population type survives what would be the asymptotic fractions? etc. This kind of analysis is common in various other types of networks (e.g., wireless networks (e.g., [10]), biological networks (e.g., [11])), but is relatively less studied in the context of financial networks (e.g., [6]). We are interested in asymptotic outcome of these kinds of dynamics (if there exists one) and study the influence of various network parameters on the same. We also study the outcome after finite number of rounds.

Dynamics: After each round, a random number of new agents join the network and a random set of agents switch their strategies. Let \mathcal{N}_t be the number of new agents joining the network in round t . Each of them contact two random agents, uniformly from the ones that participated in the previous round. If both the contacted agents belong to the same group, the new agent adapts the strategy of that group. Otherwise, *the new agent adapts the strategy of the agent that had higher return in previous round*. The contacted agents may not reveal complete information, which can influence the choices of the new agent. We assume $\{\mathcal{N}_t\}_t$ are i.i.d and that $P(\mathcal{N}_t < \bar{\mathcal{N}}) = 1$ for some constant $\bar{\mathcal{N}} < \infty$.

Let $(n_1(t), n_2(t), n(t) = n_1(t) + n_2(t))$ represent respective sizes of groups G_1, G_2 and total population. Let $\epsilon_t := n_1(t)/n(t)$ represent the fraction belonging to G_1 . The

conditional probability and expectation, conditioned on ϵ_t , is denoted by P_{ϵ_t} and E_{ϵ_t} . A new agent joins G_1 , if it contacts both the agents from G_1 (which happens with conditional probability ϵ_t^2). In case it contacted one agent from each group (which happens w.p. $2\epsilon_t(1 - \epsilon_t)$), it would join G_1 w.p. $q_{\epsilon_t} := P_{\epsilon_t}(R^1 \geq R^2)$; observe q_{ϵ_t} is the probability that the return of agent from G_1 is superior. Then the expected number of agents ξ_t that join G_1 among \mathcal{N}_t , conditioned on ϵ_t , is given by:

$$E_{\epsilon_t}[\xi_t] = E_{\epsilon_t}[E[\xi_1(t)|\mathcal{N}_t]] = E[\mathcal{N}_t] (\epsilon_t^2 + 2\epsilon_t(1 - \epsilon_t)q_{\epsilon_t}).$$

This is the case when the contacted agents reveal correct information. *Due to incorrect information, any new agent can make an erroneous decision w.p. $(1 - b_n)$.* Thus,

$$\begin{aligned} E_{\epsilon_t}[\xi_t] &= E[\mathcal{N}_t](\epsilon_t^2 + \epsilon_t(1 - \epsilon_t)\hat{q}_{\epsilon_t}), \text{ with,} \\ \hat{q}_{\epsilon_t} &:= b_n q_{\epsilon_t} + (1 - b_n)(1 - q_{\epsilon_t}). \end{aligned} \quad (12)$$

A random i.i.d. number (\mathcal{S}_t) of the existing agents may attempt to switch their strategy. Each of them contact another agent (sampled uniformly) to compare with their experience. *The strategy of the former is modified to that of the contacted agent's only if the (revealed) return of the latter is better.*

Any agent attempting to switch, is from G_2 w.p. $1 - \epsilon_t$; and it switches to G_1 only if the contacted agent is from G_1 and if the corresponding return is better. Further due to incorrect information, there can be an erroneous decision w.p. $(1 - b_s)$. In all, any agent among \mathcal{S}_t can switch from G_2 to G_1 with (conditional) probability²:

$$(1 - \epsilon_t)\epsilon_t \hat{q}_{\epsilon_t}^s \text{ where } \hat{q}_{\epsilon_t}^s := (b_s q_{\epsilon_t} + (1 - b_s)(1 - q_{\epsilon_t})). \quad (13)$$

Let $\Xi_1(t)$ be the number of agents among \mathcal{S}_t that switch from G_2 to G_1 , similarly define $\Xi_2(t)$. From (13), computing as before, $E_{\epsilon_t}[\Xi_1(t)] = \epsilon_t(1 - \epsilon_t)\hat{q}_{\epsilon_t}^s E[\mathcal{S}_t]$. In similar lines,

$$E_{\epsilon_t}[\Xi_2(t)] = \epsilon_t(1 - \epsilon_t) (b_s(1 - q_{\epsilon_t}) + (1 - b_s)q_{\epsilon_t}) E[\mathcal{S}_t].$$

Thus the expected number of changes in G_1 are:

$$\begin{aligned} E_{\epsilon_t}[\Xi_1(t) - \Xi_2(t)] &= \epsilon_t(1 - \epsilon_t)(2\hat{q}_{\epsilon_t}^s - 1)E[\mathcal{S}_t], \\ &= \epsilon_t(1 - \epsilon_t)(2b_s - 1)(2q_{\epsilon_t} - 1)E[\mathcal{S}_t]. \end{aligned}$$

Analysis: The evolution after round t is summarized by:

$$\begin{aligned} n_1(t+1) &= n_1(t) + \xi_t + \Xi_1(t) - \Xi_2(t) \text{ and,} \\ n(t+1) &= n(t) + \mathcal{N}_t. \end{aligned} \quad (14)$$

Let ϵ_0 and n_0 respectively represent the initial fraction and total population in the system. Define $\psi_t := n(t)/(t+n_0)$ to represent the rate at which population evolves, and observe:

²Given that the agent switching is from G_1 , the probability that the contacted agent is from G_1 is actually $\epsilon_t n(t)/(n(t) - 1) \approx \epsilon_t$, the approximation improves as $n(t) \uparrow$.

$$\psi_{t+1} = \psi_t + \frac{1}{t + n_0 + 1} (\mathcal{N}_t - \psi_t). \quad (15)$$

Define $W_{t+1} = \xi_t + \Xi_1(t) - \Xi_2(t)$. Recall $\epsilon_t = n_1(t)/n(t)$ and so,

$$\epsilon_{t+1} = \frac{n(t)\epsilon_t + W_{t+1}}{n(t) + \mathcal{N}_t} = \epsilon_t + \frac{1}{t + n_0 + 1} \frac{W_{t+1} - \mathcal{N}_t\epsilon_t}{\psi_{t+1}}.$$

Define $\Theta_t = [\epsilon_t, \psi_t]^T$. One can rewrite the update equations,

$$\begin{aligned} \Theta_{t+1} &= \Theta_t + \gamma_t Y_{t+1}, \text{ with, } \gamma_t := \frac{1}{t + n_0 + 1} \text{ and} \\ Y_{t+1} &:= \left[\frac{W_{t+1} - \mathcal{N}_t\epsilon_t}{\psi_{t+1}}, \mathcal{N}_t - \psi_t \right]^T. \end{aligned} \quad (16)$$

The conditional expectation of Y_{t+1} with respect to the σ -algebra $\mathcal{G}_t := \sigma(\epsilon_s, \psi_s, s \leq t)$, is given by the following:

$$\begin{aligned} E[Y_{t+1}|\mathcal{G}_t] &= \bar{\mathbf{g}}(\epsilon_t, \psi_t) + [e_t, 0], \text{ with } \bar{\mathbf{g}} = [\bar{g}_\epsilon, \bar{g}_\psi], \text{ where,} \\ \bar{g}_\epsilon(\epsilon, \psi) &:= \frac{\beta\epsilon(1-\epsilon)(2q_\epsilon - 1)}{\psi}, \quad \bar{g}_\psi(\epsilon, \psi) = E[\mathcal{N}] - \psi, \\ e_t &:= E\left[\left(\frac{1}{\psi_{t+1}} - \frac{1}{\psi_t}\right)\left(W_{t+1} - \mathcal{N}_{t+1}\epsilon_t\right)\middle|\mathcal{G}_t\right], \text{ and,} \\ \beta &:= \left[(2b_n - 1)E[\mathcal{N}] + (2b_s - 1)E[\mathcal{S}]\right]. \end{aligned} \quad (17)$$

Stochastic approximation methods are well known techniques to study random trajectory $\{(\psi_t, \epsilon_t)\}$ given in (16) by approximating it with an ODE formed using the above conditional expectations:

$$\begin{aligned} \dot{\epsilon}_t &= \bar{g}_\epsilon(\epsilon, \psi), \text{ and,} \\ \dot{\psi} &= E[\mathcal{N}] - \psi \text{ for } \epsilon \in [0, 1], \psi > 0. \end{aligned} \quad (18)$$

In Theorem 2 (given later) we will prove that the above ODE indeed approximates (16), by showing that the difference term e_t would converge to 0 (see proof of Theorem 2). Further, the right-hand side of the ODE is not continuous, because of discrete-valued mapping $\epsilon \mapsto q_\epsilon$, which can take values only among 0, $(1 - \delta)$ and 1. We nevertheless prove the approximation result and derive further results using [14, Theorem 2.2, pp. 131]. Since $\bar{g}_\epsilon(\cdot)$ is only measurable in ϵ , the results cannot be applied directly. We provide the required justifications while proving Theorem 2.

Prior to that we study the function $\epsilon \mapsto q_\epsilon$, with an aim to analyze ODE (17); this study utilizes the asymptotic approximation derived in the previous section. The following results are required for analyzing the ODE in Corollary 1 (provided after Theorem 1).

Theorem 1 *Assume $w(1 + d) \geq v$ for the financial network.*

- a) *There exist an $0 \leq \bar{\epsilon} \leq 1$ such that, $q_\epsilon = 1 - \delta$ for all $\epsilon < \bar{\epsilon}$ and $q_\epsilon = 1$ for all $\epsilon \geq \bar{\epsilon}$.*

b) We have $\bar{\epsilon} < 1$ if and only if (with $w_{dv} := w(1+d) - v$)

$$\begin{aligned} \frac{w(r_b - d) - w_{dv}}{w(1 + 2\alpha\delta - \alpha)} &> \frac{(2u - r_s - r_b)}{2\alpha(1 - \delta)} \text{ or} \\ \frac{w(r_b - d) - w_{dv}}{w(1 + 2\alpha\delta - \alpha)} &> \frac{2(u - d)}{1 + \alpha} \text{ and } \frac{u - r_b}{u - d} < \alpha(1 - \delta). \end{aligned} \quad (19)$$

c) The $\bar{\epsilon}$ can be derived as zero of the equation:

$$\begin{aligned} \epsilon^2 + m_1\epsilon + m_2 = 0, \text{ with, } m_2 &= \frac{w\alpha\delta(u - r_b) + (2\alpha - 1)(1 - \delta)w_{dv}}{m_3} \\ m_1 &= \frac{w(u - r_b)(1 - \alpha + \alpha\delta) + w\alpha\delta(u - r_s)}{m_3} \\ &\quad + \frac{(1 - \delta)(w_{dv} - w(r_b - d)(2\alpha - 1))}{m_3}, \end{aligned}$$

where $m_3 = w(u - r_s)(1 - \alpha(1 - \delta)) - w(1 - \delta)(r_b - d)$. ■

Clearly, from ODE (18), the solution of ψ component is $\psi(t) = E[\mathcal{N}] + (\psi_0 - E[\mathcal{N}])e^{-t}$, with $\psi_0 = 1$. Further using Theorem 1, the ϵ -component of ODE (18) simplifies to:

$$\begin{aligned} \dot{\epsilon}_t &= \frac{\kappa\epsilon(1 - \epsilon)}{E[\mathcal{N}] + (\psi_0 - E[\mathcal{N}])e^{-t}} \text{ with,} \\ \kappa &= \begin{cases} \beta(1 - 2\delta) & \text{if } \epsilon_t \leq \bar{\epsilon} \\ \beta & \text{if } \epsilon_t \geq \bar{\epsilon}. \end{cases} \end{aligned} \quad (20)$$

This ODE has unique solution for any initial condition ϵ_0 :

$$\epsilon_t = \frac{\epsilon_0(E[\mathcal{N}]e^t + 1 - E[\mathcal{N}])^{\kappa/E[\mathcal{N}]}}{1 - \epsilon_0 + \epsilon_0(E[\mathcal{N}]e^t + 1 - E[\mathcal{N}])^{\kappa/E[\mathcal{N}]}}. \quad (21)$$

The above immediately implies the following result:

Corollary 1 [ODE Analysis] Assume initial condition $\epsilon_0 \in (0, 1)$. Then with $\bar{\epsilon}$ as in Theorem 1,

a) When $\beta > 0$ and $\delta > 1/2$, we have the following convergence of the unique ODE solution (as $t \rightarrow \infty$):

$$\epsilon_t \rightarrow 0 \text{ if } \epsilon_0 < \bar{\epsilon} \text{ and } \epsilon_t \rightarrow 1 \text{ if } \epsilon_0 \geq \bar{\epsilon}. \quad (22)$$

b) When $\beta < 0$, $\delta > 1/2$ and $\bar{\epsilon} \in \{0, 1\}$, then as $t \rightarrow \infty$

$$\epsilon_t \rightarrow 1 \text{ if } \bar{\epsilon} = 1 \text{ and } \epsilon_t \rightarrow 0 \text{ if } \bar{\epsilon} = 0.$$

c) When $\delta < 1/2$ $\epsilon_t \rightarrow 0$ if $\beta < 0$, else $\epsilon_t \rightarrow 1$ as $t \rightarrow \infty$. ■

From (14)-(16), $\epsilon_t \in [0, 1]$ for all t and all sample paths. The ψ component can be bounded a.s. by law of large numbers. By Corollary 1 the combined domain of attraction of the asymptotically stable attractors among $\{0, 1\}$ is $[0, 1]$ and so the dynamics trivially visits the combined domain infinitely often, under the assumptions of the Corollary. *We now have the main result (proof is in Appendix):*

Theorem 2 *Assume $w(1 + d) \geq v$. The sequence $\{\epsilon_t\}$ generated by (14)-(16) converges almost surely to limit points $\{\epsilon^*\}$ of the ODE (18), obtained in Corollary 1, under the respective conditions provided in the Corollary. ■*

Remarks We will show below that the limit points of Theorem 2 are ESS and thus the dynamics settles to a pure ESS ($\epsilon^* \in \{0, 1\}$), further when switching is predominant.

The settling point (ESS), depends upon network parameters and the initial proportion of agents. There is a possibility of the dynamics converging to 1, i.e., $\epsilon_t \rightarrow 1$, only when the conditions of Theorem 1.(b) are satisfied. In all other cases, the system converges to pure ESS with all risky agents.

Under the conditions of Theorem 1.(a), when $\epsilon > \bar{\epsilon}$ the financial network could have entered systemic risk regime (scenario in which majority of the agents have defaulted), if further $\epsilon > \bar{\epsilon}_2$ by Lemma 4 provided in Appendix. But the *rational agents averted such a catastrophic event by all of them choosing the risk-free strategy whenever $\beta > 0$; by Corollary 1, $\epsilon_t \rightarrow 1$ for such initial conditions. In fact we noticed that the system always avoids systemic risk regime even in test cases with $\beta < 0$, in Section 6 on numerical study; furthermore, this was true even when $w(1 + d) < v$, but we need the obvious condition $v < w(1 + u)$.*

Our theoretical study is under the assumption that agents are always able to repay the taxes/senior debt (even with downward return, $w(1 + d) > v$). One can easily extend this study for the case when $w(1 + d) < v < w(1 + u)$, we conjecture that (21) again represents the ODE solution. We have few such examples in Figure 1 where the ODE solution well matches the random trajectory.

Mixed Limit: Consider the case with $w(1 + d) \geq v$, $\beta < 0$ and $0 < \bar{\epsilon} < 1$. From (21), it is clear that ODE solution trajectory is increasing when below $\bar{\epsilon}$ and decreasing when above $\bar{\epsilon}$. Thus it wanders around $\bar{\epsilon}$, as $t \rightarrow \infty$. *We conjecture that the finance dynamics converges to this intermediate limit point.* We indeed illustrate the same using numerical examples in Section 6 (see Figure 1 and Table 1).

Finite Round Approximation: Using similar arguments as in Theorem 2, one can show that the sequence $\{\epsilon_t\}_{t \geq \bar{t}}$ can be approximated by ODE solution uniformly over any finite time horizon i.e., for any $T > 0$ (e.g., as in [12, Theorem 1]), as $\bar{t} \rightarrow \infty$. To be more precise, for any $k \geq 1$, we have that:

$$\epsilon_{l+k} \approx \frac{\epsilon_0 (E[\mathcal{N}]e^{t_{k,l}} + 1 - E[\mathcal{N}])^{\kappa/E[\mathcal{N}]}}{1 - \epsilon_0 + \epsilon_0 (E[\mathcal{N}]e^{t_{k,l}} + 1 - E[\mathcal{N}])^{\kappa/E[\mathcal{N}]}} \text{, with,} \quad (23)$$

$t_{k,l} = \sum_{j=l+1}^{l+k} \frac{1}{j+n_0+l}$, when $(l + n_0)$ is sufficiently large; thus one can estimate the approximate fraction of entities that adapt risk-free strategy after $l + k$ number of rounds. In

Section 6, we observe that the approximation is good for majority of cases, even for $l \approx 1$ and n_0 in few hundreds.

Average Dynamics: In our previous work [7], we consider the analysis with a large number of samples/observations; we would like to summarize relevant results of [7], with an aim to compare them with the dynamics of the current paper.

Any new agent at first contacts two random (uniformly sampled) agents of the previous round. If both the contacted agents belong to the same group, the new agent adapts the strategy of that group. When it contacts agents from both the groups *it investigates more before making a choice*; the new agent observes significant portion of the network, in that, it obtains a good estimate of the average utility of agents belonging to both the groups. It adapts the strategy of the group with maximum (estimated) average utility. Say it observes the average of each group with an error that is normally distributed with mean equal to the expected return $\phi_i(\epsilon) = E[R_i^k]$ with $k = 1, 2$ and variance proportional to the size of the group, i.e., it observes (here $\mathcal{U}(0, \sigma^2)$ is a zero mean Gaussian random variable with variance σ^2)

$\hat{\phi}_i(\epsilon) = \phi_i(\epsilon) + \mathcal{U}_i$ with $\mathcal{U}_1 \sim \mathcal{U}\left(0, \frac{1}{\bar{c}\epsilon}\right)$ and $\mathcal{U}_2 \sim \mathcal{U}\left(0, \frac{1}{\bar{c}(1-\epsilon)}\right)$, for some \bar{c} large. The dynamics converges to the following limit points (more details of the dynamics and proof can be found in [7]):

Lemma 2 Define $\bar{r}_r := u\delta + d(1 - \delta)$. Assume $\epsilon_0 \in (0, 1)$. There exists a $\bar{\delta} < 1$ (depends upon the instance of the problem) such that the following are valid for all $\delta \geq \bar{\delta}$:

- a) If $\bar{r}_r > r_b > r_s$ then $\epsilon_t \rightarrow 0$ almost surely.
- b) If $\phi_1(\epsilon) > \phi_2(\epsilon)$ for all ϵ then $\epsilon_t \rightarrow 1$ almost surely.
- c) When $r_b > \bar{r}_r > r_s$, and case (b) is negated there exists a unique zero ϵ^* of the equation $\phi_1(\epsilon) - \phi_2(\epsilon) = 0$ and

$$\epsilon_t \rightarrow \epsilon^* \text{ almost surely; further for } \delta \rightarrow 1, \epsilon^* \rightarrow \frac{r_b - \bar{r}_r}{\bar{r}_r - r_s}. \blacksquare$$

Remarks: From Theorem 2 and Corollary 1, under the corresponding hypothesis, the random dynamics always converge to either ‘risky’ or ‘risk-free’ agents, i.e., pure strategies (with $\beta > 0$). On the other hand, the average dynamics, as in Lemma 2, either converges to pure or mixed strategies further based on the parameters. Thus there is a big difference in the settling behaviour depending upon the number of (perfect) observations; *when agents observe sparsely, the network eventually settles to one of the two strategies, and if they observe more samples there is a possibility of the emergence of mixed limit*. On the other hand, even with sparse observations one can have mixed limit ($\bar{\epsilon}$ of Theorems 1-2) due to imperfect information (see also mixed ESS related discussions in Section 5).

5 Limit points and ESS

We will now show that the limit points of the ODE (18), which are also the (a.s.) limits of the replicator dynamics (by Theorem 2) are indeed evolutionary stable, *under predominant*

switching, i.e., when $\text{sign}(\beta) = \text{sign}(2b_s - 1)$. We will also show that the mixed limits are evolutionary stable.

The utility (a.k.a. fitness) of any agent is $+1$ if it encounters opposite group agent and if its return is better; it is -1 , if the opponent has better return. Thus the utility of agent using risk-free strategy, conditioned on $\epsilon_t = \epsilon$:

$$\begin{aligned} u_1(\epsilon) &= (1 - \epsilon) ((b_s q_\epsilon + (1 - b_s) q_\epsilon^c) - (b_s q_\epsilon^c + (1 - b_s) q_\epsilon)) \\ &= (1 - \epsilon)(2q_\epsilon - 1)(2b_s - 1), \text{ with, } q_\epsilon^c := 1 - q_\epsilon. \end{aligned} \quad (24)$$

Similarly that of the agent using risky-investments equals:

$$u_2(\epsilon) = -\epsilon(2q_\epsilon - 1)(2b_s - 1). \quad (25)$$

Mixed ESS:

Intuitively a strategy (chosen by majority) is called mixed ESS if a small fraction of mutants using some other strategy are eventually wiped out. We provide the precise definition directly in our finance network context and show that all our limits (18) are indeed mixed ESS.

For such definitions one needs to consider mixed strategies; ϵ is a mixed strategy, if an agent invests in risk-free or risky assets respectively with probabilities ϵ and $1 - \epsilon$. Each agent adapting a mixed strategy ϵ is equivalent to ϵ fraction of them investing in risk-free assets (see e.g., [15]); for large networks this equivalence is simply due to law of large numbers. Consider a mixture of population, where x (a small) fraction of mutants use ϵ while the remaining use $\bar{\epsilon}$ (intermediate limit in our case). With such a population mix, the fraction of agents investing in risk-free assets is given by, $\epsilon_x := x\epsilon + (1 - x)\bar{\epsilon}$. Let $u(\epsilon, \epsilon_x)$, $u(\bar{\epsilon}, \epsilon_x)$ respectively represent the utilities of the mutants and the other agents (non-mutants), against population mix, ϵ_x .

Definition 1 A strategy $\bar{\epsilon}$ is mixed ESS if for any $\epsilon \neq \bar{\epsilon}$, there exist $\bar{x} > 0$ such that $u(\epsilon, \epsilon_x) < u(\bar{\epsilon}, \epsilon_x)$ for all $x \leq \bar{x}$.

From (24) -(25) the utility of the mutant, i.e., the agent using mixed strategy ϵ , is given by:

$$\begin{aligned} u(\epsilon, \epsilon_x) &= \epsilon u_1(\epsilon_x) + (1 - \epsilon) u_2(\epsilon_x) \\ &= \epsilon(1 - \epsilon_x)(2q_{\epsilon_x} - 1)(2b_s - 1) - (1 - \epsilon)\epsilon_x(2q_{\epsilon_x} - 1)(2b_s - 1). \end{aligned}$$

One can compute $u(\bar{\epsilon}, \epsilon_x)$ similarly, and the difference in the two utilities:

$$u(\epsilon, \epsilon_x) - u(\bar{\epsilon}, \epsilon_x) = (\epsilon - \bar{\epsilon})(2q_{\epsilon_x} - 1)(2b_s - 1). \quad (26)$$

For predominant switching, the mixed limit is possible only when $\beta < 0$, i.e., if $2b_s - 1 < 0$. Further when $\epsilon < \bar{\epsilon}$, we have $\epsilon_x < \bar{\epsilon}$ for any $x \in (0, 1)$; thus from Lemma 3, $q_{\epsilon_x} = 1 - \delta$ and so RHS of (26) is negative. When $\epsilon > \bar{\epsilon}$, again from Lemma 3, $q_{\epsilon_x} = 1$ and the RHS is negative. Thus $\bar{\epsilon}$ is mixed ESS, with any \bar{x} .

Next consider pure limits $\epsilon^* \in \{0, 1\}$. We will indeed show that there are again Mixed ESS (stable against mutants using mixed strategies), which also implies stability against pure strategies. We again consider predominant switching.

From Theorem 1.(a), we have $\epsilon^* = 0$ with $\beta > 0$, (i.e., $(2b_s - 1) > 0$) and $\delta > 1/2$, only when $\bar{\epsilon} > 0$. Thus, for any mutant using (mixed) strategy $\epsilon > 0$, by Lemma 3, one can choose small enough \bar{x} such that $q_{\epsilon x} = 1 - \delta$ for all $x \leq \bar{x}$; basically we require $\epsilon \bar{x} < \bar{\epsilon}$. Then as in (26), for all such x ,

$$u(\epsilon, \epsilon x) - u(\epsilon^*, \epsilon x) = u(\epsilon, \epsilon x) - u(0, \epsilon x) = \epsilon(1 - 2\delta)(2b_s - 1) < 0.$$

Similarly when $\epsilon^* = 1$, the only possible mutant strategies are $\epsilon < 1$, again for small enough x we have $q_{\epsilon x + (1-x)} = 1$; the rest of the arguments are similar. One can prove other pure limits (e.g., with $\beta < 0$) are mixed ESS in a similar way.

When the dynamics are dominated by addition of new agents, then limit points need not be evolutionary stable. That is anticipated as the dynamics are not due to the existing agents; further practically one might have very few additions in comparison to the changes in strategies of the agents.

Evolutionary stable strategy for average dynamics: The ODE for average dynamics is given by (see [7])

$$\dot{\epsilon}_t = \epsilon(1 - \epsilon)(2g(\epsilon) - 1) \text{ where } g(\epsilon) := \int_{-\infty}^{(\phi_1(\epsilon) - \phi_2(\epsilon))\sqrt{\bar{c}\epsilon(1-\epsilon)}} e^{-x^2/2} \frac{dx}{\sqrt{2\pi}}.$$

In [7] we found the limit points of the ODE and omitted the details regarding them being evolutionary stable. These limit points are also summarized in Lemma 2. We will show below that these limit points are indeed evolutionary stable as in the above paragraphs, but now with the (approximate) expected return (10) being the utility of any agent.

Mixed ESS with $\epsilon^* = \frac{r_b - \bar{r}_r}{\bar{r}_r - r_s}$: We again consider a mixture of population, where x (a small) fraction of mutants use ϵ while the remaining use ϵ^* . From (10) the utility of the mutant, i.e., the agent using mixed strategy ϵ , is given by:

$$u(\epsilon, \epsilon_x) = \epsilon\phi_1(\epsilon_x) + (1 - \epsilon)\phi_2(\epsilon_x) \text{ where, } \epsilon_x = x\epsilon + (1 - x)\epsilon^*.$$

One can compute $u(\epsilon^*, \epsilon_x)$ similarly, and the difference in the two utilities:

$$u(\epsilon^*, \epsilon_x) - u(\epsilon, \epsilon_x) = (\epsilon^* - \epsilon)(\phi_1(\epsilon_x) - \phi_2(\epsilon_x)). \quad (27)$$

When $\epsilon < \epsilon^*$, we have $\epsilon_x < \epsilon^*$ for any $x \in (0, 1)$; thus from the proof of [7, Corollary 1, pp. 227], $\phi_1(\epsilon_x) > \phi_2(\epsilon_x)$ (when $\delta \geq \bar{\delta}$), and hence the RHS of (27) is positive. Similarly, the RHS is positive even when $\epsilon > \epsilon^*$. Hence $\epsilon^* = \frac{r_b - \bar{r}_r}{\bar{r}_r - r_s}$ is a mixed ESS under the conditions of Lemma 2(c).

Pure ESS with $\epsilon^* = 0$: For any mutant using (mixed) strategy $\epsilon > 0$, by Lemma 2(a) and (27), for all such x ,

$$u(\epsilon^*, \epsilon_x) - u(\epsilon, \epsilon_x) = \epsilon(\phi_2(\epsilon_x) - \phi_1(\epsilon_x)) > 0.$$

Therefore $\epsilon^* = 0$ is an ESS under the conditions of Lemma 2(a).

Pure ESS with $\epsilon^* = 1$: Similarly when $\epsilon^* = 1$, the only possible mutant strategies are $\epsilon < 1$, again by Lemma 2.(b) and (27), for all such x ,

$$u(\epsilon^*, \epsilon x) - u(\epsilon, \epsilon x) = (1 - \epsilon)(\phi_1(\epsilon x) - \phi_2(\epsilon x)) > 0.$$

Therefore $\epsilon^* = 1$ is an ESS under the conditions of Lemma 2(b).

Thus the replicator dynamics either settles to a pure strategy ESS or mixed ESS (see Lemma 2(c)), depending upon the parameters of the network; after a large number of rounds, either the fraction of agents following one of the strategies converges to one or zero or the system reaches a mixed ESS which balances the expected returns of the two groups.

6 Numerical Observations

This section reinforces our theoretical findings using the estimates obtained from Monte-Carlo (MC) simulations. For each run of the simulation: a) a sequence of binomial shocks, sequence of users joining, random sampling, etc. are realized; b) at each round of each run, the fixed point equations are solved using iterative methods to obtain the clearing vector; and c) at the end of each round the comparisons of random samples are made using the numerical estimates to continue the dynamics as in Section 4.

In Table 1 we consider the first example with asymptotic limits. Each run of the simulation is considered for $T = 1000$ number of rounds and with initial population size $n_0 = 300$. We set $E[\mathcal{N}_t] = 1$ and $E[\mathcal{S}_t] = 10$ while the rest of the details are in the table itself. The results are tabulated for various configurations. The configurations are specified in the first two columns, ϵ_{Th}^* is the theoretical limit obtained using ODE (20), and ϵ_{MC}^* is the MC based estimate. We also included $\bar{\epsilon}$ of Theorem 1. The first observation is that the theoretical ODE-limits well match the MC-based limits.

Configuration $(b_s, b_n, \alpha, \delta)$	ϵ_0	ϵ_{Th}^*	ϵ_{MC}^*	$\bar{\epsilon}$
(0.9, 0.9, 0.95, 0.85)	0.85	1	0.9866	0.8350
(0.9, 0.9, 0.95, 0.85)	0.75	0.0004	0.0008	0.8350
(0.4, 0.4, 0.95, 0.85)	0.8	0.8350	0.8367	0.8350
(0.9, 0.9, 0.95, 0.45)	0.6	1	0.9985	0.2233
(0.15, 0.15, 0.95, 0.45)	0.2	0.0749	0.0649	0.2233

Table 1: MC estimates versus ODE-limits: $v = 20, w = 70, (u, r_s, r_b, d) = (0.15, 0.1, 0.11, -0.6)$.

The first two rows consider a configuration with $\beta > 0$ and $\bar{\epsilon} < 1$. Results well corroborate with Corollary 1.(a); when initial $\epsilon_0 = 0.75 < \bar{\epsilon}$, the random trajectory converges to 0; the limit is 1 if $\epsilon_0 = 0.85$. The third row considers the same configuration, except for $b_n = b_s = 0.4$. With this partial information ($\beta < 0$), we have the conjectured mixed ESS: the trajectory wanders around $\bar{\epsilon}$. In fact we noticed similar convergence around $\bar{\epsilon}$ for many other cases.

In the last two rows, a case with large probability of economic shocks is considered ($\delta = 0.45$). Depending upon the quality of information ($\beta < 0$ or $\beta > 0$), the dynamics converge to a pure limit ($\epsilon_t \rightarrow 1$ or $\epsilon_t \rightarrow 0$ respectively). This reinforces the results of Corollary 1. (c).




Figure 1: Imperfect ($\beta < 0$) versus Perfect ($\beta > 0$) Information




Figure 2: Predominant switching with $E[S]=28$ and $E[N] = .2$

In figures 1-2 we illustrate the accuracy of finite horizon approximation by plotting ODE-based trajectories (23) and the corresponding MC trajectories for four configurations.

The two curves close-in as the number of rounds increase. This is true even for the case in Figure 1, where we have $w(1 + d) < v$, which is not covered by theory.

In the Figure 1, we have a case with a comparable fraction of new agents ($E[\mathcal{N}] = 20$ and $E[\mathcal{S}] = 28$). Here we compared the scenarios with and without perfect information. When $\beta > 0$ (red curves) the dynamics converge towards $\epsilon^* = 0$, i.e., all the agents settle to ‘risky’ strategy. With erroneous information, i.e., with $\beta < 0$, the dynamics (blue curves) settle to mixed fraction $\bar{\epsilon}$. In both the scenarios, dynamics avoid the systemic risk regime ($P_d = 1$ only for $\epsilon > \bar{\epsilon}_2$ and $\bar{\epsilon} \leq \bar{\epsilon}_2$, see Lemmas 3, 4).

In Figure 2 we consider the scenarios with perfect information and predominant switching; we observe that the dynamics settle to a pure ESS, i.e., $\epsilon^* = 0$ or $\epsilon^* = 1$, depending upon ϵ_0 .

7 Conclusions

We consider a financial network wherein the agents are interconnected via liability edges/connections. There are two types of agents; one group lends to others and invests in risk-free projects, while the second group borrows/lends and invests in risky ventures. These agents try to adapt their strategies based on their experiences and observations. Some new agents may also join the network. Thus we have a sequence of random networks evolving with time, where the strategic agents are changing their connections locally to improve their returns. We analyze such evolution and establish the emergence of evolutionary stable strategies. Towards this, we reduced the analysis of the evolution of this complex time-varying financial system to that of an appropriate ordinary differential equation (ODE); some recent results helped reduce the large dimensional random fixed point equations to simpler representations, which are instrumental in deriving the ODE.

Using the attractors of the resulting ODE, we showed that with perfect information, the replicator dynamics converges to one of the two pure evolutionary stable strategies, i.e., configurations with all ‘risky’ or all ‘risk-free’ agents. With imperfect information, dynamics can settle to a mixed limit. In all the cases, the dynamics averted ‘systemic risk event’, where most agents default. We also performed Monte-Carlo simulations to reaffirm the theoretical findings.

Author contributions

All authors contributed to the study conception and design. The first draft of the manuscript was written by Indrajit Saha, while, Veeraruna Kavitha helped in improving the manuscript. All authors read and approved the final manuscript.

Data availability

Data sharing not applicable to this article as no datasets were generated or analysed during the current study. We generated synthetic data which is included in the article itself.

8 Appendix

8.1 Details of Asymptotic Approximation

The clearing vector (5), defined using equations (2)-(5), can be viewed as the solution of random fixed point equations, which depend upon the realizations of the economic shocks $\{K_i\}_{i \in G_2}$ to the network. We obtain approximate clearing vector by applying the single group results of [8, Corollary 1 and Subsection 4.2] only to group G_2 . Towards this we consider a fictitious big node (like in [8]) and from each node $j \in G_2$ there is a dedicated fraction $(1 - c_\epsilon)$ with $c_\epsilon := \frac{\alpha + \alpha\epsilon}{\alpha + \epsilon}$ directed towards the fictitious node. This financial system is exactly similar to the graphical model described by [8, equations (1)-(6) and (40)-(41)], after the following mapping details:

$$\begin{aligned} G_i &= K_i, \eta_i^{bs} = 0 \text{ (a.s.)}, W_{j,i} = \frac{L_{ji}}{y} = \frac{I_{ji}c_\epsilon}{np_{ss}(1 - \epsilon)}, p_{c_1} = p_{c_2} = 0, \\ p_1 = p_2 &= p_{ss}, \text{ and } p_1^{sb} = p_2^{sb} = 1 - c_\epsilon. \end{aligned} \quad (28)$$

With the above mapping details the required assumptions of [8] are satisfied: assumption **B.1** is immediately satisfied (see (5)), assumption **B.3** is satisfied with $\sigma = 1$ and with any $0 \leq \varsigma < 1$ (as the fixed point equations do not depend upon x_b). The weight factors are as in [8, Subsection 4.2] and hence assumption **B.2** is not required. Finally, the assumption **B.4** is satisfied with $\rho = 1$. Hence by [8, Corollary 1] (as $0 < c_\epsilon < 1$ with $\epsilon \in (0, 1)$), the solution of the random fixed point equations (5) can be approximated using that corresponding to the limit system given in [8, Corollary 1]. Thus we have the convergence provided³ in equations (7)-(9) for any $\epsilon \in (0, 1)$ for G_2 (as the network size increases to infinity). This in turn provides the convergence results for G_1 .

Network with only risky agents ($\epsilon = 0$): The total amount lend by any agent to its neighbours equals $\approx w\alpha/(1 - \alpha)$, where the approximation is again accurate at limit. In a similar way the total amount borrowed by any agent also equals $\approx w\alpha/(1 - \alpha)$. Thus any agent invests $\approx w$ (their initial wealth) in risky assets. In this case the limit aggregate clearing vector (8) reduces to the following:

$$\bar{x}^\infty = E \left[\min \left\{ \left(K_i + \bar{x}^\infty - v \right)^+, y \right\} \right]. \quad (29)$$

When $w(1 + d) \geq v$ it is easy to observe that $\bar{x}^\infty = y$ is the unique solution of the above equation. Hence $P_d = 0$. Therefore $R_i^2 = K_i - v$ at limit for any i . For this case [8,

³One can partially justify similar approximation for systems with $\epsilon \in \{0, 1\}$ using [8, Theorem 1 and Subsection 4.2].

Corollary 1] is not applicable, however [8, Theorem 1] (applied to single group as in [8, subsection 4.2 1], partially justifies the above approximation.

Network with only risk-free agents ($\epsilon = 1$): On the other hand with $\epsilon = 1$, all are risk-free agents and they invest completely in risk-free assets. Thus the return of any agent $i \in G_1$ equals, $R_i^1 = w(1 + r_s) - v$.

8.2 Proofs

Proof of Lemma 1: We consider the following scenarios with $v < k_d$. The average clearing vector for the group G_2 agents satisfies (see (8)):

$$\begin{aligned} \bar{x}^\infty = & \min \left\{ \left(k_d + c_\epsilon \bar{x}^\infty - v \right)^+, y \right\} (1 - \delta) \\ & + \min \left\{ \left(k_u + c_\epsilon \bar{x}^\infty - v \right)^+, y \right\} \delta. \end{aligned} \quad (30)$$

Case 1: First consider the case when downward shock can be absorbed i.e., default probability is $P_d = 0$. If we have $k_d + yc_\epsilon - v \geq y$ then the average clearing vector $\bar{x}^\infty = y\delta + y(1 - \delta) = y$, and the above condition simplifies to the bound:

$$k_d - v + yc_\epsilon \geq y \implies c_\epsilon \geq \frac{y - w}{y}.$$

Case 2: Consider the case in which only the agents that receive shock will default, i.e., when $P_d = 1 - \delta$. The corresponding average clearing vector equals:

$$\bar{x}^\infty = y\delta + (w + c_\epsilon \bar{x}^\infty)(1 - \delta) \text{ and satisfies } k_d - v + c_\epsilon \bar{x}^\infty < y, k_u - v + c_\epsilon \bar{x}^\infty \geq y.$$

In this case the average clearing vector reduces to $\bar{x}^\infty = \frac{y\delta + w(1 - \delta)}{1 - c_\epsilon(1 - \delta)}$, and using the same in the bounds we have:

$$\frac{y - \bar{w}}{y - (1 - \delta)(\bar{w} - \underline{w})} \leq c_\epsilon < \frac{y - w}{y}.$$

Case 3 (Systemic-risk regime): Consider the case in which all the agents default i.e., when $P_d = 1$. In this we first calculate \bar{x}^∞ which is obtained by solving following fixed point equation: If we have $k_u - v + c_\epsilon \bar{x}^\infty < y$ then from (30) the average clearing vector reduces to:

$$\bar{x}^\infty = (k_d - v + c_\epsilon \bar{x}^\infty)(1 - \delta) + (k_u - v + c_\epsilon \bar{x}^\infty)\delta = \frac{E[W]}{1 - c_\epsilon}.$$

Substituting \bar{x}^∞ we have the required bound: $c_\epsilon < \frac{y - \bar{w}}{y - (1 - \delta)(\bar{w} - \underline{w})}$. ■

Proof of Theorem 1.(a): By Lemma 3, the mapping $\epsilon \mapsto q_\epsilon$ is monotone; there exists $\bar{\epsilon}$, with $\bar{\epsilon}_1 < \bar{\epsilon} \leq \bar{\epsilon}_2$, such that $q_\epsilon = 1 - \delta$ for all $\epsilon \leq \bar{\epsilon}_2$ and equals 1 for the rest; here $\bar{\epsilon}_1, \bar{\epsilon}_2$ are given by Lemma 4.

Proof of part (b) From Lemma 3, $\bar{\epsilon} < 1$ if and only if the conditions of part (b) are satisfied.

Proof of part (c) By the proof of Lemma 3, $g(\epsilon)$ definition, $\bar{\epsilon}$ equals the zero of the following (when condition (19) is satisfied) $\epsilon^2 + m_1\epsilon + m_2 = 0$, with, m_1, m_2 be an appropriate constants. ■

Lemma 3 *There exists a unique $\bar{\epsilon}$, with $\bar{\epsilon}_1 \leq \bar{\epsilon} \leq \bar{\epsilon}_2$ and with $\{\bar{\epsilon}_i\}$ as in Lemma 4, such that the $q_\epsilon := E[R^1 \geq R^2]$ satisfies the following threshold property:*

$$q_\epsilon = \begin{cases} 1 - \delta & \text{if } \epsilon < \bar{\epsilon} \\ 1 & \text{if } \epsilon \geq \bar{\epsilon}. \end{cases} \quad (31)$$

Further $\bar{\epsilon} < 1$ if and only if equation (19) is satisfied.

Proof: Case A: When $P_d = 0$ for some ϵ , the returns of the agents are given by the following (recall $r_b \geq r_s > d, w(1+d) \geq v$):

$$R^1(\epsilon) = \left(w\epsilon(1+r_s) + \frac{(1-\alpha)(1-\epsilon)}{(\alpha+\epsilon)}y - v \right)^+ \quad (32)$$

$$= (w\epsilon(1+r_s) + w(1-\epsilon)(1+r_b) - v)^+ = (w(1+r_b) - v + w\epsilon(r_s - r_b)), \text{ and}$$

$$R^2(\epsilon) = \left(K_i + \frac{\alpha(1+\epsilon)}{\alpha+\epsilon}y - v - y \right)^+ = (K_i - w\epsilon(1+r_b) - v)^+ \quad (33)$$

$$= \begin{cases} R_u^2 & \text{w.p. } \delta & \text{where } R_u^2 := w(1+u) - v + w\epsilon(u - r_b) \\ (R_d^2)^+ & \text{w.p. } 1 - \delta & \text{where } R_d^2 := w(1+d) - v + w\epsilon(d - r_b). \end{cases}$$

Hence with upward movement, $R^2 - R^1 = R_u^2 - R^1 = w(u - r_b) + w\epsilon(u - r_s) > 0$. It is also clear that $(R_d^2(\epsilon))^+ - R^1(\epsilon) < 0$. Thus $q_\epsilon = 1 - \delta$, when $P_d = 0$. By Lemma 4, this regime lasts for all ϵ satisfying $0 \leq \epsilon \leq \bar{\epsilon}_1$.

Case B: When $P_d = 1 - \delta$, by Lemma 4, we have $\bar{\epsilon}_1 < \epsilon \leq \bar{\epsilon}_2$. In this case we have $R_u^2 > 0$, but $R^1(\epsilon)$ can be positive or zero. Since $R_d^2 = 0$, we have that $q_\epsilon = P(R_1(\epsilon) \geq R_2(\epsilon)) \geq 1 - \delta$. Further in this case,

$$R^1(\epsilon) = \left(w\epsilon(1+r_s) + \frac{(1-\alpha)(1-\epsilon)}{(\alpha+\epsilon)}\bar{x}^\infty - v \right)^+, \text{ and,}$$

$$R_u^2 = \left(w(1+\epsilon)(1+u) + \frac{\alpha(1+\epsilon)}{\alpha+\epsilon}\bar{x}^\infty - v - y \right).$$

And hence

$$\begin{aligned} R_u^2(\epsilon) - R^1(\epsilon) &\leq w(1+u) + w\epsilon(u - r_s) - y + \bar{x}^\infty \left(\frac{2\alpha + \epsilon - 1}{\alpha + \epsilon} \right) \\ &= w(1+u) + w\epsilon(u - r_s) + (\bar{x}^\infty - y) - \bar{x}^\infty \left(\frac{1-\alpha}{\alpha+\epsilon} \right) \\ &= w(u - r_b) + w\epsilon(u - r_s) + (\bar{x}^\infty - y) \left(1 - \frac{1-\alpha}{\alpha+\epsilon} \right). \end{aligned} \quad (34)$$

If the upper bound on RHS of (34) is negative then clearly $R_u^2(\epsilon) < R^1(\epsilon)$. When the RHS is positive and $R^1(\epsilon) = 0$, then clearly $R_u^2(\epsilon) > R^1(\epsilon)$. On the other hand, when RHS

is positive and $R^1(\epsilon) > 0$, then the RHS is the exact value and not the upper bound, and hence again $R_u^2(\epsilon) > R^1(\epsilon)$. Thus in all, $R_u^2(\epsilon) > R^1(\epsilon)$ if and only if the RHS of (34) is positive.

Hence with $P_d = 1 - \delta$, we compute the following and derive the required analysis by checking the negative/positive sign of the RHS of (34). First observe from Lemma 1 that

$$(\bar{x}^\infty - y) \left(1 - \frac{1 - \alpha}{\alpha + \epsilon} \right) = (1 - \delta) \frac{w(1 + d) - v - w\epsilon(r_b - d)}{\epsilon + \alpha\delta - \alpha\epsilon(1 - \delta)} (2\alpha + \epsilon - 1) \quad (35)$$

and consider the following function constructed (basically the denominator in the above is positive and multiply it with remaining terms of the RHS) using the RHS of (34):

$$\begin{aligned} g(\epsilon) &:= w(u - r_b)(\epsilon + \alpha\delta - \alpha\epsilon(1 - \delta)) + w\epsilon(u - r_s)(\epsilon + \alpha\delta - \alpha\epsilon(1 - \delta)) \\ &\quad + (2\alpha - 1 + \epsilon)(1 - \delta)(w(1 + d) - v - w\epsilon(r_b - d)), \\ g'(\epsilon) &= w(u - r_b)(1 - \alpha(1 - \delta)) + w(u - r_s)(2\epsilon + \alpha\delta - 2\alpha\epsilon(1 - \delta)) \\ &\quad + (1 - \delta) \left(w(1 + d) - v - w\epsilon(r_b - d) - w(r_b - d)(2\alpha - 1 + \epsilon) \right), \\ g'' &= 2w(u - r_s)(1 - \alpha(1 - \delta)) - 2w(1 - \delta)(r_b - d), \\ g(\bar{\epsilon}_1) &:= w(u - r_b)(\bar{\epsilon}_1 + \alpha\delta - \alpha\bar{\epsilon}_1(1 - \delta)) + w\bar{\epsilon}_1(u - r_s)(\bar{\epsilon}_1 + \alpha\delta - \alpha\bar{\epsilon}_1(1 - \delta)) > 0, \\ g(1) &= w(1 + 2\alpha\delta - \alpha)(2u - r_s - r_b) + 2\alpha(1 - \delta)(w(1 + d) - v - w(r_b - d)). \end{aligned} \quad (36)$$

Observe that $g(\cdot)$ is concave in ϵ if $(u - r_s)(1 - \alpha(1 - \delta)) < (1 - \delta)(r_b - d)$, else it is convex in ϵ .

Sub-case 1: Consider the regime when $(u - r_s)(1 - \alpha(1 - \delta)) < (1 - \delta)(r_b - d)$. Observe that $g(\bar{\epsilon}_1) > 0$ (by definition of $\bar{\epsilon}_1$). By concavity of $g(\cdot)$, the function (and hence the RHS of (34)) can at maximum change the sign one time. Further when the RHS is zero, clearly $q_\epsilon = 1$. In other words there exists $\bar{\epsilon}$ with $\bar{\epsilon}_1 \leq \bar{\epsilon} \leq \bar{\epsilon}_2$, such that $q_\epsilon = 1 - \delta$ for all $\epsilon < \bar{\epsilon}$ and equals 1 when $\bar{\epsilon} \leq \epsilon < \bar{\epsilon}_2$.

Sub-case 2: Consider the regime $(u - r_s)(1 - \alpha(1 - \delta)) \geq (1 - \delta)(r_b - d)$. With this, it is easy to verify that $g(1) > 0$. With $g(1) > 0$ then we have $g'(\bar{\epsilon}_1) > 0$, because $2w(u - r_s)(1 - \alpha(1 - \delta)) + (1 - \alpha)(1 - 2\alpha(1 - \delta)) \geq 0$. Once again, $g(\bar{\epsilon}_1) > 0$. In this case due to convexity, $g(\epsilon)$ do not changes sign for all $\bar{\epsilon}_1 \leq \epsilon \leq \bar{\epsilon}_2$, and hence $q_\epsilon = 1 - \delta$ for all $\bar{\epsilon}_1 \leq \epsilon \leq \bar{\epsilon}_2$.

Case C: When $P_d = 1$, then $R^2 = 0$ a.s., while $R^1 \geq 0$, thus $q_\epsilon = 1$ and this is for all $\epsilon > \bar{\epsilon}_2$. Thus the threshold property. Further one can have $\bar{\epsilon} < 1$, either when $g(1) < 0$ (which also ensures $\bar{\epsilon}_1 < 1$) or when $\bar{\epsilon}_2 < 1$ and hence the result. ■

Lemma 4 Define $\bar{\epsilon}_1 := \frac{w(1+d)-v}{w(r_b-d)}$. There exists $\bar{\epsilon}_2 \geq \bar{\epsilon}_1$, which is strictly greater when $\bar{\epsilon}_1 < 1$ such that

$$P_d = \begin{cases} 0 & \text{if } 0 \leq \epsilon \leq \bar{\epsilon}_1 \\ 1 - \delta & \text{if } \bar{\epsilon}_1 < \epsilon \leq \bar{\epsilon}_2, \\ 1 & \text{if } \epsilon > \bar{\epsilon}_2. \end{cases} \quad (37)$$

Further $\bar{\epsilon}_2 < 1$ if and only if the second line of equation (19) is satisfied.

Proof: First consider the following function of ϵ , constructed using the bound a_1 and c_ϵ (the denominators are positive and then by making the denominator common in term $c_\epsilon - a_1$) of Lemma 1:

$$h(\epsilon) = \left(\alpha + \alpha\epsilon \right) y - \left(y - k_d + v \right) \left(\alpha + \epsilon \right) = (\alpha + \epsilon) \left(w(1+d) - v + w\epsilon(d - r_b) \right).$$

From Lemma 1, if $h(\epsilon) \geq 0$ then $P_d = 0$ and clearly this regime lasts for all ϵ , with $0 \leq \epsilon \leq \bar{\epsilon}_1$. Further more, beyond $\bar{\epsilon}_1$, $P_d > 0$.

Next consider the following function of ϵ , constructed similarly using the bound a_2 and c_ϵ of Lemma 1:

$$\begin{aligned} f(\epsilon) &= \left(\alpha + \alpha\epsilon \right) \left(y - (1 - \delta)w(1 + \epsilon)(u - d) \right) - \left(y - k_u + v \right) \left(\alpha + \epsilon \right) \\ &= (\alpha + \epsilon) \left(w(1+d) - v - w\epsilon(r_b - d) \right) + \left((\alpha + \alpha\epsilon)\delta + (1 - \alpha)\epsilon \right) w(u - d) \end{aligned} \quad (38)$$

From Lemma 1, if $f(\epsilon) \geq 0$ then $P_d \leq 1 - \delta$ and if $f(\epsilon) < 0$ then $P_d = 1$. Therefore it suffices to study this function. By (38), $f(0)$, $f(1)$ and its derivatives are given by:

$$\begin{aligned} f(0) &= \alpha \left(w(1+d) - v + w\delta(u - d) \right) > 0, \\ f(1) &= (1 + \alpha) \left((w(1+d) - v) - w(r_b - d) \right) + w \left(4\alpha\delta + 2(1 - \alpha) \right) (u - d), \\ f''(\epsilon) &= 2w \left(u - r_b - \alpha(u - d)(1 - \delta) \right) \text{ for all } \epsilon \left[\text{as } d - r_b = u - r_b - (u - d) \right], \text{ and,} \\ f'(0) &= w \left(\alpha(u - r_b) - (2\alpha(1 - \delta) - 1)(u - d) \right) + w(1+d) - v. \end{aligned}$$

It is clear that $f(0) > 0$. When the second derivative $f'' < 0$, the function is concave in ϵ , then clearly $f(\epsilon) \geq 0$ for all $\epsilon \leq \bar{\epsilon}_2$ (for some $0 < \bar{\epsilon}_2 \leq 1$) and $f(\epsilon) < 0$ for the rest, where $\bar{\epsilon}_2 < 1$ if and only if $f(1) < 0$.

Now consider the case with $f'' \geq 0$, and then f is convex (or linear). Under this condition, clearly the first derivative $f'(0) > 0$ (as $\alpha^2 - 2\alpha(1 - \delta) + 1 > 0$). Thus, $f(0) > 0$ implies $f(\epsilon) > 0$ for all $\epsilon \leq \bar{\epsilon}_2$ and so $P_d \leq (1 - \delta)$ for all $\epsilon \leq \bar{\epsilon}_2$, where $\bar{\epsilon}_2 = 1$.

In all, we have the existence of an $\bar{\epsilon}_2$ such that $f(\epsilon) \geq 0$ (and hence $P_d \leq 1 - \delta$) if and only if $\epsilon \leq \bar{\epsilon}_2$. Also observe from the second equality of (38) that, at $\epsilon = \bar{\epsilon}_1$, we have $f(\bar{\epsilon}_1) > 0$. Thus $\bar{\epsilon}_1 < \bar{\epsilon}_2$, whenever $\bar{\epsilon}_1 < 1$. Recall $P_d = 0$ if and only if $\epsilon \leq \bar{\epsilon}_1$. Hence we have (37).

Further $\bar{\epsilon}_2 < 1$ if and only if $f(1) < 0$ (and $f'' < 0$) which is equivalent to the second line of (19). ■

We first begin with some definitions.

Definition 2 Asymptotically stable (Attractor): A set A is said to be Asymptotically stable in the sense of Lyapunov, which we refer as attractor, if there exist a neighbourhood (called domain of attraction) starting in which the ODE trajectory converges to A as time progresses (e.g., [14]).

Definition 3 Equicontinuous in extended sense ([14]): Suppose that for each n , $h_n(\cdot)$ is on the \mathcal{R}^r -valued measurable function on $(-\infty, \infty)$ and $\{h_n(0)\}$ is bounded. Also suppose that for each T and $\varepsilon > 0$ there is a $\kappa > 0$ such that

$$\limsup_n \sup_{0 \leq t-s \leq \kappa, t \leq T} \|h_n(t) - h_n(s)\| \leq \varepsilon.$$

Then we say that $\{h_n(\cdot)\}$ is equicontinuous in the extended sense. By [14, Theorem 2.2], there exists a sub sequence that converges to some continuous limit function.

Proof of Theorem 2: We prove the result using [14, Theorem 2.2, pp. 131], as $\bar{g}_\varepsilon(\cdot)$ is only measurable. Towards this, we first need to prove (a.s.) equicontinuity in the extended sense of the following sequence of two-dimensional functions defined for each n (for any $t \geq 0$):

$$[\varepsilon^n(t), \psi^n(t)] := [\varepsilon_n, \psi_n] + \sum_{i=n}^{m(t_n+t)-1} \gamma_i Y_i, \quad m(t) := \max \left\{ n : \sum_{k=0}^{n-1} \gamma_k \leq t \right\}.$$

This proof goes through almost exactly as in the proof of [14, Theorem 2.1, pp. 127] and we follow exactly the same pattern. We begin with discussing some initial steps: i) the random vector Y_t depends on (ε_t, ψ_t) and $(W_{t+1}, \mathcal{N}_{t+1})$; ii) observe $\sup_t E|Y_t|^2 < \infty$, which is trivially true by law of large numbers (LLN); iii) clearly $\varepsilon_t \leq 1$ and $\psi_t \leq 1 + \bar{N}$ for all t and all sample paths; iv) we have $E[Y_t | \mathcal{G}_t] = \bar{g}(\varepsilon_t, \psi_t) + [e_t, 0]$ where,

$$e_t := E \left[\left(\frac{1}{\psi_{t+1}} - \frac{1}{\psi_t} \right) (W_{t+1} - \mathcal{N}_{t+1} \varepsilon_t) \middle| \mathcal{G}_t \right],$$

and v) the projection term in [14], $Z_t \equiv 0$.

We further require to handle the difference term e_t . We will now show that the error term e_t converges to zero in the limit and continue with the rest of the proof thereafter. Towards this, observe that $|\psi_t - \mathcal{N}_t| \leq \psi_t + \bar{N}$ where \bar{N} is such that $P(\mathcal{N} \leq \bar{N}) = 1$. Thus from (15),

$$\left| \frac{1}{\psi_{t+1}} - \frac{1}{\psi_t} \right| = \left| \frac{\gamma_t (\psi_t - \mathcal{N}_t)}{\psi_t \psi_{t+1}} \right| \leq \frac{\gamma_t}{\psi_{t+1}} \left(1 + \frac{\bar{N}}{\psi_t} \right). \quad (39)$$

Consider the (almost sure) sample paths in which $\psi_t \rightarrow E[\mathcal{N}]$ by LLN, as one can rewrite:

$$\psi_t = \frac{n(t)}{t + n_0} = \frac{n_0 + \sum_{k=1}^t \mathcal{N}_k}{t + n_0}. \quad (40)$$

For such sample paths (i.e., almost surely), $\psi_t \geq \varepsilon$ for all t for some appropriate $\varepsilon > 0$ and hence using (39)

$$|e_t| \leq \frac{\bar{s} \gamma_t}{\varepsilon} E \left[\left| W_{t+1} - \mathcal{N}_{t+1} \varepsilon_t \right| \middle| \mathcal{G}_t \right] \leq \frac{(2E[\mathcal{N}] + E[\mathcal{S}]) \bar{s}}{\varepsilon} \gamma_t \quad \text{a.s., where } \bar{s} := 1 + \frac{\bar{N}}{\varepsilon}$$

and recall $W_{t+1} = \xi_t + \Xi_1(t) - \Xi_2(t)$.

Thus $e_t \rightarrow 0$ and $\sum_t |e_t| \gamma_t < \infty$ a.s. The update equation (starting at n and for any $t \geq 0$) can be written as below (as in [14]):

$$\begin{aligned} [\epsilon^n(t), \psi^n(t)] &= [\epsilon_n, \psi_n] + \int_0^t \bar{\mathbf{g}}(\epsilon^n(s), \psi^n(s)) ds + S^n(t) + \rho^n(t) \\ \text{with, } \rho^n(t) &:= \sum_{i=n}^{m(t+t_n)-1} \gamma_i \bar{\mathbf{g}}(\epsilon_n, \psi_n) - \int_0^t \bar{\mathbf{g}}(\epsilon^n(s), \psi^n(s)) ds, \text{ and} \\ S^n(t) &:= \sum_{i=n}^{m(t+t_n)-1} \gamma_i \Delta S_i = S_{m(t+t_n)-1} - S_n, \Delta S_n = Y_n - \bar{\mathbf{g}}(\epsilon_n, \psi_n). \end{aligned}$$

Observe in the above that $\epsilon^n(t_k) = \epsilon_k$ for any $k > n$, where $t_k := \sum_{i < k} \gamma_i$. By LLN $\psi_k \rightarrow E[\mathcal{N}]$ in almost all sample paths and the idea is to show the equicontinuity of the functions $\{\epsilon^n(\cdot), \psi^n(\cdot)\}_n$ for those sample paths; this guarantees the existence of limit function (along a sub-sequence) as in [14], and proceeding further as in [14] we can show that the limit function satisfies the ODE (17).

The arguments required to show the equicontinuity in extended sense are exactly as in [14], because of the following: (i) $\{S_n\}$ is a martingale and using well known Martingale inequality (as in [14]), $\lim_m P\{\sup_{j \geq m} |S_j - S_m| \geq \mu\} = 0$, for any μ , because

$$\sum_i \gamma_i^2 < \infty \text{ and } E[\Delta S_i \Delta S_j] = E[E[\Delta S_i \Delta S_j | \mathcal{G}_j]] = 0, \forall j < i;$$

(ii) recall $\psi_k \rightarrow E[\mathcal{N}]$ as $k \rightarrow \infty$ for chosen sample paths; there exists a $C_g, \bar{k} < \infty$ such that $|\bar{g}_\epsilon(\epsilon_k, \psi_k)| \leq |\beta|/\psi_k < C_g$, $|E[\mathcal{N}] - \psi_k| \leq C_g$ and hence $\sup_t |\rho^k(t)| \leq 2C_g \gamma_k$ for all $k \geq \bar{k}$; (iii) the sequence $\{[\epsilon^n(0), \psi^n(0)]\}_{n \geq \bar{k}} = \{[\epsilon_n, \psi_n]\}_{n \geq \bar{k}}$ is bounded a.s. by $[1, 1 + \bar{\mathcal{N}}]$; and (iv) finally for any $t \geq t'$:

$$\begin{aligned} &\left| \int_0^t \bar{g}_\epsilon(\epsilon^n(s), \psi^n(s)) ds - \int_0^{t'} \bar{g}_\epsilon(\epsilon^n(s), \psi^n(s)) ds \right| \\ &\leq (t - t') \sup_{\epsilon \in [0, 1], k \geq n} |\bar{g}_\epsilon(\epsilon, \psi_k)| = C_g(t - t'). \end{aligned}$$

In Corollary 1 we have identified the attractors of (17), and showed that the combined domain of attraction is the whole $[0, 1] \times [0, \bar{C}]$ (for any $\bar{C} < \infty$). Choose \bar{C} such that the dynamics visits $[0, 1] \times [0, \bar{C}]$ infinitely often (possible because $P(\mathcal{N}_t < \bar{\mathcal{N}}) = 1$) and hence converges (a.s.) to one of the limit points of Theorem 1 and β by [14, Theorem 2.2, pp. 131]. ■

References

- [1] Acemoglu, Daron, Asuman Ozdaglar, and Alireza Tahbaz-Salehi. "Systemic risk and stability in financial networks." *American Economic Review* 105, no. 2 (2015): 564-608.

- [2] Allen, Franklin, and Douglas Gale. "Financial contagion." *Journal of political economy* 108, no. 1 (2000): 1-33.
- [3] Eisenberg, Larry, and Thomas H. Noe. "Systemic risk in financial systems." *Management Science* 47, no. 2 (2001): 236-249.
- [4] Kavitha, Veeraruna, Indrajit Saha, and Sandeep Juneja. "Random Fixed Points, Limits and Systemic risk." In *2018 IEEE Conference on Decision and Control (CDC)*, pp. 5813-5819. IEEE, 2018.
- [5] Yang, Ke, Kun Yue, Hong Wu, Jin Li, and Weiyi Liu. "Evolutionary analysis and computing of the financial safety net." In *International Workshop on Multi-disciplinary Trends in Artificial Intelligence*, pp. 255-267. Springer, Cham, 2016.
- [6] Li, Honggang, Chensheng Wu, and Mingyu Yuan. "An evolutionary game model of financial markets with heterogeneous players." *Procedia Computer Science* 17 (2013): 958-964.
- [7] Saha, Indrajit, and Veeraruna Kavitha. "Financial replicator dynamics: emergence of systemic-risk-averting strategies." In *International Conference on Network Games, Control and Optimization*, pp. 211-228. Springer, Cham, 2021.
- [8] Saha, Indrajit, and Veeraruna Kavitha. "Random fixed points, systemic risk and resilience of heterogeneous financial network." *arXiv preprint arXiv:2112.03598* (2021).
- [9] Smith, J. M. P. G. R., and George R. Price. "The logic of animal conflict." *Nature* 246, no. 5427 (1973): 15-18.
- [10] Tembine, Hamidou, Eitan Altman, Rachid El-Azouzi, and Yezekael Hayel. "Evolutionary games in wireless networks." *IEEE Transactions on Systems, Man, and Cybernetics, Part B (Cybernetics)* 40, no. 3 (2009): 634-646.
- [11] Miekisz, Jacek. "Evolutionary game theory and population dynamics." In *Multiscale problems in the life sciences*, pp. 269-316. Springer, Berlin, Heidelberg, 2008.
- [12] Singh, Vartika, Khushboo Agarwal, and Veeraruna Kavitha. "Evolutionary Vaccination Games with premature vaccines to combat ongoing deadly pandemic." In *EAI International Conference on Performance Evaluation Methodologies and Tools*, pp. 185-206. Springer, Cham, 2021.
- [13] Glasserman, Paul, and H. Peyton Young. "How likely is contagion in financial networks?." *Journal of Banking & Finance* 50 (2015): 383-399.
- [14] Kushner, Harold, and G. George Yin. *Stochastic approximation and recursive algorithms and applications*. Vol. 35. Springer Science & Business Media, 2003.
- [15] Easley, David, and Jon Kleinberg. *Networks, crowds, and markets: Reasoning about a highly connected world*. Cambridge university press, 2010.