

# RELAXED WYTHOFF HAS ALL BEATTY SOLUTIONS

JON KAY & GEREMÍAS POLANCO

ABSTRACT. In this paper, we modify a generalization of Wythoff’s game due to Fraenkel to accommodate non-monotonic settings. In doing so, we construct a class of Wythoff games covering the class of all Beatty sequences. In the appendix, we show that if a solution set for one of Fraenkel’s generalized games is known to be Beatty, then this generalized Fraenkel game is equivalent to a  $t$ -Wythoff game.

## 1. SHORT HISTORICAL INTRODUCTION

One of the most influential games in Combinatorial Game Theory is Nim, which many believe originated in China under the translated name “picking stones” [8]. Nim became popular in modern mathematics when Charles Leonard Bouton (1869–1922) from Harvard University published a complete solution of the game in the *Annals of Mathematics* in 1901 [2]. This event marked the birth of combinatorial game theory, a subject greatly influenced by the Sprague-Grundy theory that was independently discovered in 1935 by the German mathematician Roland Percival Sprague (1894–1967) [16], and in 1939 by the English mathematician and statistician Patrick Michael Grundy (1917–1959) [7]. The mathematical foundation of the subject was presented by the English mathematician John Horton Conway (1937–2020) in the book “On Numbers and Games” (1976) [10]. Conway can be considered the creator of Combinatorial Game Theory. This creatorship is shared and further established by Conway and his coauthors: American mathematician Elwyn Ralph Berlekamp (1940–2019) (a sharp individual who with a Ph.D. in electrical engineering, became a research mathematician while working at Bell Labs [13]), and the English mathematician Richard Kenneth Guy (1916–2020). They published the triply coauthored series: “Winnings Ways for your Mathematical Plays” starting in 1982 [14].

Just a few years after the appearance of Nim, in 1907, the Dutch mathematician Willem Abraham Wythoff, in the paper “A modification of the game of Nim” [17], published a complete solution to another subtraction game that since then has been called Wythoff’s game.

Wythoff’s game starts with two piles, labeled  $A$  and  $B$ , of finitely many items. Two players alternate, and at each turn the current player may either

- (1) remove any amount of items from pile  $A$ ,
- (2) remove any amount of items from pile  $B$ ,
- (3) or remove an equal amount of items from piles  $A$  and  $B$  simultaneously.

The player who leaves both piles empty wins. The differences between Wythoff’s game and Nim are that in Nim there may be more than two piles, and players are permitted to remove items from only one pile at a time. It is for the second of these two reasons that the third

---

*Date:* August 2, 2022.

move listed above will frequently be called the *Wythoff move*. The first and second moves will sometimes be called *Nim moves*.

From this definition arises a complete set of *winning positions*: game positions for which one player may force the other player to lose no matter how the other player progresses through the game. The strategic game positions are presented as follows:

the pair  $(a, b)$  is a winning position if and only if  $(a, b) \in \{([n\phi], [n\phi^2]) \mid n \in \mathbb{N}\}$ ,

where  $[\cdot]$  denotes the floor function,  $\phi = (1 + \sqrt{5})/2$ , and pairs denote the count of items in piles  $A$  and  $B$  in increasing order. Taking the left coordinate of these pairs yields a sequence of integers  $\{[n\phi]\}$  called the Beatty sequence of slope  $\phi$  and similarly for the right coordinate we obtain the Beatty sequence of slope  $\phi^2$ . It is no coincidence that  $1/\phi + 1/\phi^2 = 1$ , for Lord Rayleigh discovered in 1894 this condition dictates that the sequences  $\{[n\phi]\}$  and  $\{[n\phi^2]\}$  exactly cover the positive integers [15]. Beatty sequences have a rich literature in connection with a vast number of topics in number theory and combinatorics (see for instance, [1], and [12] and the references therein).

In 1959, Connell [3] defined a parametric family of  $t$ -Wythoff games, defining an alternate Wythoff move in which a player who removes  $a$  items from pile  $A$  and  $b$  items from pile  $B$  may do so only if  $|b - a| < t$  for some fixed positive integer  $t$ . The winning positions of this game are given by the Beatty sequences with slopes  $\alpha$  and  $\beta$  where

$$(1) \quad \alpha = \frac{2 - t + \sqrt{t^2 + 4}}{2} \quad \frac{1}{\alpha} + \frac{1}{\beta} = 1.$$

Note that  $\alpha = \phi$  is the special case of  $t = 1$ , inducing the original Wythoff game. We like to call this parametric family of irrational  $\alpha$ 's the *Wythoff golden family* of irrationals because they generalize the golden ratio and give solutions to the  $t$ -Wythoff game. It is easy to show that irrationals in the Wythoff golden family have a fractional part that is invariant under passing to the Beatty conjugate. This has important implications about the paper we refer to next.

Another central figure in the theory of Wythoff-like games is the Israeli mathematician Aviezri S. Fraenkel. He is a prolific author who has made significant contributions to combinatorial game theory, and has a skill in studying games related to Wythoff's. A quick search of his publications reveals more than ten mentions of Wythoff's game in the titles, and more than two dozen mentions and/or treatments of the game in the body of his papers. Aware that the mentioning of these facts does not do justice to his contributions, we iterate one of his generalizations of Wythoff's game in this paper.

In Fraenkel's 2004 paper, "New Games Related to Old and New Sequences," he defines a broader class of Wythoff games, again altering only the Wythoff move. In these games, a player who removes  $a$  items from pile  $A$  and  $b$  items from pile  $B$  may do so only if  $|b - a|$  is bounded by a supplied function  $f$  of the current game state. Setting  $f \equiv 1$  or  $f \equiv t$  are the special cases of the original Wythoff and  $t$ -Wythoff, respectively. Fraenkel computes the winning positions of an arbitrary game (subject to certain hypotheses on the constraint function  $f$ ) in a masterful way, appealing to the minimum excluded function fundamental to the theory of computation. To make this precise, we repeat the theorem in context.

**Definition 1** (Fraenkel's General two-pile subtraction game). *A general two-pile subtraction game is played with standard Nim moves, and a Wythoff move from  $(x_0, y_0)$  to  $(x_1, y_1)$  is*

valid if and only if the following inequality is satisfied

$$(2) \quad |(y_0 - y_1) - (x_0 - x_1)| < f(x_1, y_1, x_0)$$

for a supplied constraint function  $f$  which has the properties stated in the following theorem.

**Theorem 1** (Fraenkel's Master Theorem). *Let  $S = \{(a_k, b_k)\}_{k=0}^{\infty}$  where  $a_0 = b_0 = 0$  and*

$$(3) \quad a_{n+1} = \text{mex}\{a_k, b_k \mid k < n + 1\}$$

$$(4) \quad b_{n+1} = f(a_n, b_n, a_{n+1}) + b_n + a_{n+1} - a_n$$

*If  $f$  is positive, monotone, and semi-additive, then  $S$  is the set of  $P$ -positions of a general 2-pile subtraction game with constraint function  $f$ .*

**Remark 1.** *In this theorem,  $P$ -positions are what we refer to as winning positions, and we will sometimes use the terms interchangeably.*

As expected, the winning positions are complementary sequences, and the Master Theorem gives sufficient conditions on the function  $f$  for which an appropriate two-pile subtraction game has its winning positions given by the above recurrence. If the winning positions are *a priori* known to be Beatty sequences, then the monotonicity hypothesis is necessary, which is proven in the second appendix of this paper. The third appendix provides motivation for an appropriate modification of Fraenkel's game that we call the *Relaxed Wythoff*.

## 2. STATEMENT OF THE MAIN THEOREM AND OUTLINE OF THE PAPER

Relaxed Wythoff is a modification of the Fraenkel game which we now define. To perform the variation, we modify the Wythoff move of a general two-pile subtraction game by removing the absolute value symbol in equation 2 (this is equation 8 in Frankel's paper [6]) as follows:

**Definition 2** (Relaxed Wythoff). *A two-pile subtraction game is played with standard Nim moves, and a Wythoff move from  $(x_0, y_0)$  to  $(x_1, y_1)$  is valid if and only if the following inequality is satisfied*

$$(5) \quad (y_0 - y_1) - (x_0 - x_1) < f(x_1, y_1, x_0)$$

*where  $f$  is a non-negative function satisfying  $f(0, 0, 1) \neq 0$  and  $x_0 \leq y_0$ .*

From this modification we obtain the main result of this paper:

**Theorem 2.** *Relaxed Wythoff surjects onto the set of complementary Beatty sequences. In other words, given any pair of complementary Beatty sequences  $A$  and  $B$ , there exists a Relaxed Wythoff game whose winning positions are given by the pairs  $(a_n, b_n)$  where  $a_n \in A$  and  $b_n \in B$ , for all integers  $n \geq 0$ .*

**Remark 2.** *In the second appendix it is shown that if the winning positions arising in Fraenkel's Master theorem are complementary Beatty sequences, it is necessary that the constraint function be constant. The first appendix is devoted to studying a condition equivalent to the constraint function being constant in a Beatty setting, which is used in the next theorem.*

Before going any further, we want to say a few words about two games in the literature that accomplish similar goals of this paper, namely, producing winning positions that are *general* complementary Beatty sequences, not necessarily within the Wythoff golden family. An *invariant* game is one “for which each allowed move is independent of the position it is played from” [4]. A game which is not invariant is called a variant game. A class of invariant games is defined in [9], and it is elaborated in a follow up paper [5] that the computational complexity depends heavily on the size of the piles, making it hard to play the game by hand. For this reason, in the follow up paper, the authors define a class of games with simpler rules.

We want to highlight that our approach sacrifices invariance, but the game we present is a natural generalization of the general two pile game that closely resembles the original Wythoff game, (except obviously for the absolute value symbol in the game constraint). Moreover, the game is easy to play by hand.

The rest of the paper is organized as follows. In section 3, we discuss the motivation for our approach and demonstrate its limitation. In section 4, we remedy this limitation by defining a new class of games with a modified Wythoff move. In section 5 we demonstrate the main result of this paper, which is that this class of games has winning positions in the form of any desired pair of complementary Beatty sequences. Concluding remarks in section 6 are given. The first appendix contains a result which is used to study the behavior of a function related to Fraenkel’s Master Theorem. The second appendix proves that the general two-pile subtraction games defined by Fraenkel have almost no solutions which are complementary Beatty sequences, ultimately showing that no novel complementary Beatty sequences arise from these types of games, except for the ones in the Wythoff golden family. The final appendix is a general account of the ideas and methodology of this paper.

### 3. FRAENKEL’S MASTER THEOREM DOES NOT SURJECT IN A BEATTY SETTING

In the second appendix, the following theorem is proven by playing the game by hand inductively. The proof we give here provides a hint for finding a general framework to find Wythoff moves by studying a certain second difference as in the first appendix.

**Theorem 3** (Nonsurjectivity). *The complementary Beatty sequences arising in Fraenkel’s Master Theorem are those in the Wythoff golden family only.*

*Proof.* Suppose a function  $f$  is supplied to the theorem such that the resulting winning positions are complementary Beatty sequences

$$(6) \quad a_n = [\alpha n] \quad b_n = [\beta n] \quad \forall n$$

where  $1/\alpha + 1/\beta = 1$ . Inspecting equation 4 in theorem 1 shows that we can perform the inversion

$$(7) \quad f(a_n, b_n, a_{n+1}) = (b_{n+1} - a_{n+1}) - (b_n - a_n)$$

$$(8) \quad = (b_{n+1} - b_n) - (a_{n+1} - a_n).$$

Utilizing the integer and fractional part, we can rewrite

$$(9) \quad b_{n+1} = [\beta(n+1)] = [([\beta] + \{\beta\})(n+1)] = [[\beta](n+1) + \{\beta\}(n+1)]$$

$$(10) \quad = [\beta](n+1) + [\{\beta\}(n+1)].$$

Then the first difference becomes

$$(11) \quad [\beta(n+1)] - [\beta n] = ([\beta](n+1) - [\{\beta\}(n+1)]) - ([\beta]n - [\{\beta\}n])$$

$$(12) \quad = [\beta] + g_{1/\{\beta\}}(n)$$

where  $g_\rho(n) = [\rho(n+1)] - [\rho n]$  for real  $\rho$ . The supplied constraint function becomes

$$(13) \quad f = ([\beta] + g_{1/\{\beta\}}(n)) - ([\alpha] + g_{1/\{\alpha\}}(n))$$

$$(14) \quad = g_{1/\{\beta\}}(n) - g_{1/\{\alpha\}}(n) + [\beta] - [\alpha].$$

The part of this sum which varies is solely the difference

$$(15) \quad d_\alpha(n) := g_{1/\{\beta\}}(n) - g_{1/\{\alpha\}}(n).$$

In the first appendix it is shown that this difference  $d_\alpha$  is constant if and only if there exists a positive integer  $t$  such that

$$(16) \quad \alpha = \frac{2 - t + \sqrt{t^2 + 4}}{2}.$$

This means the Master Theorem accounts for a countable family of games with Beatty solutions, namely, those in the Wythoff golden family only. Therefore, almost every pair of complementary Beatty sequences does not arise from the theorem.  $\square$

#### 4. RELAXED WYTHOFF

Relaxed Wythoff is a modification of the general Fraenkel game which we now define. To perform the variation, we modify the Wythoff move of a general two-pile subtraction game by removing the absolute value symbol in equation 2 (this is equation 8 in Frankel's paper [6]) as follows:

**Definition 3** (Relaxed Wythoff). *A two-pile subtraction game is played with standard Nim moves, and a Wythoff move from  $(x_0, y_0)$  to  $(x_1, y_1)$  is valid if and only if the following inequality is satisfied*

$$(17) \quad (y_0 - y_1) - (x_0 - x_1) < f(x_1, y_1, x_0)$$

where  $f$  is a non-negative function satisfying  $f(0, 0, 1) \neq 0$  and  $x_0 \leq y_0$ .

The removal of the absolute value symbol is *equivalent* to imposing a certain growth condition on the winning positions, namely that any winning position  $(a, b)$  satisfies

$$(18) \quad b - a > \max(b_k - a_k)$$

where the maximum is taken over the winning positions smaller than  $(a, b)$ . The derivation of this coercive estimate is elaborated in the third appendix. By not taking the absolute value, we are able to relax two hypotheses on the constraint function as elaborated in appendix two, namely that of monotonicity (equivalent to constancy in a Beatty setting) and with it sub-additivity. The final assumption that the initial value of the constraint function is non-zero guarantees that the left and right coordinates of a fixed non-trivial winning position never coincide.

This results in a game for which we can make different looking moves. The removal from pile  $A$  is allowed to be *indefinitely more* than the removal from pile  $B$ . Note that we are working under convention that pile  $A$  is smaller than pile  $B$ .

To characterize the solutions in a computable fashion, we introduce the following lemma.

**Lemma 1** (Double Mex Lemma). *If a two-pile subtraction game has a set of winning positions which exactly cover the natural numbers, then they may be reconstructed with the recurrence*

$$(19) \quad (\widetilde{a}_0, \widetilde{b}_0) := (0, 0)$$

$$(20) \quad \widetilde{a}_{n+1} := \text{mex}\{a_k, b_k \mid k < n + 1\}$$

$$(21) \quad \widetilde{b}_{n+1} := \text{mex}\{b \mid (\widetilde{a}_{n+1}, b) \text{ can move to some } (a_k, b_k)\}$$

*Proof.* Suppose the winning positions of a subtraction game on two piles are known to equal

$$(22) \quad P = \{(a_n, b_n)\}_{n=0}^{\infty}.$$

The game finishes at  $(0, 0)$ , so  $a_0 = b_0 = 0$ . The winning positions cover the natural numbers, so we can discern that  $a_1 = 1$ . Now proceed by induction.

The first reconstructed winning position is computed by the mex:

$$(23) \quad \widetilde{a}_1 = \text{mex}\{0\} = 1$$

$$(24) \quad \widetilde{b}_1 = \text{mex}\{b \mid (1, b) \text{ can move to some } (a_k, b_k)\}$$

It is clear that  $\widetilde{a}_1 = a_1$ , since they both equal one. It follows from the uniqueness of a winning strategy that  $\widetilde{b}_1 = b_1$ . The inductive step is performed similarly.  $\square$

This lemma applies to any two-pile subtraction game whose  $P$  positions exactly cover the natural numbers, because we made reference only to  $P$  positions in the above proof. Moreover, the mex in both coordinates indicates the tacit assumption that the winning positions *exactly* cover the natural numbers. We are now free to remove the absolute value symbol from Fraenkel's game and utilize the lemma to determine the winning positions of Relaxed Wythoff. We first prove the following:

**Lemma 2.** *Let  $f$  be a function satisfying the hypotheses indicated by the definition of the Relaxed Wythoff game. If the game is played with  $f$  as its constraint function, then the winning positions of this game exactly cover the natural numbers.*

*Proof.* We first prove the existence of a winning strategy by induction with base case  $(a_0, b_0) = (0, 0)$ . Suppose the winning positions have been found up to  $(a_{n-1}, b_{n-1})$ . The next winning position can be found by first evaluating the mex

$$(25) \quad a_n = \text{mex}\{a_k, b_k \mid k < n\}$$

to obtain the left-coordinate; this method is indicated by the Nim moves. Now we need to find  $\widetilde{b}$  such that a player cannot move  $(a_n, \widetilde{b})$  to one of the known winning positions  $(a_k, b_k)$ ,  $k < n$ ; otherwise, the pair would be a losing position.

Consider the set  $S = \{f(a_k, b_k, a_n) \mid k < n\}$  and evaluate  $s = \max S$ . Because  $S$  is finite, the plane

$$(26) \quad z - y + x = s + a_n \quad (x, y) = (a_k, b_k) \quad k < n$$

encloses finitely many points below it, and thus infinitely many positive integers  $z$  satisfy

$$(27) \quad z - y + x \geq s + a_n \quad (x, y) = (a_k, b_k) \quad k < n.$$

Equivalently, we can see that

$$(28) \quad s \leq z - y - (a_n - x) = z - b_k - (a_n - a_k) \quad k < n.$$

Setting  $\tilde{b}_n$  to be the minimum of all such  $z$ , the above equation implies

$$(29) \quad \tilde{b}_n - b_k - (a_n - a_k) \geq f(a_k, b_k, a_n) \quad \text{for all } k < n$$

and thus  $(a_n, \tilde{b}_n)$  cannot move to one of the known winning positions, indicating that this pair is itself also a winning position.

Now we prove the winning positions exactly cover the natural numbers. Notice that any natural number which never appears in the right coordinate of a winning position automatically will lie in the left coordinate of some winning position because the Nim type moves are resolved by putting a mex in the the left coordinates. Therefore it is sufficient to show that no pair of non-trivial winning positions  $(a, b)$  satisfy  $a = b$ .

Suppose the winning positions are determined up to  $(a_{n-1}, b_{n-1})$ . An inductive hypothesis asserts that  $a_k \neq b_k$  for all  $k < n$ , so let us make a step to show that the next winning position  $(a_n, b_n)$  satisfies  $a_n \neq b_n$ . Suppose  $(a_n, b_n)$  has no move to a lower winning position  $(a_k, b_k)$ . Then

$$(30) \quad (b_n - b_n) - (a_n - a_k) = (b_n - a_n) - (b_k - a_k) \geq f(a_k, b_k, a_n) \quad \text{for all } k < n.$$

In case  $b_n = a_n$ , a contradiction arises immediately, since  $-(b_k - a_k)$  is negative and  $f$  is non-negative. Therefore,  $a_n \neq b_n$ , and we can tell that the winning positions exactly cover the natural numbers.  $\square$

**Theorem 4** (Winning Positions of Relaxed Wythoff). *Let  $f$  be a function satisfying the hypotheses indicated by the Relaxed Wythoff game. If the game is played with  $f$  as its constraint function, then the winning positions of this game are given by the recurrence:*

$$(31) \quad (a_0, b_0) := (0, 0)$$

$$(32) \quad a_{n+1} := \text{mex}\{a_k, b_k \mid k < n + 1\}$$

$$(33) \quad b_{n+1} := a_{n+1} + f(a_n, b_n, a_{n+1}) + b_n - a_n$$

*Proof.* Let  $f$  be a function as described. Suppose the winning positions are known up to  $(a_n, b_n)$ . The next winning position will then be computed with the lemma:

$$(34) \quad a_{n+1} = \text{mex}\{a_k, b_k \mid k < n\}$$

$$(35) \quad b_{n+1} = \text{mex}\{b \mid (a_n, b) \text{ can move to some } (a_k, b_k)\}$$

If  $(a_{n+1}, b)$  can move to some  $(a_k, b_k)$ , then we know

$$(36) \quad (b - b_k) - (a_{n+1} - a_k) < f(a_k, b_k, a_{n+1}).$$

By the definition of the mex operator,  $b_{n+1}$  is not one of these  $b$ 's, and thus it must satisfy the opposite of this inequality, namely

$$(37) \quad b_{n+1} \geq a_{n+1} + f(a_k, b_k, a_{n+1}) + b_k - a_k \quad \forall k < n + 1$$

Since the inequality holds for all  $k$ , we may as well set  $k = n$  and then resolve the inequality to an equality by a minimality argument:

$$(38) \quad b_{n+1} = a_{n+1} + f(a_n, b_n, a_{n+1}) + b_n - a_n,$$

completing the proof.  $\square$

## 5. RELAXED WYTHOFF IS SURJECTIVE IN A BEATTY SETTING

For the main result, it is now demonstrated that the class of winning positions arising from gameplay under a certain class of functions is in correspondence with the set of complementary Beatty sequences.

**Theorem 5.** *Relaxed Wythoff surjects onto the set of complementary Beatty sequences. In other words, given any pair of complementary sequences  $A$  and  $B$  there is a function  $f$  satisfying the hypotheses indicated by the Relaxed Wythoff game such that if the game is played with constraint function  $f$ , then the winning positions are given by  $(a_n, b_n)$  where  $a_n \in A$  and  $b_n \in B$ .*

*Proof.* Defining  $A = \{[n\alpha]\}$  and  $B = \{[n\beta]\}$ , where irrationals  $1 < \alpha < 2 < \beta$  satisfy  $1/\alpha + 1/\beta = 1$ , gives a fully general setting. As in the non-surjectivity proof of the Master Theorem, the appropriate constraint function ought to be

$$(39) \quad f(a_k, b_k, [n\alpha]) = ([n\beta] - [(n-1)\beta]) - ([n\alpha] - [(n-1)\alpha]) \quad k < n$$

Because we have relaxed so many conditions above, all we need to do is prove that this function is non-negative and that  $f(0, 0, 1) \neq 0$ .

Rewriting again as in equation 14, the constraint function becomes

$$(40) \quad f(\cdot, \cdot, [n\alpha]) = g_{1/\{\beta\}}(n) - g_{1/\{\alpha\}}(n) + [\beta] - [\alpha] = d_\alpha(n) + [\beta] - [\alpha]$$

The functions  $g_{1/\{\beta\}}$  and  $g_{1/\{\alpha\}}$  are indicator functions, as elaborated in the first appendix, so it is evident that their difference is bounded below by  $-1$ . Moreover, we have set  $1 < \alpha < 2$ , so its conjugate  $\beta$  is subject to the bound  $\beta \geq 2$ . Therefore,  $[\beta] - [\alpha] \geq 1$  and thus the function  $f$  is always non-negative. To verify that  $f(0, 0, 1) \neq 0$ , just remember that  $[\beta] - [\alpha] \geq 1$  evaluate:

$$(41) \quad f(0, 0, 1) = ([\beta] - [0\beta]) - ([\alpha] - [0\alpha]) = [\beta] - [\alpha] \geq 1.$$

Therefore, the hypotheses on  $f$  are satisfied, so the winning positions induced by the function may now be evaluated using theorem 4 and it is seen directly that the winning positions are the complementary Beatty sequences with lesser slope  $\alpha$ .  $\square$

## 6. CONCLUSION

Wythoff's game and its successors have an intimate connection to Beatty sequences, explored first by Wythoff and later by other authors (see for instance [4], [6], [9]). Here, we defined a new game and revealed structure behind these fascinating sequences. The primary mathematical content is the double mex lemma, an identification of a necessary and sufficient condition for Fraenkel's Master Theorem to behave in a Beatty setting, and the analysis of the  $d_\alpha$  function which finds an equivalent condition. Once the monotonicity hypothesis was identified as limiting the range of the theorem, we were able to evade this limitation by restructuring the Wythoff move, allowing us to prove a version of the master theorem for Relaxed Wythoff, demonstrating a relation to the complete class of complementary Beatty sequences. We note that in [4] and [9] an invariant version of this game is played. The present paper gives a non invariant and simple approach that is naturally connected with the original Wythoff's game and with Frankel's generalization of Wythoff.

## 7. APPENDIX: CHARACTERIZING THE MONOTONICITY OF $d_\alpha$

In this appendix, we present a result that is used in section 3. Let  $1 < \alpha < 2 < \beta$  satisfy  $1/\alpha + 1/\beta = 1$  and define

$$(42) \quad d_\alpha(n) := g_{1/\{\beta\}}(n) - g_{1/\{\alpha\}}(n)$$

This function bifurcates as  $\alpha$  varies as follows:

**Lemma 3.** *If  $d_\alpha$  is monotonic, then  $d_\alpha \equiv 0$ . Otherwise,  $d_\alpha$  oscillates.*

*Proof.* Suppose  $d_\alpha$  is monotonic. Let  $X = \{k/\{\alpha\}\}_{k=0}^\infty$  and  $Y = \{k/\{\beta\}\}_{k=0}^\infty$ . From Chapter 9 of [1], recall that

$$(43) \quad g_{1/\{\alpha\}}(n) = \begin{cases} 1 & n \in X \\ 0 & n \notin X \end{cases} \quad g_{1/\{\beta\}}(n) = \begin{cases} 1 & n \in Y \\ 0 & n \notin Y \end{cases}$$

We expand the difference of the indicator functions to obtain:

$$(44) \quad d_\alpha(n) = \begin{cases} 0 & n \in (X \cap Y) \cup (X^c \cap Y^c) \\ 1 & n \in X^c \cap Y \\ -1 & n \in X \cap Y^c \end{cases}$$

We finish the proof by ruling out the last two cases. These cases arise when  $X^c \cap Y$  or  $X \cap Y^c$  are nonempty. Since the sets  $X$  and  $Y$  are Beatty sequences, if one of these intersections is non-empty then the intersection actually has infinite size, a classical result which can be found in [11]. This shows that  $d_\alpha$  oscillates between at least two of the three values, violating monotonicity. Therefore, the intersections defining the second two values of the trichotomy are empty, indicating that  $d_\alpha \equiv 0$ .  $\square$

**Remark 3.** *The oscillation referred to in this theorem is aperiodic, as is typical in the study of Beatty sequences and characteristic words.*

We can also compute the value  $\alpha$  if we know the function  $d_\alpha$  is constant.

**Lemma 4.** *If  $d_\alpha$  is monotonic, then  $\alpha$  is in the Wythoff golden family.*

*Proof.* Adopt the notation from the trichotomy equation 44. The previous lemma shows  $X^c \cap Y = X \cap Y^c = \emptyset$ , so that  $X = Y$ . Therefore, we shall have  $\{\alpha\} = \{\beta\}$ . Some quick algebra shows  $\alpha$  is in the Wythoff golden family. To demonstrate, select  $t$  such that  $[\beta] = t + 1$ . Because  $\{\alpha\} = \alpha - [\alpha] = \alpha - 1$  and  $\{\beta\} = \beta - [\beta]$ , it follows that

$$(45) \quad \alpha - [\alpha] = \beta - [\beta],$$

and thus we obtain

$$(46) \quad \alpha - 1 = \frac{1}{1 - \frac{1}{\alpha}} - (t + 1).$$

From this we see that

$$(47) \quad \alpha = \frac{2 - t + \sqrt{t^2 + 4}}{2}.$$

Thus  $\alpha$  is in the Wythoff golden family with the parameter  $t = [\beta] - [\alpha]$ .  $\square$

## 8. APPENDIX II: NECESSITY OF CONSTANCY IN A BEATTY SETTING

The following theorem characterizes the Beatty winning positions arising from generalized two-pile subtraction games as in definition 1.

**Theorem 6.** *If the winning positions from a generalized two-pile subtraction game defined above are known to be complementary Beatty sequences, then the sequence with smaller slope belongs to the Wythoff golden family.*

*Proof.* Let  $f$  be a positive non-constant monotone function. Suppose the winning positions of a generalized two-pile subtraction game under the constraint function  $f$  are complementary Beatty sequences

$$(48) \quad \{([n\alpha'], [n\beta'])\} \quad \text{with} \quad \frac{1}{\alpha'} + \frac{1}{\beta'} = 1.$$

We will use the double mex lemma to determine the winning positions of the game in definition 1 under the constraint function  $f$  first in the case that  $f$  increases and second in the case that  $f$  decreases, both times finding a contradiction by inspecting the difference of successive left or right values of the computed winning positions.

Select the least  $n > 0$  such that the difference  $f(n+1) - f(n) = d$  is positive. By minimality, there exists  $t > 0$  such that  $f \equiv t$  up to  $n$  and the winning positions up to  $n$  are initial segments of the complementary Beatty sequences with slopes  $1 < \alpha < 2 < \beta$  where

$$(49) \quad \alpha = \frac{2-t+\sqrt{t^2+4}}{2} \quad \frac{1}{\alpha} + \frac{1}{\beta} = 1.$$

Write the first  $n$  winning positions.

$$(50) \quad \begin{array}{cccc} 0 & 0 & 0 & 0 \\ [\alpha'] & [\beta'] & [\alpha] & [\beta] \\ \vdots & \vdots & \vdots & \vdots \\ [n\alpha'] & [n\beta'] & [n\alpha] & [n\beta] \end{array} =$$

Now we can find the next winning position by the double mex lemma which gives the first coordinate of the winning position

$$(51) \quad a = [(n+1)\alpha]$$

and verbally describes the second coordinate of the winning position

$$(52) \quad b = \text{mex}\{b \mid (a, b) \text{ can move to some } (a_k, b_k)\}.$$

From the structure of complementary Beatty sequences with lesser slope in the Wythoff golden family, we know that

$$(53) \quad b_k - a_k = kt.$$

To compute this  $b$ , note that  $(a, b)$  has no move to the previous pairs. Therefore  $b$  is the least  $b$  satisfying

$$(54) \quad |(b-a) - (b_k - a_k)| \geq t + d \quad \forall k = 1, 2, \dots$$

Rearranging the inequality shows

$$(55) \quad (b - a) - kt \geq k + d \iff b \geq [(n + 1)\alpha] + kt + d$$

$$(56) \quad \iff b \geq [(n + 1)\beta] + d$$

so that  $b = [(n + 1)\beta] + d$ , leaving another row of winning positions:

$$(57) \quad \begin{array}{cccc} 0 & 0 & 0 & 0 \\ [\alpha'] & [\beta'] & [\alpha] & [\beta] \\ \vdots & \vdots & \vdots & \vdots \\ [n\alpha'] & [n\beta'] & [n\alpha] & [n\beta] \\ [(n + 1)\alpha'] & [(n + 1)\beta'] & [(n + 1)\alpha] & [(n + 1)\beta] + d \end{array} .$$

From the last row in equation (57) we can see

$$(58) \quad d = [(n + 1)\beta'] - [(n + 1)\beta].$$

Because  $[n\beta] = [n\beta']$  we can also see that

$$(59) \quad \beta' \in \left[ \frac{[n\beta]}{n}, \frac{[n\beta] + 1}{n} \right)$$

from which it follows that

$$(60) \quad (n + 1)\beta' \geq \frac{[n\beta](n + 1)}{n} = [n\beta] + \frac{[n\beta]}{n}$$

Expand

$$(61) \quad [n\beta] = [n([\beta] + \{\beta\})] = n[\beta] + [n\{\beta\}] = n[\beta] + n\{\beta\} - \{n\{\beta\}\} = n\beta - \{n\{\beta\}\}$$

leaving us with

$$(62) \quad (n + 1)\beta' \geq [n\beta] + \beta - \frac{\{n\{\beta\}\}}{n}.$$

Therefore depending on  $n$  we can see that

$$(63) \quad [(n + 1)\beta'] \in \{[n\beta] + [\beta] - 1, [n\beta] + [\beta]\}$$

leading us to finally determine  $d$  in terms of  $\beta$ :

$$(64) \quad d = [n\beta] + [\beta] - [(n + 1)\beta] \quad \text{or} \quad d = [n\beta] + [\beta] - [(n + 1)\beta] - 1.$$

We then obtain

$$(65) \quad [n\beta] = [n([\beta] + \{\beta\})] = n[\beta] + [n\{\beta\}]$$

$$(66) \quad [(n + 1)\beta] = [(n + 1)([\beta] + \{\beta\})] = (n + 1)[\beta] + [(n + 1)\{\beta\}],$$

and thus we have

$$(67) \quad d = [n\{\beta\}] - [(n + 1)\{\beta\}] \quad \text{or} \quad d = [n\{\beta\}] - [(n + 1)\{\beta\}] - 1.$$

It is clear that  $(n + 1)\{\beta\} > n\{\beta\}$ , so  $[(n + 1)\{\beta\}] \geq [n\{\beta\}]$ , in which case  $d$  is not positive, contradicting our hypothesis.

For decreasing  $f$ , select the least  $r > 0$  such that  $f(r) - f(r + 1) = d$ . This assumption forces  $f(r) = t \geq 1$  since  $f(r + 1) \geq 0$ . We will determine some winning positions similar to the previous case. The first few winning positions arise as the initial segments of known

Beatty sequences with lesser slope in the golden irrational family. The next winning position has left coordinate  $a = \lfloor (n+1)\alpha \rfloor$  and the right coordinate is given verbally

$$(68) \quad b = \text{mex}\{b \mid (a, b) \text{ can move to some } (a_k, b_k)\}.$$

If  $f(r+1) = 0$ , then  $b = a$  because the Wythoff move is blocked by the strict inequality against zero. The other case is that  $f(r+1) = t - d > 0$ . Note that  $t \geq 2$  at this point because we have already examined the case  $t = 1$  and  $d = 1$ . We will now argue that the next winning position is  $(a, b) = (a, a+1)$ . Suppose  $(a, b) \rightarrow (a_k, b_k)$  is a valid move. Then

$$(69) \quad |(b - b_k) - (a - a_k)| < t - d$$

Evaluating this absolute value is not hard since we figured out some details already:

$$(70) \quad |(b - b_k) - (a - a_k)| = |(b - a) - (b_k - a_k)| = |1 - kt| = kt - 1$$

Thus we can figure out  $k$ :

$$(71) \quad kt - 1 < t - d \implies kt < t - d + 1 \implies k < (t - d + 1)/t = 1 - (d - 1)/t < 1$$

We might now believe  $k = 0$ , but this move is easy to disprove by the same argument that  $(1, 2)$  does not move to  $(0, 0)$  in the classical Wythoff game. Therefore the next winning position is  $(a, a+1)$  which indicates the winning positions are Beatty if and only if  $a = 1$ , a reduction to the classical Wythoff game. Therefore,  $f$  does not decrease.

We have shown that the supplied constraint function neither increases nor decreases when the winning positions are complementary Beatty sequences which means that the lesser slope lies in the Wythoff golden family as in equation 49.  $\square$

### 9. APPENDIX III: DERIVING THE MODIFICATION

To see where the modification should come from, study the winning positions arising from a generalized two-pile subtraction game by applying the wildly useful lemma

$$(72) \quad a_{n+1} = \text{mex}\{a_i, b_i \mid i < n + 1\}$$

$$(73) \quad b_{n+1} = \text{mex}\{b \mid |(b - b_i) - (a_{n+1} - a_i)| < f(-)\}$$

The action of the absolute value symbol is

$$(74) \quad |(b - b_i) - (a_{n+1} - a_i)| = \begin{cases} (b - b_i) - (a_{n+1} - a_i) & b - a_{n+1} > \max(b_i - a_i) \\ (a_{n+1} - a_i) - (b - b_i) & \text{otherwise} \end{cases}$$

The first case is the case which generates Beatty sequences, because the winning positions are monotonic in both coordinates, which is equivalent to the stated growth condition. Therefore, we release the difference from the absolute value symbol when we define Relaxed Wythoff.

## 10. ACKNOWLEDGMENTS

This work was completed under the care and supervision of several institutions and advisors. Gratefully, we acknowledge Hampshire College and The Five College Consortium for providing the authors with a place to meet each other and explore mathematics in an experimental fashion. Amherst College and Smith College provided new academic homes for the second author during turmoil at Hampshire College and afterwards. Once the first author completed his studies at Hampshire College, he was given a host of new ideas at The University of California, Santa Cruz, particularly from Dr. François Monard. The first author also thanks The Ross Program and Dr. Jim Fowler for providing a motivating and hospitable work environment.

## REFERENCES

- [1] J-P. Allouche and J. Shallit. *Automatic Sequences*. Cambridge University Press, 2003.
- [2] C. L. Bouton. Nim, A Game with a Complete Mathematical Theory. *Annals of Mathematics*, 3(1/4):35–39, 1901–1902.
- [3] I. G. Connell. A Generalization of Wythoff’s Game\*. *Canadian Mathematical Bulletin*, 2(3):181–190.
- [4] E. Duchêne and M. Rigo. Invariant games. *Theoretical Computer Science*, 411(34–36):3169–3180, 2010.
- [5] M. Fisher and U. Larsson. Chromatic Nim finds a game for your solution. *Games of No Chance 5*. MSRI Publications, 70:321–330. Cambridge University Press, 2019.
- [6] A. S. Fraenkel. New Games Related to Old and New Sequences. *Advances in Computer Games*. The International Federation for Information Processing, 135:367–382, 2003.
- [7] L. S. Goddard. “Patrick Michael Grundy.” *Journal of the London Mathematical Society*, s1-35:377–379, 1960.
- [8] A. H. Jorgensen. Context and Driving Forces in the Development of the Early Computer Game Nimbi. *IEEE Annals of the History of Computing*, 31(3):44–53, 2009.
- [9] U. Larsson, P. Hegarty, and A. S. Fraenkel. Invariant and dual subtraction games resolving the Duchêne–Rigo conjecture. *Theoretical Computer Science*, 412(8–10):729–735, 2011.
- [10] J. Lurie. “J. H. Conway. *On numbers and games*.” *The Journal of Symbolic Logic*, 63(4):1602–1604, 1998.
- [11] I. Niven. Diophantine Approximations. *Interscience Tracts in Pure and Applied Mathematics*, vol. 14. Interscience Publishers, 1963.
- [12] G. Polanco. The logarithm of irrational numbers and Beatty sequences. *Acta Arithmetica*, 179:101–123, 2017.
- [13] R. Sanders. “Elwyn Berlekamp, game theorist and coding pioneer, dies at 78.” *Berkeley News*. The University of California, Berkeley, 2019.
- [14] L. Schneider. “Winning Ways for your Mathematical Plays.” *The Mathematics Teacher; Reston*, 95(2):158, 2002.
- [15] I. J. Schoenberg. The location of frets on a guitar and theorem of Lord Rayleigh. In *Mathematical Time Exposures*, page 40. The Mathematical Association of America, 1982.
- [16] R. Sprague. Über mathematische Kampfspiele. *Tôhoku Mathematical Journal*, 41:438–444, 1935.
- [17] W. A. Wythoff. A Modification of the game of Nim. *Nieuw Archiefvoor Wiskunde* 7(2):199–202, 1907.