

ON THE GENUS OF PROJECTIVE CURVES NOT CONTAINED IN HYPERSURFACES OF GIVEN DEGREE

VINCENZO DI GENNARO

ABSTRACT. Fix integers $r \geq 4$ and $i \geq 2$ (for $r = 4$ assume $i \geq 3$). Assuming that the rational number s defined by the equation $\binom{i+1}{2}s + (i+1) = \binom{r+i}{i}$ is an integer, we prove an upper bound for the genus of a reduced and irreducible complex projective curve in \mathbb{P}^r , of degree $d \gg s$, not contained in hypersurfaces of degree $\leq i$. It turns out that this bound coincides with the Castelnuovo's bound for a curve of degree d in \mathbb{P}^{s+1} . We prove that the bound is sharp if and only if there exists an integral surface $S \subset \mathbb{P}^r$ of degree s , not contained in hypersurfaces of degree $\leq i$. Such a surface, if existing, is necessarily the isomorphic projection of a rational normal scroll surface of degree s in \mathbb{P}^{s+1} . The existence of such a surface S is known for $i = 2$ and $i = 3$. It follows that, when $i = 2$ or $i = 3$, the bound is sharp, and the extremal curves are isomorphic projection in \mathbb{P}^r of Castelnuovo's curves of degree d in \mathbb{P}^{s+1} .

Keywords: Projective curve. Castelnuovo-Halphen Theory. Quadric and cubic hypersurfaces. Projection of a rational normal scroll surface. Maximal rank.

MSC2010: Primary 14N15; Secondary 14N25; 14M05; 14J26; 14J70

1. INTRODUCTION

The aim of this note is to improve [3, Proposition 1.3 and Proposition 1.4], and to simplify the proof of [3, Theorem 1.2]. In fact, we are going to prove the following Theorem 1.1. We refer to Remark 1.2 below for some comments on the claim.

Theorem 1.1. *Fix integers $r \geq 4$ and $i \geq 2$ (for $r = 4$ assume $i \geq 3$). Assume that the rational number s defined by the equation*

$$(1) \quad \binom{i+1}{2}s + (i+1) = \binom{r+i}{i}$$

is an integer. Let $C \subset \mathbb{P}^r$ be a reduced and irreducible complex curve, of degree d and arithmetic genus $p_a(C)$, not contained in a hypersurface of degree $\leq i$. Assume $d \gg s$, and divide $d - 1 = ms + \epsilon$, $0 \leq \epsilon \leq s - 1$. Then

$$(2) \quad p_a(C) \leq \binom{m}{2}s + m\epsilon.$$

Moreover, this bound is sharp if and only if there exists an integral surface in \mathbb{P}^r of degree s , not contained in hypersurfaces of degree $\leq i$. In this case, every extremal curve D is not a.C.M., and it is contained in a flag $S \subset T \subset \mathbb{P}^r$, where S is a surface of degree s , uniquely determined by D , not contained in hypersurfaces of degree i , and T is a hypersurface of \mathbb{P}^r of degree $i + 1$. Furthermore, S is the isomorphic projection in \mathbb{P}^r of a rational normal scroll surface S' of degree s in

\mathbb{P}^{s+1} , and, via this isomorphism, D corresponds to a Castelnuovo's curve of S' of degree d .

Remark 1.2. (i) For instance, if $i = 2$, the number s defined by the equation (1) is an integer if and only if r is not divisible by 3 [1, p. 484, line 12 from below]. If $i = 3$, the number s is an integer if and only if $r = 1, 2, 9, 10, 11, 18, 19, 27, 29$ modulo 36. When $r = 4$, the number s is an integer if and only if $i = 1, 2, 5, 10$ modulo 12. When $r = 5$, the number s is an integer if and only if $i = 1, 2, 7, 11, 13, 17, 22, 23, 26, 31, 37, 38, 41, 43, 46, 47, 53, 58$ modulo 60.

(ii) For the case $r = 4$ and $i = 2$, we refer to [3, Theorem 1.1].

(iii) The bound appearing in (2) is the Castelnuovo's bound for a non-degenerate integral curve of degree d in \mathbb{P}^{s+1} [4, p. 87, Theorem (3.7)].

(iv) Recall that $h^0(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(i)) = \binom{r+i}{i}$, and that, if $S' \subseteq \mathbb{P}^{s+1}$ is a rational normal scroll surface of degree σ , then $h^0(S', \mathcal{O}_{S'}(i)) = \binom{i+1}{2}\sigma + (i+1)$. Moreover, notice that for an integral surface in \mathbb{P}^r ($r \geq 4$) of degree σ , not contained in a hypersurface of degree $\leq i$, one has $\binom{i+1}{2}\sigma + (i+1) \geq \binom{r+i}{i}$ [3, (11)]. When the number s defined by (1) is not an integer, then one may define s differently as the minimal integer σ such that $\binom{i+1}{2}\sigma + (i+1) \geq \binom{r+i}{i}$ (compare with [3, (11)]). For instance, when $i = 2$, we have $3s + 3 = \binom{r+2}{2}$ if r is not divisible by 3, and $3s + 1 = \binom{r+2}{2}$ otherwise. More generally, divide

$$\binom{r+i}{i} - (i+1) = \alpha \binom{i+1}{2} + \beta, \quad 0 \leq \beta \leq \binom{i+1}{2} - 1.$$

Then one has $s = \alpha$ if $\beta = 0$, and $s = \alpha + 1$ if $\beta > 0$. We hope to give some information in the case $\beta > 0$ in a forthcoming paper.

(v) Let S be an integral surface in \mathbb{P}^r , not contained in hypersurfaces of degree $\leq i$, of degree s given by the equation (1). In view of Lemma 2.1 (see below), S is the isomorphic projection in \mathbb{P}^r of a rational normal scroll surface $S' \subset \mathbb{P}^{s+1}$ of degree s . Therefore, since $h^0(\mathbb{P}^r, \mathcal{I}_S(i)) = 0$, the restriction map $h^0(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(i)) \rightarrow h^0(S, \mathcal{O}_S(i))$ is bijective (compare with (iv) above). It follows that S is of maximal rank (compare with [1, p. 482, line 14 from below]). We deduce that *the existence of an integral surface in \mathbb{P}^r , not contained in hypersurfaces of degree $\leq i$, of degree s given by the equation (1), is equivalent to the existence of a rational normal scroll surface of degree s in \mathbb{P}^{s+1} which can be projected isomorphically into \mathbb{P}^r as a surface of maximal rank.* By [1, Theorem 2] one knows that, for $i = 2$ and $i = 3$, a general projection in \mathbb{P}^r of a smooth rational normal scroll surface of degree s , is of maximal rank. Therefore, at least for $i = 2$ and $i = 3$, the bound (2) is sharp. We do not know whether there are other values of i for which this is true.

(vi) Taking into account the proof of Theorem 1.1 (see below), one may explicit the assumption $d \gg s$. In fact, an elementary computation, that we omit, proves that it suffices to assume $d > 179$ in the case $i = 2$ and $r = 5$, and

$$d > \max \left\{ \frac{2s}{r-2} \prod_{j=1}^{r-2} [(r-1)!s]^{\frac{1}{r-1-j}}, \frac{4s}{r-2} (s+1)^3 \right\}$$

otherwise (compare with [3, Section 2, (v), (vi), (vii)]). Notice that, for $r \geq 4$ and $i \geq 2$, one has $\frac{4s}{r-2}(s+1)^3 \geq is$.

In view of previous Remark 1.2, (v), we have:

Corollary 1.3. *If $i = 2$ or $i = 3$, the bound (2) is sharp.*

2. THE PROOF OF THEOREM 1.1

Lemma 2.1. *Fix integers $r \geq 4$ and $i \geq 2$ (for $r = 4$ assume $i \geq 3$). Assume that the rational number s defined by the equation (1) is an integer. Let $S \subset \mathbb{P}^r$ be an integral surface of degree s , not contained in a hypersurface of \mathbb{P}^r of degree $\leq i$. Then S is the isomorphic projection in \mathbb{P}^r of a rational normal scroll surface $S' \subset \mathbb{P}^{s+1}$ of degree s .*

Proof. It suffices to prove that $h^0(S, \mathcal{O}_S(1)) = s + 2$ (notice that $s \geq 5$).

Let $\Sigma \subset \mathbb{P}^{r-1}$ be a general hyperplane section of S . For every $j \geq 1$ one has $h^0(\Sigma, \mathcal{O}_\Sigma(j)) \leq 1 + js$ [3, (10)]. Hence, from the natural exact sequence $0 \rightarrow \mathcal{O}_S(j-2) \rightarrow \mathcal{O}_S(j-1) \rightarrow \mathcal{O}_\Sigma(j-1) \rightarrow 0$, we obtain $h^0(S, \mathcal{O}_S(1)) \leq s + 2$, and:

$$h^0(S, \mathcal{O}_S(i-1)) \leq h^0(S, \mathcal{O}_S(1)) + \sum_{j=2}^{i-1} [1 + js] \leq i + \binom{i}{2}s.$$

We deduce that if $h^0(S, \mathcal{O}_S(i-1)) = i + \binom{i}{2}s$, then $h^0(S, \mathcal{O}_S(1)) = s + 2$. Therefore, it suffices to prove that $h^0(S, \mathcal{O}_S(i-1)) = i + \binom{i}{2}s$. Since $h^0(\mathbb{P}^r, \mathcal{I}_S(i-1)) = 0$, from the natural exact sequence $0 \rightarrow \mathcal{I}_S(i-1) \rightarrow \mathcal{O}_{\mathbb{P}^r}(i-1) \rightarrow \mathcal{O}_S(i-1) \rightarrow 0$, we get:

$$h^1(\mathbb{P}^r, \mathcal{I}_S(i-1)) = h^0(S, \mathcal{O}_S(i-1)) - \binom{r+i-1}{i-1} \leq i + \binom{i}{2}s - \binom{r+i-1}{i-1}.$$

Hence, it suffices to prove that

$$(3) \quad h^1(\mathbb{P}^r, \mathcal{I}_S(i-1)) = i + \binom{i}{2}s - \binom{r+i-1}{i-1}.$$

To this purpose notice that, since $h^0(\mathbb{P}^r, \mathcal{I}_S(i)) = 0$, from the natural exact sequence $0 \rightarrow \mathcal{I}_S(i-1) \rightarrow \mathcal{I}_S(i) \rightarrow \mathcal{I}_{\Sigma, \mathbb{P}^{r-1}}(i) \rightarrow 0$, we have:

$$(4) \quad h^0(\mathbb{P}^{r-1}, \mathcal{I}_{\Sigma, \mathbb{P}^{r-1}}(i)) \leq h^1(\mathbb{P}^r, \mathcal{I}_S(i-1)) \leq i + \binom{i}{2}s - \binom{r+i-1}{i-1}.$$

Now, from the natural exact sequence $0 \rightarrow \mathcal{I}_{\Sigma, \mathbb{P}^{r-1}}(i) \rightarrow \mathcal{O}_{\mathbb{P}^{r-1}}(i) \rightarrow \mathcal{O}_\Sigma(i) \rightarrow 0$, we get:

$$\begin{aligned} h^1(\mathbb{P}^{r-1}, \mathcal{I}_{\Sigma, \mathbb{P}^{r-1}}(i)) &= h^0(\mathbb{P}^{r-1}, \mathcal{I}_{\Sigma, \mathbb{P}^{r-1}}(i)) - \binom{r+i-1}{r-1} + h^0(\Sigma, \mathcal{O}_\Sigma(i)) \\ &\leq i + \binom{i}{2}s - \binom{r+i-1}{i-1} - \binom{r+i-1}{r-1} + (1 + is), \end{aligned}$$

and this number is equal to 0 in view of the equation (1). It follows that:

$$h^0(\mathbb{P}^{r-1}, \mathcal{I}_{\Sigma, \mathbb{P}^{r-1}}(i)) = i + \binom{i}{2}s - \binom{r+i-1}{i-1}.$$

And so from (4) we obtain (3). □

Proof of Theorem 1.1. First we notice that C cannot be contained in a surface of degree $< s$, because every such a surface is contained in a hypersurface of degree i [3, (11)]. On the other hand, if C is not contained in a surface of degree $< s + 1$, then, since $d \gg s$, by [2, loc. cit.] we have (compare with [3, Section 2, (v), (vi), (vii)]):

$$p_a(C) \leq \frac{d^2}{2(s+1)} + O(d) < \frac{d^2}{2s} + O(d) = \binom{m}{2} s + m\epsilon.$$

Therefore, in order to prove Theorem 1.1, we may assume C is contained in a surface S of degree s , not contained in a hypersurface of degree $\leq i$. By previous Lemma 2.1, we know that S is an isomorphic projection of a rational normal scroll surface $S' \subset \mathbb{P}^{s+1}$. Hence, C is isomorphic to a curve $C' \subseteq S'$ of degree d , in particular $p_a(C) = p_a(C')$. Since $d \gg s$, by Bezout's theorem C' is non-degenerate. Therefore, the bound (2) for $p_a(C)$ follows from Castelnuovo's bound for $p_a(C')$ [4, loc. cit.]. Observe that previous argument shows also that if the bound (2) is sharp, then every extremal curve $D \subset \mathbb{P}^r$ is contained in a surface $S \subset \mathbb{P}^r$ of degree s , not contained in a hypersurface of degree $\leq i$. Conversely, if such a surface S exists, with notations as before, let $D' \subset S'$ be a Castelnuovo's curve of degree d . Let $D \subset S$ be the projection of D' in S . Then D is an extremal curve, and so the bound (2) is sharp. In fact, $p_a(D) = p_a(D')$. Moreover, since $d > is$, D cannot be contained in a hypersurface of degree i by Bezout's theorem (compare with Remark 1.2, (vi)). Now, let $D \subset \mathbb{P}^r$ be an extremal curve. We just proved that D is contained in a surface S of degree s , a fortiori not contained in a hypersurface of degree $\leq i$. The surface S is an isomorphic projection of a rational normal scroll surface $S' \subset \mathbb{P}^{s+1}$, and, as before, via this isomorphism, D corresponds to a Castelnuovo's curve of S' . Since $d \gg s$, again by Bezout's theorem, the surface S of degree s containing D is unique. By [3, (11)] and the definition of s , it follows that S is contained in a hypersurface T of degree $i + 1$. Moreover, D cannot be a.C.M.. Otherwise, $H^1(\mathbb{P}^r, \mathcal{I}_S(1)) = 0$ [3, Section 2, (vii)], and so the restriction map $H^0(\mathbb{P}^r, \mathcal{I}_S(i)) \rightarrow H^0(\mathbb{P}^{r-1}, \mathcal{I}_{\mathbb{P}^{r-1}, \Sigma}(i))$ is onto ($\Sigma :=$ a general hyperplane section of S). This is impossible, because, in view of [3, (10)], Σ is contained in a hypersurface of degree i , and S is not. \square

REFERENCES

- [1] Ballico, E. - Ellia, Ph.: *On Projections of Ruled and Veronese Surfaces*, Journal of Algebra, **121**, 477-487 (1989).
- [2] Chiantini, L. - Ciliberto, C. - Di Gennaro, V.: *The genus of projective curves*, Duke Math. J., **70**(2), 229-245 (1993).
- [3] Di Gennaro, V.: *The genus of curves in \mathbb{P}^4 and \mathbb{P}^5 not contained in quadrics*, Rendiconti del Circolo Matematico di Palermo, Series 2 (2022), Online first articles, 29 July 2022, <https://doi.org/10.1007/s12215-022-00788-x>
- [4] Eisenbud, D. - Harris, J.: *Curves in Projective Space*, Sém. Math. Sup. 85 Les Presses de l'Université de Montréal, 1982.

UNIVERSITÀ DI ROMA "TOR VERGATA", DIPARTIMENTO DI MATEMATICA, VIA DELLA RICERCA SCIENTIFICA, 00133 ROMA, ITALY.

Email address: digennar@mat.uniroma2.it