

# Padovan and Perrin Numbers as Sums of Two Jacobsthal Numbers

Mustafa Ismail<sup>1</sup>, Ahmed Gaber<sup>2</sup> and M. Anwar<sup>3</sup> <sup>123</sup>

Department of Mathematics, University of Ain Shams

Faculty of science, Egypt

e-mails: [mostafaesmail\\_p@sci.asu.edu.eg](mailto:mostafaesmail_p@sci.asu.edu.eg)<sup>1</sup>,

[a.gaber@sci.asu.edu.eg](mailto:a.gaber@sci.asu.edu.eg)<sup>2</sup>,

[mohammed.anwar@hotmail.com](mailto:mohammed.anwar@hotmail.com), [mohamedanwar@sci.asu.edu.eg](mailto:mohamedanwar@sci.asu.edu.eg)<sup>3</sup>

## Abstract

Let  $\{P_k\}_{k \geq 0}$  be the Padovan sequence defined by  $P_k = P_{k-2} + P_{k-3}$  with initial values are  $P_0 = P_1 = P_2 = 1$ . Let  $\{R_k\}_{k \geq 0}$  be the Perrin sequence defined by  $R_k = R_{k-2} + R_{k-3}$  with initial values are  $R_0 = 3, R_1 = 0, R_2 = 2$ . And let  $\{J_n\}_{n \geq 0}$  be the Jacobsthal sequence defined by  $J_n = 2J_{n-1} + J_{n-2}$  with initials  $J_0 = 0, J_1 = 1$ . In this paper we determine all Padovan and Perrin numbers which are sum of two Jacobsthal number.

**Keywords:** Padovan sequence, Perrin sequence, Jacobsthal sequence, linear form in logarithm, reduction method.

**2020 Mathematics Subject Classification:** 11B39, 11D72, 11J70, 11J86.

## 1 Introduction

The Padovan sequence  $\{P_k\}$  is defined by  $P_k = P_{k-2} + P_{k-3}$  with initial values are  $P_0 = P_1 = P_2 = 1$ . The first few values of  $P_k$  are

$$1, 1, 1, 2, 2, 3, 4, 5, 7, 9, 12, 16, 21, 28, 37, 49, 65, 86, 114, 151, \dots$$

The numbers  $P_k$  appear as the integer sequence (OEIS [A000931](#)) from [11].

The sequence was described by Ian Stewart in his Scientific American column Mathematical Recreations in June 1996 [14]. Some properties of padovan sequence and generalizations can be found in [5][10][16]. Diophantine equations involving padovan sequence have been studied by many mathematicians. For example, In 2019, García Lomeli and Hernández Santos solve Pillai's problem with Padovan numbers and powers of two [1]. Also García Lomeli, Hernández Santos and Zacateca determine all repdigits in base 10 that can be written as a sum of two Padovan numbers[3]. In 2020 Mahadi Ddamulira determine all positive square-free integers  $d$  such that the Pell equations  $x^2 - dy^2 = N$  with  $N \in \{\pm 1, \pm 4\}$  have at least two positive integer solutions  $(x, y)$  and  $(x', y')$  such that both  $x$  and  $x'$  are sums of two Padovan numbers [8]. In 2021 Mahadi Ddamulira determine all Padovan numbers that are concatenations of two distinct repdigits [7]. In 2022 García Lomeli, Hernández Santos and Florian luca determine

Fibonacci numbers as sums of two padovan numbers [2]. The Perrin sequence  $\{R_k\}$  is a linear recurrent sequence defined by  $R_k = R_{k-2} + R_{k-3}$  with initials  $R_0 = 3, R_1 = 0, R_2 = 2$ . Although the sequence is named after R. Perrin who studied it in 1899, it had been explored earlier, in 1876, by Édouard Lucas. Perrin numbers were studied by many mathematicians in the nineteenth century. The following is the list of few Perrin numbers:

$$3, 0, 2, 3, 2, 5, 5, 7, 10, 12, 17, 22, 29, 39, 51, 68, 90, 119, 158, 209, \dots$$

The numbers  $P_k$  appear as the integer sequence (OEIS [A001608](#)) from [12]. The Jacobsthal sequence  $\{J_n\}$  is a linear recurrent sequence defined by  $J_n = 2J_{n-1} + J_{n-2}$  with initials  $J_0 = 0, J_1 = 1$ . The first few values of  $J_n$  are:

$$0, 1, 1, 3, 5, 11, 21, 43, 85, 171, 341, 683, 1365, 2731, 5461, 10923, 21845, 43691, \dots$$

The numbers  $J_n$  appear as the integer sequence (OEIS [A001045](#)) from [13]. Properties of Jacobsthal sequence and generalization found in [15][6]. In this paper we determine all Padovan numbers which are sum of two Jacobsthal number, *i.e.*, we solve the following equations

$$P_k = J_n + J_m. \tag{1.1}$$

And

$$R_k = J_n + J_m. \tag{1.2}$$

**Theorem 1.1.** *The only non-negative solutions  $(k, n, m)$  which satisfy equation (1.1) and  $n \geq m$  belongs to the set*

$$\left\{ \begin{array}{ccccc} (0, 1, 0), & (1, 1, 0), & (2, 1, 0), & (3, 2, 1), & (4, 2, 1), \\ (5, 3, 0), & (6, 3, 2), & (6, 3, 1), & (7, 4, 0), & (10, 5, 1), \\ (10, 5, 2), & (11, 5, 4), & (12, 6, 0), & (17, 8, 1), & (17, 8, 2) \end{array} \right\}.$$

**Theorem 1.2.** *The only non-negative solutions  $(k, n, m)$  which satisfy equation (1.2) and  $n \geq m$  belongs to the set*

$$\left\{ \begin{array}{ccccc} (0, 3, 1), & (1, 0, 0), & (2, 2, 1), & (2, 1, 1), & (3, 3, 1), \\ (4, 2, 1), & (4, 1, 1), & (5, 4, 0), & (6, 4, 0), & (8, 4, 4), \\ (9, 5, 1), & (9, 5, 2), & (11, 6, 1), & (11, 6, 2), & (16, 8, 4) \end{array} \right\}.$$

## 2 PRELIMINARY RESULTS

### 2.1 Padovan Sequence

Here, we recall some important properties of the Padovan sequence  $\{a_n\}_{n \geq 0}$ . The characteristic equation

$$\Psi(x) := x^3 - x - 1 = 0,$$

has roots  $\alpha, \beta, \gamma = \bar{\beta}$ ,

$$\alpha = \frac{r_1 + r_2}{6}, \quad \beta = \frac{-(r_1 + r_2) + \sqrt{-3}(r_1 - r_2)}{12}; \tag{2.1}$$

and

$$r_1 = \sqrt[3]{108 + 12\sqrt{69}} \quad \text{and} \quad r_2 = \sqrt[3]{108 - 12\sqrt{69}}. \quad (2.2)$$

Furthermore, the Binet formula is given by

$$P_k = a\alpha^k + b\beta^k + c\gamma^k \quad \text{for all } k \geq 0, \quad (2.3)$$

where

$$a = \frac{\alpha + 1}{(\alpha - \beta)(\alpha - \gamma)}, \quad b = \frac{\beta + 1}{(\beta - \alpha)(\beta - \gamma)}, \quad c = \frac{\gamma + 1}{(\gamma - \alpha)(\gamma - \beta)} = \bar{b}. \quad (2.4)$$

The minimal polynomial of  $a$  over the integers is given by

$$23x^3 - 23x^2 + 6x - 1,$$

has zeros  $a, b, c$  with  $|a|, |b|, |c| < 1$ . Numerically, the following estimates holds:

$$\begin{aligned} 1.32 < \alpha < 1.33; \\ 0.86 < |\beta| = |\gamma| = \alpha^{-\frac{1}{2}} < 0.87; \\ 0.72 < a < 0.73; \\ 0.24 < |b| = |c| < 0.25. \end{aligned} \quad (2.5)$$

From (2.1), (2.2) and (2.5), we can prove that

$$|P_k - a\alpha^k| = |b\beta^k + c\gamma^k| < \frac{1}{\alpha^{\frac{k}{2}}} \quad \text{for all } k \geq 1. \quad (2.6)$$

Furthermore, by induction one can prove that

$$\alpha^{k-3} \leq P_k \leq \alpha^{k-1} \quad \text{holds for all } k \geq 1. \quad (2.7)$$

## 2.2 Perrin Sequence

The Perrin sequence is a linear recurrent sequence defined by  $R_k = R_{k-2} + R_{k-3}$  for  $k > 2$  with initials  $R_0 = 3, R_1 = 1$ . We can use principle of mathematical induction to prove that

$$\alpha^{k-2} \leq R_k \leq \alpha^{k+1}. \quad (2.8)$$

The characteristic equation of Perrin sequence is the same of The characteristic equation of Padovan sequence

$$\Psi(x) := x^3 - x - 1 = 0,$$

and has roots  $\alpha, \beta, \gamma = \bar{\beta}$  Furthermore, the Binet formula is given by

$$P_k = \alpha^k + \beta^k + \gamma^k \quad \text{for all } k \geq 0, \quad (2.9)$$

## 2.3 Jacobsthal Sequence

The Jacobsthal sequence is an additive sequence similar to the Fibonacci sequence, defined by the recurrence  $J_n = J_{n-1} + 2J_{n-2}$  with  $J_0 = 0, J_1 = 1$ . The characteristic equation is

$$P(x) = x^2 - x - 2.$$

The Binet formula for the Jacobsthal sequence is given by

$$J_n = \frac{2^n - (-1)^n}{3}. \quad (2.10)$$

**Lemma 2.1.**

$$2^{n-2} \leq J_n \leq 2^{n-1} \text{ for all } n \geq 1. \quad (2.11)$$

*Proof.* Using the principle of mathematical induction.  $\square$

## 2.4 Linear forms in logarithms.

Let  $\eta$  be an algebraic number of degree  $d$  with minimal polynomial over  $\mathbb{Z}$

$$a_0x^d + a_1x^{d-1} + \dots + a_d = a_0 \prod_{i=1}^d (x - \eta^{(i)}),$$

where the leading coefficient  $a_0$  is positive and the  $\eta^{(i)}$ 's are the conjugates of  $\eta$ . Then the logarithmic height of  $\eta$  is given by

$$h(\eta) := \frac{1}{d} \left( \log a_0 + \sum_{i=1}^d \log (\max \{ |\eta^{(i)}|, 1 \}) \right)$$

The following are some of the properties of the logarithmic height function  $h(\cdot)$ :

$$\begin{aligned} h(\eta_1 \pm \eta_2) &\leq h(\eta_1) + h(\eta_2) + \log 2; \\ h(\eta_1 \eta_2^{\pm 1}) &\leq h(\eta_1) + h(\eta_2); \\ h(\eta^s) &= |s| h(\eta) \quad (s \in \mathbb{Z}). \end{aligned}$$

**Theorem 2.2.** Let  $\eta_1, \dots, \eta_l$  be positive real algebraic numbers in a real algebraic number field  $\mathbb{L} \subset \mathbb{R}$  of degree  $D$ ,  $b_1, \dots, b_l$  be a non zero integers, and assume that

$$\Lambda_1 := \eta_1^{b_1} \dots \eta_l^{b_l} - 1 \neq 0.$$

Then,

$$\log |\Lambda| > -1.4 \cdot 30^{l+3} \cdot l^{4.5} \cdot D^2 \cdot (1 + \log D) \cdot (1 + \log B) A_1 \dots A_l, \quad (2.12)$$

where

$$B \geq \max\{|b_1|, \dots, |b_l|\},$$

and

$$A_i \geq \max\{Dh(\eta_i), |\log \eta_i|, 0.16\}, \text{ for all } i = 1, \dots, l.$$

*Proof.* The proof can be found in [9].  $\square$

## 2.5 Reduction Procedure

During the calculations, we get upper bounds on our variables which are too large, thus we need to reduce them. To do so, we use a slight variation of result due to Dujella and pethö . For a real number  $X$ , we put  $||X|| := \min\{|X - n| : n \in \mathbb{Z}\}$  for the distance from  $X$  to the nearest integer.

**Lemma 2.3.** *Let  $M$  be a positive integer. Let  $\tau, \mu, A > 0, B > 1$  be given real numbers. Assume  $\frac{p}{q}$  is a convergent of  $\tau$  such that  $q > 6M$  and  $\epsilon := ||\mu q|| - M||\tau q|| > 0$ . If  $(n, m, \omega)$  is a positive solution to the inequality*

$$0 < |n\tau - m + \mu| < \frac{A}{B^\omega}$$

with  $n \leq M$ , then

$$\omega < \frac{\log\left(\frac{Aq}{\epsilon}\right)}{\log B}.$$

*Proof.* Dujella and pethö proved this result in [4]. □

**Legendre theorem of continued fraction.**

**Theorem 2.4.** *Let  $x$  be a real number and let  $p, q \in \mathbb{Z}$  If*

$$\left| \frac{p}{q} - x \right| < \frac{1}{2q^2}$$

then  $\frac{p}{q}$  is a convergent continued fraction of  $x$ .

## 3 Proof of theorem 1.1

*Proof.* We ran a computation for  $k \leq 850$  in sage and got only the indicated solutions. We now assume that  $k > 850, n \geq 10$  and from the symmetry of equation (1.1) we can assume that  $n \geq m$ .

### 3.1 bounding $m, n$ and $k$ .

We use the inequality (2.7)(2.11) to get a relation between  $k, n$ . Thus we have,

$$\alpha^{k-3} \leq P_k = J_n + J_m \leq 2^n \quad \text{and} \quad 2^{n-2} \leq J_n + J_m = P_k \leq \alpha^{k-1}. \quad (3.1)$$

This implies that,

$$(n-2) \frac{\log 2}{\log \alpha} + 1 \leq k \leq n \frac{\log 2}{\log \alpha} + 3. \quad (3.2)$$

In particular, we take  $k < 3n$ . Here we replace the Padovan and Jacobsthal sequences in (1.1) by their corresponding Binet formulas as follows.

$$a\alpha^k + b\beta^k + c\gamma^k = \frac{2^n - (-1)^n}{3} + \frac{2^m - (-1)^m}{3}. \quad (3.3)$$

Then,

$$\left| a\alpha^k - \frac{2^n}{3} \right| = \left| \frac{2^m}{3} - \frac{((-1)^n + (-1)^m)}{3} - b\beta^k - c\gamma^k \right|. \quad (3.4)$$

Using equation (2.6) we get,

$$\left| a\alpha^k - \frac{2^n}{3} \right| < \frac{4 \cdot 2^m}{3}. \quad (3.5)$$

Thus,

$$\left| \frac{3a\alpha^k}{2^n} - 1 \right| < \frac{4}{2^{n-m}}. \quad (3.6)$$

Now we apply matveev theorem(2.2) for the following data:

$$\Lambda_1 = 3a\alpha^k 2^{-n} - 1, \quad l = 3, \quad \eta_1 = 3a, \quad \eta_2 = \alpha, \quad \eta_3 = 2, \quad b_1 = 1, \quad b_2 = k, \quad b_3 = -n.$$

First we show that  $\Lambda_1 \neq 0$ . If  $\Lambda_1 = 0$ , then  $3a\alpha^k = 2^n$ . Consider the automorphism  $\sigma$  such that  $\sigma(a\alpha) = b\beta$ . Then  $|3b\beta^k| = 2^n$ . But  $|3b\beta^k| < 3$ , then  $2^n < 3$  which is a contradiction since  $n \geq 10$ . Then we take  $\mathbb{L} = \mathbb{Q}(\alpha)$ , for which  $D = 3$ . Since,

$$h(\eta_1) \leq h(3) + h(a) \leq \log 3 + \frac{1}{3} \log 23;$$

$$h(\eta_2) = \frac{1}{3} \log \alpha;$$

$$h(\eta_3) = \log 2.$$

We take,

$$A_1 = 3 \log 3 + \log 23, \quad A_2 = \log \alpha, \quad \text{and} \quad A_3 = 3 \log 2.$$

Finally we take  $B = 3n$ . Thus theorem(2.2) gives

$$\log |\Lambda_1| > -1.4 \times 30^6 \times 3^{4.5} \times 9 \times (1 + \log 3)(1 + \log 3n)(3 \log 3 + \log 23)(3 \log 2 \log \alpha),$$

which can simplify to,

$$\log |\Lambda_1| > -4 \times 10^{14}(1 + \log 3n). \quad (3.7)$$

Then also from (3.6) we have,

$$\log |\Lambda_1| < \log 4 + (m - n) \log 2. \quad (3.8)$$

Thus by comparing inequalities in (3.7) and (3.8) we get,

$$(n - m) \log 2 - \log 4 < 4 \times 10^{14}(1 + \log 3n). \quad (3.9)$$

Hence,

$$m \log 2 > n \log 2 - 4 \times 10^{14}(1 + \log 3n) - \log 4. \quad (3.10)$$

Now we write equation(1.1) as,

$$a\alpha^k - \frac{2^n(1 + 2^{m-n})}{3} = -b\beta^k - c\gamma^k - \frac{(-1)^n - (-1)^m}{3}, \quad (3.11)$$

which can be written as follows,

$$\left| \frac{3a\alpha^k 2^{-n}}{1+2^{m-n}} - 1 \right| = \left| \frac{3 \cdot 2^{-n}}{1+2^{m-n}} \left( -b\beta^k - c\gamma^k - \frac{(-1)^n - (-1)^m}{3} \right) \right|. \quad (3.12)$$

Using equation(2.6) and the fact  $\frac{(-1)^n - (-1)^m}{3} \leq \frac{2}{3}$  and  $m \leq n$ . Then we get,

$$\left| \frac{3a\alpha^k 2^{-n}}{1+2^{m-n}} - 1 \right| < \frac{5}{2^m}. \quad (3.13)$$

Let  $\Lambda_2 = \frac{3a}{1+2^{m-n}}\alpha^k 2^{-n} - 1$ . Hence,

$$\log |\Lambda_2| < \log 5 - m \log 2. \quad (3.14)$$

Now we apply matveev theorem(2.2) for the following data:

$$\eta_1 = \frac{3a}{1+2^{m-n}}, \quad \eta_2 = \alpha, \quad \eta_3 = 2, \quad l = 3, \quad b_1 = 1, \quad b_2 = k, \quad b_3 = -n.$$

First we show that  $\Lambda_2 \neq 0$ . If  $\Lambda_2 = 0$ , then  $3a\alpha^k = 2^n + 2^m$ . Consider the automorphism  $\sigma$  such that  $\sigma(a\alpha) = b\beta$ . Then  $|3b\beta^k| = 2^n + 2^m$ . But  $|3b\beta^k| < 3$ , then  $2^n + 2^m < 3$  which is a contradiction since  $n \geq 10$ . Then we take  $\mathbb{L} = \mathbb{Q}(\alpha)$ , for which  $D = 3$ . Since,

$$h(\eta_1) \leq h(3) + h(a) + h(1+2^{m-n}) \leq \log 3 + \frac{1}{3} \log 23 + (n-m) \log 2 + \log 2;$$

$$h(\eta_2) = \frac{1}{3} \log \alpha;$$

$$h(\eta_3) = \log 2.$$

We take,

$$A_1 = 3 \log 3 + \log 23 + 3(n-m) \log 2 + 3 \log 2, \quad A_2 = \log \alpha, \quad \text{and} \quad A_3 = 3 \log 2.$$

Finally we take  $B = 3n$ . Thus theorem (2.2) gives

$$\log |\Lambda_2| > -1.4 \times 30^6 \times 3^{4.5} \times 9 \times (1 + \log 3)(1 + \log 3n)(3 \log 3 + \log 23 + 3 \log 2 + 3(n-m) \log 2)(3 \log 2 \log \alpha),$$

Then using equation(3.9),(3.10),(3.14) and some computation we get,

$$n \log 2 < 4.2 \times 10^{14}(1 + \log 3n) + 2 \times 10^{27}(1 + \log 3n)^2 + 3. \quad (3.15)$$

Using mathematica we can find

$$n < 2 \times 10^{31}. \quad (3.16)$$

### 3.2 Reducing bound on $n$ .

Now we will reduce the above bound on  $n$ . To do this, we first assume that  $n - m \geq 20$  and we put

$$\Gamma_1 = \log(3a) + k \log \alpha - n \log 2.$$

Then we have from equation(3.6) that,

$$\Lambda_1 = e^{\Gamma_1} - 1 < \frac{4}{2^{m-n}} < \frac{1}{4}, \quad (3.17)$$

which implies that

$$|\Gamma_1| < \frac{1}{2}. \quad (3.18)$$

Then  $|\Gamma_1| < 2|e^{\Gamma_1} - 1|$ . Therefore we get

$$|\Gamma_1| < \frac{8}{2^{n-m}}. \quad (3.19)$$

We observe that  $\Gamma_1 \neq 0$  since  $\Lambda_1 \neq 0$ . Then

$$0 < \left| \frac{\log 3a}{\log 2} - n + k \left( \frac{\log \alpha}{\log 2} \right) \right| < \frac{8}{2^{n-m} \log 2} < \frac{12}{2^{n-m}}. \quad (3.20)$$

We apply lemma (2.3) with  $M = 6 \times 10^{31}$  ( $M > 3n > k$ ),  $\tau = \frac{\log \alpha}{\log 2}$ ,  $\mu = \frac{\log 3a}{\log 2}$ ,  $A = 12$ ,  $B = 2$ . Write  $\tau$  as a continued fraction  $[a_0, a_1, \dots]$  we get  $q_{85} = 6926604615162884914996228107542442207397 > 2 \times 10^{32} > 6M$ . We compute

$$\epsilon = ||\mu q_{81}|| - M ||\tau q_{85}|| > 0.3.$$

Thus by lemma (2.3), we get  $n - m \leq 137$ . Now we put

$$\Gamma_2 = \log \left( \frac{3a}{1 + 2^{m-n}} \right) + k \log \alpha - n \log 2.$$

Then we have from equation(3.13) that,

$$\Lambda_2 = e^{\Gamma_2} - 1 < \frac{5}{2^m} < \frac{1}{4}, \quad (3.21)$$

which implies that

$$|\Gamma_2| < \frac{1}{2}. \quad (3.22)$$

Then  $|\Gamma_2| < 2|e^{\Gamma_2} - 1|$ . Therefore we get

$$|\Gamma_2| < \frac{10}{2^m}. \quad (3.23)$$

We observe that  $\Gamma_2 \neq 0$  since  $\Lambda_2 \neq 0$ . Then

$$0 < \left| \frac{\log \left( \frac{3a}{1+2^{m-n}} \right)}{\log 2} - n + k \left( \frac{\log \alpha}{\log 2} \right) \right| < \frac{15}{2^m}. \quad (3.24)$$

We apply lemma (2.3) with  $M = 6 \times 10^{31}$  ( $M > 3n > k$ ),  $\tau = \frac{\log \alpha}{\log 2}$ ,  $\mu = \frac{\log \left( \frac{3a}{1+2^{m-n}} \right)}{\log 2}$ ,  $A = 15$ ,  $B = 2$ . Write  $\tau$  as a continued fraction  $[a_0, a_1, \dots]$  we get  $q_{85} = 67577997293290973143551202848941006 > 2 \times 10^{32} > 6M$ . We compute every  $\epsilon$  such that  $n - m < 137$

$$\epsilon = ||\mu q_{85}|| - M ||\tau q_{85}|| > 0.003.$$

Thus by lemma (2.3), we get  $m < 140$  so  $n < 275$  and  $k < 850$ . We used Sage to solve (1.1) if  $m < 144$  so  $n < 281$  and  $k < 850$ . We get the indicated solution in theorem (1.1). Then the proof is completed.  $\square$



## 4 Proof of theorem 1.2

*Proof.* We ran a computation for  $k \leq 870$  in sage and got only the indicated solutions. We now assume that  $k > 870, n \geq 10$  and from the symmetry of equation (1.2) we can assume that  $n \geq m$ .

### 4.1 bounding $m, n$ and $k$ .

We use the inequality (2.11)(4.13) to get a relation between  $k, n$ . Thus we have,

$$\alpha^{k-2} \leq R_k = J_n + J_m \leq 2^n \text{ and } 2^{n-2} \leq J_n + J_m = P_k \leq \alpha^{k+1}. \quad (4.1)$$

This implies that,

$$(n-2) \frac{\log 2}{\log \alpha} - 1 \leq k \leq n \frac{\log 2}{\log \alpha} + 2. \quad (4.2)$$

In particular, we take  $k < 3n$ . Here we replace the Perrin and Jacobsthal sequences in (1.2) by their corresponding Binet formulas as follows.

$$\alpha^k + \beta^k + \gamma^k = \frac{2^n - (-1)^n}{3} + \frac{2^m - (-1)^m}{3}. \quad (4.3)$$

Then,

$$\left| \alpha^k - \frac{2^n}{3} \right| = \left| \frac{2^m}{3} - \frac{((-1)^n + (-1)^m)}{3} - \beta^k - \gamma^k \right|. \quad (4.4)$$

Using equation(2.5) we get,

$$\begin{aligned} \left| \alpha^k - \frac{2^n}{3} \right| &\leq \frac{2^m}{3} + \frac{2}{3} + 2|\beta^k| \\ &< \frac{2^m}{3} + \frac{2^m}{3} + \frac{4 \cdot 2^m}{3} \frac{4 \cdot 2^m}{3}. \end{aligned}$$

Then

$$\left| \alpha^k - \frac{2^n}{3} \right| < \frac{6 \cdot 2^m}{3} \quad (4.5)$$

Thus,

$$\left| \frac{3\alpha^k}{2^n} - 1 \right| < \frac{6}{2^{n-m}}. \quad (4.6)$$

Now we apply matveev theorem(2.2) for the following data:

$$\Lambda_3 = 3\alpha^k 2^{-n} - 1, \quad l = 3, \quad \eta_1 = 3, \quad \eta_2 = \alpha, \quad \eta_3 = 2, \quad b_1 = 1, \quad b_2 = k, \quad b_3 = -n.$$

First we show that  $\Lambda_3 \neq 0$ . If  $\Lambda_3 = 0$ , then  $3\alpha^k = 2^n$ . Consider the automorphism  $\sigma$  such that  $\sigma(\alpha) = \beta$ . Then  $|3\beta^k| = 2^n$ . But  $|3\beta^k| < 3$ , then  $2^n < 3$  which is a contradiction since  $n \geq 10$ . Then we take  $\mathbb{L} = \mathbb{Q}(\alpha)$ , for which  $D = 3$ . Since,

$$h(\eta_1) = \log 3;$$

$$h(\eta_2) = \frac{1}{3} \log \alpha;$$

$$h(\eta_3) = \log 2.$$

We take,

$$A_1 = 3 \log 3, \quad A_2 = \log \alpha, \quad \text{and} \quad A_3 = 3 \log 2.$$

Finally we take  $B = 3n$ . Thus theorem(2.2) gives

$$\log |\Lambda_3| > -1.4 \times 30^6 \times 3^{4.5} \times 9 \times (1 + \log 3)(1 + \log 3n)(3 \log 3)(3 \log 2 \log \alpha),$$

which can simplify to,

$$\log |\Lambda_3| > -6 \times 10^{12}(1 + \log 3n). \quad (4.7)$$

Then also from (4.6) we have,

$$\log |\Lambda_3| < \log 6 + (m - n) \log 2. \quad (4.8)$$

Thus by comparing inequalities in (4.7) and (4.8) we get,

$$(n - m) \log 2 - \log 6 < 6 \times 10^{12}(1 + \log 3n). \quad (4.9)$$

Hence,

$$m \log 2 > n \log 2 - 6 \times 10^{12}(1 + \log 3n) - \log 6. \quad (4.10)$$

Now we write equation(1.2) as,

$$\alpha^k - \frac{2^n(1 + 2^{m-n})}{3} = -\beta^k - \gamma^k - \frac{(-1)^n - (-1)^m}{3}, \quad (4.11)$$

which can be written as follows,

$$\left| \frac{3\alpha^k 2^{-n}}{1 + 2^{m-n}} - 1 \right| = \left| \frac{3 \cdot 2^{-n}}{1 + 2^{m-n}} \left( -\beta^k - \gamma^k - \frac{(-1)^n - (-1)^m}{3} \right) \right|. \quad (4.12)$$

Using equation(2.5) and the fact  $\frac{(-1)^n - (-1)^m}{3} \leq \frac{2}{3}$  and  $m \leq n$ . Then we get,

$$\left| \frac{3\alpha^k 2^{-n}}{1 + 2^{m-n}} - 1 \right| < \frac{8}{2^m}. \quad (4.13)$$

Let  $\Lambda_4 = \frac{3}{1 + 2^{m-n}} \alpha^k 2^{-n} - 1$ . Hence,

$$\log |\Lambda_4| < \log 8 - m \log 2. \quad (4.14)$$

Now we apply matveev theorem(2.2) for the following data:

$$\eta_1 = \frac{3}{1 + 2^{m-n}}, \quad \eta_2 = \alpha, \quad \eta_3 = 2, \quad l = 3, \quad b_1 = 1, \quad b_2 = k, \quad b_3 = -n.$$

First we show that  $\Lambda_4 \neq 0$ . If  $\Lambda_4 = 0$ , then  $3\alpha^k = 2^n + 2^m$ . Consider the automorphism  $\sigma$  such that  $\sigma(\alpha) = \beta$ . Then  $|3\beta^k| = 2^n + 2^m$ . But  $|3\beta^k| < 3$ , then  $2^n + 2^m < 3$  which is a contradiction since  $n \geq 10$ . Then we take  $\mathbb{L} = \mathbb{Q}(\alpha)$ , for which  $D = 3$ . Since,

$$h(\eta_1) \leq h(3) + h(1 + 2^{m-n}) \leq \log 3 + (n - m) \log 2 + \log 2;$$

$$h(\eta_2) = \frac{1}{3} \log \alpha;$$

$$h(\eta_3) = \log 2.$$

We take,

$$A_1 = 3 \log 3 + 3(n - m) \log 2 + 3 \log 2, \quad A_2 = \log \alpha, \quad \text{and} \quad A_3 = 3 \log 2.$$

Finally we take  $B = 3n$ . Thus theorem (2.2) gives

$$\log |\Lambda_4| > -1.4 \times 30^6 \times 3^{4.5} \times 9 \times (1 + \log 3)(1 + \log 3n)(3 \log 3 + 3 \log 2 + 3(n - m) \log 2)(3 \log 2 \log \alpha),$$

Then using equations (4.9),(4.10),(4.14) and some computation we get,

$$n \log 2 < 2.1 \times 10^{13}(1 + \log 3n) + 3 \times 10^{25}(1 + \log 3n)^2 + 4. \quad (4.15)$$

Using mathematica we can find

$$n < 3 \times 10^{29}. \quad (4.16)$$

## 4.2 Reducing bound on $n$ .

Now we will reduce the above bound on  $n$ . To do this, we first assume that  $n - m \geq 20$  and we put

$$\Gamma_3 = \log(3) + k \log \alpha - n \log 2.$$

Then we have from equation(4.6) that,

$$\Lambda_3 = e^{\Gamma_3} - 1 < \frac{6}{2^{m-n}} < \frac{1}{4}, \quad (4.17)$$

which implies that

$$|\Gamma_3| < \frac{1}{2}. \quad (4.18)$$

Then  $|\Gamma_3| < 2 |e^{\Gamma_3} - 1|$ . Therefore we get

$$|\Gamma_3| < \frac{12}{2^{n-m}}. \quad (4.19)$$

We observe that  $\Gamma_3 \neq 0$  since  $\Lambda_3 \neq 0$ . Then

$$0 < \left| \frac{\log 3}{\log 2} - n + k \left( \frac{\log \alpha}{\log 2} \right) \right| < \frac{12}{2^{n-m} \log 2} < \frac{18}{2^{n-m}}. \quad (4.20)$$

We apply lemma (2.3) with  $M = 9 \times 10^{29}$  ( $M > 3n > k$ ),  $\tau = \frac{\log \alpha}{\log 2}$ ,  $\mu = \frac{\log 3}{\log 2}$ ,  $A = 18$ ,  $B = 2$ . Write  $\tau$  as a continued fraction  $[a_0, a_1, \dots]$  we get  $q_{85} = 6926604615162884914996228107542442207397 > 6 \times 10^{30} > 6M$ . We compute

$$\epsilon = ||\mu q_{81}|| - M ||\tau q_{85}|| > 0.3.$$

Thus by lemma (2.3), we get  $n - m \leq 142$ . Now we put

$$\Gamma_4 = \log \left( \frac{3}{1 + 2^{m-n}} \right) + k \log \alpha - n \log 2.$$

And let  $m > 70$  then we have from equation(4.13) that,

$$\Lambda_4 = e^{\Gamma_4} - 1 < \frac{8}{2^m} < \frac{1}{4}, \quad (4.21)$$

which implies that

$$|\Gamma_4| < \frac{1}{2}. \quad (4.22)$$

Then  $|\Gamma_4| < 2|e^{\Gamma_4} - 1|$ . Therefore we get

$$|\Gamma_4| < \frac{16}{2^m}. \quad (4.23)$$

We observe that  $\Gamma_4 \neq 0$  since  $\Lambda_4 \neq 0$ . Then

$$0 < \left| \frac{\log\left(\frac{3}{1+2^{m-n}}\right)}{\log 2} - n + k \left( \frac{\log \alpha}{\log 2} \right) \right| < \frac{24}{2^m}. \quad (4.24)$$

We apply lemma (2.3) with  $M = 9 \times 10^{29}$  ( $M > 3n > k$ ),  $\tau = \frac{\log \alpha}{\log 2}$ ,  $\mu = \frac{\log\left(\frac{3}{1+2^{m-n}}\right)}{\log 2}$ ,  $A = 24$ ,  $B = 2$ . Write  $\tau$  as a continued fraction  $[a_0, a_1, \dots]$  we get  $q_{85} = 67577997293290973143551202848941006 > 3 \times 10^{30} > 6M$ . We compute every  $\epsilon$  such that  $n - m < 142$  except if  $n - m = 1$

$$\epsilon = ||\mu q_{85}|| - M ||\tau q_{85}|| > 0.003.$$

Thus by lemma (2.3), we get  $m < 145$  so  $n < 287$  and  $k < 870$ . We used Sage to solve (1.2) if  $m < 145$  so  $n < 287$  and  $k < 870$ . We get the indicated solution in theorem (1.2). If  $n - m = 1$  we get  $\epsilon$  always negative. So we solve equation(1.2) if  $n - m = 1$ . In this case equation(1.2) can be written as  $R_k = J_m + J_{m+1}$  and can be reduced as

$$R_k = 2^m. \quad (4.25)$$

Then  $k < 3m$  and from equation(4.16) we get  $m < 3 \times 10^{29}$ . As the same way of proving  $\Lambda_3 < \frac{6}{2^{m-n}}$  we get

$$\alpha^{k2^{-m}} - 1 < \frac{6}{2^m},$$

Thus

$$\left| k \frac{\log \alpha}{\log 2} - m \right| < \frac{12}{2^m}.$$

If  $m > 10$  using  $k < 3m$  then  $\frac{12}{2^m} < \frac{1}{2k^2}$ . From Legendre's theorem, this implies that  $\frac{m}{k}$  is a convergent of  $\frac{\log \alpha}{\log 2}$ . We know that  $k < M$  then using computation with sage gives that

$$q_{69} \leq M < q_{70} \quad \text{and} \quad b := \max\{a_i : i = 1, 2, \dots, 72\} = 80.$$

Then, from the properties of the convergents of a continued fraction we get

$$\frac{1}{(b+2)l} < \left| k \frac{\log \alpha}{\log 2} - m \right| < \frac{12}{2^m}.$$

Thus

$$2^m < 12 \cdot 82 \cdot M.$$

Then  $m \leq 109$ . We used Sage to solve equation (4.25) if  $m < 299$  and we get the indicated solutions of theorem (1.2). Then the theorem (1.2) is proved.  $\square$

## References

- [1] G. L. A. Cecilia and H. H. SANTOS(2019) *Pillai's problem with Padovan numbers and powers of two*, *Revista Colombiana de Matemáticas* Volume 55, 1, 1-14.
- [2] G. L. A. Cecilia, H. H. Santos and L. Florian (2022) *Fibonacci numbers as sums of two Padovan numbers*, *Afrika Matematika*, Volume 33, 1, 1-10.
- [3] L. A. Cecilia, H. H. Santos and Z. Zac (2019) *Repdigits as sums of two Padovan numbers*, *Journal of Integer Sequences*, Volume 22, 2,3.
- [4] Dujella and A.Petho (1998) *A generalization of a theorem of Baker and Davenport*, *The Quarterly Journal of Mathematics*, Volume 49, 195, 291-301.
- [5] A. Faisantn (2019), *ON THE PADOVAN SEQUENCE*, hal-02131654.
- [6] T. Koshy (2017) *Foibanacci and Lucas Numbers with Applications* , Volume 2, Second edition, Hoboken, New Jersey, 433-490.
- [7] D. Mahadi (2021) *Padovan numbers that are concatenations of two distinct repdigits*, *Mathematica Slovaca*, Volume 71, 2, 275-284.
- [8] D. Mahadi (2020) *On the  $x$ -coordinates of Pell equations that are products of two Padovan numbers*, *INTEGERS: Electronic Journal of Combinatorial Number Theory*, Volume 20.
- [9] M. Matveev (2000) *An explicit lower bound for a homogeneous rational linear form in the logarithms of algebraic numbers. II*, Volume 64, 6, 1217.
- [10] S. Ramaswamy(2021), *Properties of Padovan Sequence*, *Turkish Journal of computer and Mathematics Education*.
- [11] N. J. A. Sloane, *Sequence A000931 (Padovan numbers )*". *The On-Line Encyclopedia of Integer Sequences. OEIS Foundation*.
- [12] N. J. A. Sloane, *Sequence A001608 (Perrin numbers )*". *The On-Line Encyclopedia of Integer Sequences. OEIS Foundation*.
- [13] N. J. A. Sloane, *Sequence A014551 (Jacobsthal–Lucas numbers)*". *The On-Line Encyclopedia of Integer Sequences. OEIS Foundation*.
- [14] I. Stewart(1996), *Tales of a Neglected Number Scientific American*, No. 6, 92-93.
- [15] A. F. Torunbalcı (2018) *On generalizations of the Jacobsthal sequence*, *Notes on number theory and discrete mathematics*, Volume 24, 1, pp. 120-135.
- [16] S. Yüksel (2021), *On generalized Padovan numbers*, *Preprints*.