

INVOLUTORY PERMUTATION AUTOMORPHISMS OF BINARY LINEAR CODES

FATMA ALTUNBULAK AKSU, ROGHAYEH HAFEZIEH AND İPEK TUVAY

ABSTRACT. We investigate the properties of binary linear codes of even length whose permutation automorphism group is a cyclic group generated by an involution. Up to dimension or co-dimension 4, we show that there is no quasi group code whose permutation automorphism group is isomorphic to C_2 . By generalizing the method we use to prove this result, we obtain results on the structure of putative extremal self-dual $[72, 36, 16]$ and $[96, 48, 20]$ codes in the presence of an involutory permutation automorphism.

1. INTRODUCTION

A *binary linear code* C of length n is a subspace of \mathbb{F}_2^n . The symmetric group S_n acts on \mathbb{F}_2^n by permuting the set of its coordinates. The group of permutations which sends C onto itself is called *the permutation automorphism group of C* and is denoted by $\text{PAut}(C)$. The permutation automorphism group of a code carries a lot of information about the algebraic structure of the code. For instance, if there is a non-trivial $G \leq \text{PAut}(C)$ for a binary linear code C , the problem of determining C by the help of G is a well-known and difficult problem in coding theory (see for example [4]).

Our intention to attack this problem is to investigate the structure of binary linear codes in the presence of an involutory permutation automorphism. In [1], a method for constructing self-dual codes with an involutory permutation automorphism group is given. Later in [2], extremal doubly even self-dual binary codes of length $24m$ with an involutory permutation automorphism are considered. It is shown in [2, Theorem 5.3] that a putative self-dual $[72, 36, 16]$ code can not have an involutory automorphism with some fixed points. Moreover, there are five candidates for the permutation automorphism group of a putative self-dual $[72, 36, 16]$ code and two of the candidates are C_2 and $C_2 \times C_2$ (see [4, Theorem 6.3]). Inspired by all of these, our aim is to deduce some structural results on binary linear codes having an involutory permutation automorphism. We first discuss linear codes with an arbitrary involutory permutation automorphism, then we focus on the ones with involutory fixed point free permutation automorphisms. The fixed subcode associated to a permutation automorphism gives a lot of information about the code itself, as we can see in [1, 2, 5]. In particular, the fixed subcode of an involutory permutation automorphism plays a central role in our arguments.

Borello and Willems, in [7], give an intrinsic description of linear codes of length n whose permutation automorphism group contains a free subgroup of S_n . These codes are called *quasi group codes*. Abelian codes, group codes, quasi-cyclic and quasi-abelian codes belong to the class of quasi group codes. Hence, this approach gives a unified way to consider these different class of codes. In this paper, we show that there is no quasi group code C of even length $n > 2$ and dimension k or $n - k$ where $1 \leq k \leq 4$ where $\text{PAut}(C) \cong C_2$ (see Corollary 4.6). On the other hand, by the methods we develop to prove this result, we obtain some results on the structures of putative extremal self-dual $[72, 36, 16]$ and $[96, 48, 20]$ codes.

The organization of the paper is as follows. In Section 2, we give the background and some lemmas concerning the dimension of the fixed subspace of an involutory permutation automorphism. In Section 3, we show that if a binary linear code C of even length with dimension or co-dimension

Date: August 2, 2022.

2010 Mathematics Subject Classification. 94B05, 11T71, 20B05.

Key words and phrases. Group code, Quasi group code, permutation automorphism of a code.

2 satisfies $\text{PAut}(C) \cong C_2$, then its length is 2 or 4. We also determine all binary linear codes C of length 4 with $\text{PAut}(C) \cong C_2$ up to permutation equivalence. Section 4 deals with binary linear codes of even length with involutory fixed point free permutation automorphisms with dimension 3 and dimension 4. Also, there are some important corollaries together with a conjecture in this section. In Section 5, we give a generalization of our idea that is used in the proof of Theorem 4.4 and we present the applications on putative extremal self-dual [72, 36, 16] and [96, 48, 20] codes.

2. BACKGROUND AND LEMMAS

In this paper, we deal with binary linear codes. Recall that a *binary linear code* C of length n is a subspace of \mathbb{F}_2^n . An element $c = c_1c_2 \dots c_n \in C$ is called a *codeword* of C and its *Hamming weight* is defined to be

$$\text{wt}(c) = |\{i \in \{1, \dots, n\} \mid c_i \neq 0\}|.$$

The *minimum weight* of C is defined by $\text{wt}(C) = \min\{\text{wt}(c) \mid c \in C \setminus \{0\}\}$. If C is a binary linear code with length n , dimension k and minimum weight d , then C is called a *binary* $[n, k, d]$ *code*. For a code $C \subseteq \mathbb{F}_2^n$ and a non-negative integer i , let $A_i(C)$ denote the number of codewords in C of weight i . The sequence $(A_0(C), A_1(C), \dots, A_n(C))$ is called *the weight distribution* of C .

There is a natural action of S_n on \mathbb{F}_2^n induced by the action of S_n on the set $\{1, \dots, n\}$. More precisely, for $c = c_1c_2 \dots c_n \in \mathbb{F}_2^n$ and $\beta \in S_n$,

$$c^\beta = c_{\beta^{-1}(1)}c_{\beta^{-1}(2)} \dots c_{\beta^{-1}(n)}.$$

For a linear code C , set $C^\beta = \{c^\beta \mid c \in C\}$. It is easy to see that C^β is also a linear code. A permutation $\beta \in S_n$ is called a *permutation automorphism* or simply an *automorphism* of C if $C^\beta = C$. The stabilizer

$$\text{PAut}(C) = \{\beta \in S_n \mid C^\beta = C\}$$

is called the *permutation automorphism group* of C .

For a linear code C of length n , if there exists a subgroup G of $\text{PAut}(C)$ which is transitive and free, then C is called a *group code* or a *G-code*. This characterization is given first in [6] then in [7], where a weaker notion is introduced. C is called a *quasi group code* or a *quasi-G code* if there is a free subgroup G inside $\text{PAut}(C)$.

Two linear codes C_1 and C_2 of length n are called *permutation equivalent* if there is a $\beta \in S_n$ such that $C_1^\beta = C_2$. In this case, note that $\text{PAut}(C_2) = \text{PAut}(C_1)^\beta$. So if two linear codes are permutation equivalent, then their permutation automorphism groups are conjugate in S_n . In this paper, we denote the k -cycles in S_n as (a_1, a_2, \dots, a_k) where $a_1, a_2, \dots, a_k \in \{1, 2, \dots, n\}$.

For a permutation automorphism β of a binary linear code C , the fixed point set of β , which we denote by $F_\beta(C)$, is defined as

$$F_\beta(C) = \{c \in C \mid c^\beta = c\}.$$

Note that, this set is closed under addition, so $F_\beta(C)$ is a subspace of C . This is called the *fixed subcode* of C under β .

We begin with an observation that holds for any permutation automorphism of a binary linear code which is an involution. It turns out that there is a relationship between the dimension of the code and its fixed subcode under that involution. The notation $\lceil x \rceil$ denotes the smallest integer greater or equal to x .

Lemma 2.1. *Let C be a binary linear code of length n and dimension k . Let $\beta \in \text{PAut}(C)$ be an involution and $f = \dim(F_\beta(C))$. We have that $f \geq (k - f)$ or equivalently $f \geq \lceil k/2 \rceil$.*

Proof. If $C = F_\beta(C)$, the claim holds trivially. So assume that $C \neq F_\beta(C)$. Let's start with a basis \mathcal{B}_1 of $F_\beta(C)$ and then extend it to a basis \mathcal{B} of C . Let

$$\mathcal{B} \setminus \mathcal{B}_1 = \{w_{(1)}, \dots, w_{(k-f)}\}$$

and for any $i = 1, \dots, k - f$ define $x_{(i)} = w_{(i)} + w_{(i)}^\beta$. Then since $w_{(i)} \notin F_\beta(C)$, we have that $x_{(i)} \neq 0$. Also $x_{(i)} = x_{(i)}^\beta$, so that $x_{(i)} \in F_\beta(C)$. If for some $\lambda_i \in \mathbb{F}_2$, $\sum_{i=1}^{k-f} \lambda_i x_{(i)} = 0$, it follows

that

$$\sum_{i=1}^{k-f} \lambda_i w_{(i)} = \left(\sum_{i=1}^{k-f} \lambda_i w_{(i)} \right)^\beta$$

and so $\sum_{i=1}^{k-f} \lambda_i w_{(i)} \in F_\beta(C) \cap \langle \mathcal{B} \setminus \mathcal{B}_1 \rangle$. However, $F_\beta(C) \cap \langle \mathcal{B} \setminus \mathcal{B}_1 \rangle = \{0\}$, which implies that $\sum_{i=1}^{k-f} \lambda_i w_{(i)} = 0$. Hence by the linear independence of the set $\mathcal{B} \setminus \mathcal{B}_1$, it follows that $\lambda_i = 0$ for $i = 1, \dots, k-f$. Therefore, $\{x_{(1)}, \dots, x_{(k-f)}\}$ is a linearly independent subset of $F_\beta(C)$, which yields that $f \geq (k-f)$. \square

Lemma 2.2. *Suppose that C is a binary linear code of dimension k and length $n \geq 4$. If $\text{PAut}(C) = \langle \beta \rangle$, for $\beta = (1, 2)(3, 4) \dots (t, t+1)$ where t is an odd integer such that $3 \leq t \leq (n-1)$ is an arbitrary odd integer, then $\dim(F_\beta(C)) \leq (k-1)$. In particular, if $k \geq 2$ we have that $\lfloor k/2 \rfloor \leq \dim(F_\beta(C)) \leq (k-1)$.*

Proof. Let $f = \dim(F_\beta(C))$. If $f = k$, it follows that $F_\beta(C) = C$. Then every element c of C satisfy $c_i = c_{i+1}$ for all odd $i \in \{1, 2, \dots, t\}$. But then the transposition $(i, i+1) \in \text{PAut}(C)$ for any odd $i \in \{1, 2, \dots, t\}$. This implies that $\text{PAut}(C) \neq \langle \beta \rangle$, a contradiction, so $f \neq k$ and the result follows from Lemma 2.1. \square

3. CODES OF EVEN LENGTH WITH LOWER DIMENSIONS OR CO-DIMENSIONS

We begin with observations on permutation automorphisms of binary linear codes of even length whose dimension or co-dimension is equal to 1.

Proposition 3.1. *Let C be a binary linear code of length $n = 2m$ where $m \geq 2$. Assume that $\dim(C) = 1$ or $\dim(C) = n-1$. Then the following hold.*

- (i) *We have that $\text{PAut}(C) \not\cong C_2$.*
- (ii) *If $\text{wt}(C)$ is even and $\text{wt}(C) \neq n$, then C is a quasi group code which is not a group code.*
- (iii) *Assume that $n = 2^r$ (or equivalently $m = 2^{r-1}$) and $\text{wt}(C) \neq n$. Then C is a quasi group code which is not a group code if and only if $\text{wt}(C)$ is even.*

Proof. If $\dim(C) = n-1$, then $\dim(C^\perp) = 1$ and $\text{PAut}(C) = \text{PAut}(C^\perp)$. Hence it is enough to prove all assertions for 1-dimensional linear codes. Assume that $\dim(C) = 1$ and the weight of C is equal to d . It follows that C is generated by v where v has exactly d number of 1's. Then since $\text{PAut}(C) = \text{stab}_{S_n}(v)$, we have that

$$\text{PAut}(C) \cong S_d \times S_{n-d}.$$

Hence we have that $|\text{PAut}(C)| > 2$. So the assertion (i) is proved.

If $\text{wt}(C) = d$ is even, there is a fixed point free involution γ in S_d , and since $n-d$ is also even, there is a fixed point free involution δ in S_{n-d} . Then we have that $\langle \gamma\delta \rangle \leq \text{PAut}(C)$ and it follows that C is a quasi- $\langle \gamma\delta \rangle$ code. From the structure of $\text{PAut}(C)$, it is easy to see that we can not find a transitive subgroup, hence C is not a group code which finishes the proof of (ii).

Now let us prove the third assertion. Suppose that C is a quasi- H code for $H \leq \text{PAut}(C)$. Then it follows that the order of H is a divisor of $n = 2^r$ and H is a free subgroup of S_n . So there is a fixed point free involution $\gamma \in H$. It follows that γ fixes v and so the number of 1's in v should be even since γ is an involution. The converse direction follows from (ii). \square

Now let us investigate linear codes with dimension or co-dimension equal to 2.

Theorem 3.2. *Let C be a binary linear code of length $n = 2m$ where $m \geq 3$. If $\dim(C) = 2$ or $\dim(C) = n-2$, then $\text{PAut}(C) \not\cong C_2$.*

Proof. Suppose to the contrary that C is a binary linear code of length $n \geq 6$ and $\text{PAut}(C) = \langle \beta \rangle$ where β is an involution in S_n . It is enough to prove the theorem when $\dim(C) = 2$ since $\text{PAut}(C) = \text{PAut}(C^\perp)$ and $\dim(C^\perp) = n - \dim(C)$.

Since any conjugate of a permutation automorphism of C is a permutation automorphism of the permutation equivalent code of C , it is enough to prove the result by considering all possible

involutions $\beta \in S_n$ with different cycle structures. Thus we can assume without loss of generality that $\beta = (1, 2)(3, 4) \dots (t, t+1)$ where $1 \leq t \leq (n-1)$ is an arbitrary odd integer. Let us first assume that $3 \leq t \leq (n-1)$. Then by Lemma 2.2, the dimension f of $F_\beta(C)$ is equal to 1. So

$$C = \{0, x, w, w^\beta\}$$

where $x \in F_\beta(C)$ and $w \notin F_\beta(C)$ and $w^\beta = w + x$. Since $x \in F_\beta(C)$, we know that $x_i = x_{i+1}$ for all odd i where $i \leq t$. If there is an odd integer $j \leq t$ such that $x_j = x_{j+1} = 0$, then since $w^\beta = w + x$, it implies that $w_j = w_{j+1}$. Hence the transposition $(j, j+1)$ fixes all elements on C , so it lies in $\text{PAut}(C)$, which is a contradiction. If for any odd integer $i \leq t$ we have $x_i = x_{i+1} = 1$, then since $w^\beta = w + x$, it implies that $w_i \neq w_{i+1}$ for any odd integer $i \leq t$. Then since $t \geq 3$, there exist two different integers $k, l \leq t+1$ such that $w_k = w_l$ so that $(w^\beta)_k = (w^\beta)_l$. Then it follows that the transposition (k, l) fixes all elements in C , so $(k, l) \in \text{PAut}(C)$. But since (k, l) is different from β this is a contradiction.

Now suppose $t = 1$, then $\beta = (1, 2)$. Suppose that $m \geq 4$. Let u and w be two non-zero elements of C . We claim that there exist at least two integers $k, l \in \{3, 4, \dots, n\}$ such that $u_k = u_l$ and $w_k = w_l$. If this is not true, since $n \geq 8$, there exist at least three elements $k, t, l \in \{3, 4, \dots, n\}$ such that

$$u_k = u_t \text{ and } w_k \neq w_t, \quad u_t = u_l \text{ and } w_t \neq w_l$$

But since these elements take only the values of 0 and 1, it would imply that $u_k = u_l$ and $w_k = w_l$. It follows that the transposition (k, l) which is different from $\beta = (1, 2)$ fixes u, w and $u + w$ hence is in $\text{PAut}(C)$, which is a contradiction.

Suppose that $t = 1$ and $m = 3$, that is $\beta = (1, 2)$. Let u and w be two non-zero elements of C . If we can find $k, l \in \{3, 4, 5, 6\}$ such that $u_k = u_l$ and $w_k = w_l$ as in the previous paragraph then we are done. So let us assume that such k and l do not exist. Then it follows that there exist exactly two integers $k, l \in \{3, 4, 5, 6\}$ with the property that $u_k = u_l = 1$ and that $k', l' \in \{3, 4, 5, 6\}$ with the property that $u_{k'} = u_{l'} = 0$. This implies that $w_k \neq w_l$ and $w_{k'} \neq w_{l'}$. If $u_1 = 1$, then $(1, k)$ and $(1, l)$ fixes u . Since $w_k \neq w_l$, we have that either $w_1 = w_k$ or $w_1 = w_l$. If $w_1 = w_k$, then the transposition $(1, k)$ fixes w and if $w_1 = w_l$ then $(1, l)$ fixes w . If $u_1 = 0$, then both $(1, k')$ and $(1, l')$ fixes u . Since $w_{k'} \neq w_{l'}$, we have that either $w_1 = w_{k'}$ or $w_1 = w_{l'}$, similarly choose either of the transposition $(1, k')$ or $(1, l')$ which fixes w in this case. Since in either case we find a transposition in $\text{PAut}(C)$ which is different from β , we arrive to a contradiction. This finishes the proof. □

Remark 3.3. The lower bound for m in Theorem 3.2 is sharp. Indeed, we show that there exists a linear code of length 4 whose permutation automorphism group is isomorphic to C_2 in the proof of the following result. When the length is equal to 2, then $C = \mathbb{F}_2^2$ and $\text{PAut}(C) = C_2$ in this case.

Proposition 3.4. *Let C be a binary linear code of length 4 and $\dim(C) = 2$. Let $\beta \in S_4$ be a transposition. Then $\text{PAut}(C) = \langle \beta \rangle$ if and only if for every $c \in C$ we have $c^\beta = c$ and $A_i(C) = 1$ for $i = 0, 1, 2, 3$ and $A_4(C) = 0$.*

Proof. Without loss of generality we can assume that $\beta = (1, 2)$.

Suppose that $\text{PAut}(C) = \langle \beta \rangle$. We first claim that $F_\beta(C) = C$. If $F_\beta(C) \neq C$, then the dimension of $F_\beta(C)$ is equal to 1. Let x be the non-zero element of $F_\beta(C)$, then

$$C = \{0, x, w, w^\beta\}$$

where $w \notin F_\beta(C)$ and $x = w + w^\beta$. Since $w \notin F_\beta(C)$, we have that $w_1 \neq w_2$, so that $x_1 = x_2 = 1$. Moreover, since $x = w + w^\beta$ we have that $x_3 = x_4 = 0$. Hence, $x = 1100$. If $w_3 = w_4$ then the transposition $(3, 4)$ fixes w and w^β , as well as x . So we get a contradiction. So $w_3 \neq w_4$. Since we also have that $w_1 \neq w_2$, we have that either $w_1 = w_3$ or $w_1 = w_4$. Assume without loss of generality that $w_1 = 1$. Then if $w_1 = w_3$ or $w_1 = w_4$, then $(1, 3)$ or $(1, 4)$ fixes w respectively. In the first case $(w^\beta)_2 = (w^\beta)_3 = 1$, so that $(w^\beta)^{(1,3)} = 1100 = x$ and $x^{(1,3)} = w^\beta$. In the second case $(w^\beta)_2 = (w^\beta)_4 = 1$, so that $(w^\beta)^{(1,4)} = 1100 = x$ and $x^{(1,4)} = w^\beta$. In both of the cases, we obtain

a transposition in $\text{PAut}(C)$ which is different from β . So we have that $F_\beta(C) = C$, or equivalently for all $c \in C, c^\beta = c$.

Our next claim is $1111 \notin C$. Assume to the contrary that $1111 \in C$. Then there are non-zero elements $u, w \in C$ such that $u + w = 1111$. There are $k, l \in \{2, 3, 4\}$ such that $u_k = u_l$. But since $u + w = 1111$, we have that $w_k = w_l$. Thus (k, l) which is different from β lies in $\text{PAut}(C)$, a contradiction. So $A_4(C) = 0$.

Since $c^\beta = c$ for all $c \in C$, we have that $c_1 = c_2$ for all $c \in C$. If $c_1 = c_2 = 0$ for all $c \in C$, then there is an element $v \in C$ such that $v_3 \neq v_4$ (because otherwise there are less than 4 elements). But this implies that there is $u \in C$ such that $u \neq v$ with $u_3 \neq u_4$. Then the involution $(3, 4)$ flips u and v and it fixes $u + v$. So $(3, 4) \in \text{PAut}(C)$, a contradiction. So there is at least one element $w \in C$ such that $w_1 = w_2 = 1$. Then it follows that $w = 1110$ or $w = 1101$ since $1111 \notin C$. If $w = 1110$, then the other elements of C are 1100 and 0010 . If $w = 1101$, then the other elements of C are 1100 and 0001 . Hence, $A_i(C) = 1$ for $i = 0, 1, 2, 3$.

Suppose that for any $c \in C$ we have $c^\beta = c$ and $A_i(C) = 1$ for $i = 0, 1, 2, 3$ and $A_4(C) = 0$. Then we have that $c_1 = c_2$ for any $c \in C$ and $1111 \notin C$. Then since there exists a unique element with weight 1, there is an element $u \in C$ such that $u = 0010$ or $u = 0001$. Moreover, since there is a unique element with weight equal to 3, we have that either 1110 or 1101 is in C . Then we have that $C = \{0000, 0010, 1100, 1110\}$ or $C = \{0000, 0001, 1100, 1101\}$. Now it is not difficult to see that $\text{PAut}(C) = \langle \beta \rangle$. \square

4. FIXED POINT FREE AUTOMORPHISMS

Throughout this section let C be a binary linear code of length $n = 2m$ and let $\sigma \in \text{PAut}(C)$ be a fixed point free involutory permutation. Then without loss of generality we can assume

$$\sigma = \prod_{1 \leq i \leq n, i \text{ odd}} (i, i+1) = (1, 2)(3, 4) \dots (n-1, n).$$

Then it is easy to see that $c = c_1 c_2 \dots c_n \in F_\sigma(C)$ if and only if $c_i = c_{i+1}$ for all odd $i \in \{1, 2, \dots, n\}$. For an element $c \in F_\sigma(C)$ let us define the set of all odd integers in the support of c as T_c , that is

$$T_c := \{1 \leq i \leq n \mid i \text{ is odd and } c_i c_{i+1} = 11\}.$$

Lemma 4.1. *Let C be a binary linear code of length $n = 2m$ with $\sigma = (1, 2)(3, 4) \dots (n-1, n) \in \text{PAut}(C)$. Let $w \in C \setminus F_\sigma(C)$ and $y := w + w^\sigma \in F_\sigma(C)$. Assume that x is a non-zero element in $F_\sigma(C)$ such that $T_x \subseteq T_y$, then the involution $\alpha_x = \prod_{i \in T_x} (i, i+1)$ in S_n satisfies $w^{\alpha_x} = w + x$.*

Proof. Since $w^\sigma = w + y$ and $T_x \subseteq T_y$, it is easy to see that if $i \in T_x$ then $x_i x_{i+1} = 11$ and $w_i \neq w_{i+1}$. So $(w+x)_i (w+x)_{i+1} = w_{i+1} w_i$ when $i \in T_x$. Moreover, if $i \notin T_x$ then $x_i x_{i+1} = 00$, so that $(w+x)_i (w+x)_{i+1} = w_i w_{i+1}$. As a result, for any odd i , we have that

$$(w+x)_i (w+x)_{i+1} = \begin{cases} w_{i+1} w_i & \text{if } i \in T_x \\ w_i w_{i+1} & \text{if } i \notin T_x \end{cases}.$$

On the other hand, by the definition of α_x , for any odd i we have that

$$(w^{\alpha_x})_i (w^{\alpha_x})_{i+1} = \begin{cases} w_{i+1} w_i & \text{if } i \in T_x \\ w_i w_{i+1} & \text{if } i \notin T_x \end{cases}.$$

These altogether imply that $w^{\alpha_x} = w + x$. \square

Remark 4.2. Note that if $x = 11 \dots 11$, then $\alpha_x = \sigma$ in the lemma above.

The following result is a kind of generalization of Lemma 2.2 in the presence of an involutory fixed point free permutation automorphism. By Proposition 3.1, there is no binary linear code with length 4 and dimension 3 whose permutation automorphism group is isomorphic to C_2 . Note also that $\text{PAut}(\mathbb{F}_2^4) = S_4$. As a consequence of these, the result below is concerned with codes of length at least 6.

Theorem 4.3. *Suppose that C is a binary linear code of dimension $k \geq 3$ and length $n = 2m$ with $m \geq 3$ such that $\text{PAut}(C) = \langle \sigma \rangle$ for $\sigma = (1, 2)(3, 4) \dots (n-1, n)$, then $\dim(F_\sigma(C)) \neq k-1$ so that $\lceil k/2 \rceil \leq \dim(F_\sigma(C)) \leq k-2$. In particular, $k \neq 3$ or equivalently there is no 3-dimensional linear code C of even length with the property that $\text{PAut}(C) = \langle \sigma \rangle$.*

Proof. By Lemma 2.2, it is enough to show that $F_\sigma(C)$ can not be $(k-1)$ -dimensional. Suppose to the contrary that $F_\sigma(C)$ is $(k-1)$ -dimensional. Let $w \in C \setminus F_\sigma(C)$, then $w^\sigma \neq w$ and $w + w^\sigma \in F_\sigma(C)$.

If for some odd i , $w_i = w_{i+1}$ then the transposition $(i, i+1)$ fixes w as well as any element of $F_\sigma(C)$. Moreover, since $F_\sigma(C)$ is $(k-1)$ -dimensional any element of C can be written as $\lambda_1 w + \lambda_2 c$ where $c \in F_\sigma(C)$ and $\lambda_1, \lambda_2 \in \mathbb{F}_2$. Hence, $(i, i+1)$ fixes every element of C , that is $\text{PAut}(C) \neq \langle \sigma \rangle$, a contradiction.

Hence, for all odd $i \in \{1, 2, \dots, n\}$, $w_i \neq w_{i+1}$, and it follows that

$$(w + w^\sigma)_i = (w + w^\sigma)_{i+1} = w_i + w_{i+1} = 1,$$

that is $w + w^\sigma = y = (11 \dots 11)$. Now, since $\dim(F_\sigma(C)) = k-1 \geq 2$, there is a non-zero element $x \in F_\sigma(C)$ such that $x \neq y$, so $x \neq (11 \dots 11)$. Then T_x is non-empty, since x is a non-zero element of $F_\sigma(C)$. Also since $x \neq (11 \dots 11)$, we have that T_x is strictly contained T_y and hence the permutation

$$\alpha_x = \prod_{i \in T_x} (i, i+1)$$

is an involution in S_n which is different from σ . Hence, from Lemma 4.1, we have that $w^{\alpha_x} = w + x$. Since any element of C can be written as $\lambda_1 w + \lambda_2 c$ where $c \in F_\sigma(C)$ and $\lambda_1, \lambda_2 \in \mathbb{F}_2$ and that α_x fixes any element of $F_\sigma(C)$, we deduce that $\alpha_x \in \text{PAut}(C)$, which contradicts with the fact that $\text{PAut}(C) = \langle \sigma \rangle$. This finishes the proof. \square

Theorem 4.4. *There is no 4-dimensional binary linear code C of length $n = 2m$ with the property that $\text{PAut}(C) = \langle \sigma \rangle$ where $\sigma = (1, 2)(3, 4) \dots (n-1, n)$.*

Proof. Suppose to the contrary that C is a 4-dimensional binary linear code of length $n = 2m$ with the property that $\text{PAut}(C) = \langle \sigma \rangle$. If $n = 4$, we have that $C = \mathbb{F}_2^4$ and $\text{PAut}(C) = S_4$, so we have that $n = 2m \geq 6$. Furthermore, by Theorem 4.3, we have that $\dim(F_\sigma(C)) = 2$. Let $\{x, y\}$ be a basis for $F_\sigma(C)$ and extend it to a basis

$$\mathcal{B} = \{x, y, w, u\}$$

of C . Then $w + w^\sigma$ and $u + u^\sigma$ lie in $F_\sigma(C)$. If $w + w^\sigma = u + u^\sigma$, then we have that $w + u \in F_\sigma(C) \cap \langle w, u \rangle = \{0\}$, which implies that $w = u$. So we have that $w + w^\sigma \neq u + u^\sigma$. Hence without loss of generality, assume that $w + w^\sigma = x$ and $u + u^\sigma = y$. Then $i \in T_x$ if and only if $w_i \neq w_{i+1}$. Similarly, $i \in T_y$ if and only if $u_i \neq u_{i+1}$.

Note that $T_x \neq T_y$ since $x \neq y$. Set $T := T_x \cup T_y$.

Case 1: Assume that T is not equal to the set of all odd integers inside $\{1, 2, \dots, n\}$. Set

$$\alpha = \prod_{i \in T} (i, i+1).$$

Since T is non-empty and not equal to the set of all odd integers inside $\{1, 2, \dots, n\}$, α is an involution different from σ and it fixes both x and y . Moreover,

$$(w^\alpha)_i (w^\alpha)_{i+1} = \begin{cases} w_{i+1} w_i & \text{if } i \in T \\ w_i w_{i+1} & \text{if } i \notin T \end{cases}.$$

Since any $i \in T_x \iff w_i \neq w_{i+1}$, we have that

$$(w+x)_i (w+x)_{i+1} = \begin{cases} w_{i+1} w_i & \text{if } i \in T_x \\ w_i w_{i+1} = w_{i+1} w_i & \text{if } i \in T \setminus T_x \\ w_i w_{i+1} & \text{if } i \notin T \end{cases}.$$

Hence, $w^\alpha = w + x$. Similarly, it is easy to see that $u^\alpha = u + y$. Therefore, $\alpha \in \text{PAut}(C)$ which is a contradiction.

Case 2: Assume that T is equal to the set of all odd integers inside $\{1, 2, \dots, n\}$.

Case 2(a): One of $T_x \setminus T_y$ or $T_y \setminus T_x$ is an empty set (note that if both of these sets are empty then $T_x = T_y$ which is not possible). Without loss of generality assume that $T_x \setminus T_y = \emptyset$. Then T_x is strictly smaller than $T_y = T$. Consider the involution

$$\alpha_x = \prod_{i \in T_x} (i, i+1).$$

Then α_x is a non-trivial involution different from σ . We have that α_x fixes x and y . Since $T_x \subset T_y$, by Lemma 4.1, we have that $w^{\alpha_x} = w + x$ and $u^{\alpha_x} = u + x$. Therefore, $\alpha_x \in \text{PAut}(C)$ which is a contradiction.

Case 2(b): Both $T_x \setminus T_y$ and $T_y \setminus T_x$ has a unique element. Then it follows that $T_x \cap T_y$ has $m - 2 \geq 1$ elements.

Case 2(b) (i): Assume $m = 3$ or equivalently $T_x \cap T_y$ has a unique element. Then the length of the code is $2m = 6$ and so $\dim(C^\perp) = 6 - 4 = 2$. But this contradicts with Theorem ??, since $\text{PAut}(C) = \text{PAut}(C^\perp)$.

Case 2(b) (ii): Assume $T_x \cap T_y = \{i, j\}$ with the property that $u_i = u_j$ and $w_i \neq w_j$. Let $T_x \setminus T_y = \{k\}$ and $T_y \setminus T_x = \{l\}$. Set

$$\alpha = (k, k+1)(i, j)(i+1, j+1).$$

Then α is a non-trivial involution different from σ . Since $x_i x_{i+1} = 11 = x_j x_{j+1}$ and $y_i y_{i+1} = 11 = y_j y_{j+1}$, we have that α fixes x and y . Moreover, by the assumption on u , it follows that $u_{i+1} = u_{j+1}$ and $u_k = u_{k+1}$ and so we have that α fixes u . Since $k, i, j \in T_x$, we have that $w_k \neq w_{k+1}$ and also $w_i \neq w_j \implies w_j = w_{i+1}$ and $w_{j+1} = w_i$, we have that for any odd r

$$(w^\alpha)_r (w^\alpha)_{r+1} = \begin{cases} w_{k+1} w_k & \text{if } r = k \\ w_j u_{j+1} & \text{if } r = i \\ w_i w_{i+1} & \text{if } r = j \\ w_r w_{r+1} & \text{otherwise} \end{cases}$$

$$(w+x)_r (w+x)_{r+1} = \begin{cases} w_{k+1} w_k & \text{if } r = k \\ w_j u_{j+1} & \text{if } r = i \\ w_i w_{i+1} & \text{if } r = j \\ w_r w_{r+1} & \text{otherwise} \end{cases}.$$

Therefore, we get that $w^\alpha = w + x$. So $\alpha \in \text{PAut}(C)$ and we have a contradiction.

Case 2(b) (iii): Assume $T_x \cap T_y$ has at least two elements i and j with the property that $u_i = u_j$ and $w_i = w_j$. Set

$$\alpha = (i, j)(i+1, j+1).$$

Then since $x_i x_{i+1} = 11 = x_j x_{j+1}$ and $y_i y_{i+1} = 11 = y_j y_{j+1}$, we have that α fixes x and y . Since i and j satisfy $u_i = u_j$ and $w_i = w_j$, then it follows that $u_{i+1} = u_{j+1}$ and $w_{i+1} = w_{j+1}$ since $u_i \neq u_{i+1}$, $u_j \neq u_{j+1}$. So α fixes both w and u . Hence, $\alpha \in \text{PAut}(C)$, a contradiction since α is an involution which is different from σ .

Case 2(b) (iv): Assume $T_x \cap T_y$ has at least two elements i and j with the property that $u_i \neq u_j$ and $w_i \neq w_j$. Set

$$\alpha = (i, j+1)(i+1, j).$$

Then since $x_i x_{i+1} = 11 = x_j x_{j+1}$ and $y_i y_{i+1} = 11 = y_j y_{j+1}$, we have that α fixes x and y . Since i and j satisfy $u_i \neq u_j$ and $w_i \neq w_j$, then it follows that $u_i = u_{j+1}$ and $w_i = w_{j+1}$ since $u_i \neq u_{i+1}$, $u_j \neq u_{j+1}$. Similarly, $u_{i+1} = u_j$ and $w_{i+1} = w_j$. So α fixes both w and u . Hence,

$\alpha \in \text{PAut}(C)$, a contradiction since α is an involution which is different from σ .

Case 2(b) (v): Assume $T_x \cap T_y$ has at least three elements. Suppose there is no pair of elements which satisfies Case 2(b)(iii) and Case 2(b)(iv). But this is not possible. Indeed, say $i, j, k \in T_x \cap T_y$ such that $u_i = u_j$ and $w_i \neq w_j$. If $u_i = u_k$ and $w_i \neq w_k$, then we have that $u_j = u_k$ and $w_j = w_k$, hence we are in Case 2(d)(ii). If $u_i \neq u_k$ and $w_i = w_k$, then we have that $u_j \neq u_k$ and $w_j \neq w_k$, hence we are in Case 2(d)(iii). So there is a pair of elements which satisfies Case 2(b)(iii) and Case 2(b)(iv) and we are done by using the same methods in these cases.

Case 2(c): Assume that one of $T_x \setminus T_y$ or $T_y \setminus T_x$ has at least three elements. Without loss of generality assume that $T_x \setminus T_y$ has at least three elements. Among these elements there exist $k, l \in T_x \setminus T_y$ such that $u_k u_{k+1} = u_l u_{l+1}$ (since $k, l \notin T_y$ we have that $u_k u_{k+1}, u_l u_{l+1} \in \{00, 11\}$). Since $k, l \in T_x$ we have that $w_k \neq w_{k+1}$ and $w_l \neq w_{l+1}$. Let us choose $k_1 \in \{k, k+1\}$ and $l_1 \in \{l, l+1\}$ such that $w_{k_1} = w_{l_1}$. Then $\alpha = (k_1, l_1)$ fixes w and u . It is not difficult to see that α fixes x and y also since $k, l \in T_x \setminus T_y$. So $\alpha \in \text{PAut}(C)$ and this is a contradiction since α is an involution different from σ .

Note that the same proof works in the case that $T_x \setminus T_y = \{k, l\}$ has exactly two elements and $u_k u_{k+1} = u_l u_{l+1}$.

Case 2(d): Assume that one of $T_x \setminus T_y$ or $T_y \setminus T_x$ has exactly two elements which does not satisfy the condition in the previous sentence. Suppose that $T_x \setminus T_y = \{k, l\}$ and $u_k u_{k+1} \neq u_l u_{l+1}$. Since $k, l \in T_x$, we have that $x_k x_{k+1} = 11 = x_l x_{l+1}$ and $w_k \neq w_{k+1}$ and $w_l \neq w_{l+1}$. Also, since $k, l \notin T_y$, we have that $y_k y_{k+1} = 00 = y_l y_{l+1}$ and $u_k u_{k+1}, u_l u_{l+1} \in \{00, 11\}$. Let $k_1 \in \{k, k+1\}$ and $l_1 \in \{l, l+1\}$ such that $w_{k_1} = w_{l_1}$ so that $w_{\sigma(k_1)} = w_{\sigma(l_1)}$. Consider the involution

$$\alpha = \left(\prod_{i \in T_y \setminus T_x} (i, i+1) \right) (k_1, l_1) (\sigma(k_1), \sigma(l_1)).$$

It is easy to see that α is a non-trivial involution which is different from σ . Moreover, α fixes x and y . Since $w_i = w_{i+1}$ for all $i \in T_y \setminus T_x$, it is easy to see that α fixes w . For any odd r , we have that

$$(u^\alpha)_r (u^\alpha)_{r+1} = \begin{cases} u_{r+1} u_r & \text{if } r \in T_y \setminus T_x \\ u_r u_{r+1} & \text{if } r \in T_x \cap T_y \\ u_l u_{l+1} = u_{l+1} u_l & \text{if } r = k \\ u_k u_{k+1} = u_{k+1} u_k & \text{if } r = l \end{cases}.$$

Note that for odd r , we have that $(x+y)_r (x+y)_{r+1} = 00$ if and only if $r \in T_x \cap T_y$. Also for $r \in T_y \setminus T_x$, we have that $u_r \neq u_{r+1}$ and $u_k u_{k+1} + 11 = u_l u_{l+1}$ since $u_k u_{k+1} \neq u_l u_{l+1}$. As a result, we have that

$$(u+x+y)_r (u+x+y)_{r+1} = \begin{cases} u_{r+1} u_r & \text{if } r \in T_y \setminus T_x \\ u_r u_{r+1} & \text{if } r \in T_x \cap T_y \\ u_l u_{l+1} & \text{if } r = k \\ u_k u_{k+1} & \text{if } r = l \end{cases}.$$

Hence, we deduce that $u^\alpha = u+x+y$ and it follows that $\alpha \in \text{PAut}(C)$. However, α is an involution different from σ so we have a contradiction.

Since we get a contradiction in all possible cases, the proof is finished. \square

Corollary 4.5. *There is no binary linear code C of dimension k and length $n = 2m$ with $m \geq 2$ such that $\text{PAut}(C) = \langle \sigma \rangle$ where $k = i$ or $k = n - i$ for $i \in \{1, 2, 3, 4\}$ and $\sigma = (1, 2)(3, 4) \dots (n-1, n)$.*

Proof. Follows from Proposition 3.1, Theorem 3.2, Theorem 4.3 and Theorem 4.4 since $\text{PAut}(C) = \text{PAut}(C^\perp)$ and $\dim(C) = n - \dim(C^\perp)$. \square

Corollary 4.6. *There is no binary quasi group code C of dimension k and length $n = 2m$ with $m \geq 2$ such that $\text{PAut}(C) \cong C_2$ where $k = i$ or $k = n - i$ for $i \in \{1, 2, 3, 4\}$.*

Proof. Suppose that C is a binary quasi group code such that $\text{PAut}(C) \cong C_2$. Then $\text{PAut}(C) = \langle \gamma \rangle$ where γ is a fixed point free involution in S_n . Then γ and σ are conjugate elements of S_n , that is $\sigma = \beta^{-1}\gamma\beta$ for some $\beta \in S_n$. Let $C_1 = C^\beta$, then $\text{PAut}(C_1) = \langle \sigma \rangle$. But then Corollary 4.5 gives us a contradiction. \square

Corollary 4.7. *There is no binary linear code C of length $2 < n = 2m \leq 8$ with $\text{PAut}(C) = \langle \sigma \rangle$ where $\sigma = (1, 2)(3, 4) \dots (n - 1, n)$.*

Proof. Follows from Corollary 4.5. \square

Corollary 4.8. *There is no binary quasi group code C of length $2 < n = 2m \leq 8$ with $\text{PAut}(C) \cong C_2$.*

Proof. Follows from Corollary 4.6. \square

Our computer calculations give us the impression that the generalization of Corollary 4.8 may be possible. But we can not give a proof of it, as can be seen from the proof of Theorem 4.4 there will be many cases to consider. We are claiming the following is true.

Conjecture 4.9. *There is no binary quasi group code C of length $n = 2m > 2$ such that $\text{PAut}(C) \cong C_2$.*

5. A GENERALIZATION AND AN APPLICATION TO SOME PUTATIVE EXTREMAL SELF-DUAL BINARY CODES

In this section, we present a generalization of the ideas that are used in the proof of Theorem 4.4. By applying this generalization to the putative extremal self-dual binary [72, 36, 16] code and putative extremal self-dual binary [96, 48, 20] code, we obtain results concerning the structure of these codes.

Let C be a binary linear code of length $n = 2m$ and $\sigma = (1, 2)(3, 4) \dots (n - 1, n) \in \text{PAut}(C)$. We have that the fixed subcode $F_\sigma(C)$ is a subcode of C . So there is a subcode $N_\sigma(C)$ such that

$$C = F_\sigma(C) \oplus N_\sigma(C).$$

Let $\{w_{(1)}, \dots, w_{(k-f)}\}$ be a basis of $N_\sigma(C)$ where $k = \dim(C)$ and $f = \dim(F_\sigma(C))$. Then by Lemma 2.1, we have that $f \geq (k - f)$. On the other hand, by the proof of Lemma 2.1, there is a linearly independent subset $\{x_{(1)}, \dots, x_{(k-f)}\}$ of $F_\sigma(C)$ where $x_{(j)} = w_{(j)} + w_{(j)}^\sigma$ for $j = 1, \dots, k - f$. Now extend $\{x_{(1)}, \dots, x_{(k-f)}\}$ to a basis $\{x_{(1)}, \dots, x_{(f)}\}$ of $F_\sigma(C)$. Set

$$T(\sigma) := T_{x_{(1)}} \cup T_{x_{(2)}} \cup \dots \cup T_{x_{(k-f)}}$$

where $T_{x_{(j)}}$ is defined as in Section 4. Then note that if $i \notin T(\sigma)$ then $(x_{(j)})_i(x_{(j)})_{i+1} = 00$ for any $j = 1, \dots, k - f$. So if $i \notin T(\sigma)$ the equation $x_{(j)} = w_{(j)} + w_{(j)}^\sigma$ implies that $(w_{(j)})_i = (w_{(j)})_{i+1}$ for any $j = 1, \dots, k - f$.

Theorem 5.1. *Let C be a binary linear code of length $n = 2m$ where $m \geq 2$ such that $\sigma = (1, 2)(3, 4) \dots (n - 1, n) \in \text{PAut}(C)$. Suppose that $T(\sigma) \neq \{1, 3, \dots, n - 1\}$. Then we have that $\text{PAut}(C) \neq \langle \sigma \rangle$. In particular, $\text{PAut}(C)$ has automorphisms which have some fixed points and that $C_2 \times C_2 \leq \text{PAut}(C)$.*

Proof. Let C be a binary linear code of dimension k . Let $\beta = \prod_{i \notin T(\sigma)} (i, i + 1)$. Since $T(\sigma) \neq \{1, 3, \dots, n - 1\}$, the permutation β is a non-trivial involution. Also if $i \notin T(\sigma)$ then $(x_{(j)})_i(x_{(j)})_{i+1} = 00$ and $(w_{(j)})_i = (w_{(j)})_{i+1}$ for any $j = 1, \dots, k - f$. Hence β fixes all elements in $F_\sigma(C)$ and $N_\sigma(C)$ so it lies inside $\text{PAut}(C)$. It is clear that β has some fixed points and $\langle \sigma, \beta \rangle \cong C_2 \times C_2$ \square

Corollary 5.2. *Let C be a binary self-dual [72, 36, 16] code. If $\sigma = (1, 2)(3, 4) \dots (71, 72) \in \text{PAut}(C)$, then $T(\sigma)$ is equal to the set of all odd integers between 1 and 71.*

Proof. Suppose to the contrary that $T(\sigma) \neq \{1, 3, \dots, 71\}$. Then by Theorem 5.1, there is an involution in $\text{PAut}(C)$ which is not fixed point free. But this contradicts with [2, Theorem 5.3]. So $T_\sigma = \{1, 3, \dots, 71\}$. □

Corollary 5.3. *Let C be a binary self-dual [96, 48, 20] code. If $\sigma = (1, 2)(3, 4) \dots (95, 96) \in \text{PAut}(C)$, then $T(\sigma)$ is equal to the set of all odd integers between 1 and 95.*

Proof. Suppose to the contrary that $T(\sigma) \neq \{1, 3, \dots, 95\}$. Then by Theorem 5.1, there is an involution in $\text{PAut}(C)$ which is not fixed point free. But this contradicts with [1, Theorem 8]. So $T(\sigma) = \{1, 3, \dots, 95\}$. □

REFERENCES

- [1] S. Bouyuklieva. A method for constructing self-dual codes with an automorphism of order 2. *IEEE Transactions on Information Theory* **46** (2), (2000), 496–504.
- [2] S. Bouyuklieva. On the automorphisms of order 2 with fixed points for the extremal self-dual codes of length 24m. *Des. Codes Cryptogr.* **25**, (2002), 5–13.
- [3] M. Borello. The automorphism group of a self-dual [72, 36, 16] code is not an elementary abelian group of order 8. *Finite Fields Appl.* **25** (2014), 1–7.
- [4] M. Borello. On automorphism groups of binary linear codes. *Topics in Finite Fields, Contemp. Math.* **632** (2015), 29–41.
- [5] M. Borello, J. de la Cruz. Some new results on the self-dual [120, 60, 24] code. *Finite Fields Appl.* **50** (2018), 17–34.
- [6] J. J. Bernal, A. del Rio, J. J. Simón. An intrinsical description of group codes. *Des. Codes Cryptogr.* **51.3**, (2009), 289–300.
- [7] M. Borello, W. Willems. On the algebraic structure of quasi-group codes. to appear in *J. Algebra and its Appl.* (2022), arXiv:1912.09167.

MIMAR SINAN FINE ARTS UNIVERSITY, DEPARTMENT OF MATHEMATICS, 34380, BOMONTI, ŞİŞLİ, ISTANBUL, TURKEY

Email address: fatma.altunbulak@msgsu.edu.tr

GEBZE TECHNICAL UNIVERSITY, DEPARTMENT OF MATHEMATICS, 41400, GEBZE, KOCAELI, TURKEY

Email address: roghayeh@gtu.edu.tr

MIMAR SINAN FINE ARTS UNIVERSITY, DEPARTMENT OF MATHEMATICS, 34380, BOMONTI, ŞİŞLİ, ISTANBUL, TURKEY

Email address: ipek.tuvay@msgsu.edu.tr