

PROPORTIONAL FAIR DIVISION OF MULTI-LAYERED CAKES

A PREPRINT

 **Mohammad Azharuddin Sanpui**
 Department of Mathematics
 Indian Institute of Technology Kharagpur
 India
 azharuddinsanpui123@gmail.com

ABSTRACT

We study the multi-layered cake cutting problem, where the multi-layered cake is divided among agents proportionally. This problem was initiated by Hosseini et al.(2020) under two constraints, one is contiguity and the other is feasibility. Basically we will show the existence of proportional multi-allocation for any number of agents with any number of preferences that satisfies contiguity and feasibility constraints using the idea of switching point for individual agent and majority agents. First we show that exact feasible multi-allocation is guaranteed to exist for two agents with two types of preferences. Second we see that we always get an envy-free multi-allocation that satisfies the feasibility and contiguity constraints for three agent with two types of preferences such that each agent has a share to each layer even without the knowledge of the unique preference of the third agent. Using the switching point property we will prove the existence of proportional multi-allocation that is feasible and contiguous over 3-layered cake for $n \geq 3$ agents and generalize this idea when the number of layers is odd and number of agents always greater than number of layers. We also show the existence of proportional multi-allocation that satisfies the feasibility and contiguity constraints over $2^a P^b$ -layered cake for $n \geq 2^a P^b$ agents where $a \in \mathbb{Z}_+$, $b \in \mathbb{Z}_+ \cup \{0\}$ and P is prime. We further prove the existence of proportional multi-allocation over m' -layered cake for n' agents ensures the existence of proportional multi-allocation over $2^a m'^b$ -layered cake for $2^a n'^b$ agents where $n' \geq m'$, $a \in \mathbb{Z}_+$ and $b \in \{0, 1\}$.

Keywords Proportional Cake Cutting · Multi-layered Cake · Social Choice Theory

1 Introduction

In our daily lives, there are many examples of time scheduling whereby we set our time in such a manner that we can do our daily essential work. Consider a group of students of a university who want to use multiple facilities, such as a seminar room or an indoor games room. The start and closed times of the two facilities are the same. Each student of the group has a different preferred time duration to get each facility room and everyone is also willing to take both facilities.

The problem of fair division of a divisible heterogenous resource among different agents with their different preferences over divisible resource has been studied in the classical cake cutting model (Steinhaus [1949],Brams and Jones [2006],Procaccia [2013],Kurokawa et al. [2013]). We assume a cake as the unit interval $[0, 1]$ that is divided among agents where each agents has different perference over the entire cake. The concept of fair division was given by the three mathematicians, Hugo Steinhaus, Bronis law Knaster and Stefan Banach to a meeting in the Scottish Cafe in Lvov (in Poland) to the end of the world war II. Many cake cutting techniques have been developed over few decades; specially, two types fair division: proportional division where each agent gets his share at least $\frac{1}{n}$, n is the number of agents and envy-free division where each agent i receives an allocated piece of cake that has highly valuation than others' piece of cake with respect to his valuation (Saberri et al. [2012],Cloutier et al. [2010]).

In the above example of taking multiple facilities, we cannot think this problem as a single layered cake cutting problem. Because here we have to divided two time intervals independently in such a way that the each agent can get both facilities. Now the question is how to allocate the time intervals of the facilities according to their preferences in a fair

way. That means every student can get each facility and their time intervals for each facility never overlap. Adopting the above constraints, Hosseini, Igarshi, and Searns have initiated the multi-layered cake cutting problem in 2020. In the multi-layered cake cutting problem, We see how to solve this problem. We consider each facility as a divisible heterogeneous layer of a multi-layered cake. Every student has additive preference, called *valuation* over disjoint (non-overlapping) intervals. Note that the valuation of the same part of the cake for different students can be very different. A division of a multi-layered cake is feasible if no student's time intervals contain overlapping intervals. The division is contiguous if each student gets a contiguous time interval for each facility. Some work has been done on dividing multiple cakes among agents (Nicolas Lebert and Frédéric Meunier and Quentin Carbonneaux [2013], Cloutier et al. [2010], Segal-Halevi [2021]). But these works are not consider as multi-layered cake cutting under feasible and contiguous constraints.

1.1 Problem definition

Let $\mathcal{N} = \{1, 2, \dots, n\}$ be a set of n agents and $\mathcal{C} = (\mathcal{L}_1, \mathcal{L}_2, \dots, \mathcal{L}_m)$ be an m -layered cake where for each l , \mathcal{L}_l is l -th layer of the cake and each \mathcal{L}_l is a subinterval of the unit interval $\mathcal{I} = [0, 1] \subset \mathbb{R}$. The valuation function of the agent a_i for layer \mathcal{L}_l is a function representing the preference of the agent over different parts of \mathcal{L}_l denoted by $V_{il} : [0, 1] \rightarrow \mathbb{R}_0^+$ where $a_i = i$ for $i \in \mathcal{N}$. Furthermore, to make sure that the preference of each agent for the whole cake is 1, the valuation function is normalized, with $\sum_{1 \leq l \leq m} V_{il}(\mathcal{L}_l) = 1$. A division of a multi-layered cake is said to be *feasible* if no agent's share contain overlapping intervals, and *contiguous* if each allocated piece of a layer is contiguous. Intuitively, this means that from each layer, an agent is going to get a continuous piece. Furthermore, an agent is never going to get pieces from two different layers that overlap vertically. This condition is natural due to the example given for scheduling previously. For example, if a student is already using a seminar room, he cannot use the indoor games room at the same time.

Assume that V_i and V_{il} are the valuation functions over the entire cake \mathcal{C} and l -th layer cake \mathcal{L}_l corresponding to each agent a_i . Now we want to find a complete multi-allocation $\mathcal{P} = (\mathcal{P}_1, \mathcal{P}_2, \dots, \mathcal{P}_n)$ over the entire cake \mathcal{C} that satisfies contiguity and feasibility conditions where each $\mathcal{P}_i = (P_{i1}, P_{i2}, \dots, P_{im}) \subset \mathbb{R}^m$, \mathcal{P}_i is called a layered piece fixed to the agent a_i and $\bigcup_{i=1}^n P_{il} = \mathcal{L}_l$ for each layer l . That means corresponding to each agent a_i and each layer l , P_{il} is a contiguous piece of l -th layered cake \mathcal{L}_l and for each layered piece \mathcal{P}_i , any two pieces of cake, P_{ip} and P_{iq} never overlap where $p \neq q$.

We have the following different types of fairness notions for cake cutting.

A multi-allocation $\mathcal{P} = (\mathcal{P}_1, \mathcal{P}_2, \dots, \mathcal{P}_n)$ of an m -layered cake \mathcal{C} for n agents is said to be **proportional** if for every $i \in \mathcal{N}$, $V_i(\mathcal{P}_i) = \sum_{l=1}^m V_{il}(P_{il}) = \sum_{l=1}^m \sum_{I \in P_{il}} V_{il}(I) \geq \frac{1}{n}$.

A multi-allocation $\mathcal{P} = (\mathcal{P}_1, \mathcal{P}_2, \dots, \mathcal{P}_n)$ of an m -layered cake \mathcal{C} for n agents is said to be **envy-free** if for any two different $i, j \in \mathcal{N}$, $V_i(\mathcal{P}_i) \geq V_j(\mathcal{P}_j) \Rightarrow \sum_{k=1}^m V_{ik}(P_{ik}) \geq \sum_{k=1}^m V_{jk}(P_{jk})$.

A multi-allocation $\mathcal{P} = (\mathcal{P}_1, \mathcal{P}_2, \dots, \mathcal{P}_n)$ of an m -layered cake \mathcal{C} for n agents is said to be **exact** if for any two different $i, j \in \mathcal{N}$, $V_i(\mathcal{P}_j) = \frac{1}{n} \Rightarrow \sum_{k=1}^m V_{ik}(P_{ik}) = \frac{1}{n}$.

Intuitively, envy-freeness means that no agent will want to exchange his share of the cake with that of some other agent.

1.2 Our Results

Specially We focus on proportional multi-allocation of a multi-layered cake, under feasibility and contiguity constraints. We show the existence of proportional multi-allocation over 3-layered cake for 3 agents that is feasible and contiguous using the switching point property. we also see that we always get an envy-free multi-allocation that satisfies the feasibility and contiguity constraints for three agent with two types of preferences such that each agent has a share to each layer even without the knowledge of the unique preference of the third agent. We further show the existence of proportional multi-allocation over 3-layered cake of $n \geq 3$ agents that is feasible and contiguity.

Agents(n)	Layers(m)	Fairness	feature
2	2	Exact	Multi-allocation satisfies feasibility constraint only
3	2	EF	Every agent has a share to each layer
3	3	Prop	Use switching point property
$n \geq 3$	3	Prop	Use switching point property
$n \geq 2^a 3^b$	$2^a 3^b, a \in \mathbb{Z}_+, b \in \{0, 1\}$	Prop	Use switching point property
any $n \geq m$	$2^a P^b$	Prop	$a \in \mathbb{Z}_+, b \in \mathbb{Z}_+ \cup \{0\}$ P is prime
$2^a n'^b$	$2^a m'^b$	Prop	$a \in \mathbb{Z}_+, b \in \mathbb{Z}_+ \cup \{0\}$, P is prime †

† assumes that an m' -layered cake with n' agents has a multi-allocation that is contiguous and feasible. Note that when $m > n$, then no feasible multi-allocation exists.

2 Preliminaries

Naturally, in the cake cutting problem the cake is the unit interval $[0, 1]$ and a piece of cake is finite union of disjoint subintervals of $[0, 1]$ (Dubins and Spanier [1961], Even and Paz [1984], Robertson and Webb [1998]). If $x, y \in \mathbb{R}$ and $x \leq y$, then $[x, y] = \{r | x \leq r \leq y\}$ is called an interval. If $x, y \in [0, 1]$, then $[x, y]$ is called a subinterval of the unit interval $[0, 1]$. A subinterval of $[0, 1]$ is called a contiguous piece of the cake. Specially two types fairness on the allocation of cakes are very well, one is proportional and another is envy-free (Tasnádi [2003], Aziz and Mackenzie [2020], Haris Aziz and Simon Mackenzie [2016]).

We consider the multi-layered cake cutting model that was initiated by Hosseini, Igarashi, and Searns (Hosseini et al. [2020]). In the multi-layered cake cutting problem, the number of layers and the number of agents are given. The setting of the model includes a set $\mathcal{N} = \{1, 2, \dots, n\}$ of agents and a set $\mathcal{M} = \{1, 2, \dots, m\}$ of layers. An m -layered cake is denoted by $\mathcal{C} = (\mathcal{L}_l)_{l \in \mathcal{M}}$ where \mathcal{L}_l is an interval such that $\mathcal{L}_l \subseteq [0, 1]$ for each $l \in \mathcal{M}$. Corresponding to each $l \in \mathcal{M}$, we call \mathcal{L}_l as l -th layer cake and l as the l -th layer.

Corresponding to each l -th layer, each agent $i \in \mathcal{N}$ is endowed with a non-negative integrable density function $v_{il} : \mathcal{L}_l \rightarrow \mathbb{R}_0^+$ where $l \in \mathcal{M}$. The valuation function of the agent a_i for layer \mathcal{L}_l is a function representing the preference of the agent over different parts of \mathcal{L}_l denoted by $V_{il} : [0, 1] \rightarrow \mathbb{R}_0^+$. If P is a piece of cake of l -th layer, then $V_{il}(P)$ represents the value assigned to it by the agent a_i i.e., $V_{il}(P) = \sum_{I \in P} \int_{x \in I} v_{il}(x) dx$. We consider the total valuation of each agent a_i over the entire cake is 1 i.e., $\sum_{1 \leq l \leq m} V_{il}(\mathcal{L}_l) = 1$.

A sequence $P = (P_l)_{l \in \mathcal{M}}$ of pieces of each layer $l \in \mathcal{M}$ of multi-layered cake is called a layered piece. A layered piece $P = (P_l)_{l \in \mathcal{M}}$ is said to be contiguous if for each $l \in \mathcal{M}$, P_l is a *contiguous* piece of the layer l . A layered piece $P = (P_l)_{l \in \mathcal{M}}$ is said to be *non-overlapping*, if for any two pieces from different layers never overlap i.e., for any two different layers $l, l' \in \mathcal{M}$ and for any $I \in P_l$ and $I' \in P_{l'}$ either $I \cap I' = \emptyset$ or $I \cap I' = \{b\}$ where b is the endpoint of both I and I' . We assume the valuation functions related to layers of each agent a_i are additive over layers and written as $V_i(P) = \sum_{1 \leq l \leq m} V_{ij}(P_j)$ where V_i is the valuation function of each agent i over the entire cake where $i \in \mathcal{N}$.

Let P and P' are two layered pieces. If for any $i \in \mathcal{N}$, $V_i(P) \geq V_i(P')$ then it is said that the agent a_i weakly prefers the layered piece P to the layered piece P' . A multi-allocation $\mathcal{P} = (\mathcal{P}_1, \mathcal{P}_2, \dots, \mathcal{P}_n)$ is a partition of the multi-layered cake where each $\mathcal{P}_i = (P_{ij})_{j \in \mathcal{M}}$ is a layered piece of the cake is assigned to agent i where $i \in \mathcal{N}$. Corresponding to a multi-allocation \mathcal{P} and an agent $i \in \mathcal{N}$, the valuation of each agent i is $V_i(\mathcal{P}_i) = \sum_{j \in \mathcal{M}} V_{ij}(P_{ij})$. A multi-allocation \mathcal{P} is said to be complete if for each $j \in \mathcal{M}$, $\bigcup_{i=1}^n P_{ij} = \mathcal{C}_j$. A multi-allocation $\mathcal{P} = (\mathcal{P}_1, \mathcal{P}_2, \dots, \mathcal{P}_n)$ is said to be

- contiguous if for each $i \in \mathcal{N}$, \mathcal{P}_i is contiguous.
- feasible if for each $i \in \mathcal{N}$, \mathcal{P}_i is non-overlapping.

Note that throughout in this paper proportional multi-allocation means that multi-allocation always satisfies contiguity and feasibility constraints and $n \geq m$, due to feasibility constraint.

Fairness notions. Different types of fairness are defined on the resource allocation (Berliant et al. [1992], Thomson [2011]). Cake cutting is a problem of the allocation of divisible resource and there are many fairness defined on the allocation of cake cutting problem (Segal-Halevi and Nitzan [2019], Segal-Halevi [2021]).

A multi-allocation \mathcal{P} of an m -layered cake \mathcal{C} for n agents is said to be

- *proportional* if for every $i \in \mathcal{N}$, $V_i(\mathcal{P}_i) = \sum_{j=1}^m V_{ij}(P_{ij}) = \geq \frac{1}{n}$
- *envy-free* if for any $i, j \in \mathcal{N}$, $V_i(\mathcal{P}_i) \geq V_j(\mathcal{P}_j)$ i.e., $\sum_{k=1}^m V_{ik}(\mathcal{P}_{ik}) \geq \sum_{k=1}^m V_{jk}(\mathcal{P}_{jk})$
- *equitable* if for any $i, j \in \mathcal{N}$, $V_i(\mathcal{P}_i) = V_j(\mathcal{P}_j)$ i.e., $\sum_{k=1}^m V_{ik}(\mathcal{P}_{ik}) = \sum_{k=1}^m V_{jk}(\mathcal{P}_{jk})$
- *exact* if for any $i, k \in \mathcal{N}$, $V_i(\mathcal{P}_k) = \frac{1}{n}$ i.e., $\sum_{1 \leq j \leq m} V_{ij}(P_{kj}) = \frac{1}{n}$

The m (even)-layered cut. For m -layered cake \mathcal{C} (m is even), Hosseini, Igarashi, and Searns have defined a particular way to partition the entire multi-layered cake \mathcal{C} into diagonal pieces that follows feasibility constraint. For any $x \in \mathcal{I} = [0, 1]$, defined :

1. $LR(x, \mathcal{C}) = (\bigcup_{l=1}^{\frac{m}{2}} \mathcal{L}_l \cap [0, x]) \cup (\bigcup_{l=\frac{m}{2}+1}^m \mathcal{L}_l \cap [x, 1])$
2. $RL(x, \mathcal{C}) = (\bigcup_{l=1}^{\frac{m}{2}} \mathcal{L}_l \cap [x, 1]) \cup (\bigcup_{l=\frac{m}{2}+1}^m \mathcal{L}_l \cap [0, x])$

Consider $\mathcal{S}_1 = \{1, 2, \dots, \frac{m}{2}\}$ and $\mathcal{S}_2 = \{\frac{m}{2} + 1, \frac{m}{2} + 2, \dots, m\}$.

- $LR(x, \mathcal{C})$ consists of all subintervals of type $[0, x]$ of each layer $l \in \mathcal{S}_1$ and subintervals of type $[x, 1]$ of each layer $l \in \mathcal{S}_2$.

- $RL(x, \mathfrak{C})$ consists of all subintervals of type $[0, x]$ of each layer $l \in \mathcal{S}_2$ and subintervals of type $[x, 1]$ of each layer $l \in \mathcal{S}_1$.




Figure 1: Diagonal pieces when $x = \frac{1}{2}$ and $m = 4$.

Computational model. In the cake cutting problem there are different types computational model. These are Robertson–Webb query model (Robertson and Webb [1998]), Direct revelation model (Menon and Larson [2017]), Moving-knife model (Austin [1982]), Simultaneous queries model (Balkanski et al. [2014]) etc. Robertson-Webb query model plays an important role in the cake cutting problem. This model has been generalized to two-dimensional cakes (Segal-Halevi et al. [2015]) and multi-dimensional cakes (Cseh and Fleiner [2018]). Following Robertson–Webb query model, Hosseini, Igarashi, and Searns have introduced a new computational model in the multi-layered cake cutting problem where there are two types of queries, one is a short knife and the other is long knife.

1. **Short Knife:** This query is applied on a layer of a multi-layered cake \mathfrak{C} .

- Short evaluation query:** For any known interval $[x, y]$ of the l -th layered cake \mathcal{L}_l of the m -layered cake \mathfrak{C} , $eval_l(i, x, y)$ implies the happiness $V_{ij}([x, y])$ of the agent a_i in the interval $[x, y]$ of l -th layer cake \mathcal{L}_l . Here $[x, y] \subseteq \mathcal{L}_l \subseteq [0, 1]$.
- Short cut query:** For any known point x on the l -th layered cake \mathcal{L}_j and $p \in [0, 1]$, $cut_l(i, x, p)$ indicates the minimum point y on the l -th layered cake \mathcal{L}_l for which $V_{ij}([x, y]) = p$.




Figure 2: Short knife is applied on first layer such that $V_{i1}(\mathcal{L}_1 \cap [x, y]) = p_i$ where $i \in \mathcal{N}$.

2. **Long Knife:** This query is applied on the entire multi-layered cake \mathfrak{C} .

- Long evaluation query:** For any known point $x \in [0, 1]$, $eval(i, x)$ implies the happiness $V_i(LR(x))$ of the agent a_i for $LR(x)$.
- Long cut query:** For any known $p \in [0, 1]$, $cut(i, p)$ indicates the minimum point x for which the happiness of the agent a_i should be p for cake piece $LR(x)$ if p exists.




Figure 3: Long knife is applied over entire cake at $x = \frac{1}{2}$ such that $V_i(LR(x, \mathfrak{C})) = p_i$ where $i \in \mathcal{N}$.

Switching point. Now we will look for a point in the interval $x \in [0, 1]$ which divides the entire m -layered cake \mathfrak{C} into two pairs of diagonal pieces, $LR(x)$ and $RL(x)$, such that both have the same valuation for a particular agent i.e., If there exists a point $x \in [0, 1]$, for which the valuation of the agent a_i for two pairs of diagonal pieces is same i.e. $V_i(LR(x)) = V_i(RL(x))$. Then the point x is called a switching point over the entire cake \mathfrak{C} for the agent a_i . Existence of such type point was introduced by Hosseini, Igarashi, and Searns. The value of an agent a_i over the entire m -layered cake is $V_i(\mathfrak{C})$ where $i \in \mathcal{N}$. Assume that the number of layer is even.

So $V_i(LR(x)) + V_i(RL(x)) = V_i(\mathfrak{C})$ i.e., either $V_i(LR(x)) \geq \frac{V_i(\mathfrak{C})}{2}$ and $V_i(RL(x)) \leq \frac{V_i(\mathfrak{C})}{2}$ or, $V_i(LR(x)) \leq \frac{V_i(\mathfrak{C})}{2}$ and $V_i(RL(x)) \geq \frac{V_i(\mathfrak{C})}{2}$. Consider a function $s : [0, 1] \rightarrow \mathbb{R}_0^+$ such that $s(x) = V_i(LR(x))$ for $x \in [0, 1]$. Since $s(x) = \sum_{1 \leq l \leq \frac{m}{2}} V_{il}(\mathcal{C}_l \cap [0, x]) + \sum_{\frac{m}{2} + 1 \leq l \leq m} V_{il}(\mathcal{C}_l \cap [x, 1])$ and each function V_{il} is continuous. So s is continuous on the interval $[0, 1]$. Since each V_{il} is non-atomic. Then either $s(0) \geq \frac{V_i(\mathfrak{C})}{2}$ and $s(1) \leq \frac{V_i(\mathfrak{C})}{2}$ or, $s(0) \leq \frac{V_i(\mathfrak{C})}{2}$ and $s(1) \geq \frac{V_i(\mathfrak{C})}{2}$. So there exists a point $y \in [0, 1]$ such that $s(y) = \frac{V_i(\mathfrak{C})}{2}$ by the intermediate theorem for continuous functions. This point y is the switching point for the agent a_i over the entire cake where $i \in \mathcal{N}$.

If \mathcal{S} is the set of majority agents, that means $|\mathcal{S}| \geq \lceil \frac{n}{2} \rceil$. Let \mathcal{A}_p and \mathcal{A}_q be two layered contiguous pieces and \mathcal{S} be the majority set. If every agent $i \in \mathcal{S}$, weakly prefers \mathcal{A}_p to \mathcal{A}_q . Then we call that the majority weakly prefer \mathcal{A}_p to \mathcal{A}_q . we denote it by $\mathcal{A}_p \stackrel{m}{\succeq} \mathcal{A}_q$. A point $x \in [0, 1]$ is said to be a majority switching point over the m -layered cake \mathfrak{C} , if $LR(x) \stackrel{m}{\succeq} RL(x)$ and $RL(x) \stackrel{m}{\succeq} LR(x)$.

3 Exact multi-layered cake cutting

First, we will look into the problem of obtaining complete exact multi-allocation over two layered cake for two agents that satisfies feasibility constraints. Here we will use the same idea of Austin moving-knife procedure for two agents where two knives are used. The main mathematical tool used by Austin moving-knife procedure is the intermediate value theorem. Basically we achieve our goal using the intermediate value theorem through the continuous movement of two knives. Here we define two different types of pieces (similar as diagonal pieces) on the entire cake according to the position of these two knives. The feature of these two knives are almost same as long knife. When the position of one knife is at the one of the ends of the unit interval and the other's is at any interior of the unit interval, then both of the knives are equivalent to a long knife. Both of the knives work on the entire cake similar as long knife. Let us explain how we define two different pieces while we moves two knives over the entire two layered cake:

Let K_1 and K_2 be two knives and x and y be the positions of the knives K_1 and K_2 respectively at an instant. Let us assume these two knives divide the entire cake into two portions $TLR(x, y) = ([0, x] \cap \mathcal{L}_1) \cup ([x, y] \cap \mathcal{L}_2) \cup ([y, 1] \cap \mathcal{L}_1)$ and $TRL(x, y) = ([0, x] \cap \mathcal{L}_2) \cup ([x, y] \cap \mathcal{L}_1) \cup ([y, 1] \cap \mathcal{L}_2)$ where $x \leq y$. When $x = 0$ and $0 < y < 1$ then $TLR(0, y) = RL(y)$ and $TRL(0, y) = LR(y)$. When $0 < x < 1$ and $y = 1$ then $TLR(x, 1) = LR(x)$ and $TRL(x, 1) = RL(x)$.




Figure 4: Examples of partitions for two pairs $(x = 0, y = \frac{1}{2})$ and $(x = \frac{1}{2}, y = 1)$.

Theorem 1. *There exists an exact complete multi-allocation over any two layered cake for two agents that satisfies the feasibility constraints.*

Proof. The Proof is almost similar as Austin moving-knife procedure for $n = 2$ agents. In a way analogous to the Austin-moving knife procedure we move two knives over a single layered cake in such a way that the value of the piece between two knives is always half with respect to agent 1. The second agent will say to stop the movement of knives while he thinks value of the piece between knives is half. This situation must happen due to intermediate value theorem. The initial position of first knife is at left end point of the unit interval and that of second knife is at the point which divides the cake into two equal valued pieces with respect to agent 1. In the similar way of Austin moving-knife procedure we start to move the knives K_1 and K_2 from the positions at 0 and s_1 respectively such that the valuation of agent 1 on $TLR(0, s_1)$ is half. This implies s_2 is the switching point of agent 1. We continuously move these two knives in such a way that the valuation of agent 1 on $TLR(x, y)$ is always half where $x \leq y$. The equality $x = y$ happens only when both have the same switching point. While we continuously move these two knives always keeping the value half on the piece $TLR(x, y)$ with respect to agent 1, then agent 2 will say "stop" at some point when he thinks the value on the piece $TLR(x, y)$ is half. The situation will happen due to the intermediate value theorem.

	0	$x = x^*$	$y = y^*$	1
\mathcal{L}_1	TLR	TRL	TLR	
\mathcal{L}_2	TRL	TLR	TRL	

Figure 5: When the exactness happens at $x = x^*$ and $y = y^*$ where $TRL(x^*, y^*) = TRL$ and $TLR(x^*, y^*) = TLR$.

□

4 Envy-free multi-layered cake cutting

In the literature of cake-cutting, there is no protocol for contiguous allocation with $n \geq 3$ agents without mild constraints on agents' preferences (Su [1999]). In multi-layered cake-cutting, Hosseini, Igarashi, and Searns proved the existence envy-free multi-allocation over 2-layered cake with at most two preferences for 2 and 3 agents respectively. Igarashi and Meunier proved the existence of envy-free multi-allocation over m -layered cake for n agents using combinatorial topology where n is a prime power and $m \leq n$. Here we will see that there exists a complete multi-allocation over 2-layered cake of 3 agents satisfying non-overlapping and contiguous constraints and each agent has a share to each layer, where the number of preferences is at most two. We show the existence of this multi-allocation only using the valuation function of majority agents.

Theorem 2. *There exists an envy-free complete multi-allocation $\mathcal{P} = (\mathcal{P}_1, \mathcal{P}_2, \mathcal{P}_3)$ satisfying feasibility and contiguity conditions for $n = 3$ and $m = 2$, where two agents share the same valuation function, can always be found even without the knowledge of the unique valuation function of the third agent.*

Proof. In the present scenario, two agents share the same valuation function, and the third agent has a unique valuation function.

Let p and q be two agents such that they have same valuation function $V_p = V_q = V$. Here the number of layers of the multi-layered cake \mathcal{C} is $m = 2$. and $\mathcal{C} = (\mathcal{L}_l)_{l \in \mathcal{M}}$ where $\mathcal{M} = \{1, 2\}$. Since $V(\mathcal{C}) = V(\mathcal{L}_1) + V(\mathcal{L}_2) = 1$, so either $V(\mathcal{L}_1) \leq \frac{1}{2}$ and $V(\mathcal{L}_2) \geq \frac{1}{2}$ or $V(\mathcal{L}_1) \geq \frac{1}{2}$ and $V(\mathcal{L}_2) \leq \frac{1}{2}$. So we get a switching point x for agent p (or q). By definition of switching point, $V(LR(x)) = V(RL(x)) = \frac{1}{2}$. Again $V(LR(x)) = V(\mathcal{L}_1 \cap [0, x]) + V(\mathcal{L}_2 \cap [x, 1]) = \frac{1}{2}$. Here we find out a complete multi-allocation $\mathcal{P} = (\mathcal{P}_1, \mathcal{P}_2, \mathcal{P}_3)$ such that $V(\mathcal{P}_i) = \frac{1}{3}$ for all $i \in \mathcal{N} = \{1, 2, 3\}$ and $\mathcal{P}_i = (P_{i1}, P_{i2})$. Where each P_{i1} and P_{i2} are contiguous pieces of layers 1 and 2 respectively and $P_{i1}, P_{i2} \neq \emptyset$ for each $i \in \mathcal{N}$. In this protocol, we use two short knives and one long knife to get a complete multi-allocation that satisfies both of two conditions, contiguity and feasibility, and allocates a non-zero piece to every agent from both layers.

First we give two short knives to the agent p (or q) to cut the 1st layer of the cake \mathcal{C} into a pair of pieces $\mathcal{L}_1 \cap [0, y]$ and $\mathcal{L}_1 \cap [y, 1]$ and 2nd layer of the cake \mathcal{L}_2 into a pair of pieces $\mathcal{L}_2 \cap [0, z]$ and $\mathcal{L}_2 \cap [z, 1]$ such that $V(\mathcal{L}_1 \cap [0, y] \cup \mathcal{L}_2 \cap [z, 1]) = V(\mathcal{L}_1 \cap [0, y]) + v(\mathcal{L}_2 \cap [z, 1]) = \frac{1}{3}$ where $0 < y < x < z < 1$. This type is possible because of switching point x . So $V(\mathcal{L}_1 \cap [y, 1] \cup \mathcal{L}_2 \cap [0, z]) = V(\mathcal{L}_1 \cap [y, 1]) + V(\mathcal{L}_2 \cap [0, z]) = \frac{2}{3}$.i.e. either $V(\mathcal{L}_1 \cap [y, 1]) \geq \frac{1}{3}$ and $V(\mathcal{L}_2 \cap [0, z]) \leq \frac{1}{3}$ or, $V(\mathcal{L}_1 \cap [y, 1]) \leq \frac{1}{3}$ and $V(\mathcal{L}_2 \cap [0, z]) \geq \frac{1}{3}$.

Consider a new cake $\mathcal{C}' = (\mathcal{L}'_1, \mathcal{L}'_2)$ where $\mathcal{L}'_1 = \mathcal{L}_1 \setminus [y, 1]$ and $\mathcal{L}'_2 = \mathcal{L}_2 \setminus [0, z]$. Now we have $V(\mathcal{C}') = \frac{2}{3}$ i.e. $V(\mathcal{L}'_1) + V(\mathcal{L}'_2) = \frac{2}{3}$. This ensures the existence of switching point c over the new cake \mathcal{C}' for the agent p (or q). So we get $V(LR(c, \mathcal{C}')) = V(RL(c, \mathcal{C}')) = \frac{1}{3}$. Again $LR(c, \mathcal{C}') = \mathcal{L}'_1 \cap [0, c] \cup \mathcal{L}'_2 \cap [c, 1]$ and $RL(c, \mathcal{C}') = \mathcal{L}'_1 \cap [c, 1] \cup \mathcal{L}'_2 \cap [0, c]$. Let $\mathcal{P}_1 = (\mathcal{L}_1 \setminus [0, y], \mathcal{L}_2 \setminus [z, 1])$, $\mathcal{P}_2 = (\mathcal{L}'_1 \cap [0, c], \mathcal{L}'_2 \cap [c, 1])$, and $\mathcal{P}_3 = (\mathcal{P}'_1 \cap [c, 1], \mathcal{L}'_2 \cap [0, c])$. Clearly, each P_{il} is a contiguous piece for each $i \in \mathcal{N} = \{1, 2, 3\}$ and each $l \in \mathcal{M} = \{1, 2\}$ and also P_{i1} and P_{i2} are non-overlapping for each $i \in \mathcal{N} = \{1, 2, 3\}$. Agent $r \neq p, q$ selects a layered piece \mathcal{P}_r of most valuation. Since each layered piece has same valuation for agents p and q . So there exists an envy-free complete multi-allocation $\mathcal{P} = (\mathcal{P}_1, \mathcal{P}_2, \mathcal{P}_3)$ that satisfied both conditions, feasibility and contiguity for 3 agents with at most two preferences.

	0	y	c	x	z	1
\mathcal{L}_1	P_{11}	P_{21}		P_{31}		
\mathcal{L}_2		P_{32}	P_{22}	P_{12}		

Figure 5: Envy-free multi-allocation \mathcal{P}

□

5 Proportional multi-layered cake cutting

Our aim is to find the existence of proportional multi-allocation over multi-layered cake when the number of layers is odd and the number of agents is greater than or equal to the number of agents. When the number of layers is a prime power and the number of agents is equal to the number of agents, then we always get a proportional multi-allocation that is feasible and contiguous Igarashi and Meunier [2021]. Using the idea of switching point property, we will show that there exists a proportional multi-allocation over 3-layered cake for $n \geq 3$ agents that satisfy both of the two constraints. We also show that there exists a proportional multi-allocation that is feasible and contiguous over $2^a P^b$ -layered cake for $n \geq 2^a P^b$ agents where $a \in \mathbb{Z}_+$, $b \in \mathbb{Z}_+ \cup \{0\}$ and P is a prime. The existence of a proportional multi-allocation over m -layered cake for $n \geq m$ agents implies the existence of a proportional multi-allocation over any $2^a m$ -layered cake for $2^a n$ agents $a \in \mathbb{Z}_+$. Here we also observe that a proportional multi-allocation always exists where $n > m$ and $m = 5, 7, 9, 11, 13, 15, 17, 19, 21$ and so.

The m -layered cut. Here we define m -layered cut when the number m of layers is odd. Let $m = 2k + 1$ where $k \in \mathbb{Z}_+$. Suppose $\mathfrak{C} = (\mathcal{L}_l)_{l \in \mathcal{M}}$ be an m -layered cake where m is odd and \mathcal{L}_l is the l -th layer of the cake. Define a new cake $\mathfrak{C}^* = (\mathcal{L}_l^*)_{l \in \mathcal{M}^*}$ where $\mathcal{L}_1^* = \mathcal{L}_1 \cup \mathcal{L}_2$, i.e, we consider both of the two layers \mathcal{L}_1 and \mathcal{L}_2 as the first layer \mathcal{L}_1^* of the new cake \mathfrak{C}^* , and $\mathcal{L}_l^* = \mathcal{L}_{l+1}$ for $l = 2, 3, \dots, 2k$ and $\mathcal{M}^* = \{1, 2, \dots, 2k\}$. Now we have a $2k$ -layered cake \mathfrak{C}^* , since the number of layers of the new cake \mathfrak{C}^* is even. Now we can use m -layered cut formula defined in the preliminaries. Now as per their definition, we get the diagonal pieces $LR(x)$ and $RL(x)$ of the new cake \mathfrak{C}^* . So for every $x \in [0, 1]$, we get

$$\begin{aligned} LR(x, \mathfrak{C}^*) &= \left(\bigcup_{l=1}^k \mathcal{L}_l^* \cap [0, x] \right) \cup \left(\bigcup_{l=k+1}^{2k} \mathcal{L}_l^* \cap [x, 1] \right). \\ RL(x, \mathfrak{C}^*) &= \left(\bigcup_{l=1}^k \mathcal{L}_l^* \cap [x, 1] \right) \cup \left(\bigcup_{l=k+1}^{2k} \mathcal{L}_l^* \cap [0, x] \right). \end{aligned} \quad (1)$$

After putting $\mathcal{L}_1^* = \mathcal{L}_1 \cup \mathcal{L}_2$, $\mathcal{L}_l^* = \mathcal{L}_{l+1}$ for $j = 2, 3, \dots, 2k$ to (1), we get

$$\begin{aligned} LR(x, \mathfrak{C}^*) &= \left(\bigcup_{l=1}^{k+1} \mathcal{L}_l \cap [0, x] \right) \cup \left(\bigcup_{l=k+2}^{2k+1} \mathcal{L}_l \cap [x, 1] \right). \\ RL(x, \mathfrak{C}^*) &= \left(\bigcup_{l=1}^{k+1} \mathcal{L}_l \cap [x, 1] \right) \cup \left(\bigcup_{l=k+2}^{2k+1} \mathcal{L}_l \cap [0, x] \right). \end{aligned} \quad (2)$$

Define,

$$\begin{aligned} LR(x, \mathfrak{C}) &= LR(x, \mathfrak{C}^*) = \left(\bigcup_{l=1}^{k+1} \mathcal{L}_l \cap [0, x] \right) \cup \left(\bigcup_{l=k+2}^{2k+1} \mathcal{L}_l \cap [x, 1] \right). \\ RL(x, \mathfrak{C}) &= RL(x, \mathfrak{C}^*) = \left(\bigcup_{l=1}^{k+1} \mathcal{L}_l \cap [x, 1] \right) \cup \left(\bigcup_{l=k+2}^{2k+1} \mathcal{L}_l \cap [0, x] \right). \end{aligned} \quad (3)$$

- If the number of layers is odd, i.e, $m = 2k + 1$ Then the merge of the diagonal piece $LR(x, \mathfrak{C})$ (or $RL(x, \mathfrak{C})$) is defined as $(\mathcal{L}'_l)_{l \in \mathcal{M}'}$ (or $(\mathcal{L}''_l)_{l \in \mathcal{M}'}$) where $\mathcal{M}' = \{1, 2, \dots, k + 1\}$, $\mathcal{L}'_1 = \mathcal{L}_1 \cap [0, x]$ (or $\mathcal{L}'_1 = \mathcal{L}_1 \cap [x, 1]$) and $\mathcal{L}'_l = \mathcal{L}_l \cap [0, x] \cup \mathcal{L}_{k+l} \cap [x, 1]$ (or $\mathcal{L}'_l = \mathcal{L}_l \cap [x, 1] \cup \mathcal{L}_{k+l} \cap [0, x]$) for $l \in \mathcal{M}' \setminus \{1\}$.

Lemma 3. Let \mathfrak{C} be an m -layered cake, where $m = 2k + 1, k \in \mathbb{Z}_+$, and $LR(x, \mathfrak{C})$ and $RL(x, \mathfrak{C})$ are the diagonal pieces. Suppose \mathfrak{C}' (or, \mathfrak{C}'') is a $(k + 1)$ -layered cake obtained from the merge of $LR(x, \mathfrak{C})$ (or, $RL(x, \mathfrak{C})$). If a layered piece \mathcal{P}_l of \mathfrak{C}' (or \mathfrak{C}'') is contiguous and non-overlapping then it is also contiguous and non-overlapping layered piece of \mathfrak{C} .

Proof. Let $\mathcal{P}_i = (P_1, P_2, \dots, P_{k+1})$ be a contiguous and non-overlapping layered piece of \mathfrak{C}' . This implies each P_l is a contiguous piece of the layer \mathcal{L}'_l for $l \in \mathcal{M}'$. Since $\mathcal{L}'_1 = \mathcal{L}_1 \cap [0, x]$ and $\mathcal{L}'_l = \mathcal{L}_l \cap [0, x] \cup \mathcal{L}_{k+l} \cap [x, 1]$

for $l \in \mathcal{M} \setminus \{1\}$. So we get $P_1 = (\mathcal{L}_1 \setminus [0, x]) \cap P_1$ and $P_l = (\mathcal{L}_l \cap [0, x] \cup \mathcal{L}_{k+l} \cap [x, 1]) \cap P_l = (\mathcal{C}_j \cap [0, x]) \cap P_l \cup (\mathcal{C}_{k+l} \cap [x, 1]) \cap P_l$ for $l \in \{2, 3, \dots, k+1\}$. Let $P'_1 = P_1$, $P'_l = (\mathcal{L}_l \setminus [0, x]) \cap P_l$ for $l \in \{2, 3, \dots, k+1\}$ and $P'_{k+l} = (\mathcal{L}_l \setminus [x, 1]) \cap P_l$ for $l \in \{2, k, \dots, k+1\}$. Since P_1, P_2, \dots , and P_{k+1} are non-overlapping and contiguous pieces so $P'_1, P'_2, \dots, P'_{2k}$ and P'_{2k+1} are also contiguous and non-overlapping pieces. So $\mathcal{P}'_l = (P'_1, P'_2, \dots, P'_{2k+1})$ is also a contiguous and non-overlapping layered piece of the original cake \mathcal{C} . \square

The existence of a proportional division of multi-layered cake has been proved by Igarashi and Meunier using combinatorial topology over m -layered cake for n agents where $m \leq n$ and n is prime power (Igarashi and Meunier [2021]). Now we will prove the existence of a proportional multi-allocation over 3-layered cake for 3 agents using the switching point property. Following this process we also get a proportional division of 4-layered cake for 4 agents without using the majority switching point criterion but the process quite large.

Theorem 4. *There exists a proportional complete multi-allocation over 3-layered cake for 3 agents that satisfies feasibility and contiguity constraints.*

Proof. Let $\mathcal{C} = (\mathcal{L}_l)_{l \in \mathcal{M}}$ be an 3-layered cake and V_1, V_2 and V_3 are the valuation functions of agents 1, 2 and 3 respectively. So for each $i \in \mathcal{N}$ and each $l \in \mathcal{M}$ either $V_i(\mathcal{L}_l) \geq \frac{1}{3}$ or $V_i(\mathcal{L}_l) \leq \frac{1}{3}$. For every agent $i \in \mathcal{N}$, we get two pair of layers $(\mathcal{L}_p, \mathcal{L}_q)$ and $(\mathcal{L}_q, \mathcal{L}_r)$ such that in each pair one layer has valuation at most $\frac{1}{3}$ and another layer has valuation at least $\frac{1}{3}$ where $p, q, r \in \mathcal{M}$. So by the pigeon hole theory, we get a pair $(\mathcal{L}_r, \mathcal{L}_s)$ of layers where the above valuation condition is satisfied by at least two valuation functions where $r, s \in \mathcal{M}$. Without loss of generality we assume that the valuation function V_1 satisfies the valuation condition for the pair $(\mathcal{L}_1, \mathcal{L}_2)$ of layers. So we get a point y such that $V_1(LR(\mathcal{C}_1 \cup \mathcal{C}_2, y)) = \frac{1}{3}$. Consider a new cake $\mathcal{C}' = (\mathcal{L}'_1, \mathcal{L}'_2)$ where $\mathcal{L}'_1 = \mathcal{L}_2 \cap [0, y] \cup \mathcal{L}_1 \cap [y, 1]$ and $\mathcal{L}'_2 = \mathcal{L}_3$. So the agent 1 has the valuation $\frac{2}{3}$ over the new cake \mathcal{C}' . Again by the switching point criterion, we get a point z such that $V_1(LR(\mathcal{C}', z)) = V_1(RL(\mathcal{C}', z)) = \frac{1}{3}$ where either $0 \leq y \leq z \leq 1$ or, $0 \leq z \leq y \leq 1$. Now we get a multi-allocation $\mathcal{P} = (\mathcal{P}_1, \mathcal{P}_2, \mathcal{P}_3)$ where $V_1(\mathcal{P}_1) = V_1(\mathcal{P}_2) = V_3(\mathcal{P}_3) = \frac{1}{3}$. According to the position of y and z , layered pieces \mathcal{P}_i are different. Using this multi-allocation \mathcal{P} , we find the required multi-allocation \mathcal{P}^* .

Case 1: When $z \leq y$, then $\mathcal{P}_1 = (\mathcal{L}_1 \cap [0, y], \mathcal{L}_2 \cap [y, 1], \emptyset)$, $\mathcal{P}_2 = (\emptyset, \mathcal{L}_2 \cap [0, z], \mathcal{L}_3 \cap [z, 1])$ and $\mathcal{P}_3 = (\mathcal{L}_1 \cap [y, 1], \mathcal{L}_2 \cap [z, y], \mathcal{L}_3 \cap [0, z])$. If we get two different layered pieces \mathcal{P}_p and \mathcal{P}_q such that $V_2(\mathcal{P}_p) \geq \frac{1}{3}$ and $V_3(\mathcal{P}_q) \geq \frac{1}{3}$ where $p \neq q \in \{1, 2, 3\}$. Then we are done. If not. So we get only one layered piece \mathcal{P}_r that has valuation greater than $\frac{1}{3}$ for two agents 2 and 3 where $r \in \{1, 2, 3\}$.




Figure 6: Multi-allocation \mathcal{P} when $z \leq y$.

Subcase i: When $r = 1$, then allocate the layered piece \mathcal{P}_2 to the agent 1. Define a new cake $\mathcal{C}^* = (\mathcal{L}_1^*, \mathcal{L}_2^*)$ where $\mathcal{L}_1^* = \mathcal{L}_1$ and $\mathcal{L}_2^* = \mathcal{L}_3 \cap [0, z] \cup \mathcal{L}_2 \cap [z, 1]$. Now the agents 2 and 3 have valuations at least $\frac{2}{3}$ over the new cake. Now after applying the cut-and-choose protocol for the agents 2 and 3 over the new cake \mathcal{C}^* , we get the required proportional multi-allocation that satisfies the conditions, feasibility and contiguity. By the switching point criterion, there exists a point x such that $V_2(LR(x, \mathcal{C}^*)) = V_2(RL(x, \mathcal{C}^*)) = \frac{V_2(\mathcal{C}^*)}{2} > \frac{1}{3}$. Agent 3 chooses a weakly preferred layered piece among the diagonal pieces. According to the positions of x and z , we get two types multi-allocation. Without loss of generality we consider that agent 2 prefers $LR(x, \mathcal{C}^*)$ and agent 3 prefers $RL(x, \mathcal{C}^*)$.

When $x \leq z$, then we get required multi-allocation $\mathcal{P}^* = (\mathcal{P}_1^*, \mathcal{P}_2^*, \mathcal{P}_3^*)$ where $\mathcal{P}_1^* = (\emptyset, \mathcal{L}_2 \cap [0, z], \mathcal{L}_3 \cap [z, 1])$, $\mathcal{P}_2^* = (\mathcal{L}_1 \cap [0, x], \mathcal{L}_2 \cap [z, 1], \mathcal{L}_3 \cap [x, z])$ and $\mathcal{P}_3^* = (\mathcal{L}_1 \cap [x, 1], \emptyset, \mathcal{L}_3 \cap [0, x])$.

	0	x	z	1
\mathcal{L}_1	P_{21}^*	P_{31}^*		
\mathcal{L}_2	P_{12}^*		P_{22}^*	
\mathcal{L}_3	P_{33}^*	P_{23}^*	P_{13}^*	
	$\mathfrak{C} = (\mathcal{L}_1, \mathcal{L}_2, \mathcal{L}_3)$			

Figure 7: Proportional multi-allocation \mathcal{P}^* when $x \leq z$.

When $z \leq x$, then $\mathcal{P}_2^* = (\mathcal{L}_1 \cap [0, x], \mathcal{L}_2 \cap [x, 1], \emptyset)$, $\mathcal{P}_3^* = (\mathcal{L}_1 \cap [x, 1], \mathcal{L}_2 \cap [z, x], \mathcal{L}_3 \cap [0, z])$ and \mathcal{P}_1^* remains same.

	0	z	x	1
\mathcal{L}_1	P_{21}^*		P_{31}^*	
\mathcal{L}_2	P_{12}^*	P_{32}^*	P_{22}^*	
\mathcal{L}_3	P_{33}^*		P_{13}^*	
	$\mathfrak{C} = (\mathcal{L}_1, \mathcal{L}_2, \mathcal{L}_3)$			

Figure 8: Proportional multi-allocation \mathcal{P}^* when $z \leq x$.

Subcase ii: When $r = 2$ or 3 , then allocate the layered piece \mathcal{P}_1 to the agent 1. Define a new cake $\mathfrak{C}^* = (\mathcal{L}_1^*, \mathcal{L}_2^*)$ where $\mathcal{L}_1^* = \mathcal{L}_2 \cap [0, y] \cup \mathcal{L}_1 \cap [y, 1]$ and $\mathcal{L}_2^* = \mathcal{L}_3$. Now the agents 2 and 3 have valuations over the new cake \mathfrak{C}^* at least $\frac{2}{3}$. Similar as subcase i (Case i), we get two types multi-allocation. When $x \leq y$, then we get a multi-allocation $\mathcal{P}^* = (\mathcal{P}_1^*, \mathcal{P}_2^*, \mathcal{P}_3^*)$ where $\mathcal{P}_1^* = \mathcal{P}_1$, $\mathcal{P}_2^* = (\emptyset, \mathcal{L}_2 \cap [0, x], \mathcal{L}_3 \cap [x, 1])$ and $\mathcal{P}_3^* = (\mathcal{L}_1 \cap [y, 1], \mathcal{L}_2 \cap [x, y], \mathcal{L}_3 \cap [0, x])$.

	0	x	y	1
\mathcal{L}_1	P_{11}^*		P_{31}^*	
\mathcal{L}_2	P_{22}^*	P_{32}^*		P_{12}^*
\mathcal{L}_3	P_{33}^*	P_{23}^*		
	$\mathfrak{C} = (\mathcal{L}_1, \mathcal{L}_2, \mathcal{L}_3)$			

Figure 9: Proportional multi-allocation \mathcal{P}^* when $x \leq y$.

When $y \leq x$, then we get a multi-allocation $\mathcal{P}^* = (\mathcal{P}_1^*, \mathcal{P}_2^*, \mathcal{P}_3^*)$ where $\mathcal{P}_1^* = \mathcal{P}_1$, $\mathcal{P}_2^* = (\mathcal{L}_1 \cap [y, x], \mathcal{L}_2 \cap [0, y], \mathcal{L}_3 \cap [x, 1])$ and $\mathcal{P}_3^* = (\mathcal{L}_1 \cap [x, 1], \emptyset, \mathcal{L}_3 \cap [0, x])$.

	0	y	x	1
\mathcal{L}_1	P_{11}^*		P_{21}^*	P_{31}^*
\mathcal{L}_2	P_{22}^*		P_{12}^*	
\mathcal{L}_3	P_{33}^*			P_{23}^*

$\mathfrak{C} = (\mathcal{L}_1, \mathcal{L}_2, \mathcal{L}_3)$

Figure 10: Proportional multi-allocation \mathcal{P}^* when $y \leq x$.

Case 2: When $y \leq z$, then $\mathcal{P}_1 = (\mathcal{L}_1 \cap [0, y], \mathcal{L}_2 \cap [y, 1], \emptyset)$, $\mathcal{P}_2 = (\mathcal{L}_1 \cap [y, z], \mathcal{L}_2 \cap [0, y], \mathcal{L}_3 \cap [z, 1])$ and $\mathcal{P}_3 = (\mathcal{L}_1 \cap [z, 1], \emptyset, \mathcal{L}_3 \cap [0, z])$. Now if we get two layered pieces \mathcal{P}_p and \mathcal{P}_q such that $V_2(\mathcal{P}_p) \geq \frac{1}{3}$ and $V_3(\mathcal{P}_q) \geq \frac{1}{3}$ where $p \neq q \in \{1, 2, 3\}$. Then we are done. Otherwise we get only one layered piece \mathcal{P}_r that has valuation at least $\frac{1}{3}$ for two agents 2 and 3 where $r \in \{1, 2, 3\}$.

	0	y	z	1
\mathcal{L}_1	P_{11}		P_{21}	P_{31}
\mathcal{L}_2	P_{22}		P_{12}	
\mathcal{L}_3	P_{33}			P_{23}

$\mathfrak{C} = (\mathcal{L}_1, \mathcal{L}_2, \mathcal{L}_3)$

Figure 11: Multi-allocation \mathcal{P} when $y \leq z$.

Subcase i: When $r = 1$, then allocate the layered piece \mathcal{P}_3 to agent 1. Now define a new cake $\mathfrak{C}^* = (\mathcal{L}_1^*, \mathcal{L}_2^*)$ where $\mathcal{L}_1^* = \mathcal{L}_1 \cap [0, z] \cup \mathcal{L}_3 \cap [z, 1]$ and $\mathcal{L}_2^* = \mathcal{L}_2$. Another two agents have valuations at least $\frac{2}{3}$ over the new cake \mathfrak{C}^* . Now if we apply the cut-and-choose protocol over the new cake for two agents 2 and 3, then we get the required multi-allocation for three agents. Let x be the switching point over the new cake for the agent 2. So we have $V_2(LR(\mathfrak{C}^*, x)) = V_2(RL(\mathfrak{C}^*, x)) = \frac{V_2(\mathfrak{C}^*)}{2} > \frac{1}{3}$. Without loss of generality we consider that agent 2 prefers $LR(\mathfrak{C}^*, x)$ and agent 3 prefers another diagonal piece. According to the positions of x and z , we get two types multi-allocation.

When $x \leq z$, the required multi-allocation is $\mathcal{P}^* = (\mathcal{P}_1^*, \mathcal{P}_2^*, \mathcal{P}_3^*)$ where $\mathcal{P}_1^* = (\mathcal{L}_1 \cap [z, 1], \emptyset, \mathcal{L}_3 \cap [0, z])$, $\mathcal{P}_2^* = (\mathcal{L}_1 \cap [0, x], \mathcal{L}_2 \cap [x, 1], \emptyset)$ and $\mathcal{P}_3^* = (\mathcal{L}_1 \cap [x, z], \mathcal{L}_2 \cap [0, x], \mathcal{L}_3 \cap [z, 1])$.

	0	x	z	1
\mathcal{L}_1	P_{21}^*		P_{31}^*	P_{11}^*
\mathcal{L}_2	P_{32}^*		P_{22}^*	
\mathcal{L}_3	P_{13}^*			P_{33}^*

$\mathfrak{C} = (\mathcal{L}_1, \mathcal{L}_2, \mathcal{L}_3)$

Figure 12: Proportional multi-allocation \mathcal{P}^* when $x \leq z$.

When $z \leq x$, the layered pieces of required multi-allocation are $\mathcal{P}_1^* = (\mathcal{L}_1 \cap [z, 1], \emptyset, \mathcal{L}_3 \cap [0, z])$, $\mathcal{P}_2^* = (\mathcal{L}_1 \cap [0, z], \mathcal{L}_2 \cap [x, 1], \mathcal{L}_3 \cap [z, x])$ and $\mathcal{P}_3^* = (\emptyset, \mathcal{L}_2 \cap [0, x], \mathcal{L}_3 \cap [x, 1])$.




Figure 13: Proportional multi-allocation \mathcal{P}^* when $z \leq x$.

Subcase ii: When $r = 2$ or 3 , we allocate the layered piece \mathcal{P}_1 to agent 1.. Define a new cake $\mathfrak{C}^* = (\mathcal{L}_1^*, \mathcal{L}_2^*)$ where $\mathcal{L}_1^* = (\mathcal{L}_1 \cap [0, y] \cup \mathcal{L}_2 \cap [y, 1])$ and $\mathcal{L}_2^* = \mathcal{L}_3$. So the agents 2 and 3 have valuations over the new cake \mathfrak{C}^* at least $\frac{2}{3}$. Similar as subcase i (case ii), we get two types multi-allocation.

When $x \leq y$, then the multi-allocation is $\mathcal{P}^* = (\mathcal{P}_1^*, \mathcal{P}_2^*, \mathcal{P}_3^*)$ where $\mathcal{P}_1^* = (\mathcal{L}_1 \cap [0, y], \mathcal{L}_2 \cap [y, 1], \emptyset)$, $\mathcal{P}_2^* = (\emptyset, \mathcal{L}_2 \cap [0, x], \mathcal{L}_3 \cap [x, 1])$ and $\mathcal{P}_3^* = (\mathcal{L}_1 \cap [y, 1], \mathcal{L}_2 \cap [x, y], \mathcal{L}_3 \cap [0, x])$.




Figure 14: Proportional multi-allocation \mathcal{P}^* when $x \leq y$.

When $y \leq x$, the layered pieces of required multi-layered cake are $\mathcal{P}_1^* = (\mathcal{L}_1 \cap [0, y], \mathcal{L}_2 \cap [y, 1], \emptyset)$, $\mathcal{P}_2^* = (\mathcal{L}_1 \cap [y, x], \mathcal{L}_2 \cap [0, y], \mathcal{L}_3 \cap [x, 1])$ and $\mathcal{P}_3^* = (\mathcal{L}_1 \cap [x, 1], \emptyset, \mathcal{L}_3 \cap [0, x])$.




Figure 15: Proportional multi-allocation \mathcal{P}^* when $y \leq x$.

So for any case, we get a proportional multi-allocation \mathcal{P}^* that satisfies the both of two conditions, contiguity and feasibility. \square

Now we will show the existence of proportional multi-allocation when $n \geq m$ and $m = 3$ that is not proved using switching point property. This problem was solved by Igarashi and Meunie using combinatorial topology. Using the following way we can observe that we always get a proportional multi-allocation for pairs (n, m) where $n > m$ and $m = 5, 15, 17, 19, 21, 23, 29, 31$, and so on. Basically we use the concept of majority switching point and the definition of merge of diagonal pieces when the number of layers is odd. Then consider the merge of the diagonal pieces as a new cake where the first layer of the new cake is a subinterval. After getting the new cake we will check that there exists a proportional multi-allocation over the new cake for the majority. Until now we don't know the existence of proportional multi-allocation over $P_1 P_2$ -layered cake for n agents where $n \geq P_1 P_2$ and P_1 and P_2 are two distinct prime numbers where both are not equal to 2. Suppose $m = 15$ then using the following way we can show the existence of proportional multi-allocation for $n > 15$ agents. Also we can see for the pairs (n, m) where $n > m$ and $m = 21, 33, 35$ and so on.

Theorem 5. *There exists a proportional complete multi-allocation over a 3-layered cake \mathcal{C} with n agents satisfies feasibility and contiguity conditions where $n \geq 3$.*

Proof. Here we prove the existence of the required multi-allocation for two cases: n is even or odd where $n \geq 4$. We have shown the proof for $n = 3$.

Case 1: When n is even say $n = 2k$, $n \in \mathbb{Z}_+ \setminus \{1\}$. We cannot apply majority switching point property directly, because the number of layers is 3(odd). We apply this property after considering the first and second layers of the cake, namely \mathcal{L}_1 and \mathcal{L}_2 , as a single layer of the cake. Let us consider a new cake $\mathcal{C}' = (\mathcal{L}'_1, \mathcal{L}'_2)$ where $\mathcal{L}'_1 = \mathcal{L}_1 \cup \mathcal{L}_2, \mathcal{L}'_2 = \mathcal{L}_3$ and $\mathcal{C} = (\mathcal{L}_1, \mathcal{L}_2, \mathcal{L}_3)$. Since the number of layers is even and also the number of agents is even. Then we get a majority switching point $x \in [0, 1]$. Now we can partition \mathcal{N} into two sets, \mathcal{N}_1 and \mathcal{N}_2 such that each agent $i \in \mathcal{N}_1$ weakly prefers $LR(x)$ to $RL(x)$ and each agent $i \in \mathcal{N}_2$ weakly prefers $RL(x)$ to $LR(x)$ where $|\mathcal{N}_1| = |\mathcal{N}_2| = \frac{|\mathcal{N}|}{2}$. Here we have $LR(x, \mathcal{C}') = (\mathcal{L}_1 \cup \mathcal{L}_2) \cap [0, x] \cup \mathcal{L}_3 \cap [x, 1]$ and $RL(x, \mathcal{C}') = \mathcal{L}_3 \cap [0, x] \cup (\mathcal{L}_1 \cup \mathcal{L}_2) \cap [x, 1]$. Now we consider two other 2-layered cakes \mathcal{C}^* and \mathcal{C}^{**} where $\mathcal{C}^* = (\mathcal{L}_1 \cap [0, x], \mathcal{L}_2 \cap [0, x] \cup \mathcal{L}_3 \cap [x, 1])$ and $\mathcal{C}^{**} = (\mathcal{L}_1 \cap [x, 1], \mathcal{L}_3 \cap [0, x] \cup \mathcal{L}_2 \cap [x, 1])$. So we get two 2-layered cakes, \mathcal{C}^* and \mathcal{C}^{**} and for each 2-layered cake, the number of agents is $k = \frac{n}{2}$ where $k = \frac{n}{2} \geq 2$. Now we get a proportional multi-allocation \mathcal{P}^i for the set \mathcal{N}_i of agents that is feasible and contiguous follows from *theorem 5* by Hosseini et al. [2020] where $i \in \{1, 2\}$. Together with multi-allocations \mathcal{P}^1 and \mathcal{P}^2 , we get a proportional multi-allocation \mathcal{P} over the given cake for n agents that is feasible and contiguous. Pictures are shown in figure 15.




Figure 16

Case 2: When n is odd say $n = 2k + 1$, $k \in \mathbb{Z}_+ \setminus \{1\}$. If we reduce this case to case 1, we are done. Ask every agent a_i to mark the point y_i over the top layered cake \mathcal{C}_1 such that the valuation over contiguous piece $\mathcal{L}_1 \cap [0, y_i]$ is at least as highly valued as proportional value $\frac{1}{n}$. We cut a piece $\mathcal{C}_1 \cap [0, y]$ and allocate it to an agent say p

where $y = y_p = \min\{y_1, y_2, y_3, \dots, y_n\}$. Without loss of generality we assume that $p = 1$. Consider a new cake $\mathcal{C}^* = (\mathcal{L}_1^*, \mathcal{L}_2^*, \mathcal{L}_3^*)$ together with $2k$ where $\mathcal{L}_1^* = \mathcal{L}_1 \cap [y, 1]$, $\mathcal{L}_2^* = \mathcal{L}_2$ and $\mathcal{L}_3^* = \mathcal{L}_3$. Now we have a cake \mathcal{C}^* together with $2k$ agents and each agent a_i has a valuation at least $\frac{n-1}{n} V_i(\mathcal{C}) = \frac{2k}{2k+1} V_i(\mathcal{C})$ over the cake \mathcal{C}^* . We get a proportional multi-allocation \mathcal{P} with $2k$ agents over \mathcal{C}^* that is contiguous and feasible follows from case 1. Together with the allocated piece $\mathcal{L}_1 \cap [0, y]$ and the multi-allocation \mathcal{P} , we get a proportional multi-allocation \mathcal{P}^* over \mathcal{C} for n agents that is feasible and contiguous. Pictures are shown in figure 16.




Figure 17

□

We can generalize the idea of above theorem and the generalized theorem is:

Theorem 6. *If \mathcal{C} is a $(2k + 1)$ -layered cake and $n > m$. Then there exists a proportional multi-allocation over \mathcal{C} for n agents if there exists a proportional multi-allocation over $(k + 1)$ -layered cake for the set of at least $(k + 1)$ agents.*

Now we will prove the existence of a proportional multi-allocation for $n = m$ where m is of the form $2^a 3^b$ for some $a \in \mathbb{Z}_+$ and $b \in \{0, 1\}$. Since the existence of majority switching point permits us to divide into two subproblems. We will repeat this procedure until the number of layers of each subproblem becomes either 2 or 3.

Theorem 7. *There exists a proportional multi-allocation that satisfies feasibility and contiguity constraints over any m -layered cake for $n = m$ agents where m is of the form $2^a 3^b$ for $a \in \mathbb{Z}_+$ and $b \in \{0, 1\}$.*

The proof of the above theorem is followed from the proof of the theorem 4 and the proof of the theorem 4 of the paper by [Hosseini et al., 2020].

Theorem 8. *There exists a proportional multi-allocation that satisfies feasibility and contiguity constraints over any m -layered cake for $n \geq m$ agents where m is of the form $2^a 3^b$ for $a \in \mathbb{Z}_+$ and $b \in \{0, 1\}$.*

The proof of the above theorem is followed from the proof of the theorem 5 of the paper by Hosseini et al. [2020].

First Hosseini, Igarashi, and Searns proved the existence of proportional multi-allocation over 2^a agents for n agents where $n \geq 2^a$ and $a \in \mathbb{Z}_+$. Igarashi and Meunier showed the existence of proportional multi-allocation over P^a -layered cake for P^a agents where $a \in \mathbb{Z}_+$ and P is a prime. Using above two result we will show the existence of proportional multi-allocation over $2^a P^b$ -layered cake for n agents $n \geq 2^a P^b$ and $a, b \in \mathbb{Z}_+$ and P is a prime.

Theorem 9. *If the number of agents and the number of layers are equal and is of the form $2^a P^b$ where $a, b \in \mathbb{Z}_+ \cup \{0\}$ and P is prime. Then we always get a proportional multi-allocation that is contiguous and feasible.*

Proof. W.l.o.g. we consider that each layer piece \mathcal{L}_l is a copy of the unit interval. Here we will show the existence of the required multi-allocation without defining any recursive algorithm. We also prove this using recursive algorithm to *alternative proof*.

Case i: When $m = n = 1$. Allocate the cake \mathcal{C} to the single agent.

Case ii: When $b = 0$. Then cut-and-choose algorithm ensures the existence of proportional multi-allocation.

Case iii: When $a = 0$. Then the proposition 4 of the paper by Igarashi and Meunier [2021] ensures the existence of the required multi-allocation.

Case iv: If we can prove the existence of the required multi-allocation whenever a and b are positive .i.e, $a, b \neq 0$. We are done. Let $m = n = 2^a P^b$ where $a, b \in \mathbb{Z}_+$ and P is prime. Since the number of layer is even, we can use the majority switching point criterion on the cake and also on the merge of its diagonal pieces while its number of layer is even. Continuing this process until we get 2^a cakes that are obtained from merging of diagonal pieces and each cake has $m^{(a)} = P^b$ layers together with $n^{(a)} = P^b$ agents . So we get a proportional multi-allocation over P^b -layered cake for $n^{(a)} = P^b$ agents that satisfies two conditions, contiguity and feasibility followed by *case iii*. Combining all the multi-allocations we get a proportional multi-allocation over the original cake that is contiguous and feasible followed from the corollary 1 of the paper by Hadi et al(2020).

Alternative proof. Assume that each layer of the original cake is a copy of the unit interval. Now design a recursive algorithm $\mathcal{A}l$ is mentioned in the following:

Input: A cake \mathcal{C}' having $|\mathcal{M}'|$ layered together a subset \mathcal{N}' of agents with $|\mathcal{N}'| \geq 2$ and a valuation scheme $(V_i)_{i \in \mathcal{N}'}$.

Output: A proportional feasible multi-allocation of the given cake to the agents.

Here $m = n$. If $m = n = 1$, then allocate the cake to the single agent. If $m = n = 2$, then the algorithm is same as the cut-and-choose algorithm. If $m = n = P^b$ where $b \neq 0$, then we run the procedure as described in the proposition 2 in the paper by Igarashi and Meunier [2021] where P is prime. If $m = n = 2^a$ where $a \neq 0$, then we run the procedure in the *theorem 4* in the paper of Hadi et al (2020). If $a, b \in \mathbb{Z}_+$, the designed algorithm searches a majority switching point $x \in [0, 1]$ over the cake \mathcal{C}' . Let $\mathcal{I}_1 = LR(x, \mathcal{C}')$ and $\mathcal{I}_2 = RL(x, \mathcal{C}')$. Since m is even. So by the definition of a majority switching point we can partition the set \mathcal{N}' of agents into two subsets \mathcal{N}_1 and \mathcal{N}_2 such that $|\mathcal{N}_1| = |\mathcal{N}_2| = \frac{|\mathcal{N}'|}{2}$ where each agent of the subset \mathcal{N}_1 of agents weakly prefers \mathcal{I}_1 to \mathcal{I}_2 and also each agent of the subset \mathcal{N}_2 of agents weakly prefers \mathcal{I}_2 to \mathcal{I}_1 . Then apply $\mathcal{A}l$ to the new cake \mathcal{C}^i with the set \mathcal{N}_i of agents for $i = 1, 2$, respectively where \mathcal{C}^i is the merge of the diagonal piece \mathcal{I}_i for $i = 1, 2$. Now we prove that by induction on the exponential of a , that a proportional complete multi-allocation \mathcal{P} is returned as an output that satisfies the two conditions, contiguity and feasibility. If $a = 1, b \geq 1$ then $m = n = 2P^b$ (even). So we get a majority switching point x with two subsets \mathcal{N}_1 and \mathcal{N}_2 of agents with $|\mathcal{N}_1| = |\mathcal{N}_2| = P^b$ such that each agent of the set \mathcal{N}_i of agents weakly prefers \mathcal{I}_i to \mathcal{I}_j where $\mathcal{I}_1 = LR(x), \mathcal{I}_2 = RL(x), i, j \in \{1, 2\}$ and $i \neq j$. Now applying the algorithm $\mathcal{A}l$ to the cake \mathcal{C}^i is obtained from the merge of \mathcal{I}_i with the set \mathcal{N}_i of agents, we get a multi-allocation that is contiguous and feasible followed by the *proposition 2* in the paper by Igarashi and Meunier [2021] where $i \in \{1, 2\}$. Since every non-overlapping and contiguous layered piece of the diagonal pieces is also non-overlapping and contiguous layered piece of the original cake. Thus the algorithm gives a proportional complete multi-allocation that is contiguous and feasible as an output. Assume that the algorithm $\mathcal{A}l$ gives the required multi-allocation for the values of a upto k .i.e, $1 \leq a \leq k$. Let $a = k + 1, b \geq 1$, then $m = n = 2^{k+1}P^b$ (even). So we get a majority switching point x with two subsets \mathcal{N}_1 and \mathcal{N}_2 of agents with $|\mathcal{N}_1| = |\mathcal{N}_2| = 2^k P^b$ such that each agent of the set \mathcal{N}_i of agents weakly prefers \mathcal{I}_i to \mathcal{I}_j where $\mathcal{I}_1 = LR(x), \mathcal{I}_2 = RL(x), i, j \in \{1, 2\}$ and $i \neq j$. Now applying the algorithm $\mathcal{A}l$ to the cake \mathcal{C}^i is obtained from the merge of \mathcal{I}_i with the set \mathcal{N}_i of agents, we get a multi-allocation that is contiguous and feasible followed by induction hypothesis where $i \in \{1, 2\}$. Since every non-overlapping and contiguous layered piece of the diagonal pieces is also non-overlapping and contiguous layered piece of the original cake. Thus the algorithm gives a proportional complete multi-allocation that is contiguous and feasible as an output. □

Theorem 10. *There exists a proportional multi-allocation over m -layered cake for any number $n \geq m$ of agents that is feasible and contiguous where $m = 2^a P^b$; $a, b \in \mathbb{Z}_+ \cup \{0\}$ and P is prime.*

Proof. We design a recursive algorithm $\mathcal{A}l$ is mentioned in the following:

Input: A cake \mathcal{C}' having $|\mathcal{M}'|$ -layered together a subset \mathcal{N}' of agents with $|\mathcal{N}'| \geq 2$ and a valuation scheme $(V_i)_{i \in \mathcal{N}'}$.

Output: A proportional feasible multi-allocation of the given cake to the agents. When $n = m$, then we think that the algorithm is similar to the algorithm described in the above *theorem 6*. When $a = 0, b \neq 0$, then we apply the algorithm in the proof of theorem 5 by Hosseini et al. [2020]. When $b = 0, a \neq 0$, then we apply the algorithm in the proof of the proposition 2 by Igarashi and Meunier [2021]. If $n > m$ and for some integer $a \geq 1$ and $b \geq 1$, then the algorithm finds a layer \mathcal{L}_l such that some agents have valuation at least $\frac{1}{n}$ over \mathcal{L}_l . Without loss of generality we assume that $l = 1$. Ask every agent a_i to mark the point y_i over the top layered cake \mathcal{C}_1 such that the valuation over contiguous piece $\mathcal{L}_1 \cap [0, y_i]$ is at least as highly valued as proportional value $\frac{1}{n}$. The algorithm $\mathcal{A}l$ allocate the piece $Y = \mathcal{C}_1 \cap [0, y]$ to

an agent say p where $y = y_p = \min\{y_1, y_2, y_3, \dots, y_n\}$. Now apply the algorithm \mathcal{Al} to a cake $\mathcal{C}' = (\mathcal{L}_l)_{l \in \mathcal{M}'}$ is obtained from the remaining piece of the cake \mathcal{C} with the subset $\mathcal{N}' \setminus \{p\}$ of agents where $\mathcal{L}'_1 = \mathcal{C}_1 \setminus Y$ and $\mathcal{L}'_l = \mathcal{L}_l$ for $l \neq 1$. Using induction on \mathcal{N}' we will show that the multi-allocation $\mathcal{P} = (\mathcal{P}_1, \mathcal{P}_2, \dots, \mathcal{P}_n)$ return by \mathcal{Al} as an output that is contiguous and feasible and satisfies proportionality. When $|\mathcal{N}'| = m$, this is true by the above theorem. Assume that the claim holds for $m \leq |\mathcal{N}'| \leq k - 1$. Now we will show that the claim also holds for $|\mathcal{N}'| = k$. Since $k > m$. So by predefined condition we consider that an agent k gets the contiguous piece Y has a value at least proportional share. Now the valuation of each agent a_i of the set $\mathcal{N}' \setminus \{k\}$ of agents over the remaining cake is at least $\frac{|\mathcal{N}'|-1}{|\mathcal{N}'|} V_i(\mathcal{C}')$. By the induction hypothesis, each agent a_i gets his allocated piece \mathcal{P}_i that has valuation at least $\frac{1}{|\mathcal{N}'|} V_i(\mathcal{C}')$ for $i \neq k$. So the multi-allocation \mathcal{P} is proportional as well as feasible and contiguous by induction hypothesis. \square

Follows from above theorems we generalize the following theorem.

Theorem 11. *If corresponding to a m' -layered cake with n' agents, there exists a proportional complete multi-allocation that is feasible and contiguous. Then we always get a proportional multi-allocation over $2^a m'^b$ -layered cake with $2^a n'^b$ agents that is feasible and contiguous where $n' \geq m'$, $b \in \{0, 1\}$ and $a \in \mathbf{Z}_+ \cup \{0\}$.*

Proof. Without loss of generality we assume that each layer \mathcal{L}_j is a copy of the unit interval $[0, 1]$. We design a recursive algorithm \mathcal{Al} is mentioned in the following:

Input: A cake \mathcal{C}' having $|\mathcal{M}'|$ layered together a subset \mathcal{N}' of agents with $|\mathcal{N}'| \geq 2$ and a valuation scheme $(V_i)_{i \in \mathcal{N}'}$
Output: A proportional contiguous multi-allocation of the given cake to the agents.

We consider the case when $a \neq 0$, then the algorithm \mathcal{Al} searches a majority switching point $x \in [0, 1]$. Let $\mathcal{I}_1 = LR(x, \mathcal{C}')$ and $\mathcal{I}_2 = RL(x, \mathcal{C}')$. Since m is even. We get two subsets \mathcal{N}_1 and \mathcal{N}_2 of \mathcal{N}' of agents such that $|\mathcal{N}_1| = |\mathcal{N}_2| = \frac{|\mathcal{N}'|}{2}$ where each agent of the subset \mathcal{N}_i of agents weakly prefers \mathcal{I}_i to \mathcal{I}_j where $i, j \in \{1, 2\}$ and $i \neq j$. We will show that a proportional feasible multi-allocation \mathcal{P} is returned as output by the algorithm \mathcal{Al} using multi-dimensional induction on the exponential of a and b . If $a = b = 0$, then the algorithm \mathcal{Al} allocates the entire cake to the single agent. Since $m = n = 1$. If $a = 0, b = 1$ then $m = m'$ and $n = n'$. When $n = n'$ and $m = m'$, we run the procedure to get the required multi-allocation as the given condition that there exists a multi-allocation corresponding to a m' -layered cake with n' agents that is feasible and contiguous. If $a = 1, b = 0$, then $m = n = 2$. we run the cut-and-choose algorithm. If $a = 1, b = 1$, then $m = 2m'$ and $n = 2n'$. Since m and n both are even. So by the definition of a majority switching, we get a majority switching point $x \in [0, 1]$ and two subsets \mathcal{N}_1 and \mathcal{N}_2 of \mathcal{N}' of agents with $|\mathcal{N}_1| = |\mathcal{N}_2| = \frac{|\mathcal{N}'|}{2}$ such that each agent of the subset \mathcal{N}_i of agents weakly prefers \mathcal{I}_i to \mathcal{I}_j where $\mathcal{I}_1 = LR(x, \mathcal{C}')$, $\mathcal{I}_2 = RL(x, \mathcal{C}')$, $i, j \in \{1, 2\}$ and $i \neq j$. Now applying the algorithm \mathcal{Al} to the cake \mathcal{C}^i is obtained from the merge of \mathcal{P}_i with the set \mathcal{N}_1 of agents, we get a multi-allocation \mathcal{P}^i over \mathcal{C}^i to the set \mathcal{N}_i of agents that is feasible followed by the condition when $a = 0, b = 1$ where $i \in \{1, 2\}$. Since every non-overlapping and contiguous layered piece of the diagonal pieces is also non-overlapping and contiguous layered piece of the original cake. Thus the algorithm gives a proportional complete multi-allocation that is contiguous and feasible as an output. If $a \geq 1$ and $b \in \{0, 1\}$ then $n = 2^a n'^b$. If we can show the existence of the required multi-allocation by induction on the exponential a . Then we are done by multidimensional induction. Let we get a proportional complete multi-allocation that is feasible for the values of a where $1 \leq a \leq k$. Suppose that $a = k + 1$. Then we have an $2^{k+1} m'^b$ -layered cake \mathcal{C}' with $2^{k+1} n'^b$ agents. Since n and m is even. So by the definition of a majority switching, we get a majority switching point $x \in [0, 1]$ and two subsets \mathcal{N}_1 and \mathcal{N}_2 of \mathcal{N}' of agents with $|\mathcal{N}_1| = |\mathcal{N}_2| = \frac{|\mathcal{N}'|}{2} = 2^k n'^b$ such that each agent of the subset \mathcal{N}_i of agents weakly prefers \mathcal{I}_i to \mathcal{I}_j where $\mathcal{I}_1 = LR(x, \mathcal{C}')$, $\mathcal{I}_2 = RL(x, \mathcal{C}')$, $i, j \in \{1, 2\}$ and $i \neq j$. Now applying the algorithm \mathcal{Al} to $2^k m'^b$ cake \mathcal{C}^i is obtained from the merge of \mathcal{I}_i with the set \mathcal{N}_i of agents, we get a multi-allocation \mathcal{P}^i over \mathcal{C}^i to the set \mathcal{N}_i of agents that is feasible followed by the induction hypothesis where $i \in \{1, 2\}$. Since every non-overlapping and contiguous layered piece of the diagonal pieces is also non-overlapping and contiguous layered piece of the original cake. Thus the algorithm gives a proportional complete multi-allocation over the entire cake of agents that is contiguous and feasible as an output \square

6 Acknowledgements

The author would like to thank Bodhayan Roy for his suggestions which have improved the presentation of this paper significantly.

References

- Hugo Steinhaus. Sur la division pragmatique. *Econometrica*, 17:315–319, 1949.
- Steven J. Brams and Michael A. Jones. Better ways to cut a cake. *Notices of the American Mathematical Society*, 53: 1314–1321, 2006.
- Ariel D. Procaccia. Cake cutting: not just child’s play. *Commun. ACM*, 56:78–87, 2013.
- David Kurokawa, John K. Lai, and Ariel D. Procaccia. How to cut a cake before the party ends. *Proceedings of the AAAI Conference on Artificial Intelligence*, 2013.
- Amin Saberi, Xiaotie Deng, and Qi Qi. Algorithmic solutions for envy-free cake cutting. *Operations Research*, 60(6): 1461–76, 2012.
- John Cloutier, Kathryn L. Nyman, and Francis Edward Su. Two-player envy-free multi-cake division. *Math. Soc. Sci.*, 59:26–37, 2010.
- Nicolas Lebert and Frédéric Meunier and Quentin Carbonneaux. Envy-free two-player m-cake and three-player two-cake divisions. *Operations Research Letters*, 41(6):607–610, 2013.
- Erel Segal-Halevi. Fair multi-cake cutting. *Discret. Appl. Math.*, 291:15–35, 2021.
- Lester Eli Dubins and Edwin Henry Spanier. How to cut a cake fairly. *The American Mathematical Monthly*, 68(1): 1–17, 1961.
- Shimon Even and Azaria Paz. A note on cake cutting. *Discret. Appl. Math.*, 7:285–296, 1984.
- Jack Robertson and William Webb. *Cake-Cutting Algorithms Be Fair If You Can*. CRC Press, 1998.
- Attila Tasnádi. A new proportional procedure for the n-person cake-cutting problem. *Economics Bulletin*, 4:1–3, 2003.
- Haris Aziz and Simon Mackenzie. A bounded and envy-free cake cutting algorithm. *Communications of the ACM*, 63: 119 – 126, 2020.
- Haris Aziz and Simon Mackenzie. A discrete and bounded envy-free cake cutting protocol for any number of agents. In *Proceedings of the 57th Symposium on Foundations of Computer Science (FOCS)*, pages 416–427, 2016.
- Hadi Hosseini, Ayumi Igarashi, and Andrew searns. Fair division of time: Multi-layered cake cutting. In *IJCAI*, 2020.
- Marcus Berliant, William Thomson, and Karl Dunz. On the fair division of a heterogeneous commodity. *Journal of Mathematical Economics*, 21(3):201–216, 1992.
- William Thomson. Fair allocation rules. *Handbook of Social Choice and Welfare*, 2:393–506, 2011.
- Erel Segal-Halevi and Shmuel Nitzan. Fair cake-cutting among families. *Social Choice and Welfare*, 53:709–740, 2019.
- Vijay Krishna Menon and K. Larson. Deterministic, strategyproof, and fair cake cutting. In *IJCAI*, 2017.
- A. Keith Austin. Sharing a cake. *The Mathematical Gazette*, 66(437):212–215, 1982.
- Eric Balkanski, Simina Brânzei, David Kurokawa, and Ariel D. Procaccia. Simultaneous cake cutting. In *AAAI*, 2014.
- Erel Segal-Halevi, Shmuel Nitzan, Avinatan Hassidim, and Yonatan Aumann. Fair and square: Cake-cutting in two dimensions. *ArXiv*, abs/1409.4511, 2015.
- Ágnes Cseh and Tamás Fleiner. The complexity of cake cutting with unequal shares. *ACM Transactions on Algorithms (TALG)*, 16:1 – 21, 2018.
- Francis Edward Su. Rental harmony: Sperner’s lemma in fair division. *American Mathematical Monthly*, 106:930–942, 1999.
- Ayumi Igarashi and Frédéric Meunier. Envy-free division of multi-layered cakes. In *WINE*, 2021.