

Geometrical and analytical results for Einstein solitons

Enrique Fernando López Agila¹ and José Nazareno Vieira Gomes²

ABSTRACT. We compute a lower bound for the scalar curvature of a gradient Einstein soliton under a certain assumption on its potential function. We establish an asymptotic behavior of the potential function on a noncompact gradient shrinking Einstein soliton. As a result, we obtain the finiteness of its fundamental group and its canonical weighted volume and we prove the validity of the weak maximum principle at infinity for a drifted Laplacian. We also prove some geometrical and analytical results for constructing gradient Einstein solitons that are realized as warped metrics, and we give a few explicit examples.

1. Introduction

Our objective here is to prove some geometrical and analytical results for gradient ρ -Einstein solitons that have not yet been addressed in the literature. A Riemannian metric g on an n -dimensional smooth manifold M^n , with $n \geq 3$, is a gradient ρ -Einstein soliton if there exists a smooth function $\varphi : M^n \rightarrow \mathbb{R}$ such that the Ricci tensor of g is given by

$$Ric + \nabla^2 \varphi = (\lambda + \rho S)g, \quad (1.1)$$

for some real numbers $\rho \neq 0$ and λ . Here, $\nabla^2 \varphi$ is the Hessian of φ , and $S = tr(Ric)$ is the scalar curvature of g . The function φ is called the potential function of a gradient ρ -Einstein soliton (M^n, g, φ) . The soliton is trivial whenever $\nabla \varphi$ is parallel. The reason for considering $n \geq 3$ is that, for $n = 2$, Eq. (1.1) reduces to the gradient Yamabe solitons equation, which are special solutions to the Yamabe flow. In this case, we refer the reader to Daskalopoulos and Sesum [10].

It is known that a gradient ρ -Einstein soliton arises as a self-similar solution of the Ricci Bourguignon flow [5], namely:

$$\frac{\partial}{\partial t} g(t) = -2(Ric - \rho S g(t)),$$

with $g(0) = g$ on a smooth manifold M^n , which has a unique solution for a positive time interval, when $\rho < \frac{1}{2(n-1)}$ and M^n is compact, for any initial Riemannian metric g , see Catino et al. [7, Theorem 2.1]. A gradient ρ -Einstein soliton is steady

2010 *Mathematics Subject Classification.* Primary 53C21, 53C25; Secondary 53C15.
Key words and phrases. Einstein soliton; warped metric; volume growth.

for $\lambda = 0$, shrinking for $\lambda > 0$ and expanding for $\lambda < 0$. By a complete gradient ρ -Einstein soliton (M^n, g, φ) , we mean a geodesically complete Riemannian manifold (M^n, g) with a potential function φ such that $\nabla\varphi$ is complete.

It is also known that any n -dimensional compact gradient ρ -Einstein soliton, with $\rho \leq \frac{1}{2(n-1)}$, is either shrinking of positive scalar curvature or the soliton is trivial. Moreover, any compact gradient Schouten soliton (i.e., $\rho = \frac{1}{2(n-1)}$) or traceless Ricci soliton (i.e., $\rho = \frac{1}{n}$) is trivial, see Catino and Mazzieri [8, Theorem 3.1]. It is also trivial any complete gradient steady Schouten soliton, see [8, Theorem 1.5]. The examples we present in Section 4 include two nontrivial three-dimensional complete noncompact traceless Ricci solitons. The reason why they are of dimension three comes from an analytical argument, see Examples 1 and 2. The first one of them is steady and the second one is shrinking.

Catino and Mazzieri also proved that any n -dimensional complete gradient Schouten soliton which is either shrinking or steady has nonnegative scalar curvature as a special case of a more general result on complete ancient solutions for the Schouten flow, see [8, Proposition 5.1 and Corollary 5.2]. Very recently, Borges [4, Theorem 1.1] showed that is bounded the scalar curvature S of any geodesically complete noncompact gradient Schouten soliton which is either shrinking or expanding, more precisely, it satisfies $0 \leq \lambda S \leq 2(n-1)\lambda^2$.

Example 4 is a nontrivial incomplete noncompact gradient Schouten soliton with unbounded negative scalar curvature, which can be either steady, shrinking or expanding, and then it shows that the assumption of completeness is essential for the results by Catino and Mazzieri, and by Borges.

In our first theorem, we give our contribution for this subject by considering a special class of gradient ρ -Einstein solitons. More precisely, we prove:

Theorem 1. *Let (M^n, g, φ) be a geodesically complete gradient ρ -Einstein soliton with scalar curvature bounded from below and $0 < \rho < \frac{1}{2(n-1)}$. Suppose there exists a point $p \in M^n$ such that for all $r(x) \geq r_0$,*

$$\varphi(x) \geq -Ar(x)^2 + B, \quad (1.2)$$

for some constants $A \geq 0$ and B , where $r(x)$ is the distance function from p and r_0 is a sufficiently large positive integer. Then,

$$S(x) \geq \min \left\{ 0, \frac{\lambda n}{1 - \rho n} \right\}.$$

The assumption on the potential function in (1.2) occurs for any geodesically complete noncompact gradient shrinking Schouten soliton, see Borges [4, Theorem 1.2]. For the case of gradient Ricci solitons, see Cao and Zhou [6, Theorem 1.1] and also Munteanu and Sesum [14, Corollary 5.2]. In Remark 1 we show that (1.2) is equivalent to an asymptotic behavior of the potential function.

The proof of Theorem 1 follows from an appropriate bound for the volume growth of geodesic balls together with the weak maximum principle at infinity for a drifted Laplacian, see Lemmas 1 and 2.

Having proved Theorem 1, we work to establish an asymptotic behavior of the potential function on a geodesically complete noncompact gradient shrinking ρ -Einstein soliton. For this, we follow the approach by Cao and Zhou from which we know that the potential function φ of an arbitrary complete noncompact gradient shrinking (i.e., $\lambda > 0$) Ricci soliton (M^n, g) has an asymptotic behavior similar

to the potential function of a Gaussian shrinking soliton. Indeed, from the proof of Proposition 2.1 in [6], we can conclude that for every minimizing unit speed geodesic $\gamma : [0, \infty) \rightarrow M^n$ emanating from a point $p \in M^n$ the potential function satisfies

$$\lim_{t \rightarrow +\infty} \frac{1}{t} \frac{d\varphi}{dt}(t) = \lambda.$$

In our second theorem, we obtain a similar result for the potential function of a geodesically complete noncompact gradient shrinking ρ -Einstein soliton with nonnegative scalar curvature and $\rho > 0$. Notice that, in the shrinking case, the assumptions in Theorem 1 imply that the scalar curvature is nonnegative. Also, the Borges' result before mentioned guarantees that any geodesically complete noncompact gradient shrinking Schouten soliton has nonnegative scalar curvature.

Theorem 2. *Let (M^n, g, φ) be a geodesically complete noncompact gradient shrinking ρ -Einstein soliton with nonnegative scalar curvature and $\rho > 0$. Then:*

- (1) *For any fixed point $p \in M^n$, and every minimizing unit speed geodesic $\gamma : [0, \infty) \rightarrow M^n$ emanating from p , we obtain*

$$\limsup_{t \rightarrow +\infty} \frac{1}{t} \frac{d\varphi}{dt} \geq \lambda \tag{1.3}$$

uniformly, i.e., it does not depend on the initial direction of the geodesic.

- (2) *M^n has a finite fundamental group, and the canonical weighted volume $\text{vol}_\varphi(M)$ is finite. In particular, the weak maximum principle at infinity for the φ -Laplacian holds on M^n .*

As an application of Theorem 2, we prove a compactness criteria for a class of gradient shrinking ρ -Einstein solitons.

Corollary 1. *Let (M^n, g, φ) be a geodesically complete gradient shrinking ρ -Einstein soliton with nonnegative scalar curvature and $\rho > 0$. Suppose there exists a point $p \in M^n$ such that for all $r(x) \geq r_0$, $|\nabla\varphi(x)| \leq (\lambda - \varepsilon)r(x) + c$, for some positive constants c and ε , with $\varepsilon \leq \lambda$, where $r(x)$ is the distance function from p and r_0 is a sufficiently large positive integer. Then M^n is a compact manifold.*

The proof of Theorem 2 follows from a more general setting that we discuss in Section 5, which is motivated by the construction of gradient ρ -Einstein solitons that are realized as warped metrics $g = g_B + f^2 g_F$ on the product manifold $B^n \times F^m$, see Section 3. For simplicity, we say that $B^n \times_f F^m$ is a *gradient ρ -Einstein soliton warped product* with a potential function η . The notation $\tilde{\psi}$ stands for the lift of a smooth function ψ on B^n to $B^n \times F^m$. The main result of this subject is:

Theorem 3. *Let $B^n \times_f F^m$ be a geodesically complete gradient ρ -Einstein soliton warped product with a nonconstant warping function. Suppose one of the following conditions hold:*

- (1) *$m \neq 2$ or $\rho \neq -1/2$;*
(2) *f is bounded.*

Then, the scalar curvature of the fiber is constant. Moreover, the potential function $\eta = \tilde{\varphi}$ and $\lambda + \rho S_g = \tilde{\Lambda}$, for some smooth functions φ and Λ on B^n .

Theorem 3 is simpler under the assumption that the base is compact, see Proposition 1. We also prove the necessary and sufficient conditions to construct a gradient ρ -Einstein soliton warped product, see Propositions 2 and 3. We highlight that

Examples 1, 2 and 4 above mentioned are of gradient ρ -Einstein soliton warped products. We establish the sign of the scalar curvature of the fiber of a gradient ρ -Einstein soliton warped product which is either shrinking or steady, provided the warping function reaches a minimum, see Corollary 2. We also establish a compactness criteria for a geodesically complete gradient shrinking ρ -Einstein soliton warped product by means of an asymptotic behavior of the potential function, see Corollary 3.

2. Lower bound for the scalar curvature

The goal of this section is to compute a lower bound for the scalar curvature of a geodesically complete gradient ρ -Einstein soliton by means of the volume growth of geodesic balls. Let $(M^n, g, e^{-h}dvol)$ be a weighted Riemannian manifold, where $dvol$ denotes the volume form on Riemannian manifold (M^n, g) and $e^{-h}dvol$ its corresponding weighted volume form, for some $h \in C^\infty(M)$. We recall the more appropriate tensor in this setting, the so-called Bakry-Emery Ricci and denoted by

$$Ric_h = Ric + \nabla^2 h.$$

Let $B(p, R) \subset M^n$ be the metric ball of radius $R > 0$ and centered at $p \in M^n$ with boundary $\partial B(p, R)$. We set

$$vol_h(B(p, R)) = \int_{B(p, R)} e^{-h} dvol \quad \text{and} \quad vol_h(\partial B(p, R)) = \int_{\partial B(p, R)} e^{-h} dvol_{n-1},$$

where $dvol_{n-1}$ stands for the area form on $\partial B(p, R)$. We denote the drifted Laplacian by

$$\Delta_h u = e^h \operatorname{div} (e^{-h} \nabla u),$$

and we also refer to it as the h -Laplacian.

To estimate the scalar curvature an approach is to look for conditions for the validity of the weak maximum principle at infinity for some drifted Laplacian, since the validity of this principle is closely related to the volume growth of geodesic balls.

We first give certain conditions to obtain an appropriate bound for the volume growth of geodesic balls. For this, we combine two classical tools in Riemannian geometry. The first one of them is known as the Bochner technique, which we apply to the gradient of the distance function to get a Riccati inequality. The second one of them is the first variation of the area formula, which allows us to estimate the weighted volume form in polar coordinates.

Lemma 1. *Let (M^n, g, φ) be a geodesically complete gradient ρ -Einstein soliton with scalar curvature bounded from below and $0 < \rho < \frac{1}{2(n-1)}$. Suppose there exists a point $p \in M^n$ such that the potential function satisfies Ineq. (1.2). Then, for any $r_0 > 0$ there exist constants A_0 , B_0 and C_0 , which depend on r_0 , such that for all $R \geq r_0$,*

$$vol_{\hat{\varphi}}(B(p, R)) \leq A_0 + B_0 \int_{r_0}^R e^{C_0 t^2} dt,$$

where $\hat{\varphi} = \frac{1}{1-2\rho(n-1)}\varphi$.

PROOF. Let $p \in M^n$ be a point such that the potential function φ satisfies Ineq. (1.2). Let $B(p, R) \subset M^n$ be a geodesic ball of radius $R > 0$ and centered

at p . Consider the distance function $r(x)$ from p , and apply the classical Bochner formula for $r(x)$ to obtain

$$0 = |\nabla^2 r|^2 + Ric(\partial r, \partial r) + g(\nabla \Delta r, \partial r).$$

By Schwarz inequality

$$0 \geq \frac{(\Delta r)^2}{n-1} + Ric(\partial r, \partial r) + g(\nabla \Delta r, \partial r).$$

Define $\hat{\varphi} = \frac{1}{1-2\rho(n-1)}\varphi$, $z = \Delta r$ and $z_{\hat{\varphi}} = z - \langle \nabla \hat{\varphi}, \partial r \rangle$. Then $\dot{z} = g(\nabla \Delta r, \partial r)$ and $\dot{z}_{\hat{\varphi}} = \dot{z} - \nabla^2 \hat{\varphi}(\partial r, \partial r)$. Thus, the previous inequality becomes

$$\dot{z}_{\hat{\varphi}} \leq -\frac{z^2}{n-1} - Ric_{\hat{\varphi}}(\partial r, \partial r),$$

which is known as a Riccati inequality. Use the gradient ρ -Einstein soliton equation (1.1) to obtain

$$\dot{z}_{\hat{\varphi}} \leq -\frac{z^2}{n-1} - (\lambda + \rho S) - 2\rho(n-1)\nabla^2 \hat{\varphi}(\partial r, \partial r).$$

By assumption $S \geq a$, for some constant a , and since $\rho > 0$, one has

$$\dot{z}_{\hat{\varphi}} \leq -\lambda + \rho|a| - 2\rho(n-1)\nabla^2 \hat{\varphi}(\partial r, \partial r).$$

Integrating twice give us

$$\int_{r_0}^r z_{\hat{\varphi}}(t) dt \leq C_0 r^2 + B_1, \quad (2.1)$$

for some constants C_0 and B_1 , where we used the assumptions (1.2) and $\rho < \frac{1}{2(n-1)}$.

Now, consider the annular set $B(p, r_0, R) = B(p, R) \setminus B(p, r_0)$, and note that

$$vol_{\hat{\varphi}}(B(p, R)) = vol_{\hat{\varphi}}(B(p, r_0)) + vol_{\hat{\varphi}}(B(p, r_0, R)).$$

Next, define $A_0 = vol_{\hat{\varphi}}(B(p, r_0))$ so that it is enough to control $vol_{\hat{\varphi}}(B(p, r_0, R))$. For it, use exponential polar coordinate around p and write the volume element $dvol = \mathcal{A}(r, \theta) dr \wedge d\theta_{n-1}$, where $d\theta_{n-1}$ is the standard volume element on the unit sphere S^{n-1} . Define $\mathcal{A}_{\hat{\varphi}}(r, \theta) = e^{-\hat{\varphi}} \mathcal{A}(r, \theta)$, and observe that by the first variation of the area formula, one has

$$\frac{\mathcal{A}_{\hat{\varphi}}(r, \theta)}{\mathcal{A}_{\hat{\varphi}}(r_0, \theta)} = e^{\int_{r_0}^r z_{\hat{\varphi}}(t, \theta) dt}$$

for $r \geq r_0 > 0$, see, e.g., Zhu [18] or Wei and Wylie [17]. So, (2.1) implies that

$$\mathcal{A}_{\hat{\varphi}}(r, \theta) \leq \mathcal{A}_{\hat{\varphi}}(r_0, \theta) e^{C_0 r^2 + B_1}.$$

Whence

$$\begin{aligned} vol_{\hat{\varphi}}(B(p, r_0, R)) &= \int_{r_0}^R \int_{S^{n-1}} \mathcal{A}_{\hat{\varphi}}(r, \theta) d\theta dr \\ &\leq \int_{r_0}^R \int_{S^{n-1}} \mathcal{A}_{\hat{\varphi}}(r_0, \theta) e^{C_0 r^2 + B_1} d\theta dr \\ &= \mathcal{A}_{\hat{\varphi}}(r_0) \int_{r_0}^R e^{C_0 r^2 + B_1} dr, \end{aligned}$$

where

$$A_{\hat{\varphi}}(r_0) = \int_{S^{n-1}} \mathcal{A}_{\hat{\varphi}}(r_0, \theta) d\theta.$$

This is enough to conclude the proof of the lemma. \square

We now guarantee the validity of the weak maximum principle at infinite for a drifted Laplacian on a class of gradient ρ -Einstein solitons. Recall that the weak maximum principle at infinity for the drifted Laplacian Δ_h holds on M^n if given a C^2 function $u : M^n \rightarrow \mathbb{R}$ satisfying $\sup_M u = u^* < +\infty$, there exists a sequence $\{x_n\} \subset M^n$ along which

$$(i) \ u(x_n) \geq u^* - \frac{1}{n} \quad \text{and} \quad (ii) \ \Delta_h u(x_n) \leq \frac{1}{n}.$$

We know that the weak maximum principle at infinite is valid for the operator Δ_φ on any geodesically complete gradient Ricci soliton with potential function φ , see Pigola, Rimoldi and Setti [15, Corollary 10]. Here, as an application of Lemma 1, we obtain:

Lemma 2. *Let (M^n, g, φ) be a geodesically complete gradient ρ -Einstein soliton with scalar curvature bounded from below and $0 < \rho < \frac{1}{2(n-1)}$. Suppose there exists a point $p \in M^n$ such that the potential function satisfies Ineq. (1.2). Then, the weak maximum principle at infinity for the $\hat{\varphi}$ -Laplacian holds on M^n , where $\hat{\varphi} = \frac{1}{1-2\rho(n-1)}\varphi$.*

PROOF. Lemma 1 implies that

$$\frac{r}{\log \text{vol}_{\hat{\varphi}}(B_r)} \notin L^1(+\infty).$$

Indeed, it is enough to note that $\frac{r}{\log(A_0 + B_0 \int_{r_0}^r e^{C_0 t^2} dt)} = o\left(\frac{1}{r}\right)$ as $r \rightarrow +\infty$. So, by Theorem 9 in [15], the weak maximum principle at infinity is valid for the $\hat{\varphi}$ -Laplacian on M^n . \square

We conclude this section by proving our first main result.

2.1. Proof of Theorem 1.

PROOF. We know from [8] that for any gradient ρ -Einstein soliton is true that

$$\left(\rho(1-n) + \frac{1}{2}\right) \Delta S - \frac{1}{2} g(\nabla S, \nabla \varphi) = \lambda S + \rho S^2 - |\text{Ric}|^2.$$

As $|\text{Ric}|^2 \geq \frac{S^2}{n}$, we get

$$\Delta_{\hat{\varphi}} S \leq \xi_1 S + \xi_2 S^2, \tag{2.2}$$

where

$$\hat{\varphi} = \frac{1}{1-2\rho(n-1)}\varphi, \quad \xi_1 = \frac{2\lambda}{1-2\rho(n-1)} \quad \text{and} \quad \xi_2 = \frac{2(\rho - \frac{1}{n})}{1-2\rho(n-1)}.$$

Considering the function $u = -S$, we have

$$\Delta_{\hat{\varphi}} u \geq \xi_1 u - \xi_2 u^2.$$

The assumption about S implies that $u^* = \sup_M u < +\infty$, and then by Lemma 2 we can apply the weak maximum principle at infinity to obtain

$$0 \geq \xi_1 u^* - \xi_2 (u^*)^2,$$

whence

$$u \leq \max \left\{ 0, -\frac{\xi_1}{\xi_2} \right\} = \left\{ 0, \frac{\lambda n}{1 - \rho n} \right\}.$$

Therefore

$$S(x) \geq \min \left\{ 0, \frac{\lambda n}{1 - \rho n} \right\}.$$

This finishes the proof. \square

Remark 1. *The following assumption is equivalent to Ineq. (1.2). Suppose there exists a point $p \in M^n$ such that for every minimizing unit speed geodesic $\gamma : [0, +\infty) \rightarrow M^n$ emanating from p ,*

$$\liminf_{t \rightarrow +\infty} \frac{\varphi(\gamma(t))}{t^2} \geq -A, \quad (2.3)$$

for some constant $A \geq 0$. Indeed, if φ satisfies (1.2), then (2.3) is trivially satisfied. Conversely, if φ satisfies (2.3), then for $\varepsilon = 1$ there is a positive integer r_0 so that for all $t \geq r_0$,

$$\frac{\varphi(\gamma(t))}{t^2} \geq -A - 1.$$

Thus, for $r(x) \geq r_0$,

$$\varphi(x) \geq -(A + 1)r^2(x).$$

This implies (1.2).

3. Gradient Einstein soliton warped products

Given two Riemannian manifolds (B^n, g_B) and (F^m, g_F) as well as a positive smooth warping function f on B^n , let us consider on the product manifold $B^n \times F^m$ the warped metric

$$g = \pi_1^* g_B + (f \circ \pi_1)^2 \pi_2^* g_F,$$

where π_1 and π_2 are the natural projections on B^n and F^m , respectively. Under these conditions, the product manifold is called the *warped product* of B^n and F^m . The notation $\tilde{\varphi} = \varphi \circ \pi_1$ stands for the lift of a smooth function φ on B^n to $B^n \times F^m$. If f is a constant, then $(B^n \times F^m, g)$ is a *standard Riemannian product*.

In this section, we study gradient ρ -Einstein solitons that are realized as warped metrics g on $B^n \times F^m$. For simplicity, we say that $B^n \times_f F^m$ is a *gradient ρ -Einstein soliton warped product* with a potential function η , and g is a *gradient ρ -Einstein soliton warped metric*. We denote by Ric_g the Ricci tensor of the warped metric g , while Ric_B and Ric_F are the Ricci tensors of g_B and g_F , respectively. We shall use similar notation for the geometric objects of g , g_B and g_F .

We recall that a warped product is a geodesically complete manifold for all warping functions if and only if both base and fiber are geodesically complete manifolds, see Bishop and O'Neill [2, Lemma 7.2].

It is known that the potential function of a gradient Ricci soliton warped product with a nonconstant warping function depends only on the base, see Borges and Tenenblat [3, Corollary 2.2]. Moreover, the scalar curvature of the fiber is constant, see Feitosa, Freitas Filho and Gomes [11]. Notice that our Theorem 3 is a similar result for a gradient ρ -Einstein soliton warped product with a nonconstant warping function.

We begin with the case of compact base. In this case, the idea of the proof is analogous to Borges and Tenenblat's result or, alternatively, see Lemma 4.1 in the paper by the second author in joint work with Marrocos and Ribeiro [13].

Proposition 1. *Let $B^n \times_f F^m$ be a gradient ρ -Einstein soliton warped product with a nonconstant warping function and compact base. Then, the scalar curvature of the fiber is constant. Moreover, the potential function $\eta = \tilde{\varphi}$ and $\lambda + \rho S_g = \tilde{\Lambda}$ for some smooth functions φ and Λ on B^n .*

PROOF. In order to apply this proof in the more general case of geodesically complete gradient ρ -Einstein soliton warped products, we shall use the compactness assumption just in the last part of the proof.

Let g be a gradient ρ -Einstein soliton warped metric on $B^n \times F^m$, then

$$\text{Ric}_g + \nabla_g^2 \eta = (\rho S_g + \lambda)g, \quad (3.1)$$

for some constants $\rho \neq 0$ and λ . So, by using Lemmas 7.3 and 7.4 in Bishop and O'Neill [2], we obtain

$$0 = (\nabla_g^2 \eta)(X, V) = X(V(\eta)) - (\nabla_X V)(\eta) = X(V(\eta)) - \frac{X(f)}{f}V(\eta),$$

for all X, V horizontal and vertical vector fields, respectively. This nullity implies

$$X \left(V \left(\frac{\eta}{f} \right) \right) = X \left(\frac{V(\eta)}{f} \right) = \frac{1}{f} \left[X(V(\eta)) - \frac{X(f)}{f}V(\eta) \right] = 0.$$

Therefore, the function $V \left(\frac{\eta}{f} \right)$ depends only on the fiber F^m . Thus, without loss of generality, we can write $\eta = \varphi + fh$, for some smooth functions φ on B^n and h on F^m , which has been the main idea of Borges and Tenenblat.

Next we observe that, for horizontal vector fields, equation (3.1) becomes

$$\text{Ric}_B(X, Y) - \frac{m}{f} \nabla_B^2 f(X, Y) + \nabla_B^2 \varphi(X, Y) + h \nabla_B^2 f(X, Y) = (\lambda + \rho S_g)g_B(X, Y) \quad (3.2)$$

which follows from Lemma 7.4 in [2] or, alternatively, [16, Corollary 43]. This lemma also implies

$$S_g = S_B - \frac{2m}{f} \Delta_B f + \frac{S_F}{f^2} - m(m-1) \frac{|\nabla_B f|^2}{f^2}. \quad (3.3)$$

From equations (3.2) and (3.3), we get

$$V(h) \nabla_B^2 f(X, Y) = \rho V \left(\frac{S_F}{f^2} \right) g_B(X, Y), \quad (3.4)$$

We now take traces in (3.4) to obtain

$$V(h) \Delta_B f = \rho \frac{V(S_F)}{f^2} n.$$

Since B^n is compact, Stokes' theorem implies $V(S_F) = 0$, and then S_F is constant. Now, using that f is nonconstant, from (3.4) we obtain $V(h) = 0$, i.e., h is constant. We conclude that $\lambda + \rho S_g = \tilde{\Lambda}$ and $\eta = \tilde{\varphi}$ for some functions $\varphi, \Lambda \in C^\infty(B^n)$. \square

The proof of Proposition 1 allows a generalization to geodesically complete gradient ρ -Einstein soliton warped products $B^n \times_f F^m$ with nonconstant warping function f . For this, we recall that ∇f cannot be parallel, see Proposition 3.6

in [3]. In particular, if the scalar curvature of the fiber is constant, then we obtain immediately from (3.4) the same result of Proposition 1.

We are now ready to prove the main result of this section.

3.1. Proof of Theorem 3.

PROOF. Here, we shall follow the notations and the results obtained as in Proposition 1. The proof requires a more careful analysis of equation (3.4).

We start by noting that the scalar curvature of the fiber is nonconstant if and only if h is nonconstant, which follows immediately from (3.4) and the fact that ∇f cannot be parallel, see [3, Proposition 3.6].

Now we claim that if the scalar curvature is nonconstant, then the following properties hold:

- (i) $\nabla_B^2 f = \frac{a}{f^2} g_B$, for some constant $a \neq 0$;
- (ii) $\rho = -\frac{1}{2}$ and $m = 2$.

Indeed, take a horizontal vector field X such that $g_B(X, \nabla f) = 0$. From (3.4), we get

$$V(h)\nabla_B^2 f(\nabla_B f, X) = 0.$$

Since h is nonconstant, there exists a vertical vector field V such that $V(h) \neq 0$, and then $\nabla_B^2 f(\nabla_B f, X) = 0$. Therefore, there exists a smooth function ψ on B^n so that $\nabla_B^2 f(\nabla_B f) = \psi \nabla_B f$. From (3.4), one has

$$\left(V(h)\psi - \rho \frac{V(S_F)}{f^2} \right) |\nabla_B f|^2 = 0. \quad (3.5)$$

Consider the closed set $A = \{x \in B^n; |\nabla_B f|(x) = 0\}$. Note that, the set $\text{int}(A)$ is empty or, equivalently, A^C is a dense set in B^n . In fact, if there exists $x_0 \in \text{int}(A)$, then $\nabla_B^2 f(x_0) = 0$ that implies S_F is constant from (3.4), which is a contradiction. From (3.5) and by continuity we have that

$$V(h)\psi - \rho \frac{V(S_F)}{f^2} = 0$$

on $B^n \times F^m$. So,

$$V(h)X(\psi f^2) = 0.$$

As $V(h) \neq 0$ it follows that $\psi f^2 = a$, for some constant $a \neq 0$, and then

$$aV(h) = \rho V(S_F). \quad (3.6)$$

Again from (3.4)

$$V(h) \left(\nabla_B^2 f - \frac{a}{f^2} g_B \right) = 0,$$

we conclude the proof of item (i).

To prove item (ii), we first obtain from item (i), the identity

$$|\nabla f|^2 + \frac{2a}{f} = b, \quad (3.7)$$

for some constant b .

Since $\eta = \varphi + fh$, and by using the ρ -Einstein soliton equation on F^m (see [16, Corollary 43]), we obtain

$$\begin{aligned} Ric_F - \left(\frac{\Delta_B f}{f} + (m-1) \frac{|\nabla_B f|^2}{f^2} \right) f^2 g_F + f \nabla_F^2 h - f \left(\nabla_B f(\varphi) + h |\nabla_B f|^2 \right) g_F \\ = (\lambda + \rho S_g) f^2 g_F. \end{aligned} \quad (3.8)$$

Keep in mind that the scalar curvature S_g is given by (3.3), and recall the second contracted Bianchi identity $div_F Ric_F = \frac{\nabla_F S_F}{2}$. Thus, we use (3.8) to obtain

$$\frac{\nabla_F S_F}{2} + f div_F \nabla_F^2 h - f |\nabla_B f|^2 \nabla_F h = \rho \nabla_F S_F. \quad (3.9)$$

Substituting (3.7) into (3.9), we have

$$\left(\frac{1}{2} - \rho \right) \nabla_F S_F + f div_F \nabla_F^2 h - f \left(b - \frac{2a}{f} \right) \nabla_F h = 0.$$

On the other hand (3.6) implies $\nabla_F S_F = \frac{a}{\rho} \nabla_F h$, and then

$$a \left(\frac{1}{2\rho} + 1 \right) \nabla_F h + f \left(div_F \nabla_F^2 h - b \nabla_F h \right) = 0.$$

Since f is nonconstant and it depends only on B^n , we get

$$a \left(\frac{1}{2\rho} + 1 \right) \nabla_F h = 0.$$

Hence, $\rho = -1/2$. Let us now prove that $m = 2$. We take the trace of equation (3.8) and we use the value of ρ to get

$$\begin{aligned} S_F - \left(\frac{\Delta_B f}{f} + (m-1) \frac{|\nabla_B f|^2}{f^2} \right) f^2 m + f \Delta_F h - f \left(\nabla_B f(\varphi) + h |\nabla_B f|^2 \right) m \\ = \left(\lambda - \frac{S_g}{2} \right) f^2 m. \end{aligned}$$

Whence

$$\left(1 + \frac{m}{2} \right) \nabla_F S_F + f \nabla_F \Delta_F h - f |\nabla_B f|^2 m \nabla_F h = 0.$$

Substituting $\nabla_F S_F = -2a \nabla_F h$ and (3.7) into the previous equation, we obtain

$$a(-2+m) \nabla_F h + f \left(\nabla_F \Delta_F h - bm \nabla_F h \right) = 0.$$

Hence, $m = 2$, and we finish the proof of item (ii). Note that item (ii) proves item (1) of the theorem.

For item (2), it is enough to prove that f does not satisfy item (i) when it is bounded.

Without loss of generality, assume $a > 0$, and take a unit vector $v \in T_p B$ such that $df_p v > 0$, $p \in B^n$. Let γ be the unit speed geodesic on B^n such that $\gamma(0) = p$ and $\dot{\gamma}(0) = v$. Define $\psi(t) = f(\gamma(t))$. Then, from (i) one has

$$\ddot{\psi}(t) = \nabla_B^2 f(\dot{\gamma}(t), \dot{\gamma}(t)) = \frac{a}{\psi^2(t)} > 0.$$

Thus, $\dot{\psi}(t) > 0$ for all $t \geq 0$ and ψ is an increasing function on $[0, +\infty)$ and bounded, since f is bounded. Hence ψ has a horizontal asymptote. In particular,

$$0 = \lim_{t \rightarrow +\infty} \dot{\psi}(t) = \lim_{t \rightarrow +\infty} \left(\dot{\psi}(0) + \int_0^t \frac{a}{\psi^2(s)} ds \right) > \dot{\psi}(0) = df_p(v) > 0,$$

which is a contradiction. This completes the proof of the theorem. \square

3.2. Necessary and sufficient conditions to construct a gradient Einstein soliton warped product. We start with some known result on gradient Ricci almost soliton. Recall that a Riemannian manifold (M^n, g) is a gradient Ricci almost soliton if there exist two smooth functions η and τ on M^n satisfying

$$Ric + \nabla^2 \eta = \tau g,$$

From Theorem 3, it is clear that a gradient ρ -Einstein soliton warped product is a very special case of gradient Ricci almost soliton warped product.

Let us consider a gradient Ricci almost soliton warped product $B^n \times_f F^m$, with $\eta = \tilde{\varphi}$ and $\tau = \tilde{\Lambda}$, for some smooth functions φ and Λ on B^n . The second author in joint work with Feitosa, Freitas Filho and Pina [12] established the necessary and sufficient conditions for constructing such a soliton. More precisely, the next two equations

$$Ric_B + \nabla_B^2 \varphi - \frac{m}{f} \nabla_B^2 f = \Lambda g_B, \quad (3.10)$$

$$-2\Lambda d\varphi + d \left((2 - m - n)\Lambda + |\nabla_B \varphi|^2 - \Delta_B \varphi - \frac{m}{f} \nabla_B \varphi(f) \right) = 0, \quad (3.11)$$

must be satisfied on B^n . Moreover, $Ric_F = \mu g_F$, with constant μ given by

$$\mu = \Lambda f^2 + f \Delta_B f + (m - 1) |\nabla_B f|^2 - f \nabla_B \varphi(f). \quad (3.12)$$

They proved that equation (3.12) is a consequence of (3.10) and (3.11). Now, we look for a necessary condition for Λ to be a linear function of the scalar curvature S_g of $B^n \times_f F^m$, which satisfies equation (3.3) on B^n . Taking the trace of (3.10), we get

$$S_B = n\Lambda - \Delta_B \varphi + \frac{m}{f} \Delta_B f. \quad (3.13)$$

From equations (3.3), (3.12) and (3.13), we obtain

$$S_g = (n + 1)\Lambda - \left(\Delta_B \varphi + \frac{\nabla_B \varphi(f)}{f} \right) - (m - 1) \left(\frac{\Delta_B f}{f} + (m - 1) \frac{|\nabla_B f|^2}{f^2} \right). \quad (3.14)$$

Since we are looking for conditions for Λ to be a linear function of S_g up to a constant, the condition that arises naturally from (3.14) is

$$\alpha \nabla_B \Lambda = \nabla_B \left(\Delta_B \varphi + \frac{\nabla_B \varphi(f)}{f} + (m - 1) \left(\frac{\Delta_B f}{f} + (m - 1) \frac{|\nabla_B f|^2}{f^2} \right) \right), \quad (3.15)$$

for some constant $\alpha \neq n + 1$.

The reason why we take $\alpha \neq n + 1$ comes from the fact that when $\alpha = n + 1$, then the scalar curvature S_g is constant. Indeed, by (3.15) we get

$$(n + 1)\Lambda = \Delta_B \varphi + \frac{\nabla_B \varphi(f)}{f} + (m - 1) \left(\frac{\Delta_B f}{f} + (m - 1) \frac{|\nabla_B f|^2}{f^2} \right) + c,$$

for some constant c . Substituting this last equation into (3.14), we obtain $S_g = c$. In particular, this reduces the problem of construction of the gradient ρ -Einstein soliton warped products to the case of gradient Ricci solitons, and then Eq. (3.15) is unnecessary, see [11, Theorem 3].

First, we prove the necessary conditions for constructing a gradient ρ -Einstein soliton warped product.

Proposition 2. *Let $B^n \times_f F^m$ be a gradient ρ -Einstein soliton warped product with potential function $\tilde{\varphi}$ and soliton function $\tilde{\Lambda} = \lambda + \rho\tilde{S}_g$. Then equations (3.10), (3.11) and (3.15) are satisfied on (B^n, g_B) , and $Ric_F = \mu g_F$, where μ is the constant given by Eq. (3.12).*

PROOF. Equation (3.10) and the fact that $Ric_F = \mu g_F$ with constant μ given by (3.12), follow directly by Proposition 2 in [12], while (3.11) is equation (1.3) in [12] on B^n . Taking the covariant derivative of equation (3.14), we conclude that (3.15) is satisfied for $\alpha = n + 1 - \frac{1}{\rho}$. \square

Second, we prove the sufficient conditions for constructing a gradient ρ -Einstein soliton warped product.

Proposition 3. *Let (B^n, g_B) be a Riemannian manifold with three smooth functions $f > 0, \Lambda$, and φ satisfying equations (3.10), (3.11) and (3.15). Take the constant μ satisfying Eq. (3.12) and a Riemannian manifold (F^m, g_F) with Ricci tensor $Ric_F = \mu g_F$. Then $B^n \times_f F^m$ is a gradient ρ -Einstein soliton warped product with potential function $\tilde{\varphi}$, soliton function $\tilde{\Lambda}$ and $\rho = \frac{1}{n+1-\alpha}$.*

PROOF. Since f, φ and Λ satisfy (3.10) and (3.11) on B^n , and $Ric_F = \mu g_F$ with μ given by (3.12), Theorem 1 in [12] guarantees that $B^n \times_f F^m$ is a gradient Ricci almost soliton warped product with potential function $\tilde{\varphi}$ and soliton function $\tilde{\Lambda}$. In particular, Eq. (3.14) is valid, and then by (3.15) we get

$$S_g = (n + 1 - \alpha)\Lambda + c,$$

for some constant c . Now, it is enough to define $\rho = \frac{1}{n+1-\alpha}$ and $\lambda = \Lambda - \frac{1}{n+1-\alpha}S_g$ to obtain a gradient ρ -Einstein soliton warped product. \square

Next, we give an immediately application of Proposition 2.

Corollary 2. *Let $B^n \times_f F^m$ be a gradient ρ -Einstein soliton warped product which is either shrinking or steady, with potential function $\tilde{\varphi}$, nonconstant warping function and nonnegative scalar curvature. If f reaches a minimum and $\rho > 0$, then the fiber has positive scalar curvature. In particular, F^m is a compact manifold.*

PROOF. By Proposition 2, $Ric_F = \mu g_F$, where μ is the constant given by (3.12) on B^n , i.e.,

$$\frac{\mu}{f} - \Lambda f = \Delta_B f + \frac{m-1}{f} |\nabla_B f|^2 - \nabla_B \varphi(f).$$

By assumption $\rho > 0$, $S_g \geq 0$ and $\lambda > 0$ or $\lambda = 0$ that imply $\Lambda = \lambda + \rho S_g \geq 0$. If $\mu \leq 0$ we have

$$0 \geq \Delta_B f + \frac{m-1}{f} |\nabla_B f|^2 - \nabla_B \varphi(f).$$

If f reaches a minimum, then it must be a constant by the strong maximum principle, which is a contradiction. So, $\mu > 0$ and then F^m is compact, since it is an Einstein manifold. \square

4. Examples

In this section, we present some examples of gradient ρ -Einstein soliton warped metrics. For this purpose, we use Proposition 3 and we consider bases that are conformal to Euclidean spaces \mathbb{R}^n in the same way as in [12, Proposition 4].

Example 1. Let us consider the hyperbolic space \mathbb{H}^n , namely, the open half space $x_n > 0$ with the metric $\coth^2(x_n)\delta_{ij}$. The functions

$$\begin{aligned} f &= \coth(x_n), \\ \varphi &= \frac{2}{3}(-2 + m + n) \log(\cosh(x_n)) \quad \text{and} \\ \Lambda &= -\frac{1}{3 \cosh^4(x_n)}(2(-2 + m + n) + (1 + m + n) \cosh(2x_n)) \end{aligned}$$

satisfy equations (3.10) and (3.11) on \mathbb{H}^n . Besides, the constant μ given by Eq. (3.12) is null. Theorem 1 in [12] guarantees that, for any geodesically complete Ricci flat Riemannian manifold F^m , the warped metric g with warping function f , potential function $\tilde{\varphi}$ and soliton function $\tilde{\Lambda}$ is a geodesically complete gradient Ricci almost soliton on $\mathbb{H}^n \times F^m$. The proof is a straightforward computation. Moreover, from (3.3) its scalar curvature is

$$S_g = -\frac{(-1 + m + n)(-2 + m + n + 2 \cosh(2x_n))}{\cosh^4(x_n)}. \quad (4.1)$$

The special case of ρ -Einstein solitons occurs when $n = 1$ and $m = 2$, since Eq. (3.15) is satisfied for $\alpha = -1$ on \mathbb{H}^1 , so that $\mathbb{H}^1 \times_f F^2$ is a nontrivial complete gradient steady traceless Ricci soliton warped product. Indeed, the completeness of $\nabla\varphi$ is a straightforward computation. Using the divergent curve criterion we prove that $\mathbb{H}^1 \times_f F^2$ is geodesically complete. For this, it is enough to prove that the metric $\coth^2(t)dt^2$ is geodesically complete, since the fiber is chosen to be geodesically complete. Let $\gamma : [t_0, +\infty) \rightarrow \mathbb{H}^1$ be a divergent curve. First, we assume $\gamma(t) \rightarrow +\infty$ as $t \rightarrow +\infty$, and we suppose without loss of generality that $\gamma'(t) > 0$, $\forall t \geq t_0$. Then

$$\begin{aligned} \int_{t_0}^t \coth(t)|\gamma'(t)|dt &= \int_{t_0}^t \coth(t)\gamma'(t)dt \\ &= \gamma(t) \coth(t) - \gamma(t_0) \coth(t_0) + \int_{t_0}^t \frac{\gamma(t)}{\sinh^2(t)}dt \\ &\geq \gamma(t) \coth(t) - \gamma(t_0) \coth(t_0). \end{aligned}$$

Hence

$$\lim_{t \rightarrow +\infty} \int_{t_0}^t \coth(t)|\gamma'(t)|dt = +\infty.$$

The case $\gamma(t) \rightarrow 0$ as $t \rightarrow \infty$ is similar, and then we obtain the required completeness of $\mathbb{H}^1 \times_f F^2$.

Example 2. Consider the metric $\cosh^2(x_n)\delta_{ij}$ on \mathbb{R}^n . The functions

$$\begin{aligned} f &= \cosh(x_n), \\ \varphi &= \frac{1}{12}(-2 + m + n)(8 \log(\cosh(x_n)) + \cosh(2x_n)) \quad \text{and} \\ \Lambda &= -\frac{3 - (-2 + m + n) \sinh^4(x_n)}{3 \cosh^4(x_n)} \end{aligned}$$

satisfy equations (3.10) and (3.11) on \mathbb{R}^n . Besides, the constant μ given by Eq. (3.12) is null. Again, Theorem 1 in [12] guarantees that, for any geodesically complete Ricci flat Riemannian manifold F^m , the warped metric g with warping function f , potential function $\tilde{\varphi}$ and soliton function $\tilde{\Lambda}$ is a complete gradient Ricci almost soliton on $\mathbb{R}^n \times F^m$. The proof is a straightforward computation. Moreover, from (3.3)

its scalar curvature is

$$S_g = -\frac{(-1+m+n)(6-m-n+(-2+m+n)\cosh(2x_n))}{2\cosh^4(x_n)}.$$

Again, the special case of ρ -Einstein solitons occurs when $n = 1$ and $m = 2$, since Eq. (3.15) is satisfied for $\alpha = -1$ on $(\mathbb{R}, \cosh^2(t)dt^2)$, so that $\mathbb{R} \times_f F^2$ is a nontrivial complete gradient shrinking traceless Ricci soliton warped product with $\lambda = 1/3$. We can prove its completeness by an analogous argument as in Example 1.

Example 3. Let \mathbb{R}_+^n , $n \geq 3$, be a Euclidean half space with the canonical coordinates $x = (x_1, \dots, x_n)$ such that $x_n > 0$ and the metric $\frac{1}{x_n}\delta_{ij}$. The functions

$$f = \frac{1}{x_n}, \quad \varphi = \frac{2(2-m-n)}{3} \ln x_n \quad \text{and} \quad \Lambda = \frac{2(5-m-4n)x_n^2}{3}$$

satisfy equations (3.10), (3.11) and (3.15) on \mathbb{R}_+^n for

$$\alpha = -\frac{17+3m^2-18n+4n^2+2m(-8+5n)}{2(4n+m-5)}.$$

The constant μ given by Eq. (3.12) is $\mu = \frac{5-m-4n}{3}$. Proposition (3) guarantees that, for any Einstein manifold F^m , with $\text{Ric}_F = \mu g_F$, the warped metric g with warping function f , potential function $\tilde{\varphi}$ and soliton function $\tilde{\Lambda}$ is a gradient ρ -Einstein steady soliton warped metric on $\mathbb{R}_+^n \times F^m$, with $\rho = \frac{1}{n+1-\alpha}$. Moreover, from (3.3) its scalar curvature is

$$S_g = -\frac{1}{3} (7+3m^2-20n+12n^2+2m(-7+6n)) x_n^2.$$

In this case, the warped metric g is incomplete, since the curve $\gamma : [1, +\infty) \rightarrow (0, \dots, 0, t)$ on \mathbb{R}_+^n satisfies

$$\lim_{r \rightarrow +\infty} \int_1^r \frac{1}{t^2} dt = 1.$$

As we already observed in the introduction, the next example shows that the assumption of completeness is essential for the results by Catino and Mazzieri [8, Theorem 1.5] and by Borges [4, Theorem 1.1]

Example 4. Let \mathbb{R}^n be a Euclidean space with coordinates $x = (x_1, \dots, x_n)$ and metric $e^{2\xi}\delta_{ij}$, where $\xi = \sum_{i=1}^n \alpha_i x_i$, $\alpha_i \in \mathbb{R}$ with $\sum_i \alpha_i^2 = 1$ and $n \geq 3$. The functions

$$f = e^\xi, \quad \varphi = \frac{c}{2} e^{2\xi} - \frac{(2-m-n)}{2} \xi \quad \text{and} \quad \Lambda = c + \frac{(2-m-n)}{2} e^{-2\xi}$$

satisfy equations (3.10), (3.11) and (3.15) on \mathbb{R}^n for $\alpha = -2m - n + 3$ and any constant c . The constant μ given by Eq. (3.12) is null. Proposition (3) guarantees that, for any Ricci flat Riemannian manifold F^m , the warped metric g with warping function f , potential function $\tilde{\varphi}$ and soliton function $\tilde{\Lambda}$ is a gradient Schouten soliton warped metric on $\mathbb{R}^n \times F^m$ which can be either steady, shrinking or expanding, since $\lambda = c$. Moreover, from (3.3) its scalar curvature is

$$Sg = -(m+n-2)(m+n-1)e^{-2\xi}. \quad (4.2)$$

To show that the warped metric g is incomplete, we can use a similar argument as in Example 3.

5. Asymptotic behavior of the potential function

In this section, we prove Theorem 2. We start by proving an interesting result for a class of noncompact geodesically complete Riemannian manifold (B^n, g_B) . For this, we consider the following modification of the m -Bakry-Emery Ricci tensor, which stems from the construction of both gradient Ricci solitons (see [11, 13]) and gradient ρ -Einstein solitons (see Proposition 2) that are realized as warped metrics:

$$Ric_{h,\varphi}^m := Ric + \nabla^2 \varphi + \nabla^2 h - \frac{1}{m} dh \otimes dh = Ric + \nabla^2 \varphi - \frac{m}{f} \nabla^2 f,$$

where m is a positive integer number, φ and h are smooth functions on B^n and $f = e^{-h/m}$.

The next proposition is motivated by the results proved in Section 3.

Proposition 4. *Let (B^n, g_B) be a noncompact geodesically complete Riemannian manifold that admits two smooth functions h and φ such that*

$$Ric_{h,\varphi}^m \geq \lambda, \quad (5.1)$$

for some positive constant λ . Fix a point $p \in B^n$, and take a minimizing unit speed geodesic $\gamma : [0, \infty) \rightarrow B^n$ emanating from p . It is true that

$$\limsup_{t \rightarrow +\infty} \frac{1}{t} \frac{d\varphi}{dt} \geq \lambda,$$

uniformly, i.e., it does not depend on the initial direction of the geodesic.

PROOF. The main idea is based on the proof of Proposition 2.1 in [6], in our case has been necessary to control the extra term $\nabla^2 h - \frac{1}{m} dh \otimes dh$.

Take $r(x)$ the distance function from a point $p \in B^n$, and consider $\gamma : [0, r(x)] \rightarrow B^n$ a minimizing unit speed geodesic, with $\gamma(0) = p$. By the second variation of the distance function, we have

$$0 \leq \int_0^r \left(\dot{\phi}^2(t)(n-1) - \phi^2(t) Ric(\gamma', \gamma') \right) dt, \quad (5.2)$$

for every smooth piecewise function $\phi(t)$ on $[0, r]$ with $\phi(0) = \phi(r) = 0$. We choose $\phi(t)$ given by

$$\phi(t) = \begin{cases} t, & t \in [0, 1], \\ 1, & t \in [1, r-1], \\ r-t, & t \in [r-1, r]. \end{cases} \quad (5.3)$$

Using (5.1), (5.2), (5.3), and integration by parts, we get

$$\begin{aligned} 0 &\leq \int_0^r \left(\dot{\phi}^2(n-1) - \phi^2 Ric(\dot{\gamma}, \dot{\gamma}) \right) dt \\ &\leq \int_0^r \left(\dot{\phi}^2(n-1) + \phi^2 \left(-\lambda + \frac{d^2 h}{dt^2} - \frac{1}{m} \left(\frac{dh}{dt} \right)^2 + \frac{d^2 \varphi}{dt^2} \right) \right) dt \\ &= \int_0^r \left(\dot{\phi}^2(n-1) - 2\phi \dot{\phi} \frac{dh}{dt} + \phi^2 \left(-\lambda - \frac{1}{m} \left(\frac{dh}{dt} \right)^2 + \frac{d^2 \varphi}{dt^2} \right) \right) dt \\ &\leq \int_0^r \left(\dot{\phi}^2(n-1) + 2 \left| \phi \dot{\phi} \frac{dh}{dt} \right| + \phi^2 \left(-\lambda - \frac{1}{m} \left(\frac{dh}{dt} \right)^2 + \frac{d^2 \varphi}{dt^2} \right) \right) dt. \end{aligned} \quad (5.4)$$

Applying the elementary inequality $2\sqrt{ab} \leq a + b$, for $a = m\dot{\phi}^2$ and $b = \frac{1}{m}\phi^2 \left(\frac{dh}{dt}\right)^2$ we have

$$2 \int_0^r \left| \phi \dot{\phi} \frac{dh}{dt} \right| dt \leq \int_0^r \left(m\dot{\phi}^2 + \frac{1}{m}\phi^2 \left(\frac{dh}{dt} \right)^2 \right) dt.$$

Combining this latter equation and (5.4) we get

$$\begin{aligned} 0 &\leq \int_0^r \left(\dot{\phi}^2(m+n-1) + \phi^2 \left(-\lambda + \frac{d^2\varphi}{dt^2} \right) \right) dt \\ &= 2(m+n-1) - \lambda \left(r - \frac{1}{3} \right) + \int_0^r \phi^2 \frac{d^2\varphi}{dt^2} dt. \end{aligned} \quad (5.5)$$

Analyzing

$$\begin{aligned} \int_0^r \phi^2 \frac{d^2\varphi}{dt^2} dt &= \int_0^1 \left(-2\phi \dot{\phi} \frac{d\varphi}{dt} \right) dt + \int_{r-1}^r \left(-2\phi \dot{\phi} \frac{d\varphi}{dt} \right) dt \\ &\leq \sup_{x \in B(p,1)} |\nabla\varphi| + 2 \int_{r-1}^r (r-t) \frac{d\varphi}{dt} dt \\ &\leq \sup_{x \in B(p,1)} |\nabla\varphi| + \sup_{t \in [r-1, r]} \frac{d\varphi}{dt}. \end{aligned} \quad (5.6)$$

On the other hand,

$$\frac{1}{r} \sup_{t \in [r-1, r]} \frac{d\varphi}{dt} \leq \sup_{t \in [r-1, r]} \frac{1}{t} \frac{d\varphi}{dt} \leq \sup_{t \geq r-1} \frac{1}{t} \frac{d\varphi}{dt}.$$

Combining this later inequality, (5.5) and (5.6), we obtain

$$\lambda \leq \limsup_{t \rightarrow +\infty} \frac{1}{t} \frac{d\varphi}{dt}.$$

This completes the proof of the theorem. \square

Now we are in a position to prove an asymptotic behavior of the potential function on a noncompact gradient shrinking ρ -Einstein soliton of nonnegative scalar curvature and $\rho > 0$.

5.1. Proof of Theorem 2.

PROOF. By assumption $S \geq 0$ and $\rho > 0$, then

$$Ric + \nabla^2\varphi \geq \lambda g.$$

As $\lambda > 0$ by Proposition 4 we conclude (1.3), and applying Corollary 5.1 in [17], we obtain that M^n has a finite fundamental group and that $vol_\varphi(M)$ is finite. In particular,

$$\frac{r}{\log vol_\varphi(B_r)} \notin L^1(+\infty).$$

Hence, by Theorem 9 in [15] the weak maximum principle at infinity for the φ -Laplacian holds on M^n . \square

5.2. Proof of Corollary 1.

PROOF. The proof goes by contradiction. Suppose M^n is noncompact. Theorem 2 guarantees that for any minimizing unit speed geodesic $\gamma : [0, +\infty) \rightarrow M^n$ emanating from $p \in M^n$, we get

$$\limsup_{t \rightarrow +\infty} \frac{|\nabla\varphi(\gamma(t))|}{t} \geq \limsup_{t \rightarrow +\infty} \frac{1}{t} \frac{d\varphi}{dt} \geq \lambda,$$

On the other hand, by assumption

$$\limsup_{t \rightarrow +\infty} \frac{|\nabla\varphi(\gamma(t))|}{t} \leq \lambda - \varepsilon.$$

This leads us to a contradiction. \square

Corollary 3. *Let $B^n \times_f F^m$ be a geodesically complete gradient shrinking ρ -Einstein soliton warped product with nonnegative scalar curvature, potential function $\tilde{\varphi}$ and $\rho > 0$. Suppose there exists a point $p \in B^n$ such that for all $r(x) \geq r_0$, $|\nabla\varphi(x)| \leq (\lambda - \varepsilon)r(x) + c$, for some positive constants c and ε , with $\varepsilon \leq \lambda$, where $r(x)$ is the distance function from p and r_0 is a sufficiently large positive integer. Then $B^n \times_f F^m$ is a compact manifold.*

PROOF. Since $S_g \geq 0$, we get

$$\text{Ric}_B + \nabla^2\varphi - \frac{m}{f}\nabla^2 f = (\lambda + \rho S_g)g_B \geq \lambda g_B.$$

First, we conclude that B^n is compact by contradiction from Proposition 4 and an analogous argument to the proof of Corollary 1. Now we apply the weak maximum principle to f in (3.12) to conclude that $\mu > 0$. Thus, by Bonnet-Myers theorem F^m is compact. Therefore $B^n \times_f F^m$ is compact. \square

In the more general setting of gradient Ricci almost solitons, we have:

Corollary 4. *Let $B^n \times_f F^m$ be a geodesically complete gradient Ricci almost soliton warped product with potential function $\tilde{\varphi}$ and soliton function $\tilde{\lambda}$ satisfying $\lambda^* = \inf_{x \in B} \lambda(x) > 0$. Suppose there exists a point $p \in B^n$ such that for all $r(x) \geq r_0$, $|\nabla\varphi(x)| \leq (\lambda^* - \varepsilon)r(x) + c$, for some positive constants c and ε , with $\varepsilon \leq \lambda^*$, where $r(x)$ is the distance function from p and r_0 is a sufficiently large positive integer. Then $B^n \times_f F^m$ is a compact manifold.*

PROOF. Applying Proposition 2 in [12], we have the proof analogous to the demonstration of Corollary 3. \square

Remark 2. *Note that Corollary 3 has the following alternative proof. First, we observe that, in the same way as in the Ambrose compactness criteria [1], we affirm that if for every geodesic ray $\gamma : [1, +\infty) \rightarrow B^n$, we have*

$$\lim_{t \rightarrow +\infty} \int_1^t \text{Ric}_h^m(\gamma', \gamma') d\xi = +\infty,$$

then B^n is compact, see Cavalcante, Oliveira and Santos [9]. Now we can prove that, under the hypotheses of Corollary 3, this criteria is satisfied on B^n , and then B^n is compact, while the compactness of the fiber F^m follows again by Bonnet-Myers theorem, so $B^n \times_f F^m$ is compact.

Acknowledgements

Enrique F. L. Agila has been partially supported by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Grant 001. José N. V. Gomes has been partially supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), of the Ministry of Science, Technology and Innovation of Brazil, Grant 310458/2021-8.

References

- [1] W. Ambrose, A theorem of Myers, *Duke Math. J.* 24 (3) (1957) 345-348.
- [2] R.L. Bishop and B. O'Neill, Manifolds of negative curvature, *Trans. Amer. Math. Soc.* 145 (1969) 1-49.
- [3] V. Borges and K. Tenenblat, Ricci almost solitons on semi-Riemannian warped products, *Math. Nachr.* 295 (1) (2022) 22-43.
- [4] V. Borges, On complete gradient Schouten solitons, *Nonlinear Anal.* 221 (2022), online version.
- [5] J. Bourguignon, Ricci curvature and Einstein metrics, *Global Differential Geometry and Global Analysis* (Berlin, 1979), *Lecture Notes in Math.*, vol. 838, Springer (Berlin, 1981), pp. 42-63.
- [6] H.-D. Cao and D. Zhou, On complete gradient Shrinking Ricci solitons, *J. Differential Geom.* 85 (2) (2010) 175-186.
- [7] G. Catino, L. Cremaschi, Z. Djadli, C. Mantegazza and L. Mazzieri, The Ricci-Bourguignon flow, *Pacific J. Math.* 287 (2) (2017) 337-370.
- [8] G. Catino and L. Mazzieri, Gradient Einstein solitons, *Nonlinear Anal.* 132 (2016) 66-94.
- [9] M. Cavalcante, J. Oliveira and M. Santos, Compactness in weighted manifolds and applications, *Results Math.* 68 (1) (2015) 143-156.
- [10] P. Daskalopoulos and N. Sesum, The classification of locally conformally flat Yamabe solitons, *Adv. Math.* 240 (2013) 346-369.
- [11] F. Feitosa, A. Freitas Filho and J. Gomes, On the construction of gradient Ricci soliton warped product, *Nonlinear Anal.* 161 (2017) 30-43.
- [12] F. Feitosa, A. Freitas Filho, J. Gomes and R. Pina, Gradient Ricci almost soliton warped product, *J. Geom. Phys.* 143 (2019) 22-32.
- [13] J. Gomes, M. Marrocos and A. Ribeiro, A note on gradient Ricci soliton warped metrics, *Math. Nachr.* 294 (10) (2021) 1879-1888.
- [14] O. Munteanu and N. Sesum, On gradient Ricci solitons, *J. Geom. Anal.* 23 (2) (2013) 539-561.
- [15] S. Pigola, M. Rimoldi and A. Setti, Remarks on non-compact gradient Ricci solitons, *Math. Z.* 268 (3) (2011) 777-790.
- [16] B. O'Neill, *Semi-Riemannian geometry with applications to relativity*, Academic press, London 1983.
- [17] G. Wei and W. Wylie, Comparison geometry for the Bakry-Emery Ricci tensor, *J. Differential Geom.* 83 (2) (2009) 337-405.
- [18] S. Zhu, *The comparison geometry of Ricci curvature*, *Comparison geometry* (Berkeley, CA, 1993-94), *Math. Sci. Res. Inst. Publ.*, vol. 30, Cambridge Univ. Press, Cambridge, (1997), pp. 221-262.

¹INSTITUTO DE MATEMÁTICA, ESTATÍSTICA E COMPUTAÇÃO CIENTÍFICA, UNIVERSIDADE ESTADUAL DE CAMPINAS, SÃO PAULO, BRAZIL.

²DEPARTAMENTO DE MATEMÁTICA, CENTRO DE CIÊNCIAS EXATAS E TECNOLOGIA, UNIVERSIDADE FEDERAL DE SÃO CARLOS, SÃO PAULO, BRAZIL.

Email address: ¹elopez@ime.unicamp.br

Email address: ²jnvgomes@ufscar.br; jnvgomes@pq.cnpq.br

URL: ¹<https://www.ime.unicamp.br>

URL: ²<https://www2.ufscar.br>