

# Extensions of metric Hom-Jacobi-Jordan algebras

Nejib Saadaoui

Université de Gabès, Institut Supérieur d'Informatique et de Multimédia de Gabès,  
Campus universitaire-BP 122, 6033 Cité El Amel 4, Gabès, TUNISIE.

September 13, 2022

## Abstract

The main purpose of this paper is to provide a second cohomology group of a (metric) Hom-Jacobi algebra with coefficients in a given representation. Moreover, we show that second cohomology group classifies abelian extensions of a (metric) Hom-Jacobi algebra algebra by a representation.

**Keywords:** Hom-Jacobi-Jordan algebras, split extensions, twofold extensions, quadratic extensions.

## Introduction

The concept of a Hom-Lie algebra was studied first, from mathematical viewpoint by Hartwig, Larsson, and Silvestrov in [5], which is a non-associative algebra satisfying the skew symmetry and the  $\sigma$ -twisted Jacobi identity. When  $\sigma = id$ , the Hom-Lie algebras degenerate to exactly the Lie algebras. In this sense, it is natural to seek for possible generalizations of known theories from Lie to Hom-Lie algebras: Hom-Lie algebras structures was defined in [3, 6]. As well as, derivations, representations and cohomology of Hom-Lie algebras was studied in [9]. Cohomology and deformations of Hom-Lie algebras was studied in [8, 1]. The Jacobi Jordan algebra is introduced by Dietrich Burde and Alice Fialowski in [4] which is a non-associative algebra satisfying the symmetric condition and the Jacobi identity algebras.

The difference with Lie algebras is thus shown in the symmetrization and therefore view to the importance of Hom algebras in several domains it is

important to generalize the results determined in Jacobi-Jordan algebras to Hom -Jacobi- Jordan algebras. Cyrille started this generalization in [2], in this paper we are interested to studied the second group of cohomology of these algebras and its relation with extensions. The paper is organized as follows. In the first section, we recall the definition of Hom-Jacobi-Jordan algebras and we give the classification of Hom-Jacobi-Jordan algebras of dimension 2. we search the conditions for the direct sum of two vector spaces  $M$  and  $V$  to be a Hom-Jacobi-Algebra. During, this investigation, we discover the definitions of representation of 2-cocycle. looking about conditions to have two direct sums  $(M, d, \gamma)$  and  $(M, d', \gamma)$  of Hom-Jacobi-Algebra are isomorphic, we find the definition of the operator of 1-cobords as well as the space of 2-cobords. In order to classify these Hom Jacobi algebras of type direct sum, we need to add certain conditions on this isomorphism, which brings us to define the extensions we generalize a known result of Lie case, that is the class of the extensions is in bijection with the second cohomology group. In the last of this section, we are interested in a particular example of extensions, called central extension and we give the central extensions of 2 dimensional Jacobi-Algebras. In section 2, we introduce metric Hom-Jacobi-Jordan algebras. In section 3 we provide the second cohomology group of a metric Hom-Jacobi-Jordan algebra with coefficients in a given representation. In section 4, we define the twofold extensions. Section 5 deals with quadratic extensions of metric Hom-Jacobi-Jordan algebras. We show that the second cohomology group classifies abelian extensions of a metric Hom-Jacobi-Jordan algebra.

## 1 Hom-Jacobi-Jordan algebras

In this section, first we recall some basic facts about Hom-Jacobi-Jordan algebras, we classify 2-dimensional Hom-Jacobi-Jordan algebras, we define a second cohomology groupe with coefficients in a given representation and then in a trivial, adjoint, coadjoint representation. we define a c-Hom-Jacobi-algebra. Started by a direct sum of two vectoriel space  $J$  and  $V$ , we define representation, second space of cocyle, second space of cobord, second group of cohomology and extension of Hom-Jacobi-Jordan algebra. We show that the class of the extensions of a Hom Jordan Jacobi algebra  $J$  by  $V$  is in bijection with the second cohomology group of  $J$  on  $V$ . Moreover, we give all 2-dimensional extension of Hom-Jacobi-Jordan algebra by 2- dimensional abelian Hom-Jacobi-Jordan algebra.

**Definition 1.1.** ([2]) A Hom-Jacobi-Jordan algebra is a triple  $(J, \delta, \alpha)$ , where  $J$  is a vector space equipped with a symmetric bilinear map  $\delta: J \times J \rightarrow J$

and a linear map  $\alpha: J \rightarrow J$  such that

$$\delta(\alpha(x), \delta(y, z)) + \delta(\alpha(y), \delta(z, x)) + \delta(\alpha(z), \delta(x, y)) = 0 \quad (1.1)$$

for all  $x, y, z$  in  $J$ , which is called Hom-Jacobi identity.

We recover Jacobi-Jordan algebras when the linear map  $\delta$  is the identity map.

A Hom-Jacobi-Jordan-algebra is called abelian if the bilinear map  $\delta$  vanishes for all elements in  $J$ .

A Hom-Jacobi-Jordan-algebra is called a multiplicative Hom-Jacobi-Jordan if  $\alpha$  is an algebraic morphism with

$$\alpha(\delta(x, y)) = \delta(\alpha(x), \alpha(y)) \quad (1.2)$$

for any  $x, y \in J$ .

Two Hom-Jacobi-Jordan algebra  $(J, \delta, \alpha)$  and  $(J', \delta', \alpha')$  are said to be isomorphic if there exists a bijective algebra morphism  $\phi: J \rightarrow J'$  compatible with  $\alpha$  and  $\alpha'$ , i.e

$$\phi(\delta(x, y)) = \delta(\phi(x), \phi(y))' \text{ and } \phi \circ \alpha = \alpha' \circ \phi. \quad (1.3)$$

In general for  $n$ -dimensional case in terms of structure constants we have:

$$\delta(e_i, e_j) = \sum_{s=1}^n d[i, j][s] e_s \quad \text{and} \quad \alpha(e_j) = \sum_{s=1}^n a_{sj} e_s. \quad (1.4)$$

Substituting (1.4) in the symmetry identity  $\delta(e_i, e_j) = \delta(e_j, e_i)$  yields

$$d[i, j][s] = d[j, i][s]. \quad (1.5)$$

Substituting (1.4) in the Hom-Jacobi-Jordan identity (1.1) yields

$$\sum_{1 \leq s, l \leq n} (a_{si} d[j, k][l] + a_{sj} d[k, i][l] + a_{sk} d[i, j][l]) d[s, l][r] = 0. \quad (1.6)$$

Substituting (1.4) in the multiplicativity conditions (1.2) yields

$$\sum_{1 \leq k \leq n} d[i, j][k] a_{sk} = \sum_{1 \leq p, q \leq n} a_{pi} a_{qj} d[p, q][s] \quad (1.7)$$

Let  $(J, \delta, \alpha)$  be  $n$ -dimensional Hom-Jacobi-Jordan algebra with ordered basis  $(e_1, \dots, e_n)$  and  $J'$  be  $n$ -dimensional vector spaces with ordered basis

$(e'_1, \dots, e'_n)$ . Let  $f: J \rightarrow J'$  be an isomorphism map. Let  $\alpha' = f\alpha f^{-1}$ . We set with respect to a basis  $(e'_1, \dots, e'_n)$ :

$$f(e_j) = \sum_{i=1}^n f_{ij}e'_i, \quad (1.8)$$

$$\delta'(e'_i, e'_j) = \sum_{k=1}^n d'[i, j][k]e'_k, \quad i, j \in \{1, \dots, n\}. \quad (1.9)$$

Condition (1.3) translates to the following equation

$$\sum_{k=1}^n d[i, j][k]f_{sk} = \sum_{1 \leq p, q \leq n} f_{pi}f_{qj}c'[p, q][s], \quad i, j, s \in \{1, \dots, n\}. \quad (1.10)$$

Then, if the previous condition satisfied,  $J'$  is a Hom-Jacobi-Jordan algebra isomorphic to  $J$ .

For the rest of this section, we use the previous systems for find classification of 2-dimensional complex Hom-Jacobi-Jordan algebras.

The following Lemma describes the class of complex 2-dimensional Hom-Jacobi-Jordan algebras whose the matrix of  $\alpha$  is of the form  $\begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}$ .

**Lemma 1.2.** *Let  $(J, [\cdot, \cdot], \alpha)$  be 2-dimensional Hom-Jacobi-Jordan algebra with ordered basis  $(u_1, u_2)$ . Take  $\alpha(u_1) = au_1$  and  $\alpha(u_2) = bu_2$ . Then there is a basis  $(e_1, e_2)$  of  $J$  in which  $(J, [\cdot, \cdot], \alpha)$  has one of following form.*

- $J_1^1(0, b, 0): [e_1, e_1] = e_2, [e_1, e_2] = e_2,$   
 $\alpha(e_1) = 0e_1, \alpha(e_2) = be_2$
- $J_2^1(0, b, 0): [e_1, e_1] = e_2,$   
 $\alpha(e_1) = 0e_1, \alpha(e_2) = be_2$
- $J_3^1(0, b, 0): [e_1, e_1] = e_1, [e_2, e_2] = e_1$   
 $\alpha(e_1) = 0e_1, \alpha(e_2) = be_2$
- $J_4^1(0, b, 0): [e_1, e_1] = e_1,$   
 $\alpha(e_1) = 0e_1, \alpha(e_2) = be_2$
- $J_5^1(0, b, -bc): [e_1, e_1] = e_2, [e_1, e_2] = e_2, [e_2, e_2] = ce_2$   
 $\alpha(e_1) = 0e_1, \alpha(e_2) = -bce_1 + be_2$
- $J_6^1(0, b, 0): [e_1, e_1] = e_1 + e_2,$   
•  $\alpha(e_1) = 0e_1, \alpha(e_2) = be_2$

- $J_7^1(0, b, 0): [e_2, e_2] = e_1$   
 $\alpha(e_1) = 0e_1, \alpha(e_2) = be_2$
- $J_8^1(0, b, 0): [e_1, e_2] = e_2, \alpha(e_1) = 0e_1, \alpha(e_2) = be_2$
- $J_9^1(0, b, -b): [e_1, e_2] = e_2, [e_2, e_2] = e_2$   
 $\alpha(e_1) = 0e_1, \alpha(e_2) = -be_1 + be_2$
- $J_{10}^1(a, b, 0): [e_1, e_1] = e_2$   
 $\alpha(e_1) = ae_1, \alpha(e_2) = be_2$
- $J_{11}^1(a, b, 1): [e_1, e_1] = e_2,$   
 $\alpha(e_1) = ae_1, \alpha(e_2) = e_1 + be_2$

where the omitted products are being zero and  $[e_2, e_1] = [e_1, e_2]$ .

*Proof.*

**Case 1.**  $a_{11} = 0$

**Subcase 1.1.**  $d[1, 1][1] = 0$  and  $d[1, 1][2] \neq 0$ .

We solve by a computer algebra, the system of equations (1.5) and (1.6) such that:  $a_{11} = a_{12} = a_{21} = 0, d[1, 1][1] = 0$  and  $d[1, 1][2] \neq 0$  we obtain the following solutions:

- (i)  $d[1, 2][1] = 0, d[2, 1][1] = 0, d[2, 1][2] = d[1, 2][2], d[2, 2][1] = 0, d[2, 2][2] = 0.$
- (ii)  $d[1, 2][1] = 0, d[2, 1][1] = 0, d[2, 1][2] = 0, d[1, 2][2] = 0, d[2, 2][1] = 0, d[2, 2][2] = 0.$

Hence we get the following corresponding Hom-Jacobi-Jordan algebras:

$$s_1 : \begin{cases} [u_1, u_1] = d[1, 1][2]u_2 \\ [u_1, u_2] = d[1, 2][2]u_2 \\ [u_2, u_1] = d[1, 2][2]u_2 \\ [u_2, u_2] = 0 \end{cases} \quad s_2 : \begin{cases} [e_1, e_1] = d[1, 1][2]e_2 \\ [e_1, e_2] = 0 \\ [e_2, e_1] = 0 \\ [e_2, e_2] = 0 \end{cases}$$

Using  $s_1$ , take  $d[1, 1][1] = 0, d[1, 2][1] = 0, d[2, 2][1] = 0, d[2, 2][2] = 0$  and  $f_{12} = f_{21} = 0, c[1, 1][1] = 0, c[1, 1][2] = 1, c[1, 2][1] = 0, c[1, 2][2] = 1, c[2, 1][1] = 0, c[2, 1][2] = 1, c[2, 2][1] = 0, c[2, 2][2] = 0$ , in (1.10), we obtain

$f = \begin{pmatrix} d[1,2][2] & 0 \\ 0 & \frac{d[1,2][2]^2}{d[1,1][2]} \end{pmatrix}$  (the other solutions are not isomorphism). Then,  $f\alpha f^{-1} = \alpha$ . Hence  $\alpha' = \alpha$  and

$$\begin{cases} [e_1, e_1]' = e_2 \\ [e_1, e_2]' = e_2 \\ [e_2, e_1]' = e_2 \\ [e_2, e_2]' = 0 \end{cases}$$

Therefore we obtain the Hom-Jordan-Jacobi algebra  $J_1^1(0, b, 0)$ .

Using  $s_2$ , take  $d[1,1][1] = 0$ ,  $d[1,2][1] = 0$ ,  $d[1,2][2] = 0$ ,  $d[2,2][1] = 0$ ,  $d[2,2][2] = 0$  and  $f_{12} = f_{21} = 0$ ,  $c[1,1][1] = 0$ ,  $c[1,1][2] = 1$ ,  $c[1,2][1] = 0$ ,  $c[1,2][2] = 0$ ,  $c[2,1][1] = 0$ ,  $c[2,1][2] = 0$ ,  $c[2,2][1] = 0$ ,  $c[2,2][2] = 0$ , in (1.10),

we obtain  $f = \begin{pmatrix} f[1,1] & 0 \\ 0 & \frac{f[1,1]^2}{d[1,1][2]} \end{pmatrix}$  (the other solutions are not isomorphism).

Then,  $f\alpha f^{-1} = \alpha$ . Hence  $\alpha' = \alpha$  and

$$\begin{cases} [e_1, e_1]' = e_2 \\ [e_1, e_2]' = 0 \\ [e_2, e_1]' = 0 \\ [e_2, e_2]' = 0 \end{cases}$$

therefore we obtain the Hom-Jordan-Jacobi algebra  $J_2^1(0, b, 0)$ .

**Subcase 1.2.**  $d[1,1][1] \neq 0$  and  $d[1,1][2] = 0$  :

Taking  $a_{11} = a_{12} = a_{21} = 0$ ,  $d[1,1][2] = 0$  in (1.5) and (1.6). The solutions gives the following table of multiplication:

$$s_1 : \begin{cases} [u_1, u_1] = d[1,1][1]u_1 \\ [u_1, u_2] = 0 \\ [u_2, u_1] = 0 \\ [u_2, u_2] = d[2,2][1]u_1 \end{cases} \quad s_2 : \begin{cases} [u_1, u_1] = d[1,1][1]u_1 \\ [u_1, u_2] = 0 \\ [u_2, u_1] = 0 \\ [u_2, u_2] = 0 \end{cases}$$

Using  $s_1$ , setting  $d[1,1][2] = 0$ ,  $d[1,2][1] = 0$ ,  $d[1,2][2] = 0$ ,  $d[2,1][1] = 0$ ,  $d[2,1][2] = 0$ ,  $d[2,2][2] = 0$ , and  $f_{12} = f_{21} = 0$ ,  $c[1,1][2] = 1$ ,  $c[1,2][1] = 0$ ,  $c[1,2][2] = 0$ ,  $c[2,1][1] = 0$ ,  $c[2,1][2] = 0$ ,  $c[2,2][2] = 0$ , in (1.10), we

obtain  $f = \begin{pmatrix} d[1,1][1] & 0 \\ 0 & \sqrt{d[1,1][1]d[2,2][1]} \end{pmatrix}$  then  $\alpha' = f\alpha f^{-1} = \alpha$  Then,  $f\alpha f^{-1} = \alpha$ . Hence  $\alpha' = \alpha$  and

$$\begin{cases} [e_1, e_1]' = e_1 \\ [e_1, e_2]' = 0 \\ [e_2, e_1]' = 0 \\ [e_2, e_2]' = e_1 \end{cases}$$

The algebra obtained from this subcase is equivalent to  $J_3^1(0, b, 0)$ .

Using  $s_2$ , take  $d[1, 1][2] = 0$ ,  $d[1, 2][1] = 0$ ,  $d[1, 2][2] = 0$ ,  $d[2, 1][1] = 0$ ,  $d[2, 1][2] = 0$ ,  $d[2, 2][1] = 0$ ,  $d[2, 2][2] = 0$ , and  $f_{12} = f_{21} = 0$ ,  $c[1, 1][2] = 1$ ,  $c[1, 2][1] = 0$ ,  $c[1, 2][2] = 0$ ,  $c[2, 1][1] = 0$ ,  $c[2, 1][2] = 0$ ,  $c[2, 2][1] = 0$ ,  $c[2, 2][2] = 0$ , in (1.10), we obtain  $f = \begin{pmatrix} d[1, 1][1] & 0 \\ 0 & 1 \end{pmatrix}$  then  $\alpha' = f\alpha f^{-1} = \alpha$

Then,  $f\alpha f^{-1} = \alpha$ . Hence  $\alpha' = \alpha$  and

$$\begin{cases} [e_1, e_1]' = e_1 \\ [e_1, e_2]' = 0 \\ [e_2, e_1]' = 0 \\ [e_2, e_2]' = e_1 \end{cases}$$

thus the algebra  $J_4^1(0, b, 0)$  is obtained.

**Subcase 1.3.**  $d[1, 1][1] \neq 0$  and  $d[1, 1][2] \neq 0$  We obtain:

$$s_1 : \begin{cases} [u_1, u_1] = d[1, 1][1]u_1 + d[1, 1][2]u_2 \\ [u_1, u_2] = \frac{d[2, 1][1]}{d[1, 1][1]}(d[1, 1][1]u_1 + d[1, 1][2]u_2) \\ [u_2, u_1] = \frac{d[2, 1][1]}{d[1, 1][1]}(d[1, 1][1]u_1 + d[1, 1][2]u_2) \\ [u_2, u_2] = -\frac{d[2, 1][1]}{d[1, 1][2]}(d[1, 1][1]u_1 + d[1, 1][2]u_2) \end{cases} \quad s_2 : \begin{cases} [u_1, u_1] = d[1, 1][1]u_1 + d[1, 1][2]u_2 \\ [u_1, u_2] = 0 \\ [u_2, u_1] = 0 \\ [u_2, u_2] = 0 \end{cases}$$

Let  $v_1 = d[1, 1][1]u_1 + d[1, 1][2]u_2$ . Hence, the matrix of  $\alpha$  in the basis  $(u_1, v_1)$  is  $\beta = \begin{pmatrix} 0 & -a_{2,2}d[1, 1][1] \\ 0 & a_{22} \end{pmatrix}$  and a simple calcul gives that the bracket has the following form

$$s_1 : \begin{cases} [u_1, u_1] = v_2 \\ [u_1, v_2] = \lambda v_2 \\ [v_2, u_1] = \lambda v_2 \\ [v_2, v_2] = \lambda c v_2 \end{cases} \quad s_2 : \begin{cases} [u_1, u_1] = v_2 \\ [u_1, v_2] = 0 \\ [v_2, u_1] = 0 \\ [v_2, v_2] = 0 \end{cases}$$

Now, take in (1.10),  $d[1, 1][1] = 0$ ,  $d[1, 1][2] = 1$ ,  $d[1, 2][1] = 0$ ,  $d[1, 2][2] = \lambda$ ,  $d[2, 1][1] = 0$ ,  $d[2, 1][2] = \lambda$ ,  $d[2, 2][1] = 0$ ,  $d[2, 2][2] = \lambda c$ ,  $f_{12} = f_{21} = 0$ . We obtain  $c[1, 1][1] = 0$ ,  $c[1, 1][2] = \frac{f_{22}}{f_{11}}$ ,  $c[1, 2][1] = 0$ ,  $d[1, 2][2] = \frac{\lambda}{f_{11}}$ ,  $c[2, 1][1] = 0$ ,  $c[2, 1][2] = \frac{\lambda}{f_{11}}$ ,  $d[2, 2][1] = 0$ ,  $d[2, 2][2] = \frac{\lambda a_{2,2}}{f_{22}}$ . Then, with  $f_{11} = \lambda$  and  $f_{22} = \lambda^2$ , we obtain

$$\begin{cases} [u_1, u_1]' = v_2 \\ [u_1, v_2]' = v_2 \\ [u_2, u_1]' = v_2 \\ [e_2, e_2]' = \frac{c}{\lambda} v_2 \end{cases}$$

and  $\beta' = f\beta f^{-1} = \begin{pmatrix} 0 & -\frac{c}{\lambda}a_{22} \\ 0 & a_{22} \end{pmatrix}$  that is  $J_5^1(0, b, -bc)$ .

For  $s_2$  we take  $v_1 = \frac{1}{d[1,1][1]}u_1$  and  $v_2 = \frac{d[1,1][2]}{d[1,1][1]^2}u_2$ . Hence, the matrix of  $\alpha$  in the basis  $(v_1, v_2)$  is  $\beta = \begin{pmatrix} 0 & 0 \\ 0 & a_{22} \end{pmatrix} = \alpha$  and a simple calcul gives that the bracket has the following form

$$\left\{ \begin{array}{l} [v_1, v_1] = v_1 + v_2 \\ [v_1, v_2] = 0 \\ [v_2, v_1] = 0 \\ [v_2, v_2] = 0 \end{array} \right.$$

Hence we obtain  $J_6^1(0, b, 0)$ .

**Subcase 1.4.**  $[e_1, e_1] = 0$  and  $d[2, 2][1] = 0, d[2, 2][2] \neq 0$ :

The solution is trivial.

**Subcase 1.5.**  $[e_1, e_1] = 0$  and  $d[2, 2][1] \neq 0, d[2, 2][2] = 0$ :

The Hom-Jacobi-Jordan algebra obtained from this case is equivalent to  $J_7^1(0, b, 0)$ .

**Subcase 1.6.**  $[e_1, e_1] = 0$  and  $[e_2, e_2] = 0$ ,

The Hom-Jacobi-Jordan algebra obtained from this case is equivalent to  $J_8^1(0, b, 0)$ .

**Subcase 1.7.**  $[e_1, e_1] = 0$  and  $[e_2, e_2] \neq 0$ ,

The Hom-Jacobi-Jordan algebra obtained from this case is equivalent to  $J_9^1(0, b, 0)$ .

**Case 2.**  $a_{11} \neq 0, a_{22} \neq 0$ ,

**Subcase 2.1.**  $d[1, 1][1] = 0, d[1, 1][2] \neq 0$ . We obtain  $J_{10}^2(a, b, 0)$ .

**Subcase 2.2.**  $d[1, 1][1] = 0, d[1, 1][2] \neq 0$

We obtain  $J_{10}^2(a, b, 0)$ .

**Subcase 2.3.**  $d[1, 1][1] \neq 0, d[1, 1][2] = 0$ . We obtain the zero solution.

**Subcase 2.4.**  $d[1, 1][1] \neq 0, d[1, 1][2] \neq 0$ .

We obtain  $J_{11}^2(a, b, 1)$ .

**Subcase 2.5.**  $d[1, 1][1] = 0, d[1, 1][2] = 0$ .

We obtain  $J_{10}^2(b, a, 0)$



■

The following Lemma describes the class of complex 2-dimensional Hom-Jacobi-Jordan algebras whose the matrix of  $\alpha$  is of the form  $\begin{pmatrix} a & 1 \\ 0 & a \end{pmatrix}$ .

**Lemma 1.3.** *Let  $(J, [\cdot, \cdot], \alpha)$  be 2-dimensional Hom-Jacobi-Jordan algebra with ordered basis  $(u_1, u_2)$ . Take  $\alpha(u_1) = au_1$  and  $\alpha(u_2) = u_1 + au_2$ . Then there is a basis  $(e_1, e_2)$  of  $J$  in which  $(J, [\cdot, \cdot], \alpha)$  has one of following form.*

- $J_{12}^2(0, 0, 1): [e_1, e_1] = e_2,$   
 $\alpha(e_1) = 0, \alpha(e_2) = e_1,$
- $J_{13}^2(0, 0, c): [e_1, e_1] = e_2, [e_2, e_2] = e_2$   
 $\alpha(e_1) = 0, \alpha(e_2) = ce_1,$
- $J_{15}^2(0, 0, 1): [e_2, e_2] = e_1,$   
 $\alpha(e_1) = 0, \alpha(e_2) = e_1,$
- $J_{16}^2(0, 0, c): [e_1, e_2] = e_1, [e_2, e_2] = e_1,$   
 $\alpha(e_1) = 0, \alpha(e_2) = ce_1,$
- $J_{17}^2(0, 0, 1): [e_2, e_2] = e_2,$   
 $\alpha(e_1) = 0, \alpha(e_2) = e_1,$
- $J_{18}^2(0, 0, 1): [e_1, e_2] = e_1,$   
 $\alpha(e_1) = 0, \alpha(e_2) = e_1,$
- $J_{19}^2(a, a, 1): [e_1, e_1] = e_2,$   
 $\alpha(e_1) = ae_1, \alpha(e_2) = e_1 + ae_2,$
- $J_{20}^2(a, a, 1): [e_2, e_2] = e_1,$   
 $\alpha(e_1) = ae_1, \alpha(e_2) = e_1 + ae_2,$

Combining these two previous lemma we get the following Theorem.

**Theorem 1.4.** *The isomorphism classes of 2-dimensional Hom-Jacobi-Jordan algebra can be represented in Lemma 1.2 and Lemma 1.3.*

## 1.1 The second cohomology group of Hom-Jacobi-Jordan algebras

Let  $J$  and  $V$  two vectors spaces and let  $\alpha: J \rightarrow J, \beta: V \rightarrow V$  be to linear maps. For convenience, we denotes

$$\begin{aligned} M &= J \times V = V \times J = J \oplus V = JV = VJ \\ &= \{(x, v) \mid x \in J, v \in V\} = \{(v, x) \mid x \in J, v \in V\} \end{aligned}$$

$$M = J \times V = V \times J = J \oplus V = \{(x, v) \mid x \in J, v \in V\} = \{(v, x) \mid x \in J, v \in V\}.$$

Define the linear map  $\gamma: M \rightarrow M, \gamma(x + v) = \alpha(x) + \beta(v)$ , for all  $x \in J, v \in V$ . Let  $d: M \times M \rightarrow M$  be a symmetric bilinear map. We will determine the necessary and sufficient conditions for  $M$  to be a Hom-Jordan-Jacobi algebras and  $V$  is an abelian ideal of  $M$  (i.e  $d(x, v) \in V$  for all  $x \in J, v \in V$ ). these conditions lead us to define the second cohomology group of Hom-Jacobi-Jordan algebras.

Since  $d \in S^2(M)$ , Then there exists  $\delta \in S^2(J), \lambda \in S^2(JV, V), \lambda' \in S^2(JV, J), \alpha \in S^2(J, V), \mu \in S^2(V)$  such that  $d = \delta + \lambda + \lambda' + \mu + \alpha$ . We assume that  $\lambda'$  and  $\mu$  are zero (that is because  $V$  is an abelian ideal of  $M$ ). For the multiplicative Hom-Jacobi-Jordan for an element  $(x, y) \in J^2$ :

$$\gamma \circ d(x, y) = d \circ \gamma(x, y) \iff \alpha(\delta(x, y)) + \beta(\theta(x, y)) = \delta(\alpha(x), \alpha(y)) + \theta(\alpha(x), \alpha(y)).$$

Hence

$$\alpha(\delta(x, y)) = \delta(\alpha(x), \alpha(y))$$

and

$$\beta(\theta(x, y)) = \theta(\alpha(x), \alpha(y)).$$

The last equality is called 2-Hom-cochain condition.

**Definition 1.5.** A symmetric bilinear map with values in  $V$  is defined to be a 2-Hom-cochain  $\theta \in S^2(J, V)$  such it is compatible with  $\alpha$  and  $\beta$  in the sense that  $\beta \circ \theta = \theta \circ \alpha$ . Denote  $C_{\alpha, \beta}^2(J, V)$  the set of 2-Hom-cochain:

$$C_{\alpha, \beta}^2(J, V) = \{\theta \in S^2(J, V) \mid \beta \circ \theta = \theta \circ \alpha\}.$$

For the multiplicative Hom-Jacobi-Jordan for an element  $(x, v) \in JV$ :

$$\gamma \circ d(x, v) = d \circ \gamma(x, v) \iff \beta(\lambda(x, v)) = \lambda(\alpha(x), \beta(v)).$$

We define the linear map  $\rho: J \rightarrow \text{End}(V)$  by  $\rho(x)(v) = \lambda(x, v)$ . Thus

$$\beta(\rho(x)v) = \rho(\alpha(x))\beta(v) \tag{1.11}$$

For the Hom-Jacobi identity for an element  $(x, y, v) \in J^2V$ :

$$\begin{aligned} & d(\gamma(x), d(y, v)) + d(\gamma(y), d(v, y)) + d(\gamma(v), d(x, y)) \\ &= \lambda(\alpha(x), \lambda(y, v)) + \lambda(\alpha(y), \lambda(v, x)) + \lambda(\beta(v), \delta(x, y)) \\ &= \rho(\alpha(x) \circ \rho(y)v) + \lambda(\alpha(y), \lambda(v, x)) + \lambda(\beta(v), \delta(x, y)) \end{aligned}$$

Therefore

$$\begin{aligned} & d(\gamma(x), d(y, v)) + d(\gamma(y), d(v, y)) + d(\gamma(v), d(x, y)) = 0 \\ \iff & \rho(\delta(x, y))\beta(v), = -\rho(\alpha(x)) \circ \rho(y)v - \rho(\alpha(y)) \circ \rho(x)v. \end{aligned} \quad (1.12)$$

The Equations (1.11) and (1.35) are called representation conditions.

**Definition 1.6.** A representation of a Hom-Jacobi-Jordan algebra  $(J, \delta, \alpha)$  on a vector space  $V$  with respect to  $\beta \in \text{End}(V)$  is a linear map  $\rho: J \rightarrow \text{End}(V)$ , such that representation conditions are satisfied i.e:

$$\rho(\alpha(x)) \circ \beta = \beta \circ \rho(x) \quad (1.13)$$

$$\rho(\delta(x, y)) \circ \beta = -\rho(\alpha(x))\rho(y) - \rho(\alpha(y))\rho(x). \quad (1.14)$$

for all  $x, y \in J$ . We denote a representation of a Hom-Jacobi-Jordan algebra  $(J, \delta, \alpha)$  by a triple  $(V, \rho, \beta)$ .

For the Hom-Jacobi identity for an element  $(x, y, z) \in J^3$ :

$$\begin{aligned} & d(\gamma(x), d(y, z)) + d(\gamma(y), d(z, y)) + d(\gamma(z), d(x, y)) \\ &= \delta(\alpha(x), \delta(y, z)) + \delta(\alpha(y), \delta(z, x)) + \delta(\alpha(z), \delta(x, y)) \\ &+ \theta(\alpha(x), \delta(y, z)) + \theta(\alpha(y), \delta(z, x)) + \theta(\alpha(z), \delta(x, y)) \\ &+ \rho(\alpha(x))\theta(y, z) + \rho(\alpha(y))\theta(z, y) + \rho(\alpha(z))\theta(x, y) \end{aligned} \quad (1.15)$$

Hence

$$\begin{aligned} & d(\gamma(x), d(y, z)) + d(\gamma(y), d(z, y)) + d(\gamma(z), d(x, y)) = 0 \\ \iff & \underbrace{\delta(\alpha(x), \delta(y, z)) + \delta(\alpha(y), \delta(z, x)) + \delta(\alpha(z), \delta(x, y))}_{\in J} \\ & + \underbrace{\theta(\alpha(x), \delta(y, z)) + \theta(\alpha(y), \delta(z, x)) + \theta(\alpha(z), \delta(x, y))}_{\in V} \\ & + \underbrace{\rho(\alpha(x))\theta(y, z) + \rho(\alpha(y))\theta(z, y) + \rho(\alpha(z))\theta(x, y)}_{\in V} = 0. \end{aligned}$$

Thus

$$\begin{aligned}
& \left( d(\gamma(x), d(y, z)) + d(\gamma(y), d(z, y)) + d(\gamma(z), d(x, y)) = 0 \right) \\
\iff & \left( \delta(\alpha(x), \delta(y, z)) + \delta(\alpha(y), \delta(z, x)) + \delta(\alpha(z), \delta(x, y)) = 0 \text{ and} \right. \\
& \quad \theta(\alpha(x), \delta(y, z)) + \theta(\alpha(y), \delta(z, x)) + \theta(\alpha(z), \delta(x, y)) \\
& \quad \left. + \rho(\alpha(x))\theta(y, z) + \rho(\alpha(y))\theta(z, y) + \rho(\alpha(z))\theta(x, y) = 0 \right). \quad (1.16)
\end{aligned}$$

The latest equality is called 2-cocycle condition. Any Hom 2-cochain satisfies the 2-cocycle condition is called 2-cocycle of  $J$  on  $V$ . The space of all 2-cocycles of  $J$  on  $V$  is denoted  $Z^2(J, V)$ .

**Definition 1.7.** We call 2-coboundary operator of Hom-Jacobi-Jordan algebra  $J$  the map

$$d^2 : C_{\alpha, \beta}^2(J, V) \rightarrow S^3(J, V), \quad f \mapsto d^2(f)$$

defined by

$$\begin{aligned}
d^2(f)(x, y, z) = & f(\delta(x, y), \alpha(z)) + f(\delta(x, z), \alpha(y)) + f(\alpha(x), \delta(y, z)) \\
& + \rho(\alpha(x))f(y, z) + \rho(\alpha(y))f(x, z) + \rho(\alpha(z))f(x, y). \quad (1.17)
\end{aligned}$$

Clearly, a 2-Hom-cochain  $f$  is a 2-cocycle if and only if  $d^2(f) = 0$ . Hence,  $\ker(d^2) = Z^2(J, V)$ .

We summarize the main facts in the theorem below.

**Theorem 1.8.**  *$(M, d, \gamma)$  is a multiplicative Hom-Jacobi-Jordan algebra and  $V$  is an abelian ideal of  $M$  if and only if the following conditions are holds:*

- (1)  $\lambda' = \mu = 0$ ;
- (2)  $(J, \delta, \alpha)$  is a Hom-Jacobi-Jordan algebra;
- (3) the linear map  $\rho: J \rightarrow \text{End}(V)$ ,  $x \mapsto \lambda(x, \cdot)$  define a representation of  $J$  on  $V$ ;
- (4)  $\theta$  is a 2-cocycle on  $V$  (i.e  $\theta \in Z^2(J, V)$  ).

Let  $d' = d + \theta'$  where  $\theta'$  is a 2-Hom-cochain. We assume that  $(M, d, \gamma)$  and  $(M, d', \gamma)$  are two isomorphic Hom-Jacobi-Jordan algebras. Then there exist an isomorphism  $\Phi: (M, d, \gamma) \rightarrow (M, d', \gamma)$  such that  $\Phi(d(x, y)) = d'(\Phi(x), \Phi(y))$ .

For all  $x \in J$ , let  $\Phi(x) = s(x) + i(x)$  where  $s(x) \in J$  and  $i(x) \in V$  and  $\Phi(v) = s'(v) + i'(v)$  where  $s'(v) \in J$  and  $i'(v) \in V$ . We have

$$\begin{aligned}
\Phi(d(x, y)) = d'(\Phi(x), \Phi(y)) &\iff \Phi(\delta(x, y) + \theta(x, y)) = d'(s(x) + i(x), s(y) + i(y)) \\
&\iff \Phi(\delta(x, y) + \theta(x, y)) = d(s(x) + i(x), s(y) + i(y)) + \theta'(s(x), s(y)) \\
&\iff s(\delta(x, y)) + i(\delta(x, y)) = \delta(s(x), s(y)) + \rho(s(x))i(y) + \rho(s(y))i(x) + \theta(s(x), s(y)) \\
&\quad + \theta'(s(x), s(y)) \\
&\iff \begin{cases} s(\delta(x, y)) = \delta(s(x), s(y)) \\ i(\delta(x, y)) = \rho(s(x))i(y) + \rho(s(y))i(x) + \theta(s(x), s(y)) + \theta'(s(x), s(y)) \end{cases}
\end{aligned}$$

Define  $f: s(J) \rightarrow V$  by

$$f(s(x)) = i(x).$$

for all  $x \in J$ . Then

$$\begin{aligned}
f(s(\delta(x, y))) &= \delta(s(x), s(y)) + \rho(s(x))i(y) + \rho(s(y))i(x) + \theta(s(x), s(y)) \\
&\quad + \theta'(s(x), s(y)) \\
\iff f(\delta(s(x), s(y))) &= \rho(s(x))f(s(y)) + \rho(s(y))f(s(x)) + \theta(s(x), s(y)) \\
&\quad + \theta'(s(x), s(y)) \\
\iff \theta'(s(x), s(y)) + \theta(s(x), s(y)) &= f(\delta(s(x), s(y))) - \rho(s(x))f(s(y)) - \rho(s(y))f(s(x)) \\
&\iff \theta'(a, b) + \theta(a, b) = d^1(f)(a, b)
\end{aligned}$$

where  $a = s(x)$ ,  $b = s(y)$  and

$$d^1(f)(a, b) = f(\delta(a, b)) - \rho(a)f(b) - \rho(b)f(a) \quad (1.18)$$

We say that  $\theta' + \theta$  satisfies the 2-cobord condition.

We have

$$\theta'(a, b) = d^1(f)(a, b) - \theta(a, b) \quad (1.19)$$

Moreover, since  $\Phi$  is an isomorphism of Hom-Jacobi-Jordan algebra, we have  $\Phi\gamma(x) = \gamma(\Phi(x))$ . Hence

$$s(\alpha(x)) = \alpha(s(x)) \quad (1.20)$$

and

$$i(\alpha(x)) = \beta(i(x)). \quad (1.21)$$

for all  $x \in J$ .

Applying  $f$  to the latest equality, we obtain  $f(s(\alpha(x))) = \beta f(s(x))$ . Using (1.20), we have  $f(\alpha(s(x))) = \beta f(s(x))$ . Hence

$$f(\alpha(a)) = \beta f(a). \quad (1.22)$$

Then we say that  $f$  is a 1-Hom cochain. We assume that  $s(J) = J$  and we denote  $C_{\alpha,\beta}^1(J, V)$  the space of all 1-Hom cochain of  $J$  on  $V$ .

Since  $(M, d, \gamma)$  and  $(M, d', \gamma)$  are two Hom-Jacobi-Jordan algebras, using Theorem 1.8,  $\theta$  and  $\theta'$  are 2-cocyle. Thus with (1.19), we deduce that  $d^1(f)$  is a 2-cocyle. that is  $d^2(d^1(f))(a, b, c) = 0$  for all  $a, b, c \in s(J)$

Bringing these results together we have

**Definition 1.9.** We call 1-coboundary operator of Hom-Jacobi-Jordan algebra  $J$  the map

$$d^1 : C_{\alpha,\beta}^1(J, V) \rightarrow S^2(J, V), \quad f \mapsto d^1 f$$

defined by

$$d^1(f)(x, y) = f(\delta(x, y)) - \rho(x)f(y) - \rho(y)f(x). \quad (1.23)$$

**Proposition 1.10.**

$$d^2 \circ d^1 = 0.$$

*Proof.* We Have

$$d^1(f)(x, y) = f(\delta(x, y)) - \rho(x)f(y) - \rho(y)f(x).$$

Then

$$\begin{aligned} & d^2 \circ d^1(f)(x, y, z) \\ &= d^1(f)([x, y], \alpha(z)) + d^1(f)\delta(\delta(x, z), \alpha(y)) + d^1(f)(\alpha(x), \delta(y, z)) \\ &+ \rho(\alpha(x))d^1(f)(y, z) + \rho(\alpha(y))d^1(f)(x, z) + \rho(\alpha(z))d^1(f)(x, y) \\ &= f((\delta(x, y), \alpha(z))) + f(\delta(x, z), \alpha(y)) + f(\delta(y, z), \alpha(x)) \end{aligned} \quad (1.24)$$

$$- \rho(\delta(x, y))f(\alpha(z)) - \rho(\alpha(x))\rho(y)f(z) - \rho(\alpha(y))\rho(x)f(z) \quad (1.25)$$

$$- \rho(\delta(x, z))f(\alpha(y)) - \rho(\alpha(x))\rho(z)f(y) - \rho(\alpha(z))\rho(x)f(y) \quad (1.26)$$

$$- \rho(\delta(y, z))f(\alpha(x)) - \rho(\alpha(z))\rho(y)f(x) - \rho(\alpha(y))\rho(z)f(x) \quad (1.27)$$

$$- \rho(\alpha(z))f(\delta(x, y)) + \rho(\alpha(z))f(\delta(x, y)) - \rho(\alpha(y))f(\delta(x, z)) \quad (1.28)$$

$$+ \rho(\alpha(y))f(\delta(x, z)) - \rho(\alpha(x))f(\delta(y, z)) + \rho(\alpha(x))f(\delta(y, z)) \quad (1.29)$$

By the Hom-Jacobi identity, we obtain that Eq.(1.24)=0.

Since  $f \circ \alpha = \beta \circ f$  and the linear map  $\rho$  satisfies (1.14), we have that Eq.(1.25)=0, Eq.(1.26)=0 and Eq.(1.27)=0. The sum of the other six items is zero obviously. Therefore, we have  $d^2 \circ d^1(f)(x, y, z) = 0$ . The proof is completed. ■

We have with respect to the cohomology defined by the coboundary operators

$$d^k : C_{\alpha,\beta}^k(J, V) \rightarrow C_{\alpha,\beta}^{k+1}(J, V), \quad (k \in \{1; 2\}) :$$

- The 2-cocycles space is defined as  $Z^2(J, V) = \ker d^2$ .
- The 2-coboundary space is defined as  $B^2(J, V) = \text{Im } d^1$ .
- The  $2^{\text{th}}$  cohomology space is the quotient  $H^2(\mathcal{G}) = Z^2(J, V)/B^2(J, V)$ .

## 1.2 The trivial representation of Jacobi-Jordan algebras

Let  $[\cdot, \cdot]_V : J \times V \rightarrow \mathbb{R}$  be the zero map. Obviously,  $[\cdot, \cdot]_V$  is a representation of the Jacobi-Jordan algebra  $J$  respect to any  $\beta \in \text{End}(V)$ . We call this representation the trivial representation of Hom-Jacobi-Jordan algebra  $J$ . The corresponding coboundary operators  $d_T^k : S^k(J, V) \rightarrow S^{k+1}(J, V)$  is given by

$$d_T^k(f)(x_1, \dots, x_{k+1}) = \sum_{1 \leq i < j \leq k+1} f(\alpha^{k-1}(x_1), \dots, [x_i, x_j], \dots, \widehat{x}_j, \dots, \alpha^{k-1}(x_{k+1})) \quad (1.30)$$

where  $k \in \{1, 2, 3\}$  and  $\alpha^0 = \text{Id}_J$  (i.e  $\alpha^0(x) = x$  for all  $x \in J$ ).

A straightforward calculation shows that  $d_T^2 \circ d_T^1 = 0$ . We denote by  $H^2(J, V)$  the corresponding 2nd cohomology groups.

## 1.3 The adjoint representation of Jacobi-Jordan algebras

Since  $(J, [\cdot, \cdot], \alpha)$  is a Hom-Jacobi-Jordan algebra, the Hom-Jacobi condition (1.1) may be written for  $x, y, z \in J$

$$[[x, y], \alpha(z)] = -[\alpha(x), [y, z]] - [\alpha(y), [z, x]],$$

Then the triple  $(J, [\cdot, \cdot], \alpha)$  determines a representation of the Hom-Jacobi-Jordan algebra  $J$ . We call that representation adjoint representation of the Hom-Jacobi-Jordan algebra. The corresponding 1-coboundary and 2-coboundary are defined as follows.

$$d_a^1(f)(x, y) = f([x, y]) - [f(x), y] - [x, f(y)]$$

and

$$\begin{aligned} d_a^2(f)(x, y, z) &= f([x, y], \alpha(z)) + f([x, z], \alpha(y)) + f(\alpha(x), [y, z]) \\ &\quad + [f(x, y), \alpha(z)] + [f(x, z), \alpha(y)] + [\alpha(x), f(y, z)]. \end{aligned}$$

## 1.4 The coadjoint representation of Jacobi-Jordan algebras

Let  $J^*$  be the dual vector space of  $J$ . We define the linear map  $\alpha': J^* \rightarrow J^*$  by

$$\alpha'(Z) = Z(\alpha(\cdot)) \quad (1.31)$$

and the bilinear map  $[\cdot, \cdot]': J \times J^* \rightarrow J^*$  by

$$[x, Z]' = Z([x, \cdot]).$$

Let  $Z \in J^*$ ,  $x, y \in J$ . We compute the right hand side of the identity (1.14)

$$\begin{aligned} -[\alpha(x), [y, Z]'] - [\alpha(y), [x, Z]'] &= -[\alpha(x), Z([y, \cdot])]' - [\alpha(y), Z([x, \cdot])]' \\ &= -Z([y, [\alpha(x), \cdot]]) - Z([x, [\alpha(y), \cdot]]) \end{aligned}$$

On the other hand, we have

$$[[x, y], \alpha'(Z)]' = [[x, y], Z(\alpha(\cdot))]' = Z(\alpha([x, y, \cdot]))$$

Therefore, with the above notation, we have the following proposition:

**Proposition 1.11.** *( $J^*$ ,  $[\cdot, \cdot]'$ ,  $\alpha'$ ) is a representation of  $J$  if and only if*

$$\alpha([x, y, z]) = -[y, [\alpha(x), z]] - [x, [\alpha(y), z]] \quad (1.32)$$

for all  $x, y, z \in J$ .

Associated to the representation  $\rho'$ , the coboundary operator  $d^1$  (resp.  $d^2$ ) defined in (1.23) (resp. (1.17)) are given by

$$d^{*1}(f)(x, y) = f([x, y]) - f(y)([x, \cdot]) - f(x)([y, \cdot])$$

and

$$\begin{aligned} d^{*2}g(x, y, z) &= f([x, y], \alpha(z)) + g([x, z], \alpha(y)) + g([y, z], \alpha(x)) \\ &\quad + g(x, y)([\alpha(z), \cdot]) + g(x, z)([\alpha(y), \cdot]) + g(y, z)([\alpha(x), \cdot]). \end{aligned}$$

Hence, by Theorem 1.10 we have

**Proposition 1.12.**

$$d^{*2} \circ d^{*1} = 0$$



## 1.5 The real representations

Let  $f \in C_{\alpha, \alpha'}^1(J, J^*)$  where  $\alpha'$  is defined in (1.31). For all  $x, t \in J$ , we denote

$$f(x)(t) = f(x, t).$$

We have

$$\begin{aligned} f \circ \alpha(x)(t) = \alpha' \circ f(x)(t) &\iff f(\alpha(x))(t) = f(x)(\alpha(t)) \\ &\iff f(\alpha(x), t) = f(x, \alpha(t)). \end{aligned}$$

Hence, a 2-Hom-cochain on  $J$  with values in  $\mathbb{R}$  is defined to be a bilinear form  $f$  such that

$$f(\alpha(x), t) = f(x, \alpha(t)).$$

for all  $x, t \in J$ . Denote by  $C^2(J, \mathbb{R})$  by the vector space of 2-Hom-cochains. Now, Let  $f \in S^2(J, \mathbb{R})$ . For all  $x, y, t \in J$ , we denote

$$f(x, y)(t) = f(x, y, t).$$

Then, a 3-Hom-cochain on  $J$  with values in  $\mathbb{R}$  is defined to be a trilinear form  $\gamma$  such that

$$\gamma(x, y, t) = \gamma(y, x, t).$$

for all  $x, y, t \in J$ . Denote by  $C^3(J, \mathbb{R})$  by the vector space of 3-Hom-cochains. The set of 4-Hom-cochain is given by the vector space

$$C^4(J, \mathbb{R}) = \{\gamma \in \text{Hom}(J^4, \mathbb{R}) \mid \gamma(x, y, z, \cdot) \in S^3(J, J^*)\}$$

Define  $d^2: C^2(J, \mathbb{R}) \rightarrow C^3(J, \mathbb{R})$  by

$$d_r^2 f(x, y, t) = f([x, y], t) - f(y, [x, t]) - f(x, [y, t]). \quad (1.33)$$

and define  $d_r^3: C^3(J, \mathbb{R}) \rightarrow C^4(J, \mathbb{R})$  by

$$\begin{aligned} d_r^3 \gamma(x, y, z, t) &= \gamma([x, y], \alpha(z), t) + \gamma([x, z], \alpha(y), t) + \gamma([y, z], \alpha(x), t) \\ &+ \gamma(x, y, [\alpha(z), t]) + \gamma(y, z, [\alpha(x), t]) + \gamma(x, z, [\alpha(y), t]) \end{aligned} \quad (1.34)$$

**Theorem 1.13.** *With the above notations, we have  $d_r^3 \circ d_r^2 = 0$*

*Proof.* For any  $f \in C^{k+1}(J)$  ( $k \in \{1, 2\}$ ) we denote by  $f'$  a symmetric  $k$ -linear map satisfies

$$f'(x_1, \dots, x_k)(t) = f(x_1, \dots, x_k, t).$$

We have,

$$\begin{aligned}
d^3(d_r^2 f)(x, y, z, t) &= d_r^2 f([x, y], \alpha(z), t) + d_r^2 f([x, z], \alpha(y), t) + d_r^2 f([y, z], \alpha(x), t) \\
&\quad + d_r^2 f(x, y, [\alpha(z), t]) + d_r^2 f(y, z, [\alpha(x), t]) + d_r^2 f(x, z, \delta([\alpha(y), t])) \\
&= (d_r^2 f)'([x, y], \alpha(z))(t) + (d_r^2 f)'([x, z], \alpha(y))(t) + (d_r^2 f)'([y, z], \alpha(x))(t) \\
&\quad + (d_r^2 f)'(x, y)([\alpha(z), \cdot])(t) + (d_r^2 f)'(x, z)([\alpha(y), \cdot])(t) + (d_r^2 f)'(y, z)([\alpha(x), \cdot])(t) \\
&= d^{*2}((d_r^2 f)')(x, y, z)(t)
\end{aligned}$$

and

$$\begin{aligned}
(d_r^2 f)'(x, y)(t) &= f([x, y], t) - f(y, [x, t]) - f(x, [y, t]) \\
&= f'([x, y])(t) - f'(y)([x, t]) - f'(x)([y, t]) \\
&= d^{*1}(f')(x, y)(t)
\end{aligned}$$

Hence, by Proposition 1.12, one can deduce  $d_r^3 \circ d_r^2(f)(x, y, z, t) = 0$ . ■

## 1.6 Cohomology of c-Hom-Jacobi-Jordan algebras

**Definition 1.14.** A c-Hom-Jacobi-Jordan algebra is a triple  $(J, [\cdot, \cdot], \alpha)$  consisting of a vector space  $J$ , a symmetric bilinear map (bracket)  $[\cdot, \cdot]: J \times J \rightarrow J$  and a linear map  $\alpha: J \rightarrow J$  satisfying the following aHom-Jacobi identity:

$$\alpha([\![x, y], z\!]) = -[y, [\alpha(x), z]] - [x, [\alpha(y), z]]$$

**Definition 1.15.** A representation of a aHom-Jacobi-Jordan algebra  $(J, [\cdot, \cdot], \alpha)$  on a vector space  $V$  with respect to  $\beta \in \text{End}(V)$  is a bilinear map  $[\cdot, \cdot]_V: J \times V \rightarrow V$ , such that for all  $x, y \in J, v \in V$ , the following equalities are satisfied:

$$\beta(\rho(\delta(x, y))v) = -\rho(x) \circ \rho(\alpha(y))v - \rho(y) \circ \rho(\alpha(x))v. \quad (1.35)$$

$$\beta(\rho(x)\rho(y)v) = -\rho(y)\rho(x)\beta(v) - \rho(\delta(\alpha(y), x))v. \quad (1.36)$$

Define  $d^1: C_{\alpha, \beta}^1(J, V) \rightarrow C_{\alpha, \beta}^2(J, V)$  by

$$d^1(f)(x, y) = f(\delta(x, y)) - \rho(x)f(y) - \rho(y)f(x).$$

and  $d^2: C_{\alpha, \beta}^2(J, V) \rightarrow C^3(J, V)$  by

$$\begin{aligned}
d_c^2(f)(x, y, z) &= \theta(x, \delta(\alpha(y), z)) + \theta(y, \delta(z, \alpha(x)) + \beta(\theta(z, \delta(x, y))) \quad (1.37) \\
&\quad + \rho(x)\theta(\alpha(y), z) + \rho(y)\theta(z, \alpha(y)) + \beta(\rho(z)\theta(x, y)).
\end{aligned}$$

**Theorem 1.16.** We have  $d_c^2 \circ d^1 = 0$ .

*Proof.* Similar to Proof of Theorem 1.12 (we use  $f \circ \alpha = \beta \circ f$  and the linear map  $\rho$  satisfies (1.35) and (1.36)). ■

## 2 Extensions of Hom-Jacobi-Jordan algebras

**Definition 2.1.** Let  $(J, \delta, \alpha)$ ,  $(V, \mu, \beta)$  be two Hom-Jacobi-Jordan algebras. An extension of a Hom-Lie algebra  $J$  by  $V$  is an exact sequence

$$0 \longrightarrow (V, \mu, \beta) \xrightarrow{i} (M, d, \gamma) \xrightarrow{\pi} (J, \delta, \alpha) \longrightarrow 0$$

satisfying  $\alpha_M \circ i = i \circ \beta$  and  $\alpha \circ \pi = i \circ \beta$ .

Two extensions

$$\begin{array}{ccccc} 0 & \longrightarrow & V & \xrightarrow{i} & (M, d) & \xrightarrow{\pi} & J & \longrightarrow & 0 \\ & & \downarrow id_V & & \downarrow \Phi & & \downarrow id_J & & \\ 0 & \longrightarrow & V & \xrightarrow{i'} & (M', d') & \xrightarrow{\pi'} & J & \longrightarrow & 0 \end{array}$$

are equivalent if there exist an isomorphism  $\Phi : (M, d) \rightarrow (M', d')$ , such that  $\Phi \circ i = i'$  and  $\pi' \circ \Phi = \pi$ .

**Definition 2.2.** An extension

$$0 \longrightarrow (V, \mu, \beta) \xrightarrow{i} (M, d, \gamma) \xrightarrow{\pi} (J, \delta, \alpha) \longrightarrow 0$$

is called:

- (1) trivial if there exists an ideal  $I$  complementary to  $\ker \pi$ ,
- (2) split if there exists a Hom-subalgebra  $S \subset M$  complementary to  $\ker \pi$ ,
- (3) central if the  $\ker \pi$  is contained in the center  $Z(M)$  of  $L$ . That is  $d(i(V), M) = 0$ .
- (4) abelian if  $\mu = 0$ .

Define the following standard split extension by

$$E_0 : 0 \longrightarrow (V, \beta) \xrightarrow{i_0} (J \oplus V, d, \alpha + \beta) \xrightarrow{\pi_0} (J, \delta, \alpha) \longrightarrow 0,$$

$$E'_0 : 0 \longrightarrow (V', \beta') \xrightarrow{i'_0} (J' \oplus V', d', \alpha' + \beta') \xrightarrow{\pi'_0} (J', \delta', \alpha') \longrightarrow 0,$$

where  $i_0(v) = v$ ,  $\pi_0(x + v) = x$ .  $i'_0(v') = v'$ ,  $\pi'_0(x' + v') = x'$ .

We assume that  $E_0$  and  $E'_0$  are equivalent. Then there exist an isomorphism of Hom-Jacobi-Jordan algebra

$$\Phi : (J \oplus V, d, \alpha + \beta) \rightarrow (J' \oplus V', d', \alpha' + \beta')$$

Satisfies  $\Phi \circ i_0 = i' \circ \Phi$  and  $\pi' \circ \Phi = \Phi \circ \pi_0$ .

We set  $\Phi = s + i$  and  $s = s_1 + i_1$ ,  $i = s_2 + i_2$ , where  $s: L \rightarrow L' \oplus V'$ ,  $s_1: L \rightarrow L'$ ,  $i_1: L \rightarrow V'$ ,  $i: V \rightarrow L' \oplus V'$ ,  $s_2: V \rightarrow L'$ ,  $i_2: V \rightarrow V'$ . We denote  $\Phi(v) = v'$  and  $\Phi(x) = x'$ .

We have  $i(v) = \Phi(v) = \Phi(i_0(v)) = i'(\Phi(v)) = i'(v') = v'$

and  $x' = \Phi(x) = \Phi(\pi_0(x)) = \pi'(\Phi(x)) = \pi'(s(x)) = \pi'(s_1(x) + i_1(x)) = s_1(x)$ .

Hence  $i_1(x) = s(x) - x'$ ,  $i(v) = i_2(v) = v'$ ,  $s_2 = 0$ . We have

$$\begin{aligned} & d'(\Phi(x+v), \Phi(y+w)) = d'(x'+v', y'+w') \\ & = d'(s_1(x) + i_1(x) + i(v), s_1(y) + i_1(y) + i(w)) \\ = & \delta'(s_1(x), s_1(y)) + \theta'(s_1(x), s_1(y)) + \lambda'_l(s_1(x), i_1(y) + i(w)) + \lambda'_r(i_1(x) + i(v), s_1(y)) \\ & = \delta'(s_1(x), s_1(y)) + \theta'(s_1(x), s_1(y)) \\ & + \rho'(s_1(x))i_1(y) + \rho'(s_1(x))i(w) + \rho'(s_1(y))i_1(x) + \rho'(s_1(y))i(v) \end{aligned}$$

Moreover,

$$\begin{aligned} & d'(\Phi(x+v), \Phi(y+w)) \\ & = \Phi(d(x+v, y+w)) \\ & = \Phi(\delta(x, y) + \theta(x, y)) + \lambda_l(x, w) + \lambda_r(v, y) + \mu(v, w) \\ & = s_1 \circ \delta(x, y) + i_1 \circ \delta(x, y) + i_2 \circ \theta(x, y) + i_2 \circ \rho(x)w + i_2 \circ \rho(y)v. \end{aligned}$$

So,

$$s_1 \circ \delta(x, y) = \delta'(s_1(x), s_1(y)) \quad (2.1)$$

$$\begin{aligned} i_1 \circ \delta(x, y) + i_2 \circ \theta(x, y) &= \theta'(s_1(x), s_1(y)) + \rho'(s_1(x))i_1(y) \\ &+ \rho'(s_1(y))i_1(x) \end{aligned} \quad (2.2)$$

$$i \circ \rho(x)w = \rho'(s_1(x))i(w) \quad (2.3)$$

$$i \circ \rho(y)v = \rho'(s_1(y))i(v) \quad (2.4)$$

Define  $h: L' \rightarrow V'$  by  $h(s_1(x)) = i_1(x)$  then

$$d^1(h)(s_1(x), s_1(y)) = -h(\delta'(s_1(x), s_1(y))) + \rho'(s_1(x))h(s_1(y)) + \rho'(s_1(y))h(s_1(x)).$$

Hence, by (2.2), we obtain

$$i_2 \circ \theta(x, y) = \theta'(s_1(x), s_1(y)) + d^1(h)(s_1(x), s_1(y)). \quad (2.5)$$

**Lemma 2.3.** *Let two equivalent standard split abelian extension*

$$\begin{array}{ccccccc} 0 & \longrightarrow & (V, \mu, \beta) & \xrightarrow{i_0} & (J \oplus V, d, \alpha + \beta) & \xrightarrow{\pi_0} & (J, \delta, \alpha) \longrightarrow 0 \\ & & \varphi \downarrow & & \Phi \downarrow & & s \downarrow \\ 0 & \longrightarrow & (V', \mu', \beta') & \xrightarrow{i'_0} & (J \oplus V, d, \alpha + \beta) & \xrightarrow{\pi'_0} & (L', \delta', \alpha', \beta') \longrightarrow 0 \end{array}$$

where  $d = \delta + \rho + \theta$  and  $d' = \delta' + \rho' + \theta'$ . Then,  $i \circ \theta$  is a 2-cocycle of  $L'$  on  $V'$  and it is equivalent to  $\theta'$  (i.e.  $\overline{i \circ \theta} = \bar{\theta}'$ ).

*Proof.* By (2.5), we have  $i_2 \circ \theta(x, y) - \theta'(s_1(x), s_1(y)) \in B^2(J', V')$ . Hence  $\overline{i \circ \theta} = \bar{\theta}'$ .  $\blacksquare$

**Lemma 2.4.** *If  $\theta$  and  $\theta'$  are two equivalent 2 cocycle of  $J$  on  $V$ . Then, the extensions  $(J \oplus V, d, \alpha + \beta)$  and  $(J \oplus V, d', \alpha + \beta)$  are equivalent.*

*Proof.* Let  $\theta$  and  $\theta'$  are two equivalent 2 cocycle of  $J$  on  $V$ . Then, there exist  $h \in C^1(J, V)$  satisfies  $\theta' = \theta + d^1(h)$ . Define the linear map  $\Phi: J \oplus V \rightarrow J \oplus V$  by  $\Phi(x + v) = x - h(x) + v$ . Clearly,  $\Phi$  is an isomorphism. Moreover, we have

$$\begin{aligned} d'(\Phi(x + v), \Phi(y + w)) &= d'(x - h(x) + v, y - h(y) + w) \\ &= \delta(x, y) + \rho(x)w + \rho(y)v - \rho(x)h(y) - \rho(y)h(x) + \theta'(x, y) \\ &= \delta(x, y) + \rho(x)w + \rho(y)v - \rho(x)h(y) - \rho(y)h(x) + \theta(x, y) + d^1(h)(x, y) \\ &= \delta(x, y) + \rho(x)w + \rho(y)v + \theta(x, y) - h(\delta(x, y)) \\ &= \Phi(d(x + v, y + w)). \end{aligned}$$

Hence,  $\Phi: J \oplus V \rightarrow J \oplus V$  is an isomorphism of Hom-Jacobi-Jordan algebra. Therefore, the following diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & (V, \mu, \beta) & \xrightarrow{i_0} & (J \oplus V, d, \alpha + \beta) & \xrightarrow{\pi_0} & (J, \delta, \alpha) \longrightarrow 0 \\ & & \text{id}_V \downarrow & & \Phi \downarrow & & \text{id}_J \downarrow \\ 0 & \longrightarrow & (V, \mu, \beta) & \xrightarrow{i_0} & (J \oplus V, d', \alpha + \beta) & \xrightarrow{\pi_0} & (J, \delta, \alpha) \longrightarrow 0 \end{array}$$

gives, the extensions  $(J \oplus V, d, \alpha + \beta)$  and  $(J \oplus V, d', \alpha)$  are equivalent.  $\blacksquare$

**Theorem 2.5.** *The set  $\text{Ext}(J, V)$  of equivalence classes of split extensions of  $(J, \delta, \alpha)$  by an abelian Hom-Jordan-Jacobi algebra  $(V, \beta_V)$  is one-to-one correspondence with  $Z^2(J, V)/B^2(J, V)$ , that is*

$$\text{Ext}(J, V) \cong Z^2(J, V)/B^2(J, V).$$

## 2.1 Central extensions of two dimensional Hom-Jacobi-Jordan Algebras

Let  $(J, \delta, \alpha)$  be a Hom-Jacobi-Jordan algebra, and  $V$  a vector space. Any 2-cocycle  $\theta \in Z^2(J, V)$ , where  $V$  is viewed as a trivial  $J$ -module, leads to a central extension

$$0 \longrightarrow V \xrightarrow{i} M_\theta \xrightarrow{\pi} J \longrightarrow 0$$

of  $J$  by  $V$  as follow: As a vector space  $M_\theta = J \oplus V$ , the bracket  $d$  on  $M_\theta$  is given by

$$d(x + v, y + w) = \delta(x, y) + \theta(x, y) \text{ for } x, y \in J \text{ and } v, w \in V.$$

The isomorphism class of this extension depends only on the class of  $\theta$  in  $H^2(J, V)$ .

Suppose that  $V$  is finite-dimensional with ordered basis  $(v_1, \dots, v_m)$ . The 2-cocycle condition (1.16) translate to the following system

$$\sum_{1 \leq r, l \leq n} a_{li} C_{jk}^r m_{lj}^s + a_{lj} C_{ki}^r m_{lk}^s + a_{lk} C_{ij}^r m_{li}^s = 0 \text{ for } i, j, k \in \{1, \dots, n\}, s \in \{1, \dots, m\}. \quad (2.6)$$

where  $\theta(e_i, e_j) = \sum_{s=1}^m m_{ij}^s v_s$ . Since  $\theta$  is symmetric, we have

$$m_{ij}^s = m_{ji}^s \text{ for } i, j \in \{1, \dots, n\}. \quad (2.7)$$

The 2-cobord condition (1.18) translate to the following system

$$\sum_{1 \leq r \leq n} b_{sr} C_{ij}^r = m_{ij}^s \text{ for } i, j \in \{1, \dots, n\}, s \in \{1, \dots, m\}. \quad (2.8)$$

where  $f(e_r) = \sum_{s=1}^m b_{sr} v_s$ .

Now we assume that  $m = 2$ . We solve the system formed by (2.6) and (2.7) we obtain the space of 2-cocycle of a 2-dimensional Hom-Jordan-Jacobi algebra  $J_i^j(a, b, c)$ . We solve the system (2.8) we obtain the space of 2-cobord of a Hom-Jordan-Jacobi algebra  $J_i^j(a, b, c)$ . Then, we obtain the following table:

- $J_1^1(0, b, 0)$ :  $[e_1, e_1] = e_2, [e_1, e_2] = e_2$   
 $\alpha(e_1) = 0e_1, \alpha(e_2) = be_2$   
 $Z^2(J_1^1(0, b, 0))$ :  $\theta(e_1, e_1) = a_1v_1 + a_2v_2, \theta(e_1, e_2) = b_1v_1 + b_2v_2,$   
 $B^2(J_1^1(0, b, 0))$ :  $\varphi(e_1, e_1) = d_1v_1 + d_2v_2, \varphi(e_1, e_2) = d_1v_1 + d_2v_2,$   
 $Ext(J_1^1(0, b, 0))$ :  $d(e_1, e_1) = e_2, d(e_1, e_2) = e_2 + b_1v_1 + b_2v_2$
- $J_2^1(0, b, 0)$ :  $[e_1, e_1] = e_2, \alpha(e_1) = 0e_1, \alpha(e_2) = be_2$   
 $Z^2$ :  $\theta(e_1, e_1) = a_1v_1 + a_2v_2, \theta(e_1, e_2) = b_1v_1 + b_2v_2,$   
 $B^2$ :  $\varphi(e_1, e_1) = d_1v_1 + d_2v_2$   
 $Ext(J_2^1(0, b, 0))$ :  $d(e_1, e_1) = e_2, d(e_1, e_2) = b_1v_1 + b_2v_2$

- $J_3^1(0, b, 0)$ :  $[e_1, e_1] = e_1, [e_2, e_2] = e_1, \alpha(e_1) = 0e_1, \alpha(e_2) = be_2$   
 $Z^2: \theta(e_1, e_1) = a_1v_1 + a_2v_2, \theta(e_2, e_2) = c_1v_1 + c_2v_2,$   
 $B^2: \varphi(e_1, e_1) = d_1v_1 + d_2v_2, \varphi(e_2, e_2) = d_1v_1 + d_2v_2$   
 $Ext(J_3^1(0, b, 0)): d(e_1, e_1) = e_1, d(e_2, e_2) = e_1 + c_1v_1 + c_2v_2$
- $J_4^1(0, b, 0)$ :  $[e_1, e_1] = e_1, \alpha(e_1) = 0e_1, \alpha(e_2) = be_2$   
 $Z^2: \theta(e_1, e_1) = a_1v_1 + a_2v_2, \theta(e_2, e_2) = c_1v_1 + c_2v_2,$   
 $B^2: \varphi(e_1, e_1) = d_1v_1 + d_2v_2$   
 $Ext(J_4^1(0, b, 0)): d(e_1, e_1) = e_1, d(e_2, e_2) = c_1v_1 + c_2v_2$
- $J_5^1(0, b, -bc)$ :  $[e_1, e_1] = e_2, [e_1, e_2] = e_2, [e_2, e_2] = ce_2$   
 $\alpha(e_1) = 0e_1, \alpha(e_2) = -bce_1 + be_2$   
 $Z^2: \theta(e_1, e_1) = a_1v_1 + a_2v_2, \theta(e_1, e_2) = b_1v_1 + b_2v_2, \theta(e_2, e_2) = c(b_1v_1 + b_2v_2),$   
 $B^2(J_5^1(0, b, -bc))$ : trivial  
 $Ext(J_5^1(0, b, -bc)): d(e_1, e_1) = e_2 + a_1v_1 + a_2v_2, d(e_1, e_2) = e_2 + b_1v_1 + b_2v_2,$   
 $d(e_2, e_2) = ce_2 + c(b_1v_1 + b_2v_2),$
- $J_6^1(0, b, 0)$ :  $[e_1, e_1] = e_1 + e_2, \alpha(e_1) = 0e_1, \alpha(e_2) = be_2$   
 $Z^2: \theta(e_1, e_1) = a_1v_1 + a_2ev_2, \theta(e_1, e_2) = b_1v_1 + b_2v_2, \theta(e_2, e_2) = b_1v_1 + b_2v_2,$   
 $B^2: \varphi(e_1, e_1) = d_1v_1 + d_2v_2,$   
 $Ext(J_6^1(0, b, 0)): d(e_1, e_1) = e_1 + e_2, d(e_1, e_2) = b_1v_1 + b_2v_2, d(e_2, e_2) =$   
 $b_1v_1 + b_2v_2.$
- $J_7^1(0, b, 0)$ :  $[e_2, e_2] = e_1$   
 $\alpha(e_1) = 0e_1, \alpha(e_2) = be_2$   
 $Z^2: \theta(e_1, e_1) = a_1v_1 + a_2v_2, \theta(e_2, e_2) = c_1v_1 + c_2v_2,$   
 $B^2: \varphi(e_2, e_2) = f_1v_1 + f_2v_2$   
 $Ext(J_7^1(0, b, 0)): d(e_2, e_2) = e_1.$
- $J_8^1(0, b, 0)$ :  $[e_1, e_2] = e_2, \alpha(e_1) = 0e_1, \alpha(e_2) = be_2$   
 $Z^2: \theta(e_1, e_1) = a_1e_1 + a_2e_2, \theta(e_1, e_2) = b_1e_1 + b_2e_2,,$   
 $B^2: \varphi(e_1, e_2) = g_1v_1 + g_2v_2,$   
 $Ext(J_8^1(0, b, 0)): d(e_1, e_1) = a_1v_1 + a_2v_2, d(e_1, e_2) = e_2$
- $J_9^1(0, b, -b)$ :  $[e_1, e_2] = e_2, [e_2, e_2] = e_2$   
 $\alpha(e_1) = 0e_1, \alpha(e_2) = -be_1 + be_2$   
 $Z^2: \theta(e_1, e_1) = a_1v_1 + a_2v_2, \theta(e_1, e_2) = b_1v_1 + b_2v_2, \theta(e_2, e_2) = b_1v_1 + b_2v_2,$   
 $B^2: \varphi(e_1, e_2) = g_1v_1 + g_2v_2, \varphi(e_2, e_2) = g_1v_1 + g_2v_2$   
 $Ext(J_9^1(0, b, -b)): d(e_1, e_1) = a_1v_1 + a_2v_2, d(e_1, e_2) = e_2, d(e_2, e_2) = e_2.$
- $J_{10}^1(a, b, 0)$ :  $[e_1, e_1] = e_2$   
 $\alpha(e_1) = ae_1, \alpha(e_2) = be_2$   
 $Z^2: \theta(e_1, e_1) = a_1v_1 + a_2v_2,$

$$B^2: \varphi(e_1, e_1) = d_1v_1 + d_2v_2$$

$$Ext(J_{10}^1(a, b, 0)): d(e_1, e_1) = e_2$$

- $J_{11}^1(a, b, 1): [e_1, e_1] = e_2, \alpha(e_1) = ae_1, \alpha(e_2) = e_1 + be_2$   
 $Z^2: \theta(e_1, e_1) = a_1v_1 + a_2v_2,$   
 $B^2: \varphi(e_1, e_1) = d_1v_1 + d_2v_2,$   
 $Ext(J_{11}^1(a, b, 1)): d(e_1, e_1) = e_2.$
- $J_{12}^2(0, 0, 1): [e_1, e_1] = e_2$   
 $\alpha(e_1) = 0e_1, \alpha(e_2) = e_1$   
 $Z^2: \theta(e_1, e_1) = a_1v_1 + a_2v_2, \theta(e_2, e_2) = c_1v_1 + c_2v_2,$   
 $B^2: \varphi(e_1, e_1) = d_1v_1 + d_2v_2,$   
 $Ext(J_{12}^2(0, 0, 1)): d(e_1, e_1) = e_2, d(e_2, e_2) = c_1v_1 + c_2v_2$
- $J_{13}^2(0, 0, c): [e_1, e_1] = e_2, [e_2, e_2] = e_2$   
 $\alpha(e_1) = 0e_1, \alpha(e_2) = ce_1$   
 $Z^2(J_{13}^2(0, 0, c)): \theta(e_1, e_1) = a_1v_1 + a_2v_2, \theta(e_2, e_2) = c_1v_1 + c_2v_2,$   
 $B^2(J_{13}^2(0, 0, c)): \varphi(e_1, e_1) = d_1v_1 + d_2v_2, \varphi(e_2, e_2) = d_1v_1 + d_2v_2,$   
 $Ext(J_{13}^2(0, 0, c)): d(e_1, e_1) = e_2, d(e_2, e_2) = e_2 + c_1v_1 + c_2v_2$
- $J_{14}^2(0, 0, 1): [e_2, e_2] = e_1$   
 $\alpha(e_1) = 0e_1, \alpha(e_2) = e_1$   
 $Z^2(J_{14}^2(0, 0, 1)): \theta(e_1, e_2) = b_1v_1 + b_2v_2, \theta(e_2, e_2) = c_1v_1 + c_2v_2,$   
 $B^2(J_{14}^2(0, 0, 1)): \varphi(e_2, e_2) = f_1v_1 + f_2v_2, ,$   
 $Ext(J_{14}^2(0, 0, 1)): d(e_1, e_2) = b_1v_1 + b_2v_2, d(e_2, e_2) = e_1$
- $J_{15}^2(0, 0, c): [e_1, e_2] = e_1, [e_2, e_2] = e_1$   
 $\alpha(e_1) = 0e_1, \alpha(e_2) = ce_1$   
 $Z^2(J_{15}^2(0, 0, c)): \theta(e_1, e_2) = b_1v_1 + b_2v_2, \theta(e_2, e_2) = c_1v_1 + c_2v_2,$   
 $B^2(J_{15}^2(0, 0, c)): \varphi(e_1, e_2) = g_1v_1 + g_2v_2, \varphi(e_2, e_2) = g_1v_1 + g_2v_2,$   
 $Ext(J_{15}^2(0, 0, c)): d(e_1, e_2) = e_1, d(e_2, e_2) = e_1 + c_1v_1 + c_2v_2,$
- $J_{16}^2(0, 0, 1): [e_2, e_2] = e_2$   
 $\alpha(e_1) = 0e_1, \alpha(e_2) = e_1$   
 $Z^2(J_{16}^2(0, 0, 1)): \theta(e_1, e_1) = a_1v_1 + a_2v_2, \theta(e_2, e_2) = c_1v_1 + c_2v_2,$   
 $B^2(J_{16}^2(0, 0, 1)): \varphi(e_2, e_2) = h_1v_1 + h_2v_2,$   
 $Ext(J_{16}^2(0, 0, 1)): d(e_1, e_1) = a_1v_1 + a_2v_2, d(e_2, e_2) = e_2,$
- $J_{17}^2(0, 0, 1): [e_1, e_2] = e_1$   
 $\alpha(e_1) = 0e_1, \alpha(e_2) = e_1$   
 $Z^2(J_{17}^2(0, 0, 1)): \theta(e_1, e_2) = b_1v_1 + b_2v_2, \theta(e_2, e_2) = c_1v_1 + c_2v_2,$   
 $B^2(J_{17}^2(0, 0, 1)): \varphi(e_1, e_2) = g_1v_1 + g_2v_2,$   
 $Ext(J_{17}^2(0, 0, 1)): d(e_1, e_2) = e_1, d(e_2, e_2) = c_1v_1 + c_2v_2$



- $J_{18}^2(a, a, 1)$ :  $[e_1, e_1] = e_2$   
 $\alpha(e_1) = ae_1, \alpha(e_2) = e_1 + ae_2$   
 $Z^2(J_{18}^2(a, a, 1))$ :  $\theta(e_1, e_1) = a_1v_1 + a_2v_2$  ,  
 $B^2(J_{18}^2(a, a, 1))$ :  $B^2$ :  $\varphi(e_1, e_1) = d_1v_1 + d_2ev_2$ ,  
 $Ext(J_{18}^2(a, a, 1))$ :  $d(e_1, e_1) = e_2$
- $J_{19}^2(a, a, 1)$ :  $[e_2, e_2] = e_1$   
 $\alpha(e_1) = ae_1, \alpha(e_2) = e_1 + ae_2$   
 $Z^2(J_{19}^2(a, a, 1))$ :  $\theta(e_2, e_2) = c_1v_1 + c_2v_2$  ,  
 $B^2(J_{19}^2(a, a, 1))$ :  $\varphi(e_2, e_2) = h_1v_1 + h_2v_2$ ,  
 $Ext(J_{19}^2(a, a, 1))$ :  $d(e_2, e_2) = e_1$  .

### 3 Metric Hom-Jacobi-Jordan algebra

**Definition 3.1.** A metric Hom-Jacobi-Jordan algebra is a 4-tuple  $(J, [\cdot, \cdot], \alpha, B)$ , where  $J$  is a vector space,  $B, [\cdot, \cdot]: J \times J \rightarrow J$  are two symmetric bilinear maps and  $\alpha: J \rightarrow J$  be a linear map satisfying the following identities:

$$[\alpha(x), [y, z]] + [\alpha(z), [x, y]] + [\alpha(y), [z, x]] = 0, \quad (3.1)$$

$$\alpha([ [x, y], z ]) + [y, [\alpha(x), z]] + [x, [\alpha(y), z]] = 0 \quad (3.2)$$

$$B(x, [y, z]) = B([x, y], z) \quad (3.3)$$

$$B(\alpha(x), y) = B(x, \alpha(y)) \quad (3.4)$$

for any  $x, y, z \in J$ .

**Definition 3.2.** Let  $(\mathfrak{J}, B)$  be a metric Hom-Jacobi-Jordan algebra.

1. An ideal  $I$  of  $\mathfrak{J}$  is said to be nondegenerate if  $B|_{I \times I}$  is nondegenerate.
2. Let  $I$  be an ideal of  $\mathfrak{J}$ . The orthogonal  $I^\perp$  of  $I$ , with respect to  $B$ , is defined by

$$I^\perp = \{x \in \mathfrak{J} \mid B(x, y) = 0 \forall y \in I\}.$$

3. An ideal  $I$  is isotropic if  $I \subset I^\perp$ .

**Proposition 3.3.** Let  $J$  be a Hom-Jacobi-Jordan algebra and  $B: J \times J \rightarrow \mathbb{R}$  a nondegenerate bilinear form satisfies  $B(\alpha(x), y) = B(x, \alpha(y))$ . Define a trilinear form  $\gamma: J^3 \rightarrow \mathbb{R}$  by  $\gamma(x, y, z) = B([x, y], z)$ . Then,  $(J, B)$  is a metric Hom-Jacobi-Jordan if and only if the trilinear form  $\gamma$  is symmetric and it satisfies  $d_r^3 \gamma = 0$  where  $d_r^3$  is defined in (1.34).

*Proof.* We assume that  $(J, B)$  is a metric Hom-Jacobi-Jordan. Then

$$\begin{aligned}\gamma(x, y, z) &= B([x, y], z) = B([y, x], z) = \gamma(y, x, z) \\ \gamma(x, y, z) &= B([x, y], z) = B(z, [x, y]) = B(z, [y, x]) = \gamma(z, y, x).\end{aligned}$$

Similar we verify that  $\gamma(x, y, z) = \gamma(x, z, y)$ ,  $\gamma(x, y, z) = \gamma(y, z, x)$  and  $\gamma(x, y, z) = \gamma(z, x, y)$ . Hence  $\gamma \in C^3(J, \mathbb{R})$ . Moreover, we have

$$\begin{aligned}d_r^3\gamma(x, y, z, t) &= \gamma([x, y], \alpha(z), t) + \gamma([x, z], \alpha(y), t) + \gamma([y, z], \alpha(x), t) \\ &\quad + \gamma(x, y, [\alpha(z), t]) + \gamma(y, z, [\alpha(x), t]) + \gamma(x, z, [\alpha(y), t]) \\ &= B([[x, y], \alpha(z)], t) + B([[x, z], \alpha(y)], t) + B([\alpha(x), [y, z]], t)\end{aligned}\tag{3.5}$$

$$+ B([x, y], [\alpha(z), t]) + B([y, z], [\alpha(x), t]) + B([x, z], [\alpha(y), t])\tag{3.6}$$

By the Hom-Jacobi identity, we obtain that Eq.(3.5)=0.

By the fact that invariance of  $B$  with respect to the bracket  $[\cdot, \cdot]$ , i.e  $B(x, [y, z]) = B([x, y], z)$ , we get

$$(3.6) = B(x, [y, [\alpha(z), t]]) + B([[y, z], t], \alpha(x)) + B(x, [z, [\alpha(y), t]])$$

By (3.4), we have  $B([[y, z], t], \alpha(x)) = B(\alpha([[y, z], t]), x)$ . Hence,

$$(3.6) = B(x, [y, [\alpha(z), t]]) + B(x, \alpha([[y, z], t])) + B(x, [z, [\alpha(y), t]])\tag{3.7}$$

Therefore, By (3.2), we have (3.6) = 0.

However,  $d_r^3\gamma(x, y, z, t) = 0$ .

Conversely, we assume that  $\gamma \in S^3(\mathfrak{J}, \mathbb{R})$  and  $d_r^3\gamma = 0$ . Then, we have

$$\gamma(x, y, z) = \gamma(y, z, x) \implies B([x, y], z) = B([y, z], x) \implies B([x, y], z) = B(x, [y, z]).$$

So, Eq.(3.3) holds.

Now, we prove the symmetric condition for  $[\cdot, \cdot]$ :

By  $\gamma \in S^3(J, \mathbb{R})$ , we have  $\gamma(x, y, z) = \gamma(y, x, z)$ , Hence  $B([x, y], z) = B([y, x], z)$ .

Since  $B$  is nondegenerate one can deduce  $[x, y] = [y, x]$ .

We have

$$(3.6) = B([[x, y], \alpha(z)], t) + B([[y, z], \alpha(x)], t) + B([[x, z], \alpha(y)], t).$$

Hence  $d^3\gamma(x, y, z, t) = 2\left(B([[x, y], \alpha(z)], t) + B([[y, z], \alpha(x)], t) + B([[x, z], \alpha(y)], t)\right)$ .

Since  $d^3\gamma = 0$  and  $B$  is nondegenerate, we obtain the Eq. (3.1).

Then, (3.5) = 0. So, by (3.7), we obtain the condition (3.2). ■

## 4 Cohomology of metric Hom-Jacobi-Jordan-algebra

Define a bilinear multiplication on  $C^p(J, \mathfrak{a}) \times C^q(J, \mathfrak{a})$  by

$$B(f \wedge g)(x_1, \dots, x_{p+q}) = \sum_{\sigma \in Sh(p,q)} B(f(x_{\sigma(1)}, \dots, x_{\sigma(p)}), g(x_{\sigma(p+1)}, \dots, x_{\sigma(p+q)})) \quad (4.1)$$

where  $Sh(p, q)$  are the permutations in  $\mathfrak{S}_{p+q}$  which are increasing on the first  $p$  and the last  $q$  elements.

**Proposition 4.1.** *Let  $f \in S^2(J, \mathfrak{a})$  and  $g \in S^1(J, \mathfrak{a})$ . We have*

$$d_r^3 B(f \wedge g)(x, y, z, t) = B(d_T^2 f, g \circ \alpha)(x, y, z, a) + B(f \circ \alpha, d_T^1 g)(x, y, z, a) \quad (4.2)$$

for any  $x, y, z, a \in J$  and  $t = \alpha(a)$ .

*Proof.* We take  $\gamma = B(f \wedge g)$ . Then, by (1.34) and (3.4), we obtain

$$\begin{aligned} d_r^3 \gamma(x, y, z, t) &= \gamma([x, y], \alpha(z), \alpha(a)) + \gamma([x, z], \alpha(y), \alpha(a)) + \gamma([y, z], \alpha(x), \alpha(a)) \\ &\quad + \gamma(x, y, [\alpha(z), \alpha(a)]) + \gamma(y, z, [\alpha(x), \alpha(a)]) + \gamma(x, z, [\alpha(y), \alpha(a)]) \\ &= \gamma([x, y], \alpha(z), \alpha(a)) + \gamma([x, z], \alpha(y), \alpha(a)) + \gamma([y, z], \alpha(x), \alpha(a)) \\ &\quad + \gamma(\alpha(x), \alpha(y), [z, a]) + \gamma(\alpha(y), \alpha(z), [x, a]) + \gamma(\alpha(x), \alpha(z), [y, a]). \end{aligned}$$

Hence, we replace  $\gamma$  by  $B(f \wedge g)$ , we obtain (4.2). ■

Let  $M = J \oplus V$  be a Hom-Jacobi-Jordan algebra with a structure  $d$  such that  $V$  is an abelian ideal of  $M$ . Then by Proposition 1.8,  $d = \delta + \lambda + \theta$ , where  $(J, \delta)$  is a Hom-Jacobi-Jordan algebra.  $\lambda(x, v) = \lambda(v, x) = \rho(x)v$ ,  $\rho$  is a representation of  $J$  on  $V$  and  $\theta$  is a 2-cocycle of  $J$  on  $V$ . Let  $N = M \oplus J^*$  and  $d': N^2 \rightarrow N$  a bilinear symmetric map such that  $d'(a, b) = d(a, b)$  for all  $a, b \in \mathfrak{m}$ . We assume that  $d'(Z, v) = 0$  and we use the representation  $\rho': \mathfrak{m} \rightarrow \text{End}(J^*)$  defined by

$$\rho'(x + v)(Z) = Z(\delta(x, \cdot)). \quad (4.3)$$

Then there exists bilinear maps  $\gamma': \mathfrak{m} \times \mathfrak{m} \rightarrow J^*$  such that

$$d'(x, y) = \delta(x, y) + \theta(x, y) + \gamma'(x, y); \quad (4.4)$$

$$d'(x, v) = \rho(x)v + \gamma'(x, v); \quad (4.5)$$

$$d'(v, w) = \gamma'(v, w); \quad (4.6)$$

$$d'(Z, x) = Z(\delta(x, \cdot)); \quad (4.7)$$

$$d'(Z_1, v + Z_2) = 0. \quad (4.8)$$

Let  $B: \mathfrak{n}^2 \rightarrow \mathbb{R}$  be a nondegenerate bilinear form. then there exists bilinear forms  $B_J: \mathfrak{J}^2 \rightarrow \mathbb{R}$ ,  $B_\rho: J \times \mathfrak{a} \rightarrow \mathbb{R}$ ,  $B_\alpha: \mathfrak{J}^2 \rightarrow \mathbb{R}$ ,  $B_{J^*}: \mathfrak{n} \times J^* \rightarrow \mathbb{R}$  such that  $B = B_J + B_\rho + B_\alpha + B_{J^*}$ . We assume that  $B_J = 0$ . We assume that  $B_{J^*}(x + v + Z', Z) = Z(x)$ . Define a trilinear form  $\gamma$  on  $\mathfrak{n}$  by  $\gamma_{\mathfrak{n}}(a, b, c) = B(d'(a, b), c)$  for all  $a, b, c \in \mathfrak{n}$ . We assume that

$$B_\rho(\theta(x, y), z) = B(x, \theta(y, z)) \quad (4.9)$$

for all  $x, y, z \in J$ .

We want to find conditions on  $\gamma'$  so that  $M$  is a metric Hom-Jacobi-Jordan algebra. By Proposition 3.3,  $\gamma' \in S^3(M, \mathbb{R})$  and  $d_r^3 = 0$ .

Let  $\gamma(x, y, z) = \gamma'(x, y)(z)$ . Then, by (4.4), we obtain

$$B(d'(x, y), z) = B(x, d'(y, z)) \iff \gamma \in S^3(J, \mathbb{R}).$$

By (4.5), (4.4) and  $\gamma'(x, y)(v) = 0$  (because  $\gamma'(x, y) \in J^*$ ), we obtain

$$\begin{aligned} B(d'(y, x), v) &= B(y, d'(x, v)) \\ \iff \gamma'(x, v)(y) &= B_\rho(\delta(y, x), v) + B_\alpha(\theta(y, x), v) - B_\rho(\rho(y)v, x) \end{aligned}$$

Hence

$$\gamma'(x, v) = B_\rho(\delta(\cdot, x), v) + B_\alpha(\theta(\cdot, x), v) - B_\rho(\rho(\cdot)v, x). \quad (4.10)$$

By (4.5), (4.6), we have

$$B(d'(x, v), w) = B(x, d'(v, w)) \iff \gamma'(v, w)(x) = B_\alpha(\rho(x)v, w).$$

Hence

$$\gamma'(v, w) = B_\alpha(\rho(\cdot)v, w). \quad (4.11)$$

Therefore, we obtain

$$\begin{aligned} d'(x, y) &= \delta(x, y) + \theta(x, y) + \gamma(x, y, \cdot); \\ d'(x, v) &= \rho(x)v + B_\rho(\delta(\cdot, x), v) + B_\alpha(\theta(\cdot, x), v) - B_\rho(\rho(\cdot)v, x); \\ d'(v, w) &= B_\alpha(\rho(\cdot)v, w); \\ d'(Z, x) &= Z(\delta(x, \cdot)); \\ d'(Z_1, v + Z_2) &= 0. \end{aligned} \quad (4.12)$$

We define  $\gamma_{\mathfrak{n}}: \mathfrak{n}^3 \rightarrow \mathbb{R}$  by

$$\gamma_{\mathfrak{n}}(a, b, c) = B_{\mathfrak{n}}(d'(a, b), c)$$

for all  $a, b, c \in \mathfrak{n}$ .

If  $\gamma_{\mathfrak{n}} \in S^3(\mathfrak{n}, \mathbb{R})$ , we have  $\gamma_{\mathfrak{n}}(x, y, v) = \gamma_{\mathfrak{n}}(x, v, y)$  and  $\gamma_{\mathfrak{n}}(x, v, y) = \gamma_{\mathfrak{n}}(y, v, x)$ . Therefore,

$$\gamma'(y, v)(x) - \gamma'(x, v)(y) = -B_{\rho}(\rho(x)v, y) + B_{\rho}(\rho(y)v, x) \quad (4.13)$$

and

$$\gamma'(x, v)(y) - \gamma'(y, v)(x) = -B(\rho(x)v, y) + B(\rho(y)v, x) \quad (4.14)$$

One can deduce  $0 = (4.14) + (4.13) = 2\left(B(\rho(y)v, x) - B(\rho(x)v, y)\right)$ . Hence

$$B(\rho(x)v, y) = B(\rho(y)v, x)$$

and  $\gamma'(x, v)(y) = \gamma'(y, v)(x)$ .

Use  $d' \in S^2(\mathfrak{n})$ , we obtain

$$B_{\mathfrak{a}}(\rho(\cdot)v, w) = B_{\mathfrak{a}}(\rho(\cdot)w, v).$$

We deduce the following definitions of quadratic representation.

**Definition 4.2.** Let  $J$  be a Hom-Jacobi-Jordan algebra.  $(\rho, \mathfrak{a}, B_{\mathfrak{a}}, B_{\rho})$  is called Quadratic representation (or  $J$ -module), if  $(\mathfrak{a}, B_{\mathfrak{a}})$  is an abelian metric Hom-Jacobi-Jordan algebra,  $\rho: J \rightarrow \text{Hom}(\mathfrak{a})$  a representation of the Hom-Jacobi-Jordan algebra  $J$  on  $\mathfrak{a}$ , and  $B_{\rho}: J \times \mathfrak{a} \rightarrow \mathbb{R}$  a symmetric bilinear form which is symmetric, i.e

$$B_{\mathfrak{a}}(\rho(x)(v), w) = B_{\mathfrak{a}}(v, \rho(x)(w)) \quad (4.15)$$

$$B_{\rho}(\rho(x)(v), y) = B_{\rho}(x, \rho(y)(v)) \quad (4.16)$$

for all  $x, y \in J$  and  $v, w \in \mathfrak{a}$ .

Recall that if  $\mathfrak{n}$  is a metric Hom-Jacobi-Jordan algebra. Then  $\gamma'$  is a 2-cocycle on  $\mathfrak{m}$ . We have

$$\begin{aligned} d^2\gamma'(x, y, z)(t) &= \gamma'(d(x, y), \alpha(z))(t) + \gamma'(d(x, z), \alpha(y))(t) + \gamma'(d(y, z), \alpha(x))(t) \\ &\quad + \gamma'(x, y)(d(\alpha(z), t)) + \gamma'(y, z)(d(\alpha(x), t)) + \gamma'(x, z)(d(\alpha(y), t)). \end{aligned}$$

By  $d(x, y) = [x, y] + \theta(x, y)$ ,  $\gamma'(x, y)(v) = 0$ , (4.10) and (4.11),  $\theta \circ \alpha = \beta \circ \alpha$ ,  $d^2\theta = 0$  and  $B_{\mathfrak{a}}(\beta(v), w) = B_{\mathfrak{a}}(v, \beta(w))$  we obtain

$$d^{*2}\gamma'(x, y, z)(t) = d_c^3\gamma(x, y, z, a) + \frac{1}{2}B_{\mathfrak{a}}(\theta \wedge (\theta \circ \alpha))(x, y, z, a) + B_{\rho}((\delta \circ \alpha) \wedge \theta)(x, y, z, a) \quad (4.17)$$

where  $t = \alpha(a)$ .

Moreover, we have

$$\begin{aligned} d_m^2 \gamma'(x, y, v)(t) &= \gamma'(d(x, y), \beta(v))(t) + \gamma'(d(x, v), \alpha(y))(t) + \gamma'(d(y, v), \alpha(x))(t) \\ &\quad + \rho'(\beta(v))\gamma'(x, y)(t) + \rho'(\alpha(x))\gamma'(y, v)(t) + \rho'(\alpha(y))\gamma'(x, v)(t) \end{aligned}$$

By (4.10) and  $B_\rho = 0$ , we have  $\gamma'(x, v) = B_\alpha(\theta(\cdot, x), v)$ . Using (4.3) and (4.15) we obtain

$$d_m^2 \gamma'(x, y, v)(t) = B_\alpha(d_c^2 \theta(x, y, t), v). \quad (4.18)$$

We deduce the following definitions.

**Definition 4.3.**  $(\theta, \gamma)$  is called a quadratic 2-cochain if  $\theta \in C^2(J_c, \mathfrak{a})$  and  $\gamma \in C^2(J, J^*)$  such that the trilinear map  $\gamma'$  defined by  $\gamma'(x, y, z) = \gamma(x, y)(z)$  is symmetric. We define a linear map by  $d_Q^2: C_Q^2(\mathfrak{J}, \mathfrak{a}) \rightarrow C^3(\mathfrak{J}, \mathfrak{a})$  by

$$d_Q^2(\theta, \gamma) = \left( d_c^2 \theta, d_r^3 \gamma + \frac{1}{2} B_\alpha(\theta \wedge (\theta \circ \alpha)) \right) \quad (4.19)$$

$(\theta, \gamma)$  is called a quadratic 2-cocycle of  $\mathfrak{J}$  on  $\mathfrak{a}$  if and only if  $d_Q^2(\theta, \gamma) = 0$ . We denote  $Z_Q^2(\mathfrak{J}, \mathfrak{a})$  the space of all quadratic 2-cocycle on  $\mathfrak{a}$ .

**Proposition 4.4.**  $(\mathfrak{n}, d')$  is a metric Hom-Jacobi-Jordan algebra and  $J^*$  is an abelian ideal of  $\mathfrak{n}$  if and only if  $(\theta, \gamma)$  is a quadratic 2-cocycle.

*Proof.* We assume that  $(\mathfrak{n}, d')$  is a metric Hom-Jacobi-Jordan algebra and  $J^*$  is an abelian ideal of  $\mathfrak{n}$ . We have  $d' = d + \gamma'$ . Hence, using Theorem 1.8, we obtain  $d_m^2 \gamma' = 0$ . So;  $d_m^2 \gamma'(x, y, z)(t) = 0$  and  $d_m^2 \gamma'(x, y, v)(t) = 0$ .

Hence, by (4.17) (resp.(4.18) ),  $B_\alpha$  is nondegenerate and  $B_\rho = 0$ , we obtain

$$d_r^3 \gamma(x, y, z, t) + \frac{1}{2} B_\alpha(\theta \wedge (\theta \circ \alpha)) + B_\rho((\delta \circ \alpha) \wedge \theta)(x, y, z, a) = 0 \quad (4.20)$$

$$d_c^2 \theta(x, y, z) = 0 \quad (4.21)$$

By (4.20) and (4.21), we have  $d_Q^2(\theta, \gamma) = 0$

Conversely, we assume that  $(\theta, \gamma)$  is a quadratic 2-cocycle.

Let  $\gamma_n = B(d'(a, b), c)$ . Using Theorem 3.3, we show that  $\gamma_n$  is symmetric and  $d^3 \gamma_n = 0$ .

We have

$$\gamma_n(x, y, z) = B(d'(x, y), z) = B(\delta(x, y) + \theta(x, y) + \gamma(x, y, \cdot), z) = \gamma(x, y, z)$$

Since  $\gamma \in S^3(J)$  we have the restriction of  $\gamma_n$  to  $J^3$  is symmetric.

$$\begin{aligned}\gamma_n(x, y, v) &= B(d'(x, y), v) = B_a(\theta(x, y), v) \\ \gamma_n(x, v, y) &= B(d'(x, v), y) = B_a(\theta(x, y), v).\end{aligned}$$

Therefore, using  $d'(x, y) = d'(y, x)$  and  $d'(x, v) = d'(v, x)$ , one can deduce the restriction of  $\gamma_n$  to  $J^2 \times V$  is symmetric.

$$\begin{aligned}\gamma_n(x, v, w) &= B(d'(x, v), w) = B_a(\rho(x)w, v) \\ \gamma_n(v, w, x) &= B(d'(v, w), x) = B_a(\rho(x)v, w)\end{aligned}$$

Hence the restriction of  $\gamma_n$  to  $J \times V^2$  is symmetric.

$$\gamma_n(v, w, u) = B(d'(v, w), u) = B(B_a(\rho(\cdot)v, w), u) = 0$$

Hence the restriction of  $\gamma_n$  to  $V^3$  is symmetric.

Now we show that  $d^3\gamma_n = 0$ .

We have

$$d_r^3\gamma_n(x, y, z, t) = 2d_r^3\gamma(x, y, z, t) + B_a(\theta \wedge (\theta \circ \alpha))(x, y, z, a) = 0$$

$$\begin{aligned}\frac{1}{2}d_r^3\gamma_n(x, y, z, v) &= B_a(\theta(\delta(x, y), \alpha(z)), v) + B_a(\theta(\delta(x, z), \alpha(y)), v) + B_a(\theta(\delta(y, z), \alpha(x)), v) \\ &\quad + B(\rho(\alpha(z))\theta(x, y), v) + B(\rho(\alpha(x))\theta(y, z), v) + B(\rho(\alpha(y))\theta(x, z), v) \\ &= \frac{1}{2}B_a(d_c^2\theta(x, y, z), v) = 0.\end{aligned}$$

$$\begin{aligned}&d_r^3\gamma_n(x, y, u, v) \\ &= \gamma_n(d'(x, y), \beta(u), v) + \gamma_n(d'(x, u), \alpha(y), v) + \gamma_n(d'(y, u), \alpha(x), v) \\ &\quad + \gamma_n(d'(\alpha(x), v), y, u) + \gamma_n(d'(\alpha(y), v), x, u) + \gamma_n(d'(\beta(u), v), x, y) \\ &= B_n\left(u, \beta(\rho(\delta(x, y))v)\right) + B_a(\rho(\alpha(y))\rho(x)u, v) + B_a(\rho(\alpha(x))\rho(y)u, v) \\ &\quad + B\left(\rho(y)\rho(\alpha(x))v, u\right) + B\left(\rho(x)\rho(\alpha(y))v, u\right) + B_a(u, (\beta\rho(\delta(x, y))v)).\end{aligned}$$

Therefore by (1.35),  $d_r^3\gamma_n(x, y, u, v) = 0$ . ■

## 4.1 Quadratic 2-cobord

In this subsection, we aim, similar to Lie case, define a quadratic 2-cobord. Let  $(\theta, \gamma)$  and  $(\theta', \gamma')$  two quadratic 2-cochain such that  $d_Q^2(\theta, \gamma) = d_Q^2(\theta', \gamma')$ . Then,

$$d_c^2\theta = d_c^2\theta' \quad (4.22)$$

and

$$d_r^3\gamma + \frac{1}{2}B(\theta \wedge (\theta \circ \alpha)) = d_r^3\gamma' + \frac{1}{2}B(\theta' \wedge (\theta' \circ \alpha)) \quad (4.23)$$

(4.22) we get, there exist a 1-cochain  $\tau$  satisfies

$$\theta' = \theta + d_c^1\tau \quad (4.24)$$

Thus, using (4.23), we have

$$\begin{aligned} d_r^3\gamma &= d_r^3\gamma' + \frac{1}{2}B\left((\theta + d^1\tau) \wedge ((\theta + d^1\tau) \circ \alpha)\right) \\ &= d_r^3\gamma' + \frac{1}{2}B(\theta \wedge (\theta \circ \alpha)) + \frac{1}{2}B(\theta \wedge (d^1\tau \circ \alpha)) + \frac{1}{2}B(d^1\tau \wedge (\theta \circ \alpha)) \\ &\quad + \frac{1}{2}B(d^1\tau \wedge (d^1\tau \circ \alpha)) - \frac{1}{2}B(\theta \wedge (\theta \circ \alpha)) \\ &= d_r^3\gamma' + \frac{1}{2}B(\theta \wedge (d^1\tau \circ \alpha)) + \frac{1}{2}B(d^1\tau \wedge (\theta \circ \alpha)) \\ &\quad + \frac{1}{2}B(d^1\tau \wedge (d^1\tau \circ \alpha)). \end{aligned}$$

We have

$$\begin{aligned} B(f \wedge (g \circ \alpha))(x, y, z, t) &= B(f(x, y), g(\alpha(z), \alpha(t))) + \dots \\ &= B(f(x, y), \beta(g(z, t))) + \dots \\ &= B(\beta(f(x, y)), g(z, t)) + \dots \\ &= B(f(\alpha(x), \alpha(y)), g(z, t)) + \dots \\ &= B((f \circ \alpha) \wedge g)(x, y, z, t) \end{aligned}$$

Then,

$$\begin{aligned} d^3\gamma &= d^3\gamma' + \frac{1}{2}B((\theta \circ \alpha) \wedge d^1\tau) + \frac{1}{2}B(d^1\tau \wedge (\theta \circ \alpha)) \\ &\quad + \frac{1}{2}B(d^1\tau \wedge (d^1\tau \circ \alpha)) \\ &= d^3\gamma' + B(d^1\tau \wedge (\theta \circ \alpha)) + \frac{1}{2}B(d^1\tau \wedge (d^1\tau \circ \alpha)) \end{aligned}$$



$$\begin{aligned}
& d_Q^2(\theta, \gamma) = d_Q^2(\theta', \gamma') \\
& \iff \begin{cases} d_c^2\theta = d_c^2\theta' \\ d^3\gamma + \frac{1}{2}B(\theta \wedge \theta) = d^3\gamma' + \frac{1}{2}B(\theta' \wedge \theta') \end{cases} \\
& \iff \begin{cases} \theta' = \theta + d^1\tau \\ d^3\gamma + \frac{1}{2}B(\theta \wedge \theta) = d^3\gamma' + \frac{1}{2}B((\theta + d^1\tau) \wedge (\theta + d^1\tau)) \end{cases} \\
& \iff \begin{cases} \theta' = \theta + d_c^1\tau \\ d^3\gamma = d^3\gamma' + B(d^1\tau \wedge (\theta \circ \alpha)) + \frac{1}{2}B(d^1\tau \wedge (d^1\tau \circ \alpha)) \end{cases}
\end{aligned}$$

By Proposition 4.1 and Proposition 1.10 , we have

$$d^3B(d^1\tau \wedge \tau) = B(d^2 \circ d^1\tau \wedge (\tau \circ \alpha)) + B((d^1\tau \circ \alpha) \wedge d^1\tau) = B((d^1\tau \circ \alpha) \wedge d^1\tau) \quad (4.25)$$

From this and (4.1), we obtain

$$d^3\gamma' = d^3\gamma - \frac{1}{2}d^3B(\tau \wedge d^1\tau) - B(d^1\tau \wedge (\theta \circ \alpha)). \quad (4.26)$$

Since  $d^3B(\theta \wedge \tau) = B((d^2\theta \circ \alpha) \wedge \tau) + B(\theta \wedge (d^1\tau \circ \alpha))$ , we can deduce the following result.

$$d^3\gamma' = d^3\gamma - \frac{1}{2}d^3B(\tau \wedge d^1\tau) - d^3B(\theta \wedge \tau). \quad (4.27)$$

From this and (4.24) we obtain the following result

**Lemma 4.5.**

$$d_Q^2(\theta, \gamma) = d_Q^2(\alpha', \gamma') \iff \begin{cases} \theta' = \theta + d_c^1\tau \\ d_r^3\gamma' = d_r^3\gamma - \frac{1}{2}d_r^3B(\tau \wedge d^1\tau) - d^3B(\tau \wedge \theta) \end{cases} \quad (4.28)$$

Using the previous Lemma and Proposition 4.1, we obtain

**Proposition 4.6.** *If  $d^2\theta = 0$  we have*

$$d_Q^2(\theta, \gamma) = d_Q^2(\alpha', \gamma') \iff \begin{cases} \theta' = \theta + d^1\tau \\ \gamma' = \gamma + d_r^2\sigma - B(\tau \wedge (\theta + \frac{1}{2}d^1\tau)) \end{cases}$$

Moreover, with (4.28) we can give the following definition

**Definition 4.7.** Define a linear map  $d_Q^1: C_Q^1(\mathfrak{J}, \mathfrak{a}) \rightarrow C_Q^2(\mathfrak{J}, \mathfrak{a})$  by

$$d_Q^1(\tau, \sigma) = \left( d_c^1 \tau, d_c^2 \sigma - B(\tau \wedge \left( \theta + \frac{1}{2} d^1 \tau \right)) \right)$$

A quadratic 2-cochain  $(\theta, \gamma)$  is called 2-cobord if and only if there exist a quadratic 1-cochain  $(\tau, \sigma)$  satisfies  $d_Q^1(\tau, \sigma) = (\theta, \gamma)$ . Denote  $B_Q^2(\mathfrak{J}, \mathfrak{a})$  the space of all quadratic 2-cobord.

**Proposition 4.8.** *Any quadratic 2-cobord is a quadratic 2-cocycle.*

*Proof.* We set  $\theta = d^1 \tau$  and  $\gamma = d^2 \sigma - \frac{1}{2} B_{\mathfrak{a}}(d^1 \tau \wedge \tau)$ . Using Proposition 4.2, we have  $d^3 \gamma = -\frac{1}{2} B_{\mathfrak{a}}(d^1 \tau \wedge (d^1 \tau \circ \alpha))$ . Hence, by (4.19)

$$\begin{aligned} d_Q^2(\theta, \gamma) &= (d_c^2 \theta, d_r^3 \circ d_r^2 \sigma - \frac{1}{2} B_{\mathfrak{a}}(d^1 \tau \wedge (d^1 \tau \circ \alpha)) + \frac{1}{2} B(d^1 \tau \wedge (d^1 \tau \circ \alpha))) \\ &= (0, 0). \end{aligned}$$

■

**Definition 4.9.** We call by the  $2^{th}$  quadratic cohomology group of the metric Jacobi-Jordan algebra  $\mathfrak{J}$  with on  $\mathfrak{a}$ , the quotient

$$H_Q^2(\mathfrak{J}, \mathfrak{a}) = Z_Q^2(\mathfrak{J}, \mathfrak{a}) / B_Q^2(\mathfrak{J}, \mathfrak{a})$$

## 5 Twofold extension

Let  $\rho$  be a representation of a Hom-Jacobi-Jordan algebra  $(J, \delta)$  on a vector space  $\mathfrak{a}$  and  $B_{\mathfrak{a}}: \mathfrak{a}^2 \rightarrow \mathbb{R}$  be a bilinear non degenerate form. we set  $\mathfrak{n} = J \oplus \mathfrak{a} \oplus J^*$

Let  $(\theta, \gamma) \in C_Q^2(J, \mathfrak{a})$ . Define a bilinear map  $d: \mathfrak{n}^2 \rightarrow \mathfrak{n}$  by

$$\begin{aligned} d(x, y) &= \delta(x, y) + \theta(x, y) + \gamma(x, y, \cdot); \\ d(x, v) &= \rho(x)v + B_{\mathfrak{a}}(\theta(\cdot, x), v); \\ d(v, w) &= B_{\mathfrak{a}}(\rho(\cdot)v, w); \\ d(Z, x) &= Z(\delta(x, \cdot)); \\ d(Z_1, v + Z_2) &= 0. \end{aligned}$$

We assume that  $(\theta, \gamma)$  is a quadratic 2-cocycle. Then, using the Proposition 4.4,  $(\mathfrak{n}, d, B_{\mathfrak{n}})$  is a metric Hom-Jacobi-Jordan algebra.

**Definition 5.1.** We denote the metric Hom-Jordan-Jacobi algebra  $(\mathfrak{n}, d, B_{\mathfrak{n}})$  constructed above by  $\mathfrak{d}_{\theta, \gamma}(\mathfrak{a}, J, \rho)$  and call it a twofold extension.

**Definition 5.2.** Two Twofold extensions  $\mathfrak{d}_{\theta,\gamma} := \mathfrak{d}_{\theta,\gamma}(\mathfrak{a}, J, \rho)$  and  $\mathfrak{d}_{\theta',\gamma'} := \mathfrak{d}_{\theta',\gamma'}(\mathfrak{a}, J, \rho)$  are equivalent if and only if there exist an isomorphism  $\Phi : J \oplus \mathfrak{a} \oplus J^* \rightarrow J \oplus \mathfrak{a} \oplus J^*$ , such  $\Phi$  an isometry,  $\Phi(J^*) \subset J^*$  and the following extensions

$$\begin{array}{ccccc} 0 & \longrightarrow & \mathfrak{a} & \xrightarrow{i} & J \oplus \mathfrak{a} & \xrightarrow{\pi} & J & \longrightarrow & 0 \\ & & & \downarrow id_{\mathfrak{a}} & & \downarrow \Phi' & & \downarrow id_J & \\ 0 & \longrightarrow & \mathfrak{a} & \xrightarrow{i} & J \oplus \mathfrak{a} & \xrightarrow{\pi} & J & \longrightarrow & 0 \end{array}$$

are equivalent, where  $\Phi'$  is the projection of  $\Phi$  on  $J \oplus \mathfrak{a}$ .

**Lemma 5.3.** Let  $\mathfrak{d}_{\theta,\gamma} := \mathfrak{d}_{\theta,\gamma}(\mathfrak{a}, J, \rho)$  and  $\mathfrak{d}_{\theta',\gamma'} := \mathfrak{d}_{\theta',\gamma'}(\mathfrak{a}, J, \rho)$  two Twofold extensions such that  $d_Q^2(\theta, \gamma) = d_Q^2(\theta', \gamma')$ . Then, the Twofold extensions  $\mathfrak{d}_{\theta,\gamma} := \mathfrak{d}_{\theta,\gamma}(\mathfrak{a}, J, \rho)$  and  $\mathfrak{d}_{\theta',\gamma'} := \mathfrak{d}_{\theta',\gamma'}(\mathfrak{a}, J, \rho)$  are equivalent.

*Proof.* Define the linear map  $\Phi : J \oplus \mathfrak{a} \oplus J^* \rightarrow J \oplus \mathfrak{a} \oplus J^*$  by

$$\Phi(x + v + Z) = x + v + Z - \tau(x) - \frac{1}{2}B_{\mathfrak{a}}(\tau(x), \tau(\cdot)) + B_{\mathfrak{a}}(v, \tau(\cdot)).$$

We have

$$\begin{aligned} d(x, y) &= \delta(x, y) + \theta(x, y) + \gamma(x, y, \cdot); \\ d(x, v) &= \rho(x)v + B_{\mathfrak{a}}(\alpha(\cdot, x), v); \\ d(v, w) &= B_{\mathfrak{a}}(\rho(\cdot)v, w); \\ d(Z, x) &= Z(\delta(x, \cdot)); \\ d(Z_1, v + Z_2) &= 0. \end{aligned}$$

Using Proposition 4.6, we have  $\theta' = \theta + d^1\tau$  and  $\gamma' = \gamma + d^2\sigma - B(\tau \wedge (\theta + \frac{1}{2}d^1\tau))$ . Hence, the structure  $d'$  of the Twofold extension  $\mathfrak{d}_{\theta',\gamma'} := \mathfrak{d}_{\theta',\gamma'}(\mathfrak{a}, J, \rho)$  is given by

$$\begin{aligned} d'(x, y) &= \delta(x, y) + \theta(x, y) + d^1\tau(x, y) + \gamma(x, y, \cdot) \\ &\quad + d_r^2\sigma(x, y, \cdot) - B((\theta + \frac{1}{2}d^1\tau) \wedge \tau)(x, y, \cdot); \\ d'(x, v) &= \rho(x)v + B_{\mathfrak{a}}(\theta(\cdot, x) + d^1\tau(\cdot, x), v); \\ d'(v, w) &= B_{\mathfrak{a}}(\rho(\cdot)v, w); \\ d'(Z, x) &= Z(\delta(x, \cdot)); \\ d'(Z_1, v + Z_2) &= 0. \end{aligned}$$

We have

$$\begin{aligned}
\Phi(d'(x, y)) &= \delta(x, y) + \theta(x, y) + d^1\tau(x, y) + \gamma(x, y, \cdot) \\
&\quad + d^2\sigma(x, y, \cdot) - B_a\left(\left(\theta + \frac{1}{2}d^1\tau\right) \wedge \tau\right)(x, y, \cdot) \\
&\quad - \tau(\delta(x, y)) - \sigma(\delta(x, y), \cdot) + \frac{1}{2}B_a(\tau(\delta(x, y)), \tau(\cdot)) \\
&\quad + B_a\left(\left(\theta(x, y) + d^1\tau(x, y)\right), \tau(\cdot)\right)
\end{aligned}$$

Hence, by (1.33), (4.1) and (4.15), we obtain

$$\begin{aligned}
\Phi(d'(x, y)) &= \delta(x, y) + \theta(x, y) + \gamma(x, y, \cdot) - \rho(x)\tau(y) - \rho(y)\tau(x) \\
&\quad - \sigma(y, \delta(x, \cdot)) - \sigma(x, \delta(y, \cdot)) \\
&\quad - B_a(\theta(x, \cdot), \tau(y)) - B_a(\theta(y, \cdot), \tau(x)) \\
&\quad - \frac{1}{2}B_a\left(\tau(\delta(x, \cdot)), \tau(y)\right) - \frac{1}{2}B_a\left(\tau(\delta(y, \cdot)), \tau(x)\right) + B_a\left(\rho(\cdot)\tau(x), \tau(y)\right).
\end{aligned}$$

In other hand, we have

$$\begin{aligned}
d(\Phi(x), \Phi(y)) &= d\left(x - \tau(x) - \sigma(x, \cdot) - \frac{1}{2}B_a(\tau(x), \tau(\cdot)), y - \tau(y) - \sigma(y, \cdot) - \frac{1}{2}B_a(\tau(y), \tau(\cdot))\right) \\
&= \delta(x, y) + \theta(x, y) + \gamma(x, y, \cdot) - \rho(x)\tau(y) - B_a(\theta(\cdot, x), \tau(y)) \\
&\quad - \sigma(y, \delta(x, \cdot)) - \frac{1}{2}B_a\left(\tau(y), \tau(\delta(x, \cdot))\right) - \rho(y)\tau(x) - B_a(\theta(\cdot, y), \tau(x)) \\
&\quad + B_a(\rho(\cdot)\tau(x), \tau(y)) - \sigma(x, \delta(y, \cdot)) - \frac{1}{2}B_a\left(\tau(x), \tau(\delta(y, \cdot))\right).
\end{aligned}$$

Therefore,  $\Phi(d'(x, y)) = d(\Phi(x), \Phi(y))$ .

Similar, we show that  $\Phi(d'(x, w)) = d(\Phi(x), \Phi(w))$ ,  $\Phi(d'(x, Z)) = d(\Phi(x), \Phi(Z))$ ,  $\Phi(d'(v, w)) = d(\Phi(v), \Phi(w))$ .

$$\begin{aligned}
B(\Phi(x), \Phi(y)) &= B\left(x - \tau(x) - \frac{1}{2}B_a(\tau(x), \tau(\cdot)), y - \tau(y) - \frac{1}{2}B_a(\tau(y), \tau(\cdot))\right) \\
&= B_a(\tau(x), \tau(y)) - \frac{1}{2}B_a(\tau(y), \tau(x)) - \frac{1}{2}B_a(\tau(x), \tau(y)) \\
&= 0 = B(x, y)
\end{aligned}$$

$$\begin{aligned}
B(\Phi(x), \Phi(v)) &= B\left(x - \tau(x) - \frac{1}{2}B_a(\tau(x), \tau(\cdot)), v + B_a(v, \tau(\cdot))\right) \\
&= -B_a(\tau(x), v) + B_a(v, \tau(x)) = 0
\end{aligned}$$

$$\begin{aligned} B(\Phi(u), \Phi(v)) &= B(u + B_{\mathfrak{a}}(u, \tau(\cdot)), v + B_{\mathfrak{a}}(v, \tau(\cdot))) \\ &= B_{\mathfrak{a}}(u, v) \end{aligned}$$

■

**Lemma 5.4.** *Let  $\mathfrak{d}_{\theta, \gamma} := \mathfrak{d}_{\theta, \gamma}(\mathfrak{a}, J, \rho)$  and  $\mathfrak{d}_{\theta', \gamma'} := \mathfrak{d}_{\theta', \gamma'}(\mathfrak{a}, J, \rho)$  be two equivalent Twofold extensions. Then, the quadratic 2-cocycle  $(\theta - \theta', \gamma - \gamma')$  is trivial.*

*Proof.* Let  $\Phi(x) = f(x) + \tau(x) + \zeta(x)$  where  $f: J \rightarrow J$ ,  $\tau: J \rightarrow \mathfrak{a}$  and  $\zeta: J \rightarrow J^*$ . Using  $\pi \circ \Phi' = \pi$ , we obtain  $f(x) = x$ . Then

$$\Phi(x) = x + \tau(x) + \zeta(x).$$

Let  $\Phi(v) = g(v) + h(v) + \eta(v)$ , where  $g: \mathfrak{a} \rightarrow J$ ,  $h: \mathfrak{a} \rightarrow \mathfrak{a}$  and  $\eta: \mathfrak{a} \rightarrow J^*$ . Using  $\Phi' \circ i = i$ , we obtain  $g(v) = 0$  and  $h(v) = v$ . Then  $\Phi(v) = v + \eta(v)$ . Using  $B(v, x) = B(\Phi(v), \Phi(x))$ , we obtain  $\eta(v)(x) = -B_{\mathfrak{a}}(v, \tau(x))$ . With  $\Phi$  is an isometry and  $\Phi(J^*) \subset J^*$  we obtain  $\Phi(Z) = Z$ .

Using  $B(\Phi(x), \Phi(y)) = B(x, y)$  we obtain  $B_{\mathfrak{a}}(\tau(x), \tau(y)) = -\zeta(x)(y) - \zeta(y)(x)$ . Since  $\zeta(x)(y) = \zeta(y)(x)$ , we obtain  $\zeta(x, y) = -\frac{1}{2}B_{\mathfrak{a}}(\tau(x), \tau(y))$ .

By  $\Phi(d(x, y)) = d'(\Phi(x), \Phi(y))$ , we obtain

$$\theta(x, y) = \theta'(x, y) - \tau(\delta(x, y)) + \rho(x)\tau(y) + \rho(y)\tau(x) = \theta'(x, y) - d^1\tau(x, y)$$

and

$$\gamma(x, y, \cdot) = \gamma'(x, y, \cdot) - B_{\mathfrak{a}}\left(\left(\theta' + \frac{1}{2}d(-\tau)\right) \wedge (-\tau)\right)(x, y, \cdot).$$

Hence,

$$\begin{cases} \alpha = \alpha' + d^1(-\tau) \\ \gamma = \gamma'(x, y, \cdot) - B_{\mathfrak{a}}\left(\left(\alpha' + \frac{1}{2}d(-\tau)\right) \wedge (-\tau)\right)(x, y, \cdot) \end{cases}$$

Using Proposition 1.8, we have  $d_c^2\theta = 0$ .

Therefore, using Proposition 4.6, we have  $d_Q^2(\theta, \gamma) = d_Q^2(\theta', \gamma')$  ■

**Theorem 5.5.** *The set  $Ext(\mathfrak{J}, \mathfrak{a})$  of equivalence classes of Twofold extensions  $(\mathfrak{J}, \delta, \alpha)$  by an abelian  $\mathfrak{a}$  is one-to-one correspondence with  $Z_Q^2(\mathfrak{J}, \mathfrak{a})/B^2(\mathfrak{J}, \mathfrak{a})$ , that is*

$$Ext(\mathfrak{J}, \mathfrak{a}) \cong H_Q^2(\mathfrak{J}, \mathfrak{a}).$$

## 6 Quadratic extensions

Let  $(J, \delta, \alpha)$  be a Hom-Jacobi-Jordan algebra and  $(\mathfrak{a}, \rho, B_{\mathfrak{a}})$  a symmetric  $J$ -module.

**Definition 6.1.** A quadratic extension of  $J$  by  $\mathfrak{a}$  is given by a quadruple  $(\mathfrak{J}, I, i, \pi)$ , where

- $(\mathfrak{J}, d, \gamma)$  is a metric Hom-Jacobi-Jordan algebra;
- $I$  is an isotropic ideal in  $\mathfrak{J}$  and;
- $i$  and  $\pi$  are Hom-Jacobi-Jordan homomorphisms constituting an exact sequence of Jacobi-Jordan algebras

$$0 \longrightarrow \mathfrak{a} \xrightarrow{i} \mathfrak{J}/I \xrightarrow{\pi} J \longrightarrow 0,$$

which is consistent with the representation  $\rho$  of  $J$  on  $\mathfrak{a}$  in the following sense:

$$i(\rho(x)(v)) = d(\tilde{x}, i(v)) \in i(\mathfrak{a})$$

holds for all  $\tilde{x} \in \mathfrak{J}/I$  with  $\pi(\tilde{x}) = x$ . In addition we require that  $i(\mathfrak{a}) = I^{\perp}/I$  and that  $i: \mathfrak{a} \rightarrow I^{\perp}/I$  is an isometry.

**Proposition 6.2.** Let  $\mathfrak{d}_{\theta, \gamma} := \mathfrak{d}_{\theta, \gamma}(\mathfrak{a}, J, \rho)$  be a Twofold extension. Define the linear map  $i_0: \mathfrak{a} \rightarrow \mathfrak{d}/J^*$  by  $i_0(v) = v + J^*$  and the linear map  $\pi_0: \mathfrak{d}/J^* \rightarrow J$  by  $\pi_0(x + v + J^*) = x$ . Then the quadruple  $(\mathfrak{d}_{\alpha, \gamma}, J^*, i_0, \pi_0)$  define a quadratic extension.

*Proof.* Define the symmetric bilinear map  $\tilde{d}: \mathfrak{n}/J^* \times \mathfrak{n}/J^* \rightarrow \mathfrak{n}/J^*$  by

$$\begin{aligned} \tilde{d}(x + J^*, y + J^*) &= \delta(x, y) + \theta(x, y) + J^* \\ \tilde{d}(x + J^*, v + J^*) &= \rho(x)v + J^*. \end{aligned}$$

Then  $d(x + J^*, i_0(v) + J^*) = d(x + J^*, v + J^*) = \rho(x)v + J^* = i_0(\rho(x)v)$ .

Moreover, we have

$$i_0(v) = v + J^*, B(v, Z) = 0 \text{ then, } i_0(\mathfrak{a}) \subset J^{*\perp}.$$

$$a \in J^{*\perp} \implies B(a, Z) = 0 \implies a \in \mathfrak{a} \oplus J^*.$$

$$B(i_0(v), i_0(w)) = B(v + J^*, w + J^*) = 0.$$

■

In the rest of this paper,  $J$  is assumed to be finite-dimensional.

**Proposition 6.3.** *Let*

$$0 \longrightarrow \mathfrak{a} \xrightarrow{i} \mathfrak{J}/I \xrightarrow{\pi} J \longrightarrow 0, \quad (6.1)$$

be an extension of  $J$  by  $\mathfrak{a}$  such that  $i: \mathfrak{a} \rightarrow i(\mathfrak{a})$  is an isometry. Then, the quadruple  $(\mathfrak{J}, I, i, \pi)$  define a quadratic extension if and only if the following sequence define an extension of  $\mathfrak{J}/I$  by  $J^*$ :

$$0 \longrightarrow J^* \xrightarrow{\tilde{\pi}^*} \mathfrak{J} \xrightarrow{\pi_n} \mathfrak{J}/I \longrightarrow 0, \quad (6.2)$$

where  $\pi_n$  is the natural projection  $\mathfrak{J} \rightarrow \mathfrak{J}/I$ ,  $\tilde{\pi} = \pi \circ \pi_n$ ,  $\tilde{\pi}^*$  the dual map of  $\tilde{\pi}$  where we identify  $J^*$  with  $J$ .

*Proof.* Since  $J$  has a finite dimensional,

$$(I^\perp)^\perp = I$$

,

$$Im(\tilde{\pi}^*) = (\ker(\tilde{\pi}))^\perp$$

,

$$\ker(\tilde{\pi}^*) = (Im(\tilde{\pi}))^\perp$$

$$\tilde{d}(\bar{x}, \bar{y}) = d(x, y) + I.$$

We assume that  $(\mathfrak{J}, I, i, \pi)$  is a quadratic extension. Then,  $\ker(\pi) = Im(i) = I^\perp/I$ . Moreover, for any  $x \in J$ , we have

$$\begin{aligned} x \in \ker(\tilde{\pi}) &\Leftrightarrow \pi \circ \pi_n(x) = 0 \Leftrightarrow \pi(\bar{x}) = 0 \Leftrightarrow \bar{x} \in \ker(\pi) \\ &\Leftrightarrow \bar{x} \in I^\perp/I \\ &\Leftrightarrow x \in I^\perp \end{aligned}$$

Then  $\ker(\tilde{\pi}) = I^\perp$ . Hence,  $(\ker(\tilde{\pi}))^\perp = I$ . Since  $Im(\tilde{\pi}^*) = (\ker(\tilde{\pi}))^\perp$ , one can deduce  $Im(\tilde{\pi}^*) = I$ . So  $Im(\tilde{\pi}^*) = \ker(\pi_n)$  and the sequence

$$0 \longrightarrow J^* \xrightarrow{\tilde{\pi}^*} \mathfrak{J} \xrightarrow{\pi_n} \mathfrak{J}/I \longrightarrow 0,$$

define an extension of  $\mathfrak{J}/I$  by  $J^*$ .

Conversely, we assume that the following sequence

$$0 \longrightarrow J^* \xrightarrow{\tilde{\pi}^*} \mathfrak{J} \xrightarrow{\pi_n} \mathfrak{J}/I \longrightarrow 0,$$

define an extension. Then,  $Im(\tilde{\pi}^*) = \ker(\pi_n) = I$ . We have  $Im(\tilde{\pi}^*) = (\ker(\tilde{\pi}))^\perp$ . Then  $(\ker(\tilde{\pi}))^\perp = I$ . So,  $\ker(\tilde{\pi}) = I^\perp$ . Hence,  $I \subset I^\perp$ .

$$\begin{aligned} \tilde{x} \in \ker(\pi) &\iff \pi(\tilde{x}) = 0_J \iff \pi(\pi_n(x)) = 0_J \\ &\iff x \in \ker(\tilde{\pi}) \iff x \in I^\perp \iff \tilde{x} \in I^\perp/I. \end{aligned}$$

Hence  $Im(i) = \ker(\pi) = I^\perp/I$ . This completes the proof. ■

**Definition 6.4.** Two quadratic extensions  $(\mathfrak{J}_1, I_1, i_1, \pi_1)$ ,  $(\mathfrak{J}_2, I_2, i_2, \pi_2)$  of  $J$  by  $\mathfrak{a}$  are called to be equivalent if there exists an isomorphism of metric Lie algebras  $\Phi: \mathfrak{J}_1 \rightarrow \mathfrak{J}_2$  which maps  $i_1$  onto  $i_2$  and satisfies  $\overline{\Phi} \circ i_1 = i_2$  and  $\pi_2 \circ \overline{\Phi} = \pi_1$ , where  $\overline{\Phi}: \mathfrak{J}_1/I_1 \rightarrow \mathfrak{J}_2/I_2$  is the induced map.

**Proposition 6.5.** Any quadratic extension  $(\mathfrak{J}, I, i, \pi)$  is equivalent to a Twofold extension  $(\mathfrak{d}_{\theta, \gamma}, J^*, i_0, j_0)$ .

*Proof.* Let

$$0 \longrightarrow \mathfrak{a} \xrightarrow{i} \mathfrak{J}/I \xrightarrow{\pi} J \longrightarrow 0,$$

be the extension of  $J$  defined in (6.1). Then there exist a metric Hom-Jacobi-Jordan subalgebra  $H \subset \mathfrak{J}/I$  complementary to  $\ker \pi$ . Since  $Im i = \ker \pi$ , we have  $\mathfrak{J}/I = H \oplus i(\mathfrak{a})$ . The map  $\pi_{/H}: H \rightarrow J$  (resp  $k: \mathfrak{a} \rightarrow i(\mathfrak{a})$ ) defined by  $\pi_{/H}(\overline{x}) = \pi(\overline{x})$  (resp.  $k(v) = i(v)$ ) is bijective, its inverse  $s$  (resp.  $l$ ). Considering the map  $\Phi: J \oplus \mathfrak{a} \rightarrow \mathfrak{J}/I$  defined by  $\Phi(x + v) = s(x) + i(v)$ , it is easy to verify that  $\Phi$  is an isomorphism,  $i(\mathfrak{a})$  is an ideal of  $\mathfrak{J}/I$  and

$$\mathfrak{J}/I = s(J) \oplus i(\mathfrak{a}). \quad (6.3)$$

Now, we consider the extension

$$0 \longrightarrow J^* \xrightarrow{\tilde{\pi}^*} \mathfrak{J} \xrightarrow{\pi_n} \mathfrak{J}/I \longrightarrow 0,$$

be the extension defined in (6.2). Then there exist a metric Jacobi-Jordan subalgebra  $H' \subset \mathfrak{J}$  complementary to  $\ker \pi_n$ . Since  $Im \tilde{\pi}^* = \ker \pi_n = I$ , we have  $\mathfrak{J} = H' \oplus I$ . The restriction of  $\pi_n$  to  $H'$ ,  $\pi': H' \rightarrow \mathfrak{J}/I$  (resp  $k': J^* \rightarrow \tilde{\pi}^*(J^*)$ ) defined by  $\pi' = \pi_n(x)$  (resp.  $k'(Z) = \tilde{\pi}^*(Z)$ ) is bijective, its inverse  $s'$  (resp.  $l'$ ). Considering the map  $\Phi': \mathfrak{J}/I \oplus J^* \rightarrow \mathfrak{J}$  defined by  $\Phi'(\overline{x} + Z) = s'(\overline{x}) + i'(Z)$ , it is easy to verify that  $\Phi'$  is an isomorphism,  $\tilde{\pi}^*(J^*)$  is an ideal of  $\mathfrak{J}/I$  and  $\mathfrak{J} = s'(\mathfrak{J}/I) \oplus \tilde{\pi}^*(J^*)$ . We can write  $\pi_n: s'(\mathfrak{J}/I) \rightarrow \mathfrak{J}/I$  and  $\pi: s(J) \rightarrow J$ . Hence  $\tilde{\pi}^*(J^*) = (s' s(J))^*$



Using (6.3) and  $\tilde{\pi}^*(J^*) = (s's(J))^*$ , we obtain  $\mathfrak{J} = s's(J) \oplus s'i(\mathfrak{a}) \oplus (s's(J))^*$ .

Then, we have

$$\begin{aligned} d'(s's(x), s's(y)) &= \delta(s's(x), s's(y)) + \theta(s's(x), s's(y)) + \gamma(s's(x), s's(y), \cdot); \\ d'(s's(x), s'i(v)) &= \rho(s's(x))v + B_\rho(s'i(v), \alpha(s's(x), \cdot)); \\ d'(s'i(v), s'i(w)) &= B_\alpha(\rho(\cdot)(s'i(v)), s'i(w)); \\ d'(Z, s's(x)) &= Z(\delta(s's(x), \cdot)); \\ d'(Z_1, s'i(v) + Z_2) &= 0. \end{aligned}$$

for all  $x \in \mathfrak{J}$ ,  $v \in \mathfrak{a}$ ,  $Z \in \mathfrak{J}^*$ .

Now, define the linear map  $\Psi: J \oplus \mathfrak{a} \oplus J^* \rightarrow \mathfrak{J}$  by  $\Psi(x + v + Z) = s's(x) + s'i(v) + (s's)^*(Z)$  and the bilinear map  $d: J \oplus \mathfrak{a} \oplus J^* \rightarrow J \oplus \mathfrak{a} \oplus J^*$  by

$$d(x+v+Z, y+w+Z') = \Psi^{-1}(d'(s's(x)+s'i(v)+(s's)^*(Z), s's(y)+s'i(w)+(s's)^*(Z'))).$$

Then

$$\begin{aligned} d'(\Psi(x + v + Z), \Psi(y + w + Z')) &= d'(s's(x) + s'i(v) + (s's)^*(Z), s's(y) + s'i(w) + (s's)^*(Z')) \\ &= \Psi(d(x + v + Z, y + w + Z')). \end{aligned}$$

Moreover, we have  $\bar{\Psi} \circ i_0(v) = i(v)$  and  $\pi \circ \bar{\Psi}(\bar{x}) = \pi \circ s(x) = x = \pi_0(x)$ . ■

The upshot of this section is to show the following theorem:

**Theorem 6.6.** *The set  $Ext_Q(J, \mathfrak{a}, \rho)$  of equivalence classes of quadratic extensions of  $J$  by an abelian  $\mathfrak{a}$  is in bijection with  $H_Q^2(\mathfrak{J}, \mathfrak{a}, \rho)$ .*

*Proof.*  $(\mathfrak{J}_1, I_1, i_1, \pi_1)$  and  $(\mathfrak{J}_2, I_2, i_2, \pi_2)$  two equivalent quadratic extensions of a Jacobi-Jordan algebra  $(J, \delta)$  by an abelian Jacobi-Jordan algebra  $\mathfrak{a}$ . Then, by **Proposition 6.5**,  $(\mathfrak{J}_1, I_1, i_1, \pi_1)$  is equivalent to a Twofold extension  $\mathfrak{d}_{\theta_1, \gamma_1}(\mathfrak{a}, J_1, \rho)$  and  $(\mathfrak{J}_2, I_2, i_2, \pi_2)$  is equivalent to a Twofold extension  $\mathfrak{d}_{\theta_2, \gamma_2}(\mathfrak{a}, J_2, \rho)$ . So,  $\mathfrak{d}_{\alpha_1, \gamma_1}(\mathfrak{a}, J_1, \rho)$  and  $\mathfrak{d}_{\alpha_2, \gamma_2}(\mathfrak{a}, J_2, \rho)$  are equivalent. Therefore by Theorem 5.5,  $\theta_1$  and  $\theta_2$  are equivalent. ■

## References

- [1] Ammar, F., Ejbehi, Z., Makhlouf, A.: *Cohomology and Deformations of Hom-algebras*. Journal of Lie Theory **21** (4), 813 – 836 (2011)
- [2] C. E. Haliya and G. D. Houndedji "Hom-Jacobi-Jordan and Hom-antissociative algebras with symmetric invariant nondegenerate bilinear forms." Quasigroups and Related Systems. 29, (2021), 61-88

- [3] Casas, J. M., Insua, M. A., Pacheco Rego, N.: *On universal central extensions of Hom-Lie algebras*. Hacettepe Journal of Mathematics and Statistics **44** (2) , 277 – 288 (2015)
- [4] D. Burde and A. Fialowski. "Jacobi–Jordan algebras." Linear Algebra and its Applications. 459, (2014), 586–594.
- [5] Hartwig, J.T., Larsson, D., Silvestrov, S.D.: *Deformations of Lie algebras using  $\sigma$ -derivations*. Journal of Algebra **295**(2), 314 – 361. (2006).
- [6] Jin, Q., Li, X.: *Hom-Lie algebra structures on semi-simple Lie algebras*. Journal of Algebra **319** , 1398 – 1408 (2008)
- [7] Makhlouf, A., Silvestrov, S.: *Hom-algebra structures*. Journal of Generalized Lie Theory and Applications **2** (2), 51 – 64 (2008)
- [8] Makhlouf, A., Silvestrov, S.: *Notes on 1-parameter formal deformations of Hom-associative and Hom-Lie algebras* . Forum Math **22** (4), 715–739 (2010)
- [9] Sheng, Y.: *Representations of Hom-Lie algebras*. Algebras and Representation Theory **15**(6), 1081–1098 (2012)