

# QUANTUM TELEPORTATION IN THE COMMUTING OPERATOR FRAMEWORK

ALEXANDRE CONLON, JASON CRANN, DAVID W. KRIBS, AND RUPERT H. LEVENE

**ABSTRACT.** We introduce a notion of teleportation scheme between subalgebras of semi-finite von Neumann algebras in the commuting operator model of locality. Using techniques from subfactor theory, we present unbiased teleportation schemes for relative commutants  $N' \cap M$  of a large class of finite-index inclusions  $N \subseteq M$  of tracial von Neumann algebras, where the unbiased condition means that no information about the teleported observables are contained in the classical communication sent between the parties. For a large class of subalgebras  $N$  of matrix algebras  $M_n(\mathbb{C})$ , including those relevant to hybrid classical/quantum codes, we show that any tight teleportation scheme for  $N$  necessarily arises from an orthonormal unitary Pimsner-Popa basis of  $M_n(\mathbb{C})$  over  $N'$ , generalising work of Werner [91]. Combining our techniques with those of Brannan-Ganesan-Harris [22], we compute quantum chromatic numbers for a variety of quantum graphs arising from finite-dimensional inclusions  $N \subseteq M$ .

## 1. INTRODUCTION

Quantum teleportation [12], the transfer of qubits between two separated parties using preshared entanglement and local operations and classical communication, is a quintessential protocol in quantum information. It and its variants are used in a multitude of scenarios, including quantum error correction [13], quantum cryptography [44] and universal quantum computation [45].

Mathematically, the protocol involves three single qubit quantum systems,  $A_0 \otimes A_1 \otimes B = \mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$ , with Alice ( $A$ ) having access to the first two systems and Bob ( $B$ ) having access to the third. The composite system  $A_1 B$  is entangled in the (maximally entangled) Bell state  $|\beta_{00}\rangle = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$ . The entire two qubit Bell basis for  $A_1 B$  is then obtained as  $|\beta_{ij}\rangle = (1 \otimes X^i Z^j)|\beta_{00}\rangle$ , for  $0 \leq i, j \leq 1$  and  $X, Z$  denoting the usual single qubit Pauli operators (in this case acting on the third qubit). One can then verify that for an arbitrary single qubit  $|\psi\rangle \in \mathbb{C}^2$ , we have

$$|\psi\rangle|\beta_{00}\rangle = \frac{1}{2}(|\beta_{00}\rangle|\psi\rangle + |\beta_{01}\rangle(X|\psi\rangle) + |\beta_{10}\rangle(Z|\psi\rangle) + |\beta_{11}\rangle(XZ|\psi\rangle)).$$

Thus,  $A$  proceeds by measuring the first two qubit composite system  $A_0 A_1$  in its Bell basis  $\{|\beta_{ij}\rangle\}$ , which is given as a quantum measurement on the combined three qubit system by the family of projection operators  $P_{ij} \otimes 1$ ,  $0 \leq i, j \leq 1$ , with  $P_{ij} = |\beta_{ij}\rangle\langle\beta_{ij}|$  on  $A_0 A_1$  and  $1$  the identity operator on  $B$ . The party  $A$  then communicates the result  $(i, j)$  of this measurement to  $B$ , who then implements the unitary reversal  $(X^i Z^j)^*$  on the third system to obtain the state  $|\psi\rangle$ , and this completes the protocol.

Using the maximally entangled state  $|\psi_n\rangle := \frac{1}{\sqrt{n}} \sum_{i=0}^{n-1} |ii\rangle \in \mathbb{C}^n \otimes \mathbb{C}^n$  and the generalised Pauli  $X$  and  $Z$  operators on  $\mathbb{C}^n$ , the procedure generalises verbatim to states in  $\mathbb{C}^n$ . This latter protocol was put into the larger context of *teleportation schemes* by Werner [91], which allowed for broader possible implementations by the parties. Specifically, a teleportation scheme for  $\mathbb{C}^n$  consists of a triple  $(\omega, \{F_i\}_{i \in I}, \{T_i\}_{i \in I})$  where  $\omega$  is a density on  $\mathbb{C}^n \otimes \mathbb{C}^n$  (entangled resource state),  $\{F_i\}_{i \in I}$  is a positive operator-valued measure (POVM) on  $\mathbb{C}^n \otimes \mathbb{C}^n$  (Alice's measurement system) and  $\{T_i\}_{i \in I}$

2010 *Mathematics Subject Classification.* 46L10, 46L30, 46N50, 47L90, 81P40, 81P45, 81R15.

*Key words and phrases.* quantum teleportation, local operations and classical communication, von Neumann algebra, subfactors, quantum graphs, quantum chromatic numbers.

are unital completely positive (UCP) maps on  $M_n(\mathbb{C})$  (Bob's quantum channels in the Heisenberg picture) for which

$$\mathrm{tr}(\rho B) = \sum_{i \in I} \mathrm{tr}((\rho \otimes \omega)(F_i \otimes T_i(B))), \quad \forall \rho, B \in M_n(\mathbb{C}).$$

The scheme  $(\omega, \{F_i\}_{i \in I}, \{T_i\}_{i \in I})$  is *tight* [91] if  $|I| = n^2$ , that is, the amount of classical signals communicated coincides with the dimension of the algebra to be teleported: in the Heisenberg picture, Bob's observable algebra ( $M_n(\mathbb{C})$ ) is teleported to Alice's local observable algebra. Werner established a correspondence between tight teleportation schemes for  $M_n(\mathbb{C})$  and unitary error bases of  $M_n(\mathbb{C})$  [91], that is, orthonormal bases of unitaries with respect to the (normalised) Hilbert-Schmidt inner product (a prime example given by the generalised Pauli operators).

Rather than teleportation of a full system, one can imagine scenarios in which it is desirable to teleport quantum information encoded into subsystems of a full system or even hybrid forms of classical and quantum information. This could arise for instance with subsystem codes used in quantum algorithms and fault tolerant quantum computing architectures [1, 6, 61, 63, 74, 78] or hybrid codes used for the simultaneous transmission of classical and quantum information in communication schemes [15, 16, 24, 32, 46, 62, 65, 71, 72]. Such scenarios admit convenient mathematical descriptions in the Heisenberg picture, and as such, can be studied through generalisations of teleportation schemes to the commuting operator framework, in which locality is modelled by commuting algebras of observables [47]. Indeed, the mathematical origins of quantum theory [87] together with recent advances in non-local games (e.g., [34, 48, 54, 59, 75, 79]) and the increasing number of connections between quantum information and quantum field theory (e.g., [4, 25, 42, 49, 52]) continue to motivate the study/extension of central results in quantum information to the commuting operator framework. Examples include entropy theory (e.g., [17, 39, 41, 50, 51, 67]), quantum error correction (e.g., [14, 28]), the theory of local operations (e.g., [29, 40, 85]) and entanglement in quantum field theory (e.g., [52] and the references therein).

In this paper we continue this line of work by introducing a general notion of teleportation scheme in the setting of semi-finite von Neumann algebras and studying analogues of tightness and related properties in this setting. Our main examples utilize subfactor theory and deepen the connection between standard teleportation of observables in  $M_n(\mathbb{C})$  and Jones' basic construction for the inclusion  $\mathbb{C} \subseteq M_n(\mathbb{C})$  (see, e.g., [92]). In particular, in section 3 we establish unbiased teleportation protocols for relative commutants  $N' \cap M$  of (finite-index) inclusions  $N \subseteq M$  which admit orthonormal Pimsner-Popa bases in the unitary normaliser of  $N$ . See Theorem 3.7 for details.

Restricting attention to the tripartite system  $M_n(\mathbb{C}) \otimes M_n(\mathbb{C}) \otimes M_n(\mathbb{C})$ , in section 4 we generalise Werner's characterization of tight teleportation schemes [91], showing that, for a large class of (unital  $*$ -)subalgebras  $N \subseteq M_n(\mathbb{C})$ , any tight (and faithful) teleportation scheme of  $N$  between the first and third subsystems necessarily arises from a unitary orthonormal Pimsner-Popa basis for  $M_n(\mathbb{C})$  over  $N'$ . See Theorem 4.1 for a precise statement. Our result applies, in particular, to homogeneous subalgebras  $N$ , which model hybrid classical/quantum codes in the Heisenberg picture [15, 24, 46, 71, 72].

It is known that unitary error bases generate quantum-to-classical graph homomorphisms [22, 70], and therefore can be used to estimate/calculate various chromatic numbers of quantum graphs [22]. Based on our generalised teleportation schemes and techniques from [22], in section 5 we calculate various chromatic numbers for quantum graphs arising from inclusions  $N \subseteq M$  on a finite-dimensional Hilbert space  $H$ . In particular, when  $N$  is a factor, we show that

$$\chi_q(N', M, \mathcal{B}(H)) = \chi_{qc}(N', M, \mathcal{B}(H)) = [M : N],$$

where  $[M : N]$  is the index of  $N$  and  $M$ , and  $\chi_q$  and  $\chi_{qc}$  are the quantum and quantum commuting chromatic numbers of the quantum graph  $(N', M, \mathcal{B}(H))$ , respectively. This generalises the case  $N = \mathbb{C}$  established in [22]. Also, when the inclusion  $N \subseteq M$  admits an orthonormal Pimsner-Popa

basis in the unitary normaliser of  $N$ , we show that

$$\chi_{loc}(M, N', \mathcal{B}(L^2(M, \tau))) = \chi_q(M, N', \mathcal{B}(L^2(M, \tau))) = \chi_{qc}(M, N', \mathcal{B}(L^2(M, \tau))) = [M : N],$$

where  $\chi_{loc}$  is the local chromatic number.

Several natural lines of investigation are suggested by this work, including connections with weak Hopf  $C^*$ -algebras and the structure of depth-2 subfactors [73], as well as teleportation in monoidal categories [3]. The final outlook section elaborates on these and other connections left for future work.

## 2. PRELIMINARIES

In this section we outline relevant preliminaries from the theory of tracial von Neumann algebras, including Jones' basic construction, Pimsner-Popa bases, and notions of entanglement in the commuting operator framework.

**2.1. Jones' Basic Construction.** Let  $M$  be a tracial von Neumann algebra, that is, a finite von Neumann algebra with fixed normal faithful tracial state  $\tau$ . Inclusions of von Neumann subalgebras  $N \subseteq M$  will always be assumed unital.

The GNS construction of  $(M, \tau)$  yields the Hilbert space  $L^2(M, \tau)$ , the GNS map  $\Lambda_\tau : M \rightarrow L^2(M, \tau)$  and the (faithful) representation  $\pi_\tau : M \rightarrow \mathcal{B}(L^2(M, \tau))$ , where

$$\pi_\tau(x)\Lambda_\tau(y) = \Lambda_\tau(xy), \quad x, y \in M.$$

We often simply write  $x$  for  $\pi_\tau(x)$  when convenient. The inner product on  $L^2(M, \tau)$  satisfies

$$\langle \Lambda_\tau(x), \Lambda_\tau(y) \rangle = \tau(y^*x), \quad x, y \in M.$$

The adjoint operation in  $M$  yields a conjugate linear isometry  $J$  on  $L^2(M, \tau)$  via

$$J\Lambda_\tau(x) = \Lambda_\tau(x^*), \quad x \in M.$$

The associated (anti)representation of right multiplication  $\pi_\tau^r : M \rightarrow \mathcal{B}(L^2(M, \tau))$  is given by

$$\pi_\tau^r(x)\Lambda_\tau(y) = \Lambda_\tau(yx), \quad x, y \in M.$$

One easily sees that  $\pi_\tau^r(x) = J\pi_\tau(x)^*J$  for every  $x$  in  $M$ , and that  $\pi_\tau^r$  is  $*$ -preserving by traciality. It follows that  $\pi_\tau(M)' = \pi_\tau^r(M)$  (see, e.g., [81, Theorem 2.22]).

For a von Neumann subalgebra  $N \subseteq M$ , let  $L^2(N, \tau) = \overline{\Lambda_\tau(N)}$  be the associated closed subspace of  $L^2(M, \tau)$ . The orthogonal projection  $e_N$  onto  $L^2(N, \tau)$  induces the unique  $\tau$ -preserving faithful normal conditional expectation  $E_N : M \rightarrow N$  via

$$e_N\Lambda_\tau(x) = \Lambda_\tau(E_N(x)), \quad x \in M.$$

(See [82, Theorem IX.4.2] for a more general result.) The projection  $e_N$  is commonly known as the *Jones projection* for the inclusion  $N \subseteq M$ . The von Neumann subalgebra  $M_1 := \langle M, e_N \rangle$  of  $\mathcal{B}(L^2(M, \tau))$  generated by  $M$  and  $e_N$  is the result of the basic construction of the inclusion  $N \subseteq M$ . We list some standard facts (see, e.g., [57, §3.1]):

- (1)  $e_N x e_N = E_N(x) e_N$ ,  $x \in M$ ;
- (2)  $E_N(axb) = a E_N(x) b$ ,  $a, b \in N$ ,  $x \in M$ ;
- (3)  $e_N \in N'$ ;
- (4)  $J e_N = e_N J$ ;
- (5)  $M_1 = J N' J$ .

It follows that the map

$$\gamma_0 := \pi_\tau^r|_{N' \cap M} : N' \cap M \rightarrow M' \cap M_1$$

is an anti-isomorphism.

The algebra  $M_1$  has a canonical faithful semi-finite normal trace  $\text{tr}_1$  determined by  $\text{tr}_1(xe_N y) = \tau(xy)$ ,  $x, y \in M$  [77, §1.1.2]. The trace  $\tau$  is *Markov* for the inclusion  $N \subseteq M$  if  $\text{tr}_1$  is finite, and

$\tau_1 := \text{tr}_1(1)^{-1}\text{tr}_1$  has  $\tau_1|_M = \tau$ . In this case, we may iterate the basic construction to obtain  $M \subseteq M_1 \subseteq M_2$ , where  $M_2$  is the von Neumann subalgebra of  $\mathcal{B}(L^2(M_1, \tau_1))$  generated by  $M_1$  and  $e_M$ , the Jones projection for the inclusion  $M \subseteq M_1$ . As above,  $M_2 = J_{\tau_1} M' J_{\tau_1}$ , and

$$\gamma_1 := \pi_{\tau_1}^r|_{M' \cap M_1} : M' \cap M_1 \rightarrow M'_1 \cap M_2$$

is an anti-isomorphism. The composition

$$\Gamma := \gamma_1 \circ \gamma_0 : N' \cap M \rightarrow M'_1 \cap M_2$$

is therefore a  $*$ -isomorphism, known as the *canonical shift* (see e.g., [58, Proposition 2.22]). One may continue to iterate the basic construction, yielding an increasing sequence of finite von Neumann algebras  $N \subseteq M \subseteq M_1 \subseteq \dots$  known as the Jones tower of the inclusion  $N \subseteq M$ . In this paper we will only be concerned with the first two iterations.

**2.2. Bases.** Let  $N \subseteq M$  be an inclusion of finite von Neumann algebras. A finite subset  $B = \{\lambda_i \mid i = 1, \dots, d\} \subseteq M$  is a (left) *Pimsner–Popa basis*, or simply *basis*, for  $M$  over  $N$  if either of the following equivalent conditions hold:

- (1)  $\sum_{i=1}^d \lambda_i^* e_N \lambda_i = 1$ ;
- (2)  $x = \sum_{i=1}^d E_N(x \lambda_i^*) \lambda_i$  for all  $x \in M$ .

When  $E_N(\lambda_i \lambda_j^*) = \delta_{i,j} 1$ , we say that  $\{\lambda_i\}_{i=1}^d$  is an *orthonormal basis* of  $M$  over  $N$  (compare with [77], wherein orthonormality allows  $E_N(\lambda_i \lambda_i^*)$  to be a projection in  $N$ ). In this case,  $B$  forms an orthonormal basis of  $M$  as a (left) Hilbert  $N$ -module, with respect to the  $N$ -valued inner product  $\langle x, y \rangle_N = E_N(x y^*)$ . This notion of basis was introduced in [76] in the setting of  $II_1$ -factors, and was later generalised (see, e.g., [77, 83]).

Following the terminology of [58], we call an inclusion  $N \subseteq M$  *strongly Markov* if the trace  $\tau$  is Markov and there exists a finite Pimsner–Popa basis for  $M$  over  $N$ . In this case, the element

$$\sum_i \lambda_i^* \lambda_i \in \mathbb{R}^+ 1$$

is independent of the Pimsner–Popa basis. Indeed, by [77, §1.1.4]  $E_M(e_N) = \alpha 1$  for some scalar  $\alpha > 0$  (where  $E_M$  is the  $\tau_1$ -preserving conditional expectation  $M_1 \rightarrow M$ ), so that

$$\sum_i \lambda_i^* \lambda_i = \alpha^{-1} \sum_i E_M(\lambda_i^* e_N \lambda_i) = \alpha^{-1} 1.$$

The associated scalar  $\alpha^{-1}$  is the (Watatani) index  $[M : N]$  of  $M$  in  $N$  [83]. Thus, we have

$$E_M(e_N) = [M : N]^{-1} 1. \quad (1)$$

It follows that

$$e_N e_M e_N = [M : N]^{-1} e_N, \quad e_M e_N e_M = [M : N]^{-1} e_M, \quad [M : N] = [M_1 : M]. \quad (2)$$

For example, a finite-index inclusion  $N \subseteq M$  of  $II_1$  factors is strongly Markov with the index coinciding with the Jones index [56]. Also, a connected (i.e.,  $Z(N) \cap Z(M) = \mathbb{C}$ ) inclusion  $N \subseteq M$  of finite-dimensional  $C^*$ -algebras is strongly Markov and the index coincides with  $\|\Lambda_N^M\|^2$ , the square of the (operator) norm of the inclusion matrix  $\Lambda_N^M$ .

**Example 2.1.** Let  $N = \mathbb{C} \subseteq M$ . Then  $E_N = \tau(\cdot)1$  is the “completely depolarising channel”. The Jones projection satisfies

$$e_N \Lambda_\tau(x) = \Lambda_\tau(E_N(x)) = \tau(x) \Lambda_\tau(1) = \Lambda_\tau(1) \Lambda_\tau(1)^* \Lambda_\tau(x), \quad x \in M.$$

Hence,  $e_N = \Lambda_\tau(1) \Lambda_\tau(1)^*$ .

When  $M = M_n(\mathbb{C})$ ,  $\Lambda_{\tau_n}(1) = n^{-1/2} \sum_{i=0}^{n-1} |ii\rangle =: \psi_n$  is the maximally entangled state. In this case, any orthonormal basis of  $M_n(\mathbb{C})$  with respect to  $\tau_n$  (i.e., the normalised Hilbert–Schmidt inner

product) will form an orthonormal Pimsner–Popa basis for the inclusion  $\mathbb{C} \subseteq M_n(\mathbb{C})$ . A natural choice is the image of the Weyl representation

$$W : \mathbb{Z}_n \times \mathbb{Z}_n \ni (k, l) \mapsto V^l U^k \in M_n(\mathbb{C}),$$

where  $U$  and  $V$  are the translation and multiplication operators associated with the standard basis  $\{|k\rangle\}_{k \in \mathbb{Z}_n}$  of  $\mathbb{C}^n$  (also known as generalised Pauli operators):

$$U|k\rangle = |k+1\rangle, \quad V|k\rangle = e^{2\pi i k/n}|k\rangle, \quad k \in \mathbb{Z}_n. \quad (3)$$

One easily verifies that

$$\tau_n(W(z')^* W(z)) = \delta_{z', z}, \quad z, z' \in \mathbb{Z}_n^2.$$

**Example 2.2.** Let  $N = \ell_n^\infty$  (diagonals) inside  $M = M_n(\mathbb{C})$ . Then the translation operators  $\{U^k \mid k \in \mathbb{Z}_n\}$  with  $U$  defined as in Example 2.1 form an orthonormal basis for  $M$  over  $N$ . Indeed, the decomposition

$$x = \sum_{k \in \mathbb{Z}_n} E_N(x(U^k)^*) U^k$$

corresponds to breaking  $x$  into the sum of its diagonals.

Another basis of  $M_n(\mathbb{C})$  over  $\ell_n^\infty$  is  $\{\frac{1}{\sqrt{n}}|\chi\rangle\langle\chi| \mid \chi \in \widehat{\mathbb{Z}_n}\}$ . This “character basis” of  $M_n(\mathbb{C})$  over  $\ell_n^\infty$  has the nice property that each  $|\chi\rangle$  (suitably normalized) acts as a trace vector for  $\ell_n^\infty$ .

**Example 2.3.** The first part of Example 2.2 generalizes naturally to  $M \subseteq G \rtimes M$ , where  $G$  is a finite group acting on a von Neumann algebra  $M$ . In that case  $\{(\lambda_s \otimes 1) \mid s \in G\}$  is a basis of  $G \rtimes M$  over  $M$  via the usual “Fourier” decomposition

$$x = \sum_{s \in G} E_M(x(\lambda_s \otimes 1)^*) (\lambda_s \otimes 1), \quad x \in G \rtimes M,$$

where  $E_M : G \rtimes M \rightarrow M$  is the canonical conditional expectation satisfying

$$E_M\left(\sum_{s \in G} x_s (\lambda_s \otimes 1)\right) = x_e,$$

for any finitely supported function  $x : G \rightarrow M$ .

**2.3. Entanglement.** The entanglement of  $\psi_n = n^{-1/2} \sum_{i=0}^{n-1} |ii\rangle \in \mathbb{C}^n \otimes \mathbb{C}^n$  is expressed through the relation

$$\pi_{\tau_n}(x)\psi_n = (x \otimes 1)\psi_n = (1 \otimes x^t)\psi_n = \gamma_0(x)\psi_n, \quad x \in M_n(\mathbb{C}). \quad (4)$$

From a mathematical perspective, the heart of the standard teleportation protocol (see below) is the following double application of (4):

$$(|\psi_n\rangle\langle\psi_n| \otimes 1)(x \otimes 1 \otimes 1)(1 \otimes |\psi_n\rangle\langle\psi_n|) = (|\psi_n\rangle\langle\psi_n| \otimes 1)(1 \otimes 1 \otimes x)(1 \otimes |\psi_n\rangle\langle\psi_n|), \quad x \in M_n(\mathbb{C}). \quad (5)$$

Viewing  $\psi_n$ , or rather its associated density, as the Jones projection for  $\mathbb{C} \subseteq M_n(\mathbb{C})$ , these manifestations of entanglement generalize naturally, as is well-known (see [18, Lemma 2.4], for example).

**Lemma 2.4.** *Let  $N \subseteq M$  be an inclusion of finite von Neumann algebras. For any  $x \in N' \cap M$ ,*

$$xe_N = \pi_\tau^r(x)e_N = \gamma_0(x)e_N.$$

*Proof.* For any  $y \in M$ ,

$$\begin{aligned} xe_N \Lambda_\tau(y) &= x \Lambda_\tau(E_N(y)) = \Lambda_\tau(x E_N(y)) = \Lambda_\tau(E_N(y)x) \\ &= \pi_\tau^r(x) \Lambda_\tau(E_N(y)) = \gamma_0(x) e_N \Lambda_\tau(y). \end{aligned}$$

□

**Lemma 2.5.** *Let  $N \subseteq M$  be a strongly Markov inclusion. For any  $x \in N' \cap M$ ,*

$$e_N x e_M = e_N \Gamma(x) e_M.$$

*Proof.* Taking the adjoint of the relation from Lemma 2.4 we have  $e_N x = e_N \gamma_0(x)$  with  $\gamma_0(x) \in M' \cap M_1$ . Applying Lemma 2.4 to the inclusion  $M \subseteq M_1$ , we have  $\gamma_0(x)e_M = \gamma_1(\gamma_0(x))e_M = \Gamma(x)e_M$ . The result follows.  $\square$

Lemma 2.4 implies that any unit vector  $\psi \in L^2(N, \tau)$  is a perfectly correlated/EPR state with respect to the commuting algebras  $N' \cap M$  and  $M' \cap M_1$ , meaning that any self-adjoint  $x \in N' \cap M$  has an ‘‘EPR double’’  $x' \in M' \cap M_1$  for which

$$\langle (x - x')^2 \psi, \psi \rangle = 0.$$

Indeed, by Lemma 2.4, for  $x' = \gamma_0(x)$ ,

$$\langle (x - x')^2 \psi, \psi \rangle = \|(x - x')\psi\|^2 = \|(x - x')e_N \psi\|^2 = 0.$$

For details on perfect correlation see [5] for the type I case and [90] for the general von Neumann algebraic setting (both works of course building on the seminal paper [37] of Einstein–Podolski–Rosen). More generally, if a unitary  $u$  belongs to the normaliser

$$\mathcal{N}_M(N) := \{u \in \mathcal{U}(M) \mid u^* N u = N\},$$

then  $u(N' \cap M)u^* = N' \cap M$  and  $u^* \psi$  is also an EPR state with respect to the same commuting algebras:

$$\gamma_0(uxu^*)u^* \psi = u^* \gamma_0(uxu^*) \psi = u^* \gamma_0(uxu^*) e_N \psi = u^*(uxu^*) \psi = xu^* \psi, \quad x \in N' \cap M. \quad (6)$$

In other words, the EPR double of  $x \in N' \cap M$  relative to the state  $u^* \psi$  is  $\gamma_0(uxu^*) \in M' \cap M_1$ . Moreover, the restricted vector state  $\omega_{u^* \psi}|_{N' \cap M}$  is tracial, which is often viewed as a form of maximal entanglement in the commuting operator framework (see e.g., [60, §V.A] or [29, §6]). Explicitly, for  $x, y \in N' \cap M$ , we have

$$\begin{aligned} \omega_{u^* \psi}(xy) &= \langle xyu^* \psi, u^* \psi \rangle = \langle x\gamma_0(yu^*)u^* \psi, u^* \psi \rangle \\ &= \langle \gamma_0(yu^*)xu^* \psi, u^* \psi \rangle = \langle xu^* \psi, \gamma_0(yu^*)u^* \psi \rangle \\ &= \langle xu^* \psi, y^* u^* \psi \rangle = \langle yxu^* \psi, u^* \psi \rangle. \end{aligned}$$

### 3. TELEPORTATION SCHEMES FOR SEMI-FINITE VON NEUMANN ALGEBRAS

The standard teleportation protocol fits naturally into the framework of the basic construction for  $\mathbb{C} \subseteq M_n(\mathbb{C})$ . Indeed, iterating the construction gives

$$\underbrace{\mathbb{C}}_N \subseteq \underbrace{M_n(\mathbb{C})}_M \subseteq \underbrace{M_n(\mathbb{C}) \otimes M_n(\mathbb{C})}_{M_1} \subseteq \underbrace{M_n(\mathbb{C}) \otimes M_n(\mathbb{C}) \otimes M_n(\mathbb{C})}_{M_2}.$$

From the operator algebraic perspective, the observable algebra of the global system of the protocol is  $M_2$ , while Alice and Bob’s observable algebras are  $M_1 \cong M_n(\mathbb{C}) \otimes M_n(\mathbb{C}) \otimes 1$  and  $M'_1 \cap M_2 \cong 1 \otimes 1 \otimes M_n(\mathbb{C})$ , respectively. The first Jones projection  $e_N \in M_1$  is the (rank-1 projection onto the) maximally entangled state  $\psi_n$ , while the second Jones projection  $e_M \in M_2$  is  $1 \otimes |\psi_n\rangle\langle\psi_n|$  (since  $M = M_n(\mathbb{C}) \otimes 1$  when viewed as a subalgebra of  $M_1 = M_n(\mathbb{C}) \otimes M_n(\mathbb{C})$ ). Note that  $e_M$  is precisely the entangled resource shared by Alice and Bob. Let  $B = \{u_i\}_{i=1}^{n^2}$  be any unitary orthonormal basis of  $M_n(\mathbb{C})$  with respect to the normalised Hilbert–Schmidt inner product. Then  $B$  is a Pimsner–Popa basis for  $\mathbb{C} \subseteq M_n(\mathbb{C})$ , and Alice’s local measurement in the associated teleportation protocol is with respect to the projection-valued measure (PVM)  $\{P_i := (u_i^* \otimes 1)e_N(u_i \otimes 1)\} \subseteq M_1$ . Bob’s operations are local conjugations by the  $u_i$  on the third tensor factor, i.e., automorphisms of  $M'_1 \cap M_2$ .

In the Heisenberg picture, the (standard) teleporation identity associated to the unitary basis  $B$  is

$$x = \sum_{i=1}^{n^2} (\text{id} \otimes \text{tr} \otimes \text{tr})((1 \otimes \psi_{\tau_n} \psi_{\tau_n}^*)(P_i \otimes u_i x u_i^*)), \quad x \in M_n(\mathbb{C}).$$



From this perspective, Bob's observable algebra  $M'_1 \cap M_2$  is teleported to Alice's local algebra  $M = M_n(\mathbb{C}) \otimes 1 \otimes 1 = N' \cap M$ . Recasting the above identity inside  $M_2$ ,

$$x \otimes 1 \otimes 1 = n^2 \sum_{i=1}^{n^2} E_M(e_M(P_i \otimes u_i x u_i^*)) = \Gamma^{-1}(1 \otimes 1 \otimes x),$$

where  $E_M = (\text{id} \otimes \tau_n \otimes \tau_n)$  is the unique trace-preserving conditional expectation from  $M_2$  onto  $M$ , and  $\Gamma$  is the canonical shift. Thus, from this perspective, the celebrated teleportation identity is an LOCC implementation of (the inverse of) the canonical shift associated to the inclusion  $\mathbb{C} \subseteq M_n(\mathbb{C})$ . This observation, together with the framework of [91] motivated the definition to follow. We first recall the commuting model of local operations and classical communication (LOCC) recently developed in [29].

Given two commuting von Neumann subalgebras  $A$  and  $B$  of an ambient von Neumann algebra  $M$ , a (one-way, right) LOCC operation is a normal, unital completely positive (UCP) map  $\Phi : M \rightarrow M$  of the form  $\Phi = \sum_{i=1}^{\infty} S_i \circ T_i$  (point weak\*-convergent), where  $S_i : M \rightarrow M$  is a normal CP  $B$ -bimodule map satisfying  $S_i(A) \subseteq A$ , and  $T_i : M \rightarrow M$  are normal UCP  $A$ -bimodule maps satisfying  $T_i(B) \subseteq B$ . (Note that the invariance conditions  $S_i(A) \subseteq A$  and  $T_i(B) \subseteq B$  were automatic in [29] since they considered  $M = \mathcal{B}(H)$ , and  $B = A'$ , in which case bimodularity implies invariance.) We also recall that  $A \vee B$  denotes the von Neumann algebra generated by  $A \cup B$ .

**Definition 3.1.** Let  $A$  and  $B$  be two commuting von Neumann subalgebras of a tracial von Neumann algebra  $(M, \tau)$ . Suppose that  $A$  contains von Neumann subalgebras  $A_0, A_1$ , for which there exist anti-isomorphisms  $\gamma_0 : A_0 \rightarrow A_1$  and  $\gamma_1 : A_1 \rightarrow B$ . A *teleportation scheme* for  $A_0$  relative to  $A, B \subseteq A \vee B$  consists of the following:

- a  $\tau$ -density operator  $\omega$  in  $A'_0 \cap M$ ;
- a collection  $\{T_i\}_{i \in I}$  of normal UCP  $A$ -bimodule maps on  $A \vee B$  for which  $T_i(B) \subseteq B$ ,  $i \in I$ .
- a POVM  $\{F_i\}_{i \in I}$  in  $A$  such that  $\sum_{i \in I} \text{Ad}(F_j) \circ T_i$  is a one-way right LOCC map relative to  $A, B \subseteq A \vee B$ , and

$$\sum_{i \in I} E_{A_0}(F_i T_i(\Gamma(a))\omega) = a, \quad a \in A_0, \quad (7)$$

where  $\Gamma : A_0 \rightarrow B$  is the  $*$ -isomorphism  $\Gamma = \gamma_1 \circ \gamma_0$ , and  $E_{A_0}$  is the  $\tau$ -preserving conditional expectation from  $M$  onto  $A_0$ , and the series converges in the weak\* topology.

The scheme  $(\omega, \{T_i\}_{i \in I}, \{F_i\}_{i \in I})$  is

- *faithful* if  $\tau(F_i \rho \omega) > 0$  for all  $i \in I$  and  $\tau$ -densities  $\rho \in A_0$ .
- *minimal* if  $\omega \in A_1 \vee B$  and  $\{F_i\}_{i \in I} \subseteq A_0 \vee A_1$ .

When  $|I| < \infty$ , the scheme  $(\omega, \{T_i\}_{i \in I}, \{F_i\}_{i \in I})$  is

- *tight* if  $\dim(A_0) = |I|$ ,
- *unbiased* if  $\tau(F_i \rho \omega) = |I|^{-1}$  for all  $i \in I$  and  $\tau$ -densities  $\rho \in A_0$ .

**Remark 3.2.**

- (1) As in [91], tightness means the amount of classical signals sent from Alice to Bob equals the dimension of the algebra to be teleported.
- (2) Minimality implies that the resource state  $\omega$  lives in the “smallest” algebra possessing a density which entangles  $A_1$  and  $B$ . Similar remark for Alice's POVM  $\{F_i\}$ .
- (3) The unbiasedness property ensures that for any input state  $\rho$  from  $A_0$ , Alice's local measurement result is uniformly random, so that no information about  $\rho$  is contained in the classical information sent to Bob.
- (4) Definition 3.1 readily generalises to semi-finite von Neumann algebras  $(M, \tau)$  with a normal conditional expectation onto  $A_0$ . For instance, when the restriction of the (normal semi-finite faithful) trace  $\tau$  to  $A_0$  is semi-finite (in which case there is a unique normal

$\tau$ -preserving conditional expectation from  $M$  onto  $A_0$  by Takesaki's theorem [82, Theorem IX.4.2]). One can also envision similarly defined schemes beyond semi-finite von Neumann algebras, although we will not pursue them in this paper.

**Example 3.3.** Let  $M = M_n(\mathbb{C}) \otimes M_n(\mathbb{C}) \otimes M_n(\mathbb{C})$ ,

$$A_0 = M_n(\mathbb{C}) \otimes 1 \otimes 1, \quad A_1 = 1 \otimes M_n(\mathbb{C}) \otimes 1, \quad B = 1 \otimes 1 \otimes M_n(\mathbb{C}).$$

We recover the standard teleportation scheme with

- $\omega = n^2(1 \otimes |\psi_n\rangle\langle\psi_n|) \in A_1 \vee B$  (the factor  $n^2$  is to ensure normalisation with respect to  $\tau_n \otimes \tau_n$ );
- $T_i = \text{Ad}(1 \otimes 1 \otimes u_i)$ , where  $\{u_i\}_{i=1}^{n^2}$  is any unitary orthonormal basis of  $M_n(\mathbb{C})$ ;
- $\{F_i = (u_i^* \otimes 1)|\psi_n\rangle\langle\psi_n|(u_i \otimes 1)\}_{i=1}^{n^2} \subseteq A_0 \vee A_1$ .

**Example 3.4.** Recently, Huang studied teleportation in a  $II_1$ -factor setting, showing the existence of certain (finite-dimensional) matrix subalgebras and analogues of the standard teleportation protocol for them [53]. We sketch how his protocol fits into our general framework, referring the reader to [53, §IV] for details.

Let  $M$  be a  $II_1$ -factor on  $H$  with cyclic and separating trace vector  $\psi \in H$ , that is,  $\tau(x) = \langle x\psi, \psi \rangle$ ,  $x \in M$  (where  $\tau$  is the unique trace of  $M$ ). Let  $N = M'$ , and let  $\gamma_1(x) = Jx^*J$  denote the canonical anti-isomorphism between  $M$  and  $N$ , where  $J$  is the modular conjugation associated to  $(M, \psi)$ . The global system is  $N \overline{\otimes} \mathcal{B}(H)$ , where Alice's observable algebra is  $N \otimes M$  and Bob's is  $1 \otimes N$ . For a natural number  $n \in \mathbb{N}$ , Huang takes a PVM  $\{P_j\}_{j=0}^{n-1}$  in  $M$  consisting of equivalent projections satisfying  $\tau(P_j) = \frac{1}{n}$  for all  $j$  (which exists in  $II_1$ -factors), and defines an associated family  $\{V_{j,k} \mid j, k = 0, \dots, n-1\} \subseteq M$  of partial isometries which form matrix units for a subalgebra  $A_1$  of  $M$  isomorphic to  $M_n(\mathbb{C})$  [53, Equation (6)]. Let  $W_{j,k} = \gamma(V_{k,j})$ ,

$$\Psi_{j,k} = \frac{1}{N} \sum_{\mu, \nu} e^{\frac{2\pi i k(\nu - \mu)}{n}} W_{\mu+j, \nu+j} \otimes V_{\mu, \nu}, \quad \text{and} \quad U_{j,k} = \sum_{\mu, \nu=0}^{n-1} e^{\frac{2\pi i k \nu}{n}} \delta_{\mu, \nu+j} V_{\mu, \nu},$$

where addition of the indicies is modulo  $n$ . Then  $\{\Psi_{j,k} \mid j, k = 0, \dots, n-1\}$  is PVM in  $N \otimes M$  (Alice's algebra) and  $U_{j,k}$  are unitaries in  $M$ . With  $A_0 = B = \text{span}\{W_{j,k} \mid j, k = 0, \dots, n-1\} \subseteq N$ ,

$$\gamma_0 : A_0 \otimes 1 \ni a \otimes 1 \mapsto 1 \otimes \gamma(a) \in 1 \otimes A_1,$$

and  $\Gamma = (\text{id} \otimes \gamma_1) \circ \gamma_0 : A_0 \otimes 1 \rightarrow 1 \otimes B$ , one can show that

$$\sum_{j,k=0}^{n-1} (\text{id} \otimes \omega_\psi)(\Psi_{j,k}(\text{id} \otimes \text{Ad}(\gamma(U_{j,k}))) (\Gamma(a))) = a, \quad a \in A_0. \quad (8)$$

Thus, Huang's protocol fits into our framework for semi-finite von Neumann algebras viewing  $(\text{id} \otimes \omega_\psi)$  as a normal conditional expectation  $N \overline{\otimes} \mathcal{B}(H) \rightarrow N$  (and further composing with the unique normal trace-preserving conditional expectation  $E_{A_0} : N \rightarrow A_0$  on either side of (8)).

Given the discussion at the beginning of the section, it is natural to consider teleportation schemes for more general inclusions  $N \subseteq M$  in connection with the basic construction, in particular, when  $N$  and/or  $M$  are not necessarily factors. We first examine the natural "direct sum protocol" when  $N = \mathbb{C}$ , and  $M$  is finite-dimensional.

Suppose  $M \cong \bigoplus_{j=1}^m M_{n_j}(\mathbb{C})$ . The (unique) Markov trace  $\tau$  for the inclusion  $\mathbb{C} \subseteq M$  is

$$\tau = \frac{1}{\dim M} \sum_{j=1}^m n_j \text{tr}_{n_j},$$



where  $\text{tr}_{n_j}$  is the unnormalized trace on  $M_{n_j}(\mathbb{C})$ . It follows that  $L^2(M, \tau) \cong \bigoplus_{j=1}^m \mathbb{C}^{n_j} \otimes \mathbb{C}^{n_j}$ , and that  $M$  is represented on  $L^2(M, \tau)$  as  $M = \bigoplus_{j=1}^m M_{n_j}(\mathbb{C}) \otimes 1_{n_j}$ . The anti-isomorphism  $\gamma_0 : M \rightarrow M'$  is then simply

$$\gamma_0(\bigoplus_j x_j \otimes 1_{n_j}) = \bigoplus_j 1_{n_j} \otimes x_j^t,$$

where  $t$  denotes transposition.

Since  $M_1 = \langle M, e_{\mathbb{C}} \rangle = \mathcal{B}(L^2(M, \tau))$ , the extended trace  $\tau_1$  is the unique normalised trace, and  $L^2(M_1, \tau_1) \cong L^2(M, \tau) \otimes L^2(M, \tau)$ . The conjugation  $J_1$  is then the tensor flip (plus complex conjugation) and we have

$$M_2 = J_1 M' J_1 = J_1(M' \otimes \mathcal{B}(L^2(M, \tau))) J_1 = \mathcal{B}(L^2(M, \tau)) \otimes M'.$$

In the Proposition below, we view  $M_2 = \mathcal{B}(L^2(M, \tau)) \otimes M'$  as a tripartite system, consisting of the (anti-)isomorphic commuting subalgebras  $M \otimes 1$ ,  $M' \otimes 1$  and  $1 \otimes M'$ . Alice has access to the first two, and Bob has access to the third. Here, the pertinent trace is  $\tau_2 := \tau_1 \otimes \tau'$ , where  $\tau'(y) = \tau(Jy^*J)$ ,  $y \in M'$ .

**Proposition 3.5.** *Let  $M$  be a finite-dimensional  $C^*$ -algebra, and  $M_1$  and  $M_2$  be the result of the iterated basic construction of the inclusion  $\mathbb{C} \subseteq M$  with respect to the Markov trace  $\tau$ . Then there exists a tight, minimal teleportation scheme  $(\omega, \{T_i\}_{i=1}^d, \{F_i\}_{i=1}^d)$  for  $M$  relative to*

$$\mathcal{B}(L^2(M, \tau)) \otimes 1, 1 \otimes M' \subseteq (M_2, \tau_2).$$

*Proof.* As above, we take  $M \cong \bigoplus_{j=1}^m M_{n_j}(\mathbb{C})$  so that  $M$  is represented on  $L^2(M, \tau)$  as  $M = \bigoplus_{j=1}^m M_{n_j}(\mathbb{C}) \otimes 1_{n_j}$ .

Let  $W_j : \mathbb{Z}_{n_j} \times \mathbb{Z}_{n_j} \rightarrow \mathcal{U}(M_{n_j}(\mathbb{C}))$  denote the Weyl representation of  $\mathbb{Z}_{n_j} \times \mathbb{Z}_{n_j}$ , and  $|\psi_{n_j}\rangle$  denote the maximally entangled state in  $\mathbb{C}^{n_j} \otimes \mathbb{C}^{n_j}$  so that  $|\psi_{n_j}\rangle\langle\psi_{n_j}|$  is the Jones projection for the inclusion  $\mathbb{C} \subseteq M_{n_j}(\mathbb{C})$  (cf. Example 2.1). Define  $F_{j,z_j} \in M \vee M' \subseteq M_1$  by

$$F_{j,z_j} = 0_{n_1} \oplus \dots \oplus 0_{n_{j-1}} \oplus (W_j(z_j)^* \otimes 1_{n_j}) |\psi_{n_j}\rangle\langle\psi_{n_j}| (W_j(z_j) \otimes 1_{n_j}) \oplus 0_{n_{j+1}} \oplus \dots \oplus 0_{n_m},$$

for  $j = 1, \dots, n$  and  $z_j \in \mathbb{Z}_{n_j}^2$ . It follows that,  $\sum_{z_j \in \mathbb{Z}_{n_j}^2} F_{j,z_j} = 1_{n_j} \otimes 1_{n_j}$  for each  $j = 1, \dots, n$ . Hence,  $\{F_{j,z_j} \mid j = 1, \dots, n, z_j \in \mathbb{Z}_{n_j}^2\}$  is a POVM in  $M_1$  with cardinality  $d := \dim M$ . Put

$$\tilde{W}_j(z_j) = 1_{n_1} \oplus \dots \oplus 1_{n_{j-1}} \oplus W_j(z_j) \oplus 1_{n_{j+1}} \oplus \dots \oplus 1_{n_m},$$

which is a unitary in  $M$ , and define the automorphism  $T_{j,z_j} : M_2 \rightarrow M_2$  by

$$T_{j,z_j}(y) = \Gamma(\tilde{W}_j(z_j)) y \Gamma(\tilde{W}_j(z_j)^*), \quad y \in M_2.$$

Clearly, each  $T_{j,z_j}$  leaves  $M'_1 \cap M_2 = \Gamma(M)$  invariant, and is an  $A$ -bimodule map, where  $A = \mathcal{B}(L^2(M, \tau)) \otimes 1$ .

Finally, let  $\omega := (\dim M)e_M$ . Then  $\omega \in M' \cap M_2 = M' \otimes M'$ . Noting that  $\dim M = [M : \mathbb{C}]$ , the Markov property (1) implies that

$$\tau_2(\omega) = \dim M (\tau_1 \otimes \tau')(e_M) = (\dim M) \tau_1(E_{M_1}(e_M)) = \tau_1(1) = 1,$$

so  $\omega$  is a  $\tau_2$ -density. We show that

$$x = E_M \left( \sum_{j=1}^m \sum_{z_j \in \mathbb{Z}_{n_j}^2} F_{j,z_j} T_{j,z_j}(\Gamma(x)) \omega \right), \quad x \in M,$$

where  $E_M \equiv E_{M_1 \rightarrow M} \circ E_{M_2 \rightarrow M_1}$  is the unique trace preserving conditional expectation from  $M_2$  to  $M$ , and  $\Gamma$  is the canonical shift from  $M$  to  $M'_1 \cap M_2$ . This will establish the Proposition, as tightness and minimality are clear by construction.

First consider the  $T_{j,z_j}(\Gamma(x))e_M$  term in the above equation (pulling the constant off  $\omega$ ).

$$T_{j,z_j}(\Gamma(x))e_M = \Gamma\left(\tilde{W}_j(z_j)x\tilde{W}_j(z_j)^*\right)e_M = \gamma_0\left(\tilde{W}_j(z_j)x\tilde{W}_j(z_j)^*\right)e_M,$$

the last equality following from Lemma 2.4. Write  $x = \bigoplus_j(x_j \otimes 1_{n_j})$ . Then multiplying the above by  $F_{j,z_j}$ , we see that

$$\begin{aligned} & F_{j,z_j}\gamma_0\left(\tilde{W}_j(z_j)x\tilde{W}_j(z_j)^*\right)e_M \\ &= F_{j,z_j}\gamma_0\left((x_1 \otimes 1_{n_1}) \oplus \cdots \oplus (W_j(z_j)x_jW_j(z_j)^* \otimes 1_{n_j}) \oplus \cdots \oplus (x_m \otimes 1_{n_m})\right)e_M \\ &= \left((W_j(z_j)^* \otimes 1_{n_j})|\psi_{n_j}\rangle\langle\psi_{n_j}|(W_j(z_j) \otimes 1_{n_j})\right)\left(1_{n_j} \otimes \overline{W_j(z_j)}x_j^tW_j(z_j)^t\right)e_M \\ &= \left((W_j(z_j)^* \otimes 1_{n_j})|\psi_{n_j}\rangle\langle\psi_{n_j}|(1_{n_j} \otimes \overline{W_j(z_j)}x_j^tW_j(z_j)^t)(W_j(z_j) \otimes 1_{n_j})\right)e_M \\ &= (W_j(z_j)^* \otimes 1_{n_j})|\psi_{n_j}\rangle\langle\psi_{n_j}|((\overline{W_j(z_j)}x_j^tW_j(z_j)^t)^tW_j(z_j) \otimes 1_{n_j})e_M \quad (\text{Lemma 2.4 with } |\psi_{n_j}\rangle\langle\psi_{n_j}|) \\ &= (W_j(z_j)^* \otimes 1_{n_j})|\psi_{n_j}\rangle\langle\psi_{n_j}|(W_j(z_j)x_j \otimes 1_{n_j})e_M \\ &= F_{j,z_j}xe_M \end{aligned}$$

Hence,

$$\begin{aligned} E_M\left(\sum_{j=1}^m \sum_{z_j \in \mathbb{Z}_{n_j}^2} F_{j,z_j}T_{j,z_j}(\Gamma(x))\omega\right) &= \dim ME_M\left(\sum_{j=1}^m \sum_{z_j \in \mathbb{Z}_{n_j}^2} F_{j,z_j}xe_M\right) \\ &= x \dim ME_M(e_M) \\ &= x, \end{aligned}$$

where the last equality uses the Markov property

$$E_M(e_M) = E_M(E_{M_1}(e_M)) = \dim M^{-1} = [M : \mathbb{C}]^{-1}. \quad \square$$

**Remark 3.6.** Proposition 3.5 shows that the ‘‘direct sum’’ of the standard protocols allows one to teleport any finite-dimensional algebra  $M$ . Moreover, the scheme is tight in the sense that the number of classical messages sent by Alice to Bob coincides with the dimension of the algebra to be teleported. However, one drawback of this approach is that Alice’s measurement results are correlated with the location of the state within  $M = \bigoplus_j M_{n_j}(\mathbb{C})$ . Indeed, if  $\rho \in M$  is a density living entirely in one summand, say  $M_{n_j}(\mathbb{C})$ , then the probability of Alice measuring outcome  $(i, z_i)$ ,  $i \neq j$ , is  $\tau(F_{i,z_i}\rho\omega) = 0$ .

It could be desirable that Alice’s measurement result, i.e., the classical information sent to Bob, contains no information about the state to be teleported. This is precisely the unbiased condition, and one can imagine this feature to be important in potential applications of the teleportation schemes introduced here; for instance, in the context of quantum privacy applications such as those considered in [2, 10, 11, 19, 26, 30, 27, 38, 55, 64, 66]. If  $\mathbb{C} \subseteq M$  admits an orthonormal basis (with respect to the Markov trace) consisting of unitaries, then we can obtain an unbiased scheme. In fact, the same is true for more general inclusions  $N \subseteq M$ .

**Theorem 3.7.** *Let  $N \subseteq M$  be a strongly Markov inclusion of tracial von Neumann algebras. Suppose there is an orthonormal Pimsner–Popa basis  $\{u_i\}_{i=1}^d$  for  $M/N$  inside the normaliser  $\mathcal{N}_M(N)$ , that  $N' \cap M$  is finite-dimensional, and that  $N' \cap M \subseteq L \subseteq N'$  for some injective factor  $L$ . Then there exists an unbiased teleportation scheme  $(\omega, \{F_i\}_{i=1}^d, \{T_i\}_{i=1}^d)$  for  $N' \cap M$ , with respect to  $M_1, M'_1 \cap M_2 \subseteq M_1 \vee (M'_1 \cap M_2)$ .*

In the proof, the following Lemma will be used to achieve the locality condition between Alice's and Bob's operations in Definition 3.1. It is essentially an application of the periodicity of the tower of the basic construction (see, e.g., [76, Proposition 1.5], or [57, Proposition 4.3.7]). We include details for the convenience of the reader (and since we do not restrict to  $II_1$  factors).

**Lemma 3.8.** *Let  $N \subseteq M$  be a strongly Markov inclusion of tracial von Neumann algebras. Suppose there is an orthonormal Pimsner–Popa basis  $\{u_i\}_{i=1}^d$  for  $M$  over  $N$  in the normaliser  $\mathcal{N}_M(N)$ . Then there exist unitaries  $\{v_i\}_{i=1}^d \subseteq M_2$  satisfying*

$$v_i \Gamma(x) v_i^* = \Gamma(u_i x u_i^*), \quad i = 1, \dots, d, \quad x \in N' \cap M.$$

*If, in addition,  $N' \cap M$  is finite-dimensional and  $N' \cap M \subseteq L \subseteq N'$  for some injective factor  $L$ , then  $\text{Ad}(v_i)|_{M'_1 \cap M_2}$  extends to a  $M_1$ -bimodule  $*$ -automorphism of  $M_1 \vee (M'_1 \cap M_2)$ .*

*Proof.* Define  $\varphi : M_d(M) \rightarrow M_2$  by

$$\varphi([x_{i,j}]) = [M : N] \sum_{i,j=1}^d u_i^* e_N x_{i,j} e_M e_N u_j, \quad x \in M_d(M).$$

Using orthonormality of  $\{u_i\}$  together with the relations  $e_M(\cdot)e_M = E_M(\cdot)e_M$  and  $e_N(\cdot)e_N = E_N(\cdot)e_N$  on  $M_1$  and  $M$ , respectively, it follows that  $\varphi$  is multiplicative:

$$\begin{aligned} \varphi([x_{i,j}])\varphi([y_{i,j}]) &= [M : N]^2 \sum_{i,j,k,l=1}^d u_i^* e_N x_{i,j} e_M (e_N u_j u_k^* e_N y_{k,l}) e_M e_N u_l \\ &= [M : N]^2 \sum_{i,j,k,l=1}^d u_i^* e_N x_{i,j} E_M(e_N u_j u_k^* e_N y_{k,l}) e_M e_N u_l \\ &= [M : N]^2 \sum_{i,j,k,l=1}^d u_i^* e_N x_{i,j} E_M(E_N(u_j u_k^*) e_N) y_{k,l} e_M e_N u_l \\ &= [M : N]^2 \sum_{i,j,l=1}^d u_i^* e_N x_{i,j} E_M(e_N) y_{j,l} e_M e_N u_l \\ &= [M : N] \sum_{i,j,l=1}^d u_i^* e_N x_{i,j} y_{j,l} e_M e_N u_l \\ &= \varphi([x_{i,j}][y_{k,l}]). \end{aligned}$$

Also, as  $e_M \in M'$ ,

$$\varphi([x_{i,j}])^* = [M : N] \sum_{i,j=1}^d u_j^* e_N e_M x_{i,j}^* e_N u_i = [M : N] \sum_{i,j=1}^d u_i^* e_N x_{j,i}^* e_M e_N u_j = \varphi([x_{i,j}]^*).$$

Also, the relation  $e_N e_M e_N = [M : N]^{-1} e_N$  (see (2)) together with the basis property implies that  $\varphi$  is unital.

For each  $i = 1, \dots, d$ , define  $v_i := \varphi(\text{diag}(u_i, u_i, \dots, u_i)) \in M_2$ . Then

$$v_i = [M : N] \sum_{j=1}^d u_j^* e_N u_i e_M e_N u_j$$

is unitary as the image of a unitary in  $M_d(M)$  under a unital  $*$ -homomorphism.

Let  $x \in N' \cap M$ . By Lemma 2.5 we have  $e_N \Gamma(x) e_M = e_N x e_M$ . Also, by orthonormality and the fact that  $\Gamma(x) \in M'_1 \cap M_2$ , for each  $j, k = 1, \dots, d$  we have

$$e_N u_j \Gamma(x) u_k^* e_N = E_N(u_j u_k^*) e_N \Gamma(x) = \delta_{j,k} e_N \Gamma(x).$$

Thus,

$$\begin{aligned} v_i \Gamma(x) v_i^* &= [M : N]^2 \sum_{j,k=1}^d u_j^* e_N u_i e_M (e_N u_j \Gamma(x) u_k^* e_N) e_M u_i^* e_N u_k \\ &= [M : N]^2 \sum_{j=1}^d u_j^* e_N u_i e_M (e_N \Gamma(x) e_M) u_i^* e_N u_j \\ &= [M : N]^2 \sum_{j=1}^d u_j^* e_N u_i e_M (e_N x e_M) u_i^* e_N u_j \\ &= [M : N]^2 \sum_{j=1}^d u_j^* e_N u_i (e_M e_N e_M) x u_i^* e_N u_j \\ &= [M : N] \sum_{j=1}^d u_j^* e_N (u_i e_M) x u_i^* e_N u_j \\ &= [M : N] \sum_{j=1}^d u_j^* e_N e_M (u_i x u_i^*) e_N u_j \\ &= [M : N] \sum_{j=1}^d u_j^* e_N e_M \Gamma(u_i x u_i^*) e_N u_j \\ &= [M : N] \left( \sum_{j=1}^d u_j^* e_N e_M e_N u_j \right) \Gamma(u_i x u_i^*) \\ &= \left( \sum_{j=1}^d u_j^* e_N u_j \right) \Gamma(u_i x u_i^*) \\ &= \Gamma(u_i x u_i^*). \end{aligned}$$

Now, suppose, in addition, that  $N' \cap M$  is finite-dimensional and  $N' \cap M \subseteq L \subseteq N'$  for some injective factor  $L$ . Then  $L_0 := J_1 J_0(L) J_0 J_1 \subseteq M'_1$  is an injective factor such that  $M'_1 \cap M_2 \subseteq L_0 \subseteq M'_1$ . Multiplication induces a  $*$ -isomorphism  $m : L'_0 \otimes_{\min} L_0 \cong C^*(L'_0, L_0)$ , the  $C^*$ -subalgebra of  $\mathcal{B}(L^2(M_1, \tau_1))$  generated by  $L_0$  and  $L'_0$  [35, Corollary 4.6]. By injectivity of the minimal tensor product,  $M_1 \otimes_{\min} (M'_1 \cap M_2) \subset L'_0 \otimes_{\min} L_0$ . It follows that the restriction of  $m$  induces a  $*$ -isomorphism

$$m : M_1 \otimes_{\min} (M'_1 \cap M_2) \cong C^*(M_1, M'_1 \cap M_2).$$

Note also that by finite-dimensionality of  $M'_1 \cap M_2 = \Gamma(N' \cap M)$ ,

$$M_1 \otimes_{\min} (M'_1 \cap M_2) = M_1 \overline{\otimes} (M'_1 \cap M_2),$$

where  $\overline{\otimes}$  is the von Neumann tensor product. Since  $M_1$  is weakly closed and commutes with the finite-dimensional  $C^*$ -algebra  $M'_1 \cap M_2$ , by considering a system of matrix units for the latter algebra we can see that  $C^*(M_1, M'_1 \cap M_2)$  is weakly closed, and hence coincides with  $M_1 \vee (M'_1 \cap M_2)$ . The desired extension of  $\text{Ad}(v_i)$  is then the composition

$$M_1 \vee (M'_1 \cap M_2) \cong M_1 \overline{\otimes} (M'_1 \cap M_2) \xrightarrow{(\text{id} \otimes \text{Ad}(v_i))} M_1 \overline{\otimes} (M'_1 \cap M_2) \cong M_1 \vee (M'_1 \cap M_2).$$

Note that the extension is necessarily an  $M_1$ -bimodule  $*$ -automorphism.  $\square$

*Proof of Theorem 3.7.* Let  $\omega := [M : N]e_M$ . Then  $\omega$  is a  $\tau$ -density in  $M' \cap M_2$  by the Markov property. For each  $i = 1, \dots, d$ , let  $F_i = u_i^* e_N u_i$ . Then  $\{F_i\}_{i=1}^d$  is a PVM in  $N' \cap M_1$ . Finally, let  $v_i := \varphi(\text{diag}(u_i, u_i, \dots, u_i)) \in M_2$  be the unitary associated to  $u_i$  from Lemma 3.8, and define  $T_i$  to be the  $M_1$ -bimodule extension of  $\text{Ad}(v_i)|_{M'_1 \cap M_2}$  to  $M_1 \vee (M'_1 \cap M_2)$ . It follows that  $\sum_{i=1}^d \text{Ad}(F_i) \circ T_i$  is a one-way (right) LOCC operation relative to  $M_1, M'_1 \cap M_2 \subseteq M_1 \vee (M'_1 \cap M_2)$ .

Fix  $x \in N' \cap M$ . We will show that

$$x = \sum_{i=1}^d E_{N' \cap M}(F_i T_i(\Gamma(x))\omega),$$

so that  $(\omega, \{F_i\}_{i=1}^d, \{T_i\}_{i=1}^d)$  forms a teleportation scheme. By Lemma 3.8

$$T_i(\Gamma(x)) = v_i \Gamma(x) v_i^* = F_i \Gamma(u_i x u_i^*).$$

Applying Lemma 2.4 twice, we see that

$$\begin{aligned} E_{N' \cap M}(F_i T_i(\Gamma(x))\omega) &= [M : N] E_{N' \cap M}(F_i T_i(\Gamma(x))e_M) \\ &= [M : N] E_{N' \cap M}(F_i \Gamma(u_i x u_i^*)e_M) \\ &= [M : N] E_{N' \cap M}(F_i \gamma_0(u_i x u_i^*)e_M) \\ &= [M : N] E_{N' \cap M}(u_i^* e_N u_i \gamma_0(u_i x u_i^*)e_M) \\ &= [M : N] E_{N' \cap M}(u_i^* e_N \gamma_0(u_i x u_i^*)u_i e_M) \\ &= [M : N] E_{N' \cap M}(u_i^* e_N (u_i x u_i^*)u_i e_M) \\ &= [M : N] E_{N' \cap M}(u_i^* e_N u_i x e_M) \\ &= [M : N] E_{N' \cap M}(F_i e_M)x. \end{aligned}$$

Summing over  $i$ , and applying the Markov property

$$E_{N' \cap M}(e_M) = E_{N' \cap M}(E_{M_1}(e_M)) = [M_1 : M]^{-1} 1 = [M : N]^{-1} 1$$

gives the identity.

It remains to show the unbiased property, that is,  $\tau(F_i \rho \omega) = [M : N]^{-1}$  for all  $i$  and  $\tau$ -densities  $\rho \in N' \cap M$ . For any such  $\rho$ ,  $F_i \rho \in M_1$ , so that  $\tau(F_i \rho e_M) = [M : N]^{-1} \tau(F_i \rho)$ . Thus,

$$\begin{aligned} \tau(F_i \rho \omega) &= [M : N] \tau(F_i \rho e_M) = \tau(u_i^* e_N u_i \rho) \\ &= \tau(e_N(u_i \rho u_i^*)) = [M : N]^{-1} \tau(u_i \rho u_i^*) \\ &= \frac{1}{[M : N]}. \end{aligned}$$

$\square$

**Remark 3.9.** The unbiased scheme in Theorem 3.7 is not necessarily tight. For example, tightness would imply  $[M : N] = \dim(N' \cap M)$ , which, for a finite-index inclusion  $N \subseteq M$  of  $II_1$ -factors, means that the inclusion has depth 1 [43, Theorem 4.6.3(vii)].

The main assumption in Theorem 3.7 was the existence of a Pimsner–Popa basis  $\{u_i\}$  for  $M$  over  $N$  inside the normaliser  $\mathcal{N}_M(N)$ . In this case, the completeness relation  $\sum_i u_i^* e_N u_i = 1$  corresponds to the decomposition

$$L^2(M, \tau) = \bigoplus_i u_i^* L^2(N, \tau) \quad (9)$$

of  $L^2(M, \tau)$  into maximally entangled subspaces with respect to the commuting subsystems  $N' \cap M$  and  $M' \cap M_1$  (see Section 2.3). When  $N = \mathbb{C} \subseteq M_n(\mathbb{C}) = M$ , the normaliser assumption holds

trivially, and the decomposition (9) simply corresponds to an orthonormal basis of maximally entangled vectors of  $L^2(M_n(\mathbb{C}), \tau_n) = \mathbb{C}^n \otimes \mathbb{C}^n$ , i.e., Alice's local measurement in teleportation. Moreover, the normalisation property implies that the decomposition (9) is one of  $N$ -bimodules, suggesting a potential connection with categorical approaches to teleportation [3], which we leave for future investigations (see Outlook section).

**Example 3.10.** Recall that the hypotheses of Theorem 3.7 require the existence of an orthonormal Pimsner-Popa basis in the normaliser. This holds in the following cases:

- (1) Any inclusion  $M \subseteq G \rtimes M$ , where  $G$  is a finite group with a trace-preserving action on a tracial von Neumann algebra  $M$  (see Example 2.3).
- (2) Any finite-index regular inclusion  $N \subseteq M$  of  $II_1$ -factors for which  $N' \cap M$  is either simple or commutative [8, 9]. (Follow the proof of [9, Theorem 3.21].) If [9, Conjecture 3.20] is true, the same holds for any regular inclusion of finite index.
- (3) Any inclusion  $N \subseteq M_n(\mathbb{C})$  with  $N = \bigoplus_{j=1}^k M_l(\mathbb{C})$  homogeneous. (This may be seen as a special case of (1).) Homogeneous algebras of this type can be used to model hybrid quantum codes [15, 24, 46, 71, 72], so our scheme from Theorem 3.7 could theoretically be used to teleport hybrid quantum codes in an unbiased manner.

We end this section with a partial converse of Example 3.10 (3).

**Proposition 3.11.** *A multiplicity-free inclusion  $N \subseteq M_n(\mathbb{C})$  admits a Pimsner-Popa basis in  $\mathcal{N}_{M_n(\mathbb{C})}(N)$  if and only if  $N$  is homogeneous.*

*Proof.* If  $N \subseteq M_n(\mathbb{C})$  is multiplicity free and  $N$  is homogeneous, then  $n = kl$  and  $N = \bigoplus_{j=1}^k M_l(\mathbb{C}) \cong \ell_k^\infty \otimes M_l(\mathbb{C})$ . The existence of the required basis follows from Example 3.10 (3).

Suppose  $N = \bigoplus_{j=1}^k M_{n_j}(\mathbb{C}) \subseteq M_n(\mathbb{C})$  is a multiplicity-free inclusion. Let  $z_1, \dots, z_k$  be the minimal central projections of  $N$ , indexed such that  $z_j N \cong M_{n_j}(\mathbb{C})$ . It follows that  $E_N = \sum_{j=1}^k z_j(\cdot)z_j$  and that  $e_N = \sum_{j=1}^k z_k \otimes z_k \in \mathcal{B}(\mathbb{C}^n \otimes \mathbb{C}^n)$ .

Let  $\{\lambda_i\}_{i=1}^d$  be any Pimsner-Popa basis for  $M_n(\mathbb{C})/N$ . Then

$$\begin{aligned} 1_n \otimes 1_n &= \sum_{i=1}^d (\lambda_i^* \otimes 1_n) e_N (\lambda_i \otimes 1_n) \\ &= \sum_{i=1}^d (\lambda_i^* \otimes 1_n) \left( \sum_{j=1}^k z_j \otimes z_j \right) (\lambda_i \otimes 1_n) \\ &= \sum_{j=1}^k \left( \sum_{i=1}^d \lambda_i^* z_j \lambda_i \right) \otimes z_j. \end{aligned}$$

If each  $\lambda_i$  is unitary, then we can multiply by  $1_n \otimes z_j$  and take unnormalised traces to obtain

$$nn_j = \text{tr}(1_n \otimes z_j) = \text{tr} \left( \left( \sum_{i=1}^d \lambda_i^* z_j \lambda_i \right) \otimes z_j \right) = dn_j^2,$$

so  $n_j = n/d$  for every  $j$ , hence  $N$  is homogeneous.  $\square$

#### 4. RIGIDITY OF TELEPORTATION FOR FINITE-DIMENSIONAL INCLUSIONS

Werner [91] established a one-to-one correspondence between tight teleportation schemes for the tripartite system  $M_n(\mathbb{C}) \otimes M_n(\mathbb{C}) \otimes M_n(\mathbb{C})$  and orthonormal bases of unitaries of  $M_n(\mathbb{C})$  (a.k.a, unitary error bases). In this section we generalize Werner's result to inclusions of the form  $N \subseteq M_n(\mathbb{C})$ .



**Theorem 4.1.** *Let  $N \subseteq (M_n(\mathbb{C}), \tau)$  be an inclusion such that  $\tau|_{N'}$  is the Markov trace for  $\mathbb{C} \subseteq N'$ . Put  $A_0 := N' \otimes 1 \otimes 1$ ,  $A_1 := 1 \otimes N' \otimes 1$  and  $B = 1 \otimes 1 \otimes N'$ . If  $(\omega, \{F_i\}_{i=1}^d, \{T_i\}_{i=1}^d)$  is a tight, minimal, faithful teleportation scheme for  $A_0$ , with respect to*

$$M_n(\mathbb{C}) \otimes M_n(\mathbb{C}) \otimes 1, 1 \otimes 1 \otimes N' \subseteq M_n(\mathbb{C}) \otimes M_n(\mathbb{C}) \otimes N',$$

*then there exist (1) an orthonormal basis  $\{u_i\}_{i=1}^d$  of  $M_n(\mathbb{C})$  over  $N$  in  $\mathcal{N}_{M_n(\mathbb{C})}(N)$ , (2) a unitary  $u \in \mathcal{N}_{M_n(\mathbb{C})}(N)$ , (3) a positive invertible operator  $z \in \mathcal{Z}(N)$  such that*

- $\omega = [M_n(\mathbb{C}) : N](1 \otimes z^{1/2}u)e_N(1 \otimes u^*z^{1/2})$ ;
- $F_i = (u_i^*u \otimes 1)e_N(u^*u_i \otimes 1)$ ,  $i = 1, \dots, d$ ;
- $T_i(x) = u_i x u_i^*$ ,  $x \in N'$ .

The proof requires several preparations. We follow the same general strategy as in [91], although our setting does not allow us to work at the level of individual Kraus operators. Instead, we mostly argue at the level of CP maps, which, incidentally, allows us to circumnavigate (the analogue of) [91, Proposition 3], albeit we begin with the a priori stronger assumption of faithfulness.

The first Lemma is a simple (known) observation.

**Lemma 4.2.** *Let  $N \subseteq (M, \tau)$  be an inclusion of finite von Neumann algebras. Write  $\mathcal{N}_M(N) := \{u \in \mathcal{U}(M) \mid u^*Nu = N\}$ , and  $\mathcal{N}_M(E_N) := \{u \in \mathcal{U}(M) \mid u^*E_N(x)u = E_N(u^*xu), x \in M\}$ . Then we have the equality*

$$\mathcal{N}_M(N) = \mathcal{N}_M(E_N).$$

Moreover, any  $u \in \mathcal{N}_M(N)$  satisfies

$$ue_Nu^* = \gamma_0(u)e_N\gamma_0(u^*).$$

*Proof.* The inclusion  $\mathcal{N}_M(E_N) \subseteq \mathcal{N}_M(N)$  follows easily from the definitions. Given  $u \in \mathcal{N}_M(N)$ , it follows that  $\text{Ad}(u) \circ E_N \circ \text{Ad}(u^*)$  is a  $\tau$ -preserving conditional expectation from  $M$  to  $N$ . By uniqueness,  $\text{Ad}(u) \circ E_N \circ \text{Ad}(u^*) = E_N$ , which is the desired normalisation property.

If  $u \in \mathcal{N}_M(N) = \mathcal{N}_M(E_N)$ , then for all  $x \in M$ ,

$$\begin{aligned} ue_Nu^*\Lambda_\tau(x) &= \Lambda_\tau(ue_N(u^*x)) = \Lambda_\tau(ue_N(u^*x)u^*u) = \Lambda_\tau(E_N(xu^*)u) \\ &= \gamma_0(u)\Lambda_\tau(E_N(xu^*)) = \gamma_0(u)e_N\Lambda_\tau(xu^*) \\ &= \gamma_0(u)e_N\gamma_0(u^*)\Lambda_\tau(x). \end{aligned}$$

□

The next two Lemmas generalise [91, Lemma 7] and [91, Lemma 2], respectively.

**Lemma 4.3.** *Let  $M$  be a finite-dimensional  $C^*$ -algebra and let  $\{T_i\}_{i=1}^d$  be a family of CP maps  $T_i : M \rightarrow M$  such that  $\sum_{i=1}^d T_i = \text{id}_M$ . If  $M = \bigoplus_{j=1}^m z_j M$ , where  $z_1, \dots, z_m$ , are the minimal central projections of  $M$ , then for each  $i, j$  there exists  $\mu_i^j \in [0, \infty)$  such that*

$$T_i|_{z_j M} = \mu_i^j \text{id}_{z_j M}.$$

*Proof.* Let  $p$  a minimal projection of  $M$ . Then,

$$0 \leq T_i(p) \leq \sum_i T_i(p) = p,$$

$T_i(p) = pT_i(p)p \in pMp = \mathbb{C}p$  by minimality. It follows that  $T_i(z_j M) \subseteq z_j M$ , for each  $i$  and  $j$ . Then for each  $j$ ,  $T_i|_{z_j M} = z_j M \rightarrow z_j M$  is CP and  $\sum_i T_i|_{z_j M} = \text{id}_{z_j M}$ . By [91, Lemma 7], each  $T_i|_{z_j M} = \mu_i^j \text{id}_{z_j M}$ . □

**Remark 4.4.** The coefficients  $\mu_i^j$  depend on the summand  $j$ , in general, and can be zero even if  $T_i \neq 0$ . For instance, consider  $T_1, T_2 : \ell_2^\infty \rightarrow \ell_2^\infty$  given by

$$T_1(x, y) = (2^{-1}x, 0), \quad T_2(x, y) = (2^{-1}x, y), \quad (x, y) \in \ell_2^\infty.$$

Then  $T_1$  and  $T_2$  are CP and  $T_1 + T_2 = \text{id}_{\ell_2^\infty}$ . The associated coefficients are  $\mu_1^1 = \frac{1}{2}$ ,  $\mu_1^2 = 0$ ,  $\mu_2^1 = \frac{1}{2}$ ,  $\mu_2^2 = 1$ .

**Lemma 4.5.** *Let  $N \subseteq (M, \tau)$  be a strongly Markov inclusion of finite-dimensional von Neumann algebras. A basis  $\{\lambda_i\}_{i=1}^d$  for  $M$  over  $N$  is orthonormal (i.e.,  $E_N(\lambda_i \lambda_j^*) = \delta_{i,j} 1$ ) if and only if,*

$$d = \frac{\dim M}{\dim N}.$$

*Proof.* Assume  $d = \dim M / \dim N$ , and let  $\{e_n \mid n = 1, \dots, \dim N\}$  be an orthonormal basis for  $L^2(N, \tau)$ . Then  $e_N = \sum_{n=1}^{\dim N} |e_n\rangle\langle e_n|$ , so that

$$1_{L^2(M, \tau)} = \sum_{i=1}^d \lambda_i^* e_N \lambda_i = \sum_{i=1}^d \sum_{n=1}^{\dim N} \lambda_i^* |e_n\rangle\langle e_n| \lambda_i.$$

Since  $d \dim N = \dim M = \dim(L^2(M, \tau))$ , by [91, Lemma 2], the set  $\{\lambda_i^* |e_n\rangle \mid i = 1, \dots, d; n = 1, \dots, \dim N\}$ , is an orthonormal basis of  $L^2(M, \tau)$ . In particular, the subspaces  $\lambda_i^* L^2(N, \tau)$  are orthogonal,  $\{\lambda_i^* e_N \lambda_i\}_{i=1}^d$  are mutually orthogonal projections and each  $\lambda_i^*$  acts isometrically on  $L^2(N, \tau)$ . Hence, for every  $\xi, \eta \in L^2(M, \tau)$

$$\begin{aligned} \langle E_N(\lambda_i \lambda_j^*) e_N \xi, \eta \rangle &= \langle e_N \lambda_i \lambda_j^* e_N \xi, \eta \rangle \\ &= \langle \lambda_j^* e_N \xi, \lambda_i^* e_N \eta \rangle \\ &= \delta_{i,j} \langle \lambda_i^* e_N \xi, \lambda_i^* e_N \eta \rangle \\ &= \delta_{i,j} \langle e_N \xi, e_N \eta \rangle. \end{aligned}$$

Since the map  $N \ni y \rightarrow y e_N \in \mathcal{B}(L^2(M, \tau))$  is injective (by faithfulness of  $\tau$ ),  $E_N(\lambda_i \lambda_j^*) = \delta_{i,j} 1$ .

Conversely, suppose  $\{\lambda_i\}_{i=1}^d$  is orthonormal. Then,

$$(\lambda_i^* e_N \lambda_i)(\lambda_j^* e_N \lambda_j) = \lambda_i E_N(\lambda_i \lambda_j^*) e_N \lambda_j = \delta_{i,j} \lambda_i^* e_N \lambda_i,$$

so  $\lambda_i^* e_N \lambda_i$  are mutually orthogonal projections, and  $\lambda_i^*$  is isometric on  $L^2(N, \tau)$  :

$$\begin{aligned} \langle \lambda_i^* e_N \xi, \lambda_i^* e_N \eta \rangle &= \langle e_N \lambda_i \lambda_i^* e_N \xi, \eta \rangle \\ &= \langle E_N(\lambda_i \lambda_i^*) e_N \xi, \eta \rangle \\ &= \langle e_N \xi, \eta \rangle \\ &= \langle e_N \xi, e_N \eta \rangle. \end{aligned}$$

It follows that each projection  $\lambda_i^* e_N \lambda_i$  has rank  $\dim N$ . The identity  $1_{L^2(M, \tau)} = \sum_{i=1}^d \lambda_i^* e_N \lambda_i$  then implies that  $\dim M = d \dim N$ .  $\square$

The next Lemma will produce useful decompositions for the operations in the teleportation scheme of Theorem 4.1.

**Lemma 4.6.** *Let  $N \subseteq M$  be a strongly Markov inclusion with basis  $\{\lambda_i\}_{i=1}^d$ . Then for each positive element  $x_1 \in N' \cap M_1$ , there exists  $\{a_i\}_{i=1}^d$  in  $M$  such that*

$$x_1 = \sum_{i=1}^d a_i^* e_N a_i \quad \text{and} \quad \sum_{i=1}^d a_i^* (N' \cap M) a_i \subseteq N' \cap M.$$

Moreover, the CP map  $\sum_i a_i^*(\cdot)a_i$  on  $N' \cap M$  is independent of the chosen basis  $\{\lambda_i\}_{i=1}^d$ . If, in addition,  $M$  admits a unitary basis  $\{u_i\}_{i=1}^d$  over  $N$  in  $\mathcal{N}_M(N)$ , then we can choose the  $a_i$  so that  $a_i^*(N' \cap M)a_i, a_i(N' \cap M)a_i^* \subseteq N' \cap M$ .

*Proof.* By positivity of  $x_1$ , we get  $x_1 = \sqrt{x_1} \cdot \sqrt{x_1} = \sum_{i=1}^d \sqrt{x_1} \lambda_i^* e_N \lambda_i \sqrt{x_1}$ . Since  $\sqrt{x_1} \lambda_i^* \in M_1$ , there exists a unique element  $a_i^* \in M$  such that  $a_i^* e_N = \sqrt{x_1} \lambda_i^* e_N$  [57, Lemma 4.3.1, Remark 4.3.2(a)], namely  $a_i^* = [M : N]E_M(\sqrt{x_1} \lambda_i^* e_N)$ . The first equation follows.

As for the inclusion, noting that  $a_i^* = [M : N]E_M(a_i^* e_N)$ , for  $x \in N' \cap M$  we have

$$a_i^* x a_i = [M : N]E_M(a_i^* e_N) x a_i = [M : N]E_M(a_i^* e_N x a_i) = [M : N]E_M(a_i^* e_N a_i \gamma_0(x)),$$

and  $\sum_{i=1}^d a_i^* e_N a_i = x_1$ , so  $\sum_{i=1}^d a_i^* x a_i = [M : N]E_M(x_1 \gamma_0(x))$  which belongs to  $N' \cap M$  since  $x_1, \gamma_0(x) \in N'$ . If  $\{\mu_j\}$  was another basis for  $M$  over  $N$ , and  $b_j^* = [M : N]E_M(\sqrt{x_1} \mu_j^* e_N)$ , then decomposing  $\lambda_i$  relative to  $\{\mu_j\}$ , we see that

$$\begin{aligned} a_i^* &= [M : N]E_M(\sqrt{x_1} \lambda_i^* e_N) \\ &= [M : N] \sum_j E_M(\sqrt{x_1} \mu_j^* E_N(\mu_j \lambda_i^*) e_N) \\ &= [M : N] \sum_j E_M(\sqrt{x_1} \mu_j^* e_N) E_N(\mu_j \lambda_i^*) \\ &= \sum_j b_j^* E_N(\mu_j \lambda_i^*). \end{aligned}$$

Hence, for every  $x \in N' \cap M$ , we have

$$\begin{aligned} \sum_i a_i^* x a_i &= \sum_i \sum_{j,k} b_j^* E_N(\mu_j \lambda_i^*) x E_N(\lambda_i \mu_k^*) b_k \\ &= \sum_i \sum_{j,k} b_j^* E_N(\mu_j \lambda_i^* E_N(\lambda_i \mu_k^*)) x b_k \\ &= \sum_{j,k} b_j^* E_N(\mu_j \mu_k^*) x b_k \\ &= [M : N] \sum_{j,k} E_M(\sqrt{x_1} \mu_j^* e_N) E_N(\mu_j \mu_k^*) x b_k \\ &= [M : N] \sum_{j,k} E_M(\sqrt{x_1} \mu_j^* E_N(\mu_j \mu_k^*) e_N) x b_k \\ &= [M : N] \sum_k E_M(\sqrt{x_1} \mu_k^* e_N) x b_k \\ &= \sum_k b_k^* x b_k. \end{aligned}$$

Now, suppose, in addition, that each  $\lambda_i = u_i \in \mathcal{N}_M(N)$  and  $x_1 \in (N' \cap M_1)^+$ . Then defining the  $a_i$  as above, for every  $x \in N' \cap M$  and  $y \in N$ , we have

$$\begin{aligned}
y(a_i^* x a_i) &= [M : N]^2 y E_M(\sqrt{x_1} u_i^* e_N) x E_M(e_N u_i \sqrt{x_1}) \\
&= [M : N]^2 E_M(y \sqrt{x_1} u_i^* e_N) x E_M(e_N u_i \sqrt{x_1}) \quad (\text{bimodule property of } E_M) \\
&= [M : N]^2 E_M(\sqrt{x_1} (y u_i^*) e_N) x E_M(e_N u_i \sqrt{x_1}) \quad (\sqrt{x_1} \in N') \\
&= [M : N]^2 E_M(\sqrt{x_1} u_i^* (u_i y u_i^*) e_N) x E_M(e_N u_i \sqrt{x_1}) \\
&= [M : N]^2 E_M(\sqrt{x_1} u_i^* e_N) x E_M(e_N (u_i y u_i^*) u_i \sqrt{x_1}) \quad (u_i y u_i^* \in N) \\
&= [M : N]^2 E_M(\sqrt{x_1} u_i^* e_N) x E_M(e_N u_i y \sqrt{x_1}) \\
&= [M : N]^2 E_M(\sqrt{x_1} u_i^* e_N) x E_M(e_N u_i \sqrt{x_1}) y \\
&= (a_i^* x a_i) y.
\end{aligned}$$

Similarly,

$$\begin{aligned}
y(a_i x a_i^*) &= [M : N]^2 y E_M(e_N u_i \sqrt{x_1}) x E_M(\sqrt{x_1} u_i^* e_N) \\
&= [M : N]^2 E_M(e_N (y u_i) \sqrt{x_1}) x E_M(\sqrt{x_1} u_i^* e_N) \\
&= [M : N]^2 E_M(e_N u_i (u_i^* y u_i) \sqrt{x_1}) x E_M(\sqrt{x_1} u_i^* e_N) \\
&= [M : N]^2 E_M(e_N u_i \sqrt{x_1}) x E_M(\sqrt{x_1} (u_i^* y u_i) u_i^* e_N) \quad (u_i^* y u_i \in N) \\
&= [M : N]^2 E_M(e_N u_i \sqrt{x_1}) x E_M(\sqrt{x_1} u_i^* y e_N) \\
&= (a_i x a_i^*) y.
\end{aligned}$$

□

Finally, we require the following equivalence with the trace assumption in Theorem 4.1, which will be used in conjunction with tightness to deduce orthonormality of the constructed basis.

**Lemma 4.7.** *Let  $N \subseteq (M_n(\mathbb{C}), \tau)$  be an inclusion such that  $N = \bigoplus_{j=1}^J M_{n_j}(\mathbb{C}) \otimes 1_{m_j}$ . Then  $\tau|_{N'}$  is the Markov trace for the inclusion  $\mathbb{C} \subseteq N'$  if and only if*

$$\frac{n_j}{m_j} = \frac{n}{\dim N'}, \quad j = 1, \dots, J.$$

When this is the case,  $(\text{id} \otimes \tau)e_N = (\tau \otimes \text{id})e_N = [M_n(\mathbb{C}) : N]^{-1}1$ .

*Proof.* Since  $N' = \bigoplus_{j=1}^J 1_{n_j} \otimes M_{m_j}(\mathbb{C})$ , the trace vector associated to  $\tau|_{N'}$  is  $\frac{1}{n}(n_1, \dots, n_J)$ . Since the Markov trace on  $N'$  has trace vector  $\frac{1}{\dim N'}(m_1, \dots, m_J)$ , the first claim follows.

Let  $M_1 = \langle M_n(\mathbb{C}), e_N \rangle$  be the result of the basic construction. Then (abusing notation)  $M_1 = JN'J = M_n(\mathbb{C}) \otimes N'$  and the canonical conditional expectation  $E_{M_n(\mathbb{C})} : M_1 = M_n(\mathbb{C}) \otimes N' \rightarrow M_n(\mathbb{C})$  is simply  $(\text{id} \otimes \tau_{N'})$ , where  $\tau_{N'}$  is the Markov trace for the inclusion  $\mathbb{C} \subseteq N'$ . Thus, assuming  $\tau|_{N'} = \tau_{N'}$ , the Markov property implies that

$$(\text{id} \otimes \tau)e_N = E_M(e_N) = [M_n(\mathbb{C}) : N]^{-1}1.$$

As  $e_N = J e_N J$ , we also have

$$(\tau \otimes \text{id})e_N = (\tau \otimes \text{id})J e_N J = \overline{(\text{id} \otimes \tau)e_N} = [M_n(\mathbb{C}) : N]^{-1}1.$$

□

We are now in position to prove the main result of the section.

*Proof of Theorem 4.1.* Throughout the proof we let  $M := M_n(\mathbb{C})$ . Transposition/complex conjugation on  $M_n(\mathbb{C})$  will be taken relative to a block-diagonalising basis for  $N'$ .

Suppose  $(\omega, \{F_i\}_{i=1}^d, \{T_i\}_{i=1}^d)$  is a tight, minimal faithful teleportation scheme for  $N'$ , with respect to the bipartite system

$$M_n(\mathbb{C}) \otimes M_n(\mathbb{C}) \otimes 1, 1 \otimes 1 \otimes N' \subseteq M_n(\mathbb{C}) \otimes M_n(\mathbb{C}) \otimes N',$$

where we take  $A_0 = N' \otimes 1 \otimes 1$  and  $A_1 = 1 \otimes N' \otimes 1$  inside  $A = M_n(\mathbb{C}) \otimes M_n(\mathbb{C}) \otimes 1$ , and  $B = 1 \otimes 1 \otimes N'$ . Since  $T_i$  is an  $A$ -bimodule map, it must be supported solely on Bob's algebra  $B$ . We therefore view each  $T_i$  as a UCP map on  $N'$ .

Tightness means  $d = \dim N'$ , and minimality ensures that

- $\{F_i\}_{i=1}^d \subseteq A_0 \vee A_1 = N' \otimes N' \otimes 1$ , and
- $\omega \in A_1 \vee B = 1 \otimes N' \otimes N'$ .

The teleportation identity then reads

$$x = \sum_{i=1}^d (E_{N'} \otimes \tau \otimes \tau)((F_i \otimes 1)(1 \otimes 1 \otimes T_i(x))(1 \otimes \omega)), \quad x \in N'. \quad (10)$$

Fixing a Pimsner-Popa basis  $\{\lambda_\alpha\}$  for  $M$  over  $N$  (which always exists for connected inclusions of finite-dimensional  $C^*$ -algebras [7]), by Lemma 4.6, there exist families  $\{w_\alpha\}, \{a_{i,\beta}\}$  in  $M$  (indexed by the same set) satisfying

$$\omega = \sum_{\alpha} (w_\alpha^* \otimes 1)e_N(w_\alpha \otimes 1), \quad F_i = \sum_{\beta} (a_{i,\beta}^* \otimes 1)e_N(a_{i,\beta} \otimes 1), \quad i = 1, \dots, d.$$

Expanding the argument of  $(E_{N'} \otimes \tau \otimes \tau)$  from equation (10), we can rearrange and apply entanglement of  $e_N$  (Lemma 2.4) as follows

$$\begin{aligned} & (F_i \otimes 1)(1 \otimes 1 \otimes T_i(x))(1 \otimes \omega) \\ &= \sum_{\alpha, \beta} ((a_{i,\beta}^* \otimes 1)e_N(a_{i,\beta} \otimes 1) \otimes 1)(1 \otimes 1 \otimes T_i(x))(1 \otimes ((w_\alpha^* \otimes 1)e_N(w_\alpha \otimes 1))) \\ &= \sum_{\alpha, \beta} ((a_{i,\beta}^* \otimes 1 \otimes 1)(e_N \otimes 1)(a_{i,\beta} \otimes w_\alpha^* \otimes 1)(1 \otimes (1 \otimes T_i(x)e_N(w_\alpha \otimes 1)))) \\ &= \sum_{\alpha, \beta} ((a_{i,\beta}^* \otimes 1 \otimes 1)(e_N \otimes 1)(a_{i,\beta} \otimes w_\alpha^* T_i(x)^t \otimes 1)(1 \otimes (e_N(w_\alpha \otimes 1)))). \end{aligned}$$

But then, as  $\tau|_{N'}$  is the Markov trace for  $\mathbb{C} \subseteq N'$ , Lemma 4.7 implies

$$\begin{aligned} & (E_{N'} \otimes \tau \otimes \tau)((F_i \otimes 1)(1 \otimes 1 \otimes T_i(x))(1 \otimes \omega)) \\ &= \sum_{\alpha, \beta} (E_{N'} \otimes \tau \otimes \tau)((a_{i,\beta}^* \otimes 1 \otimes 1)(e_N \otimes 1)(a_{i,\beta} \otimes w_\alpha^* T_i(x)^t \otimes 1)(1 \otimes (e_N(w_\alpha \otimes 1)))) \\ &= [M : N]^{-1} \sum_{\alpha, \beta} (E_{N'} \otimes \tau)((a_{i,\beta}^* \otimes 1)e_N(a_{i,\beta} \otimes w_\alpha^* T_i(x)^t w_\alpha)) \\ &= [M : N]^{-1} (E_{N'} \otimes \tau)((\Psi_i \otimes \text{id})(e_N)(1 \otimes \Phi(T_i(x)^t))), \end{aligned}$$

where  $\Phi := \sum_{\alpha} w_{\alpha}^*(\cdot)w_{\alpha}$  and  $\Psi_i := \sum_{\beta} a_{i,\beta}^*(\cdot)a_{i,\beta}$  are CP maps  $N' \rightarrow N'$  (by Lemma 4.6). Applying entanglement of  $e_N$  and the Markov property once again, we have

$$\begin{aligned}
& (E_{N'} \otimes \tau \otimes \tau)((F_i \otimes 1)(1 \otimes 1 \otimes T_i(x))(1 \otimes \omega)) \\
&= [M : N]^{-1}(E_{N'} \otimes \tau)((\Psi_i \otimes \text{id})(e_N(1 \otimes \Phi(T_i(x)^t)))) \\
&= [M : N]^{-1}(E_{N'} \otimes \tau)((\Psi_i \otimes \text{id})(e_N(\Phi(T_i(x)^t)^t \otimes 1))) \\
&= [M : N]^{-2}E_{N'}(\Psi_i(\Phi(T_i(x)^t)^t)) \\
&= [M : N]^{-2}\Psi_i(\Phi(T_i(x)^t)^t) \\
&= [M : N]^{-2}\Psi_i(\Psi(T_i(x))),
\end{aligned}$$

where  $\Psi := t \circ \Phi \circ t = \sum_{\alpha} w_{\alpha}^t(\cdot)\overline{w_{\alpha}}$  is a CP map  $N' \rightarrow N'$  (as  $N'$  is invariant under transposition). Thus, we have shown

$$\sum_{i=1}^d [M : N]^{-2}\Psi_i \circ \Psi \circ T_i = \text{id}_{N'}.$$

Hence, by Lemma 4.3, if  $N' = \bigoplus_{j=1}^J z_j N'$ , for minimal central projections  $z_1, \dots, z_J$ , then for each  $i$ , there exist  $\mu_i^j, \dots, \mu_i^J \geq 0$  such that

$$[M : N]^{-2}\Psi_i \circ \Psi \circ T_i|_{z_j N'} = \mu_i^j \text{id}_{z_j N'}.$$

Let  $\sigma_i := \sum_j \mu_i^j z_j \in \mathcal{Z}(N)$ . Then for every  $x \in N'$ ,

$$[M : N]^{-2}\Psi_i \circ \Psi \circ T_i(x) = \sum_{j=1}^J [M : N]^{-2}\Psi_i \circ \Psi \circ T_i(z_j x) = \sum_{j=1}^J \mu_i^j z_j x = \sigma_i x. \quad (11)$$

In particular,  $\sigma_i = [M : N]^{-2}\Psi_i \circ \Psi \circ T_i(1) = [M : N]^{-2}\Psi_i(\Psi(1))$ . Next, we show that  $\sigma_i$  is invertible. Since the scheme is faithful, the element

$$\rho_i := (\text{id} \otimes \tau \otimes \tau)((F_i \otimes 1)(1 \otimes \omega)) \in N'$$

satisfies

$$\tau(\rho \rho_i) = (\tau \otimes \tau \otimes \tau)((F_i \otimes 1)(\rho \otimes 1 \otimes 1)(1 \otimes \omega)) > 0$$

for all  $\tau$ -densities  $\rho \in N'$ . Thus,  $\rho_i$  is invertible in  $N'$ . In fact,  $\rho_i = \sigma_i$ : expanding the operators  $F_i$  and  $\omega$  once again using Lemma 4.6 and performing similar manipulations using the entanglement and Markov property of  $e_N$ , we see that

$$\begin{aligned}
\rho_i &= \sum_{\alpha, \beta} (\text{id} \otimes \tau \otimes \tau)((a_{i,\beta}^* \otimes 1)e_N(a_{i,\beta} \otimes 1) \otimes 1)(1 \otimes (w_{\alpha}^* \otimes 1)e_N(w_{\alpha} \otimes 1)) \\
&= (\text{id} \otimes \tau \otimes \tau)((\Psi_i \otimes \text{id})(e_N) \otimes 1)(1 \otimes (\Phi \otimes \text{id})(e_N)) \\
&= [M : N]^{-1}(\text{id} \otimes \tau)((\Psi_i \otimes \text{id})(e_N)(1 \otimes \Phi(1))) \quad (\text{trace out } 3^{\text{rd}} \text{ leg}) \\
&= [M : N]^{-1}(\text{id} \otimes \tau)((\Psi_i \otimes \text{id})(e_N(1 \otimes \Phi(1)))) \\
&= [M : N]^{-1}(\text{id} \otimes \tau)((\Psi_i \otimes \text{id})(e_N(\Phi(1)^t \otimes 1))) \quad (\Phi(1) \in N') \\
&= [M : N]^{-2}\Psi_i(\Phi(1)^t) \\
&= [M : N]^{-2}\Psi_i(\Psi(1)).
\end{aligned}$$

Hence,

$$\sigma_i = [M : N]^{-2}\Psi_i \circ \Psi \circ T_i(1) = [M : N]^{-2}\Psi_i(\Psi(1)) = \rho_i$$

is a positive invertible element of  $\mathcal{Z}(N)$ . Hence,  $[M : N]^{-2}\sigma_i^{-1/2}\Psi_i(\Psi(\cdot))\sigma_i^{-1/2}$  is a UCP map  $N' \rightarrow N'$  which is a left (hence two-sided) inverse to  $T_i$  by equation (11) (and finite-dimensionality of  $N'$ ). Then  $T_i$  is a unital complete order isomorphism of the unital  $C^*$ -algebra  $N'$ , so is necessarily a



\*-automorphism (see, e.g., [36, Corollary 5.2.3]).  $T_i$  is therefore the restriction of a \*-automorphism of  $M$  to  $N'$  (by the proof of [43, Proposition 2.3.3], for instance), that is,  $T_i(x) = u_i x u_i^*$  for some unitary  $u_i \in \mathcal{N}_M(N') = \mathcal{N}_M(N)$ . As shown above,

$$(E_{N'} \otimes \tau \otimes \tau)((F_i \otimes 1)(1 \otimes 1 \otimes T_i(x))(1 \otimes \omega)) = [M : N]^{-2} \Psi_i(\Psi(T_i(x))) = \sigma_i x, \quad x \in N', \quad (12)$$

so we have

$$\sigma_i^{1/2} u_i^* x u_i \sigma_i^{1/2} = (E_{N'} \otimes \tau \otimes \tau)((F_i \otimes 1)(1 \otimes 1 \otimes x)(1 \otimes \omega)), \quad x \in N'.$$

Summing over  $i$ , and using the fact that  $\{F_i\}_{i=1}^d$  is a POVM,

$$\sum_{i=1}^d \sigma_i^{1/2} u_i^* x u_i \sigma_i^{1/2} = (E_{N'} \otimes \tau \otimes \tau)((1 \otimes 1 \otimes x)(1 \otimes \omega)) = \tau(x(\tau \otimes \text{id})(\omega))1, \quad x \in N'. \quad (13)$$

Put  $z := (\tau \otimes \text{id})(\omega) \in N'$ . The above relation together with the fact that  $e_N \in N' \cap M_1 = N' \otimes N'$  implies

$$\sum_{i=1}^d (1 \otimes \sigma_i^{1/2} u_i^*) e_N (1 \otimes u_i \sigma_i^{1/2}) = (\text{id} \otimes \tau)(e_N(1 \otimes z)) \otimes 1 = (\text{id} \otimes \tau)(e_N(z^t \otimes 1)) \otimes 1 = [M : N]^{-1} z^t \otimes 1.$$

Tracing out the right hand side, and using the fact that  $u_i \sigma_i u_i^* \in N'$ ,

$$[M : N]^{-1} z^t = \sum_{i=1}^d (\text{id} \otimes \tau)(e_N(1 \otimes u_i \sigma_i u_i^*)) = \sum_{i=1}^d (\text{id} \otimes \tau)(e_N(\overline{u_i} \sigma_i^t u_i^t \otimes 1)) = [M : N]^{-1} \sum_{i=1}^d \overline{u_i} \sigma_i^t u_i^t,$$

implying  $z = \sum_{i=1}^d u_i \sigma_i u_i^*$  is a positive invertible element of  $\mathcal{Z}(N)$ . But then by (13)

$$\sum_{i=1}^d \sigma_i^{1/2} u_i^* z^{-1/2} x z^{-1/2} u_i \sigma_i^{1/2} = \tau(x)1, \quad x \in N',$$

from which it follows that

$$\sum_{i=1}^d (\sigma_i^{1/2} u_i^* z^{-1/2} \otimes 1) e_N (z^{-1/2} u_i \sigma_i^{1/2} \otimes 1) = 1 \otimes (\tau \otimes \text{id})(e_N) = [M : N]^{-1} 1 \otimes 1,$$

i.e.,  $\{\sqrt{[M : N]} z^{-1/2} u_i \sigma_i^{1/2}\}_{i=1}^d$  forms a basis for  $M$  over  $N$ . However,  $\tau|_{N'}$  is the Markov trace for  $\mathbb{C} \subseteq N'$ , so by Lemma 4.7 we have

$$\frac{n_j}{m_j} = \frac{n}{\dim N'}, \quad j = 1, \dots, J,$$

where  $N = \bigoplus_{j=1}^J M_{n_j}(\mathbb{C}) \otimes 1_{m_j}$  is the decomposition induced from the inclusion  $N \subseteq M_n(\mathbb{C})$ . Hence,

$$\dim N = \sum_{j=1}^J n_j^2 = \sum_{j=1}^J \frac{m_j^2 n^2}{(\dim N')^2} = \frac{\dim N' \cdot n^2}{(\dim N')^2} = \frac{\dim M}{\dim N'}.$$

By tightness, we therefore have  $d = \dim N' = \frac{\dim M}{\dim N'}$ , which entails the orthonormality of the basis  $\{\sqrt{[M : N]} z^{-1/2} u_i \sigma_i^{1/2}\}_{i=1}^d$  by Lemma 4.5. As  $u_i \sigma_i u_i^* \in \mathcal{Z}(N)$ , we get

$$1 = [M : N] E_N(z^{-1/2} u_i \sigma_i u_i^* z^{-1/2}) = [M : N] z^{-1/2} u_i \sigma_i u_i^* z^{-1/2},$$

that is,  $u_i \sigma_i u_i^* = [M : N]^{-1} z$ . But then the basis elements are

$$\sqrt{[M : N]} z^{-1/2} u_i \sigma_i^{1/2} = \sqrt{[M : N]} z^{-1/2} (u_i \sigma_i^{1/2} u_i^*) u_i = u_i,$$

so  $\{u_i\}_{i=1}^d$  is an orthonormal basis of  $M$  over  $N$  inside  $\mathcal{N}_M(N)$ .

Next, consider the decomposition of  $\omega$  induced by  $\{u_i\}_{i=1}^d$  as in Lemma 4.6, that is

$$\omega = \sum_i (w_i^* \otimes 1) e_N(w_i \otimes 1)$$

with each  $w_i^* w_i, w_i w_i^* \in N'$ , where  $w_i^* = [M : N] E_M(\sqrt{\omega} u_i^* e_N)$ . Also by Lemma 4.6, the associated CP map  $\Phi : N' \rightarrow N'$  (defined above) satisfies  $\Phi = \sum_i w_i^*(\cdot) w_i$ , so that  $\Psi = t \circ \Phi \circ t = \sum_i w_i^t(\cdot) \overline{w_i}$ .

Let  $\tilde{\Psi} = [M : N]^{-1} \Psi \circ \text{Ad}(z^{-1/2})$ . From equation (12) it follows that

$$\Psi_i(\tilde{\Psi}(x)) = [M : N]^{-1} \Psi_i(\Psi(z^{-1/2} x z^{-1/2})) = [M : N] (\sigma_i^{1/2} u_i^* z^{-1/2}) x (z^{-1/2} u_i \sigma_i^{1/2}) = u_i^* x u_i \quad (14)$$

for every  $x \in N'$ . Moreover,  $\tilde{\Psi}$  is  $\tau$ -preserving on  $N'$ :

$$\begin{aligned} \tau(\tilde{\Psi}(x)) &= [M : N]^{-1} \tau(\Psi(z^{-1/2} x z^{-1/2})) \\ &= [M : N]^{-1} \sum_{i=1}^d \tau(w_i^t z^{-1/2} x z^{-1/2} \overline{w_i}) \\ &= [M : N]^{-1} \sum_{i=1}^d \tau(z^{-1/2} x z^{-1/2} \overline{w_i} w_i^t) \\ &= \sum_{i=1}^d \tau(z^{-1/2} x z^{-1/2} (\tau \otimes \text{id})(e_N(1 \otimes \overline{w_i} w_i^t))) \\ &= \sum_{i=1}^d \tau(z^{-1/2} x z^{-1/2} (\tau \otimes \text{id})(e_N(w_i w_i^* \otimes 1))) \quad (w_i w_i^* \in N') \\ &= \sum_{i=1}^d \tau(z^{-1/2} x z^{-1/2} (\tau \otimes \text{id})((w_i^* \otimes 1) e_N(w_i \otimes 1))) \\ &= \tau(z^{-1/2} x z^{-1/2} (\tau \otimes \text{id})(\omega)) \\ &= \tau(z^{-1/2} x z^{-1/2} z) \\ &= \tau(x). \end{aligned}$$

Thus,  $T_i \circ \Psi_i = (\tilde{\Psi})^{-1}$  (by equation (14)) is  $\tau$ -preserving, implying that  $\Psi_i$  is  $\tau$ -preserving as  $T_i$  is. Hence, the adjoint maps  $\tilde{\Psi}^*, \Psi_i^*, T_i^* \in \mathcal{CP}(N')$ , defined relative to  $\tau$ , are all UCP, and satisfy  $\tilde{\Psi}^* \circ \Psi_i^* \circ T_i^* = \text{id}_{N'}$ . It follows that  $\tilde{\Psi}^*$  and  $\Psi_i^*$  are  $*$ -automorphisms of  $N'$ , so there exist  $u, v_1, \dots, v_d \in \mathcal{N}_M(N') = \mathcal{N}_M(N)$  (Lemma 4.2) such that

$$\tilde{\Psi}^*(x) = u x u^*, \quad \text{and} \quad \Psi_i^*(x) = v_i x v_i^*, \quad x \in N'.$$

Hence,  $\Psi(x) = [M : N] \tilde{\Psi}(z^{1/2} x z^{1/2}) = [M : N] u^* z^{1/2} x z^{1/2} u$ , so that

$$\Phi(x) = t \circ \Psi \circ t(x) = [M : N] u^t z^{1/2} x z^{1/2} \overline{u},$$

and therefore

$$\omega = (\Phi \otimes \text{id})(e_N) = [M : N] (u^t z^{1/2} \otimes 1) e_N(z^{1/2} \overline{u} \otimes 1) = [M : N] (1 \otimes z^{1/2} u) e_N(1 \otimes u^* z^{1/2}),$$

the last equality following from Lemma 4.2 (and the entanglement of  $e_N$ ).

Finally, as  $u v_i u_i^*(\cdot) u_i v_i^* u^* = \text{id}_{N'}$ , it follows that

$$F_i = (\Psi_i \otimes \text{id})(e_N) = (v_i^* \otimes 1) e_N(v_i \otimes 1) = (u_i^* u \otimes 1) e_N(u^* u_i \otimes 1),$$

and the proof is complete.  $\square$

**Remark 4.8.** The hypothesis that  $\tau|_{N'}$  is the Markov trace for  $\mathbb{C} \subseteq N'$  is valid whenever both  $N$  and  $N'$  are homogeneous subalgebras of  $M_n(\mathbb{C})$ , that is  $N = \bigoplus_{j=1}^J M_k(\mathbb{C}) \otimes 1_l$  (where  $k, l$  are constant in  $j$ ). Indeed, in this case we have  $n = Jkl$ ,  $N' = \bigoplus_{j=1}^J 1_k \otimes M_l(\mathbb{C})$ , and  $\dim N' = Jl^2$ , so that  $\frac{k}{n} = \frac{l}{\dim N'}$ . Since the trace vector associated to  $\tau|_{N'}$  is  $\frac{1}{n}(k, \dots, k)$  and the Markov trace on  $N'$  has trace vector  $\frac{1}{\dim N'}(l, \dots, l)$ , the claim follows.

Hence, Theorem 4.1 applies in particular whenever  $N' = \bigoplus_{j=1}^J M_l(\mathbb{C})$  ( $k \equiv 1$ ). As noted above, homogeneous subalgebras model a distinguished special case of hybrid classical/quantum codes that lend themselves to explicit code constructions and analyses [24, 46, 71, 72].

**Remark 4.9.** Verdon recently generalised Werner’s characterisation of tight teleportation schemes to the setting of entanglement-invertible channels using graphical techniques [86]. One can phrase Theorem 4.1 in Verdon’s context, but it is unclear whether the explicit structure of our resulting scheme (i.e., unitary Pimsner-Popa basis in the normaliser) would follow from his characterisation. In any event, our independent work uses different techniques.

## 5. APPLICATIONS TO QUANTUM CHROMATIC NUMBERS

Quantum graphs can be studied from a variety of perspectives, including non-commutative cofusability graphs of quantum channels [33], quantum relations [88, 89], and  $C^*$ -algebras with a quantum adjacency matrix [70, 20]. See [31] for a recent survey and relations between the approaches. In this work, we follow Weaver’s approach [88, 89] via quantum relations, so that a *quantum graph* is a triple  $(\mathcal{S}, M, \mathcal{B}(H))$ , consisting of a von Neumann algebra  $M \subseteq \mathcal{B}(H)$  and a weak\* closed operator system  $\mathcal{S} \subseteq \mathcal{B}(H)$  which is an  $M'$ -bimodule.

A simple way to construct quantum graphs over a von Neumann algebra  $M$  is through inclusions: any von Neumann subalgebra  $N \subseteq M$  gives rise to a pair of quantum graphs  $(M, N', \mathcal{B}(H))$  and  $(N', M, \mathcal{B}(H))$ , whose associated bimodules are given by the inclusions  $N \subseteq M$  and  $M' \subseteq N'$ , respectively. In this section we combine some of our techniques with those of [22] to compute chromatic numbers for examples of such quantum graphs. For simplicity, we restrict attention to finite-dimensional examples and leave the infinite-dimensional generalizations to future work.

As with quantum graphs themselves, generalizations of graph theoretic parameters including chromatic numbers can be studied from a variety of perspectives. Motivated by [23, Definition 5.10] and [22, Theorem 4.7], we will use the following definition. See [22, Theorem 4.7] for the mentioned equivalence.

**Definition 5.1.** A quantum graph  $(\mathcal{S}, M, \mathcal{B}(H))$  on a finite-dimensional Hilbert space  $H$  is  $(L, c)$  *colourable*, where  $L$  is a tracial von Neumann algebra and  $c \in \mathbb{N}$ , if the following two equivalent conditions hold:

- (1) There is a UCP map  $\Phi : \ell_c^\infty \rightarrow M \otimes L$  of the form  $\Phi(\cdot) = \sum_{i=1}^m A_i^*(\cdot)A_i$  satisfying
  - (a)  $A_i(M' \otimes 1_L)A_j^* \subseteq \ell_c^\infty$  for all  $i, j$ , and
  - (b)  $A_i(\mathcal{S} \cap (M')^\perp \otimes 1_L)A_j^* \subseteq (\ell_c^\infty)^\perp$ , for all  $i, j$ .
- (2) There is a PVM  $\{P_a\}_{a=1}^c$  in  $M \otimes L$  satisfying

$$P_a((\mathcal{S} \cap (M')^\perp) \otimes 1_L)P_a = 0, \quad a = 1, \dots, c. \quad (15)$$

Orthogonal complements of  $M'$  and  $\ell_c^\infty$  are taken in  $\mathcal{B}(H)$  and  $M_c(\mathbb{C})$ , respectively.

We will then refer to either  $(L, c, \Phi)$  or  $(L, c, \{P_a\}_{a=1}^c)$  as a *colouring* of  $(\mathcal{S}, M, \mathcal{B}(H))$ .

By [22, Theorem 4.7], Definition 5.1(1) means precisely that  $(L, c, \Phi)$  forms a perfect quantum commuting strategy for the quantum-to-classical graph homomorphism game between  $(\mathcal{S}, M, \mathcal{B}(H))$  and  $(M_c(\mathbb{C}), \ell_c^\infty, M_c(\mathbb{C}))$  (the complete graph on  $c$  vertices). Condition (a) is then viewed as a type of “synchronicity” condition arising from this non-local game. See [22, §4, §5] and [23, §5] for details and related notions. One can also interpret condition (a) in terms of quantum relations: it means

that  $\Phi$  preserves the “diagonal” quantum relations within the reflexive quantum relations defined by the quantum graphs (see [89]). Condition (b) means that  $(L, c, \Phi)$  is an “entanglement assisted” quantum graph homomorphism from  $(\mathcal{S}, M, \mathcal{B}(H))$  to  $(M_c(\mathbb{C}), \ell_c^\infty, M_c(\mathbb{C}))$ . Indeed, when  $L = \mathbb{C}$ , we recover the notion of quantum graph homomorphism through pushforwards of operator systems as introduced by Stahlke [80] (see also [89]). For simplicity, we do not consider more general algebraic colourings as in [22] (which loosens restrictions on the  $*$ -algebra  $L$ ) but some of our arguments carry through verbatim to “hereditary” colourings (see [22, §5] for details on algebraic colourings).

Definition 5.1(2) is a useful reformulation that will be frequently used in the sequel.

**Definition 5.2.** Let  $(\mathcal{S}, M, \mathcal{B}(H))$  be a quantum graph on a finite-dimensional Hilbert space  $H$ .

- Its *quantum commuting chromatic number* is

$$\chi_{qc}(\mathcal{S}, M, \mathcal{B}(H)) := \min\{c \in \mathbb{N} \mid (\mathcal{S}, M, \mathcal{B}(H)) \text{ is } (L, c) \text{ colourable for some } L\}.$$

- Its *quantum chromatic number* is

$$\chi_q(\mathcal{S}, M, \mathcal{B}(H)) := \min\{c \in \mathbb{N} \mid (\mathcal{S}, M, \mathcal{B}(H)) \text{ is } (L, c) \text{ colourable with } L \text{ finite-dimensional}\}.$$

- Its *local chromatic number* is

$$\chi_{loc}(\mathcal{S}, M, \mathcal{B}(H)) := \min\{c \in \mathbb{N} \mid (\mathcal{S}, M, \mathcal{B}(H)) \text{ is } (L, c) \text{ colourable with } L = \mathbb{C}\}.$$

We sometimes refer to the corresponding sets of colourings as *qc-colourings*, *q-colourings* and *loc-colourings*, respectively.

For complete quantum graphs  $(M_n(\mathbb{C}), M, M_n(\mathbb{C}))$ , it was shown in [22, Theorem 5.6, Theorem 5.9] that

$$\chi_q(M_n(\mathbb{C}), M, M_n(\mathbb{C})) = \chi_{qc}(M_n(\mathbb{C}), M, M_n(\mathbb{C})) = \dim M.$$

Their result, which utilizes teleportation type techniques for one direction, generalises in a straightforward fashion to quantum graphs from finite-dimensional inclusions  $N \subseteq M$  with  $N$  a factor, see Theorem 5.6 below. We include details for convenience of the reader.

**Proposition 5.3.** *Let  $N \subseteq M$  be an inclusion of von Neumann algebras on a finite-dimensional Hilbert space  $H$ , with  $N$  a factor. Then*

$$\chi_q(N', M, \mathcal{B}(H)) \leq [M : N]. \quad (16)$$

*Proof.* Let  $M = \bigoplus_{j=1}^m 1_{n_j} \otimes M_{k_j}(\mathbb{C})$  be the induced decomposition from the representation  $M \subseteq \mathcal{B}(H)$ . Since  $N$  is a factor, we have  $N \cong M_d(\mathbb{C})$  for some  $d$ , and without loss of generality, there exist  $l_j \in \mathbb{N}$  for which  $k_j = l_j d$ , so that  $[M : N] = \sum_{j=1}^d l_j^2$ , and

$$M = \bigoplus_{j=1}^m 1_{n_j} \otimes M_{l_j}(\mathbb{C}) \otimes M_d(\mathbb{C}) = \left( \bigoplus_{j=1}^m 1_{n_j} \otimes M_{l_j}(\mathbb{C}) \right) \otimes M_d(\mathbb{C}).$$

The embedding  $N \subseteq M$  is then simply  $x \mapsto 1_n \otimes x$ , where  $n = \sum_{j=1}^m n_j l_j$ . For each compressed inclusion  $1_{l_j} \otimes M_d(\mathbb{C}) \subseteq M_{l_j}(\mathbb{C}) \otimes M_d(\mathbb{C})$ ,  $x \mapsto 1_{l_j} \otimes x$  (still unital), pick an orthonormal Pimsner-Popa basis  $\{u_i\}_{i=1}^{l_j^2}$  of unitaries lying in  $M_{l_j}(\mathbb{C}) \otimes 1$ , and let  $e_j \in M_{l_j}(\mathbb{C}) \otimes M_{l_j}(\mathbb{C})$  denote the Jones projection for the inclusion  $\mathbb{C} \subseteq M_{l_j}(\mathbb{C})$  (i.e., the maximally entangled state).

Set  $l = \text{lcm}(l_1, \dots, l_m)$ , and for each  $j$ , pick a unital  $*$ -homomorphism  $\pi_j : M_{l_j}(\mathbb{C}) \hookrightarrow M_l(\mathbb{C})$ . Letting  $\Sigma$  denote the tensor flip, define projections  $P_{i,j} \in M \otimes M_l(\mathbb{C})$  by

$$P_{i,j} := 1_{n_j} \otimes (\text{id}_{l_j} \otimes \text{id}_d \otimes \pi_j)(\Sigma_{23}((u_i^* \otimes 1_{l_j})e_j(u_i \otimes 1_{l_j}) \otimes 1_d)), \quad j = 1, \dots, m, \quad i = 1, \dots, l_j^2.$$

Then  $\{P_{i,j}\}$  is a PVM:

$$\begin{aligned}
\sum_{j=1}^m \sum_{i=1}^{l_j^2} P_{i,j} &= \sum_{j=1}^m \sum_{i=1}^{l_j^2} 1_{n_j} \otimes (\text{id}_{l_j} \otimes \text{id}_d \otimes \pi_j)(\Sigma_{23}((u_i^* \otimes 1_{l_j})e_j(u_i \otimes 1_{l_j}) \otimes 1_d)) \\
&= \sum_{j=1}^m 1_{n_j} \otimes (\text{id}_{l_j} \otimes \text{id}_d \otimes \pi_j)(\Sigma_{23}(1_{l_j} \otimes 1_{l_j} \otimes 1_d)) \\
&= \sum_{j=1}^m 1_{n_j} \otimes (\text{id}_{l_j} \otimes \text{id}_d \otimes \pi_j)(1_{l_j} \otimes 1_d \otimes 1_{l_j}) \\
&= \sum_{j=1}^m 1_{n_j} \otimes 1_{l_j} \otimes 1_d \otimes 1_l \\
&= 1_M \otimes 1_l.
\end{aligned}$$

Now, the relative complement

$$\begin{aligned}
N' \cap (M')^\perp &= \{X = [X_{j,j'}] \otimes 1_d \in \mathcal{B}(\oplus_{j=1}^m \mathbb{C}^{n_j} \otimes \mathbb{C}^{l_j}) \otimes 1_d \mid X \perp (\bigoplus_{j=1}^m M_{n_j}(\mathbb{C}) \otimes 1_{l_j} \otimes 1_d)\} \\
&= \{X = [X_{j,j'}] \otimes 1_d \in \mathcal{B}(\oplus_{j=1}^m \mathbb{C}^{n_j} \otimes \mathbb{C}^{l_j}) \otimes 1_d \mid (\text{id} \otimes \tau_{l_j})(X_{j,j}) = 0 \ \forall j = 1, \dots, m\},
\end{aligned}$$

where  $\tau_{l_j}$  is the normalised trace on  $M_{l_j}(\mathbb{C})$ . Since

$$(1_{n_j} \otimes e_j)(Y \otimes 1_{l_j})(1_{n_j} \otimes e_j) = (\text{id} \otimes \tau_{l_j})(Y) \otimes e_j$$

for any  $Y \in M_{n_j}(\mathbb{C}) \otimes M_{l_j}(\mathbb{C})$  (maximally entangled state is a trace vector), for any  $X = [X_{j,j'}] \otimes 1_d \in N' \cap (M')^\perp$ , and every  $j$  we have

$$\begin{aligned}
&(1_{n_j} \otimes (u_i^* \otimes 1_{l_j})e_j(u_i \otimes 1_{l_j}))([X_{k,k'}] \otimes 1_{l_j})(1_{n_j} \otimes (u_i^* \otimes 1_{l_j})e_j(u_i \otimes 1_{l_j})) \\
&= (1_{n_j} \otimes (u_i^* \otimes 1_{l_j})e_j(1_{l_j} \otimes u_i^t))(X_{j,j} \otimes 1_{l_j})(1_{n_j} \otimes (1_{l_j} \otimes \bar{u}_i)e_j(u_i \otimes 1_{l_j})) \\
&= (1_{n_j} \otimes (u_i^* \otimes 1_{l_j})e_j)(X_{j,j} \otimes 1_{l_j})(1_{n_j} \otimes (1_{l_j} \otimes e_j(u_i \otimes 1_{l_j}))) \\
&= (1_{n_j} \otimes u_i^* \otimes 1_{l_j})((\text{id}_{n_j} \otimes \tau_{l_j})(X_{j,j}) \otimes e_j)(1_{n_j} \otimes (1_{l_j} \otimes u_i \otimes 1_{l_j})) \\
&= 0.
\end{aligned}$$

Simple manipulations with the flip map  $\Sigma_{23}$  show that

$$P_{i,j}(X \otimes 1_l)P_{i,j} = 0, \quad X \in N' \cap (M')^\perp.$$

Hence,  $(M_l(\mathbb{C}), [M : N], \{P_{i,j}\})$  is a finite-dimensional colouring of  $(N', M, \mathcal{B}(H))$ , so its quantum chromatic number is at most  $[M : N]$ .  $\square$

Continuing with the proof strategy of [22, Theorem 5.9], we now show that equality holds in (16). The same argument works more generally for hereditary colourings (see [22, §5]), but for simplicity of presentation we restrict to quantum and quantum commuting colourings. We require a generalised version of [22, Lemma 5.8].

**Lemma 5.4.** *Let  $N \subseteq M$  be factors on a finite-dimensional Hilbert space  $H$ . If  $(L, c, \{P_a\}_{a=1}^c)$  is a colouring of  $(N', M, \mathcal{B}(H))$ , then for each  $a$ ,  $R_a = [M : N](E_N \otimes \text{id}_L)P_a$  is a projection such that  $\sum_{a=1}^c R_a = [M : N]1_N \otimes 1_L$ .*

*Proof.* There exist  $d, m, n \in \mathbb{N}$  such that  $H = \mathbb{C}^d \otimes \mathbb{C}^m \otimes \mathbb{C}^n$ ,  $M = 1_d \otimes M_m(\mathbb{C}) \otimes M_n(\mathbb{C})$  and  $N = 1_d \otimes 1_m \otimes M_n(\mathbb{C})$ . Then  $[M : N] = m^2$ .

Since  $P_a \in M \otimes L = 1_d \otimes M_m(\mathbb{C}) \otimes M_n(\mathbb{C}) \otimes L$ , write

$$P_a = \sum_{i,j=1}^m \sum_{k,l=1}^n 1_d \otimes e_{i,j} \otimes e_{k,l} \otimes P_{i,j,k,l}^a,$$

where  $e_{i,j} \in M_m(\mathbb{C})$  and  $e_{k,l} \in M_n(\mathbb{C})$  are matrix units and  $P_{i,j,k,l}^a \in L$ . Let  $i_0, j_0 \in \{1, \dots, m\}$ ,  $i_0 \neq j_0$ . Then

$$1_d \otimes e_{i_0, j_0} \otimes 1_n \in N' \cap (M')^\perp = \{X \in M_d(\mathbb{C}) \otimes M_m(\mathbb{C}) \otimes M_n(\mathbb{C}) \mid (\text{id}_d \otimes \tau_m \otimes \tau_n)(X) = 0\}.$$

Hence,

$$\begin{aligned} 0 &= P_a(1_d \otimes e_{i_0, j_0} \otimes 1_n \otimes 1_L)P_a \\ &= \sum_{i,j,j'=1}^m \sum_{k,k',l,l'=1}^n 1_d \otimes e_{i,j} e_{i_0, j_0} e_{i', j'} \otimes e_{k,l} e_{k', l'} \otimes P_{i,j,k,l}^a P_{i', j', k', l'}^a \\ &= \sum_{i,j'=1}^m \sum_{k,k',l'=1}^n 1_d \otimes e_{i,j'} \otimes e_{k,l'} \otimes P_{i,i_0,k,k'}^a P_{j_0, j', k', l'}^a \\ &= \sum_{i,j'=1}^m \sum_{k,l'=1}^n 1_d \otimes e_{i,j'} \otimes e_{k,l'} \otimes \left( \sum_{k'=1}^n P_{i,i_0,k,k'}^a P_{j_0, j', k', l'}^a \right), \end{aligned}$$

so that

$$\sum_{k'=1}^n P_{i,i_0,k,k'}^a P_{j_0, j', k', l'}^a = 0, \quad \forall i, j', k, l', \quad i_0 \neq j_0. \quad (17)$$

Similarly,  $1_d \otimes (e_{i_0, i_0} - e_{j_0, j_0}) \otimes 1_n \in N' \cap (M')^\perp$  and it follows that

$$\sum_{l=1}^n P_{i,i_0,k,l}^a P_{i_0, j', l, l'}^a = \sum_{l=1}^n P_{i, j_0, k, l}^a P_{j_0, j', l, l'}^a, \quad \forall i, j', k, l', i_0, j_0. \quad (18)$$

Finally, since  $P_i$  is a projection, one easily sees that

$$P_{i,j',k,l'}^a = \sum_{j=1}^m \sum_{l=1}^n P_{i,j,k,l}^a P_{j,j',l,l'}^a \quad \forall i, j', k, l'. \quad (19)$$

The conditional expectation  $E_N : M \rightarrow N$  is the (normalised) partial trace  $(\text{id}_d \otimes \tau_m \otimes \text{id}_n)$ , so that

$$R_a = m^2 (E_N \otimes \text{id}_L) P_a = m \sum_{i=1}^m \sum_{k,l=1}^n 1_d \otimes e_{k,l} \otimes P_{i,i,k,l}^a.$$



Hence,

$$\begin{aligned}
R_a^2 &= m^2 \sum_{k,l'=1}^n 1_d \otimes e_{k,l'} \otimes \left( \sum_{i,i'=1}^m \sum_{k'=1}^n P_{i,i,k,k'}^a P_{i',i',k',l'}^a \right) \\
&= m^2 \sum_{k,l'=1}^n 1_d \otimes e_{k,l'} \otimes \left( \sum_{i=1}^m \sum_{k'=1}^n P_{i,i,k,k'}^a P_{i,i,k',l'}^a \right) \quad (\text{by (17)}) \\
&= m \sum_{k,l'=1}^n 1_d \otimes e_{k,l'} \otimes \left( \sum_{i,j=1}^m \sum_{k'=1}^n P_{i,j,k,k'}^a P_{j,i,k',l'}^a \right) \quad (\text{by (18)}) \\
&= m \sum_{k,l'=1}^n 1_d \otimes e_{k,l'} \otimes \left( \sum_{i=1}^m P_{i,i,k,l'}^a \right) \quad (\text{by (19)}) \\
&= R_a.
\end{aligned}$$

That  $R_a = R_a^*$  and  $\sum_{a=1}^c R_a = m^2 1_N \otimes 1_L$  are immediate from its definition.  $\square$

**Remark 5.5.** Note that in the above proof we took  $R_a \in 1_d \otimes M_n(\mathbb{C}) \otimes L$ . Since  $P_a \in 1_d \otimes M_m(\mathbb{C}) \otimes M_n(\mathbb{C}) \otimes L$ , we could also trace out the (trivial) first leg of  $P_a$  in the definition of  $R_a$  to the same end. This will be done in the next proof.

**Theorem 5.6.** *Let  $N \subseteq M$  be an inclusion of von Neumann algebras on a finite-dimensional Hilbert space  $H$  with  $N$  a factor. Then*

$$\chi_q(N', M, \mathcal{B}(H)) = \chi_{qc}(N', M, \mathcal{B}(H)) = [M : N].$$

*Proof.* By Proposition 5.3 and the fact that  $\chi_q(N', M, \mathcal{B}(H)) \geq \chi_{qc}(N', M, \mathcal{B}(H))$ , it suffices to show that

$$\chi_{qc}(N', M, \mathcal{B}(H)) \geq [M : N].$$

Let  $M = \bigoplus_{j=1}^m 1_{n_j} \otimes M_{k_j}(\mathbb{C})$  be the induced decomposition from the representation  $M \subseteq \mathcal{B}(H)$ . As in the proof of Proposition 5.3, we may assume  $N \cong M_d(\mathbb{C})$  for some  $d \in \mathbb{N}$  and  $k_j = l_j d$  for some  $l_j \in \mathbb{N}$  so that

$$M = \bigoplus_{j=1}^m 1_{n_j} \otimes M_{l_j}(\mathbb{C}) \otimes M_d(\mathbb{C}) = \left( \bigoplus_{j=1}^m 1_{n_j} \otimes M_{l_j}(\mathbb{C}) \right) \otimes M_d(\mathbb{C}).$$

Note that  $[M : N] = \sum_{j=1}^m l_j^2$ .

Suppose  $\{P_a\}_{a=1}^c \subseteq M \otimes L$  is a  $qc$ -colouring of  $(N', M, \mathcal{B}(H))$ . Letting  $z_j$  denote the central projection of  $M$  onto the  $j^{\text{th}}$  summand  $M_{l_j}(\mathbb{C}) \otimes M_d(\mathbb{C})$ , it follows that  $\{P_a(z_j \otimes 1_L)\}_{a=1}^c \subseteq M z_j \otimes L$  is a  $qc$ -colouring of  $(z_j N' z_j, M z_j, \mathcal{B}(z_j H))$ , as  $(M z_j)' = M' z_j$  and  $X \in z_j N' z_j \cap (M' z_j)^\perp \cap \mathcal{B}(z_j H)$ , implies  $X = z_j X z_j \in N' \cap (M')^\perp$ , so that

$$P_a(z_j \otimes 1_L)((z_j N' z_j \cap (M' z_j)^\perp \mathcal{B}(z_j H)) \otimes 1_L)(z_j \otimes 1_L) P_a = 0.$$

By Lemma 5.4,  $R_a^j = l_j^2 (\tau_{n_j} \otimes \tau_{l_j} \otimes \text{id}_d \otimes \text{id}_L)((z_j \otimes 1) P_a)$  is a projection in  $M_d(\mathbb{C}) \otimes L$  satisfying  $\sum_{a=1}^c R_a^j = l_j^2 (1_d \otimes 1_L)$  (see also Remark 5.5). Moreover,  $R_a^i \perp R_a^j$  when  $i \neq j$ . To see this, first note that  $N' = \mathcal{B}(\bigoplus_j \mathbb{C}^{n_j} \otimes \mathbb{C}^{l_j}) \otimes 1_d$ , so taking a matrix unit  $(e_{i_0}^i \otimes e_{j_0}^i)(e_{k_0}^j \otimes e_{l_0}^j)^* \otimes 1_d \in N' \cap (M')^\perp$  ( $i \neq j$ ), and writing

$$P_a(z_j \otimes 1_L) = \sum_{k,l=1}^{l_j} 1_{n_j} \otimes e_{k,l}^j \otimes P_{a,k,l}^j,$$

with  $P_{a,k,l}^j \in M_d(\mathbb{C}) \otimes L$ , we have

$$\begin{aligned}
0 &= P_a((e_{i_0}^i \otimes e_{j_0}^i)(e_{k_0}^j \otimes e_{l_0}^j)^* \otimes 1_d \otimes 1_L)P_a \\
&= P_a(z_i \otimes 1_L)((e_{i_0}^i \otimes e_{j_0}^i)(e_{k_0}^j \otimes e_{l_0}^j)^* \otimes 1_d \otimes 1_L)(z_j \otimes 1_L)P_a \\
&= e_{i_0}^i (e_{k_0}^j)^* \otimes \sum_{k,l=1}^{l_i} \sum_{k',l'=1}^{l_j} e_{k,l}^i e_{i_0}^i (e_{l_0}^j)^* e_{k',l'}^j \otimes P_{a,k,l}^i P_{a,k',l'}^j \\
&= e_{i_0}^i (e_{k_0}^j)^* \otimes \sum_{k=1}^{l_i} \sum_{l'=1}^{l_j} e_k^i (e_{l'}^j)^* \otimes P_{a,k,i_0}^i P_{a,l_0,l'}^j.
\end{aligned}$$

Hence,  $P_{a,k,i_0}^i P_{a,l_0,l'}^j = 0$  for all  $k, i_0, l_0, l'$  whenever  $i \neq j$ , from which the claim  $R_a^i \perp R_a^j$  is easily deduced. It follows that  $R_a := \sum_{j=1}^m R_a^j$  is a projection in  $M_d(\mathbb{C}) \otimes L$  satisfying

$$\sum_{a=1}^c R_a = \sum_{a=1}^c \sum_{j=1}^m R_a^j = \sum_{j=1}^m l_j^2 (1_d \otimes 1_L) = [M : N] 1_d \otimes L_1.$$

On the other hand,  $1_d \otimes 1_L - R_a \geq 0$  so that

$$(c - [M : N]) 1_d \otimes 1_L = \sum_{a=1}^c (1_d \otimes 1_L - R_a) \geq 0,$$

implying  $c \geq [M : N]$ . □

We now combine some of our techniques from previous sections with those of [22, Theorem 5.9] to calculate  $\chi_{loc}(M, N', \mathcal{B}(L^2(M, \tau)))$  for a large class of finite-dimensional inclusions  $N \subseteq M$ .

**Theorem 5.7.** *Let  $N \subseteq M$  be a strongly Markov inclusion of finite-dimensional von Neumann algebras which admits an orthonormal Pimsner-Popa basis  $\{u_i\}$  for  $M/N$  in  $\mathcal{N}_M(N)$ . Then*

$$\chi_{loc}(M, N', \mathcal{B}(L^2(M, \tau))) = \chi_q(M, N', \mathcal{B}(L^2(M, \tau))) = \chi_{qc}(M, N', \mathcal{B}(L^2(M, \tau))) = [M : N].$$

*Proof.* First, by the Markov property, the cardinality of  $\{u_i\}$  is the index  $[M : N]$ :

$$1 = \tau_1(1) = \sum_i \tau_1(u_i^* e_N u_i) = \sum_i \tau_1(e_N) = \frac{|\{u_i\}|}{[M : N]}.$$

It suffices to show  $\chi_{loc}(M, N', \mathcal{B}(L^2(M, \tau))) \leq [M : N]$  and  $\chi_{qc}(M, N', \mathcal{B}(L^2(M, \tau))) \geq [M : N]$ .

$\chi_{loc}(M, N', \mathcal{B}(L^2(M, \tau))) \leq [M : N]$ : Let  $\tau_d$  denote the unique tracial state on  $\mathcal{B}(L^2(M, \tau))$ . Recall that  $\mathcal{N}_M(N) = \mathcal{N}_M(E_N)$  (Lemma 4.2), so that each  $u_i$  normalises  $E_N$ , and that  $\{u_i^* \psi_n \mid i = 1, \dots, [M : N], n = 1, \dots, \dim N\}$  forms an orthonormal basis of  $L^2(M, \tau)$  whenever  $\{\psi_n\}$  is an orthonormal basis of  $L^2(N, \tau)$  (see the proof of Lemma 4.5). It follows that  $E_N$  is  $\tau_d$ -invariant:

given  $x \in M$ , we have

$$\begin{aligned}
\tau_d(x) &= \frac{1}{\dim M} \sum_{n=1}^{\dim N} \sum_{i=1}^{[M:N]} \langle x u_i^* \psi_n, u_i^* \psi_n \rangle \\
&= \frac{1}{\dim M} \sum_{n=1}^{\dim N} \sum_{i=1}^{[M:N]} \langle e_N u_i x u_i^* e_N \psi_n, \psi_n \rangle \\
&= \frac{1}{\dim M} \sum_{n=1}^{\dim N} \sum_{i=1}^{[M:N]} \langle E_N(u_i x u_i^*) \psi_n, \psi_n \rangle \\
&= \frac{1}{\dim M} \sum_{n=1}^{\dim N} \sum_{i=1}^{[M:N]} \langle u_i E_N(x) u_i^* \psi_n, \psi_n \rangle \\
&= \tau_d(E_N(x)).
\end{aligned}$$

Combined with the faithfulness of  $\tau_d$ , it follows that

$$M \cap N^\perp = \{x \in M \mid \tau_d(xy) = 0, \forall y \in N\} = \{x \in M \mid E_N(x) = 0\} = \text{Ker}(E_N).$$

Now, let  $P_i := u_i^* e_N u_i$ . Then  $\{P_i\}_{i=1}^{[M:N]}$  is a PVM in  $N'$  since  $e_N \in N'$ , and each  $u_i$  normalises  $N'$ . To show that  $\{P_i\}_{i=1}^{[M:N]}$  is an *loc*-colouring of the quantum graph  $(M, N', \mathcal{B}(L^2(M, \tau)))$ , by (15), it suffices to show that  $P_i x P_i = 0$  for all  $x$  in  $M \cap ((N')^\perp)^\perp = M \cap N^\perp$ . But for any  $x \in M \cap N^\perp = \text{Ker}(E_N)$ ,

$$\begin{aligned}
P_i x P_i &= (u_i^* e_N u_i) x (u_i^* e_N u_i) \\
&= u_i^* e_N (u_i x u_i^*) e_N u_i \\
&= u_i^* E_N(u_i x u_i^*) e_N u_i \\
&= u_i^* u_i E_N(x) u_i^* e_N u_i \\
&= 0.
\end{aligned}$$

$\chi_{qc}(M, N', \mathcal{B}(L^2(M, \tau))) \geq [M : N]$ : Suppose  $\{P_a\}_{a=1}^c \subseteq N' \otimes L$  is a *qc*-colouring with  $L$  a tracial von Neumann algebra. Then  $P_a(x \otimes 1_L)P_a = 0$  for all  $x \in M \cap N^\perp = \text{Ker}(E_N)$ , implying  $P_a(x \otimes 1_L)P_a = P_a(E_N(x) \otimes 1_L)P_a$  for all  $x \in M$ . In particular,

$$P_a(u_i u_j^* \otimes 1_L)P_a = P_a(E_N(u_i u_j^*) \otimes 1_L)P_a = \delta_{i,j} P_a.$$

Hence, for each  $a$ ,  $\{(u_i^* \otimes 1_L)P_a(u_i \otimes 1_L)\}_{i=1}^{[M:N]}$  is a family of mutually orthogonal projections in  $N' \otimes L$ . By (the left basis version of) [58, Proposition 2.24], the map

$$E_{M'} : N' \ni y \mapsto \frac{1}{[M : N]} \sum_{i=1}^{[M:N]} u_i^* y u_i \in M'$$

is a conditional expectation (unique with respect to canonical traces on  $N'$  and  $M'$ ). Define  $R_a := [M : N](E_{M'} \otimes \text{id}_L)P_a$ . Then

$$R_a = \sum_{i=1}^{[M:N]} (u_i^* \otimes 1_L)P_a(u_i \otimes 1_L)$$

is a projection in  $M' \otimes L$  satisfying  $\sum_{a=1}^c R_a = [M : N]1_{M'} \otimes 1_L$ . But then,

$$(c - [M : N])1_{M'} \otimes 1_L = \sum_{a=1}^c (1_{M'} \otimes 1_L - R_a) \geq 0,$$

forcing  $c \geq [M : N]$ . □

**Remark 5.8.** Note that Theorem 5.7 does not contradict [22, Theorem 5.11] which forbids finite local chromatic number for complete quantum graphs of the form  $(M_n(\mathbb{C}), M, M_n(\mathbb{C}))$  with  $M$  non-abelian. There, the operator system  $\mathcal{S} = M_n(\mathbb{C})$  coincides with the algebra of bounded operators on the representation space  $\mathbb{C}^n$ , whereas in Theorem 5.7, the operator system  $\mathcal{S} = M$  is not the full algebra  $\mathcal{B}(L^2(M, \tau))$  (unless  $M = \mathbb{C}$ ).

## 6. OUTLOOK

In this work, we introduced a model of quantum teleportation in the commuting operator framework, deepened connections with subfactor theory and generalised Werner’s characterisation of tight teleportation schemes. Several natural lines of investigation are left for future work, including

- (1) rigidity of teleportation for more general inclusions  $N \subseteq M$ ;
- (2) further connections with subfactor theory, depth-2 inclusions [73], weak  $C^*$ -Hopf algebras [68, 69, 73], and categorical approaches to quantum teleportation [3];
- (3) superdense coding in the commuting operator framework, building on [53];
- (4) colourings of infinite quantum graphs from finite-index inclusions of  $II_1$  factors;
- (5) connections with quantum automorphism groups [21, §7].

*Acknowledgements.* The second author was partially supported by the NSERC Discovery Grant RGPIN-2017-06275, and would like to thank Michael Brannan, Samuel Harris and Ivan Todorov for helpful discussions. The third author was partially supported by the NSERC Discovery Grant RGPIN-2018-400160, and would like to thank Rajesh Pereira for helpful discussions.

## REFERENCES

- [1] P. Aliferis and A.W. Cross, *Subsystem fault tolerance with the Bacon-Shor code*, Phys. Rev. Lett. 98 (2007), 220502.
- [2] A. Ambainis, M. Mosca, A. Tapp and R. de Wolf, *Private quantum channels*, 41st Annual Symposium on Foundations of Computer Science (Redondo Beach, CA, 2000), 547-553, IEEE Comput. Soc. Press, Los Alamitos, CA, 2000.
- [3] S. Abramsky and B. Coecke, *A categorical semantics of quantum protocols*. Proceedings of the 19th Annual IEEE Symposium on Logic in Computer Science (2004), 415-425.
- [4] A. Almheiri, X. Dong X and D. Harlow, *Bulk locality and quantum error correction in AdS/CFT*. J. High Energy Phys. 04 (2015), 163.
- [5] R. Arens and V. S. Varadarajan, *On the concept of Einstein-Podolsky-Rosen states and their structure*. J. Math. Phys. 41 (2000), 638-651.
- [6] D. Bacon, *Operator quantum error-correcting subsystems for self-correcting quantum memories*, Phys. Rev. A 73 (2006), 012340.
- [7] K. C. Bakshi, *On Pimsner-Popa bases*. Proc. Indian Acad. Sci. Math. Sci. 127 (2017), no. 1, 117-132.
- [8] K. C. Bakshi and V. P. Gupta, *On orthogonal systems, two sided bases and regular subfactors*. New York J. Math. 26 (2020), 817-835.
- [9] K. C. Bakshi and V. P. Gupta, *A few remarks on Pimsner-Popa bases and regular subfactors of depth 2*. Glasgow Math. J. (2021) 1-17.
- [10] S.D. Bartlett, P. Hayden and R.W. Spekkens, *Random subspaces for encryption based on a private shared Cartesian frame*, Phys. Rev. A 72 (5), (2005) 052329.
- [11] S.D. Bartlett, T. Rudolph and R.W. Spekkens, *Decoherence-full subsystems and the cryptographic power of a private shared reference frame*, Phys. Rev. A 70 (3), (2004) 032307.
- [12] C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres and W. K. Wootters, *Teleporting an unknown quantum state via dual classical and Einstein-Podolsky-Rosen channels*. Phys. Rev. Lett. 70 (1993), no. 13, 1895-1899.
- [13] C. H. Bennett, D. P. DiVincenzo, J. A. Smolin and W. K. Wootters, *Mixed-state entanglement and quantum error correction*. Phys. Rev. A (3) 54 (1996), no. 5, 3824-3851.
- [14] C. Bény, A. Kempf and D. W. Kribs, *Quantum error correction on infinite-dimensional Hilbert spaces*. J. Math. Phys. 50 (2009), no. 6, 062108, 24 pp.

- [15] C. Bény, A. Kempf and D.W. Kribs, *Generalization of quantum error correction via the Heisenberg picture*, Phys. Rev. Lett. 98 (2007), 100502.
- [16] C. Bény, A. Kempf and D.W. Kribs, *Quantum error correction of observables*, Phys. Rev. A 76 (2007), 042303.
- [17] M. Berta, F. Furrer and V. B. Scholz, *The smooth entropy formalism for von Neumann algebras*, J. Math. Phys. 57 (2016), no. 1, 015213, 25 pp.
- [18] D. Bisch, *Bimodules, higher relative commutants and the fusion algebra associated to a subfactor*. The Fields Institute for Research in Mathematical Sciences Communications Series, 13 (1997), 13-63.
- [19] P.O. Boykin and V. Roychowdhury, *Optimal encryption of quantum bits*, Phys. Rev. A 67 (2003), 042317.
- [20] M. Brannan, A. Chirvasitu, K. Eifler, S. J. Harris, V. Paulsen, X. Su and M. Wasilewski, *Bigalois extensions and the graph isomorphism game*. Comm. Math. Phys. 375 (2020), no. 3, 1777-1809.
- [21] M. Brannan, K. Eifler, C. Voigt and M. Weber, *Quantum Cuntz-Krieger algebras*. Trans. Amer. Math. Soc. to appear. arXiv:2009.09466.
- [22] M. Brannan, P. Ganesan and S. J. Harris, *The quantum-to-classical graph homomorphism game*. arXiv:2009.07229v2.
- [23] M. Brannan, S. J. Harris, I. G. Todorov and L. Turowska, *Synchronicity for quantum non-local games*. arXiv:2106.11489.
- [24] N. Cao, D.W. Kribs, C.-K. Li, M. Nelson, Y.-T. Poon and B. Zeng, *Quantum error correction of observables*, Phys. Rev. A 76 (2007), 042303.
- [25] M. Cha, P. Naaijken and B. Nachtergaele, *On the stability of charges in infinite quantum spin systems*. Comm. Math. Phys. 373 (2020), no. 1, 219-264.
- [26] A. Church, D.W. Kribs, R. Pereira and S. Plosker, *Private quantum channels, conditional expectations, and trace vectors*, Quant. Inf. & Comp. 11 (2011), 774-783.
- [27] R. Cleve, D. Gottesman, and H.-L. Lo, *How to share a quantum secret*, Phys. Rev. Lett. 83 (1999), 648.
- [28] J. Crann, D. W. Kribs, R. H. Levene and I. G. Todorov, *Private algebras in quantum information and infinite-dimensional complementarity*. J. Math. Phys. 57 (2016), no. 1, 015208.
- [29] J. Crann, D. W. Kribs, R. H. Levene and I. G. Todorov, *State convertibility in the von Neumann algebra framework*. Comm. Math. Phys. 378 (2020), no. 2, 1123-1156.
- [30] C. Crepeau, D. Gottesman and A. Smith, *Secure multi-party quantum computing*, 34th Annual Symposium on Theory of Computing (ACM, Montreal) (2002), 643.
- [31] M. Daws, *Quantum graphs: different perspectives, homomorphisms and quantum automorphisms*. arXiv:2203.08716.
- [32] I. Devetak and P.W. Shor, *The capacity of a quantum channel for simultaneous transmission of classical and quantum information*, Comm. Math. Phys. 256 (2005), 287-303.
- [33] R. Duan, S. Severini and A. Winter, *Zero-error communication via quantum channels, noncommutative graphs, and a quantum Lovász number*. IEEE Trans. Inform. Theory 59 (2013), no. 2, 1164-1174.
- [34] K. Dykema, V. Paulsen and J. Prakash, *Non-closure of the set of quantum correlations via graphs*. Comm. Math. Phys. 365 (2019), no. 3, 1125-1142.
- [35] E. G. Effros and C. E. Lance, *Tensor products of operator algebras*. Adv. Math. 25 (1977), no. 1, 1-34.
- [36] E. G. Effros and Z.-J. Ruan, *Operator spaces*. London Mathematical Society Monographs. New Series, 23. The Clarendon Press, Oxford University Press, New York, 2000.
- [37] A. Einstein, B. Podolsky and N. Rosen, *Can quantum-mechanical description of physical reality be considered complete?* Phys. Rev. 47 (1935), 777-780.
- [38] L. Fiedler, P. Naaijken and T.J. Osborne, *Jones index, secret sharing and total quantum dimension*, New J. Phys. 19 (2017), 023039.
- [39] P. Ganesan, L. Gao, S. K. Pandey and S. Plosker, *Quantum majorization on semi-finite von Neumann algebras*. J. Funct. Anal. 279 (2020), no. 7, 108650, 49 pp.
- [40] L. Gao, M. Junge and N. LaRacuenta, *Unifying entanglement with uncertainty via symmetries of observable algebras*. arXiv:1710.10038.
- [41] L. Gao, M. Junge and N. LaRacuenta, *Relative entropy for von Neumann subalgebras*. Internat. J. Math. 31 (2020), no. 6, 2050046, 35 pp.
- [42] E. Gesteau and M. J. Kang, *Nonperturbative gravity corrections to bulk reconstruction*. arXiv:2112.12789.
- [43] F. M. Goodman, P. de la Harpe and V. F. R. Jones, *Coxeter Graphs and Towers of Algebras*. Mathematical Sciences Research Institute Publications, 14. Springer-Verlag New York Inc., 1989.
- [44] G. Gordon and G. Rigolin, *Quantum cryptography using partially entangled states*. Opt. Commun. 283 (2010), 184.
- [45] D. Gottesman and I. Chuang, *Demonstrating the viability of universal quantum computation using teleportation and single-qubit operations*. Nature 402 (1999), 390.
- [46] M. Grassl, S. Lu and B. Zeng, *Codes for simultaneous transmission of quantum and classical information*, 2017 IEEE International Symposium on Information Theory (ISIT) (2017), 1718-1722.

- [47] R. Haag and D. Kastler, *An algebraic approach to quantum field theory*. J. Math. Phys. 5 (1964), 848-861.
- [48] U. Haagerup and M. Musat, *An asymptotic property of factorizable completely positive maps and the Connes embedding problem*. Comm. Math. Phys. 338 (2015), 721-752.
- [49] D. Harlow, *The Ryu-Takayanagi formula from quantum error correction*. Comm. Math. Phys. 354 (2017), 865-912.
- [50] F. Hiai, *Quantum  $f$ -divergences in von Neumann algebras I. Standard  $f$ -divergences*. J. Math. Phys. 59 (2018), no. 10, 102202, 27 pp.
- [51] F. Hiai, *Quantum  $f$ -divergences in von Neumann algebras II. Maximal  $f$ -divergences*. J. Math. Phys. 60 (2019), no. 1, 012203, 30 pp.
- [52] S. Hollands and K. Sanders, *Entanglement measures and their properties in quantum field theory*. Springer Briefs in Mathematical Physics 34, 2018. Springer International Publishing.
- [53] S. Huang, *Dense coding and quantum teleportation using type  $II_1$  von Neumann algebras*. J. Math. Phys. 60 (2019), 082107.
- [54] Z. Ji, A. Natarajan, T. Vidick, J. Wright and H. Yuen,  *$MIP^* = RE$* , arXiv:2001.04383.
- [55] T. Jochym-O'Connor, D.W. Kribs, R. Laflamme and S. Plosker, *Private quantum subsystems*, Phys. Rev. Lett. 111, (2013) 030502.
- [56] V. F. R. Jones, *Index for subfactors*. Invent. Math. 72 (1983), no. 1, 1-25.
- [57] V. F. R. Jones and V. S. Sunder, *Introduction to subfactors*. London Mathematical Society Lecture Note Series, 234. Cambridge University Press, Cambridge, 1997.
- [58] V. F. R. Jones and D. Penneys, *The embedding theorem for finite depth subfactor planar algebras*. Quantum Topol. 2 (2011), no. 3, 301-337.
- [59] M. Junge, M. Navascues, C. Palazuelos, D. Perez-Garcia, V. B. Scholz and R. F. Werner. *Connes' embedding problem and Tsirelson's problem*, J. Math. Phys. 52 (2011), no. 1, 012102.
- [60] M. Keyl, D. Schlingemann and R. F. Werner, *Infinitely entangled states*. Quantum Inf. Comput. 3 (2003), no. 4, 281-306.
- [61] A. Klappenecker and P.K. Sarvepalli, *Clifford code constructions of operator quantum error-correcting codes*, IEEE Trans. Inf. Thy. 54 (2008), 5760-5765.
- [62] I. Kremsky, M.-H. Hsieh and T.A. Brun, *Classical enhancement of quantum-error-correcting codes*, Phys. Rev. A 78 (2008), 012341.
- [63] D.W. Kribs, R. Laflamme and D. Poulin, *Unified and generalized approach to quantum error correction*, Phys. Rev. Lett. 94 (2005), 180501.
- [64] D.W. Kribs, J. Levick, M. Nelson, R. Pereira and M. Rahaman, *Approximate quasi-orthogonality of operator algebras and relative quantum privacy*, Rep. Math. Phys. 87 (2021), 167-181.
- [65] G. Kuperberg, *The capacity of hybrid quantum memory*, IEEE Trans. Inf. Thy. 49 (2003), 1465-1473.
- [66] J. Levick, D.W. Kribs and R. Pereira, *Quantum privacy and Schur product channels*, Rep. Math. Phys. 80 (2017), 333-347.
- [67] R. Longo and F. Xu, *Relative entropy in CFT*. Adv. Math. 337 (2018), 139-170.
- [68] A. Molnar, A. R. de Alarcón, J. Garre-Rubio, N. Schuch, J. Ignacio Cirac and D. Pérez-Garcia, *Matrix product operator algebras I: representations of weak Hopf algebras and projected entangled pair states*. arXiv:2204.05940.
- [69] A. R. de Alarcón, J. Garre-Rubio, A. Molnar and D. Pérez-Garcia, *Matrix Product Operator Algebras II: Phases of Matter for 1D Mixed States*. arXiv:2204.06295.
- [70] B. Musto, D. Reutter and D. Verdon. *A compositional approach to quantum functions*. J. Math. Phys. 59 (2018), no. 8, 081706, 42 pp.
- [71] A. Nemeč and A. Klappenecker, *Hybrid codes*, 2018 IEEE International Symposium on Information Theory (ISIT) (2018), 796-800.
- [72] A. Nemeč and A. Klappenecker, *Infinite Families of Quantum-Classical Hybrid Codes*, IEEE Trans. Inf. Thy. 67 (2021), 2847-2856.
- [73] D. Nikshych and L. Vainerman, *A characterization of depth-2 subfactors of  $II_1$  factors*. J. Funct. Anal. 171 (2000), no. 2, 278-307.
- [74] O. Oreshkov, T.A. Brun and D.A. Lidar, *Fault-tolerant holonomic quantum computation*, Phys. Rev. Lett. 102 (2009), 070502.
- [75] N. Ozawa, *About the Connes embedding conjecture: algebraic approaches*. Jpn. J. Math. 8 (2013), no. 1, 147-183.
- [76] M. Pimsner and S. Popa, *Entropy and index for subfactors*. Ann. Sci. Ecole Norm. Sup. (4) 19 (1986), no. 1, 57-106.
- [77] S. Popa, *Classification of amenable subfactors of type II*. Acta Math. 172 (1994), no. 2, 163-255.
- [78] D. Poulin, *Stabilizer formalism for operator quantum error correction*, Phys. Rev. Lett. 95 (2005), 230504.
- [79] W. Slofstra, *Tsirelson's problem and an embedding theorem for groups arising from non-local games*. J. Amer. Math. Soc. 33 (2020), no. 1, 1-56.

- [80] D. Stahlke, *Quantum zero-error source-channel coding and non-commutative graph theory*, IEEE Trans. Inform. Theory 62 (2016), no. 1, 554-577.
- [81] M. Takesaki, *Theory of Operator Algebras I*. Encyclopedia of Mathematical Sciences 124, Springer-Verlag Berlin–Heidelberg–New York, 2001.
- [82] M. Takesaki, *Theory of Operator Algebras II*. Encyclopedia of Mathematical Sciences 125, Springer-Verlag Berlin–Heidelberg–New York, 2003.
- [83] Y. Watatani, *Index for  $C^*$ -subalgebras*. Mem. Amer. Math. Soc. 83 (1990), no. 424, vi+117 pp.
- [84] I. G. Todorov and L. Turowska, *Quantum no-signalling correlations and non-local games*. arXiv:2009.07016.
- [85] R. Verch and R. F. Werner, *Distillability and positivity of partial transposes in general quantum field systems*. Rev. Math. Phys. 17 (2005), no. 5, 545-576.
- [86] D. Verdon, *Entanglement-invertible channels*. arXiv:2204.04493.
- [87] J. von Neumann, *Mathematical foundations of quantum mechanics*. Princeton University Press, 1955.
- [88] N. Weaver, *Quantum relations*. Mem. Amer. Math. Soc. 215 (2012), no. 1010, v-vi, 81-140.
- [89] N. Weaver, *Quantum graphs as quantum relations*. J. Geom. Anal. 31 (2021), no. 9, 9090-9112.
- [90] R. F. Werner, *EPR states for von Neumann algebras*. arXiv:quant-ph/9910077.
- [91] R. F. Werner, *All teleportation and dense coding schemes*. J. Phys. A 34 (2001), no. 35, 7081-7094.
- [92] Y. Zhang, *Teleportation, braid group and Temperley-Lieb algebra*. J. Phys. A 39 (2006), no. 37, 11599-11622.

SCHOOL OF MATHEMATICS & STATISTICS, CARLETON UNIVERSITY, OTTAWA, ON, CANADA H1S 5B6  
*Email address:* alexandreconlon@cmail.carleton.ca

SCHOOL OF MATHEMATICS & STATISTICS, CARLETON UNIVERSITY, OTTAWA, ON, CANADA H1S 5B6  
*Email address:* jasoncramm@cunet.carleton.ca

DEPARTMENT OF MATHEMATICS & STATISTICS, UNIVERSITY OF GUELPH, GUELPH, ON, CANADA N1G 2W1  
*Email address:* dkribs@uoguelph.ca

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY COLLEGE DUBLIN, BELFIELD, DUBLIN 4, IRELAND  
*Email address:* rupert.levene@ucd.ie