

# Coboundary Expansion of Sheaves on Graphs and Weighted Mixing Lemmas

Uriya A. First  
Department of Mathematics  
University of Haifa

Tali Kaufman  
Computer Science Department,  
Bar-Ilan University

August 4, 2022

## Abstract

We study the coboundary expansion of graphs, but instead of using  $\mathbb{F}_2$  as the coefficient group when forming the cohomology, we use a *sheaf* on the graph. We prove that if the graph under discussion is a good expander, then it is also a good coboundary expander relative to any *constant* augmented sheaf (equivalently, relative to any coefficient group  $R$ ); this, however, may fail for *locally constant* sheaves. We moreover show that if we take the quotient of a constant augmented sheaf on an excellent expander graph by a “small” subsheaf, then the quotient sheaf is still a good coboundary expander.

Along the way, we prove a new version of the Expander Mixing Lemma applying to  $r$ -partite weighted graphs.

## 1 Introduction

### 1A Coboundary and Cosystolic Expanders

#### Expander Graphs

Informally, a finite graph is called an expander if relatively many edges cross between every set of vertices and its complement. Since their introduction in the 1970s by Pinsker, expander graph have been widely studied and found numerous applications in many areas of mathematics and computer science.

If  $X$  is a graph and  $w : X \rightarrow \mathbb{R}_+$  is a function assigning non-negative weights to the vertices and edges of  $X$  (one can take  $w \equiv 1$  to ignore the weights), then the expansion of  $(X, w)$  is quantified by its *Cheeger constant*,

$$h(X, w) = \min_{\emptyset \neq S \subsetneq V} \frac{w(E(S, V - S))}{\min\{w(S), w(V - S)\}}. \quad (1.1)$$

Here,  $V$  is the vertex set of  $X$  and  $E(A, B)$  denotes the set of edges with one vertex in  $A$  and the other in  $B$ . One says that  $(X, w)$  is an  $\varepsilon$ -expander if  $h(X) \geq \varepsilon$ .

#### Coboundary Expansion

Meshulam–Wallach [26] and Gromov [10], following the earlier work of Linial–Meshulam [19], observed that the  $\varepsilon$ -expansion condition for graphs can be restated in terms of cohomology with  $\mathbb{F}_2$ -coefficients, and thus generalized to higher dimensions if  $X$  is a simplicial complex, rather than a graph.

Recall that if  $X$  is a simplicial complex and  $i \in \mathbb{N} \cup \{-1, 0\}$ , then the set of  *$i$ -cochains* on  $X$  with  $\mathbb{F}_2$ -coefficients is  $C^i = C^i(X, \mathbb{F}_2) := \mathbb{F}_2^{X(i)}$ , where  $X(i)$  is the set of  $i$ -dimensional faces.<sup>1</sup> That is, an  $i$ -cochain is an assignment of an element in  $\mathbb{F}_2$  to every  $i$ -face of  $X$ . The coboundary map  $d_i : C^i \rightarrow C^{i+1}$  is defined by

$$d_i f(y) = \sum_{x \text{ is an } i\text{-face of } y} f(x)$$

<sup>1</sup>The simplicial complex  $X$  also includes a single empty face, of dimension  $-1$ .

for all  $f \in C^i$  and  $y \in X(i+1)$ , and the  $\mathbb{F}_2$ -vector spaces of  $i$ -boundaries and  $i$ -cocycles are  $B^i = B^i(X, \mathbb{F}_2) = \text{im } d_{i-1}$  and  $Z^i = Z^i(X, \mathbb{F}_2) = \ker d_i$ , respectively. A standard computation shows that  $d_i \circ d_{i-1} = 0$  and thus  $B^i \subseteq Z^i$ . The quotient space  $Z^i/B^i$  is  $H^i(X, \mathbb{F}_2)$ , which is the *reduced* singular cohomology of  $X$  with  $\mathbb{F}_2$ -coefficients.

Coboundary expansion measures the expansion of  $i$ -cochains under  $d_i$ , taking into account that  $B^i$  must be mapped to 0. Formally,  $X$  is called an  $\varepsilon$ -coboundary expander in dimension  $i$  if

$$\|d_i f\|_{\text{Ham}} \geq \varepsilon d_{\text{Ham}}(f, B^i) \quad \forall f \in C^i,$$

where  $\|\cdot\|_{\text{Ham}}$  and  $d_{\text{Ham}}$  are the Hamming norm and distance (in  $\mathbb{F}_2^{X(i)}$  or  $\mathbb{F}_2^{X(i+1)}$ ), respectively. There is an evident variant of this definition to *weighted* simplicial complexes, see [24], for instance.

For example, if  $f \in C^0$  is supported on  $A \subseteq X(0)$ , then the support of  $d_0 f$  is precisely the set of edges leaving  $A$ . In addition,  $B^0$  consists of the two vectors  $(1, \dots, 1)$  and  $(0, \dots, 0)$ . Consequently,  $X$  is an  $\varepsilon$ -coboundary expander in dimension 0 if and only if  $X$  is an  $\varepsilon$ -expander. Coboundary expansion is therefore a high-dimensional version of expansion in graphs.

## Applications of Coboundary Expansion

Coboundary expansion is a natural extension of graph-expansion to higher dimensions, and its study is part of the recent trend of *high dimensional-expanders*; see [21] for a survey. In addition, it was used by Gromov [10] to get bounds on the minimal amount of overlapping forced by mapping a simplicial complex into  $\mathbb{R}^d$ . Coboundary expansion is also strongly related to *property testing* in theoretical computer sciences [14]. Indeed, it measures how well one could test whether  $f \in C^i$  is a coboundary by probing just a few coordinates from  $f$ . Finally, as we shall shortly recall, coboundary expansion *at the links* can be used as a mean to establish *cosystolic expansion*.

## Cosystolic Expansion

Cosystolic expansion is a more lax version of coboundary expansion which was introduced in [5], [13], [7] in order to extend the reach of Gromov's aforementioned work. It is also a form of property testing, the property now being whether  $f \in C^i$  is an  $i$ -cocycle, and has applications in the construction of *quantum CSS codes*; see [14]. Formally, a simplicial complex  $X$  is an  $(\varepsilon, \delta)$ -cosystolic expander in dimension  $i$  if

- (1)  $\|d_i f\|_{\text{Ham}} \geq \varepsilon d_{\text{Ham}}(f, Z^i)$  for all  $f \in C^i$ , and
- (2)  $\|g\|_{\text{Ham}} \geq \delta$  for all  $g \in Z^i - B^i$ .

If  $H^i(X, \mathbb{F}_2) = 0$ , then this is equivalent to  $X$  being an  $\varepsilon$ -coboundary expander.

The works of Kaufman–Kazhdan–Lubotzky [13], Evra–Kaufman [7] and Kaufman–Mass [15], [16] showed that if  $\dim X \geq i + 2$  and all the *proper links* of  $X$  are sufficiently good coboundary expanders in a range of dimensions, then  $X$  is an  $(\varepsilon, \delta)$ -cosystolic expander in dimension  $i$ , with  $\varepsilon, \delta$  depending on the suppressed expansion constants. Coboundary expansion is therefore an essential ingredient in establishing cosystolic expansion.

## Other Coefficients

Meshulam and Wallach [26] observed that the definition of coboundary expansion makes sense if we replace  $\mathbb{F}_2$  with any abelian group of coefficients  $R$ .<sup>2</sup> (In fact, for  $i \in \{0, 1\}$ , one could even consider non-abelian coefficient groups as in [3].) The expansion constants attached to different coefficient groups are *a priori* not related.

In the recent paper [8], the authors introduced *sheaves* on simplicial complexes, which may be regarded as a vast generalization of (abelian) coefficient groups, and noted that coboundary expansion and cosystolic expansion make sense in this much broader context.

The purpose of this work is to study coboundary expansion of graphs, but in the more general context of sheaves. However, before discussing that, we first survey the known examples of coboundary expanders.

<sup>2</sup>One then needs to introduce signs in the definition of the coboundary map  $d_i$ , see §4C.

## Known Constructions of Coboundary Expanders

In contrast to expander graphs, examples of simplicial complexes  $X$  which are good cosystolic expanders in all dimensions  $i \in \{0, \dots, \dim X - 1\}$  are much less common. In fact, for dimension  $d > 2$ , it is widely open whether there is an infinite family of  $d$ -dimensional simplicial complexes of bounded degree<sup>3</sup> which are  $\varepsilon$ -coboundary expanders in dimensions  $0, \dots, d - 1$ .

The known constructions of coboundary expanders can be divided into two families: random and explicit. The known random constructions are the following.

- (r1) Linial–Meshulam [19] and Meshulam–Wallach [26] considered a generalization of the Erdős–Rényi model for random graphs to  $k$ -dimensional simplicial complexes: one starts with a the complete  $(k - 1)$ -dimensional simplicial complex on  $n$  vertices and each  $k$ -simplex is added with a fixed probability  $p \in [0, 1]$ . They showed that if  $p \geq \frac{k \log n + \omega(n)}{n}$  with  $\omega(n) \rightarrow \infty$ , and  $R$  is a finite abelian group, then there is  $\varepsilon > 0$  such that this model almost surely produces an  $\varepsilon$ -cosystolic expander in dimension  $k - 1$  relative to the coefficient group  $R$ . (The work [19] considered the case  $k = 2$  and  $R = \mathbb{F}_2$ , and the general case is [26].) Note that taking  $p = 1$  implies that the complete  $k$ -dimensional complex on  $n$  vertices is a good coboundary expander in dimensions  $k - 1$  relative to any finite coefficient group  $R$ .
- (r2) Dotterrer and Kahle [4] considered the more general random model where one begins with a  $k$ -dimensional simplicial complex  $X$  and dilutes  $X$  by keeping every  $k$ -face of  $X$  with probability  $p \in [0, 1]$ . They showed that if  $p$  is above a certain threshold depending the  $(k - 1)$ -th coboundary expansion of  $X$ , then there is  $\varepsilon > 0$  such that the resulting diluted complex is almost surely an  $\varepsilon$ -coboundary expander in dimension  $k - 1$  w.r.t. the coefficient group  $\mathbb{F}_2$ . They also show that their result is effective if  $X$  is a complete  $(k + 1)$ -partite simplicial complex on  $(k + 1)n$  vertices, the  $k$ -dimensional skeleton of a  $n$ -dimensional simplex (recovering (r1)), or a the  $k$ -dimensional skeleton of an  $n$ -dimensional cross-polytope.
- (r3) Lubotzky–Meshulam [23] (in dimension 2) and Lubotzky–Luria–Rosenthal [22] (in dimension  $> 2$ ), showed that for every  $k \geq 1$  and  $\varepsilon > 0$ , there is  $N = N(k)$  such that gluing  $N$  random *Steiner systems* on  $n$  elements (chosen according to a particular distribution) on the complete  $(k - 1)$ -dimensional simplicial complex on those elements will almost surely result in an  $\varepsilon$ -coboundary expander in dimension  $k - 1$ , where  $\varepsilon > 0$  depends on  $k$ ; the coefficient group is  $\mathbb{F}_2$ .

We now list the known explicit constructions of coboundary expanders. Mind that most of these results impose weights on the simplicial complex at hand.

- (e0) By definition, expander graphs are good coboundary expanders in dimension 0 w.r.t. the coefficient group  $\mathbb{F}_2$ . Surprisingly, the case of a general coefficient group has not been considered so far. We will show below that expander graphs have good coboundary expansion in dimension 0 with respect to any coefficient group  $R$  (Theorem 5.2, Corollary 5.3).
- (e1) Meshulam–Wallach [26] (see (r1)) and Gromov [10] showed independently that a simplex on  $n + 1$  elements is an  $\frac{n+1}{n-k}$ -cosystolic expander in dimension  $k$  w.r.t. the coefficient group  $\mathbb{F}_2$ .<sup>4</sup> Similar bounds for the  $n$ -dimensional cross-polytope and  $(k + 1)$ -partite complex on  $(k + 1)n$  vertices appear in Dotterrer–Kahle [4] (see (r2)).
- (e2) The second author, Kazhdan and Lubotzky [13, Corollary 8.1] showed that some infinite families of *Ramanujan complexes* (in the sense of [25]) of dimension 3 are  $\varepsilon$ -coboundary expanders in dimensions 0 and 1 w.r.t. the coefficient group  $\mathbb{F}_2$ . This result relies on Serre’s conjecture on the congruence subgroup property.
- (e3) Lubotzky, Moses and Meshulam [24] showed that there is a uniform lower bound on the  $k$ -dimensional coboundary expansion w.r.t. the coefficient group  $\mathbb{F}_2$  of finite *spherical buildings* of a fixed dimension  $d \geq k + 1$  admitting a strongly transitive action. A similar uniform bound applying to all abelian coefficient groups was proved by the second author and D. Mass [15, Theorem 4.1]. This was also shown to hold for finite buildings of type  $A_3$  with non-abelian coefficients groups by Dinur and Meshulam [3].

<sup>3</sup>The degree of a simplicial complex  $X$  is the maximal number of faces containing a vertex of  $X$ .

<sup>4</sup>The weights on the faces of the  $(n + 1)$ -simplex are given by  $w(x) = \binom{n+1}{\dim x + 1}^{-1}$ .

- (e4) The second author and Oppenheim [17] proved a lower bound on the  $\mathbb{F}_2$ -coefficients  $k$ -dimensional coboundary expansion of any simplicial complex  $X$  by means of certain global invariants, generalizing the diameter of a graph. In the context of spherical buildings, this recovers the results of [24] (cf. (e3)). They apply this result to give examples of coboundary expanders arising from certain *coset geometries*.

## 1B Sheaves

### Sheaves

Sheaves on simplicial complexes were introduced by the authors in [8] as discrete versions of sheaves on topological spaces, and with the purpose of constructing *2-query locally testable codes*. Here, we shall regard them simply as a generalizing coefficient groups in the context of cohomology.

In more detail, an *augmented sheaf* on a simplicial complex  $X$  consists of

- (1) an abelian group  $\mathcal{F}(x)$  for every face  $x \in X$  (including the empty face),
- (2) a group homomorphism  $\text{res}_{y \leftarrow x}^{\mathcal{F}} : \mathcal{F}(x) \rightarrow \mathcal{F}(y)$  for every pair of faces  $x, y \in X$  with  $x \subsetneq y$ ,

subject to the requirement  $\text{res}_{z \leftarrow y}^{\mathcal{F}} \circ \text{res}_{y \leftarrow x}^{\mathcal{F}} = \text{res}_{z \leftarrow x}^{\mathcal{F}}$  whenever  $x \subsetneq y \subsetneq z$ . The maps  $\text{res}_{y \leftarrow x}^{\mathcal{F}}$  are called the *restriction maps* of  $\mathcal{F}$ . A sheaf is an augmented sheaf with  $\mathcal{F}(\emptyset) = 0$ ; informally, this means that the empty face is ignored in the definition.

If  $R$  is an abelian group, then we can define an augmented sheaf  $\mathcal{F}$  on  $X$  by setting  $\mathcal{F}(x) = R$  for all  $x \in X$ , and letting all the restriction maps be  $\text{id}_R$ . This augmented sheaf is called the *constant augmented sheaf* associated to  $R$  and denoted  $R_+$ . If we were to set  $\mathcal{F}(\emptyset) = 0$  instead of  $\mathcal{F}(\emptyset) = R$ , the result would be the *constant sheaf* associated to  $R$ .

### Coboundary and Cosystolic Expansion of Sheaves

If  $\mathcal{F}$  is an augmented sheaf on  $X$ , then the  $i$ -cochains of  $X$  with coefficients in  $\mathcal{F}$  are members of  $C^i(X, \mathcal{F}) = \prod_{x \in X} \mathcal{F}(x)$ ; here  $i \in \mathbb{N} \cup \{0, -1\}$ . Similarly to the case of  $\mathbb{F}_2$ -coefficients, one can define the  $i$ -th coboundary map  $d_i : C^i(X, \mathcal{F}) \rightarrow C^{i+1}(X, \mathcal{F})$  by<sup>5</sup>

$$(d_i f)(y) = \sum_{x \text{ is an } i\text{-face of } y} \text{res}_{y \leftarrow x}^{\mathcal{F}}(f(x))$$

and consider the abelian groups of  $i$ -coboundaries and  $i$ -cochains, denoted  $B^i(X, \mathcal{F})$  and  $Z^i(X, \mathcal{F})$ , respectively. The notions of  $\varepsilon$ -coboundary expansion and  $(\delta, \varepsilon)$ -cosystolic expansion in dimension  $i$  extend naturally to pairs  $(X, \mathcal{F})$  consisting of a simplicial complex and an augmented sheaf. The Hamming distance on  $C^i(X, \mathcal{F})$  is defined by  $d_{\text{Ham}}(f, g) = \#\{x \in X(i) : f(x) \neq g(x)\}$ , or as the weight of the latter set in case weights are present.

The discussion at 1A can now be regarded as addressing the special case of coboundary and cosystolic expansion of constant augmented sheaves, particularly that of  $(\mathbb{F}_2)_+$ .

## 1C Purpose of This Paper

This paper concerns with finding new examples of triples  $(X, \mathcal{F})$  consisting of a weighted simplicial complex  $X$  and a sheaf  $\mathcal{F}$  which together form a good coboundary expander in dimension 0. As in the case of constant augmented sheaves, such examples can be used to produce pairs  $(X, \mathcal{F})$  which are good cosystolic expanders, which in turn give rise to *locally testable codes*; see [8].

We focus particularly on the following general question: Suppose that  $(X, \mathcal{F})$  is known to be a good coboundary expander in dimension 0. Does  $(X, \mathcal{F})$  remain a good coboundary expander in dimension 0 if we make mild modifications to  $\mathcal{F}$ ?

Broadly speaking, we show that if  $\mathcal{F}$  is a constant augmented sheaf and the modification consists of taking a *quotient* of  $\mathcal{F}$  by a “small” *subsheaf*, then the answer is “yes”, i.e., the quotient is a good coboundary expander. This is the first example of non-constant sheaves with good coboundary expansion. On the other

<sup>5</sup>Signs should be introduced if some of the groups  $\mathcal{F}(x)$  are not elementary abelian 2-groups; see §4B.

hand, if we modify the same  $\mathcal{F}$  by changing a few of its restriction maps, then this may result in poor coboundary expansion even if the underlying graph and  $\mathcal{F}$  are both excellent expanders.

In addition, we show that any constant augmented sheaf  $R_+$  on an  $\varepsilon$ -expander graph is a good coboundary expander in dimension 0; as noted earlier, this was known only in the case of  $(\mathbb{F}_2)_+$  (equiv. for  $\mathbb{F}_2$  coefficients). Moreover, we relate the coboundary expansion of different locally constant sheaves, showing that if  $R$  and  $S$ , then the 0-dimensional coboundary expansion of  $(X, R_+)$  is at most twice the 0-dimensional coboundary expansion of  $(X, S_+)$ .

Along the way, we prove a fine version of the Expander Mixing Lemma for weighted  $r$ -partite expander graphs, which may be of independent interest.

## 1D Main Results

All of our results implicitly assume that a weight function on the simplicial complex or graph under discussion has been specified; see §2B for details. A weighted graph is said to an  $\varepsilon$ -expander if  $h(X, w) \geq \varepsilon$  (see (1.1)), and a  $[\mu, \lambda]$ -spectral expander, or just  $[\mu, \lambda]$ -expander for short, if all eigenvalues of its weighed adjacency operator (see §2C) except the largest — which is always 1 — live in the interval  $[\mu, \lambda]$ . For example, if  $X$  is a  $k$ -regular graph, and its (non-weighted) adjacency matrix is  $A$ , then  $X$  is a  $[\mu, \lambda]$ -expander if and only if all eigenvalues of  $A$  except  $k$  live in the interval  $[\mu k, \lambda k]$ .

### Constant and Locally Constant Sheaves on Graphs

We show that if  $X$  is an  $\varepsilon$ -expander, then  $(X, R_+)$  is a  $\frac{\varepsilon}{2}$ -coboundary expander in dimension 0 for every nontrivial abelian group  $R$  (Theorem 5.2 and (2.2)). Similarly, if  $X$  is a  $[-1, \lambda]$ -spectral expander, then  $(X, R_+)$  is a  $(1 - \lambda)$ -coboundary expander in dimension 0 (Corollary 5.2). Moreover, if  $S$  is another nontrivial abelian group and  $(X, R_+)$  is known to be an  $\varepsilon$ -coboundary expander in dimension 0, then  $(X, S_+)$  is an  $\frac{\varepsilon}{2}$ -coboundary expander in dimension 0 (Corollary 5.4).

The sheaves most resembling the constant augmented sheaves on  $X$  are the *locally constant sheaves*, also known as *local systems* (cf. [12, §2]). Specifically, a sheaf  $\mathcal{F}$  on a graph  $X$  is called locally constant if  $\text{res}_{e \leftarrow v} : \mathcal{F}(v) \rightarrow \mathcal{F}(e)$  is an isomorphism for every  $e \in X(1)$ ,  $v \in X(0)$  with  $v \subseteq e$ . (We have  $\mathcal{F}(\emptyset) = 0$  because  $\mathcal{F}$  is sheaf, rather than an augmented sheaf.) Contrary to the expectations, we show in Example 5.6 that every expander graph admits a locally constant sheaf  $\mathcal{F}$  such that  $(X, \mathcal{F})$  is a poor coboundary expander in dimension 0. This sheaf is obtained by mildly modifying the restriction maps of a constant sheaf, showing that such modifications could drastically affect the coboundary expansion.

### Quotients of Constant Augmented Sheaves

Let  $R$  be an abelian group and consider the augmented sheaf  $R_+$  on a graph  $X$ . For every vertex  $v \in X(0)$  and edge  $e \in X(1)$ , pick subgroups  $R_v, R_e \subseteq R$ . We use this data to define a subsheaf  $\mathcal{C}$  of  $R_+$  as follows:

1.  $\mathcal{C}(\emptyset) = 0$ ,
2.  $\mathcal{C}(v) = R_v$  for all  $v \in X(0)$ , and
3.  $\mathcal{C}(e) = R_u + R_v + R_e$  for all  $e \in X(1)$ , where  $u, v$  are the vertices of  $e$ .

This is the smallest subsheaf  $\mathcal{C}$  of  $R_+$  with  $R_x \subseteq \mathcal{C}(x)$  for all  $x \in X(0) \cup X(1)$ . If the groups  $\{R_x\}_{x \in X(0) \cup X(1)}$  are “small” with respect to  $R$ , then  $\mathcal{F} = R_+/\mathcal{C}$  can be regarded as a “small” modification of  $R_+$ .

Suppose that the following conditions holds:

- (1) For every subgraph  $Y$  of  $X$  which is either a cycle of length  $\leq \lceil \frac{2}{3}|X(0)| \rceil$  or a path of a length  $\leq 2$  (see §2A), the summation map  $\bigoplus_{y \in Y(0) \cup Y(1)} R_y \rightarrow R$  is injective.
- (2) For every distinct  $u, v \in X(0)$ , we have  $R_u \cap R_v = 0$ .

We show in Theorem 6.1, we that if  $X$  is a  $[-\lambda, \lambda]$ -expander, then  $(X, \mathcal{F})$  is an  $\varepsilon$ -coboundary expander in dimension 0 where  $\varepsilon = \frac{2}{5} - O(\lambda) - O(t)$ . Here,  $t$  is a certain constant depending on how uniform the weights

on  $X$  are; it equals  $\frac{1}{k}$  if  $X$  is a  $k$ -regular graph. We also prove a version of this result to  $(r+1)$ -partite graphs — Theorem 6.2 — which allows us to ignore the eigenvalue  $-\frac{1}{r}$  that is always present in the  $r$ -partite case.

The proof of theorems 6.1 and 6.2 is quite involved. We do not know if the linear independence condition (1) can be relaxed in general, but we expect it could be. See Section 8 for further discussion about this question.

### The Case of Buildings

Recall that finite *spherical buildings* are known to be excellent coboundary expanders with respect to every constant augmented sheaf. Refining results of Evra and the second author [7], we show that their underlying graphs are also excellent partite spectral expanders (Theorem 7.2), and thus our Theorem 6.2 can be applied to them. Letting  $X$  denote the 1-dimensional skeleton of a  $q$ -thick finite spherical building of dimension  $r$ , and letting  $R$  and  $\mathcal{F}$  be as above, we show in Corollary 7.6 that  $(X, \mathcal{F})$  is a  $(\frac{2r}{5r+2} - O(\frac{1}{\sqrt{q}}))$ -cosystolic expander in dimension 0, provided  $(*)$  holds.

### Mixing Lemmas for Weighted and Partite Graphs

As a means to establish the previous results, we prove a fine version of the Expander Mixing Lemma applying to weighted graphs (Theorem 3.2), and another version applying to weighted  $r$ -partite graphs (Theorem 3.2). These results, which we could not find in the literature, will likely be useful elsewhere.

## 1E Implications to Locally Testable Codes

Generalizing the works [13], [7], [15] (which address the case of constant sheaves in our terminology), the authors showed in [8] that if  $(X, \mathcal{F})$  is a sheaved simplicial complex such that the restriction of  $\mathcal{F}$  to every proper link of  $X$  is a good coboundary expander in a range of dimensions, then  $(X, \mathcal{F})$  is a good cosystolic expander in a desired dimension  $i$ . In order to apply this result, one needs a source of examples of sheaved simplicial complexes that are good coboundary expanders, which is precisely what this paper provides, albeit in dimension 0. As shown in [8], the 0-dimensional case already has implications to the open problem of constructing *2-query locally testable codes*, and our results here play a critical role in the constructions of such codes suggested in *op. cit.* Similarly, higher dimensional generalizations of our results may have implications to the construction of *quantum CSS codes*.

## 1F Outline

The paper is organized as follows: Section 2 is preliminary and recalls necessary facts about simplicial complexes, weight functions, spectral expansion and partite graphs. In Section 3 we prove weighted versions of the Cheeger Inequality and the Expander Mixing Lemma. Section 4 recalls sheaves on graphs, their cohomology and their expansion. In Section 5, we show that an expander graph with a constant augmented sheaf is a good coboundary expander in dimension 0, but that this may fail for locally constant sheaves. The subject matter of Section 6 is proving that taking a quotient of a constant augmented sheaf by a small subsheaf still results in a good coboundary expander in dimension 0, provided the underlying graph is a sufficiently good expander. This result is applied to finite spherical buildings in Section 7. Finally, in Section 8, we raise some questions about possible extensions of our results.

## Acknowledgments

Tali Kaufman's research is supported by an ERC grant and an ISF grant.

## 2 Preliminaries

### 2A Simplicial Complexes

If not indicated otherwise, simplicial complexes and graphs are assumed to be finite. A graph is a 1-dimensional simplicial complex. In particular, we do not allow graphs to have loops or double edges (but double edges

can be accounted for using weight functions discussed in §2B).

Let  $X$  be a simplicial complex. We write  $V(X)$  for the vertex set of  $X$  and  $X(i)$  ( $-1 \leq i \in \mathbb{Z}$ ) for the set of  $i$ -faces of  $X$ . Note that a vertex and a 0-face are not the same thing, namely,  $v \in V(x)$  if and only if  $\{v\} \in X(0)$ . When there is no risk of confusion, we will abuse the notation and treat vertices as 0-faces and vice versa. The  $k$ -dimensional skeleton of  $X$  is the simplicial complex  $X(\leq k) := \bigcup_{i=-1}^k X(i)$ . Given  $z \in X$  and  $A \subseteq X$ , we write

$$\begin{aligned} A_{\supseteq z} &= \{y \in A : y \supseteq z\}, \\ A_{\subseteq z} &= \{y \in A : y \subseteq z\}, \\ A_z &= \{y - z \mid y \in A_{\supseteq z}\}. \end{aligned}$$

For example, if  $x \in X(0)$  is a 0-face, then  $X(1)_{\supseteq x}$  is the set of edges (i.e. 1-faces) containing  $x$ , and  $X(1)_x$  is the set 0-faces adjacent to  $x$ . Also, for any  $z \in X$ , the set  $X_z$  is the *link* of  $X$  at  $z$ ; it is a simplicial complex.

An ordered edge in  $X$  is a pair  $(u, v)$  such that  $\{u, v\}$  is an edge in  $X$ . The set of ordered edges in  $X$  is denoted  $X_{\text{ord}}(1)$ . If  $A, B \subseteq X(0)$ , then we write  $E(A, B)$  for the set of edges in  $X$  with one vertex in  $A$  and the other in  $B$ . Likewise,  $E_{\text{ord}}(A, B)$  is the set of ordered edges  $(u, v) \in X_{\text{ord}}(1)$  with  $\{u\} \in A$  and  $\{v\} \in B$ . We also let  $E(A) = E(A, A)$  and  $E_{\text{ord}}(A) = E_{\text{ord}}(A, A)$ .

The simplicial complex  $X$  is said to be *pure* of dimension  $d$  ( $0 \leq d \in \mathbb{Z}$ ) if every face of  $X$  is contained in a  $d$ -face. We then say that  $X$  is a  $d$ -*complex* for short. The  $k$ -dimensional skeleton of a  $d$ -complex is a  $k$ -complex for all  $k \in \{0, \dots, d\}$ .

Suppose now that  $X$  is a graph and let  $Y \subseteq X$  be a subset. We say that  $Y$  is a *cycle* of length  $\ell$  ( $\ell \geq 3$ ) if  $Y$  is a subgraph of  $X$  that is isomorphic to the cycle graph on  $\ell$  vertices. Given  $x, y \in X(0)$  ( $x = y$  is allowed), we say that  $Y$  a *closed path*<sup>6</sup> of length  $\ell$  ( $\ell \geq 0$ ) from  $x$  to  $y$  if  $Y$  is a subgraph of  $X$  such that  $x, y \in Y(0)$ ,  $Y(0) - \{x, y\}$  contains exactly  $\ell - 1$  distinct 0-faces  $x_1, \dots, x_{\ell-1}$ , and  $Y(1)$  consists of exactly  $\ell$  edges which are  $x \cup x_1, x_1 \cup x_2, \dots, x_{\ell-2} \cup x_{\ell-1}, x_{\ell-1} \cup y$ . An *open path* from  $x$  to  $y$  is a subset  $Z \subseteq X$  of the form  $Y - \{x, y\}$ , where  $Y$  is a path from  $x$  to  $y$  in  $X$ ; the length of  $Z$  is defined to be the length of  $Y$ . An open path  $Z$  is *not* a subgraph of  $X$ , but we shall still write  $Z(i) = Z \cap X(i)$  for  $i \in \{0, 1\}$ .

The following easy lemma, whose proof is left to the reader, will be needed in the sequel.

**Lemma 2.1.** *Let  $X$  be a graph, let  $X'$  be a subgraph of  $X$  and let  $C$  be a cycle in  $X$  meeting  $X'$ . Then  $C - X'$  is a disjoint union of open paths.*

## 2B Weights

A weight function on  $d$ -complex  $X$  is a function  $w : X \rightarrow \mathbb{R}_+$  such that

$$(W1) \quad \sum_{y \in X(d)} w(y) = 1,$$

$$(W2) \quad w(x) = \binom{d+1}{\dim x + 1}^{-1} \sum_{y \in X(d)_{\supseteq x}} w(y) \text{ for all } x \in X \text{ with } \dim x < d.$$

We then say that  $(X, w)$  is a *weighted  $d$ -complex*; if  $d = 1$  we say that  $X$  is a *weighted graph*. If we strengthen (W1) to  $w(y) = |X(d)|^{-1}$  for all  $y \in X(d)$ , then this recovers the weight functions considered in [24] and [15].

Given a weight function  $w : X \rightarrow \mathbb{R}_+$  and  $A \subseteq X$ , we write  $w(A) = \sum_{a \in A} w(a)$ . A similar convention applies to subsets of  $X_{\text{ord}}(1)$ , where the weight of an ordered edge is defined to be the weight of its underlying unordered edge.

The conditions (W1) and (W2) imply that  $w$  defines a probability measure on  $X(d)$ , and that for  $x \in X(i)$  with  $i \in \{-1, \dots, d-1\}$ , the value  $w(x)$  is the probability of obtaining  $x$  by choosing a  $d$ -face  $y \in X(d)$  according to  $w$  and then choosing an  $i$ -face of  $y$  randomly uniformly. Consequently,  $w(X(i)) = 1$  for all  $i \in \{-1, \dots, d-1\}$ . It is also straightforward to check that (W2) implies

$$w(X(\ell)_{\supseteq x}) = \binom{\ell+1}{k+1} w(x) \tag{2.1}$$

for all  $-1 \leq k \leq \ell \leq d$  and  $x \in X(k)$ . As a result, if  $(X, w)$  is a weighted  $d$ -complex, then for every  $k \in \{0, \dots, d\}$ , its  $k$ -dimensional skeleton  $(X(\leq k), w|_{X(\leq k)})$  is a weighted  $k$ -complex.

<sup>6</sup>Here, ‘‘closed’’ should be understood as topologically closed, rather than having the same start and end point.

**Example 2.2.** (i) Let  $X$  be a  $d$ -complex. The *canonical weight function*  $w_X : X \rightarrow \mathbb{R}_+$  is defined by putting  $w_X(y) = \frac{1}{|X(d)|}$  for all  $y \in X(d)$  and defining  $w_X$  on the other faces using the formula in (W2).

(ii) If  $X$  is a  $k$ -regular graph on  $n$  vertices, then  $X$  is a pure 1-dimensional simplicial complex, and the canonical weight function assigns every edge of  $X$  the weight  $\frac{2}{kn}$ , every vertex of  $X$  the weight  $\frac{1}{n}$  and the value 1 to the empty face.

(iii) If  $X$  is a  $d$ -complex, and  $0 \leq k < d$ , then  $(X(\leq k), w_X|_{X(\leq k)})$  is a weighted  $k$ -complex, but  $w_X|_{X(\leq k)}$  is in general different from  $w_{X(\leq k)}$ .

**Remark 2.3.** In [27, Definition 2.1] and [11, Definition 3.2], a balanced weight function on a  $d$ -complex is defined to be a function  $m : X \rightarrow \mathbb{R}_+$  such that  $m(x) = (d - \dim x)! \cdot m(X(d) \supseteq x)$  for all  $x \in X$ . If  $(X, w)$  in a weighted  $d$ -complex in our sense, then  $m : X \rightarrow \mathbb{R}_+$  defined by  $m(x) = \frac{(d+1)!}{(\dim x + 1)!} w(x)$  is a balanced weight function in the sense of [27] and [11].

## 2C Expansion of Weighted Graphs

Let  $(X, w)$  be a weighted graph. Given  $i \in \{0, 1\}$ , write  $C^i(X, \mathbb{R})$  for the set of functions  $f : X(i) \rightarrow \mathbb{R}$ . We endow  $C^i(X, \mathbb{R})$  with the inner product defined by

$$\langle f, g \rangle = \frac{1}{(i+1)!} \sum_{x \in X(i)} f(x)g(x)w(x)$$

for all  $f, g \in C^i(X, \mathbb{R})$ . Given  $A \subseteq X(0)$ , we write  $1_A$  for the function in  $C^0(X, w)$  taking the value 1 on  $A$  and 0 on  $X(0) - A$ .

The *weighted adjacency operator* of  $(X, w)$ , denoted  $\mathcal{A} = \mathcal{A}_{X,w}$ , and the *weighted Laplacian* of  $(X, w)$ , denoted  $\Delta = \Delta_{X,w}$ , are the linear operators from  $C^0(X, \mathbb{R})$  to itself defined by

$$\begin{aligned} (\mathcal{A}f)(x) &= \sum_{e \in X(1) \supseteq x} \frac{w(e)}{2w(x)} f(e-x), \\ (\Delta f)(x) &= f(x) - \sum_{e \in X(1) \supseteq x} \frac{w(e)}{2w(x)} f(e-x), \end{aligned}$$

for all  $f \in C^0(X, \mathbb{R})$  and  $x \in X(0)$ . We have  $\mathcal{A} = \text{id} - \Delta$ . For example, if  $X$  is a  $k$ -regular graph and  $w$  is the canonical weight function of  $X$ , then  $\mathcal{A}_{X,w}$  is the ordinary adjacency operator of  $X$  scaled by  $\frac{1}{k}$ , cf. Example 2.2(ii).

Fix a linear ordering  $L$  on  $V(X)$ . Given an edge  $e = \{u, v\} \in X(1)$  with  $u < v$  relative to  $L$ , we write  $e^+ = \{v\}$  and  $e^- = \{u\}$ . The *0-coboundary map* is the linear operator  $d_0 = d_0^L : C^0(X, \mathbb{R}) \rightarrow C^1(X, \mathbb{R})$  defined by

$$(d_0 f)(e) = f(e^+) - f(e^-),$$

for all  $f \in C^0(X, \mathbb{R})$ ,  $e \in X(1)$ . The weighted 1-*boundary map* is the dual operator  $d_0^* : C^1(X, \mathbb{R}) \rightarrow C^0(X, \mathbb{R})$  relative to the inner products of  $C^0(X, \mathbb{R})$  and  $C^1(X, \mathbb{R})$ .

**Lemma 2.4.** *Under the previous assumptions:*

(i)  $\Delta = d_0^* d_0$ . In particular,  $\Delta$  is positive semidefinite and  $\mathcal{A}$  is self-adjoint.

(ii)  $\text{Spec } \Delta \subseteq [0, 2]$  and  $\text{Spec } \mathcal{A} \subseteq [-1, 1]$ .

(iii)  $\Delta 1_{X(0)} = 0$  and  $\mathcal{A} 1_{X(0)} = 1_{X(0)}$ .

*Proof.* (i) This follows from [27, Proposition 2.11]. (Consult Remark 2.3. For a general weighted  $d$ -complex  $(X, w)$ , the inner product of  $C^i(X, \mathbb{R})$  used in *op. cit.* is given by  $\langle f, g \rangle = \frac{(d+1)!}{(i+1)!} \sum_{x \in X(i)} f(x)g(x)w(x)$ . The Laplacian  $\Delta$  is denoted  $\Delta_0^+$  in *op. cit.*)



(ii) By (i), it is enough to prove that  $\|\mathcal{A}\| \leq 1$ . Let  $f \in C^0(X, \mathbb{R})$  and  $x \in X(0)$ . Then  $\sum_{e \in X(1) \supseteq x} \frac{w(e)}{2w(x)} = \frac{2w(x)}{2w(x)} = 1$ , so by Jensen's inequality,

$$\left[ \sum_{e \in X(1) \supseteq x} \frac{w(e)}{2w(x)} f(e-x) \right]^2 \leq \sum_{e \in X(1) \supseteq x} \frac{w(e)}{2w(x)} [f(e-x)]^2.$$

Using this, we get

$$\begin{aligned} \|\mathcal{A}f\|^2 &= \sum_{x \in X(0)} w(x) \left[ \sum_{e \in X(1) \supseteq x} \frac{w(e)}{2w(x)} f(e-x) \right]^2 \\ &\leq \sum_{x \in X(0)} w(x) \sum_{e \in X(1) \supseteq x} \frac{w(e)}{2w(x)} [f(e-x)]^2 = \sum_{x \in X(0)} \sum_{e \in X(1) \supseteq x} \frac{w(e)}{2} f(e-x)^2 \\ &= \sum_{y \in X(0)} \sum_{e \in X(1) \supseteq y} \frac{w(e)}{2} f(y)^2 = \sum_{y \in X(0)} w(y) f(y)^2 = \|f\|^2, \end{aligned}$$

which is what we want.

(iii) Let  $x \in X(0)$ . Then  $(\mathcal{A}1_{X(0)})(x) = \sum_{e \in X(1) \supseteq x} \frac{w(e)}{2w(x)} = 1 = 1_{X(0)}(x)$ .  $\square$

Let  $(X, w)$  be a weighted graph. We define

$$C_{\circ}^0(X, \mathbb{R}) = 1_{X(0)}^{\perp} = \{f \in C^0(X, \mathbb{R}) : \sum_{x \in X(0)} w(x) f(x) = 0\}.$$

By Lemma 2.4,  $\Delta$  and  $\mathcal{A}$  take  $C_{\circ}^0(X, \mathbb{R})$  to itself. Given  $\mu \leq \lambda$  in  $\mathbb{R}$ , we say that  $(X, w)$  is a  $[\mu, \lambda]$ -spectral expander, or just  $[\mu, \lambda]$ -expander for short, if  $\text{Spec}(\mathcal{A}|_{C_{\circ}^0(X, \mathbb{R})}) \subseteq [\mu, \lambda]$ .

**Remark 2.5.** (i) At this level of generality, a weighted graph can be a  $[\mu, \lambda]$ -expander even when both  $\mu$  and  $\lambda$  are negative. For example, consider a complete graph  $X$  on  $n+1$  vertices with its canonical weight function  $w = w_X$ . It is well-known that eigenvalues of  $\mathcal{A}_{X, w}$  are  $1, -\frac{1}{n}, \dots, -\frac{1}{n}$  (including multiplicities), so  $(X, w)$  is a  $[-\frac{1}{n}, -\frac{1}{n}]$ -expander.

(ii) We shall see below that if  $(X, w)$  is a bipartite weighted graph, then  $-1 \in \text{Spec} \mathcal{A}_{X, w}$ , so such a weighted graph is a  $[\mu, \lambda]$ -expander only when  $\mu \leq -1$ .

We now recall Kaufman and Oppenheim's version of the Cheeger Inequality for graphs [18, Theorem 4.4(1)], which relates  $[-1, \lambda]$ -expansion and the Cheeger constant  $h(X, w)$  (see the Introduction). To that end, it is convenient to introduce the following variation on the Cheeger constant  $h(X, w)$ , namely,

$$h'(X, w) = \min_{\emptyset \neq A \subsetneq X(0)} \frac{w(E(A, X(0) - A))}{2w(A)w(X(0) - A)}$$

(this is denoted  $h_G$  in *op. cit.*). Informally,  $h'(X, w)$  is minimum possible ratio between the weight of the edges leaving  $A$ , and the expected weight if  $(X, w)$  were to behave like a random graph. Since  $\min\{\alpha, 1 - \alpha\} \leq 2\alpha(1 - \alpha) \leq 2 \min\{\alpha, 1 - \alpha\}$  for all  $\alpha \in [0, 1]$ , we have

$$\frac{1}{2}h(X, w) \leq h'(X, w) \leq h(X, w). \quad (2.2)$$

**Theorem 2.6** ([18, Theorem 4.4(1)]). *Let  $(X, w)$  be a weighted graph which is also a  $[-1, \lambda]$ -expander. Then*

$$h'(X, w) \geq 1 - \lambda.$$

*That is, for every  $A \subseteq X$ , we have  $w(E(A, X(0) - A)) \geq (1 - \lambda) \cdot 2w(A)(1 - w(A))$ .*

A theorem of Friedland and Nabben [9, Theorem 2.1] implies a converse to the theorem, namely, if  $h(X, w) \geq \varepsilon$ , then  $(X, w)$  is a  $[-1, \sqrt{1 - \frac{\varepsilon^2}{4}}]$ -expander. (Specifically, assuming  $V(X) = \{1, \dots, n\}$ , apply [9, Theorem 2.1] with  $w_{i,j} = w(\{i, j\})$  and  $d_i = 2w(\{i\})$ . The numbers  $\delta_i$  defined *op. cit.* are  $2w(\{i\})$  in our notation, and constant  $i(X, w)$  considered there equals  $\frac{1}{2}h(X, w)$  in our notation.)

## 2D Partite Simplicial Complexes

Recall that an  $(r+1)$ -partite simplicial complex is a tuple  $(X, V_0, \dots, V_r)$  such that  $X$  is a simplicial complex,  $V_0, \dots, V_r$  is a partition of the set of vertices  $V(X)$  (in particular,  $V_i \neq \emptyset$  for all  $i$ ), and every face  $x \in X$  contains at most one vertex from each  $V_i$ , i.e.,  $|x \cap V_i| \leq 1$  for all  $i \in \{0, \dots, r\}$ . We then write  $X_{\{i\}}$  for the set of 0-faces having their vertex in  $V_i$ . We say that  $(X, V_0, \dots, V_r)$  is pure if every face of  $X$  is contained in an  $(r+1)$ -face; in this case  $\dim X = r$ . A bipartite graph is just 2-partite simplicial complex.

A *weighted  $(r+1)$ -partite simplicial complex* is a tuple  $(X, w, V_0, \dots, V_r)$  such that  $(X, V_0, \dots, V_r)$  is a pure  $(r+1)$ -partite simplicial complex and  $w$  is a weight function on the  $r$ -complex  $X$ .

When there is no risk of confusion, we will sometimes abbreviate the partition  $(V_0, \dots, V_r)$  from the notation, writing simply that  $X$  is an  $(r+1)$ -partite complex, or that  $(X, w)$  is a weighted  $(r+1)$ -partite simplicial complex.

**Lemma 2.7.** *Let  $(X, w, V_0, \dots, V_r)$  be a weighted pure  $(r+1)$ -partite simplicial complex. Then:*

(i)  $w(X_{\{i\}}) = \frac{1}{r+1}$  for all  $i \in \{0, \dots, r\}$ .

(ii)  $w(E(X_{\{i\}}, X_{\{j\}})) = \frac{2}{r(r+1)}$  for all distinct  $i, j \in \{0, \dots, r\}$ .

(iii) *The subspace  $L$  of  $C^0(X, \mathbb{R})$  spanned by  $1_{X_{\{0\}}}, \dots, 1_{X_{\{r\}}}$  is invariant under  $\mathcal{A} = \mathcal{A}_{X,w}$ . The  $r+1$  eigenvalues of  $\mathcal{A}$  on this subspace are  $1, -\frac{1}{r}, \dots, -\frac{1}{r}$  (including multiplicities). If the link  $X_z$  is connected for all  $z \in X(\leq r-2)$ , then  $L$  is the sum of the 1-eigenspace and  $-\frac{1}{r}$ -eigenspace of  $\mathcal{A}$ .*

*Proof.* We prove (i) and (ii) together. Given  $I \subseteq \{0, \dots, r\}$ , let  $X_I$  denote the set of  $(|I|-1)$ -faces of  $X$  having a vertex in  $V_i$  for all  $i \in I$ . We claim that  $w(X_I) = \binom{r+1}{|I|}^{-1}$ . Indeed,

$$\begin{aligned} w(X_I) &= \sum_{x \in X_I} w(x) = \sum_{x \in X_I} \binom{r+1}{|I|}^{-1} \sum_{y \in X(r+1) \supseteq x} w(y) \\ &= \binom{r+1}{|I|}^{-1} \sum_{y \in X(r+1)} \sum_{x \in (X_I) \subseteq y} w(y) = \binom{r+1}{|I|}^{-1} \sum_{y \in X(r+1)} 1 \cdot w(y) \\ &= \binom{r+1}{|I|}^{-1} w(X(r+1)) = \binom{r+1}{|I|}^{-1}, \end{aligned}$$

where the fourth equality holds because every  $(r+1)$ -face contains exactly one face in  $X_I$ . (i) and (ii) now follow by taking  $I = \{i\}$  and  $I = \{i, j\}$ .

Part (iii) follows from [27, Proposition 5.2] and its proof.  $\square$

Given a weighted pure  $(r+1)$ -partite simplicial complex  $(X, w, V_0, \dots, V_r)$ , we write  $C_{\diamond}^0(X, \mathbb{R})$  for  $L^\perp$ , where  $L = \text{span}_{\mathbb{R}}\{1_{X_{\{0\}}}, \dots, 1_{X_{\{r\}}}\}$ . In view of Lemma 2.7(iii), when regarding the spectrum of  $\mathcal{A} = \mathcal{A}_{X,w}$  on  $C^0(X, \mathbb{R})$ , it is reasonable to set aside the eigenvalues occurring on the subspace  $L$ . Thus, we say that  $(X, w, V_0, \dots, V_r)$  is an  $(r+1)$ -partite  $[\mu, \lambda]$ -expander if the spectrum of  $\mathcal{A}$  on  $C_{\diamond}^0(X, \mathbb{R})$  is contained in the interval  $[\mu, \lambda]$ .

Oppenheim [27, Lemma 5.5]<sup>7</sup> observed that if  $(X, w)$  is a weighted  $r$ -partite  $[-1, \lambda]$ -expander with  $0 \leq \lambda \leq \frac{1}{r}$ , then  $(X, w)$  is in fact an  $r$ -partite  $[-r\lambda, \lambda]$ -expander. In fact, when  $(X, w)$  is a weighted bipartite graph,  $\text{Spec } \mathcal{A}_{X,w}$  is symmetric around 0; this follows from the following lemma, whose straightforward proof is left to the reader.

**Lemma 2.8.** *Let  $(X, w, V_0, V_1)$  be a weighted bipartite graph, let  $\lambda \in \mathbb{R}$  and let  $f \in C_{\diamond}^0(X, \mathbb{R})$  be a  $\lambda$ -eigenfunction of  $\mathcal{A} = \mathcal{A}_{X,w}$ . Define  $f' \in C_{\diamond}^0(X, \mathbb{R})$  by  $f'(x) = f(x)$  if  $x \in X_{\{0\}}$  and  $f'(x) = -f(x)$  otherwise. Then  $\mathcal{A}f' = -\lambda f'$ .*

<sup>7</sup>There is a typo in this source: the last inequality should be " $1 + \frac{1}{n}(1 - \lambda(X)) \leq \kappa(X) \leq 1 + n(1 - \lambda(X))$ ".

### 3 Mixing Lemmas for Weighted Graphs

In this section, we prove two fine versions of the Expander Mixing Lemma applying to weighted graphs, and  $(r+1)$ -partite weighted simplicial complexes, respectively. The former result is well-known for non-weighted graphs, but we were unable to find the finer weighted analogue considered here in the literature. Both results will be needed in the sequel to establish our main results about coboundary expansion.

Recall that given a weighted graph  $(X, w)$  and  $A \subseteq X(0)$ , we write  $1_A \in C^0(X, \mathbb{R})$  for the function taking the value 1 on  $A$  and 0 elsewhere.

**Lemma 3.1.** *Let  $(X, w)$  be a weighted graph and let  $A, B \subseteq X(0)$ . Then*

$$\langle \mathcal{A}_{X,w} 1_A, 1_B \rangle = \frac{1}{2} w(E_{\text{ord}}(A, B)).$$

*Proof.* By unfolding the definitions,  $\langle \mathcal{A}1_A, 1_B \rangle$  evaluates to

$$\begin{aligned} \sum_{x \in X(0)} (\mathcal{A}1_A)(x) 1_B(x) w(x) &= \sum_{x \in X(0)} \sum_{e \in X(1) \supseteq x} \frac{w(e)}{2w(x)} 1_A(e-x) 1_B(x) w(x) \\ &= \frac{1}{2} \sum_{x \in X(0)} \sum_{e \in X(1) \supseteq x} 1_A(e-x) 1_B(x) w(e) = \frac{1}{2} w(E_{\text{ord}}(A, B)). \quad \square \end{aligned}$$

**Theorem 3.2** (Weighted Expander Mixing Lemma). *Let  $(X, w)$  be a weighted graph, which is also a  $[\mu, \lambda]$ -expander. Let  $A, B \subseteq X(0)$  and put  $\alpha = w(A)$ ,  $\beta = w(B)$ . Then:*

$$(i) \quad \left| \frac{1}{2} w(E_{\text{ord}}(A, B)) - \alpha\beta \right| \leq \max\{|\lambda|, |\mu|\} \sqrt{\alpha\beta(1-\alpha)(1-\beta)}.$$

$$(ii) \quad \mu\alpha(1-\alpha) \leq w(E(A)) - \alpha^2 \leq \lambda\alpha(1-\alpha).$$

*Proof.* By assumption, the eigenvalues of  $\mathcal{A}$  on  $C^0_{\circ}(X, \mathbb{R})$  live in  $[\mu, \lambda]$ . As a result, for every  $f, g \in C^0_{\circ}(X, \mathbb{R})$ , we have

$$\mu \|f\|^2 \leq \langle \mathcal{A}f, f \rangle \leq \lambda \|f\|^2, \quad (3.1)$$

$$|\langle \mathcal{A}f, g \rangle| \leq \max\{|\lambda|, |\mu|\} \|f\| \|g\|. \quad (3.2)$$

Put  $f := 1_A - \alpha 1_{X(0)}$  and  $g := 1_B - \beta 1_{X(0)}$ . As in the proof of Theorem 2.6,  $f, g \in C^0_{\circ}(X, \mathbb{R})$  and  $\|f\|^2 = \alpha(1-\alpha)$ ,  $\|g\|^2 = \beta(1-\beta)$ . As a result,

$$\begin{aligned} \langle \mathcal{A}1_A, 1_B \rangle &= \langle \mathcal{A}(\alpha 1_{X(0)} + \mathcal{A}f, \beta 1_{X(0)} + g) \rangle = \langle \mathcal{A}(\alpha 1_{X(0)}), \beta 1_{X(0)} \rangle + \langle \mathcal{A}f, g \rangle \\ &= \alpha\beta \langle 1_{X(0)}, 1_{X(0)} \rangle + \langle \mathcal{A}f, g \rangle = \alpha\beta + \langle \mathcal{A}f, g \rangle, \end{aligned}$$

where in the third equality we used Lemma 2.4(iii). By Lemma 3.1,  $\langle \mathcal{A}1_A, 1_B \rangle = \frac{1}{2} w(E_{\text{ord}}(A, B))$ , so we get

$$\frac{1}{2} w(E_{\text{ord}}(A, B)) - \alpha\beta = \langle \mathcal{A}f, g \rangle. \quad (3.3)$$

Now, by (3.1),

$$\left| \frac{1}{2} w(E_{\text{ord}}(A, B)) - \alpha\beta \right| \leq \max\{|\lambda|, |\mu|\} \|f\| \|g\| = \max\{|\lambda|, |\mu|\} \sqrt{\alpha\beta(1-\alpha)(1-\beta)},$$

which proves (i). Also, taking  $A = B$  in (3.3) and using (3.2) gives

$$\mu\alpha(1-\alpha) = \mu \|f\|^2 \leq \frac{1}{2} w(E_{\text{ord}}(A)) - \alpha^2 \leq \lambda \|f\|^2 = \lambda\alpha(1-\alpha),$$

and (ii) follows because  $\frac{1}{2} w(E_{\text{ord}}(A)) = w(E(A))$ . □

Recall from Lemma 2.7 that if  $(X, w, V_0, \dots, V_r)$  is a weighted pure  $(r+1)$ -partite simplicial complex, then  $-\frac{1}{r}$  is an eigenvalue of  $\mathcal{A}$ . As a result, the constant  $\max\{|\lambda|, |\mu|\}$  appearing in Theorem 3.2(i) is at least  $\frac{1}{r}$ . In particular, when  $X$  is a bipartite graph, this constant is 1, and Theorem 3.2(i) gives almost no information about  $w(E_{\text{ord}}(A, B))$ . We remedy this in the following theorem.

**Theorem 3.3** (Expander Mixing Lemma for Weighted  $(r+1)$ -Partite Graphs). *Let  $(X, w)$  be a weighted pure  $(r+1)$ -partite simplicial complex that is an  $(r+1)$ -partite  $[-\lambda, \lambda]$ -expander. Let  $A, B \subseteq X(0)$  and put  $\alpha = w(A)$ ,  $\beta = w(B)$ . Then:*

(i) *If there are  $T, S \subseteq \{0, \dots, r\}$  such that  $A \subseteq \cup_{i \in T} X_{\{i\}}$ ,  $B \subseteq \cup_{j \in S} X_{\{j\}}$  and  $S \cap T = \emptyset$ , then*

$$|\frac{1}{2}w(E(A, B)) - \frac{r+1}{r}\alpha\beta| \leq \lambda(r+1)\sqrt{\alpha\beta(\frac{|T|}{r+1} - \alpha)(\frac{|S|}{r+1} - \beta)}.$$

(ii) *In general,  $\frac{1}{2}w(E_{\text{ord}}(A, B)) \leq \frac{r+1}{r}\alpha\beta + \lambda r\sqrt{\alpha\beta(1-\alpha)(1-\beta)}$ .*

*Proof.* For  $i \in \{0, \dots, r\}$ , let  $A_i = A \cap X_{\{i\}}$ ,  $B_i = B \cap X_{\{i\}}$  and put  $\alpha_i = w(A_i)$ ,  $\beta_i = w(B_i)$ . Define  $f_i = 1_{A_i} - (r+1)\alpha_i 1_{X_{\{i\}}}$  and  $g_i = 1_{B_i} - (r+1)\beta_i 1_{X_{\{i\}}}$ . Since  $\text{supp}(f_i) \subseteq X_{\{i\}}$ , we have  $\langle 1_{X_{\{j\}}}, f_i \rangle = 0$  for all  $j \in \{0, \dots, r\} - \{i\}$ , whereas by Lemma 2.7(i), we also have  $\langle 1_{X_{\{i\}}}, f_i \rangle = \langle 1_{X_{\{i\}}}, 1_{A_i} \rangle - (r+1)\alpha_i \langle 1_{X_{\{i\}}}, 1_{X_{\{i\}}} \rangle = w(A_i) - (r+1)\alpha_i \frac{1}{r+1} = 0$ , so  $f_i \in C_{\diamond}^0(X, \mathbb{R})$ . Likewise,  $g_i \in C_{\diamond}^0(X, \mathbb{R})$ . We further note that

$$\begin{aligned} \|f_i\|^2 &= w(A_i)(1 - (r+1)\alpha_i)^2 + w(X_{\{i\}} - A_i)((r+1)\alpha_i)^2 \\ &= \alpha_i(1 - (r+1)\alpha_i)^2 + (\frac{1}{r+1} - \alpha_i)(r+1)^2\alpha_i^2 = \alpha_i(1 - (r+1)\alpha_i). \end{aligned}$$

Fix some distinct  $i, j \in \{0, \dots, r\}$ . By Lemmas 3.1 and Lemma 2.7(ii), we have  $\langle \mathcal{A}1_{\{X_i\}}, 1_{\{X_j\}} \rangle = \frac{1}{2}w(E(X_i, X_j)) = \frac{1}{2} \cdot \frac{2}{r(r+1)} = \frac{1}{r(r+1)}$ . Now, using Lemma 3.1 again, we find that

$$\begin{aligned} \frac{1}{2}w(E(A_i, B_j)) &= \langle \mathcal{A}1_{A_i}, 1_{B_j} \rangle \\ &= \langle \mathcal{A}((r+1)\alpha_i 1_{X_{\{i\}}}) + \mathcal{A}f_i, (r+1)\beta_j 1_{X_{\{j\}}} + g_j \rangle \\ &= \langle (r+1)\alpha_i \mathcal{A}1_{X_{\{i\}}}, (r+1)\beta_j 1_{X_{\{j\}}} \rangle + \langle f_i, g_j \rangle \\ &= (r+1)^2\alpha_i\beta_j \cdot \frac{1}{r(r+1)} + \langle \mathcal{A}f_i, g_j \rangle = \frac{r+1}{r}\alpha_i\beta_j + \langle \mathcal{A}f_i, g_j \rangle. \end{aligned} \tag{3.4}$$

Since  $(X, w)$  is an  $(r+1)$ -partite  $[-\lambda, \lambda]$ -expander, we have

$$\langle \mathcal{A}f_i, g_j \rangle \leq \lambda \|f_i\| \|g_j\| = \lambda \sqrt{\alpha_i\beta_i(1 - (r+1)\alpha_i)(1 - (r+1)\beta_j)}.$$

Together with (3.4), this implies that

$$|\frac{1}{2}w(E(A_i, B_j)) - \frac{r+1}{r}\alpha_i\beta_j| \leq \lambda \sqrt{\alpha_i\beta_i(1 - (r+1)\alpha_i)(1 - (r+1)\beta_j)}. \tag{3.5}$$

It is routine, yet tedious, to check that the real two-variable function  $h(x, y) = \sqrt{xy(1 - (r+1)x)(1 - (r+1)y)}$  is concave on  $[0, \frac{1}{r+1}] \times [0, \frac{1}{r+1}]$ .<sup>8</sup> Thus, by Jensen's inequality, for any sequence of points  $\{(x_k, y_k)\}_{k=1}^t$  in the square  $[0, \frac{1}{r+1}]^2$ , we have  $\sum_{k=1}^t h(x_k, y_k) \leq th(\sum_k \frac{x_k}{t}, \sum_k \frac{y_k}{t})$ . We apply this with together with (3.5) to prove (i) and (ii).

<sup>8</sup>Indeed, writing  $k = r+1$ , the determinant of the Hessian matrix of  $h$  evaluates to  $\frac{k^3}{4}[\frac{k}{1-kx} + \frac{1}{x} + \frac{k}{1-ky} + \frac{1}{y} - 4k]$ , which is positive on the interior of  $[0, \frac{1}{r+1}]^2$ . In addition,  $\frac{\partial^2 h}{(\partial x)^2} = -\frac{\sqrt{y(1-ky)}}{[x(1-kx)]^{3/2}}$  is negative on that domain, so the Hessian matrix is negative semidefinite.

To prove (i), we consider the points  $\{(\alpha_i, \beta_j)\}_{i \in S, j \in T}$ . By (3.5), Jensen's inequality and our assumptions on  $A$  and  $B$ , we have

$$\begin{aligned}
\left| \frac{1}{2}w(E(A, B)) - \frac{r+1}{r}\alpha\beta \right| &= \left| \sum_{i \in T} \sum_{j \in S} \frac{1}{2}w(E(A_i, B_j)) - \frac{r+1}{r} \sum_{i \in T} \sum_{j \in S} \alpha_i \beta_j \right| \\
&\leq \sum_{i \in T} \sum_{j \in S} \left| \frac{1}{2}w(E(A_i, B_j)) - \frac{r+1}{r}\alpha_i \beta_j \right| \\
&\leq \lambda \sum_{i \in T} \sum_{j \in S} \sqrt{\alpha_i \beta_j (1 - (r+1)\alpha_i)(1 - (r+1)\beta_j)} \\
&\leq \lambda |T| |S| \sqrt{\frac{\alpha}{|T|} \frac{\beta}{|S|} \left(1 - \frac{(r+1)\alpha}{|T|}\right) \left(1 - \frac{(r+1)\beta}{|S|}\right)} \\
&= \lambda(r+1) \sqrt{\alpha\beta \left(\frac{|T|}{r+1} - \alpha\right) \left(\frac{|S|}{r+1} - \beta\right)},
\end{aligned}$$

which is what we want.

To prove (ii), we consider all the points  $(\alpha_i, \beta_j)$  with  $i, j \in \{0, \dots, r\}$  and  $i \neq j$ . By (3.5) and Jensen's inequality, we have

$$\begin{aligned}
\frac{1}{2}w(E_{\text{ord}}(A, B)) &= \frac{1}{2} \sum_{i \neq j} w(E(A_i, B_j)) \\
&\leq \sum_{i \neq j} \left[ \frac{r+1}{r} \alpha_i \beta_j + \lambda \sqrt{\alpha_i \beta_j (1 - (r+1)\alpha_i)(1 - (r+1)\beta_j)} \right] \\
&\leq \frac{r+1}{r} \sum_{i, j} \alpha_i \beta_j + \lambda r(r+1) \sqrt{\frac{\alpha}{r+1} \frac{\beta}{r+1} \left(1 - \frac{(r+1)\alpha}{r+1}\right) \left(1 - \frac{(r+1)\beta}{r+1}\right)} \\
&= \frac{r+1}{r} \alpha\beta + \lambda r \sqrt{\alpha\beta(1-\alpha)(1-\beta)},
\end{aligned}$$

so we are done.  $\square$

**Remark 3.4.** In the notation of Theorem 3.3, if we write  $\alpha_i = w(A \cap X_{\{i\}})$  and  $\beta_i = w(B \cap X_{\{i\}})$  for  $i \in \{0, \dots, r\}$ , then the proof of (ii) actually gives the stronger inequality

$$\left| \frac{1}{2}w(E_{\text{ord}}(A, B)) - \frac{r+1}{r}\alpha\beta + \frac{r+1}{r} \sum_{i=0}^r \alpha_i \beta_i \right| \leq \lambda r \sqrt{\alpha\beta(1-\alpha)(1-\beta)}.$$

## 4 Sheaves on Graphs

We recall from [8] the definition of augmented sheaves on simplicial complexes, and their cohomology and coboundary expansion, focusing particularly in the case of augmented sheaves on graphs.

### 4A Sheaves on Graphs

Let  $X$  be a simplicial complex. Recall from the Introduction that an *augmented sheaf*  $\mathcal{F}$  on  $X$  consists of

- (1) an abelian group  $\mathcal{F}(x)$  for every  $x \in X$  (including the empty face), and
- (2) a group homomorphism  $\text{res}_{y \leftarrow x}^{\mathcal{F}} : \mathcal{F}(x) \rightarrow \mathcal{F}(y)$  for all  $x \subsetneq y \in X$

such that

$$\text{res}_{z \leftarrow y}^{\mathcal{F}} \circ \text{res}_{y \leftarrow x}^{\mathcal{F}} = \text{res}_{z \leftarrow x}^{\mathcal{F}} \tag{4.1}$$

whenever  $x \subsetneq y \subsetneq z \in X$ . We will usually drop the superscript  $\mathcal{F}$  from  $\text{res}_{y \leftarrow x}^{\mathcal{F}}$  when there is no risk of confusion.

A *sheaf* on  $X$  is an augmented sheaf  $\mathcal{F}$  with  $\mathcal{F}(\emptyset) = 0$ . In this case, if  $X$  is a graph, then condition (4.1) always holds.

**Example 4.1.** Let  $X$  be a simplicial complex and let  $R$  be an (additive) abelian group.

The *constant augmented sheaf* associated to  $R$  is the augmented sheaf  $\mathcal{F}$  on  $X$  determined by  $\mathcal{F}(x) = R$  for all  $x \in X$  and  $\text{res}_{y \leftarrow x}^{\mathcal{F}} = \text{id}_R$  for all  $x \subsetneq y \in X$ . We denote this  $\mathcal{F}$  by  $R_+$ .

The *constant sheaf* associated to  $R$  is the sheaf  $\mathcal{F}'$  determined by setting  $\mathcal{F}'(x) = R$  for  $x \in X - \{\emptyset\}$ ,  $\text{res}_{y \leftarrow x}^{\mathcal{F}'} = \text{id}_R$  for all  $\emptyset \neq x \subsetneq y \in X$ ,  $\mathcal{F}'(\emptyset) = 0$  and  $\text{res}_{y \leftarrow \emptyset}^{\mathcal{F}'} = 0$ . Abusing the notation, we will sometimes use  $R$  to denote the constant sheaf associated to  $R$ .

There are obvious notions of subsheaves, quotient sheaves, and homomorphisms between sheaves, see [8]. A sheaf  $\mathcal{F}$  on a graph  $X$  is said to be *constant* if it is isomorphic to the constant sheaf associated to some abelian group  $R$ . A sheaf  $\mathcal{F}$  is called *locally constant* if all the restriction map  $\text{res}_{y \leftarrow x}^{\mathcal{F}} : \mathcal{F}(x) \rightarrow \mathcal{F}(y)$  is an isomorphism for all  $\emptyset \neq x \subsetneq y \in X$ .

## 4B Sheaf Cohomology

Let  $\mathcal{F}$  be an augmented sheaf on a simplicial complex  $X$ . Fix a linear ordering  $L$  on the vertices of  $X$  and, for every  $i \in \mathbb{N} \cup \{-1, 0\}$ , let  $C^i(X, \mathcal{F})$  denote  $\prod_{x \in X(i)} \mathcal{F}(x)$ . Elements of  $C^i(X, \mathcal{F})$  are called  *$i$ -chains* with coefficients in  $\mathcal{F}$ . Given  $f \in C^i(X, \mathcal{F})$ , we write the  $x$ -component of  $f$  as  $f(x)$ . Writing  $x = \{v_0, \dots, v_i\}$  with  $v_0 < v_1 < \dots < v_i$  relative to  $L$ , we let  $x_j$  denote  $x - \{v_j\}$ . As in [8], the  $i$ -coboundary map  $d_i : C^i(X, \mathcal{F}) \rightarrow C^{i+1}(X, \mathcal{F})$  is defined by

$$(d_i f)(y) = \sum_{j=0}^{i+1} (-1)^j \text{res}_{y \leftarrow x_j} f(x_j)$$

for all  $f \in C^i(X, \mathcal{F})$ ,  $y \in X(i+1)$ . For example,  $d_{-1}$  and  $d_0$  are given by:

$$\begin{aligned} (d_{-1} f)(v) &= \text{res}_{v \leftarrow \emptyset} f(\emptyset) & \forall v \in X(0) \\ (d_0 f)(e) &= \text{res}_{e \leftarrow e^+} f(e^+) - \text{res}_{e \leftarrow e^-} f(e^-) & \forall e \in X(1), \end{aligned}$$

where here, as in §2C,  $e^+$  is the larger 0-face of  $e$  and  $e^-$  is a smaller 0-face of  $e$ , relative to  $L$ . We have  $d_i \circ d_{i-1} = 0$ . As usual, the  $i$ -coboundaries,  $i$ -cocycles and  $i$ -th cohomology of  $(X, \mathcal{F})$  are defined to be

$$B^i(X, \mathcal{F}) := \text{im } d_{i-1}, \quad Z^i(X, \mathcal{F}) := \text{im } d_i, \quad H^i(X, \mathcal{F}) = \frac{Z^i(X, \mathcal{F})}{B^i(X, \mathcal{F})},$$

respectively.

For example, if  $\mathcal{F} = R_+$  for an abelian group  $R$  (see Example 4.1), then  $B^0(X, R_+)$  is the set of constant functions from  $X(0)$  to  $R$ ,  $Z^0(X, R_+)$  is the set of functions  $f : X(0) \rightarrow R$  which are constant on each connected component of  $X$  and  $H^0(X, R_+)$  is isomorphic to  $R^{|\pi_0(X)|-1}$  and coincides with the reduced singular cohomology group  $\tilde{H}^0(X, R)$ .

## 4C Coboundary and Cosystolic Expansion

Let  $(X, w)$  be a weighted graph (see 2B) and let  $\mathcal{F}$  be an augmented sheaf on  $X$ . For  $i \in \{-1, 0, 1\}$ , the  *$w$ -support norm* on  $C^i(X, \mathcal{F})$  is the function  $\|\cdot\|_w : C^i(X, \mathcal{F}) \rightarrow \mathbb{R}$  defined by

$$\|f\|_w = w(\text{supp } f),$$

where  $\text{supp } f = \{x \in X(i) : f(x) \neq 0\}$ . This is a norm on the abelian group  $C^i(X, \mathcal{F})$  in the sense of [8], i.e.,  $\|0\|_w = 0$ ,  $\|-f\|_w = \|f\|_w$  and  $\|f+g\|_w \leq \|f\|_w + \|g\|_w$  for all  $f, g \in C^0(X, \mathcal{F})$ .<sup>9</sup> Given  $f \in C^0(X, \mathcal{F})$ , we put

$$\begin{aligned} \|f + B^0(X, \mathcal{F})\|_w &= \min\{\|f'\|_w \mid f' \in f + B^0(X, \mathcal{F})\}, \\ \|f + Z^0(X, \mathcal{F})\|_w &= \min\{\|f'\|_w \mid f' \in f + Z^0(X, \mathcal{F})\}. \end{aligned}$$

<sup>9</sup>Warning: When  $\mathcal{F}$  is the constant sheaf  $\mathbb{R}$ , the norm  $\|\cdot\|_w$  is *not* the norm induced from the inner products we defined on  $C^0(X, \mathbb{R})$  and  $C^1(X, \mathbb{R})$  in §2C. Moreover,  $(C^i(X, \mathbb{R}), \|\cdot\|_w)$  is not a normed  $\mathbb{R}$ -vector space.

Let  $\varepsilon, \delta \in [0, \infty)$ . Following [8]<sup>10</sup>, we say that  $(X, w, \mathcal{F})$  is an  $\varepsilon$ -coboundary expander in dimension 0 if

$$(B) \|d_1 f\|_w \geq \varepsilon \|f + B^0(X, \mathcal{F})\|_w \text{ for all } f \in C^0(X, \mathcal{F}),$$

and we call  $(X, w, \mathcal{F})$  an  $(\varepsilon, \delta)$ -cosystolic expander in dimension 0 if

$$(C1) \|d_1 f\|_w \geq \varepsilon \|f + Z^0(X, \mathcal{F})\|_w \text{ for all } f \in C^0(X, \mathcal{F}), \text{ and}$$

$$(C2) \|f\|_w \geq \delta \text{ for all } f \in Z^0(X, \mathcal{F}) - B^0(X, \mathcal{F}).$$

Note that both definitions do not depend on the linear ordering on  $V(X)$  chosen to define  $d_0$ . If (B) holds for some  $\varepsilon > 0$ , then  $H^0(X, \mathcal{F}) = 0$ .

**Example 4.2.** Let  $(X, w)$  be a weighted graph, and let  $\mathcal{F}$  be the augmented sheaf  $(\mathbb{F}_2)_+$  (Example 4.1). We noted in the introduction that  $(X, w, \mathcal{F})$  is an  $\varepsilon$ -coboundary expander in dimension 0 if and only if its Cheeger constant  $h(X, w)$  is at least  $\varepsilon$ . If  $X$  is moreover  $k$ -regular,  $w$  is its canonical weight function (Example 2.2), and we consider the non-weighted Cheeger constant

$$\tilde{h}(X) = \min_{\emptyset \neq S \subsetneq X(0)} \frac{|E(S, X(0) - S)|}{\min\{|S|, |X(0) - S|\}},$$

then this is also equivalent to  $\tilde{h}(X) \geq \frac{k\varepsilon}{2}$ .

The triple  $(X, w, (\mathbb{F}_2)_+)$  is an  $(\varepsilon, \delta)$ -cosystolic expander in dimension 0 if and only if each connected component of  $X$  is an  $\varepsilon$ -expander in the sense of the Introduction, and the weight of the vertices of every connected component is at least  $\delta$ .

The main results of the following sections — Theorems 5.2, 6.1, 6.2 and Corollary 7.6 — will only concern with coboundary expansion in dimension 0. However, they can be converted to statements about cosystolic expansion by means of the following remark.

**Remark 4.3.** Let  $(X, w)$  be a weighted graph, let  $\mathcal{F}$  be an augmented sheaf on  $X$  and let  $\varepsilon > 0$ . Denote by  $\mathcal{F}_0$  the subsheaf of  $\mathcal{F}$  determined by  $\mathcal{F}_0(\emptyset) = 0$  and  $\mathcal{F}_0(x) = \mathcal{F}(x)$  for  $x \neq \emptyset$ , and put  $\delta = \max\{\|d_{-1} f\|_w \mid f \in \mathcal{F}(\emptyset)\}$ . If  $(X, w, \mathcal{F})$  is an  $\varepsilon$ -coboundary expander in dimension 0, then  $(X, w, \mathcal{F}_0)$  is a  $(\varepsilon, \delta)$ -coboundary expander in dimension 0. The converse holds if  $H^0(X, \mathcal{F}) = 0$ .

## 5 Coboundary Expansion of Constant Sheaves on Graphs

We now show that if a weighted graph  $(X, w)$  is a  $[-1, \lambda]$ -spectral expander (see §2C), then  $(X, w, \mathcal{F})$  is a good cosystolic expander in dimension 0 for every constant sheaf  $\mathcal{F}$ . This is well-known when  $X$  is a  $k$ -regular graph with its canonical weight function and  $\mathcal{F} = \mathbb{F}_2$  (notation as in Example 4.1); use Example 4.2 and the Cheeger Inequality for Graphs ([2, Theorem 1.2.3] or Theorem 2.6 above).

In contrast, we show that under mild assumptions,  $X$  admits locally constant sheaves  $\mathcal{F}$  such that  $(X, w, \mathcal{F})$  has poor coboundary expansion in dimension 0.

**Lemma 5.1.** *Suppose that  $\alpha_0, \dots, \alpha_t \in [0, 1]$  satisfy  $\alpha_0 \geq \max\{\alpha_1, \dots, \alpha_t\}$  and  $\sum_{i=0}^t \alpha_i \leq 1$ . Then  $\sum_{i=0}^t \alpha_i(1 - \alpha_i) \geq \sum_{i=1}^t \alpha_i$ .*

*Proof.* After rearranging, the inequality becomes  $\sum_{i=1}^t \alpha_i^2 \leq \alpha_0(1 - \alpha_0)$ . Since  $\alpha_0 \geq \max\{\alpha_1, \dots, \alpha_t\}$ , we have  $\sum_{i=1}^t \alpha_i^2 \leq \sum_{i=1}^t \alpha_0 \alpha_i \leq \alpha_0(1 - \alpha_0)$ .  $\square$

Recall from §2C that  $h'(X, w) = \min_{\emptyset \neq A \subsetneq X(0)} \frac{w(E(A, X(0) - A))}{2w(A)w(X(0) - A)}$ .

**Theorem 5.2.** *Let  $(X, w)$  be a weighted graph and let  $R$  be a nontrivial (additive) abelian group. Then:*

(i)  $(X, w, R_+)$  is an  $h'(X, w)$ -coboundary expander in dimension 0.

(ii) If  $(X, w, R_+)$  is an  $\varepsilon$ -coboundary expander in dimension 0, then  $h(X, w) \geq \varepsilon$  and  $h'(X, w) \geq \frac{\varepsilon}{2}$ .

<sup>10</sup>This definition slightly differs from the one in [8] if  $\mathcal{F}(x) = 0$  for some nonempty faces  $x \in X$ .

If  $X$  is connected, then Remark 4.3, and (i) imply that  $(X, w, R)$  is an  $(h'(X, w), 1)$ -cosystolic expander in dimension 0.

*Proof.* (i) Given  $a \in R$ , write  $f_a$  for element of  $C^0(X, R_+)$  defined by  $f_a(x) = a$  for all  $x \in X(0)$ . We abbreviate  $\|\cdot\|_w$  to  $\|\cdot\|$  and  $h'(X, w)$  to  $h'$ .

Let  $f \in C^0(X, R_+)$ . We need to show that  $\|d_0 f\| \geq h' \|f + B^k(X, G_+)\|$ , or equivalently, that  $\|d_0 f\| \geq h' \|f - f_a\|$  for some  $a \in R$ . Let  $a_0 = 0, a_1, \dots, a_t \in R$  be the values attained by  $f$  together with 0, and write  $A_i = \{x \in X(0) : f(x) = a_i\}$  and  $\alpha_i = w(A_i)$ . Then  $\sum_{i=0}^t \alpha_i = 1$  and  $\sum_{i=1}^t \alpha_i = \|f\|$ . Choose  $j \in \{0, \dots, t\}$  such that  $\alpha_j = \max\{\alpha_0, \dots, \alpha_t\}$ . By replacing  $f$  with  $f - f_{a_j}$ , we may assume that  $j = 0$ .

Let  $e \in X(1)$  and let  $u, v$  be the 0-faces of  $e$ . If  $u \in A_i$  and  $v \in A_j$  for distinct  $i$  and  $j$ , then  $f(x) \in \{a_i - a_j, a_j - a_i\}$ , and otherwise  $f(x) = 0$ . This means that  $\|d_0 f\| = \sum_{0 \leq i < j \leq t} w(E(A_i, A_j))$ .

Writing  $A_i^c := X(0) - A_i$ , we have  $\sum_{0 \leq i < j \leq t} w(E(A_i, A_j)) = \sum_{i=0}^t \frac{1}{2} w(E(A_i, A_i^c))$ . By the definition of  $h' = h'(X, w)$ ,

$$w(E(A_i, A_i^c)) \geq h' \cdot 2\alpha_i(1 - \alpha_i)$$

Thus,  $\|d_0 f\| \geq h' \sum_{i=0}^t \alpha_i(1 - \alpha_i) \geq h' \sum_{i=1}^t \alpha_i = h' \|f\|$ , where the second inequality is Lemma 5.1.

(ii) Fix a nontrivial element  $a \in R$ . For every  $A \subseteq X(0)$ , let  $f_A \in C^0(X, R_+)$  denote the function from  $X$  to  $R$  taking the value  $a$  on  $A$  and  $0_R$  elsewhere. Writing  $A^c := X(0) - A$ , we have  $\text{supp } d_0 f_A = E(A, A^c)$  and  $\|f_A + B^0(X, R_+)\| = \min\{w(A), w(A^c)\}$ . Since  $(X, w, R_+)$  is an  $\varepsilon$ -coboundary expander in dimension 0, we have  $w(E(A, A^c)) \geq \varepsilon \min\{w(A), w(A^c)\}$ . As  $A \subseteq X(0)$  was arbitrary, this means that  $h(X, w) \geq \varepsilon$ , and by (2.2),  $h'(X, w) \geq \frac{\varepsilon}{2}$ .  $\square$

**Corollary 5.3.** *Let  $(X, w)$  be a weighted graph which is a  $[-1, \lambda]$ -expander, and let  $R$  be a nontrivial abelian group. Then  $(X, w, R_+)$  is a  $(1 - \lambda)$ -coboundary expander in dimension 0.*

*Proof.* This follows from Theorems 2.6 and 5.2(i).  $\square$

**Corollary 5.4.** *Let  $(X, w)$  be a weighted graph, let  $R, S$  be nontrivial abelian groups and let  $\varepsilon \in [0, 2]$ . If  $(X, w, R_+)$  is an  $\varepsilon$ -coboundary expander in dimension 0, then  $(X, w, S_+)$  is an  $\frac{\varepsilon}{2}$ -expander in dimension 0.*

*Proof.* Combine parts (i) and (ii) of Theorem 5.2.  $\square$

**Example 5.5.** Let  $X$  be a complete graph on 3 vertices and let  $w$  be its canonical weight function. It is routine to check that the coboundary expansion of  $(X, w, (\mathbb{F}_2)_+)$  in dimension 0 is 2 while the coboundary expansion of  $(X, w, R_+)$  is  $\frac{3}{2}$  for every abelian group  $R$  admitting at least 3 elements. (The latter can be checked either directly, or using Corollary 5.3 and the fact that  $(X, w)$  is a  $[-1, -\frac{1}{2}]$ -expander.)

We now demonstrate that locally constant sheaves on good expander graphs may have poor coboundary expansion.

**Example 5.6.** Let  $(X, w)$  be a connected weighted graph such that every vertex in  $X$  is contained in at least 2 edges. Let  $e_0$  be an edge of minimal weight in  $X(1)$ , and let  $v_0$  be one of the 0-faces of  $e_0$ . Let  $\mathbb{F}$  be a field more than 2 elements, and let  $\alpha \in \mathbb{F} - \{0, 1\}$ . We define a locally constant sheaf  $\mathcal{F}$  on  $X$  as follows: put  $\mathcal{F}(x) = \mathbb{F}$  for all  $x \in X - \{v_0\}$ , and for all  $\emptyset \neq v \subsetneq e \in X(1)$ , let

$$\text{res}_{e \leftarrow v}^{\mathcal{F}} = \begin{cases} \text{id}_{\mathbb{F}} & (e, v) \neq (e_0, v_0) \\ \alpha \text{id}_{\mathbb{F}} & (e, v) = (e_0, v_0). \end{cases}$$

We claim that the coboundary expansion of  $(X, w, \mathcal{F})$  in dimension 0 is at most  $w(e_0) \leq \frac{1}{|X(1)|}$ . In particular, Theorem 5.2(i) and Corollary 5.3 fail for locally constant sheaves  $\mathcal{F}$ . To see this, note first that  $Z^0(X, \mathcal{F}) = 0$ ; this can be seen by inspecting the values of  $f \in Z^0(X, \mathcal{F})$  on a cycle subgraph containing  $e_0$  (such a cycle exists by our assumption on  $X$ ) and using the connectivity of  $X$ . Thus,  $\|f + B^0(X, \mathcal{F})\|_w = \|f\|_w$  for all  $f \in C^0(X, \mathcal{F})$ . Taking  $f = (1_{\mathbb{F}})_{x \in X(0)}$ , we get  $\text{supp}(d_0 f) = \{e_0\}$ , so  $\|d_0 f\|_w = w(e_0)$  while  $\|f + B^0(X, \mathcal{F})\|_w = 1$ .



## 6 Coboundary Expansion of Quotients of Constant Sheaves on Graphs

In this section, we show that taking the quotient of a constant augmented sheaf on a weighted graph by a “small” subsheaf still results in a good coboundary expander, provided that the weighted graph is a good enough spectral expander.

Given an abelian group  $R$  and a collection of subgroups  $\{R_i\}_{i \in I}$ , we say that  $\{R_i\}_{i \in I}$  are *linearly disjoint* (in  $R$ ) if the summation map  $(r_i)_{i \in I} \mapsto \sum_{i \in I} r_i : \bigoplus_{i \in I} R_i \rightarrow R$  is injective. For example, if  $R$  is a vector space over a field  $\mathbb{F}$  and  $R_i = \mathbb{F}v_i$  for some  $v_i \in R$ , then  $\{R_i\}_{i \in I}$  are linearly disjoint if and only if the vectors  $\{v_i\}_{i \in I}$  (including multiplicities) are linearly independent in  $R$ .

**Theorem 6.1.** *Let  $(X, w)$  be a weighted graph with  $n$  vertices and let  $R$  be an (additive) abelian group. Suppose that we are given subgroups  $R_x$  of  $R$  for every nonempty  $x \in X$ , and set  $R_\emptyset = \{0_R\}$ . Define a subsheaf  $\mathcal{C}$  of  $R_+$  by setting  $\mathcal{C}(x) = \sum_{y \subseteq x} R_y$  ( $x \in X$ ), and put*

$$t = \max\left\{\frac{w(e)}{w(x)} \mid x \in X(0), e \in X(1)_{\supseteq x}\right\} \quad \text{and} \quad s = \max\{w(e) \mid e \in X(1)\}.$$

Suppose that

- (1) for every subgraph  $Y$  of  $X$  which is either a cycle of length  $\leq \lceil \frac{2}{3}|X(0)| \rceil$  or a path of a length  $\leq 2$  (in the sense of §2A), the subgroups  $\{R_y\}_{y \in Y}$  are linearly disjoint in  $R$ , and
- (2) for every distinct  $u, v \in X(0)$ , the subgroups  $R_u$  and  $R_v$  are linearly disjoint.

If  $X$  is a  $[\mu, \lambda]$ -spectral expander ( $\mu, \lambda \in \mathbb{R}$ ), then  $(X, w, R_+/\mathcal{C})$  is an  $\varepsilon$ -coboundary expander in dimension 0 for

$$\varepsilon = \frac{2 - 4\lambda - 4 \max\{|\lambda|, |\mu|\} - 5t - 2s}{5 - 2\lambda}.$$

**Theorem 6.2.** *Let  $(X, w)$  be an  $(r+1)$ -partite weighted simplicial complex. Let  $R, \{R_x\}_{x \in X}, \mathcal{C}, s, t$  be as in Theorem 6.1 and assume that conditions (1) and (2) of that theorem are fulfilled. If  $(X, w)$  is an  $(r+1)$ -partite  $[\mu, \lambda]$ -expander (see §2D) with  $\lambda \geq -\frac{1}{r}$ , then  $(X, w, R_+/\mathcal{C})$  is an  $\varepsilon$ -coboundary expander in dimension 0 for*

$$\varepsilon = \frac{2r - 4r\lambda - 4r^2 \max\{|\lambda|, |\mu|\} - (5r+2)t - 2rs}{5r+2 - 2r\lambda}.$$

In both theorems, if all edges in  $X$  have the same weight, or if all the subgroups  $\{R_v\}_{v \in V(x)}$  are linearly disjoint, then we can eliminate the term  $-2s$ , resp.  $-2rs$ , in the numerator of  $\varepsilon$ ; see Remark 6.11 below. We also note that elementary analysis shows that artificially increasing  $\lambda$  will only decrease the resulting coboundary expansion  $\varepsilon$ . For a statement regarding the cosystolic expansion of the sheaf  $R/\mathcal{C}$ , see Corollary 6.12 below.

**Example 6.3.** Suppose that  $X$  is a connected  $k$ -regular graph on  $n$  vertices and  $w$  is its canonical weight function (Example 2.2). Let  $R, \{R_x\}_{x \in X}, \mathcal{C}, t$  and  $s$  be as in Theorem 6.1. Then  $t = \frac{2/kn}{1/n} = \frac{2}{k}$  and  $s = \frac{2}{kn}$ ; in fact, we can ignore  $s$  because all edges have the same weight. Let  $\rho \in [0, 1]$  be a number with  $|\lambda| \leq \rho$  for any eigenvalue  $\lambda \neq \pm 1$  of  $\mathcal{A}_{X,w}$ . Recall that  $X$  is called a *Ramanujan graph* if we can take  $\rho = \frac{2\sqrt{k-1}}{k}$ ; see [2] (for instance) for details and motivation for this definition.

If  $X$  is not bipartite, then  $(X, w)$  is a  $[-\rho, \rho]$ -expander (see §2C), and Theorem 6.1 implies that  $(X, w, R_+/\mathcal{C})$  is an  $\varepsilon$ -coboundary expander in dimension 0 for  $\varepsilon = \frac{2-8\rho-10/k}{5} = \frac{2}{5} - \frac{8}{5}\rho - \frac{2}{k}$ . If  $X$  is a Ramanujan graph, for instance, then  $\varepsilon \geq \frac{2}{5} - \frac{16\sqrt{k-1}}{5k} - \frac{2}{k} = \frac{2}{5} - O\left(\frac{1}{\sqrt{k}}\right)$ .

If  $X$  is bipartite, then  $(X, w)$  is a 2-partite  $[-\rho, \rho]$ -expander and Theorem 6.2 implies that  $(X, w, R_+/\mathcal{C})$  is an  $\varepsilon$ -coboundary expander in dimension 0 for  $\varepsilon = \frac{2-8\rho-14/k}{7} = \frac{2}{7} - \frac{8}{7}\rho - \frac{2}{k}$ . Again, taking  $X$  to be a bipartite Ramanujan graph gives  $\varepsilon \geq \frac{2}{7} - \frac{16\sqrt{k-1}}{7k} - \frac{2}{k} = \frac{2}{7} - O\left(\frac{1}{\sqrt{k}}\right)$ .

The rest of this section is dedicated to proving Theorems 6.1 and 6.2. We prove both theorems together in a series of lemmas.

**Lemma 6.4.** *Let  $(X, w)$  be a weighted graph and let  $\mathcal{F}$  be an augmented sheaf on  $X$ . Let  $\varepsilon \in \mathbb{R}_+$ ,  $\lambda \in \mathbb{R}$  and suppose that  $(X, w)$  is a  $[-1, \lambda]$ -expander, and for every  $v \in X(0)$  and  $h \in \mathcal{F}(v) - \{0\}$ , we have*

$$w(\{e \in X(1)_{\supseteq v} : \text{res}_{e \leftarrow v} h(v) \neq 0\}) \geq \varepsilon w(v).$$

Let  $f \in C^0(X, \mathcal{F})$  and  $\alpha \in [0, \infty)$ . If  $\|f\|_w \leq \alpha$ , then

$$(\varepsilon - 2\lambda - (2 - 2\lambda)\alpha)\|f\|_w \leq \|d_0 f\|_w.$$

*Proof.* Write  $A = \text{supp } f$  and  $A^c = X(0) - A$ . Since decreasing  $\alpha$  increases the left hand side of the desired inequality, it is enough to prove the lemma for  $\alpha = \|f\|_w$ .

Fix some  $v \in X(0)$  with  $f(v) \neq 0$ . We claim that for every  $e \in X(1)_{\supseteq v}$  with  $\text{res}_{e \leftarrow v} f(v) \neq 0$ , at least one of the following hold:

- (i)  $e \in E(A)$ ,
- (ii)  $e \in \text{supp}(d_0 f) \cap E(A, A^c)$ .

Indeed, we have  $(d_0 f)(e) = \pm \text{res}_{e \leftarrow v} f(v) \mp \text{res}_{e \leftarrow e-v} f(e-v)$ . If  $f(e-v) \neq 0$ , then  $v$  and  $e-v$  are in  $\text{supp } f$ , and  $e \in E(A)$ . Otherwise  $f(e-v) = 0$ , so  $(d_0 f)(e) = \pm \text{res}_{e \leftarrow v} f(v) \neq 0$  and  $e \in \text{supp}(d_0 f) \cap E(A, A^c)$ . This means that

$$w(\{e \in X(1)_{\supseteq v} : \text{res}_{e \leftarrow v} f(v) \neq 0\}) \leq w(E(A)_{\supseteq v}) + w([\text{supp}(d_0 f) \cap E(A, A^c)]_{\supseteq v}).$$

By assumption, the left hand side of the last inequality is at least  $\varepsilon w(v)$ . Summing over all  $v \in \text{supp } f$ , we get

$$\begin{aligned} \varepsilon \|f\|_w &\leq \sum_{v \in A} w(E(A)_{\supseteq v}) + \sum_{v \in A} w([\text{supp}(d_0 f) \cap E(A, A^c)]_{\supseteq v}) \\ &\leq 2w(E(A)) + w(\text{supp } d_0 f) \leq 2(\alpha^2 + \lambda\alpha(1 - \alpha)) + \|d_0 f\|_w. \end{aligned}$$

Here, the second inequality holds because every edge in  $E(A)$  is counted exactly twice and every edge in  $\text{supp}(d_0 f)$  is counted at most once, whereas the third inequality follows from Theorem 3.2(ii). By rearranging, we find that

$$\|d_0 f\|_w \geq \varepsilon \|f\|_w - 2\alpha^2 - 2\lambda\alpha(1 - \alpha) = (\varepsilon - 2\lambda - 2\alpha + 2\lambda\alpha)\|f\|_w. \quad \square$$

**Lemma 6.5.** *Let  $(X, w), R, \{R_x\}_{x \in X}, \mathcal{C}, t$  be as in Theorem 6.1 or Theorem 6.2, and let  $\mathcal{F} = R_+/\mathcal{C}$ . If  $\{R_y\}_{y \in Y}$  are linearly disjoint in  $R$  for every path  $Y \subseteq X$  of length 2, then for every  $x \in X(0)$  and  $h \in \mathcal{F}(x) - \{0\}$ , we have*

$$w(\{e \in X(1)_{\supseteq x} : \text{res}_{e \leftarrow x} h \neq 0\}) \geq (2 - t)w(x).$$

*Proof.* Let  $x \in X(0)$  and  $h \in \mathcal{F}(x) - \{0\}$ . If  $\text{res}_{e \leftarrow x} h \neq 0$  for all  $e \in X(1)_{\supseteq x}$ , then  $w(\{e \in X(1)_{\supseteq x} : \text{res}_{e \leftarrow x} h \neq 0\}) = w(X(1)_{\supseteq x}) = 2w(x)$  because  $w$  is a weight function (see §2B), and the lemma holds.

Suppose now that there exists  $y \in X(1)_{\supseteq x}$  such that  $\text{res}_{y \leftarrow x} h = 0$ . We claim that  $\text{res}_{z \leftarrow x} h \neq 0$  for all  $z \in X(1)_{\supseteq x}$  different from  $y$ . Fix such  $z$  and let  $x' = y - x$  and  $x'' = z - x$ . Then  $Y = \{\emptyset, x, x', x'', y, z\}$  is a path of length 2 in  $X$ , meaning that  $R_\emptyset = 0, R_x, R_{x'}, R_{x''}, R_y, R_z$  are linearly disjoint in  $R$ . Choose  $g \in R$  with  $h(x) = g + R_x$  (note that  $\mathcal{C}(x) = R_x + R_\emptyset = R_x$ ). Since  $\text{res}_{y \leftarrow x} h = 0$ , we have  $g \in R_y + R_x + R_{x'}$ . Likewise, if  $\text{res}_{z \leftarrow x} h = 0$ , then  $g \in R_z + R_x + R_{x''}$ . Consequently,  $g \in (R_y + R_x + R_{x'}) \cap (R_z + R_x + R_{x'') = R_x = \mathcal{C}(x)$ , but this contradicts our assumption that  $h = g + \mathcal{C}(x) \neq 0$  in  $\mathcal{F}(x)$ . We conclude that  $\{e \in X(1)_{\supseteq x} : \text{res}_{e \leftarrow x} h \neq 0\} = X(1)_{\supseteq x} - \{y\}$ . Since  $w(X(1)_{\supseteq x}) - w(y) = 2w(x) - w(y) \geq 2w(x) - tw(x) = (2 - t)w(x)$ , we are done.  $\square$

**Lemma 6.6.** *Let  $(X, w)$  be a weighted graph with  $n$  vertices. Define  $t$  and  $s$  as in Theorem 6.1. and let  $T$  be a subgraph of  $X$ .*

- (i) *If  $w(T(1)) \geq t$ , then  $T$  contains a cycle.*
- (ii) *If  $w(T(1)) \geq t + s$ , then  $T$  contains a cycle of length  $\leq \lceil \frac{2}{3}n \rceil$ .*

*Proof.* (i) It is enough to show that if  $T$  contains no cycles, then  $w(T(1)) < t$ . In this case,  $T$  is a forest, i.e., every connected component of  $T$  is a tree. Choose roots for the trees in  $T$  and denote the set of roots by  $R \subseteq V(X)$ . For  $v \in V(X) - R$ , we let  $p(v) \in V(X)$  denote the parent of  $v$  in  $T$ . Then  $w(T(1)) = \sum_{v \in V(X) - R} w(\{v, p(v)\}) \leq \sum_{v \in V(X) - R} tw(\{v\}) < \sum_{x \in X(0)} tw(x) = t$ .

(ii) By (i),  $T$  contains a cycle  $C_1$ . Choose an edge  $e_1$  in  $C_1$ . Then  $w(e_1) \leq s$ . Let  $T' = T - \{e_1\}$ , i.e., the graph obtained from  $T$  by removing the edge  $e_1$ . Then  $w(T'(1)) = w(T(1)) - w(e_1) \geq t + s - s = t$ , so by (i),  $T'$  contains another cycle,  $C_2$ , and  $e_1 \notin C_2$ . If  $C_1(0) \cap C_2(0) = \emptyset$ , then one of  $C_1, C_2$  has less than  $\frac{1}{2}n$  vertices, and is therefore the required cycle.

Suppose now that  $C_1(0) \cap C_2(0) \neq \emptyset$ . By Lemma 2.1,  $C_1 - C_2$  is a nonempty disjoint union of open paths. Let  $Z$  be one these paths, and let  $x, y$  denote its end points. If  $x = y$ , then  $C_1$  and  $C_2$  share exactly one vertex. Thus,  $|C_1(0)| + |C_2(0)| \leq n + 1$ , and again, one of  $C_1, C_2$  is the required cycle. Assume  $x \neq y$ . Then  $x, y \in C_2(0)$ . Let  $P$  and  $Q$  denote the two paths from  $x$  to  $y$  contained in  $C_2$  and put  $p = |P(0)|$ ,  $q = |Q(0)|$ ,  $z = |Z(0)|$ . Since  $P(0) - \{x, y\}$ ,  $Q(0)$  and  $Z(0)$  are pairwise disjoint, we have  $p + q + z \leq n + 2$ . Let  $R_1 := P \cup Q$ ,  $R_2 := P \cup Z$  and  $R_3 := Q \cup Z$ . Then  $R_1, R_2, R_3$  are cycles, and  $|R_1(0)| + |R_2(0)| + |R_3(0)| = (p + q - 2) + (p + z) + (q + z) = 2(p + q + z) - 2 \leq 2(n + 1)$ . This means that there is  $i \in \{1, 2, 3\}$  such that  $|R_i(0)| \leq \lfloor \frac{2}{3}(n + 1) \rfloor = \lceil \frac{2}{3}n \rceil$ , so we are done.  $\square$

**Remark 6.7.** The proof of Lemma 6.6(ii) also shows that the girth of a graph with  $n$  vertices and  $n + 1$  edges is at most  $\lceil \frac{2}{3}n \rceil$ . This bound is tight, e.g., consider a graph  $X$  obtained by gluing 3 closed paths of length  $k$  or  $k - 1$  at their endpoints.

**Lemma 6.8.** *Let  $X$  be a connected graph which is a union of its cycle subgraphs. Then every two vertices in  $X$  can be connected by a path of length  $\leq \frac{2}{3}(|X(0)| - 1)$ .*

*Proof.* Write  $n = |X(0)|$ , and let  $x, y \in X(0)$  be distinct 0-faces. Since  $X$  is connected, there is path from  $x$  to  $y$ , and our assumption on  $X$  implies that this path is contained in a union of cycles. This means that there are cycles  $R_1, \dots, R_t$  such that  $x \in R_1(0)$ ,  $y \in R_t(0)$  and there exists  $x_i \in R_i(0) \cap R_{i+1}(0)$  for all  $i \in \{1, \dots, t - 1\}$ . Set  $x_0 = x$  and  $x_t = y$ . We prove the lemma by induction on  $t$ .

If  $t = 1$ , then  $x, y \in R_1$ , so there is a path from  $x$  to  $y$  of length at most  $\lfloor \frac{1}{2}|R_1(0)| \rfloor \leq \lfloor \frac{1}{2}n \rfloor \leq \frac{2}{3}(n - 1)$  (because  $R_1$ , and hence  $X$ , has at least 3 vertices). Suppose henceforth that  $t > 1$ .

If  $R_i \cap R_j \neq \emptyset$  for some  $i, j \in \{1, \dots, t\}$  with  $i + 2 \leq j$ , then we can remove  $R_{i+1}, \dots, R_{j-1}$  from  $R_1, \dots, R_t$  and finish by the induction hypothesis. We may therefore assume that  $R_i \cap R_j = \emptyset$  whenever  $i + 2 \leq j$ .

Next, if  $|R_i(0) \cap R_{i+1}(0)| = 1$  for all  $i \in \{1, \dots, t - 1\}$ , Then  $R_i(0) \cap R_{i+1}(0) = \{x_i\}$  for all  $i \in \{1, \dots, t - 1\}$ . Since  $R_i \cap R_j = \emptyset$  when  $i + 2 \leq j$ , this means that the sets  $R_1(0) - \{x_1\}, \dots, R_t(0) - \{x_t\}$  are pairwise disjoint. Thus, writing  $r_i = |R_i(0)| - 1$ , we have  $\sum_{i=1}^t r_i \leq |X(0) - \{x_t\}| = n - 1$ . Let  $i \in \{1, \dots, t\}$ . Since  $R_i$  is a cycle with  $r_i + 1$  vertices, there is a closed path  $P_i$  from  $x_{i-1}$  to  $x_i$  of length at most  $\lfloor \frac{1}{2}(r_i + 1) \rfloor \leq \frac{2}{3}r_i$  (because  $r_i \geq 2$ ). The union  $P_1 \cup \dots \cup P_t$  is a path from  $x_0 = x$  to  $x_t = y$  of length at most  $\sum_{i=1}^t \frac{2}{3}r_i \leq \frac{2}{3}(n - 1)$ .

Finally, if there is  $i \in \{1, \dots, t - 1\}$  such that  $R_i$  and  $R_{i+1}$  share at least 2 vertices, then  $R_i - R_{i+1}$  is a disjoint union of open paths and each of these paths has distinct end points (cf. Lemma 2.1). Of these open paths, let  $P$  be denote the one containing  $x_{i-1}$  and let  $z, w$  be its endpoints. Then  $z, w \in R_{i+1}(0)$ . Let  $Q$  denote a closed path from  $z$  to  $w$  in  $R_{i+1}$  which also includes  $x_{i+1}$ . Then  $R' := P \cup Q$  is a cycle containing both  $x_{i-1}$  and  $x_{i+1}$ . We replace  $R_i, R_{i+1}$  with  $R'$  and proceed by induction on  $t$ .  $\square$

**Lemma 6.9.** *Let  $X$  be a cycle graph, let  $R, \{R_x\}_{x \in X}$  and  $\mathcal{C}$  be as in Theorem 6.1 and suppose that all the  $\{R_x\}_{x \in X}$  are linearly disjoint in  $R$ . Then for every  $f \in Z^0(X, R_+/\mathcal{C})$ , there exists  $h \in R$  such that  $f(v) = h + R_v$  for all  $v \in X(0)$ .*

*Proof.* Let  $v_0, \dots, v_{\ell-1}$  be the 0-faces of  $X$  and let  $e_0, \dots, e_{\ell-1}$  be the edges of  $X$ . We may choose the numbering so that  $v_i, v_{(i+1) \bmod \ell} \subseteq e_i$  for all  $i$ . We write  $v_\ell = v_0$  for convenience.

Let  $f \in Z^0(X, R_+/\mathcal{C})$ , and choose  $g \in C^0(X, R_+)$  projecting onto  $f$ , i.e.,  $f(v) = g(v) + R_v$  for all  $v \in X(0)$ . For every  $i \in \{0, \dots, \ell - 1\}$ , write  $g_i = g(v_i)$ , and let  $g'_i = g_i - g_0$ . Unfolding the definitions, the assumption  $d_0 f = 0$  is equivalent to having  $g'_{i+1} - g'_i = g_{i+1} - g_i \in R_{v_i} + R_{v_{i+1}} + R_{e_i}$  for all  $i \in \{0, \dots, \ell - 1\}$ .

We claim that there exist  $c_0 \in R_{v_0}$ ,  $c_1, \tilde{c}_1 \in R_{v_1}$ ,  $\dots$ ,  $c_{\ell-2}, \tilde{c}_{\ell-2} \in R_{v_{\ell-2}}$ ,  $\tilde{c}_{\ell-1} \in R_{v_{\ell-1}}$  and  $u_i \in R_{e_i}$  ( $i \in \{0, \dots, \ell - 2\}$ ) such that  $g'_i = c_0 + u_0 + c_1 + \dots + c_{i-1} + u_{i-1} + \tilde{c}_i$  for all  $i \in \{1, \dots, \ell - 1\}$ . The proof is by induction on  $i$ . For the case  $i = 1$ , note that  $g'_1 = g_1 - g_0 \in R_{v_0} + R_{v_1} + R_{e_0}$ , so we can choose

$c_0 \in R_{v_0}$ ,  $\tilde{c}_1 \in R_{v_1}$  and  $u_0 \in R_{e_0}$  such that  $g'_1 = c_0 + u_0 + \tilde{c}_1$ . Suppose now that  $i \in \{1, \dots, \ell - 2\}$  and  $c_0, \tilde{c}_1, c_1, \dots, \tilde{c}_{i-1}, c_{i-1}, \tilde{c}_i$  and  $u_0, \dots, u_{i-1}$  were chosen so that  $g'_j = c_0 + u_0 + \dots + c_{j-1} + u_{j-1} + \tilde{c}_j$  for all  $j \in \{1, \dots, i\}$ . Then

$$g'_{i+1} - c_0 - u_0 - \dots - c_{i-1} - u_{i-1} - \tilde{c}_i = g'_{i+1} - g'_i \in R_{v_i} + R_{v_{i+1}} + R_{e_i}.$$

Since  $R_{v_0}, \dots, R_{v_{i+1}}, R_{e_0}, \dots, R_{e_i}$  are linearly disjoint in  $R$ , we can choose  $c_i \in R_{v_i}$ ,  $u_i \in R_{e_i}$  and  $\tilde{c}_{i+1} \in R_{v_{i+1}}$  such that  $g'_{i+1} = c_0 + u_0 + \dots + c_{i-1} + u_{i-1} + c_i + u_i + \tilde{c}_{i+1}$ , hence our claim.

Now, we have  $c_0 + u_0 + \dots + c_{\ell-2} + u_{\ell-2} + \tilde{c}_{\ell-1} = g'_{\ell-1} = g_{\ell-1} - g_0 \in R_{v_{\ell-1}} + R_{v_0} + R_{e_{\ell-1}}$ , so again, since  $\{R_x\}_{x \in X(0) \cup X(1)}$  are linearly disjoint, we must have  $c_1 = \dots = c_{\ell-2} = 0$  and  $u_0 = \dots = u_{\ell-2} = 0$ . This means that  $g'_i = c_0 + \tilde{c}_i$  for all  $i \in \{1, \dots, \ell - 1\}$ . Define  $h = g_0 + c_0$ . Then for all  $i \in \{1, \dots, \ell - 1\}$ , we have  $g_i = g'_i + g_0 = h + \tilde{c}_i \in h + R_{v_i}$ , and  $g_0 = h - c_0 \in h + R_{v_0}$ . This means that  $f(v_i) = h + R_{v_i}$  for all  $i$ , so we proved the existence of  $h$ .  $\square$

We are now ready to prove the following key lemma.

**Lemma 6.10.** *Under the assumptions of Theorem 6.1, put  $\mathcal{F} = R_+/\mathcal{C}$  and let  $f \in C^0(X, \mathcal{F})$  and  $\beta \in [0, 1]$ . If  $\|d_0 f\|_w \leq \beta$ , then*

$$\|f + B^0(X, \mathcal{F})\|_w < \frac{2}{3} + \frac{1}{3}[\beta + t + s + \lambda + 2 \max\{|\lambda|, |\mu|\}].$$

If, instead, we use the assumptions of Theorem 6.2, then

$$\|f + B^0(X, \mathcal{F})\|_w < \frac{2r+2}{3r+2} + \frac{r}{3r+2}[\beta + t + s + \lambda + 2r \max\{|\lambda|, |\mu|\}].$$

*Proof. Step 1.* Let  $n = |X(0)|$ . We call a subgraph  $Y$  of  $X$  an  $f$ -blob, or just blob for short, if:

- (b1)  $Y$  is connected and equals to the union of its cycle subgraphs, and
- (b2) there exists  $g \in G$  such that  $f(v) = g + R_v$  for all  $v \in Y(0)$ .

Note that condition (b1) implies that  $|Y(0)| > 2$ , because a cycle has at least 3 vertices. We denote an element  $g \in G$  as in (b2) by  $g_Y$ ; we will see below that  $g_Y$  is uniquely determined by  $Y$ .

Denote the set of  $f$ -blobs by  $\mathcal{B}$ . By Lemma 6.9 and assumption (1) of Theorem 6.1, every cycle  $Y \subseteq X$  of length  $\lceil \frac{2}{3}n \rceil$  or less is a blob.

Note that if there exists a blob  $Y$  with  $w(Y(0)) \geq \alpha$ , then  $\|f + B^0(X, \mathcal{F})\| \leq 1 - \alpha$ . Indeed, writing  $g = g_Y$  and  $f' = f - d_{-1}g$ , we have  $\|f + B^0(X, \mathcal{F})\|_w \leq \|f'\|_w \leq 1 - \alpha$ , because  $f'$  vanishes on  $Y(0)$ . We will prove the lemma by showing that there exists a sufficiently large blob.

*Step 2.* Observe that if  $Y$  and  $Z$  are blobs such that  $|Y(0) \cap Z(0)| > 1$ , then  $Y \cup Z$  is also a blob. Indeed, (b1) holds for  $Y \cup Z$  because it holds for  $Y$  and  $Z$ , and  $Y \cap Z \neq \emptyset$ . To see that (b2) holds, fix a choice of  $g_Y$  and  $g_Z$ . Then for every  $v \in Y(0) \cap Z(0)$ , we have  $g_Y + R_v = f(v) = g_Z + R_v$ , so  $g_Y - g_Z \in R_v$ . Choosing distinct  $u, v \in Y(0) \cap Z(0)$ , we get  $g_Y - g_Z \in R_u \cap R_v = 0$ , because  $R_u$  and  $R_v$  are linearly disjoint (assumption (2) of Theorem 6.1). This means that  $g_Y = g_Z$  and (b2) holds for  $Y \cup Z$  by taking  $g := g_Y = g_Z$ .

Applying the previous paragraph with  $Y = Z$  shows that  $g_Y$  is uniquely determined by  $Y$ .

*Step 3.* Write  $M = \bigcup_{Y \in \mathcal{B}} Y$ . Then  $M$  is a subgraph of  $X$ .

We claim that  $M(1) \subseteq X(1) - \text{supp}(d_0 f)$ . To show this, it is enough to prove that  $Y(1) \subseteq X(1) - \text{supp}(d_0 f)$  for any blob  $Y$ . Let  $e$  be an edge in  $Y$  and let  $u$  and  $v$  be its vertices. Then  $f(u) = g_Y + R_u$  and  $f(v) = g_Y + R_v$ . As a result,  $(d_0 f)(e) = g_Y - g_Y + (R_u + R_v + R_e) = 0$  in  $\mathcal{F}(e)$ , meaning that  $e \notin \text{supp}(d_0 f)$ , hence our claim.

We observed in Step 1 that every cycle of length  $\leq \lceil \frac{2}{3}n \rceil$  in  $X$  is a blob, and thus contained in  $M$ . It follows that the graph underlying  $X(1) - \text{supp}(d_0 f) - M(1)$  contains no cycles of length  $\leq \lceil \frac{2}{3}n \rceil$ . Thus, by Lemma 6.6,  $w(X(1) - \text{supp}(d_0 f) - M(1)) < t + s$ . Since  $\|d_0 f\|_w \leq \beta$ , it follows that

$$w(M(1)) > 1 - \beta - t - s.$$

*Step 4.* A blob is called maximal if it is not properly contained in any other blob. Write  $\mathcal{M}$  for the set of maximal blobs. Since every blob is contained in a maximal blob,  $M = \bigcup_{Y \in \mathcal{M}} Y$ . Step 2 tells us that for every  $Y \in \mathcal{M}$  and  $Z \in \mathcal{B}$ , either  $Y = Z$ , or  $|Y(0) \cap Z(0)| \leq 1$ .

Let  $N$  denote the set of 0-faces of  $X$  belonging to more than one blob in  $\mathcal{M}$ . We define a graph  $\Gamma$  as follows: The vertices of  $\Gamma$  are  $\mathcal{M} \cup N$  and the edges of  $\Gamma$  are pairs  $\{x, Y\}$  such that  $x \in N$ ,  $Y \in \mathcal{M}$  and  $x \in Y(0)$ . See Figure 1 for an illustration.




Figure 1: An illustration of the collection of blobs  $\mathcal{M}$  (left), the associated graph  $\Gamma$  (middle), and the partition  $\{Y^* \mid Y \in \mathcal{M}\}$  (right). The blobs are labelled A–E. The black vertices are those in living in  $N$ . The set of roots taken on the right is  $\mathcal{R} = \{A\}$ .

We claim that the graph  $\Gamma$  has no cycles. For the sake of contradiction, suppose otherwise. Then there exist  $\ell \geq 2$  and distinct  $x_0, \dots, x_{\ell-1} \in N$ ,  $Y_0, \dots, Y_{\ell-1} \in \mathcal{M}$  such that  $\{x_i\} = Y_i \cap Y_{i+1}$  for all  $i \in \{0, \dots, \ell-1\}$  (with the convention that  $Y_\ell = Y_0$ ). By applying Lemma 6.8 to  $Y_i$ , we see that there is a closed path  $P_i \subseteq Y_i$  of length  $\leq \frac{2}{3}(|Y_i(0)| - 1)$  from  $x_{i-1}$  to  $x_i$ . Since the sets  $Y_0(0) - \{x_0\}, \dots, Y_{\ell-1}(0) - \{x_{\ell-1}\}$  are pairwise disjoint, the union  $P = \bigcup_{i=0}^{\ell-1} P_i$  is a cycle of length at most  $\frac{2}{3} \sum_{i=0}^{\ell-1} (|Y_i(0)| - 1) \leq \frac{2}{3}n$ . This means that  $P$  is a blob — see Step 1. By construction,  $P$  and each of the  $Y_i$  share at least two 0-faces, namely,  $x_i$  and  $x_{i+1}$ , so we must have  $Y_0 = \dots = Y_{\ell-1} = P$ . This contradicts our assumption that  $Y_0, \dots, Y_{\ell-1}$  are distinct.

*Step 5.* By the previous step,  $\Gamma$  is a forest. We choose a set of roots  $\mathcal{R}$  for the trees in  $\Gamma$ . By the definition of  $N$ , every connected component of  $\Gamma$  contains a vertex in  $\mathcal{M}$ , so we may choose  $\mathcal{R}$  to be contained in  $\mathcal{M}$ . We denote the parent of every every  $Y \in \mathcal{M} - \mathcal{R}$  by  $x(Y)$ . Since  $x(Y)$  is not a root, it has a parent, which we denote by  $g(Y)$  (the grandparent of  $Y$ ). For every blob  $Y \in \mathcal{M}$ , define:

$$Y^* = \begin{cases} Y(0) & Y \in \mathcal{R} \\ Y(0) - \{x(Y)\} & Y \in \mathcal{M} - \mathcal{R}. \end{cases}$$

We claim that  $\{Y^* \mid Y \in \mathcal{M}\}$  is a partition of  $M(0)$ . To see this, note first that  $\bigcup_{Y \in \mathcal{M}} Y^* \subseteq \bigcup_{Y \in \mathcal{M}} Y(0) = M(0)$ . To see that  $M(0) \subseteq \bigcup_{Y \in \mathcal{M}} Y^*$ , observe that every  $x \in M(0)$  belongs to  $Y(0)$  for some  $Y \in \mathcal{M}$ . If  $Y \in \mathcal{R}$ , then  $Y^* = Y(0)$  and  $x \in Y^*$ . Otherwise,  $Y^* = Y(0) - \{x(Y)\}$ , so  $x \in Y^*$  provided  $x \neq x(Y)$ . If  $x = x(Y)$ , then  $x \in g(Y)(0)$  by the definition of  $\Gamma$ . Since  $g(Y)$  is the parent of  $x$  in  $\Gamma$ , the 0-face  $x$  cannot be the parent of  $g(Y)$ , so  $x \neq x(g(Y))$ , which means that  $x \in g(Y)^*$ . This proves that  $x \in \bigcup_{Y \in \mathcal{M}} Y^*$ . To finish, we need to show that  $Y^* \cap Z^* = \emptyset$  for any distinct  $Y, Z \in \mathcal{M}$ . If there is  $x \in Y^* \cap Z^*$ , then  $x \in N$  and  $x$  is adjacent to  $Y$  and  $Z$  in  $\Gamma$ . This means that at least one of  $Y, Z$  is not a parent of  $x$ ; say it is  $Y$ . Then  $x$  must be the parent of  $Y$ , which means that  $x \notin Y - \{x(Y)\} = Y^*$ , a contradiction.

Next, let  $e \in M(1)$ . We claim that one of the following holds:

- (i)  $e \in E(Y^*)$  for some  $Y \in \mathcal{M}$ ,
- (ii)  $e \in E(Y^*, g(Y)^*)$  for some  $Y \in \mathcal{M} - \mathcal{R}$ .

Indeed, let  $y, z$  denote the 0-faces of  $e$ . Since  $M = \bigcup_{Y \in \mathcal{M}} Y$ , there is  $Y \in \mathcal{M}$  such that  $e \in Y(1)$ , and thus  $y, z \in Y(0)$ . If both  $y$  and  $z$  are different from  $x(Y)$ , then  $y, z \in Y^*$  and (i) holds. Otherwise, exactly one of  $y, z$  equals  $x(Y)$ , say  $y = x(Y)$  and  $z \in Y^*$ . Then  $y$  is the parent of  $Y$  in  $\Gamma$ . As explained in the previous paragraph,  $y \in g(Y)$  and  $y$  is not a parent of  $g(Y)$ , so  $y \in g(Y)^*$  and (ii) holds.

*Step 6.* At this point, we assume we are in the setting of Theorem 6.1. For every  $Y \in \mathcal{M}$ , put  $\alpha_Y = w(Y^*)$  and set

$$\alpha = \max\{\alpha_Y \mid Y \in \mathcal{M}\}.$$

Since  $\{Y^* \mid Y \in \mathcal{M}\}$  is a partition of  $M(0)$  (Step 5), we have  $\sum_{Y \in \mathcal{M}} \alpha_Y = w(M(0)) \leq 1$ .

Let  $g^{-1}(Y)$  denote the set of blobs  $Z \in \mathcal{M}$  with  $g(Z) = Y$ . By Step 3 and the last paragraph of Step 5, we have

$$1 - \beta - t - s < w(M(1)) \leq \sum_{Y \in \mathcal{M}} w(E(Y^*)) + \sum_{Y \in \mathcal{M}} w(E(Y^*, \bigcup_{Z \in g^{-1}(Y)} Z^*)).$$

Put  $\theta = \max\{|\mu|, |\lambda|\}$ . By Theorem 3.2, the right hand side is at most

$$\begin{aligned} & \sum_{Y \in \mathcal{M}} (\alpha_Y^2 + \lambda \alpha_Y) + \sum_{Y \in \mathcal{M}} (2\alpha_Y \sum_{Z \in g^{-1}(Y)} \alpha_Z + 2\theta \sqrt{\alpha_Y \sum_{Z \in g^{-1}(Y)} \alpha_Z}) \\ & \leq \sum_{Y \in \mathcal{M}} \alpha_Y (\alpha + \lambda) + \sum_{Y \in \mathcal{M}} (2\alpha_Y \sum_{Z \in g^{-1}(Y)} \alpha_Z + \theta (\alpha_Y + \sum_{Z \in g^{-1}(Y)} \alpha_Z)) \\ & = \sum_{Y \in \mathcal{M}} \alpha_Y (\alpha + \lambda) + \theta \sum_{Y \in \mathcal{M}} \alpha_Y + \sum_{Y \in \mathcal{M}} \sum_{Z \in g^{-1}(Y)} (2\alpha_Y + \theta) \alpha_Z \\ & \leq \alpha + \lambda + \theta + \sum_{Z \in \mathcal{M} - \mathcal{R}} \alpha_Z (2\alpha_{g(Z)} + \theta) \\ & \leq \alpha + \lambda + \theta + (2\alpha + \theta) = 3\alpha + 2\theta + \lambda. \end{aligned}$$

Thus,  $1 - \beta - t - s < 3\alpha + 2\theta + \lambda$ , and by rearranging, we get

$$\alpha > \frac{1}{3} - \frac{\beta}{3} - \frac{t}{3} - \frac{s}{3} - \frac{\lambda + 2 \max\{|\lambda|, |\mu|\}}{3}.$$

By the definition of  $\alpha$ , there exists a blob  $Y$  with  $w(Y(0)) \geq w(Y^*) = \alpha$ . As explained in Step 1, it follows that  $\|f + B^0(X, \mathcal{F})\|_w \leq 1 - \alpha$ , so this completes the proof of the lemma under the assumptions of Theorem 6.1.

*Step 7.* Finally, suppose that we are in the setting of Theorem 6.2. As in Step 6, we find that

$$1 - \beta - t - s < \sum_{Y \in \mathcal{M}} w(E(Y^*)) + \sum_{Y \in \mathcal{M}} w(E(Y^*, \bigcup_{Z \in g^{-1}(Y)} Z^*)).$$

Using Theorem 3.3(ii) and Theorem 3.2(ii) (the assumption  $\lambda \geq -\frac{1}{r}$  guarantees that  $(X, w)$  is a  $[-1, \lambda]$ -expander if we forget the  $(r+1)$ -partite structure of  $X$ ) we see that the right hand side is at most

$$\sum_{Y \in \mathcal{M}} (\alpha_Y^2 + \lambda \alpha_Y) + \sum_{Y \in \mathcal{M}} (\frac{2r+2}{r} \alpha_Y \sum_{Z \in g^{-1}(Y)} \alpha_Z + 2r\theta \sqrt{\alpha_Y \sum_{Z \in g^{-1}(Y)} \alpha_Z})$$

and a computation similar to the one in Step 6 shows that this expression is bounded by

$$\alpha + \lambda + r\theta + (\frac{2r+2}{r} \alpha + r\theta) = \frac{3r+2}{r} \alpha + 2r\theta + \lambda.$$

As a result,

$$\alpha > \frac{r}{3r+2} [1 - \beta - t - s - \lambda - 2r \max\{|\lambda|, |\mu|\}]$$

and we conclude the proof as in Step 6.  $\square$

We are now ready to prove Theorems 6.1 and 6.2.

*Proof of Theorem 6.1.* Fix some  $\beta \in [0, 1]$ , and let  $f \in C^0(X, R_+/\mathcal{C})$  be an 0-cochain such that  $\|f\|_w = \|f + B^0(X, R_+/\mathcal{C})\|_w$ . If  $\|d_0 f\|_w \geq \beta$ , then we have  $\|d_0 f\|_w \geq \beta \|f\|_w$ . On the other hand, if  $\|d_0 f\|_w < \beta$ , then by Lemma 6.10, we have  $\|f\|_w = \|f + B^0(X, G_+/\mathcal{C})\|_w \leq \frac{2}{3} + \frac{1}{3}[\beta + t + s + \lambda + 2 \max\{|\lambda|, |\mu|\}]$ . By Lemmas 6.4 and 6.5 this means that

$$\begin{aligned} \|d_0 f\|_w & \geq \left[ 2 - t - 2\lambda - (2 - 2\lambda) \left[ \frac{2}{3} + \frac{\beta}{3} + \frac{t}{3} + \frac{s}{3} + \frac{\lambda + 2 \max\{|\lambda|, |\mu|\}}{3} \right] \right] \|f\|_w \\ & \geq \left[ \left[ \frac{2}{3} - \frac{4}{3}\lambda - \frac{4}{3} \max\{|\lambda|, |\mu|\} - \frac{5}{3}t - \frac{2}{3}s \right] - \left( \frac{2-2\lambda}{3} \right) \beta \right] \|f\|_w \end{aligned}$$

As a result,  $(X, w, R_+/\mathcal{C})$  is an  $\varepsilon$ -coboundary expander in dimension 0 for

$$\varepsilon = \min\left\{ \beta, \left[ \frac{2}{3} - \frac{4}{3}\lambda - \frac{4}{3} \max\{|\lambda|, |\mu|\} - \frac{5}{3}t - \frac{2}{3}s \right] - \left( \frac{2-2\lambda}{3} \right) \beta \right\}$$

The right hand side is minimized when  $\varepsilon = \beta = \frac{2-4\lambda-4 \max\{|\lambda|, |\mu|\}-5t-2s}{5-2\lambda}$ , and theorem follows.  $\square$

*Proof of Theorem 6.2.* Similarly to the proof of Theorem 6.1, we find that for every  $\beta \in [0, 1]$ , the triple  $(X, w, R_+/C)$  is an  $\varepsilon$ -coboundary expander in dimension 0 for

$$\varepsilon = \min\left\{\beta, \left[\frac{2r}{3r+2} - \frac{4r}{3r+2}\lambda - \frac{4r^2}{3r+2}\theta - \frac{5r+2}{3r+2}t - \frac{2r}{3r+2}s\right] - \frac{(2-2\lambda)r}{3r+2}\beta\right\},$$

where  $\theta = \max\{|\lambda|, |\mu|\}$ . The minimum of this expression is attained for  $\varepsilon = \beta = \frac{2r-4r\lambda-4r^2\theta-(5r+2)t-2rs}{5r+2-2r\lambda}$ , hence the theorem.  $\square$

**Remark 6.11.** In Theorems 6.1 and 6.2, if all edges in  $X$  have the same weight, or if *all* the subgroups  $\{R_x\}_{x \in X}$  are linearly disjoint, then we can eliminate the terms  $-2s$ , resp.  $-2rs$ , in the expression for  $\varepsilon$ , or equivalently take  $s = 0$ .

Indeed, if all edges in  $X$  have the same weight, then the assertion  $w(M(1)) > 1 - \beta - t - s$  in Step 3 of the proof of Lemma 6.10 implies that  $w(M(1)) \geq 1 - \beta - t$ . Likewise, if all the subgroups  $\{R_x\}_{x \in X}$  of  $R$  are linearly disjoint, then in the same place in the proof, we can apply Lemma 6.6(i) instead of Lemma 6.6(ii) and get that  $w(M(1)) > 1 - \beta - t$ . Carrying the entire proof of the Lemma 6.10 using the inequality  $w(M(1)) \geq 1 - \beta - t$  in place of  $w(M(1)) > 1 - \beta - t - s$  allows us to eliminate  $s$  at the cost of replacing the strict inequalities in the proof with non-strict inequalities. This, in turn, eliminates  $s$  from Theorems 6.1 and 6.2.

**Corollary 6.12.** *Let  $X, w, R, \{R_x\}_{x \in X}, C, \varepsilon$  be as in Theorem 6.1 or Theorem 6.2. Put  $\eta = \max\{w(x) \mid x \in X(0)\}$  (we always have  $\eta \leq \frac{\varepsilon}{t}$ ). Then, regarding  $C$  as a subsheaf of the constant sheaf  $R$  on  $X$ , the triple and  $(X, w, R/C)$  is an  $(\varepsilon, 1 - \eta)$ -cosystolic expander in dimension 0.*

*Proof.* By Theorems 6.1 and 6.2 and Remark 4.3, it is enough to show that for all  $f \in R = C_{-1}(X, R_+/C)$ , we have  $\|d_{-1}f\| \geq 1 - \eta$ . Observe that  $(d_{-1}f)(v) = f + R_v \in R/R_v$  for all  $v \in X(0)$ . Thus,  $\|d_{-1}f\| = 1 - w(A)$ , where  $A$  is the set of 0-faces  $v$  such that  $f \in R_v$ . Since every two of the groups  $\{R_v\}_{v \in X(0)}$  are linearly disjoint,  $A$  is either empty or a singleton, so  $w(A) \leq \eta$  and the lemma follows. (It is worth noting that, if  $R_v \neq 0$  for all  $v$ , then by choosing  $f \in R_v$  with  $w(v)$  maximal, we get  $\|d_{-1}f\| = 1 - \eta$ , so the constant  $1 - \eta$  cannot be lowered in general.)  $\square$

## 7 The Case of Finite Buildings

We now apply Corollary 5.3, and Theorem 6.2 to the 0-skeleton of finite buildings admitting a strongly transitive group action, giving upper bounds on the coboundary expansion in terms of the *thickness* and the *type* of the building. This is the main reason why we have taken care to treat the case of  $(r + 1)$ -partite weighted simplicial complexes. Applications of the results of this section to locally testable codes appear in [8].

We refer the reader to [1] for an extensive discussion of buildings. Here we satisfy with saying that a building is a possibly-infinite simplicial complex  $X$  equipped with a collection  $\mathcal{E}$  of subcomplexes called *apartments* satisfying a list of axioms. All the apartments of  $X$  are isomorphic to each other and to a *Coxeter complex*  $\Sigma$ . The data of  $\Sigma$  (up to isomorphism) is encoded by a *Coxeter diagram*  $T$ , which is a finite graph with edges labelled by elements from the set  $\{3, 4, 5, \dots\} \cup \{\infty\}$  (unlabelled edges are given the label 3 by default). We write  $T = T(X)$  and say that  $T$  is the *type* of  $X$ . There is a labelling of the vertices of  $X$  by the vertices of  $T$  making  $X$  into a pure  $(r + 1)$ -partite simplicial complex, where  $r = \dim X = |V(T)| - 1$ . The type of a face  $z \in X$  is the set  $t(z) \subseteq V(T)$  consisting of the types of the vertices of  $z$ . If  $\dim z \leq \dim X - 1$ , then the link  $X_z$  is also a building and its type  $T(X_z)$  is the Coxeter diagram obtained from  $T = T(X)$  by removing the vertices in  $t(z)$ .

Let  $B$  denote the  $V(T) \times T(V)$  real matrix determined by

$$B_{u,v} = \begin{cases} 1 & u = v \\ -\cos \frac{\pi}{m} & \{u, v\} \in T(1) \text{ and } m \text{ is the label of } \{u, v\} \\ 0 & \{u, v\} \notin T(1). \end{cases}$$

The building  $X$  is called *spherical* if  $B$  is positive definite, or equivalently, if its apartments are finite. Following [1, Chapter 10], we call  $X$  *affine* if  $B$  is positive semidefinite and  $\text{rank } B = |V(T)| - 1$  (consult [1,

Proposition 10.44]). See [1, p. 50, Remark 10.33(b)] for a complete list of the possible spherical and affine Coxeter diagrams. If  $X$  is an affine building and  $z$  is a face with  $0 \leq \dim z < \dim X$ , then the link  $X_z$  is a spherical building.

A building  $X$  of dimension  $r$  is called  $q$ -thick if every  $(r-1)$ -dimensional face of  $X$  is contained in at least  $q$   $r$ -faces. We say that  $X$  is thick if it is 3-thick.

Recall from [1, §6.1.1] that an  $r$ -dimensional building  $X$  is said to possess a *strongly transitive action* if there is a group  $G$  acting on  $X$  via type-preserving simplicial automorphisms such that  $G$  takes apartments to apartments and acts transitively on the set of pairs  $(A, x)$  consisting of an apartment  $A \in \mathcal{E}$  and an  $r$ -face  $x$  in  $A$ . Since  $G$  is type preserving, this means that  $G$  acts transitively on the set of faces of a fixed type  $t \subseteq T(X)(0)$ . Moreover, for every  $z \in X$  of dimension  $< \dim X - 1$ , the building  $X_z$  also possesses a strongly transitive action. It follows from Tits' classification of thick spherical and affine buildings, see [1, Chapter 9, §11.9] for a survey, that all thick finite buildings of dimension  $\geq 2$  and all locally-finite thick affine buildings of dimension  $\geq 3$  admit a strongly transitive action.

**Example 7.1.** Let  $\mathbb{F}$  be a field and let  $n \in \mathbb{N}$ . The incidence complex of nontrivial subspaces of  $\mathbb{F}^{n+1}$ , denoted  $A_n(\mathbb{F})$ , is an  $(n-1)$ -dimensional building of type  $A_n$ , where  $A_n$  is the Coxeter diagram consisting of a single path of length  $n-1$  with all edges labeled 3. In more detail, the vertices of  $A_n(\mathbb{F})$  are the nontrivial subspaces of  $\mathbb{F}^{n+1}$  and its faces are the sets of vertices which are totally ordered by inclusion. The apartments of  $A_n(\mathbb{F})$  are induced from bases of  $\mathbb{F}^{n+1}$  as follows: If  $E$  is a basis of  $\mathbb{F}^{n+1}$ , then the collection of faces  $x = \{V_0, \dots, V_i\} \in A_n(\mathbb{F}^n)$  for which each  $V_j$  is spanned by a subset of  $E$  is an apartment.

The group  $\mathrm{GL}_{n+1}(\mathbb{F})$  acts on  $A_n(\mathbb{F})$  via its standard action on  $\mathbb{F}^{n+1}$ . This action is strongly transitive because  $\mathrm{GL}_{n+1}(\mathbb{F})$  acts transitively on the set of bases of  $\mathbb{F}^{n+1}$ .

When  $n = 2$ , the graph  $A_2(\mathbb{F})$  is nothing but the incidence graph of points and lines in the 2-dimensional projective space over  $\mathbb{F}$ .

Let  $X$  be a building of type  $T = T(X)$ . We define

$$m(X) = m(T) = \max \{2\} \cup \{n \mid T \text{ has an edge labelled } n\}$$

with the convention that  $m(X) = m(T) = \infty$  if  $T$  has an edge labelled  $\infty$ . Observe that  $m(X) \geq m(X_z)$  for every face  $z \in X$  of dimension  $< \dim X - 1$ . This definition is motivated by the following theorem, which we derive from results of Evra–Kaufman [7] (see also [6]) and Oppenheim [27].

**Theorem 7.2.** *Let  $X$  be a (finite)  $r$ -dimensional simplicial complex such that one of the following hold:*

- (1)  $X$  is a  $q$ -thick spherical building admitting a strongly transitive action;
- (2) the universal covering of  $X$  is a  $q$ -thick affine building of dimension  $\geq 2$  admitting a strongly transitive action.

*In both cases,  $X$  has the structure of a pure  $(r+1)$ -partite simplicial complex. Let  $w$  denote the canonical weight function of  $X$  (Example 2.2), let  $T$  denote the Coxeter diagram of the building mentioned in (1) or (2) and put  $m = m(T)$ . Then  $(X, w)$  is an  $(r+1)$ -partite  $[-r\lambda, \lambda]$ -expander for*

$$\lambda = \frac{\sqrt{m-2}}{\sqrt{q} - (r-1)\sqrt{m-2}},$$

*provided  $q \geq r^2(m-2)$ .*

*Proof.* Suppose first that  $r = 1$ . Then only case (1) is possible. Thus,  $X$  is a 1-dimensional building admitting a strongly transitive action by a group  $G$ . The Coxeter graph of  $X$  consists of two vertices and one edge labeled  $m$ , so each apartment in  $X$  is a cycle graph of length  $2m$ . We label the vertices of  $T = T(X)$  by 0 and 1 and write  $X_{\{i\}}$  ( $i \in \{0, 1\}$ ) for the 0-faces of type  $\{i\}$  in  $X$ . Since  $G$  acts transitively on  $X_{\{0\}}$  and  $X_{\{1\}}$ ,  $X$  is a biregular graph. Write  $n_i = |X_{\{i\}}|$  and let  $k_i$  denote the number of edges containing a 0-face in  $X_{\{i\}}$ . Then  $|X(1)| = n_0 k_0 = n_1 k_1$ , which means that  $w(e) = \frac{1}{n_0 k_0} = \frac{1}{n_1 k_1}$  for all  $e \in X(1)$  and  $w(x) = \frac{1}{2n_i}$  for all  $x \in X_{\{i\}}$ . We may assume without loss of generality that  $k_0 \leq k_1$ . Note also that  $q \leq k_0$ . We now adapt the proofs of [7, Propositions 5.21, 5.22] to our *weighted graph* situation, and also improve the expansion constants, in order to show that  $(X, w)$  is a bipartite  $[-\sqrt{\frac{m-2}{q}}, \sqrt{\frac{m-2}{q}}]$ -expander.



Let  $f' \in C^0_\diamond(X, \mathbb{R})$  (notation as in §2D) denote a nonzero eigenfunction of  $\mathcal{A} := \mathcal{A}_{X,w}$  with eigenvalue  $\lambda$ . We need to prove that  $\lambda^2 \leq \frac{m-2}{q}$ . It is enough to consider the case  $\lambda \neq 0$ . In this case,  $f'$  cannot vanish on  $X_{\{0\}}$ . Thus, we may choose  $s \in X_{\{0\}}$  with  $f'(s) \neq 0$  and put  $K = \{g \in G : gs = s\}$ . By applying Lemma 2.8 to  $f'$ , we see that there exists  $f'' \in C^0_\diamond(X, \mathbb{R})$  with  $\mathcal{A}f'' = -\lambda f''$  and  $f''(s) \neq 0$ .

Given 0-faces  $x, y \in X(0)$ , we denote the  $K$ -orbit of  $x \in X(0)$  by  $[x]$ , and write  $d(x, y)$  for the length of the shortest path from one  $x$  to  $y$  in  $X$ . We claim that following hold (cf. [7, Definition 5.20]):

- (i) The number of  $K$ -orbits in  $X(0)$  is at most  $m + 1$ .
- (ii) For all  $i \in \{0, 1\}$ , the 0-faces in  $X_{\{i\}}$  of maximal distance from  $s$  form a  $G$ -orbit. This maximal distance is  $m - ((m + i) \bmod 2)$ .
- (iii) If  $x \in X(0) - \{s\}$  is not of maximal distance from  $s$ , then there is exactly one 0-face  $y$  adjacent to  $x$  with  $d(s, y) < d(s, x)$ .

This is similar to the proof of [7, Propositions 5.22]: To see (i), fix an apartment  $E$  containing  $s$  and apply [7, Lemma 5.16] to conclude that every  $K$ -orbit in  $X(0)$  meets  $E$ . Let  $x, y \in E(0)$  be the 0-faces adjacent to  $s$ . By the strong transitivity of the  $G$ -action, there is  $g \in G$  such that  $g(E) = E$  and  $g\{s, x\} = g\{s, y\}$ . Since  $G$  preserves types,  $g(s) = s$ , so  $g$  is a reflection of the  $2m$ -cycle  $E$  fixing  $s$ . This means that all except possibly 2  $K$ -orbits meet  $E$  in at least two vertices, and (i) follows. The first assertion of (ii) is shown exactly as in *op. cit.*, and the second assertion follows from the fact that  $s \in X_{\{0\}}$  and each apartment is a cycle of length  $2m$ . As for (iii), let  $D$  be the set of 0-faces  $y \in X(0)$  adjacent to  $x$  with  $d(s, y) < d(s, x)$ . It is shown in the proof [7, Propositions 5.22] that there is an apartment  $E$  of  $X$  with  $s, x \in E$  and  $D \subseteq E$ . Since  $E$  is a cycle graph,  $D$  must be a singleton.

Next, let us regard  $\mathcal{A} = \mathcal{A}_{X,w}$  as an operator from the complex vector space  $C^0(X, \mathbb{C})$  to itself; this does not affect the spectrum of  $\mathcal{A}$ . Put  $Y = K \backslash X$  and let  $C^0(Y, \mathbb{C})$  denote the set of functions from  $Y$  to  $\mathbb{C}$  (note that  $Y$  is not a graph in general). We define a linear operator  $\mathcal{B} : C^0(Y, \mathbb{C}) \rightarrow C^0(Y, \mathbb{C})$  by

$$(\mathcal{B}g)[x] = \sum_{y \in X(1)_x} \frac{w(x \cup y)}{2w(x)} g[y].$$

Note that this does not depend on the representative  $x$  in the orbit  $[x]$ .

We claim that  $\text{Spec } \mathcal{B} \subseteq \text{Spec } \mathcal{A}$ ; in particular,  $\text{Spec } \mathcal{B} \subseteq \mathbb{R}$ . Indeed, given  $\mu \in \mathbb{C}$  and  $g \in C^0(Y, \mathbb{C})$  with  $\mathcal{B}g = \mu g$ , the function  $f \in C^0(X, \mathbb{C})$  defined by  $f(x) = g[x]$  satisfies  $\mathcal{A}f = \mu f$  because, for all  $x \in X(0)$ ,

$$(\mathcal{A}f)(x) = \sum_{y \in X(0)_x} \frac{w(x \cup y)}{w(x)} f(y) = \sum_{y \in X(0)_x} \frac{w(x \cup y)}{w(x)} g[y] = (\mathcal{B}g)[x] = \mu g[x] = \mu f(x).$$

We further claim that the  $\mathcal{A}$ -eigenvalue  $\lambda$  corresponding to  $f' \in C^0_\diamond(X, \mathbb{R})$  is in  $\text{Spec } \mathcal{B}$ . Indeed, define  $g' \in C^0(Y, \mathbb{C})$  by  $g'[y] = \sum_{k \in K} f'(ky)$  and note that  $g'[s] = |K|f'(s) \neq 0$ . Then, for every  $x \in X(0)$ ,

$$\begin{aligned} (\mathcal{B}g')[x] &= \sum_{y \in X(1)_x} \frac{w(x \cup y)}{2w(x)} g'[y] = \sum_{y \in X(1)_x} \sum_{k \in K} \frac{w(x \cup y)}{2w(x)} f'(ky) \\ &= \sum_{k \in K} \sum_{y \in X(1)_x} \frac{w(kx \cup ky)}{2w(kx)} f'(ky) = \sum_{k \in K} \sum_{y \in X(1)_{kx}} \frac{w(kx \cup y)}{2w(kx)} f'(y) \\ &= \sum_{k \in K} (\mathcal{A}f)(kx) = \sum_{k \in K} \lambda f(kx) = \lambda g'[x]. \end{aligned}$$

Thus,  $\lambda g' = g'$  and  $\lambda \in \text{Spec } \mathcal{B}$ . Applying this argument for the  $A$ -eigenfunctions  $f''$ ,  $1_{X(0)}$  and  $1_{X_{\{0\}}} - 1_{X_{\{1\}}}$  shows that we also have  $1, -1, -\lambda \in \text{Spec } \mathcal{B}$ . Since  $\lambda \neq -\lambda$  (because  $\lambda \neq 0$ ) and  $\text{Spec } \mathcal{B} \subseteq \mathbb{R}$ , this means that

$$2 + 2\lambda^2 = (-1)^2 + (-\lambda)^2 + \lambda^2 + 1^2 \leq \text{Tr}(\mathcal{B}^2). \quad (7.1)$$

We now bound  $\text{Tr}(\mathcal{B}^2)$  from above. Fix a set of representatives  $U$  for  $K \backslash X(0)$  and write  $1_{[x]} \in C^0(Y, \mathbb{C})$  for the characteristic function of  $\{[x]\}$ . Then

$$\begin{aligned} \text{Tr}(\mathcal{B}^2) &= \sum_{x \in U} (\mathcal{B}^2 1_{[x]})(x) = \sum_{x \in U} \sum_{y \in X(1)_x} \frac{w(x \cup y)}{2w(x)} (\mathcal{B} 1_{[x]})(y) \\ &= \sum_{x \in U} \sum_{y \in X(1)_x} \sum_{z \in X(1)_y} \frac{w(x \cup y)w(y \cup z)}{4w(x)w(y)} 1_{[x]}(z) = \frac{1}{k_0 k_1} \sum_{x \in U} |L(x)|, \end{aligned}$$

where  $L(x)$  is the set of triples  $(x, y, z) \in X(0)^3$  with  $\{x, y\}, \{y, z\} \in X(1)$  and  $z \in [x]$ . Let  $s'$  be the unique 0-face in  $U$  with  $d(s, s') = m$  and let  $s''$  be the unique 0-face in  $U$  with  $d(s, s'') = m - 1$ ; they exist by (ii) above. Put  $\epsilon = m \bmod 2$  and note that  $s' \in X_{\{\epsilon\}}$ . We analyze the size of  $L(x)$  by splitting into four cases:

- I)  $x = s'$ : Let  $(s', y, z) \in L(s')$ . There are  $k_\epsilon$  possibilities for  $y$ . Since  $d(s, y) = m - 1 > 0$ , there is some  $z' \in X(1)_y$  with  $d(s, z') < d(s, y) < d(s, s') = d(s, z)$ , so  $z' \neq z$ . This means that, for each  $y$ , there are at most  $k_{1-\epsilon} - 1$  possibilities for  $z$ . Thus,  $|L(s')| \leq k_\epsilon(k_{1-\epsilon} - 1) = k_0 k_1 - k_{1-\epsilon}$ .
- II)  $x = s''$ :  $|L(s'')| \leq k_0 k_1$ ; in fact, this holds for any  $x \in U$ .
- III)  $x = s$ : Since  $[s] = \{s\}$ , we have  $L(s) = \{(s, y, s) \mid y \in X(1)_s\}$ , so  $|L(s)| = k_0$ .
- IV)  $x \neq s, s', s''$ : Write  $t(x) = \{i\}$ . If  $(x, y, z) \in L(x)$ , then both  $x$  and  $y$  are not of maximum distance from  $s$  in  $X(0)$ . Noting that  $d(s, x) = d(s, z)$  (because  $z \in [x]$ ), (iii) implies that  $y$  is uniquely determined by  $x$  if  $d(s, y) < d(s, x)$  and  $z$  is uniquely determined by  $y$  in if  $d(s, y) > d(s, z)$ . Thus,  $|L(x)| \leq 1 \cdot k_{1-i} + k_i \cdot 1 = k_0 + k_1$ .

By (i),  $|U| \leq m + 1$ , so we conclude that

$$\text{Tr}(\mathcal{B}^2) \leq \frac{(k_0 k_1 - k_{1-\epsilon}) + k_0 k_1 + k_0 + (m - 2)(k_0 + k_1)}{k_0 k_1} \leq 2 + \frac{2(m - 2)}{q},$$

where the inequality holds because  $q \leq k_0 \leq k_{1-\epsilon}$ . Combining this with (7.1) gives the desired conclusion

$$\lambda^2 \leq \frac{m - 2}{q}.$$

This proves the theorem when  $r = 1$ .

Suppose now that  $r > 1$ . Assumptions (1) and (2) imply that all the positive-dimensional links of  $X$  are connected, and for every  $z \in X(r - 2)$ , the complex  $X_z$  is a 1-dimensional spherical building admitting a strongly transitive action and having  $m(X_z) \leq m$ . Since we proved the theorem when  $r = 1$ , the weighted graph  $(X_z, w_{X_z})$  is a 2-partite  $[-\sqrt{\frac{m-2}{q}}, \sqrt{\frac{m-2}{q}}]$ -expander. Moreover, our assumption  $q \geq r^2(m - 2)$  implies that  $\sqrt{\frac{m-2}{q}} \leq \frac{1}{r}$ . Thus, by a theorem of Oppenheim [27, Corollary 5.6]<sup>11</sup> (see also the formula at the end of [27, Theorem 1.4]), if  $(X, w)$  is an  $(r + 1)$ -partite  $[-r\lambda, \lambda]$ -expander for

$$\lambda = 1 - \frac{r(1 - \sqrt{\frac{m-2}{q}}) - (r - 1)}{(r - 1)(1 - \sqrt{\frac{m-2}{q}}) - (r - 2)} = \frac{\sqrt{m - 2}}{\sqrt{q} - (r - 1)\sqrt{m - 2}}. \quad \square$$

**Example 7.3.** Let  $\mathbb{F}_q$  be a finite field with  $q$  elements and let  $X = A_2(\mathbb{F}_q)$  (notation as in Example 7.1). Then  $X$  is a  $(q + 1)$ -thick building admitting a strongly transitive action, and  $m(X) = m(A_2) = 3$ . Thus, by Theorem 7.2,  $(X, w_X)$  is a bipartite  $[-\frac{1}{\sqrt{q+1}}, \frac{1}{\sqrt{q+1}}]$ -expander. This agrees with the well-known fact that  $\text{Spec}(\mathcal{A}_{X,w}) = \{\pm 1, \pm \frac{\sqrt{q}}{q+1}\}$  with 1 and  $-1$  occurring with multiplicity 1. (In fact, the counting argument in cases I)–IV) in the proof can be slightly improved to make the bounds match.)

<sup>11</sup>There is a typo in [27, Corollary 5.6]: the expression “ $1 - (n - k)f^{n-k-2}(\lambda)$ ” should be “ $1 + (n - k - 1)f^{n-k-2}(\lambda)$ ”.

**Corollary 7.4.** *Let  $X, w, r, q, m$  be as in Theorem 7.2 and assume  $q \geq r^2(m-2)$ . Then  $(X, w, R_+)$  is a  $(1 - \frac{\sqrt{m-2}}{\sqrt{q-(r-1)\sqrt{m-2}}})$ -coboundary expander in dimension 0 for every nontrivial abelian group  $R$ .*

*Proof.* This follows from Theorem 7.2 and Corollary 5.3.  $\square$

**Lemma 7.5.** *Let  $X$  be a pure  $r$ -dimensional (finite) simplicial complex and let  $w$  be its canonical weight function. Suppose that  $X$  is  $q$ -thick, i.e., every  $(r-1)$ -face of  $X$  is contained in at least  $q$   $r$ -faces. Then, for every  $e \in X(1)$  and  $x \in X(0)_{\subseteq e}$ , we have  $\frac{w(e)}{w(x)} \leq \frac{2}{q+r-1}$ .*

*Proof.* Fix  $e \in X(1)$  and let  $x, y$  be the 0-faces of  $e$ . It follows readily from the defining properties of  $w$  in §2B that  $\frac{w(e)}{w(x)} = \frac{2|X(r)_{\supseteq e}|}{r|X(r)_{\supseteq x}|}$ . Let  $c \in X(r)_{\supseteq e}$ . By assumption, the set  $N(c) := X(r)_{\supseteq c-y} - \{c\}$  has at least  $q-1$  members. Note that the members of  $N(c)$  are  $r$ -faces containing  $x$  but not  $e$ . Let  $c' \in X(r)_{\supseteq x} - X(r)_{\supseteq e}$ . If  $c \in X(r)_{\supseteq e}$  is an  $r$ -face such that  $c' \in N(c)$ , then  $c \cup c' = c' \cup y$ , which means that  $e \subseteq c \subseteq c' \cup y$ . Since  $|c' \cup y| = r+2$  and  $|e| = 2$ , there are at most  $r$  faces  $c \in X(r)_{\supseteq e}$  with  $c' \in N(c)$ . As a result,

$$r(|X(r)_{\supseteq x}| - |X(r)_{\supseteq e}|) \geq \sum_{c \in X(r)_{\supseteq e}} |N(c)| \geq (q-1)|X(r)_{\supseteq e}|.$$

By rearranging, we find that  $\frac{|X(r)_{\supseteq e}|}{|X(r)_{\supseteq x}|} \leq (1 + \frac{q-1}{r})^{-1} = \frac{r}{q+r-1}$ , so  $\frac{w(e)}{w(x)} \leq \frac{2}{q+r-1}$ .  $\square$

**Corollary 7.6.** *Let  $X, w, r, q, m$  be as in Theorem 7.2 and assume  $q \geq r^2(m-2)$ . Let  $R$  be an abelian group and let  $\{R_x\}_{x \in X - \{\emptyset\}}$  be subgroups of  $R$  satisfying conditions (1) and (2) of Theorem 6.1 (after setting  $R_\emptyset = \{0_R\}$ ), e.g., such that the summation map  $\bigoplus_{x \in X} R_x \rightarrow R$  is injective. For every  $x \in X$ , put  $\mathcal{C}(x) = \sum_{y \subset x} R_y$ . Then  $\mathcal{C}$  is a subsheaf of  $R_+$  and  $(X, w, R_+/\mathcal{C})$  an  $\varepsilon$ -coboundary expander in dimension 0 for*

$$\varepsilon = \frac{2r}{5r+2} - \frac{(4r^3+4r)\sqrt{m-2}}{(5r+2)(\sqrt{q}-(r-1)\sqrt{m-2})} - \frac{14r+4}{(5r+2)(q+r-1)}.$$

*Proof.* By Theorem 7.2,  $(X, w)$  is an  $(r+1)$ -partite  $[-r\lambda, \lambda]$ -expander for  $\lambda = \frac{\sqrt{m-2}}{q-(r-1)\sqrt{m-1}}$ , and by Lemma 7.5, we have  $t := \max\{\frac{w(e)}{w(x)} \mid e \in X(1), x \in X(0)_{\subseteq e}\} \leq \frac{2}{q+r-1}$ . In particular,  $s := \max\{w(e) \mid e \in X(1)\} \leq t = \frac{2}{q+r-1}$ . Now apply Theorem 6.2. (The expansion constant guaranteed by the theorem is slightly better than the simpler expression used in the corollary.)  $\square$

**Remark 7.7.** When  $r = \dim X > 1$ , the constants  $t$  and  $s$  of Theorem 6.2 are often much smaller than the bounds used in the proof of Corollary 7.6. Using a better upper bound on  $t$  and  $s$  will decrease the absolute value of right fraction in the expression for  $\varepsilon$  in Corollary 7.6. Also, when  $r = \dim X = 1$ , we may eliminate  $s$  by Remark 6.11, thus changing the right fraction in the expression for  $\varepsilon$  to  $-t = -\frac{2}{(q+r-1)}$ .

**Example 7.8.** The 0-dimensional coboundary expansion  $\varepsilon$  provided by Corollary 7.6 and Remark 7.7 is detailed in the following table for some spherical  $q$ -thick buildings of dimensions 1 and 2 admitting a strongly transitive action.

$\dim X$	$T(X)$	$\varepsilon$	$\varepsilon > 0$ if
1	$A_2$	$\frac{2}{7} - \frac{8}{7\sqrt{q}} - \frac{2}{q}$	$q \geq 29$
	$C_2$	$\frac{2}{7} - \frac{8\sqrt{2}}{7\sqrt{q}} - \frac{2}{q}$	$q \geq 45$
	$G_2$	$\frac{2}{7} - \frac{16}{7\sqrt{q}} - \frac{2}{q}$	$q \geq 78$
2	$A_3$	$\frac{1}{3} - \frac{10}{3(\sqrt{q}-1)} - \frac{8}{3(q+1)}$	$q \geq 136$
	$C_3$	$\frac{1}{3} - \frac{10\sqrt{2}}{3(\sqrt{q}-\sqrt{2})} - \frac{8}{3(q+1)}$	$q \geq 257$

## 8 Further Questions

We finish with several questions about possible extensions of Theorems 6.1 and 6.2.

If not indicated otherwise,  $(X, w)$  is assumed to be a weighted graph (resp.  $(r + 1)$ -partite weighted simplicial complex) which is a  $[-\lambda, \lambda]$ -expander (resp.  $(r + 1)$ -partite  $[-\lambda, \lambda]$ -expander) for some  $\lambda > 0$ . We let  $R$  be a nontrivial abelian group,  $\{R_x\}_{x \in X}$  be subgroups of  $R$  with  $R_\emptyset = 0$ , and define the subsheaf  $\mathcal{C}$  of  $R_+$  as in Theorem 6.1.

As  $\lambda$  approaches 0, the lower bound on the 0-dimensional coboundary expansion of  $(X, w, R_+/\mathcal{C})$  provided by Theorem 6.1 (resp. Theorem 6.2) approaches  $\frac{2}{5}$  (resp.  $\frac{2r}{5r+2}$ ). We expect that this could be improved.

**Question 8.1.** *Provided that all the  $\{R_x\}_{x \in X}$  are linearly disjoint in  $R$ , is it the case that the coboundary expansion of  $(X, w, R_+/\mathcal{C})$  in dimension 0 approaches 1 as  $\lambda \rightarrow 0$ ?*

Next, we ask whether condition (1) of Theorem 6.1 can be relaxed.

**Question 8.2.** *Let  $m : \mathbb{N} \rightarrow \mathbb{N} \cup \{0\}$  be a function satisfying  $0 \leq m(n) \leq n$  for all  $n \in \mathbb{N}$ . Assuming  $\lambda$  is fixed and sufficiently small, does the coboundary expansion of  $(X, w, R_+/\mathcal{C})$  in dimension 0 remains bounded away from 0 if (instead of condition (1) of Theorem 6.1) we require that every  $m(|X(0)|)$  of the  $\{R_x\}_{x \in X}$  are linearly disjoint in  $R$ ? More specifically:*

- (a) *Can we take  $m = o(n)$ ? Can we take  $m = O(1)$ ?*
- (b) *What if we also require that  $R_x = 0$  when  $x$  is not a 0-face?*

The motivation behind (b) is that in [8], we only need Theorems 6.1 and 6.2 in the special case where  $R_x = 0$  for all  $x \in X - X(0)$ . Also, if one allows  $R_x$  to be nonzero when  $x$  is an edge, then it is impossible to take  $m = o(\log n)$ . Indeed, suppose that  $X$  is a graph on  $n$  vertices, let  $\mathbb{F}$  be a finite field, and let  $R$  be an  $\mathbb{F}$ -vector space of dimension  $\geq n$ . Let  $\{f(v)\}_{v \in X(0)}$  be linearly independent vectors in  $R$ . For every edge  $e \in X(1)$ , set  $R_e = \mathbb{F}(f(u) - f(v))$ , where  $u, v$  are the 0-faces of  $e$ , and set  $R_u = 0$  for all  $u \in X(0)$ . Then, by design,  $f \in Z^0(X, R_+/\mathcal{C})$ , which means that  $H^0(X, R_+/\mathcal{C}) \neq 0$ , and therefore  $(X, w, R_+/\mathcal{C})$  is not an  $\varepsilon$ -coboundary expander in dimension 0 for any  $\varepsilon > 0$ . It is not difficult to check that if  $g$  is the girth of  $X$  (i.e., the length of the smallest cycle in  $X$ ), then every  $g - 1$  of the  $\{R_x\}_{x \in X}$  are linearly disjoint. Since for every prime number  $p$ , there exist infinitely many non-partite (resp. bipartite)  $(p + 1)$ -regular Ramanujan graphs of girth  $\frac{2}{3} \log_{p+1} n + O(1)$  (resp.  $\frac{4}{3} \log_{p+1} n + O(1)$ ), see [20], the answer to Question 8.2 is “no” if we take  $m = o(\log n)$ . However, the answer might still be “yes” under the assumptions of (b).

Finally, we ask whether Theorems 6.1 and 6.2 extend to higher dimensions.

**Question 8.3.** *Suppose that  $(X, w)$  is a weighted simplicial complex (resp.  $(r + 1)$ -partite weighted simplicial complex), and we are given subgroups  $\{R_x\}_{x \in X}$  of  $R$  which are all linearly disjoint. Form the subsheaf  $\mathcal{C}$  of  $R_+$  as in Theorem 6.1 and suppose that the underlying graph of  $X_z$  is a  $[-\lambda, \lambda]$ -expander (resp.  $(\dim X - \dim z)$ -partite  $[-\lambda, \lambda]$ -expander) for every  $z \in X$  with  $\dim z \leq \dim X - 2$ . Provided  $\lambda$  is sufficiently small, is there an  $\varepsilon > 0$ , depending only on  $\lambda$ ,  $\dim X$  and  $r$ , such that  $(X, w, R_+/\mathcal{C})$  is an  $\varepsilon$ -coboundary expander in dimensions  $0, \dots, \dim X - 1$ ?*

## References

- [1] Peter Abramenko and Kenneth S. Brown. *Buildings*, volume 248 of *Graduate Texts in Mathematics*. Springer, New York, 2008. Theory and applications.
- [2] Giuliana Davidoff, Peter Sarnak, and Alain Valette. *Elementary number theory, group theory, and Ramanujan graphs*, volume 55 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 2003.
- [3] Irit Dinur and Roy Meshulam. Near coverings and cosystolic expansion – an example of topological property testing, 2019. Preprint. arXiv:1909.08507.
- [4] Dominic Dotterrer and Matthew Kahle. Coboundary expanders. *J. Topol. Anal.*, 4(4):499–514, 2012.

- [5] Dominic Dotterrer, Tali Kaufman, and Uli Wagner. On expansion and topological overlap. *Geom. Dedicata*, 195:307–317, 2018.
- [6] Shai Evra and Tali Kaufman. Bounded degree cosystolic expanders of every dimension. In *STOC'16—Proceedings of the 48th Annual ACM SIGACT Symposium on Theory of Computing*, pages 36–48. ACM, New York, 2016.
- [7] Shai Evra and Tali Kaufman. Bounded degree cosystolic expanders of every dimension. 2017. Summary appeared in STOC'16—Proceedings of the 48th Annual ACM SIGACT Symposium on Theory of Computing.
- [8] Uriya A. First and Tali Kaufman. On good 2-query locally testable codes from sheaves on high dimensional expanders. 2022. Preprint.
- [9] Shmuel Friedland and Reinhard Nabben. On Cheeger-type inequalities for weighted graphs. *J. Graph Theory*, 41(1):1–17, 2002.
- [10] Mikhail Gromov. Singularities, expanders and topology of maps. Part 2: From combinatorics to topology via algebraic isoperimetry. *Geom. Funct. Anal.*, 20(2):416–526, 2010.
- [11] Danijela Horak and Jürgen Jost. Spectra of combinatorial Laplace operators on simplicial complexes. *Adv. Math.*, 244:303–336, 2013.
- [12] Bruce W. Jordan and Ron Livné. Ramanujan local systems on graphs. *Topology*, 36(5):1007–1024, 1997.
- [13] Tali Kaufman, David Kazhdan, and Alexander Lubotzky. Isoperimetric inequalities for Ramanujan complexes and topological expanders. *Geom. Funct. Anal.*, 26(1):250–287, 2016.
- [14] Tali Kaufman and Alexander Lubotzky. High dimensional expanders and property testing. In *ITCS'14—Proceedings of the 2014 Conference on Innovations in Theoretical Computer Science*, pages 501–506. ACM, New York, 2014.
- [15] Tali Kaufman and David Mass. Cosystolic expanders over any abelian group. *Electronic Colloquium on Computational Complexity (ECCC)*, 25:134, 2018.
- [16] Tali Kaufman and David Mass. Unique-Neighbor-Like Expansion and Group-Independent Cosystolic Expansion. In Hee-Kap Ahn and Kunihiko Sadakane, editors, *32nd International Symposium on Algorithms and Computation (ISAAC 2021)*, volume 212 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 56:1–56:17, Dagstuhl, Germany, 2021. Schloss Dagstuhl – Leibniz-Zentrum für Informatik.
- [17] Tali Kaufman and Izhar Oppenheim. Coboundary and cosystolic expansion from strong symmetry, 2021.
- [18] Tali Kaufman and Izhar Oppenheim. High dimensional expansion implies amplified local testability. *CoRR*, abs/2107.10488, 2021.
- [19] Nathan Linial and Roy Meshulam. Homological connectivity of random 2-complexes. *Combinatorica*, 26(4):475–487, 2006.
- [20] A. Lubotzky, R. Phillips, and P. Sarnak. Ramanujan graphs. *Combinatorica*, 8(3):261–277, 1988.
- [21] Alexander Lubotzky. High dimensional expanders. In *Proceedings of the International Congress of Mathematicians—Rio de Janeiro 2018. Vol. I. Plenary lectures*, pages 705–730. World Sci. Publ., Hackensack, NJ, 2018.
- [22] Alexander Lubotzky, Zur Luria, and Ron Rosenthal. Random Steiner systems and bounded degree coboundary expanders of every dimension. *Discrete Comput. Geom.*, 62(4):813–831, 2019.
- [23] Alexander Lubotzky and Roy Meshulam. Random Latin squares and 2-dimensional expanders. *Adv. Math.*, 272:743–760, 2015.

- [24] Alexander Lubotzky, Roy Meshulam, and Shahar Mozes. Expansion of building-like complexes. *Groups Geom. Dyn.*, 10(1):155–175, 2016.
- [25] Alexander Lubotzky, Beth Samuels, and Uzi Vishne. Ramanujan complexes of type  $\tilde{A}_d$ . *Israel J. Math.*, 149:267–299, 2005. Probability in mathematics.
- [26] R. Meshulam and N. Wallach. Homological connectivity of random  $k$ -dimensional complexes. *Random Structures Algorithms*, 34(3):408–417, 2009.
- [27] Izhar Oppenheim. Vanishing of cohomology and property (T) for groups acting on weighted simplicial complexes. *Groups Geom. Dyn.*, 9(1):67–101, 2015.