

Black hole solutions as topological thermodynamic defects

Shao-Wen Wei^{1,2} *, Yu-Xiao Liu^{1,2} †, Robert B. Mann³ ‡

¹Lanzhou Center for Theoretical Physics, Key Laboratory of Theoretical Physics of Gansu Province, School of Physical Science and Technology, Lanzhou University, Lanzhou 730000, People's Republic of China,

²Institute of Theoretical Physics & Research Center of Gravitation, Lanzhou University, Lanzhou 730000, People's Republic of China,

³ Department of Physics & Astronomy, University of Waterloo, Waterloo, Ont. Canada N2L 3G1
(Dated: August 4, 2022)

In this work, employing the generalized off-shell free energy, we treat black hole solutions as defects in the thermodynamic parameter space. The results show that the positive and negative winding numbers corresponding to the defects indicate the local thermodynamical stable and unstable black hole solutions, respectively. The topological number W defined as the sum of the winding numbers for all the black hole branches at an arbitrary given temperature is found to be a universal number independent of the black hole parameters. Moreover, this topological number only depends on the thermodynamic asymptotic behaviors of the black hole temperature at small and large black hole limits. Different black hole systems are characterized by three classes via this topological number. This number could help us well understanding the black hole thermodynamics, and further shed new light on the fundamental nature of quantum gravity.

PACS numbers: 04.70.Dy, 04.70.Bw, 05.70.Ce

Introduction— Black holes predicted by general relativity play a central role in modern physics. Observations of binary black hole mergers by the LIGO/Virgo collaboration [1] and reconstruction of event-horizon-scale images of M87* by the Event Horizon Telescope (EHT) [2] have opened new windows to study the strong gravitational nature of black holes. Alternatively, seeking underlying characteristic properties of black holes is also quite valuable, since they can yield features that can be tested by further astronomical observations.

One such approach is topology: by ignoring detailed structure, generic properties of a system can be discerned. This approach has been extensively employed to study many physical phenomena such as magnetic monopoles, the quantum Hall effect, and superconductivity. A key concept is that of defects, which are generally thought to be related to zero points of a field at $\vec{x} = \vec{z}$

$$\phi(\vec{x})|_{\vec{x}=\vec{z}} = 0. \quad (1)$$

in a space, and which can reveal certain properties of field configurations. In different dimensions, a defect can be a point charge, a string, or even a domain wall. The most simple topological quantity associated with the zero point of a field is its winding number. Equipped with it, the nature of a system possessing defects can be determined.

Of particular recent interest has been a special vector constructed in the coordinate space of a black hole spacetime [3, 4] by making use of null geodesics. The light ring of a black hole is located exactly at the zero point of this vector field. After calculating the winding number corresponding to this zero point, a four-dimensional stationary

nonextremal spinning black hole in asymptotically flat spacetime with a topologically spherical Killing horizon was shown to allow at least one standard light ring outside the horizon for each sense of rotation [4]. This treatment was generalized to nonspinning black holes with more than one zero point being discovered [5]. The value of the winding number was found to be related to the stability of the light rings [6]. This study was further extended to the thermodynamic critical points of black holes [7, 8] and the timelike circular orbit [9].

In general relativity, a black hole solution satisfies the Einstein field equation, which we reformulate as

$$\mathcal{E}_{\mu\nu} \equiv G_{\mu\nu} - \frac{8\pi G}{c^4} T_{\mu\nu} = 0. \quad (2)$$

We propose, analogous to Eq. (1), that a physical black hole solution is a zero point of the tensor field $\mathcal{E}_{\mu\nu}$, at which all its components vanish. Although the Einstein equations also admit other defect-like solutions, such like cosmic strings and branes, we here only focus on black holes. As a result, we can endow a black hole solution with a topological charge as well. Adopting this topological argument, we can then study the local and global particular properties of a black hole. Different black hole solutions are also expected to be divided into different classes.

Realization in thermodynamics— As a starting point to realize the idea that a black hole solution possesses a topological charge, we begin with black hole thermodynamics.

In 1977, Gibbons and Hawking [10] proposed that the partition function of a canonical ensemble for black holes can be evaluated with its Euclidean action in the form of the gravitational path integral. In “zero-loop” approximation, it reads

$$\mathcal{Z} = e^{-\beta F} = \int D[g] e^{-\frac{\mathcal{I}}{\hbar}} \sim e^{-\frac{\mathcal{I}}{\hbar}}, \quad (3)$$

*E-mail: weishw@lzu.edu.cn

†E-mail: liuyx@lzu.edu.cn

‡E-mail: rbmann@uwaterloo.ca

where F and \mathcal{I} is the free energy and Euclidean action of the black hole. The period β of the Euclidean time is the inverse of the black hole temperature. For a Schwarzschild black hole one has $\beta = 8\pi M$. Following this approach, it was found that negative heat capacity and imaginary energy fluctuations were produced. Soon afterwards, these shortcomings were solved by York [11] by imagining that the black hole is placed inside a cavity. Fixing the temperature T of the cavity surface, the results show that the heavy-mass black hole branch with mass $M > \sqrt{3}/8\pi T$ is thermodynamically stable, and thus the corresponding partition function is well defined.

In order to describe a black hole with an arbitrary mass, York defined a generalized free energy following the standard definition of the free energy while with the mass and temperature being two independent variables [11]. This naturally extends one more dimension for the thermodynamic parameter space of the black hole. Furthermore, the generalized free energy reduces to the standard one when the relation between the black hole mass and temperature is satisfied [12].

Inspired by this, we would like to introduce the following generalized free energy

$$\mathcal{F} = E - \frac{S}{\tau}, \quad (4)$$

for a black hole system with energy E and entropy S . The parameter τ is an extra variable having the dimension of time, and can be thought as the inverse temperature of the cavity enclosing the black hole. Here we let the time parameter τ vary freely instead of the mass [11]. In general this generalized free energy is off-shell except at

$$\tau = T^{-1}, \quad (5)$$

at which the black hole solution satisfies the Einstein field equations (2). We emphasize that the generalized free energy is well defined, meaning that the black hole system is uniquely determined by fixing τ , the black hole horizon r_h and other black hole parameters. Recently a similar generalized free energy (4) has been used to study the dynamic evolution of Hawking-Page phase transitions and small-large black hole phase transitions [13–15], and black hole triple points with oscillatory behavior [16].

We now describe the explicit construction for the vector ϕ in (1) via a thermodynamic approach. We define

$$\phi = \left(\frac{\partial \mathcal{F}}{\partial r_h}, -\cot \Theta \csc \Theta \right) \quad (6)$$

where, inspired by the axis limit [4], we introduce a parameter $0 \leq \Theta \leq \pi$ for convenience. At $\Theta = 0, \pi$, the component ϕ^Θ diverges and thus the direction of the vector is outwards.

More importantly, the zero points of ϕ correspond to the conditions $\Theta = \pi/2$ and $\tau = T^{-1}$ in (5). This confirms that the black hole solution exactly meets the zero point of the vector ϕ . Therefore, from the viewpoint of

topology, we can endow each black hole solution with a topological charge by using ϕ .

Following Duan's ϕ -mapping topological current theory [17, 18], we can introduce the topological current as

$$j^\mu = \frac{1}{2\pi} \epsilon^{\mu\nu\rho} \epsilon_{ab} \partial_\nu n^a \partial_\rho n^b, \quad \mu, \nu, \rho = 0, 1, 2, \quad (7)$$

where $\partial_\nu = \frac{\partial}{\partial x^\nu}$ and $x^\nu = (\tau, r_h, \Theta)$. The unit vector is defined as $n^a = \frac{\phi^a}{\|\phi\|}$ ($a = 1, 2$) with $\phi^1 = \phi^{r_h}$ and $\phi^2 = \phi^\Theta$. Here the parameter τ can be considered as a time parameter of the topological defect. Moreover, it is easy to check that this topological current is conserved, i.e., $\partial_\mu j^\mu = 0$. By making use of the Jacobi tensor $\epsilon^{ab} J^\mu \left(\frac{\phi}{x} \right) = \epsilon^{\mu\nu\rho} \partial_\nu \phi^a \partial_\rho \phi^b$ and the two-dimensional Laplacian Green function $\Delta_{\phi^a} \ln \|\phi\| = 2\pi \delta^2(\phi)$, the topological current can be reexpressed as [5]

$$j^\mu = \delta^2(\phi) J^\mu \left(\frac{\phi}{x} \right). \quad (8)$$

Obviously, j^μ is nonzero only at $\phi^a(x^i) = 0$, and we denote its i -th solution as $\vec{x} = \vec{z}_i$. The density of the topological current is then [19]

$$j^0 = \sum_{i=1}^N \beta_i \eta_i \delta^2(\vec{x} - \vec{z}_i). \quad (9)$$

The positive Hopf index β_i counts the number of the loops that ϕ^a makes in the vector ϕ space when x^μ goes around the zero point z_i , and the Brouwer degree $\eta_i = \text{sign}(J^0(\phi/x)_{z_i}) = \pm 1$. Given a parameter region Σ , the corresponding topological number can be obtained

$$W = \int_{\Sigma} j^0 d^2x = \sum_{i=1}^N \beta_i \eta_i = \sum_{i=1}^N w_i, \quad (10)$$

where w_i is the winding number for the i -th zero point of ϕ contained in Σ . If two given loops $\partial\Sigma$ and $\partial\Sigma'$ enclose the same zero point of ϕ , they possess the same winding number. Alternatively, if there is no zero point in the enclosed region, we will have $W = 0$. If Σ is the neighbourhood of a zero point of ϕ , it will yield local topological properties, whereas if Σ is the entire parameter space, the global topological W number will be revealed.

In the above approach, the isolated zero points require the Jacobian $J_0(\phi/x) \neq 0$. If this condition is violated, the defect bifurcates [20]. Considering the x^1 direction, this bifurcation phenomenon occurs when

$$\frac{dx^1}{dx^0} = \frac{dr_h}{d\tau} = \infty. \quad (11)$$

Via Taylor expansion near the bifurcation point, one obtains

$$\tau - \tau^* = \frac{1}{2} \frac{d^2\tau}{dr_h^2} \Big|_{r_h^*} (r_h - r_h^*)^2 + \mathcal{O}(r_h - r_h^*)^3, \quad (12)$$

where τ^* and r_h^* take values at the bifurcation point. For $\frac{d^2\tau}{dr_h^2}|_{r_h^*} > 0$, it denotes a generation point, otherwise it is an annihilation one.

Local and global topological properties—Following Eq. (3), one can obtain the free energy by calculating the first-order Euclidean Einstein action

$$\mathcal{I} = -\frac{1}{16\pi} \int \sqrt{g} R d^4x + \frac{1}{8\pi} \oint \sqrt{\gamma} K d^3x - \mathcal{I}_{\text{subtract}} \quad (13)$$

where K is the trace of the extrinsic curvature, γ_{ij} is the induced metric on the boundary, and $\mathcal{I}_{\text{subtract}}$ is the subtraction term.

For the Schwarzschild black hole, one easily has $E = \partial_\beta \mathcal{I} = M$ and $S = (\beta \partial_\beta \mathcal{I} - \mathcal{I}) = 4\pi M^2$ [10]. Accordingly, the black hole thermodynamic relations can be deduced straightforwardly. For the Schwarzschild black hole, we obtain the generalized free energy from the definition (4)

$$\mathcal{F} = \frac{r_h}{2} - \frac{\pi r_h^2}{\tau}, \quad (14)$$

where $r_h = 2M$ has been used. Then the components of the constructed vector ϕ can be calculated

$$\phi^{r_h} = \frac{1}{2} - \frac{2\pi r_h}{\tau}, \quad (15)$$

$$\phi^\Theta = -\cot \Theta \csc \Theta. \quad (16)$$

We show the unit vector field n on a portion of the Θ - r_h plane in Fig. 1(a) for the Schwarzschild black hole with $\tau = 4\pi r_0$ with r_0 an arbitrary length scale set by the size of a cavity surrounding the black hole.

From the figure, the zero point is located at $r_h/r_0 = 1$ and $\Theta = \pi/2$. For a black hole, $\tau = 1/T_{BH} = 8\pi M$, giving $r_0 = 2M$ yielding $r_h = 2M$ as expected. Since the winding number w is independent of these loops enclosing the zero point, we can calculate it for an arbitrary loop; for example see C_1 given in Fig. 1(a). Performing the calculation, we obtain the winding number $w = -1$. If we choose an alternate orientation convention by adding a minus sign in ϕ^{r_h} in (6), we obtain $w = 1$ instead. However the winding numbers of other types of black holes will likewise be changed.

Considering the gravitational action together with the electromagnetic field, it is easy to find $E = \frac{r_h^2 + Q^2}{2r_h}$ and $S = \pi r_h^2$ for the Reissner-Nordström (RN) black hole. Thus the generalized free energy is

$$\mathcal{F} = \frac{r_h^2 + Q^2}{2r_h} - \frac{\pi r_h^2}{\tau}. \quad (17)$$

Note that for the RN black hole, the radius of the horizon has a lower bound $r_h = Q$, corresponding to an extremal black hole with vanishing temperature. However, here τ can be thought as the inverse temperature of the cavity. Thus r_h is independent of τ , and can vary freely, ignoring the lower bound. Employing it, we plot the unit vector field n in Fig. 1(b) for arbitrarily chosen typical values $\tau/r_0 = 34.48$ and $Q/r_0 = 1$.




FIG. 1: The red arrows represent the unit vector field n on a portion of the r_h - Θ plane. The zero points (ZPs) marked with black dots are at $(r_h, \Theta) = (1, \frac{\pi}{2})$, $(1.46, \frac{\pi}{2})$, and $(2.15, \frac{\pi}{2})$, for ZP_1 , ZP_2 , and ZP_3 , respectively. The blue contours C_i are closed loops enclosing the zero points. (a) The unit vector field for the Schwarzschild black hole with $\tau/r_0 = 4\pi$. (b) The unit vector field for the RN black hole with $\tau/r_0 = 34.48$ and $Q/r_0 = 1$.

We find two zero points, ZP_2 and ZP_3 , at $r_h/r_0 = 1.46$ and 2.15 respectively. A detailed calculation shows that their respective winding numbers are $w = 1$ and -1 . We also obtain the heat capacities $C_Q/r_0^2 = 18.01$ and -64.81 at constant charge $Q/r_0 = 1$ for the respective positive and negative zero points. Noting that each zero point of the unit vector has a winding number 1 or -1 , from this we conjecture that winding number is related to local thermodynamic stability, with positive/negative values corresponding to stable/unstable black hole solutions.

Turning to global properties of the topology, if we choose the region Σ as the whole parameter space or the loop $\partial\Sigma$ as the boundary of the parameter space, i.e., $(0 < \Theta < \pi) \cup (0 < r_h < \infty)$, we can obtain the topological number W for a black hole solution. This global property can be used to classify different black hole solutions. Since we here take the axis limit, where the direction of the vector ϕ is up at $\Theta = \pi$ and down at $\Theta = 0$, the value of the topological number W actually depends on the direction of the vector at $r_h = 0$ and ∞ .

This suggests that black hole solutions sharing the same behavior at $r_h = 0$ and ∞ possess the same topological number, and thus are topologically equivalent.

By solving the equation $\phi^{r_h} = 0$, we can obtain a curve in the $r_h - \tau$ plane. The results are

$$\tau = 4\pi r_h \quad (18)$$

and

$$\tau = \frac{4\pi r_h^3}{r_h^2 - Q^2} \quad (19)$$

for the Schwarzschild and RN black holes, respectively. To show zero points of the component ϕ^{r_h} , we plot these two curves in Fig. 2. For large τ (e.g. $\tau = \tau_2$) there are respectively one and two intersection points for the Schwarzschild and RN black holes. The intersection points exactly satisfy the condition (5), and therefore denote the on-shell black hole solutions with temperature $T = \tau^{-1}$. In contrast to the Schwarzschild black hole, for $\tau < \tau_c$, the two intersection points for the RN black hole coincide and then disappear. Based on the local property of a zero point, we have the topological number $W = -1$ for the Schwarzschild black hole, while $W = 1 - 1 = 0$ for the RN black hole with the charge $Q/r_0 = 1$. In particular, at the point $\tau_c = 6\sqrt{3}\pi Q$, it is easy to get $\frac{d^2\tau}{dr_h^2} = 6\sqrt{3}\pi/Q > 0$ for the RN black hole. This suggests that τ_c is a generation point, which can also be observed in Fig. 2.

To investigate whether or not W depends on the charge, we examine the behavior of the curve $\tau(r_h)$ in Eq. (19) at the limit $r_h \rightarrow 0$ and ∞ . Obviously, for a non-vanishing charge Q , the vanishing/diverging behaviour of τ as $r_h \rightarrow 0/\infty$ does not change, and so W remains the same for different values of Q . For the Schwarzschild black hole with $Q = 0$, the behaviour of $\tau(r_h)$ for small r_h differs from the charged case, and so the topological numbers W for the two black hole solutions are different.

Another interesting black hole solution is the charged Reissner-Nordström anti-de Sitter (RN-AdS) black hole, which exhibits a small-large black hole phase transition reminiscent of the liquid-gas phase transition of van der Waals phase transition [21]. In the extended phase space, the cosmological constant is treated as the pressure P of the system [22]. The free energy for this black hole system is

$$\mathcal{F} = \frac{8\pi Pr_h^4 + 3r_h^2 + 3Q^2}{6r_h} - \frac{\pi r_h^2}{\tau}. \quad (20)$$

Taking the pressure $Pr_0^2 = 0.0022$ to be smaller than its critical value, we exhibit zero points of ϕ^{r_h} in the $r_h - \tau$ plane in Fig. 3 with $Q/r_0 = 1$. Quite different from the Schwarzschild and RN black holes shown in Fig. 2, we observe that there are three black hole branches for $\tau_a < \tau < \tau_b$, one small black hole branch for $\tau < \tau_a$ and one large black hole branch for $\tau > \tau_b$. Computing the winding number for these black hole branches, we find




FIG. 2: Zero points of the vector ϕ shown in the $r_h - \tau$ plane. The blue dashed and red solid lines are for the Schwarzschild black hole (Sch BH) and RN black hole (BH) with $Q/r_0 = 1$. The black dot with $\tau_c = 6\sqrt{3}\pi Q$ denotes the generation point for the RN black hole. At $\tau = \tau_1$, there is one Schwarzschild black hole, and at $\tau = \tau_2$, there are one Schwarzschild black hole and two RN black holes.




FIG. 3: Zero points of ϕ^{r_h} shown in the $r_h - \tau$ plane for the RN-AdS black hole with $Pr_0^2 = 0.0022$ and $Q/r_0 = 1$. The black solid, blue dashed, and red solid lines are for the small, intermediate, and large black hole branches, respectively. Black and blue dots are the annihilation and generation points. Different color regions have different number of the black hole branches. However their W number is constant and equals 1.

that the small and large black hole branches have $w = 1$, while the intermediate black hole branch has $w = -1$. Therefore, at this pressure the RN-AdS black hole always has $W = 1 - 1 + 1 = 1$, which is independent of τ . Since the pressure P is positive for the RN-AdS black hole, it does not affect the asymptotic behavior of τ at small and large r_h . As a result, the topological number W is always unity for the RN-AdS black hole.

Note that for these values of Pr_0^2 and Q/r_0 , one generation point and one annihilation point can be found at $\tau/r_0 = \tau_a/r_0 = 25.84$ and $\tau/r_0 = \tau_b/r_0 = 27.62$, respectively. If the pressure is larger than its critical value, r_h/r_0 will be a monotonically increasing function of τ/r_0 , and thus no bifurcation phenomenon will be observed. However the topological number W still keeps the same.

Conclusions— Summarizing our results in Table I, we

	Sch BH	RN BH	RN-AdS BH
W	-1	0	1
annihilation point	0	1	1 or 0
generation point	0	0	1 or 0

TABLE I: The topological number W , numbers of annihilation and generation points for the Schwarzschild, RN, and RN-AdS black holes.

have constructed a universal topological number W by treating the black hole solution as a defect from the viewpoint of thermodynamics. Different black hole solutions are characterized by different topological numbers and belong to different topological classes.

Locally, each zero point of the vector field defined in (6) with the generalized free energy exactly corresponds to one on-shell black hole solution. Thus each black hole solution is endowed with one winding number. Our study shows that the thermodynamic stability of the black hole is indicated from the value of the winding number. Positive/negative winding number corresponds to thermodynamically stable/unstable black hole solutions. Of particular interest are the annihilation and generation points, which may be quite important for the time evolution of a black hole placed in a cavity.

When considering the full parameter space, the topo-

logical number W provides us with a global topological property of a black hole solution. We have shown that the Schwarzschild black hole, RN black hole, and RN-AdS black hole have respectively $W = -1, 0,$ and $1,$ independent of the black hole parameters. Since the topological number W is the sum of the winding numbers for the zero points of the vector, and it is only dependent on the behavior of the curve $\tau(r_h)$ at small and large r_h limits, we conjectured that the topological number W can take three values, $-1, 0,$ and $1.$ This suggests that other black hole solutions shall be divided into three characteristic topological classes, a conjecture in need of further confirmation.

Our topological approach classifies each black hole solution into certain classes sharing the similar thermodynamic properties. It provides a considerable material for the topology of black hole thermodynamics. We expect to uncover the deeper nature of other black hole solutions via the topological approach.

Acknowledgements.—This work was supported by the National Natural Science Foundation of China (Grants No. 12075103, No. 11875151, and No. 12047501), the 111 Project (Grant No. B20063), and the National Key Research and Development Program of China (Grant No. 2020YFC2201503) and by the Natural Sciences and Engineering Research Council of Canada.

-
- [1] B. P. Abbott et al (Virgo, LIGO Scientific), *Observation of gravitational waves from a binary black hole merger*, Phys. Rev. Lett. **116**, 061102 (2016), [arXiv:1602.03837 [gr-qc]].
- [2] K. Akiyama et al. (Event Horizon Telescope), *First M87 Event Horizon Telescope Results. I. The Shadow of the Supermassive Black Hole*, Astrophys. J. **875**, L1 (2019), [arXiv:1906.11238 [astro-ph.GA]].
- [3] P. V. P. Cunha, E. Berti, and C. A. R. Herdeiro, *Light ring stability in ultra-compact objects*, Phys. Rev. Lett. **119**, 251102 (2017), [arXiv:1708.04211 [gr-qc]].
- [4] P. V. P. Cunha and C. A. R. Herdeiro, *Stationary Black Holes and Light Rings*, Phys. Rev. Lett. **124**, 181101 (2020), [arXiv:2003.06445 [gr-qc]].
- [5] S.-W. Wei, *Topological charge and black hole photon spheres*, Phys. Rev. D **102**, 064039 (2020), [arXiv:2006.02112 [gr-qc]].
- [6] M. Guo and S. Gao, *Universal Properties of Light Rings for Stationary Axisymmetric Spacetimes*, Phys. Rev. D **103**, 104031 (2021), [arXiv:2011.02211 [gr-qc]].
- [7] S.-W. Wei and Y.-X. Liu, *Topology of black hole thermodynamics*, Phys. Rev. D **105**, 104003 (2022) [arXiv:2112.01706 [gr-qc]].
- [8] M. B. Ahmed, D. Kubiznak, and R. B. Mann, *Vortex/anti-vortex pair creation in black hole thermodynamics*, [arXiv:2207.02147 [hep-th]].
- [9] S.-W. Wei and Y.-X. Liu, *Topology of equatorial timelike circular orbits around stationary black holes*, [arXiv:2207.08397 [gr-qc]].
- [10] G. W. Gibbons and S. W. Hawking, *Action integrals and partition functions in quantum gravity*, Phys. Rev. D **15**, 2752 (1977).
- [11] J. W. York, *Black-hole thermodynamics and the Euclidean Einstein action*, Phys. Rev. D **33**, 2092 (1986).
- [12] B. F. Whiting, *Black holes and gravitational thermodynamics*, Class. Quantum Grav. **7**, 15 (1990).
- [13] R. Li and J. Wang, *Thermodynamics and kinetics of Hawking-Page phase transition*, Phys. Rev. D **102**, 024085 (2020).
- [14] R. Li, K. Zhang and J. Wang, *Thermal Dynamic Phase Transition of Reissner-Nordström Anti-de Sitter Black Holes on Free Energy Landscape*, J. High Energ. Phys. **2020**, 90 (2020), [arXiv:2008.00495 [hep-th]].
- [15] S.-W. Wei, Y.-Q. Wang, and Y.-X. Liu, *Dynamic properties of thermodynamic phase transition for five-dimensional neutral Gauss-Bonnet AdS black hole on free energy landscape*, Nucl. Phys. B **976**, 115692 (2022), [arXiv:2009.05215 [gr-qc]].
- [16] S.-W. Wei, Y.-Q. Wang, Y.-X. Liu, and R. B. Mann, *Observing dynamic oscillatory behavior of triple points among black hole thermodynamic phase transitions*, Sci. China Phys. Mech. Astron. **64**, 270411 (2021), [arXiv:2102.00799 [gr-qc]].
- [17] Y. S. Duan and M. L. Ge, *SU(2) gauge theory and electrodynamics of N moving magnetic monopoles*, Sci. Sin. **9**, 1072 (1979).
- [18] Y. S. Duan, *The structure of the topological current*, SLAC-PUB-3301, (1984).

- [19] J. A. Schouton, *Tensor analysis for Physicists* (Clarendon, Oxford, 1951).
- [20] L.-B. Fu, Y.-S. Duan, and H. Zhang, *Evolution of the Chern-Simons vortices*, Phys. Rev. D **61**, 045004 (2000), [arXiv:hep-th/0112033].
- [21] D. Kubiznak and R. B. Mann, *P-V criticality of charged AdS black holes*, J. High Energy Phys. **1207**, 033 (2012), [arXiv:1205.0559[hep-th]].
- [22] D. Kastor, S. Ray, and J. Traschen, *Enthalpy and the mechanics of AdS black holes*, Class. Quant. Grav. **26**, 195011 (2009), [arXiv:0904.2765[hep-th]].