

Asymptotic freeness through unitaries generated by polynomials of Wigner matrices

Félix Parraud
KTH Royal Institute of Technology
parraud@kth.se

Kevin Schnelli
KTH Royal Institute of Technology
schnelli@kth.se

Abstract

We study products of functions applied in self-adjoint polynomials in deterministic matrices and independent Wigner matrices; we compute the deterministic approximations of such products and control the fluctuations. We focus on minimizing the assumption of smoothness on those functions while optimizing the error term with respect to N , the size of the matrices. As an application, we build on the idea that the long-time Heisenberg evolution associated to Wigner matrices generates asymptotic freeness as first shown in [8]. More precisely given P a self-adjoint non-commutative polynomial and Y^N a d -tuple of independent Wigner matrices, we prove that the quantum evolution associated to the operator $P(Y^N)$ yields asymptotic freeness for large times.

1 Introduction and main results

In this paper, we consider polynomials in several independent Wigner matrices Y_1^N, \dots, Y_d^N . In the early nineties, in his seminal work [36], Voiculescu showed that the fitting framework to understand spectral properties of such polynomials is Free Probability, a non-commutative probability theory with a notion of freeness analogous to independence in classical probability. For Gaussian Wigner matrices he proved that for any given collection of polynomials P_1, \dots, P_k and $i_j \in [1, d]$, almost surely,

$$\lim_{N \rightarrow \infty} \operatorname{tr}_N (P_1(Y_{i_1}^N) \cdots P_k(Y_{i_k}^N)) = \tau(P_1(x_{i_1}) \cdots P_k(x_{i_k})), \quad (1.1)$$

where tr_N denotes the normalized trace on $\mathbb{M}_N(\mathbb{C})$; $x_1, \dots, x_d \in \mathcal{C}_d$ is a system of d free semicircular variables and τ is the trace on the \mathcal{C}^* -algebra \mathcal{C}_d ; see Definition 2.7. This result was generalized in numerous ways, to begin with in [13], the author extended this result to non-Gaussian Wigner matrices coupled with deterministic matrices under certain assumptions; see also Theorem 5.4.5 of [3] for a proof without those assumptions on the deterministic matrices. It is further possible to consider continuous functions but only by approximating them with polynomials. For such results, one usually first proves the convergence in expectation, and then uses concentration inequalities to establish the almost sure convergence. One of the limitations of such combinatorial methods is that they are not well-suited to obtain estimates on the convergence rate in (1.1). Strong quantitative estimates for smooth functions

Date: August 3, 2022.

Keywords: Asymptotic freeness, concentration inequalities, quantum evolution.

MSC 2020: 46L54, 60B20, 15B52.

F.P. is supported by the Knut and Alice Wallenberg Foundation.

K.S. is supported in parts by the Swedish Research Council and the Knut and Alice Wallenberg Foundation.

were first obtained in [19] by Haagerup and Thorbjørnsen in the Gaussian case. They studied not only the convergence of the trace but also of the operator norm. These results were extended to include deterministic matrices in [28, 33], and to more general Wigner matrices in [1, 4]. Those papers however focus on studying the expectation rather than proving almost sure estimates. While in the Gaussian case measure concentration inequalities are usually sufficient to deduce almost sure results, one does not necessarily have such tools for general Wigner matrices. There are however ways around this problem, most notably [8]. We will give more details about their results and methods below. Finally, beyond the question of polynomials, it is also worth noting that the case of non-commutative rational functions was tackled in [10, 37].

The main result of our paper is an estimate for the trace as well as the matrix entries of any products of sufficiently smooth functions applied in polynomials in deterministic and independent Wigner matrices. We are in particular interested in optimizing the error term with respect to not only N but also the derivatives of our functions. This will be especially useful for Theorem 1.2 later below.

Theorem 1.1. *Let the following objects be given,*

- $Y^N = (Y_1^N, \dots, Y_d^N)$ independent real symmetric or complex Hermitian Wigner matrices of size N satisfying Assumption 2.1 below,
- $A^N = (A_1^N, \dots, A_q^N)$ deterministic matrices of size N , such that $\sup_{1 \leq i \leq q, N \in \mathbb{N}^*} \|A_i^N\| < \infty$,
- $x = (x_1, \dots, x_d)$ a system of free semicircular variables, free from A^N , i.e. they belong to the free product $\mathcal{C}_d(x) * \mathbb{M}_N(\mathbb{C})$ where \mathcal{C}_d is the C^* -algebra generated by x (see Definition 2.7 below),
- f_1, \dots, f_k , $k \geq 1$, functions such that either $f_i = \text{id}_{\mathbb{R}}$ or there exist a complex-valued measure μ_i such that

$$\forall t \in \mathbb{R}, \quad f_i(t) = \int_{\mathbb{R}} e^{ity} d\mu_i(y), \quad (1.2)$$

- P_1, \dots, P_k non-commutative polynomials, such that whenever $f_i \neq \text{id}_{\mathbb{R}}$, P_i is self-adjoint (see Sub-section 2.3).

Then with the convention $\|f_i\|_4 = 1$ if $f_i = \text{id}_{\mathbb{R}}$, and otherwise

$$\|f_i\|_4 := \int_{\mathbb{R}} (1 + y^4) d|\mu_i|(y), \quad (1.3)$$

we have the following result. For any $\varepsilon > 0$, with high probability,

$$\begin{aligned} & \text{tr}_N (f_1(P_1(Y^N, A^N)) \cdots f_k(P_k(Y^N, A^N))) \\ &= \tau_N (f_1(P_1(x, A^N)) \cdots f_k(P_k(x, A^N))) + \mathcal{O} \left(N^\varepsilon \frac{\max_i \|f_i\|_4}{N} \right), \end{aligned} \quad (1.4)$$

where tr_N is the normalized trace on $\mathbb{M}_N(\mathbb{C})$, while τ_N is the trace on the free product $\mathcal{C}_d * \mathbb{M}_N(\mathbb{C})$; see Definition 2.7. Moreover for $\mathbf{x}, \mathbf{y} \in \mathbb{C}^N$, we also have with high probability that

$$\begin{aligned} & \langle \mathbf{x}, f_1(P_1(Y^N, A^N)) \cdots f_k(P_k(Y^N, A^N)) \mathbf{y} \rangle \\ &= \langle \mathbf{x}, E_{\mathbb{M}_N(\mathbb{C})} [f_1(P_1(x, A^N)) \cdots f_k(P_k(x, A^N))] \mathbf{y} \rangle + \mathcal{O} \left(N^\varepsilon \frac{\max_i \|f_i\|_4}{\sqrt{N}} \|\mathbf{x}\|_2 \|\mathbf{y}\|_2 \right), \end{aligned} \quad (1.5)$$

where $E_{\mathbb{M}_N(\mathbb{C})}$ is the conditional expectation from $\mathcal{C}_d * \mathbb{M}_N(\mathbb{C})$ to $\mathbb{M}_N(\mathbb{C})$ (see Definition 2.8 and the remarks that follow, notably for some special cases where this conditional expectation is easy to compute). Finally if for every i , Y_i^N is a GUE or GOE random matrix, then one can replace $\|f_i\|_4$ by $\|f_i\|_2$ in all of the previous formulas.

Theorem 1.1 calls for the following remarks:

- The norm $\|\cdot\|_4$ is the one associated to the fourth Wiener space $W_4(\mathbb{R})$; we refer to [31], Section 3.2, for more information on the topic. In particular, if f and its Fourier transform are integrable, then f

satisfies Assumption (1.2) with the measure $d\mu(y) = \widehat{f}(y) dy$. Then assuming that $\|f\|_4$ is finite implies that f is four times differentiable and

$$\|f\|_4 = \int_{\mathbb{R}} (1 + y^4) |\widehat{f}(y)| dy = \int_{\mathbb{R}} |\widehat{f}(y)| + |\widehat{f^{(4)}}(y)| dy.$$

Thus heuristically $\|f\|_4$ is related to the fourth derivative of f .

- Note that in Theorem 1.1, one can take the functions $(f_i)_{i \in [1, k]}$ to depend on N although this implies that the norm in the error term will also depend on N . This allows us for example to consider mesoscopic test functions such as for $E \in \mathbb{R}$ and g_i satisfying Assumptions 1.2 and 1.3,

$$f_i : x \mapsto g_i(N^\alpha(x - E)).$$

However the error term will be of order $N^{4\alpha + \varepsilon - 1} \max_i \|g_i\|_4$ instead of $N^{\varepsilon - 1} \max_i \|f_i\|_4$ and the deterministic approximation will depend on N .

- The estimates in (1.4) and (1.5) are optimal in terms of the powers in N since the crucial concentration results in Propositions 3.1 and 3.3 below are themselves optimal with respect to N . However, we believe that the optimal power in $y = 1 + \max |y_i|$ in Proposition 3.1 and 3.3 can be improved to two, as it is the case for Gaussian ensembles, but see also Theorem 2.6 in [8] for $k \leq 4$. Thus in turn we expect that the estimates in (1.4) and (1.5) hold for functions in the second Wiener space; i.e. with $\|f\|_2$ instead of $\|f\|_4$.
- Lastly, we remark that the Wigner matrices Y^N can be both real symmetric or complex Hermitian, or a mix of them, and we do not need the matrix entries to have the same law.

One can notably compare Theorem 1.1 with Theorem 2.6 of [8] and Corollary 2.7 of [9]. The biggest difference is that we consider any polynomials P_1, \dots, P_k whereas in [8] the authors considered monomials of degree 1, i.e. $P_i(Y^N, A^N) = Y_{j_i}^N$ for some $j_i \in [1, d]$, or $P_i(Y^N, A^N) = A_{j_i}^N$ for some $j_i \in [1, q]$. For such products the key result to prove Theorem 2.6 in [8] is a multi-resolvent local law (Theorem 3.4 of their paper): Let $G^N(z) := (Y^N - z)^{-1}$, $z \in \mathbb{C} \setminus \mathbb{R}$, denote the resolvent or Green function of a single Wigner matrix Y^N . Then the Green function $G^N(z)$ is well approximated by $m_{sc}(z)I^N$ with $m_{sc}(z)$ the Stieltjes transform of the semicircle law, in averaged sense [14, 17, 18] as well as in isotropic sense [6, 24], for $|\Im z| \gg N^{-1}$, i.e. down to local scales slightly above the typical eigenvalue spacing. Theorem 3.4 in [8] gives the deterministic approximation for multi-resolvents of the form $G^N(z_1)A_1^N G^N(z_2)A_2^N \dots A_{k-1}^N G^N(z_k)A_k^N$ with optimal error terms and on local scales; where (A_i^N) are as in Theorem 1.1. The Helffer-Sjöstrand calculus allows then to extend this local law to observables of the form $f_1(Y^N)A_1^N f_2(Y^N) \dots f_k(Y^N)A_k^N$, for Sobolev functions (f_i) , in averaged and isotropic sense, yielding the results in Theorem 2.6 of [8]. Yet, local laws for a single Green function of a polynomial in Wigner matrices [2, 15, 29] are difficult to derive, partly due to hard to check stability conditions stemming from the linearization trick [19]. In particular for our desired results the local laws needed to be established up to the spectral edges, which for general polynomials seems challenging, not to speak of multi-resolvent local laws.

We circumvent this difficulty by working with the Fourier transform instead of the Stieltjes transform. A main novelty are concentration estimates for quantities of the form $e^{iP_1 y_1} R_1 \dots e^{iP_k y_k} R_k$ with (P_i) and (R_i) polynomials in Y^N and A^N , and $(y_i) \in \mathbb{R}^k$; see Propositions 3.1 and 3.3 below. These concentration results are obtained by a long-time continuous interpolation between Wigner matrices and the GUE applied to the non-commutative setting using recursive moment estimates [20, 27]; see [25, 26] for long-time interpolations for Green functions of single Wigner matrices. This strategy does not rely on the linearization of non-commutative polynomials and hence avoids stability issues. The concentration estimates in Propositions 3.1 and 3.3 are then combined with results in [33] to replace GUE matrices with semicircular variables by interpolating them with the help of free stochastic calculus. This strategy of interpolating between random matrices and free variables was first developed in [11] before being refined to get a better estimate of the remainder term in [33]. Via the Fourier transform our results in Theorem 1.1 follow for functions in the fourth Wiener space.

Another difference to [8] is that our method focuses on algebraic rather than combinatorial aspects. In [8] the main results, including the multi-resolvent local law, are stated using advanced combinatorics, such as free cumulant functions, partial traces or the Kreweeras complement. A key insight of [8] was

to connect resolvent expansions to non-crossing partitions and free cumulants. In this paper, we rely instead on the algebraic formalism of free probability, e.g. free products of C^* -algebras or conditional expectations, that allows us to formulate the convergence results in (1.4) and (1.5) in all generality, i.e. with P_i arbitrary polynomials in Y^N and A^N . In Subsection 2.2, we outline how to pass from an algebraic to a combinatorial formulation in some special cases.

Motivations for Theorem 1.1 come from free probability and mathematical physics. As explained in [8], the RAGE theorem (see [12], Theorem 5.8) states that given H a self-adjoint operator on an infinite dimensional Hilbert space, ϕ a state in the continuous spectral subspace of H , and C a compact operator, then the Heisenberg time evolution of C vanishes on ϕ when t goes to infinity, more precisely, the Cesaro mean of $\langle \phi, e^{itH} C e^{-itH} \phi \rangle$ vanishes for large t . Thus heuristically the RAGE theorem describes the asymptotic behavior of the Heisenberg time evolution under some assumptions on C and ϕ . For example one may consider the operator $H = P(x_1, \dots, x_d)$ where P is a polynomial and (x_i) are free semicircular variables. It is well-known that the spectrum of a polynomial in independent Wigner matrices behaves similarly to the one of the same polynomial evaluated in free semicircular variables, see for example Theorem 5.4.2 in [3]. Consequently one expects that up to an error which depends on N , the Heisenberg time evolution associated with the operator $H = P(Y_1^N, \dots, Y_d^N)$ will behave similarly. And indeed, from Theorem 1.1, we get that if P is such a polynomial, then for t and N large enough, we have for any bounded sequence of deterministic matrices (B^N) and (C^N) , as well as vectors $\mathbf{x}, \mathbf{y} \in \mathbb{C}^N$, that

$$\langle \mathbf{x}, e^{itP(Y^N)} C^N e^{-itP(Y^N)} \mathbf{y} \rangle \simeq \text{tr}_N(C^N) \times \langle \mathbf{x}, \mathbf{y} \rangle, \quad (1.6)$$

$$\text{tr}_N \left(e^{itP(Y^N)} C^N e^{-itP(Y^N)} B^N \right) \simeq \text{tr}_N(C^N) \times \text{tr}_N(B^N). \quad (1.7)$$

That the quantities on the right sides do not necessarily converge towards 0 comes from the fact that (C^N) may not behave asymptotically as a compact operator, indeed one could for example take (C^N) the identity matrices. Equations (1.6) and (1.7) indicate decays of correlations under the long-time Heisenberg evolution, a phenomenon often referred to as thermalization in [8]. More generally, the authors showed in Corollary 2.12 of [8], that the long-time Heisenberg evolution associated to a Wigner matrix generates asymptotic freeness.

The first link between Free Probability and Random Matrix Theory was made by Voiculescu in the nineties. In [36] he showed that the trace of any polynomial in GUE random matrices converges towards the trace of the same polynomial but evaluated in free semicircular operators. He introduced the notion of asymptotic freeness accordingly: A sequence of families of random matrices (X_1^N, \dots, X_d^N) is asymptotically free if for every collection of polynomials $P_1, \dots, P_n \in \mathbb{C}[X]$ such that $\text{tr}_N(P_j(X_{i_j}^N))$ converges towards 0, then if $i_j \neq i_{j+1}$ for every j , we have that

$$\lim_{N \rightarrow \infty} \text{tr}_N(P_1(X_{i_1}^N) \cdots P_n(X_{i_n}^N)) = 0.$$

In particular, if (A^N) and (B^N) are asymptotically free, then

$$\lim_{N \rightarrow \infty} \text{tr}_N \left((A^N - \text{tr}_N(A^N)) (B^N - \text{tr}_N(B^N)) \right) = 0.$$

Consequently, Equation (1.7) is a corollary of the asymptotic freeness of $e^{itP(Y^N)} C^N e^{-itP(Y^N)}$ and B^N .

Given two sequences of real numbers (u_N) and (v_N) , we write $u_N \ll v_N$ if u_N/v_N converges towards 0.

Theorem 1.2. *Let the collection of deterministic matrices $A^N = (A_1^N, \dots, A_q^N)$ and of Wigner matrices $Y^N = (Y_1^N, \dots, Y_d^N)$ be given as in Theorem 1.1. Besides we assume that for every i , A_i^N converges in distribution towards a non-commutative random variable a_i ; see Definition 2.5. Further, let $y_1^N \leq \dots \leq y_k^N \in \mathbb{R}$ be such that for any $i < k$, $1 \ll y_{i+1}^N - y_i^N \ll N^{1/4}$, and let P be a non-constant self-adjoint non-commutative polynomial in d variables. Then with*

$$a_i^N := e^{iy_i^N P(Y^N)} A_i^N e^{-iy_i^N P(Y^N)},$$

almost surely the family of non-commutative random variables (a_1^N, \dots, a_k^N) converges jointly in distribution towards (a_1, \dots, a_k) where the (a_i) are free. Moreover, if for every i , Y_i^N is a GUE or GOE random matrix, then one can replace $N^{1/4}$ by $N^{1/2}$ in the previous assumptions, i.e. one can assume that for any $i < k$, $1 \ll y_{i+1}^N - y_i^N \ll N^{1/2}$.

One can notably compare this theorem with Corollary 2.12 of [8]. The equivalents of Equations (1.4) and (1.5) are given in Theorem 2.6 of [8] (respectively Corollary 2.7 of [9]), there the error term depends on the k -th Sobolev norm (respectively $k/2$) of the functions (f_i) where k is the number of functions considered. In the case of the function $f_y : t \in \mathbb{R} \mapsto e^{iyt}$, the k -th Sobolev norm of f_y is of order y^k whereas if we use Theorem 1.1, we will get that $\|f_y\|_4$ is of order y^4 . While the former is better for smaller k , this dependency in k requires one to assume in Corollary 2.12 of [8] that $1 \ll y_{i+1}^N - y_i^N \ll N^{1/k}$ for any k , which can be improved to $N^{2/k}$ thanks to Remark 2.8 of [9]. This prevents the formulation of a result of convergence in distribution as usually defined in Free Probability; see Definition 2.5 below.

One can also wonder about the thermalization decay rates in Theorem 1.2. Indeed, given a polynomial P as defined in this theorem, the convergence rates depend on the behavior of the Fourier transform of the limiting spectral measure of P evaluated in free semicircular variables. For example, in the case of a single Wigner matrix, the spectral measure at the limit is the semicircle law which has square root decay at the spectral edges, hence this yields the exponent $\delta_P = 3/2$ in the equation below; as in Corollary 2.10 of [8]. For general P , thanks to Theorem 1.1(5-6) of [35], one can show that there exists a constant $\delta_P > 0$, such that for any polynomial Q , one has that for any $\varepsilon > 0$, with high probability,

$$\tau_N(Q(a_1^N, \dots, a_k^N)) = \tau(Q(a_1, \dots, a_k)) + \mathcal{O}\left(\frac{N^\varepsilon}{N} \max_{i \neq j} |y_i^N - y_j^N|^4 + \frac{1}{\min_{i \neq j} |y_i^N - y_j^N|^{\delta_P}}\right).$$

This estimate can be further improved depending on the polynomial Q , as in Corollary 2.11 of [8]. However it is not easy to compute the exponent δ_P associated to a given polynomial, thus we use the Riemann-Lebesgue lemma in the last step in the proof of Theorem 1.2. This is why we do not give the precise thermalization decay rates unlike Corollaries 2.9, 2.10 and 2.11 in [8]. However given a specific polynomial P , it is in principle possible to use the algorithm of Theorem 4.1 in [5] to compute the spectral distribution of a given polynomial in semicircular variables. This can be used in lieu of Riemann-Lebesgue.

We conclude this section by summarizing the organization of the paper. In Section 2, we first recall definitions from random matrix theory and free probability theory. Then in Subsection 2.3, we introduce the formalism suited to handle polynomials in non-commutative random variables and their derivatives. In Subsection 2.4, we recall the Schwinger-Dyson equations for Gaussian Wigner matrices as well as the cumulant expansions which can be viewed as a generalization of the Schwinger-Dyson equations for Wigner matrices. In Section 3, we derive key concentration estimates for the trace of products of exponentials of polynomials in Wigner matrices. This is accomplished by a continuous interpolation between Wigner matrices and GUE matrices in combination with a recursive moment estimate. In Section 4, we then combine the concentration estimates from Section 3 with the main results from [33] to prove our main results in Theorem 1.1 and Theorem 1.2.

2 Framework and standard properties

2.1 Definitions from Random Matrix Theory

In this subsection we define the main object of our study, the Wigner matrix. We make the assumption that the matrix entries are independent and have finite moments to every order, however we do not need to assume that they have the same law.

Definition 2.1. *We say that a square random matrix Y of size N is a Wigner matrix if it is a Hermitian or symmetric matrix whose entries are independent up to the Hermitian symmetry and such that for any i, j , $\mathbb{E}[Y_{i,j}] = 0$, and additionally if $i \neq j$, $\mathbb{E}[|Y_{i,j}|^2] = N^{-1}$. Besides we assume that for any $p \in \mathbb{N}$,*

$$\sup_{N \in \mathbb{N}, 1 \leq i, j \leq N} \mathbb{E} \left[\left| \sqrt{N} Y_{i,j} \right|^p \right] < \infty. \quad (2.1)$$

The assumption on the large moments being finite is natural as we are working with polynomials. Although since in Theorem 1.1 we give equations which hold true with high probability rather than estimates on the expectation, there might be a way around this assumption by using the truncation method (see Section 2 of [1]), but this would reflect in the error terms of Equations (1.4) and (1.5).

Besides, we use two specific types of Wigner matrices whose entries are all Gaussian, the Gaussian Unitary Ensemble (GUE) and the Gaussian Orthogonal Ensemble (GOE).

Definition 2.2. A GUE random matrix X^N of size N is a Hermitian matrix whose entries are random variables with the following laws:

- For $1 \leq i \leq N$, the random variables $\sqrt{N}X_{i,i}^N$ are independent centered Gaussian random variables of variance 1.
- For $1 \leq i < j \leq N$, the random variables $\sqrt{2N} \Re X_{i,j}^N$ and $\sqrt{2N} \Im X_{i,j}^N$ are independent centered Gaussian random variables of variance 1, independent of $(X_{i,i}^N)_i$.

Definition 2.3. A GOE random matrix X^N of size N is a symmetric matrix whose entries are random variables with the following laws:

- For $1 \leq i \leq N$, the random variables $\sqrt{\frac{N}{2}}X_{i,i}^N$ are independent centered Gaussian random variables of variance 1.
- For $1 \leq i < j \leq N$, the random variables $\sqrt{N}X_{i,j}^N$ are independent centered Gaussian random variables of variance 1, independent of $(X_{i,i}^N)_i$.

Finally we conclude this subsection by defining a notation that we will use regularly in the rest of the paper.

Definition 2.4. Given a sequence of events $(B_N)_{N \geq 1}$, we say that they happen with high probability if for any $k > 0$, we have that

$$\mathbb{P}(B_N) \geq 1 - N^{-k},$$

for N large enough.

2.2 Definitions from Free Probability Theory

In order to be self-contained, we begin by recalling the following definitions from free probability.

Definition 2.5. • A C^* -probability space $(\mathcal{A}, *, \tau, \|\cdot\|)$ is a unital C^* -algebra $(\mathcal{A}, *, \|\cdot\|)$ endowed with a state τ , i.e. a linear map $\tau : \mathcal{A} \rightarrow \mathbb{C}$ satisfying $\tau(1_{\mathcal{A}}) = 1$ and $\tau(a^*a) \geq 0$ for all $a \in \mathcal{A}$. In this paper we always assume that τ is a trace, i.e. that it satisfies $\tau(ab) = \tau(ba)$ for any $a, b \in \mathcal{A}$. An element of \mathcal{A} is called a non-commutative random variable. We will always work with a faithful trace, namely, for $a \in \mathcal{A}$, $\tau(a^*a) = 0$ if and only if $a = 0$.

- Let $\mathcal{A}_1, \dots, \mathcal{A}_n$ be $*$ -subalgebras of \mathcal{A} , having the same unit as \mathcal{A} . They are said to be free if for all k , for all $a_i \in \mathcal{A}_{j_i}$ such that $j_1 \neq j_2, j_2 \neq j_3, \dots, j_{k-1} \neq j_k$:

$$\tau\left((a_1 - \tau(a_1))(a_2 - \tau(a_2)) \cdots (a_k - \tau(a_k))\right) = 0. \quad (2.2)$$

Families of non-commutative random variables are said to be free if the $*$ -subalgebras they generate are free.

- Let $A = (a_1, \dots, a_k)$ be a k -tuple of non-commutative random variables. The joint distribution of the family A is the linear form $\mu_A : P \mapsto \tau[P(A, A^*)]$ on the set of polynomials in $2k$ non-commutative variables. By convergence in distribution, for a sequence of families of variables $(A_N)_{N \geq 1} = (a_1^N, \dots, a_k^N)_{N \geq 1}$ in C^* -algebras $(\mathcal{A}_N, *, \tau_N, \|\cdot\|)$, we mean the pointwise convergence of the map

$$\mu_{A_N} : P \mapsto \tau_N(P(A_N, A_N^*)).$$

- A family of non-commutative random variables $x = (x_1, \dots, x_d)$ is called a free semicircular system if the non-commutative random variables are free, self-adjoint ($x_i = x_i^*$), and for all $k \in \mathbb{N}$ and $i \in [1, d]$, one has

$$\tau(x_i^k) = \int_{\mathbb{R}} t^k d\sigma(t),$$

with $d\sigma(t) = \frac{1}{2\pi} \sqrt{4 - t^2} \mathbf{1}_{|t| \leq 2} dt$ the semicircle distribution.

It is important to note that thanks to [30, Theorem 7.9], which we recall next, one can consider free copies of any non-commutative random variable.

Theorem 2.6. *Let $(\mathcal{A}_i, \phi_i)_{i \in I}$ be a family of C^* -probability spaces such that the functionals $\phi_i : \mathcal{A}_i \rightarrow \mathbb{C}$, $i \in I$, are faithful traces. Then there exist a C^* -probability space (\mathcal{A}, ϕ) with ϕ a faithful trace, and a family of norm-preserving unital $*$ -homomorphism $W_i : \mathcal{A}_i \rightarrow \mathcal{A}$, $i \in I$, such that:*

- $\phi \circ W_i = \phi_i$, $\forall i \in I$.
- The unital C^* -subalgebras $W_i(\mathcal{A}_i)$, $i \in I$, form a free family in (\mathcal{A}, ϕ) .

We will usually denote \mathcal{A} by $\ast_{i \in I} \mathcal{A}_i$ or simply $\mathcal{A}_1 \ast \mathcal{A}_2$ when I only has two elements.

Let us finally fix a few notations concerning the spaces and traces that we use in this paper.

Definition 2.7. • $\mathbb{M}_N(\mathbb{C})$ is the set of matrices of size N with coefficients in \mathbb{C} .

- (\mathcal{A}_N, τ_N) is the free product $\mathbb{M}_N(\mathbb{C}) \ast \mathcal{C}_d$ of $\mathbb{M}_N(\mathbb{C})$ with \mathcal{C}_d the C^* -algebra generated by a free semicircular system $x = (x_1, \dots, x_d)$, that is the C^* -probability space built in Theorem 2.6. Note that when restricted to $\mathbb{M}_N(\mathbb{C})$, τ_N is just the regular normalized trace on matrices, in this case we will denote it by tr_N . The restriction of τ_N to \mathcal{C}_d is denoted as τ . Note that one can view this space as the limit of a matrix space, we refer to Proposition 3.5 from [11].
- Tr_N is the non-normalized trace on $\mathbb{M}_N(\mathbb{C})$, while tr_N is the normalized one.
- We denote $E_{r,s} \in \mathbb{M}_N(\mathbb{C})$ the matrix with entries equal to 0 except in (r, s) where it is equal to 1.

In order to interpret Theorem 1.1, we need to define the conditional expectation first.

Definition 2.8. *Let \mathcal{A} be a C^* -algebra and \mathcal{B} be a unital C^* -algebra, then $E : \mathcal{A} \rightarrow \mathcal{B}$ is said to be a conditional expectation if it is a linear map such that:*

- E is positive, i.e. for every $a \in \mathcal{A}$, there exist $b \in \mathcal{B}$ such that $E[a^*a] = b^*b$.
- E is a $\mathcal{B} - \mathcal{B}$ -bimodule map, i.e. for every $a \in \mathcal{A}$, $b_1, b_2 \in \mathcal{B}$, one has $E[b_1 a b_2] = b_1 E[a] b_2$.
- For every $b \in \mathcal{B}$, $E[b] = b$.

It is well-known that we have the following property.

Proposition 2.9. *Let (\mathcal{A}, τ) be a W^* -algebra, i.e. a von Neumann algebra with a faithful normal trace τ , and \mathcal{B} a W^* -subalgebra. Then there always exists a conditional expectation $E : \mathcal{A} \rightarrow \mathcal{B}$. Besides given $a \in \mathcal{A}$, it is characterized by the fact that for any $b \in \mathcal{B}$, $\tau(E[a]b) = \tau(ab)$. Finally, for every element $a \in \mathcal{A}$, we have that $\|E[a]\| \leq \|a\|$, where $\|\cdot\|$ is the operator norm.*

With those definitions and properties, one can deduce the following which gives us some understanding on Equation (1.5).

Remark 2.10. • While the C^* -algebra \mathcal{A}_N built in Definition 2.7 is not necessarily a von Neumann algebra, one can always replace it by its enveloping von Neumann algebra. Hence since $\mathbb{M}_N(\mathbb{C})$ is a von Neumann algebra, there exist a conditional expectation $E_{\mathbb{M}_N(\mathbb{C})}$ from \mathcal{A}_N to $\mathbb{M}_N(\mathbb{C})$. That being said, it is not really necessary to understand the methods used to define the conditional expectation to read this paper. The rest of this remark will provide more details on how to estimate this quantity most of the time.

- Thanks to Proposition 2.9 we can deduce several properties of the conditional expectation in Equation (1.5), first and foremost one has the following bound

$$\begin{aligned} & \left| \langle \mathbf{x}, E_{\mathbb{M}_N(\mathbb{C})} [f_1(P_1(x, A^N)) \cdots f_k(P_k(x, A^N))] \mathbf{y} \rangle \right| \\ & \leq \|E_{\mathbb{M}_N(\mathbb{C})} [f_1(P_1(x, A^N)) \cdots f_k(P_k(x, A^N))]\| \|\mathbf{x}\|_2 \|\mathbf{y}\|_2 \\ & \leq \|f_1(P_1(x, A^N))\| \cdots \|f_k(P_k(x, A^N))\| \|\mathbf{x}\|_2 \|\mathbf{y}\|_2. \end{aligned}$$

Besides if $f_i = \text{id}_{\mathbb{R}}$, then $\|f_i(P_i(x, A^N))\| = \|P_i(x, A^N)\|$, and otherwise

$$\|f_i(P_i(x, A^N))\| \leq \sup_{x \in \mathbb{R}} |f(x)| \leq \int_{\mathbb{R}} 1 \, d|\mu_i| \leq \|f_i\|_4.$$

Consequently as long as our functions do not depend on N , the first order term in Equation (1.5) is also independent of N except with respect to the norm of the matrices A^N and the one of the vectors \mathbf{x}, \mathbf{y} which is to be expected.

- It is also worth noting that since

$$\begin{aligned} & \langle \mathbf{x}, E_{\mathbb{M}_N(\mathbb{C})} [f_1(P_1(x, A^N)) \cdots f_k(P_k(x, A^N))] \mathbf{y} \rangle \\ &= \text{Tr}_N (E_{\mathbb{M}_N(\mathbb{C})} [f_1(P_1(x, A^N)) \cdots f_k(P_k(x, A^N))] \mathbf{y} \mathbf{x}^*) \\ &= N \tau_N (f_1(P_1(x, A^N)) \cdots f_k(P_k(x, A^N)) \mathbf{y} \mathbf{x}^*), \end{aligned}$$

it is not necessary to compute the conditional expectation of $f_1(P_1(x, A^N)) \cdots f_k(P_k(x, A^N))$ to determine the first order term of Equation (1.5). However, the formulation with the conditional expectation renders it clear that this term is of order one.

- It is possible to use advanced combinatorics associated with free probability to compute the leading terms on the right of (1.4) and (1.5), especially when for every i such that $f_i \neq \text{id}_{\mathbb{R}}$, P_i is a polynomial in x rather than (x, A^N) ; see Section 2 of [8]. For example with s a single semicircular variable, we have that

$$\tau_N (f_1(s) A_1^N \cdots f_k(s) A_k^N) = \sum_{\pi \in \text{NCP}[k]} \langle A_1^N, \dots, A_k^N \rangle_{K(\pi)} \prod_{B \in \pi} \text{sc}_o[B],$$

where $\text{NCP}[k]$ is the set of non-crossing partitions of $[[1, k]]$, $\langle A_1^N, \dots, A_k^N \rangle_{K(\pi)}$ is a quantity that depends only on the matrices A_1^N, \dots, A_k^N and the Kreweras complement $K(\pi)$ of π , whereas the free cumulant function $\text{sc}_o[\cdot]$ depends on the functions f_1, \dots, f_k and the semicircular distribution. We refer to [30] for an introduction to the combinatorics of free probability.

- If one does not have any matrices A^N , then since $f_1(P_1(x)) \cdots f_k(P_k(x))$ are free from $\mathbb{M}_N(\mathbb{C})$, we have that for any $b \in \mathbb{M}_N(\mathbb{C})$,

$$\tau_N (f_1(P_1(x)) \cdots f_k(P_k(x)) b) = \tau_N (b) \times \tau (f_1(P_1(x)) \cdots f_k(P_k(x))).$$

Hence, $E_{\mathbb{M}_N(\mathbb{C})} [f_1(P_1(x)) \cdots f_k(P_k(x))] = \tau (f_1(P_1(x)) \cdots f_k(P_k(x)))$ and

$$\langle \mathbf{x}, E_{\mathbb{M}_N(\mathbb{C})} [f_1(P_1(x)) \cdots f_k(P_k(x))] \mathbf{y} \rangle = \langle \mathbf{x}, \mathbf{y} \rangle \times \tau (f_1(P_1(x)) \cdots f_k(P_k(x))).$$

2.3 Noncommutative polynomials and derivatives

Let $\mathcal{A}_{d,2r} = \mathbb{C}\langle X_1, \dots, X_d, Z_1, \dots, Z_{2r} \rangle$ be the set of non-commutative polynomials in $d+2r$ variables. We set $q = 2r$ to simplify notations. We denote by $\deg M$ the total degree of M (that is the sum of its degree in each letter $X_1, \dots, X_d, Z_1, \dots, Z_{2r}$). Let us now define several maps that are frequently used in Operator Algebra. First, for $A, B, C \in \mathcal{A}_{d,q}$, let

$$A \otimes B \# C := ACB, \quad A \otimes B \tilde{\#} C := BCA, \quad m(A \otimes B) := BA. \quad (2.3)$$

We define an involution $*$ on $\mathcal{A}_{d,q}$ by $X_i^* = X_i$, $Z_i^* = Z_{i+r}$ if $1 \leq i \leq r$, $Z_i^* = Z_{i-r}$ else, and then we extend it to $\mathcal{A}_{d,q}$ by linearity and the formula $(\alpha PQ)^* = \bar{\alpha} Q^* P^*$. $P \in \mathcal{A}_{d,q}$ is said to be self-adjoint if $P^* = P$. Self-adjoint polynomials have the property that if $x_1, \dots, x_d, z_1, \dots, z_r$ are elements of a \mathcal{C}^* -algebra such that x_1, \dots, x_d are self-adjoint, then so is $P(x_1, \dots, x_d, z_1, \dots, z_r, z_1^*, \dots, z_r^*)$.

Definition 2.11. If $1 \leq i \leq d$, one defines $\partial_i : \mathcal{A}_{d,q} \longrightarrow \mathcal{A}_{d,q} \otimes \mathcal{A}_{d,q}$ the non-commutative derivative with respect to X_i as such. First on a monomial $M \in \mathcal{A}_{d,q}$, one sets

$$\partial_i M = \sum_{A, B \in \mathcal{A}_{d,q} \text{ such that } M = AX_i B} A \otimes B,$$

and then extend it by linearity to all polynomials. We can also define ∂_i by induction with the formulas,

$$\begin{aligned} \forall P, Q \in \mathcal{A}_{d,q}, \quad \partial_i(PQ) &= (\partial_i P)(1 \otimes Q) + (P \otimes 1)(\partial_i Q), \\ \forall i, j, \quad \partial_i X_j &= \delta_{i,j} 1 \otimes 1, \quad \partial_i Z_j = 0. \end{aligned} \tag{2.4}$$

Similarly, with m as in (2.3), one defines the cyclic derivative $D_i : \mathcal{A}_{d,q} \longrightarrow \mathcal{A}_{d,q}$ for $P \in \mathcal{A}_{d,q}$ by

$$D_i P = m \circ \partial_i P.$$

In this paper we need to work not only with polynomials but also with more general functions, since we will work with the Fourier transform we introduce the following space.

Definition 2.12. We set $\mathcal{S} = \{R \in \mathcal{A}_{d,q} \mid R^* = R\}$, then we denote

$$\mathcal{F}_{d,q} = \mathbb{C}\langle (E_R)_{R \in \mathcal{S}}, X_1, \dots, X_d, Z_1, \dots, Z_{2r} \rangle,$$

the set of polynomials in $X_1, \dots, X_d, Z_1, \dots, Z_{2r}$ as well as in a family of variable indexed by \mathcal{S} . Then given $y = (x_1, \dots, x_d, z_1, \dots, z_r, z_1^*, \dots, z_r^*)$ elements of a \mathcal{C}^* -algebra, one can define by induction the evaluation of an element of $\mathcal{F}_{d,q}$ in y by following the following rules:

- $\forall Q \in \mathcal{A}_{d,q}$, $Q(y)$ is defined as usual,
- $\forall Q_1, Q_2 \in \mathcal{F}_{d,q}$, $(Q_1 + Q_2)(y) = Q_1(y) + Q_2(y)$, $(Q_1 Q_2)(y) = Q_1(y) Q_2(y)$,
- $\forall R \in \mathcal{S}$, $E_R(y) = e^{iR(y)}$.

One can extend the involution $*$ from $\mathcal{A}_{d,q}$ to $\mathcal{F}_{d,q}$ by setting $(E_R)^* = E_{(-R)}$, and then again we have that if $Q \in \mathcal{F}_{d,q}$ is self-adjoint, then so is $Q(y)$. Finally in order to make notations more transparent, we will usually write e^{iR} instead of E_R .

Note that for technical reasons which are explained in Remark 2.10 of [33], one cannot view $\mathcal{F}_{d,q}$ as a subalgebra of the set of formal power series in $X_1, \dots, X_d, Z_1, \dots, Z_{2r}$. Hence why we need to introduce the notation E_R .

Now, as we will see in Proposition 2.15, a natural way to extend the definition of ∂_i (and D_i) to $\mathcal{F}_{d,q}$ is by setting

$$\partial_i e^{iR} = \mathbf{i} \int_0^1 (e^{i\alpha R} \otimes 1) \partial_i R (1 \otimes e^{i(1-\alpha)R}) d\alpha. \tag{2.5}$$

However we cannot define the integral properly on $\mathcal{F}_{d,q} \otimes \mathcal{F}_{d,q}$, but after evaluating our polynomials in a matrix space this is not a problem anymore. Indeed a tensor of matrix spaces is also a matrix space and hence the integral of continuous functions on this space is well-defined. Thus we define the non-commutative differential on $\mathcal{F}_{d,q}$ as following.

Definition 2.13. For $\alpha \in [0, 1]$, we define $\partial_{\alpha,i} : \mathcal{F}_{d,q} \rightarrow \mathcal{F}_{d,q} \otimes \mathcal{F}_{d,q}$ as the map which satisfies (2.4) and is such that for any $R \in \mathcal{A}_{d,q}$ self-adjoint,

$$\partial_{\alpha,i} e^{iR} = \mathbf{i} (e^{i\alpha R} \otimes 1) \partial_i R (1 \otimes e^{i(1-\alpha)R}), \quad D_{\alpha,i} = m \circ \partial_{\alpha,i}.$$

Then, given $y = (y_1, \dots, y_{d+q})$ elements of $M_N(\mathbb{C})$, we define for any $Q \in \mathcal{F}_{d,q}$,

$$\partial_i Q(y) = \int_0^1 \partial_{\alpha,i} Q(y) d\alpha, \quad D_i Q(y) = \int_0^1 D_{\alpha,i} Q(y) d\alpha.$$

Note that for any $P \in \mathcal{A}_{d,q}$, since $\int_0^1 1d\alpha = 1$, we do also have with $\partial_i Q$ defined as in Definition 2.11, that

$$\partial_i Q(y) = \int_0^1 \partial_{\alpha,i} Q(y) d\alpha.$$

Thus Definition 2.13 extends indeed the definition of ∂_i from $\mathcal{A}_{d,q}$ to $\mathcal{F}_{d,q}$. Besides, it also means that we can define rigorously the composition of those maps. Since the map $\partial_{\alpha,i}$ goes from $\mathcal{F}_{d,q}$ to $\mathcal{F}_{d,q} \otimes \mathcal{F}_{d,q}$ it is very easy to do so. For example one defines the following operator that we use later on.

Definition 2.14. Let $Q \in \mathcal{F}_{d,q}$, given $y = (y_1, \dots, y_{d+q})$ elements of a C^* -algebra, let $i \in [1, d]$, with \circ the composition of operator we define

$$\partial_i \circ D_i Q(y) = \int_{[0,1]^2} \partial_{\alpha_2,i} \circ D_{\alpha_1,i} Q(y) d\alpha_1 d\alpha_2.$$

2.4 The Schwinger-Dyson equations and their generalization

A tool that we use repeatedly with Gaussian random matrices are the so-called Schwinger-Dyson equations. It is a consequence of Gaussian integration by parts which can be summarized into the following formula. If Z is a centered Gaussian random variable with variance one and f a C^1 function, then

$$\mathbb{E}[Zf(Z)] = \mathbb{E}[\partial_Z f(Z)]. \quad (2.6)$$

From there on, we deduce the Schwinger-Dyson equations in the following proposition. For more information about these equations and their applications, we refer to [3], Lemma 5.4.7.

Proposition 2.15. Let X^N be a GOE random matrix of size N , A^N deterministic matrices, $Q \in \mathcal{F}_{1,q}$, then

$$\mathbb{E}\left[\mathrm{tr}_N(X^N Q(X^N, A^N))\right] = \mathbb{E}\left[\mathrm{tr}_N^{\otimes 2}(\partial_1 Q(X^N, A^N))\right] + \frac{1}{N} \mathbb{E}\left[\mathrm{tr}_N(h(\partial_1 Q(X^N, A^N)))\right], \quad (2.7)$$

where h is the linear map such that $h(A \otimes B) = A^T B$ with A^T is the transpose of A .

Proof. Let us first assume that $Q \in \mathcal{A}_{1,q}$. One can write $X^N = \frac{1}{\sqrt{N}}(x_{r,s})_{1 \leq r,s \leq N}$ and thus

$$\begin{aligned} \mathbb{E}\left[\mathrm{tr}_N(X^N Q(X^N))\right] &= \frac{1}{N^{3/2}} \sum_{r,s} \mathbb{E}\left[x_{r,s} \mathrm{Tr}_N(E_{r,s} Q(X^N, A^N))\right] \\ &= \frac{1}{N^{3/2}} \sum_{r,s} \mathbb{E}\left[\mathrm{Tr}_N(E_{r,s} \partial_{x_{r,s}} Q(X^N, A^N))\right] \\ &= \frac{1}{N^2} \sum_{r,s} \mathbb{E}\left[\mathrm{Tr}_N(E_{r,s} \partial_1 Q(X^N, A^N) \# (E_{s,r} + E_{r,s}))\right] \\ &= \mathbb{E}\left[\mathrm{tr}_N^{\otimes 2}(\partial_1 Q(X^N, A^N))\right] + \frac{1}{N} \mathbb{E}\left[\mathrm{tr}_N(h(\partial_1 Q(X^N, A^N)))\right], \end{aligned}$$

where notably we used that for any matrices $A, B \in \mathbb{M}_N(\mathbb{C})$,

$$\sum_{1 \leq i,j \leq N} A_{i,j} B_{i,j} = \mathrm{Tr}_N(A^T B).$$

If $Q \in \mathcal{F}_{d,q}$, then the proof is pretty much the same but we need to use Duhamel's formula (for a very similar proof see [34], Proposition 2.2) which states that for any matrices A and B ,

$$e^B - e^A = \int_0^1 e^{\alpha B} (B - A) e^{(1-\alpha)A} d\alpha. \quad (2.8)$$

Thus this allows us to prove that for any self-adjoint polynomials $P \in \mathcal{A}_{d,q}$,

$$\partial_{x_{r,s}} e^{iP(X^N)} = i \int_0^1 e^{i\alpha P(X^N)} \partial_1 P(X^N) \# E_{s,r} e^{i(1-\alpha)P(X^N)} d\alpha,$$

and the conclusion follows. \square

In the case of GUE random matrices we have an even shorter formula. The following Proposition is actually Proposition 2.23 of [33] whose proof is quite similar to the one of Proposition 2.15.

Proposition 2.16. *Let X^N be a GUE random matrix of size N , A^N deterministic matrices, $Q \in \mathcal{F}_{1,q}$, then*

$$\mathbb{E} \left[\text{tr}_N (X^N Q(X^N, A^N)) \right] = \mathbb{E} \left[\text{tr}_N^{\otimes 2} (\partial_1 Q(X^N, A^N)) \right]. \quad (2.9)$$

For studying Wigner matrices, there exist a more general formula, the cumulant expansion. Its usefulness in random matrix theory was recognized in [23] and has widely been used since, *e.g.* [7, 16, 20, 21, 22, 27, 32]. We will use a specific version with an explicit expression for the remainder. To do so we follow the proof of Proposition 3.1 of [32] but we do not upper bound the remainder.

Proposition 2.17. *Let u, v be real random variables such that $\mathbb{E}[|u|^{p+2}] < \infty$ and $\mathbb{E}[|v|^{p+2}] < \infty$ for some integer p . Let $(\kappa_{n,m})_{n,m \in \mathbb{N}}$ be the joint cumulants of u and v , *i.e.* the numbers that satisfy*

$$\log \mathbb{E} [e^{itu+isv}] = \sum_{0 \leq n+m \leq p} \frac{\kappa_{n,m}}{n!m!} (\mathbf{it})^n (\mathbf{is})^m + o(\max\{|t|, |s|\}^p).$$

Then for any function $\Phi : \mathbb{R}^2 \rightarrow \mathbb{C}$ of the class \mathcal{C}^{p+1} , we have that

$$\mathbb{E} [u\Phi(u, v)] = \sum_{0 \leq a+b \leq p} \frac{\kappa_{a+1,b}}{a!b!} \times \mathbb{E} [\partial_u^a \partial_v^b \Phi(u, v)] + \varepsilon_{p+1}, \quad (2.10)$$

with

$$\varepsilon_{p+1} = \mathbb{E} [u g_{0,0}(u, v)] - \sum_{0 \leq a+b \leq p} \frac{\kappa_{a+1,b}}{a!b!} \times \mathbb{E} [g_{a,b}(u, v)],$$

where we defined

$$g_{a,b} : (X_1, X_2) \in \mathbb{R}^2 \mapsto \frac{1}{(p-a-b)!} \sum_{\varepsilon \in \{1,2\}^{p+1-a-b}} X_{\varepsilon_1} \cdots X_{\varepsilon_{p+1-a-b}} \int_0^1 \partial_{\varepsilon_1} \cdots \partial_{\varepsilon_{p+1-a-b}} \partial_1^a \partial_2^b \Phi(tX_1, tX_2) (1-t)^{p-a-b} dt.$$

Proof. Since we have that

$$\mathbb{E} [ue^{itu+isv}] = \mathbb{E} [e^{itu+isv}] \times (-\mathbf{i}\partial_t) (\log \mathbb{E} [e^{itu+isv}]),$$

then with $\mu_{n,m} = \mathbb{E}[u^n v^m]$, this implies that for $n+m \leq p$,

$$\mu_{n+1,m} = \sum_{\substack{0 \leq a \leq n \\ 0 \leq b \leq m}} \frac{\kappa_{a+1,b}}{a!b!} \times \frac{n!m!}{(n-a)!(m-b)!} \mu_{n-a,m-b}.$$

Consequently for any polynomial P of degree smaller than p , we have that

$$\begin{aligned} \mathbb{E}[uP(u, v)] &= \sum_{\substack{0 \leq a < \infty \\ 0 \leq b < \infty}} \frac{\kappa_{a+1,b}}{a!b!} \times \mathbb{E} [\partial_u^a \partial_v^b P(u, v)] \\ &= \sum_{0 \leq a+b \leq p} \frac{\kappa_{a+1,b}}{a!b!} \times \mathbb{E} [\partial_u^a \partial_v^b P(u, v)]. \end{aligned}$$

For $X_1, X_2 \in \mathbb{R}$, let $f : t \in [0, 1] \mapsto \Phi(tX_1, tX_2)$, then thanks to Taylor's theorem, we have that

$$f(1) = \sum_{j=0}^p \frac{f^{(j)}(0)}{j!} + \frac{1}{p!} \int_0^1 f^{(p+1)}(t)(1-t)^p dt.$$

Consequently there exists a polynomial π_p of degree at most p such that

$$\Phi(X_1, X_2) = \pi_p(X_1, X_2) + \frac{1}{p!} \sum_{\varepsilon \in \{1,2\}^{p+1}} X_{\varepsilon_1} \cdots X_{\varepsilon_{p+1}} \int_0^1 \partial_{\varepsilon_1} \cdots \partial_{\varepsilon_{p+1}} \Phi(tX_1, tX_2) (1-t)^p dt \quad (2.11)$$

$$= \pi_p(X_1, X_2) + g_{0,0}(X_1, X_2).$$

Thus we get that

$$\begin{aligned} \mathbb{E}[u\Phi(u, v)] &= \mathbb{E}[u\pi_p(u, v)] + \mathbb{E}[ug_{0,0}(u, v)] \\ &= \sum_{0 \leq a+b \leq p} \frac{\kappa_{a+1,b}}{a!b!} \times \mathbb{E}[\partial_u^a \partial_v^b \pi_p(u, v)] + \mathbb{E}[ug_{0,0}(u, v)]. \end{aligned}$$

But thanks to Equation (2.11), we have that

$$\begin{aligned} &\partial_1(\Phi(X_1, X_2) - \pi_p(X_1, X_2)) \\ &= \frac{1}{p!} \sum_{\substack{\varepsilon \in \{1,2\}^{p+1} \\ j \in [1,p+1]}} X_{\varepsilon_1} \cdots X_{\varepsilon_{j-1}} \partial_1(X_{\varepsilon_j}) X_{\varepsilon_{j+1}} \cdots X_{\varepsilon_{p+1}} \int_0^1 \partial_{\varepsilon_1} \cdots \partial_{\varepsilon_{p+1}} \Phi(tX_1, tX_2) (1-t)^p dt \\ &\quad + \frac{1}{p!} \sum_{\varepsilon \in \{1,2\}^{p+1}} X_{\varepsilon_1} \cdots X_{\varepsilon_{p+1}} \int_0^1 \partial_{\varepsilon_1} \cdots \partial_{\varepsilon_{p+1}} \partial_1 \Phi(tX_1, tX_2) \times t(1-t)^p dt \\ &= \frac{1}{p!} \sum_{\varepsilon \in \{1,2\}^p} X_{\varepsilon_1} \cdots X_{\varepsilon_p} \int_0^1 \partial_{\varepsilon_1} \cdots \partial_{\varepsilon_p} \partial_1 \Phi(tX_1, tX_2) (1-t)^p dt \times (p+1) \\ &\quad + \frac{1}{p!} \sum_{\varepsilon \in \{1,2\}^p} X_{\varepsilon_1} \cdots X_{\varepsilon_p} \int_0^1 \frac{d}{dt} (\partial_{\varepsilon_1} \cdots \partial_{\varepsilon_p} \partial_1 \Phi(tX_1, tX_2)) \times t(1-t)^p dt \\ &= \frac{1}{p!} \sum_{\varepsilon \in \{1,2\}^p} X_{\varepsilon_1} \cdots X_{\varepsilon_p} \int_0^1 \partial_{\varepsilon_1} \cdots \partial_{\varepsilon_p} \partial_1 \Phi(tX_1, tX_2) \times (1-t)^{p-1} \times ((1-t)(p+1) - (1-t) + pt) dt \\ &= \frac{1}{(p-1)!} \sum_{\varepsilon \in \{1,2\}^p} X_{\varepsilon_1} \cdots X_{\varepsilon_p} \int_0^1 \partial_{\varepsilon_1} \cdots \partial_{\varepsilon_p} \partial_1 \Phi(tX_1, tX_2) \times (1-t)^{p-1} dt. \end{aligned}$$

Hence by induction we get that $\partial_1^a \partial_2^b (\Phi - \pi_p) = g_{a,b}$. Note that we used that Φ was of class \mathcal{C}^{p+2} in this computation but by an argument of density we only need Φ to be of class \mathcal{C}^{p+1} eventually. Thus in conclusion,

$$\mathbb{E}[u\Phi(u, v)] = \sum_{0 \leq a+b \leq p} \frac{\kappa_{a+1,b}}{a!b!} \times \mathbb{E}[\partial_u^a \partial_v^b \Phi(u, v)] + \mathbb{E}[ug_{0,0}(u, v)] - \sum_{0 \leq a+b \leq p} \frac{\kappa_{a+1,b}}{a!b!} \times \mathbb{E}[g_{a,b}(u, v)].$$

This proves Equation (2.10). □

3 Reduction of the problem to the case of GUE matrices

3.1 Concentration of the trace

In order to prove Theorem 1.1, the first step is to reduce the proof to the case of GUE random matrices. To do so we prove two concentration results. First for the trace in Proposition 3.1, then for the scalar product in Proposition 3.3.

Proposition 3.1. *Given*

- Y^N a d -tuple of independent Wigner matrices as in Definition 2.1,
- X^N a d -tuple of independent GUE random matrices, independent from Y^N ,
- A^N deterministic matrices such that $\sup_{1 \leq i \leq q, N \in \mathbb{N}^*} \|A_i^N\| < \infty$.

Then with $y_i \in \mathbb{R}$, and $y = 1 + \max_i |y_i|$, P_1, \dots, P_k non-commutative self-adjoint polynomials, R_1, \dots, R_k non-commutative polynomials, if $Q = e^{iP_1 y_1} R_1 \dots e^{iP_k y_k} R_k$, we have for any $\varepsilon > 0$ that with high probability,

$$\mathrm{tr}_N(Q(Y^N, A^N)) = \mathbb{E}[\mathrm{tr}_N(Q(X^N, A^N))] + \mathcal{O}\left(N^\varepsilon \frac{y^4}{N}\right). \quad (3.1)$$

Besides if Y^N a d -tuple of independent GOE or GUE matrices, then one can replace y^4 by y^2 in the previous equality.

Remark 3.2. We suspect that it should be possible to improve $\|\cdot\|_4$ to $\|\cdot\|_2$ in the previous theorem but at the moment we can only do so for GOE and GUE matrices. This comes from certain inequalities that one could maybe improve by pushing the cumulant expansion further and proceeding by induction; see e.g. how we upper bound the term (3.18) or inequality (3.17).

Proof. Our first step is to define

$$M_N := \mathrm{tr}_N(Q(Y^N, A^N) - \mathbb{E}[Q(X^N, A^N)]).$$

Then we want to study the moment of M_N in order to use Markov inequality. To do so we proceed by induction, we are going to prove that for any $n \in \mathbb{N}$, for any $\varepsilon > 0$, there exists a constant $C_{n,\varepsilon}$ such that

$$\mathbb{E}[|M_N|^n] \leq C_{n,\varepsilon} \left(\frac{y^4}{N^{1-\varepsilon}}\right)^n. \quad (3.2)$$

We show at the end of the proof that this inequality is true for $n = 1, 2, 3$. Let us now pick $n \geq 4$ and assume that Equation (3.2) is true for moments smaller than n . Then we set for $t \in \mathbb{R}^+$,

$$X_t^N = (Y^N e^{-t/2} + X^N (1 - e^{-t})^{1/2}, A^N),$$

so that we have the following equality,

$$\begin{aligned} & \frac{d}{dt} \mathbb{E} \left[\mathrm{tr}_N(Q(X_t^N)) \times M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] \\ &= -\frac{e^{-t}}{2} \sum_{1 \leq s \leq d} \mathbb{E} \left[\mathrm{tr}_N \left(D_s Q(X_t^N) \left(e^{t/2} Y_s^N - \frac{1}{(1 - e^{-t})^{1/2}} X_s^N \right) \right) \times M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] \\ &= -\frac{e^{-t}}{2} \sum_{1 \leq s \leq d} \mathbb{E} \left[\left(e^{t/2} \mathrm{tr}_N(D_s Q(X_t^N) Y_s^N) - \mathrm{tr}_N^{\otimes 2}(\partial_s D_s Q(X_t^N)) \right) \times M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right], \end{aligned}$$

where we used the Schwinger-Dyson equations, i.e. Proposition 2.16, to get the last line. Since

$$\begin{aligned} \mathbb{E} \left[\mathrm{tr}_N(Q(X_0^N)) \times M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] &= \mathbb{E} \left[\mathrm{tr}_N(Q(Y^N)) \times M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right], \\ \lim_{t \rightarrow \infty} \mathbb{E} \left[\mathrm{tr}_N(Q(X_t^N)) \times M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] &= \mathbb{E} \left[\mathrm{tr}_N(Q(X^N)) \right] \times \mathbb{E} \left[M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right], \end{aligned}$$

we have that

$$\mathbb{E}[|M_N|^n] = \frac{1}{2} \sum_{1 \leq s \leq d} \int_0^\infty e^{-t} \mathbb{E} \left[\left(e^{t/2} \mathrm{tr}_N(D_s Q(X_t^N) Y_s^N) - \mathrm{tr}_N^{\otimes 2}(\partial_s D_s Q(X_t^N)) \right) \times M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] dt. \quad (3.3)$$

Next we want to use the cumulant expansion, that is Proposition 2.17 with $u = \Re((Y_s^N)_{i,j})$, $v = \Im((Y_s^N)_{i,j})$ and

$$\Phi : (\Re((Y_s^N)_{i,j}), \Im((Y_s^N)_{i,j})) \mapsto e^{t/2} \mathrm{tr}_N(D_s Q(X_t^N) E_{i,j}) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}}.$$

In particular if $i \neq j$, we have that

$$\begin{aligned} \partial_u \Phi(u, v) &= \text{tr}_N \left((\partial_s D_s Q(X_t^N) \# (E_{i,j} + E_{j,i})) E_{i,j} \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \\ &\quad + \left(\frac{n}{2} - 1 \right) \text{tr}_N \left(D_s Q(X_t^N) E_{i,j} \right) \text{tr}_N \left(D_s Q(X_0^N) (E_{i,j} + E_{j,i}) \right) M_N^{\frac{n}{2}-2} \overline{M_N^{\frac{n}{2}}} \\ &\quad + \frac{n}{2} \text{tr}_N \left(D_s Q(X_t^N) E_{i,j} \right) \text{tr}_N \left(D_s Q^*(X_0^N) (E_{i,j} + E_{j,i}) \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}-1}}, \end{aligned} \quad (3.4)$$

and

$$\begin{aligned} \partial_v \Phi(u, v) &= \mathbf{i} \text{tr}_N \left((\partial_s D_s Q(X_t^N) \# (E_{i,j} - E_{j,i})) E_{i,j} \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \\ &\quad + \mathbf{i} \left(\frac{n}{2} - 1 \right) \text{tr}_N \left(D_s Q(X_t^N) E_{i,j} \right) \text{tr}_N \left(D_s Q(X_0^N) (E_{i,j} - E_{j,i}) \right) M_N^{\frac{n}{2}-2} \overline{M_N^{\frac{n}{2}}} \\ &\quad + \mathbf{i} \frac{n}{2} \text{tr}_N \left(D_s Q(X_t^N) E_{i,j} \right) \text{tr}_N \left(D_s Q^*(X_0^N) (E_{i,j} - E_{j,i}) \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}-1}}. \end{aligned} \quad (3.5)$$

Further if $i = j$, then $\partial_u \Phi(u, v)$ is defined similarly but with $E_{i,i}$ instead of $E_{i,j} + E_{j,i}$, and $\partial_v \Phi(u, v) = 0$. Besides with u and v defined as such, their cumulants satisfy

$$\begin{aligned} \kappa_{1,0} &= \frac{1}{2} \Re \mathbb{E}[(Y_s^N)_{i,j}] = 0, \\ \kappa_{0,1} &= \frac{1}{2\mathbf{i}} \Im \mathbb{E}[(Y_s^N)_{i,j}] = 0, \\ \kappa_{2,0} + \kappa_{0,2} &= \mathbb{E}[|(Y_s^N)_{i,j}|^2], \\ \kappa_{2,0} - \kappa_{0,2} + 2\mathbf{i}\kappa_{1,1} &= \mathbb{E}[(Y_s^N)_{i,j}^2]. \end{aligned} \quad (3.6)$$

In particular, as we set in Definition 2.1, if $i \neq j$, $\kappa_{2,0} + \kappa_{0,2} = 1/N$. Thus by using Proposition 2.17 with $p = 2$, we get that

$$\begin{aligned} &e^{t/2} \mathbb{E} \left[(Y_s^N)_{i,j} \times \text{tr}_N \left(D_s Q(X_t^N) E_{i,j} \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] \\ &= \frac{1}{N} \mathbb{E} \left[\text{tr}_N \left((\partial_s D_s Q(X_t^N) \# E_{j,i}) E_{i,j} \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] \\ &\quad + \frac{1}{N} \left(\frac{n}{2} - 1 \right) \mathbb{E} \left[\text{tr}_N \left(D_s Q(X_t^N) E_{i,j} \right) \text{tr}_N \left(D_s Q(X_0^N) E_{j,i} \right) M_N^{\frac{n}{2}-2} \overline{M_N^{\frac{n}{2}}} \right] \\ &\quad + \frac{n}{2N} \mathbb{E} \left[\text{tr}_N \left(D_s Q(X_t^N) E_{i,j} \right) \text{tr}_N \left(D_s Q^*(X_0^N) E_{j,i} \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}-1}} \right] \\ &\quad + \left(\mathbb{E}[(Y_s^N)_{i,j}^2] - \frac{\mathbf{1}_{i=j}}{N} \right) \mathbb{E} \left[\text{tr}_N \left((\partial_s D_s Q(X_t^N) \# E_{i,j}) E_{i,j} \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] \\ &\quad + \left(\frac{n}{2} - 1 \right) \left(\mathbb{E}[(Y_s^N)_{i,j}^2] - \frac{\mathbf{1}_{i=j}}{N} \right) \mathbb{E} \left[\text{tr}_N \left(D_s Q(X_t^N) E_{i,j} \right) \text{tr}_N \left(D_s Q(X_0^N) E_{i,j} \right) M_N^{\frac{n}{2}-2} \overline{M_N^{\frac{n}{2}}} \right] \\ &\quad + \frac{n}{2} \left(\mathbb{E}[(Y_s^N)_{i,j}^2] - \frac{\mathbf{1}_{i=j}}{N} \right) \mathbb{E} \left[\text{tr}_N \left(D_s Q(X_t^N) E_{i,j} \right) \text{tr}_N \left(D_s Q^*(X_0^N) E_{i,j} \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}-1}} \right] \\ &\quad + \frac{1}{N^{3/2}} H_{i,j}^{N,n} + R_{i,j}^{N,n}, \end{aligned} \quad (3.7)$$

where $H_{i,j}^{N,n}$ is the term coming from the second order derivative (i.e. $\partial_u^2 \Phi(u, v)$, $\partial_v^2 \Phi(u, v)$ and $\partial_u \partial_v \Phi(u, v)$), and $R_{i,j}^{N,n}$ is the remainder ε_3 , consequently this term correspond to the third order derivatives. Then one can use the following inequality. Given two matrices A and B of size N ,

$$\sum_{1 \leq i, j \leq N} |A_{i,j} B_{i,j}| \leq \sqrt{\sum_{1 \leq i, j \leq N} |A_{i,j}|^2} \sqrt{\sum_{1 \leq i, j \leq N} |B_{i,j}|^2} = \sqrt{\text{Tr}_N(AA^*) \text{Tr}_N(BB^*)} \leq N \|A\| \|B\|. \quad (3.8)$$

Consequently thanks to Equation (2.1), after summing over i and j , one can bound the third, fourth, sixth and seventh line in Equation (3.7) by

$$\frac{y^2}{N^2} \mathbb{E} \left[|M_N|^{n-2} \|A_1(Z^N)\| \dots \|A_l(Z^N)\| \right], \quad (3.9)$$

for some polynomials A_1 to A_k and $Z^N = (X^N, Y^N, A^N)$. Similarly after summing over i and j , one can bound the fifth line in Equation (3.7) by

$$\frac{y^2}{N} \mathbb{E} [|M_N|^{n-1} \|A_1(Z^N)\| \dots \|A_l(Z^N)\|]. \quad (3.10)$$

In order to apply our induction hypothesis, we need to first remove the term $\|A_1(Z^N)\| \dots \|A_l(Z^N)\|$ inside the expectation. To do so, let us first remark that since given a self-adjoint operator T , we have that $\|e^{iT}\| = 1$, consequently with R_1, \dots, R_k as in Proposition 3.1, we have for every $\varepsilon > 0$,

$$\begin{aligned} & \mathbb{E} [|M_N|^{n-1} \|A_1(Z^N)\| \dots \|A_l(Z^N)\|] \\ & \leq N^\varepsilon \mathbb{E} [|M_N|^{n-1}] + 2^{n-1} \mathbb{E} \left[\|R_1(X^N, A^N)\|^{n-1} \|R_1(Y^N, A^N)\|^{n-1} \dots \|R_k(X^N, A^N)\|^{n-1} \right. \\ & \quad \left. \|R_k(Y^N, A^N)\|^{n-1} \times \|A_1(Z^N)\| \dots \|A_l(Z^N)\| \mathbf{1}_{\|A_1(Z^N)\| \dots \|A_l(Z^N)\| \geq N^\varepsilon} \right], \end{aligned} \quad (3.11)$$

Then we use the following inequalities, first for any non-negative random variable A and B we have that

$$\begin{aligned} \mathbb{E}[AB \mathbf{1}_{B \geq N^\varepsilon}] & \leq N^\varepsilon \mathbb{E}[B \mathbf{1}_{B \geq N^\varepsilon}] + \mathbb{E}[AB \mathbf{1}_{A \geq N^\varepsilon} \mathbf{1}_{B \geq N^\varepsilon}] \\ & \leq N^\varepsilon \mathbb{E}[B \mathbf{1}_{B \geq N^\varepsilon}] + \mathbb{E}[AB \mathbf{1}_{AB \geq N^\varepsilon}]. \end{aligned} \quad (3.12)$$

Secondly we have that for any $g > 1$,

$$\begin{aligned} \mathbb{E}[B \mathbf{1}_{B \geq N^\varepsilon}] & = N^\varepsilon \mathbb{P}(B \geq N^\varepsilon) + \int_{N^\varepsilon}^{\infty} \mathbb{P}(B \geq \delta) d\delta \\ & \leq \frac{1}{N^{\varepsilon(g-1)}} \mathbb{E}[B^g] \left(1 + \frac{1}{g-1} \right). \end{aligned} \quad (3.13)$$

Finally one can use the fact that for any $g \in \mathbb{N}$,

$$\begin{aligned} & \mathbb{E} \left[\|A_1(Z^N)\|^g \dots \|A_l(Z^N)\|^g \right] \\ & \leq \mathbb{E} \left[\|A_1(Z^N)\|^{gl} \right]^{1/l} \dots \mathbb{E} \left[\|A_l(Z^N)\|^{gl} \right]^{1/l} \\ & \leq \mathbb{E} \left[\text{Tr}_N \left(A_1(Z^N)^{2gl} \right) \right]^{1/(2l)} \dots \mathbb{E} \left[\text{Tr}_N \left(A_k(Z^N)^{2gl} \right) \right]^{1/(2l)} \\ & \leq NC_g. \end{aligned} \quad (3.14)$$

Indeed the fact that the expectation of the normalized trace of a polynomial in independent Wigner matrices converges and hence is bounded is a well-known fact; see the proof of Theorem 5.4.2 in [3] for example.

Consequently by picking g large enough, one has that for every $\varepsilon > 0$, there exists a constant $C_{n,\varepsilon}$ such that the terms in Equations (3.9) and (3.10) are bounded respectively by

$$C_{n,\varepsilon} N^\varepsilon \frac{y^2}{N^2} \mathbb{E} [|M_N|^{n-2}] + \frac{y^2}{N^n} \quad \text{and} \quad C_{n,\varepsilon} N^\varepsilon \frac{y^2}{N} \mathbb{E} [|M_N|^{n-1}] + \frac{y^2}{N^n},$$

where as defined in proposition 3.3, $y = 1 + \max_i |y_i|$. From there on we can apply our induction hypothesis to bound the terms in Equations (3.9) and (3.10) by

$$C_{n,\varepsilon} \times \left(\frac{y^A}{N^{1-\varepsilon}} \right)^n,$$

for some constant $C_{n,\varepsilon}$. Consequently, after summing over i, j , Equation (3.7) simplifies into

$$\begin{aligned} & e^{t/2} \mathbb{E} \left[\times \text{tr}_N (D_s Q(X_t^N) Y_s^N) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] \\ & = \mathbb{E} \left[\text{tr}_N^{\otimes 2} (\partial_s D_s Q(X_t^N)) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] \end{aligned} \quad (3.15)$$

$$+ \frac{1}{N^{3/2}} \sum_{1 \leq i, j \leq N} H_{i,j}^{N,n} + \sum_{1 \leq i, j \leq N} R_{i,j}^{N,n} + \mathcal{O} \left(\left(\frac{y^4}{N^{1-\varepsilon}} \right)^n \right).$$

Next we want to tackle the term $H_{i,j}^{N,n}$. Thanks to assumption (2.1), $H_{i,j}^{N,n}$ can be bounded by a linear combination of terms of the form

$$\begin{aligned} & \mathbb{E} [|\operatorname{tr}_N(A F_1 B F_2 C F_3)| \times |M_N|^{n-1}], \\ & n \mathbb{E} [|\operatorname{tr}_N(A F_1 B F_2) \operatorname{tr}_N(C F_3)| \times |M_N|^{n-2}], \\ & n^2 \mathbb{E} [|\operatorname{tr}_N(A F_1) \operatorname{tr}_N(B F_2) \operatorname{tr}_N(C F_3)| \times |M_N|^{n-3}], \end{aligned}$$

with $F_1, F_2, F_3 \in \{E_{i,j}, E_{j,i}\}$ and A, B, C are elements of $\mathcal{F}_{d,q}$ evaluated in X_t^N . Besides thanks once again to assumption (2.1), the mixed cumulant of the real and imaginary part of $(Y_s^N)_{i,j}$ can be uniformly bounded over i, j and N , so one can sum $H_{i,j}^{N,n}$ over i, j and by using this kind of inequalities

$$\begin{aligned} \sum_{1 \leq i, j \leq N} |A_{i,j} B_{i,j} C_{i,j}| & \leq \sqrt{\sum_{1 \leq i, j \leq N} |A_{i,j} B_{i,j}|^2} \sqrt{\sum_{1 \leq i, j \leq N} |C_{i,j}|^2} \\ & \leq \sqrt{\operatorname{Tr}_N(A^* A) \operatorname{Tr}_N(C^* C)} \|B\| \\ & \leq N \|A\| \|B\| \|C\|, \end{aligned} \quad (3.16)$$

and

$$\begin{aligned} \sum_{1 \leq i, j \leq N} |A_{i,i} B_{j,j} C_{i,j}| & \leq \sqrt{\sum_{1 \leq i, j \leq N} |A_{i,i} B_{j,j}|^2} \sqrt{\sum_{1 \leq i, j \leq N} |C_{i,j}|^2} \\ & \leq \sqrt{\operatorname{Tr}_N(A^* A) \operatorname{Tr}_N(B^* B) \operatorname{Tr}_N(C^* C)} \\ & \leq N^{3/2} \|A\| \|B\| \|C\|, \end{aligned} \quad (3.17)$$

one has the following bounds for some constant C_n , and polynomials A_1 to A_l ,

$$\sum_{i,j} |H_{i,j}^{N,n}| \leq C_n \frac{y^3}{N} \mathbb{E} \left[\left(|M_N|^{n-1} + \frac{1}{N} |M_N|^{n-2} + \frac{1}{N^2} |M_N|^{n-3} \right) \|A_1(Z^N)\| \dots \|A_l(Z^N)\| \right].$$

Consequently, similarly to Equation (3.11), for any $\varepsilon > 0$,

$$\begin{aligned} \sum_{i,j} |H_{i,j}^{N,n}| & \leq C_n \frac{y^3}{N^{1-\varepsilon}} \mathbb{E} \left[|M_N|^{n-1} + \frac{1}{N} |M_N|^{n-2} + \frac{1}{N^2} |M_N|^{n-3} \right] \\ & \quad + 2^{n-1} C_n \frac{y^3}{N} \mathbb{E} \left[\|R_1(X^N, A^N)\|^{n-1} \|R_1(Y^N, A^N)\|^{n-1} \dots \|R_k(X^N, A^N)\|^{n-1} \right. \\ & \quad \left. \|R_k(Y^N, A^N)\|^{n-1} \|A_1(Z^N)\| \dots \|A_l(Z^N)\| \mathbf{1}_{\|A_1(Z^N)\| \dots \|A_l(Z^N)\| \geq N^\varepsilon} \right]. \end{aligned}$$

Then by using Equations (3.12),(3.13),(3.14), one has that there exists a constant $C_{n,\varepsilon}$ such that

$$\sum_{i,j} |H_{i,j}^{N,n}| \leq C_{n,\varepsilon} \left(\frac{y^3}{N^{1-\varepsilon}} \mathbb{E} \left[|M_N|^{n-1} + \frac{1}{N} |M_N|^{n-2} + \frac{1}{N^2} |M_N|^{n-3} \right] + \frac{y^3}{N^n} \right).$$

Hence by using our induction hypothesis, we have that for some constant $C_{n,\varepsilon}$,

$$\sum_{i,j} |H_{i,j}^{N,n}| \leq C_{n,\varepsilon} \frac{y^3}{N^{1-\varepsilon}} \times \left(\frac{y^4}{N^{1-\varepsilon}} \right)^{n-1}.$$

It remains to bound $R_{i,j}^{N,n}$. Since we have an explicit expression for the remainder in Proposition 2.17, we let $M_N^{t,i,j,s}$ be defined just like M_N but with the entries (j, i) and (i, j) of the matrix Y_s^N multiplied by t . Then one has that $R_{i,j}^{N,n}$ is bounded by a linear combination of terms of the form,

$$\frac{y^4}{N^{2-r/2}} \mathbb{E} \left[|(Y_s^N)_{i,j}|^r \int_0^1 \frac{|M_N^{t,i,j,s}|^{n-m}}{N^m} dt \|A_1(Z^N)\| \dots \|A_l(Z^N)\| \right], \quad (3.18)$$

where $r, m \in \llbracket 1, 4 \rrbracket$, and A_1 to A_l are polynomials. Besides we have that,

$$|M_N^{t,i,j,s} - M_N| \leq \frac{ty \left(|(Y_s^N)_{j,i}| + |(Y_s^N)_{i,j}| \right)}{N} \|B_1(Z^N)\| \dots \|B_{l'}(Z^N)\|, \quad (3.19)$$

for some polynomials B_1 to $B_{l'}$. Hence since Y is Hermitian, if n is an integer, one can bound (3.18) by a linear combination of terms of the form, for $g \in [0, n - m]$,

$$\frac{y^{4+g}}{N^{2-r/2}} \mathbb{E} \left[|(Y_s^N)_{i,j}|^{r+g} \frac{|M_N|^{n-m-g}}{N^{m+g}} \|A_1(Z^N)\| \dots \|A_l(Z^N)\| \left(\|B_1(Z^N)\| \dots \|B_{l'}(Z^N)\| \right)^g \right],$$

which can in turn be bounded by terms of the form

$$\begin{aligned} & \frac{y^{4+g}}{N^{2+(1-\varepsilon)g/2-\varepsilon r/2}} \mathbb{E} \left[\frac{|M_N|^{n-m-g}}{N^{m+g}} \|C_1(Z^N)\| \dots \|C_l(Z^N)\| \right] \\ & + \frac{y^{4+g}}{N^{2-r/2+m+g}} \mathbb{E} \left[\mathbf{1}_{|(Y_s^N)_{i,j}| \geq N^{\varepsilon-1/2}} |(Y_s^N)_{i,j}|^{r+g} \|\tilde{C}_1(Z^N)\| \dots \|\tilde{C}_{l'}(Z^N)\| \right], \end{aligned}$$

where C_i and \tilde{C}_i are polynomials. Thus by using equations (3.12), (3.13) and (3.14), as well as assumption (2.1), we can bounded this term by

$$C_{n,\varepsilon} \left(\frac{y^{4+g}}{N^{2+(1-\varepsilon)g/2-\varepsilon r/2+m+g}} \mathbb{E} [|M_N|^{n-m-g}] + \frac{y^{4+g}}{N^{n+2}} \right),$$

for some constant $C_{n,\varepsilon}$. Thus by applying directly our induction hypothesis, we get that $R_{i,j}^{N;n}$ is bounded by $N^{-2} \times (y^4 N^{\varepsilon-1})^n$ up to a multiplicative constant. Hence returning to Equation (3.15), we get that

$$\begin{aligned} & e^{t/2} \mathbb{E} \left[\text{tr}_N (D_s Q(X_t^N) Y_s^N) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] \\ & = \mathbb{E} \left[\text{tr}_N^{\otimes 2} (\partial_s D_s Q(X_t^N)) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] + \mathcal{O} \left(\left(\frac{y^4}{N^{1-\varepsilon}} \right)^n \right) \end{aligned}$$

Consequently by plugging this estimate back into (3.3), we have that $\mathbb{E}[|M_N|^n]$ is of order $\left(\frac{y^4}{N^{1-\varepsilon}} \right)^n$ for any ε .

Finally, in the case of $n = 1, 2, 3$, we proceed as such. Since $M_N^2 = M_N \overline{M_N}$, one can use the same proof as previously to prove that

$$\mathbb{E}[|M_N|^2] \leq C_{n,\varepsilon} \left(\frac{y^4}{N^{1-\varepsilon}} \mathbb{E}[|M_N|] + \frac{y^8}{N^{(1-\varepsilon)2}} \right).$$

But by Jensen's inequality, $\mathbb{E}[|M_N|]^2 \leq \mathbb{E}[|M_N|^2]$, consequently

$$\mathbb{E}[|M_N|]^2 \leq C_{n,\varepsilon} \left(\frac{y^4}{N^{1-\varepsilon}} \mathbb{E}[|M_N|] + \frac{y^8}{N^{(1-\varepsilon)2}} \right).$$

Hence necessarily our induction hypothesis is true for $n = 1$, and thus $n = 2$ as well. We proceed similarly for $n = 3$. Thus for any $\delta > 0$, we have for any n ,

$$\mathbb{P} \left(|\text{tr}_N (Q(Y^N, A^N)) - \mathbb{E}[\text{tr}_N (Q(X^N, A^N))]| \geq \delta \right) \leq \mathbb{E}[|M_N|^n] \delta^{-n} \leq C_{n,\varepsilon} \left(\frac{y^4}{N^{1-\varepsilon}} \right)^n \delta^{-n}$$

Consequently by Borel-Cantelli, for any $\varepsilon > 0$, we have with high probability that

$$\text{tr}_N (Q(Y^N, A^N)) = \mathbb{E}[\text{tr}_N (Q(X^N, A^N))] + \mathcal{O} \left(\frac{y^4}{N^{1-\varepsilon}} \right),$$

which proves Equation (3.1).

Note that in the case where Y_s^N is a GOE random matrix, then by using Proposition 2.15 (with a slight modification to take into account the term $M_N^{\frac{n}{2}-1}\overline{M_N^{\frac{n}{2}}}$) one has that

$$\begin{aligned}
& e^{t/2} \mathbb{E} \left[\text{tr}_N (D_s Q(X_t^N) Y_s^N) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] \\
&= \frac{1}{N} \mathbb{E} \left[\text{tr}_N^{\otimes 2} (\partial_s D_s Q(X_t^N)) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] \\
&\quad + \frac{1}{N^2} \left(\frac{n}{2} - 1 \right) \mathbb{E} \left[\text{tr}_N (D_s Q(X_t^N) D_s Q(X_0^N)) M_N^{\frac{n}{2}-2} \overline{M_N^{\frac{n}{2}}} \right] \\
&\quad + \frac{n}{2N^2} \mathbb{E} \left[\text{tr}_N \left(D_s Q(X_t^N) (D_s Q(X_0^N))^* \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}-1}} \right] \\
&\quad + \frac{1}{N} \mathbb{E} \left[\text{tr}_N (h(\partial_s D_s Q(X_t^N))) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] \\
&\quad + \frac{1}{N^2} \left(\frac{n}{2} - 1 \right) \mathbb{E} \left[\text{tr}_N \left((D_s Q(X_t^N))^T D_s Q(X_0^N) \right) M_N^{\frac{n}{2}-2} \overline{M_N^{\frac{n}{2}}} \right] \\
&\quad + \frac{n}{2N^2} \mathbb{E} \left[\text{tr}_N \left((D_s Q(X_t^N))^T (D_s Q(X_0^N))^* \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}-1}} \right], \tag{3.20}
\end{aligned}$$

where A^T is the transpose of A , and h is the linear map such that $h(A \otimes B) = A^T B$. Thus assuming that $\mathbb{E}[|M_N|^k] = \mathcal{O}(y^{2k} N^{-k(1-\varepsilon)})$ for $k < n$, then one proves that

$$\begin{aligned}
& e^{t/2} \mathbb{E} \left[\text{tr}_N (D_s Q(X_t^N) Y_s^N) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] \\
&= \mathbb{E} \left[\text{tr}_N^{\otimes 2} (\partial_s D_s Q(X_t^N)) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] + \mathcal{O} \left(\left(\frac{y^2}{N^{1-\varepsilon}} \right)^n \right).
\end{aligned}$$

Hence by reinjecting in Equation (3.3), we prove by induction that $\mathbb{E}[|M_N|^n] = \mathcal{O}(y^{2n} N^{-n(1-\varepsilon)})$ for any n . The case where Y_s^N is a GUE random matrix is even simpler. Indeed we compute that

$$\begin{aligned}
& e^{t/2} \mathbb{E} \left[\text{tr}_N (D_s Q(X_t^N) Y_s^N) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] \\
&= \frac{1}{N} \mathbb{E} \left[\text{tr}_N^{\otimes 2} (\partial_s D_s Q(X_t^N)) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] \\
&\quad + \frac{1}{N^2} \left(\frac{n}{2} - 1 \right) \mathbb{E} \left[\text{tr}_N (D_s Q(X_t^N) D_s Q(X_0^N)) M_N^{\frac{n}{2}-2} \overline{M_N^{\frac{n}{2}}} \right] \\
&\quad + \frac{n}{2N^2} \mathbb{E} \left[\text{tr}_N \left(D_s Q(X_t^N) (D_s Q(X_0^N))^* \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}-1}} \right],
\end{aligned}$$

and the result follows. \square

3.2 Concentration of the scalar product

Next we prove a similar concentration result for the scalar product. The main difference is that instead of having a concentration of order N^{-1} , it is only of order $N^{-1/2}$.

Proposition 3.3. *Given*

- Y^N a d -tuple of independent Wigner matrices as in Definition 2.1,
- X^N a d -tuple of independent GUE random matrices, independent from Y^N ,
- A^N deterministic matrices such that $\sup_{1 \leq i \leq q, N \in \mathbb{N}^*} \|A_i^N\| < \infty$.

Then with $\mathbf{x}, \mathbf{y} \in \mathbb{C}^N$, $y_i \in \mathbb{R}$, and $y = 1 + \max_i |y_i|$, P_1, \dots, P_k non-commutative self-adjoint polynomials, R_1, \dots, R_k non-commutative polynomials, if $Q = e^{iP_1 y_1} R_1 \dots e^{iP_k y_k} R_k$, we have for any $\varepsilon > 0$ that with high probability,

$$\langle \mathbf{x}, Q(Y^N, A^N) \mathbf{y} \rangle = \mathbb{E} [\langle \mathbf{x}, Q(X^N, A^N) \mathbf{y} \rangle] + \mathcal{O} \left(N^\varepsilon \frac{y^4}{\sqrt{N}} \|\mathbf{x}\|_2 \|\mathbf{y}\|_2 \right). \tag{3.21}$$

Besides if Y^N a d -tuple of independent GOE or GUE matrices, then one can replace y^4 by y^2 in the previous equality.

Proof. First one has that for any matrix $R \in \mathbb{M}_N(\mathbb{C})$, $\langle \mathbf{x}, R\mathbf{y} \rangle = \text{Tr}_N(R\mathbf{y}\mathbf{x}^*)$. Consequently, with $P = \mathbf{y}\mathbf{x}^*$, as previously we set

$$M_N := \text{Tr}_N \left((Q(Y^N, A^N) - \mathbb{E}[Q(X^N, A^N)])P \right),$$

and we want to study the moments of M_N in order to use Markov's inequality, which we again do by induction. We want to prove that for any $n \in \mathbb{N}, \varepsilon > 0$ there exist a constant $C_{n,\varepsilon}$ such that

$$\mathbb{E}[|M_N|^n] \leq C_{n,\varepsilon} \left(\frac{y^4}{N^{1/2-\varepsilon}} \|P\| \right)^n. \quad (3.22)$$

One can always assume that $y^4 \leq \sqrt{N}$, otherwise, since

$$|M_N| \leq (\|Q(Y^N, A^N)\| + \mathbb{E}[\|Q(X^N, A^N)\|]) \|P\|,$$

we immediately get Equation (3.22) thanks to Equations (3.13) and (3.14). We will show at the end of the proof why (3.22) holds for $n = 1, 2, 3$. First just like in Equation (3.3), we have that for $n \in \mathbb{N}, n \geq 4$,

$$\begin{aligned} \mathbb{E}[|M_N|^n] &= \frac{1}{2} \sum_{1 \leq s \leq d} \int_0^\infty e^{-t} \mathbb{E} \left[\left(e^{t/2} \text{Tr}_N \left(\partial_s Q(X_t^N) \# P \times Y_s^N \right) \right. \right. \\ &\quad \left. \left. - N^{-1} \text{Tr}_N^{\otimes 2} \left(\partial_s (\partial_s Q(X_t^N) \# P) \right) \right) \times M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] dt. \end{aligned} \quad (3.23)$$

Then we want to use the cumulant expansion with $u = \Re((Y_s^N)_{i,j})$ and $v = \Im((Y_s^N)_{i,j})$, and

$$\Phi : (\Re((Y_s^N)_{i,j}), \Im((Y_s^N)_{i,j})) \longmapsto e^{t/2} \text{Tr}_N \left(\partial_s Q(X_t^N) \# P \times E_{i,j} \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}}.$$

In particular if $i \neq j$, we have that

$$\begin{aligned} \partial_u \Phi(u, v) &= \text{Tr}_N \left(\left(\partial_s \left(\partial_s Q(X_t^N) \# P \right) \# (E_{i,j} + E_{j,i}) \right) E_{i,j} \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \\ &\quad + \left(\frac{n}{2} - 1 \right) \text{Tr}_N \left(\left(\partial_s Q(X_t^N) \# P \right) E_{i,j} \right) \text{tr}_N \left(\left(\partial_s Q(X_0^N) \# P \right) (E_{i,j} + E_{j,i}) \right) M_N^{\frac{n}{2}-2} \overline{M_N^{\frac{n}{2}}} \\ &\quad + \frac{n}{2} \text{Tr}_N \left(\left(\partial_s Q(X_t^N) \# P \right) E_{i,j} \right) \text{tr}_N \left(\left(\partial_s Q(X_0^N) \# P \right)^* (E_{i,j} + E_{j,i}) \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}-1}}, \end{aligned} \quad (3.24)$$

and

$$\begin{aligned} \partial_v \Phi(u, v) &= \mathbf{i} \text{Tr}_N \left(\left(\partial_s \left(\partial_s Q(X_t^N) \# P \right) \# (E_{i,j} - E_{j,i}) \right) E_{i,j} \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \\ &\quad + \mathbf{i} \left(\frac{n}{2} - 1 \right) \text{Tr}_N \left(\left(\partial_s Q(X_t^N) \# P \right) E_{i,j} \right) \text{tr}_N \left(\left(\partial_s Q(X_0^N) \# P \right) (E_{i,j} - E_{j,i}) \right) M_N^{\frac{n}{2}-2} \overline{M_N^{\frac{n}{2}}} \\ &\quad + \mathbf{i} \frac{n}{2} \text{Tr}_N \left(\left(\partial_s Q(X_t^N) \# P \right) E_{i,j} \right) \text{tr}_N \left(\left(\partial_s Q(X_0^N) \# P \right)^* (E_{i,j} - E_{j,i}) \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}-1}}. \end{aligned} \quad (3.25)$$

Moreover if $i = j$, then $\partial_u \Phi(u, v)$ is defined similarly but with $E_{i,i}$ instead of $E_{i,j} + E_{j,i}$, and $\partial_v \Phi(u, v) = 0$. By using again Equations (3.6) and Proposition 2.17 with $p = 1$, we get that

$$\begin{aligned} &e^{t/2} \mathbb{E} \left[(Y_s^N)_{i,j} \times \text{Tr}_N \left(\partial_s (Q(X_t^N) \# P) E_{i,j} \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] \\ &= \frac{1}{N} \mathbb{E} \left[\text{Tr}_N \left(\left(\partial_s \left(\partial_s Q(X_t^N) \# P \right) \# E_{j,i} \right) E_{i,j} \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] \\ &\quad + \frac{1}{N} \left(\frac{n}{2} - 1 \right) \mathbb{E} \left[\text{Tr}_N \left(\left(\partial_s Q(X_t^N) \# P \right) E_{i,j} \right) \text{Tr}_N \left(\left(\partial_s Q(X_0^N) \# P \right) E_{j,i} \right) M_N^{\frac{n}{2}-2} \overline{M_N^{\frac{n}{2}}} \right] \\ &\quad + \frac{n}{2N} \mathbb{E} \left[\text{Tr}_N \left(\left(\partial_s Q(X_t^N) \# P \right) E_{i,j} \right) \text{Tr}_N \left(\left(\partial_s Q^*(X_0^N) \# P \right) E_{j,i} \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}-1}} \right] \\ &\quad + \left(\mathbb{E}[(Y_s^N)_{i,j}^2] - \frac{1_{i=j}}{N} \right) \mathbb{E} \left[\text{tr}_N \left(\left(\partial_s (\partial_s Q(X_t^N) \# P) \# E_{i,j} \right) E_{i,j} \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] \end{aligned} \quad (3.26)$$

$$\begin{aligned}
& + \left(\frac{n}{2} - 1\right) \left(\mathbb{E}[(Y_s^N)_{i,j}^2] - \frac{\mathbf{1}_{i=j}}{N} \right) \mathbb{E} \left[\text{tr}_N \left((\partial_s Q(X_t^N) \tilde{\#} P) E_{i,j} \right) \text{tr}_N \left((\partial_s Q(X_0^N) \tilde{\#} P) E_{i,j} \right) M_N^{\frac{n}{2}-2} \overline{M_N^{\frac{n}{2}}} \right] \\
& + \frac{n}{2} \left(\mathbb{E}[(Y_s^N)_{i,j}^2] - \frac{\mathbf{1}_{i=j}}{N} \right) \mathbb{E} \left[\text{tr}_N \left((\partial_s Q(X_t^N) \tilde{\#} P) E_{i,j} \right) \text{tr}_N \left((\partial_s Q(X_0^N) \tilde{\#} P)^* E_{i,j} \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}-1}} \right] \\
& + R_{i,j}^{N,n},
\end{aligned}$$

where $R_{i,j}^{N,n}$ is the remainder ε_2 , consequently this term correspond to the second order derivatives. Then one can use the following inequality. Given two matrices A and B of size N and rank 1,

$$\sum_{1 \leq i,j \leq N} |A_{i,j} B_{i,j}| \leq \sqrt{\sum_{1 \leq i,j \leq N} |A_{i,j}|^2} \sqrt{\sum_{1 \leq i,j \leq N} |B_{i,j}|^2} = \sqrt{\text{Tr}_N(AA^*) \text{Tr}_N(BB^*)} \leq \|A\| \|B\|. \quad (3.27)$$

Consequently thanks to Equation (2.1), one can bound the third, fourth, sixth and seventh line of Equation (3.26) by

$$\frac{y^2 \|P\|^2}{N} \mathbb{E} \left[|M_N|^{n-2} \|A_1(Z^N)\| \dots \|A_l(Z^N)\| \right], \quad (3.28)$$

for some polynomials A_1 to A_l and $Z^N = (X^N, Y^N, A^N)$. Similarly one can bound the fifth line by

$$\frac{y^2 \|P\|}{\sqrt{N}} \mathbb{E} \left[|M_N|^{n-1} \|A_1(Z^N)\| \dots \|A_l(Z^N)\| \right]. \quad (3.29)$$

One now wants to remove the term $\|A_1(Z^N)\| \dots \|A_l(Z^N)\|$ inside the expectation in order to use our induction hypothesis. To do so, let us first remark that given a self-adjoint operator T , we have that $\|e^{iT}\| = 1$, consequently with R_1, \dots, R_k as in Proposition 3.3, we have for every $\varepsilon > 0$,

$$\begin{aligned}
& \mathbb{E} \left[|M_N|^{n-1} \|A_1(Z^N)\| \dots \|A_l(Z^N)\| \right] \\
& \leq N^\varepsilon \mathbb{E} \left[|M_N|^{n-1} \right] + 2^{n-1} \mathbb{E} \left[\left\| R_1(X^N, A^N) \right\|^{n-1} \left\| R_1(Y^N, A^N) \right\|^{n-1} \dots \left\| R_k(X^N, A^N) \right\|^{n-1} \right. \\
& \quad \left. \left\| R_k(Y^N, A^N) \right\|^{n-1} \|A_1(Z^N)\| \dots \|A_l(Z^N)\| \mathbf{1}_{\|A_1(Z^N)\| \dots \|A_l(Z^N)\| \geq N^\varepsilon} \right] \|P\|^{n-1}, \quad (3.30)
\end{aligned}$$

Consequently by using Equations (3.12), (3.13) and (3.14), one has that for every $\varepsilon > 0$, there exists a constant $C_{n,\varepsilon}$ such that the terms in Equations (3.28) and (3.29) are bounded respectively by

$$C_{n,\varepsilon} N^\varepsilon \frac{y^2 \|P\|^2}{N} \mathbb{E} \left[|M_N|^{n-2} \right] + \frac{y^2 \|P\|^n}{N^n} \quad \text{and} \quad C_{n,\varepsilon} N^\varepsilon \frac{y^2 \|P\|}{N} \mathbb{E} \left[|M_N|^{n-1} \right] + \frac{y^2 \|P\|^n}{N^n}.$$

From there on one can apply our induction hypothesis to bound the terms in Equations (3.28) and (3.29) by

$$C_{n,\varepsilon} \times \left(\frac{y^4 \|P\|}{N^{1/2-\varepsilon}} \right)^n,$$

for some constant $C_{n,\varepsilon}$. Thus after summing over i, j , Equation (3.26) simplifies into

$$\begin{aligned}
& e^{t/2} \mathbb{E} \left[\text{Tr}_N \left(\partial_s(Q(X_t^N) \tilde{\#} P) Y_s^N \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] \\
& = \frac{1}{N} \mathbb{E} \left[\text{Tr}_N^{\otimes 2} \left(\partial_s \left(\partial_s Q(X_t^N) \tilde{\#} P \right) \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] \\
& \quad + \sum_{1 \leq i,j \leq N} R_{i,j}^{N,n} + \mathcal{O} \left(\left(\frac{y^4 \|P\|}{N^{1/2-\varepsilon}} \right)^n \right), \quad (3.31)
\end{aligned}$$

Since we have an explicit expression of the remainder $R_{i,j}^{N,n}$ in Proposition 2.17, we define $M_N^{t,i,j,s}$ just like M_N but with the entries (i, j) and (j, i) of the matrix Y_s^N multiplied by t . Then we have that $R_{i,j}^{N,n}$ is bounded by a linear combination of terms of the form,

$$\frac{y^3}{N^{3/2-r/2}} \mathbb{E} \left[|(Y_s^N)_{i,j}|^r \int_0^1 \left| (A_1^{t,i,j,s})_{e_1, e_2} \dots (A_3^{t,i,j,s})_{e_5, e_6} \right| |M_N^{t,i,j,s}|^{n-m} dt \right], \quad (3.32)$$

where $r, m \in \llbracket 1, 3 \rrbracket$, $e_1, \dots, e_6 \in \{i, j\}$, and $A_1^{t,i,j,s}$ to $A_3^{t,i,j,s}$ are matrices which are evaluation of polynomials in $\mathcal{F}_{d,q}$. Besides thanks to P , m of those matrices have rank 1. Furthermore the indices e_1, \dots, e_6 come from the derivatives with respect to u and v which makes $E_{i,j}$ and $E_{j,i}$ appear, coupled with the fact that $\text{Tr}_N(AE_{i,j}) = A_{j,i}$. Consequently half of the indices e_l are equal to i and the other half to j . This fact will come in handy in the next few computations.

Now one proceeds as such, one can find matrices A_1^1, A_1^2 , polynomials B_1 to B_l and C_1 to $C_{l'}$ such that

$$\begin{aligned} |(A_1^{t,i,j,s})_{e_1, e_2}| &\leq |(A_1^1)_{e_1, e_2}| + |y(Y_s^N)_{i,j}| |(A_1^2)_{\tilde{e}_1, \tilde{e}_2}| \|B_1(Z^N)\| \dots \|B_l(Z^N)\| \\ &\quad + |y(Y_s^N)_{i,j}|^2 \|C_1(Z^N)\| \dots \|C_{l'}(Z^N)\| \|P\|^{h_1}, \end{aligned}$$

where either $\tilde{e}_1 = e_1$ or $\tilde{e}_2 = e_2$. Besides if A_1^1 is of rank 1, then so is A_1^2 and $h_1 = 1$ (otherwise $h_1 = 0$). One can also find polynomials D_1 to D_l such that

$$|M_N^{t,i,j,s} - M_N| \leq |y(Y_s^N)_{i,j}| \|D_1(Z^N)\| \dots \|D_l(Z^N)\| \|P\|,$$

which means that

$$|M_N^{t,i,j,s}|^{n-m} \leq \sum_{i=1}^{n-m} \binom{n-m}{p} |y(Y_s^N)_{i,j}|^p \|D_1(Z^N)\|^p \dots \|D_l(Z^N)\|^p \|P\|^p |M_N|^{n-m-p}.$$

Thus thanks to the estimates in (3.12), (3.13), (3.14) as well as Assumption (2.1), one can bound (3.32) by a linear combination of terms of the form

$$\begin{aligned} C_{n,\varepsilon} N^\varepsilon \frac{y^{3+p} \|P\|^p}{N^{(3+p)/2}} \mathbb{E} \left[\left(|(A_1^1)_{e_1, e_2}| + \frac{y}{\sqrt{N}} |(A_1^2)_{\tilde{e}_1, \tilde{e}_2}| + \frac{y^2 \|P\|^{h_1}}{N} \right) \right. \\ \times \left(|(A_2^1)_{e_3, e_4}| + \frac{y}{\sqrt{N}} |(A_2^2)_{\tilde{e}_3, \tilde{e}_4}| + \frac{y^2 \|P\|^{h_2}}{N} \right) \\ \times \left. \left(|(A_3^1)_{e_5, e_6}| + \frac{y}{\sqrt{N}} |(A_3^2)_{\tilde{e}_5, \tilde{e}_6}| + \frac{y^2 \|P\|^{h_3}}{N} \right) |M_N|^{n-m-p} \right]. \end{aligned} \quad (3.33)$$

Besides if A, B, C are matrices of size N , assuming that g of them are of rank 1, one has that

$$\begin{aligned} \sum_{1 \leq i, j \leq N} |A_{i,j} B_{i,j} C_{i,j}| &\leq \sqrt{\sum_{1 \leq i, j \leq N} |A_{i,j} B_{i,j}|^2} \sqrt{\sum_{1 \leq i, j \leq N} |C_{i,j}|^2} \\ &\leq \sqrt{\text{Tr}_N(C^* C)} \times \max\{\|B\| \sqrt{\text{Tr}_N(A^* A)}, \|A\| \sqrt{\text{Tr}_N(B^* B)}\} \\ &\leq N^{(1-g/2) \wedge 0} \|A\| \|B\| \|C\|, \end{aligned} \quad (3.34)$$

and

$$\begin{aligned} \sum_{1 \leq i, j \leq N} |A_{i,i} B_{j,j} C_{i,j}| &\leq \sqrt{\sum_{1 \leq i, j \leq N} |A_{i,i} B_{j,j}|^2} \sqrt{\sum_{1 \leq i, j \leq N} |C_{i,j}|^2} \\ &\leq \sqrt{\text{Tr}_N(A^* A) \text{Tr}_N(B^* B) \text{Tr}_N(C^* C)} \\ &\leq N^{(3-g)/2} \|A\| \|B\| \|C\|. \end{aligned} \quad (3.35)$$

Similarly, we have that

$$\begin{aligned} \sum_{1 \leq i, j \leq N} |A_{i,j} B_{i,j} C_{i,i}| &\leq N^{(3-g)/2} \|A\| \|B\| \|C\|, \\ \sum_{1 \leq i, j \leq N} |A_{j,j} B_{i,i} C_{i,i}| &\leq N^{2-g/2} \|A\| \|B\| \|C\|, \\ \sum_{1 \leq i, j \leq N} |A_{j,i} B_{i,i} C_{i,i}| &\leq N^{2-g/2} \|A\| \|B\| \|C\|, \end{aligned} \quad (3.36)$$

$$\sum_{1 \leq i, j \leq N} |A_{i,i} B_{i,i} C_{i,i}| \leq N^{2-(g \wedge 2)/2} \|A\| \|B\| \|C\|.$$

If we only have two matrices, g of them being of rank 1,

$$\begin{aligned} \sum_{1 \leq i, j \leq N} |A_{i,j} B_{i,j}| &\leq N^{1-g/2} \|A\| \|B\|, \\ \sum_{1 \leq i, j \leq N} |A_{i,j} B_{i,i}| &\leq N^{(3-g)/2} \|A\| \|B\|, \\ \sum_{1 \leq i, j \leq N} |A_{i,i} B_{i,i}| &\leq N^{2-g/2} \|A\| \|B\|. \end{aligned} \quad (3.37)$$

Further, with one matrix, with $g = 1$ if it is of rank 1 and 0 otherwise,

$$\begin{aligned} \sum_{1 \leq i, j \leq N} |A_{i,j}| &\leq N^{(3-g)/2} \|A\|, \\ \sum_{1 \leq i, j \leq N} |A_{i,i}| &\leq N^{2-g/2} \|A\|. \end{aligned} \quad (3.38)$$

Thus thanks to Equations (3.34) and (3.35), we have that

$$\frac{y^3}{N^{3/2}} \sum_{i,j} |(A_1^1)_{e_1, e_2} (A_2^1)_{e_3, e_4} (A_3^1)_{e_5, e_6}| \leq \frac{y^3}{N^{m/2}} \|A_1^1\| \|A_2^1\| \|A_3^1\|.$$

Thanks to the first two line of Equation (3.36), we also get that

$$\frac{y^4}{N^2} \sum_{i,j} |(A_1^2)_{\tilde{e}_1, \tilde{e}_2} (A_2^2)_{e_3, e_4} (A_3^1)_{e_5, e_6}| \leq \frac{y^4}{N^{m/2}} \|A_1^2\| \|A_2^1\| \|A_3^1\|.$$

With all of Equation (3.36), since we assumed that $y^4 \leq \sqrt{N}$, we have that

$$\begin{aligned} \frac{y^5}{N^{5/2}} \sum_{i,j} |(A_1^2)_{\tilde{e}_1, \tilde{e}_2} (A_2^2)_{\tilde{e}_3, \tilde{e}_4} (A_3^1)_{e_5, e_6}| &\leq \frac{y^5}{N^{(1+m)/2}} \|A_1^2\| \|A_2^2\| \|A_3^1\| \leq \frac{y}{N^{m/2}} \|A_1^2\| \|A_2^2\| \|A_3^1\|, \\ \frac{y^6}{N^3} \sum_{i,j} |(A_1^2)_{\tilde{e}_1, \tilde{e}_2} (A_2^2)_{\tilde{e}_3, \tilde{e}_4} (A_3^1)_{\tilde{e}_5, \tilde{e}_6}| &\leq \frac{y^6}{N^{1+(m \wedge 2)/2}} \|A_1^2\| \|A_2^2\| \|A_3^2\| \leq \frac{y^2}{N^{m/2}} \|A_1^2\| \|A_2^2\| \|A_3^2\|. \end{aligned}$$

Then thanks to Equation (3.37), we also have

$$\begin{aligned} \frac{y^5}{N^{5/2}} \sum_{i,j} |(A_2^1)_{e_3, e_4} (A_3^1)_{e_5, e_6}| &\leq \frac{y^5}{N^{(1+m \wedge 2)/2}} \|A_2^1\| \|A_3^1\| \leq \left(\frac{y^4}{N^{1/2}} \right)^m \|A_2^1\| \|A_3^1\|, \\ \frac{y^6}{N^3} \sum_{i,j} |(A_2^2)_{\tilde{e}_3, \tilde{e}_4} (A_3^1)_{e_5, e_6}| &\leq \frac{y^6}{N^{1+(m \wedge 2)/2}} \|A_2^2\| \|A_3^1\| \leq \left(\frac{y^4}{N^{1/2}} \right)^m \|A_2^2\| \|A_3^1\|, \\ \frac{y^7}{N^{7/2}} \sum_{i,j} |(A_2^2)_{\tilde{e}_3, \tilde{e}_4} (A_3^2)_{\tilde{e}_5, \tilde{e}_6}| &\leq \frac{y^7}{N^{(3+m \wedge 2)/2}} \|A_2^2\| \|A_3^2\| \leq \left(\frac{y^4}{N^{1/2}} \right)^m \|A_2^2\| \|A_3^2\|. \end{aligned}$$

Consequently, thanks again to Equation (3.12), (3.13), (3.14), we get that for some constant $C_{n,\varepsilon}$, after summing over i and j , the term in Equation (3.33) is bounded by

$$C_{n,\varepsilon} \left(\frac{y^4}{N^{1/2-\varepsilon}} \right)^{m+p} \mathbb{E} [|M_N|^{n-m-p}] \|P\|^{m+p}.$$

Thus from Equations (3.31) and (3.32) combined with the above estimate, after summing over i, j , we get that

$$e^{t/2} \mathbb{E} \left[\text{Tr}_N \left(\partial_s (Q(X_t^N) \# P) Y_s \right) M_N^{\frac{3}{2}-1} \overline{M_N^{\frac{3}{2}}} \right]$$

$$= \frac{1}{N} \mathbb{E} \left[\text{Tr}_N^{\otimes 2} \left(\partial_s \left(\partial_s Q(X_t^N) \tilde{\#} P \right) \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] + \mathcal{O} \left(\left(\frac{y^4}{N^{1/2-\varepsilon}} \|P\| \right)^n \right),$$

where in the last line we used the induction hypothesis coupled with the fact that the non-normalized trace of a rank 1 matrix is smaller than its norm. Hence by plugging this estimate back into Equation (3.23), we get that $\mathbb{E}[|M_N|^n]$ is of order $\left(\frac{y^4 \|P\|}{N^{1-\varepsilon}} \right)^n$ for any ε .

Finally in the case of $n = 1, 2, 3$, we proceed as such. Since $M_N^2 = M_N \overline{M_N}$, one can use the same proof as previously to prove that

$$\mathbb{E}[|M_N|^2] \leq C_{n,\varepsilon} \left(\frac{y^4}{N^{1/2-\varepsilon}} \mathbb{E}[|M_N|] \|P\| + \frac{y^8}{N^{(1/2-\varepsilon)2}} \|P\|^2 \right).$$

But by Jensen's inequality, $\mathbb{E}[|M_N|]^2 \leq \mathbb{E}[|M_N|^2]$, consequently

$$\mathbb{E}[|M_N|]^2 \leq C_{n,\varepsilon} \left(\frac{y^4}{N^{1/2-\varepsilon}} \|P\| \mathbb{E}[|M_N|] + \frac{y^8}{N^{(1/2-\varepsilon)2}} \|P\|^2 \right).$$

Hence necessarily our induction hypothesis is true for $n = 1$, and thus $n = 2$ as well. We proceed similarly for $n = 3$. Thus for any $\delta > 0$, we have for any n ,

$$\mathbb{P} \left(|\langle \mathbf{x}, Q(Y^N, A^N) \mathbf{y} \rangle - \mathbb{E}[\langle \mathbf{x}, Q(X^N, A^N) \mathbf{y} \rangle]| \geq \delta \right) \leq \mathbb{E}[|M_N|^n] \delta^{-n} \leq C_{n,\varepsilon} \left(\frac{y^4 \|P\|}{N^{1/2-\varepsilon}} \right)^n \delta^{-n}$$

Consequently by Borel Cantelli for any $\varepsilon > 0$, we have with high probability that

$$\langle \mathbf{x}, Q(Y^N, A^N) \mathbf{y} \rangle = \mathbb{E}[\langle \mathbf{x}, Q(X^N, A^N) \mathbf{y} \rangle] + \mathcal{O} \left(\frac{y^4 \|P\|}{N^{1/2-\varepsilon}} \right).$$

Hence the conclusion since $\|P\| = \|\mathbf{x}\|_2 \|\mathbf{y}\|_2$.

In the case where Y_s^N is a GOE random matrix, then then by using Proposition 2.15 (with a slight modification to take into account the term $M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}}$), Equation (3.26) simplifies into

$$\begin{aligned} & e^{t/2} \mathbb{E} \left[\text{Tr}_N \left(\partial_s(Q(X_t^N) \tilde{\#} P) Y_s^N \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] \\ &= \frac{1}{N} \mathbb{E} \left[\text{Tr}_N^{\otimes 2} \left(\partial_s \left(\partial_s Q(X_t^N) \tilde{\#} P \right) \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] \\ &+ \frac{1}{N} \left(\frac{n}{2} - 1 \right) \mathbb{E} \left[\text{Tr}_N \left(\left(\partial_s Q(X_t^N) \tilde{\#} P \right) \left(\partial_s Q(X_0^N) \tilde{\#} P \right) \right) M_N^{\frac{n}{2}-2} \overline{M_N^{\frac{n}{2}}} \right] \\ &+ \frac{n}{2N} \mathbb{E} \left[\text{Tr}_N \left(\left(\partial_s Q(X_t^N) \tilde{\#} P \right) \left(\partial_s Q(X_0^N) \tilde{\#} P \right)^* \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}-1}} \right] \\ &+ \frac{1}{N} \mathbb{E} \left[\text{Tr}_N \left(h \left(\partial_s(Q(X_t^N) \tilde{\#} P) \right) \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] \\ &+ \frac{1}{N} \left(\frac{n}{2} - 1 \right) \mathbb{E} \left[\text{Tr}_N \left(\left(\partial_s(Q(X_t^N) \tilde{\#} P) \right)^T \left(\partial_s(Q(X_0^N) \tilde{\#} P) \right) \right) M_N^{\frac{n}{2}-2} \overline{M_N^{\frac{n}{2}}} \right] \\ &+ \frac{n}{2N} \mathbb{E} \left[\text{Tr}_N \left(\left(\partial_s(Q(X_t^N) \tilde{\#} P) \right)^T \left(\partial_s(Q(X_0^N) \tilde{\#} P) \right)^* \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}-1}} \right], \end{aligned}$$

where A^T is the transpose of A , and h is the linear map such that $h(A \otimes B) = A^T B$. Thus assuming that $\mathbb{E}[|M_N|^k] = \mathcal{O}(y^{2k} N^{-k(1-\varepsilon)/2} \|P\|^k)$ for $k < n$, then one proves that

$$\begin{aligned} & e^{t/2} \mathbb{E} \left[\text{Tr}_N \left(\partial_s(Q(X_t^N) \tilde{\#} P) Y_s \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] \\ &= \frac{1}{N} \mathbb{E} \left[\text{Tr}_N^{\otimes 2} \left(\partial_s \left(\partial_s Q(X_t^N) \tilde{\#} P \right) \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] + \mathcal{O} \left(\left(\frac{y^2}{N^{1/2-\varepsilon}} \|P\| \right)^n \right), \end{aligned}$$

and hence by using Equation (3.23), one concludes by induction that $\mathbb{E}[|M_N|^n] = \mathcal{O}(y^{2n} N^{-n(1-\varepsilon)/2} \|P\|^n)$ for any n . The case where Y_s^N is a GUE random matrix is even simpler. Indeed we compute that

$$e^{t/2} \mathbb{E} \left[\text{Tr}_N \left(\partial_s(Q(X_t^N) \tilde{\#} P) Y_s^N \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right]$$

$$\begin{aligned}
&= \frac{1}{N} \mathbb{E} \left[\text{Tr}_N^{\otimes 2} \left(\partial_s \left(\partial_s Q(X_t^N) \tilde{\#} P \right) \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}}} \right] \\
&\quad + \frac{1}{N} \left(\frac{n}{2} - 1 \right) \mathbb{E} \left[\text{Tr}_N \left(\left(\partial_s Q(X_t^N) \tilde{\#} P \right) \left(\partial_s Q(X_0^N) \tilde{\#} P \right) \right) M_N^{\frac{n}{2}-2} \overline{M_N^{\frac{n}{2}}} \right] \\
&\quad + \frac{n}{2N} \mathbb{E} \left[\text{Tr}_N \left(\left(\partial_s Q(X_t^N) \tilde{\#} P \right) \left(\partial_s Q(X_0^N) \tilde{\#} P \right)^* \right) M_N^{\frac{n}{2}-1} \overline{M_N^{\frac{n}{2}-1}} \right],
\end{aligned}$$

and the result follows. \square

4 Proof of the main theorems

Proof of Theorem 1.1. To prove Equation (1.4), we combine Proposition 3.1 with Lemma 3.6 of [33]. Indeed, with Q defined as in Proposition 3.1, we have for any $\varepsilon > 0$, that with high probability

$$\text{tr}_N(Q(Y^N, A^N)) = \mathbb{E}[\text{tr}_N(Q(X^N, A^N))] + \mathcal{O}\left(N^\varepsilon \frac{y^4}{N}\right). \quad (4.1)$$

Whereas if we use Lemma 3.6 of [33] (with $n = 0$), where $x = (x_1, \dots, x_d)$ are free semicircular variables, free from $A^N \in \mathbb{M}_N(\mathbb{C})^q$, we get that

$$\begin{aligned}
\mathbb{E}[\text{tr}_N(Q(X^N, A^N))] &= \tau_N(Q(x, A^N)) \\
&\quad + \frac{1}{N^2} \int_0^\infty \int_0^t \left(\int_{[0,1]^4} e^{-r-t} \mathbb{E}[\tau_N(R_{\alpha,\beta,\delta,\gamma,r,t})] \, d\alpha d\beta d\gamma d\delta \right) dr dt,
\end{aligned} \quad (4.2)$$

where $R_{\alpha,\beta,\delta,\gamma,r,t}$ is such that for some constant C independent of $\alpha, \beta, \delta, \gamma, r, t, y$ and N ,

$$\mathbb{E}[\|R_{\alpha,\beta,\delta,\gamma,r,t}\|] \leq Cy^4.$$

Hence we have that

$$\mathbb{E}[\text{tr}_N(Q(X^N, A^N))] = \tau_N(Q(x, A^N)) + \mathcal{O}\left(\frac{y^4}{N^2}\right). \quad (4.3)$$

Finally with f_1 to f_k defined as in Theorem 1.1, we set

$$\tilde{f}_i : t \mapsto \begin{cases} e^{\mathbf{i}y_i t} & \text{if } f_i \neq \text{id}_{\mathbb{R}}, \\ t & \text{else.} \end{cases}$$

Then with $Q = \tilde{f}_1(P_1) \dots \tilde{f}_k(P_k)$, and μ_i the Dirac measure in 1 if $f_i = \text{id}_{\mathbb{R}}$, thanks to the functional calculus we have

$$f_1(P_1(Y^N, A^N)) \dots f_k(P_k(Y^N, A^N)) = \int_{\mathbb{R}^k} Q(Y^N, A^N) \, d\mu_1(y_1) \dots d\mu_k(y_k). \quad (4.4)$$

Thus by combining Equations (4.1) and (4.3), we get Equation (1.4) after integrating over y .

Besides in the case where we are working only with GUE and GOE random matrices, then one can replace y^4 by y^2 in Equation (4.1). Then if $y^2 \leq N$, Equation (4.2) let us conclude, whereas if $y^2 \geq N$, then we simply use the fact that

$$\text{tr}_N(Q(Y^N, A^N)) - \tau_N(Q(x, A^N)) = \mathcal{O}(1) = \mathcal{O}\left(\frac{y^2}{N}\right).$$

To prove Equation (1.5), we also start by using Proposition 3.3 to prove that for any $\varepsilon > 0$, with high probability

$$\langle \mathbf{x}, Q(Y^N, A^N) \mathbf{y} \rangle = \mathbb{E}[\langle \mathbf{x}, Q(X^N, A^N) \mathbf{y} \rangle] + \mathcal{O}\left(N^\varepsilon \frac{y^4}{\sqrt{N}} \|\mathbf{x}\|_2 \|\mathbf{y}\|_2\right), \quad (4.5)$$

where one can replace y^4 by y^2 in the case where we are working only with GUE and GOE random matrices. Then we need to estimate $\mathbb{E}[\langle \mathbf{x}, Q(X^N, A^N) \mathbf{y} \rangle]$. Thanks again to Lemma 3.6 of [33], we get that

$$\mathbb{E}[\langle \mathbf{x}, Q(X^N, A^N) \mathbf{y} \rangle]$$

$$\begin{aligned}
&= \mathbb{E} [\text{Tr}_N (Q(X^N, A^N) \mathbf{y} \mathbf{x}^*)] \\
&= N \tau_N (Q(x, A^N) \mathbf{y} \mathbf{x}^*) + \frac{1}{N} \int_0^\infty \int_0^t \left(\int_{[0,1]^4} e^{-r-t} \mathbb{E} [\tau_N (R_{\alpha, \beta, \delta, \gamma, r, t} \mathbf{y} \mathbf{x}^*)] d\alpha d\beta d\gamma d\delta \right) dr dt \\
&= \text{Tr}_N (E_{\mathbb{M}_N(\mathbb{C})} [Q(x, A^N)] \mathbf{y} \mathbf{x}^*) \\
&\quad + \frac{1}{N^2} \int_0^\infty \int_0^t \left(\int_{[0,1]^4} e^{-r-t} \mathbb{E} [\text{Tr}_N (E_{\mathbb{M}_N(\mathbb{C})} [R_{\alpha, \beta, \delta, \gamma, r, t}] \mathbf{y} \mathbf{x}^*)] d\alpha d\beta d\gamma d\delta \right) dr dt \\
&= \langle \mathbf{x}, E_{\mathbb{M}_N(\mathbb{C})} [Q(x, A^N)] \mathbf{y} \rangle \\
&\quad + \frac{1}{N^2} \int_0^\infty \int_0^t \left(\int_{[0,1]^4} e^{-r-t} \mathbb{E} [\text{Tr}_N (E_{\mathbb{M}_N(\mathbb{C})} [R_{\alpha, \beta, \delta, \gamma, r, t}] \mathbf{y} \mathbf{x}^*)] d\alpha d\beta d\gamma d\delta \right) dr dt,
\end{aligned}$$

where $R_{\alpha, \beta, \delta, \gamma, r, t}$ is once again such that for some constant C independent of $\alpha, \beta, \delta, \gamma, r, t, y$ and N ,

$$\mathbb{E} [\|R_{\alpha, \beta, \delta, \gamma, r, t}\|] \leq C y^4.$$

Consequently since $P = y \mathbf{x}^*$ is of rank 1 and $\|P\| = \|\mathbf{x}\|_2 \|\mathbf{y}\|_2$,

$$\mathbb{E} [\langle \mathbf{x}, Q(X^N, A^N) \mathbf{y} \rangle] = \langle \mathbf{x}, E_{\mathbb{M}_N(\mathbb{C})} [Q(x, A^N)] \mathbf{y} \rangle + \mathcal{O} \left(\frac{y^4}{N^2} \|\mathbf{x}\|_2 \|\mathbf{y}\|_2 \right).$$

The rest of the proof follows as for Equation (1.4) with the difference that after integrating over y , we use the fact that $b \mapsto E_{\mathbb{M}_N(\mathbb{C})}[b]$ is a continuous linear functional to switch the integral and the conditional expectation. \square

Proof of Theorem 1.2. Let Q_1 to Q_p be non commutative polynomials, $i_1, \dots, i_p \in \llbracket 1, k \rrbracket$ be such that for every j , $\tau(Q_j(a_{i_j})) = 0$ and if $j < p$, $i_j \neq i_{j+1}$. Then if we set $u_{2j-1}^N := e^{i(y_{i_j}^N - y_{i_{j-1}}^N)P(x)}$ (with the convention $i_0 = i_p$) and $u_{2j}^N = Q_j(A_{i_j}^N)$, one can apply Theorem 1.1 to obtain that with high probability

$$\tau_N (Q_1(a_{i_1}^N) \dots Q_p(a_{i_p}^N)) = \tau_N (u_1^N u_2^N \dots u_{2p-1}^N u_{2p}^N) + \mathcal{O} \left(\frac{1}{N} \max_{i \neq j} |y_i^N - y_j^N|^4 \right).$$

Further, if Y^N is a family of GUE and GOE matrices, then

$$\tau_N (Q_1(a_{i_1}^N) \dots Q_p(a_{i_p}^N)) = \tau_N (u_1^N u_2^N \dots u_{2p-1}^N u_{2p}^N) + \mathcal{O} \left(\frac{1}{N} \max_{i \neq j} |y_i^N - y_j^N|^2 \right).$$

Since by assumption for any i, j , $|y_i^N - y_j^N| \ll N^{1/2}$ in the case of GUE and GOE matrices, and $|y_i^N - y_j^N| \ll N^{1/4}$ in all generality, we get in both cases that with high probability

$$\tau_N (Q_1(a_{i_1}^N) \dots Q_p(a_{i_p}^N)) = \tau_N (u_1^N u_2^N \dots u_{2p-1}^N u_{2p}^N) + o(1).$$

Thus the former equality is true almost surely by Borel-Cantelli.

But then thanks to Proposition 11.4 of [30], with $\text{NCP}[2p]$ the set of non-crossing partition of $\llbracket 1, 2p \rrbracket$ (i.e. the set of partitions of $\llbracket 1, 2p \rrbracket$ such that one cannot find a, c in a first block as well as b, d in another block such that $a < b < c < d$), one can write

$$\tau_N (u_1^N \dots u_{2p}^N) = \sum_{\pi \in \text{NCP}[k]} \kappa_\pi (u_1^N, u_2^N, \dots, u_{2p-1}^N, u_{2p}^N),$$

where $\kappa_\pi (u_1^N, u_2^N, \dots, u_{2p-1}^N, u_{2p}^N)$ are the so-called free cumulants. We do not need to define those in details in this proof, however it is important to note that $\kappa_\pi (u_1^N, u_2^N, \dots, u_{2p-1}^N, u_{2p}^N)$ is a linear combination of product of traces such that u_i^N and u_j^N can only be in the same trace if i and j belong to the same block in the partition π . Besides thanks to Theorem 11.16 of [30], if i, j belongs to the same block in the partition π and u_i^N is free from u_j^N , then $\kappa_\pi (u_1^N, u_2^N, \dots, u_{2p-1}^N, u_{2p}^N) = 0$. Consequently the elements u_j^N for odd j cannot be in the same trace as those for even j . Thus by a quick induction, one

can see that necessarily for $\kappa_\pi(u_1^N, u_2^N, \dots, u_{2p-1}^N, u_{2p}^N)$ to be non zero, there has to be a block in the partition π which consists of a single element which is not 1 (this is important since we did not assume that $i_1 \neq i_p$, and hence one could have $u_1^N = 1$). Then since for any j ,

$$\lim_{N \rightarrow \infty} \tau_N(u_{2j}^N) = \tau(Q_j(a_{i_j})) = 0,$$

we only need to show that for any j , $\lim_{N \rightarrow \infty} \tau \left(e^{i(y_{i_j}^N - y_{i_{j-1}}^N)P(x)} \right) = 0$. But since by assumption $i_j \neq i_{j-1}$, we have that $|y_{i_j}^N - y_{i_{j-1}}^N| \gg 1$. Consequently, all we need to prove is that $\lim_{|t| \rightarrow \infty} \tau(e^{itP(x)}) = 0$. However thanks to [35], Theorem 1.1, (6), there exists $f \in L^1(\mathbb{R})$ such that

$$\tau(e^{itP(x)}) = \int_{\mathbb{R}} e^{ita} f(a) da.$$

Thus thanks to the Riemann-Lebesgue lemma we have that

$$\lim_{N \rightarrow \infty} \tau_N(u_{2j-1}^N) = 0.$$

Hence we proved that for any $i_1, \dots, i_p \in [1, k]$ such that for $j < p$, $i_j \neq i_{j+1}$ and non commutative polynomials Q_j , such that for every j , $\tau(Q_j(a_{i_j})) = 0$,

$$\lim_{N \rightarrow \infty} \tau_N \left(Q_1(a_{i_1}^N) \dots Q_p(a_{i_p}^N) \right) = 0.$$

Hence (a_1^N, \dots, a_k^N) converges in distribution towards the free family (a_1, \dots, a_k) . \square

Acknowledgements

We thanks László Erdős for comments on the first draft. F.P. is supported by the Knut and Alice Wallenberg Foundation. K.S. is supported in parts by the Swedish Research Council (VR-2017-05195, VR-2021-04703), and the Knut and Alice Wallenberg Foundation. This material is based upon work supported by the National Science Foundation under Grant No. DMS-1928930 while K.S participated in a program hosted by the Mathematical Sciences Research Institute in Berkeley, CA, during the Fall 2021 semester.

References

- [1] Anderson, G. W. (2013). Convergence of the largest singular value of a polynomial in independent Wigner matrices. *The Annals of Probability*, 41(3B), 2103-2181.
- [2] Anderson, G. W. (2015). A local limit law for the empirical spectral distribution of the anticommutator of independent Wigner matrices. *Annales de l'IHP Probabilités et statistiques*, 51(3), 809-841.
- [3] Anderson, G. W., Guionnet, A., & Zeitouni, O. (2010). *An introduction to random matrices* (No. 118). Cambridge university press.
- [4] Belinschi, S. T., & Capitaine, M. (2017). Spectral properties of polynomials in independent Wigner and deterministic matrices. *Journal of Functional Analysis*, 273(12), 3901-3963.
- [5] Belinschi, S. T., Mai, T., & Speicher, R. (2017). Analytic subordination theory of operator-valued free additive convolution and the solution of a general random matrix problem. *Journal für die reine und angewandte Mathematik* (Crelles Journal), 2017(732), 21-53.
- [6] Bloemendal, A., Erdős, L., Knowles, A., Yau, H. T., & Yin, J. (2014). Isotropic local laws for sample covariance and generalized Wigner matrices. *Electronic Journal of Probability*, 19, 1-53.
- [7] Boutet, D. M. A., & Khorunzhy, A. (1999). Asymptotic distribution of smoothed eigenvalue density. II. Wigner random matrices. *Random Operators and Stochastic Equations*, 7(2), 149-168.

- [8] Cipolloni, G., Erdős, L., & Schröder, D. (2022). Thermalisation for Wigner matrices. *Journal of Functional Analysis*, 282(8), 109394.
- [9] Cipolloni, G., Erdős, L., & Schröder, D. (2021). Optimal multi-resolvent local laws for Wigner matrices. *arXiv preprint arXiv:2112.13693*.
- [10] Collins, B., Mai, T., Miyagawa, A., Parraud, F., & Yin, S. (2021). Convergence for noncommutative rational functions evaluated in random matrices. *arXiv preprint arXiv:2103.05962*.
- [11] Collins, B., Guionnet, A., & Parraud, F. (2022). On the operator norm of non-commutative polynomials in deterministic matrices and iid GUE matrices. *Cambridge Journal of Mathematics*, 10(1), 195–260.
- [12] Cycon, H. L., Froese, R. G., Kirsch, W., & Simon, B. (2009). *Schrödinger operators: With application to quantum mechanics and global geometry*. Springer.
- [13] Dykema, K. (1993). On certain free product factors via an extended matrix model. *Journal of functional analysis*, 112(1), 31-60.
- [14] Erdős, L., Knowles, A., Yau, H. T., & Yin, J. (2013). The local semicircle law for a general class of random matrices. *Electronic Journal of Probability*, 18, 1-58.
- [15] Erdős, L., Krüger, T., & Nemish, Y. (2020). Local laws for polynomials of Wigner matrices. *Journal of Functional Analysis*, 278(12), 108507.
- [16] Erdős, L., Krüger, T., & Schröder, D. (2019). Random matrices with slow correlation decay. In *Forum of Mathematics, Sigma* (Vol. 7). Cambridge University Press.
- [17] Erdős, L., Schlein, B., & Yau, H. T. (2009). Local semicircle law and complete delocalization for Wigner random matrices. *Communications in Mathematical Physics*, 287(2), 641-655.
- [18] Erdős, L., Yau, H. T., & Yin, J. (2012). Rigidity of eigenvalues of generalized Wigner matrices. *Advances in Mathematics*, 229(3), 1435-1515.
- [19] Haagerup, U., & Thorbjørnsen, S. (2005). A new application of random matrices: $\text{Ext}(C_{red}^*(\mathbb{F}_2))$ is not a group. *Annals of mathematics*, 162(2), 711-775.
- [20] He, Y., & Knowles, A. (2017). Mesoscopic eigenvalue statistics of Wigner matrices. *The Annals of Applied Probability*, 27(3), 1510-1550.
- [21] He, Y., & Knowles, A. (2021). Fluctuations of extreme eigenvalues of sparse Erdős-Rényi graphs. *Probability Theory and Related Fields*, 180(3), 985-1056.
- [22] Huang, J., Landon, B., & Yau, H. T. (2020). Transition from Tracy-Widom to Gaussian fluctuations of extremal eigenvalues of sparse Erdős-Rényi graphs. *The Annals of Probability*, 48(2), 916-962.
- [23] Khorunzhy, A. M., Khoruzhenko, B. A., & Pastur, L. A. (1996). Asymptotic properties of large random matrices with independent entries. *Journal of Mathematical Physics*, 37(10), 5033-5060.
- [24] Knowles, A., & Yin, J. (2013). The isotropic semicircle law and deformation of Wigner matrices. *Communications on Pure and Applied Mathematics*, 66(11), 1663-1749.
- [25] Knowles, A., & Yin, J. (2017). Anisotropic local laws for random matrices. *Probability Theory and Related Fields*, 169(1), 257-352.
- [26] Lee, J. O., & Schnelli, K. (2015). Edge universality for deformed Wigner matrices. *Reviews in Mathematical Physics*, 27(08), 1550018.
- [27] Lee, J. O., & Schnelli, K. (2018). Local law and Tracy-Widom limit for sparse random matrices. *Probability Theory and Related Fields*, 171(1), 543-616.
- [28] Male, C. (2012). The norm of polynomials in large random and deterministic matrices. *Probability Theory and Related Fields*, 154(3), 477-532.

- [29] Nemish, Y. (2017). Local law for the product of independent non-Hermitian random matrices with independent entries. *Electronic Journal of Probability*, 22, 1-35.
- [30] Nica, A., & Speicher, R. (2006). *Lectures on the combinatorics of free probability* (Vol. 13). Cambridge University Press.
- [31] Nikitopoulos, E. A. (2020). Noncommutative C^k Functions and Fréchet Derivatives of Operator Functions. *arXiv preprint arXiv:2011.03126*.
- [32] Lytova, A., & Pastur, L. (2009). Central limit theorem for linear eigenvalue statistics of random matrices with independent entries. *The Annals of Probability*, 37(5), 1778-1840.
- [33] Parraud, F. (2020). Asymptotic expansion of smooth functions in polynomials in deterministic matrices and iid GUE matrices. *arXiv preprint arXiv:2011.04146*.
- [34] Parraud, F. (2022). On the operator norm of non-commutative polynomials in deterministic matrices and iid Haar unitary matrices. *Probability Theory and Related Fields*, 182(3), 751–806.
- [35] Shlyakhtenko, D., & Skoufranis, P. (2015). Freely independent random variables with non-atomic distributions. *Transactions of the American Mathematical Society*, 367(9), 6267-6291.
- [36] Voiculescu, D. (1991). Limit laws for random matrices and free products. *Inventiones mathematicae*, 104(1), 201-220.
- [37] Yin, S. (2018). Non-commutative rational functions in strong convergent random variables. *Advances in Operator Theory*, 3(1), 178-192.