

# Homomorphisms and principal congruences of bounded lattices. III. The Independence Theorem

G. GRÄTZER

ABSTRACT. A new result of G. Czédli states that for an ordered set  $P$  with at least two elements and a group  $G$ , there exists a bounded lattice  $L$  such that the ordered set of principal congruences of  $L$  is isomorphic to  $P$  and the automorphism group of  $L$  is isomorphic to  $G$ .

I provide an alternative proof utilizing a result of mine with J. Sichler from the late 1960-s.

## 1. Introduction

For a bounded lattice  $L$ , let  $\text{Princ } L$  denote the ordered set of principal congruences of  $L$ . I proved in [14] that  $\text{Princ } L$  can be characterized as a bounded ordered set.<sup>1</sup> Let  $\text{Aut } L$  denote the group of automorphisms of a lattice  $L$ . G. Czédli [3] proved that these two structures are independent.

**Theorem 1** (Independence Theorem). *Let  $P$  be a bounded ordered set with at least two elements and let  $G$  be a group. Then there exists a bounded lattice  $L$  such that  $\text{Princ } L$  is order isomorphic to  $P$  and  $\text{Aut } L$  is group isomorphic to  $G$ .*

I will sketch a new approach to this result.

Note that Czédli proved more than what is stated in Theorem 1. He constructed a lattice  $L$  that is selfdual and of length 16. To prove this, he had to work much harder than I do in this note.

**Notation.** We use the notation as in [15]. You can find the complete

*Part I. A Brief Introduction to Lattices and Glossary of Notation* of [15] at

[tinyurl.com/lattices101](http://tinyurl.com/lattices101)

## 2. Families of rigid lattices

We call a family  $\{K_i \mid i \in I\}$  of bounded lattices *mutually rigid* if  $K_i$  has no  $\{0, 1\}$ -embedding into  $K_j$  for  $i \neq j \in I$ . Of course, all  $K_i$ ,  $i \in I$  are rigid—they only have the trivial automorphism.

---

2010 *Mathematics Subject Classification*: Primary: 06B10.

*Key words and phrases*: bounded lattice, congruence, principal, order, automorphism.

<sup>1</sup>See the references for all the papers that built on this result.

Our proof of the Independence Theorem is based on the following result, see G. Grätzer and J. Sichler [21].

**Theorem 2.** *For every cardinal  $\mathfrak{m}$ , there exists a mutually rigid family  $\{K_i \mid i \in I\}$  of simple bounded lattices satisfying  $|I| = \mathfrak{m}$ .*

Theorem 2 is a very special case of the result of G. Grätzer and J. Sichler [21], which deals with endomorphism semigroups. However, Theorem 2 adds “simple”, and we will now justify it.

Let  $G$  be an infinite connected graph in which every element is in some cycle of odd length. Let  $\text{Lat } G$  denote the free bounded lattice generated by  $G$  with respect to the condition that all edges  $\{x, y\}$  of  $G$  be complemented pairs in  $\text{Lat } G$ . It follows from Corollary 567 in LTF, [13] (originally, C. C. Chen and G. Grätzer [1] and G. Grätzer [11]), that a pair of elements  $\{x, y\} \neq \{0, 1\}$  is complementary in  $\text{Lat } G$  iff  $\{x, y\}$  is an edge of  $G$ . Let  $G$  be a rigid graph. In G. Grätzer and J. Sichler [21], we proved that  $\text{Lat } G$  is a rigid lattice.

Now we extend  $\text{Lat } G$  to a lattice  $\text{Lat}^+ G$ . For every  $a \in L$  with  $0 < a$ , we add an atom  $p_a < a$  so that

$$p_a \vee x = a, \quad (1)$$

$$p_a \wedge x = 0 \quad (2)$$

for all  $0 < x < a$ . We observe that the new lattice  $\text{Lat}^+ G$  is rigid, because we can recognize the elements of  $G$  as complemented elements that are not atoms or dual atoms.

We claim that  $\text{Lat}^+ G$  is simple. Indeed, if  $\alpha > \mathbf{0}$  is a congruence of  $\text{Lat}^+ G$ , then there are  $u < v \in \text{Lat}^+ G$  such that

$$u \equiv v \pmod{\alpha}. \quad (3)$$

By (2),

$$0 \equiv p_v \pmod{\alpha}. \quad (4)$$

It follows by (1) that

$$1 \equiv p_1 \pmod{\alpha} \quad (5)$$

and therefore,

$$0 \equiv x \pmod{\alpha} \quad (6)$$

for all  $x \in \text{Lat}^+ G - \{1, p_1\}$ .

Take an edge  $\{x_1, x_2\}$  of  $G$ . Applying (6) twice, we obtain that

$$0 \equiv x_i \pmod{\alpha} \quad (7)$$

for  $i = 1, 2$ . Joining these two congruences, we conclude that

$$0 \equiv 1 \pmod{\alpha}, \quad (8)$$

that is  $\alpha = \mathbf{1}$ , and so  $\text{Lat}^+ G$  is simple.

So  $\text{Lat}^+ G$  is simple and rigid.

By Z. Hedrlín and A. Pultr [23], we can take a family  $\{G_i \mid i \in I\}$  of mutually rigid, connected graphs with  $|I| = \mathfrak{m}$ , in which every element is in

some cycle of odd length. Then the family  $\{\text{Lat}^+ G_i \mid i \in I\}$  is rigid and simple; in fact, it is mutually rigid for the same reason as we argued before.

### 3. Proving the Independence Theorem

Let  $P$  be the order in Theorem 1. Let 0 and 1 denote the zero and unit of  $P$ , respectively.

We start out by constructing the lattice Frame  $P$ , as in [14], see also [15].

Let Frame  $P$  consist of the elements  $o, i$  and the elements  $a_p, b_p$  for every  $p \in P$ , where  $a_p \neq b_p$  for every  $p \in P^-$  and  $a_0 = b_0, a_1 = b_1$ . These elements are ordered and the lattice operations are formed as in Figure 1.




FIGURE 1. The lattice Frame  $P$

We are going to construct a lattice  $K$  as an extension of Frame  $P$ . For  $p < q \in P^-$ , we add five elements to the sublattice  $\{o, a_p, b_p, a_q, b_q, i\}$  of Frame  $P$  as illustrated in Figure 2. In  $K$ , the principal congruence  $\text{con}(a_p, b_p)$  represents  $p \in P^-$ .

Now we prove Theorem 1.

Let  $P$  and  $G$  be given as in this theorem. The lattice  $K$  we have just constructed satisfies that  $\text{Princ } K \cong P$ . However,  $\text{Aut } K \cong \text{Aut } P$ .

Let  $I = P^-$ . For every  $[o, a_p]$ , we insert the lattice  $K_p$ , provided by Theorem 2, into  $[o, a_p] \subseteq K$  identifying the bounds, obtaining the lattice  $\bar{K}$ . We then have  $\text{Princ } \bar{K} \cong P$  and  $\text{Aut } \bar{K}$  is rigid.

Now take the Frucht lattice, Frucht  $G$  (see [15], [13], R. Frucht [9] and [10]); it is a lattice of length 3, satisfying  $\text{Aut}(\text{Frucht } G) \cong G$ .

We obtain the lattice  $L$  by forming the disjoint union of  $\bar{K}$  and Frucht  $G$  and identifying the bounds, see Figure 3. The diagram does not show  $K_p$ .




FIGURE 2. The sublattice




FIGURE 3. The final step

## REFERENCES

- [1] Chen, C. C. and Grätzer, G.: On the construction of complemented lattices. *J. Algebra* **11**, 56–63 (1969)
- [2] Czédli, G.: Representing a monotone map by principal lattice congruences. *Acta Mathematica Hungarica* **147** 12–18 (2015)
- [3] Czédli, G.: An independence theorem for ordered sets of principal congruences and automorphism groups of bounded lattices. *Acta Sci. Math (Szeged)* **82**, 3–18 (2016)
- [4] Czédli, G.: Large sets of lattices without order embeddings. *Communications in Algebra* **44** 668–679 (2016)
- [5] Czédli, G.: Cometic functors and representing order-preserving maps by principal lattice congruences. *Algebra Universalis*
- [6] Czédli, G.: The ordered set of principal congruences of a countable lattice. *Algebra Universalis* **75**, 351–380 (2016)
- [7] Czédli, G.: Representing some families of monotone maps by principal lattice congruences. *Algebra Universalis* **77**, 51–77 (2017)
- [8] Czédli, G., Grätzer, G., and Lakser, H.: Principal congruences of planar semimodular lattices. *Algebra Universalis*

- [9] Frucht, R.: Herstellung von Graphen mit vorgegebener abstrakter Gruppe. *Compos. Math.* **6**, 239–250 (1938)
- [10] Frucht, R.: Lattices with a given group of automorphisms, *Canad. J. Math.* **2**, 417–419 (1950)
- [11] Grätzer, G.: A reduced free product of lattices. *Fund. Math.* **73**, 21–27 (1971/72)
- [12] Grätzer, G.: The Congruences of a Finite Lattice. A *Proof-by-Picture* Approach. Birkhäuser, Boston (2006)
- [13] Grätzer, G.: Lattice Theory: Foundation. Birkhäuser Verlag, Basel (2011)
- [14] Grätzer, G.: The order of principal congruences of a bounded lattice. *Algebra Universalis* **70**, 95–105 (2013)
- [15] Grätzer, G.: The Congruences of a Finite Lattice. A *Proof-by-Picture* Approach, second edition. Birkhäuser, Basel (2016)
- [16] Grätzer, G.: Homomorphisms and principal congruences of bounded lattices. I. Isotone maps of principal congruences. *Acta Sci. Math. (Szeged)* **82**, 353–360 (2016)
- [17] Grätzer, G.: Homomorphisms and principal congruences of bounded lattices. II. Sketching the proof for sublattices. *Algebra Universalis*
- [18] Grätzer, G. and Lakser, H.: Notes on the set of principal congruences of a finite lattice. I. Some preliminary results.
- [19] Grätzer, G. and Lakser, H.: Notes on the set of principal congruences of a finite lattice. II. A combinatorial condition for representability.
- [20] Grätzer, G. and Lakser, H.: Notes on the set of principal congruences of a finite lattice. III. Revisiting the representation theorem of finite distributive lattices with principal congruences.
- [21] Grätzer, G. and Sichler, J.: On the endomorphism semigroup (and category) of bounded lattices. *Pacif. J. Math.* **35**, 639–647 (1970)
- [22] Grätzer, G. and Wehrung, F. eds.: Lattice Theory: Special Topics and Applications. Volume 1. Birkhäuser Verlag, Basel, 2014
- [23] Hedrlín, Z. and Pultr, A.: Relations (graphs) with given finitely generated semigroups. *Monatsh. Math.* **68**, 213–217 (1964)

G. GRÄTZER

Department of Mathematics, University of Manitoba, Winnipeg, MB R3T 2N2, Canada  
e-mail, G. Grätzer: [gratzer@me.com](mailto:gratzer@me.com)