

AN EXAMPLE OF A DOMAIN IN A COMPLEX FLAG MANIFOLD

NOBUTAKA BOUMUKI

ABSTRACT. This paper shows an example of a connected open neighborhood of a compact connected complex submanifold in a complex flag manifold.

1. A COUNTER-EXAMPLE

We shall give an example of a complex flag manifold $X = G_{\mathbb{C}}/B$ of simply connected complex semisimple Lie group $G_{\mathbb{C}}$, a closed connected complex (Lie) subgroup $Q \subsetneq G_{\mathbb{C}}$, a compact connected complex submanifold Z in X , and a domain D in X such that

- (c1) $B \subset Q$,
- (c2) Z is contained in a fiber of the natural projection $\text{Pr} : G_{\mathbb{C}}/B \rightarrow G_{\mathbb{C}}/Q$, $gB \mapsto gQ$,
- (c3) $Z \subset D$, and
- (c4) $\mathcal{O}(D) \cong \mathbb{C}$, i.e., all holomorphic functions on D are constant.

This (c2) and Definition 1 in [6, p.92] imply that Z is *degenerate* in X , and hence (c2) and (c4) tell us that the two conditions (1) and (16) in Theorem 2 [6, pp.92–93] are not equivalent to each other. That has effects on some papers (e.g. [2], [3], [5], [7], etc).

Using the notation in Helgason [4, pp.444–446], we give

Example 1. Let $G_{\mathbb{C}} := Sp(2, \mathbb{C})$, $G := Sp(1, 1)$, and $\mathfrak{g}_{\mathbb{C}} := \mathfrak{sp}(2, \mathbb{C})$. Define two subalgebras $\mathfrak{b}, \mathfrak{q} \subset \mathfrak{g}_{\mathbb{C}}$ and a compact connected subgroup $S \subset G$ by

$$\mathfrak{b} := \left\{ \left(\begin{array}{cccc} z_1 & 0 & 0 & 0 \\ w_1 & z_2 & 0 & 0 \\ w_2 & w_3 & -z_1 & -w_1 \\ w_3 & w_4 & 0 & -z_2 \end{array} \right) \middle| z_1, z_2, w_1, w_2, w_3, w_4 \in \mathbb{C} \right\},$$

$$\mathfrak{q} := \left\{ \left(\begin{array}{cccc} z_1 & 0 & 0 & 0 \\ w_1 & z_2 & 0 & z_3 \\ w_2 & w_3 & -z_1 & -w_1 \\ w_3 & z_4 & 0 & -z_2 \end{array} \right) \middle| z_1, z_2, z_3, z_4, w_1, w_2, w_3 \in \mathbb{C} \right\};$$

$$S := \left\{ \left(\begin{array}{cccc} x & 0 & 0 & 0 \\ 0 & y_1 & 0 & y_2 \\ 0 & 0 & 1/x & 0 \\ 0 & y_3 & 0 & y_4 \end{array} \right) \middle| x \in U(1), \begin{pmatrix} y_1 & y_2 \\ y_3 & y_4 \end{pmatrix} \in Sp(1) \right\} (\cong U(1) \times Sp(1)).$$

Denote by B and Q the normalizers of \mathfrak{b} and \mathfrak{q} in $G_{\mathbb{C}}$, respectively, and note that B is a Borel subgroup of $G_{\mathbb{C}}$ (because \mathfrak{b} coincides with the direct sum $\mathfrak{h} \oplus \bigoplus_{\alpha < 0} \mathfrak{g}^{\alpha}$ of the Cartan subalgebra \mathfrak{h} and all the negative-root subspaces $\mathfrak{g}^{\alpha} \subset \mathfrak{g}_{\mathbb{C}}$ corresponding to [4, pp.189–190], The algebra

2020 *Mathematics Subject Classification.* Primary 32M10; Secondary 17B20.

Key words and phrases. domain, complex flag manifold, degenerate, constant holomorphic function.

$\mathfrak{c}_n = \mathfrak{sp}(n, \mathbb{C})$ in case $n = 2$]) and that Q is a parabolic subgroup of $G_{\mathbb{C}}$ satisfying (c1) $B \subset Q$. In addition, denote by Z (resp. D) the orbit of the subgroup $S \subset G_{\mathbb{C}}$ (resp. $G \subset G_{\mathbb{C}}$) through the origin $o \in G_{\mathbb{C}}/B$, where we remark that $G_{\mathbb{C}}$ holomorphically acts on $G_{\mathbb{C}}/B$ in a natural way. Then, (c3) $Z \subset D$ holds, and one obtains

- $$(i) \ S \cap B = G \cap B = \left\{ \left(\begin{array}{cccc} x & 0 & 0 & 0 \\ 0 & y & 0 & 0 \\ 0 & 0 & 1/x & 0 \\ 0 & 0 & 0 & 1/y \end{array} \right) \middle| x, y \in U(1) \right\} \cong U(1) \times U(1),$$
- (ii) $Z = SB/B \cong S/(S \cap B) \cong (U(1) \times Sp(1))/(U(1) \times U(1))$, $\dim_{\mathbb{R}} Z = 2$,
(iii) $D = GB/B \cong G/(G \cap B) \cong Sp(1, 1)/(U(1) \times U(1))$, $\dim_{\mathbb{R}} D = 8 = \dim_{\mathbb{R}} G_{\mathbb{C}}/B$,
(iv) $SB \subset Q$,
(v) $Z = SB/B \subset Q/B = \text{Pr}^{-1}(o)$

by a direct computation. This (v) implies that (c2) holds, and the (iii) yields (c4) $\mathcal{O}(D) \cong \mathbb{C}$ because D is a flag domain in $G_{\mathbb{C}}/B$ and the Lie group $G = Sp(1, 1)$ is not of Hermitian type (e.g. the proof of Lemma 3.4 in [1, p.5]).

REFERENCES

- [1] N. Boumuki, *Homogeneous pseudo-Kähler manifolds having non-constant holomorphic functions*, In: N. Koike, M. S. Tanaka, and K. Baba (eds.) Proceedings of the conference “Submanifold Geometry and Lie Group Actions 2020”, pp.1–12. URL: <https://www.rs.tus.ac.jp/kurando.baba/conference/record/2020/boumuki-abstract.pdf>
- [2] N. Boumuki, *On the dimensions of vector spaces concerning holomorphic vector bundles over elliptic orbits*, Tsukuba J. Math. (to appear).
- [3] T. Hayama, A. Huckleberry, and Q. Latif, *Pseudoconcavity of flag domains: the method of supporting cycles*, Math. Ann., **375**, (2019), 671–685.
- [4] S. Helgason, *Differential geometry, Lie groups, and symmetric spaces*, Corrected reprint of the 1978 original, Graduate Studies in Mathematics 34, American Mathematical Society, Providence, RI, 2001.
- [5] A. Huckleberry, *Cycle connectivity and automorphism groups of flag domains*, Developments and retrospectives in Lie theory, pp.113–126, Dev. Math., **37**, Springer, Cham, 2014.
- [6] J. Kollár, *Neighborhoods of subvarieties in homogeneous spaces*, Hodge theory and classical algebraic geometry, pp.91–107, Contemp. Math., **647**, Amer. Math. Soc., Providence, RI, 2015.
- [7] Y.-C. Tu, *Higher ramification loci over homogeneous spaces*, Manuscripta Math., **148**, (2015), 235–241.

DIVISION OF MATHEMATICAL SCIENCES, FACULTY OF SCIENCE AND TECHNOLOGY
OITA UNIVERSITY, 700 DANNOHARU, OITA-SHI, OITA 870-1192, JAPAN
Email address: boumuki@oita-u.ac.jp