

CLASSIFICATION OF COHEN–MACAULAY t –SPREAD LEXSEGMENT IDEALS VIA SIMPLICIAL COMPLEXES

Marilena Crupi, Antonino Ficarra

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCES, PHYSICS AND EARTH SCIENCES,
UNIVERSITY OF MESSINA, VIALE FERDINANDO STAGNO D’ALCONTRES 31, 98166 MESSINA, ITALY

E-mail addresses: mcrupi@unime.it; antficarra@unime.it

Abstract

We study the minimal primary decomposition of completely t –spread lexsegment ideals via simplicial complexes. We determine some algebraic invariants of such a class of t –spread ideals. Hence, we classify all t –spread lexsegment ideals which are Cohen–Macaulay.

INTRODUCTION

In combinatorial commutative algebra, one often associates to a combinatorial object X a suitable monomial ideal I_X in a polynomial ring $S = K[x_1, \dots, x_n]$ in finitely many variables with coefficients in a field K . Distinguished combinatorial properties of the object X are reflected by algebraic properties of the ideal I_X , and conversely. One says that X is Cohen–Macaulay if the ring S/I_X is Cohen–Macaulay. A typical task is to classify all Cohen–Macaulay objects X of a given class. For instance, in combinatorics, to a simplicial complex Δ on the vertex set $[n]$ one associates the Stanley–Reisner ideal I_Δ , *i.e.*, the ideal generated by all squarefree monomials $x_{i_1} \cdots x_{i_r}$, $r \leq n$, such that $\{i_1, \dots, i_r\} \notin \Delta$. A simplicial complex Δ is said Cohen–Macaulay if the ideal I_Δ is Cohen–Macaulay, *i.e.*, S/I_Δ is a Cohen–Macaulay ring. To classify all Cohen–Macaulay simplicial complexes is an hopeless task. However, many results can be found, for instance, in [18].

The aim of this article is to classify all Cohen–Macaulay t –spread lexsegment ideals for $t \geq 1$.

Let $S = K[x_1, \dots, x_n]$ be the polynomial ring in n variables with coefficients over a fixed field K . Let $t \geq 0$ be an integer. A monomial $u = x_{i_1} x_{i_2} \cdots x_{i_d}$ is t –spread if

Keywords: Betti numbers, t –spread ideals, primary decomposition, simplicial complexes, Cohen–Macaulay rings.

2020 Mathematics Subject Classification: 05E40, 13B25, 13D02, 16W50.

$i_{j+1} - i_j \geq t$, for all $j = 1, \dots, d - 1$. If $t \geq 1$, a t -spread monomial is a squarefree monomial. A monomial ideal is t -spread if it is generated by t -spread monomials. Such a topic introduced in [13] has immediately captured the attention of many algebraists (see, for instance, [1, 2, 4, 3, 5, 6, 11, 15]). Let $t \geq 1$. A t -spread ideal I of S is said a t -spread lexsegment ideal if for all t -spread monomials $u \in I$ and all t -spread monomials $v \in S$ with $\deg(u) = \deg(v)$ and such that $v \geq_{\text{slex}} u$, then $v \in I$ [6], where \geq_{slex} is the squarefree lexicographic order [8, 18]. Therefore, for $t = 1$, such a definition coincides with the *classical* definition of squarefree lexsegment ideal [8]. Let $t \geq 1$ and let $M_{n,d,t}$ be the set of all t -spread monomials of S of degree d . Assume that $M_{n,d,t}$ is endowed with the squarefree lexicographic order \geq_{slex} . Let $u, v \in M_{n,d,t}$. If $u \geq_{\text{slex}} v$, the set $\mathcal{L}_t(u, v) = \{w \in M_{n,d,t} : u \geq_{\text{slex}} w \geq_{\text{slex}} v\}$ is called an *arbitrary t -spread lexsegment*. If u is the maximum monomial of $M_{n,d,t}$, or v is the minimum monomial of $M_{n,d,t}$, we say that $\mathcal{L}_t(u, v)$ is an *initial* or a *final t -spread lexsegment*, respectively. The ideal generated by an arbitrary t -spread lexsegment is called *arbitrary t -spread lexsegment ideal*. It is clear that the notion of initial t -spread lexsegment ideal coincides with the notion of t -spread lexsegment ideal in [6]. An arbitrary t -spread lexsegment ideal $I = (\mathcal{L}_t(u, v))$ is said *completely t -spread lexsegment* if $I = (\mathcal{L}_t(\max(M_{n,d,t}), v)) \cap (\mathcal{L}_t(u, \min(M_{n,d,t})))$, [15]. One can easily observe that every initial and every final t -spread lexsegment ideal is a completely t -spread lexsegment, but not all arbitrary t -spread lexsegment ideals are completely t -spread lexsegment ideals. In [15], the authors characterized all completely t -spread lexsegment ideals and classified all completely t -spread lexsegment ideals with a linear resolution. The case $t = 0$ has been faced by De Negri and Herzog in [10].

In this article we determine the minimal primary decomposition of completely t -spread lexsegment ideals and some algebraic invariants of such a class of ideals by tools from the simplicial complex theory. Hence, we classify all t -spread lexsegment ideals which are Cohen–Macaulay. The case $t = 0$ has been considered in [14] and the case $t = 1$ has been analyzed in [9].

The article is organized as follows. Section 1 contains preliminary notions and results. In Section 2, we explicitly compute the minimal primary decomposition for initial and final t -spread lexsegment ideals (Theorems 2.1, 2.7). As a consequence we are able to compute the standard primary decomposition for the class of completely t -spread lexsegments (Theorem 2.12). Moreover, we obtain formulas for the Krull dimension, the projective dimension and the depth of S/I when I is an initial or a final t -spread lexsegment ideal. In Section 3, we classify all t -spread lexsegment ideals which are Cohen–Macaulay by the results stated in Section 2. The notion of Betti splitting will be a crucial tool. Finally, Section 4 contains our conclusions and perspectives.

All the examples are constructed by means of *Macaulay2* packages one of which developed in [1]. Furthermore, some functions described in [1] have allowed us to test

classes of Cohen–Macaulay t –spread lexsegment ideals whose determination has been fundamental for the development of Section 3.

1. PRELIMINARIES

Let $S = K[x_1, \dots, x_n]$ be the polynomial ring in n indeterminates with coefficients in a field K . S is a \mathbb{N} –graded K –algebra with $\deg(x_i) = 1$ for all i .

If I is a monomial ideal of S , we denote by $G(I)$ the unique minimal set of monomial generators of I . Moreover, we set $G(I)_j = \{u \in G(I) : \deg(u) = j\}$.

Given a non empty subset $A \subseteq [n]$, we set $\mathbf{x}_A = \prod_{i \in A} x_i$. Moreover, we set $\mathbf{x}_\emptyset = 1$. Every monomial $w \in S$ can be written as $w = x_{i_1} x_{i_2} \cdots x_{i_d}$, with sorted indexes $1 \leq i_1 \leq i_2 \leq \cdots \leq i_d \leq n$. We define the *support* of w as follows:

$$\text{supp}(w) = \{j : x_j \text{ divides } w\} = \{i_1, i_2, \dots, i_d\}.$$

With the notation above, $\min(w) = \min\{i : i \in \text{supp}(u)\} = i_1$, $\max(w) = \max\{i : i \in \text{supp}(u)\} = i_d$. Moreover, we set $\max(1) = \min(1) = 0$. If $i_1 \neq i_2 \neq \dots \neq i_d$ we say that u is *squarefree*. In such a case, u can be written as $u = \mathbf{x}_A$ with $A = \text{supp}(u)$.

We quote next definition from [13].

Definition 1.1 Given $n \geq 1, t \geq 0$ and $u = x_{i_1} x_{i_2} \cdots x_{i_d} \in S$, with $1 \leq i_1 \leq i_2 \leq \dots \leq i_d \leq n$, we say that u is t –spread if $i_{j+1} - i_j \geq t$, for all $j = 1, \dots, d-1$. We say that a monomial ideal I of S is t –spread if it is generated by t –spread monomials.

We note that any monomial ideal of S is a 0–spread monomial ideal, and any squarefree monomial ideal of S is a 1–spread monomial ideal.

Let $n, d, t \geq 1$. We denote by $M_{n,d,t}$ the set of all t –spread monomials of S of degree d . Throughout this paper we tacitly assume that $n \geq 1 + (d-1)t$, otherwise $M_{n,d,t} = \emptyset$. From [13], we have that $|M_{n,d,t}| = \binom{n-(t-1)(d-1)}{d}$. The monomial ideal generated by $M_{n,d,t}$ is denoted by $I_{n,d,t}$ and is called the t –spread Veronese ideal of degree d of the polynomial ring $S = K[x_1, \dots, x_n]$ [13].

From now on we consider $t \geq 1$ and assume that $M_{n,d,t}$ is endowed with the *squarefree lexicographic order*, \geq_{slex} , [18]. Recall that given $u = x_{i_1} x_{i_2} \cdots x_{i_d}, v = x_{j_1} x_{j_2} \cdots x_{j_d} \in M_{n,d,t}$, with $1 \leq i_1 < i_2 < \cdots < i_d \leq n$, $1 \leq j_1 < j_2 < \cdots < j_d \leq n$, then $u >_{\text{slex}} v$ if

$$i_1 = j_1, \dots, i_{s-1} = j_{s-1} \text{ and } i_s < j_s$$

for some $1 \leq s \leq d$.

Let U be a not empty subset of $M_{n,d,t}$. We denote by $\max(U)$ ($\min(U)$, respectively) the maximum (minimum, respectively) monomial $w \in U$, with respect to \geq_{slex} . One has

$$\begin{aligned}\max(M_{n,d,t}) &= x_1 x_{1+t} x_{1+2t} \cdots x_{1+(d-1)t}, \\ \min(M_{n,d,t}) &= x_{n-(d-1)t} x_{n-(d-2)t} \cdots x_{n-t} x_n.\end{aligned}$$

In [15], the following definitions have been introduced.

Definition 1.2 Let $u, v \in M_{n,d,t}$, $u \geq_{\text{slex}} v$. The set

$$\mathcal{L}_t(u, v) = \{w \in M_{n,d,t} : u \geq_{\text{slex}} w \geq_{\text{slex}} v\}$$

is called an *arbitrary t -spread lexsegment set*, or simply a *t -spread lexsegment set*. The set

$$\mathcal{L}_t^i(v) = \{w \in M_{n,d,t} : w \geq_{\text{slex}} v\} = \mathcal{L}_t(\max(M_{n,d,t}), v)$$

is called an *initial t -spread lexsegment* and the set

$$\mathcal{L}_t^f(u) = \{w \in M_{n,d,t} : w \leq_{\text{slex}} u\} = \mathcal{L}_t(u, \min(M_{n,d,t}))$$

is called a *final t -spread lexsegment*.

One can observe that, if $u, v \in M_{n,d,t}$, with $u \geq_{\text{slex}} v$, then $\mathcal{L}_t(u, v) = \mathcal{L}_t^i(v) \cap \mathcal{L}_t^f(u)$.

Definition 1.3 A t -spread monomial ideal I of S is called an *arbitrary t -spread lexsegment ideal*, or simply a *t -spread lexsegment ideal* if it is generated by an (arbitrary) t -spread lexsegment.

A t -spread lexsegment ideal $I = (\mathcal{L}_t(u, v))$ is a *completely t -spread lexsegment ideal* if

$$I = J \cap T,$$

where $J = (\mathcal{L}_t^i(v))$ and $T = (\mathcal{L}_t^f(u))$.

In [15], all completely t -spread lexsegment ideals have been characterized.

Now, we recall some important notions that will be useful in the sequel.

A t -spread monomial ideal I is a *t -spread strongly stable ideal* if for all t -spread monomial $u \in I$, all $j \in \text{supp}(u)$ and all $1 \leq i < j$ such that $x_i(u/x_j)$ is t -spread, it follows that $x_i(u/x_j) \in I$.

Every initial t -spread lexsegment ideal I of S is a t -spread strongly stable ideal. Hence, in order to compute the graded Betti numbers of I one may use the Ene, Herzog, Qureshi's formula ([13, Corollary 1.12]):

$$\beta_{i,i+j}(I) = \sum_{u \in G(I)_j} \binom{\max(u) - t(j-1) - 1}{i}. \quad (1)$$

A t -spread final lexsegment ideal I of S is also a t -spread strongly stable ideal, but with the order of the variables reversed, *i.e.*, $x_n > x_{n-1} > \cdots > x_1$. Hence, in order to compute their graded Betti numbers one may use the following *modified* Ene, Herzog, Qureshi's formula:

$$\beta_{i,i+j}(I) = \sum_{u \in G(I)_j} \binom{n - \min(u) - t(j-1)}{i}. \quad (2)$$

Let $>_{\text{lex}}$ be the usual *lexicographic order* on S , with $x_1 > x_2 > \cdots > x_n$, [18]. The next theorem collects some results from [15] that will be pivotal for the aim of this article.

Theorem 1.4 [15]. *Given $n, d, t \geq 1$, let $u = x_{i_1}x_{i_2} \cdots x_{i_d}$ and $v = x_{j_1}x_{j_2} \cdots x_{j_d}$ be t -spread monomials of degree d of S such that $u \geq_{\text{slex}} v$.*

- (1) *Let $I = (\mathcal{L}_t(u, v))$. The following conditions are equivalent:*
 - (a) *I is a completely t -spread lexsegment ideal, *i.e.*, $I = J \cap T$;*
 - (b) *for every $w \in M_{n,d,t}$ with $w <_{\text{slex}} v$ there exists an integer $s > i_1$ such that x_s divides w and $x_{i_1}(w/x_s) \leq_{\text{lex}} u$.*
- (2) *Let $I = (\mathcal{L}_t(u, v))$ be a completely t -spread lexsegment ideal with $\min(v) > \min(u) = 1$. I has a linear resolution if and only if one of the following conditions hold:*
 - (i) $i_2 = 1 + t$;
 - (ii) $i_2 > 1 + t$ and for the largest $w \in M_{n,d,t}$, $w <_{\text{lex}} v$, we have $x_1(w/x_{\max(w)}) \leq_{\text{lex}} x_1x_{i_2-t}x_{i_3-t} \cdots x_{i_d-t}$.
- (3) *Let $I = (\mathcal{L}_t(u, v))$ be a completely t -spread lexsegment ideal and suppose $I = (\mathcal{L}_t(u, v))$ has a linear resolution. Then, for all $i \geq 0$,*

$$\beta_i(I) = \sum_{w \in \mathcal{L}_i^f(u)} \binom{n - \min(w) - (d-1)t}{i} - \sum_{\substack{w \in \mathcal{L}_i^f(v) \\ w \neq v}} \binom{\max(w) - (d-1)t - 1}{i}.$$

We close the section recalling some notions about simplicial complexes.

Given $n \in \mathbb{N}$, we set $[n] = \{1, 2, \dots, n\}$. We recall that a *simplicial complex* on the vertex set $[n]$ is a family of subsets of $[n]$ such that

- $\{i\} \in \Delta$ for all $i \in [n]$, and

- if $F \subseteq [n]$, $G \subseteq F$, we have $G \in \Delta$.

The dimension of Δ is the number $d = \max\{|F| - 1 : F \in \Delta\}$. Any $F \in \Delta$ is called a *face* and $|F| - 1$ is the *dimension* of F . A *facet* of Δ is a maximal face with respect to the inclusion. The set of facets of Δ is denoted by $\mathcal{F}(\Delta)$.

It is well known that for any squarefree ideal (1–spread ideal) I of S there exists a unique simplicial complex Δ on $[n]$ such that $I = I_\Delta$, where

$$I_\Delta = (\mathbf{x}_F : F \subseteq [n], F \notin \Delta)$$

is the Stanley–Reisner ideal of Δ [18]. In the sequel when we say that a simplicial complex Δ is associated to a squarefree ideal I of S , we mean that $I_\Delta = I$.

A simplicial complex Δ on $[n]$ is called *pure* if its facets have the same dimension. We say that Δ is *Cohen–Macaulay* if $K[\Delta] = S/I_\Delta = K[x_1, \dots, x_n]/I_\Delta$ is a Cohen–Macaulay ring. If Δ is Cohen–Macaulay then Δ is pure (see, for instance, [18]).

It is well known that any monomial ideal (not necessarily squarefree) has a unique minimal primary decomposition. We refer to it as the *standard primary decomposition*. Moreover, for a squarefree monomial ideal, its standard primary decomposition is $I = \bigcap_{\mathfrak{p} \in \text{Min}(I)} \mathfrak{p}$, [18, Corollary 1.3.6], where $\text{Min}(I)$ is the set of minimal primes of I . Every minimal prime is a monomial prime ideal, *i.e.*, $\mathfrak{p} = (x_{i_1}, x_{i_2}, \dots, x_{i_s})$, for some $A = \{i_1 < i_2 < \dots < i_s\} \subseteq [n]$. To denote such an ideal we also use the notation \mathfrak{p}_A .

Proposition 1.5 [18, Lemma 1.5.4] *The standard primary decomposition of I_Δ is*

$$I_\Delta = \bigcap_{F \in \mathcal{F}(\Delta)} \mathfrak{p}_{[n] \setminus F}.$$

Given two positive integers $j \geq k$, we set $[j, k] = \{\ell \in \mathbb{N} : j \leq \ell \leq k\}$ and we associate to a t –spread monomial two special sets of integers.

Let $t \geq 1$ and let $w = x_{\ell_1} x_{\ell_2} \cdots x_{\ell_d}$ be a t –spread monomial of S of degree d .

If $\max(w) \leq n + 1 - t$, we define the t –spread support of w as follows:

$$\text{supp}_t(w) = \bigcup_{r=1}^d [\ell_r, \ell_r + (t - 1)]; \quad (3)$$

whereas, if $\min(w) \geq t$, we define the t –spread cosupport of w as follows:

$$\text{cosupp}_t(w) = \bigcup_{r=1}^d [\ell_r - (t - 1), \ell_r]. \quad (4)$$

Moreover, we set $\text{supp}_t(1) = \text{cosupp}_t(1) = \emptyset$.

Remark 1.6 Let $w = x_2x_5x_{10}x_{13} \in S = K[x_1, \dots, x_{13}]$ be a 3–spread monomial. One can note that $\max(w) = 13 > 13 + 1 - 3 = 11$. Furthermore, $\text{supp}_3(w/x_{\max(w)}) = \{2, 3, 4, 5, 6, 7, 10, 11, 12\}$ and $\text{cosupp}_3(w/x_{\min(w)}) = \{3, 4, 5, 8, 9, 10, 11, 12, 13\}$.

Observe that for all $w \in M_{n,d,t}$, we can always compute $\text{supp}_t(w/x_{\max(w)})$ and $\text{cosupp}_t(w/x_{\min(w)})$.

In [13, Theorem 2.3], Ene, Herzog, Qureshi computed the standard primary decomposition of the t –spread Veronese ideal of degree d , $I_{n,d,t}$,

$$I_{n,d,t} = \bigcap_{D \in \mathcal{D}} \mathfrak{p}_{[n] \setminus D}, \quad (5)$$

where

$$\mathcal{D} = \left\{ \bigcup_{r=1}^{d-1} [\ell_r, \ell_r + (t-1)] \subseteq [n] : \ell_{r+1} - \ell_r \geq t, r = 1, \dots, d-2 \right\}.$$

Using our notation, each set $D \in \mathcal{D}$ may be written as $\text{supp}_t(x_{\ell_1} \cdots x_{\ell_{d-1}})$, with $x_{\ell_1} \cdots x_{\ell_{d-1}} \in M_{n+1-t, d-1, t}$. Hence, the primary decomposition (5) becomes

$$I_{n,d,t} = \bigcap_{w \in M_{n+1-t, d-1, t}} \mathfrak{p}_{[n] \setminus \text{supp}_t(w)}. \quad (6)$$

In particular, $I_{n,d,t}$ is an *height–unmixed* ideal, *i.e.*, all its minimal primes have the same height, $n - (d-1)t$ [13, Theorem 2.3]. Furthermore, $I_{n,d,t}$ is a completely t –spread lexsegment ideal [15].

2. THE PRIMARY DECOMPOSITION OF COMPLETELY t –SPREAD LEXSEGMENT IDEALS

In this section we study the primary decomposition of a completely t –spread lexsegment ideal of $S = K[x_1, \dots, x_n]$. The case $t = 1$ has been analyzed in [19]. The sets defined in (3) and (4) will be pivotal for our aim.

We start the section with some comments and remarks.

Let $n, d, t \geq 1$, $u = x_{i_1}x_{i_2} \cdots x_{i_d}$ and $v = x_{j_1}x_{j_2} \cdots x_{j_d}$ monomials of $M_{n,d,t}$, with $u \geq_{\text{slex}} v$. Set $L = \mathcal{L}_t(u, v)$, $I = (L)$. We assume that I is a completely t –spread lexsegment ideal, *i.e.*, $I = J \cap T$, with $J = (\mathcal{L}_t^i(v))$ and $T = (\mathcal{L}_t^f(u))$.

- If $u = v$, $I = (L) = (u)$ is a principal ideal, and its standard primary decomposition is $I = (x_{i_1}) \cap \cdots \cap (x_{i_d})$. Therefore, we may assume $u >_{\text{slex}} v$.

- If $\deg(u) = \deg(v) = d = 1$, then $I = (L) = (x_{i_1}, x_{i_1+1}, \dots, x_{j_1-1}, x_{j_1}) = \mathfrak{p}_{[i_1, j_1]}$ is the standard primary decomposition of I . Thus, we may assume $d \geq 2$.
- If I is initial, *i.e.*, $I = J = (\mathcal{L}_t^i(v))$, we may assume $\min(v) = j_1 \geq 2$, otherwise

$$J = (x_1) \cap (\mathcal{L}_t(x_{1+t} \cdots x_{1+(d-1)t}, v/x_1)),$$

and to determine the standard primary decomposition of J , it suffices to determine that of $(\mathcal{L}_t(x_{1+t} \cdots x_{1+(d-1)t}, v/x_{j_1}))$ which is an initial t -spread lexsegment ideal in the polynomial ring $K[x_{1+t}, \dots, x_n]$ with fewer indeterminates than S .

- If I is final, *i.e.*, $I = T = (\mathcal{L}_t^f(u))$, we can assume $\min(u) = i_1 = 1$. Indeed, if $i_1 > 1$, none of the variables x_1, \dots, x_{i_1-1} divides any minimal monomial generator of T . So computing the standard primary decomposition of T is equivalent to computing that of $T \cap K[x_{i_1}, \dots, x_n]$ in the polynomial ring $K[x_{i_1}, \dots, x_n]$ with fewer indeterminates than S .

Hence, from now on, we assume that $I = (\mathcal{L}_t(u, v)) = J \cap T$, with $u >_{\text{slex}} v$, $\min(u) = 1$, $\min(v) \geq 2$ and $\deg(u) = \deg(v) = d \geq 2$.

To compute the standard primary decomposition of $I = J \cap T$, we can proceed as follows. Firstly, we determine the decomposition of J , and if $I = J$ we are done. Secondly, we determine that of T , and if $I = T$ we are done. Finally, knowing the primary decompositions of J and T we take into account the intersection $J \cap T = I$, and deleting the non minimal primes, we obtain the standard primary decomposition of I in the general case.

2.1. The initial t -spread lexsegment case

In this subsection we analyze the case of the standard primary decomposition of an initial t -spread lexsegment ideal of $S = K[x_1, \dots, x_n]$.

Theorem 2.1 *Let $v = x_{j_1} x_{j_2} \cdots x_{j_d}$ be a t -spread monomial of degree $d \geq 2$ of S with $j_1 \geq 2$ and let $J = (\mathcal{L}_t^i(v))$ be an initial t -spread lexsegment ideal of S . Then, the standard primary decomposition of J is*

$$J = \bigcap_{p=1}^d \mathfrak{p}_{F_p} \cap \bigcap_{F \in \mathcal{F}} \mathfrak{p}_{[n] \setminus F},$$

where

$$\begin{aligned} F_p &= [j_p] \setminus \text{supp}_t(x_{j_1} x_{j_2} \cdots x_{j_{p-1}}), \quad p = 1, \dots, d, \\ \mathcal{F} &= \{ \text{supp}_t(w) : w \in M_{n+1-t, d-1, t}, \quad w >_{\text{slex}} v/x_{\max(v)} \}. \end{aligned}$$

Proof. Let Δ be the simplicial complex on the vertex set $[n]$ associated to J . From Proposition 1.5, we have to prove that the facets of Δ are exactly the sets $[n] \setminus F_p$, $p = 1, \dots, d$, together with the sets of \mathcal{F} , *i.e.*,

$$\mathcal{F}(\Delta) = \{[n] \setminus F_p : p = 1, \dots, d\} \cup \mathcal{F}. \quad (7)$$

Observe that if $G \in \Delta$, $G \neq \emptyset$, then G is a facet of Δ if and only if $G \cup \{i\} \notin \Delta$, for all $i \in [n] \setminus G$, *i.e.*, if and only if $\mathbf{x}_{G \cup \{i\}} \in J$, for all $i \in [n] \setminus G$.

We show that each $F \in \mathcal{F}$ is a facet of Δ . Let $F \in \mathcal{F}$, then

$$\begin{aligned} F &= \text{supp}_t(x_{\ell_1} \cdots x_{\ell_{d-1}}) = \bigcup_{r=1}^{d-1} [\ell_r, \ell_r + (t-1)] \\ &= \{\ell_1, \ell_1 + 1, \dots, \ell_1 + (t-1), \dots, \ell_{d-1}, \ell_{d-1} + 1, \dots, \ell_{d-1} + (t-1)\}, \end{aligned}$$

with $x_{\ell_1} \cdots x_{\ell_{d-1}} \in M_{n+1-t, d-1, t}$ and $x_{\ell_1} \cdots x_{\ell_{d-1}} >_{\text{slex}} v/x_{\max(v)} = x_{j_1} \cdots x_{j_{d-1}}$.

Clearly, $\mathbf{x}_F \notin J$, since \mathbf{x}_F is not a multiple of any t -spread monomial of degree d of S . To prove that F is a facet, it suffices to show that $\mathbf{x}_{F \cup \{i\}} \in J$, for all $i \in [n] \setminus F$.

Let $i \in [n] \setminus F$. We have:

- if $i < \ell_1$, then $\mathbf{x}_{F \cup \{i\}}$ is divided by the t -spread monomial $x_i x_{\ell_1 + (t-1)} \cdots x_{\ell_{d-1} + (t-1)}$;
- if $\ell_j + (t-1) < i < \ell_{j+1}$, for some $1 \leq j \leq d-2$, then $\mathbf{x}_{F \cup \{i\}} = \mathbf{x}_F x_i$ is divided by the t -spread monomial $(\prod_{r=1}^j x_{\ell_r}) x_i (\prod_{r=j+1}^{d-1} x_{\ell_r + (t-1)})$;
- if $\ell_{d-1} + (t-1) < i$, then $\mathbf{x}_{F \cup \{i\}}$ is divided by the monomial $x_{\ell_1} \cdots x_{\ell_{d-1}} x_i$.

More in detail, in every case, $\mathbf{x}_{F \cup \{i\}}$ is a multiple of a t -spread monomial of degree d strictly greater than v , with respect to $>_{\text{slex}}$. Therefore $\mathbf{x}_{F \cup \{i\}} \in J$ and F is a facet.

Now, we determine all the facets of Δ . Let $G \in \Delta$ and define the following integers:

$$\ell_1 = \min(G), \text{ and}$$

$$\ell_i = \min \{r \in G : r \geq \ell_{i-1} + t\}, \text{ for } i \geq 2.$$

Let ℓ_k be the last element of the sequence $\ell_1 < \ell_2 < \cdots$.

If $\ell_i \leq j_i$ for all i , then $k \leq d-1$, otherwise $x_{\ell_1} x_{\ell_2} \cdots x_{\ell_d} \in J$. Against the fact that $G \in \Delta$. Then,

$$\begin{aligned} G &\subseteq \bigcup_{i=1}^k [\ell_i, \ell_i + (t-1)] \cup \bigcup_{i=k+1}^{d-1} [j_i, j_i + (t-1)] \\ &= \text{supp}_t(x_{\ell_1} \cdots x_{\ell_k} x_{j_{k+1}} \cdots x_{j_{d-1}}) = F \in \mathcal{F}. \end{aligned}$$

Assume there exists an integer $1 \leq p \leq d$ such that $\ell_p > j_p$ and let p be minimal for such a property. By the meaning of p , for $q < p$, $\ell_q \leq j_q$. We need to distinguish two cases.

CASE 1. There exists an integer $q < p$ such that $\ell_q < j_q$.

In such a case, $x_{\ell_1}x_{\ell_2}\cdots x_{\ell_{p-1}} >_{\text{slex}} x_{j_1}x_{j_2}\cdots x_{j_{p-1}}$. Moreover, $k \leq d-1$, otherwise $x_{\ell_1}x_{\ell_2}\cdots x_{\ell_{p-1}}x_{\ell_p}\cdots x_{\ell_d} >_{\text{slex}} v$ and so $\mathbf{x}_G \in J$. A contradiction. Hence,

$$G \subseteq \bigcup_{i=1}^k [\ell_i, \ell_i + (t-1)]$$

and we can quickly find a facet $F \in \mathcal{F}$ which contains $\bigcup_{i=1}^k [\ell_i, \ell_i + (t-1)]$ and consequently G .

CASE 2. For all integers $q < p$, $\ell_q = j_q$.

In such a case, since $\ell_p > j_p$, we have $x_{j_1}\cdots x_{j_p} >_{\text{slex}} x_{\ell_1}\cdots x_{\ell_p}$, and

$$G \subseteq \bigcup_{i=1}^{p-1} [j_i, j_i + (t-1)] \cup [j_p + 1, n] = [n] \setminus F_p$$

and $[n] \setminus F_p$ is clearly a facet. Finally, all the facets of Δ are those described in (7). \square

Remark 2.2 We note that for an initial t -spread lexsegment ideal $J = I_\Delta$, all the facets $F \in \mathcal{F}(\Delta)$ have cardinality $|F| \geq (d-1)t$. Indeed, from Theorem 2.1, $\mathcal{F}(\Delta) = \{[n] \setminus F_p : p = 1, \dots, d\} \cup \mathcal{F}$. If $F \in \mathcal{F}$, then $|F| = (d-1)t$. Otherwise, if $F = [n] \setminus F_p$ for some p , then $|F| = |[n] \setminus F_p| = n - |F_p| = n - (j_p - (p-1)t)$. We observe that $j_p \leq n - (d-p)t$. Thus

$$|F| = n + (p-1)t - j_p \geq n + (p-1)t - n + (d-p)t = (d-1)t,$$

as desired.

We illustrate the previous result with an example.

Example 2.3 Let $v = x_2x_5x_7 \in S = \mathbb{Q}[x_1, \dots, x_7]$ be a 2-spread monomial and consider the initial 2-spread ideal $J = (\mathcal{L}_2^i(v))$ of S :

$$J = (x_1x_3x_5, x_1x_3x_6, x_1x_3x_7, x_1x_4x_6, x_1x_4x_7, x_1x_5x_7, x_2x_4x_6, x_2x_4x_7, x_2x_5x_7).$$

We have $n = 7, d = 3, t = 2$. The monomials $w \in M_{n+1-t, d-1, t} = M_{6, 2, 2}$ such that $w >_{\text{slex}} v/x_{\max(v)} = x_2x_5$ are the following

$$x_1x_3, \quad x_1x_4, \quad x_1x_5, \quad x_1x_6, \quad x_2x_4.$$

Therefore,

$$\begin{aligned}\mathcal{F} &= \{\text{supp}_2(x_1x_3), \text{supp}_2(x_1x_4), \text{supp}_2(x_1x_5), \text{supp}_2(x_1x_6), \text{supp}_2(x_2x_4)\} \\ &= \{\{1, 2, 3, 4\}, \{1, 2, 4, 5\}, \{1, 2, 5, 6\}, \{1, 2, 6, 7\}, \{2, 3, 4, 5\}\}.\end{aligned}$$

Moreover, $v = x_{j_1}x_{j_2}x_{j_3} = x_2x_5x_7$, hence

$$\begin{aligned}F_1 &= [j_1] \setminus \text{supp}_2(1) = [2] \setminus \emptyset = \{1, 2\}, \\ F_2 &= [j_2] \setminus \text{supp}_2(x_{j_1}) = [5] \setminus \{2, 3\} = \{1, 4, 5\}, \\ F_3 &= [j_3] \setminus \text{supp}_2(x_{j_1}x_{j_2}) = [7] \setminus \{2, 3, 5, 6\} = \{1, 4, 7\}.\end{aligned}$$

Finally, the standard decomposition of J is

$$\begin{aligned}J &= (x_1, x_2) \cap (x_1, x_4, x_5) \cap (x_1, x_4, x_7) \cap (x_1, x_6, x_7) \cap (x_3, x_4, x_5) \\ &\quad \cap (x_3, x_4, x_7) \cap (x_3, x_6, x_7) \cap (x_5, x_6, x_7).\end{aligned}$$

Here are some corollaries of Theorem 2.1.

Corollary 2.4 *In the hypotheses of Theorem 2.1, $\text{pd}(S/J) = n - (d-1)t$, $\text{depth}(S/J) = (d-1)t$ and $\dim(S/J) = n - j_1$.*

Proof. Since J is a t -spread strongly stable ideal, by (1), $\text{pd}(S/J) = \text{pd}(J) + 1 = n - (d-1)t$. So, by the Auslander–Buchsbaum formula, $\text{depth}(S/J) = \text{depth}(S) - \text{pd}(S/J) = n - (n - (d-1)t) = (d-1)t$.

Let Δ be the simplicial complex associated to J . By Theorem 2.1, the facets of Δ are those described in (7), and so, since $j_1 \leq n - (d-1)t$, and $j_1 + (i-1)t \leq j_i$, for $i = 2, \dots, d$, we have

$$\begin{aligned}\text{height}(J) &= \min \{n - (d-1)t, |F_\ell| : \ell = 1, \dots, d\} \\ &= \min \{n - (d-1)t, j_1, j_2 - t, \dots, j_d - (d-1)t\} = j_1.\end{aligned}$$

Therefore, $\dim(K[\Delta]) = \dim(S/J) = n - j_1$. \square

Corollary 2.5 *An initial t -spread lexsegment ideal generated in degree $d \geq 2$ is Cohen–Macaulay if and only if it is the t -spread Veronese ideal of degree d .*

Proof. Let $J = (\mathcal{L}_t^i(v))$, $v = x_{j_1}x_{j_2} \cdots x_{j_d}$ and let Δ be the simplicial complex associated to J . Assume J is Cohen–Macaulay, then Δ is pure. By Theorem 2.1, the facets of Δ are those described in (7), and so Δ is pure if and only if

$$(d-1)t = n - j_1 = n - (j_2 - t) = \cdots = n - (j_d - (d-1)t).$$

Hence $v = x_{n-(d-1)t}x_{n-(d-2)t} \cdots x_{n-t}x_n$ and $J = I_{n,d,t}$. Conversely, $I_{n,d,t}$ is Cohen–Macaulay. Indeed, $\dim(S/J) = \text{depth}(S/J) = (d-1)t$ (Corollary 2.4). \square

2.2. The final t -spread lexsegment case

In this subsection we determine the standard primary decomposition of a final t -spread lexsegment ideal of $S = K[x_1, \dots, x_n]$.

The next result will be crucial.

Proposition 2.6 *Let u be a t -spread monomial of degree d of S such that $\min(u) = 1$ and let $T = (\mathcal{L}_t^f(u))$ be a final t -spread lexsegment of S . Let Δ be the simplicial complex on $[n]$ associated to T . Then $|F| \in \{(d-1)t, 1 + (d-1)t\}$, for all $F \in \mathcal{F}(\Delta)$.*

Proof. Since we may see T as an initial t -spread lexsegment ideal with the order on the variables reversed $x_n > x_{n-1} > \dots > x_1$, then each facet of Δ has cardinality $\geq (d-1)t$ (Remark 2.2). To prove that $|F| \leq 1 + (d-1)t$ for all $F \in \mathcal{F}(\Delta)$, it is enough to show that each subset $G \subseteq [n]$ with cardinality $|G| = 2 + (d-1)t$ does not belong to Δ , i.e., $\mathbf{x}_G \in T$.

Let $G \subseteq [n]$ with $|G| = 2 + (d-1)t$. Define

$$\ell_1 = \min\{g \in G : g > 1\}, \text{ and}$$

$$\ell_j = \min\{k \in G : k \geq \ell_{j-1} + t\}, \text{ for } j \geq 2.$$

Since $|G| = 2 + (d-1)t$, the set $\{g \in G : g > 1\}$ has at least $1 + (d-1)t$ elements. Hence, the sequence of integers $\ell_1 < \ell_2 < \dots$ has at least d terms. Therefore, $H = \{\ell_1, \ell_2, \dots, \ell_d\} \subseteq G$ and $u >_{\text{slex}} \mathbf{x}_H$. Indeed, $\min(u) = 1$ and $\min(\mathbf{x}_H) = \ell_1 > 1$ and so $\mathbf{x}_H \in T$. It follows that $\mathbf{x}_G \in T$, as desired. \square

In the next theorem, we assume that $T \neq I_{n,d,t}$, as the case of the Veronese ideal has been covered in Theorem 2.1.

Theorem 2.7 *Let $u = x_{i_1} x_{i_2} \dots x_{i_d}$ be a t -spread monomial of degree $d \geq 2$ of S with $i_1 = 1$ and let $T = (\mathcal{L}_t^f(u))$ be a final t -spread lexsegment ideal of S , with $T \neq I_{n,d,t}$. Then, the standard primary decomposition of T is*

$$T = \bigcap_{G \in \mathcal{G}} \mathfrak{p}_{[n] \setminus G} \cap \bigcap_{H \in \mathcal{H}} \mathfrak{p}_{[n] \setminus H},$$

where

$$\begin{aligned} \mathcal{G} &= \{\text{cosupp}_t(w) \cup \{1\} : x_1 w \in M_{n,d,t}, w >_{\text{slex}} u/x_1\}, \\ \mathcal{H} &= \{\text{cosupp}_t(w) : x_1 w \in M_{n,d,t}, w \leq_{\text{slex}} u/x_1\}. \end{aligned}$$

Proof. Let Δ be a simplicial complex on the vertex set $[n]$ associated to T . By Proposition 1.5, we have to prove that

$$\mathcal{F}(\Delta) = \mathcal{G} \cup \mathcal{H}. \quad (8)$$

By Proposition 2.6, the facets of Δ have cardinality $(d-1)t$ or $1+(d-1)t$.

Let $G \in \mathcal{F}(\Delta)$ such that $|G| = 1+(d-1)t$. We prove that $G \in \mathcal{G}$. First of all, $\min(G) = 1$. Indeed, if $\min(G) \geq 2$, then setting $s_1 = \min(G)$, and

$$s_j = \min \{s \in G : s \geq s_{j-1} + t\}, \text{ for } j \geq 2,$$

the sequence $s_1 < s_2 < \dots$ has at least d elements, otherwise $|G| < 1+(d-1)t$. Hence, $\{s_1, s_2, \dots, s_d\} = U \subseteq G$ and $\mathbf{x}_U \in T$, as $\min(\mathbf{x}_U) = \min(G) \geq 2 > 1 = \min(u)$. It follows that $G \notin \Delta$. A contradiction. Therefore, $\min(G) = 1$.

Consider the following integers:

$$\ell_d = \max(G), \text{ and}$$

$$\ell_j = \max \{ \ell \in G : \ell \leq \ell_{j+1} - t \}, \text{ for } j < d.$$

The sequence $\ell_d > \ell_{d-1} > \dots > \ell_k$ has at least d terms, otherwise $|G| < 1+(d-1)t$. Moreover it has at most d terms, otherwise $\ell_1 > 1$, $x_{\ell_1} x_{\ell_2} \dots x_{\ell_d} \in T$ and then $G \notin \Delta$. A contradiction. Hence, $k = 1$.

Finally,

$$\begin{aligned} G \subseteq F &= \{ \ell_d, \ell_d - 1, \dots, \ell_d - (t-1), \dots, \ell_2, \ell_2 - 1, \dots, \ell_2 - (t-1), \ell_1 \} \\ &= \bigcup_{r=2}^d [\ell_r - (t-1), \ell_r] \cup \{ \ell_1 \} = \text{cosupp}_t(x_{\ell_2} \dots x_{\ell_d}) \cup \{ \ell_1 \}. \end{aligned}$$

Moreover $1+(d-1)t = |G| \leq |F| = 1+(d-1)t$ and $\ell_1 = 1$, and so $G = F \in \mathcal{G}$. Clearly, G is a facet. Indeed $\mathbf{x}_G \notin T$.

Now, let us determine the facets of Δ with cardinality $(d-1)t$. Let $H \in \mathcal{F}(\Delta)$ with $|H| = (d-1)t$. We prove that $H \in \mathcal{H}$. Consider the following integers

$$\ell_d = \max(H), \text{ and}$$

$$\ell_j = \max \{ \ell \in H : \ell \leq \ell_{j+1} - t \}, \text{ for } j < d.$$

The sequence $\ell_d > \ell_{d-1} > \dots > \ell_k$ has at least $d-1$ terms, otherwise $|H| < (d-1)t$, against our assumption. In fact, we have $k = 2$. Indeed, suppose $k = 1$. If $\ell_1 = 1$, then $H \subseteq G \in \mathcal{G}$. A contradiction since H is a facet. On the other hand, if $\ell_1 > 1$, then $x_{\ell_1} x_{\ell_2} \dots x_{\ell_d} \in T$, and $H \notin \Delta$. A contradiction.

Finally, setting $U = \{\ell_2, \dots, \ell_d\} \subseteq H$, if $x_1 \mathbf{x}_U >_{\text{slex}} u$, then H is contained in a facet $G \in \mathcal{G}$. A contradiction since H is a facet. If $x_1 \mathbf{x}_U \leq_{\text{slex}} u$, then $H \in \mathcal{H}$ and H is clearly a facet. The proof is complete. \square

Example 2.8 Let $u = x_1 x_4 x_6$ be a 2–spread monomial of degree 3 of $S = \mathbb{Q}[x_1, \dots, x_7]$. Consider the ideal $T = (\mathcal{L}_2^f(u))$ of S :

$$T = (x_1 x_4 x_6, x_1 x_4 x_7, x_1 x_5 x_7, x_2 x_4 x_6, x_2 x_4 x_7, x_2 x_5 x_7, x_3 x_5 x_7).$$

The monomials $w \in M_{7,2,2}$ such that $x_1 w \in M_{7,3,2}$ and $w >_{\text{slex}} u/x_1 = x_4 x_6$ are

$$x_3 x_5, \quad x_3 x_6, \quad x_3 x_7;$$

whereas, the monomials $w \in M_{7,2,2}$ such that $x_1 w \in M_{7,3,2}$ and $w \leq_{\text{slex}} u/x_1 = x_4 x_6$ are

$$x_4 x_6, \quad x_4 x_7, \quad x_5 x_7.$$

Therefore,

$$\begin{aligned} \mathcal{G} &= \{ \text{cosupp}_2(x_3 x_5) \cup \{1\}, \text{cosupp}_2(x_3 x_6) \cup \{1\}, \text{cosupp}_2(x_3 x_7) \cup \{1\} \} \\ &= \{ \{1, 2, 3, 4, 5\}, \{1, 2, 3, 5, 6\}, \{1, 2, 3, 6, 7\} \}, \\ \mathcal{H} &= \{ \text{cosupp}_2(x_4 x_6), \text{cosupp}_2(x_4 x_7), \text{cosupp}_2(x_5 x_7) \} \\ &= \{ \{3, 4, 5, 6\}, \{3, 4, 6, 7\}, \{4, 5, 6, 7\} \}. \end{aligned}$$

Thus, the standard primary decomposition of T is

$$T = (x_4, x_5) \cap (x_4, x_7) \cap (x_6, x_7) \cap (x_1, x_2, x_3) \cap (x_1, x_2, x_5) \cap (x_1, x_2, x_7).$$

Here are some corollaries of Theorem 2.7.

Corollary 2.9 *Let $T = (\mathcal{L}_t^f(u))$ be a final t –spread lexsegment ideal of S . Then $\text{pd}(S/T) = n - (d - 1)t$, $\text{depth}(S/T) = (d - 1)t$ and*

$$\dim(S/T) = \begin{cases} (d - 1)t & \text{if } u = x_1 x_{1+t} \cdots x_{1+(d-1)t}, \\ 1 + (d - 1)t & \text{otherwise.} \end{cases}$$

Proof. The ideal T is t –spread strongly stable but with the order on the variables reversed, $x_n > x_{n-1} > \cdots > x_1$. Hence, by (2), as $\min(u) = 1$ and $u \in T$, $\text{pd}(S/T) = \text{pd}(T) + 1 = n - 1 - (d - 1)t + 1 = n - (d - 1)t$. By the Auslander–Buchsbaum formula, $\text{depth}(S/T) = (d - 1)t$.

Suppose $T \neq I_{n,d,t}$. Let Δ the simplicial complex associated to T . By Theorem 2.7, the facets of Δ are those in (8). Now, \mathcal{H} is always non empty, as $u \in \mathcal{L}_t^f(u)$. Moreover, \mathcal{G} is non empty too, as $u \neq \min(M_{n,d,t})$, otherwise $T = I_{n,d,t}$. Thus

$$\begin{aligned} \text{height}(T) &= \min \{ \text{height}(\mathfrak{p}_{[n] \setminus G}), \text{height}(\mathfrak{p}_{[n] \setminus H}) : G \in \mathcal{G}, H \in \mathcal{H} \} \\ &= \min \{ n - (d-1)t, n - 1 - (d-1)t \} = n - 1 - (d-1)t. \end{aligned}$$

Thus, $\dim(K[\Delta]) = 1 + (d-1)t$.

Otherwise, if $T = I_{n,d,t}$, i.e., $u = \max(M_{n,d,t}) = x_1 x_{1+t} \cdots x_{1+(d-1)t}$, we have $\dim(K[\Delta]) = (d-1)t$ (Corollary 2.4). \square

Corollary 2.10 *A final t -spread lexsegment ideal of S generated in degree $d \geq 2$ is Cohen–Macaulay if and only if it is the t -spread Veronese ideal of degree d .*

Proof. Let $T = (\mathcal{L}_t^f(u))$ and let Δ be the simplicial complex associated to T . Assume T is Cohen–Macaulay, then Δ is pure. By contradiction, suppose $T \neq I_{n,d,t}$. By Theorem 2.7, the facets of Δ are those described in (8). The families \mathcal{G} and \mathcal{H} are both non empty, the first as $u \in \mathcal{L}_t^f(u)$, the second as $u \neq \max(M_{n,d,t})$. But, any $G \in \mathcal{G}$ has cardinality $|G| = 1 + (d-1)t$ and any $H \in \mathcal{H}$ has cardinality $|H| = (d-1)t$. So Δ should not be pure. A contradiction. Thus $T = I_{n,d,t}$. The converse easily follows. \square

Remark 2.11 A simplicial complex Δ is pure if and only if the Stanley–Reisner ideal I_Δ is unmixed, in the sense that all associated prime ideals have the same height (see, for instance, [18]). One can observe that the proofs of Corollary 2.5 and Corollary 2.10 have pointed out that the classification for initial Cohen–Macaulay t -spread lexsegment ideal and for final Cohen–Macaulay t -spread lexsegment ideal is equivalent to the classification for initial unmixed t -spread lexsegment ideal and for final unmixed t -spread lexsegment ideal.

2.3. The case of a completely t -spread lexsegment ideal

The aim of this subsection is to determine the minimal primary decomposition of a completely t -spread lexsegment ideal $I = (\mathcal{L}_t(u, v))$ of $S = K[x_1, \dots, x_n]$, $u, v \in M_{n,d,t}$, $u >_{\text{slex}} v$ with $\min(u) = 1$ and $\min(v) \geq 2$.

Theorem 2.12 *Let $u = x_{i_1} x_{i_2} \cdots x_{i_d}, v = x_{j_1} x_{j_2} \cdots x_{j_d}$ be t -spread monomials of degree $d \geq 2$ of S with $i_1 = 1$ and $j_1 \geq 2$. Let $I = (\mathcal{L}_t(u, v))$ be a completely t -spread lexsegment ideal of S , $I \neq I_{n,d,t}$. Then, the standard primary decomposition of I is*

$$I = \bigcap_{G \in \mathcal{G}} \mathfrak{p}_{[n] \setminus G} \cap \bigcap_{p \in \mathcal{I}} \mathfrak{p}_{F_p} \cap \bigcap_{F \in \mathcal{F}} \mathfrak{p}_{[n] \setminus F},$$

where F_p ($p = 1, \dots, d$) and \mathcal{G} are the families described in Theorems 2.1 and 2.7,

- $\mathcal{I} = [d] \setminus \{p \in [d] : |[n] \setminus F_p| = (d-1)t \text{ and } v/x_{j_p} >_{\text{slex}} u/x_1\}$,
- $\tilde{\mathcal{F}}$ is the family consisting of the sets $F = \text{supp}_t(w) \in \mathcal{F}$ with $w = x_{\ell_1} \cdots x_{\ell_{d-1}}$ such that
 - $1 \notin F$ and $x_{\ell_1+(t-1)} \cdots x_{\ell_{d-1}+(t-1)} \leq_{\text{slex}} u/x_1$, or
 - $1 \in F$ and $F \cup \{j\} \notin \mathcal{G}$, for all $j \notin F$,

where \mathcal{F} is the family described in Theorem 2.1.

Proof. By hypothesis, $I = J \cap T$ with $J = (\mathcal{L}_t^i(v))$ and $T = (\mathcal{L}_t^f(u))$. By Theorems 2.1 and 2.7 we have

$$I = T \cap J = \bigcap_{G \in \mathcal{G}} \mathfrak{p}_{[n] \setminus G} \cap \bigcap_{H \in \mathcal{H}} \mathfrak{p}_{[n] \setminus H} \cap \bigcap_{p=1}^d \mathfrak{p}_{F_p} \cap \bigcap_{F \in \mathcal{F}} \mathfrak{p}_{[n] \setminus F}. \quad (9)$$

To find the standard primary decomposition of I , it suffices to delete from this presentation those primes that are not minimal. Equivalently, if Δ is the simplicial complex on vertex set $[n]$ associated to I , we must determine the facets of Δ . By (9) we have that

$$\mathcal{F}(\Delta) \subseteq \mathcal{G} \cup \mathcal{H} \cup \{[n] \setminus F_p : p = 1, \dots, d\} \cup \mathcal{F}. \quad (10)$$

where $\mathcal{G}, \mathcal{H}, F_p$ ($p = 1, \dots, d$), \mathcal{F} are the families described in Theorems 2.1 and 2.7.

CLAIM. $\mathcal{F}(\Delta) = \mathcal{G} \cup \{[n] \setminus F_\ell : \ell \in \mathcal{I}\} \cup \tilde{\mathcal{F}}$.

PROOF OF THE CLAIM. We analyze the four families in (10), and in so doing we determine the facets of Δ .

- Let $G \in \mathcal{G}$. Since $I \subseteq T$ and $\mathbf{x}_G \notin T$, it follows that $\mathbf{x}_G \notin I$, i.e., $G \in \Delta$. Let us prove that G is a facet. Indeed, $G \not\subseteq F, H$, for all $F \in \mathcal{F}$ and all $H \in \mathcal{H}$. Indeed, $|F| = |H| = (d-1)t < 1 + (d-1)t = |G|$. Moreover, $G \not\subseteq [n] \setminus F_p$, as $1 \in G$ and $1 \in F_p$. In fact, $\min(v) = j_1 \geq 2$.
- Let $H \in \mathcal{H}$. Arguing as before, $H \in \Delta$. We show that we can “eliminate” the family \mathcal{H} . In fact, we can write $H = \text{cosupp}_t(x_{\ell_2} \cdots x_{\ell_d})$, with $x_1 x_{\ell_2} \cdots x_{\ell_d} \in M_{n,d,t}$ and $x_{\ell_2} \cdots x_{\ell_d} \leq_{\text{slex}} u/x_1$. Let

$$\tilde{H} = \{\ell_2 - (t-1), \ell_3 - (t-1), \dots, \ell_d - (t-1)\} \subseteq H.$$

Setting $h_i = \ell_{i+1} - (t-1)$, for $i = 1, \dots, d-1$, then $\tilde{H} = \{h_1 < h_2 < \cdots < h_{d-1}\}$. We distinguish three cases.

- CASE 1. Let $\mathbf{x}_{\tilde{H}} >_{\text{slex}} v/x_{\max(v)}$. Since $\text{supp}_t(\mathbf{x}_{\tilde{H}}) = \text{cosupp}_t(x_{\ell_2} \cdots x_{\ell_d}) = H$, then $H \in \mathcal{F}$ (we will determine when $H \in \mathcal{F}$ is a facet of Δ later).
- CASE 2. Let $\mathbf{x}_{\tilde{H}} <_{\text{slex}} v/x_{\max(v)}$. Then there exists an integer $s \in [d-1]$ such that $j_1 = h_1, \dots, j_{s-1} = h_{s-1}$ and $j_s < h_s$. Hence

$$\begin{aligned} H &\subseteq \{j_1, \dots, j_1 + (t-1), \dots, j_{s-1}, \dots, j_{s-1} + (t-1), j_s + 1, \dots, n\} \\ &= [n] \setminus F_s = F. \end{aligned}$$

Moreover, $\mathbf{x}_F \notin I$, as $I \subseteq J$ and $\mathbf{x}_F \notin J$. So H is included in $F \in \Delta$.

- CASE 3. Let $\mathbf{x}_{\tilde{H}} = v/x_{\max(v)}$. Then we have $H \subseteq [n] \setminus F_d = F$, and also $F \in \Delta$. Thus H is included in the face $F \in \mathcal{F}$.
- Let $F = [n] \setminus F_p$, some $1 \leq p \leq d$. Since $I \subseteq J$ and $\mathbf{x}_F \notin J$, then $\mathbf{x}_F \notin I$ and so $F \in \Delta$. If $|F| \geq 1 + (d-1)t$, then F is a facet as F is not contained in any other set of the families \mathcal{F}, \mathcal{H} . Moreover, $F \neq G$ for all $G \in \mathcal{G}$, as $1 \in G \setminus F$. Suppose $|F| = (d-1)t$. Then F is not a facet if and only if $F \subsetneq G$, for some $G \in \mathcal{G}$. Therefore, if and only if $F = G \setminus \{1\}$, for some $G \in \mathcal{G}$. On the other hand, $G = \{\ell_d, \ell_d - 1, \dots, \ell_d - (t-1), \dots, \ell_2, \ell_2 - 1, \dots, \ell_2 - (t-1), 1\}$ is characterized by the conditions $x_1 x_{\ell_2} \cdots x_{\ell_d} \in M_{n,d,t}$ and $x_1 x_{\ell_2} \cdots x_{\ell_d} >_{\text{slex}} u$. We observe that $\text{supp}(v/x_{j_p}) \subseteq F = [n] \setminus F_p$. Hence, if $F \subseteq G$, then

$$v/x_{j_p} \geq_{\text{slex}} x_{\ell_2} \cdots x_{\ell_d} >_{\text{slex}} u/x_1.$$

Thus, if $F \subseteq G$, then $v/x_{j_p} >_{\text{slex}} u/x_1$. On the other hand, if $v/x_{j_p} \leq_{\text{slex}} u/x_1$, then $x_{\ell_2} \cdots x_{\ell_d} \leq_{\text{slex}} v/x_{j_p} \leq_{\text{slex}} u/x_1$ and G is not a facet. Finally, we have verified that $F = [n] \setminus F_p$ is a facet if and only if $p \in \mathcal{I}$.

- Let $F \in \mathcal{F}$. Clearly $\mathbf{x}_F \notin I$ and so $F \in \Delta$. We have that

$$F = \{\ell_1, \ell_1 + 1, \dots, \ell_1 + (t-1), \dots, \ell_{d-1}, \ell_{d-1} + 1, \dots, \ell_{d-1} + (t-1)\},$$

with $x_{\ell_1} \cdots x_{\ell_{d-1}} \in M_{n+1-t,d-1,t}$ and $x_{\ell_1} \cdots x_{\ell_{d-1}} >_{\text{slex}} v/x_{\max(v)}$. If F is not a facet, then $F \subsetneq G$, for some $G \in \mathcal{G}$.

- CASE 1. Let $1 \notin F$. Then we have $F = G \setminus \{1\}$. G is characterized by the condition $x_{\ell_1+(t-1)} \cdots x_{\ell_{d-1}+(t-1)} >_{\text{slex}} u/x_1$. Hence, in such a case, F is a facet if and only if $x_{\ell_1+(t-1)} \cdots x_{\ell_{d-1}+(t-1)} \leq_{\text{slex}} u/x_1$.
- CASE 2. Let $1 \in F$. Set $\ell = \ell_j + t$, if $j = \min \{j \in [d-2] : \ell_j + (t-1) < \ell_{j+1}\}$ does exist, otherwise set $\ell = \ell_{d-1} + t$. Then $F \subseteq F \cup \{\ell\}$.

Finally F is a facet if and only if $F \cup \{\ell\} \notin \mathcal{G}$. Thus, if and only if

$$\begin{aligned} x_{\ell_1+t} \cdots x_{\ell_{j-1}+t} x_{\ell_j+t} \cdots x_{\ell_{d-1}+(t-1)} &\leq_{\text{slex}} u/x_1, & \text{if } \ell = \ell_j + t, \\ x_{\ell_1+t} \cdots x_{\ell_{j-1}+t} x_{\ell_j+t} \cdots x_{\ell_{d-1}+t} &\leq_{\text{slex}} u/x_1, & \text{if } \ell = \ell_{d-1} + t. \end{aligned}$$

The claim follows. \square

Example 2.13 Let $u = x_1x_4x_6$ and $v = x_2x_5x_7$ be 2–spread monomials of degree 3 of $S = \mathbb{Q}[x_1, \dots, x_7]$. Consider the t –spread lexsegment ideal

$$I = (\mathcal{L}_2(u, v)) = (x_1x_4x_6, x_1x_4x_7, x_1x_5x_7, x_2x_4x_6, x_2x_4x_7, x_2x_5x_7)$$

of S . Firstly, we verify that I is a completely 2–spread lexsegment ideal, *i.e.*, $I = J \cap T$, where J and T are the 2–spread lexsegment ideals in Examples 2.3 and 2.8, respectively. By [15, Theorem 3.7. (b)], I is a completely 2–spread lexsegment ideal if and only if for all $w \in M_{7,3,2}$ with $w <_{\text{slex}} v = x_2x_5x_7$, there exists $s \in \text{supp}(w)$ such that $x_{\min(u)}w/x_s = x_1w/x_s \leq_{\text{slex}} u = x_1x_4x_6$.

We have $w = x_3x_5x_7$ and we may choose $s = \min(w)$. Thus, I is a completely 2–spread lexsegment ideal.

Setting, $u = x_{i_1}x_{i_2}x_{i_3} = x_1x_{i_2}x_{i_3}$ and $v = x_{j_1}x_{j_2}x_{j_3} = x_2x_{j_2}x_{j_3}$, we have

$$\mathcal{I} = [3] \setminus \{p \in [3] : |[7] \setminus F_p| = 2 \cdot 2 \text{ and } u/x_1 = x_4x_6 <_{\text{slex}} v/x_{j_p} = x_2x_5x_7/x_{j_\ell}\}.$$

The only sets F_p with $|[7] \setminus F_p| = 2 \cdot 2 = 4$ are $F_2 = \{1, 4, 5\}$ and $F_3 = \{1, 4, 7\}$ (Example 2.3). Moreover, $v/x_{j_2} = x_2x_7 >_{\text{slex}} x_4x_6$ and $v/x_{j_3} = x_2x_5 >_{\text{slex}} x_4x_6$. Thus, $\mathcal{I} = [3] \setminus \{2, 3\} = \{1\}$.

From Examples 2.3 and 2.8, we have:

$$\mathcal{G} = \{\{1, 2, 3, 4, 5\}, \{1, 2, 3, 5, 6\}, \{1, 2, 3, 6, 7\}\}$$

and

$$\mathcal{F} = \{\{1, 2, 3, 4\}, \{1, 2, 4, 5\}, \{1, 2, 5, 6\}, \{1, 2, 6, 7\}, \{2, 3, 4, 5\}\}.$$

Let $F \in \mathcal{F}$ such that $1 \notin F$. Then $F = \{2, 3, 4, 5\} = \text{supp}_2(x_2x_4)$. We have that $x_3x_5 >_{\text{slex}} u/x_1 = x_4x_6$, so $F \notin \tilde{\mathcal{F}}$.

Let $F \in \mathcal{F}$ such that $1 \in F$. Then $F \in \tilde{\mathcal{F}}$ if and only if $F \not\subseteq G$, for all $G \in \mathcal{G}$. Since such sets do not exist, $\tilde{\mathcal{F}} = \emptyset$.

Finally, the standard primary decomposition of I is

$$I = (x_1, x_2) \cap (x_4, x_5) \cap (x_4, x_7) \cap (x_6, x_7).$$

3. COHEN–MACAULAY t –SPREAD LEXSEGMENT IDEALS

In Section 2, we have seen that if I is an initial or a final t –spread lexsegment ideal of $S = K[x_1, \dots, x_n]$, $t \geq 1$, then I is Cohen–Macaulay if and only if $I = I_{n,d,t}$ is the Veronese t –spread ideal of degree d of S (Corollaries 2.5, 2.10). In this section, we consider the more general problem of classifying all Cohen–Macaulay t –spread lexsegment ideals $I = (\mathcal{L}_t(u, v))$ of S , with $t \geq 1$. The case $t = 1$ has been examined also in [9] by using Serre’s condition (S_2).

As in Section 2, we may assume that $I = (\mathcal{L}_t(u, v))$ with $u >_{\text{slex}} v$, $\min(u) = 1$, $\min(v) \geq 2$ and $\deg(u) = \deg(v) = d \geq 2$. In order to achieve our purpose, we will distinguish two cases: $\min(v) = 2$ and $\min(v) > 2$.

First, we note that we can always suppose $n > 2 + (d-1)t$. Indeed, for $n = 1 + (d-1)t$, there is only one t –spread lexsegment ideal, namely $I = (x_1 x_{1+t} \cdots x_{1+(d-1)t}) = I_{1+(d-1)t,d,t}$ which is always a Cohen–Macaulay ideal; whereas, for $n = 2 + (d-1)t$, $I = (\mathcal{L}_t(u, v))$ with $v = x_2 x_{2+t} \cdots x_{2+(d-1)t}$, necessarily. In fact, in such a case there does exist only one t –spread monomial whose minimum is equal to 2. Hence, I is final and therefore Cohen–Macaulay if and only if $I = I_{2+(d-1)t,d,t}$ (Corollary 2.10).

In what follows we denote the squarefree lex order \geq_{slex} simply by \geq .

3.1. The height two case

In this subsection we study the Cohen–Macaulayness of a t –spread lexsegment ideal $I = (\mathcal{L}_t(u, v))$ of S with $\min(v) = 2$. We will distinguish two cases: $3 + (d-1)t \leq n \leq 3 + (2d-3)t$ and $n \geq 4 + (2d-3)t$.

We start with the case $3 + (d-1)t \leq n \leq 3 + (2d-3)t$. The notion of *Betti splitting* [12] (see also [16] and the reference therein) will be crucial.

Let I, P, Q be monomial ideals of $S = K[x_1, \dots, x_n]$ such that $G(I)$ is the disjoint union of $G(P)$ and $G(Q)$. We say that $I = P + Q$ is a *Betti splitting* if

$$\beta_{i,j}(I) = \beta_{i,j}(P) + \beta_{i,j}(Q) + \beta_{i-1,j}(P \cap Q), \quad \text{for all } i, j.$$

If $I = P + Q$ is a Betti splitting [16, Corollary 2.2], then

$$\text{pd}(I) = \max\{\text{pd}(P), \text{pd}(Q), \text{pd}(P \cap Q) + 1\}. \tag{11}$$

Lemma 3.1 [16, Corollary 2.7] *Let I be a monomial ideal of S . Let Q be the ideal generated by all elements of $G(I)$ divisible by x_i and let P be the ideal generated by all other elements of $G(I)$. If the ideal Q has a linear resolution, then $I = P + Q$ is a Betti splitting.*

For I a monomial ideal of S , we set $\text{gcd}(I) = \text{gcd}(u : u \in G(I))$.

Theorem 3.2 *Let $u = x_{i_1}x_{i_2} \cdots x_{i_d}$, $v = x_{j_1}x_{j_2} \cdots x_{j_d}$ be t -spread monomials of degree $d \geq 2$ of S with $i_1 = 1$, $j_1 = 2$ and let I be the t -spread lexsegment ideal $I = (\mathcal{L}_t(u, v))$ of S . Assume $3 + (d - 1)t \leq n \leq 3 + (2d - 3)t$ and set*

$$P = (\mathcal{L}_t(u, x_1x_{n-(d-2)t} \cdots x_{n-t}x_n)), \quad Q = (\mathcal{L}_t(x_2x_{2+t} \cdots x_{2+(d-1)t}, v)).$$

Then I is Cohen–Macaulay if and only if $\gcd(I) = 1$ and $P \cap Q$ is a principal ideal.

Proof. Let I be Cohen–Macaulay and let Δ be the simplicial complex on the vertex set $[n]$ associated to I . Then Δ is pure. Since each monomial in $G(I)$ has minimum equal to 1 or 2, the set $[n] \setminus \{1, 2\}$ is a facet of Δ . Thus, Δ must be pure of dimension $\dim(\Delta) = n - 3$.

Now, we prove that $v \in \{v_1, v_2\}$, where

$$\begin{aligned} v_1 &= \prod_{s=0}^{d-1} x_{2+st}, \\ v_2 &= \left(\prod_{s=0}^{d-2} x_{2+st} \right) x_{3+(d-1)t}. \end{aligned} \tag{12}$$

First, note that $v_1 > v_2$ are the greatest monomials of $M_{n,d,t}$ with minimum 2. So $v_1 \in I$. Let $E = \{\max(w) : w \in \mathcal{L}_t(v_1, v)\}$ and set $k = \max(E)$. Then $k \geq 2 + (d - 1)t$. Indeed $v_1 \in I$. If $v = v_1$, there is nothing to prove. Assume $v > v_1$, then $|E| \geq 2$. We show that $v = v_2$. The monomial prime ideal $\mathfrak{p}_{\{1\} \cup E}$ contains I , and $\text{height}(\mathfrak{p}_{\{1\} \cup E}) = 1 + |E| \geq 3$. Since I is Cohen–Macaulay, thus height–unmixed of height two, there must be a minimal prime $\mathfrak{p} \in \text{Min}(I)$ of height 2 that contains properly $\mathfrak{p}_{\{1\} \cup E}$. For all $j \in E$, $\mathfrak{p}_{\{1\} \cup (E \setminus \{j\})}$ does not contain I . Indeed, the monomial $(\prod_{s=0}^{d-2} x_{2+st})x_j \in \mathcal{L}_t(v_1, v)$ is not in $\mathfrak{p}_{\{1\} \cup (E \setminus \{j\})}$. Hence, x_1 can be omitted and \mathfrak{p}_E must contain I . One can quickly observe that no variable x_j , $j \in E$, can be omitted again. Hence, \mathfrak{p}_E is a minimal prime with $\text{height}(\mathfrak{p}_E) = |E| = 2$, *i.e.*, $k = 2 + (d - 1)t$. So $v \in \{v_1, v_2\}$ as desired.

For $\ell = 0, \dots, d - 1$, let us define the following monomials:

$$\begin{aligned} u_\ell &= x_1 \left(\prod_{s=\ell}^{d-2} x_{n-st-1} \right) \left(\prod_{s=0}^{\ell-1} x_{n-st} \right) \\ &= x_1 x_{n-(d-2)t-1} \cdots x_{n-\ell t-1} x_{n-(\ell-1)t} \cdots x_{n-t} x_n. \end{aligned} \tag{13}$$

One can observe that $\mathcal{L}_t(u_0, u_{d-1}) = \{u_0 > u_1 > \cdots > u_{d-1}\}$ and furthermore u_{d-1} is the smallest monomial of $M_{n,d,t}$ whose minimum is equal to 1. Thus $u > u_{d-1} > v$ and $u_{d-1} \in I$. We prove that $u = u_\ell$, for some $\ell \in \{0, \dots, d - 1\}$.

Indeed, writing $u = x_1x_{i_2} \cdots x_{i_d}$, we note that $i_2 \leq n - (d - 2)t$. The condition $u \leq u_0$ is equivalent to $i_2 \geq n - (d - 2)t - 1$. As before, we consider a suitable monomial prime ideal that contains I , namely $\mathfrak{p}_{[i_2, n-(d-2)t] \cup \{2\}}$. None of x_j with $j \in [i_2, n - (d - 2)t]$ can be omitted from $\mathfrak{p}_{[i_2, n-(d-2)t] \cup \{2\}}$. Thus, $\mathfrak{p}_{[i_2, n-(d-2)t]}$ must be a minimal prime of height two. Hence, $i_2 = n - (d - 2)t - 1$, as desired.

By what shown so far, if I is Cohen–Macaulay, then $I = (\mathcal{L}_t(u_\ell, v_k))$, for some $\ell \in \{0, \dots, d-1\}$ and $k \in \{1, 2\}$. On the other hand, I is Cohen–Macaulay if and only if $\dim(S/I) = \text{depth}(S/I)$. We note that $\dim(S/I) = n-2$. Indeed, we have shown previously that $\text{height}(I) = 2$. By the Auslander–Buchsbaum formula, $\text{depth}(S/I) = n - \text{pd}(S/I) = n-1 - \text{pd}(I)$. Hence, I is Cohen–Macaulay if and only if $\text{height}(I) = 2$ and $\text{pd}(I) = 1$.

Hence, in order to establish the theorem we need to prove the next facts.

CLAIM 1. $\text{height}(I) = 2$ if and only if $\gcd(I) = \gcd(w : w \in G(I)) = 1$.

PROOF OF CLAIM 1. Indeed, $\mathfrak{p}_{[2]} = (x_1, x_2)$ is a minimal prime of I having height two, so $\text{height}(I) \leq \text{height}(\mathfrak{p}_{[2]}) = 2$. Moreover, $\text{height}(I) = 1$ if and only if there is a minimal prime \mathfrak{q} of I whose height is 1, *i.e.*, there exists a variable x_s dividing all generators of I . The claimed statement follows.

CLAIM 2. $\text{pd}(I) = 1$ if and only if the ideal $(\mathcal{L}_t(u, u_{d-1})) \cap (\mathcal{L}_t(v_1, v)) = P \cap Q$ is a principal ideal.

PROOF OF CLAIM 2. We can observe that

$$P = (w \in G(I) : \min(w) = 1), \quad Q = (w \in G(I) : \min(w) = 2).$$

Clearly $I = P + Q$. We claim that $I = P + Q$ is a Betti splitting of I . Indeed using Lemma 3.1, it suffices to note that $Q = (w \in G(I) : x_2 \text{ divides } w)$ and that Q has a linear resolution. For all monomials $w \in G(I)$ with $\min(w) = 1$, $\min(w/x_1) \geq n - (d-2)t - 1 > 2$, as by hypothesis $n \geq 3 + (d-1)t > 3 + (d-2)t$. Moreover Q is an equigenerated initial t -spread lexsegment ideal in $K[x_2, \dots, x_n]$, so it has a linear resolution. Hence, by (11),

$$\text{pd}(I) = \max\{\text{pd}(P), \text{pd}(Q), \text{pd}(P \cap Q) + 1\}.$$

We have $\text{pd}(P) \leq 1$. Indeed, P is isomorphic to the final t -spread lexsegment ideal $(\mathcal{L}_t(u/x_1, x_{n-(d-2)t} \cdots x_{n-t}x_n))$ generated in degree $d-1$, and formula (2) gives

$$\text{pd}(I) = \begin{cases} n - (n - (d-2)t - 1) - (d-2)t = 1 & \text{if } u > u_{d-1}, \\ n - (n - (d-2)t) - (d-2)t = 0 & \text{if } u = u_{d-1}. \end{cases}$$

Analogously, as Q is an initial t -spread lexsegment ideal in $K[x_2, \dots, x_n]$, $\text{pd}(Q) = 0$ if $v = v_1$, or $\text{pd}(Q) = 1$ if $v = v_2$. Thus, (11) implies that $\text{pd}(I) = 1$ if and only if $\text{pd}(P \cap Q) = 0$, *i.e.*, if and only if $P \cap Q$ is a principal ideal. \square

The next result points out that for $n \gg 0$, precisely $n \geq 4 + (2d-3)t$, the Cohen–Macaulay t -spread lexsegment ideals with $j_1 = 2$ are just complete intersection ideals.

Theorem 3.3 *Let $u = x_{i_1}x_{i_2} \cdots x_{i_d}$ and $v = x_{j_1}x_{j_2} \cdots x_{j_d}$ be t -spread monomials of degree $d \geq 2$ of S with $i_1 = 1$, $j_1 = 2$, and let I be the t -spread lexsegment ideal $I = (\mathcal{L}_t(u, v))$ of S . Assume $n \geq 4 + (2d-3)t$, then the following conditions are equivalent:*

(a) I is Cohen–Macaulay;

(b) $u = x_1x_{n-(d-2)t} \cdots x_{n-t}x_n$, $v = x_2x_{2+t} \cdots x_{2+(d-1)t}$.

Proof. Let Δ be the simplicial complex on vertex set $[n]$ associated to I .

(a) \implies (b). If $I = I_\Delta$ is Cohen–Macaulay, then Δ is pure. The set $[n] \setminus \{1, 2\}$ is a facet of Δ . Indeed, any monomial $w \in I$ has $1 \in \text{supp}(w)$ or $2 \in \text{supp}(w)$. Therefore, Δ pure implies $\dim(\Delta) = n - 3$. Consider the set $G = [n] \setminus \{2, i_2, i_2 + 1, \dots, n - (d - 2)t\}$. Observe that $G \in \Delta$, indeed $\mathfrak{p}_{[n] \setminus G} = (x_2, x_{i_2}, x_{i_2+1}, \dots, x_{n-(d-2)t})$ is a monomial prime ideal that contains I . Now, the condition Δ is pure implies that there exists a minimal prime $\mathfrak{p} \in \text{Min}(I)$ of height two such that $\mathfrak{p} \subseteq \mathfrak{p}_{[n] \setminus G}$. The t -spread monomial $w = x_2x_{2+t} \cdots x_{2+(d-1)t} \in I$ as $u > w \geq v$. We have $x_2 \in \mathfrak{p}$. In fact, if $x_2 \notin \mathfrak{p}$, then $\mathfrak{p} \subseteq \mathfrak{p}_{[i_2, n-(d-2)t]}$. We must have $\mathfrak{p} = \mathfrak{p}_{[i_2, n-(d-2)t]}$. Indeed, if we omit some x_j , $j \in [i_2, n - (d - 2)t]$, then we can find a monomial $z \in G(I)$, with $\min(z) = 1$ and $\min(z/x_1) = j$, and this monomial does not belong to \mathfrak{p} , a contradiction. Δ pure implies $\text{height}(\mathfrak{p}) = 2$, thus $i_2 = n - (d - 2)t - 1$. But $[n - (d - 2)t - 1, n - (d - 2)t] \cap \text{supp}(w) = \emptyset$. Indeed, $n \geq 4 + (2d - 3)t$, so

$$\begin{aligned} n - (d - 2)t - 2 &\geq 4 + (2d - 3)t - (d - 2)t - 2 = 2 + (d - 1)t \\ &= \max(w). \end{aligned} \tag{14}$$

Thus $\max(w) < n - (d - 2)t - 1$. This implies that $w \in I \setminus \mathfrak{p}$, a contradiction. Hence, $x_2 \in \mathfrak{p}$. Therefore, $\text{height}(\mathfrak{p}) = 2$ and $\mathfrak{p} = (x_2, x_{i_2})$. The monomial $z = x_1x_{n-(d-2)t} \cdots x_{n-t}x_n$ belongs to I . We have $z \in \mathfrak{p}$ if and only if x_{i_2} divides z . Thus, as $1 + t \leq i_2 \leq n - (d - 2)t$, we have $i_2 = n - (d - 2)t$ and $u = x_1x_{n-(d-2)t} \cdots x_{n-t}x_n$. Let $r = \max\{\max(w) : w \in G(I) \text{ and } \min(w) = 2\}$. Observe that $r \geq 2 + (d - 1)t$, as $x_2x_{2+t} \cdots x_{2+(d-1)t} \in I$. The ideal $\mathfrak{q} = (x_1, x_{2+(d-1)t}, x_{3+(d-1)t}, \dots, x_r)$ is a minimal prime of I . The condition Δ pure implies $\text{height}(\mathfrak{q}) = 2$ and $r = 2 + (d - 1)t$. Thus, $\max(v) = 2 + (d - 1)t$ and $v = x_2x_{2+t} \cdots x_{2+(d-1)t}$.

(b) \implies (a). Let $u = x_1x_{n-(d-2)t} \cdots x_{n-t}x_n$ and $v = x_2x_{2+t} \cdots x_{2+(d-1)t}$. By (14), we have $\text{supp}(u) \cap \text{supp}(v) = \emptyset$. Moreover $I = (u, v)$. Thus, I is a complete intersection and so it is Cohen–Macaulay. \square

Remark 3.4 One can observe that under the hypotheses of Theorem 3.3, the classification for Cohen–Macaulay t -spread lexsegment ideal is equivalent to the classification for unmixed t -spread lexsegment ideal (see also, Remark 2.11).

3.2. The general case

In this subsection, we classify all Cohen–Macaulay t -spread lexsegment ideals $I = (\mathcal{L}_t(u, v))$ of $S = K[x_1, \dots, x_n]$ with $\min(v) > 2$.

3.2.1 The underlying idea behind the classification

Let $I = (\mathcal{L}_t(u, v))$ be a Cohen–Macaulay t -spread lexsegment ideal of S with $u = x_{i_1}x_{i_2}\cdots x_{i_d}$, $v = x_{j_1}x_{j_2}\cdots x_{j_d} \in M_{n,d,t}$ such that $\min(u) = i_1 = 1$, $\min(v) = j_2 > 2$, $d \geq 2$. If Δ is the simplicial complex on the vertex set $[n]$ associated to I , then Δ is pure. This is a key property in order to get the classification. First, note that since $\min(u) = 1$, then

$$u \in \mathcal{L}_t(x_1x_{1+t}\cdots x_{1+(d-1)t}, x_1x_{n-(d-2)t}x_{n-(d-3)t}\cdots x_{n-t}x_n).$$

Assume $u = x_1x_{1+t}\cdots x_{1+(d-1)t} = \max(M_{n,d,t})$, then $I = (\mathcal{L}_t(u, v)) = (\mathcal{L}_t^i(v))$ is an initial t -spread lexsegment ideal. Hence, Δ is Cohen–Macaulay if and only if $I = I_{n,d,t}$ (Corollary 2.5).

Now, let $u < x_1x_{1+t}\cdots x_{1+(d-1)t}$. Then, $F = [1 + (d-1)t] \in \Delta$. Moreover, $[1 + (d-1)t] \cup \{k\} \notin \Delta$, for all $k \notin [1 + (d-1)t]$. In fact, $j_1 > 2$ implies that $w = x_2x_{2+t}\cdots x_{2+(d-2)t}x_k \in I = I_\Delta$, thus $G = \text{supp}(w) \notin \Delta$ and $G \subseteq F \cup \{k\}$. Therefore, F is a facet of Δ and by the purity of Δ we have that

$$\dim(\Delta) = |F| - 1 = (d-1)t. \quad (15)$$

We show that $v \leq x_{n-1-(d-1)t}x_{n-1-(d-2)t}\cdots x_{n-1-t}x_{n-1}$. Suppose on the contrary that $v > x_{n-1-(d-1)t}x_{n-1-(d-2)t}\cdots x_{n-1-t}x_{n-1}$, then $A = [n-1-(d-1)t, n]$ is a face of Δ and $\dim(\Delta) \geq \dim A = |A| - 1 = (d-1)t + 1 > (d-1)t = \dim(\Delta)$, a contradiction. Thus, $v \leq x_{n-1-(d-1)t}x_{n-1-(d-2)t}\cdots x_{n-1-t}x_{n-1}$. Hence,

$$v \in \{v_0 > v_1 > \cdots > v_{d-1} > v_d = \min(M_{n,d,t})\},$$

where

$$v_\ell = \left(\prod_{s=\ell}^{d-1} x_{n-st-1} \right) \left(\prod_{s=0}^{\ell-1} x_{n-st} \right), \quad \ell \in \{0, \dots, d\}. \quad (16)$$

We can observe that $v \neq v_d$. Otherwise, $I = (\mathcal{L}_t(u, v)) = (\mathcal{L}_t^f(u))$ is a final t -spread lexsegment ideal and so I is Cohen–Macaulay if and only if I is the t -spread Veronese ideal $I_{n,d,t}$ (Corollary 2.10). Thus $u = \max(M_{n,d,t})$, against the assumption that $u < \max(M_{n,d,t})$.

Finally, for v there do exist three admissible choices:

- $v = v_{d-1}$,
- $t = 1$ and $v = v_\ell$, for $\ell \in \{0, \dots, d-2\}$,
- $t > 1$ and $v = v_\ell$, for $\ell \in \{0, \dots, d-2\}$.

In particular, if $v = v_\ell = \left(\prod_{s=\ell}^{d-1} x_{n-st-1}\right) \left(\prod_{s=0}^{\ell-1} x_{n-st}\right)$ ($\ell \in \{0, \dots, d-2\}$), then

$$H = [n - (d-1)t - 1, n] \setminus \{n - \ell t - 1\} \in \Delta. \quad (17)$$

Moreover, $|H| = 1 + (d-1)t$ and Δ pure implies that $H \cup \{1\} \notin \Delta$, i.e., $\mathbf{x}_{H \cup \{1\}} \in I = I_\Delta$.

3.2.2 The classification

Now we are in position to prove the main result in the article.

Theorem 3.5 *Let $u = x_{i_1}x_{i_2}\dots x_{i_d}$, $v = x_{j_1}x_{j_2}\dots x_{j_d}$ be t -spread monomials of degree $d \geq 2$ of S with $i_1 = 1$, $j_1 > 2$ and let I be the t -spread lexsegment ideal $I = (\mathcal{L}_t(u, v))$ of S . Assume $I \neq I_{n,d,t}$. Then I is Cohen–Macaulay if and only if one of the following conditions holds true*

(a) $u = x_1x_{n-(d-2)t} \cdots x_{n-t}x_n$ and $v = x_{n-(d-1)t-1} \left(\prod_{s=0}^{d-2} x_{n-st}\right)$;

(b) $t = 1$ and for some integer $\ell \in \{0, \dots, d-2\}$,

$$u = x_1 \left(\prod_{s=\ell+1}^{d-2} x_{n-s-1}\right) \left(\prod_{s=0}^{\ell} x_{n-s}\right), \quad v = \left(\prod_{s=\ell}^{d-1} x_{n-s-1}\right) \left(\prod_{s=0}^{\ell-1} x_{n-s}\right);$$

(c) $t > 1$ and for some integer $\ell \in \{0, \dots, d-2\}$,

$$\begin{aligned} u &\in \mathcal{L}_t \left(x_1 \left(\prod_{s=\ell}^{d-2} x_{n-st-1}\right) \left(\prod_{s=0}^{\ell-1} x_{n-st}\right), x_1 \left(\prod_{s=0}^{d-2} x_{n-st}\right) \right), \\ v &= \left(\prod_{s=\ell}^{d-1} x_{n-st-1}\right) \left(\prod_{s=0}^{\ell-1} x_{n-st}\right). \end{aligned}$$

Proof. (a) Let $v = v_{d-1} = x_{n-(d-1)t-1} \left(\prod_{s=0}^{d-2} x_{n-st}\right)$. The only monomial $w \in M_{n,d,t}$ with $w < v$ is $w = \min(M_{n,d,t}) = \left(\prod_{s=0}^{d-1} x_{n-st}\right)$. Since $x_1(w/x_{n-(d-1)t}) \leq u$, then I is a completely t -spread lexsegment ideal (Theorem 1.4(2)). Moreover, I has a linear resolution (Theorem 1.4(2)). Indeed, setting $u = x_1x_{i_2}\dots x_{i_d}$, we have $i_2 > 1 + t$ and

$$x_1x_{i_2-t}x_{i_3-t}\dots x_{i_d-t} \geq x_1x_{n-(d-1)t}x_{n-(d-2)t}\dots x_{n-t} = x_1(w/x_{\max(w)}).$$

Note that the Cohen–Macaulayness of Δ implies $\dim(S/I) = \text{depth}(S/I)$. From (15), $\dim(S/I) = n - \text{height}(I) = 1 + (d-1)t = \text{depth}(S/I)$.

On the other hand, by the Auslander–Buchsbaum formula, $\text{depth}(S/I) = n - \text{pd}(S/I) = n - 1 - \text{pd}(I)$. Thus, S/I is Cohen–Macaulay if and only if $\text{pd}(I) = n - (d-1)t - 2$. Setting $a = n - 1 - (d-1)t$ and $b = a + d$, we can note that $\text{pd}(I) \leq n - (d-1)t - 1$ if and only if (Theorem 1.4(3))

$$\begin{aligned} \beta_a(I) = \beta_{a,b}(I) &= \sum_{w \in \mathcal{L}_t^f(u)} \binom{n - \min(w) - (d-1)t}{n - 1 - (d-1)t} - \sum_{\substack{w \in \mathcal{L}_t^f(v) \\ w \neq v}} \binom{\max(w) - (d-1)t - 1}{n - 1 - (d-1)t} \\ &= |\{w \in \mathcal{L}_t^f(u) : \min(w) = 1\}| - |\{w \in \mathcal{L}_t^f(v) \setminus \{v\} : \max(w) = n\}| \\ &= |\{w \in \mathcal{L}_t^f(u) : \min(w) = 1\}| - 1 = 0, \end{aligned}$$

i.e., $u = x_1 \left(\prod_{s=0}^{d-2} x_{n-st} \right)$. Condition (a) follows.

Conversely, let $u = x_1 x_{n-(d-2)t} \cdots x_{n-t} x_n$ and $v = x_{n-(d-1)t-1} \left(\prod_{s=0}^{d-2} x_{n-st} \right)$. We prove that $\dim(S/I) = \text{depth}(S/I)$.

First, using the same arguments as before, we can verify that I is a completely t -spread lexsegment ideal which has a linear resolution.

Setting $a = n - (d-1)t - 1$ and $c = a - 1 = n - (d-1)t - 2$, we have $m = |\{w \in \mathcal{L}_t^f(u) : \min(w) = 2\}| > 1$ and

$$\begin{aligned} \beta_c(I) &= \sum_{w \in \mathcal{L}_t^f(u)} \binom{n - \min(w) - (d-1)t}{n-2-(d-1)t} - \sum_{\substack{w \in \mathcal{L}_t^f(v) \\ w \neq v}} \binom{\max(w) - (d-1)t - 1}{n-2-(d-1)t} \\ &= \binom{a}{a-1} + m \binom{a-1}{a-1} - \binom{a}{a-1} = m > 0. \end{aligned}$$

by Theorem 1.4(3). Thus, $\text{pd}(I) = c = n - (d-1)t - 2$. Hence, from the Auslander–Buchsbaum formula, $\text{depth}(S/I) = 1 + (d-1)t$.

Let us prove that $\dim(S/I) = 1 + (d-1)t$. It is sufficient to verify that $\dim(\Delta) = (d-1)t$. Using the notation of Theorem 2.12, since I is a completely t -spread lexsegment ideal, we have $\mathcal{F}(\Delta) = \mathcal{G} \cup \{[n] \setminus F_p : p \in \mathcal{I}\} \cup \tilde{\mathcal{F}}$. If $F \in \mathcal{G}$, then $|F| = 1 + (d-1)t$. If $F \in \tilde{\mathcal{F}}$, then $|F| = (d-1)t < 1 + (d-1)t$. Finally, we check that $\mathcal{I} = \{1\}$ and $|[n] \setminus F_1| = 1 + (d-1)t$. Indeed, for $p = 1$, $[n] \setminus F_1 = [j_1 + 1, n] = [n - (d-1)t, n]$ has cardinality $1 + (d-1)t$, so $1 \in \mathcal{I}$. Let $2 \leq p \leq d$, then

$$\begin{aligned} [n] \setminus F_p &= \bigcup_{i=1}^{p-1} [j_i, j_i + (t-1)] \cup [j_p + 1, n] \\ &= \bigcup_{i=1}^{p-1} [j_i, j_i + (t-1)] \cup [n - (d-p)t + 1, n] \end{aligned}$$

has cardinality $(p-1)t + n + 1 - (n - (d-p)t + 1) = (d-1)t$ and moreover,

$$\min(v/x_{j_p}) = \min(v) = n - (d-2)t - 1 < n - (d-2)t - 1 \leq \min(u/x_1).$$

Thus $v/x_{j_p} > u/x_1$ and $p \notin \mathcal{I}$ and so $\mathcal{I} = \{1\}$ and $|[n] \setminus F_1| = 1 + (d-1)t$. Hence, $\dim(\Delta) = (d-1)t$ and consequently $\dim(S/I) = n - \text{height}(I) = 1 + (d-1)t = \text{depth}(S/I)$ and I is Cohen–Macaulay.

(b) Let $t = 1$. Let us consider the set H defined in (17). We note that the smallest 1-spread (squarefree) monomial of degree d that we can “extract” from $H \cup \{1\} =$

$\{1, n-d, n-d+1, \dots, n-\ell-2, n-\ell, \dots, n-1, n\}$ is

$$\begin{aligned} z &= x_1 x_{n-d+1} \cdots x_{n-\ell-2} \cdot x_{n-\ell} \cdots x_{n-1} x_n \\ &= x_1 \left(\prod_{s=\ell+1}^{d-2} x_{n-s-1} \right) \left(\prod_{s=0}^{\ell} x_{n-s} \right). \end{aligned}$$

Thus, we must have $z \in I$, and so $u \geq z$, *i.e.*,

$$u \in \mathcal{L}_t \left(x_1 \left(\prod_{s=\ell}^{d-2} x_{n-s-2} \right) \left(\prod_{s=0}^{\ell-1} x_{n-s-1} \right), x_1 \left(\prod_{s=\ell}^{d-2} x_{n-s-1} \right) \left(\prod_{s=0}^{\ell+1} x_{n-s} \right) \right).$$

Let $v = v_\ell$, for some $\ell \in \{0, \dots, d-2\}$. The monomials $w \in M_{n,d,1}$ with $w < v$ are $v_{\ell+1} > \cdots > v_d$. Since, for $r = \ell+1, \dots, d$,

$$x_1(v_r/x_{n-(d-1)}) \leq x_1 \left(\prod_{s=\ell+1}^{d-2} x_{n-s-1} \right) \left(\prod_{s=0}^{\ell} x_{n-s} \right) \leq u,$$

I is a completely 1-spread lexsegment ideal. Moreover, I has a linear resolution. Indeed, setting $u = x_1 x_{i_2} \cdots x_{i_d}$, we have $i_2 > 1+t=2$ and

$$x_1 x_{i_2-1} x_{i_3-1} \cdots x_{i_d-1} \geq x_1 \left(\prod_{s=\ell+1}^{d-2} x_{n-s-2} \right) \left(\prod_{s=0}^{\ell} x_{n-s-1} \right) > x_1(v_{\ell+1}/x_{\max(w)}).$$

From (15), $\dim(S/I) = 1+(d-1)t = d$ and, since we are assuming I Cohen–Macaulay, then $\text{depth}(I) = d$. On the other hand from the Auslander–Buchsbaum formula, S/I is Cohen–Macaulay if and only if $\text{pd}(I) = n-d-1$. Setting $a = n-d$ and $b = a+d$, $\text{pd}(I) \leq n-d$, S/I is Cohen–Macaulay if and only if

$$\begin{aligned} \beta_a(I) &= \beta_{a,b}(I) = |\{w \in \mathcal{L}_t^f(u) : \min(w) = 1\}| - |\{w \in \mathcal{L}_t^f(v) \setminus \{v\} : \max(w) = n\}| \\ &= |\{w \in \mathcal{L}_t^f(u) : \min(w) = 1\}| - |\{v_{\ell+1} > \cdots > v_d\}| \\ &= |\{w \in \mathcal{L}_t^f(u) : \min(w) = 1\}| - (d-\ell) = 0, \end{aligned}$$

i.e., $u = x_1 \left(\prod_{s=\ell+1}^{d-2} x_{n-s-1} \right) \left(\prod_{s=0}^{\ell} x_{n-s} \right)$. The assertion (b) follows.

Conversely, let $u = x_1 \left(\prod_{s=\ell+1}^{d-2} x_{n-s-1} \right) \left(\prod_{s=0}^{\ell} x_{n-s} \right)$ and $v = v_\ell$, for some $\ell \in \{0, \dots, d-2\}$. We compute $\dim(S/I)$ and $\text{depth}(S/I)$.

Using the notation of Theorem 2.12, $\mathcal{F}(\Delta) = \mathcal{G} \cup \{[n] \setminus F_p : p \in \mathcal{I}\} \cup \tilde{\mathcal{F}}$. We show that $\dim(\Delta) = d$. If $F \in \mathcal{G}$, then $|F| = d$. Arguing as in the proof of condition (b), one can observe that $\mathcal{I} = [d-\ell]$ and for $p \in \mathcal{I}$, $|[n] \setminus F_p| = d$. Furthermore, for any $F \in \mathcal{F}$, $|F| = d-1 < d$. Hence $\dim(\Delta) = d$ and so $\dim(S/I) = d$.

Setting $a = n-d$ and $c = a-1 = n-d-1$, we have $|\{w \in \mathcal{L}_t^f(u) : \min(w) = 1\}| = d-\ell$, $m = |\{w \in \mathcal{L}_t^f(u) : \min(w) = 2\}| > 1$ and

$$\begin{aligned} \beta_c(I) &= \sum_{w \in \mathcal{L}_t^f(u)} \binom{n - \min(w) - d + 1}{n - d - 1} - \sum_{\substack{w \in \mathcal{L}_t^f(v) \\ w \neq v}} \binom{\max(w) - d}{n - d - 1} \\ &= (d-\ell) \binom{a}{a-1} + m \binom{a-1}{a-1} - (d-\ell) \binom{a}{a-1} = m > 0. \end{aligned}$$

Thus, $\text{pd}(I) = n - d - 1$ and from the Auslander–Buchsbaum formula $\text{depth}(I) = d$ and I is Cohen–Macaulay. \square

For the proof of the last case (c), we need some more preparations.

First, recall that if $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ is a short exact sequence of finitely generated graded S -modules, then $\text{pd}(M) \leq \max\{\text{pd}(L), \text{pd}(N)\}$. Moreover, given a monomial ideal I of S and a monomial u of S , there exists the short exact sequence

$$0 \rightarrow S/(I : u) \rightarrow S/I \rightarrow S/(I, u) \rightarrow 0.$$

Therefore $\text{pd}(S/I) \leq \max\{\text{pd}(S/(I : u)), \text{pd}(S/(I, u))\}$.

Lemma 3.6 *Let I, J equigenerated monomial ideals of S having initial degree d . Suppose $J \subseteq I$, then*

$$\beta_{i, i+d}(J) \leq \beta_{i, i+d}(I),$$

for all $i \geq 0$. Moreover, if J has a linear resolution, $\text{pd}(I) \geq \text{pd}(J)$.

Proof. The first assertion follows from [18, Lemma 10.3.5]. For the second assertion, assume J has a linear resolution, and let $\text{pd}(J) = s$. Then $\beta_s(J) = \beta_{s, s+d}(J) \leq \beta_{s, s+d}(I)$. Finally, $\text{pd}(I) \geq s = \text{pd}(J)$. \square

Now we are in position to prove the last statement of Theorem 3.5.

(c) Let $t \geq 2$. From (17), the smallest t -spread monomial we can “extract” from $H \cup \{1\}$ is

$$z = x_1 x_{n-(d-2)t} x_{n-(d-3)t} \cdots x_{n-t} x_n.$$

Since z is the smallest t -spread monomial of S with minimum equal to 1, then $z \in I$, and there are no further conditions to impose on the monomial u . Thus

$$u \in \mathcal{L}_t(x_1(\prod_{s=\ell}^{d-2} x_{n-st-2})(\prod_{s=0}^{\ell-1} x_{n-st-1}), x_1(\prod_{s=0}^{d-2} x_{n-st})). \quad (18)$$

Setting

$$u_p = x_1 \left(\prod_{s=p}^{d-2} x_{n-st-1} \right) \left(\prod_{s=0}^{p-1} x_{n-st} \right), \quad p = \ell, \dots, d-1,$$

we have that

$$\mathcal{L}_t(x_1(\prod_{s=\ell}^{d-2} x_{n-st-1})(\prod_{s=0}^{\ell-1} x_{n-st}), x_1(\prod_{s=0}^{d-2} x_{n-st})) = \{u_\ell > u_{\ell+1} > \cdots > u_{d-1}\}.$$

We prove that if

$$u > u_\ell = x_1(\prod_{s=\ell}^{d-2} x_{n-st-1})(\prod_{s=0}^{\ell-1} x_{n-st}), \quad (19)$$

then I is not Cohen–Macaulay.

By hypothesis $v = v_\ell$ ($\ell = 0, \dots, d-2$) and from (19)

$$u \geq x_1 \left(\prod_{s=\ell-1}^{d-2} x_{n-st-1} \right) \left(\prod_{s=0}^{\ell-2} x_{n-st} \right),$$

where $x_1 \left(\prod_{s=\ell-1}^{d-2} x_{n-st-1} \right) \left(\prod_{s=0}^{\ell-2} x_{n-st} \right)$ is the greatest monomial with minimum equal to 1 preceding u_ℓ , with respect to the squarefree lex order. The monomials $w \in M_{n,d,t}$ smaller than v are $v_{\ell+1} > \dots > v_d$ and for each v_q , $q = \ell+1, \dots, d$, we have

$$u \geq x_1 \left(\prod_{s=\ell-1}^{d-2} x_{n-st-1} \right) \left(\prod_{s=0}^{\ell-2} x_{n-st} \right) = x_1(v_{\ell+1}/x_{\min(v_{\ell+1})}) > x_1(v_q/x_{\min(v_q)}).$$

Hence, setting $u = x_1 x_{i_2} \cdots x_{i_d}$, we have $i_2 > 1+t$ and

$$\begin{aligned} x_1 x_{i_2-t} \cdots x_{i_d-t} &\geq x_1 x_{n-(d-1)t-1} \cdots x_{n-\ell t-1} \cdot x_{n-(\ell-1)t} \cdots x_{n-t} \\ &= x_1(v_\ell/x_{\max(v_\ell)}) \geq x_1(v_{\ell+1}/x_{\max(v_{\ell+1})}). \end{aligned}$$

Therefore, I is a completely t -spread lexsegment ideal with a linear resolution. Setting $a = n-1-(d-1)t$ and $b = a+d$, we have

$$\begin{aligned} \beta_a(I) = \beta_{a,b}(I) &= |\{w \in \mathcal{L}_t^f(u) : \min(w) = 1\}| - |\{w \in \mathcal{L}_t^f(v) \setminus \{v\} : \max(w) = n\}| \\ &\geq |\{u > u_\ell > \dots > u_{d-1}\}| - (d-\ell) = d+1-\ell - (d-\ell) = 1. \end{aligned}$$

It follows that $\text{pd}(I) = n-1-(d-1)t$ and consequently I is not Cohen–Macaulay. Condition (c) follows.

Conversely, assume (c) holds. We prove that $\dim(S/I) = 1+(d-1)t$. This is equivalent to verify that $\dim(\Delta) = (d-1)t$. For this aim it suffices to show the following facts:

- (i) there exists $F \in \mathcal{F}(\Delta)$ with cardinality $|F| = 1+(d-1)t$;
- (ii) for any $A \subseteq [n]$ with $|A| = 2+(d-1)t$, then $A \notin \Delta$.

(i) Let us consider $F = [n-(d-1)t, n]$. By the structure of v , it follows that $F \in \Delta$ and furthermore $|F| = 1+(d-1)t$, as desired.

(ii) Let $A \subseteq [n]$ with $|A| = 2+(d-1)t$. We have $1 \leq \min(A) \leq n-(d-1)t-1$. If $\min(A) = n-(d-1)t-1$, then $A = [n-(d-1)t-1, n] \notin \Delta$, as

$$x_{n-(d-1)t-1} x_{n-(d-2)t-1} \cdots x_{n-t-1} x_{n-1} \in I.$$

If $1 < \min(A) < n-(d-1)t-1$, setting $\ell_1 = \min(A)$ and $\ell_r = \min\{\ell \in A : \ell \geq \ell_{r-1} + t\}$, for $r \geq 2$, the sequence $L : \ell_1 > \ell_2 > \dots$ has at least d terms, and $\mathbf{x}_L \in I$. Thus, $A \notin \Delta$.

If $\min(A) = 1$, we consider the sequence L above defined. If $\mathbf{x}_L \leq u$, then $A \notin \Delta$. Otherwise, set $B = A \setminus \{1\}$.

If $\min(B) = n - (d-1)t$, then $A = \{1\} \cup B = \{1\} \cup [n - (d-1)t, n] \notin \Delta$, as $x_1 x_{n-(d-2)t} \cdots x_{n-t} x_n \in I$. If $\min(B) < n - (d-1)t - 1$, setting $\nu_1 = \min(B)$, and $\nu_r = \min\{\nu \in B : \nu \geq \nu_{r-1} + t\}$, for $r \geq 2$, the sequence $N : \nu_1 > \nu_2 > \cdots$ has at least d terms, and $\mathbf{x}_N \in I$, implying that $A \notin \Delta$.

Let $\min(B) = n - (d-1)t - 1$. Then $A = \{1\} \cup B = \{1\} \cup [n - (d-1)t - 1, n] \setminus \{m\}$, for some $m > n - (d-1)t - 1$. We can write $m = n - pt - q$, with $0 \leq p \leq d-3$ and $0 \leq q \leq t-1$. If $q \geq 2$, then $\text{supp}(v) \subseteq B \subseteq A$, and $A \notin \Delta$. If $q = 1$, then $\{1, n - (d-2)t, \dots, n-t, n\} \subseteq A$, and $x_1 x_{n-(d-2)t} \cdots x_{n-t} x_n \in I$, thus $A \notin \Delta$. If $q = 0$, we consider p . If $p < \ell - 1$, the structure of v implies that $\text{supp}(v) \subseteq A$, and once again $A \notin \Delta$. If $p \geq \ell - 1$, then $C = \{n - kt - 1 : k = 0, \dots, d-1\} \subseteq A$, and $\mathbf{x}_C > v$, so $\mathbf{x}_C \in I$, implying $A \notin \Delta$. Finally, in all possible cases, $A \notin \Delta$, and (ii) follows.

Finally, $\dim(S/I) = 1 + (d-1)t$.

Now, set $I_{u_p} = (\mathcal{L}_t(u_p, v))$, $p = \ell, \dots, d-1$. We prove that I_{u_p} is Cohen–Macaulay, *i.e.*, $\text{pd}(I_{u_p}) = n - 2 - (d-1)t$.

First, assume $p = \ell$. We have

$$u = u_\ell = x_1(v_\ell/x_{\min(v_\ell)}) > x_1(v_q/x_{\min(v_q)}), \quad q = \ell + 1, \dots, d$$

and setting $u = x_1 x_{i_2} \cdots x_{i_d}$, it follows that

$$x_1 x_{i_2-t} \cdots x_{i_d-t} = x_1(v_\ell/x_{\max(v_\ell)}).$$

Thus, I_{u_ℓ} is a completely t -spread lexsegment ideal with a linear resolution. Set $a = n - 1 - (d-1)t$, $b = a + d$, we have

$$\begin{aligned} \beta_a(I_{u_\ell}) &= \beta_{a,b}(I_{u_\ell}) = |\{w \in \mathcal{L}_t^f(u) : \min(w) = 1\}| - |\{w \in \mathcal{L}_t^f(v) \setminus \{v\} : \max(w) = n\}| \\ &= |\{u_\ell > u_{\ell+1} > \cdots > u_{d-1}\}| - |\{v_{\ell+1} > \cdots > v_d\}| = 0, \end{aligned}$$

and, as $m = |\{w \in \mathcal{L}_t^f(u) : \min(w) = 2\}| > 1$, then

$$\begin{aligned} \beta_{a-1}(I_{u_\ell}) &= \sum_{w \in \mathcal{L}_t^f(u)} \binom{n - \min(w) - (d-1)t}{n - 2 - (d-1)t} - \sum_{\substack{w \in \mathcal{L}_t^f(v) \\ w \neq v}} \binom{\max(w) - (d-1)t - 1}{n - 2 - (d-1)t} \\ &= (d-\ell) \binom{a}{a-1} + m \binom{a-1}{a-1} - (d-\ell) \binom{a}{a-1} = m > 0. \end{aligned}$$

Thus, $\text{pd}(I_{u_\ell}) = a - 1 = n - (d-1)t - 2$, and I_{u_ℓ} is Cohen–Macaulay, as desired.

Now let $p = \ell + 1$. Firstly, observe that $J = (\mathcal{L}_t(x_2 x_{2+t} \cdots x_{2+(d-1)t}, v)) \subseteq I_{u_{\ell+1}}$. Moreover, J is an initial t -spread lexsegment ideal in $K[x_2, \dots, x_n]$ with $\text{pd}(J) =$

$n - 2 - (d - 1)t$. Thus J has a linear resolution, and Lemma 3.6 implies $\text{pd}(I_{u_{\ell+1}}) \geq n - 2 - (d - 1)t$. For the other inequality, consider the short exact sequence:

$$0 \rightarrow S/(I_{u_{\ell+1}} : u_{\ell}) \rightarrow S/I_{u_{\ell+1}} \rightarrow S/(I_{u_{\ell+1}}, u_{\ell}) \rightarrow 0.$$

Observe that $S/(I_{u_{\ell+1}}, u_{\ell}) = S/I_{u_{\ell}}$, and $\text{pd}(S/I_{u_{\ell}}) = \text{pd}(I_{u_{\ell}}) + 1 = n - 1 - (d - 1)t$. Let us verify that $I_{u_{\ell+1}} : u_{\ell} = \mathfrak{p}_{[2, n - (d - 1)t - 1] \cup \{n - \ell t\}}$ is a complete intersection and $\text{pd}(S/(I_{u_{\ell+1}} : u_{\ell})) = n - 1 - (d - 1)t$.

By [18, Proposition 1.2.2], a set of generators for $I_{u_{\ell+1}} : u_{\ell}$ is given by

$$\{w/\text{gcd}(w, u_{\ell}) : w \in G(I_{u_{\ell+1}})\}.$$

If $w = u_p$, $p = \ell + 1, \dots, d - 1$, we have $u_p/\text{gcd}(u_p, u_{\ell}) = x_{n - (p - 1)t} \cdots x_{n - \ell t}$ and for $p = \ell + 1$, $u_{\ell+1}/\text{gcd}(u_{\ell+1}, u_{\ell}) = x_{n - \ell t}$, thus $x_{n - \ell t}$ divides all these generators.

If $w \in \mathcal{L}_t(x_2 x_{2+t} \cdots x_{2+(d-1)t}, v)$, then $\min(w) \in [2, n - (d - 1)t - 1]$. Note that $[2, n - (d - 1)t - 1] \cap \text{supp}(u_{\ell}) = \emptyset$, thus $x_{\min(w)}$ divides $w/\text{gcd}(w, u_{\ell})$ and for $z = x_{\min(w)}(u_{\ell}/x_1) \in G(I_{u_{\ell+1}})$, we have that $z/\text{gcd}(z, u_{\ell}) = x_{\min(w)} \in I_{u_{\ell+1}} : u_{\ell}$.

Finally, we have verified that $I_{u_{\ell+1}} : u_{\ell} = \mathfrak{p}_{[2, n - (d - 1)t - 1] \cup \{n - \ell t\}}$ and consequently $\text{pd}(S/(I_{u_{\ell+1}} : u_{\ell})) = n - 1 - (d - 1)t$.

Hence,

$$\text{pd}(S/I_{u_{\ell+1}}) \leq \max \{\text{pd}(S/I_{u_{\ell}}), \text{pd}(S/(I_{u_{\ell+1}} : u_{\ell}))\} = n - 1 - (d - 1)t,$$

so $\text{pd}(I_{u_{\ell+1}}) \leq n - 2 - (d - 1)t$. Thus $\text{pd}(I_{u_{\ell+1}}) = n - 2 - (d - 1)t$ and $I_{u_{\ell+1}}$ is Cohen–Macaulay. For $p \in \{\ell + 2, \dots, d\}$ the same arguments work and the proof of Theorem 3.5 is complete.

4. CONCLUSIONS AND PERSPECTIVES

In this article, we have investigated the Cohen–Macaulayness of t –spread lexsegment ideals using the theory of simplicial complexes. The completely t –spread lexsegment ideals have played an essential role. Non–completely t –spread lexsegment ideals are much less understood. Here are some possible questions to investigate.

Question 4.1 Determine the standard primary decomposition of non–completely t –spread lexsegment ideals.

Question 4.2 Classify the pure simplicial complexes associated to t –spread lexsegment ideals.

In [19], Olteanu classified all *sequentially Cohen–Macaulay* squarefree completely lexsegment ideals. To the best of our knowledge, a classification for squarefree non–completely lexsegment ideals is unknown. More generally, we ask the following

Question 4.3 Classify all sequentially Cohen–Macaulay t –spread lexsegment ideals.

Finally, due to many examples that we performed and our results on completely t –spread lexsegment ideals, we are led to conjecture the following:

Conjecture 4.4 Let I be a t –spread lexsegment ideal generated in degree $d \geq 2$. Then $\dim(S/I) \geq (d - 1)t$.

Note that Conjecture 4.4 is trivially true when $t = 1$. Indeed, if Δ is the simplicial complex associated to I , then each set $A \subseteq [n]$, with $|A| = d - 1$, is in Δ . Thus $\dim(S/I) \geq d - 1$. Conjecture 4.4 is also true for completely t –spread lexsegment ideals by virtue of Theorems 2.1, 2.7, 2.12.

Acknowledgement. We thank the the referee for his/her helpful suggestions that allowed us to improve the quality of the paper.

REFERENCES

- [1] L. AMATA. *Computational methods for t –spread monomial ideals*, available at arXiv preprint <https://arxiv.org/abs/2110.11801> [math.AC].
- [2] L. AMATA, M. CRUPI, A. FICARRA. *Upper bounds for extremal Betti numbers of t –spread strongly stable ideals*, Bull. Math. Soc. Sci. Math. Roumanie, **65** (113), No. 1 (2022), 13–34.
- [3] L. AMATA, M. CRUPI, A. FICARRA. *Projective dimension and Castelnuovo–Mumford regularity of t –spread ideals*, International Journal of Algebra and Computation **32**, No. 4 (2022), 837–858.
- [4] L. AMATA, A. FICARRA, M. CRUPI. *A numerical characterization of extremal Betti numbers of t –spread strongly stable ideals*, J. Algebraic Combin. **32**(4) (2022), 837–858.
- [5] C. ANDREI, V. ENE, B. LAJMIRI. *Powers of t –spread principal Borel ideals*, Archiv der Mathematik **112** (6) (2019), 587–597.
- [6] C. ANDREI–CIOBANU, *Kruskal–Katona Theorem for t –spread strongly stable ideals*, Bull.Math. Soc. Sci. Math. Roumanie (N.S.) **62**(110)(2) (2019), 107–122.
- [7] A. ARAMOVA, E. DE NEGRI, J. HERZOG, *Lexsegment ideals with a linear resolution*, Illinois J. Math. **42**(3) (1998), 509–52.
- [8] A. ARAMOVA, J. HERZOG, T. HIBI. *Squarefree lexsegment ideals*. Math.Z. **228** (1998), 353–378.
- [9] V. BONANZINGA, L. SORRENTI, N. TERAJ, *Pure and Cohen–Macaulay simplicial complexes associated with squarefree lexsegment Ideals*, Comm. Alg. **40**(11) (2012), 4195–4214
- [10] E. DE NEGRI, J. HERZOG, *Completely lexsegment ideals*, Proc. Amer. Math. Soc., **126** (12) (1998), 3467–3473.

- [11] R. DINU, *Gorenstein T -spread Veronese algebras*, Osaka J. Math. **57**(4) (2020), 935–947.
- [12] S. ELIAHOU, M. KERVAIRE, *Minimal resolutions of some monomial ideals*, J. Algebra 129 (1990), 1–25.
- [13] V. ENE, J. HERZOG, A. A. QURESHI, *t -spread strongly stable monomial ideals*, Comm. Algebra **47**(12), 5303–5316 (2019).
- [14] V. ENE, A. OLTEANU, L. SORRENTI, *Properties of lexsegment ideals*, Osaka J. Math. **47** (1) (2010) 67–87.
- [15] A. FICARRA, M. CRUPI, *Shifting operations and completely t -spread lexsegment ideals*, Comm. Alg. **50**(8) (2022), 3320–3337.
- [16] C. A. FRANCISCO, H. T. HA, A. VAN TUYL, *Splittings of monomial ideals*, Proc. Amer. Math. Soc., **137** (10) (2009), 3271–3282.
- [17] D. R. GRAYSON, M. E. STILLMAN, *Macaulay2, a software system for research in algebraic geometry*. Available at <http://www.math.uiuc.edu/Macaulay2>.
- [18] J. HERZOG, T. HIBI, *Monomial ideals*, Graduate texts in Mathematics **260**, Springer–Verlag, 2011.
- [19] O. OLTEANU, *Classes of sequentially Cohen–Macaulay squarefree monomial ideals*, Algebra Colloq. **21** (04) (2011), 575–590.