

# CAPACITY OF THE RANGE OF RANDOM WALK: THE LAW OF THE ITERATED LOGARITHM

AMIR DEMBO AND IZUMI OKADA

**ABSTRACT.** We establish both the lim sup and the lim inf law of the iterated logarithm (LIL), for the capacity of the range of a simple random walk in any dimension  $d \geq 3$ . While for  $d \geq 4$ , the order of growth in  $n$  of such LIL at dimension  $d$  matches that for the volume of the random walk range in dimension  $d - 2$ , somewhat surprisingly this correspondence breaks down for the capacity of the range at  $d = 3$ . We further establish such LIL for the Brownian capacity of a 3-dimensional Brownian sample path and novel, sharp moderate deviations bounds for the capacity of the range of a 4-dimensional simple random walk.

## 1. INTRODUCTION AND MAIN RESULTS

Let  $\tau_A$  denote the first positive hitting time of a finite set  $A$  by a simple random walk (SRW) on  $\mathbb{Z}^d$ , denoted hereafter  $(S_m)_{m \geq 0}$ . Recall that the corresponding (Newtonian) capacity is given for  $d \geq 3$ , by

$$\text{Cap}(A) := \sum_{x \in A} P^x(\tau_A = \infty) = \lim_{|z| \rightarrow \infty} \frac{P^z(\tau_A < \infty)}{G(0, z)}.$$

The asymptotics of the capacity  $R_n := \text{Cap}(\mathcal{R}_n)$  of the random walk range  $\mathcal{R}_n := \{S_1, \dots, S_n\}$  is relatively trivial for  $d = 2$  (for then  $R_n = \frac{2+\alpha(1)}{\pi} \log(\text{diam} \mathcal{R}_n)$ , see [23, Lemma 2.3.5]). In contrast, for  $d \geq 3$  such asymptotic is of an on going interest. Indeed, applying ergodic theory, over 50 years ago Jain and Orey [20] established the strong law

$$\lim_{n \rightarrow \infty} \frac{R_n}{n} = \alpha_d \quad \text{a.s.}, \quad \text{for all } d \geq 3,$$

where  $\alpha_d > 0$  iff  $d \geq 5$ . Recall Green's function for the 3-dimensional Brownian motion

$$G_B(x, y) := \int_0^\infty (2\pi t)^{-d/2} e^{-|x-y|^2/(2t)} dt = \frac{1}{2\pi} |x - y|^{-1},$$

and the corresponding Brownian capacity of  $D \subset \mathbb{R}^3$ ,

$$\text{Cap}_B(D)^{-1} := \inf \left\{ \int \int G_B(x, y) \mu(dx) \mu(dy) : \mu(D) = 1 \right\}.$$

---

*Date:* August 2, 2022.

*2010 Mathematics Subject Classification.* 60F15, 60G50.

*Key words and phrases.* Capacity, random walk, Brownian motion, law of iterated logarithm.

Research supported in part by NSF grant DMS-1954337 (A.D.) and by JSPS KAKENHI Grant-in-Aid for Early-Career Scientists (No. JP20K14329) and JSPS Overseas Research Fellowship (I.O.).

More recently, Chang [13] showed that for  $d = 3$ ,

$$\frac{R_n}{\sqrt{n}} \xrightarrow{\mathcal{D}} \frac{1}{3\sqrt{3}} \text{Cap}_B(B[0, 1]),$$

whereas Asselah et al [4] showed that in this case further, for some  $C$  finite,

$$C^{-1} \sqrt{n} \leq E[R_n] \leq C \sqrt{n}.$$

We denote throughout by  $\bar{X}$  the centering  $\bar{X} = X - E[X]$  of a generic random variable  $X$ . In higher dimensions  $d \geq 4$ , the centered capacity  $\bar{R}_n$ , converges after proper scaling to a non-degenerate limit law, which is Gaussian iff  $d \geq 5$  (see [6] for  $d = 4$  and [4, 32] for  $d \geq 5$ ). For  $d \geq 5$ , estimates of the corresponding large and moderate deviations are provided in [1] (but they are not sharp enough to imply a LIL), while the central limit theorem (CLT) is further established in [16] for  $R_n$  and a class of symmetric  $\alpha$ -stable walks, provided  $d > 5\alpha/2$ . We also note in passing the asymptotics in [9, 7, 8] of  $R_n$  for a critical branching random walk on  $\mathbb{Z}^d$ ,  $d \geq 3$ , conditional to have total population size  $n$ .

In view of these works, a natural question, which we fully resolve here, is to determine the almost sure fluctuations of  $n \mapsto R_n$  for the srw, in the form of some LIL (possibly after centering  $R_n$  when  $d \geq 4$ ). Specifically, using hereafter  $\log_k a = \log(\log_{k-1} a)$  for  $k \geq 2$ , with  $\log_1 a$  for the usual logarithm, here is our first main result, about the srw in  $\mathbb{Z}^3$ .

**Theorem 1.1.** *For  $d = 3$ , almost surely,*

$$(1.1) \quad \limsup_{n \rightarrow \infty} \frac{R_n}{h_3(n)} = 1, \quad \liminf_{n \rightarrow \infty} \frac{R_n}{\hat{h}_3(n)} = 1,$$

where

$$(1.2) \quad h_3(n) := \frac{\sqrt{6}\pi}{9} (\log_3 n)^{-1} \sqrt{n \log_2 n}, \quad \hat{h}_3(n) := \frac{\sqrt{6}\pi^2}{9} \sqrt{n (\log_2 n)^{-1}}.$$

Utilizing (3.21), we also get from Theorem 1.1 the following consequence about the Brownian capacity of the 3-dimensional (Brownian) sample path.

**Corollary 1.2.** *For  $d = 3$ , almost surely,*

$$\limsup_{n \rightarrow \infty} \frac{\text{Cap}_B(B[0, n])}{3\sqrt{3}h_3(n)} = 1, \quad \liminf_{n \rightarrow \infty} \frac{\text{Cap}_B(B[0, n])}{3\sqrt{3}\hat{h}_3(n)} = 1.$$

We next provide the LIL for the centered capacity  $\bar{R}_n$  of the range, first in case of the srw on  $\mathbb{Z}^4$  and then for srw on  $\mathbb{Z}^d$ ,  $d \geq 5$ .

**Theorem 1.3.** *For  $d = 4$ , almost surely,*

$$(1.3) \quad \limsup_{n \rightarrow \infty} \frac{\bar{R}_n}{h_d(n)} = 1, \quad \liminf_{n \rightarrow \infty} \frac{\bar{R}_n}{\hat{h}_d(n)} = -1,$$

where for some non-random  $0 < c_\star < \infty$ ,

$$(1.4) \quad h_4(n) := \frac{\pi^2 n \log_3 n}{8 (\log n)^2}, \quad \hat{h}_4(n) := c_\star \frac{n \log_2 n}{(\log n)^2}.$$

**Theorem 1.4.** *For any  $d \geq 5$ , the LIL-s (1.3) hold almost surely, now with*

$$(1.5) \quad h_d(n) = \hat{h}_d(n) := \sigma_d \sqrt{2n(1 + 1_{\{d=5\}} \log n) \log_2 n}, \quad d \geq 5,$$

where the non-random, finite  $\sigma_d^2 > 0$  are given by the leading asymptotic of  $\text{var}(R_n)$  (c.f. [32, Thm. A] for  $\sigma_5$  and [4, Thm. 1.1] for  $\sigma_d$ ,  $d \geq 6$ ).

**Remark 1.5.** *Our proof of Theorem 1.4 via Skorokhod embedding, also yields Strassen's LIL for the a.s. set of limit points in  $C([0, 1])$  of the functions  $\{t \mapsto h_d(n)^{-1} \bar{R}_n\}$ , for any  $d \geq 5$ .*

For SRW on  $\mathbb{Z}^4$  it is shown in [6, Thm. 1.2] that the moment generating function (MGF),

$$(1.6) \quad M_G(\lambda) := E[e^{\lambda \gamma_G([0, 1]^2)}],$$

of the non-Gaussian limit of  $((\log n)^2/n) \bar{R}_n$ , is infinite for all large enough  $\lambda$ . En route to Theorem 1.3, we establish the following complementary exponential tail result for  $\gamma_G([0, 1]^2)$ , which is potentially of independent interest (see also Lemma 4.6, where we establish the uniform in  $n$  boundedness of the MGF of  $((\log n)^2/n) \bar{R}_n$ ).

**Proposition 1.6.** *For  $M_G(\cdot)$  of (1.6), the value of*

$$\lambda_o = \sup\{\lambda > 0 : M_G(\lambda) < \infty\}$$

*is strictly positive and finite.*

We note that  $R_n \approx nE[\hat{P}^{S_{n/2}}(\hat{\tau}_{\mathcal{R}_n} = \infty)]$  at any fixed  $n \gg 1$  and  $d \geq 3$ , where  $\hat{P}$  and  $\hat{\tau}_A$  denote the law and the first hitting time by an i.i.d. copy of the srw. Similarly, the volume of  $\mathcal{R}_n$  in any dimension ( $d \geq 1$ ), is approximately  $nP(\tau_0 > n)$ . It has been observed before, see for example [4, Sec. 6], that the typical order of growth of  $E[\hat{P}^{S_{n/2}}(\hat{\tau}_{\mathcal{R}_n} = \infty)]$  at any  $d \geq 3$ , matches that of  $P(\tau_0 > n)$  at  $d' = d - 2$ , yielding the same order of growth in  $n$  for  $R_n$  at  $d \geq 3$  and for the volume of  $\mathcal{R}_n$  at  $d' = d - 2$ . In Theorems 1.3 and 1.4, our LIL for  $d \geq 4$  adheres to such a match with the scale for the LIL of the volume of  $\mathcal{R}_n$  at  $d' = d - 2$  (see [10, 11, 21] for the latter LIL at any  $d' \geq 2$ , as well as the limit distribution results for the volume of the Wiener sausage at  $d' \geq 2$ , and the corresponding LIL at  $d' \geq 3$ , in [25] and [15, 17], respectively). In contrast, this relation *breaks down at the lim sup LIL for  $d = 3$* , with the appearance of the novel factor  $(\log_3 n)^{-1}$  in Theorem 1.1. Nevertheless, even at  $d = 3$  the relevant deviations of  $R_n$  are due to those in the diameter of  $\mathcal{R}_n$ , except that the upper tails for these two variables differ in their growth rates. Specifically, our proofs in Sections 3.1 and 3.3 yield the following (sharper) result.

**Proposition 1.7.** *Let  $M_n := \max_{1 \leq i \leq n} |S_i|$ . For SRW of  $\mathbb{Z}^3$  and any  $\epsilon > 0$ ,*

$$P(\{R_n \geq (1 - \epsilon)h_3(n)\} \cap \{M_n \geq (1 - \epsilon)\psi(n)\} \text{ i.o.}) = 1,$$

$$P(\{R_n \leq (1 + \epsilon)\hat{h}_3(n)\} \cap \{M_n \leq (1 + \epsilon)\hat{\psi}(n)\} \text{ i.o.}) = 1,$$

where  $\psi(n) := \sqrt{(2/3)n \log_2 n}$  and  $\hat{\psi}(n) := \pi \sqrt{(1/6)n(\log_2 n)^{-1}}$ .

Proposition 1.7 implies that almost surely, the lim sup (resp. lim inf) of  $R_n$  are essentially attained simultaneously with those for  $M_n$ , since for  $d = 3$ , almost surely,

$$(1.7) \quad \limsup_{n \rightarrow \infty} \frac{M_n}{\psi(n)} = 1, \quad \liminf_{n \rightarrow \infty} \frac{M_n}{\hat{\psi}(n)} = 1.$$

Indeed, by the invariance principle it suffices for proving (1.7) to show the equivalent a.s. statement for 3-dimensional Bessel process, and the latter follows by mimicking the proof of Chung's one-dimensional LIL (see [14]), starting with the estimate (3.24). We note in passing that in addition to explicitly characterizing the value of  $c_\star$  in Theorem 1.3, the following analog of Proposition 1.7 in case  $d \geq 4$ , is also still open.

**Open problem 1.8.** Consider the SRW  $S_i = (S_i^1, \dots, S_i^d) \in \mathbb{Z}^d$ ,  $d \geq 4$ . For  $d' = d - 2$ , let  $\hat{S}_i^{d'} = (S_i^1, \dots, S_i^{d'}, 0, 0)$  and  $V_{d'}(n) = |\{\hat{S}_1^{d'}, \dots, \hat{S}_n^{d'}\}|$ . Pick non-random  $\psi_{d'}(n)$  such that a.s.

$$\limsup_{n \rightarrow \infty} \frac{\bar{V}_{d'}(n)}{\psi_{d'}(n)} = 1.$$

We then conjecture that for  $h_d(n)$  of Theorems 1.3-1.4 and any  $\epsilon > 0$ ,

$$P(\{\bar{R}_n \geq (1 - \epsilon)h_d(n)\} \cap \{\bar{V}_{d'}(n) \geq (1 - \epsilon)\psi_{d'}(n)\} \text{ i.o.}) = 1.$$

While we consider throughout only the discrete time SRW whose increments are the  $2d$  neighbors of the origin in  $\mathbb{Z}^d$ , due to sharp concentration of Poisson variables, all our results apply also for the continuous time SRW with i.i.d. Exponential(1) clocks and up to the scaling  $n \mapsto (1 - \rho)n$ , also to the  $\rho$ -lazy discrete time SRW. By definition of  $R_n$ , our results apply to any random walk on a group with a finite symmetric set of generators, whose words are isomorphic to those of the SRW (e.g. an invariance of our results under any non-random, invertible affine transformation of the walk). We note in passing the recent work [28] on the strong law for any symmetric random walk on a group of growth index  $d$  and the corresponding CLT in case  $d \geq 6$ , suggesting the possibility of a future extension of our LIL-s in this context.

Beyond the intrinsic interest in  $R_n$ , its asymptotic is also relevant for the study of intersections between two independent random walks (e.g., see [23, Ch. 3]). Similarly, [2, 3] utilize bounds on  $R_n$  to gain insights about the so called Swiss cheese picture for  $d = 3$ . Further, to understand Sznitman's [33] random interlacement model, one may use moment estimates for the capacity of the union of ranges (c.f. [13] and the references therein). Finally, the capacity equals the summation of all entries of the inverse of the (positive definite) Green's function matrix (see (2.2)), a point of view which [29] uses, for  $d = 2$ , to estimate the geometry of late points of the walk.

As for the organization of this paper, we prove Theorem 1.1 in Section 3, relying on novel relations between the capacity and Green's function which we explore in Section 2. Our proof of Theorem 1.1 further indicates that the lim sup-LIL is due to exceptional time where  $\mathcal{R}_n$  has a cylinder-like shape, with one dimension being about  $\psi(n)$  while the other two are  $O(\psi(n)/(\log_2 n))$  (see Lemma 2.1 and Section 3.2). In contrast, the lim inf-LIL seems to be due to times where the shape of  $\mathcal{R}_n$  is close enough to a ball of radius  $\hat{\psi}(n)$  to approximately match the capacity of such a ball (see (3.31) and (3.32)).

Sections 4 and 5 are devoted to the proofs of Theorems 1.3 and 1.4, respectively. Our proofs rely on the decomposition (4.2) of  $R_{n_k}$  as the sum of  $k$  independent variables  $\{U_j\}$  which are the capacities of the walk restricted to the  $k$  parts of a partition of  $[1, n_k]$ , minus some random  $\Delta_{n_k, k} \geq 0$  (which ties all these parts together). For any  $d \geq 5$  the effect of  $\Delta_{n_k, k}$  on the LIL is negligible, so upon coupling  $R_{n_k}$  with a one-dimensional Brownian motion, we immediately get the LIL for the former out of the standard LIL for the latter. As seen in Section 4, the situation is *way more delicate* for  $d = 4$ , where  $E\Delta_{n_k} \approx h_4(n)$  dominates for a suitable

slowly growing  $k = k_n$  the fluctuations of the i.i.d.  $\{U_j\}$ . The lim sup-LIL is then due to the exceptional (random) sequence  $\{n_k\}$  where  $\Delta_{n_k, k} = o(h_4(n_k))$ , while the lim inf-LIL is due to the exceptional  $\{n_k\}$  for which  $\Delta_{n_k, k} \approx \hat{h}_4(n_k) \gg E\Delta_{n_k, k}$ . Indeed, whereas Theorem 1.3 is proved via the framework developed in [11, Section 4] for the LIL for the volume of  $\mathcal{R}_n$  in the planar case ( $d' = 2$ ), special care is needed here in order to establish tight control on the moderate deviations of  $\Delta_{n, k}$  and  $U_j$  in case  $d = 4$  (c.f. Lemmas 4.1, 4.3, 4.6 and 4.7, which may be of independent interest).

## 2. CAPACITY GEOMETRY AND GREEN'S FUNCTION

The following asymptotic for the 3-dimensional capacity of cylinder-like domains (which we prove at the end of this section), is behind the factor  $(\log_3 n)^{-1}$  in the lim sup-LIL of (1.1).

**Lemma 2.1.** *For  $m \geq 1$  and  $r \geq k \in \mathbb{N}$ , let*

$$C_m(\ell, r) := (\ell\mathbb{Z})^3 \cap \{(x_1, x_2, x_3) : x_1^2 + x_2^2 \leq r^2, 1 \leq x_3 \leq m\}.$$

*Fix  $b < 2/3$ ,  $r_m = o(m)$ ,  $r_m \uparrow \infty$ . If  $C_m(1, r_m) \supseteq C_m \supseteq C_m(\ell, r_m)$  for some  $\ell \leq r_m^b$ , then*

$$(2.1) \quad \lim_{m \rightarrow \infty} \frac{\text{Cap}(C_m)}{m(\log(m/r_m))^{-1}} = \frac{\pi}{3}.$$

**Remark 2.2.** *In the sequel we prove a stronger result, namely that the upper bound in (2.1) holds as soon as  $C_m$  is contained in a union  $C_m^*(r_m)$  of at most  $m/r_m$  balls  $\mathbb{B}(z_i, r_m)$  of radius  $r_m$  in  $\mathbb{Z}^3$ , of centers such that  $|z_{i+1} - z_i| \leq r_m$  for  $1 \leq i < m/r_m$  (where  $C_m(1, r)$  is merely one possible choice for  $C_m^*(r)$ ).*

Indeed, in Section 3 we will see that lim sup of  $R_n$  is roughly attained on the event  $\{S_n^i \geq \psi(n)\}$  for  $\psi(n)$  of Proposition 1.7, with  $\mathcal{R}_n$  then having approximately the shape of such  $C_m$  for  $m = \psi(n)$ , and  $r_m = cm/\log_2 n$ , hence from Lemma 2.1 we find that

$$R_n \approx \text{Cap}(C_m) \approx \frac{\pi}{3} m(\log(m/r_m))^{-1} \approx \frac{\pi}{3} \psi(n)(\log_3 n)^{-1},$$

which is precisely  $h_3(n)$  of Theorem 1.1.

We proceed with two lemmas relating the capacity of srw with its Green's function,

$$G(x, y) = \sum_{i=0}^{\infty} P^x(S_i = y).$$

To this end, partition  $\Omega$  by the last time the walk visits  $X = \{x_1 \neq x_2 \cdots \neq x_j\}$ , to get that

$$(2.2) \quad 1 = \sum_{\ell=1}^j G(x_i, x_\ell) P^{x_\ell}(\tau_X = \infty), \quad \forall 1 \leq i \leq j.$$

**Lemma 2.3.** *For any set  $X = \{x_1, \dots, x_j\}$ , with  $\{x_j\}$  not necessarily disjoint,*

$$(2.3) \quad \frac{j}{\max_{1 \leq \ell \leq j} \{\sum_{i=1}^j G(x_i, x_\ell)\}} \leq \text{Cap}(X) \leq \frac{j}{\min_{1 \leq \ell \leq j} \{\sum_{i=1}^j G(x_i, x_\ell)\}}.$$

*Proof.* The set  $X$  of size  $|X| = k \leq j$  consists wlog of disjoint points  $\hat{X} = \{\hat{x}_1 \neq \hat{x}_2 \cdots \neq \hat{x}_k\}$ , where  $\hat{x}_v$  appears  $m_v \geq 1$  times in  $X$  (and  $\sum_{v \leq k} m_v = j$ ). Setting  $v(\ell)$  for the index of  $x_\ell$  in  $\hat{X}$  and  $q_v := (m_v)^{-1} P^{\hat{x}_v}(\tau_{\hat{X}} = \infty)$ , we see that

$$(2.4) \quad \text{Cap}(X) = \text{Cap}(\hat{X}) = \sum_{v=1}^k P^{\hat{x}_v}(\tau_{\hat{X}} = \infty) = \sum_{\ell=1}^j q_{v(\ell)}$$

and moreover, summing (2.2) over  $i \leq j$ , we get that

$$(2.5) \quad j = \sum_{i=1}^j \sum_{v=1}^k G(x_i, \hat{x}_v) P^{\hat{x}_v}(\tau_{\hat{X}} = \infty) = \sum_{\ell=1}^j q_{v(\ell)} \sum_{i=1}^j G(x_i, x_\ell).$$

The bounds of (2.3) are an immediate consequence of (2.4) and (2.5).  $\blacksquare$

**Lemma 2.4.** For  $Z_1 = \{x_1, \dots, x_{j_1}\}$ ,  $Z_2 = \{x_{j_1+1}, \dots, x_{j_1+j_2}\}$  with  $\{x_i\}$  not necessarily disjoint,

$$\text{Cap}(Z_1 \cup Z_2) \leq \text{Cap}(Z_2) + \frac{j_1 + j_2}{\min_{x \in Z_1 \setminus Z_2} \left\{ \sum_{i=1}^{j_1+j_2} G(x_i, x) \right\}}.$$

*Proof.* Since  $\tau_{Z_1 \cup Z_2} \leq \tau_{Z_2}$ , it follows that

$$(2.6) \quad \text{Cap}(Z_1 \cup Z_2) \leq \text{Cap}(Z_2) + \sum_{x \in Z_1 \setminus Z_2} P^x(\tau_{Z_1 \cup Z_2} = \infty).$$

For  $\hat{X}$  enumerating the *disjoint* points in  $Z_1 \cup Z_2$ ,  $v(\ell)$ ,  $q_v$  as in Lemma 2.3, we have that

$$\begin{aligned} \sum_{x \in Z_1 \setminus Z_2} P^x(\tau_{Z_1 \cup Z_2} = \infty) &= \sum_{\ell=1}^{j_1+j_2} q_{v(\ell)} \mathbf{1}_{\{\hat{x}_{v(\ell)} \in Z_1 \setminus Z_2\}}, \\ j_1 + j_2 &= \sum_{\ell=1}^{j_1+j_2} q_{v(\ell)} \sum_{i=1}^{j_1+j_2} G(x_i, x_\ell). \end{aligned}$$

Combining these identities with (2.6) yields the stated upper bound.  $\blacksquare$

**Remark 2.5.** In particular, applying Lemma 2.4 for

$$Z_1 = \bigcup_{i \in (j, J-j]} \hat{Z}_i, \quad Z_2 = \bigcup_{i \in [1, j] \cup (J-j, J]} \hat{Z}_i,$$

we have that for any  $\hat{Z}_i \subset \mathbb{Z}^d$ ,  $2j < J$ ,

$$\text{Cap}(Z_1 \cup Z_2) \leq \text{Cap}(Z_2) + \frac{\sum_{i=1}^J |\hat{Z}_i|}{\min_{x \in Z_1} \sum_{i=1}^J \sum_{y \in \hat{Z}_i} G(x, y)}.$$

*Proof of Lemma 2.1.* By the monotonicity of  $A \mapsto \text{Cap}(A)$ , it suffices to provide a uniform in  $\ell \leq r_m^b$  lower bound on  $\text{Cap}(C_m(\ell, r_m))$  and a matching upper bound on  $\text{Cap}(C_m^*(r_m))$ . With  $|C_m(\ell, r_m)| = (1 + o(1))\pi m r_m^2 \ell^{-3}$ , we get such a lower bound from Lemma 2.3, upon showing that for SRW on  $\mathbb{Z}^3$ ,

$$(2.7) \quad \sum_{y \in C_m(\ell, r_m)} G(x, y) \leq 3(1 + o(1))r_m^2 \ell^{-3} \log(m/r_m), \quad \forall \ell \leq r_m^b, \quad \forall x \in C_m(\ell, r_m).$$

With  $b < 2/3$ , the right-side of (2.7) diverges in  $m$  for any  $\ell \leq r_m^b$ . Further, with  $m/r_m \uparrow \infty$  and  $G(x, y)$  bounded, it suffices to sum in (2.7) over  $|x - y| \geq r_m^{2/3} \gg \ell$  and use the asymptotics

$$(2.8) \quad G(x, y) = \frac{3 + o(1)}{2\pi} |x - y|^{-1}$$

(for example, see [23, Thm. 1.5.4]). Setting  $u_m = m$ , we have for any  $v_m \uparrow \infty$  and  $r \in [r_m^{2/3}, v_m r_m]$ , at most  $C r^2 \ell^{-3}$  points  $y \in C_m(\ell, r_m)$  with  $|x - y| \in [r, r + 1]$ , while for each  $r \in [v_m r_m, u_m]$  there are at most  $(2\pi + o(1)) r_m^2 \ell^{-3}$  such points in  $C_m(\ell, r_m)$ . Thus, taking  $v_m^2 \ll \log(m/r_m)$  yields

$$(2.9) \quad \begin{aligned} \sum_{y \in C_m} G(x, y) &\leq \frac{3 + o(1)}{2\pi \ell^3} \left[ C \int_{r_m^{2/3}}^{v_m r_m} r dr + 2\pi r_m^2 \int_{v_m r_m}^{u_m} r^{-1} dr \right] \\ &= (3 + o(1)) r_m^2 \ell^{-3} \log(u_m / (r_m v_m)), \end{aligned}$$

from which (2.7) immediately follows. Turning to upper bound on  $\text{Cap}(C_m^*(r_m))$ , take now  $u_m := (m/r_m)^{1-\epsilon_m}$  and  $v_m := (m/r_m)^{\epsilon_m} \uparrow \infty$  for some  $\epsilon_m \rightarrow 0$ , splitting  $C_m^*(r_m)$  to  $Q_1 \cup Q_2$ , where

$$Q_1 := \bigcup_{i \in (u_m, (m/r_m) - u_m)} \mathbb{B}(z_i, r_m), \quad Q_2 := \bigcup_{i \notin (u_m, (m/r_m) - u_m)} \mathbb{B}(z_i, r_m).$$

Note that  $C_m^*(r_m)$  has at most  $(4\pi/3 + o(1)) r_m^2 m$ , possibly overlapping, points. Thus, combining Lemma 2.4 with the upper bound of Lemma 2.3, we get the upper bound of (2.1), once we show that for some  $\delta_m \rightarrow 0$ ,

$$(2.10) \quad \sum_{y \in C_m^*(r_m)} G(x, y) \geq (4 + \delta_m) r_m^2 \log(m/r_m), \quad \forall x \in Q_1,$$

$$(2.11) \quad \sum_{y \in Q_2} G(x, y) \geq \frac{|Q_2|}{m \delta_m} \log(m/r_m), \quad \forall x \in Q_2.$$

Fixing  $x \in \mathbb{B}(z_i, r_m) \subset Q_1$ , consider only the contribution to the LHS of (2.10) from all points  $y \in \mathbb{B}(z_j, r_m)$  with  $|j - i| \in [v_m, u_m]$ . For such a pair  $|y - x| \leq (|j - i| + 3)r_m$ , hence by (2.8),

$$G(x, y) \geq \frac{3 + o(1)}{2\pi} r_m^{-1} |j - i|^{-1},$$

resulting with

$$\sum_{y \in C_m^*(r_m)} G(x, y) \geq 2 \frac{3 + o(1)}{2\pi} \frac{|\mathbb{B}(0, r_m)|}{r_m} \sum_{j=v_m}^{u_m} j^{-1} = (4 + o(1)) r_m^2 \log(u_m/v_m),$$

which for our choices of  $u_m$  and  $v_m$  is as stated in (2.10) (for some  $\delta_m \rightarrow 0$ , uniformly over  $x \in Q_1$ ). Further,  $Q_2$  consists of two sets with an equal number of elements, each of diameter at most  $(1 + o(1))u_m r_m$ . Thus, we get by (2.8) that for some  $c > 0$ ,

$$\sum_{y \in Q_2} G(x, y) \geq c |Q_2| (u_m r_m)^{-1}, \quad \forall x \in Q_2,$$

and (2.11) follows upon choosing  $\epsilon_m \rightarrow 0$  slow enough so that  $(m/r_m)^{\epsilon_m} \gg \log(m/r_m)$ .  $\blacksquare$



3. LIL FOR SRW ON  $\mathbb{Z}^3$ : PROOF OF THEOREM 1.1

To ease the presentation we omit hereafter the integer-part symbol  $\lceil \cdot \rceil$  and divide the section to four parts, establishing the lower and then upper bounds, first for the lim sup-LIL of (1.1) and then for the lim inf-LIL of (1.1).

## 3.1. The lower bound in the limsup-LIL. Recall

$$\psi(t) := \sqrt{(2/3)t \log_2 t}, \quad h_3(t) := \frac{\pi}{3} \psi(t) (\log_3 t)^{-1},$$

of Proposition 1.7 and Theorem 1.1, respectively, and for the srw  $(S_m)$  on  $\mathbb{Z}^3$  consider the corresponding events

$$A_t := \{S_t^1 \geq \psi(t)\}, \quad A_t^* := A_t \cap \left\{ \sum_{\ell=0}^t G(0, S_\ell) \geq \frac{(1+4\delta)}{2} \frac{t}{h_3(t)} \right\}.$$

Our next lemma whose proof is deferred is key to the lower bound in our lim sup-LIL.

**Lemma 3.1.** *For srw  $(S_m)$  on  $\mathbb{Z}^3$  and any  $\delta > 0$ , there exists  $\zeta_t \rightarrow 0$  when  $t \rightarrow \infty$ , such that*

$$(3.1) \quad P(A_t^*) \leq 2\zeta_t P(A_t).$$

Given  $\epsilon > 0$ , we choose  $q > 1$  large and  $\delta > 0$  small so for  $t_n := q^n - q^{n-1}$  and all large  $n$ ,

$$(3.2) \quad \frac{1-\delta}{1+4\delta} h_3(t_n) \geq (1-\epsilon) h_3(q_n).$$

We then partition  $\mathbb{Z}_+$  to disjoint intervals  $I_n := (q^{n-1}, q^n] \cap \mathbb{Z}$  of length  $t$  and set the events

$$\hat{A}_n := \{S_{q^n}^1 - S_{q^{n-1}}^1 \geq \psi(t_n)\}, \quad \hat{H}_n(i) := \left\{ \hat{V}_n(i) \geq (1+4\delta) \frac{t_n}{h_3(t_n)} \right\},$$

where

$$\hat{V}_n(i) := \sum_{\ell \in I_n} G(S_i, S_\ell).$$

Setting

$$H_t(i) := \left\{ \sum_{\ell=1}^t G(S_i, S_\ell) \geq (1+4\delta) \frac{t}{h_3(t)} \right\}, \quad i \in [1, t],$$

note that by the stationarity of the srw increments,  $(\hat{A}_n, \hat{H}_n(q^{n-1} + i)) \stackrel{d}{=} (A_{t_n}, H_{t_n}(i))$ . Further, the event  $A_t$  is invariant to any permutation of the srw increments  $\{X_j, j \leq t\}$ , whereas given  $A_t$  the event  $H_t(i)$  depends only on  $\{S_\ell - S_i, \ell \in [1, t]\}$ . Further, the permuted increments  $\hat{X}_j = X_{(i+j) \bmod t}$  result with  $\hat{S}_\ell = S_{\ell+i} - S_i$  for all  $\ell \in [1, t-i]$ , whereas  $\hat{X}_j = X_{(i+1-j) \bmod t}$  result with  $\hat{S}_\ell = S_i - S_{i-\ell}$  for all  $\ell \in [1, i]$ . Since  $G(x, y) = G(-x, -y) \geq 0$ , it thus follows that conditional on  $A_t$  the random sum in each of the events  $H_t(i)$  is stochastically dominated by twice the random sum in the event  $A_t^*$ . Consequently, (3.1) yields that

$$(3.3) \quad \max_{i \in I_n} \{P(\hat{H}_n(i) | \hat{A}_n)\} \leq 2\zeta_{t_n} \rightarrow 0.$$



Next, consider the independent events  $G_n := \{|\Lambda_n| \geq (1 - \delta)t_n\}$ , where  $\Lambda_n$  is the subset of all those  $i \in I_n$  for which  $\hat{H}_n(i)$  does not hold. From Markov's inequality and (3.3) it follows that

$$(3.4) \quad P(G_n^c | \hat{A}_n) = P(|I_n| - |\Lambda_n| \geq \delta t_n | \hat{A}_n) \leq \frac{1}{\delta t_n} \sum_{i \in I_n} P(\hat{H}_n(i) | \hat{A}_n) \leq \frac{2\zeta_n}{\delta} \rightarrow 0.$$

Setting  $x(t) := \sqrt{2 \log_2 t}$ , note that  $(S_m^1)$  is the partial sum of  $\{-1, 0, 1\}$ -valued, zero-mean, i.i.d. variables of variance  $1/3$ . With

$$\bar{\Phi}(x) := \int_x^\infty \frac{e^{-u^2/2}}{\sqrt{2\pi}} du,$$

we have by the asymptotic normality of the moderate deviations for such partial sums (see [30, Thm. VIII.2.1]), that

$$(3.5) \quad \begin{aligned} P(\hat{A}_n) &= P(A_{t_n}) = P\left((t_n/3)^{-1/2} S_{t_n}^1 \geq x(t_n)\right) \\ &= (1 + o(1))\bar{\Phi}(x(t_n)) \geq c_1 x(t_n)^{-1} e^{-x(t_n)^2/2} \geq \frac{c_2}{n \sqrt{\log n}}, \end{aligned}$$

for some positive  $c_1$  and  $c_2 = c_2(q)$ . Hence, by (3.4), for all  $n$  large enough,

$$P(G_n) \geq P(\hat{A}_n \cap G_n) \geq \frac{1}{2} P(\hat{A}_n) \geq \frac{c_2}{2n \sqrt{\log n}}.$$

Having  $\{G_n\}$  independent with  $\sum_n P(G_n) = \infty$ , we deduce by the second Borel-Cantelli lemma that a.s. the events  $G_n$  hold for infinitely many values of  $n$ . Since

$$\hat{\mathcal{R}}_{q^n} := \{S_i\}_{i \in \Lambda_n} \subseteq \mathcal{R}_{q^n},$$

we have by the monotonicity of  $A \mapsto \text{Cap}(A)$ , the non-negativity of  $G(x, y)$ , Lemma 2.3 and the definition of  $\Lambda_n \subseteq I_n$ , that

$$R_{q^n} \geq \text{Cap}(\hat{\mathcal{R}}_{q^n}) \geq \frac{|\Lambda_n|}{\max_{i \in \Lambda_n} \{\hat{V}_n(i)\}} \geq \frac{|\Lambda_n| h_3(t_n)}{(1 + 4\delta)t_n}.$$

Consequently, in view of (3.2) we have on the event  $G_n$  that

$$R_{q^n} \geq \frac{1 - \delta}{1 + 4\delta} h_3(t_n) \geq (1 - \epsilon) h_3(q_n),$$

which since  $G_n$  holds infinity often, establishes the lower bound in the  $\limsup$ -LIL (along the sub-sequence  $q^n$ , and with  $\mathcal{R}_{q^n} \supseteq \hat{\mathcal{R}}_{q^n}$  of roughly the shape of  $C_{\psi(t_n)}$  of Lemma 2.1).

To prove Lemma 3.1, setting

$$V_I := 1_{A_t} \sum_{\ell \in I} G(0, S_\ell), \quad I \subseteq [0, t] \cap \mathbb{Z},$$

note that Lemma 3.1 is an immediate consequence of the following.

**Lemma 3.2.** *Fixing  $\delta \in (0, 1)$ , for  $\gamma_t := t(\log_2 t)^{-1}(\log_3 t)^{3/2}$  and some  $\zeta_t \rightarrow 0$ ,*

$$(3.6) \quad P\left(V_{[0, \gamma_t]} \geq \frac{\delta t}{h_3(t)}\right) \leq \zeta_t P(A_t),$$

$$(3.7) \quad P\left(V_{(\gamma_t, t]} \geq \frac{(1 + 2\delta)t}{2h_3(t)}\right) \leq \zeta_t P(A_t).$$

Our next lemma provides sharp tail estimates for the path of the one-dimensional walk  $(S_m^1)$ , which are key for proving (3.6) and (3.7), respectively.

**Lemma 3.3.** *Fixing  $\delta \in (0, 1)$ , for some  $C < \infty$ ,  $\zeta_t \rightarrow 0$  and all  $t$  large enough,*

$$(3.8) \quad \sup_{\ell \leq \gamma_t} \{P(S_{t-\ell}^1 \geq \psi(t) - \sqrt{\ell})\} \leq CP(A_t),$$

$$(3.9) \quad P(A_t \cap L_t^c) \leq 4\zeta_t P(A_t), \quad L_t := \bigcap_{\ell \in (\gamma_t, t]} \left\{ S_\ell^1 \geq \frac{\psi(t)\ell}{(1+\delta)t} \right\}.$$

*Proof of Lemma 3.3.* Let  $\eta_t := \gamma_t/t = (\log_3 t)^{3/2}/(\log_2 t)$ , setting  $x(t, r) := (x(t) - \sqrt{3r})/\sqrt{1-r}$  for  $r \leq \eta_t$  and  $x(t) = x(t, 0) := \sqrt{2\log_2 t}$ . Then, in view of the uniform Gaussian approximation of (3.5), we get (3.8) once we show that uniformly in  $r \leq \eta_t$ , the standard Gauss measure of  $[x(t, r), \infty)$ , is at most  $C$  times the Gauss measure of  $[x(t), \infty)$ . Note that  $\eta_t \rightarrow 0$  and  $x(t) \rightarrow \infty$  as  $t \rightarrow \infty$ , hence  $x(t, r)/x(t) \rightarrow 1$  uniformly in  $r \leq \eta_t$ . It thus remains only to show that for some  $C < \infty$  and all  $t$  large enough,

$$(3.10) \quad \inf_{r \leq \eta_t} \{x(t, r)^2 - x(t)^2\} \geq -2 \log C.$$

Next, setting  $u = x(t)\sqrt{r}$ , our expression for  $x(t, r)$  is such

$$(1-r)[x(t, r)^2 - x(t)^2] = u^2 - 2\sqrt{3}u + 3r \geq -3,$$

yielding (3.10) and thereby also (3.8).

Next, setting  $s_j := jt/(\log_2 t)$ , we partition  $(\gamma_t, t]$  into the disjoint intervals  $J_j = (s_j, s_{j+1}]$ ,  $j \in [(\log_3 t)^{3/2}, \log_2 t)$ . We likewise partition the events  $A_t \cap L_t^c$  according to whether the stopping time  $\tau_t := \inf\{\ell \geq \gamma_t : S_\ell^1 < \frac{\psi(t)\ell}{(1+\delta)t}\}$  equals  $\gamma_t$  or alternatively which interval  $J_j$  contains  $\tau_t$ . Note that if  $\tau_t > s_j$  then  $S_{s_j}^1 \geq \frac{\psi(t)s_j}{(1+\delta)t}$  and conditioning on the srw filtration at  $\tau_t \in J_j$ , we get by the strong Markov property (and i.i.d. increments), of the srw, that

$$P(A_t \cap \{\tau_t \in J_j\}) \leq q_t(s_j, 0) \sup_{s \in J_j} p_t(t-s, 0),$$

where

$$q_t(s_j, y) := P\left((S_{s_j}^1)_+ \geq \frac{\psi(t)s_j}{(1+\delta)t} - y\right), \quad p_t(r, y) := P\left(S_r^1 \geq \frac{\psi(t)(\delta t + r)}{(1+\delta)t} + y\right)$$

and  $(S_{s_j}^1)_+ := S_{s_j}^1 \vee 0$ . The only other way for the event  $L_t^c$  to occur, is by having

$$S_{\gamma_t}^1 < \frac{\psi(t)\eta_t}{1+\delta} := \Delta_t \log_2 t.$$

Partitioning to  $\{S_{\gamma_t}^1 \in I_i\}$ , for  $I_i := [(\log_2 t - i)\Delta_t - \Delta_t, (\log_2 t - i)\Delta_t)$  when  $0 \leq i < \log_2 t$ , and  $I_{\log_2 t} := (-\infty, 0)$ , yields that

$$P(A_t \cap \{S_{\gamma_t}^1 \in I_i\}) \leq q_t(\gamma_t, i\Delta_t + \Delta_t) p_t(t - \gamma_t, i\Delta_t)$$

and consequently,

$$\begin{aligned} P(A_t \cap L_t^c) &\leq \sum_i P(A_t \cap \{S_{\gamma_i}^1 \in I_i\}) + \sum_j P(A_t \cap \{\tau_t \in J_j\}) \\ &\leq (1 + \log_2 t) \left[ \sup_{y \in [0, \Delta_t \log_2 t]} \{q_t(\gamma_t, y + \Delta_t) p_t(t - \gamma_t, y)\} + \sup_{s \in J_j, j \geq (\log_3 t)^{3/2}} q_t(s_j, 0) p_t(t - s, 0) \right]. \end{aligned}$$

As  $P(A_t) = p_t(t, 0)$ , we thus get (3.9), if for some  $\zeta_t \rightarrow 0$

$$(3.11) \quad q_t(s_j, 0) p_t(t - s, 0) \leq \frac{\zeta_t}{\log_2 t} p_t(t, 0), \quad \forall j \geq (\log_3 t)^{3/2}, s \in J_j,$$

$$(3.12) \quad q_t(\gamma_t, y + \Delta_t) p_t(t - \gamma_t, y) \leq \frac{2\zeta_t}{\log_2 t} p_t(t, 0), \quad \forall y \leq \Delta_t \log_2 t.$$

Starting with (3.11), note that  $s_j \geq s_1 \gg 1$  and  $p_t(r, y) > 0$  only if  $r \geq \delta\psi(t)/2$ . Thus, a uniform normal approximation applies for both  $q_t(s_j, 0)$  and  $p_t(t - s, 0)$ , and with all arguments of  $\bar{\Phi}(\cdot)$  being only  $O(x(t))$  for  $x(t) = \sqrt{2 \log_2 t}$ , such approximations hold, as in (3.5), with  $o(1)$  relative error. Further, note that  $u = s/t \in [\eta_t, 1)$  with  $s_j/t \geq u - \varepsilon_t$  where  $\varepsilon_t := (\log_2 t)^{-1}$ . Hence, by the preceding, we get (3.11) upon showing that

$$(3.13) \quad \sup_{u \in [\eta_t, 1)} \left\{ \bar{\Phi}\left(\frac{x(t)}{1 + \delta} \sqrt{u - \varepsilon_t}\right) \bar{\Phi}\left(\frac{x(t)(1 + \delta - u)}{(1 + \delta) \sqrt{1 - u}}\right) \right\} \leq \frac{\zeta_t}{\log_2 t} \bar{\Phi}(x(t)).$$

Upon taking the logarithm of both sides of (3.13), recalling that  $|\log \bar{\Phi}(y) + \log y + y^2/2|$  is bounded at  $y \rightarrow \infty$ , that  $x(t)^2 \varepsilon_t$  is bounded and  $\eta_t \geq 2\varepsilon_t$ , we find (after some algebra), that it suffices for (3.13) to have for some  $\tilde{\zeta}_t \rightarrow 0$  as  $t \rightarrow \infty$ ,

$$(3.14) \quad \sup_{u \in [\eta_t, 1)} \left\{ -(\log_2 t) \theta(u) + \frac{1}{2} \log(u^{-1} - 1) \right\} \leq \log \tilde{\zeta}_t - \frac{1}{2} \log_3 t,$$

where

$$\theta(u) := \frac{u}{(1 + \delta)^2} + \frac{(1 + \delta - u)^2}{(1 + \delta)^2(1 - u)} - 1 = \frac{u}{1 - u} \frac{\delta^2}{(1 + \delta)^2}.$$

Since  $u \mapsto \theta(u)$  is non-decreasing, the supremum on the left-side of (3.14) is attained at  $u = \eta_t$ , where for large  $t$  it is at most  $-\frac{\delta^2}{4} (\log_3 t)^{3/2}$ . This is more than enough for (3.14) to hold, thereby establishing (3.11). We next turn to (3.12), which analogously to (3.13) is a consequence of

$$(3.15) \quad \sup_{v \in [0, 1]} \left\{ \bar{\Phi}\left(\frac{x(t)}{1 + \delta} \sqrt{\eta_t}(v - \varepsilon_t)\right) \bar{\Phi}\left(\frac{x(t)(1 + \delta - v\eta_t)}{(1 + \delta) \sqrt{1 - \eta_t}}\right) \right\} \leq \frac{\zeta_t}{\log_2 t} \bar{\Phi}(x(t)).$$

Restricting to  $v \leq 1/2$ , we bound the left-most term by one and take the logarithm of both sides to find that (3.15) holds because for  $v \leq 1/2$ ,

$$\frac{(1 + \delta - v\eta_t)^2}{(1 + \delta)^2(1 - \eta_t)} - 1 \geq \frac{\eta_t \delta}{1 + \delta}$$

and  $x(t)^2 \eta_t = 2(\log_3 t)^{3/2} \gg \log_3 t$ . To complete the proof of (3.15), it thus suffices (similarly to (3.14)), to have for some  $\tilde{\zeta}_t \rightarrow 0$ , that

$$(3.16) \quad \sup_{v \in [1/2, 1]} \left\{ -(\log_2 t) \theta_t(v) \right\} \leq \log \tilde{\zeta}_t - \log_3 t,$$

where it is easy to check that

$$\theta_t(v) := \frac{\eta_t v^2}{(1+\delta)^2} + \frac{(1+\delta - v\eta_t)^2}{(1+\delta)^2(1-\eta_t)} - 1 \geq \theta_t(1) \geq \frac{\eta_t \delta^2}{(1+\delta)^2}.$$

The preceding suffices for (3.16) and thereby for (3.12), thus completing the proof.  $\blacksquare$

*Proof of Lemma 3.2.* Starting with (3.6), note that for some  $C_1 < \infty$  and any  $\ell \geq 0$ ,

$$E[G(0, S_\ell)] = \sum_{i=\ell}^{\infty} P^0(S_i = 0) \leq C_1(1 + \sqrt{\ell})^{-1}.$$

Further, recall from (2.8) that  $G(0, y) \leq C_2/(1 + |y|)$  for some  $C_2 < \infty$  and all  $y \in \mathbb{Z}^3$ . Hence,

$$(3.17) \quad E[G(0, S_\ell)1_{A_t}] \leq C_2(1 + \sqrt{\ell})^{-1}P(A_t) + \sum_{|y| \leq \sqrt{\ell}} G(0, y)P(\{S_\ell = y\} \cap A_t).$$

With  $(S_\ell, S_t - S_\ell) \stackrel{d}{=} (S_\ell, \hat{S}_{t-\ell})$  for srw  $(\hat{S}_m)$  which is independent of  $(S_m)$ , if  $y \in \mathbb{Z}^3$  is such that  $|y^1| \leq |y| \leq \sqrt{\ell}$ , then

$$P(\{S_\ell = y\} \cap A_t) \leq P(S_\ell = y)P(\hat{S}_{t-\ell}^1 \geq \psi(t) - \sqrt{\ell}).$$

Therefore, thanks to (3.8), for any  $\ell \leq \gamma_t$  the right-most term in (3.17) is at most,

$$E[G(0, S_\ell)]P(\hat{S}_{t-\ell}^1 \geq \psi(t) - \sqrt{\ell}) \leq C_1(1 + \sqrt{\ell})^{-1}CP(A_t).$$

We thus deduce from (3.17), that for some  $C_3, C_4 < \infty$ ,

$$E[V_{[0, \gamma_t]}] = \sum_{\ell=0}^{\gamma_t} E[G(0, S_\ell)1_{A_t}] \leq C_3P(A_t) \sum_{\ell=0}^{\gamma_t} (1 + \sqrt{\ell})^{-1} \leq C_4P(A_t) \sqrt{\gamma_t}.$$

Consequently, by Markov's inequality and our choice of  $\gamma_t$ ,

$$P\left(V_{[0, \gamma_t]} \geq \frac{\delta t}{h_3(t)}\right) \leq \zeta_t P(A_t),$$

where our choice of  $\gamma_t$  results with  $\zeta_t := C_4 h_3(t) \sqrt{\gamma_t}/(\delta t) \rightarrow 0$  as  $t \rightarrow \infty$ .

Turning to (3.7), recall our choice of  $\gamma_t$  implying that

$$\sum_{\ell \in (\gamma_t, t]} \frac{1}{\ell} \leq \log(t/\gamma_t) \leq \log_3 t.$$

Further, with  $h_3(t) = (\pi/3)\psi(t)(\log_3 t)^{-1}$ , it follows that  $t/(2h_3(t)) = \frac{3}{2\pi}(t/\psi(t))(\log_3 t)$ . Consequently, for all  $t$  large enough, the event  $L_t$  of (3.9) implies, thanks to (2.8), that

$$V_{(\gamma_t, t]} = \sum_{\ell \in (\gamma_t, t]} G(0, S_\ell) \leq \frac{3 + o(1)}{2\pi} \sum_{\ell \in (\gamma_t, t]} \|S_\ell^1| + 1|^{-1} \leq \frac{3 + o(1)}{2\pi} \frac{(1+\delta)t}{\psi(t)} (\log_3 t) \leq \frac{(1+2\delta)t}{2h_3(t)}.$$

Since the same conclusion applies when  $A_t^c$  holds (in which case  $V_{(\gamma_t, t]} = 0$ ), we see that (3.7) is an immediate consequence of (3.9).  $\blacksquare$

**3.2. The upper bound in the limsup-LIL.** Recall that  $R_n$  is non-decreasing. Further, for any  $t_n = q^n$ ,  $q > 1$  we have that eventually  $h_3(t_n)/h_3(t_{n-1}) \leq q$ . It thus suffices to prove the upper bound in our lim sup-LIL only along each such sequence  $t_n$  (thereafter taking  $q \downarrow 1$  to complete the proof). To this end, fix  $q > 1$  and  $\delta, \eta > 0$ , setting  $m := (1 + \delta)^3 \psi(t_n)$  and  $r_m = m/b_m \uparrow \infty$  for  $b_m := 2\eta(1 + \delta)^6 (\log_2 t_n)$  (so  $r_m := \psi(t_n)/(2\eta(1 + \delta)^3 \log_2 t_n)$ ). We aim to cover  $\mathcal{R}_{t_n}$  by the union  $C_m^*(r_m)$  of  $b_m$  balls of radius  $r_m$  each, with the centers of consecutive balls at most  $r_m$  apart. Indeed, as shown in Section 2 (see Remark 2.2), this would yield  $R_{t_n} \leq (1 + \delta + o(1))^3 h_3(t_n)$ , so we then conclude by taking  $\delta \downarrow 0$  and  $q \downarrow 1$ .

Specifically, starting at  $T_0 = 0$ , set the increasing stopping times

$$T_i := \inf\{k > T_{i-1} : |S_k - S_{T_{i-1}}| > r_m - 1\}, \quad \forall i \geq 1,$$

noting that the event  $\{T_{b_m} \geq t_n\}$  implies the aforementioned containment  $\mathcal{R}_{t_n} \subseteq C_m^*(r_m)$ . Further, with  $\exp(-(1 + \delta) \log_2 t_n) \leq Cn^{-(1+\delta)}$  summable, upon employing the first Borel-Cantelli lemma, it remains only to establish the following key lemma.

**Lemma 3.4.** *For any  $q > 1$  and small  $\delta > 0$ , there exist  $\eta > 0$  and  $C < \infty$  such that*

$$(3.18) \quad P(T_{b_m} < t_n) \leq C \exp(-(1 + \delta) \log_2 t_n) \quad \forall n.$$

*Proof.* By the strong Markov property and the independence of increments of the walk, we see that  $T_{b_m}$  is the sum of  $b_m$  i.i.d. copies of the first exit time  $T_1$  of the (discrete) ball  $\mathbb{B}(0, r_m - 1)$ , by the 3D-srw. Since  $r_m \gg 1$ , we may wlog replace  $T_1$  by the first exit time of 3D Brownian motion from  $\mathbb{B}(0, \sqrt{3}r_m)$ , which can thereby, due to Brownian scaling, be further replaced by  $3r_m^2 \bar{T}_1$ , with  $\bar{T}_1 := \inf\{t \geq 0 : |B_t| \geq 1\}$  the Brownian hitting time of the unit sphere  $\mathbb{S}^2$ . Note that  $t_n = 3r_m^2 \eta b_m$  and as  $b_m/(2\eta) = 3m^2/(2t_n) = (1 + \delta)^6 (\log_2 t_n)$ , it suffices to show that for some  $\eta = \eta(\delta) > 0$  and all  $m$ ,

$$(3.19) \quad P\left(\frac{1}{b_m} \sum_{i=1}^{b_m} \bar{T}_i < \eta\right) \leq e^{-(1-\delta)^3 b_m/(2\eta)},$$

where  $\bar{T}_i$  are i.i.d. copies of  $\bar{T}_1$ . To this end, covering  $\mathbb{S}^2$  by  $c_\delta$  balls of radius  $\delta$  each, centered at some  $\theta_i \in \mathbb{S}^2$ , we have by the triangle inequality that

$$\max_i \{\langle \theta_i, B_{\bar{T}_1} \rangle\} \geq 1 - \delta.$$

Hence, fixing  $\lambda > 0$ , we get upon applying Doob's optional stopping theorem for the martingale  $M_t = \sum_i \exp(\lambda \langle \theta_i, B_t \rangle - \lambda^2 t/2)$  at the stopping time  $\bar{T}_1$ , that

$$c_\delta = M_0 = E[M_{\bar{T}_1}] \geq e^{\lambda(1-\delta)} E[e^{-\lambda^2 \bar{T}_1/2}].$$

Consequently, by Markov's inequality, we have for any  $\eta, \lambda, \delta > 0$  and integer  $b \geq 1$ , that

$$(3.20) \quad P\left(\frac{1}{b} \sum_{i=1}^b \bar{T}_i < \eta\right) \leq e^{\lambda^2 b \eta/2} E[e^{-\lambda^2 \bar{T}_1/2}]^b \leq (c_\delta e^{\lambda^2 \eta/2 - \lambda(1-\delta)})^b.$$

Taking the optimal  $\lambda = (1 - \delta)/\eta$  it is easy to check that for  $\eta \leq \eta(\delta) = \delta(1 - \delta)^2/(2 \log c_\delta)$ , the LHS of (3.20) is at most  $\exp(-(1 - \delta)^3 b/(2\eta))$ . We thus got (3.19) for any  $\delta > 0$  provided  $\eta \leq \eta(\delta)$ , thereby completing the proof.  $\blacksquare$

**Remark 3.5.** One has for any  $\delta > 0$  small and all  $t$  large enough, the classical bound

$$P(\max_{1 \leq k \leq t} |S_k| \geq (1 + \delta)^3 \psi(t)) \leq Ce^{-(1+\delta) \log_2 t}.$$

We need in (3.18) a stronger result, since for any  $b_m$  and  $r_m \gg 1$ ,

$$\{\max_{1 \leq k \leq t_n} |S_k| \geq b_m r_m\} \subset \{T_{b_m} < t_n\},$$

and while  $b_m r_m = (1 + \delta)^3 \psi(t_n)$ , our crude use of  $\delta$ -cover of  $\mathbb{S}^2$  in proving Lemma 3.4 requires us to also have  $b_m / (\log_2 t_n) \rightarrow 0$  as  $\delta \downarrow 0$ .

**3.3. The upper bound in the liminf-LIL.** For any  $A \subset \mathbb{R}^3$  and  $r > 0$ , let

$$\text{Nbd}(A, r) := \bigcup_{x \in A} \mathbb{B}(x, r)$$

denote the  $r$ -blowup of  $A$ . Utilizing [13], we first relate  $R_n$  with a suitable Brownian capacity, as stated next.

**Lemma 3.6.** We can couple the srw with a 3D Brownian motion  $(B_t, t \geq 0)$ , such that

$$(3.21) \quad \lim_{n \rightarrow \infty} \frac{R_n}{\text{Cap}_B(B[0, n/3])} = \frac{1}{3} \quad \text{a.s.}$$

and for any  $\delta \in (0, 1/2)$ ,

$$(3.22) \quad \lim_{n \rightarrow \infty} \frac{R_n}{\text{Cap}_B(\text{Nbd}(B[0, n/3], n^{1/2-\delta}))} = \frac{1}{3} \quad \text{a.s.}$$

*Proof.* The results were essentially shown in [13]. Indeed, [13, (4.15)] shows that (3.21) holds when each ratio is restricted to the events  $E_n$ , while it is also shown that a.s.  $E_n$  holds for all sufficiently large  $n$  (combine [13, (4.2)] with Borel-Cantelli). Hence (3.21) also holds without such a restriction. Turning to show (3.22), let  $\tilde{P}$  denote the probability of an independent Brownian motion  $(\tilde{B}_t)$ . Fixing  $\delta \in (0, 1/2)$  and some  $y_n \in \mathbb{Z}^3$  such that  $|y_n| = n^{1/2+\delta}$ , we similarly obtain (after dispensing of events  $E_n$ ), that by the same argument as in [13, (4.4)], a.s. one has for all large  $n$ ,

$$\begin{aligned} & \text{Cap}_B(\text{Nbd}(B[0, n/3], n^{1/2-\delta})) \\ &= (2\pi + o(1))n^{1/2+\delta} \tilde{P}(\text{Nbd}(B[0, n/3], n^{1/2-\delta}) \cap (y_n + \tilde{B}[0, \infty)) \neq \emptyset | B[0, n/3]). \end{aligned}$$

By [13, (4.13) & (4.4)], a.s., the latter expression is for all  $n$  large  $(1 + o(1))\text{Cap}_B(B[0, n/3])$ . Thus,

$$\lim_{n \rightarrow \infty} \frac{\text{Cap}_B(B[0, n/3])}{\text{Cap}_B(\text{Nbd}(B[0, n/3], n^{1/2-\delta}))} = 1 \quad \text{a.s.},$$

which in view of (3.21) completes the proof of (3.22).  $\blacksquare$

Proceeding to show the upper bound in the lim inf-LIL, let  $r(n) := \pi \sqrt{n/(2 \log_2 n)}$ , that is  $r(n) = \sqrt{3} \hat{\psi}(n)$  for  $\hat{\psi}(\cdot)$  of Proposition 1.7. Recall that by [18, Lemma 1.1] or [22], and Brownian scaling, for some  $c > 0$  and all  $t, r > 0$ ,

$$(3.23) \quad P(\sup_{s \in [0, t]} |B_s| \leq r) \geq 2ce^{-\frac{\pi^2 t}{2r^2}}.$$

We have used in (3.23) also that the largest eigenvalue of the Dirichlet Laplacian in the unit ball in  $\mathbb{R}^d$  is  $-j^2$ , where  $j = j_{(d-2)/2,1}$  denotes the first positive zero of the Bessel function of the first kind with index  $(d-2)/2$ , and in particular that  $j_{1/2,1} = \pi$  (see [34, Page 490]). Considering (3.23) for  $r = r(s_n)$ ,  $s_n = n^n$  and the Brownian increments in the disjoint intervals  $I_n = [s_{n-1}, s_n)$  of length  $t_n = s_n - s_{n-1}$ , result with

$$(3.24) \quad P\left(\sup_{t \in I_n} \{|B_t - B_{s_{n-1}}|\} \leq r(s_n)\right) \geq c \exp(-\log_2 s_n) = \frac{c}{n \log n}.$$

Thus, thanks to the independence of Brownian increments on the disjoint  $I_n$ , we get from the second Borel-Cantelli lemma that

$$(3.25) \quad P\left(\liminf_{n \rightarrow \infty} (r(s_n))^{-1} \sup_{t \in I_n} \{|B_t - B_{s_{n-1}}|\} \leq 1\right) = 1.$$

Further, as  $\sqrt{s_{n-1}(\log_2 s_{n-1})} = o(r(s_n))$ , by Kinchin's LIL for the Brownian motion,

$$(3.26) \quad P\left(\limsup_{n \rightarrow \infty} (r(s_n))^{-1} \sup_{t \leq s_{n-1}} |B_t| = 0\right) = 1.$$

Combining (3.25) and (3.26), we deduce that

$$P\left(\liminf_{n \rightarrow \infty} (r(s_n))^{-1} \sup_{t < s_n} |B_t| \leq 1\right) = 1.$$

This of course implies that also

$$(3.27) \quad P\left(\liminf_{n \rightarrow \infty} (r(n))^{-1} \sup_{t < n} |B_t| \leq 1\right) = 1.$$

Recall that for any  $r > 0$  one has that  $r^{-1} \text{Cap}_B(\mathbb{B}(0, r)) = \text{Cap}_B(\mathbb{B}(0, 1)) = 2\pi$  ( $= \kappa_1$  on [12, Page 356]). By (3.27), for any  $\epsilon > 0$ , a.s.  $B[0, n] \subset \mathbb{B}(0, (1+\epsilon)r(n))$  for infinitely many values of  $n$ , in which case also  $\text{Cap}_B(B[0, n]) \leq 2\pi(1+\epsilon)r(n)$ . That is,

$$(3.28) \quad P\left(\text{Cap}_B(B[0, n]) \leq 2\pi(1+\epsilon)r(n) \quad \text{i.o.}\right) = 1.$$

By Brownian scaling, the sequence  $\{\sqrt{3} \text{Cap}_B(B[0, n/3])\}$  has the same law as the sequence  $\{\text{Cap}_B(B[0, n])\}$ . Thus, in view of (3.21), we can also construct a coupling so that for any  $\epsilon > 0$ , we have that a.s.

$$R_n \leq \frac{(1+\epsilon)}{3\sqrt{3}} \text{Cap}_B(B[0, n]),$$

for all  $n$  large enough. With  $r(n) = \frac{3\sqrt{3}}{2\pi} \hat{h}_3(n)$ , it thus follows from (3.28) that

$$P\left(R_n \leq (1+\epsilon)^2 \hat{h}_3(n) \quad \text{i.o.}\right) = 1,$$

and taking  $\epsilon \downarrow 0$  establishes the stated upper bound in our lim inf-LIL.



**3.4. The lower bound of the liminf-LIL.** Fixing  $a > 0$ , we have by [12, (1.4)], that for any  $f(t) \uparrow \infty$  such that  $f(t) = o(t^{2/3})$ ,

$$(3.29) \quad \lim_{t \rightarrow \infty} \frac{f(t)}{t} \log P\left(|\text{Nbd}(B[0, t], a)| \leq \left(\frac{\pi}{\sqrt{2}}\right)^3 f(t)^{3/2} \omega_3\right) = -1,$$

where  $\omega_3$  denotes the volume of the unit ball (using here that the largest eigenvalue of the Dirichlet Laplacian in the *unit volume ball* in  $\mathbb{R}^3$  is  $-\omega_3^{2/3} \pi^2$ ). Fixing  $\delta \in (0, 1/2)$  as in (3.22), Brownian scaling by time factor  $3n^{2\delta-1}$  yields equality in distribution between the sequences

$$|\text{Nbd}(B[0, n/3], n^{1/2-\delta})| \stackrel{d}{=} 3^{-3/2} n^{3/2-3\delta} |\text{Nbd}(B[0, n^{2\delta}], 3^{-1/2})|.$$

Thus, considering (3.29) for  $a = 3^{-1/2}$ ,  $t = n^{2\delta}$  and  $f(t) = (1 - \epsilon)^3 n^{2\delta} (\log_2 n)^{-1}$ , we arrive at

$$\begin{aligned} & P\left(|\text{Nbd}(B[0, n/3], n^{1/2-\delta})| \leq (1 - \epsilon)^2 \hat{\psi}(n)^3 \omega_3\right) \\ &= P\left(|\text{Nbd}(B[0, n^{2\delta}], 3^{-1/2})| \leq (1 - \epsilon)^2 \left(\frac{\pi}{\sqrt{2}}\right)^3 (n^{2\delta} (\log_2 n)^{-1})^{3/2} \omega_3\right) \\ &\leq C \exp(-(1 - \epsilon)^{-2} (\log_2 n)). \end{aligned}$$

Considering  $n_k = q^k$ , we get by the first Borel-Cantelli lemma, that for fixed  $q > 1$  and  $\epsilon > 0$ ,

$$(3.30) \quad \liminf_{k \rightarrow \infty} \frac{|\text{Nbd}(B[0, n_k/3], n_k^{1/2-\delta})|}{\omega_3 \hat{\psi}(n_k)^3} \geq (1 - \epsilon)^2 \quad \text{a.s.}$$

With  $n \mapsto |\text{Nbd}(B[0, n/3], n^{1/2-\delta})|$  monotone increasing and  $\hat{\psi}(q^k)/\hat{\psi}(q^{k-1}) \rightarrow 1$  as  $k \rightarrow \infty$  followed by  $q \downarrow 1$ , we deduce from (3.30) that

$$(3.31) \quad \liminf_{n \rightarrow \infty} \frac{|\text{Nbd}(B[0, n/3], n^{1/2-\delta})|}{\omega_3 \hat{\psi}(n)^3} \geq 1 \quad \text{a.s.}$$

Next, recall the Poincaré-Carleman-Szegö theorem [31], that for any  $r > 0$ ,

$$(3.32) \quad \inf_{|A|=\omega_3 r^3} \{\text{Cap}_B(A)\} = \text{Cap}_B(\mathbb{B}(0, r)) = r \text{Cap}_B(\mathbb{B}(0, 1)) = 2\pi r.$$

Recall that  $\hat{h}_3(n) = \frac{2\pi}{3} \hat{\psi}(n)$ . Hence, by (3.32) for  $A = |\text{Nbd}(B[0, n/3], n^{1/2-\delta})|$ , we have in view of (3.30) that

$$\liminf_{n \rightarrow \infty} \frac{\text{Cap}_B(|\text{Nbd}(B[0, n/3], n^{1/2-\delta})|)}{3\hat{h}_3(n)} \geq 1, \quad \text{a.s.},$$

which together with (3.22) yields the stated lower bound for the lim inf-LIL of  $R_n$  in  $\mathbb{Z}^3$ .

#### 4. LIL FOR SRW ON $\mathbb{Z}^4$ : PROOF OF THEOREM 1.3

Hereafter we consider for integers  $0 \leq a \leq b \leq c$ , the random variables

$$(4.1) \quad R_{a,b} := \text{Cap}(\mathcal{R}(a, b)), \quad V_{a,b,c} := R_{a,b} + R_{b,c} - R_{a,c} \geq 0.$$

Note that by shift invariance  $R_{a,b} \stackrel{d}{=} R_{0,b-a} = R_{b-a}$  and  $R_{a,b}$  is independent of  $R_{b,c}$  (due to the independence of increments). In particular, for any increasing  $\{n_k\}$  starting at  $n_0 = 0$ , one has

the decomposition

$$(4.2) \quad R_{n_k} := \sum_{j=1}^k U_j - \Delta_{n_k, k},$$

in terms of the independent variables  $U_j := R_{n_{j-1}, n_j}$  and the sum of non-negative variables

$$(4.3) \quad \Delta_{n_k, k} = \sum_{j=1}^{k-1} V_{n_{j-1}, n_j, n_k} = \sum_{j=1}^{k-1} V_{0, n_j, n_{j+1}}.$$

As we shall see, the fluctuation in  $\sum_j U_j$  is negligible for the LIL-s of Theorem 1.3 where  $E\Delta_{n_k, k} \approx h_4(n_k)$ , the lim sup-LIL being due to the exceptional times with  $\Delta_{n_k, k} = o(E\Delta_{n_k, k})$ , while the lim inf-LIL is due to the exceptional times where  $\Delta_{n_k, k} \approx \hat{h}_4(n_k) \gg E\Delta_{n_k, k}$ . In contrast, we show in Section 5 that  $\Delta_{n_k, k}$  has a negligible effect when  $d \geq 5$ , where the LIL follows the usual pattern for sums of independent variables (namely, that of the LIL for a Brownian motion).

**4.1. The lower bound in the limsup-LIL.** Recall Green's function for the 4-dimensional Brownian motion,

$$(4.4) \quad G_\beta(x, y) = \frac{1}{2\pi^2} |x - y|^{-2}.$$

We start with a key lemma that improves upon the upper bound of [6, Prop. 4.1].

**Lemma 4.1.** *Suppose  $(S_m)$  and  $(\tilde{S}_m)$  are two independent SRW on  $\mathbb{Z}^4$ , while  $(\beta_s, s \geq 0)$ ,  $(\tilde{\beta}_s, s \geq 0)$  are two independent, standard 4-dimensional Brownian motions. Let*

$$X_n := \frac{1}{n} \sum_{i, \ell \in [1, n]} G(S_i, \tilde{S}_\ell), \quad Y := \int_0^1 \int_0^1 G_\beta(\beta_s, \tilde{\beta}_t) ds dt.$$

Then, for some  $C < \infty$  and any  $p, n \in \mathbb{N}$ ,

$$(4.5) \quad E[X_n^p] \leq C^p p!, \quad E[Y^p] \leq C^p p!.$$

One immediate consequence of (4.5) is that for any  $c < 1/C$ ,

$$(4.6) \quad \sup_n E[e^{cX_n}] < \infty, \quad E[e^{cY}] < \infty.$$

*Proof of Proposition 1.6.* This is a direct consequence of Lemma 4.1. Indeed, recall from [6, Prop. 4.5], that the limit in  $L^2$ ,

$$\gamma_G([0, 1]^2) := 2 \sum_{n \geq 1} \sum_{k=1}^{2^{n-1}} (\alpha(A_k^n) - E\alpha(A_k^n))$$

exists, where for any  $n \geq 1$  and  $k \leq 2^{n-1}$ ,

$$(4.7) \quad \alpha(A_k^n) := \int_{A_k^n} G_\beta(\beta_s, \beta_t) ds dt,$$

$$A_k^n := [(2k-2)2^{-n}, (2k-1)2^{-n}) \times ((2k-1)2^{-n}, 2k2^{-n}].$$

Note that  $\{\alpha(A_k^n)\}_{k \geq 1}$  are i.i.d. for any fixed  $n \geq 1$ , with  $\alpha(A_k^n) \stackrel{d}{=} 2^{-n}Y$ . That is,  $\{\alpha(A_k^n)\}$  satisfies [26, properties (i) and (ii)]. Upon replacing the upper bound of [26, Lemma 2] by (4.5), it is not hard to check that the proof of [26, Thm. 1] yields here that the moment generating function  $M_G(\lambda)$  of (1.6) is finite for sufficiently small  $\lambda > 0$ . By Hölder's inequality, and in view of [6, Thm. 1.2], it follows that  $\lambda_o \in (0, \infty)$ , as claimed. ■

The following elementary bounds are needed for proving Lemma 4.1.

**Lemma 4.2.** *There exists  $C < \infty$  such that for any  $t > 0$  and  $x, y \in \mathbb{R}^4$ ,*

$$(4.8) \quad E[|\beta_t - x|^{-2}] \leq C \min\{t^{-1}, |x|^{-2}\} \leq Ct^{-1/2}|x|^{-1},$$

$$(4.9) \quad E[|\beta_t - x|^{-1}|\beta_t - y|^{-1}] \leq Ct^{-1/2}(|x| \vee |y|)^{-1} \leq 2Ct^{-1/2}|y - x|^{-1},$$

$$(4.10) \quad E[|\beta_t - x|^{-2}|\beta_t - y|^{-1}] \leq 2Ct^{-1/2}|x|^{-1}|y - x|^{-1}.$$

Similarly, for  $|\cdot|_+ = |\cdot| \vee 1$ , any  $i \geq 0$  and  $x, y \in \mathbb{Z}^4$ ,

$$(4.11) \quad E[|S_i - x|_+^{-2}] \leq C \min\{|i|_+^{-1}, |x|_+^{-2}\} \leq C|i|_+^{-1/2}|x|_+^{-1},$$

$$(4.12) \quad E[|S_i - x|_+^{-1}|S_i - y|_+^{-1}] \leq C|i|_+^{-1/2}(|x|_+ \vee |y|_+)^{-1} \leq 2C|i|_+^{-1/2}|y - x|_+^{-1},$$

$$(4.13) \quad E[|S_i - x|_+^{-2}|S_i - y|_+^{-1}] \leq 2C|i|_+^{-1/2}|x|_+^{-1}|y - x|_+^{-1}.$$

*Proof.* Denoting by  $\phi_s(x) := (2\pi s)^{-2} \exp(-\frac{|x|^2}{2s})$  the density at  $x$  of the Gaussian law of  $\beta_s$ , we get from (4.4) after change of variables that

$$E[|\beta_t - x|^{-2}] = 2\pi^2 E[G_\beta(\beta_t, x)] = 2\pi^2 \int_t^\infty \phi_s(x) ds = t^{-1} \varphi_1(|x|^2/t) = |x|^{-2} \varphi_2(t/|x|^2),$$

for the finite decreasing functions  $\varphi_1(r) := \frac{1}{2} \int_1^\infty u^{-2} e^{-r/(2u)} du$ ,  $\varphi_2(r) := \frac{1}{2} \int_r^\infty u^{-2} e^{-1/(2u)} du$ . Thus, (4.8) holds for any  $C \geq \varphi_1(0) \vee \varphi_2(0)$ . Next, by Cauchy-Schwarz and (4.8),

$$E(|\beta_t - x|^{-1}|\beta_t - y|^{-1}) \leq (E[|\beta_t - x|^{-2}])^{1/2} (E[|\beta_t - y|^{-2}])^{1/2} \leq Ct^{-1/2}|y|^{-1}.$$

Exchanging  $x$  with  $y$  yields the first inequality in (4.9), whereby the second inequality follows (as  $|x - y| \leq |x| + |y| \leq 2|x| \vee |y|$ ). Now, by the triangle inequality, for  $\beta_t \neq x \neq y$ ,

$$|y - x| |\beta_t - x|^{-2} |\beta_t - y|^{-1} \leq |\beta_t - x|^{-1} (|\beta_t - x|^{-1} + |\beta_t - y|^{-1}),$$

so taking the expectation and using (4.8) and (4.9) to bound the RHS, results with (4.10).

With  $S_0 = 0$ , clearly (4.11) holds at  $i = 0$ , while for  $i \geq 1$ , recall [23, Thm. 1.2.1 and 1.5.4] that for some  $C$  finite and any  $x \in \mathbb{Z}^4$ ,

$$(4.14) \quad P(S_i = x) \leq Ci^{-2} \left[ e^{-2|x|^2/i} + (|x|^2 \vee i)^{-1} \right],$$

$$(4.15) \quad C^{-1}|x|_+^{-2} \leq G(0, x) \leq C|x|_+^{-2}.$$

By (4.15),

$$E[|S_i - x|_+^{-2}] \leq CE[G(S_i, x)] = C \sum_{\ell=i}^{\infty} P(S_\ell = x).$$

Further, for some  $C$  finite and all  $i \geq 1$ ,  $x \in \mathbb{Z}^4$ ,

$$\sum_{\ell=i}^{\infty} \ell^{-2} (|x|^2 \vee \ell)^{-1} \leq C i^{-1} (|x|^2 \vee i)^{-1} \leq C \min\{i^{-1}, |x|_+^{-2}\}.$$

Thus, in view of (4.14), the same computation as for (4.8) yields the first inequality of (4.11). The second inequality of (4.11) follows (as  $a \wedge b \leq \sqrt{ab}$  for any  $a, b > 0$ ), and since the strictly positive  $|\cdot|_+$  satisfies the triangle inequality on  $\mathbb{Z}^4$ , we get first (4.12), and then (4.13), by the same reasoning that led to (4.9) and (4.10), respectively.  $\blacksquare$

*Proof of Lemma 4.1.* From (4.13), for any  $p \geq 2$ ,  $1 \leq s_1 \leq \dots \leq s_p$  and  $\{y_1, \dots, y_p\} \subset \mathbb{Z}^4$ ,

$$\begin{aligned} & E \left[ \prod_{i=1}^{p-1} |S_{s_i} - y_i|_+^{-2} |S_{s_{p-1}} - y_p|_+^{-1} \right] \\ &= E \left[ \prod_{i=1}^{p-2} |S_{s_i} - y_i|_+^{-2} |S_{s_{p-1}} - S_{s_{p-2}} - (y_{p-1} - S_{s_{p-2}})|_+^{-2} |S_{s_{p-1}} - S_{s_{p-2}} - (y_p - S_{s_{p-2}})|_+^{-1} \right] \\ &\leq C |s_{p-1} - s_{p-2}|_+^{-1/2} |y_p - y_{p-1}|_+^{-1} E \left[ \prod_{i=1}^{p-2} |S_{s_i} - y_i|_+^{-2} |S_{s_{p-2}} - y_{p-1}|_+^{-1} \right], \end{aligned}$$

where we set throughout  $s_0 = 0$  and  $y_0 = 0$ . Thus, by induction on  $p \geq 1$ ,

$$(4.16) \quad E \left[ \prod_{i=1}^{p-1} |S_{s_i} - y_i|_+^{-2} |S_{s_{p-1}} - y_p|_+^{-1} \right] \leq C^{p-1} \prod_{i=1}^{p-1} |s_i - s_{i-1}|_+^{-1/2} \prod_{i=1}^p |y_i - y_{i-1}|_+^{-1}.$$

Next, note that by (4.11), for any  $p \geq 1$ ,

$$\begin{aligned} E \left[ \prod_{i=1}^p |S_{s_i} - y_i|_+^{-2} \right] &= E \left[ \prod_{i=1}^{p-1} |S_{s_i} - y_i|_+^{-2} |S_{s_p} - S_{s_{p-1}} - (y_p - S_{s_{p-1}})|_+^{-2} \right] \\ &\leq C |s_p - s_{p-1}|_+^{-1/2} E \left[ \prod_{i=1}^{p-1} |S_{s_i} - y_i|_+^{-2} |S_{s_{p-1}} - y_p|_+^{-1} \right]. \end{aligned}$$

Hence, setting  $t_0 = 0$ , in view of (4.16) and the independence of  $(S_i)$  and  $(\tilde{S}_i)$ ,

$$(4.17) \quad E \left[ \prod_{i=1}^p |S_{s_i} - \tilde{S}_{t_i}|_+^{-2} \right] \leq C^p \prod_{i=1}^p |s_i - s_{i-1}|_+^{-1/2} E \left[ \prod_{i=1}^p |\tilde{S}_{t_i} - \tilde{S}_{t_{i-1}}|_+^{-1} \right].$$

Suppose that  $t_{\sigma(1)} \leq \dots \leq t_{\sigma(p)}$  for some permutation  $\sigma$  of  $\{1, \dots, p\}$ . Then, conditioning on  $(\tilde{S}_i, i \leq t_{\sigma(p-1)})$ , we get by (4.12) and the independence of increments, that when  $\sigma(p) = \ell$ ,

$$E \left[ \prod_{i=1}^p |\tilde{S}_{t_i} - \tilde{S}_{t_{i-1}}|_+^{-1} \right] \leq C |t_{\sigma(p)} - t_{\sigma(p-1)}|_+^{-1/2} E \left[ \prod_{i=1}^{\ell-1} |\tilde{S}_{t_i} - \tilde{S}_{t_{i-1}}|_+^{-1} |\tilde{S}_{t_{\ell+1}} - \tilde{S}_{t_{\ell-1}}|_+^{-1} \prod_{i=\ell+2}^p |\tilde{S}_{t_i} - \tilde{S}_{t_{i-1}}|_+^{-1} \right].$$

Such  $\sigma$  is a bijection from  $\{1, \dots, p-1\}$  to  $\{1, \dots, \ell-1, \ell+1, \dots, p\}$ , setting  $\sigma(0) = 0$  and continuing inductively according to the values of  $\sigma(j)$ , for  $j = p-1, \dots, 1$ , yields that

$$(4.18) \quad E \left[ \prod_{i=1}^p |\tilde{S}_{t_i} - \tilde{S}_{t_{i-1}}|_+^{-1} \right] \leq C^p \prod_{j=1}^p |t_{\sigma(j)} - t_{\sigma(j-1)}|_+^{-1/2}.$$

Combining (4.17) and (4.18), we conclude that for any non-decreasing  $(s_i)$  and  $(t_{\sigma(j)})$ ,

$$(4.19) \quad E\left[\prod_{i=1}^p |S_{s_i} - \tilde{S}_{t_i}|_+^{-2}\right] \leq C^{2p} \prod_{i=1}^p |s_i - s_{i-1}|_+^{-1/2} \prod_{j=1}^p |t_{\sigma(j)} - t_{\sigma(j-1)}|_+^{-1/2}.$$

There are at most  $k^p$  words of length  $p$  composed of  $k$  distinguished symbols. Hence, we deduce from (4.19) that for any  $n, p \in \mathbb{N}$ , by (4.15),

$$(4.20) \quad E[X_n^p] \leq \frac{1}{n^p} C^p E\left[\sum_{i,\ell \in [1,n]} |S_i - \tilde{S}_\ell|_+^{-2}\right]^p \leq C^{2p} \left[\sum_{k=1}^p k^p n^{-(p-k)/2} J_{k,n}\right]^2,$$

$$J_{k,n} := \frac{1}{n^k} \sum_{1 \leq s_1 < \dots < s_k \leq n} \prod_{i=1}^k \left(\frac{s_i}{n} - \frac{s_{i-1}}{n}\right)^{-1/2}.$$

Considering  $\theta_i = s_i/n$ , we see that  $\{J_{k,n}\}$  are for each  $k \in \mathbb{N}$ , the Riemann sums of

$$(4.21) \quad J_k := \int_{0=\theta_0 < \theta_1 < \dots < \theta_k \leq 1} \prod_{i=1}^k (\theta_i - \theta_{i-1})^{-1/2} d\theta_1 \cdots d\theta_k \leq C^k (k!)^{-1/2}$$

(for the inequality, see e.g. [26, proof of Lemma 2]). Note that  $J_k = (Q^k \mathbf{1})(0)$  for the positive linear operator  $(Qf)(x) = \int_0^{1-x} y^{-1/2} f(x+y) dy$  on  $C([0, 1])$ . Setting  $(y)_n = \lceil yn \rceil / n$ , we have that  $J_{k,n} = (Q_n^k \mathbf{1})(0)$  for the positive linear operators  $(Q_n f)(x) = \int_0^{1-x} (y)_n^{-1/2} f((x)_n + (y)_n) dy$ . It is easy to see that  $(Q_n f) \leq (Qf)$  are both non-increasing whenever  $f(\cdot)$  is non-increasing. By induction on  $k \geq 0$ , we thus have that  $Q_n^k \mathbf{1} \leq Q^k \mathbf{1}$ , pointwise, so in particular  $J_{k,n} \leq J_k$  for any  $k, n \in \mathbb{N}$ . Further,  $k! \geq (k/e)^k$  and  $J_{k,n} = 0$  unless  $k \leq n$ . Hence, in view of (4.20) and (4.21) we find that

$$E[X_n^p] \leq C^{2p} \left[\sum_{k=1}^p k^p k^{-(p-k)/2} J_k\right]^2 \leq C^{2p} \left[\sum_{k=1}^p k^{(p+k)/2} C^k (k/e)^{-k/2}\right]^2 \leq p^2 C^{4p} e^p p^p,$$

and the uniform moment bounds (4.5) on  $X_n$  follow. Turning to the moments of  $Y$ , note that

$$E[Y^p] \leq p! \int_{0 \leq s_1 \leq \dots \leq s_p \leq 1} \int_{0 \leq t_1 \leq \dots \leq t_p \leq 1} E\left[\sum_{\sigma} \prod_{i=1}^p |\beta_{s_i} - \tilde{\beta}_{t_{\sigma(i)}}|^{-2}\right] ds_1 \cdots ds_p dt_1 \cdots dt_p.$$

In view of (4.21), it thus suffices to show that for some  $C < \infty$  and any  $p \in \mathbb{N}$ ,

$$(4.22) \quad E\left[\sum_{\sigma} \prod_{i=1}^p |\beta_{s_i} - \tilde{\beta}_{t_{\sigma(i)}}|^{-2}\right] \leq C^p p! \prod_{i=1}^p (s_i - s_{i-1})^{-1/2} \prod_{j=1}^p (t_j - t_{j-1})^{-1/2}.$$

We can assume wlog that  $\{\beta_{s_i}, \tilde{\beta}_{t_j}, i, j \leq p\}$  are all distinct, as are the times  $\{s_i\}_{i \leq p}$  and  $\{t_j\}_{j \leq p}$ . Thus, following our proof of (4.19), while using (4.8)–(4.10) instead of (4.11)–(4.13), yields (4.22), thereby completing the proof of the lemma.  $\blacksquare$

The next lemma, whose proof is deferred to the end of this section, allows us to complete the proof of the limsup-LIL lower bound for SRW on  $\mathbb{Z}^4$ .

**Lemma 4.3.** *For any interval  $I$ , consider the range  $\mathcal{R}_I = \{S_i\}_{i \in I}$  and  $\tilde{\mathcal{R}}_I = \{\tilde{S}_i\}_{i \in I}$  of two independent SRW-s. Denoting by  $\hat{P}$  and  $\hat{\tau}_A$  the probability and the hitting time to a set  $A$  by another independent SRW ( $\tilde{S}_i$ ), for  $i \in [n_{2\alpha}, n - n_{2\alpha}]$ ,  $n_\alpha := n(\log n)^{-\alpha}$ ,  $\alpha > 0$ , let*

$$(4.23) \quad g_{n,\alpha}(i) := 1_{\{S_i \notin \mathcal{R}(i, i+n_{2\alpha})\}} \hat{P}^{S_i}(\hat{\tau}_{\mathcal{R}(i-n_{2\alpha}, i+n_{2\alpha})} = \infty),$$

$$(4.24) \quad \bar{g}_{n,\alpha} := E[g_{n,\alpha}(i)] = E[1_{\{S_{n_{2\alpha}} \notin \mathcal{R}(n_{2\alpha}, 2n_{2\alpha})\}} \hat{P}^{S_{n_{2\alpha}}}(\hat{\tau}_{\mathcal{R}_{2n_{2\alpha}}} = \infty)],$$

with  $g_{n,\alpha}(i) = 1$  whenever  $i \in [0, n_{2\alpha}] \cup (n - n_{2\alpha}, n]$ , and  $\tilde{g}_{n,\alpha}(i)$  defined analogously for the SRW ( $\tilde{S}_i$ ). We have that  $EY_{n,m} \leq C\sqrt{nm}$  for some  $C$  finite and all  $m, n \in \mathbb{N}$ , and if  $n^\epsilon \leq m \leq n$ , then further,

$$(4.25) \quad EY_{n,m} \leq C\sqrt{nm}(\log n)^{-2},$$

$$(4.26) \quad E[(Y_{n,m} - \underline{Y}_{n,m})^2] \leq Cnm(\log n)^{-\alpha/2},$$

for some  $C = C(\epsilon, \alpha) < \infty$  and any  $\epsilon > 0$ ,  $\alpha > 4$ , where

$$Y_{n,m} := \sum_{\substack{i \in [1, n], \\ \ell \in [1, m]}} g_{n,\alpha}(i) G(S_i, \tilde{S}_\ell) \tilde{g}_{m,\alpha}(\ell), \quad \underline{Y}_{n,m} = \bar{g}_{n,\alpha} \bar{g}_{m,\alpha} \sum_{\substack{i \in [1, n], \\ \ell \in [1, m]}} G(S_i, \tilde{S}_\ell).$$

Equipped with Lemma 4.1 and Lemma 4.3, we proceed to derive the limsup-LIL lower bound for SRW on  $\mathbb{Z}^4$ . Specifically, with  $n \mapsto R_n$  having a similar structure as  $|\mathcal{R}_n|$  for the SRW on  $\mathbb{Z}^2$ , we adapt the proof in [11, Prop. 4.4] of the limsup-LIL lower-bound for the latter sequence. Specifically, set  $p = [(-\kappa + \log_3 n)/(\log 2)]$  with  $\kappa < \infty$  large and  $k = 2^p$  (so  $k = \gamma \log_2 n$  for small  $\gamma \leq e^{-\kappa}$ ). Centering both sides of (4.2) for  $n_j = jm$ ,  $m = n/k$  (assumed for simplicity to be integer), we have that for i.i.d.  $U_j := R_{(j-1)m, jm}$  and the non-random  $\varphi_n := ER_n$ ,

$$(4.27) \quad \bar{R}_n = k\varphi_{n/k} - \varphi_n + \sum_{j=1}^k \bar{U}_j - \sum_{j=1}^{k-1} V_{0, jm, (j+1)m}.$$

Further, denoting by  $\theta$  the time shift  $S_i \mapsto S_{i+1}$ , we set

$$\chi(A, B) := \sum_{y \in A} \sum_{z \in B} P^y(\tau_{A \cup B} = \infty) G(y, z) P^z(\tau_B = \infty),$$

and for all  $i \in (0, n]$ ,

$$(4.28) \quad h_n(i) := 1_{\{S_i \notin \mathcal{R}(i, n)\}} \hat{P}^{S_i}(\hat{\tau}_{\mathcal{R}_n} = \infty) \leq g_{n,\alpha}(i)$$

(see (4.23)), recalling from (4.1) and [6, Prop. 1.6] that

$$(4.29) \quad \begin{aligned} 0 \leq V_{0, jm, (j+1)m} &\leq \chi(\mathcal{R}_{jm}, \mathcal{R}(jm, (j+1)m]) + \chi(\mathcal{R}(jm, (j+1)m], \mathcal{R}_{jm}) \\ &\leq 2 \sum_{y \in \mathcal{R}_{jm}} \sum_{z \in \mathcal{R}(jm, (j+1)m]} P^y(\tau_{\mathcal{R}_{jm}} = \infty) G(y, z) P^z(\tau_{\mathcal{R}(jm, (j+1)m]} = \infty) \\ &= 2 \sum_{i=1}^{jm} \sum_{\ell=1}^m h_{jm}(i) G(S_i, S_{jm+\ell}) h_m(\ell) \circ \theta_{jm} \\ &\leq 2 \sum_{i=1}^{jm} \sum_{\ell=1}^m g_{jm,\alpha}(i) G(S_i, S_{jm+\ell}) g_{m,\alpha}(\ell) \circ \theta_{jm} := 2W_j. \end{aligned}$$

Setting in addition

$$(4.30) \quad \underline{W}_j := \bar{g}_{jm,\alpha} \bar{g}_{m,\alpha} \sum_{i=0}^{m-1} \sum_{\ell=1}^m G(S_{jm-i}, S_{jm+\ell}),$$

$$(4.31) \quad \hat{W}_j := \bar{g}_{jm,\alpha} \bar{g}_{m,\alpha} \sum_{i=m}^{jm-1} \sum_{\ell=1}^m G(S_{jm-i}, S_{jm+\ell}),$$

we see that for any fixed  $j$ ,

$$(4.32) \quad W_j - \underline{W}_j - \hat{W}_j \stackrel{d}{=} Y_{jm,m} - \underline{Y}_{jm,m}.$$

Next, taking the expected value in [6, Prop. 2.3],

$$(4.33) \quad k\varphi_{n/k} - \varphi_n = \sum_{i=1}^p \sum_{j=1}^{2^{i-1}} E[\chi_n(i, j)] - \sum_{i=1}^p \sum_{j=1}^{2^{i-1}} E[\epsilon_n(i, j)].$$

As each  $\epsilon_n(i, j) \geq 0$  is bounded by the intersection of the ranges of two independent srw-s of length  $n/2^i$ , we have that  $\max_{i,j} E[\epsilon_n(i, j)] \leq \log n$  (see [23, Section 3.4]), and with at most  $2^p \leq C \log n$  such terms, we conclude that

$$(4.34) \quad \sum_{i=1}^p \sum_{j=1}^{2^{i-1}} E[\chi_n(i, j)] - C(\log n)^2 \leq k\varphi_{n/k} - \varphi_n \leq \sum_{i=1}^p \sum_{j=1}^{2^{i-1}} E[\chi_n(i, j)].$$

Further, in view of [6, Prop. 6.1], for  $\alpha(A_1^1)$  of (4.7) and  $n'_i := n2^{1-i}$ , we find that uniformly over  $i \leq p = o(\log n)$ , as  $n \rightarrow \infty$ ,

$$\begin{aligned} \frac{2(\log n'_i)^2}{\pi^4 n'_i} E[\chi_{n'_i}(1, 1)] &\rightarrow E[\alpha(A_1^1)] = \int_{A_1^1} E[G_\beta(\beta_s, \beta_t)] ds dt \\ &= \frac{1}{2\pi^2} E[|\beta_1|^{-2}] \int_{A_1^1} |t-s|^{-1} dt ds = \frac{\log 2}{4\pi^2}, \end{aligned}$$

where we have used (4.4), as well as having  $\int_{A_1^1} |t-s|^{-1} ds dt = \log 2$  and  $E|\beta_1|^{-2} = 1/2$ . As  $E[\chi_n(i, j)] = E[\chi_{n'_i}(1, 1)]$  for all  $i, j$  (see [6, Prop. 2.3]), and

$$p = (1 + o(1)) \frac{\log_3 n}{\log 2}, \quad h_4(n) = \frac{\pi^2 n \log_3 n}{8 (\log n)^2}$$

(see (1.4)), it thus follows that

$$(4.35) \quad \sum_{i=1}^p \sum_{j=1}^{2^{i-1}} E[\chi_n(i, j)] = (1 + o(1)) p \frac{\log 2}{4\pi^2} \frac{\pi^4 n}{2(\log n)^2} = (1 + o(1)) h_4(n),$$

and combining (4.34) and (4.35) we arrive at

$$(4.36) \quad k\varphi_{n/k} - \varphi_n = (1 + o(1)) h_4(n).$$



In view of (4.27), (4.29) and (4.36), we get our limsup-LIL lower bound, precisely as in [11, Proof of Prop. 4.4], once we find for any  $\varepsilon > 0$ , constants  $c_1 < \infty$ ,  $c_2 > 0$  and events

$$(4.37) \quad G_k \subseteq \bigcap_{j=1}^k \left\{ \overline{U}_j \geq -\frac{c_1 m}{(\log m)^2} \right\} := \bigcap_{j=1}^k B_j,$$

so  $P(G_k) \geq \frac{1}{4}c_2^k$  and for any  $k, m$  as above and  $\varepsilon > 0$ , the event  $G_k$  implies that

$$(4.38) \quad \sum_{j=1}^{k-1} W_j \leq 3\varepsilon \frac{n \log_3 n}{(\log m)^2}.$$

To this end, it suffices to construct events  $F_k$  such that for some  $c_3 < \infty$ ,

$$(4.39) \quad P(F_k) \geq c_2^k, \quad F_k \subseteq \left\{ \max_{j < k} \{\hat{W}_j\} \leq \frac{c_3 m}{(\log m)^2} \right\}.$$

Indeed, we shall see that  $P(C_i) \leq \frac{1}{4}c_2^k$  for  $k \leq \gamma \log_2 n$ ,  $\gamma > 0$  small and  $n \rightarrow \infty$ , where

$$C_1 := \left\{ \sum_{j \text{ odd}} \underline{W}_j > \varepsilon \frac{n \log_3 n}{(\log m)^2} \right\}, \quad C_2 := \left\{ \sum_{j \text{ even}} \underline{W}_j > \varepsilon \frac{n \log_3 n}{(\log m)^2} \right\},$$

$$C_3 := \left\{ \max_{j < k} \{W_j - \underline{W}_j - \hat{W}_j\} > \frac{m}{(\log m)^2} \right\}.$$

Taking  $G_k := F_k \cap_{i=1}^3 C_i^c$  this would imply that  $P(G_k) \geq \frac{1}{4}c_2^k$  for large  $k$ , and it is easy to check, as stated, that (4.38) then holds on such  $G_k$ .

Next, utilizing the union bound, (4.32), Markov's inequality and (4.26), we get that

$$P(C_3) \leq \sum_{j=1}^{k-1} P\left(W_j - \underline{W}_j - \hat{W}_j \geq \frac{m}{(\log m)^2}\right) = \sum_{j=1}^{k-1} P\left(Y_{jm,m} - \underline{Y}_{jm,m} \geq \frac{m}{(\log m)^2}\right)$$

$$\leq \frac{(\log m)^4}{m^2} \sum_{j=1}^{k-1} E\left[(Y_{jm,m} - \underline{Y}_{jm,m})^2\right] \leq C(\log m)^{4-\alpha/2} \sum_{j=1}^{k-1} j \leq Ck^2(\log m)^{4-\alpha/2}.$$

In particular, for  $\alpha > 8 + 2\gamma \log(1/c_2)$ ,  $k$  as above and  $n = mk \geq n_0$ , the preceding bound implies that  $P(C_3) \leq \frac{1}{4}c_2^k$ . Turning to deal with  $\underline{W}_j$  and  $\hat{W}_j$ , upon expressing (4.24) via the independent srw-s  $\hat{S}_i$ ,  $S_i^+ := S_{n'+i} - S_{n'}$  and  $S_i^- := S_{n'-i} - S_{n'}$ , at  $n' = n_{2\alpha}$ , it follows from [4, (1.4)] that

$$(4.40) \quad \begin{aligned} \bar{g}_{n,\alpha} &= E[1_{\{0 \notin \mathcal{R}_n^+\}} \hat{P}^0(\hat{\tau}_{\mathcal{R}_n^+ \cup \mathcal{R}^-([0, n'-1])} = \infty)] \\ &= P(0 \notin \mathcal{R}_{n'}^+, \hat{R}_\infty \cap (\mathcal{R}_{n'}^+ \cup \mathcal{R}^-([0, n'-1])) = \emptyset) = (1 + o(1)) \frac{\pi^2}{8} (\log n)^{-1} \end{aligned}$$

(note that  $\log n' = (1 + o(1)) \log n$ ). In view of (4.30), we note that  $\{m^{-1}(\bar{g}_{m,\alpha})^{-2} \underline{W}_j\}$  are, for odd  $j$ , independent copies of  $X_m$  of Lemma 4.1 (except for now including also  $\ell = 0$  in  $X_m$ ). It thus follows from (4.6) and (4.40) that for some  $c > 0$ , and all  $k, m$ ,

$$E\left[\exp\left(c m^{-1} (\log m)^2 \sum_{j \text{ odd}} \underline{W}_j\right)\right] \leq \exp(k/c).$$

Hence, for  $n = mk$ , one has by Markov's inequality that

$$P(C_1) \leq \exp(-\varepsilon ck \log_3 n) \exp(k/c)$$

decays as  $n \rightarrow \infty$ , faster than  $\frac{1}{4}c_2^k$ . By the same reasoning, this applies also for  $C_2$ .

Finally, in view of (4.15), (4.31) and (4.40), for some  $C < \infty$  and any  $m, j$ ,

$$(4.41) \quad \hat{W}_j \leq \frac{Cm^2}{(\log m)^2} \sum_{s=1}^{j-1} \text{dist}(\mathcal{R}((s-1)m, sm], \mathcal{R}((jm, (j+1)m]))^{-2}.$$

As in [11, Proof of Prop. 4.4], fixing a unit vector  $\mathbf{u}$ , we let  $F_j := \cap_{i=1}^j (A_i \cap B_i)$  for

$$A_i := \left\{ S_\ell : S_{im} \subset \mathbb{B}(i\sqrt{m}\mathbf{u}, \sqrt{m}/8), \mathcal{R}((i-1)m, im] \subset \mathbb{B}\left((i-\frac{1}{2})\sqrt{m}\mathbf{u}, \frac{3}{4}\sqrt{m}\right)\right\}.$$

The event  $F_k$  guarantees that the distance of  $\mathcal{R}((s-1)m, sm]$  from  $\mathcal{R}((jm, (j+1)m]$  be at least  $(j-s-1/2)\sqrt{m}$ , so (4.41) results with the RHS of (4.39) (for  $c_3 = C \sum (r-1/2)^{-2}$  finite). As for the LHS of (4.39), first note that  $\{B_j, j \geq 1\}$  are i.i.d. and for any  $\delta > 0$  there exists  $c_1 = c_1(\delta)$  finite, such  $P(B_1^c) \leq \delta$  uniformly in  $m$  (see [6, Thm. 1.2]). Further, by the invariance principle, there exists  $c_2 > 0$  such that,

$$\lim_{m \rightarrow \infty} \inf_{S_0 \in \mathbb{B}(0, \sqrt{m}/8)} \{P^{S_0}(A_1)\} = \inf_{\beta_0 \in \mathbb{B}(0, 1/4)} \{P(|\beta_1 - 2\mathbf{u}| < 1/4, \sup_{t \in [0,1]} |\beta_t - \mathbf{u}| < 3/2)\} \geq 2c_2.$$

As  $F_j$  is measurable on  $\sigma(S_i, i \leq jm)$ , by the Markov property of the srw and its independent, stationary increments, for any  $j \geq 1$ ,

$$P(A_j \cap B_j | F_{j-1}) \geq \inf_{S_0 \in \mathbb{B}(0, \sqrt{m}/8)} \{P^{S_0}(A_1)\} - P(B_1^c) \geq c_2,$$

provided  $\delta > 0$  is small enough and  $m \geq m_0$  finite, thereby establishing the LHS of (4.39).

We conclude this sub-section by proving Lemma 4.3. To this end, for  $\alpha > 0$  let

$$I_\alpha(n) := [n_\alpha, n - n_\alpha]^2 \cap \{(i, j) : j - i \geq n_\alpha\},$$

setting for  $(i_1, i_2) \in I_\alpha(n)$  and  $n' = n_{2\alpha}$ , the events

$$H_{i_1}^{(n)} := \bigcup_{|u| \leq \sqrt{n'} \log n'} H_{i_1}^{(n')}(u), \quad H_{i_1, i_2}^{(n)} := \bigcup_{|u|, |v| \leq \sqrt{n'} \log n'} H_{i_1, i_2}^{(n')}(u, v),$$

$$H_{i_1}^{(n')}(u) := \{S_{i_1} - S_{i_1 - n'} = u\}, \quad H_{i_1, i_2}^{(n')}(u, v) := H_{i_1}^{(n')}(u) \cap \{S_{i_2} - S_{i_2 - n'} + S_{i_1 + n'} - S_{i_1} = v\}.$$

Similarly,  $\tilde{H}_{\ell_1}^{(m)}$ ,  $\tilde{H}_{\ell_1, \ell_2}^{(m)}$ ,  $\tilde{H}_{\ell_1}^{(m')}(\tilde{u})$  and  $\tilde{H}_{\ell_1, \ell_2}^{(m')}(\tilde{u}, \tilde{v})$  for the srw  $(\tilde{S}_\ell)$ ,  $(\ell_1, \ell_2) \in I_\alpha(m)$  and  $m' = m_{2\alpha}$ .

**Lemma 4.4.** *Fix  $\alpha > 2$ ,  $\varepsilon > 0$  and a permutation  $\pi$  of  $\{1, 2\}$ . Then, for  $n^\varepsilon \leq m \leq n$ ,*

$$(4.42) \quad F_1(u, \tilde{u}) := E[G(S_{i_1}, \tilde{S}_{\ell_1}) | H_{i_1}^{(n')}(u) \cap H_{\ell_1}^{(m')}(\tilde{u})] = (1 + O((\log n)^{2-\alpha}))E[G(S_{i_1}, \tilde{S}_{\ell_1})],$$

$$(4.43) \quad \begin{aligned} F_2(u, v, \tilde{u}, \tilde{v}) &:= E[G(S_{i_1}, \tilde{S}_{\ell_{\pi_1}})G(S_{i_2}, \tilde{S}_{\ell_{\pi_2}}) | H_{i_1, i_2}^{(n')}(u, v) \cap \tilde{H}_{\ell_1, \ell_2}^{(m')}(\tilde{u}, \tilde{v})] \\ &= (1 + O((\log n)^{2-\alpha}))E[G(S_{i_1}, \tilde{S}_{\ell_{\pi_1}})G(S_{i_2}, \tilde{S}_{\ell_{\pi_2}})], \end{aligned}$$

uniformly over  $(i_1, i_2) \in I_\alpha(n)$ ,  $(\ell_1, \ell_2) \in I_\alpha(m)$ ,  $|u|, |v| \leq \sqrt{n'} \log n'$  and  $|\tilde{u}|, |\tilde{v}| \leq \sqrt{m'} \log m'$ .

*Proof.* For  $(i_1, i_2) \in I_\alpha(n)$ , the law of  $(S_{i_1}, S_{i_2})$  given  $H_{i_1, i_2}^{(n')}(u, v)$  is as  $(u + S_{i_1 - n'}^{(1)}, u + v + S_{i_2 - 3n'}^{(1)})$  for an independent srw  $S_i^{(1)}$ . Similarly, when  $(\ell_1, \ell_2) \in I_\alpha(m)$ , the law of  $(\tilde{S}_{\ell_1}, \tilde{S}_{\ell_2})$  given  $\tilde{H}_{\ell_1, \ell_2}^{(m')}(u, v)$  is as  $(\tilde{u} + \tilde{S}_{\ell_1 - m'}^{(1)}, \tilde{u} + \tilde{v} + \tilde{S}_{\ell_2 - 3m'}^{(1)})$ . Consequently,

$$F_1(u, \tilde{u}) = E[G(u + S_{i_1 - n'}, \tilde{u} + \tilde{S}_{\ell_1 - m'})],$$

$$F_2(u, v, \tilde{u}, \tilde{v}) = \begin{cases} E[G(u + S_{i_1 - n'}, \tilde{u} + \tilde{S}_{\ell_1 - m'})G(u + v + S_{i_2 - 3n'}, \tilde{u} + \tilde{v} + \tilde{S}_{\ell_2 - 3m'})], & \text{if } \pi_1 = 1, \\ E[G(u + S_{i_1 - n'}, \tilde{u} + \tilde{v} + \tilde{S}_{\ell_2 - 3m'})G(u + v + S_{i_2 - 3n'}, \tilde{u} + \tilde{S}_{\ell_1 - m'})], & \text{if } \pi_1 = 2. \end{cases}$$

Note that for some  $C = C(\epsilon)$  finite,  $c = c(\epsilon) > 0$ , any  $m \geq n^\epsilon$  and  $(\ell_1, \ell_2)$ ,

$$(4.44) \quad P((H_{\ell_1}^{(m)})^c) \leq P((H_{\ell_1, \ell_2}^{(m)})^c) \leq 2P(|S_{2m'}| > \sqrt{m'} \log m') \leq C e^{-c(\log n)^2},$$

with the same bound applying also for  $P((H_{i_1, i_2}^{(n)})^c)$ . Now, by (4.15) and (4.19) (at  $p = 1, 2$ ),

$$(4.45) \quad E[G(S_i, \tilde{S}_\ell)] \leq C i^{-1/2} \ell^{-1/2},$$

$$(4.46) \quad E[G(S_{i_1}, \tilde{S}_{\ell_{\pi_1}})G(S_{i_2}, \tilde{S}_{\ell_{\pi_2}})] \leq C i_1^{-1/2} (i_2 - i_1)^{-1/2} \ell_1^{-1/2} (\ell_2 - \ell_1)^{-1/2},$$

with the LHS of (4.45) and (4.46) being the expected values of  $F_1(\cdot)$  and  $F_2(\cdot)$  according to the joint law of the corresponding srw increments (for independent srw  $S_i$  and  $\tilde{S}_\ell$ ). In view of (4.44), it thus suffices to bound the maximum fluctuation of  $F_s(\cdot)$ ,  $s = 1, 2$  over  $|u|, |v| \leq \sqrt{n'} \log n'$  and  $|\tilde{u}|, |\tilde{v}| \leq \sqrt{m'} \log m'$ , by  $C(\log n)^{2-\alpha}$  times the RHS of (4.45) and (4.46), respectively. To this end, since  $F_1(\cdot)$  depends only on  $u - \tilde{u}$  and  $F_2(\cdot)$  depends only on  $u - \tilde{u}$  and  $v - \tilde{v}$  if  $\pi_1 = 1$ , or on  $u + v - \tilde{u}$  and  $v + \tilde{v}$  if  $\pi_1 = 2$ , we may wlog fix  $\tilde{u} = \tilde{v} = 0$  and consider the maximum fluctuation of

$$F_1(u) = E[G(u + S_{i_1 - n'}, \tilde{S}_{\ell_1 - m'})],$$

$$F_2(u, v) = \begin{cases} E[G(u + S_{i_1 - n'}, \tilde{S}_{\ell_1 - m'})G(v + S_{i_2 - 3n'}, \tilde{S}_{\ell_2 - 3m'})], & \text{if } \pi_1 = 1, \\ E[G(u + S_{i_1 - n'}, \tilde{S}_{\ell_2 - 3m'})G(v + S_{i_2 - 3n'}, \tilde{S}_{\ell_1 - m'})], & \text{if } \pi_1 = 2. \end{cases}$$

over  $|u|, |v| \leq 3\sqrt{n'} \log n'$ . Further, with both  $n_\alpha/n'$  and  $m_\alpha/m'$  diverging (as  $(\log n)^\alpha$ ), it follows that uniformly over  $(i_1, i_2) \in I_\alpha(n)$ ,  $(\ell_1, \ell_2) \in I_\alpha(m)$  and  $m \geq n^\epsilon$ , the RHS of (4.45) and (4.46) also bound  $F_1(0)$  and  $F_2(0, 0)$ , respectively. Consequently, it suffices to show that for some  $C$  finite and all  $|u|, |v| \leq 3\sqrt{n'} \log n'$ ,

$$(4.47) \quad |F_1(u) - F_1(0)| \leq C(\log n)^{2-\alpha} F_1(0),$$

$$(4.48) \quad |F_2(u, v) - F_2(0, 0)| \leq C(\log n)^{2-\alpha} F_2(0, 0).$$

Turning to this task, since  $t_2 := \ell_2 - \ell_1 - 2m' \geq 0$ ,  $t_3 := i_2 - i_1 - 2n' \geq 0$ ,  $G(x, y) = G(y - x, 0)$  and  $S_i \stackrel{(d)}{=} -S_i$ , we can further simplify the functions  $F_s(\cdot)$  to be

$$F_1(u) = E[G(S_{t_1}, u)],$$

$$F_2(u, v) = \begin{cases} E[G(S_{t_1}, u)G(S_{t_1+t_2+t_3}, v)], & \text{if } \pi_1 = 1, \\ E[G(S_{t_1+t_2}, u)G(S_{t_1+t_3}, v)], & \text{if } \pi_1 = 2, \end{cases}$$

where  $t_1 = i_1 + \ell_1 - n' - m'$ . Denoting by  $p_j(u) := P(S_j = u)$ , it is easy to check that

$$(4.49) \quad F_1(u) = \sum_{j_0 > t_1} p_{j_0}(u), \quad F_2(u, v) = \sum_{j_1, j_2} p_{j_1}(u) p_{j_2}(v),$$

where the sum is over  $j_1 > t_1$  and  $j_2 > t_1 + t_2 + t_3$  in case  $\pi = 1$ , and over  $j_1 > t_1 + t_2$ ,  $j_2 > t_1 + t_3$  when  $\pi_1 = 2$ . By the local CLT for the SRW on  $\mathbb{Z}^4$ , we have for some  $C < \infty$  that

$$\left| \frac{p_j(u) + p_{j+1}(u)}{p_j(0) + p_{j+1}(0)} - 1 \right| \leq \frac{C|u|^2}{9j} \leq Cn'(\log n')^2 t_1^{-1} \leq 2C(\log n)^{2-\alpha}$$

throughout the range of parameters considered here (utilizing the fact that  $j_0, j_1, j_2 \geq t_1 \geq n_\alpha/2$ ). The same bound applies with  $v$  instead of  $u$ , and plugging these bounds in (4.49) results with (4.47)-(4.48), thereby completing the proof of the lemma.  $\blacksquare$

*Proof of Lemma 4.3.* To show (4.26), fix a permutation  $\pi$  of  $\{1, 2\}$  and consider the variables

$$\begin{aligned} g_0 &:= g_{n,\alpha}(i_1)\tilde{g}_{m,\alpha}(\ell_{\pi_1})g_{n,\alpha}(i_2)\tilde{g}_{m,\alpha}(\ell_{\pi_2}), \\ g_1 &:= g_{n,\alpha}(i_1)\tilde{g}_{m,\alpha}(\ell_{\pi_1})\bar{g}_{n,\alpha}\bar{g}_{m,\alpha}, & g_2 &:= \bar{g}_{n,\alpha}\bar{g}_{m,\alpha}g_{n,\alpha}(i_2)\tilde{g}_{m,\alpha}(\ell_{\pi_2}), \\ \mathcal{G} &:= G(S_{i_1}, \tilde{S}_{\ell_{\pi_1}})G(S_{i_2}, \tilde{S}_{\ell_{\pi_2}}). \end{aligned}$$

Note that  $g_{n,\alpha}(i)$  and  $\tilde{g}_{m,\alpha}(\ell)$  are measurable on  $\mathcal{F}_i := (S_{i+j} - S_i, j \in (-n', n'])$ ,  $n' := n_{2\alpha}$ , and  $\tilde{\mathcal{F}}_\ell := (\tilde{S}_{\ell+j} - \tilde{S}_\ell, j \in (-m', m'])$ ,  $m' = m_{2\alpha}$ , respectively. Further, when  $(i_1, i_2) \in I_\alpha(n)$  and  $(\ell_1, \ell_2) \in I_\alpha(m)$ , under the event  $H_{i_1, i_2}^{(n')}(u, v) \cap \tilde{H}_{\ell_1, \ell_2}^{(m')}(\tilde{u}, \tilde{v})$  the law of  $\mathcal{G}$  given  $\mathcal{F}_{i_1}, \mathcal{F}_{i_2}, \tilde{\mathcal{F}}_{\ell_1}$  and  $\tilde{\mathcal{F}}_{\ell_2}$  is determined by  $H_{i_1, i_2}^{(n')}(u, v) \cap \tilde{H}_{\ell_1, \ell_2}^{(m')}(\tilde{u}, \tilde{v})$ . Thus, for  $s = 0, 1, 2$ , and any such  $(i_1, i_2), (\ell_1, \ell_2)$ ,

$$E[g_s \mathcal{G} 1_{H_{i_1, i_2}^{(n')}(u, v) \cap \tilde{H}_{\ell_1, \ell_2}^{(m')}(\tilde{u}, \tilde{v})}] = E[g_s 1_{H_{i_1, i_2}^{(n')}(u, v) \cap \tilde{H}_{\ell_1, \ell_2}^{(m')}(\tilde{u}, \tilde{v})} E[\mathcal{G} | H_{i_1, i_2}^{(n')}(u, v) \cap \tilde{H}_{\ell_1, \ell_2}^{(m')}(\tilde{u}, \tilde{v})]].$$

With  $g_s \geq 0$  and  $E[g_s] = \bar{g}_{n,\alpha}^2 \bar{g}_{m,\alpha}^2$  whenever  $(i_1, i_2) \in I_\alpha(n)$  and  $(\ell_1, \ell_2) \in I_\alpha(m)$ , we get from (4.43) (of Lemma 4.4), that for some universal  $C < \infty$ ,

$$\begin{aligned} & \left| E[g_s \mathcal{G} 1_{H_{i_1, i_2}^{(n)} \cap \tilde{H}_{\ell_1, \ell_2}^{(m)}}] - E[g_s 1_{H_{i_1, i_2}^{(n)} \cap \tilde{H}_{\ell_1, \ell_2}^{(m)}}] E[\mathcal{G}] \right| \\ & \leq \sum_{\substack{|u|, |v| \leq \sqrt{n'} \log n', \\ |\tilde{u}|, |\tilde{v}| \leq \sqrt{m'} \log m'}} E[g_s 1_{H_{i_1, i_2}^{(n')}(u, v) \cap \tilde{H}_{\ell_1, \ell_2}^{(m')}(\tilde{u}, \tilde{v})}] \left| E[\mathcal{G} | H_{i_1, i_2}^{(n')}(u, v) \cap \tilde{H}_{\ell_1, \ell_2}^{(m')}(\tilde{u}, \tilde{v})] - E[\mathcal{G}] \right| \\ (4.50) \quad & \leq C(\log n)^{2-\alpha} \bar{g}_{n,\alpha}^2 \bar{g}_{m,\alpha}^2 E[\mathcal{G}]. \end{aligned}$$

In addition, with  $g_s \in [0, 1]$ ,  $\mathcal{G}$  uniformly bounded and  $(\log m)/(\log n) \geq \epsilon$ , we have from (4.44) that

$$\begin{aligned} & E[g_s \mathcal{G} 1_{(H_{i_1, i_2}^{(n)} \cap \tilde{H}_{\ell_1, \ell_2}^{(m)})^c}] + E[g_s 1_{(H_{i_1, i_2}^{(n)} \cap \tilde{H}_{\ell_1, \ell_2}^{(m)})^c}] E[\mathcal{G}] \leq C_1 (P((H_{i_1, i_2}^{(n)})^c) + P((\tilde{H}_{\ell_1, \ell_2}^{(m)})^c)) \\ (4.51) \quad & \leq 2C e^{-c(\log n)^2}. \end{aligned}$$

Combining (4.50) and (4.51) for  $s = 0, 1, 2$ , we thus find for the zero-mean

$$g := g_0 - g_1 - g_2 + \bar{g}_{n,\alpha}^2 \bar{g}_{m,\alpha}^2 = (g_{n,\alpha}(i_1)\tilde{g}_{m,\alpha}(\ell_{\pi_1}) - \bar{g}_{n,\alpha}\bar{g}_{m,\alpha})(g_{n,\alpha}(i_2)\tilde{g}_{m,\alpha}(\ell_{\pi_2}) - \bar{g}_{n,\alpha}\bar{g}_{m,\alpha}),$$

that uniformly over  $\pi, (i_1, i_2) \in I_\alpha(n)$  and  $(\ell_1, \ell_2) \in I_\alpha(m)$ ,

$$(4.52) \quad |E[g \cdot \mathcal{G}]| \leq 3C(\log n)^{2-\alpha} \bar{g}_{n,\alpha}^2 \bar{g}_{m,\alpha}^2 E[\mathcal{G}] + 6C e^{-c(\log n)^2}.$$

Further, as  $|g| \leq 1$ , we have from (4.15) and (4.19) at  $p = 2$ , that for some  $C_2 < \infty$  and uniformly over all  $(i_1, i_2) \in [1, n]^2, (\ell_1, \ell_2) \in [1, m]^2$ ,

$$(4.53) \quad |E[g \cdot \mathcal{G}]| \leq E[\mathcal{G}] \leq C_2 (i_1 \wedge i_2)^{-1/2} |i_2 - i_1|_+^{-1/2} (\ell_1 \wedge \ell_2)^{-1/2} |\ell_2 - \ell_1|_+^{-1/2}.$$

Next note that from (4.40) we have for some  $C_3, C < \infty$ ,

$$\begin{aligned} \bar{g}_{n,\alpha}^2 \bar{g}_{m,\alpha}^2 \sum_{\pi} \sum_{\substack{(i_1, i_2) \in [1, n]^2, \\ (\ell_1, \ell_2) \in [1, m]^2}} E[\mathcal{G}] &\leq \frac{C_3}{(\log n)^2 (\log m)^2} \sum_{\substack{1 \leq i_1 \leq i_2 \leq n \\ 1 \leq \ell_1 \leq \ell_2 \leq m}} i_1^{-1/2} |i_2 - i_1|_+^{-1/2} \ell_1^{-1/2} |\ell_2 - \ell_1|_+^{-1/2} \\ &\leq \frac{Cnm}{(\log n)^2 (\log m)^2}. \end{aligned}$$

With the right most term of (4.52) being  $o(n^{-5})$ , it follows that the overall contribution to

$$E[(Y_{n,m} - \underline{Y}_{n,m})^2] = \frac{1}{2} \sum_{\pi} \sum_{\substack{(i_1, i_2) \in [1, n]^2, \\ (\ell_1, \ell_2) \in [1, m]^2}} E[g \cdot \mathcal{G}]$$

from  $i_1, i_2 \in [n_\alpha, n - n_\alpha]$  with  $|i_2 - i_1| \geq n_\alpha$  and  $\ell_1, \ell_2 \in [m_\alpha, m - m_\alpha]$  with  $|\ell_2 - \ell_1| \geq m_\alpha$  is at most  $O(nm(\log n)^{-2-\alpha})$ , as specified in (4.26). Further, the sum over the RHS of (4.53) under any of the following three restrictions

$$|i_2 - i_1| < n_\alpha, \quad i_1 \wedge i_2 < n_\alpha, \quad i_1 \vee i_2 > n - n_\alpha,$$

is at most  $O(nm \sqrt{n_\alpha/n}) = O(nm(\log n)^{-\alpha/2})$ . With  $(\log m)/(\log n) \geq \epsilon$ , this applies also when summing the RHS of (4.53) under each of the analogous restrictions  $|\ell_2 - \ell_1| < m_\alpha$ ,  $\ell_1 \wedge \ell_2 < m_\alpha$ , or  $\ell_1 \vee \ell_2 > m - m_\alpha$ . As  $\alpha/2 < 2 + \alpha$ , we have thus established (4.26).

Turning to (4.25), we similarly have from (4.42) of Lemma 4.4 that for some  $C < \infty$  and  $c > 0$ , uniformly over  $m \in [n^\epsilon, n]$ ,  $i \in [n_\alpha, n - n_\alpha]$  and  $\ell \in [m_\alpha, m - m_\alpha]$ ,

$$\begin{aligned} E[g_{n,\alpha}(i) \tilde{g}_{m,\alpha}(\ell) G(S_i, \tilde{S}_\ell)] &\leq [1 + C(\log n)^{2-\alpha}] \bar{g}_{n,\alpha} \bar{g}_{m,\alpha} E[G(S_i, \tilde{S}_\ell)] + 2Ce^{-c(\log m)^2} \\ (4.54) \quad &\leq C(\log n)^{-2} i^{-1/2} \ell^{-1/2} + 2Ce^{-c(\log m)^2} \end{aligned}$$

(using in the latter inequality also (4.40), (4.15) and (4.19) at  $p = 1$ ). As  $\log m \geq \epsilon \log n$ , the sum of the RHS of (4.54) over  $i \leq n$  and  $\ell \leq m$  is at most as specified (ie.  $O(\sqrt{nm}(\log n)^{-2})$ ). Further, even when  $i < n_\alpha$  or  $\ell < m_\alpha$  or  $i > n - n_\alpha$  or  $\ell > m - m_\alpha$ , we still get the bound  $Ci^{-1/2}\ell^{-1/2}$  on the LHS of (4.54). The sum of  $i^{-1/2}\ell^{-1/2}$  subject to any one of the latter four restrictions is at most  $O(\sqrt{m_\alpha n}) = O(\sqrt{nm}(\log m)^{-\alpha/2})$ , which is as required (for  $\alpha > 4$ ).

Finally, recall that for some  $C, C_3$  finite and all  $m, n \in \mathbb{N}$ ,

$$EY_{m,n} \leq \sum_{i=1}^n \sum_{\ell=1}^m E[G(S_i, \tilde{S}_\ell)] \leq C_3 \sum_{i=1}^n \sum_{\ell=1}^m i^{-1/2} \ell^{-1/2} \leq C \sqrt{nm},$$

as claimed. ■

**4.2. The upper bound in the limsup-LIL.** As in case of the capacity limsup-LIL lower bound, we adapt here the relevant element from the proof of the limsup-LIL of  $|\mathcal{R}_n|$  and srw  $\mathbb{Z}^2$ , namely [11, Prop. 4.1]. To this end, we first establish a key approximate additivity for  $\varphi_n := ER_n$ .

**Lemma 4.5.** *There exists  $c'$  finite, such that for any  $a, b \in \mathbb{N}$ ,*

$$(4.55) \quad 0 \leq \varphi_a + \varphi_b - \varphi_{a+b} \leq c' \frac{\alpha^{1/4} (a+b)}{(\log(a+b))^2},$$

where  $\varphi_n := ER_n$  and  $\alpha := \min(a, b)/(a+b)$ .

*Proof.* Starting at the expected value of (4.1), we get by the same reasoning we have used in deriving (4.29), that

$$0 \leq E[V_{0,a,a+b}] = \varphi_a + \varphi_b - \varphi_{a+b} \leq 2E \sum_{i=1}^a \sum_{\ell=1}^b g_{a,\alpha}(i) G(S_i, \tilde{S}_\ell) \tilde{g}_{b,\alpha}(\ell) = 2EY_{a,b}.$$

Assuming wlog that  $a \leq b$ , it thus suffices to verify that  $EY_{a,b} \leq Ca^{1/4}b^{3/4}/(\log b)^2$  (yielding (4.55) for some  $c'(C) < \infty$ ). Indeed, for  $a \geq \sqrt{b}$  this follows from (4.25), whereas if  $a < \sqrt{b}$  then even the bound  $EY_{a,b} \leq C\sqrt{ab}$  which we have from Lemma 4.3, suffices. ■

Utilizing Lemma 4.5, we establish sharp tail estimates for  $\max_{j \leq n} \{\bar{R}_j\}$  (in particular, improving upon [19, Lemma 2.5]).

**Lemma 4.6.** *For some  $c > 0$ ,  $C < \infty$  and all  $n$ ,*

$$(4.56) \quad E[e^{cD^{(n)}}] \leq C, \quad D^{(n)} := \frac{(\log n)^2}{n} \max_{0 \leq j \leq n} \{\bar{R}_j\}.$$

*Proof.* Following the proof of [11, Lemma 4.3], it is easy to verify that (4.56) is a direct consequence of having [11, (4.5) & (4.6)] hold in case of  $G_j^n := \frac{(\log n)^2}{n} \bar{R}_j$ ,  $j \leq n$ . Furthermore, this argument applies even if the power  $\alpha^{1/2}$  on the right-most term in [11, (4.5)] is replaced by  $\alpha^{1/4}$ . Thus, it suffices for our purpose to verify that for some  $c_1, c_2 < \infty$  and all  $a, b \geq 0$ ,

$$(4.57) \quad \bar{R}_{a+b} - \bar{R}_a - \bar{R}_b \circ \theta_a \leq c_1 \left( \frac{\min(a, b)}{a+b} \right)^{1/4} \frac{(a+b)}{(\log(a+b))^2},$$

$$(4.58) \quad E[(\bar{R}_b \circ \theta_a)^2] \leq \frac{c_2 b^2}{(\log b)^4}.$$

Setting  $c_1 = c'$ , note that we have (4.57) in view of Lemma 4.5 and the non-negativity of  $V_{0,a,a+b}$  of (4.1), whereas for (4.58) see [6, Cor. 1.5]. ■

For the limsup-LIL upper bound, by Borel-Cantelli it suffices to show that for any  $q > 1$ ,  $\gamma > 0$  and  $\varepsilon > 0$ ,

$$(4.59) \quad \sum_i P\left(\frac{(\log m)^2}{m} \max_{r_{i-1} < \ell \leq r_i} \{\bar{R}_\ell\} \geq \left(\frac{\pi^2}{8} + 2\varepsilon\right)k \log k\right) < \infty,$$

where  $r_i = q^i$ ,  $k = 2^p$  for  $p = \lceil (\log \gamma + \log_3 r_i) / \log 2 \rceil$  and  $m = \lceil r_i/k \rceil$ . Now, considering (4.2) for  $n_j = jm$ ,  $j < k'$  and  $n_{k'} = \ell$  it follows from (4.55) that

$$(4.60) \quad \frac{(\log m)^2}{m} \max_{(k'-1)m < \ell \leq k'm} \{\bar{R}_\ell\} \leq \sum_{j=1}^{k'} D_j^{(m)} + \frac{(\log m)^2}{m} (k' \varphi_m - \varphi_{k'm}) + c',$$

where  $D_j^{(m)}$  are i.i.d. copies of  $D^{(m)}$  of Lemma 4.6. With  $k' \mapsto (k' \varphi_m - \varphi_{k'm})$  non-decreasing and  $D^{(m)} \geq 0$ , the maximum over  $k' \leq k$  of the RHS of (4.60), is attained at  $k' = k$ . Further, en route to (4.36), we showed that as  $p = o(\log m) \rightarrow \infty$ ,

$$(4.61) \quad \frac{1}{k \log k} \frac{(\log m)^2}{m} (k \varphi_m - \varphi_{km}) \rightarrow \frac{\pi^2}{8}.$$

Thus, noting that (4.56) results with

$$P\left(\sum_{j=1}^k D_j^{(m)} \geq \varepsilon k \log k\right) \leq C^k e^{-\varepsilon c k (\log k)},$$

which is summable over  $i$  for our choice of  $k = \gamma \log i$ , we have established (4.59) and thereby completed the proof of the limsup-LIL.

**4.3. Non-random and positive liminf-LIL.** Setting hereafter  $\bar{h}_4(n) := n(\log_2 n)/(\log n)^2$ , we first show that the  $[-\infty, \infty]$ -valued,

$$c_\star := -\liminf_{r \rightarrow \infty} \left\{ \frac{\bar{R}_r}{\bar{h}_4(r)} \right\},$$

is non-random. Indeed, recall (2.6) that  $\text{Cap}(\mathcal{R}_r) - \text{Cap}(\mathcal{R}[k, r]) \in [0, k]$ . Thus, for any  $k$  finite, changing  $\mathcal{R}_k$  without altering  $S_k$  yields at most a difference of  $k$  in the value of  $\text{Cap}(\mathcal{R}_r)$ , implying by the Hewitt-Savage zero-one law that  $c_\star$  is non-random. To show that  $c_\star > 0$ , it suffices to establish this for the sub-sequence  $r_{j+1} = r_j + 2n_j$ , where  $n_j := 2^{j^2}$  and  $r_0 := 0$ . To this end, setting  $Q_n := |\mathcal{R}(0, n] \cap \mathcal{R}(n, 2n]|$  we have from [6, Prop. 1.6] that

$$(4.62) \quad V_{0,n,2n} \geq 2 \inf_{j, \ell \in [1, 2n]} \{G(S_j, S_\ell)\} R_n R_{n,2n} - Q_n.$$

Recall from [23, Section 3.4] that  $EQ_n \leq C_0 \log n$  for some  $C_0 < \infty$  and all  $n$ . Hence, by Markov's inequality

$$(4.63) \quad P(\hat{A}_n^c) := P(Q_n \geq (\log n)^3) \leq C_0 (\log n)^{-2}.$$

Further, recall [6, (1.4) and Cor. 1.5], that  $ER_n \geq n/(\log n)$  and  $\text{Var}(R_n) \leq C_1 n^2/(\log n)^4$  for some  $C_1 < \infty$  and all  $n$  large, in which case by Markov's inequality

$$P\left(R_n \leq \frac{n}{2 \log n}\right) \leq \left(\frac{2 \log n}{n}\right)^2 \text{Var}(R_n) \leq \frac{4C_1}{(\log n)^2}.$$

Consequently, by the union bound,

$$(4.64) \quad P(A_n^c) := P\left(\min(R_n, R_{n,2n}) \leq \frac{n}{2 \log n}\right) \leq 8C_1 (\log n)^{-2}.$$

Next, from (4.15) we have that

$$F_{k,m} := \left\{ \max_{j \leq 2km} |S_j| \leq \sqrt{m} \right\} \implies \inf_{j, \ell \in [1, 2km]} \{G(S_j, S_\ell)\} \geq (4Cm)^{-1}.$$

Setting  $c = 1/(10C)$  it thus follows from (4.62) that for all  $n = km \geq n'(C)$ , on the event  $G_n := F_{k,m} \cap \hat{A}_n \cap A_n$ ,

$$V_{0,n,2n} \geq \frac{2}{4Cm} \left(\frac{n}{2 \log n}\right)^2 - (\log n)^3 \geq \frac{cnk}{(\log n)^2}.$$

Similarly to our derivation of the LHS of (4.39), it follows from the invariance principle that  $P(F_{k,m}) \geq c_2^k$  for some  $c_2 > 0$  and any  $k, m \geq 1$ . By (4.63)-(4.64) this implies in turn that  $P(G_n) \geq \frac{1}{3} c_2^k$  for  $k = 2^p = \lceil \gamma \log_2 n \rceil$ , provided  $\gamma' := \gamma \log(1/c_2) < 2$  and  $n \geq n'$ . To summarize, we have that for  $c' = c\gamma > 0$  and all  $n \geq n'$ ,

$$P\left(V_{0,n,2n} \geq c' \bar{h}_4(n)\right) \geq (\log n)^{-\gamma'}.$$



The same applies for the mutually independent  $\{V_{r_j, r_j+n_j, r_{j+1}}\}$ , hence upon fixing  $\gamma' < 1/2$  we get by the second Borel-Cantelli lemma, that a.s.

$$(4.65) \quad \limsup_{j \rightarrow \infty} \left\{ \bar{h}_4(n_j)^{-1} V_{r_j, r_j+n_j, r_{j+1}} \right\} \geq c' > 0.$$

Now, as in (4.27), for any  $r \geq 0$ ,  $n \geq 1$ ,

$$(4.66) \quad \bar{R}_{r, r+2n} = 2\varphi_n - \varphi_{2n} + \bar{R}_{r, r+n} + \bar{R}_{r+n, r+2n} - V_{r, r+n, r+2n}.$$

Considering (4.34) for  $p = 1$  (that is,  $k = 2$ ), we see that as  $n \rightarrow \infty$ ,

$$(4.67) \quad \frac{(\log n)^2}{n} [2\varphi_n - \varphi_{2n}] \leq \frac{(\log n)^2}{n} E[\chi_{2n}(1, 1)] \rightarrow \frac{\pi^2 \log 2}{4}.$$

Further, recall (4.59) that for any  $\delta > 0$ ,

$$(4.68) \quad \sum_j P(\bar{R}_{n_j} \geq (1 + \delta)h_4(n_j)) < \infty.$$

The same applies of course also for  $\bar{R}_{n_j} \circ \theta^{r_j}$  and  $\bar{R}_{n_j} \circ \theta^{r_j+n_j}$ , so with  $h_4(n)/\bar{h}_4(n) \rightarrow 0$ , we deduce from (4.65)-(4.68) (at  $n = n_j$  and  $r = r_j$ ), that a.s.

$$\liminf_{j \rightarrow \infty} \left\{ \frac{\bar{R}_{r_j, r_{j+1}}}{\bar{h}_4(n_j)} \right\} \leq - \limsup_{j \rightarrow \infty} \left\{ \bar{h}_4(n_j)^{-1} V_{r_j, r_j+n_j, r_{j+1}} \right\} \leq -c'.$$

Now from (4.57) (at  $a = r_j$ ,  $b = 2n_j$ ),

$$\bar{R}_{r_{j+1}} \leq \bar{R}_{r_j} + \bar{R}_{r_j, r_{j+1}} + \frac{c_1 r_{j+1}}{(\log r_{j+1})^2},$$

and since  $r_{j+1} \leq 3n_j$ , dividing by  $\bar{h}_4(n_j)$  and taking limits, yields that

$$-3c_\star \leq \liminf_{j \rightarrow \infty} \left\{ \bar{h}_4(n_j)^{-1} \bar{R}_{r_{j+1}} \right\} \leq -c' + \limsup_{j \rightarrow \infty} \left\{ \bar{h}_4(n_j)^{-1} \bar{R}_{r_j} \right\}.$$

The last term is a.s. zero (as (4.68) applies also for  $\{r_j\}$  instead of  $\{n_j\}$  and  $h_4(n)/\bar{h}_4(n) \rightarrow 0$ ), so we conclude that  $c_\star \geq c'/3 > 0$ .

**4.4. Finiteness of the liminf-LIL.** We show that  $c_\star \leq c_o < \infty$  by following the proof of the upper bound of [10, Thm. 1.7] (on the liminf LIL of  $|\mathcal{R}_n|$  in  $\mathbb{Z}^2$ ), while replacing [10, Thm. 1.5] and [10, Lemma 10.3], respectively, by

$$(4.69) \quad \sup_n \left\{ (\log n)(\log_2 n)^2 P\left( -\bar{R}_n > c_o \bar{h}_4(n) \right) \right\} < \infty,$$

$$(4.70) \quad \sup_n \left\{ (\log n)^2 P\left( \max_{n/q_0 \leq k \leq n} (\bar{R}_n - \bar{R}_k) > \epsilon \bar{h}_4(n) \right) \right\} < \infty,$$

holding for some  $c_o < \infty$ , any  $\epsilon > 0$  and some  $q_0(\epsilon) > 1$ .

Similarly to  $|\mathcal{R}_n|$ , the capacity is sub-additive (see (4.1)), and upon centering satisfies (4.57), which is the analog of [10, (10.2)]. Thus, the bound (4.70) follows as in [10, Proof of Lemma 10.3], now using (4.56) to arrive at [10, (10.14)] and to bound the RHS of [10, (10.15)].

Since  $|\bar{R}_n - \bar{R}_{n'}| \leq |n - n'|$ , it suffices to prove (4.69) only for  $n = mk$ ,  $k = 2^p$ ,  $p = [(\log_2 n)/\log 2]$ . Setting such values and  $n'_u := 2^{-u}n$  for  $1 \leq u \leq p$ , we have as in (4.27), now using an alternative expression for  $\Delta_{n,k}$  of (4.2), that

$$(4.71) \quad -\bar{R}_n = \bar{\Delta}_{n,k} - \sum_{j=1}^k \bar{U}_j, \quad \Delta_{n,k} = \sum_{u=1}^p \sum_{j=1}^{2^{u-1}} V_{(2j-2)n'_u, (2j-1)n'_u, 2jn'_u},$$

with  $k$  i.i.d. copies  $\{\bar{U}_j\}$  of  $\bar{R}_m$  and the i.i.d. variables  $\{V_{(2j-2)n'_u, (2j-1)n'_u, 2jn'_u}\}$  per fixed  $u \geq 1$ . Since  $\text{Var}(\bar{R}_m) \leq C_1 m^2 / (\log m)^4$  it follows by Markov's inequality, that

$$(4.72) \quad P\left(-\sum_{j=1}^k \bar{U}_j \geq \epsilon c_o \bar{h}_4(n)\right) \leq (\epsilon c_o \bar{h}_4(n))^{-2} \frac{C_1 k m^2}{(\log m)^4} \leq \frac{2C_1}{(\epsilon c_o)^2 k (\log_2 n)^2}.$$

Setting  $c_o > (1 - \epsilon)^{-1} c_1^{-1}$ , we arrive at (4.69) out of (4.71), (4.72) and the following lemma.

**Lemma 4.7.** *For some  $c_1 > 0$  and any  $\lambda > 0$ ,*

$$(4.73) \quad \limsup_{n \rightarrow \infty} \frac{1}{\log_2 n} \log P(\bar{\Delta}_{n,k} \geq \lambda \bar{h}_4(n)) \leq -c_1 \lambda,$$

where  $\Delta_{n,k}$  are as in (4.3) for  $k = 2^p$  and  $p = [(\log_2 n)/\log 2]$ .

We note in passing [19, Lemma 2.6], which is somewhat related to Lemma 4.7. The proof of Lemma 4.7 relies in turn on our next result.

**Lemma 4.8.** *Set  $p = [(\log_2 n)]$  and for any  $r \in \mathbb{N}$ , the partition  $I_i^{(r)} := ((i-1)r, ir]$  of  $\mathbb{N}$ . Consider for  $n'_u := 2^{-u}n$  and each  $0 \leq u \leq p$ , the i.i.d. variables*

$$\alpha_j^{(n'_u)} := \frac{1}{n} \sum_{i \in I_{2^{j-1}}^{(n'_u)}} \sum_{\ell \in I_{2^j}^{(n'_u)}} G(S_i, S_\ell), \quad 1 \leq j \leq 2^{u-1}.$$

Then, for some  $c_2 > 0$ ,

$$(4.74) \quad \sup_{n \in \mathbb{N}} E[e^{c_2 \bar{\Theta}_n}] < \infty, \quad \bar{\Theta}_n := \sum_{u=1}^p \sum_{j=1}^{2^{u-1}} \alpha_j^{(n'_u)}.$$

*Proof.* With the srw having independent and symmetric increments, one easily verifies that  $\alpha_1^{(n'_u)} \stackrel{(d)}{=} 2^{-u} X_{n'_u}$ , for  $X_n$  of Lemma 4.1. Consequently, from (4.5) and (4.6) we know that

$$(4.75) \quad \varphi(\lambda) := \sup_{u, n \in \mathbb{N}} E[\exp(\lambda 2^u \bar{\alpha}_1^{(n'_u)})] \leq 1 + c \lambda^2 < \infty,$$

for some  $c < \infty$  and all  $\lambda > 0$  small enough. The uniform MGF bound of (4.74) then follows as in [26, Page 177, Proof of Thm. 1], upon setting  $\alpha_0 = \alpha_1^{(n'_0)}$ ,  $c_2 = b_\infty > 0$  of [26], and noting that (4.75) suffices in lieu of both [26, Lemma 2] and the scale invariance of [26, property (ii)].  $\blacksquare$

*Proof of Lemma 4.7.* For  $u \leq p$ , consider the i.i.d. variables  $W_{u,j} := W^{(n'_u)} \circ \theta_{(2j-2)n'_u}$ , with

$$W^{(m)} := \sum_{i, \ell \in [1, m]} g_{m, \alpha}(i) G(S_i, S_{\ell+m}) g_{m, \alpha}(\ell) \circ \theta_m$$

having the law of  $W_1$  of (4.29) and their (i.i.d.) approximations  $\underline{W}_{u,j} := n\bar{g}_{n'_u,\alpha}^2 \alpha_j^{(n'_u)}$  (for  $g_{m,\alpha}(\cdot)$  and  $\bar{g}_{m,\alpha}$  of Lemma 4.3). Setting

$$Z_u := \sum_{j=1}^{2^{u-1}} W_{u,j}, \quad \underline{Z}_u := \sum_{j=1}^{2^{u-1}} \underline{W}_{u,j},$$

it follows by Cauchy-Schwarz and (4.26) that for some  $C < \infty$ , any  $u \leq p$  and all  $n$ ,

$$(4.76) \quad E[(Z_u - \underline{Z}_u)^2] \leq 2^{2(u-1)} E[(W_{u,1} - \underline{W}_{u,1})^2] \leq Cn^2(\log n)^{-\alpha/2}.$$

In particular, taking  $\alpha > 8 + 2c_1\lambda$  yields, by the union bound and Markov's inequality that for any  $\epsilon > 0$ ,

$$(4.77) \quad \begin{aligned} P\left(\sum_{u=1}^p (Z_u - \underline{Z}_u) \geq \epsilon \bar{h}_4(n)\right) &\leq \sum_{u=1}^p P(Z_u - \underline{Z}_u \geq \epsilon p^{-1} \bar{h}_4(n)) \\ &\leq Cp^3 (\epsilon \bar{h}_4(n))^{-2} n^2 (\log n)^{-\alpha/2} \leq C' \epsilon^{-2} (\log n)^{-c_1\lambda}. \end{aligned}$$

We also find for our choice of  $k$ , see (4.61), that

$$E\Delta_{n,k} = \frac{\pi^2}{8} \bar{h}_4(n) (1 + o(1)).$$

Similarly to (4.29), we have that  $V_{(2j-2)n'_u, (2j-1)n'_u, 2jn'_u} \leq 2W_{u,j}$  for any  $u, j$ . In view of (4.71) and (4.77) it thus suffices to establish (4.73) with  $\Delta_{n,k}$  replaced by

$$2 \sum_{u=1}^p \underline{Z}_u - \frac{\pi^2}{8} \bar{h}_4(n),$$

which in view of (4.40) and the definition of  $\Theta_n$ , can be further replaced by

$$\frac{\pi^2}{8} \frac{n}{(\log n)^2} \left( \left( \frac{\pi^2}{4} + o(1) \right) \Theta_n - \log_2 n \right).$$

Moreover, by (4.76),

$$\left| E \sum_{u=1}^p Z_u - E \sum_{u=1}^p \underline{Z}_u \right| \leq O\left(\frac{pn}{(\log n)^{\alpha/4}}\right) = o(\bar{h}_4(n)),$$

and combining [6, Lemma 6.5] with the considerations as in (4.44) and after (4.54), we deduce that for large  $m$ ,

$$2E[W^{(m)}] = (1 + o(1))E[V_{0,m,2m}].$$

It then follows that

$$2 \sum_{u=1}^p EZ_u = (1 + o(1)) \frac{\pi^2}{8} \bar{h}_4(n)$$

and consequently,

$$\frac{\pi^2}{4} E\Theta_n = (1 + o(1)) \log_2 n.$$

Thus it suffices to establish (4.73) with  $\bar{\Delta}_{n,k}$  replaced by

$$\frac{\pi^4}{32} \frac{n}{(\log n)^2} \bar{\Theta}_n,$$

which in turns follows from (4.74), upon setting  $c_1 = 32\pi^{-4}c_2 > 0$ .  $\blacksquare$

### 5. LIL FOR SRW ON $\mathbb{Z}^d$ , $d \geq 5$ : PROOF OF THEOREM 1.4

Since  $R_n$  for the SRW on  $\mathbb{Z}^d$ ,  $d \geq 5$ , has similar structural properties to the size of the range of the SRW on  $\mathbb{Z}^{d-2}$ , we establish Theorem 1.4 by adapting the proof in [11] for the LIL of the latter sequence. Specifically, setting  $\rho_n := \sqrt{n \log n}$  when  $d = 5$  and otherwise  $\rho_n := \sqrt{n}$ , we have from [32, Thm. A] in case  $d = 5$ , and from [4, Lemma 3.3] when  $d \geq 6$ , that

$$(5.1) \quad \lim_{n \rightarrow \infty} \frac{1}{\rho_n} \|\bar{R}_n\|_2 = \sigma_d.$$

We proceed with the following variant of [4, Lemma 3.2].

**Lemma 5.1.** *Recall the notations of (4.1) and for any  $0 \leq a < b$ , set*

$$\tilde{V}_{a,b} := \sup_{t \geq b} \{V_{a,b,t}\}, \quad \hat{V}_{a,b} := \sup_{s \leq a} \{V_{s,a,b}\}.$$

Then, for some  $C_{d,\ell}$  finite, any  $\ell \geq 1$  and all  $a < b$ ,

$$(5.2) \quad E[\tilde{V}_{a,b}^\ell] \leq C_{d,\ell} f_d(b-a)^\ell, \quad E[\hat{V}_{a,b}^\ell] \leq C_{d,\ell} f_d(b-a)^\ell,$$

where

$$f_5(n) = \sqrt{n}, \quad f_6(n) = \log n, \quad f_d(n) = 1 \quad \forall d \geq 7.$$

*Proof.* By the shift invariance of the SRW, we may wlog set  $a = 0$ . Further, in view of [6, (2.9) & (2.11)], for a fixed set  $A$ , the function

$$B \mapsto \text{Cap}(A) + \text{Cap}(B) - \text{Cap}(A \cup B)$$

is non-decreasing (and bounded above by  $\text{Cap}(A)$ ). In particular, the value of  $\tilde{V}_{0,n}$  is attained for  $t \rightarrow \infty$ . Thus, from [4, Prop. 1.2] we arrive at

$$(5.3) \quad \tilde{V}_{0,n} \leq 2 \sum_{x \in \mathcal{R}_n} \sum_{y \in \mathcal{R}(n,\infty)} G(x,y) \stackrel{d}{=} 2 \sum_{x \in \mathcal{R}_n} \sum_{y \in \hat{\mathcal{R}}_\infty} G(x,y),$$

where  $\hat{\mathcal{R}}_\infty$ , denotes the range of an independent SRW. Similarly, the value of  $\hat{V}_{0,n}$  is attained at  $s \rightarrow -\infty$ , with the right-side of (5.3) also bounding  $\hat{V}_{0,n}$  (we then have  $\mathcal{R}(-\infty, 0]$  instead of  $\mathcal{R}(n, \infty)$  in (5.3)). Thereafter, adapting the proofs of [4, Lemma 3.1 & 3.2] yields (5.2). Indeed, with  $p_{2k}(x, y) := P^x(S_{2k} = y)$  the square of a transition probability, we have as in the proof of [4, Lemma 3.1], that for even  $k \geq 0$  and any  $a \in \mathbb{Z}^d$ ,

$$s_a := \sum_{x,y \in \mathbb{Z}^d} G(0,x)G(0,y)p_k(x,y+a) \leq s_0.$$

In case of a lazy srw, this applies for any  $k \geq 0$ , so summing over  $k \leq n$  yields that

$$(5.4) \quad \max_{a \in \mathbb{Z}^d} \left\{ \sum_{x, y \in \mathbb{Z}^d} G(0, x) G(0, y) G_n(x, y + a) \right\} = \sum_{x, y \in \mathbb{Z}^d} G(0, x) G(0, y) G_n(x, y) \\ = \sum_{x, y \in \mathbb{Z}^d} G_n(0, x) G(0, y) G(x, y) \leq C_d f_d(n),$$

where we have utilized [4, (3.4)] for the latter inequality. Further, as in [4], up to an increase of  $C_d$  value, (5.4) extends to the original srw. Now, similarly to [4, (3.5)], it follows from (5.3) that

$$E[\tilde{V}_{0,n}^\ell] \leq 2^\ell \sum_{\underline{x}, \underline{y}} E \left[ \prod_{i=1}^{\ell} L_n(x_i) \right] E \left[ \prod_{i=1}^{\ell} L_\infty(y_i) \right] \prod_{i=1}^{\ell} G(x_i, y_i),$$

where  $L_n(x)$  denotes the total srw local time at  $x \in \mathbb{Z}^d$ , during time interval  $[1, n]$  (and the same bound applies for  $E[\hat{V}_{0,n}^\ell]$ ). For  $\ell = 1$  we thus get (5.2) out of (5.4) (as  $E[L_n(x)] = G_n(0, x)$  and  $E[L_\infty(y)] = G(0, y)$ ). The general case then follows by an inductive argument, as in the proof of [4, Lemma 3.2], utilizing also that  $a = 0$  is optimal in (5.4). ■

Utilizing Lemma 5.1, we next establish the analog of [11, Lemma 3.3] for  $R_n$ .

**Lemma 5.2.** *For any  $d \geq 5$ ,  $m \geq 3$ , there exists  $c_m$  finite, such that for all  $b > a \geq 0$*

$$(5.5) \quad \|\bar{R}_b - \bar{R}_a\|_m \leq c_m \rho_{b-a}.$$

Further, for some  $\bar{c}_m$  finite and any  $\lambda > 0$ ,  $b > a \geq 0$ ,

$$(5.6) \quad P(\max_{n \in [a, b]} \{\|\bar{R}_n - \bar{R}_a\|\} > \lambda \rho_{b-a}) \leq \bar{c}_m \lambda^{-m}.$$

*Proof.* Note that

$$V_{0,a,b} = R_a + R_{a,b} - R_b \in [0, \hat{V}_{a,b}],$$

for  $\hat{V}_{a,b}$  of Lemma 5.1. In particular, for any  $m \geq 3$ ,  $b > a$ ,

$$0 \leq E[V_{0,a,b}]^m \leq E[V_{0,a,b}^m] \leq E[\hat{V}_{a,b}^m] \leq C_{d,m} f_d(b-a)^m = o(\rho_{b-a}^m).$$

We can thus replace  $R_b - R_a$  in (5.5) by  $R_{a,b}$  and thereby due to the shift invariance of the increments, set wlog  $a = 0$  (whereupon  $R_a = 0$ ). Hence, analogously to [11, (3.34)], it suffices for (5.5) to show inductively over  $\ell \geq 1$ , that  $\sup_n \{L_{n,2\ell}\}$  is finite for

$$L_{n,\ell} := \frac{1}{\rho_n} \|\bar{R}_n\|_\ell.$$

The induction basis  $\ell = 1$  is merely (5.1). Further, with

$$\lim_{n \rightarrow \infty} \rho_{2n}^{-1} \sup_{a < 2n} \|\bar{V}_{0,a,2n}\|_{2\ell} = 0,$$

by the preceding decomposition, we can and shall replace  $R_{2n}$  in the induction step, by

$$R_n + R_{n,2n} \stackrel{d}{=} R_n + \hat{R}_n,$$

where  $\hat{R}_n$  denotes the capacity of the range of an independent second srw. For any  $\ell \geq 2$ , by the induction hypothesis  $\sup_n \{L_{n,k}\}$  are finite for all  $k \leq 2(\ell - 1)$ , hence

$$\sup_{n,n'} \sum_{k=2}^{2\ell-2} \binom{2\ell}{k} L_{n,k}^k L_{n',2\ell-k}^{2\ell-k} := c_\ell < \infty.$$

Recalling that  $\rho_n \leq 2^{-1/2} \rho_{2n}$ , we thus get similarly to [11, (3.37) & (3.38)] that

$$L_{2n,2\ell} \leq o_n(1) + \left(2^{-(\ell-1)} L_{n,2\ell}^{2\ell} + 2^{-\ell} c_\ell\right)^{1/(2\ell)},$$

from which it follows as in [11] that  $\sup_j \{L_{2^j,2\ell}\}$  is finite. Finally, for any  $n \in [2^{j-1}, 2^j)$ ,  $j \geq 2$ , we have as in the preceding that

$$R_{2^j} \stackrel{d}{=} R_n + \hat{R}_{2^j-n} - V_{0,n,2^j}.$$

Upon centering, taking the  $2\ell$ -th power and isolating the  $2\ell$ -th power of  $\bar{R}_n$ , the preceding identity results with

$$[L_{2^j,2\ell} + o_j(1)]^{2\ell} + c_\ell \geq (\rho_n/\rho_{2^j})^{2\ell} L_{n,2\ell}^{2\ell} \geq 4^{-\ell} L_{n,2\ell}^{2\ell}.$$

Thus,  $\sup_n \{L_{n,2\ell}\}$  is finite as well, completing the induction step and thereby establishing (5.5). Finally, we get (5.6) out of (5.5) precisely as in deriving [11, (3.39)] out of [11, (3.40)]. ■

Centering both sides of (4.2) we arrive at

$$(5.7) \quad \bar{R}_{n_k} = \sum_{j=1}^k \bar{U}_j - \bar{\Delta}_{n_k,k},$$

with  $\Delta_{n_k,k}$  of (4.3) and the zero-mean, independent variables  $\bar{U}_j := \bar{R}_{n_{j-1},n_j}$ . Proceeding to show that  $\bar{\Delta}_{n_k,k}$  has a negligible effect on  $\bar{R}_{n_k}$ , first recall from (5.1) that

$$(5.8) \quad \lim_{j \rightarrow \infty} \frac{E[\bar{U}_j^2]}{\rho_{n_j-n_{j-1}}^2} = \sigma_d^2,$$

whereas (5.5) at  $a = n_{j-1}$ ,  $b = n_j$ , amounts to

$$(5.9) \quad E[|\bar{U}_j|^m] \leq (c_m \rho_{n_j-n_{j-1}})^m.$$

In case  $d = 5$  we take the same values of  $\alpha, \beta < 1/2$  and  $\{n_k\}$  as in the proof of [11, Thm. 2.1]. Lemma 5.1 at  $\ell = 4$  is then the analog of [11, (3.2)], and utilizing it at  $a = n_{j-1}$ ,  $b = n_j$ ,  $j < k$ , we find by following verbatim, the derivation of [11, (3.9)], that for some  $c$  finite

$$(5.10) \quad \limsup_{k \rightarrow \infty} \frac{|\bar{\Delta}_{n_k,k}|}{\sqrt{n_k} (\log n_k)^\beta} \leq c \quad \text{a.s.}$$

Thereafter, substituting (5.10) for [11, (3.9)] and (5.9) to get [11, (3.16)], by the same reasoning as in the proof of [11, Thm. 2.1], we find that a.s.

$$(5.11) \quad \lim_{k \rightarrow \infty} h_d(n_k)^{-1} [\bar{R}_{n_k} - \sigma_d B_{\rho_{n_k}^2}] = 0,$$

for some one-dimensional standard Brownian motion  $(B_t, t \geq 0)$ . As shown after [11, (3.17)] (apart from replacing [11, Lemma 3.3(b)] by (5.6)), the stated LIL is then a direct consequence of Kinchin's LIL for the latter Brownian motion.

In case  $d \geq 6$ , we take  $\{n_k\}$  again as in the proof of [11, Thm. 2.1], except that now this is done for the choice of  $\alpha = 1$ . Then, by Lemma 5.1, for  $C = C_{d,2}$  and any  $1 \leq j < k$ ,

$$\text{Var}(V_{n_{j-1}, n_j, n_k}) \leq C(\log(n_j - n_{j-1}))^2 \leq C(\log n_k)^2.$$

Consequently, for any  $\beta > 0$  and all  $k$ , by Markov's inequality and (4.3),

$$P(|\overline{\Delta}_{n_k, k}| \geq n_k^\beta) \leq n_k^{-2\beta} \text{Var}(\Delta_{n_k, k}) \leq C n_k^{-2\beta} k^2 (\log n_k)^2.$$

Since  $|\{n_k\} \cap [2^\ell, 2^{\ell+1})| \leq \ell$  for any  $\ell \geq 1$ , eventually  $k \leq (\log n_k)^2$ . Hence, by the first Borel-Cantelli lemma, we have that for any  $\beta > 0$ ,

$$(5.12) \quad \limsup_{k \rightarrow \infty} n_k^{-\beta} |\overline{\Delta}_{n_k, k}| \leq 1 \quad \text{a.s.}$$

We then get (5.11) by following, as for  $d = 5$ , the proof of [11, Thm. 2.1], utilizing again (5.8)-(5.9), while having now, via (5.12) at  $\beta < 1/2$ , a negligible contribution at scale  $\rho_n = \sqrt{n}$  (instead of (5.10) and the scale  $\sqrt{n \log n}$  throughout [11, (3.13)-(3.17)]). Finally, recall that  $n_{k+1} - n_k \leq n_k/\ell$  whenever  $n_k \in [2^\ell, 2^{\ell+1})$ . Hence, in view of (5.6) at  $m = 6$  and  $\lambda = \varepsilon h_d(n_k) / \sqrt{n_{k+1} - n_k}$ , we have that for some  $c_\varepsilon$  finite, any  $\varepsilon > 0$  and  $n_k \in [2^\ell, 2^{\ell+1})$ ,

$$P\left(\max_{n \in (n_k, n_{k+1})} \{|\overline{R}_n - \overline{R}_{n_k}|\} > \varepsilon h_d(n_k)\right) \leq c_\varepsilon \ell^{-3}.$$

With at most  $\ell$  values of such  $n_k$ , by the first Borel-Cantelli lemma, the events on the LHS a.s. occur only for finitely many values of  $k$  and the stated LIL thus follows, as before, from (5.11).

## REFERENCE

- [1] Asselah A. and Schapira B. Deviations for the capacity of the range of a random walk. *Electron. J. Probab.* 25 (2020), Paper No.154, 28 pp.
- [2] Asselah A. and Schapira B. On the nature of the Swiss cheese in dimension 3. *Ann. Probab.* 48 (2020), no. 2, 1002-1013.
- [3] Asselah A. and Schapira B. The two regimes of moderate deviations for the range of a transient walk. *Probab. Th. Rel. Fields* 180 (2021), no. 1-2, 439-465.
- [4] Asselah A., Schapira B. and Sousi P. Capacity of the range of random walk on  $\mathbb{Z}^d$ . *Trans. Amer. Math. Soc.* 370 (2018), no. 11, 7627-7645.
- [5] Asselah A., Schapira B. and Sousi P. A strong law of large numbers for the capacity of the Wiener sausage in dimension four. *Probab. Th. Rel. Fields* 173 (2019), 813-858.
- [6] Asselah A., Schapira B. and Sousi P. Capacity of the range of random walk on  $\mathbb{Z}^4$ . *Ann. Probab.* 47 (2019), 1447-1497.
- [7] Bai T. and Hu Y. Capacity of the range of branching random walks in low dimensions. Preprint, Arxiv:2104.11898.
- [8] Bai T. and Hu Y. Convergence in law for the capacity of the range of a critical branching random walk. Preprint, Arxiv:2203.03188.
- [9] Bai T. and Wan Y. Capacity of the range of tree-indexed random walk. *Ann. Appl. Probab.* 32 (2022), no. 3, 1557-1589.
- [10] Bass R. F., Chen, X. and Rosen J. Moderate deviations for the range of planar random walks. *Mem. Amer. Math. Soc.* 198 (2009), no. 929, viii+82 pp
- [11] Bass R. F. and Kumagai T. Laws of the iterated logarithm for the range of random walks in two and three dimensions. *Ann. Probab.* 30 (2002), no. 3, 1369-1396.
- [12] van den Berg M., Bolthausen E. and den Hollander F. Moderate deviations for the volume of the Wiener sausage. *Ann. Math. (2)* 153 (2001), no. 2, 355-406.
- [13] Chang Y. Two observations on the capacity of the range of simple random walk on  $\mathbb{Z}^3$  and  $\mathbb{Z}^4$ . *Electron. Com. Probab.* 22 (2017), No. 25, 9 pp.

- [14] Chung K. L. On the maximum partial sums of sequences of independent random variables. *Trans. Amer. Math. Soc.* 64 (1948), 205-233.
- [15] Csáki, E. and Hu, Y. Strong approximations of three-dimensional Wiener sausages. *Acta Math. Hungar.* 114 (2007), no. 3, 205-226.
- [16] Cygan W., Sandrić, N. and Sebek, S. Central limit theorem for the capacity of the range of stable random walks. *Stochastics* 94 (2022), no. 2, 226-247.
- [17] Gao F. Q. and Wang Y. Q. Laws of the iterated logarithm for high-dimensional Wiener sausage. *Acta Math. Sin. (Engl. Ser.)* 27 (2011), no. 8, 1599-1610.
- [18] Hariya Y. and Hasegawa K. On divergence of expectations of the Feynman-Kac type with singular potentials. *J. Math. Soc. Japan* 68 (2016), no. 3, 1271-1296.
- [19] Hutchcroft T. and Sousi P. Logarithmic corrections to scaling in the four-dimensional uniform spanning tree. Preprint, Arxiv:2010.15830.
- [20] Jain N. C. and Orey S. On the range of random walk. *Israel J. Math.* 6 (1968), 373-380.
- [21] Jain N. C. and Pruitt W. E. The law of the iterated logarithm for the range of random walk. *Ann. Math. Stat.* 43 (1972), 1692-1697.
- [22] Kent J. T. Eigenvalue expansions for diffusion hitting times. *Z. Wahrsch. Verw. Gebiete*, 52 (1980), 309-319.
- [23] Lawler G. F. *Intersections of random walks*. Second edition, Birkhauser, 1996.
- [24] Lawler G. F. and Limic V. *Random walk: a modern introduction*. Cambridge University Press, 2010.
- [25] Le Gall J.-F. Fluctuation results for the Wiener sausage. *Ann. Probab.* 16 (1988), no. 3, 991-1018.
- [26] Le Gall J.-F. Exponential moments for the renormalized self-intersection local time of planar Brownian motion. *Seminaire de Probabilities, XXVIII*, 172-180, *Lecture Notes in Math.*, 1583, Springer, Berlin, 1994.
- [27] Mörters P. and Peres Y. *Brownian motion*. Cambridge University Press, 2010.
- [28] Mrazović R., Sandrić N. and Sebek S. Capacity of the range of random walks on groups. Arxiv:2103.04588, accepted for publication in *Kyoto J. Math.*
- [29] Okada I. Geometric structures of late points of a two-dimensional simple random walk. *Ann. Probab.* 47 (2019), no. 5, 2869-2893
- [30] Petrov V. V. *Sums of independent random variables*. Springer-Verlag, Berlin, 1975.
- [31] Pólya G. and Szegő G. *Isoperimetric inequalities in mathematical physics*. *Annals of Mathematics Studies*, no. 27, Princeton University Press, Princeton, N.J., 1951.
- [32] Schapira B. Capacity of the range in dimension 5. *Ann. Probab.* 48 (2020), 2988-3040.
- [33] Sznitman A.-S. Vacant set of random interacements and percolation. *Ann. Math. (2)* 171 (2010), no. 3, 2039-2087.
- [34] Watson G. N. *A Treatise on the theory of Bessel functions*. Cambridge University Press, 1944.

AMIR DEMBO

MATHEMATICS DEPARTMENT AND STATISTICS DEPARTMENT, STANFORD UNIVERSITY, STANFORD, CA 94305, USA.

*Email address:* adembo@stanford.edu

IZUMI OKADA

FACULTY OF MATHEMATICS, KYUSHU UNIVERSITY, FUKUOKA 819-0395, JAPAN.

*Email address:* iokada@math.kyushu-u.ac.jp