

ORDERED SURPRISES AND CONDITIONAL PROBABILITY SYSTEMS*

Adam Dominiak Matthew Kovach Gerelt Tserenjigmid

First version: April 4, 2021; This version: August 9, 2022

Abstract

We study conditioning on null events, or *surprises*, and behaviorally characterize the Ordered Surprises (OS) representation of beliefs. For feasible events, our Decision Maker (DM) is Bayesian. For null events, our DM considers a hierarchy of beliefs until one is consistent with the surprise. The DM adopts this prior and applies Bayes' rule. Unlike Bayesian updating, OS is a complete updating rule: conditional beliefs are well-defined for any event. OS is (behaviorally) equivalent to the Conditional Probability System (Myerson, 1986b) and is a special case of Hypothesis Testing (Ortoleva, 2012), clarifying the relationships between the various approaches to null events.

Keywords: Uncertainty, subjective expected utility, null events, conditional probabilities system, Bayesian updating, consequentialism, dynamic consistency.

JEL-Classifications: D01, D80, D81, D83.

*Dominiak: Virginia Tech (dominiak@vt.edu); Kovach: Virginia Tech (mkovach@vt.edu); Tserenjigmid: UC Santa Cruz (gtserenj@ucsc.edu). We are very grateful to Shachar Kariv, Burkhard Schipper, and Marie-Louise Vierø for many stimulating discussions and suggestions that improved the exposition of the paper. All mistakes are our own.

1 Introduction

Decision problems under uncertainty often feature a dynamic structure. A decision maker (DM) may acquire information about the uncertainty she faces. Given this information, the DM must formulate a conditional belief that governs her conditional behavior. Bayesian subjective expected utility is the dominant theory in economics for dynamic choice problems. However, the Bayesian paradigm is incomplete; it is silent about updating on (Savage) null events (i.e., zero probability events). Our goal is to introduce a theory of conditional beliefs that are well defined for all events, thereby establishing a complete theory of updating.

We start by defining an updating rule, which is a mapping from events to conditional beliefs, and introduce two simple properties. An updating rule is *complete* if it is has full domain (it is defined for every event), and it is *concentrated* if, for each event, no states outside the event have positive probability. Complete updating rules are important, for example, because any perfect Bayesian equilibrium, including its refinements, requires one to describe equilibrium beliefs.

The most prominent complete updating rule is Myerson's Conditional Probability System (CPS)¹ (Myerson, 1986a,b), which was motivated by the sequential equilibrium of Kreps and Wilson (1982) (henceforth, KW).² KW defined a sequential equilibrium as an assessment (σ, μ) (a strategy profile σ and a system of beliefs μ) that is "consistent" and "sequentially rational." Consistency means that there is a sequence of full support beliefs converging to σ such that the corresponding sequence of belief systems (Bayesian updates) converges to μ . Sequential rationality requires that at each information set the actions prescribed by σ maximize the conditional expected payoff given (σ, μ) . Importantly, these conditions ensure that players' beliefs about play are *well-defined at every information set*, even for information sets off the equilibrium path (e.g., null events); each player has a complete theory of belief updating. Myerson (1986b) shows that consistency of beliefs in sequential equilibria is equivalent to CPS.³

As an interpretation for how beliefs evolve in a sequential equilibrium, KW suggested

¹The idea of CPS goes back to Rényi (1955).

²Sequential equilibrium is one of the most widely applied solution concepts in dynamic games. For example, it has been used to study bargaining (Rubinstein, 1985), cheap talk (Crawford and Sobel, 1982), job search (Spence, 1973), advertising (Milgrom and Roberts, 1986), dividends (John and Williams, 1985), product quality (Miller and Plott, 1985), warranties (Gal-Or, 1989), limit pricing (Milgrom and Roberts, 1982), social norms (Bernheim, 1994), lobbying (Lohmann, 1995), and many other topics.

³See also McLennan (1989) and Battigalli (1996) for a connection between CPS and sequential equilibria.

that upon reaching an information set that is incompatible with a player’s beliefs about play in the game, the player should move to their “second most likely hypothesis” and, if that fails, then they should implement their “third most likely hypothesis,” and so on. KW show that a sufficient condition for an assessment to satisfy this property is the existence of a convergent sequence of probability measures that generate the assessment.

We revisit the “ordered hierarchy of hypotheses” of KW and introduce the notion of **Ordered Surprises** (OS). OS may be viewed as the minimal extension of Bayesian theory that allows for well-defined beliefs after null events. An OS representation of a complete updating rule is given by a collection of beliefs $\mathcal{O} = \langle \mu_0, \dots, \mu_K \rangle$, where μ_0 is the initial belief and each $k > 0$ is used to form some conditional belief after some null event. Each index indicates a distribution’s order in the hierarchy.

Our main result shows that OS is equivalent to CPS ([Theorem 1](#)); an updating rule has an OS representation if and only if it is a CPS. While one direction of this equivalence was suggested by KW, to the best of our knowledge, we are the first to formally prove this equivalence (CPS must have an ordered structure). Additionally, the explicit construction of the sequential beliefs for any CPS through the OS representation is useful in practice for those who want to find all sequential equilibria in extensive-form games.

We provide simple behavioral foundations for OS, thereby establishing a simpler characterization for CPS. Our characterization ([Theorem 4](#)) relies upon a novel axiom, **Conditional Consistency**, that implies Dynamic Consistency among the ex-ante feasible events and extends this consistency to the entire hierarchy of beliefs; it imposes Dynamic Consistency within each μ_k .

To illustrate OS, imagine a coin flip. The states h and t are the usual outcomes of heads or tails, e and e' denote edges where e' has been worn thin, while l_1 and l_2 denote landing on a marked location, which yields the state space $S = \{h, t, e, e', l_1, l_2\}$. Initially, the DM has belief $\mu_0(h) = \mu_0(t) = \frac{1}{2}$, and treats the other states as null.

Suppose the DM is informed that, astonishingly, the coin did not land on a face; $A = \{e, e', l_1, l_2\}$ was realized. While our DM now has sufficient evidence to rule out her initial beliefs, it is not clear which belief she should adopt. Suppose our DM believes that the coin landing on either of the marked locations is *more impossible* than its landing on an edge. Accordingly, her conditional beliefs are $\mu_A(\{e, e'\}) = 1$ and $\mu_A(\{l_1, l_2\}) = 0$. Since the states in $\{e, e'\}$ are no longer null, they constitute a *first-order surprise*. In contrast, $\{l_1, l_2\}$, remain null and constitute a higher-order surprise, where “order” designates the point at which a state is considered possible. Our notion of OS extends this idea to up to

K orders.

We compare OS to other approaches to conditioning on null events. In [Theorem 2](#), we prove directly that OS preferences are a special case of the Hypothesis Testing Model of [Ortoleva \(2012\)](#), and thus so is CPS. In particular, for any OS given by $\langle \mu_0, \dots, \mu_K \rangle$, we construct a second-order prior over these $K + 1$ beliefs that ensures selection in the resulting HT is consistent with the OS structure. This argument cannot be reversed, and so HT is strictly more general than OS.

CPS and OS representations complete Bayes' rule by disciplining belief updating on zero-probability events. Motivated by empirical evidence on belief updating (e.g., see [Benjamin \(2019\)](#)), we propose a one-parameter non-Bayesian extension of the OS representations based on the idea of HT. We show that this non-Bayesian extension of OS representations is still a special case of HT, confirming that it is a reasonable non-Bayesian extension of CPS. This extension might be useful for defining a non-Bayesian extension of sequential equilibrium.

1.1 Related Literature

There have been various attempts to deal with conditioning on zero-probability events. In economics, the three main approaches are (i) the conditional probability system (CPS) of [Myerson \(1986a,b\)](#), (ii) the (conditional) lexicographic probability systems (C)LPS of [Blume et al. \(1991\)](#), and (iii) the hypothesis testing model (HT) of [Ortoleva \(2012\)](#). These are carefully discussed in [subsection 2.3](#), [section 4](#), and [subsection 2.4](#), respectively.

It is worth remarking that the (conditional) LPS is fundamentally distinct from CPS and HT (hence also OS). First, the CLPS rules out null events. Second, (C)LPS behavior is incompatible with SEU, while the other three are SEU models.

There is an extensive literature that applies CPSs in dynamic games with incomplete information. [Battigalli and Siniscalchi \(1999\)](#) use CPSs to describe an (epistemic) type of a player (i.e., a complete and explicit description of players' hierarchies of conditional beliefs) and show that a universal type space for CPSs always exists. In this framework, [Battigalli and Siniscalchi \(2002\)](#) provide an epistemic characterization of extensive-form rationalizability and backward induction. Recently, [Galperti \(2019\)](#) studies Bayesian persuasion using a CPS (OS with two beliefs).⁴

Finally, there is an abundant literature on (non-Bayesian) updating. Recently, [Do-](#)

⁴[Tsakas \(2018\)](#) derives the famous Agreement Theorem under a CPS.

miniak, Kovach, and Tserenjigmid (2022) show that HT admits a “minimum-distance” representation; the conditional probability μ_E that the DM selects is the element of $\Delta(E)$ that is closest to her prior μ . Hence, CPS also admits a minimum-distance representation.

2 Model

2.1 Setup

We study dynamic choice in the formal framework of Anscombe and Aumann (1963). Uncertainty is described by a nonempty and finite set of states $S = \{s, \dots, s\}$. We denote by $\Delta(S)$ the set of all probability distributions on S . Let X be a nonempty set of outcomes and $\Delta(X) := \left\{ p : X \rightarrow [0, 1] \mid \sum_{x \in \text{supp}(p)} p(x) = 1 \text{ with } |\text{supp}(p)| \in \mathbb{N} \right\}$ be the set of all (simple) lotteries over X , where $\text{supp}(p)$ denotes the support of p .

Objects of choice are (Anscombe-Aumann) acts $f : S \rightarrow \Delta(X)$ that map states to lotteries. Let \mathcal{F} denote the set of all such acts. A constant act is an act that assigns the same lottery to all states: $f(s) = p$ for all $s \in S$. Using a standard abuse of notation, we denote by $p \in \mathcal{F}$ the corresponding constant act. Thus, we can identify the set of lotteries with the constant acts. A subset $E \subseteq S$ is called an event. Denote by Σ the algebra of events generated by S . For an event E and acts f, g we denote by $f_E g$ a composite act that returns $f(s)$ for $s \in E$ and $g(s)$ otherwise.

A preference relation over \mathcal{F} , denoted by \succsim , describes a DM’s behavior. As usual, \succ and \sim are the asymmetric and symmetric parts of \succsim , respectively. For each event $E \in \Sigma$, \succsim_E denotes the conditional preference over \mathcal{F} given E . It governs the DM’s choice upon learning E . The initial preference relation (before information is revealed) is denoted by \succsim_S . The DM’s behavior is described by a family of conditional preferences $\{\succsim_E\}_{E \in \Sigma}$; one for each $E \in \Sigma$.

Given a belief $\mu \in \Delta(S)$ and an event $E \in \Sigma$ such that $\mu(E) > 0$, we let $BU(\mu, E)(A)$ denote the Bayesian update of μ given E , where $BU(\mu, E)(A) := \frac{\mu(E \cap A)}{\mu(E)}$.

Definition 1. An **updating rule** is a pair (P, \mathcal{D}) where $\mathcal{D} \subseteq \Sigma$ and P is a mapping $P : \mathcal{D} \rightarrow \Delta(S)$.

- (i) An updating rule is **complete** if it has full domain: $\mathcal{D} = \Sigma$.
- (ii) An updating rule is **concentrated** if for any $E \in \mathcal{D}$, $P(E|E) = 1$.

In other words, P is complete if the conditional probability $P(\cdot|E)$ is well-defined for each $E \in \Sigma$.

Example 1. *Since BU is only well-defined for $\mathcal{D} = \{E \in \Sigma \mid \mu(E) > 0\} \subset \Sigma$ where μ is the prior, Bayesian updating is incomplete.*

Additionally, an LPS that admits null events (e.g., the LPS is not full support), results in an incomplete updating rule.⁵

Example 2. *Let $\mu \in \Delta(S)$, $\delta \in (0, 1]$ and let P^C denote “Conservative BU,” where*

$$P^C(\cdot|E) = \begin{cases} \delta\mu + (1 - \delta)BU(\mu, E) & \text{if } \mu(E) > 0 \\ \delta\mu + (1 - \delta)\frac{1}{|E|} & \text{if } \mu(E) = 0. \end{cases}$$

The updating rule P^C , a one-parameter extension of CPS, is complete but not concentrated (the resulting preferences violate consequentialism).⁶ When $\delta = 1$, we have “stubborn” agents who never update their beliefs (similar behavior appears in Acemoglu et al. (2013)).

2.2 Ordered Surprises

In this section, we introduce OS, which captures the DM’s subjective levels of relative (im)possibility. Intuitively, an event is null if the DM assigns it zero probability. In behavioral terms, an event E is *null* if a DM is indifferent between any two acts that only differ on E .

Definition 2. *For any $A, E \in \Sigma$, an event A is \succsim_E -null if for all $f, g \in \mathcal{F}$,*

$$f_Ag \sim_E g. \tag{1}$$

Otherwise, A is \succsim_E -feasible. If A is \succsim_S -null, then it is (ex-ante) null.

We can distinguish between types of null events by comparing the ex-ante preference \succsim_S with conditional preferences \succsim_E for various events. For example, consider the initial prior μ_0 over S . We can partition S into its feasible states F_0 and null states $S \setminus F_0$. When

⁵Technically, beliefs for a conditional LPS require an expanded notion of an updating rule. See [section 4](#) for further discussion.

⁶Similar updating rules have been studied in Epstein (2006) and Kovach (2020).

it is revealed that the true state is in fact an element of $S \setminus F_0$ — the DM’s (ex-ante) feasible states have been ruled out — she formulates new beliefs: $\mu_{S \setminus F_0}$.

Some states in $S \setminus F_0$ will have positive probability, while others remain null because they are *more impossible*. The states that are now *conditionally non-null* are “first-order surprises,” while the states that remain null are “higher-order surprises.” We can proceed in this fashion, sequentially revealing each surprise order, until all states have been considered. Following this idea, we formally define our notion of Ordered Surprises for updating rules.

Definition 3. *A complete updating rule P has an **Ordered Surprises** (OS) representation if there are probability distributions $\mu_0, \dots, \mu_K \in \Delta(S)$ such that*

$$P(\cdot|E) = BU(\mu_{k^*}, E) \text{ where } k^* = \min\{k \leq K \mid \mu_k(E) > 0\},$$

for every $E \in \Sigma$. For simplicity, we may refer to an OS representation as $\mathcal{O} = \langle \mu_0, \dots, \mu_K \rangle$.

It is without loss of generality to assume that μ_0, \dots, μ_K have disjoint supports. The indices designate the order in which each prior is considered. The DM begins with her prior over S , designated by μ_0 . She is Bayesian whenever possible, and abandons μ_0 only when necessary. After she observes a null event, she considers each μ_i in order, stops at the first belief that is consistent with the information, and applies Bayes’ rule. The following example illustrates this process.

Example 3 (Coin Flip). *Consider the coin flip example from the introduction. The states are $S = \{h, t, e, e', l_1, l_2\}$, where h and t correspond to heads or tails, e and e' correspond to the coin landing on an edge, where one edge is thinner than the other, while l_1 and l_2 correspond to the coin landing on precisely marked locations. These possibilities are described by the probability distributions*

$$\mu_0(s) = \begin{cases} \frac{1}{2} & s \in \{h, t\} \\ 0 & \text{otherwise} \end{cases}; \mu_1(s) = \begin{cases} \frac{7}{8} & s = e \\ \frac{1}{8} & s = e' \\ 0 & \text{otherwise} \end{cases}; \mu_2(s) = \begin{cases} \frac{1}{2} & s \in \{l_1, l_2\} \\ 0 & \text{otherwise} \end{cases}.$$

Our DM has initial prior μ_0 (e.g., \succsim_S has an SEU representation (u, μ_0)). Suppose she observes $A = \{e', l_1, l_2\}$. Since $\mu_0(A) = 0$, Bayesian updating is not defined. When our DM admits an Ordered Surprises representation generated by $\mathcal{O} = \langle \mu_0, \mu_1, \mu_2 \rangle$, she selects a new

belief that is consistent with A . In this case,

$$P^{OS}(\cdot|A) = BU(\mu_1, A).$$

Note that A intersects two “surprise orders”: $\{e, e'\} \cap A \neq \emptyset \neq \{l_1, l_2\} \cap A$. Under the OS model, the DM selects μ_1 because it is of lower order than μ_2 and therefore takes precedence.

2.3 Conditional Probability System

Perhaps the most well-known method for handling choice conditional on (ex-ante) null-events is the conditional probability system introduced by Myerson (1986b). In this section, we show that the OS representation is equivalent to CPS.

Myerson (1986b) initially defined a conditional probability system as a collection of conditional probabilities $(p(\cdot|E))_{E \in \Sigma}$, each one on S , that jointly satisfy certain properties. We provide an equivalent definition using our semantic of complete updating rules.

Definition 4. A complete updating rule P is a conditional probability system (CPS) if it is concentrated and for all $G \subseteq F \subseteq E$ where $F \neq \emptyset$, it satisfies

$$P(G|E) = P(G|F) \times P(F|E). \quad (2)$$

When $P(F|E) \neq 0$, (2) is equivalent to Baye’s rule.

Theorem 1. A complete updating rule P is a **Conditional Probability System** if and only if it has **Ordered Surprise** representation.

The construction of the OS representation from a CPS relies on the identification of a chain of nested “surprise orders.” The initial prior is $\mu_0 = P(\cdot|S)$. We then separate S into feasible and null states (if any). Let $S_1 \subset S$ be the set of null states for $P(\cdot|S)$, then $\mu_1 = P(\cdot|S_1)$. We continue in this way until we exhaust the state space, yielding $\mathcal{O} = \langle \mu_0, \dots, \mu_K \rangle$.

Since the OS and CPS updating rules are equivalent, families of OS preferences and CPS preferences are behaviorally equivalent as well because both admit an SEU representation (see [section 3](#)).

2.4 Hypothesis Testing Model

A recent and elegant addition to the literature on updating for null events is the Hypothesis Testing model of Ortoleva (2012). In this model, a DM is described by a second-order prior ρ and a plausibility threshold ϵ . Before any information is revealed, the DM selects her initial prior, which is the most likely prior according to ρ . The DM is SEU with respect to the initial prior and applies Bayes' rule whenever the conditioning event is "expected:" its probability exceeds ϵ . When the event is "unexpected," the DM revisits her second-order beliefs and selects a (possibly) new belief according to a maximum likelihood rule. Bayes' rule is then applied to the selected belief.

Importantly, the notion of "unexpected" events is endogenously determined via $\epsilon \in [0, 1)$. When event E is realized, it is unexpected if $\mu(E) \leq \epsilon$. When $\epsilon > 0$, the DM may exhibit non-Bayesian reactions to unexpected events. When $\epsilon = 0$, only null-events are surprising and Bayes' rule is applied whenever possible.

Definition 5. *A complete updating rule P has a **Hypothesis Testing** representation if there are probability distributions $\mu_0, \dots, \mu_K \in \Delta(S)$ such that $S = \cup_{i=0}^K \text{supp}(\mu_i)$, a second-order prior $\rho \in \Delta(\{\mu_0, \dots, \mu_K\})$ satisfying $\rho(\mu_0) > \rho(\mu_k)$ for all $k > 0$, and an $\epsilon \in [0, 1)$ such that*

$$P(\cdot|E) = \begin{cases} BU(\mu_0, E) & \text{if } \mu_0(E) > \epsilon \\ BU(\mu_{k^*}, E) & \text{if } \mu_0(E) \leq \epsilon, \end{cases}$$

where $\mu_{k^*}(E)\rho(\mu_{k^*}) > \mu_j(E)\rho(\mu_j)$ for all $j \neq k^*$.

In the HT with $\epsilon = 0$, the DM uses Bayes' rule for all ex-ante feasible events. Since the DM only selects a new prior after null-events, this is intuitively similar to the OS model. We show in the following theorem that every OS representation of beliefs admits an HT representation with $\epsilon = 0$. Consequently, OS preferences are a special case of HT preferences with $\epsilon = 0$.⁷

Theorem 2. *Any complete updating rule P with an OS representation has an HT representation with $\epsilon = 0$.*

We prove this by taking an OS representation $\mathcal{O} = \langle \mu_0, \dots, \mu_K \rangle$ and constructing a second-order belief ρ such that the resulting HT representation produces the same con-

⁷Dominiak and Lee (2022) develop a solution concept for signaling games using the HT model with $\epsilon = 0$.

ditional beliefs. The key insight is that the relative probability between surprise orders must be sufficiently large so that the correct μ_k is selected. That is, if $j < l$, then $\frac{\rho(\mu_j)}{\rho(\mu_l)}$ must be large enough so that μ_l is only selected when μ_j is impossible. Viewed another way, there is an event-independent linear order that governs the selection from \mathcal{O} .

This implication cannot be reversed as the following example illustrates. Of course, this is not surprising for HT with $\epsilon > 0$. However, even when $\epsilon = 0$, HT preferences may be inconsistent with OS preferences. This is because in HT, selection of a new prior μ after an event E is jointly determined by “how likely the event is under the prior,” given by $\mu(E)$, and “how likely the DM finds the prior,” given by $\rho(\mu)$.

Example 4. Recall our coin example (Example 3), but suppose instead that P has a HT representation with $\epsilon = 0$, $\rho(\mu_0) = \frac{1}{2}$, $\rho(\mu_1) = \frac{1}{3}$, and $\rho(\mu_2) = \frac{1}{6}$. When $A = \{e', l_1, l_2\}$,

$$\mu_1(A)\rho(\mu_1) = \frac{1}{8} \cdot \frac{1}{3} < \frac{1}{6} = \mu_2(A)\rho(\mu_2),$$

since $\mu_1(A) = \frac{1}{8}$ and $\mu_2(A) = 1$. Thus, $P^{HT}(\cdot|A) = BU(\mu_2, A)$. However, when $E = \{e, e', l_1, l_2\}$,

$$\mu_1(E)\rho(\mu_1) = \frac{1}{3} > \frac{1}{6} = \mu_2(E)\rho(\mu_2)$$

since $\mu_1(E) = \mu_2(E) = 1$. Thus, $P^{HT}(\cdot|E) = BU(\mu_1, E)$. Hence, the described HT representation does not admit an OS representation because

$$P^{HT}(e' | E) = \frac{1}{8} \neq 0 \cdot \frac{1}{8} = P^{HT}(e' | A) \cdot P^{HT}(A | E),$$

violates (2) in Definition 4.

The relationship between OS, CPS, and HT is illustrated in Figure 1.




Figure 1: Relationship between OS, CPS, and HT.

2.5 ϵ -Ordered Surprises

A natural way to generalize the Ordered Surprises representation is to retain the “sequential selection” of new beliefs while incorporating the idea of “non-Bayesian reaction to unexpected events” from the HT model. To do so, we introduce ϵ -OS. Since OS is equivalent to CPS, we interpret ϵ -OS as a way to define a one-parameter non-Bayesian extension of the CPS. This extension may lead to an interesting, non-Bayesian generalization of sequential equilibria.

Definition 6. *A complete updating rule P has an ϵ -Ordered Surprise representation if there are probability distributions $\mu_0, \dots, \mu_K \in \Delta(S)$ and $\epsilon \in [0, 1)$ such that*

$$P(\cdot|E) = BU(\mu_{k^*}, E) \text{ where } k^* = \min\{k \leq K \mid \mu_k(E) > \epsilon\},$$

for every $E \in \Sigma$.

The ϵ -OS representation incorporates the key idea of HT by allowing for non-Bayesian reactions to unexpected events: $\mu_j(E) \leq \epsilon$. However, it provides additional structure to the posterior selection process. The ϵ -OS remains a special case of HT.

Theorem 3. *Any complete updating rule that has an ϵ -OS representation also has a HT representation. Moreover, if $\epsilon = 0$, then the threshold for HT representation is also zero.*

3 Ordered Surprises Preferences

In this section, we provide a behavioral characterization of OS. Say that a family of preferences $\{\succsim_E\}_{E \in \Sigma}$ admits an **Ordered Surprises** (OS) representation if there is a complete updating rule P that has an OS representation $\mathcal{O} = \langle \mu_0, \dots, \mu_K \rangle$ and each preference \succsim_E admits an SEU representation $(u_k, P(\cdot|E))$. That is, for all $f, g \in \mathcal{F}$:

$$f \succsim_E g \quad \text{if and only if} \quad \sum_{s \in E} u_k(f(s))P(s|E) \geq \sum_{s \in E} u_k(g(s))P(s|E).$$

Note that risk attitudes may depend on the order of the surprise, consistent with the approach taken by Myerson (1986b).

Our first axiom consists of several basic conditions that ensure an SEU representation.

Axiom 1 (Conditional SEU Preferences). *for every $E \in \Sigma$,*

(i) **(Weak Order)** \succsim_E is complete and transitive;

(ii) **(Continuity)** for any $f, g, h \in F$, if $f \succ_E g$ and $g \succ_E h$, then there are $\alpha, \beta \in (0, 1)$ such that $\alpha f + (1 - \alpha)h \succ_E g$ and $g \succ_E \beta f + (1 - \beta)h$.

(iii) **(Independence)** for any $f, g, h \in F$ and $\alpha \in (0, 1]$,

$$f \succsim_E g \text{ if and only if } \alpha f + (1 - \alpha)h \succsim_E \alpha g + (1 - \alpha)h$$

(v) **(Nondegeneracy)** there are $f, g \in \mathcal{F}$ such that $f \succ_E g$.

Each condition is standard and so discussion is omitted. We also assume **Consequentialism**, a well-known condition that ensures information is believed by the DM.

Axiom 2 (Consequentialism). For every $E \in \Sigma$ and all $f, g \in \mathcal{F}$, if $f(s) = g(s)$ for all $s \in E$, then $f \sim_E g$.

This axiom requires that each conditional preference \succsim_E depends solely on the conditioning event E . Combined with **Conditional SEU Preferences, Consequentialism** implies that all states in the complementary event, $S \setminus E$, are null according to \succsim_E .

Our next axiom is novel and is the key condition for an OS representation.

Axiom 3 (Conditional Consistency). For all $E \in \Sigma$, \succsim_E -feasible $A \subset E$, and $f, g, h \in \mathcal{F}$,

$$f_A h \succsim_E g_A h \text{ if and only if } f \succsim_A g.$$

Conditional Consistency implies Dynamic Consistency (DC) but also has bite on events that are (ex-ante) \succsim_S -null. For comparison, note that DC only has implications for events that are ex-ante feasible: for A that is \succsim_S -feasible, $f_A h \succsim_S g_A h$ if and only if $f \succsim_A g$. In essence, **Conditional Consistency** extends the logic of DC to all conditional preferences E and nested events that are \succsim_E -feasible (see Epstein and Breton (1993) and Ghirardato (2002) for an excellent discussion of DC).

Theorem 4. A family of preferences $\{\succsim_E\}_{E \in \Sigma}$ admits an OS representation if and only if it satisfies **Conditional SEU Preferences, Consequentialism and Conditional Consistency**.

Axiom 4 (Risk Independence). For all lotteries $p, q \in \Delta(X)$, and events $E \in \Sigma$,

$$p \succsim_S q \text{ if and only if } p \succsim_E q.$$

Proposition 1. *Suppose the family of preferences $\{\succsim_E\}_{E \in \Sigma}$ admits an OS representation. The family satisfies **Risk Independence** if and only if risk attitudes are surprise independent: for each $k > 0$, there are $\alpha_k > 0$, and β_k such that $u_k = \alpha_k u_0 + \beta_k$.*

4 Lexicographic Probability System

When modeling conditional beliefs, the notion of sequential belief selection appears frequently in the literature. Accordingly, the word “lexicographic” is often used to describe such procedures. However, “lexicographic” is used with various meanings, as these models are typically quite different. Consequently, it is useful to clarify the difference between the Conditional LPS of Blume et al. (1991) and CPS, since we show that CPS precisely uses a sequence of probability distributions.

Definition 7. *A **Lexicographic Probability System (LPS)** is a list of probability distributions (μ_0, \dots, μ_K) . The Conditional LPS (CLPS) given E is the LPS obtained from the Bayesian updates of each μ_k (such that it is well-defined): $(BU(\mu_{k_1}, E), \dots, BU(\mu_{k_J}, E))$.*

The LPS Preference \succsim^{LPS} is defined by

$$f \succsim^{LPS} g \quad \text{if and only if} \quad \left(\sum_{s \in S} u(f(s)) \mu_i(s) \right)_{i=0}^K \geq^L \left(\sum_{s \in S} u(g(s)) \mu_i(s) \right)_{i=0}^K,$$

where \geq^L is the lexicographic order on \mathbb{R}^{K+1} . The Conditional \succsim_E is defined analogously.

There are several differences between the CLPS and CPS.⁸ First, our axiomatic characterization (in addition to Myerson (1986a)) shows that CPS is compatible with SEU, while CLPS preferences are not because they violate Continuity (except for $K = 0$). Hence, from the behavioral point of view, CLPS and CPS are fundamentally different.⁹ Second, since the CLPS is defined with Bayes’ rule, it must rule out (Savage) null events to be well defined.¹⁰ In contrast, CPS allows for null events and beliefs are well-defined for all

⁸Hammond (1994) makes a mathematical connection between CPS and CLPS: some probability distributions generated by CPS also can be generated by CLPS. However, there is no behavioral connection between CLPS and CPS; there is no preference that has both CLPS and CPS representations (except SEU preferences with full support).

⁹Imposing **Risk Independence** (State Independence in Blume et al. (1991)) on CLPS preferences implies full-support. In contrast, the CPS is a complete updating rule and allows for risk preferences to depend on the surprise order.

¹⁰In an LPS with null events, the CLPS is not defined for these events. Thus, LPS updating is incomplete; there are scenarios in which some conditional probabilities are not well-defined. In contrast, the CPS is a

conditioning events.

To further clarify, the following example illustrates that CLPS rules out indifference (indifferences are resolved ex-ante), while OS allows for information to resolve indifferences. That is, OS preferences exhibit a (strictly) positive value of information (Epstein and Breton, 1993; Ghirardato, 2002): $f_{Eh} \sim g_{Eh}$ and $g \succ_E f$ for some E and some acts.

Example 5. *Since (Conditional) LPS violates continuity, it rules out indifference between distinct acts. Recall our introductory coin example and consider the following acts:*

$$f = \begin{cases} \$v & \text{if } s \in \{h, t\} \\ \$0 & \text{otherwise} \end{cases} \quad g = \begin{cases} \$1 & \text{if } s \in \{h, t\} \\ \$t & \text{if } s \in \{e, e'\} \\ \$0 & \text{otherwise} \end{cases}.$$

Under OS, $f \sim^{OS} g$ if and only if $v = 1$, $f \succ^{OS} g$ when $v > 1$, and $g \succ^{OS} f$ if $v < 1$; the value of t is irrelevant. In the LPS generated by μ_0, μ_1, μ_2 the DM is never indifferent between f and g . If $v > 1$, $f \succ^{LPS} g$ whereas if $v \leq 1$, $g \succ^{LPS} f$.

Suppose $v = 1$, and consider $E = \{e, e'\}$. Under OS, the realization of E can resolve indifference: $f \sim^{OS} g$ and $g \succ_E^{OS} f$. In contrast, under CLPS the ex-ante and conditional preferences must agree: $g \succ^{LPS} f$ and $g \succ_E^{LPS} f$. CLPS preferences can never exhibit resolution of indifference.

5 Concluding Remarks

We introduce and behaviorally characterize the Ordered Surprises representation of preferences. The OS belief structure yields a complete updating rule that is (behaviorally) equivalent to the CPS. Additionally, we show that OS is a special case of HT, clarifying the relationship between CPS and HT.

The OS representation provides a tractable way to model any type of dynamic decision problem that allows for zero-probability contingencies, including dynamic games, strategic communication, information transmission, and Bayesian persuasion. Fruitful directions for future research include extending the idea of Ordered Surprises to ambiguity or other non-SEU preferences.

complete updating rule; conditional probabilities are always well-defined.

A Proofs

A.1 Proof of Theorem 1

(\Leftarrow) Consider an updating rule P with an OS representation $\mathcal{O} = \langle \mu_0, \dots, \mu_K \rangle$. We must show that P satisfies the conditions in Definition 4. By assumption, the rule is complete.

Since P has an OS representation, let $\Sigma_0, \dots, \Sigma_K$ be the partition of Σ such that for each k , Σ_k is the collection of events for which μ_k is used for updating:

$$\Sigma_k = \{E \in \Sigma \mid k = \min\{\tilde{k} \leq K \mid \mu_{\tilde{k}}(E) > 0\}\}.$$

We first show that P is consequentialist. Pick any $E \in \Sigma$. Then for some k^* , $E \in \Sigma_{k^*}$. By assumption, $P(\cdot|E) = BU(\mu_{k^*}, E)$, which is well-defined. Hence $P(E|E) = 1$.

To show that (2) holds, take $G, F, E \in \Sigma$ such that $G \subset F \subset E$ and $F \neq \emptyset$.

Case 1: Suppose $G, F, E \in \Sigma_k$ for some $k \leq K$. Since P has an OS representation,

$$\begin{aligned} P(G|E) &= BU(\mu_k, E)(G) = \frac{\mu_k(G)}{\mu_k(E)} = \frac{\mu_k(G)}{\mu_k(E)} \times \frac{\mu_k(F)}{\mu_k(F)} = \\ &\frac{\mu_k(G)}{\mu_k(F)} \times \frac{\mu_k(F)}{\mu_k(E)} = BU(\mu_k, F)(G) \times BU(\mu_k, E)(F) = P(G|F) \times P(F|E). \end{aligned} \quad (3)$$

Case 2: For $j \in \{G, F, E\}$, let k_j denote the index for which $j \in \Sigma_{k_j}$. Note that $k_G = k_F = k_E =$ is precisely Case 1. If $P(G|E) > 0$, then $k_G = k_E$. Further, since $G \subset F$, it follows that $0 < \mu_{k_E}(G) \leq \mu_{k_E}(F)$, and so $P(F|E) > 0$ and $k_G = k_F = k_E$. Suppose $P(G|E) = 0$. Then, $k_G \neq k_E$. For (2) to hold, we must show that either $P(G|F)$ or $P(F|E)$ is zero (or both). Suppose $P(F|E) > 0$. Then,

$$P(F|E) = BU(\mu_{k_E}, E)(F) = \frac{\mu_{k_E}(F)}{\mu_{k_E}(E)} > 0.$$

Hence, $F \in \Sigma_{k_E}$, and so $k_E = k_F$. Consequently, $k_G \neq k_F$ and $P(G|F) = 0$. Hence, (2) holds.

(\Rightarrow) Let P be a CPS. We will construct an Ordered Surprises representation $\mathcal{O} = \langle \mu_0, \dots, \mu_K \rangle$.

Set $S_0 := S$ and consider $P(\cdot|S_0) \in \Delta(S_0)$. Define $\Sigma_0 := \{E \subseteq S_0 \mid P(E|S_0) > 0\}$ and $F_0 := \{s \in S_0 : P(\{s\}|S_0) > 0\}$. For each $E \in \Sigma_0$, $P(\cdot|E)$ is the Bayesian update of $P(\cdot|S_0)$ conditional on E . To see why, consider $E \in \Sigma_0$ and $G \subset S_0$. Since $G \cap E \subseteq E \subseteq$

S_0 , by (2), we have $P(G \cap E|S_0) = P(G \cap E|E) \times P(E|S_0)$. Equivalently,

$$\frac{P(G \cap E|S_0)}{P(E|S_0)} = P(G \cap E|E), \quad (4)$$

since $P(E|S_0) > 0$. Further, since P is a CPS, $P(E|E) = 1$. Hence, $P(\cdot|E)$ is the Bayesian update of $P(\cdot|S_0)$. Let $\mu_0 := P(\cdot|S_0)$. If $F_0 = S_0$ (all states are non-null), there are no surprises and we are done. However, if $F_0 \subset S_0$, there are surprises and we proceed to the next step.

Define $S_1 := S_0 \setminus F_0$, the first-order surprise. Consider $P(\cdot|S_1)$ and, as before, define the sets $\Sigma_1 := \{E \subseteq S_1 \mid P(E|S_1) > 0\}$ and $F_1 := \{s \in S_1 : P(\{s\}|S_1) > 0\}$. Since P is a CPS, $P(S_1|S_1) = 1$. By the same argument as above, for each $E \in \Sigma_1$, $P(\cdot|E)$ is the Bayesian update of $P(\cdot|S_1)$ given E . Let $\mu_1 := P(\cdot|S_1)$. If $F_1 = S_1$, there are no more surprises. If $F_1 \subset S_1$, we proceed.

In step k , we define $S_k = S_{k-1} \setminus F_{k-1}$. Consider the conditional probability $P(\cdot|S_k)$, and define $\Sigma_k = \{E \subseteq S_k : P(E|S_k) > 0\}$ and $F_k = \{s \in S_k : P(\{s\}|S_k) > 0\}$. As before, $P(\cdot|E)$ must be the Bayesian update of $p(\cdot|S_k)$ conditional on S_k . We define $\mu_k := P(\cdot|S_k)$. We proceed in this way until we can find $K \leq |S|$ such that $F_K = S_K$. Since S is finite, such a K exists.

Finally, notice that $\Sigma_0, \Sigma_1, \dots, \Sigma_K$ is a partition of Σ and that $\mathcal{O} = \langle \mu_0, \dots, \mu_K \rangle$ is an OS representation of P .

A.2 Proof of Theorem 2

Consider a complete updating rule P that admits an OS representation $\mathcal{O} = \langle \mu_0, \dots, \mu_K \rangle$. We will construct a second-order belief $\rho \in \Delta(\{\mu_0, \dots, \mu_K\})$ such that P all has a HT representation.

Without loss, $\{supp(\mu_k)\}_{k=0}^K$ forms a partition of S . For each $k \leq K$, let F_k denote the feasible states for μ_k and $\delta_k = \min_{s \in F_k} \mu_k(s) \in (0, 1)$. Define the following $K + 1$ numbers: $v(0) = 1$ and for $k \geq 1$, $v(k) = \frac{\delta_{k-1}}{2} v(k-1)$. This is a decreasing sequence with $v(k) = 2^{-k} \prod_{j=0}^{k-1} \delta_j$ for $k \geq 1$. We use these numbers to construct ρ . For each μ_k ,

$$\rho(\mu_{S_k}) = \frac{v(k)}{\sum_{j=0}^K v(j)},$$

and $\rho(\mu) = 0$ for all $\mu \in \Delta(S) \setminus \mathcal{O}$. Finally, let $\epsilon = 0$.

By construction, $\{\mu_0\} = \arg \max \rho(\mu)$. Consider any $E \in \Sigma$. If $\mu_0(E) > 0$, then $P(\cdot|E) = BU(\mu_0, E)$ since P also has an OS representation. If $E \subseteq F_k$ for some $k > 0$, then since $\mu_k(E) > 0$ and $\mu_{k'}(E) = 0$ for all $k' \neq k$, it follows that $\mu_k(E)\rho(\mu_k) > 0 = \mu_{k'}\rho(\mu_{k'})$ and $P(\cdot|E) = BU(\mu_k, E)$. The final possibility is $E \cap F_k \neq \emptyset$ for multiple k . Let $K(E)$ denote all k such that $E \cap F_k \neq \emptyset$ and let \underline{k} be the least such k . Note that $\mu_{\underline{k}}(E) \geq \delta_{\underline{k}}$ and if $k \in K(E) \setminus \underline{k}$, then $k > \underline{k}$ and

$$\frac{\rho(\mu_{\underline{k}})}{\rho(\mu_k)} = \frac{v(\underline{k})}{v(k)} = \frac{2^{-\underline{k}} \prod_{j=0}^{\underline{k}-1} \delta_j}{2^{-k} \prod_{j=0}^{k-1} \delta_j} = \frac{2^{k-\underline{k}}}{\prod_{j=\underline{k}}^{k-1} \delta_j}.$$

Thus,

$$\prod_{j=\underline{k}}^{k-1} \delta_j \rho(\mu_{\underline{k}}) = 2^{k-\underline{k}} \rho(\mu_k).$$

Since $\mu_{\underline{k}}(E) \geq \delta_{\underline{k}} \geq \prod_{j=\underline{k}}^{k-1} \delta_j$ and $\mu_k(E) \leq 1$, it follows that

$$\mu_{\underline{k}}(E)\rho(\mu_{\underline{k}}) \geq \prod_{j=\underline{k}}^{k-1} \delta_j \rho(\mu_{\underline{k}}) = 2^{k-\underline{k}} \rho(\mu_k) > \mu_k(E)\rho(\mu_k),$$

and

$$P(\cdot|E) = BU(\mu_{\underline{k}}, E),$$

where the last line holds because P admits an OS representation.

A.3 Proof of Theorem 3

Take any complete updating rule P that has an ϵ -OS representation with some $\epsilon \in [0, 1)$. Then, there are probability distributions μ_0, \dots, μ_K such that

$$P(\cdot|E) = BU(\mu_{k^*}, E) \text{ where } k^* = \min\{k \leq K \mid \mu_k(E) > \epsilon\}$$

for every $E \in \Sigma$. Let $\Sigma_0, \dots, \Sigma_K$ be a partition of Σ such that for each k , Σ_k is the collection of events for which the prior μ_k is used for updating:

$$\Sigma_k = \{E \in \Sigma \mid k = \min\{\tilde{k} \leq K \mid \mu_{\tilde{k}}(E) > \epsilon\}\}.$$

Throughout this proof, we assume that for any $k \leq K$, E_k is an element of Σ_k . Take $\bar{\rho}_0, \underline{\rho}_0, \dots, \bar{\rho}_K, \underline{\rho}_K$ with

$$\bar{\rho}_0 > \underline{\rho}_0 > \bar{\rho}_1 > \underline{\rho}_1 > \dots > \bar{\rho}_K > \underline{\rho}_K > \delta \bar{\rho}_0 > 0$$

and $\underline{\rho}_k > \bar{\rho}_k \mu_k^{E'_k}(E_k)$ for any E_k, E'_k with $\mu_k^{E'_k}(E_k) < 1$.

Let $\mu_k^E = \text{BU}(\mu_k, E)$ for any $E \in \Sigma$. Let ρ be an element of $\Delta(\{\mu_k^{E_k}\}_{k \leq K, E_k \in \Sigma_k})$ such that (i) $\rho(\mu_k^{E_k}) \in (\underline{\rho}_k, \bar{\rho}_k)$ for any $k \leq K$ and (ii) $\rho(\mu_k^{E_k}) > \rho(\mu_k^{E'_k})$ if $\mu_k^{E_k} \neq \mu_k^{E'_k}$ and $\mu_k^{E'_k}(E_k) = 1$.

Let us first show that there is ρ that satisfies (ii). Let $\mu_k^{E_k} \succ^* \mu_k^{E'_k}$ if $\mu_k^{E_k} \neq \mu_k^{E'_k}$ and $\mu_k^{E'_k}(E_k) = 1$. It is enough to show that \succ^* is acyclic. To show acyclicity, suppose that there are E_k^1, \dots, E_k^T such that $\mu_k^{E_k^t}(E_k^{t+1}) = 1$ for each $t \leq T - 1$ and $\mu_k^{E_k^T}(E_k^1) = 1$. Note that $\mu_k^{E'_k}(E_k) = 1$ is equivalent to $\text{supp}(\mu_k) \cap E'_k \subseteq E_k$. Hence, $\mu_k^{E'_k}(E_k) = 1$ implies $\text{supp}(\mu_k) \cap E'_k \subseteq \text{supp}(\mu_k) \cap E_k$. Then, $\mu_k^{E_k^t}(E_k^{t+1}) = 1$ implies $\text{supp}(\mu_k) \cap E_k^t \subseteq \text{supp}(\mu_k) \cap E_k^{t+1}$ and $\mu_k^{E_k^T}(E_k^1) = 1$ implies $\text{supp}(\mu_k) \cap E_k^T \subseteq \text{supp}(\mu_k) \cap E_k^1$. Hence, $\text{supp}(\mu_k) \cap E_k^t = \text{supp}(\mu_k) \cap E_k^{t'}$ for any t, t' ; i.e., $\mu_k^{E_k^t} = \mu_k^{E_k^{t'}}$.

We now show that P has a HT representation with (ρ, δ) when δ is large enough. Hence, we shall show that for any E_k ,

$$\rho(\mu_k^{E_k}) \mu_k^{E_k}(E_k) = \rho(\mu_k^{E_k}) > \rho(\mu_j^{E_j}) \mu_j^{E_j}(E_k) \text{ for any } j \neq k.$$

For any $j > k$, the above holds since $\rho(\mu_k^{E_k}) > \rho(\mu_j^{E_j})$. Suppose now $j < k$. In this case, $\mu_j(E_k) \leq \epsilon$ since k is the lowest index such that $\mu_k(E_k) > \epsilon$. Then, $\mu_j^{E_j}(E_k) = \text{BU}(\mu_j, E_j)(E_k) = \frac{\mu_j(E_k \cap E_j)}{\mu_j(E_j)}$. Since $\mu_j(E_k) \leq \epsilon$ and $\mu_j(E_j) > \epsilon$, there is a large enough $\delta \in [0, 1)$ such that $\mu_j^{E_j}(E_k) \leq \delta$. Hence, by the construction of ρ ,

$$\rho(\mu_k^{E_k}) > \delta \rho(\mu_j^{E_j}) \geq \rho(\mu_j^{E_j}) \mu_j^{E_j}(E_k).$$

We finally show that the HT representation correctly chooses $\mu_k^{E_k}$ among $\{\mu_k^{E'_k}\}_{E'_k}$ for each E_k . When $\mu_k^{E'_k}(E_k) < 1$, we have

$$\rho(\mu_k^{E_k}) \mu_k^{E_k}(E_k) = \rho(\mu_k^{E_k}) > \underline{\rho}_k > \bar{\rho}_k \mu_j^{E'_k}(E_k) > \rho(\mu_k^{E'_k}) \mu_k^{E'_k}(E_k).$$

When $\mu_k^{E'_k}(E_k) = 1$ and $\mu^{E_k} \neq \mu_k^{E'_k}$,

$$\rho(\mu_k^{E_k})\mu_k^{E_k}(E_k) = \rho(\mu_k^{E_k}) > \rho(\mu_k^{E'_k}) = \rho(\mu_k^{E'_k})\mu_j^{E'_k}(E_k).$$

It is immediate from the above construction of δ , $\delta = 0$ whenever $\epsilon = 0$.

A.4 Proof of Theorem 4

Necessity of the axioms is trivial, so we only prove sufficiency. Take $E \in \Sigma$ and a conditional preference relation \succsim_E .

Step 1. By **Conditional SEU Preferences**, \succsim_E admits a state-dependent SEU representation. There are a probability measure μ_E on S and, for each $s \in E$, an expected utility function U_s on $\Delta(X)$, such that for all $f, g \in \mathcal{F}$:

$$f \succsim_E g \quad \text{if and only if} \quad \sum_{s \in E} U_s(f(s))\mu_E(s) \geq \sum_{\omega \in S_E} U_s(g(s))\mu_E(s).$$

A state $s \in S$ is \succsim_E -null if and only if $\mu_E(s) = 0$. By **Consequentialism**, states outside of E are \succsim_E -null. Moreover, μ_E is unique and for each $s \in E$, U_s is unique up to a positive affine transformation.

Step 2. (U_s is state-independent on F_E). Let $F_E = \{s \in S : s \text{ is } \succsim_E\text{-feasible}\}$ be the set of feasible states according to \succsim_E . Take $A = \{s\}$ where $s \in F_E$. By Step 1, $\succsim_{\{s\}}$ is SEU with respect to U'_s and $\mu_{\{s\}}$ such that for all $f, g \in \mathcal{F}$, $f \succsim_{\{s\}} g$ if and only if $U'_s(f(s)) \geq U'_s(g(s))$ if and only if $U'_s(p) \geq U'_s(q)$ where $f(s) = p$ and $g(s) = q$. Since $A = \{s\}$ is a \succsim_E -feasible state, by **Conditional Consistency**, for all lotteries $p, q \in \Delta(X)$,

$$p_{\{s\}}h \succsim_E q_{\{s\}}h \quad \text{if and only if} \quad p \succsim_{\{s\}} q,$$

equivalently,

$$U_s(p)\mu_E(s) \succsim_E U_s(q)\mu_E(s) \quad \text{if and only if} \quad U'_s(p) \succsim_{\{s\}} U'_s(q),$$

equivalently, by Step 1,

$$U_s(p) \geq U_s(q) \quad \text{if and only if} \quad U'_s(p) \geq U'_s(q).$$

Hence, U'_s is a positive affine transformation of U_s . Hence, for each \succsim_E -feasible state s in F_E , U_s is state-independent and so without loss $U'_s = U_s$.

Step 3. (Bayesian updating). Let $A \in \Sigma$ be a \succsim_E -feasible event with $A \subseteq E$. Notice that A is \succsim_E -feasible if and only if $\mu_E(A) > 0$, or equivalently, $A \cap F_E \neq \emptyset$. Consider acts $f_A h$ and $g_A h$ such that $f_A h \succsim_E g_A h$. By Step 1,

$$\sum_{s \in S} U_s(f_A h(s)) \mu_E(s) \geq \sum_{s \in S} U_s(g_A h(s)) \mu_E(s). \quad (5)$$

By **Consequentialism**, $\mu_E(S \setminus E) = 0$. Thus,

$$\sum_{s \in E} U_s(f_A h(s)) \mu_E(s) \geq \sum_{s \in E} U_s(g_A h(s)) \mu_E(s), \quad (6)$$

or equivalently,

$$\sum_{s \in A} U_s(f(s)) \mu_E(s) \geq \sum_{s \in A} U_s(g(s)) \mu_E(s). \quad (7)$$

By **Conditional Consistency**, $f \succsim_A g$. Thus, by Steps 1 and 2,

$$\sum_{s \in A} U_s(f(s)) \mu_A(s) \geq \sum_{s \in A} U_s(g(s)) \mu_A(s). \quad (8)$$

Since Equations (7) and (8) are equivalent by **Conditional Consistency**, we have Bayesian updating

$$\mu_A(s) = \frac{\mu_E(s)}{\mu_E(A)} \text{ for each } s \in A. \quad (9)$$

Step 4. (Ordered surprises). We proceed inductively. Let $S_0 = S$, and consider \succsim_{S_0} . We let $\mu_0 = \mu_{S_0}$, and if μ_0 has full support, stop. Otherwise, let S_1 denote the set of \succsim_{S_0} -null states, and consider \succsim_{S_1} . Then, let $\mu_1 = \mu_{S_1}$. By **Consequentialism**, $\mu_1(S_0 \setminus S_1) = 0$. If $\text{supp}(\mu_1) = S_1$, stop. Otherwise, let S_2 denote the set of \succsim_{S_1} -null states. We proceed in this fashion until we reach a K such that $\text{supp}(\mu_K) = S_K$. Since S is finite, we must eventually stop.

Then $\mathcal{O} = \langle \mu_0, \dots, \mu_K \rangle$ constitutes an OS representation. To see why, note that by construction $\text{supp}(\mu_k) \cap \text{supp}(\mu_{k'}) = \emptyset$ if $k \neq k'$ and $\cup_{k=0}^K \text{supp}(\mu_k) = S$. Further, from

the preceding step, it follows that for every E ,

$$P(\cdot|E) = \mu_E = BU(\mu_{k^*}, E) = BU(\mu_{S_{k^*}}, E)$$

where $k^* = \min\{k \leq K \mid \mu_k(E) > 0\}$.

Step 5. Combining steps, it follows that for each k there is a u_k such that \succsim_E is represented by $(u_{k^*}, P(\cdot|E))$, where $P(\cdot|E) = BU(\mu_{k^*}, E)$ and $k^* = \min\{k \leq K \mid \mu_k(E) > 0\}$.

A.5 Proof of Proposition 1

Given an OS representation, fix $k > 0$ and consider some E with $\mu_k(E) > 0$. By **Risk Independence**, it follows that for every $p, q \in \Delta(X)$, $p \succsim q$ iff $p \succsim_E q$. Consequently, $u_0(p) \geq u_0(q)$ iff $u_k(p) \geq u_k(q)$, and so by standard results u_0 and u_k represent the same risk preferences.

References

- ACEMOGLU, D., G. COMO, F. FAGNANI, AND A. OZDAGLAR (2013): “Opinion Fluctuations and Disagreement in Social Networks,” *Mathematics of Operations Research*, 38, 1–27.
- ANSCOMBE, F. AND R. AUMANN (1963): “A definition of subjective probability,” *The Annals of Mathematical Statistics*, 34, 199–205.
- BATTIGALLI, P. (1996): “Strategic Independence and Perfect Bayesian Equilibria,” *Journal of Economic Theory*, 70, 201–234.
- BATTIGALLI, P. AND M. SINISCALCHI (1999): “Hierarchies of Conditional Beliefs and Interactive Epistemology in Dynamic Games,” *Journal of Economic Theory*, 88, 188–230.
- (2002): “Strong Belief and Forward Induction Reasoning,” *Journal of Economic Theory*, 106, 356–391.
- BENJAMIN, D. J. (2019): “Errors in probabilistic reasoning and judgment biases,” in *Handbook of Behavioral Economics: Applications and Foundations*, ed. by B. D. Bernheim, S. DellaVigna, and D. Laibson, North-Holland, vol. 2, chap. 2, 69–186.
- BERNHEIM, B. D. (1994): “A Theory of Conformity,” *Journal of Political Economy*, 102, 841–877.
- BLUME, L., A. BRANDENBURGER, AND E. DEKEL (1991): “Lexicographic Probabilities and Equilibrium Refinements,” *Econometrica*, 59, 81–98.

- CRAWFORD, V. P. AND J. SOBEL (1982): “Strategic Information Transmission,” *Econometrica*, 50, 1431–1451.
- DOMINIAK, A., M. KOVACH, AND G. TSERENJIGMID (2022): “Inertial Updating,” .
- DOMINIAK, A. AND D. LEE (2022): “Testing Rational Hypotheses in Signaling Games,” *working paper*.
- EPSTEIN, L. G. (2006): “An Axiomatic Model of Non-Bayesian Updating,” *The Review of Economic Studies*, 73.
- EPSTEIN, L. G. AND M. L. BRETON (1993): “Dynamically consistent beliefs must be Bayesian,” *Journal of Economic Theory*, 61, 1–22.
- GAL-OR, E. (1989): “Warranties as a Signal of Quality,” *Canadian Journal of Economics*, 22, 50–61.
- GALPERTI, S. (2019): “Persuasion: The art of changing worldviews,” *American Economic Review*, 109, 996–1031.
- GHIRARDATO, P. (2002): “Revisiting Savage in a conditional world,” *Economic Theory*, 20, 83–92.
- HAMMOND, P. (1994): “Elementary Non-Archimedean Representations of Probability for Decision Theory and Games,” in *Patrick Suppes: Scientific Philosopher. Synthese Library*, ed. by P. Humphreys, Dordrecht: Springer Netherlands, vol. 234, 25–59.
- JOHN, K. AND J. WILLIAMS (1985): “Dividends, Dilution, and Taxes: A Signalling Equilibrium,” *Journal of Finance*, 40, 1053–1070.
- KOVACH, M. (2020): “Conservative Updating,” *working paper*.
- KREPS, D. M. AND R. WILSON (1982): “Sequential Equilibria,” *Econometrica*, 50.
- LOHMANN, S. (1995): “Information, Access, and Contributions: A Signaling Model of Lobbying,” *Public Choice*, 85, 267–284.
- MCLENNAN, A. (1989): “Consistent Conditional Systems in Noncooperative Game Theory,” *International Journal of Game Theory*, 18, 141–74.
- MILGROM, P. AND J. ROBERTS (1982): “Limit Pricing and Entry under Incomplete Information: An Equilibrium Analysis,” *Econometrica*, 50, 443–459.
- (1986): “Price and Advertising Signals of Product Quality,” *Journal of Political Economy*, 94, 796–821.
- MILLER, R. M. AND C. R. PLOTT (1985): “Product Quality Signaling in Experimental Markets,” *Econometrica*, 53, 837–872.
- MYERSON, R. (1986a): “Axiomatic Foundations of Bayesian Decision Theory,” Discussion Papers 671, Northwestern University, Center for Mathematical Studies in Economics

and Management Science.

MYERSON, R. B. (1986b): “Multistage Games with Communication,” *Econometrica*, 54, 323–358.

ORTOLEVA, P. (2012): “Modeling the Change of Paradigm: Non-Bayesian Reactions to Unexpected News,” *American Economic Review*, 102, 2410–36.

RÉNYI, A. (1955): “On a new axiomatic theory of probability,” *Acta Mathematica Hungarica*, 6, 285–335.

RUBINSTEIN, A. (1985): “A Bargaining Model with Incomplete Information About Time Preferences,” *Econometrica*, 53, 1151–1172.

SPENCE, M. (1973): “Job Market Signaling,” *Quarterly Journal of Economics*, 87, 355–374.

TSAKAS, E. (2018): “Agreeing to Disagree with Conditional Probability Systems,” *The B.E. Journal of Theoretical Economics*, 18, 160–182.