

Geodesic incompleteness of some popular regular black holes

Tian Zhou* and Leonardo Modesto†

Department of Physics, Southern University of Science and Technology, Shenzhen 518055, China

Through out the study of the geodesics of some popular spherically symmetric regular black holes, we hereby prove that the Hayward's black hole and the simplest extension of the Simpson-Visser's non-analytic smooth black hole are both geodetically *incomplete*. Hence, we provide several mere modifications of such spacetimes in order to make them geodetically complete.

I. INTRODUCTION

In order to tame the black holes' singularity problem in Einstein's gravity, many regular spacetimes have been proposed regardless of an underlying dynamic [1–10], namely they are just formal solutions of the Einstein's equations for a proper choice of the energy momentum tensor that violate some or all the energy conditions. So far it turns out that the Kretschmann scalar $R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}$ and other curvature invariants are regular at the center of the proposed regular black holes. However, the geodesic completion has not been studied properly and/or completely and will be the main topic of this paper.

For the sake of simplicity, let us consider a general spherically symmetric solution in Schwarzschild coordinates, namely¹

$$ds^2 = - \left(1 - \frac{2M(r)}{r} \right) dt^2 + \frac{dr^2}{1 - \frac{2M(r)}{r}} + r^2 d\Omega^{(2)}, \quad (1)$$

where the mass is now allowed to be a function of the radial coordinate. When $M(r)$ is constant we are back to the Schwarzschild solution.

The spacetimes we will consider in this paper have regular Kretschmann scalar $R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}$ for $r \geq 0$, although for some metrics the Kretschmann invariant can still be divergent somewhere for $r < 0$. However, the curvature invariants do not have a real physical meaning and only provide a tool for a simple and preliminary analysis of the spacetime's regularity. In physics, we have to look at the geodesic equations and see whether and where they are well defined. We here remind the case of the Taub-NUT's metric that has finite Kretschmann invariant, but it is not geodetically complete. Therefore, the very basic requirement for regular spacetimes should be the completion of all causal geodesics.

For the sake of simplicity, let us here focus on massive probe particles moving in the spacetime (1). The radial geodesics satisfy the following equation in proper time,

$$\dot{x}^2 = -1 \implies g_{tt}\dot{t}^2 + g_{rr}\dot{r}^2 = -1, \quad (2)$$

where $-g_{tt} = g_{rr}^{-1} = 1 - 2M(r)/r$ and the dot represents the derivative respect to the proper time τ . Replacing the conserved quantity corresponding to the time translation invariance, namely $e \equiv -g_{tt}\dot{t}$, in (2), (where e is a dimensionless constant) we end up with the following equation for the radial motion,

$$\dot{r}^2 = e^2 - 1 + \frac{2M(r)}{r}. \quad (3)$$

Choosing now as initial radial position r_i , the proper-time for a massive particle to move from r_i to $r < r_i$ is given by

$$\tau(r) = \int_r^{r_i} dr \frac{1}{\sqrt{\dot{r}^2}}. \quad (4)$$

Since in (3) $\dot{r}^2 > 0$, the right hand side of the above radial geodesic equation (3) is only defined for values of r compatible with:

$$e^2 - 1 + \frac{2M(r)}{r} \geq 0. \quad (5)$$

If the metric we are considering is regular in $r = 0$ (most of the known regular spacetimes have a de Sitter's core in $r = 0$), we also have to check if the proper time τ to reach such point is finite or infinite. If τ is finite (like in the de-Sitter core case), then, we are forced to extend the spacetime beyond $r = 0$ to negative values of the radial coordinate r . There is nothing strange in defining the radial coordinate to be negative because in general relativity the coordinates have no physical meaning. Now for negative values of the radial coordinate two remarkable things may happen: (i) there exists at least a point r_0 in which $\dot{r}^2 \rightarrow \infty$, namely there is at least one pole, or (ii) \dot{r}^2 is always finite $\forall r$. In the latter case, \dot{r}^2 can change sign depending on the value of the energy e . Indeed, for e small enough \dot{r}^2 can be zero let's say in the point r_0 . On the other hand if e is large enough $\dot{r}^2 > 0 \forall r$. Let us expand on the points (i) and (ii).

In the first case, near the point r_0 , we have:

$$\dot{r}^2 \sim (r - r_0)^{-n}, \quad n > 0, \quad (6)$$

where the power n can be a real number. Hence, the proper time to reach the point r_0 is finite, but the geodesic equation (3) is singular $\forall e$ and we can not extend the geodesic motion beyond r_0 . Moreover, the causal structure of the spacetime does not allow the particle to bounce back because near r_0 the radial coordinate

* 11930538@mail.sustech.edu.cn

† lmodesto@sustech.edu.cn

¹ We here work in Planck units, namely the Newton's gravitational constant, the speed of the light, and the Planck's constant are: $c = G = \hbar = 1$.

is time-like, exactly like for the Schwarzschild metric for $r \gtrsim 0$.

In the second case, close to r_0 (for e small enough), we usually have:

$$\dot{r}^2 \sim (r - r_0). \quad (7)$$

Again, according to (4), the proper time to reach r_0 is finite. Remarkably, if (7) holds \dot{r}^2 changes sign in r_0 , which implies that massive particles will bounce back because of the potential barrier whether their energy is not big enough. On the other hand for e large enough $\dot{r}^2 > 0 \forall r$. It deserves to be notice that for some probe particles with special value of the energy, $\dot{r}^2 \sim (r - r_0)^n$, where the power n is an integer such that $n \geq 2$. These massive particles take an infinite amount of proper time to arrive in r_0 .

Looking at the effective potential

$$V(r) = -g_{tt}(r) = 1 - \frac{2M(r)}{r}, \quad (8)$$

if it is finite in the whole spacetime, the particles with energy large enough can travel to arbitrarily negative values of the radial coordinate r and the spacetime is geodesically complete. On the other hand, if the effective potential tends to positive infinity at a certain point r_0 , every massive particle will bounce back. However, massless particles may arrive at r_0 for a finite value of the affine parameter. Indeed, the null geodesics satisfy the equation $\dot{r}^2 = e^2$ (this is what actually happens for the Reissner-Nordström black hole in $r = 0$). Looking at the curvature invariants, the Kretschmann scalar will diverge where g_{tt} is singular and there is a curvature singularity in $r = r_0$. Therefore, it is easy to show that the spacetime is complete for time-like geodesics, but incomplete for null geodesics.

We conclude that the geodesic completion of the metric (1) is tantamount the absence of singularities in the $g_{tt}(r)$ component of the metric $\forall r \in [-\infty, +\infty]$.

In this paper, we will focus on the geodesic completeness of two very popular regular black holes and we will point out that they are actually geodesically incomplete.

II. GEODESIC INCOMPLETENESS OF HAYWARD'S BLACK HOLE

We here consider the spherically symmetric and static Hayward's metric [2], namely

$$ds^2 = -f(r)dt^2 + f^{-1}(r)dr^2 + r^2d\Omega^{(2)}, \quad (9)$$

where the function $f(r)$ takes the form

$$f(r) = 1 - \frac{2Mr^2}{r^3 + L^3}, \quad (10)$$

in which M is a constant mass and L is a length scale. The metric asymptotically approaches the Schwarzschild spacetime, but near $r = 0$ the metric shows a de Sitter core because $f(r) \sim 1 - (2M/L^3)r^2$, contrary to




FIG. 1. The left plot shows $\dot{r}^2(r)$ for $e = 2$, $M = 5$ and $L = 1$. The right plot shows the effective potential $V = -g_{tt}(r)$ for the Hayward spacetime with $M = 5$ and $L = 1$.

the Schwarzschild singularity present at the center of the black hole.

Let us now focus on the issue of the geodesic incompleteness of the Hayward's black hole. Consider a probe massive particle that falls radially into the black hole. The radial geodesic equation of motion of a test particle is given by

$$\dot{r}^2 = e^2 - 1 + \frac{2Mr^2}{r^3 + L^3}, \quad e \equiv -g_{tt}\dot{t}, \quad (11)$$

while the effective potential reads:

$$V \equiv 1 - \frac{2Mr^2}{r^3 + L^3}. \quad (12)$$

A probe particle with energy $e < 1$ can not overcome the potential barrier and will bounce back consistently with the causality structure of the spacetime. Indeed, $g_{tt} < 0$ for $r \sim 0$. On the the other hand, for $e > 1$, $\dot{r}^2 \sim e^2 - 1$ when $r \rightarrow 0$. Integrating equation (4) for the case of the Hayward's metric, we find that a probe particle can reach the point $r = 0$ in a finite amount of proper time. Therefore, massive particles can cross the region $-L \leq r \leq 0$ in finite proper time, and the point $r = -L$ turns out to be a singularity of the Hayward's metric. Indeed, the spacetime structure of the Hayward metric near $r = -L$ is very similar to the Schwarzschild one near $r = 0$. In particular, $\dot{r} \sim (r + L)^{-1}$ for $r \rightarrow -L$, and after integrating (4) in the radial coordinate r starting from a general initial position, the proper time to reach $r = -L$ turns out to be finite. Therefore, the geodesics will end in $-L$ in finite proper time and the extended Hayward spacetime is geodesically incomplete.

We here implemented the unique spacetime continuation to negative values of r consistently with the request that $f(r)$ is a C^∞ function.




FIG. 2. $\dot{r}^2(r)$ for $e = 2$, $M = 5$, and $L = 1$. From the left to the right, the values of n are 4, 5, and 6 respectively.

III. GENERALIZED HAYWARD METRIC

We hereby consider the following generalization of the Hayward's metric,

$$f(r) = 1 - \frac{2Mr^{n-1}}{r^n + L^n}, \quad n \in \mathbb{N}, \quad n \geq 3. \quad (13)$$

The metric defined in terms of the function (13) is again asymptotically Schwarzschild, but near $r = 0$, $f(r) \sim 1 - (2M/L^n)r^{n-1}$, and the Kretschmann scalar $R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} \sim r^{2n-6}$ turns out to be regular when $n \geq 3$. Therefore, for the generalized Hayward metric, there is no curvature singularity in $r \in [0, +\infty)$.

Focusing on the geodesic completeness of the metric (9) with (13), the equation of motion for radial time-like geodesics reads:

$$\dot{r}^2 = e^2 - 1 + \frac{2Mr^{n-1}}{r^n + L^n}. \quad (14)$$

For massive particles with $e^2 > 1$, the proper time to reach $r = 0$ is finite. Hence, we again need to extend the coordinate r to negative values (a plot of \dot{r}^2 as a function of r is given in Fig.2).

If n is even \dot{r}^2 is regular everywhere. However, where \dot{r}^2 is negative the probe particle does not have enough energy to overcome the potential barrier and it bounces back. On the other hand, for probe particles with energy large enough \dot{r}^2 is always positive, and the geodesics can be extended to $r = -\infty$. Since the metric (13) is asymptotically flat, the proper time to reach $r = -\infty$ is infinite. One can also show that the affine parameter to approach $r = -\infty$ diverges for massless particles. Hence, the spacetime with (13) is complete for massive as well as massless particles.

If n is odd, \dot{r}^2 has a pole at $r = -L$, and close to the pole, the spacetime structure is similar to Schwarzschild's spacetime in $r = 0$. Thus massive particles can reach the pole in a finite amount of proper time. Since these geodesics end in $r = -L$ the spacetimes with n odd are geodesically incomplete.

To summarize, the spacetime causal structure of the metrics (13) is shown in Fig. 3. For the case with odd power n , there are three roots for $f(r) = 0$, denoted as r_+ , r_- , r_+^0 , which correspond to three horizons. The region I is our asymptotically flat spacetime region and r_+




FIG. 3. Panel on the left: maximal extension of generalized Hayward's spacetime when n is an odd positive integer. It deserves to be notice that the diagram is also infinitely extendible in both the left and right directions. Panel on the right: maximal extension of generalized Hayward's spacetime when n is an even positive integer.

is the event horizon. Contrary to the Reissner-Nordström spacetime, the metric has a regular core in $r = 0$ that forces us to extend the spacetime to $r < 0$. However, it can not be extended beyond the region IV because the geodesics end at the singularity located in $r = -L$. For the case with even power n , $f(r) = 0$ has two positive roots, denoted as r_+ and r_- in Fig 3. The region I and III, namely $-\infty < r < r_-$ and $r_+ < r < \infty$, are both asymptotically flat, while the region II interconnects two asymptotic regions of the Universe. The spacetime is geodesically complete for massless as well as massive particles.

IV. A GEODETICALLY INCOMPLETE BLACK HOLE WITH MINKOWSKI CORE

In this section we focus on the regular non-analytic black hole proposed by Simpson and Visser in [3], and defined in terms of the following function,

$$f(r) = 1 - \frac{2M}{r}e^{-a/r}. \quad (15)$$

In the $r \rightarrow 0^+$ limit, the function $f(r)$ reads $f(r) \sim 1 - O(r^2)$, and the black hole metric has an asymptotic Minkowski core where the curvature invariants vanish. However, the metric in $r = 0$ is not continuous as it is easy to check computing the limit $r \rightarrow 0^-$ of the function (15) (see Fig. 4).

Let us now check if the metric is geodesically complete, namely if the proper time to reach the Minkowski's core in $r = 0$ is infinity. The geodesic equation of motion for radially falling massive particles reads:

$$\dot{r}^2 = e^2 - 1 + \frac{2M}{r}e^{-a/r}, \quad (16)$$

while the effective potential $V = f(r)$ has been plotted in Fig. 4. When the dimensionless constant $e^2 < 1$ the




FIG. 4. We here plot the function $f(r)$ for the case of $a < 2M/e$ (solid line), and $a = 2M/e$ (dashed line). In both cases $f(r)$ is not analytic in $r = 4$. We fixed the mass to the value $M = 5$.




FIG. 5. The Penrose diagram for the maximal extension of the metric (15).

probe particles cannot reach $r = 0^+$ and will bounce back. On the other hand for probe particles with $e^2 = 1$ we have $d\tau \propto r e^{a/r} dr$, and the proper time to reach $r = 0^+$ turns out to be infinite. Finally, for $e^2 > 1$, $d\tau/dr$ tends to the constant $e^2 - 1$ when $r \rightarrow 0^+$. Therefore, the latter probe particles can reach the Minkowski core in finite proper time. Since the geodesic equation (16) (see also Fig. 4) is ill-defined in $r = 0$, the geodesics that end in $r = 0$ can not be extended to negative values of r . Hence, it turns out that the metric (15) is actually geodesically incomplete whether we assume the the metric to be defined by the function (15) also for negative values of r .

Let us end this section studying the spacetime causal structure of the metric (15), which is closely related to the value of parameter a present in (15). For the case of $a < 2M/e$ (here e is the Euler's number that should not be confused with the dimensionless energy e), the equation




FIG. 6. The function $f(r)$ for the case $a < \sqrt{2/e}M$ (solid line), and $a = \sqrt{2/e}M \approx 0.858M$ (dashed line). In both plots $M = 5$.

$f(r) = 0$ have two solutions, usually denoted as r_+ and r_- . Therefore, the spacetime structure in this case is similar to Reissner-Nordström spacetime. For the case of $a = 2M/e$, the two horizons merge together, namely, $r_+ = r_- = 2M/e$. Finally, for $a > 2M/e$, there is no horizon and the metric does not describe a black hole, but a kind of soliton. The Penrose diagram for the case $a < 2M/e$ and for the extreme case $a = 2M/e$ are given in Fig. 5.

V. GEODETICALLY COMPLETE BLACK HOLE WITH MINKOWSKI CORE

In order to avoid the the geodesic incompleteness of the metric (15), we propose a slight modification to the Simpson-Visser's metric, namely

$$f(r) = 1 - \frac{2M}{r} e^{-a^2/r^2}. \quad (17)$$

The metric tends to the Schwarzschild's metric for large values of the radial coordinate r and has a Minkowski core for $r \rightarrow 0^+$. Likewise the Simpson-Visser's metric, probe massive particles can reach $r = 0$ in finite proper time, and we are again forced to make a continuation of the spacetime beyond $r = 0$. However, contrary to the metric (15), the improved metric (17) is continuous in $r = 0$, and the extension of the spacetime to negative r is doable.

A plot of the function $f(r)$ in (17) is provided in Fig. 6, which is similar to the generalized Hayward metric (13) for even powers of the integer n . Moreover, the spacetime described by the metric (17) is geodesically complete because the proper-time for massive particles to reach the edges at $r = \pm\infty$ of the spacetime is infinite.

The spacetime structure of the metric (17) is given in Fig. 7. For the case of $a^2 < 2M^2/e$, there are two




FIG. 7. The Penrose diagram for the maximally extended metric (17).

horizons, denoted by r_+ and r_- . The regions $r = \pm\infty$ represent the causal infinity of the region I and the region III respectively. The Minkowski core in $r = 0$, which is located inside the region III, is regular. For the extreme case, namely $a^2 = 2M^2/e$, the two horizons r_+ and r_- coincide, and the spacetime consists of two asymptotically flat regions.

A more general class of geodesically complete modifications of the Simpson-Visser spacetime looks like:

$$f(r) = 1 - \frac{2M}{r} e^{-\frac{a^{2n}}{r^{2n}}}, \quad n \in \mathbb{N}, \quad n \geq 1. \quad (18)$$

The continuation of the metric (1) with the function (18) assumes the same form of $f(r)$ for positive as well as negative values of the radial coordinate r . However, if we assume $f(r)$ to be a C^∞ smooth function the spacetime metric defined in terms of (18) is not unique because the function $f(r)$ is non-analytic at $r = 0$ since any derivative of $f(r)$ is zero in $r = 0$. Therefore, as an even simpler choice we could just extend the metric for $r \in (-\infty, 0)$ to be exactly Minkowski. Notice that the latter extension to Minkowski works also for the Simpson-Visser's metric (15), which in this way turns out to be geodesically complete. In other words, the simplest extension of the Simpson-Visser metric (consisting in taking the function (15) also for $r < 0$) is not geodesically complete, but its extension to Minkowski or to other smooth functions will be consistent with the geodesic completion.

Finally, we make a comment about the uniqueness issue. Indeed, multiple (in principle infinity) extensions of the metric imply multiple evolutions for point-like particles beyond $r = 0$, which boils down to the loss of predic-

tivity. In other words, $r = 0$ turns out to be a bifurcation or multifurcation point once the metric is fixed for $r > 0$ because of the ambiguous extension to negative values of the radial coordinate. However, this is not a problem related to the geodesic principle, but to the uniqueness of the metric. In this respect will be crucial to have a theory of which one of the regular spacetimes proposed in this section will be an exact solution.

VI. CONCLUSIONS

In this paper, we focused on the geodesic completeness of some popular singularity-free black holes whose regularity is based on the curvature invariants. We showed that the well known Hayward metric (10) and the simplest extension of the non-analytic metric with Minkowski core proposed by Simpson and Visser (15) are both geodesically incomplete. Hence, such spacetimes are actually pathological though $r = 0$ is a smooth core. Thus, we proposed a generalization of both the metrics of above and showed that the modified spacetimes are geodesically complete. Although, the proposed complete spacetimes are a simple extension of the Hayward's and Simpson-Visser's metrics, we think they need an analysis of the thermodynamic properties that we have not addressed in this paper. In particular, the gravitational collapse should be rethought in light of the causal structure derived in this article.

Similarly, one can show the geodesic completeness of other popular regular black holes. For example, the *Bardeen's black hole* [1], the *Nicolini-Samilagic-Spallucci's black hole* [5–8], the *Lee-Wick black hole* [11], and the *loop black hole* [9, 10] and other singularity-free spacetimes found in conformal gravity are geodesically complete [12]. As a further feature and contrary to the other regular black holes, conformal black holes, which are a prediction of local [13, 14] and nonlocal [15–19] higher derivative theories, do not show any Cauchy horizon (if not present in the solution of Einstein's theory), usually unstable [20] in the Einstein's theory of gravity.

ACKNOWLEDGMENTS

We are grateful to Daniel Terno and Cosimo Bambi for the useful comments and questions. This work has been supported by the Basic Research Program of the Science, Technology, and Innovation Commission of Shenzhen Municipality (grant no. JCYJ20180302174206969).

[1] J. M. Bardeen, in “Conference Proceedings of GR5 (Tbilisi, USSR, 1968)”, p. 174.

[2] S. A. Hayward, “Formation and evaporation of regular black holes,” *Phys. Rev. Lett.* **96**, 031103 (2006). [arXiv:gr-qc/0506126 [gr-qc]].

- [3] A. Simpson and M. Visser, “Regular black holes with asymptotically Minkowski cores,” *Universe* **6**, no.1, 8 (2019).
- [4] A. Bonanno and M. Reuter, “Renormalization group improved black hole space-times,” *Phys. Rev. D* **62**, 043008 (2000); A. Bonanno and M. Reuter, “Spacetime structure of an evaporating black hole in quantum gravity,” *Phys. Rev. D* **73**, 083005 (2006).
- [5] P. Nicolini, A. Smailagic and E. Spallucci, “Noncommutative geometry inspired Schwarzschild black hole,” *Phys. Lett. B* **632**, 547 (2006), gr-qc/0510112.
- [6] L. Modesto, J. W. Moffat and P. Nicolini, “Black holes in an ultraviolet complete quantum gravity,” *Phys. Lett. B* **695**, 397-400 (2011) [arXiv:1010.0680 [gr-qc]].
- [7] N. Burzillà, B. L. Giacchini, T. d. Netto and L. Modesto, “Higher-order regularity in local and nonlocal quantum gravity,” *Eur. Phys. J. C* **81**, no.5, 462 (2021) [arXiv:2012.11829 [gr-qc]].
- [8] B. L. Giacchini and T. de Paula Netto, “Effective delta sources and regularity in higher-derivative and ghost-free gravity,” *JCAP* **07**, 013 (2019) [arXiv:1809.05907 [gr-qc]].
- [9] L. Modesto, “Semiclassical loop quantum black hole,” *Int. J. Theor. Phys.* **49**, 1649-1683 (2010) [arXiv:0811.2196 [gr-qc]].
- [10] L. Modesto and I. Premont-Schwarz, “Self-dual Black Holes in LQG: Theory and Phenomenology,” *Phys. Rev. D* **80**, 064041 (2009) [arXiv:0905.3170 [hep-th]].
- [11] C. Bambi, L. Modesto and Y. Wang, *Phys. Lett. B* **764**, 306-309 (2017) doi:10.1016/j.physletb.2016.11.060 [arXiv:1611.03650 [gr-qc]].
- [12] C. Bambi, L. Modesto and L. Rachwał, “Spacetime completeness of non-singular black holes in conformal gravity,” *JCAP* **05**, 003 (2017) [arXiv:1611.00865 [gr-qc]].
- [13] L. Modesto and I. L. Shapiro, “Superrenormalizable quantum gravity with complex ghosts,” *Phys. Lett. B* **755**, 279-284 (2016) [arXiv:1512.07600 [hep-th]].
- [14] L. Modesto, “Super-renormalizable or finite Lee–Wick quantum gravity,” *Nucl. Phys. B* **909**, 584-606 (2016) [arXiv:1602.02421 [hep-th]].
- [15] L. Modesto, “Super-renormalizable Quantum Gravity,” *Phys. Rev. D* **86**, 044005 (2012) [arXiv:1107.2403 [hep-th]].
- [16] L. Modesto and L. Rachwał, “Nonlocal quantum gravity: A review,” *Int. J. Mod. Phys. D* **26**, no.11, 1730020 (2017)
- [17] L. Modesto and L. Rachwał, “Super-renormalizable and finite gravitational theories,” *Nucl. Phys. B* **889**, 228-248 (2014) [arXiv:1407.8036 [hep-th]].
- [18] L. Modesto, “The Higgs mechanism in nonlocal field theory,” *JHEP* **06**, 049 (2021) [arXiv:2103.05536 [hep-th]].
- [19] L. Modesto, “Nonlocal Spacetime-Matter,” [arXiv:2103.04936 [gr-qc]].
- [20] E. Poisson and W. Israel, “Internal structure of black holes,” *Phys. Rev. D* **41**, 1796-1809 (1990); A. Ori, “Inner structure of a charged black hole: An exact mass-inflation solution,” *Phys. Rev. Lett.* **67**, 789-792 (1991).