

# Presymplectic BV-AKSZ formulation of Conformal Gravity

Ivan Dneprov and Maxim Grigoriev

*Lebedev Physical Institute,  
Leninsky ave. 53, 119991 Moscow, Russia*

*Institute for Theoretical and Mathematical Physics,  
Lomonosov Moscow State University, 119991 Moscow, Russia*

## Abstract

We elaborate on the presymplectic BV-AKSZ approach to local gauge theories and apply it to conformal gravity. More specifically, we identify a compatible presymplectic structure on the minimal model of the total BRST complex of the theory and show that these structures determine a full-scale BV formulation for a specific frame-like action which seems to be not known before. Remarkably, the underlying frame-like description does not require any artificial off-shell constraints. Instead, the action becomes equivalent to the usual conformal gravity one upon gauging away all the pure gauge variables associated to the kernel of the presymplectic structure. Finally, we show how the presymplectic BV-AKSZ formalism extends to generic gauge theories.

# Contents

<b>1</b>	<b>Introduction</b>	<b>3</b>
<b>2</b>	<b>Presymplectic BV-AKSZ formulation</b>	<b>5</b>
2.1	AKSZ sigma models . . . . .	5
2.2	Presymplectic BV-AKSZ formulation . . . . .	7
2.3	Target space and the total BRST complex . . . . .	10
<b>3</b>	<b>Presymplectic AKSZ form of conformal gravity</b>	<b>11</b>
3.1	Conformal gravity . . . . .	11
3.2	Frame like formulation . . . . .	12
3.3	Jet-bundle BRST complex for CGR . . . . .	15
3.4	Minimal model of the BRST complex for CGR . . . . .	16
3.5	Presymplectic structure . . . . .	20
3.6	Symplectic quotient and BV formulation . . . . .	22
<b>4</b>	<b>General presymplectic BV-AKSZ formalism</b>	<b>25</b>
4.1	Gauge PDEs . . . . .	26
4.2	Vertical forms . . . . .	27
4.3	Presymplectic gauge PDEs . . . . .	28
4.4	Symplectic quotient and BV formulation . . . . .	29
4.5	Example: Maxwell . . . . .	30
<b>A</b>	<b>Tensor conventions</b>	<b>32</b>
<b>B</b>	<b>Invariance of the presymplectic structure</b>	<b>34</b>
<b>C</b>	<b>The kernel of the presymplectic structure</b>	<b>36</b>
<b>D</b>	<b>Metric elimination</b>	<b>39</b>

# 1 Introduction

It is well known that AKSZ sigma models [1] with finite dimensional target spaces are topological (=no local degrees of freedom). However if one replaces target space symplectic structure with possibly degenerate presymplectic one, one arrives [2] at an elegant AKSZ-like representations for the familiar 1st order Lagrangians of gauge theories and makes manifest the graded geometry underlying the frame-like formulations of gravity and higher spin fields.

It has been later realised that the target space of the presymplectic AKSZ formulation arises as an equivalent reformulation of the total BRST complex of the local gauge theory in question while the presymplectic structure can be seen as a BRST extension of the well-known on-shell presymplectic structure [3, 4, 5] (also known as the presymplectic current). Moreover, the presymplectic current itself can be seen [6] as the decent completion of the Batalin-Vilkovisky (BV) symplectic structure.

More recently, it became clear [7], that not only the Lagrangian but also the entire BV formulation [8] of a given local gauge field theory can be encoded in the presymplectic AKSZ formulation.<sup>1</sup> In particular, the target space presymplectic structure defines a presymplectic structure on the space of AKSZ fields. Taking the symplectic quotient results in the BV symplectic structure which together with the BV-BRST differential determines BV master action. This motivates that the presymplectic BV-AKSZ framework can be considered as a rather flexible and general extension of the usual BV-BRST approach. In this approach a Lagrangian local gauge theory is described in terms of fields, which can be subject to differential constraints. For instance, a formulation of the Lagrangian system in terms of the stationary surface (equation manifold in the PDE theory terminology) is a typical example of such a description.

When applied to Lagrangian gauge theories extending gravity this approach gives an interesting interpretation of the theories (and their underlying geometry, e.g. Cartan geometry) in terms of the differential graded (pre)symplectic ge-

---

<sup>1</sup>Mention also an alternative AKSZ approach [9] to general relativity based on space+time decomposition.

ometry. It is also worth mentioning recent application of presymplectic AKSZ approach to constructing Lagrangian description of interacting higher spin theories [10].

In this work we apply the full-scale presymplectic AKSZ approach to conformal gravity. The theory has been extensively studied [11, 12, 13, 14, 15], and has interesting properties, for instance it is renormalizable although non-unitary. We concentrate on the minimal version (also known as minimal model) of the jet-bundle BV-BRST complex of the theory, which can be obtained via additional reduction of the one initially put forward in [16, 17, 18] (see also [19]), and present a presymplectic structure therein which defines a full-scale BV-AKSZ description of the theory. This gives a new formulation of conformal gravity in terms of the connection of the conformal algebra, where the field content and gauge symmetries are fully determined by the underlying presymplectic  $Q$ -manifold and no artificial constraints (like the familiar torsion-free constraint of the formulation [11]) are needed (see also [20] for the discussion of frame-like formulations of conformal gravity)). In this aspect our formulation can be considered natural and arising from first principles.

The graded geometry structures identified in this work give a new perspective on the underlying conformal geometry and more generally Cartan geometry. The formulation itself can be useful in the analysis of the respective QFT as it naturally comprises the respective BV formulation and the local BRST cohomology complex.

Finally we present an extension of the presymplectic BV-AKSZ formulation to the case of not necessarily diffeomorphism-invariant systems, where the underlying  $Q$ -manifold is not a product of the AKSZ source and the target and one has to resort to more general object known as gauge PDE [21] (see also [22, 23] for earlier but less geometrical description), which can be formulated as, in general, a nontrivial  $Q$ -bundle [24] over the shifted tangent bundle over the space-time.

## 2 Presymplectic BV-AKSZ formulation

### 2.1 AKSZ sigma models

From the very beginning the AKSZ approach [1] has been formulated for Lagrangian theories. Although its extension to gauge systems at the level of equations of motion is straightforward and amounts to forgetting the symplectic structure and rephrasing everything in terms of the BRST differential rather than BV master action it is instructive to start with this more general case.

By AKSZ sigma model at the level of equations of motion we mean a pair of  $\mathbb{Z}$ -graded  $Q$ -manifolds, one of which,  $(T[1]X, d_X)$ , where  $d_X$  is the de Rham differential of  $X$  seen as a  $Q$  structure on  $T[1]X$ , is regarded as source (i.e.  $X$  is a genuine space-time) and another one  $(F, q)$  as the target. The  $\mathbb{Z}$ -grading is denoted as  $\text{gh}()$  in what follows. In particular, for  $(T[1]X, d_X)$  the ghost degree is just a form-degree.

Fields of the model are maps from the source to the target. It is convenient to introduce a trivial fiber bundle  $(E, Q) = (T[1]X, d_X) \times (F, q)$  (here we mean product of  $Q$ -manifolds, i.e.  $Q = q + d_X$  in the adapted coordinates and regard fields as section  $T[1]X \rightarrow E$ . Recall that such maps are by definition homomorphisms of the algebras of functions and, in contrast to supermaps, they preserve the degree). The generalization to the case where such a bundle is trivial only locally (but still  $Q = q + d_X$ ) is straightforward. More general situation is discussed in Section 4.

If  $\sigma : T[1]X \rightarrow E$  denote a field configuration then the equations of motion read as

$$d_X \circ \sigma^* = \sigma^* \circ Q. \quad (2.1)$$

The infinitesimal gauge variation of the configuration  $\sigma$  is given by

$$\delta\sigma^* = d_X \circ \xi^* + \xi^* \circ Q \quad (2.2)$$

where  $\xi^* : \mathcal{C}^\infty(E) \rightarrow \mathcal{C}^\infty(T[1]X)$  is a gauge parameter map, which has a degree  $-1$  and satisfies

$$\xi^*(fg) = (\xi^*(f))\sigma^*(g) + (-1)^{|f|}\sigma^*(f)\xi^*(g). \quad (2.3)$$

In a similar way one defines gauge for gauge symmetries. The above is just a coordinate-free reformulation of the equations of motion and gauge symmetries encoded in the BRST differential of the AKSZ sigma models. Details can be found in [21].

A remarkable feature of the AKSZ construction is that the full-scale BV formulation (at the level of equations of motion) is also encoded in the AKSZ data. More precisely, the space of fields, ghosts, and antifields is given by super-maps from  $T[1]X$  to  $F$  (super-sections of  $E$ ) so that the space of sections is recovered as the degree 0 component (the body) of the space of super-sections. The coordinates (fields and antifields) on the space of supermaps can be introduced as:

$$\widehat{\sigma}^*(\psi^A) = \psi^A(x) + \psi_{\mu}^A(x)\theta^{\mu} + \psi_{\mu\nu}^A(x)\theta^{\mu}\theta^{\nu} + \dots, \quad (2.4)$$

where  $\widehat{\sigma}^*$  is a pullback map induced by super-section  $\widehat{\sigma}$  and we made use of local coordinates  $\psi^A$  on  $M$  and the adapted local coordinates  $x^{\mu}, \theta^{\mu}$  induced by local coordinates  $x^{\mu}$  on  $X$ . Note that form-degree  $k$  components  $\psi_{\mu_1 \dots \mu_k}^A$  carry ghost degree  $\text{gh}(\psi^A) - k$ . The space of maps is recovered by setting to zero all the coordinates of nonvanishing degree.

In the field-theoretical terms the BRST differential of the AKSZ sigma model can be written as:

$$s = \int d^n x d^n \theta (q^A(\psi(x, \theta)) + d_X \psi^A(x, \theta)) \frac{\delta}{\delta \psi^A(x, \theta)}, \quad (2.5)$$

where  $n = \dim X$ .

If in addition to  $Q$ -structure the target space is equipped with the compatible symplectic structure  $\omega$  of degree  $\text{gh}(\omega) = n - 1$ , i.e.  $\omega$  is nondegenerate,  $d\omega = 0$  and  $L_q\omega = 0$  then the theory described by the AKSZ model is Lagrangian. More specifically, introducing  $H$  via  $dH = i_q\omega$  the action can be written as:

$$S[\sigma] = \int_{T[1]X} [\sigma^*(\chi))(d_X) + \sigma^*(H)], \quad (2.6)$$

where  $\sigma$  denotes a section:  $T[1]X \rightarrow E$  and we have introduced one form presymplectic potential  $\chi$  such that  $d\chi = \omega$ .

It turns out that just like the BRST differential the BV master action can also be immediately written just in terms of  $q, \omega, d_X$ . More precisely, it is given by

$$S[\widehat{\sigma}] = \int_{T[1]X} [\widehat{\sigma}^*(\chi))(d_X) + \widehat{\sigma}^*(H)], \quad (2.7)$$

i.e. has the same form as the classical action but with  $\sigma$  replaced with  $\widehat{\sigma}$ . It satisfies the master equation with respect to the odd Poisson bracket of degree 1 induced by  $\omega$  on the space of super-sections [1].

If the target space is finite-dimensional the AKSZ sigma model is topological (for  $n > 1$ ). However, generic diffeomorphism-invariant gauge theories can be represented as AKSZ sigma models at the price of considering infinite-dimensional targets. For instance taking as  $M$  a fiber of the jet-bundle of BV-BRST formulation for a diff-invariant gauge system and as  $q$  the restriction its BV-BRST differential to the fiber (in the diff-invariant case the BV-BRST formulation can be locally encoded in the fiber, see e.g. [25, 26, 22]) the respective AKSZ sigma model is equivalent to the initial gauge theory [22] (see also [23, 27] for the Lagrangian version). Various equivalent (AKSZ) formulations can be arrived at by equivalent reductions of the target  $Q$ -manifold.

## 2.2 Presymplectic BV-AKSZ formulation

It turns out that relaxing the nondegeneracy condition for  $\omega$  gives a natural generalization of AKSZ sigma models. By presymplectic AKSZ sigma model we mean the same data  $(T[1]X, d_X)$  and  $(F, q)$ , where  $F$  is equipped with not necessarily invertible 2-form  $\omega$  satisfying  $\text{gh}(\omega) = n - 1, L_q\omega = 0, d\omega = 0$ .

An important difference with the usual AKSZ is that we do not necessarily require that the underlying gauge system is determined by the BRST differential  $s$  given by (2.5). Instead, the gauge system is determined by the action functional (2.6), where some of the fields (on which the action doesn't really depend) are gauged away. It turns out that a variety of gauge theories can be represented in this way [2].

To make the definition of a presymplectic BV-AKSZ formulation precise let us restrict for the moment to a coordinate patch  $U \subset X$  and use the adapted coordinates  $x^\mu, \theta^\mu, \psi^A$ . Then it is convenient to represent the space of supermaps

$\text{Smaps}(T[1]U, F)$  as  $\text{Maps}(U, \bar{F})$ , where  $\bar{F} = \text{Smaps}(\mathbb{R}^n[1], F)$ . A generic supermap from  $\bar{F}$  is given by:

$$\widehat{\sigma}^*(\psi^A) = \sum_{l=0}^n \frac{1}{l!} \psi_{\mu_1 \dots \mu_l}^A \theta^{\mu_1} \dots \theta^{\mu_l}, \quad (2.8)$$

so that as coordinates on  $\bar{F}$  one can take  $\psi_{\mu_1 \dots \mu_l}^A$ ,  $l = 0, \dots, n = \dim X$ . In what follows we use  $\Psi^I$  as a collective notation for the coordinates of  $\bar{F}$ . Applying the same construction to all the coordinate charts of  $X$  results in the locally-trivial fiber bundle  $\bar{E}$  over  $X$  with a typical fiber  $\bar{F}$ .<sup>2</sup>

The 2-form  $\omega$  on  $F$  defines a 2 form  $\bar{\omega}$  on  $\bar{F}$  in the standard way:

$$\bar{\omega} = \int d^n \theta \omega_{AB}(\psi(\theta)) d\psi^A(\theta) d\psi^B(\theta), \quad (2.9)$$

where  $d\psi^A(\theta)$  can be seen as a generating function for the differentials  $d\Psi^I$ . Note that this form transforms as a density on  $U$  under the change of  $x^\mu$ -coordinates. It defines a density on  $X$  with values in the 2-forms on the fiber, which is defined globally.<sup>3</sup>

By construction  $\bar{\omega}$  is closed and  $\text{gh}(\bar{\omega}) = -1$ . In addition we assume that  $\bar{\omega}$  is regular. This may involve taking as  $\bar{F}$  only those supermaps where the rank of  $\bar{\omega}$  is constant and maximal rank (this is exactly what one naturally does in the case of GR and, as we are going to see, in the case of conformal gravity). A regular closed two form defines an involutive distribution of its zero vectors, i.e. vector fields satisfying  $i_V \bar{\omega} = 0$ . It is the standard fact that the symplectic quotient (the space of the integrable submanifolds, which exists, at least locally) is naturally a symplectic manifold which we denote  $\bar{G}$  in what follows. Moreover, it can be realised (again, at least locally) as a submanifold transversal to the kernel distribution in which case the symplectic structure coincides with the restriction of  $\bar{\omega}$  to the submanifold. In this way we have arrived at the space of fields equipped with the odd symplectic structure of ghost degree  $-1$ , just like in the case of BV formalism.

To have a complete BV formulation, in addition to the symplectic structure  $\bar{\omega}$  on the space of fields we need a master-action satisfying master equation.

<sup>2</sup>Note that although  $E$  is assumed trivial now  $\bar{E}$  can be only locally-trivial if  $TX$  is not globally-trivial.

<sup>3</sup>Another possibility is to define an  $n + 2$  form  $\omega^{\bar{E}} = (dx)^n \bar{\omega}$  on the total space of  $\bar{E}$ .

To analyse this it is convenient to consider a jet bundle  $J^\infty(\bar{E})$ , where  $\bar{E} \rightarrow X$  is the introduced above fiber bundle with the typical fiber  $\bar{F}$ . The BV-like symplectic structure is then introduced in the standard way as an  $(n, 2)$  form

$$\omega^{pBV} = (dx)^n \bar{\omega}_{IJ} d_v \Psi^I d_v \Psi^J, \quad (2.10)$$

where we used standard local coordinates on  $J^\infty(\bar{E})$ , induced by trivialisation  $\bar{E}|_U = T[1]U \times \bar{F}$ . Note that this symplectic structure is defined globally on  $\bar{E}$ . More invariant description of the underlying geometry and its generalization is presented in section 4.

When written in terms of  $J^\infty(\bar{E})$  the integrand (over  $X$ ) of the BV-like action (2.7) reads as

$$L^{BV} = K + \bar{H},$$

$$K = (dx)^n \int d^n \theta \chi_A(\psi(\theta)) \theta^\mu D_\mu \psi^A(\theta), \quad \bar{H} = (dx)^n \int d^n \theta H(\psi(\theta)), \quad (2.11)$$

where  $D_\mu$  denote total derivatives on  $J^\infty(\bar{E})$ . It turns out that the restriction of  $L^{BV}$  to the jet sub-bundle associated to the symplectic quotient  $\bar{G}$  realized as a submanifold of  $\bar{F}$  satisfies all the conditions for the BV master action.

To see this let us first introduce a natural prolongation  $\bar{q}$  of the vector field  $q$  from  $F$  to  $\bar{F}$ :

$$\bar{q}(\psi^A(\theta)) = (q\psi^A)(\theta). \quad (2.12)$$

Note that  $\bar{q}$  is a Hamiltonian vector field on  $\bar{F}$  with Hamiltonian  $\bar{H}$ , i.e.  $i_{\bar{q}} \bar{\omega} = d\bar{H}$ . Now we can introduce a vertical evolutionary vector field  $s$ :

$$s = D + \bar{q}^{pr}, \quad (2.13)$$

where  $\bar{q}^{pr}$  denotes a natural prolongation of  $\bar{q}$  to  $\bar{E}$  defined by requiring  $[Q^{pr}, D_\mu] = 0$  and  $D$  is a vertical evolutionary vector field defined through  $D\psi^A(\theta) = \theta^\mu D_\mu(\psi^A(\theta))$ .

We have the following relations [7]:

$$i_D \omega^{pBV} = d_v K + d_h(\cdot), \quad i_D i_D \omega^{pBV} = d_h(\cdot), \quad (2.14)$$

as well as

$$i_D i_{\bar{q}^{pr}} \omega^{pBV} = d_h(\cdot). \quad (2.15)$$

which can be directly checked and amount to:

$$i_s \omega^{pBV} = d_v L^{BV} + d_h(\cdot), \quad i_s i_s \omega^{pBV} = d_h(\cdot). \quad (2.16)$$

The above relations remain intact when restricted to  $J^\infty(\bar{G} \times X)$  understood as a jet subbundle of  $\bar{E}$  (recall that we assumed that the symplectic quotient  $\bar{G}$  is realised as a submanifold of  $\bar{F}$ ), where by restriction of  $s$  we mean its projection induced by the projection  $\bar{F} \rightarrow \bar{G}$  (recall that there is a natural projection to the symplectic quotient). If by some abuse of notations  $s$  denotes its projection to  $J^\infty(\bar{G} \times X)$  the restricted relations read as:

$$i_s \omega^{BV} = d_v L^{BV} + d_h(\cdot), \quad i_s i_s \omega^{BV} = d_h(\cdot), \quad (2.17)$$

where  $\omega^{BV}$  is the restriction of  $\omega^{pBV}$  to  $J^\infty(\bar{G} \times X) \subset J^\infty(\bar{E})$ . Because  $\bar{G}$  is symplectic the above relation imply the classical master equation satisfied by  $L^{BV}$ .

To summarize we have shown how a presymplectic BV-AKSZ system gives rise to a genuine BV gauge system. The only subtle point is whether a resulting BV master action is a proper solution to the master equation. This has to be addressed separately.

## 2.3 Target space and the total BRST complex

A natural question is what is the origin of the target space  $Q$ -manifold. It turns out that it is directly related to the total BRST complex of the system.

Suppose we start with a jet-bundle BV formulation. The underlying bundle is  $\mathcal{E} \rightarrow X$ , i.e. coordinates on its fibres are fields, ghosts, ghosts-for ghosts, ..., as well as antifields (For the moment we do not assume the system to be Lagrangian and hence in general there is no natural pairing between fields and antifields). The jet-space BRST complex is given by local functions on  $J^\infty(\mathcal{E})$  with the differential being the BRST differential  $s$ , i.e. a degree 1 vertical evolutionary vector field on  $J^\infty(\mathcal{E})$ .

If the system in question is diffeomorphism-invariant (i.e. space-time reparametrizations are among the gauge symmetries encoded in  $s$ ), differential  $s$  does not depend on  $x$  and hence its restriction to a finer define a  $Q$ -structure (which we

keep denoting by  $s$ ) on a typical fiber  $F$  of  $J^\infty(\mathcal{E})$ . Moreover, locally on  $X$ ,  $Q$ -manifold  $(F, s)$  is equivalent to the total BRST complex of the system, i.e. the complex of horizontal forms on  $J^\infty(\mathcal{E})$  with the differential being  $\tilde{s} = d_h + s$  and the degree being ghost degree plus form degree.

It is known [22] that in diff-invariant case the AKSZ sigma model (at the level of equations of motion) with the target  $(F, S)$  (or its equivalent reduction) and source  $T[1]X$  gives an equivalent formulation of the gauge system in question. Furthermore, the compatible presymplectic structure  $\omega$  on  $F$  can be traced back [7] (see also [6]) to the initial BV symplectic structure extended to a cocycle of the total BRST differential  $\tilde{s}$ . More detailed and general exposition of the relation is given in Section 4.

## 3 Presymplectic AKSZ form of conformal gravity

### 3.1 Conformal gravity

Conformal gravity is a well-known theory of gravity with an additional gauge invariance compared to GR: local Weyl invariance. The action for such theory is given by

$$S[g] = \int d^4x W_{\mu\nu\rho\lambda} W^{\mu\nu\rho\lambda}, \quad (3.1)$$

where  $W_{\mu\nu\rho\lambda}$  is the trace-free part of the Riemann curvature  $R_{\mu\nu\rho\lambda}$ . The indices are raised and lowered by the metric  $g_{\mu\nu}$ . Weyl invariance implies that the action only depends on the conformal class of the metric  $[g_{\mu\nu}] = g_{\mu\nu}/\sim$ , where  $g_{\mu\nu} \sim g'_{\mu\nu} = e^{2\phi} g_{\mu\nu}$ . In field theoretical terms, the infinitesimal gauge symmetries of the theory are given by

$$\delta g_{\mu\nu} = \xi^\rho \partial_\rho g_{\mu\nu} + g_{\mu\rho} \partial_\nu \xi^\rho + g_{\rho\nu} \partial_\mu \xi^\rho + 2\omega g_{\mu\nu}, \quad (3.2)$$

where  $\xi^\mu$  and  $\omega$  are parameters of the diffeomorphisms and Weyl transformations respectively.

The equation of motion determined by  $S[g]$  read as:

$$B_{\mu\nu} = 0, \quad B_{\mu\nu} = P^{\rho\lambda} W_{\mu\rho\nu\lambda} + \nabla^\rho \nabla_\rho P_{\mu\nu} - \nabla^\rho \nabla_\mu P_{\nu\rho}, \quad (3.3)$$

where  $P_{\mu\nu}$  is the Schouten tensor. (See A for definitions.) Tensor  $B_{\mu\nu}$  is known as Bach tensor and it is Weyl invariant so that, as they should be, EOMs are diffeomorphism and Weyl invariant.

### 3.2 Frame like formulation

Just like in the case of Einstein gravity where Poincare algebra is the maximal symmetry of a vacuum solution and the theory itself can be reformulated in terms of the connection of Poincare algebra, the maximal symmetry of a solution to conformal gravity is the conformal algebra  $o(d, 2)$  and the theory itself is expected to have a formulation in terms of the connection of  $o(4, 2)$ .

We use the usual basis in  $o(4, 2)$  given by translations  $P_a$ , special conformal transformations  $K_a$ , Lorentz transformations  $J_{ab}$ , and dilation  $D$ , where the commutation relation read as:

$$\begin{aligned}
&= \eta_{bc}J_{ad} + \eta_{ad}J_{bc} - \eta_{ac}J_{bd} - \eta_{bd}J_{ac}, \\
[J_{ab}, P_c] &= \eta_{bc}P_a - \eta_{ac}P_b, & [J_{ab}, K_c] &= \eta_{bc}K_a - \eta_{ac}K_b, \\
[P_a, D] &= P_a, & [K_a, D] &= -K_a, \\
[K_a, P_b] &= -2(\eta_{ab}D + J_{ab}).
\end{aligned} \tag{3.4}$$

Here  $\eta_{ab}$  denotes a constant Minkowski metric of signature if almost positive signature.

Connection 1-form  $A$  decomposes as:

$$A = e^a P_a + \omega^{ab} J_{ab} + f^a K_a + \lambda D \tag{3.5}$$

and the curvature as:

$$\Omega = T^a P_a + R^{ab} J_{ab} + S^a K_a + \Lambda D. \tag{3.6}$$

Here and in what follows we systematically omit the wedge product symbol.

The action of the conformal gravity in terms of connection of the conformal group can be written in the following form [11]:

$$S[e, f, \omega, \lambda] = \int R_{ab} R_{cd} \epsilon^{abcd}, \tag{3.7}$$

where

$$R_{ab} = \overset{(0)}{R}_{ab} + (e_a f_b - e_b f_a) \quad (3.8)$$

is the Lorenz sector part of the full conformal curvature and

$$\overset{(0)}{R}_{ab} = d\omega_{ab} + \omega_a^c \omega_{cb} \quad (3.9)$$

is the usual curvature of the Lorentz connection, and  $\epsilon^{abcd}$  is an invariant totally antisymmetric tensor and  $\epsilon^{0123} = 1$ .

Here we do not discuss global space-time geometry and hence consider Lagrangians modulo total derivatives. In particular the Gauss-Bonnet term  $\overset{(0)}{R}_{ab} \overset{(0)}{R}_{cd} \epsilon^{abcd}$  can be omitted and the action can be rewritten as:

$$\begin{aligned} S &= \int (2 \overset{(0)}{R}_{ab} (e_c f_d - e_d f_c) + (e_a f_b - e_b f_a)(e_c f_d - e_d f_c) \epsilon^{abcd} = \\ &= \int (4 \overset{(0)}{R}_{ab} e_c f_d + 4 e_a f_b e_c f_d) \epsilon^{abcd} = \\ &= 4 \int (d\omega_{ab} e_c f_d + \omega_a^k \omega_{kb} e_c f_d + e_a f_b e_c f_d) \epsilon^{abcd}. \end{aligned} \quad (3.10)$$

It turns out [11] that action (3.10) is equivalent through elimination of the auxiliary fields to the standard conformal gravity action in the metric formulation (3.1), provided one in addition imposes the torsion constraint

$$T_a = de_a + \omega_a^b e_b + e_a \lambda = 0 \quad (3.11)$$

is assumed. This constraint is in fact algebraic (allows to uniquely express  $\omega_{ab}$  in terms of  $e_a$  and  $\lambda$ ) and hence the system remains Lagrangian. Now we briefly recall how this works. Technical details can be found in Appendix A

Let us consider first equations for  $f_d$ :

$$\epsilon^{abcd} R_{ab} e_c = 0. \quad (3.12)$$

The curvature 2-form can be expanded as  $R_{ab} = R_{abcd} \xi^c \xi^d$ . Then equation (3.12) implies  $R^c{}_{bcd} = 0$  which in turn gives:

$$f = -\frac{1}{6} \overset{(0)}{R} \quad (3.13)$$

and

$$f_{b,\mu} = -\frac{1}{2}(e^d{}_{,\mu} R^c{}_{bcd} - \frac{1}{6}e_{b,\mu} R), \quad (3.14)$$

where  $f = f^a{}_\mu e_a{}^\mu$  and  $R = R^a{}_{bac} \eta^{bc}$ . One concludes that on-shell  $f_{a,\mu}$  is an auxiliary field which coincides with the Schouten tensor (expressed through the frame field  $e^a{}_\mu$ ) on shell and can be eliminated.

Taking into account  $T_a = 0$ , the Bianchi identities  $d\Omega + [A, \Omega] = 0$  for the entire the torsion sector

$$e_c R^c{}_a + e_a \Lambda = 0, \quad (3.15)$$

implies (see Appendix A for details) that the curvature in the sector of dilations vanishes

$$\Lambda = d\lambda + e_a f^a = 0. \quad (3.16)$$

Let us turn our attention to gauge transformations. Consider the following transformations:

$$\delta A = d\alpha + [A, \alpha], \quad (3.17)$$

under which  $\Omega$  transforms as:

$$\delta\Omega = [\Omega, \alpha]. \quad (3.18)$$

Note in general these are not gauge symmetries of the action. It turns out that for  $\alpha = b^a K_a$ , i.e. for  $\alpha$  non-vanishing only in the sector of "special conformal transformations", this transformation is a gauge symmetry of the action thanks to  $T^a = 0$ . These gauge transformations are algebraic in the sector of  $\lambda$  (dilation part of the connection):

$$\delta\lambda = -e^a b_a. \quad (3.19)$$

It follows  $\lambda$  is pure gauge and can always be set to zero by a gauge transformation. In this gauge the torsion free condition (3.11) implies that  $\omega$  is just the Levi-Civita connection.

To summarize: action (3.7) supplemented with the torsion constraint (3.11) is equivalent to the Weyl action through the elimination of generalized auxiliary fields (auxiliary fields + algebraic pure gauge fields). The equivalent reduction boils down to the following steps:

1. Elimination of the auxiliary field  $f^a{}_\mu$
2. Algebraically solving zero torsion condition for the Lorentz connection,  $\omega = \omega(e, \lambda)$
3. Using gauge symmetry (3.17) with  $\alpha = b^a K_a$ , which is algebraic in the sector of  $\lambda_\mu$ , to set  $\lambda_\mu = 0$ . In this way we arrived to action (3.1) modulo a topological term but expressed in terms of the frame field  $e^a{}_\mu$  rather than metric.
4. Using gauge symmetry (3.17) with  $\alpha = k^a J_{ab}$  to eliminate all the components of the frame field but the metric,<sup>4</sup> giving the initial (3.1) in terms of the metric.

### 3.3 Jet-bundle BRST complex for CGR

As we recalled in Section 2.3 the target space of the AKSZ formulation of a given diff.-invariant system originates from a fiber of the jet-bundle BV-BRST formulation of the system. Now we recall this formulation for CGR in the metric-like form and construct a particularly useful minimal model for the respective total BRST complex.

Let us restrict to local analysis and introduce the respective structures using local coordinates. The bundle  $E$  is that of the metric  $g_{\mu\nu}(x)$ , ghosts  $\xi^\mu(x)$  associated to the diffeomorphism parameters and ghost  $\lambda(x)$ , associated to the parameter of Weyl rescalings. The associated coordinates on the jet-bundle  $J^\infty(E)$  read as:

$$x^\mu \quad g_{\mu\nu|\alpha\dots}, \quad \xi^\mu|_{\alpha\dots}, \quad \lambda|_{\alpha\dots}. \quad (3.20)$$

In these coordinates total derivatives  $D_a$  have the standard form, e.g.  $D_c g_{ab} = g_{ab|c}$ .

---

<sup>4</sup>This step is analogous to the elimination of the metric from the CGR BRST complex and strictly speaking might require assumptions. See Appendix D.

The gauge part  $\gamma$  of the BRST differential is determined by

$$\begin{aligned}\gamma g_{\mu\nu} &= \xi^\rho g_{\mu\nu|\rho} + \xi^\rho|_\mu g_{\rho\nu} + \xi^\rho|_\nu g_{\rho\mu} - 2\lambda g_{\mu\nu}, \\ \gamma \xi^\mu &= \xi^\rho \xi^\mu|_\rho, \\ \gamma \lambda &= \xi^\rho \lambda|_\rho,\end{aligned}\tag{3.21}$$

together with the conditions  $[D_\mu, \gamma]$  and  $\gamma x^\mu = 0$  (i.e. it is a vertical and evolutionary vector field).

The total BRST complex is given by the local horizontal forms on  $J^\infty(E)$  (which, as before, we identify with functions on  $J^\infty(E)$  pulled back to  $T[1]X$ ). In local coordinates these are simply functions on a new variables  $\theta^\mu$ , i.e. coordinate on the fibers of  $T[1]X$ , and coordinates (3.20). The total BRST differential reads as

$$\tilde{\gamma} = \gamma + d_h, \quad d_h = \theta^\mu D_\mu\tag{3.22}$$

Its cohomology in form degree  $n + k$ ,  $k \geq 0$  are known to be locally isomorphic to the cohomology of  $\gamma$  in the space of local functionals.

The above complex is the total BRST complex for off-shell CGR, i.e. CGR without equations of motion imposed. Its on-shell version is obtained by replacing  $J^\infty(E)$  by its subbundle  $\mathcal{E}$  determined by the equations of motion  $B_{\mu\nu} = 0$  and their prolongations  $D_{(\alpha)} B_{\mu\nu} = 0$ . A possible alternative is to introduce antifields and to work in terms of the complete BRST differential  $s = \gamma + \delta$ .

Because CGR is diff invariant the total BRST complex is locally equivalent to that of local functions on the typical fiber  $F$  of  $\mathcal{E}$  equipped with  $\gamma$  restricted to the fiber. Moreover, the AKSZ sigma model with the source  $T[1]X$  and target  $(F, \gamma)$  is locally equivalent [22] to the CGR (at the level of EOMs).

### 3.4 Minimal model of the BRST complex for CGR

We are interested in the presymplectic BV-AKSZ formulation that is natural in the sense that it involves a minimal number of fields (among all equivalent formulations of this type). In particular, the resulting formulation can be regarded as a canonical one (of course there remains a freedom of field redefinition). To this end it is natural to consider as a target space a minimal  $Q$ -manifold that is equivalent to the  $(F, \gamma)$  in the sense of natural equivalence of  $Q$ -manifolds. It

is known [22] that AKSZ sigma models (at the level of equations of motion) based on equivalent target  $Q$ -manifolds are equivalent in the sense of elimination of generalized auxiliary fields. Moreover, minimal  $Q$ -manifold corresponds to frame-like formulation of the underlying model. Minimal models for various local gauge theories were discussed from different perspectives in e.g. [28, 26, 22, 29] but the idea and somewhat implicit version appeared already in [30, 31, 32].

Let us recall the notion of equivalence for  $Q$ -manifolds. First of all we need the following:

**Definition 3.1.** *A contractible  $Q$ -manifold is a  $Q$ -manifold of the form  $(T[1]V, d_V)$ , where  $V$  is a graded linear space understood as a graded manifold and  $d_V$  is a canonical differential (De Rham) on  $T[1]V$ .*

For our present purposes it is enough to consider a rather restrictive version of equivalence: a  $Q$ -manifold  $(N, q)$  is called an equivalent reduction of  $(F, \gamma)$  if  $(F, \gamma) = (N, q) \times (T[1]V, d_V)$ , i.e.  $F$  is a product  $N \times T[1]V$  and the  $\gamma$  is a product  $Q$ -structure. This generates the equivalence relation for  $Q$ -manifolds.

It follows from the definition of the equivalent reduction that there exists independent functions  $w^a, v^a$  on  $F$  such that  $\gamma w^a = v^a$  and  $(N, q)$  can be identified as a  $Q$ -submanifold determined by  $w^a = 0$  and  $v^a = 0$ . Indeed, as  $w^a, v^a$  one can take linear coordinates on  $T[1]V$  pulled back to  $F$  by the canonical projection  $F \rightarrow T[1]V$ . Moreover, one can introduce a local coordinate system  $\phi^i, w^a, v^a$  on  $F$  where

$$\gamma \phi^i = \gamma^i(\phi), \quad \gamma w^a = v^a. \quad (3.23)$$

In this form it is clear why the equivalent reduction from  $(F, \gamma)$  to  $(N, q)$  is often referred to as elimination of contractible pairs (in this form it has been extensively used in the context of local BRST cohomology for a long time, see e.g. [33, 25, 26, 34, 35]).

A particular equivalent reduction of the  $Q$ -manifold  $(F, \gamma)$  in the case of off-shell CGR has been obtained by N. Boulanger in [16, 17]. More precisely, it was shown that all the derivatives of the metric except for Weyl curvature and its independent covariant derivatives can be taken as  $w^a$  variables while

the respective  $v^a$  variables parameterize all the derivatives of ghosts of order 2 and higher. As coordinates  $\phi^i$  one takes:  $\xi^\mu, \xi_\mu^\nu$  related to  $\xi^\mu$  ghost and its 1-st derivatives,  $\lambda, \kappa_a$  related to  $\kappa$  ghost and its 1-st derivatives, and  $W_{\mu\nu\rho\sigma|\alpha\dots}$  parameterizing Weyl tensor and its independent covariant derivatives. The action of  $\gamma$  on the ghost degree 1 variables is given by

$$\begin{aligned}\gamma\xi_\mu &= \xi^\rho\xi_{\rho\mu}, \\ \gamma\xi_\mu^\nu &= \xi_\mu^\rho\xi_\rho^\nu + P_{\mu\rho}^{\lambda\nu}\kappa_\lambda\xi^\rho + \frac{1}{2}\xi^\rho\xi^\lambda W_{\mu\nu\rho\lambda}, \\ \gamma\kappa_\nu &= \xi_\nu^\rho\kappa_\rho + \frac{1}{2}\xi^\rho\xi^\lambda C_{\nu\rho\lambda}, \\ \gamma\lambda &= \kappa_\rho\xi^\rho,\end{aligned}\tag{3.24}$$

where  $P_{\mu\rho}^{\lambda\nu} = -g^{\lambda\nu}g_{\mu\rho} + \delta_\mu^\lambda\delta_\rho^\nu + \delta_\rho^\lambda\delta_\mu^\nu$ , and denotes the Cotton tensor which can be expressed in our coordinates as  $C_{\mu\nu\rho} = W^\alpha_{\mu\nu\rho|\alpha}$ . In what follows we will also denote  $C_{\mu\nu\rho|\alpha_1\dots} := W^\alpha_{\mu\nu\rho|\alpha\alpha_1\dots}$ . The action of  $\gamma$  on degree zero variables reads as

$$\gamma g_{\mu\nu} = -2\lambda g_{\mu\nu} + \xi_\mu^\rho g_{\rho\nu} + \xi_\nu^\rho g_{\rho\mu}\tag{3.25}$$

and on Weyl tensor and its covariant derivatives  $\gamma$  acts in the following way:

$$\gamma W = \xi^\mu R_\mu W + \xi^\mu_\nu \Delta^\nu_\mu W + \kappa_\mu \Gamma^\mu W,\tag{3.26}$$

where

$$\begin{aligned}R_\beta W^\mu_{\nu\rho\sigma;\alpha_1\dots\alpha_k} &= W^\mu_{\nu\rho\sigma;\alpha_1\dots\alpha_k\beta}, \\ \Delta^\beta_\gamma W^\mu_{\nu\rho\sigma;\alpha_1\dots\alpha_k} &= -\delta_\gamma^\mu W^\beta_{\nu\rho\sigma;\alpha_1\dots\alpha_k} + \delta_\nu^\beta W^\mu_{\gamma\rho\sigma;\alpha_1\dots\alpha_k} + \dots\end{aligned}\tag{3.27}$$

The action of  $\Gamma^\mu$  can be determined from the nilpotency of  $\gamma$  and can be found in [16]. For example,  $\gamma^2\kappa_a = 0$  implies the relation  $\Gamma^\kappa C_{\mu\nu\rho} = W^\kappa_{\mu\nu\rho}$  which we need in what follows. We refer to [16] for further details.

For our purposes we need to reduce the above  $Q$ -manifold even further. It turns out that under some assumption on allowed class of metric configurations one can eliminate the metric together with the symmetric component of  $\xi_{\mu\nu}$ . Indeed,

$$\gamma g_{\mu\nu} = -2\lambda g_{\mu\nu} + \xi_\mu^\rho g_{\rho\nu} + \xi_\nu^\rho g_{\rho\mu} = -2\lambda g_{\mu\nu} + 2\xi_{(\mu\nu)}.\tag{3.28}$$

We relegate the discussion of the conditions under which this gives an equivalent reduction (for instance this the case if the metric is Euclidean) to Appendix **D** and simply assume that these are fulfilled.

In a suitable coordinate system  $x^a$  we decompose  $g_{ab} = \eta_{ab} + h_{ab}$ , with  $\eta_{ab}$  being standard Minkowski metric and find

$$\gamma h_{ab} = -2\lambda g_{ab} + 2\xi_{(ab)}, \quad \xi_{(ab)} = \frac{1}{2}(\xi_a^c g_{cb} + \xi_b^c g_{ac}). \quad (3.29)$$

So that  $h_{ab}$  can be taken as  $w^a$  variables in the equivalent reduction. The reduced  $Q$ -manifold is obtained by setting  $h_{ab} = \gamma h_{ab} = 0$ , giving

$$g_{ab} = \eta_{ab}, \quad \xi_{(ab)} = \lambda \eta_{ab}. \quad (3.30)$$

In what follows we use the antisymmetric matrix  $\rho_{ab} = \frac{1}{2}(\xi_{ab} - \xi_{ba})$  to parametrize the remaining components of  $\xi_{ab}$ .

To summarize, the reduced  $Q$ -manifold is coordinatized by degree 1 coordinates  $\xi^a, \rho^{ab}$  and degree 0 coordinates  $W_{\mu(\nu\rho\lambda|\alpha\dots)}$  (though it is very convenient to work with the overcomplete set of  $W$ -coordinates  $W_{\mu\nu\rho\lambda|\alpha\dots}$  as we keep doing). The action of the reduced  $\gamma$  (denoted by  $q$  below) on the ghost degree 1 coordinates can be easily obtained from (3.24) and is given explicitly by <sup>5</sup>:

$$\begin{aligned} q\xi^a &= \rho^a_c \xi^c + \xi^a \lambda, \\ q\rho^a_b &= \rho^a_c \rho^c_b + (\xi^a \kappa_b - \xi_b \kappa^a) + \frac{1}{2} \xi^c \xi^d W^a_{bcd}, \\ q\kappa_b &= \kappa_c \rho^c_b + \lambda \kappa_b + \frac{1}{2} \xi^c \xi^d C_{bcd}, \\ q\lambda &= \kappa_c \xi^c. \end{aligned} \quad (3.31)$$

The action of  $q$  on  $W$ -coordinates is also easily inferred from (3.26). In particular, for the Weyl and Cotton tensors we explicitly get:

$$\begin{aligned} qW^a_{bcd} &= \xi^k W^a_{bcd|k} - \rho_k^a W^k_{bcd} + \rho_b^k W^a_{kcd} + \rho_c^k W^a_{bkd} + \rho_d^k W^a_{bck} + 2\lambda W^a_{bcd}, \\ qC_{abc} &= \xi^k C_{abc|k} + \rho_a^k C_{kbc} + \rho_b^k C_{akc} + \rho_c^k C_{abk} + 3\lambda C_{abk} + \kappa_k W^k_{abc}. \end{aligned} \quad (3.32)$$

It should be stressed that the second equation is not written in independent coordinates which are the properly symmetrized Weyl "derivatives". Furthermore,

---

<sup>5</sup>By setting to zero  $W$  and  $C$  one arrives at the Chevalley-Eilenberg differential for conformal algebra, however in a slightly rescaled basis comparing to (3.4)

we need an on-shell version of this  $Q$ -manifold which is obtained by first restricting from in the fiber of jet-bundle to the surface where  $B_{ab}$  and its differential consequences are set to zero. The eliminatin of contractible pairs of  $\gamma$  is not affected as equations of motion do not affect ghosts and gauge non-invariant quantities.

In this way we have arrived at the  $Q$ -manifold  $(F, q)$  which is going to serve as a target space of the presymplectic AKSZ formulation of CGR.

### 3.5 Presymplectic structure

As was discussed above the minimal  $Q$ -manifold of CGR admits a compatible presymplectic structure of ghost degree  $n - 1$ , which originates from the descent completion of the BV symplectic structure. More precisely, the minimal  $Q$ -manifold can be seen as a  $Q$ -submanifold of total BRST complex of CGR and the presymplectic structure can be obtained as the restriction of the descent completion to the submanifold.<sup>6</sup>

Instead of computing the descent completion and its transfer to the minimal  $Q$ -manifold, which can be quite involved technically, we try to find such a structure by considering a generic ansatz of the closed 2-form of ghost degree 3 and not involving higher Weyl tensors. By some trial and error one finds that the correct presymplectic structure should involve Weyl tensor and Cotton tensors and the remaining ambiguity is fixed by  $q$ -invariance. The end result reads explicitly as:

$$\begin{aligned} \omega &= \omega_W - 2\omega_C, \\ \omega_W &= d(\rho_{ab})d(W^{abnm}\epsilon_{nmpk}\xi^p\xi^k), \quad \omega_C = d(\xi_a)d(C^a_{bc}\epsilon^{bcpk}\xi_p\xi_k). \end{aligned} \quad (3.33)$$

**Proposition 3.2.** *Presymplectic structure (3.33) satisfies:  $d\omega = 0$ ,  $L_q\omega = 0$ , and  $\text{gh}(\omega) = n + 1$  and hence together with the homological vector field  $q$  it defines a presymplectic AKSZ model.*

The first and the third properties are obvious. Let us give here main points of the proof of  $L_q\omega = 0$  (details are relegated to the Appendix B). It is enough

---

<sup>6</sup>In this setup, this is just an explicit realization of the homotopy transfer.

to show that  $i_q\omega = dH$  for some  $H$ . We have the following relations:

$$i_q\omega_W = d(\rho_{al}\rho^l_b W^{abnm}\epsilon_{nmpk}\xi^p\xi^k) + (\xi_a\kappa_b - \xi_b\kappa_a)d(W^{abnm}\epsilon_{nmpk}\xi^p\xi^k) + \frac{1}{2}W_{abij}\xi^i\xi^j d(W^{abnm}\epsilon_{nmpk}\xi^p\xi^k) + d(\rho_{ab})\xi^j W^{abnm|j}\epsilon_{nmpk}\xi^p\xi^k. \quad (3.34)$$

It is nearly  $dH_W$  with  $H_W$  given by

$$H_W = \rho_{al}\rho^l_b W^{abnm}\epsilon_{nmpk}\xi^p\xi^k + 2\xi_a\kappa_b W^{abnm}\epsilon_{nmpk}\xi^p\xi^k + W_{abij}\xi^i\xi^j W^{abnm}\epsilon_{nmpk}\xi^p\xi^k, \quad (3.35)$$

with the discrepancy being

$$i_q\omega_W - dH_W = -2d(\xi_a)\kappa_b W^{abnm}\epsilon_{nmpk}\xi^p\xi^k + 2\xi_a d(\kappa_b)W^{abnm}\epsilon_{nmpk}\xi^p\xi^k + d(\rho_{ab})\xi^j W^{abnm|j}\epsilon_{nmpk}\xi^p\xi^k. \quad (3.36)$$

As for the second term  $\omega_C$  we find:

$$i_q\omega_C = \rho_{an}\xi^n d(C^a_{bc}\epsilon^{bcpk}\xi_p\xi_k) + d(\xi^a)\rho_a{}^n C_{nbc}\epsilon^{bcpk}\xi_p\xi_k + \xi_a\lambda d(C^a_{bc}\epsilon^{bcpk}\xi_p\xi_k) + d(\xi_a)\lambda C^a_{bc}\epsilon^{bcpk}\xi_p\xi_k + d(\xi_a)\xi^d C_{abc|d}\epsilon^{bcpk}\xi_p\xi_k + d(\xi^a)\kappa_k W^k_{abc}\epsilon^{bcpk}\xi_p\xi_k. \quad (3.37)$$

Introducing:

$$H_C = \rho_{an}\xi^n C^a_{bc}\epsilon^{bcpk}\xi_p\xi_k + \xi_a\lambda C^a_{bc}\epsilon^{bcpk}\xi_p\xi_k, \quad (3.38)$$

we find

$$i_q\omega_C - dH_C = -d(\rho_{an})\xi^n C^a_{bc}\epsilon^{bcpk}\xi_p\xi_k + \xi_a d(\lambda)C^a_{bc}\epsilon^{bcpk}\xi_p\xi_k + d(\xi_a)\xi^d C_{abc|d}\epsilon^{bcpk}\xi_p\xi_k - d(\xi^a)\kappa^k W_{akbc}\epsilon^{bcpk}\xi_p\xi_k, \quad (3.39)$$

so that

$$i_q(\omega_0 - 2\omega_C) = dH, \quad (3.40)$$

where the "Hamiltonian" is given by:

$$H = H_W - 2H_C = \rho_{al}\rho^l_b W^{abnm}\epsilon_{nmpk}\xi^p\xi^k + 2\xi_a\kappa_b W^{abnm}\epsilon_{nmpk}\xi^p\xi^k - 2\rho_{an}\xi^n C^a_{bc}\epsilon^{bcpk}\xi_p\xi_k - 2\xi_a\lambda C^a_{bc}\epsilon^{bcpk}\xi_p\xi_k + W_{abij}\xi^i\xi^j W^{abnm}\epsilon_{nmpk}\xi^p\xi^k. \quad (3.41)$$

We are now ready to spell out an explicit form of the action. Introducing component fields parameterizing  $\sigma : T[1]X \rightarrow F$  via

$$\begin{aligned}\sigma^*(\xi^a) &= e_\mu^a(x)\theta^\mu, & \sigma^*(\rho^{ab}) &= \omega_\mu^{ab}(x)\theta^\mu, \\ \sigma^*(\kappa^a) &= f_\mu^a(x)\theta^\mu, & \sigma^*(\lambda) &= v_\mu(x)\theta^\mu, \\ \sigma^*(W_{abcd}) &= W_{abcd}(x), & \sigma^*(C_{abc}) &= C_{abc}(x),\end{aligned}\tag{3.42}$$

where by some abuse of notations we used the same letters to denote some target space coordinates and the respective fields. It can be convenient to treat  $e, \omega, W, C$  as differential forms on  $X$ . With this identification the action (2.6) can be written as:

$$\begin{aligned}S[e, \omega, W, C] &= \int_X [(d\omega_{ab} + \omega_{ac}\omega^c_b)W^{abnm}\epsilon_{nmpk}e^pe^k + \\ &+ W_{abcd}e^ce^dW^{abnm}\epsilon_{nmpk}e^pe^k - 2(de_a + \omega_{ad}e^d)C^a_{bc}\epsilon^{bcpk}e_pe_k],\end{aligned}\tag{3.43}$$

we omit the wedge product symbol. Note that before the symplectic reduction has been done the action is to be considered as that depending on all the degree zero fields. However here, we only spelled explicitly those fields on which it depends nontrivially. In particular, thanks to the symmetry properties of  $W, C$ , fields  $v, f$  do not contribute (as we are going to see this is agreement with the fact that  $\frac{\partial}{\partial \lambda}$  and  $\frac{\partial}{\partial \kappa}$  are in the kernel of the presymplectic structure).

To make the structure of the action more explicit define

$$\begin{aligned}R_{ab} &= d\omega_{ab} + \omega_{ac}\omega^c_b + (e_af_b - e_bf_a), & T_a &= de_a + \omega_{ad}\xi^d + e_av, \\ *W^{ab} &= W^{abnm}\epsilon_{nmpk}e^pe^k, & *C^a &= C^a_{bc}\epsilon^{bcpk}e_pe_k.\end{aligned}\tag{3.44}$$

In these terms the action reads as:

$$S = \int_X [R_{ab} \wedge *W^{ab} + W_{ab} \wedge *W^{ab} - 2T_a \wedge *C^a].\tag{3.45}$$

### 3.6 Symplectic quotient and BV formulation

In order to give a consistent interpretation to the gauge theory determined by (3.43) and to construct the respective BV description we need to identify a kernel distribution for the symplectic structure  $\bar{\omega}$ , induced on the (open domain of) the space of supermaps  $Smaps(\mathbb{R}^4[1], F)$ . In the case at hand the manifold  $\bar{F}$  is

coordinatized by the degree zero coordinates introduced in (3.42) together with the remaining coordinates introduced as e.g.:

$$\widehat{\sigma}^*(\xi^a) = \sum_{l=0}^4 \frac{1}{l!} \xi_{\mu_1 \dots \mu_l}^a \theta^{\mu_1} \dots \theta^{\mu_l}, \quad (3.46)$$

with the identification  $\xi_{\mu}^a = e_{\mu}^a$ . The ghost degree of the coefficients is determined by  $\text{gh}(\theta^{\mu}) = 1$ . We then take  $\bar{F}$  to the space of supermaps with nondegenerate component  $e_{\mu}^a$  because as we are going to see these coordinates correspond to the frame field which must be invertible.<sup>7</sup>

We have the following

**Proposition 3.3.** *The presymplectic structure  $\bar{\omega}$  is regular (i.e. has a constant rank) on  $\bar{F}$ .*

The idea of the proof is taken from [7] and amounts to identifying vector fields  $X_{\alpha}$  on  $F$  such that they are in the kernel of  $\omega$  and their prolongations to  $\bar{F}$  generate the kernel at any point of the body of  $\bar{F}$ . In terms of local coordinates the prolongation of a vector field  $X$  on  $F$  can be defined in local coordinates through:

$$\bar{X}\psi^A(\theta) = X^A(\psi(\theta)), \quad X^A(\psi) = X\psi^A, \quad (3.47)$$

where in the left hand side  $\bar{X}$  acts on the coordinates on  $\bar{F}$ , which are packed in the generating function  $\psi^A(\theta)$ . It is easy to check that  $i_X\omega = 0$  implies  $i_{\widehat{X}}\bar{\omega}$ .

The rank of  $\bar{\omega}$  can't decrease in a neighbourhood of a point. Because the rank of the distribution generated by  $\bar{X}_{\alpha}$  also can not decrease one concludes that the rank is constant and the kernel distribution is generated by  $\bar{X}_{\alpha}$ . Explicit form of  $X_{\alpha}$  and the proof that their prolongations exhaust the kernel at any point of the body of  $\bar{F}$  is given in the Appendix C.

Because  $\bar{\omega}$  is regular the general construction of Section 2.2 applies and we end up with the first order BV formulation whose field-antifield space is  $\bar{G}$  which is a symplectic quotient of  $\bar{F}$ . This BV formulation has (3.43) as a classical action. To make sure we are indeed dealing with a specific BV formulation of the conformal gravity we need to make sure that (i) (3.43) reduced to

---

<sup>7</sup>This step is completely analogous to the respective considerations [7] in the case of Einstein gravity.

the symplectic quotient is equivalent to the conformal gravity and that (ii) the gauge generators encoded in the BV action take all the gauge symmetries of system into account.

(i) Because (3.43) depends on fields of ghost degree zero only it is enough to analyze the reduction only in the sector of degree zero variables. First of all we disregard all the fields associated to higher order  $W$  as (3.43) and  $\bar{\omega}$  do not involve them and hence they can be set to zero. The same applies to  $v_\mu$  and  $f^a{}_\mu$ . Among  $X_\alpha$  the only vector fields that affect the remaining ghost degree 0 variables are

$$\begin{aligned} X^{(trace)} &= \xi^a \frac{\partial}{\partial \xi_a} - 2W^{abnm} \frac{\partial}{\partial W^{abnm}} - 3C^{abc} \frac{\partial}{\partial C^{abc}}, \\ Y_d^{(antisym)} &= \epsilon_{abcd} \xi^c \frac{\partial}{\partial \rho_{ab}} + W_{dnm}{}^a \epsilon^{nmbc} \frac{\partial}{\partial C^{abc}}, \\ Y_d^{(trace)} &= \xi^c \delta_{cd}^a \frac{\partial}{\partial \rho_{ab}} - W^{adbc} \frac{\partial}{\partial C^{abc}}. \end{aligned} \quad (3.48)$$

The action of their prolongations on  $e_{a\mu}, \omega_{ab,\mu}$  read as:

$$\begin{aligned} \bar{X}^{(trace)} e^a{}_\mu &= e^a{}_\mu, \\ \bar{Y}_d^{(antisym)} \omega_{ab,\mu} &= \epsilon_{abcd} e^c{}_\mu, \quad \bar{Y}_d^{(trace)} \omega_{ab,\mu} = (e_{a,\mu} \eta_{bc} - e_{b,\mu} \eta_{ac}). \end{aligned} \quad (3.49)$$

It follows that by choosing the embedding of  $\bar{G} \subset \bar{F}$  one can assume that the determinant of  $e^a{}_\mu$  as well as totally antisymmetric and the trace components of  $\omega_{ab,\mu} e^a{}_\mu$  can be set to whatever we want. We use this to set  $\det(e) = 1$  and to set the respective components of the torsion  $T_a = de_a + \omega_{ab} e^b$  to zero. The remaining irreducible component is associate to hook-type Young tableaux and is set to zero by the equation of motion for (3.43) obtained by varying with respect to  $C_{abc}$ . It follows this component of  $\omega$  together with  $C_{abc}$  and their elimination sets  $\omega$  to the Levi-Civita connection  $\omega_{ab,\mu}(e)$ . Finally, equations for  $W_{abcd}$  can be solved with respect to it and its elimination gives:

$$S[e, \omega(e)] = \int W_{ab} \wedge *W^{ab}, \quad (3.50)$$

i.e. the standard action of GGR written in terms of the frame field. The standard argument then implies that all the components of  $e^a{}_\mu$  which do not contribute to the metric can be gauge-fixed by local Lorentz transformations (which are

algebraic) and the action can be equivalently rewritten in terms of the metric, giving the proof that we indeed obtained CGR action from the presymplectic AKSZ.

(ii) To check that the BV formulation indeed takes into account all the gauge symmetries of the classical action let us analyze the spectrum of ghost variables. Ghosts  $\lambda, \kappa^a$  are in the kernel and do not survive on  $\bar{G}$ . At the same time  $\xi^a, \rho^{ab}$  are not affected by the kernel and hence their restriction to the symplectic quotient remain independent coordinates. Their associated gauge transformations are precisely the diffeomorphisms and the local Lorentz transformations. Note that we do not see ghosts associated to Weyl transformations because this invariance is taken into account in this formulation through the kernel of the presymplectic structure and we fixed it when passing to the symplectic quotient. Note that there are also no ghosts pertaining to the special conformal sector since they are also in the kernel.

We then conclude that we have indeed reconstructed a first order BV formulation of CGR.

Let us finally mention the substantial difference between the presymplectic AKSZ formulation of CGR and the Einstein gravity. In the case of Einstein gravity one can avoid considering the entire minimal model as a target space and truncate it to just a Poincare algebra with shifted degree, see [7] for more details. The analogous truncation doesn't work in the case of CGR. However, one can still construct a presymplectic AKSZ formulation based on conformal algebra and closely related to the frame-like formulation from [11] but this requires imposing additional constraints by hands. This possibility is discussed in Appendix E.

## 4 General presymplectic BV-AKSZ formalism

Our discussion in Section 2 was limited to the case of diffeomorphism invariant systems for which AKSZ representation (at the level of EOMs) is available. Now we explain how the presymplectic BV-AKSZ formalism extends to the general situation of not-necessarily diffeomorphism-invariant systems.

## 4.1 Gauge PDEs

If we are only interested in the equations of motion and gauge symmetries (and hence disregard the Lagrangian) a data of a local gauge theory can be encoded into the following geometric object, known as gauge PDE [7] (see also [22, 23] for the earlier less geometrical version):

**Definition 4.1.** *Gauge PDE is a  $\mathbb{Z}$ -graded fiber bundle  $\pi : E \rightarrow T[1]X$  whose total space is equipped with a homological vector field  $Q$  of degree 1 satisfying:  $Q \circ \pi^* = \pi^* \circ d_X$  and  $d_X$  is the de Rham differential on  $X$  seen as a homological vector field on  $T[1]X$ . In addition it is natural to require gauge PDE to be equivalent to a non-negatively graded one.*

One often also imposes some extra conditions of a technical nature to exclude pathological examples. We assume that  $X$  is a real manifold while  $E$  (and  $T[1]X$ ) is a  $\mathbb{Z}$ -graded one with the degree denoted by  $\text{gh}()$ . Note that gauge PDEs are special case of  $Q$ -bundles [24]. However, gauge PDEs are necessarily  $\mathbb{Z}$ -graded and are typically infinite-dimensional. Further details and references can be found in [21].

By definition, solutions of a gauge PDE are  $Q$ -sections, i.e. sections satisfying  $\sigma^* \circ Q = d_X \circ \sigma^*$ . In a similar way one defines infinitesimal gauge transformation of sections.

The notion of gauge PDE is rather flexible. For instance, the standard BV formulation at the level of equations of motion is reproduced by taking as  $E$  a jet-bundle associated to the bundle of BV fields (fields, ghosts, antifields) seen as a bundle over  $T[1]X$  (i.e. functions on  $E$  are local horizontal forms on the jet-bundle) and  $Q = d_h + s$ , where  $d_h$  is a canonical horizontal differential on the jet-bundle and  $s$  is the BV-BRST differential.

Another standard example is PDE. Let  $E_X \rightarrow X$  be an equation manifold seen as a bundle over the space-time  $X$ . The Cartan distribution gives rise to a homological vector field  $d_h$  on  $E$  which is  $E_X$  pulled back to  $T[1]X$ . In a suitable local coordinate system  $x^a, \theta^a, \psi^A$ , where  $x^a, \theta^a$  are adapted local coordinates on  $T[1]X$ , induced by local coordinates  $x^a$  on  $X$ , and  $\psi^A$  originate from local coordinates on a typical fiber, one takes:

$$d_h = \theta^a D_a, \quad D_a = \frac{\partial}{\partial x^a} + \Gamma_a^A(x, \psi) \frac{\partial}{\partial \psi^A}, \quad (4.1)$$

where  $D_a$  are components of the total derivative on the equation manifold. Note that this is a generic form of the  $Q$ -structure on  $E$  if the typical fiber is a real manifold (so that  $\text{gh}(\psi^A) = 0$ ). It is easy to see that in these coordinates solutions are covariantly-constant sections in agreement with the standard picture.

From the gauge PDE perspective AKSZ sigma models discussed in Section 2 correspond to trivial (as a  $Q$ -bundle)  $Q$ -bundles  $(E, Q) = (F, q) \times (T[1]X, d_X)$ .

## 4.2 Vertical forms

The Lagrangian (or partially Lagrangian) systems can be described as gauge PDEs with extra structures. To define these we first need to recall a particular realization of vertical forms.

Let  $\pi : E \rightarrow \mathcal{X}$  be a generic fiber bundle. The exterior algebra of local forms on  $E$  contains the ideal  $\mathcal{I}$  generated by forms of the form  $\pi^*\alpha$ , where  $\alpha \in \bigwedge^k(\mathcal{X})$ ,  $k > 0$ . The algebra of vertical forms is the quotient of  $\bigwedge^\bullet(E)$  by the ideal. Note that vertical zero-forms are just zero-forms as the ideal does not contain zero-forms. It is convenient to work with vertical forms in terms of the representatives as we do in what follows.

The De Rham differential  $d$  is well defined on the equivalence classes. Understood as an operation on vertical forms it is usually called a vertical differential. Suppose that  $Y$  is a vector field on  $E$  which is related by the projection to a vector field  $y$  on  $\mathcal{X}$ , i.e.  $Y \circ \pi^* = \pi^* \circ y$ , which in turn implies  $L_Y \circ \pi^* = \pi^* \circ L_y$ . It is easy to check that  $L_Y$  is well defined on the equivalence classes and hence defines a Lie derivative on vertical forms. Despite  $i_Y$  is not well-defined on equivalence classes (unless  $Y$  is vertical) one can still employ the Cartan formula  $L_Y = i_Y d + (-1)^{|Y|} di_Y$  for representatives.

Given a local trivialization of  $E$  over  $U \subset \mathcal{X}$ , i.e.  $E|_U = U \times F$ , each equivalence class has a unique representative that vanish on any horizontal vector. In the adapted coordinates  $\psi^A, X^\alpha$ , with  $X^\alpha$  being coordinates on the base, this representative does not involve basis differential  $dX^\alpha$ . In terms of such representatives and the adapted coordinates the vertical differential can be explicitly written as  $d\psi^A \frac{\partial}{\partial \psi^A}$ . It is then clear that locally the vertical differential is acyclic.

### 4.3 Presymplectic gauge PDEs

A presymplectic BV-AKSZ system is a gauge PDE  $(E, T[1]X, Q)$  equipped with a compatible vertical presymplectic structure  $[\omega]$ , i.e. a 2-form  $\omega$  satisfying  $\text{gh}(\omega) = n - 1$ ,  $d\omega \in \mathcal{I}$ ,  $L_Q\omega = \mathcal{I}$ , where  $n = \dim X$ . In addition we assume the following technical conditions:  $i_Q d\omega \in \mathcal{I}$  and  $i_Q L_Q\omega \in \mathcal{I}$ . Here and in what follows we work in terms of representatives.

Given a presymplectic gauge PDE one can consider a natural Lagrangian system associated to it. More precisely, we first introduce covariant Hamiltonian  $\mathcal{H}$  as a solution to<sup>8</sup>

$$i_Q\omega + d\mathcal{H} \in \mathcal{I}. \quad (4.2)$$

To see that  $\mathcal{H}$  exists observe that  $di_Q\omega \in \mathcal{I}$  thanks to  $L_Q\omega \in \mathcal{I}$  and  $i_Q d\omega \in \mathcal{I}$ . Then acyclicity of  $d$  on vertical forms implies the existence of  $\mathcal{H}$  (for  $n > 1$  it exists globally). Note that  $\mathcal{H}$  is a zero form and hence is defined uniquely modulo functions of the form  $\pi^*(h)$ ,  $h \in \mathcal{C}^\infty(T[1]X)$ . As we are going to see this ambiguity doesn't affect the equations of motion.

Together with a symplectic potential  $\chi$  defined through  $\omega - d\chi \in \mathcal{I}$  this data defines a natural action functional on sections  $T[1]X \rightarrow E$ :

$$S[\sigma] = \int_{T[1]X} [\sigma^*(\chi))(d_X) + \sigma^*(\mathcal{H})]. \quad (4.3)$$

This action is a generalization of the presymplectic version [2] of the conventional AKSZ action on one hand and, in the case where  $(E, T[1]X, Q)$  is a PDE, of the so-called intrinsic action from [36] (see also [37]).

Let us check that  $S[\sigma]$  does not depend on the choice of a representative  $\chi$  of the equivalence class  $[\chi]$ . If instead of  $\chi$  we took another representative  $\chi' = \chi + \epsilon$ , where  $\epsilon \in \mathcal{I}$  and hence it can be written in local coordinates as  $\epsilon = dx^a \alpha_a + d\theta^b \beta_b$  we find:

$$(\sigma^*(\epsilon))(d_X) = \theta^a \sigma^*(\alpha_a), \quad (4.4)$$

At the same time, the variation  $\mathcal{H}_\epsilon$  of  $\mathcal{H}$  is determined by  $i_Q d\epsilon + d\mathcal{H}_\epsilon \in \mathcal{I}$ . Using  $L_Q\epsilon \in \mathcal{I}$  one finds  $di_Q\epsilon + d\mathcal{H}_\epsilon \in \mathcal{I}$  and hence  $\mathcal{H}_\epsilon + i_Q\epsilon = \pi^*(h)$ . The variation

---

<sup>8</sup>Note that here we have changed the sign convention with respect to the AKSZ setup of section 2

of the form  $\pi^*(h)$ ,  $h \in \mathcal{C}^\infty(T[1]X)$  does not change equations of motion (can be locally represented as a total derivative) while

$$\sigma^*(i_Q\epsilon) = \sigma^*(\theta^a\alpha_a) = \theta^a\sigma^*(\alpha_a), \quad (4.5)$$

where we again use adapted coordinate system  $x^a, \theta^a, \psi^A$  on  $E$ . Note that in this coordinates  $Qx^a = \theta^a$ ,  $Q\theta^a = 0$ . It follows that the action is unchanged under  $\chi \rightarrow \chi'$  if we disregard the variations of the form  $\pi^*(h)$  which are locally total derivatives.

In this way we have arrived at a natural Lagrangian system associated to a presymplectic BV-AKSZ system. However, in general this system contains an infinite amount of fields and the interpretation of the action is to be clarified. In the case where  $(E, Q)$  is a usual PDE (seen as a gauge PDE) the consistent interpretation can be arrived at by disregarding those fields which are in the kernel of the symplectic structure [38]. In this way one can often (though not always) describe Lagrangian systems in terms of the intrinsic geometry of their equation manifolds. Just like in the case of diff-invariant systems discussed in Section 2 a more refined interpretation can be given by passing to the symplectic quotient of the induced symplectic structure on the space of super-sections of  $E$ .

#### 4.4 Symplectic quotient and BV formulation

It is convenient to identify super-sections as the space of section of a bundle  $\bar{E}$  over  $X$ . More precisely, a fiber over  $p \in X$  is the space of supermaps from  $(T_pX)[1]$  to  $F$ .

To be more explicit we now work locally using a trivialization  $E|_{T[1]U} = T[1]U \times F$  over  $U \subset X$ . The fiber  $\bar{F}$  of  $\bar{E}|_U$  is a space of supermaps from  $\mathbb{R}^n[1]$  to  $F$ . Coordinates on  $\bar{F}$  are introduced as in (2.8), where we keep denoting coordinates on  $F$  by  $\psi^A$  and use  $\Psi^I$  to denote coordinates on  $\bar{F}$ .

It turns out that the presymplectic structure  $\omega$  on  $E$  gives rise to an  $(n+2)$ -form on  $\bar{E}$  of the following structure:

$$\bar{\omega}^{\bar{E}} = (dx)^n \bar{\omega}_{IJ}(\Psi, x) d\Psi^I d\Psi^J, \quad (4.6)$$

where  $\bar{\omega}_{IJ} = \bar{\omega}(\frac{\partial}{\partial\Psi^I}, \frac{\partial}{\partial\Psi^J})$ , with  $\bar{\omega}$  on each fiber given by

$$\bar{\omega} = \int d^n\theta \omega_{AB}(\psi^A(\theta), x, \theta) d\psi^A(\theta) d\psi^B(\theta). \quad (4.7)$$

In other words on a given fiber the form is still the same (2.9). It is easy to see that  $d\bar{\omega} = 0$ . An alternative language would be to identify  $\bar{\omega}$  as a vertical 2-form with values in densities on  $X$ .

We assume that the  $n + 2$  form  $\bar{\omega}$  is regular (in the sense that the rank of  $\bar{\omega}_{IJ}$  is constant) and consider the vertical kernel of  $\bar{\omega}$ . This consists of vertical vector fields  $W$  such that  $i_W\bar{\omega} = 0$ . The vertical kernel is an involutive distribution and one can (at least locally) pass to the quotient space. Because the kernel distribution is vertical the quotient is again a bundle over  $X$  (its fibers are simply the symplectic quotients of the symplectic form  $\bar{\omega}_{IJ}d\Psi^i d\Psi^j$  on  $\bar{F}$ ). Form  $\bar{\omega}$  induces a vertically-nondegenerate  $n + 2$ -form on the quotient. This gives us a fiber-bundle equipped with  $n + 2$ -form satisfying all the conditions of the BV  $n + 2$ -form. It has a natural lift to an  $(n, 2)$ -form on  $J^\infty(\bar{E})$ .

By repeating the arguments given in Section 2.2 in this more general setting one can show that BV-like action (2.7) with  $H$  replaced with  $\mathcal{H}$  induces the BV action on the symplectic quotient and it satisfies the BV master equation modulo boundary terms.

To summarize, under some regularity assumptions a presymplectic gauge PDE encodes a Lagrangian BV system. This system is not necessarily equivalent (at the level of equations of motion) to the starting point gauge PDE. However, one can always find an equivalent gauge PDE representation and a presymplectic structure defined on it, such that the equivalence holds (though there are examples [37] where minimal model does not fit).

## 4.5 Example: Maxwell

We now illustrate the construction using the example of Maxwell theory. More precisely we take the minimal model of the respective gauge PDE (the respective total space  $Q$ -manifold is known in the literature for quite some time, see e.g. [26] and refs. therein. The solutions of this gauge PDE satisfy the so-called unfolded form of Maxwell equations, see e.g. [39]). The coordinates on the

total space of  $E \rightarrow T[1]X$  are

$$\begin{aligned} & x^a, \theta^a, \quad C, \\ & F_{a|b}, \quad F_{a|bc}, \quad \dots \quad F_{a|b_1 \dots b_l}, \quad \dots \end{aligned} \quad (4.8)$$

where tensors  $F_{a|b_1 \dots b_l}$  are symmetric in the second group of indices, satisfy the Young symmetry condition  $F_{(a|b_1 \dots b_l)} = 0$ , and are totally traceless. The action of  $Q$  is given by

$$\begin{aligned} & Qx^a = \theta^a, \quad QC = -\frac{1}{2}\theta^a\theta^b \\ & F_{a|b}, \quad QF_{a|b} = \theta^c(F_{a|bc} + \dots), \quad QF_{a|bd} = \theta^c(F_{a|bdc} + \dots), \quad \dots \end{aligned} \quad (4.9)$$

where  $\dots$  denote terms maintaining the Young symmetry properties. It is clear that we have a  $Q$ -bundle structure as  $Q \circ \pi^* = \pi^* \circ d_X$ , with  $d_X = \theta^a \frac{\partial}{\partial x^a}$  and  $\pi^*(x^a) = x^a$  and  $\pi^*(\theta^a) = \theta^a$  in these coordinates.

The compatible presymplectic structure and the respective potential can be taken as [2]:

$$\omega = dC dF^{ab}(\theta)_{ab}^{(n-2)}, \quad \chi = dC F^{ab}(\theta)_{ab}^{(n-2)}. \quad (4.10)$$

Note that in checking  $L_Q\omega = 0$  one has to use the "equations of motion", i.e. that  $F_{a|bc}$  are totally traceless.

The covariant Hamiltonian  $\mathcal{H}$  defined through  $i_Q\omega + d\mathcal{H} \in \mathcal{I}$  is given by

$$\mathcal{H} = \frac{1}{2} F_{ab} F^{ab}(\theta)^{(n)} \quad (4.11)$$

Introducing component fields parameterizing section  $\sigma : T[1]X \rightarrow E$  according to

$$\sigma^*(C) = A_b(x)\theta^c, \quad \sigma^*(F_{a|b}) = F_{a|b}(x), \quad \dots \quad (4.12)$$

the action takes the form

$$S[\sigma] = \int d^n x [F^{ab}(\partial_a A_b - \partial_b A_a) + \frac{1}{2} F^{ab} F_{ab}] \quad (4.13)$$

which is a standard covariant 1st order action of the Maxwell field.

The induced symplectic structure  $\bar{\omega}$  on the fiber  $\bar{F}$  is given by

$$\bar{\omega} = dC^0 dF_{ab}^2 + 2dA_a dF_b^1 + dC_{ab}^2 dF^{a|b} \quad (4.14)$$

where the relevant components of supersection are introduced via:

$$\begin{aligned}\widehat{\sigma}^*(C) &= \overset{0}{C}(x) + A_a(x)\theta^a + \frac{1}{2}\overset{2}{C}_{ab}(x)\theta^a\theta^b \dots \\ \widehat{\sigma}^*(F^{a|b}) &= F^{a|b}(x) + \overset{1}{F}_c^{a|b}(x)\theta^c + \frac{1}{2}\overset{2}{F}_{cd}^{a|b}(x)\theta^c\theta^d + \dots\end{aligned}\tag{4.15}$$

It follows that all the fields except for

$$\overset{0}{C}, \quad C^* = F_{ab}^{a|b}, \quad A_a, \quad A_*^a = F_b^{a|b}, \quad F^{a|b}, \quad F_{ab}^* = C_{ab}\tag{4.16}$$

are in the kernel of  $\bar{\omega}$  and can be set to zero to obtain the symplectic quotient. The ghost degree of the remaining fields are given by

$$\begin{aligned}\text{gh}(A_a) &= \text{gh}(F^{a|b}) = 0, & \text{gh}(C) &= 1, \\ \text{gh}(A_*^A) &= \text{gh}(F_{ab}^*) = -1, & \text{gh}(C) &= -1.\end{aligned}\tag{4.17}$$

By inspecting the ghost degree of the remaining fields and the above symplectic structure one concludes that we have arrived at the proper field-antifield space of the BV formulation of the Maxwell field. It is straightforward to check that the BV action (2.7) with  $H$  replaced by  $\mathcal{H}$  explicitly gives a proper solution of the master equation extending the classical action (4.13). Extension to the YM theory is also straightforward.

## Acknowledgments

ID is grateful to V. Gritzaenko and A. Yan for discussions. MG wishes to thank N. Boulanger, A. Sharapov, E. Skvortsov and especially A. Kotov for useful discussions. The work was supported by the Russian Science Foundation grant 18-72-10123.

## A Tensor conventions

In this section we describe our conventions for the tensors emerging in the context of CGR and give missing details on its frame-like description.

The Bach tensor is defined as

$$B_{\mu\nu} = P^{\rho\lambda}W_{\mu\rho\nu\lambda} + \nabla^\rho\nabla_\rho P_{\mu\nu} - \nabla^\rho\nabla_\mu P_{\nu\rho}, \quad (\text{A.1})$$

where  $\nabla^\mu$  is the metric-compatible covariant derivative and  $P_{\mu\nu}$  is the Schouten tensor:

$$P_{\mu\nu} = \frac{1}{2}\left(R^{(0)\rho}{}_{\mu\rho\mu} - \frac{R^{(0)\rho\sigma}{}_{\rho\sigma}}{6}g_{\mu\nu}\right), \quad (\text{A.2})$$

$R^{(0)\rho}{}_{\mu\nu\sigma}$  is the Riemann curvature, and Weyl tensor is defined via:

$$W_{\mu\nu}{}^{\rho\sigma} = R^{(0)\rho\sigma}{}_{\mu\nu} - 4P_{[\mu}{}^{[\rho}{}_{\nu]}{}^{\sigma]}. \quad (\text{A.3})$$

Now we give missing details on the derivation of the metric-like action (3.1) from the frame-like action (3.10). Starting with the equation of motion  $\epsilon^{abcd}R_{ab}e_c = 0$  for  $f^a$ , introducing tangent components  $R_{abcd}$  via  $R_{ab} = R_{abcd}e^ce^d$ , and wedging with  $e^f$  one finds

$$(R^{ab}{}_{km})\delta_{abd}^{kmf} = 0, \quad (\text{A.4})$$

which implies

$$2R^{ab}{}_{ab}\delta_d^f + 4R^{fb}{}_{bd} = 0. \quad (\text{A.5})$$

Contracting  $f, d$  one gets  $R^{ab}{}_{ab} = 0$  and

$$R^c{}_{bcd} = 0. \quad (\text{A.6})$$

Explicitly one has

$$R^c{}_{bcd} = R^{(0)c}{}_{bcd} + (4f_{d,b} - e_{d,a}f_{a,b} - e_{a,b}f_{d,a} + e_{d,b}f) = 0, \quad (\text{A.7})$$

where  $f = f_a^a$ . It follows

$$f = -\frac{1}{6}R^{(0)} \quad (\text{A.8})$$

and

$$f_{b,\mu} = -\frac{1}{2}\left(e^d{}_\mu R^{(0)c}{}_{bcd} - \frac{1}{6}e_{b,\mu} R^{(0)}\right) \quad (\text{A.9})$$

so that  $f_\mu^b$  is indeed a Schouten tensor on shell.

Now we turn to the properties of the Weyl and Cotton tensors and their Hodge duals, which we need in the proof of the invariance of the presymplectic structure. The Hodge duals are defined as:

$$C_{abc}^* = C_a{}^{pk} \epsilon_{pkbc}, \quad W_{abcd}^* = W_{ab}{}^{pk} \epsilon_{pkcd}. \quad (\text{A.10})$$

The inverse relations read:

$$\begin{aligned} C_{abc}^* \epsilon^{bcnm} &= C_a{}^{pk} \epsilon_{pkbc} \epsilon^{bcnm} = 4C_a{}^{nm}, \\ W_{abcd}^* \epsilon^{cdnm} &= W_{ab}{}^{pk} \epsilon_{pkcd} \epsilon^{cdnm} = 4W_{ab}{}^{nm}. \end{aligned} \quad (\text{A.11})$$

Every property of Weyl and Cotton tensors has its counterpart in terms of the duals.

### 1. Trace-free and Young symmetry

$$\begin{aligned} \eta^{ac} C_{abc} &= 0 = C_{apk}^* \epsilon^{pkba}, \\ \epsilon^{abcd} C_{abc} &= 0 = C_a{}^{*pk} \epsilon_{pkbc} \epsilon^{abcd} = 2C_a{}^{*ad}, \end{aligned} \quad (\text{A.12})$$

$$\begin{aligned} \eta^{ac} W_{abcd} &= 0 = W_{abpk}^* \epsilon^{pkad}, \\ \epsilon^{abcn} W_{abcd} &= 0 = W_{ab}{}^{*pk} \epsilon_{pkcd} \epsilon^{abcn} = 2W_{ab}{}^{*ab} \delta_d^n + 8W_{ad}{}^{*an}. \end{aligned} \quad (\text{A.13})$$

### 2. Bach equation [16]

$$C_{abc|}{}^b = 0 = C_a{}^{*pk|b} \epsilon_{pkbc}. \quad (\text{A.14})$$

### 3. Definition [40] of $C$ in terms of higher Weyl-tensors:

$$W_{abcd}|^a = W_{bcda}|^a - W_{bdca}|^a = C_{bcd}. \quad (\text{A.15})$$

### 4. The Weyl tensor enjoys the following relation:

$$W_{ab}{}^{ef} \epsilon_{cdef} = W^{ef}{}_{cd} \epsilon_{abef}. \quad (\text{A.16})$$

## B Invariance of the presymplectic structure

Here we explicitly demonstrate that the presymplectic structure (3.33) is  $q$ -invariant.

Equation (3.34) is obtained in the following way:

$$\begin{aligned}
i_q \omega_W &= \rho_{al} \rho^l_b d(W^{abnm} \epsilon_{nmpk} \xi^p \xi^k) + (\xi_a \kappa_b - \xi_b \kappa_a) d(W^{abnm} \epsilon_{nmpk} \xi^p \xi^k) + \\
&\quad + \frac{1}{2} W_{abij} \xi^i \xi^j d(W^{abnm} \epsilon_{nmpk} \xi^p \xi^k) + \\
&\quad + d\rho_{ab} (\xi^j W^{abnm} |_{|j} + \rho^a_j W^{jbnm} + \rho^b_j W^{ajnm} + \rho^n_j W^{abjm} + \\
&\quad + \rho^m_j W^{abnj} + 2\lambda W^{abnm}) \epsilon_{nmpk} \xi^p \xi^k + \\
&\quad + 2d\rho_{ab} W^{abnm} \epsilon_{nmpk} (\rho^p_j \xi^j \xi^k + \xi^p \lambda \xi^k) = \\
&= d(\rho_{al} \rho^l_b W^{abnm} \epsilon_{nmpk} \xi^p \xi^k) + (\xi_a \kappa_b - \xi_b \kappa_a) d(W^{abnm} \epsilon_{nmpk} \xi^p \xi^k) + \\
&\quad + \frac{1}{2} W_{abij} \xi^i \xi^j d(W^{abnm} \epsilon_{nmpk} \xi^p \xi^k) + d(\rho_{ab}) \xi^j W^{abnm} |_{|j} \epsilon_{nmpk} \xi^p \xi^k, \quad (\text{B.1})
\end{aligned}$$

where we made use of:

$$\begin{aligned}
\rho_{al} \rho^l_b d(W^{abnm} \epsilon_{nmpk} \xi^p \xi^k) + d\rho_{ab} (\rho^a_j W^{jbnm} + \rho^b_j W^{ajnm}) \epsilon_{nmpk} \xi^p \xi^k &= \\
&= d(\rho_{al} \rho^l_b W^{abnm} \epsilon_{nmpk} \xi^p \xi^k). \quad (\text{B.2})
\end{aligned}$$

For the second piece  $\omega_C$  of the presymplectic structure we have

$$\begin{aligned}
i_q \omega_C &= \rho_{an} \xi^n d(C^a_{bc} \epsilon^{bcpk} \xi_p \xi_k) + \xi_a \lambda d(C^a_{bc} \epsilon^{bcpk} \xi_p \xi_k) + \\
&+ d(\xi_a) (\xi^d D_d C^a_{abc} + \rho_a^n C_{nbc} + \rho_b^n C_{anc} + \rho_c^n C_{abn} + 3\lambda C_{abc} + \kappa_n W^n_{abc}) \epsilon^{bcpk} \xi_p \xi_k + \\
&\quad + 2d(\xi_a) C^a_{bc} \epsilon^{bcpk} (\rho_p^l \xi_l \xi_k + \xi_p \lambda \xi_k), \quad (\text{B.3})
\end{aligned}$$

where the following cancellations can be observed:

$$3d(\xi_a) \lambda C_{abc} \epsilon^{bcpk} \xi_p \xi_k + 2d(\xi_a) C^a_{bc} \epsilon^{bcpk} \xi_p \lambda \xi_k = d(\xi_a) \lambda C_{abc} \epsilon^{bcpk} \xi_p \xi_k \quad (\text{B.4})$$

and

$$d(\xi_a) (\rho_b^n C_{anc} + \rho_c^n C_{abn}) \epsilon^{bcpk} \xi_p \xi_k + 2d(\xi_a) C^a_{bc} \epsilon^{bcpk} \rho_p^l \xi_l \xi_k = 0. \quad (\text{B.5})$$

Taking them into account we indeed arrive at (3.37). To see that terms from equations (3.36),(3.39) give (3.40) we use properties (A.13), (A.12), (A.14), (A.15), (A.16). The nontrivial calculation is to show that:

$$d(\rho^{bc}) \xi^a \xi^p \xi^k W^*_{bcpk|a} = -2d(\rho^{bc}) \xi_c \xi^p \xi^k C^*_{bpk}. \quad (\text{B.6})$$

To prove it we start with (A.15):

$$\begin{aligned} W_{abcd}{}^a &= W_{bcda}{}^a - W_{bdca}{}^a = C_{bcd}, \\ W_{cdba}{}^a - W_{cbda}{}^a &= C_{cdb}, \\ W_{dbca}{}^a - W_{dcba}{}^a &= C_{dbc}, \end{aligned} \tag{B.7}$$

$$W_{bcda}{}^a = \frac{1}{2}(C_{bcd} + C_{cdb} - C_{dbc}) = (C_{bcd} + C_{cdb}), \tag{B.8}$$

$$W_{bc}{}^{pk|a} \epsilon_{pkda} = (C_b{}^{*pk} \epsilon_{pkcd} + C_c{}^{*pk} \epsilon_{pkdb}). \tag{B.9}$$

Multiply this equation by  $\epsilon^{dnmr}$  to get

$$\delta_{pka}^{nmr} W_{bc}{}^{pk|a} = (-C_b{}^{*pk} \delta_{pkc}^{nmr} + C_c{}^{*pk} \delta_{pkb}^{nmr}), \tag{B.10}$$

$$\xi^a \xi^p \xi^k W_{bcpk|a}^* = (-\xi_c \xi^p \xi^k C_{bpk}^* + \xi_b \xi^p \xi^k C_{cpk}^*). \tag{B.11}$$

Contracting this equation with  $d\rho^{bc}$  yields the desired relation.

We conclude that

$$\begin{aligned} i_q(\omega_W - 2\omega_C) &= d(\rho_a \rho^l{}_b W^{abnm} \epsilon_{nmpk} \xi^p \xi^k) + d(2\xi_a \kappa_b W^{abnm} \epsilon_{nmpk} \xi^p \xi^k) + \\ &+ d(-2\rho_{an} \xi^n C^a{}_{bc} \epsilon^{bcpk} \xi_p \xi_k) + d(-2\xi_a \lambda C^a{}_{bc} \epsilon^{bcpk} \xi_p \xi_k) + \frac{1}{2} W_{abij} \xi^i \xi^j d(W^{abnm} \epsilon_{nmpk} \xi^p \xi^k). \end{aligned} \tag{B.12}$$

## C The kernel of the presymplectic structure

Here we study the structure of the kernel of the presymplectic structure (3.33) and show the properties needed in the proof of Proposition 3.3. We first identify explicitly vector fields on  $F$  which are in the kernel of  $\omega$  and whose prolongations generate the kernel distribution for  $\bar{\omega}$  on  $\bar{F}$ . It is convenient to group them as follows:

- Vector fields along  $\rho^{ab}$  which we group according to the homogeneity in  $\xi$ .

We have:

$$Y_{ab}^{(4)} = \xi^{(4)} \frac{\partial}{\partial \rho_{ab}}, \quad Y_l^{(3)} = \xi_d^{(3)} \frac{\partial}{\partial \rho_{dl}}. \tag{C.1}$$

Next there are fields which are second order on  $\xi$ :

$$Y_{ld}^{(2ant)} = \epsilon_{abcd} \xi^c \xi_l \frac{\partial}{\partial \rho_{ab}}, \quad Y_{ld}^{(2trace)} = \xi^c \xi_l \delta_{cd}^{ab} \frac{\partial}{\partial \rho_{ab}}. \tag{C.2}$$

Note that these are linearly dependent as  $Y_{ld}^{(2ant)} \epsilon^{ldkp} = Y^{(2trace)kp}$ . The linear in  $\xi$  ones are:

$$\begin{aligned} Y_d^{(ant)} &= \epsilon_{abcd} \xi^c \frac{\partial}{\partial \rho_{ab}} + W_{dnm}{}^a \epsilon^{nmbc} \frac{\partial}{\partial C^{abc}}, \\ Y_d^{(trace)} &= \xi^c \delta_{cd} \frac{\partial}{\partial \rho_{ab}} - W^{adbc} \frac{\partial}{\partial C^{abc}}. \end{aligned} \quad (C.3)$$

To see that  $Y_d^{(ant)}$  belong to the kernel we observe that:

$$\begin{aligned} i_{\epsilon_{abcd} \xi^c \frac{\partial}{\partial \rho_{ab}}} \omega_W &= \epsilon_{abcd} \xi^c d(W^{abnm} \epsilon_{nmpk} \xi^p \xi^k) = \\ &= \xi^c W_{cdab} d(\xi^a \xi^b) = 2W_{dcba} d(\xi^a) \xi^c \xi^b. \end{aligned} \quad (C.4)$$

This term cancels with

$$i_{Y_d^{(ant)}}(-2\omega_C) = -2(d(\xi^a) W_{dnma} \epsilon^{nmbc} \epsilon_{bcpk} \xi^p \xi^k). \quad (C.5)$$

The considerations for  $Y_d^{(trace)}$ ,  $Y_{ld}^{(2ant)}$ , and  $Y_{ld}^{(2trace)}$  are pretty much the same.

- Vector fields along  $\xi^a$ . We also group them according to the homogeneity in  $\xi$ :

$$X_a^{(4)} = \xi^{(4)} \frac{\partial}{\partial \xi_a}, \quad X^{(3)} = \xi_a^{(3)} \frac{\partial}{\partial \xi_a}. \quad (C.6)$$

Second order ones :

$$\begin{aligned} X_d^{(2ant)} &= \epsilon_{abcd} \xi^b \xi^c \frac{\partial}{\partial \xi_a} - \xi_d W^{ab}{}_{pk} \epsilon^{pknm} \frac{\partial}{\partial W^{abnm}} - \\ &\quad - 4\xi_d C^a{}_{pk} \epsilon^{pkbc} \frac{\partial}{\partial C^{abc}} - 4\xi^a C_{pkd} \epsilon^{bcpk} \frac{\partial}{\partial C^{abc}}, \\ X_d^{(2trace)} &= \xi_b \xi_c \delta_{ad} \frac{\partial}{\partial \xi_a} - \xi_d W^{abnm} \frac{\partial}{\partial W^{abnm}} - \\ &\quad - 6\xi_d C^{abc} \frac{\partial}{\partial C^{abc}}. \end{aligned} \quad (C.7)$$

To see that they are in the kernel we observe the following relations:

$$\begin{aligned} i_{\epsilon_{aljd} \xi^l \xi^j \frac{\partial}{\partial \xi_a}} \omega_W &= 2d(\rho_{ab}) W^{abnm} \epsilon_{nmpk} \epsilon^{pljd} \xi_l \xi_j \xi^k = \\ &= 2d(\rho_{ab}) W^{abnm} \delta_{nmk}^{ljd} \xi_l \xi_j \xi^k = 2d(\rho_{ab}) W^{abnm} \xi_n \xi_m \xi^d, \end{aligned} \quad (C.8)$$

$$\begin{aligned}
i_{\epsilon_{aljd}\xi^l\xi^j}\frac{\partial}{\partial\xi_a}\omega_C &= \epsilon_{aljd}\xi^l\xi^j d(C^a_{bc}\epsilon^{bcpk}\xi_p\xi_k) + 2d(\xi_a)C^a_{bc}\epsilon^{bcpk}\epsilon_{pljd}\xi^l\xi^j\xi_k = \\
&= 2\delta_{ad}^{pk}d(C^a_{bc})\epsilon^{bcpk} + 2\delta_{aljd}^{bcpk}\xi^l\xi^j C^a_{bc}d(\xi_p)\xi_k + 2d(\xi_a)C^a_{bc}\delta_{ljd}^{bck}\xi^l\xi^j\xi_k = \\
&= 4\xi^b\xi^c C^a_{bc}d(\xi_a)\xi_d + 4\xi_a C^a_{bd}\xi^b\xi^k d(\xi_k). \quad (C.9)
\end{aligned}$$

And finally the first order one is given by:

$$X^{(trace)} = \xi^a \frac{\partial}{\partial\xi_a} - 2W^{abnm} \frac{\partial}{\partial W^{abnm}} - 3C^{abc} \frac{\partial}{\partial C^{abc}}. \quad (C.10)$$

The proof that it is the kernel is based on:

$$\begin{aligned}
i_{\xi_a}\frac{\partial}{\partial\xi_a}\omega &= 2d(\rho_{ab})W^{abnm}\epsilon_{nmpk}\xi^p\xi^k - \\
&\quad - 2(\xi_a d(C^a_{bc}\epsilon^{bcpk}\xi_p\xi_k) + 2d(\xi_a)C^a_{bc}\epsilon^{bcpk}\xi_p\xi_k). \quad (C.11)
\end{aligned}$$

The first term is canceled by  $i_{-2W^{abnm}\frac{\partial}{\partial W^{abnm}}}\omega_W$ . Then  $\xi_a d(C^a_{bc})\epsilon^{bcpk}\xi_p\xi_k = 0$  due to Bianchi identity. In terms of  $C^{apk} = C^a_{bc}\epsilon^{bcpk}$  the second term gives:

$$\begin{aligned}
\xi_a C^a_{bc}\epsilon^{bcpk}d(\xi_p)\xi_k - \xi_a C^a_{bc}\epsilon^{bcpk}\xi_p d(\xi_k) &= C^{apk} (\xi_a d(\xi_p)\xi_k - \xi_a \xi_p d(\xi_k)) = \\
&= C^{apk} \xi_a d(\xi_p)\xi_k + C^{apk} \xi_p d(\xi_k)\xi_a = (C^{apk} + C^{kpa}) (\xi_a d(\xi_p)\xi_k) = \\
&= (-C^{pka}) (\xi_a d(\xi_p)\xi_k) = C^{pka} (d(\xi_p)\xi_k \xi_a). \quad (C.12)
\end{aligned}$$

Together with the third terms in (C.11) they can be taken care of by the  $-3C^{abc}\frac{\partial}{\partial C^{abc}}$ .

- Finally we have two more independent kernel vector fields

$$L = \frac{\partial}{\partial\lambda}, \quad K = \frac{\partial}{\partial\kappa}, \quad (C.13)$$

simply because  $d\kappa^a$  nor  $d\lambda$  enter (3.33).

Now we consider the symplectic structure  $\bar{\omega}$  on  $\bar{F}$  at a point  $p$ , where all the coordinates but  $e_\mu^a$  are set to zero. At this point  $\bar{\omega}$  reads as:

$$\begin{aligned}
\bar{\omega}_p &= (dw_{ab}d^2W^{abcd}_{,cd} + d\omega_{ab,c}d^1W^{abcd}_{,d} + d\omega_{ab,cd}dW^{abcd}) - \\
&\quad - 2(d\epsilon_a d^2C^{acd}_{,cd} + de_{a,c}d^1C^{acd}_{,d} + de_{a,cd}dC^{acd}), \quad (C.14)
\end{aligned}$$

in a suitable coordinate system. By inspecting the prolongation of the above kernel vector fields  $X, Y, K, L$  at this point one finds that their linear span coincides with the kernel. The argument given in the main text then implies that their prolongations generate the kernel distribution of  $\bar{\omega}$ .

## D Metric elimination

Here we discuss the assumptions under which the metric can be eliminated. As we discuss in the main text the BRST complex of CGR as presented in [16] contain ghost degree 1 elements  $\xi_{\mu}^{\nu}$  and the metric. Thanks to

$$qg_{\mu\nu} = 2\omega g_{\mu\nu} + \xi_{\mu}^{\rho} g_{\rho\nu} + \xi_{\nu}^{\rho} g_{\mu\rho} = 2\omega g_{\mu\nu} + \xi_{(\mu\nu)}, \quad (\text{D.1})$$

$g_{\mu\nu}$  and  $\xi_{(\mu\nu)}$  can be eliminated at least locally.

However,  $g_{\mu\nu}$  are not unconstrained coordinates in the sense of definition 3.1 because the determinant of the metric can not be zero. Moreover, the space is further restricted to the space of symmetric matrices of definite signature.

It turns out that in the case of Riemannian metric (i.e. Euclidean signature) the metric can be eliminated. To see this we employ Cholesky decomposition which states that every symmetric, positive-definite matrix can be uniquely decomposed as

$$g = LL^T, \quad (\text{D.2})$$

where  $L$  is a lower-triangular matrix with positive elements on the diagonal. The space of such matrices is diffeomorphic to  $R^{n(n+1)/2}$  and the concrete diffeomorphism can be obtained by taking the coordinates on  $R^{n(n+1)/2}$  to be the entries of a lower-triangular matrix  $A$  and taking  $L = e^A$ . Entries of the matrix  $A$  then become the free coordinates parameterizing the space of metrics and form contractible pairs with the symmetric part of  $\xi_{\mu\nu}$ .

In the case of pseudo-Euclidean metrics there might be no such decomposition. We will make use of the following theorem [41] Th. 4.1.1, 4.1.2

**Theorem 1.** *If all the leading principal minors of  $A \in R^{n \times n}$  are nonvanishing then there exist a unique lower triangular matrix  $L$  with positive diagonal values and a unique diagonal matrix  $D = \text{diag}(d_1, \dots, d_n)$ ,  $d_k = \pm 1$  such that  $A = LDL^T$ .*

A metric in general coordinates might not satisfy Theorem 1. For example take the metric on a plane in  $z$  coordinates:  $ds^2 = dzd\bar{z}$ . If we restrict to such metrics that admits such factorization the elimination can still be performed. This subspace of metrics includes important cases, for example all metrics close to Minkowsky one.

## E Kaku structure

The simple approach one might try is to consider just the conformal algebra with shifted grading  $g[1]$  as the target space with the canonical Chevalie-Eilenberg differential  $q_{CE}$  on it. The presymplectic potential and structure can be read off the action (3.10) presented in [11]

$$\begin{aligned}\chi &= (d\rho_{ab}\xi_c\kappa_d)\epsilon^{abcd} = (\xi_a\kappa_b d\rho_{cd})\epsilon^{abcd}, \\ \omega_{Kaku} &= d\chi = (d\xi_a\kappa_b d\rho_{cd} - \xi_a d\kappa_b d\rho_{cd})\epsilon^{abcd}.\end{aligned}\tag{E.1}$$

This structure is invariant under  $q_{CE}$  and induces the following action:

$$S = \int (d\omega_{ab}e_c f_d + \omega_a{}^k \omega_{kb} e_c f_d + e_a f_b e_c f_d)\epsilon^{abcd}.\tag{E.2}$$

This procedure reproduces the action (3.10) but for it to be equivalent to the conformal gravity in metric formulation one has to additionally impose the zero torsion condition as was discussed previously.

## References

- [1] M. Alexandrov, M. Kontsevich, A. Schwartz, and O. Zaboronsky, “The Geometry of the master equation and topological quantum field theory,” *Int.J.Mod.Phys.* **A12** (1997) 1405–1430, hep-th/9502010.
- [2] K. B. Alkalaev and M. Grigoriev, “Frame-like Lagrangians and presymplectic AKSZ-type sigma models,” *Int. J. Mod. Phys.* **A29** no. 18, (2014) 1450103, arXiv:1312.5296 [hep-th].

- [3] J. Kijowski and W. M. Tulczyjew, *A symplectic framework for field theories* . 1979.
- [4] C. Crnkovic and E. Witten, “Covariant Description Of Canonical Formalism In Geometrical Theories,”. in Three hundred years of gravitation, S. W. Hawking and W. Israel, eds., pp. 676-684. Cambridge University Press, Cambridge, 1987.
- [5] I. Khavkine, “Presymplectic current and the inverse problem of the calculus of variations,” *J. Math. Phys.* **54**, (Oct., 2012) 111502, 1210.0802.
- [6] A. A. Sharapov, “Variational tricomplex, global symmetries and conservation laws of gauge systems,” arXiv:1607.01626 [math-ph].
- [7] M. Grigoriev and A. Kotov, “Presymplectic AKSZ formulation of Einstein gravity,” *JHEP* **09** (2021) 181, arXiv:2008.11690 [hep-th].
- [8] I. Batalin and G. Vilkovisky, “Gauge Algebra and Quantization,” *Phys.Lett.* **B102** (1981) 27–31.
- [9] G. Canepa, A. S. Cattaneo, and M. Schiavina, “General Relativity and the AKSZ construction,” arXiv:2006.13078 [math-ph].
- [10] A. Sharapov and E. Skvortsov, “Higher spin gravities and presymplectic AKSZ models,” *Nucl. Phys. B* **972** (2021) 115551, arXiv:2102.02253 [hep-th].
- [11] M. Kaku, P. K. Townsend, and P. van Nieuwenhuizen, “Gauge Theory of the Conformal and Superconformal Group,” *Phys. Lett.* **B69** (1977) 304–308.
- [12] J. Maldacena, “Einstein Gravity from Conformal Gravity,” arXiv:1105.5632 [hep-th].
- [13] J. T. Trujillo, *Weyl Gravity as a Gauge Theory*. PhD thesis, Utah State U., 2013.

- [14] P. D. Mannheim, “Making the Case for Conformal Gravity,”  
*Found. Phys.* **42** (2012) 388–420, arXiv:1101.2186 [hep-th].
- [15] L. Rachwał, “Introduction to Quantization of Conformal Gravity,”  
*Universe* **8** no. 4, (2022) 225, arXiv:2204.13856 [hep-th].
- [16] N. Boulanger, “A Weyl-covariant tensor calculus,”  
*J. Math. Phys.* **46** (2005) 053508, arXiv:hep-th/0412314.
- [17] N. Boulanger, “General solutions of the Wess-Zumino consistency condition for the Weyl anomalies,” *JHEP* **07** (2007) 069,  
arXiv:0704.2472 [hep-th].
- [18] N. Boulanger, “Algebraic Classification of Weyl Anomalies in Arbitrary Dimensions,” *Phys. Rev. Lett.* **98** (2007) 261302,  
arXiv:0706.0340 [hep-th].
- [19] E. Joung, M.-g. Kim, and Y. Kim, “Unfolding conformal geometry,”  
*JHEP* **12** (2021) 092, arXiv:2108.05535 [hep-th].
- [20] C. Preitschopf and M. A. Vasiliev, “Conformal field theory in conformal space,” *Nucl.Phys.* **B549** (1999) 450–480, arXiv:hep-th/9812113.
- [21] M. Grigoriev and A. Kotov, “Gauge PDE and AKSZ-type Sigma Models,” *Fortsch. Phys.* (2019), arXiv:1903.02820 [hep-th].
- [22] G. Barnich and M. Grigoriev, “First order parent formulation for generic gauge field theories,” *JHEP* **01** (2011) 122,  
arXiv:1009.0190 [hep-th].
- [23] M. Grigoriev, “Parent formulation at the Lagrangian level,”  
*JHEP* **07** (2011) 061, arXiv:1012.1903 [hep-th].
- [24] A. Kotov and T. Strobl, “Characteristic classes associated to Q-bundles,”  
*Int. J. Geom. Meth. Mod. Phys.* **12** no. 01, (2014) 1550006,  
arXiv:0711.4106 [math.DG].
- [25] G. Barnich, F. Brandt, and M. Henneaux, “Local BRST cohomology in the antifield formalism. I. General theorems,” *Commun. Math. Phys.* **174** (1995) 57–92, hep-th/9405109.

- [26] F. Brandt, “Local BRST Cohomology and Covariance,” *Commun. Math. Phys.* **190** (1997) 459–489, hep-th/9604025.
- [27] M. Grigoriev, “Parent formulations, frame-like Lagrangians, and generalized auxiliary fields,” *JHEP* **1212** (2012) 048, arXiv:1204.1793 [hep-th].
- [28] G. Barnich, F. Brandt, and M. Henneaux, “Local BRST cohomology in Einstein Yang-Mills theory,” *Nucl. Phys.* **B455** (1995) 357–408, hep-th/9505173.
- [29] M. Grigoriev, K. Mkrtchyan, and E. Skvortsov, “Matter-free higher spin gravities in 3D: Partially-massless fields and general structure,” *Phys. Rev. D* **102** no. 6, (2020) 066003, arXiv:2005.05931 [hep-th].
- [30] R. Stora, “Algebraic structure and topological origin of anomalies,” Seminar given at Cargese Summer Inst.: Progress in Gauge Field Theory, Cargese, France, Sep 1-15, 1983.
- [31] J. Manes, R. Stora, and B. Zumino, “Algebraic Study of Chiral Anomalies,” *Commun. Math. Phys.* **102** (1985) 157.
- [32] M. A. Vasiliev, “Equations of Motion of Interacting Massless Fields of All Spins as a Free Differential Algebra,” *Phys. Lett.* **B209** (1988) 491–497.
- [33] M. Henneaux, “Elimination of the Auxiliary Fields in the Antifield Formalism,” *Phys. Lett.* **B238** (1990) 299.
- [34] G. Barnich, F. Brandt, and M. Henneaux, “Local BRST cohomology in gauge theories,” *Phys.Rept.* **338** (2000) 439–569, hep-th/0002245.
- [35] F. Brandt, “Jet coordinates for local BRST cohomology,” *Lett. Math. Phys.* **55** (2001) 149–159, arXiv:math-ph/0103006.
- [36] M. Grigoriev, “Presymplectic structures and intrinsic Lagrangians,” arXiv:1606.07532 [hep-th].

- [37] M. Grigoriev and V. Gritzaenko, “Presymplectic structures and intrinsic Lagrangians for massive fields,”  
*Nuclear Physics B* **975** (9, 2021) 115686,  
arXiv:2109.05596 [hep-th].
- [38] M. Grigoriev and A. A. Tseytlin, “On conformal higher spins in curved background,” *J. Phys.* **A50** no. 12, (2017) 125401,  
arXiv:1609.09381 [hep-th].
- [39] V. E. Lopatin and M. A. Vasiliev, “Free Massless Bosonic Fields of Arbitrary Spin in  $d$ -dimensional De Sitter Space,”  
*Mod. Phys. Lett.* **A3** (1988) 257.
- [40] N. Boulanger and J. Erdmenger, “A Classification of local Weyl invariants in  $D=8$ ,” *Class. Quant. Grav.* **21** (2004) 4305–4316,  
arXiv:hep-th/0405228 [hep-th].
- [41] G. H. Golub and C. F. Van Loan, *Matrix Computations*. The Johns Hopkins University Press, third ed., 1996.