

Born-Oppenheimer and the Geometry of Ray Space

Joseph Samuel

International Centre for Theoretical Sciences, Bangalore, India 560089
and
Raman Research Institute, Bangalore, India 560080

August 8, 2022

Abstract

We consider a simple toy model to study the Born-Oppenheimer approximation. The model describes a slow neutron interacting with a strong, spatially varying magnetic field. Regarding the neutron spin as a fast degree of freedom, we work out the slow dynamics of the motion of the neutron. As expected, the neutron experiences an effective scalar potential V_{dyn} coming from the eigenvalue of the spin and an effective magnetic field coming from the Berry phase vector potential A . We draw attention to an additional potential V_{geom} originating in the geometry of ray space and the Fubini-Study metric. While this term is sometimes mentioned in the Born-Oppenheimer literature, its geometric origin has not, to our knowledge, been appreciated. We describe a magnetic field configuration to isolate this “geometric scalar potential” in experiments with slow neutrons. We also note that there are optical analogues in which the experiments may be easier to perform.

email: sam@icts.res.in, sam@rri.res.in

1 Introduction

Berry's work[1] has illuminated the geometry underlying the ray space of quantum mechanics. The seminal ideas in [1] led to numerous experiments (for example, [2, 3, 4]), applications and generalisations[5, 6]. It was also realised that the idea had been anticipated several times[7, 8] in different contexts. Michael Berry traces the seed of the idea as far back as William Rowan Hamilton (1832) in his theoretical studies on conical double refraction and their experimental confirmation by Herbert Lloyd. One of the anticipations was due to Longuet-Higgins[9] in the context of the Born-Oppenheimer approximation which is widely used in the quantum theory of lattice dynamics and molecular physics. The Born-Oppenheimer approximation separates the fast electronic motion from the slow nuclear motion. Initially the motion of the nuclei is neglected and the frozen nuclear coordinates are treated as parameters in the electronic problem. Invoking the adiabatic theorem to allow for slow motion of the nuclei, one then uses a reduced Schrödinger equation to solve for the nuclear motion.

Let us motivate our discussion with a simple classical example . Consider a well oiled bicycle wheel¹ mounted on an axle rod so that it is free to rotate about an axis pointing along the rod. If the direction of the rod is held fixed, and the wheel spun at angular velocity ω the wheel rotates through an angle ωt in time t (neglecting friction). If the direction of the rod is slowly changed to traverse a closed circuit C in the space of directions, we may naively expect that the total angle of rotation is still ωt . However, this naive expectation is wrong. The total angle of rotation includes an additional piece given by the solid angle traversed by the circuit C . The explanation is that the wheel rotates about different axes during the circuit and such rotations do not commute. This effect is the Hannay angle[10], a classical analogue of the Berry phase. To perform this experiment, it helps to mark the rim of the bicycle wheel and use a stroboscope or camera to track the motion of the wheel.

So far we have regarded the direction of the rod as a parameter within the control of the experimenter. However, parameters are also dynamical variables, subject to the laws of motion. To realise this, let us suspend the wheel and axle from the ceiling and repeat the

¹The precession and nutation of a rapidly spinning top can also be viewed as slow modes subject to a monopole magnetic field in exactly the same way.

experiment by spinning the wheel and setting the axle pendulum in motion. We find that the trajectory of the axle resembles that of a charged particle in a magnetic field. With more work one can arrange for the axle to trace its path in a sand pit and see precessing ellipses. Reversing the sense of the spin reverses the sense of the precession. What we are seeing here, in the spirit of the Born-Oppenheimer approximation, is the back reaction of the fast mode (the spin of the bicycle wheel) on the slow modes (the oscillation of the pendulum) of the system. A less elaborate experiment, less convincing, but more in tune with the sloppy habits of theoretical physicists, is to twist a wet tea bag so its string is wound up and let it untwist while oscillating, suspended from its string. Michael Berry has been subjected to both these demonstrations on one of his visits to the Raman Research Institute.

In this paper we consider a simple quantum mechanical toy model² to study the Born-Oppenheimer approximation. This quantum model is described in section II. In section III, we derive the dynamics of the slow modes taking into account the back reaction of the fast mode. There are two effects which are well known: a scalar potential coming from the eigenvalue of the fast mode and an effective magnetic field coming from the Berry potential. We find also a third effect: an extra scalar potential, which is the main interest of this paper. In section IV we describe some geometrical aspects of ray space to bring out the geometric nature of the extra scalar potential. Section V describes magnetic field configurations which bring out the effect of different terms in the slow Hamiltonian. Section VI shows how the geometrical ideas of the toy model carry over to the more realistic model of molecular dynamics. Section VII is a concluding discussion.

Notation: We use Dirac's bracket notation for the fast mode but not for the slow modes. Projectors onto one dimensional subspaces (proportional to the vector $|e\rangle$) of the fast Hilbert space are denoted P^e and satisfy $P^e P^e = P^e$. The orthogonal projection is $Q^e = \mathbb{1} - P^e$. We will use a global space independent frame $\{|+\rangle, |-\rangle\}$ as well as a local frame $\{|e_+(\vec{R})\rangle, |e_-(\vec{R})\rangle\}$, which is space dependent. Spatial derivatives, $\frac{\partial}{\partial R^i}$ will be denoted ∇_i . The projectors P^+ and $Q^+ = P^- = \mathbb{1} - P^+$, commute with ∇_i , since the basis $\{|+\rangle, |-\rangle\}$ is independent of space.

²Another toy model has been discussed in [11]

2 A Toy Model

We consider a slow neutron in a strong external magnetic field $\vec{B}(\vec{R})$. Let us regard the spin as the fast variable and the spatial position \vec{R} of the neutron as the slow variable. The total Hamiltonian of our system is the sum

$$H_T = H_f + H_s \quad . \quad (1)$$

The fast Hamiltonian H_f is that of a spin half in a magnetic field

$$H_f = -\mu \vec{B}(\vec{R}) \cdot \vec{\sigma} \quad , \quad (2)$$

where $\vec{B}(\vec{R})$ is spatially varying magnetic field and μ is the magnetic dipole moment of the neutron. We will assume that $|\vec{B}(\vec{R})|$ does not vanish in the region of interest. In the spirit of the Born-Oppenheimer approximation, we will assume that the magnetic field $\vec{B}(\vec{R})$ varies slowly with \vec{R} . (More precisely, we would like $\frac{d\vec{B}}{dt} \cdot \vec{\nabla}(\hat{B}(\vec{R})) \ll \frac{\mu|\vec{B}(\vec{R})|}{\hbar}$, where \hat{B} is a unit vector and $\frac{d\vec{R}}{dt}$ a typical velocity of the slow neutron .)

The slow Hamiltonian H_s is given by the kinetic energy of the neutron

$$H_s = -\frac{\hbar^2}{2m} \vec{\nabla} \cdot \vec{\nabla} \quad . \quad (3)$$

We now regard \vec{R} as a parameter in the fast Hamiltonian $H_f(\vec{R})$ and diagonalise the fast Hamiltonian. (In molecular physics, this corresponds to holding the nuclear configuration fixed and solving for the electronic eigenstates.) Let the total wave function of the system be

$$\Psi(\vec{R}) = \begin{bmatrix} \psi_+(\vec{R}) \\ \psi_-(\vec{R}) \end{bmatrix} = \psi_+(\vec{R}) |+\rangle + \psi_2(\vec{R}) |-\rangle, \quad (4)$$

where

$$|+\rangle = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad |-\rangle = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \quad (5)$$

We will refer to the orthonormal basis $\{|+\rangle, |-\rangle\}$ as the global frame. We will also need a local frame adapted to the magnetic field. To do this, let us solve the eigenvalue equation

$$H_f(\vec{R}) |e(\vec{R})\rangle = \lambda |e(\vec{R})\rangle \quad (6)$$

for the fast spin degree of freedom.

This equation has two independent solutions $\{|e_+(\vec{R})\rangle, |e_-(\vec{R})\rangle\}$ (satisfying $\langle e_+(\vec{R})|e_-(\vec{R})\rangle = 0$) corresponding to the two distinct eigenvalues $\{\lambda_+ = -\mu|B|, \lambda_- = \mu|B|\}$. For normalised eigenstates $|e_\pm(\vec{R})\rangle$, equations (6) determine $|e_a(\vec{R})\rangle, a = \{+, -\}$ up to a phase factor $w_a(\vec{R}), |w_a(\vec{R})| = 1$ a complex number of modulus unity, which depends on a as well as \vec{R} . Let us make an arbitrary smooth choice of these phases³. We now have $\{|e_+(\vec{R})\rangle, |e_-(\vec{R})\rangle\}$, which defines a local orthonormal basis which is adapted to the magnetic field and provides us with a moving frame. We refer to this as the local basis.

The two orthonormal bases $|a\rangle$ and $|e_a(\vec{R})\rangle$ are related by a Unitary transformation $U(\vec{R})$

$$|e_a(\vec{R})\rangle = U(\vec{R}) |a\rangle \quad (7)$$

We can transform our original eigenvalue problem (6) by defining $\Psi(\vec{R}) = U(\vec{R})\Phi(\vec{R})$. The new eigenvalue equation is

$$\mathcal{H}_T\Phi = E\Phi, \quad (8)$$

where $\mathcal{H}_T = U^{-1}H_TU$ is given by

$$\mathcal{H}_T = \lambda_+(\vec{R})\sigma_3 - \frac{\hbar^2}{2m}U^\dagger(\vec{R})\nabla_i\nabla_iU(\vec{R}) \quad (9)$$

³This is always possible locally.

Let us now work out the dynamics governing the slow modes of the system.

3 The Reduced Hamiltonian

Let us assume that the spin is in the ground state $|+\rangle$ of the fast Hamiltonian. The adiabatic theorem assures us that (with an exponentially small chance of transition) the spin remains in the ground state. We can now pass to the reduced space by projecting the wave function and the Hamiltonian into the $|+\rangle$ subspace. $\phi(\vec{R}) = \langle + | \Phi(\vec{R}) \rangle$. The spin state can be described by the density matrix $\rho_f = (1 + \sigma_3)/2 = |+\rangle \langle +| = P^+$, where P^+ is the one dimensional projector on to the ground state. The reduced Hamiltonian for the slow degrees of freedom is $\mathcal{H}_{red} = Tr[\mathcal{H}_T \rho_f] = \langle + | \mathcal{H}_T | + \rangle$.

Writing out the reduced Hamiltonian, we have for any wave function $\xi(\vec{R})$

$$\mathcal{H}_{red}\xi(\vec{R}) = \lambda_+(\vec{R}) \langle + | \sigma_3 | + \rangle \xi(\vec{R}) - \frac{\hbar^2}{2m} \langle + | U^\dagger(\vec{R}) \nabla_i \nabla_i U(\vec{R}) | + \rangle \xi(\vec{R}), \quad (10)$$

where it is understood that the derivative operators act on everything to their right unless explicitly bracketed.

3.1 Dynamical Scalar Potential

The first term in \mathcal{H}_{red} , gives us the fast eigenvalue which gives an effective space dependent scalar potential $V_{dyn}(\vec{R}) = -\mu |\vec{B}(\vec{R})|$ for the slow modes. We call this the dynamical scalar potential as it is related to the eigenvalue of the fast Hamiltonian. This term is expected: in molecular physics, the electronic eigenvalue serves as a scalar potential in which the slow nuclei move. Adiabatically moving a neutron to a region of lower magnetic field requires the mover to do some work.

3.2 Geometric Vector Potential

The second term in $\mathcal{H}_{red}\xi(\vec{R})$ can be rewritten as

$$-\frac{\hbar^2}{2m} \langle +|U^\dagger(\vec{R})\nabla_i U(\vec{R})U^\dagger(\vec{R})\nabla_i U(\vec{R})|+ \rangle \xi(\vec{R}) \quad (11)$$

by inserting the identity $UU^\dagger = 1$ between the derivatives. It is elementary to verify that for any function $\xi(\vec{R})$

$$U^\dagger(\vec{R})\nabla_i U(\vec{R}) \xi(\vec{R}) = \nabla_i \xi(\vec{R}) + \omega_i(\vec{R})\xi(\vec{R}) \quad (12)$$

where $\omega_i(\vec{R}) = U^\dagger(\vec{R})\nabla_i U(\vec{R})$ is an effective $U(2)$ gauge field (with vanishing $U(2)$ field strength). Writing $\mathcal{D}_i = \nabla_i + \omega_i$ as the covariant derivative in the $U(2)$ gauge field, we have

$$Tr[P^+\mathcal{D}_i\mathcal{D}_i] = \langle +|\mathcal{D}_i\mathcal{D}_i|+ \rangle = \langle +|\mathcal{D}_i|+ \rangle \langle +|\mathcal{D}_i|+ \rangle + \langle +|\mathcal{D}_i P^-\mathcal{D}_i|+ \rangle, \quad (13)$$

where $P^- = (\mathbb{1} - P^+)$. For any function $\xi(\vec{R})$, $\langle +|\mathcal{D}_i|+ \rangle \xi(\vec{R}) =$

$$\nabla_i \xi(\vec{R}) + \langle +|\omega_i|+ \rangle \xi(\vec{R}) = \nabla_i \xi(\vec{R}) + [\langle +|U^\dagger(\vec{R})\nabla_i U(\vec{R})|+ \rangle] \xi(\vec{R}) = \quad (14)$$

$$\nabla_i \xi(\vec{R}) + [\langle e_+(\vec{R})|\nabla_i|e_+(\vec{R}) \rangle] \xi(\vec{R}) = \nabla_i \xi(\vec{R}) + A_i \xi(\vec{R}) \quad (15)$$

where $A_i = \langle e_+(\vec{R})|\nabla_i|e_+(\vec{R}) \rangle$ is the Berry potential, which we also refer to as A_{geom} the geometric vector potential. As a result the first term in (11) gives us

$$(\nabla_i + A_i)(\nabla_i + A_i)\xi(\vec{R}) \quad (16)$$

This term shows that the slow modes are subject to an effective magnetic field represented by Berry potential A_i . This is an expected result, which has been discussed in the molecular physics literature[12]. This is the quantum analogue of the classical effect described in the introduction of this paper.

3.3 Geometric Scalar Potential

Like the Berry potential, the second term in (13) also has a simple geometric interpretation and this is the main point of the present paper. Expanding the expression for the covariant derivative, we have

$$\langle + | (\nabla_i + \omega_i) P^- (\nabla_i + \omega_i) | + \rangle \quad (17)$$

Since ∇_i commutes with P^- , all the derivative terms drop out in (17) leaving us with $\langle + | \omega_i P^- \omega_i | + \rangle = - \langle + | \omega_i^\dagger P^- \omega_i | + \rangle$ which equals

$$- \langle + | (\nabla_i U^\dagger(\vec{R})) U(\vec{R}) P^- U^\dagger(\vec{R}) (\nabla_i U(\vec{R})) | + \rangle = - (\nabla_i \langle e_+(\vec{R}) |) U(\vec{R}) P^- U^\dagger(\vec{R}) (\nabla_i | e_+(\vec{R}) \rangle) \quad (18)$$

$$= - (\nabla_i \langle e_+(\vec{R}) |) (\mathbb{1} - | e_+(\vec{R}) \rangle \langle e_+(\vec{R}) |) (\nabla_i | e_+(\vec{R}) \rangle) = - (\nabla_i \langle e_+(\vec{R}) |) Q^{e+} (\nabla_i | e_+(\vec{R}) \rangle) \quad (19)$$

where $Q^{e+} = P^{e-} = \mathbb{1} - P^{e+}$ is the projector orthogonal to the state $| e_+(\vec{R}) \rangle$. Supplying the necessary constants, we find the final expression for the geometric scalar potential

$$V_{geom}(\vec{R}) = \frac{\hbar^2}{2m} \{ (\nabla_i \langle e_+(\vec{R}) |) (\nabla_i | e_+(\vec{R}) \rangle) - (\nabla_i \langle e_+(\vec{R}) |) | e_+(\vec{R}) \rangle \langle e_+(\vec{R}) | (\nabla_i | e_+(\vec{R}) \rangle) \}, \quad (20)$$

This potential is non-negative and vanishes only when the *direction* of \vec{B} is constant. In the next section we will bring out the geometry behind these terms in the slow Hamiltonian.

4 Geometry of Ray Space

The ray space of quantum mechanics \mathcal{R} is defined as the set of normalised states modulo multiplication by a unit complex number, a phase. For finite dimensional Hilbert spaces, this is just complex projective space $\mathbb{C}\mathbb{P}^n$. In the toy model discussed here $n = 1$ and we have $\mathbb{C}\mathbb{P}^1 = S^2$. \mathcal{R} has three intertwining structures on it: a Riemannian structure, a symplectic structure and a complex structure. These three structures combine to form a Kähler structure.

The Riemannian structure is physically accessible as the overlap between quantum states:

$|\langle\phi|\psi\rangle|^2$ represents a transition probability. Geometrically, the same quantity is the distance between rays. Passing to infinitesimally separated rays $|\phi\rangle = |\psi\rangle + \delta|\psi\rangle$, we arrive at the Fubini-Study metric [13] for normalised states.

$$ds^2 = \langle\delta\psi|\delta\psi\rangle - \langle\delta\psi|\psi\rangle\langle\psi|\delta\psi\rangle \quad (21)$$

where we choose representative states from each ray. Given a normalised $|\psi\rangle$ and two vectors $\delta_1|\psi\rangle$ and $\delta_2|\psi\rangle$, the Fubini-Study metric is

$$\gamma(\delta_1|\psi\rangle, \delta_2|\psi\rangle) = \langle\delta_1\psi|\delta_2\psi\rangle - \langle\delta_1\psi|\psi\rangle\langle\psi|\delta_2\psi\rangle \quad (22)$$

The geometric scalar potential (20) is exactly of this form with $\delta|\psi\rangle$ replaced by $\nabla_i|e_+(\vec{R})\rangle$.

Although we have used representative kets instead of rays, this is actually a metric on the ray space. This metric can also be expressed directly in terms of density matrices. If ρ is a density matrix and $\dot{\rho}$ and ρ' tangent vectors at ρ , the Fubini-Study metric is given by

$$\gamma(\dot{\rho}, \rho') = \frac{1}{2}\text{Tr}[\dot{\rho}\rho'] \quad (23)$$

The symplectic structure is physically manifested in the geometric phase which is detectable in interference phenomena. The symplectic form on ray space is the curvature of the Berry connection and acts like a magnetic field. Given any closed curve C in ray space, the Berry phase is $\exp i\phi$, where ϕ is the symplectic area of any disc that C bounds. The Berry phase for open curves can be similarly defined (provided the ends of the curve are not on orthogonal rays) by closing the ends by a geodesic curve. The geodesic in ray space is determined by the Fubini-Study metric. The work of Pancharatnam[7] in optics drew attention to the importance of geodesics. These ideas were carried over to the Hilbert space of quantum mechanics in [14].

The complex structure is intimately tied up with the symmetry of time reversal. Time reversal is an anti-unitary operator in quantum mechanics. Wigner's theorem tells us that

all ray space isometries are either unitary or anti-unitary. To discriminate between these two possibilities we use the Bargmann invariant [15, 16] which is an example of the Pancharatnam phase[7]. For Hamiltonians which are time reversal invariant, the geometric phase can only take values ± 1 and becomes a topological phase. This was the context for Longuet-Higgins discovery of conical intersections in molecular physics.

Let us now understand the geometric scalar potential in terms of the Fubini-Study metric. At every point \vec{R} , Eq. (6) determines a ray in the fast Hilbert space. In the case of the two state system we consider here the ray space is S^2 , the Bloch sphere.

$$f : \mathbb{R}^3 \mapsto S^2 \quad (24)$$

The map f can be used to pull back the Fubini-Study metric tensor to real space. This gives us a second rank tensor in real space. Contracting this tensor with the spatial metric gives us the geometric potential. Using the unit vector \hat{n} , where $\rho(\vec{R}) = \frac{1+\hat{n}(\vec{R})}{2}$, to describe a point on the Bloch sphere, f gives us $\hat{n}(\vec{R})$ and we have the pull back explicitly written as

$$\gamma_{ij} = \frac{1}{2} \text{Tr}[(\nabla_i \rho(\vec{R}))(\nabla_j \rho(\vec{R}))] = \frac{1}{4} (\nabla_i \hat{n}(\vec{R})) \cdot (\nabla_j \hat{n}(\vec{R})), \quad (25)$$

The potential V_{geom} is then given by $g^{ij}(\vec{R})\gamma_{ij}(\vec{R})$ the contraction of γ_{ij} with the metric tensor of space. Although $g_{ij} = \delta_{ij}$ is flat in this simple toy model, we include \vec{R} dependence to leave open the possibility of using generalised coordinates on the slow configuration space. The Fubini-Study metric is a Riemannian metric on the ray space and so is positive definite. However, the pull back γ_{ij} of this tensor is not a Riemannian metric on space since it may be degenerate. It is however, non-negative (positive semi definite). Similarly, the pullback of the symplectic form on the ray space, gives us a two-form on space, but this could be degenerate (or indeed vanish entirely).

5 Isolating the Scalar Geometric Potential

Putting together the work of the last sections, we find the Hamiltonian for the slow modes:

$$\mathcal{H}_{red} = V_{dyn}(\vec{R}) - \frac{\hbar^2}{2m}(\nabla_i - A_i(\vec{R}))(\nabla_i - A_i(\vec{R})) + V_{geom}(\vec{R}), \quad (26)$$

where $V_{dyn}(\vec{R}) = \lambda_+(\vec{R})$, A_i is the Berry potential and $V_{geom}(\vec{R})$ is the geometric potential.

We see that the slow modes are subject to three effects by the fast modes.

1. The first is a dynamic scalar potential $V_{dyn}(\vec{R})$ arising from the eigenvalue of the fast mode.
2. The second is modification of the kinetic energy term by the magnetic field $F = dA$ of the Berry potential. Let us be clear that this is not the same as the external $\vec{B}(\vec{R})$ we introduced; remember the neutron is neutral and insensitive to the external magnetic field $\vec{B}(\vec{R})$. Only the spin couples to $\vec{B}(\vec{R})$. What we are seeing here is *geometric magnetism*, which is distinct from the magnetic effects created by currents.
3. The third is the scalar geometric potential V_{geom} that is of interest in this paper. One could think of this as a “*geometric electricity*” in analogy with the usual scalar potential in electrostatics.

Let us look at some examples to isolate these separate effects. By choosing a magnetic field $\vec{B}(\vec{R}) = b(\vec{R})\hat{i}$ which varies only in magnitude, but not in direction, we eliminate effects 2 and 3 and isolate the dynamic scalar potential. To isolate the effect of the geometric vector potential, we consider the magnetic field $\vec{B}(\vec{R}) = B_0(\hat{i} \cos y + \hat{j} + \hat{k} \sin y)$ which has constant dynamic and geometric potentials. Here effects 1 and 3 are eliminated and only effect 2 remains.

To isolate the effect of the geometric scalar potential, we present an example in which the first two effects are absent and the third is present. This will convince the reader that the third effect is independent of the first two. It may also help an experimental detection

of the third effect if the first two effects are absent. We will now describe a magnetic field configuration $\vec{B}(\vec{R})$, (with $\vec{\nabla} \cdot \vec{B} = 0$) with the property that

- $|\vec{B}(\vec{R})|$ is constant and so $V_{dyn}(\vec{R})$ is independent of \vec{R} .
- the “magnetic field” $F = dA$ due to the Berry potential vanishes.

With $\vec{R} = (x, y, z)$, consider $\vec{B}(\vec{R}) = B_0(\cos a(y)\hat{i} + \sin a(y)\hat{k})$ where $a(y)$ is any function depending only on y . This makes the fast Hamiltonian $H_f = -\mu\vec{B}(\vec{R}) \cdot \vec{\sigma}$ real and proportional to $\cos a(y)\sigma_3 + \sin a(y)\sigma_1$, which is

$$\begin{bmatrix} \cos a(y) & \sin a(y) \\ \sin a(y) & -\cos a(y) \end{bmatrix} \quad (27)$$

The ground state of this Hamiltonian H_f is

$$|e_+(y)\rangle = \begin{bmatrix} \sin a(y)/2 \\ -\cos a(y)/2 \end{bmatrix} \quad (28)$$

It is easily checked that 1) $\lambda_+(\vec{R}) = -\mu B_0$ is actually independent of \vec{R} , so this term has no effect on the slow modes. 2) The Berry potential vanishes identically $A_i = 0$ and so there are no “geometric magnetic” effects and 3) the geometrical scalar potential V_{geom} does *not* vanish and is given by $\frac{\hbar^2}{8m}(\frac{da(y)}{dy})^2$. Note that V_{geom} is non-negative.

6 More Realistic Models

Let us carry these ideas over to the more realistic situation of molecular dynamics. The total Hamiltonian is

$$H_T = T_N + V_{NN} + T_e + V_{ee} + V_{Ne} \quad (29)$$

where the terms are, in order, T_N the nuclear kinetic energy, V_{NN} the nuclear-nuclear interaction, T_e the electronic kinetic energy, V_{ee} the electron-electron interaction and V_{Ne} the

nuclear-electronic interaction. The fast Hamiltonian consists of all those terms that depend on electronic coordinates,

$$H_f = T_e + V_{ee} + V_{Ne}(R), \quad (30)$$

where it is understood that the dependence of V_{Ne} on the nuclear coordinates R is parametric. We use the symbol R to represent a set of generalised coordinates in the nuclear configuration space. The slow Hamiltonian consists of the remaining terms

$$H_s = T_N + V_{NN} \quad (31)$$

Following the Born-Oppenheimer approach, we need to solve eq.(6) for the eigenstates $|e_n(R)\rangle$ and eigenvalues $\lambda_n(R)$. While this problem is considerably more difficult than our simple toy model, it is within the reach of physical chemists aided by modern computers. We will assume that there are no degeneracies $\lambda_n(R) \neq \lambda_m(R)$ for m, n distinct. The general ideas we used for the toy model carry over to these more realistic and difficult models. After performing a unitary transformation to the basis $\{|e_n(R)\rangle\}$, the slow mode Hamiltonian, including the back reaction of the fast modes on the slow ones is given by

$$\mathcal{H}_{red} = \tilde{T}_N + V_{NN} + V_{dyn} + V_{geom}, \quad (32)$$

where \tilde{T}_N is the nuclear kinetic energy modified by the Berry potential A , and V_{dyn} and V_{geom} are the two scalar potentials of dynamical and geometric origin. Assuming that the fast system is in the state $|e_n(R)\rangle$ with eigenvalue $\lambda_n(R)$, here are explicit expressions for these

$$V_{dyn}(R) = \lambda_n(R). \quad (33)$$

The effect of the geometric vector potential and the geometric scalar potential can be expressed in terms of the second rank tensor $\mathcal{V}_{ij} = \bar{\mathcal{V}}_{ji}$ on the nuclear configuration space

$$\mathcal{V}_{ij}(R) = \sum_{m \neq n} \frac{\langle e_n(R) | \partial_i H_f(R) | e_m(R) \rangle \langle e_m(R) | \partial_j H_f(R) | e_n(R) \rangle}{(\lambda_n(R) - \lambda_m(R))^2} \quad (34)$$

The effective magnetic field strength due to the Berry potential is given by the imaginary part of the tensor \mathcal{V} :

$$F_{ij}(R) = \frac{\mathcal{V}_{ij} - \mathcal{V}_{ji}}{2i}. \quad (35)$$

This agrees with eq.(10) of Ref. [1]. The pull-back of Fubini-Study metric on the fast ray space to the nuclear configuration space is given by the real part:

$$\gamma_{ij}(R) = \frac{\mathcal{V}_{ij} + \mathcal{V}_{ji}}{2}. \quad (36)$$

The geometric scalar potential is given by

$$V_{geom}(R) = g^{ij}(R)\gamma_{ij}(R), \quad (37)$$

where the metric g on the nuclear coordinate space has been contracted with γ . Note also that the metric on the nuclear configuration space is determined by the kinetic energy term, which weights the nuclei according to their masses. It may be advantageous to leave open the choice of coordinates on this space. For instance nuclear separations may be better coordinates than nuclear positions.

7 Conclusion

We have described a geometric effect that arises in the Born-Oppenheimer approximation in a toy model. This is a scalar potential which has its origin in the geometry of ray space, namely the Fubini-Study metric. We have given a magnetic field configuration in which this geometric potential can be isolated from the other effects. One difference between V_{geom} and the effects due to V_{dyn} and A_{geom} is that V_{geom} has the same effect on both spin states. But the other two act oppositely on the $|+\rangle$ and $|-\rangle$ states. The effect of V_{geom} on neutrons is to favour regions where the magnetic field direction is slowly varying and avoid regions of rapid variation, where the conditions of the adiabatic theorem would be violated. The advantage of a toy model is that the calculations can be done explicitly and we can learn from it.

While the more realistic models of molecular physics are not easy to solve, the same geometric ideas apply there. If one solves the fast eigenvalue problem on a computer, one can extract the slow dynamics in terms of eigenvalues and eigenstates in an entirely geometric manner. The metric on the nuclear configuration space is the one derived from the kinetic energy, which weights different nuclei according to their mass. The non-negativity of the geometric scalar potential remains true here as it follows from general arguments. The magnitude of the geometric scalar potential is smaller than the dynamical scalar potential by the ratio of the electronic mass to the nuclear mass. The effect of the geometric potential, while small, is measurable in the rotational and vibrational spectra of molecules.

In our toy model, we have treated the specific example of a neutron in a strong magnetic field. However, the mathematics we presented here has wider applications. One example is neutral cold atoms trapped using magnetic fields[17]. Such configurations are used in cold atom laboratories to achieve Bose-Einstein condensation as well as Fermi-degeneracy in dilute atomic gases. Magnetic traps only confine atoms in the weak field-seeking states. For weak magnetic field gradients and slow atoms, spin-flip transitions are suppressed by the adiabatic theorem. In practice, these transitions do occur in regions of very small magnetic fields which creates a region of trap loss in the magnetic trap due to spin flips to untrapped states and are referred to as Majorana losses. The magnetic field configuration of section V could be thought of as providing a geometric trap for atoms of all spins. This would have the effect of confining atoms to a two-dimensional plane. One could also construct geometric wave guides for atoms using other magnetic field configurations.

Another application, unrelated to quantum mechanics, is polarised waves. When polarised light propagates through a birefringent medium, the medium acts like a magnetic field. In general one could have linear, circular and even elliptical birefringence, in which two orthogonal polarisations traverse the medium at different speeds. Mathematically, this problem of polarised wave optics maps on all fours to the toy model of this paper. If one restricts to linear birefringence and arranges for a helical configuration of molecules, one can even reproduce the example of section V in which the waves are affected only by the geometric

scalar potential. Such helical configurations can be generated as supramolecular structures using liquid crystals. The ideas described here in the quantum context could be experimentally verified in the classical wave context of optics. The effect of geometric potential can be seen in an interference experiment. This experiment could also be done in a scaled up setup by replacing light by radio waves and replacing the molecular helical variation with centimetre size dielectric rods.

We hope to interest geometers as well as experimentalists in understanding the uses of this geometric scalar potential.

Acknowledgements: This paper is dedicated to Michael Berry on his 80th birthday. It was motivated by a talk given by Berry at ICTS, Bangalore, titled: The geometric phase and the separation of the world, during a program titled: Geometric Phases in Optics and Topological Matter. I thank Sanjukta Roy for clearing up my ideas on atom traps and Supurna Sinha for reading through the paper and suggesting improvements. This work was supported by a grant from the Simons Foundation (677895, R. G.).

References

- [1] Michael Victor Berry. Quantal phase factors accompanying adiabatic changes. *Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences*, 392(1802):45–57, 1984.
- [2] Raymond Y. Chiao and Yong-Shi Wu. Manifestations of berry’s topological phase for the photon. *Phys. Rev. Lett.*, 57:933–936, Aug 1986.
- [3] Rajendra Bhandari and Joseph Samuel. Observation of topological phase by use of a laser interferometer. *Phys. Rev. Lett.*, 60:1211–1213, Mar 1988.
- [4] D Chruscinski and A Jamiolkowski. *Geometric Phases in Classical and Quantum Mechanics*. Springer, 2004.
- [5] F Wilczek and A Shapere. *Geometric Phases in Physics*. WORLD SCIENTIFIC, 1989.

- [6] Y. Aharonov and J. Anandan. Phase change during a cyclic quantum evolution. *Phys. Rev. Lett.*, 58:1593–1596, Apr 1987.
- [7] S Pancharatnam. Generalized theory of interference, and its applications. part i. coherent pencils (1956) proc. In *Indian Acad. Sci. A*, volume 44, pages 247–262, 1956.
- [8] S. Ramaseshan and Rajaram Nityananda. The interference of polarised light as an early example of berry’s phase. *Current Science*, 55(24):1225–1226, 1986.
- [9] Hugh Christopher Longuet-Higgins. The intersection of potential energy surfaces in polyatomic molecules. *Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences*, 344(1637):147–156, 1975.
- [10] J H Hannay. Angle variable holonomy in adiabatic excursion of an integrable hamiltonian. *Journal of Physics A: Mathematical and General*, 18(2):221–230, feb 1985.
- [11] Gautam Gangopadhyay and Binayak Dutta-Roy. The born–oppenheimer approximation: A toy version. *American Journal of Physics*, 72(3):389–392, 2004.
- [12] C. Alden Mead and Donald G. Truhlar. On the determination of born–oppenheimer nuclear motion wave functions including complications due to conical intersections and identical nuclei. *The Journal of Chemical Physics*, 70(5):2284–2296, 1979.
- [13] Tohru Eguchi, Peter B. Gilkey, and Andrew J. Hanson. Gravitation, gauge theories and differential geometry. *Physics Reports*, 66(6):213–393, 1980.
- [14] Joseph Samuel and Rajendra Bhandari. General setting for berry’s phase. *Phys. Rev. Lett.*, 60:2339–2342, Jun 1988.
- [15] V. Bargmann. Note on wigner’s theorem on symmetry operations. *Journal of Mathematical Physics*, 5(7):862–868, 1964.
- [16] N. Mukunda and R. Simon. Quantum kinematic approach to the geometric phase. i. general formalism. *Annals of Physics*, 228(2):205–268, 1993.

- [17] David E. Pritchard. Cooling neutral atoms in a magnetic trap for precision spectroscopy.
Phys. Rev. Lett., 51:1336–1339, Oct 1983.