

# WATSON-CRICK CONJUGATES OF WORDS AND LANGUAGES

KALPANA MAHALINGAM, ANURAN MAITY

ABSTRACT. This paper is a theoretical study of notions in combinatorics of words motivated by information being encoded as DNA strands in DNA computing. We study Watson-Crick conjugates or  $\theta$ -conjugates, a generalization of the classical notions of conjugates of a word, inspired by biomolecular computing. The Watson-Crick mapping  $\theta$  is an involution that is also an antimorphism. We study some combinatorial properties of  $\theta$ -conjugates of a word. We characterize words that have the same set of  $\theta$ -conjugates. We investigate whether or not the family of certain languages is closed under  $\theta$ -conjugate operation. We also study the iterated  $\theta$ -conjugates. We then discuss the concept of  $\theta$ -conjugate-free language and some decidability problems for  $\theta$ -conjugate-freeness for different language classes.

## 1. INTRODUCTION

Encrypting information as DNA strands as in, e.g., DNA Computing, brings up for exploration new characteristics based on the specific biochemical properties of DNA molecules. Since each single-stranded DNA molecule is a sequence of adenine( $A$ ), guanine( $G$ ), cytosine( $C$ ) and thymine( $T$ ), it can be thought of as a word over the alphabet of bases  $\{A, G, T, C\}$ . Moreover, one of the major biochemical properties of DNA molecules is the Watson–Crick complementarity of the bases  $G$  and  $C$  and, respectively,  $A$  and  $T$ . Because of this property, two Watson–Crick complementary single DNA strands with opposite orientations stick together to form a DNA double strand in a process called base-pairing. Recently, there were several approaches from classical combinatorics on words to generalize this significant characteristic of DNA molecules (see [4, 9, 10]). Formally, the Watson-Crick complementarity property on strings over  $\Sigma$  is an involution mapping  $\theta$  with the additional property that  $\theta(uv) = \theta(v)\theta(u)$  for all strings  $u, v \in \Sigma^*$ .

The conjugate of a word is a very basic and important concept in combinatorics on words. The conjugacy relation defines a partition of words into equivalence classes which are also called circular words. The cyclic shift of a language  $L$ , which is the conjugacy operation on  $L$ , is defined as  $\text{shift}(L) = \{vu \mid uv \in L\}$ . Oshiba [18] proved that the regular and context-free languages are closed under the cyclic shift operation. The number of states in a deterministic finite automaton (DFA) needed to recognize the cyclic shift of an  $n$ -state DFA language has been investigated by Maslov [17]. The classical results on conjugacy are presented by Shyr [22]. The notion of  $\theta$ -conjugate of a word was defined by Kari and Mahalingam [9] to generalize the classical notions of conjugacy of words from the perspective of DNA computing. Kari and Mahalingam [9, 10] studied the properties of  $\theta$ -palindromes and they provided a complete characterization of the set of all words

---

*Key words and phrases.* Theoretical DNA computing, combinatorics on words, Watson-Crick conjugate, antimorphic involution.




FIGURE 1. Structure of conjugates and  $\theta$ -conjugates of a word

that are not Watson-Crick palindromes. They linked Watson-Crick palindromes to classical notions such as primitive words. They proved that the set of Watson-Crick palindromic words that can not be written as the product of two non-empty Watson-Crick palindromes equals the set of primitive Watson-Crick palindromes. Czeizler, Kari and Seki [4] studied the properties of  $\theta$ -primitive words. They proved the existence of a unique  $\theta$ -primitive root of a given word, and provided some constraints under which two distinct words share their  $\theta$ -primitive root. The combinatorial properties of strings in connection to partial words were investigated by Blanchet-Sadri and Luhmann [2].

In molecular biology, it is important to retrieve information from DNA sequences, align these sequences and analyse the exact orders of DNA sequences. Researchers have considered several chromosomal operations on DNA sequences including inversions, pseudo-inversions, translocations, swaps and pseudoknots in the past. Swap is an important operation for bio-sequences such as DNA or RNA sequences, and is closely related to mutations. This operation is extensively studied in [1, 11, 12, 16]. We consider a special kind of swap operation called  $\theta$ -conjugate where  $\theta$  is an antimorphic involution. In this paper, we investigate some combinatorial properties of the set of all  $\theta$ -conjugates of a word. The number of palindromes and Watson-Crick palindromes in the set of all  $\theta$ -conjugates of a word is studied by Mahalingam, Maity, and Pandoh [15]. We study the structure of  $\theta$ -conjugates of a word. We discuss the equality of two  $\theta$ -conjugates sets. Also, we examine whether or not certain family of languages is closed under  $\theta$ -conjugate operation. Then, we study iterated  $\theta$ -conjugates and  $\theta$ -conjugate-free languages.

The paper is organized as follows. In Section 3, we study some combinatorial properties of the set of all  $\theta$ -conjugates of a word. We first characterize the elements of the  $\theta$ -conjugates set. Then, we find the structure of words  $u$  and  $v$  such that  $C_\theta(u) = C_\theta(v)$ . In Section 4, we investigate the closure properties of  $\theta$ -conjugate operation. We show that the families of regular and context-sensitive languages are closed under  $\theta$ -conjugate operation and the family of context-free languages is not closed under  $\theta$ -conjugate operation. In Section 5, we define and discuss the iterated  $\theta$ -conjugates. In Section 6, we investigate some decidability problems for  $\theta$ -conjugate-free languages. We show that if  $L$  is a regular language with no  $\theta$ -palindrome, then whether  $L$  is  $\theta$ -conjugate-free or not is decidable. We also prove that it is undecidable to determine whether or not a given context-free language  $L$  is  $\theta$ -conjugate-free. We end the paper with some concluding remarks.

## 2. BASIC DEFINITIONS AND NOTATIONS

An alphabet ( $\Sigma$ ) is a finite non-empty set whose elements are called letters. A word or string over  $\Sigma$  is a finite sequence of elements of  $\Sigma$ . The zero letter sequence is called the empty word and is denoted by  $\lambda$ . The set of all finite words over  $\Sigma$  is denoted by  $\Sigma^*$ , and  $\Sigma^+ = \Sigma^* \setminus \{\lambda\}$ . A language  $L$  over  $\Sigma$  is a subset of  $\Sigma^*$ . The length of a word  $w$ , length of the sequence, is denoted by  $|w|$ .  $\Sigma^n$  and  $\Sigma^{\geq n}$  denotes the set of all words over  $\Sigma$  of length  $n$  and length greater than or equal to  $n$ , respectively. For  $a \in \Sigma$ ,  $|w|_a$  denotes the number of occurrences of  $a$  in  $w$ . The set of letters that occur in a word  $w$  and in a language  $L$  is denoted by  $\text{Alph}(w)$  and  $\text{Alph}(L) = \bigcup_{w \in L} \text{Alph}(w)$ , respectively. A word  $u$  is a factor or subword of a word  $w$  if  $w = xuy$  for  $x, y \in \Sigma^*$ . When  $x = \lambda$ ,  $u$  is called a prefix of  $w$ , and when  $y = \lambda$ ,  $u$  is called a suffix of  $w$ . For a word  $w = w_1w_2 \cdots w_n$  such that  $w_i \in \Sigma$ , the reversal of  $w$ , denoted by  $w^R$ , is the word  $w_n \cdots w_2w_1$ . A word  $w$  is a palindrome if  $w = w^R$ . Two words  $u$  and  $v$  are called conjugates of each other if there exists  $x, y \in \Sigma^*$  such that  $u = xy$  and  $v = yx$  (see Figure 1). The set of all conjugates of  $w \in \Sigma^*$  is denoted by  $C(w)$ . A word  $w \in \Sigma^*$  is called primitive if it is not a power of another word, i.e.,  $w$  is primitive if and only if  $w = x^n$  with  $n \geq 0$  and  $x \in \Sigma^*$  implies  $w = x$ . The set of all primitive words is denoted by  $Q$ . For every word  $w \in \Sigma^+$ , there exists a unique word  $\rho(w) \in \Sigma^+$ , called the *root* of  $w$ , such that  $\rho(w) \in Q$  and  $w = \rho(w)^n$  for some  $n \geq 1$ .

A function  $\psi : \Sigma^* \rightarrow \Sigma^*$  is said to be a *morphism* if  $\psi(uv) = \psi(u)\psi(v)$  for  $u, v \in \Sigma^*$ . A function  $\theta : \Sigma^* \rightarrow \Sigma^*$  is said to be an *antimorphism* if  $\theta(uv) = \theta(v)\theta(u)$  for  $u, v \in \Sigma^*$ . The functions  $\psi$  and  $\theta$  are called *involution* if  $\psi^2$  and  $\theta^2$  are identity mapping on  $\Sigma^*$ . A word  $u$  is a  $\theta$ -conjugate of another word  $w$  if  $uv = \theta(v)w$  for some  $v \in \Sigma^*$ . The set of all  $\theta$ -conjugates of  $w$  is denoted by  $C_\theta(w)$ . When  $\theta$  is an antimorphic involution, then  $C_\theta(w)$  is called the set of all Watson-Crick conjugates of  $w$ . A word  $w$  is called a  $\theta$ -palindrome if  $w = \theta(w)$ . Consider  $\Sigma = \{a, b\}$  and an antimorphic involution  $\theta$  such that  $\theta(a) = b$ . Then, the word  $aabb$  is a  $\theta$ -palindrome but not a palindrome. The set of all  $\theta$ -palindromes over  $\Sigma$  is denoted by  $P_\theta$ .

A grammar  $G = (V, T, P, S)$  is defined as a quadruple where  $V$  is a finite set of non-terminal symbols,  $T$  is a finite set of terminal symbols,  $P$  is a finite set of production rules and  $S \in V$  is the start symbol. The language generated by the grammar  $G$  is denoted by  $L(G)$ . If  $w \in L(G)$ , then  $w$  is generated by grammar  $G$ , i.e.,  $S \rightarrow^* w$ . A deterministic finite automata (DFA)  $M$  is defined by  $M = (S', \Sigma, \delta, q_0, F)$ , where  $S'$  is a finite set of states,  $\Sigma$  is an alphabet,  $\delta : S' \times \Sigma \rightarrow S'$  is a transition function,  $q_0 \in S'$  is the start state and  $F \subseteq S'$  is the set of final states. A language accepted by DFA  $M$ , denoted by  $L(M)$ , is called regular language. For all other concepts in formal language theory and combinatorics on words, the reader is referred to [13, 22].

We recall the following results from the literature which will be used in this paper. In 1962, Lyndon and Schützenberger [14] proved that for  $u, v \in \Sigma^*$ , if  $uv = vu$ , then  $u, v$  are powers of a common word. Also, they solved the equation  $uv = vw$  for  $u, v, w \in \Sigma^+$ . The results are following.

**Lemma 2.1.** [14] *Let  $u, v, w \in \Sigma^+$ .*

- If  $uv = vu$ , then  $u$  and  $v$  are powers of a common primitive word.
- If  $uv = vw$ , then for  $k \geq 0$ ,  $x \in \Sigma^+$  and  $y \in \Sigma^*$ ,  $u = xy$ ,  $v = (xy)^k x$ ,  $w = yx$ .

In 2008, Kari and Mahalingam [9] found the solution of the equation  $uv = \theta(v)u$  for an antimorphic involution  $\theta$ .

**Proposition 2.2.** [9] *If  $uv = \theta(v)u$  and  $\theta$  is an antimorphic involution, then  $u = x(yx)^i$ ,  $v = yx$  where  $i \geq 0$  and  $x, y \in P_\theta$  for  $x \in \Sigma^+$ ,  $y \in \Sigma^*$ .*

Throughout this paper, we take  $\theta$  as an antimorphic involution over  $\Sigma$ .

### 3. PROPERTIES OF $\theta$ -CONJUGATES

The conjugacy class of a word  $w$ , denoted by  $C(w)$ , is the set of all words  $yx$  such that  $w = xy$  where  $x, y \in \Sigma^*$ . The concept of conjugacy class of words have been extensively studied in [6, 7, 13]. Mahalingam, Maity, and Pandoh [15] studied the  $\theta$ -conjugates of a word and found the distribution of palindromes and  $\theta$ -palindromes in the set of all  $\theta$ -conjugates of a word. In this section, we study more combinatorial properties of the set of all  $\theta$ -conjugates of a word. We first study the structure of elements of the set  $C_\theta(w)$  where  $w \in \Sigma^*$ . Then, we characterize  $w$  and  $w'$  such that  $C_\theta(w) \cap C_\theta(w') \neq \emptyset$  where  $\emptyset$  denotes the empty set. We also find necessary and sufficient conditions for the equality of two  $\theta$ -conjugates sets.

First, we recall the following result from Kari and Mahalingam [9].

**Proposition 3.1.** [9] *Let  $u$  be a  $\theta$ -conjugate of  $w$ . Then, for an antimorphic involution  $\theta$ , there exists  $x, y \in \Sigma^*$  such that either  $u = xy$  and  $w = y\theta(x)$  or  $w = \theta(u)$ .*

Thus, for a word  $w$ ,  $C_\theta(w) = \{\theta(v)u : w = uv, u, v \in \Sigma^*\}$  (see Figure 1). We illustrate the concept of  $\theta$ -conjugates of a word with the help of some examples.

**Example 3.2.** Consider  $\Sigma = \{a, b, c, d\}$  and  $\theta$  be such that  $\theta(a) = b$ ,  $\theta(c) = d$ .

- (1) If  $w = aac$ , then  $C_\theta(w) = \{aac, daa, dba, dbb\}$ .
- (2) If  $w = abcd$ , then  $C_\theta(w) = \{abcd, cabc, cdab, cdaa\}$ .

Now, we prove that if  $u \in \Sigma^*$  and  $a \in \Sigma$ , then  $C_\theta(ua) = \{ua\} \cup \theta(a)C_\theta(u)$ .

**Lemma 3.3.** *For  $u \in \Sigma^*$  and  $a \in \Sigma$ ,  $C_\theta(ua) = \{ua\} \cup \theta(a)C_\theta(u)$ .*

*Proof.* We first show that  $C_\theta(ua) \subseteq \{ua\} \cup \theta(a)C_\theta(u)$ . Let  $x \in C_\theta(ua)$ . Then, for  $ua = u'v'$ ,  $x = \theta(v')u'$  where  $u', v' \in \Sigma^*$ . If  $v' = \lambda$ , then  $x = ua$ . If  $v' \in \Sigma^+$ , then  $v' = za$  for  $z \in \Sigma^*$  and hence,  $x = \theta(a)\theta(z)u'$ ,  $u = u'z$  and  $\theta(z)u' \in C_\theta(u)$ . Therefore,  $x \in \{ua\} \cup \theta(a)C_\theta(u)$  and  $C_\theta(ua) \subseteq \{ua\} \cup \theta(a)C_\theta(u)$ . Now, consider  $y \in \{ua\} \cup \theta(a)C_\theta(u)$ . If  $y = ua$ , then clearly  $y \in C_\theta(ua)$ . If  $y \in \theta(a)C_\theta(u)$ , then  $y = \theta(a)\theta(q)p$  for  $u = pq$ ,  $p, q \in \Sigma^*$ . This implies  $y \in C_\theta(pqa)$ , i.e.,  $y \in C_\theta(ua)$  and  $\{ua\} \cup \theta(a)C_\theta(u) \subseteq C_\theta(ua)$ . Therefore,  $C_\theta(ua) = \{ua\} \cup \theta(a)C_\theta(u)$ .  $\square$

**Corollary 3.4.** For  $v \in \Sigma^+$  and  $u \in \Sigma^*$ ,  $\theta(v)C_\theta(u) \subseteq C_\theta(uv)$ .

Also, from Proposition 2.2 and Lemma 3.3, we have the following.

**Corollary 3.5.** For  $u \in \Sigma^*$  and  $a \in \Sigma$ ,  $C_\theta(ua) = \theta(a)C_\theta(u)$  iff  $ua \in P_\theta$ .

We illustrate Lemma 3.3 and Corollary 3.5 with the help of some examples.

**Example 3.6.** Consider  $\Sigma = \{a, b, c\}$  and  $\theta$  be such that  $\theta(a) = b$ ,  $\theta(c) = c$ .

- (1) If  $w = bccb$ , then  $C_\theta(w) = \{bccb, abcc, acbc, accb, acca\}$ . Note that  $bccb$  is not a  $\theta$ -palindrome. Now,  $C_\theta(bcc) = \{bcc, cbc, ccb, cca\}$  and  $C_\theta(bccb) = \{bccb\} \cup \theta(b)C_\theta(bcc)$ .
- (2) If  $w = abcab$ , then  $C_\theta(w) = \{abcab, aabca, ababc, abcaa\}$ . Note that  $w$  is a  $\theta$ -palindrome. Now,  $C_\theta(abca) = \{abca, babc, bcab, bcaa\}$  and  $C_\theta(abcab) = \theta(b)C_\theta(abca)$ .

The conjugacy relation is an equivalence relation. Then, for  $w, w' \in \Sigma^+$ ,  $C(w) \cap C(w') \neq \emptyset$  iff  $C(w) = C(w')$ . Kari and Mahalingam [9] proved that  $\theta$ -conjugate relation is a reflexive relation but not symmetric and not transitive, i.e., not an equivalence relation. Then, for  $w = ab$ ,  $w' = aa$  and  $\theta(a) = b$ ,  $C_\theta(ab) \cap C_\theta(aa) = \{aa\}$  but  $C_\theta(aa) \neq C_\theta(ab)$ . We now characterize  $w, w'$  such that  $C_\theta(w) \cap C_\theta(w') \neq \emptyset$  and  $u, v$  such that  $C_\theta(u) = C_\theta(v)$ . We have the following results.

**Lemma 3.7.** For  $w, w' \in \Sigma^+$ ,  $C_\theta(w) \cap C_\theta(w') \neq \emptyset$  iff  $w = w'$  or  $w' \in C_\theta(w)$  or  $w' = v\gamma$ ,  $w = u\gamma$  where  $v \in C_\theta(u)$  and  $u, v, \gamma \in \Sigma^+$ .

*Proof.* Let  $w, w' \in \Sigma^+$  such that  $C_\theta(w) \cap C_\theta(w') \neq \emptyset$ . Consider  $z \in C_\theta(w) \cap C_\theta(w')$ . Then,

$$(1) \quad z = \theta(\beta)\alpha = \theta(\beta')\alpha'$$

where  $w = \alpha\beta$ ,  $w' = \alpha'\beta'$  and  $\alpha, \alpha', \beta, \beta' \in \Sigma^*$ . Without loss of generality, assume that  $|\alpha| \leq |\alpha'|$  and  $|\beta| \geq |\beta'|$ . Then, Equation (1) implies  $\alpha' = y\alpha$  and  $\beta = \theta(y)\beta'$  for  $y \in \Sigma^*$ . Thus,  $w = \alpha\beta = \alpha\theta(y)\beta'$  and  $w' = \alpha'\beta' = y\alpha\beta'$ . Now, if  $y = \lambda$ , then  $w = w' = \alpha\beta'$ . If  $\beta' = \lambda$ , then  $w' \in C_\theta(w)$ . Let  $z' = \alpha\theta(y)$ ,  $z'' = y\alpha$ ,  $y \in \Sigma^+$  and  $\beta' \in \Sigma^+$ . Then,  $w = z'\beta'$ ,  $w' = z''\beta'$  and  $z'' \in C_\theta(z')$ .

Conversely, for  $w, w' \in \Sigma^+$ , consider  $w = w'$  or  $w' \in C_\theta(w)$  or  $w' = v\gamma$ ,  $w = u\gamma$  where  $v \in C_\theta(u)$  and  $u, v, \gamma \in \Sigma^+$ . If  $w = w'$  or  $w' \in C_\theta(w)$ , then  $C_\theta(w) \cap C_\theta(w') \neq \emptyset$ . Now, consider  $w' = v\gamma$ ,  $w = u\gamma$  where  $v \in C_\theta(u)$  and  $u, v, \gamma \in \Sigma^+$ . Since,  $v \in C_\theta(u)$  and by Corollary 3.4,  $\theta(\gamma)C_\theta(u) \subseteq C_\theta(u\gamma)$ , we have,  $\theta(\gamma)v \in C_\theta(u\gamma)$ . Also,  $\theta(\gamma)v \in C_\theta(v\gamma)$ . Thus,  $\theta(\gamma)v \in C_\theta(w) \cap C_\theta(w')$ , i.e.,  $C_\theta(w) \cap C_\theta(w') \neq \emptyset$ . Hence, the proof.  $\square$

Mahalingam, Maity, and Pandoh [15] studied the equality of two  $\theta$ -conjugates sets and proved the following.

**Theorem 3.8.** [15] Let  $u, v \in \Sigma^*$  such that  $C_\theta(u) = C_\theta(v)$ .

- (1) If  $|C_\theta(u)| = 1$ , then  $v = u$ .

(2) If  $|C_\theta(u)| = 2$ , then  $v = u$  or  $v = \theta(u)$ .

(3) If  $|C_\theta(u)| \geq 3$ , then  $v = u$  or  $v = u^R$  such that  $u = a^{n_1}u'a^{n_2}$  and  $v = a^{n_2}u'a^{n_1}$  where  $a \in \Sigma$ ,  $n_1, n_2 \geq 1$ ,  $n_1 \neq n_2$ ,  $u' \in \Sigma^+$  such that  $a$  is not prefix and suffix of  $u'$ ,  $u' = u'^R$  and  $\theta(c) = c$  for all  $c \in \text{Alph}(u)$ .

With the help of few examples, we now show that the converse of the above theorem is not true.

**Example 3.9.** Let  $\Sigma = \{a, b, c\}$  and  $\theta$  be such that  $\theta(a) = b$  and  $\theta(c) = c$ .

(1) Consider  $u = ca$  and  $v = \theta(u) = bc$ . Then,  $|C_\theta(u)| = 2$  and  $cb \in C_\theta(v) \setminus C_\theta(u)$ .

(2) Consider  $u = abaa$  and  $v = u^R = aaba$ . Then,  $|C_\theta(u)| \geq 3$  and  $baab \in C_\theta(v) \setminus C_\theta(u)$ .

Hence, in both cases,  $C_\theta(u) \neq C_\theta(v)$ .

We refine Theorem 3.8 and find necessary and sufficient conditions for the equality of two  $\theta$ -conjugates sets. First, we study the structure of words  $w$  such that  $|C_\theta(w)| = 1$ ,  $|C_\theta(w)| = 2$  and  $|C_\theta(w)| \geq 3$ .

Mahalingam, Maity, and Pandoh [15] characterized the word  $w$  such that  $|C_\theta(w)| = 1$ . They proved the following result.

**Lemma 3.10.** [15] *Let  $z \in \Sigma^*$ . Then,  $|C_\theta(z)| = 1$  iff  $z = a^n$  such that  $\theta(a) = a$  for  $a \in \Sigma$ .*

We now characterize words  $u$  such that  $|C_\theta(u)| = 2$  and  $|C_\theta(u)| \geq 3$ .

**Lemma 3.11.** *For  $u \in \Sigma^*$  and  $a_1, a_2, c \in \Sigma$ ,  $|C_\theta(u)| = 2$  iff  $u$  is in one of the following forms:*

- (1)  $u = a_1$  such that  $\theta(a_1) \neq a_1$
- (2)  $u = a_1^m a_2$ ,  $m \geq 1$  such that  $\theta(a_1) = a_1$  and  $\theta(a_2) \neq a_1$
- (3)  $u = a_1 \theta(a_1)$  such that  $\theta(a_1) \neq a_1$
- (4)  $u = a_1 c a_1$  such that  $c \neq a_1$ ,  $\theta(c) = c$  and  $\theta(a_1) = a_1$ .

*Proof.* Let  $u = u_1 u_2 \cdots u_n \in \Sigma^n$  where  $u_i \in \Sigma$  for  $1 \leq i \leq n$ . Consider  $|C_\theta(u)| = 2$ . If  $n = 1$ , then  $u = u_1$  and as  $|C_\theta(u)| = 2$ ,  $\theta(u_1) \neq u_1$ . Now, consider  $n \geq 2$ . We have the following cases:

- $\theta(u_n) \neq u_1$  : Then, by Lemma 3.3, each element of  $C_\theta(u) \setminus \{u\}$  starts with  $\theta(u_n)$ . Since,  $\theta(u_n) \neq u_1$  and  $|C_\theta(u)| = 2$ , each element of  $C_\theta(u) \setminus \{u\}$  is equal to  $\theta(u)$ , i.e., for each  $x'', y'' \in \Sigma^+$  for which  $u = x'' y''$ , we have,  $\theta(y'') x'' = \theta(y'') \theta(x'')$ , i.e.,  $\theta(x'') = x''$ . This implies  $u = u_1^{n-1} u_n$  where  $\theta(u_1) = u_1$ .
- $\theta(u_n) = u_1$  : Then, for  $u_2 u_3 \cdots u_{n-1} = x$ ,  $u = u_1 x \theta(u_1)$ . We have the following cases:
  - $\theta(u_1) \neq u_1$  : Now,  $u_1 \theta(x) u_1, u_1 \theta(x) \theta(u_1) \in C_\theta(u)$ . If  $\theta(x) \neq x$ , then  $u, u_1 \theta(x) u_1$  and  $u_1 \theta(x) \theta(u_1)$  are three distinct elements of  $C_\theta(u)$ , which is a contradiction. Thus,  $\theta(x) = x$ . Let there exists  $a \in \text{Alph}(x)$  such that  $a \neq u_1$  and  $\theta(a) \neq a$ . Then, as  $\theta(x) = x$ , for  $x', z \in \Sigma^*$ ,  $u = u_1 x' a z \theta(a) \theta(x') \theta(u_1)$ . Now,  $u, u_1 x' u_1 x' a z \theta(a)$  and  $u_1 x' a \theta(z) \theta(a) \theta(x') u_1$  are three distinct elements of  $C_\theta(u)$ , which is a contradiction. So, there does not exist any  $a \in \text{Alph}(x)$  such that  $a \neq u_1$  and  $\theta(a) \neq a$ . This implies  $\theta(u_1) \notin \text{Alph}(x)$ . Let there exist  $b \in \text{Alph}(x)$  such

that  $b \neq u_1$ . Then,  $\theta(b) = b$ ,  $b \neq \theta(u_1)$  and  $u = u_1^{n_1} b z' \theta(u_1)$  for  $z' \in \Sigma^*$  and  $n_1 \geq 1$ . Now,  $u, u_1 \theta(z') u_1^{n_1} b$  and  $u_1 \theta(z') \theta(b) u_1^{n_1}$  are three distinct elements of  $C_\theta(u)$ , which is a contradiction. So, there does not exist any  $b \in \text{Alph}(x)$  such that  $b \neq u_1$ . Therefore,  $u = u_1^{n-1} \theta(u_1)$  and  $x = u_1^{n-2}$ . Since,  $\theta(x) = x$  and  $\theta(u_1) \neq u_1$ ,  $x = \lambda$ , i.e.,  $n = 2$ . Thus,  $u = u_1 \theta(u_1)$ .

–  $\theta(u_1) = u_1$  : Then,  $u = u_1 x u_1$ . Now, there exists a letter  $c \in \text{Alph}(x)$  such that  $c \neq u_1$ , otherwise,  $u = u_1^n$  with  $\theta(u_1) = u_1$ , which is a contradiction to the fact that  $|C_\theta(u)| = 2$ . Then, for  $v \in \Sigma^*$  and  $m_1 \geq 1$ ,  $u = u_1^{m_1+1} c v u_1$ . Let  $\alpha = u_1^{m_1+1} c v$  and  $\beta = u_1 \theta(v) u_1^{m_1} c$ . Since,  $\alpha, \beta \in C_\theta(u)$ ,  $\alpha \neq u$ ,  $\beta \neq u$  and  $|C_\theta(u)| = 2$ , we have,  $\alpha = \beta$ , i.e.,  $u_1^{m_1+1} c v = u_1 \theta(v) u_1^{m_1} c$ , i.e.,  $\theta(v) u_1^{m_1} c = u_1^{m_1} c v$ . If  $v \in \Sigma^+$ , then by Proposition 2.2,  $u_1^{m_1} c \in P_\theta$ , which is a contradiction as  $\theta(u_1) = u_1 \neq c$ . Thus,  $v = \lambda$  and  $u = u_1^{m_1+1} c u_1$ . Now, if  $\theta(c) \neq c$ , then  $u, u_1^{m_1+1} c$  and  $u_1 \theta(c) u_1^{m_1}$  are three distinct elements of  $C_\theta(u)$ , which is a contradiction. Thus,  $\theta(c) = c$  and  $u = u_1^{m_1+1} c u_1$ . Also, if  $m_1 \geq 2$ , then  $u, u_1^{m_1+1} c$  and  $u_1 c u_1^{m_1}$  are three distinct elements of  $C_\theta(u)$ , which is a contradiction. Therefore,  $m_1 = 1$ ,  $u = u_1 c u_1$  where  $c \neq u_1$ ,  $\theta(u_1) = u_1$  and  $\theta(c) = c$ .

The converse can be proved easily by direct computation. Hence, we omit the proof.  $\square$

Let  $D_1 = \{b^m \mid b \in \Sigma, \theta(b) = b, m \geq 0\}$ ,  $D_2 = \{b \mid b \in \Sigma, \theta(b) \neq b\}$ ,  $D_3 = \{a^m b \mid a, b \in \Sigma, \theta(a) = a, \theta(b) \neq a, m \geq 1\}$ ,  $D_4 = \{a \theta(a) \mid a \in \Sigma, \theta(a) \neq a\}$  and  $D_5 = \{a b a \mid a, b \in \Sigma, \theta(a) = a, \theta(b) = b, a \neq b\}$ . Consider  $T_e = D_1 \cup D_2 \cup D_3 \cup D_4 \cup D_5$  and  $T_o = D_2 \cup D_3 \cup D_4 \cup D_5$ . We will use  $T_e$  and  $T_o$  in the following.

From Lemmas 3.10 and 3.11, we have the structure of words  $u$  such that  $|C_\theta(u)| \geq 3$ .

**Lemma 3.12.** *For  $u \in \Sigma^*$ ,  $|C_\theta(u)| \geq 3$  iff  $u \in \Sigma^* \setminus T_e$ .*

We now characterize words  $u$  and  $v$  such that  $C_\theta(u) = C_\theta(v)$ . Using Lemma 3.10, we first discuss the case  $|C_\theta(u)| = 1$  for  $u \in \Sigma^*$  whose proof is straight forward.

**Proposition 3.13.** *Consider  $u, v \in \Sigma^*$  and  $|C_\theta(u)| = 1$ . Then,  $C_\theta(u) = C_\theta(v)$  iff  $u = v = a^n$  for  $a \in \Sigma$  with  $\theta(a) = a$  and  $n \geq 0$ .*

Now, we discuss the case  $|C_\theta(u)| = 2$  for  $u \in \Sigma^*$ .

**Proposition 3.14.** *Consider  $a, b, c \in \Sigma$ ,  $u, v \in \Sigma^k$ ,  $k \geq 1$  and  $|C_\theta(u)| = 2$ . Then,  $C_\theta(u) = C_\theta(v)$  iff  $u$  and  $v$  satisfy one of the following:*

- (1)  $u = ab$  and  $v = ba$  for  $\theta(a) = a$ ,  $\theta(b) = b$ ,  $a \neq b$
- (2)  $u = c$  and  $v = \theta(c)$  where  $\theta(c) \neq c$
- (3)  $u = v$  where  $u \in T_o$ .

*Proof.* For  $u, v \in \Sigma^k$ , consider  $C_\theta(u) = C_\theta(v)$  and  $|C_\theta(u)| = 2$ . Then, by Theorem 3.8, either  $v = u$  or  $v = \theta(u)$ . If  $u = v$ , then by Lemma 3.11,  $u \in T_o$ . Now, consider  $v \neq u$ . Then,  $v = \theta(u)$  where  $\theta(u) \neq u$ . If  $|u| = k = 1$ , then as  $|C_\theta(u)| = 2$ ,  $u = c$  and  $v = \theta(c)$  for  $c \in \Sigma$  and  $\theta(c) \neq c$ . Now, consider  $|u| = k \geq 2$ . Since,  $|C_\theta(u)| = 2$  and  $u \neq v$ , for each  $\alpha, \beta \in \Sigma^+$  for which  $u = \alpha\beta$ ,  $\theta(\beta)\alpha = z \in C_\theta(u)$  is either equal

to  $u = \alpha\beta$  or equal to  $\theta(u) = \theta(\beta)\theta(\alpha)$ . If  $z = u$ , then by Proposition 2.2,  $u$  is a  $\theta$ -palindrome, which is a contradiction to the fact that  $\theta(u) \neq u$ . Thus,  $z = \theta(u)$ , i.e.,  $\theta(\beta)\alpha = \theta(\beta)\theta(\alpha)$ , i.e.,  $\theta(\alpha) = \alpha$ . Now,  $v = \theta(u) = \theta(\beta)\theta(\alpha)$  implies  $\alpha\theta(\beta) = z' \in C_\theta(v)$ . Since,  $C_\theta(u) = C_\theta(v)$  and  $|C_\theta(u)| = 2$ ,  $z'$  is either equal to  $u$  or  $\theta(u)$ . If  $z' = \theta(u)$ , i.e.,  $\alpha\theta(\beta) = \theta(\beta)\theta(\alpha)$ , then as  $\alpha \in P_\theta$ ,  $\alpha\theta(\beta) = \theta(\beta)\alpha$ . Then, by Lemma 2.1,  $\theta(\beta) = p^i$  and  $\alpha = p^j$  for  $p \in Q$  and  $i, j \geq 1$ . Since,  $\alpha \in P_\theta$ , we have,  $p \in P_\theta$ . This implies  $\beta \in P_\theta$  and  $u \in P_\theta$ , which is a contradiction to the fact that  $u \notin P_\theta$ . Thus,  $z' = u$ , i.e.,  $\alpha\theta(\beta) = \alpha\beta$ , i.e.,  $\theta(\beta) = \beta$ . Therefore, for each  $\alpha, \beta \in \Sigma^+$  for which  $u = \alpha\beta$ , we have,  $\alpha, \beta \in P_\theta$ . This implies if  $|u| = k = 2$ , then  $u = ab$  and  $v = ba$  where  $\theta(a) = a$ ,  $\theta(b) = b$  and  $a \neq b$ . If  $|u| = k \geq 3$ , then  $u = v = d_1^k$  for  $d_1 \in \Sigma$  and  $\theta(d_1) = d_1$ , which is a contradiction to the fact that  $u \neq v$ . Therefore, if  $|C_\theta(u)| = 2$  and  $C_\theta(u) = C_\theta(v)$ , then  $u = v$  where  $u \in T_o$  or  $u = ab$  and  $v = ba$  for  $\theta(a) = a$ ,  $\theta(b) = b$ ,  $a \neq b$  or  $u = c$  and  $v = \theta(c)$  where  $\theta(c) \neq c$ . The converse is straight forward.  $\square$

We now discuss the case  $|C_\theta(u)| \geq 3$  for  $u \in \Sigma^*$ .

**Proposition 3.15.** *Consider  $u, v \in \Sigma^*$ ,  $a_1, b_1 \in \Sigma$  and  $|C_\theta(u)| \geq 3$ . Then,  $C_\theta(u) = C_\theta(v)$  iff  $u$  and  $v$  satisfy one of the following:*

- (1)  $u = a_1^{m+1}b_1a_1^m$  and  $v = a_1^mb_1a_1^{m+1}$  where  $m \geq 1$ ,  $\theta(a_1) = a_1$ ,  $\theta(b_1) = b_1$  and  $a_1 \neq b_1$
- (2)  $u = v$  where  $u \in \Sigma^* \setminus T_e$ .

*Proof.* For  $u, v \in \Sigma^*$ , consider  $|C_\theta(u)| \geq 3$  and  $C_\theta(u) = C_\theta(v)$ . Then, by Theorem 3.8, either  $u = v$  or  $u = a_1^{n_1}u'a_1^{n_2}$  and  $v = a_1^{n_2}u'a_1^{n_1}$  where  $a_1 \in \Sigma$ ,  $n_1, n_2 \geq 1$ ,  $n_1 \neq n_2$ ,  $u' \in \Sigma^+$  such that  $a_1$  is not prefix and suffix of  $u'$ ,  $u' = u'^R$  and  $\theta(c_1) = c_1$  for all  $c_1 \in \text{Alph}(u)$ . If  $u = v$ , then by Lemma 3.12,  $u \in \Sigma^* \setminus T_e$ . Now, consider  $u \neq v$ . Without loss of generality, let us take  $n_1 > n_2$ . Now, if  $|u'| \geq 2$ , then as  $u' = u'^R$ ,  $\theta(c_1) = c_1$  for all  $c_1 \in \text{Alph}(u)$  and  $a_1$  is not prefix and suffix of  $u'$ , we have,  $u' = d_1\alpha d_1$  for  $d_1 \in \Sigma$ ,  $d_1 \neq a_1$ ,  $\theta(d_1) = d_1$  and  $\alpha \in \Sigma^*$ . This implies  $u = a_1^{n_1}d_1\alpha d_1a_1^{n_2}$  and  $v = a_1^{n_2}d_1\alpha d_1a_1^{n_1}$  where  $n_1 > n_2$ . Then,  $a_1^{n_1}d_1\theta(\alpha)a_1^{n_2}d_1 \in C_\theta(v) \setminus C_\theta(u)$ , which is a contradiction to our assumption that  $C_\theta(u) = C_\theta(v)$ . Thus, as  $u' \in \Sigma^+$ ,  $|u'| = 1$ . Then,  $u = a_1^{n_1}b_1a_1^{n_2}$  and  $v = a_1^{n_2}b_1a_1^{n_1}$  where  $u' = b_1 \in \Sigma$ ,  $a_1 \neq b_1$  and  $\theta(b_1) = b_1$ . If  $n_1 - n_2 \geq 2$ , i.e.,  $n_1 - n_2 = 2 + k$  for  $k \geq 0$ , then  $u = a_1^{n_2+k+2}b_1a_1^{n_2}$  and  $v = a_1^{n_2}b_1a_1^{n_2+k+2}$ . Now,  $a_1^{k+n_2+1}b_1a_1^{n_2+1} \in C_\theta(v) \setminus C_\theta(u)$ , which is a contradiction to our assumption that  $C_\theta(u) = C_\theta(v)$ . Thus, as  $n_1 > n_2$ ,  $n_1 - n_2 = 1$  and  $u = a_1^{n_2+1}b_1a_1^{n_2}$  and  $v = a_1^{n_2}b_1a_1^{n_2+1}$ .

Conversely, if  $u = v$  where  $u \in \Sigma^* \setminus T_e$ , then obviously  $C_\theta(u) = C_\theta(v)$ . Now, consider  $u = a_1^{m+1}b_1a_1^m$  and  $v = a_1^mb_1a_1^{m+1}$  where  $m \geq 1$ ,  $\theta(a_1) = a_1$ ,  $\theta(b_1) = b_1$  and  $a_1 \neq b_1$ . Then, by direct computation,  $C_\theta(u) = C_\theta(v)$ .  $\square$

Combining Propositions 3.13, 3.14 and 3.15, we have the following theorem.

**Theorem 3.16.** *For  $u, v \in \Sigma^*$ ,  $C_\theta(u) = C_\theta(v)$  iff either  $v = u$  or  $v = \theta(u)$  such that one of the following holds:*

- (1)  $u = a_1b_1$  for  $a_1, b_1 \in \Sigma$  with  $\theta(a_1) = a_1$ ,  $\theta(b_1) = b_1$  and  $a_1 \neq b_1$ .



(2)  $u = c_1$  for  $c_1 \in \Sigma$  with  $\theta(c_1) \neq c_1$ .

(3)  $u = a_1^{m+1}b_1a_1^m$  for  $a_1, b_1 \in \Sigma$  with  $\theta(a_1) = a_1$ ,  $\theta(b_1) = b_1$ ,  $a_1 \neq b_1$  and  $m \geq 1$ .

#### 4. CLOSURE PROPERTIES OF $\theta$ -CONJUGATE

The conjugates of a language  $L$  is defined as  $C(L) = \bigcup_{w \in L} C(w)$ . Jiraskova and Okhotin [8] showed that the family of regular languages is closed under conjugate operation. Oshiba [18] proved that if  $L$  is context-free, then  $C(L)$  is context-free. Brandstadt [3] showed that the family of context-sensitive languages is closed under conjugate operation. Now, the definition of  $\theta$ -conjugates of a word can be naturally extended to a language. In this section, we first define the  $\theta$ -conjugates of a language. Then, we examine whether or not a family of languages is closed under  $\theta$ -conjugate operation.

**Definition 4.1.** For  $L \subseteq \Sigma^*$ , the  $\theta$ -conjugates of  $L$ , denoted by  $C_\theta(L)$ , is defined as

$$C_\theta(L) = \bigcup_{w \in L} C_\theta(w).$$

We now investigate the closure properties of certain well-known families of languages with respect to  $\theta$ -conjugate operation. For this we need the following result.

**Lemma 4.2.** *If for a morphic involution  $\psi$  and an antimorphic involution  $\theta$  over  $\Sigma$ ,  $\psi(a) = \theta(a)$  for all  $a \in \Sigma$ , then  $\psi(u) = \theta(u^R)$  for  $u \in \Sigma^*$ .*

*Proof.* Consider  $u = u_1u_2 \cdots u_n$  where  $u_i \in \Sigma$ . Then, as  $\psi(a) = \theta(a)$  for all  $a \in \Sigma$ ,  $\psi(u_i) = \theta(u_i)$  for all  $i$  ( $1 \leq i \leq n$ ). This implies  $\psi(u) = \psi(u_1)\psi(u_2) \cdots \psi(u_n) = \theta(u_1)\theta(u_2) \cdots \theta(u_n) = \theta(u_n u_{n-1} \cdots u_1) = \theta(u^R)$ .  $\square$

First we show that the regular languages are closed under  $\theta$ -conjugate operation.

**Theorem 4.3.** *If  $L \subseteq \Sigma^*$  is regular, then  $C_\theta(L)$  is also regular.*

*Proof.* Since  $L$  is regular there exists a deterministic finite automaton (DFA)  $A$  such that  $L(A) = L$ . Consider  $A = (Q', \Sigma, \delta, q_0, F)$  where  $Q' = \{q_0, q_1, \dots, q_n\}$  is the finite set of states,  $\Sigma$  is the input alphabet,  $\delta : Q' \times \Sigma \rightarrow Q'$  is the transition function,  $q_0$  is the initial state and  $F \subseteq Q'$  is the set of all final states.

For all  $i$  ( $0 \leq i \leq n$ ), let  $B_i = (Q', \Sigma, \delta, q_i, F)$  be an automaton with the same states, same transition functions and the same final states as  $A$ , and with the initial state  $q_i$ . Then, as  $A$  is deterministic, each  $B_i$  is deterministic for  $0 \leq i \leq n$ . Now, for each  $i$ ,  $L(B_i)$  is the regular language accepted by  $B_i$ . Since reverse of a regular language is regular [13], for each  $i$ ,  $(L(B_i))^R$  is regular. Consider a morphic involution  $\psi : \Sigma^* \rightarrow \Sigma^*$  such that  $\psi(a) = \theta(a)$  for all  $a \in \Sigma$ . Since regular languages are closed under morphism [13],  $\psi((L(B_i))^R)$  is also regular. Now, we show that  $\psi((L(B_i))^R) = \theta(L(B_i))$ . Let  $z \in \psi((L(B_i))^R)$ . Then, for some  $\alpha \in (L(B_i))^R$ ,  $z = \psi(\alpha)$ , i.e., by Lemma 4.2,  $z = \theta(\alpha^R)$ . This implies as  $\alpha^R \in L(B_i)$ ,  $z \in \theta(L(B_i))$ . Then,  $\psi((L(B_i))^R) \subseteq \theta(L(B_i))$ . Now, consider  $z' \in \theta(L(B_i))$ . Then, for some  $\beta \in L(B_i)$ ,  $z' = \theta(\beta)$ , i.e., by Lemma 4.2,  $z' = \psi(\beta^R)$ . Since,  $\beta \in L(B_i)$ ,  $z' \in \psi((L(B_i))^R)$ , i.e.,  $\theta(L(B_i)) \subseteq \psi((L(B_i))^R)$ . Therefore,

$\psi((L(B_i))^R) = \theta(L(B_i))$ . Now, as  $\theta(L(B_i)) (0 \leq i \leq n)$  is regular, for each  $i (0 \leq i \leq n)$  there exists a DFA  $B'_i$  such that  $L(B'_i) = \theta(L(B_i))$ .

Let for all  $i (0 \leq i \leq n)$ ,  $C_i = (Q', \Sigma, \delta, q_0, \{q_i\})$  be an automaton with the same states, the same transitions, and the same initial states as  $A$ , and with the only final state  $q_i$ . Then, as  $A$  is deterministic, each  $C_i$  is deterministic. By definition, a string  $w$  is in  $C_\theta(L)$  iff  $w = \theta(v)u$  for  $uv \in L$ . Consider the middle state in the accepting computation of  $A$  on  $uv$ , reached after consuming  $u$ , and let us reformulate the condition as follows: there exists a state  $q_i \in Q'$  such that the computation of  $A$  on  $u$  ends in the state  $q_i$ , while  $v$  is accepted by  $A$  starting from  $q_i$ . Then,  $\theta(v)$  is accepted by  $B'_i$  and  $u$  is accepted by  $C_i$  and

$$C_\theta(L) = \bigcup_{i=0}^n L(B'_i)L(C_i).$$

Since,  $L(B'_i)$  and  $L(C_i)$  are regular languages and concatenation of two regular languages is regular [13],  $L(B'_i)L(C_i)$  is regular. As finite union of regular languages is regular [13],  $C_\theta(L)$  is regular. □

Now, we consider the closure property for the context-free languages with respect to  $\theta$ -conjugate operation.

**Theorem 4.4.** *Context-free languages are not closed under  $\theta$ -conjugate operation.*

*Proof.* Consider the following context-free language  $L = \{a^n c^m d^m a^n \mid n, m \geq 1\}$  over  $\Sigma = \{a, b, c, d, e, f\}$ . Let  $\theta$  be such that  $\theta(a) = b$ ,  $\theta(c) = e$  and  $\theta(d) = f$ . Then,  $C_\theta(L) = \{b^i a^n c^m d^m a^{n-i} \mid n, m \geq 1, 0 \leq i \leq n\} \cup \{b^n f^j a^n c^m d^{m-j} \mid n, m \geq 1, 0 < j \leq m\} \cup \{b^n f^m e^k a^n c^{m-k} \mid n, m \geq 1, 0 < k \leq m\} \cup \{b^n f^m e^m b^l a^{n-l} \mid n, m \geq 1, 0 < l \leq n\}$ . Consider the regular language  $L' = \{b^{n_1} f^{m_1} e a^{m_2} c^{n_2} \mid n_1, m_1, m_2, n_2 \geq 2\}$ . Now, if  $C_\theta(L)$  is context-free, then as the intersection of a context-free language and a regular language is context-free [21],  $C_\theta(L) \cap L' = \{b^n f^m e a^n c^{m-1} \mid n \geq 2, m \geq 3\}$  is context-free. But by pumping lemma for context-free languages,  $\{b^n f^m e a^n c^{m-1} \mid n \geq 2, m \geq 3\}$  is not context-free which is a contradiction. Thus,  $C_\theta(L)$  is not context-free. □

We, furthermore, consider the closure property for context-sensitive languages and show that context-sensitive languages are closed under the  $\theta$ -conjugate operation. We first show that if  $L$  is context-sensitive, then  $bbC_\theta(L)$  for  $b \in \Sigma$  is context-sensitive.

**Theorem 4.5.** *If  $L \subseteq \Sigma^+$  is context-sensitive, then  $bbC_\theta(L)$  for  $b \in \Sigma$  is context-sensitive.*

*Proof.* Consider  $L$  is a context-sensitive language. Since the context-sensitive languages are closed under reversal [20], then  $L^R$  is also context-sensitive. Let  $\psi : \Sigma^* \rightarrow \Sigma^*$  be a morphic involution such that  $\psi(a) = \theta(a)$  for all  $a \in \Sigma$ . Consider

$$L' = \bigcup_{uv \in L^R} \{\psi(u)v^R\}.$$

Then, by Lemma 4.2,

$$L' = \bigcup_{uv \in L^R} \{\psi(u)v^R\} = \bigcup_{v^R u^R \in L} \{\theta(u^R)v^R\} = C_\theta(L).$$

We prove that  $bbL'$  where  $b \in \Sigma$  is context-sensitive language. Let  $G = (V, \Sigma, P, S)$  be a context-sensitive grammar for  $L^R$ . We now construct a context-sensitive grammar  $G_N$  such that  $L(G_N) = bbL'$ . Let  $V_1 = \{S', T_1, T_2, T_3, T_4, T\}$  and  $V_2 = \{X_a^o, X_a^I, X_a^D, X_a^R, X_a^T, X_a^F, \theta(X_a^o), \theta(X_a^D) \mid a \in \Sigma\}$ . Consider  $G_N = (V_N, \Sigma, P_N, S')$  where  $V_N = V \cup V_1 \cup V_2$ ,  $V \cap V_1 = \emptyset$ ,  $V \cap V_2 = \emptyset$ ,  $V_1 \cap V_2 = \emptyset$ ,  $S'$  is the start symbol and  $P_N$  is the union of the following sets :

$$\begin{aligned}
(1) \quad P_N(1) &= \{S' \rightarrow T_1 S T_2 \mid T_1 S T\} & (13) \quad P_N(13) &= \{Z X_{a_1}^I X_{a_2}^o \rightarrow X_{a_1}^I Z X_{a_2}^R \mid a_1, a_2 \in \Sigma\} \\
(2) \quad P_N(2) &= \{h(\alpha) \rightarrow h(\beta) \mid \alpha \rightarrow \beta \in P\} & (14) \quad P_N(14) &= \{Z X_{a_1}^R X_{a_2}^o \rightarrow X_{a_1}^R Z X_{a_2}^R \mid a_1, a_2 \in \Sigma\} \\
(3) \quad P_N(3) &= \{T_1 X_a^o \rightarrow \theta(X_a^D) T_1 \mid a \in \Sigma\} & (15) \quad P_N(15) &= \{Z X_{a_1}^R X_{a_2}^F \rightarrow X_{a_2}^T Z X_{a_1}^F \mid a_1, a_2 \in \Sigma\} \\
(4) \quad P_N(4) &= \{X_a^o T_2 \rightarrow T_3 X_a^F \mid a \in \Sigma\} & (16) \quad P_N(16) &= \{X_{a_1}^R X_{a_2}^T Z \rightarrow X_{a_2}^T Z X_{a_1}^o \mid a_1, a_2 \in \Sigma\} \\
(5) \quad P_N(5) &= \{X_a^o T_3 \rightarrow T_3 X_a^o \mid a \in \Sigma\} & (17) \quad P_N(17) &= \{X_{a_1}^I X_{a_2}^T Z \rightarrow X_{a_2}^D Z X_{a_1}^I \mid a_1, a_2 \in \Sigma\} \\
(6) \quad P_N(6) &= \{X_a^o T \rightarrow T_4 X_a^o \mid a \in \Sigma\} & (18) \quad P_N(18) &= \{Z X_{a_1}^I X_{a_2}^F \rightarrow X_{a_2}^D X_{a_1}^D Z \mid a_1, a_2 \in \Sigma\} \\
(7) \quad P_N(7) &= \{\theta(X_a^D) Z \rightarrow Z \theta(a) \mid a \in \Sigma\} \\
(8) \quad P_N(8) &= \{\theta(X_a^D) Z' \rightarrow Z' \theta(a) \mid a \in \Sigma\} \\
(9) \quad P_N(9) &= \{Z \rightarrow bb, Z' \rightarrow bb, Z'' \rightarrow bb\} \\
(10) \quad P_N(10) &= \{Z'' X_a^o \rightarrow X_a^D Z \mid a \in \Sigma\} \\
(11) \quad P_N(11) &= \{X_a^D Z \rightarrow Z a \mid a \in \Sigma\} \\
(12) \quad P_N(12) &= \{Z X_a^o \rightarrow Z X_a^I \mid a \in \Sigma\}
\end{aligned}$$

where  $Z = T_1 T_3$ ,  $Z' = T_1 T_2$ ,  $Z'' = T_1 T_4$  and  $h : (V \cup \Sigma)^* \rightarrow V_N$  is the morphism defined by  $h(B) = B$  for  $B \in V$  and  $h(a) = X_a^o$  for  $a \in \Sigma$ . We prove the following two claims:

**Claim 1 :**  $bbL' \subseteq L(G_N)$  : Let  $x \in bbL'$ . Then,  $x = bb\psi(y_1 y_2 \cdots y_i)(y_n y_{n-1} \cdots y_{i+1})$  where  $y_1 y_2 \cdots y_n \in L^R$  and  $y_i \in \Sigma$ . This implies as  $\psi(a) = \theta(a)$  for all  $a \in \Sigma$ ,  $x = bb\theta(y_1)\theta(y_2) \cdots \theta(y_i)(y_n y_{n-1} \cdots y_{i+1})$ . Now, we show that  $x \in L(G_N)$ . Since,  $y_1 y_2 \cdots y_n = y \in L^R$ ,  $S$  derives  $y$ . Then, by  $P_N(1)$ ,  $P_N(2)$  and definition of  $h$ ,  $S'$  derives  $T_1 X_{y_1}^o X_{y_2}^o \cdots X_{y_n}^o T_2$  or  $T_1 X_{y_1}^o X_{y_2}^o \cdots X_{y_n}^o T$ .

(we use boxes to denote the variables used in each derivation step):

Now,

$$\begin{aligned}
S' &\rightarrow^* T_1 X_{y_1}^o X_{y_2}^o \cdots \boxed{X_{y_n}^o T} \\
&\rightarrow \boxed{T_1 X_{y_1}^o} X_{y_2}^o \cdots T_4 X_{y_n}^o && \text{( by } P_N(6) \text{ )} \\
&\rightarrow^* \theta(X_{y_1}^D) \theta(X_{y_2}^D) \cdots \theta(X_{y_{n-1}}^D) T_1 T_4 X_{y_n}^o && \text{( by } P_N(3) \text{ )} \\
&= \theta(X_{y_1}^D) \theta(X_{y_2}^D) \cdots \theta(X_{y_{n-1}}^D) \boxed{Z'' X_{y_n}^o} \\
&\rightarrow \theta(X_{y_1}^D) \theta(X_{y_2}^D) \cdots \theta(X_{y_{n-1}}^D) \boxed{X_{y_n}^D Z} && \text{( by } P_N(10) \text{ )} \\
&\rightarrow^* bb\theta(y_1)\theta(y_2) \cdots \theta(y_{n-1})y_n && \text{( by } P_N(11), P_N(7) \text{ and } P_N(9) \text{ )}
\end{aligned}$$

So, for  $i = n - 1$ ,  $x \in L(G_N)$ . Now,

$$\begin{aligned}
S' &\rightarrow^* \boxed{T_1 X_{y_1}^o} X_{y_2}^o \cdots X_{y_n}^o T_2 \\
&\rightarrow^* \theta(X_{y_1}^D) \cdots \theta(X_{y_n}^D) T_1 T_2 && \text{( by } P_N(3) \text{ )} \\
&= \theta(X_{y_1}^D) \cdots \boxed{\theta(X_{y_n}^D) Z'} \\
&\rightarrow^* bb\theta(y_1) \cdots \theta(y_n) && \text{( by } P_N(8) \text{ and } P_N(9) \text{ )}
\end{aligned}$$

So, for  $i = n$ ,  $x \in L(G_N)$ . Now,

$$\begin{aligned}
S' &\rightarrow^* \boxed{T_1 X_{y_1}^o} X_{y_2}^o \cdots X_{y_n}^o T_2 \\
&\rightarrow \theta(X_{y_1}^D) \boxed{T_1 X_{y_2}^o} \cdots X_{y_n}^o T_2 && \text{( by } P_N(3) \text{ )} \\
&\rightarrow^* \theta(X_{y_1}^D) \cdots \theta(X_{y_i}^D) T_1 X_{y_{i+1}}^o \cdots \boxed{X_{y_n}^o T_2} && \text{( by } P_N(3) \text{ )} \\
&\rightarrow \theta(X_{y_1}^D) \cdots \theta(X_{y_i}^D) T_1 X_{y_{i+1}}^o \cdots \boxed{X_{y_{n-1}}^o T_3} X_{y_n}^F && \text{( by } P_N(4) \text{ )} \\
&\rightarrow^* \theta(X_{y_1}^D) \cdots \theta(X_{y_i}^D) T_1 T_3 X_{y_{i+1}}^o \cdots X_{y_n}^F && \text{( by } P_N(5) \text{ )} \\
&= \theta(X_{y_1}^D) \cdots \theta(X_{y_i}^D) \boxed{Z X_{y_{i+1}}^o} X_{y_{i+2}}^o \cdots X_{y_n}^F \\
&\rightarrow \theta(X_{y_1}^D) \cdots \theta(X_{y_i}^D) \boxed{Z X_{y_{i+1}}^I X_{y_{i+2}}^o} \cdots X_{y_n}^F && \text{( by } P_N(12) \text{ )} \\
&\rightarrow \theta(X_{y_1}^D) \cdots \theta(X_{y_i}^D) X_{y_{i+1}}^I \boxed{Z X_{y_{i+2}}^R X_{y_{i+3}}^o} \cdots X_{y_n}^F && \text{( by } P_N(13) \text{ )} \\
&\rightarrow \theta(X_{y_1}^D) \cdots \theta(X_{y_i}^D) X_{y_{i+1}}^I X_{y_{i+2}}^R \boxed{Z X_{y_{i+3}}^R X_{y_{i+4}}^o} \cdots X_{y_n}^F && \text{( by } P_N(14) \text{ )} \\
&\rightarrow^* \theta(X_{y_1}^D) \cdots \theta(X_{y_i}^D) X_{y_{i+1}}^I X_{y_{i+2}}^R X_{y_{i+3}}^R X_{y_{i+4}}^R \cdots \boxed{Z X_{y_{n-1}}^R X_{y_n}^F} && \text{( by } P_N(14) \text{ )} \\
&\rightarrow \theta(X_{y_1}^D) \cdots \theta(X_{y_i}^D) X_{y_{i+1}}^I X_{y_{i+2}}^R \cdots \boxed{X_{y_{n-2}}^R X_{y_n}^T Z} X_{y_{n-1}}^F && \text{( by } P_N(15) \text{ )} \\
&\rightarrow \theta(X_{y_1}^D) \cdots \theta(X_{y_i}^D) X_{y_{i+1}}^I X_{y_{i+2}}^R \cdots \boxed{X_{y_{n-3}}^R X_{y_n}^T Z} X_{y_{n-2}}^o X_{y_{n-1}}^F && \text{( by } P_N(16) \text{ )} \\
&\rightarrow^* \theta(X_{y_1}^D) \cdots \theta(X_{y_i}^D) \boxed{X_{y_{i+1}}^I X_{y_n}^T Z} X_{y_{i+2}}^o \cdots X_{y_{n-2}}^o X_{y_{n-1}}^F && \text{( by } P_N(16) \text{ )} \\
&\rightarrow \theta(X_{y_1}^D) \cdots \theta(X_{y_i}^D) X_{y_n}^D \boxed{Z X_{y_{i+1}}^I X_{y_{i+2}}^o} \cdots X_{y_{n-2}}^o X_{y_{n-1}}^F && \text{( by } P_N(17) \text{ )} \\
&\vdots \\
&\rightarrow^* \theta(X_{y_1}^D) \cdots \theta(X_{y_i}^D) X_{y_n}^D X_{y_{n-1}}^D \cdots \boxed{Z X_{y_{i+1}}^I X_{y_{i+2}}^F} && \text{( by prev. derivation )} \\
&\rightarrow \theta(X_{y_1}^D) \cdots \theta(X_{y_i}^D) X_{y_n}^D X_{y_{n-1}}^D \cdots X_{y_{i+2}}^D \boxed{X_{y_{i+1}}^D Z} && \text{( by } P_N(18) \text{ )} \\
&\rightarrow \theta(X_{y_1}^D) \cdots \theta(X_{y_i}^D) X_{y_n}^D X_{y_{n-1}}^D \cdots \boxed{X_{y_{i+2}}^D Z} y_{i+1} && \text{( by } P_N(11) \text{ )} \\
&\rightarrow^* \theta(X_{y_1}^D) \cdots \boxed{\theta(X_{y_i}^D) Z} y_n y_{n-1} \cdots y_{i+2} y_{i+1} && \text{( by } P_N(11) \text{ )} \\
&\rightarrow^* \boxed{Z} \theta(y_1) \cdots \theta(y_i) y_n y_{n-1} \cdots y_{i+2} y_{i+1} && \text{( by } P_N(7) \text{ )} \\
&\rightarrow bb\theta(y_1) \cdots \theta(y_i) y_n y_{n-1} \cdots y_{i+2} y_{i+1} && \text{( by } P_N(9) \text{ )}
\end{aligned}$$

Thus, for  $i < n - 1$ ,  $x \in L(G_N)$ . Therefore,  $bbL' \subseteq L(G_N)$ .

**Claim 2 :**  $L(G_N) \subseteq bbL'$  : Consider  $y' \in L(G_N)$ . We show that  $y'$  is in form  $bb\theta(u^R)v^R$  where  $uv \in L^R$ . Now,  $S'$  derives  $y'$ . Then, by  $P_N(1)$  and  $P_N(2)$ , for some  $\alpha \in L^R$ , either  $S' \rightarrow^* T_1h(\alpha)T_2 \rightarrow^* y'$  or  $S' \rightarrow^* T_1h(\alpha)T \rightarrow^* y'$ . Consider  $\alpha = a_1a_2 \cdots a_n$  where  $a_j \in \Sigma$ . Then, by definition of  $h$ ,  $S'$  derives

$$(2) \quad T_1X_{a_1}^o X_{a_2}^o \cdots X_{a_n}^o T_2 \text{ or } T_1X_{a_1}^o X_{a_2}^o \cdots X_{a_n}^o T.$$

- $S' \rightarrow^* T_1X_{a_1}^o X_{a_2}^o \cdots X_{a_n}^o T$ : Then, it generates exactly the following word:

$$\begin{aligned} S' &\rightarrow^* T_1X_{a_1}^o X_{a_2}^o \cdots X_{a_n}^o T \\ &\rightarrow^* \theta(X_{a_1}^D)\theta(X_{a_2}^D) \cdots \theta(X_{a_{n-1}}^D)T_1T_4X_{a_n}^o && \text{(by } P_N(3)\&P_N(6)\text{)} \\ &= \theta(X_{a_1}^D)\theta(X_{a_2}^D) \cdots \theta(X_{a_{n-1}}^D)Z''X_{a_n}^o \\ &\rightarrow \theta(X_{a_1}^D)\theta(X_{a_2}^D) \cdots \theta(X_{a_{n-1}}^D)X_{a_n}^D Z && \text{(by } P_N(10)\text{)} \\ &\rightarrow^* Z\theta(a_1)\theta(a_2) \cdots \theta(a_{n-1})a_n && \text{(by } P_N(11)\&P_N(7)\text{)} \\ &\rightarrow bb\theta(a_1)\theta(a_2) \cdots \theta(a_{n-1})a_n && \text{(by } P_N(9)\text{)} \end{aligned}$$

Then,  $y' = bb\theta(a_{n-1} \cdots a_2a_1)a_n \in bbL'$  where  $a_1a_2 \cdots a_n \in L^R$ .

- $S' \rightarrow^* T_1X_{a_1}^o X_{a_2}^o \cdots X_{a_n}^o T_2$ : Then, by applying  $P_N(3)$ ,  $P_N(4)$ ,  $P_N(5)$ , we have,

$$\begin{aligned} S' &\rightarrow^* T_1X_{a_1}^o X_{a_2}^o \cdots X_{a_n}^o T_2 \\ &\rightarrow^* \theta(X_{a_1}^D)\theta(X_{a_2}^D) \cdots \theta(X_{a_i}^D)T_1T_3X_{a_{i+1}}^o \cdots X_{a_{n-1}}^o X_{a_n}^F \\ &= \theta(X_{a_1}^D)\theta(X_{a_2}^D) \cdots \theta(X_{a_i}^D)ZX_{a_{i+1}}^o \cdots X_{a_{n-1}}^o X_{a_n}^F \end{aligned}$$

Now, we have the following cases:

- $i = n$ : Then,

$$\begin{aligned} S' &\rightarrow^* \theta(X_{a_1}^D)\theta(X_{a_2}^D) \cdots \theta(X_{a_n}^D)T_1T_2 \\ &= \theta(X_{a_1}^D)\theta(X_{a_2}^D) \cdots \theta(X_{a_n}^D)Z' \\ &\rightarrow^* Z'\theta(a_1)\theta(a_2) \cdots \theta(a_n) && \text{(by } P_N(8)\text{)} \\ &\rightarrow bb\theta(a_1)\theta(a_2) \cdots \theta(a_n) && \text{(by } P_N(9)\text{)} \end{aligned}$$

Then,  $y' = bb\theta(a_n \cdots a_2a_1) \in bbL'$  where  $a_1a_2 \cdots a_n \in L^R$ .

- $i = n - 1$ : Then, as  $Z = T_1T_3$ ,

$$S' \rightarrow^* \theta(X_{a_1}^D)\theta(X_{a_2}^D) \cdots \theta(X_{a_{n-1}}^D)ZX_{a_n}^F$$

which can not derive a sequence of terminals.

–  $i < n - 1$ : Then,

$$S' \rightarrow^* \theta(X_{a_1}^D)\theta(X_{a_2}^D)\cdots\theta(X_{a_i}^D)ZX_{a_{i+1}}^o \cdots X_{a_{n-1}}^o X_{a_n}^F.$$

Now, we can apply either  $P_N(7)$  or  $P_N(12)$ . If we apply  $P_N(7)$ , then we can not get a string of terminals. Thus, by applying  $P_N(12)$ , we have,

$$S' \rightarrow^* \theta(X_{a_1}^D)\theta(X_{a_2}^D)\cdots\theta(X_{a_i}^D)ZX_{a_{i+1}}^I X_{a_{i+2}}^o \cdots X_{a_{n-1}}^o X_{a_n}^F$$

Now, if we apply  $P_N(7)$ , then we can not get a string of terminals. So, we have to apply  $P_N(13)$ .

Let  $|X_{a_{i+1}}^I X_{a_{i+2}}^o \cdots X_{a_{n-1}}^o X_{a_n}^F| = m \geq 2$ . We now use induction on  $m$  to prove that

$$ZX_{a_{i+1}}^I X_{a_{i+2}}^o \cdots X_{a_{n-1}}^o X_{a_n}^F \rightarrow^* Za_n a_{n-1} \cdots a_{i+1}.$$

For  $m = 2$ ,

$$ZX_{a_{n-1}}^I X_{a_n}^F \rightarrow X_{a_n}^D X_{a_{n-1}}^D Z \rightarrow^* Za_n a_{n-1}.$$

For  $m = 3$ ,

$$\begin{aligned} \boxed{ZX_{a_{n-2}}^I X_{a_{n-1}}^o} X_{a_n}^F &\rightarrow X_{a_{n-2}}^I \boxed{ZX_{a_{n-1}}^R X_{a_n}^F} \\ &\rightarrow \boxed{X_{a_{n-2}}^I X_{a_n}^T Z} X_{a_{n-1}}^F \\ &\rightarrow X_{a_n}^D \boxed{ZX_{a_{n-2}}^I X_{a_{n-1}}^F} \\ &\rightarrow X_{a_n}^D X_{a_{n-1}}^D X_{a_{n-2}}^D Z \\ &\rightarrow^* Za_n a_{n-1} a_{n-2} \end{aligned}$$

Let the statement be true for strings with length less than  $m$ . Now,

$$\begin{aligned} \boxed{ZX_{a_{i+1}}^I X_{a_{i+2}}^o} \cdots X_{a_{n-1}}^o X_{a_n}^F &\rightarrow X_{a_{i+1}}^I \boxed{ZX_{a_{i+2}}^R X_{a_{i+3}}^o} \cdots X_{a_{n-1}}^o X_{a_n}^F \\ &\rightarrow X_{a_{i+1}}^I X_{a_{i+2}}^R ZX_{a_{i+3}}^R \cdots X_{a_{n-1}}^o X_{a_n}^F \\ &\rightarrow^* X_{a_{i+1}}^I X_{a_{i+2}}^R X_{a_{i+3}}^R \cdots \boxed{ZX_{a_{n-1}}^R X_{a_n}^F} \\ &\rightarrow X_{a_{i+1}}^I X_{a_{i+2}}^R X_{a_{i+3}}^R \cdots \boxed{X_{a_{n-2}}^R X_{a_n}^T Z} X_{a_{n-1}}^F \\ &\rightarrow X_{a_{i+1}}^I X_{a_{i+2}}^R X_{a_{i+3}}^R \cdots X_{a_{n-3}}^R X_{a_n}^T ZX_{a_{n-2}}^o X_{a_{n-1}}^F \end{aligned}$$

Here, we can apply either  $P_N(12)$  or  $P_N(16)$ . If we apply  $P_N(12)$ , then  $S'$  do not derive string of terminals. Thus, we apply  $P_N(16)$ . Then,

$$ZX_{a_{i+1}}^I X_{a_{i+2}}^o \cdots X_{a_{n-1}}^o X_{a_n}^F \rightarrow^* X_{a_{i+1}}^I X_{a_{i+2}}^R X_{a_{i+3}}^R \cdots X_{a_{n-4}}^R X_{a_n}^T ZX_{a_{n-3}}^o X_{a_{n-2}}^o X_{a_{n-1}}^F$$

Now, similar to the above, we can not apply  $P_N(12)$  here. So, we apply  $P_N(16)$  and continue the process. Then, we have,

$$ZX_{a_{i+1}}^I X_{a_{i+2}}^o \cdots X_{a_{n-1}}^o X_{a_n}^F \rightarrow^* \boxed{X_{a_{i+1}}^I X_{a_n}^T Z} X_{a_{i+2}}^o \cdots X_{a_{n-2}}^o X_{a_{n-1}}^F$$

$$\begin{aligned}
&\rightarrow X_{a_n}^D Z X_{a_{i+1}}^I X_{a_{i+2}}^o \cdots X_{a_{n-2}}^o X_{a_{n-1}}^F && \text{(by } P_N(17)) \\
&\rightarrow^* X_{a_n}^D Z a_{n-1} \cdots a_{i+1} && \text{(by induction)} \\
&\rightarrow Z a_n a_{n-1} \cdots a_{i+1} && \text{(by } P_N(11))
\end{aligned}$$

Therefore,

$$\begin{aligned}
S' &\rightarrow^* \theta(X_{a_1}^D) \theta(X_{a_2}^D) \cdots \theta(X_{a_i}^D) Z X_{a_{i+1}}^I X_{a_{i+2}}^o \cdots X_{a_{n-1}}^o X_{a_n}^F \\
&\rightarrow^* \theta(X_{a_1}^D) \theta(X_{a_2}^D) \cdots \theta(X_{a_i}^D) Z a_n a_{n-1} \cdots a_{i+1} \\
&\rightarrow^* bb\theta(a_1)\theta(a_2) \cdots \theta(a_i) a_n a_{n-1} \cdots a_{i+1}.
\end{aligned}$$

Then,  $y' = bb\theta(a_i a_{i-1} \cdots a_2 a_1) a_n a_{n-1} \cdots a_{i+1} \in bbL'$  where  $a_1 a_2 \cdots a_n \in L^R$ .

Therefore,  $L(G_N) \subseteq bbL'$ . □

For  $L \subseteq \Sigma^+$  and  $b \in \Sigma \setminus \text{Alph}(L)$ , we now prove that if  $bbL$  is context-sensitive, then  $L$  is context-sensitive. To do this we recall the following definition and a result from Rozenberg and Salomaa [20].

**Definition 4.6.** [20] Let  $k$  be a nonnegative integer. A morphism  $h, h : \Sigma^* \rightarrow \Delta^*$  is called  $k$ -erasing with respect to a language  $L \subseteq \Sigma^*$  iff for each  $w \in L$ ,

$$|w| \leq k \times |h(w)|.$$

A family  $\mathcal{L}$  of languages is closed under linear erasing iff  $h(L) \in \mathcal{L}$  for all  $L \in \mathcal{L}$  and for all  $k$ -erasing morphisms  $h, k \geq 0$ .

**Theorem 4.7.** [20] *The family of context-sensitive languages is closed under linear erasing.*

Using Theorem 4.7, we now prove that if  $bbL$  is context-sensitive for  $b \in \Sigma \setminus \text{Alph}(L)$ , then  $L$  is context-sensitive.

**Theorem 4.8.** *Consider  $L \subseteq \Sigma^+$  and  $bbL$  is a context-sensitive language for  $b \in \Sigma \setminus \text{Alph}(L)$ . Then,  $L$  is a context-sensitive language.*

*Proof.* Consider a 3-erasing morphism  $h : \Sigma^* \rightarrow \Sigma^*$  defined by  $h(b) = \lambda$  and  $h(a) = a$  for all  $a \in \Sigma \setminus \{b\}$ . Now, for each  $w \in bbL, w = bb\alpha$  for  $\alpha \in L$ . Then, as  $h(\alpha) = \alpha$  and  $h(w) = \alpha$ , we have,  $|w| \leq 3 \times |h(w)|$  and  $h(bbL) = L$ . Thus, by Theorem 4.7,  $L$  is context-sensitive. □

From Theorems 4.5 and 4.8, we have the following:

**Theorem 4.9.** *If  $L \subseteq \Sigma^+$  is context-sensitive, then  $C_\theta(L)$  is context-sensitive.*

*Proof.* Let  $\Sigma_1 = \text{Alph}(L)$ . Consider  $\Sigma' = \Sigma \cup \{b\}$  such that  $b \notin \Sigma_1 \cup \theta(\Sigma_1)$ . Now, as  $L$  is context-sensitive language over  $\Sigma'$ , by Theorem 4.5,  $bbC_\theta(L)$  is context-sensitive. Since,  $b \notin \Sigma_1 \cup \theta(\Sigma_1), b \notin \text{Alph}(C_\theta(L))$ . Then, by Theorem 4.8,  $C_\theta(L)$  is context-sensitive. Hence, the proof. □

## 5. ITERATED $\theta$ -CONJUGATES

In this section, we first define the iterated  $\theta$ -conjugates of strings and languages. Then, we characterize the elements of iterated  $\theta$ -conjugates. We also count the cardinality of iterated  $\theta$ -conjugates. Then, we study the closure property of iterated  $\theta$ -conjugate operation.

**Definition 5.1.** For a string  $w$  and a language  $L$ , we define  $C_\theta^n(w)$  recursively as  $C_\theta^0(w) = \{w\}$  and  $C_\theta^n(w) = C_\theta^{n-1}(C_\theta(w))$ , and  $C_\theta^n(L)$  recursively as  $C_\theta^0(L) = \{L\}$  and  $C_\theta^n(L) = C_\theta^{n-1}(C_\theta(L))$ . Then, we define  $C_\theta^*(w) = \bigcup_{n \geq 0} C_\theta^n(w)$  and  $C_\theta^*(L) = \bigcup_{n \geq 0} C_\theta^n(L)$ .

Since,  $w \in C_\theta(w)$  for  $w \in \Sigma^*$ , we have the following result for iterated  $\theta$ -conjugates whose proof is straight forward.

**Lemma 5.2.** For  $w \in \Sigma^*$  and  $m, k \geq 0$ ,  $C_\theta^m(w) \subseteq C_\theta^{m+k}(w)$ .

Since the conjugacy relation is an equivalence relation,  $u \in C(w)$  implies  $C(u) = C(w)$ . Then,  $C^2(w) = C(C(w)) = C(w)$ ,  $C^3(w) = C(C^2(w)) = C(w), \dots, C^n(w) = C(C^{n-1}(w))$ . But this is not true for iterated  $\theta$ -conjugates of a word. For an example,  $ba \in C_\theta^2(ab)$  but  $ba \notin C_\theta(ab)$  where  $\theta(a) = b$ . We now show that for  $w \in \Sigma^*$ ,  $C_\theta^*(w)$  is equivalent to  $\{u \mid |u|_a + |u|_{\theta(a)} = |w|_a + |w|_{\theta(a)} \text{ for all } a \in \Sigma\}$ . To prove this we need the following results.

From the definition of  $\theta$ -conjugates, it is clear that if  $u \in C_\theta^m(w)$  for  $m \geq 1$ , then for all  $a \in \text{Alph}(w)$ ,  $|u|_a + |u|_{\theta(a)} = |w|_a + |w|_{\theta(a)}$ .

**Lemma 5.3.** If  $u \in C_\theta^m(w)$  for  $m \geq 1$ , then  $|u|_a + |u|_{\theta(a)} = |w|_a + |w|_{\theta(a)}$  for all  $a \in \text{Alph}(w)$ .

We have an immediate result to Lemma 5.3.

**Corollary 5.4.** If  $u \in C_\theta^*(w)$ , then  $|u|_a + |u|_{\theta(a)} = |w|_a + |w|_{\theta(a)}$  for all  $a \in \text{Alph}(w)$ .

Consider for  $v \in \Sigma^+$  and  $a, b \in \Sigma$ , either  $|v|_a \geq 1$  or  $|v|_{\theta(a)} \geq 1$  and either  $|v|_b \geq 1$  or  $|v|_{\theta(b)} \geq 1$ , i.e.,  $|v|_a + |v|_{\theta(a)} \geq 1$  and  $|v|_b + |v|_{\theta(b)} \geq 1$ , respectively. Then, the following can be proved by direct computation.

**Lemma 5.5.** Consider  $v, z \in \Sigma^+$ ,  $|v| = 2$  and  $a, b \in \Sigma$ . If  $vz \in C_\theta^m(w)$  for  $m \geq 1$ ,  $|v|_a + |v|_{\theta(a)} \geq 1$  and  $|v|_b + |v|_{\theta(b)} \geq 1$ , then  $zab \in C_\theta^{m+4}(w)$ .

Also, if  $|u| \geq 3$  and  $uz \in C_\theta^m(w)$  for  $m \geq 1$  and  $|u|_a + |u|_{\theta(a)} \geq 1$ , then  $u'za \in C_\theta^{m+4}(w)$ . We have the following.

**Lemma 5.6.** Consider  $u, w \in \Sigma^+$ ,  $z \in \Sigma^*$ ,  $|u| \geq 3$  and  $a \in \Sigma$ . If  $uz \in C_\theta^m(w)$  for  $m \geq 1$  and  $|u|_a + |u|_{\theta(a)} \geq 1$ , then  $u'za \in C_\theta^{m+4}(w)$  for some  $u' \in \Sigma^+$ .

*Proof.* Let  $u \in \Sigma^+$  and  $uz \in C_\theta^m(w)$  where  $|u|_a + |u|_{\theta(a)} \geq 1$ , i.e., either  $|u|_a \geq 1$  or  $|u|_{\theta(a)} \geq 1$ . Then, we have two cases. If



- $|u|_a \geq 1$  : Then, for some  $\alpha, \beta \in \Sigma^*$ ,  $u = \alpha\beta$  and

$$\begin{aligned} \alpha\beta z \in C_\theta^m(w) &\implies \theta(z)\theta(\beta)\alpha \in C_\theta^{m+1}(w) \implies \theta(a)\theta(z)\theta(\beta)\alpha \in C_\theta^{m+2}(w) \\ &\implies \theta(\alpha)\beta z a \in C_\theta^{m+3}(w) \\ &\implies \theta(\alpha)\beta z a \in C_\theta^{m+4}(w) \quad (\text{ by Lemma 5.2}) \end{aligned}$$

- $|u|_{\theta(a)} \geq 1$  : Then, for some  $\alpha', \beta' \in \Sigma^*$ ,  $u = \alpha'\theta(a)\beta'$  and

$$\begin{aligned} \alpha'\theta(a)\beta' z \in C_\theta^m(w) &\implies \theta(z)\theta(\beta')\alpha' \in C_\theta^{m+1}(w) \implies \theta(\alpha')\theta(a)\theta(z)\theta(\beta') \in C_\theta^{m+2}(w) \\ &\implies \beta' z a \alpha' \in C_\theta^{m+3}(w) \\ &\implies \theta(\alpha')\beta' z a \in C_\theta^{m+4}(w). \end{aligned}$$

Hence, the proof.  $\square$

For  $w \in \Sigma$ , we can prove by direct computation,  $C_\theta^*(w) = C_\theta(w) = \{u \mid |u|_a + |u|_{\theta(a)} = |w|_a + |w|_{\theta(a)} \text{ for all } a \in \Sigma\}$ . Also, for  $w \in \Sigma^2$ , We can show by direct computation,  $C_\theta^*(w) = C_\theta^3(w) = \{u \mid |u|_a + |u|_{\theta(a)} = |w|_a + |w|_{\theta(a)} \text{ for all } a \in \Sigma\}$ . We now find  $C_\theta^*(w)$  for  $|w| \geq 3$ . From Lemmas 5.3, 5.5 and 5.6, we have the following.

**Theorem 5.7.** *Let  $w \in \Sigma^n$  and  $n \geq 3$ . Then,  $C_\theta^{4n-6}(w) = \{u \mid |u|_a + |u|_{\theta(a)} = |w|_a + |w|_{\theta(a)} \text{ for all } a \in \Sigma\}$ .*

*Proof.* Consider  $w \in \Sigma^n$ ,  $n \geq 3$  and  $T = \{u \mid |u|_a + |u|_{\theta(a)} = |w|_a + |w|_{\theta(a)} \text{ for all } a \in \Sigma\}$ . By Lemma 5.3, we have,  $C_\theta^{4n-6}(w) \subseteq T$ . Now, we prove that  $T \subseteq C_\theta^{4n-6}(w)$ . Let  $x \in T$ . Then, as  $|x|_a + |x|_{\theta(a)} = |w|_a + |w|_{\theta(a)}$  for all  $a \in \Sigma$ , we have,  $|x| = |w| = n \geq 3$ . Consider  $x = x_1 x_2 \cdots x_n$  where  $x_j \in \Sigma$  for  $1 \leq j \leq n$ . Then, as  $|x|_a + |x|_{\theta(a)} = |w|_a + |w|_{\theta(a)}$  for all  $a \in \Sigma$ , we have, for each  $i$  ( $1 \leq i \leq n$ ), either  $w = \alpha_i x_i \beta_i$  or  $w = \alpha_i \theta(x_i) \beta_i$  where  $\alpha_i, \beta_i \in \Sigma^*$ .

If  $w = \alpha_1 x_1 \beta_1$ , then  $\theta(\beta_1) \alpha_1 x_1 \in C_\theta(w)$ . Since for each  $i$  ( $2 \leq i \leq n-2$ ), either  $|\theta(\beta_1) \alpha_1|_{x_i} \geq 1$  or  $|\theta(\beta_1) \alpha_1|_{\theta(x_i)} \geq 1$ , by Lemma 5.6,  $v x_1 x_2 \cdots x_{n-2} \in C_\theta^{1+4(n-3)}(w)$ , i.e.,  $v x_1 x_2 \cdots x_{n-2} \in C_\theta^{4n-11}(w)$  where  $|v| = 2$ . Now,  $|v|_{x_n} + |v|_{\theta(x_n)} \geq 1$  and  $|v|_{x_{n-1}} + |v|_{\theta(x_{n-1})} \geq 1$ . Then, by Lemma 5.5,  $x_1 x_2 \cdots x_{n-2} x_{n-1} x_n \in C_\theta^{4n-11+4}(w)$ , i.e.,  $x_1 x_2 \cdots x_{n-2} x_{n-1} x_n \in C_\theta^{4n-7}(w)$ . Therefore, by Lemma 5.2,  $x_1 x_2 \cdots x_{n-2} x_{n-1} x_n \in C_\theta^{4n-6}(w)$ .

If  $w = \alpha_1 \theta(x_1) \beta_1$ , then  $\theta(\beta_1) x_1 \alpha_1 \in C_\theta(w)$  and  $\theta(\alpha_1) \theta(\beta_1) x_1 \in C_\theta^2(w)$ . Then, similar to the above,  $x_1 x_2 \cdots x_{n-2} x_{n-1} x_n \in C_\theta^{2+4(n-3)+4}(w)$ , i.e.,  $x_1 x_2 \cdots x_{n-2} x_{n-1} x_n \in C_\theta^{4n-6}(w)$ . Therefore,  $T \subseteq C_\theta^{4n-6}(w)$ .

Hence, the proof.  $\square$

From Corollary 5.4 and Theorem 5.7, we have the following result.

**Theorem 5.8.** *Let  $w \in \Sigma^n$  where  $n \geq 3$ . Then,  $C_\theta^*(w) = C_\theta^{4n-6}(w) = \{u \mid |u|_a + |u|_{\theta(a)} = |w|_a + |w|_{\theta(a)} \text{ for all } a \in \Sigma\}$ .*

Now, for  $w \in \Sigma^n$ ,  $C_\theta^*(w) = \{u \mid |u|_a + |u|_{\theta(a)} = |w|_a + |w|_{\theta(a)} \text{ for all } a \in \Sigma\}$ . Then, using basic counting principles, we have the following formula for the total number of elements in  $C_\theta^*(w)$  for  $w \in \Sigma^n$ .

**Theorem 5.9.** Let  $w \in \Sigma^n$  and  $x_1, x_2, \dots, x_k \in \text{Alph}(w)$  such that for  $j \neq i$ ,  $x_j \neq x_i$  and  $x_j \neq \theta(x_i)$ . Consider  $x_r \neq \theta(x_r)$  for  $1 \leq r \leq t \leq k$  and  $x_r = \theta(x_r)$  for  $t+1 \leq r \leq k$ . Let for  $1 \leq r \leq t$ ,  $|w|_{x_r} + |w|_{\theta(x_r)} = n_r$  and for  $t+1 \leq r \leq k$ ,  $|w|_{x_r} = m_r$ . If  $\sum_{r=1}^t n_r = n'$ ,  $\sum_{r=t+1}^k m_r = n''$  and  $n' + n'' = n$ , then

$$|C_\theta^*(w)| = \binom{n}{n_1 \ n_2 \ \dots \ n_t \ m_{t+1} \ m_{t+2} \ \dots \ m_k} \times 2^{n'}.$$

We illustrate Theorem 5.7 with the help of an example.

**Example 5.10.** Consider  $w = cag$  where  $\theta(a) = b$ ,  $\theta(c) = d$  and  $\theta(g) = h$ . Then,  $C_\theta(w) = \{hca, cag, hbd, hbc\}$  and  $C_\theta^2(w) = \{bdg, cag, hbc, bdh, cah, dhh, dah, bhc, chb, hca, hbd, dag\}$ . Similarly, we can find  $C_\theta^3(w)$ ,  $C_\theta^4(w)$ ,  $C_\theta^5(w)$  and  $C_\theta^6(w)$ . Now, one can observe that  $hac \in C_\theta^6(w) \setminus C_\theta^5(w)$ ,  $|C_\theta^6(w)| = 48$  and  $C_\theta^6(w) = C_\theta^m(w)$  for all  $m \geq 6$ . Hence,  $C_\theta^*(w) = C_\theta^6(w)$ .

Now, we study the closure property of iterated  $\theta$ -conjugate operation.

**Lemma 5.11.** Regular languages and context-free languages are not closed under iterated  $\theta$ -conjugate operation. Furthermore, the iterated  $\theta$ -conjugates of a regular language is not necessarily context-free.

*Proof.* Consider a regular language  $L = \{(bce)^*\}$  over  $\{b, c, d, e, f\}$  where  $\theta(b) = b$ ,  $\theta(c) = d$  and  $\theta(e) = f$ . Since,  $C_\theta^*(w) = \{u \mid |u|_a + |u|_{\theta(a)} = |w|_a + |w|_{\theta(a)} \text{ for all } a \in \Sigma\}$ ,  $C_\theta^*(L) = \bigcup_{w \in L} \{u \mid |u|_a + |u|_{\theta(a)} = |w|_a + |w|_{\theta(a)} \text{ for all } a \in \Sigma\}$ . Then,  $C_\theta^*(L) \cap \{b^{m_1}c^{m_2}e^{m_3} \mid m_i \geq 0\} = \{b^n c^n e^n \mid n \geq 0\}$  which is not context-free. Note that  $\{b^{m_1}c^{m_2}e^{m_3} \mid m_i \geq 0\}$  is a regular language. Since the regular languages and the context-free languages are closed under intersection with regular languages, the claim holds.  $\square$

## 6. $\theta$ -CONJUGATE-FREE LANGUAGES

In this section, we define a  $\theta$ -conjugate-free language  $L$ , where there are no two distinct strings in  $L$  such that one is a  $\theta$ -conjugate of the other. Then, we investigate some decidability problems for  $\theta$ -conjugate-freeness for different language classes.

**Definition 6.1.** A language  $L$  is  $\theta$ -conjugate-free if for all  $w \in L$ ,  $C_\theta(w) \cap L = \{w\}$ .

**6.1. Decidability of theta-conjugate-freeness.** Endrullis, Shallit, and Smith [5] proved that whether or not a regular language contains two distinct conjugates is decidable. Also, it is easy to prove whether or not a context-free language containing two distinct conjugates is undecidable. We now investigate the decidability problem for  $\theta$ -conjugate-freeness of  $L$  when  $L$  is either regular or context-free.

First we show that when  $L$  is a regular language with no  $\theta$ -palindromes, then  $L$  is  $\theta$ -conjugate-free or not is decidable.

**Theorem 6.2.** If  $L$  is a regular language with no  $\theta$ -palindrome, then  $L$  is  $\theta$ -conjugate-free or not is decidable.

*Proof.* Consider  $L$  is a regular language with no  $\theta$ -palindromes. From Proposition 3.3, we know that  $C_\theta(u'a) = \{u'a\} \cup \theta(a)C_\theta(u')$  for  $u' \in \Sigma^*$  and  $a \in \Sigma$ . Then,  $C_\theta(L) = \bigcup_{w \in L} C_\theta(w) = L \cup \bigcup_{u'a \in L} \theta(a)C_\theta(u')$  where  $u' \in \Sigma^*$  and  $a \in \Sigma$ . Consider  $T = \bigcup_{u'a \in L} \theta(a)C_\theta(u')$ . Then,  $C_\theta(L) = L \cup T$ . From Theorem 4.3, it is easy to prove that if  $L$  is regular, then  $T$  is also regular.

Now, we prove that  $L$  is  $\theta$ -conjugate-free iff  $L \cap T = \emptyset$ . Let  $L$  is  $\theta$ -conjugate-free. Then, for  $w \in L$ ,  $C_\theta(w) \cap L = \{w\}$ . If  $x \in L \cap T$ , then  $x \in L$  and  $x \in T$ , i.e.  $x \in \theta(a)C_\theta(z)$  for some  $za \in L$ . Then,  $za \in L$  and  $x \in C_\theta(za) \cap L$ . Now, if  $x = za$ , then for  $\alpha \in \Sigma^+$  and  $\beta \in \Sigma^*$ ,  $x = \theta(\alpha)\beta$ ,  $za = \beta\alpha$  and  $\theta(\alpha)\beta = \beta\alpha$  which implies  $x$  and  $za$  are  $\theta$ -palindromes which is a contradiction to the fact that  $L$  do not contain any  $\theta$ -palindrome. Thus,  $x \neq za$ , i.e.,  $\{x, za\} \subseteq C_\theta(za) \cap L$  which implies  $L$  is not  $\theta$ -conjugate-free which is a contradiction. Hence,  $L \cap T = \emptyset$ . Now, consider  $L \cap T = \emptyset$ . If  $w, w' \in L$  such that  $w \neq w'$  and  $w' \in C_\theta(w)$ , then  $w' \in T$  which implies  $w' \in L \cap T$ , a contradiction. Thus,  $L$  is  $\theta$ -conjugate-free. Therefore,  $L$  is  $\theta$ -conjugate-free iff  $L \cap T = \emptyset$ . Now,  $L$  and  $T$  are both regular languages and since we can check the intersection emptiness of two regular languages, we can decide whether or not  $L$  is  $\theta$ -conjugate-free.  $\square$

Now, we discuss the decidability result for a context-free language. First we recall the following undecidability result. An instance of the Post's Correspondence Problem (PCP) [19] consists of  $n \in \mathbb{N}$  and two ordered  $n$ -tuples of strings  $(U, V)$ , where  $U = (u_0, u_1, \dots, u_{n-1})$  and  $V = (v_0, v_1, \dots, v_{n-1})$ ,  $u_i, v_i \in \Sigma^*$ ,  $0 \leq i \leq n-1$ . A solution for the PCP instance  $(U, V)$  is a sequence of integers  $i_1, i_2, \dots, i_k$ ,  $0 \leq i_j \leq n-1$ ,  $j = 1, 2, \dots, k$ ,  $k \geq 1$ , such that

$$u_{i_1}u_{i_2} \cdots u_{i_k} = v_{i_1}v_{i_2} \cdots v_{i_k}.$$

**Proposition 6.3.** [19] *The decision problem of determining whether or not a given PCP instance has a solution is unsolvable.*

Now, we prove that deciding the  $\theta$ -conjugate-freeness of a given context-free language is undecidable by reducing PCP to this problem.

**Theorem 6.4.** *It is undecidable to determine whether or not a given context-free language  $L$  is  $\theta$ -conjugate-free.*

*Proof.* Let  $\Sigma$  be alphabet and  $(U, V)$  be an instance of PCP problem where  $U = (u_0, u_1, \dots, u_{n-1})$ ,  $V = (v_0, v_1, \dots, v_{n-1})$  and  $u_i, v_i \in \Sigma^*$ . Assume that the symbols  $\#, \%, \$, \&$  are distinct and not in  $\Sigma$ . Consider  $\Sigma' = \Sigma \cup \{\#, \%, \$, \&\}$  and  $\theta : \Sigma'^* \rightarrow \Sigma'^*$  be an antimorphic involution such that for all  $a \in \Sigma$ ,  $\theta(a) \in \Sigma$  and for all  $b \in \{\#, \%, \$, \&\}$ ,  $\theta(b) \in \{\#, \%, \$, \&\}$ .

We define a grammar  $G = (N, \Sigma', P, S)$  where  $N = \{S, N_U, N_V\}$  is the set of non-terminals,  $\Sigma'$  is the set of terminals,  $S$  is the start symbol and  $P$  is the set of derivation rules which includes the following rules:

- $S \rightarrow \%\$N_U\#\& \mid \theta(\&)\theta(\#)N_V\$\%$
- $N_U \rightarrow \%\$N_Uu_i \mid \%u_i$
- $N_V \rightarrow \theta(v_i)N_V\$\% \mid \theta(v_i)\%$

for  $i \in \{0, 1, \dots, n-1\}$ . Now, any string of  $L(G)$  is in one of the following forms :

$$(3) \quad (\%\$)^{l+1}\%u_{i_0}u_{i_1} \cdots u_{i_l}\#\& = \alpha \text{ (say)}$$

or

$$(4) \quad \theta(\&)\theta(\#)\theta(v_{i_1}) \cdots \theta(v_{i_l})\theta(v_{i_0})(\%\$)^{l+1}\% = \beta \text{ (say)}.$$

We now show that  $L(G)$  is not  $\theta$ -conjugate-free if and only if the PCP instance  $(U, V)$  has a solution.

First we prove that if the PCP instance  $(U, V)$  has a solution, then  $L(G)$  is not  $\theta$ -conjugate-free. Assume that the PCP instance  $(U, V)$  has a solution. Then, there is a sequence of integers  $i_0, i_1, \dots, i_k$  such that  $u_{i_0}u_{i_1} \cdots u_{i_k} = v_{i_0}v_{i_1} \cdots v_{i_k}$ . Then,  $L(G)$  contains the following elements:

$$\beta' = \theta(\&)\theta(\#)\theta(v_{i_0}v_{i_1} \cdots v_{i_k})(\%\$)^{k+1}\% = \theta(\&)\theta(\#)\theta(u_{i_0}u_{i_1} \cdots u_{i_k})(\%\$)^{k+1}\%$$

and

$$\alpha' = (\%\$)^{k+1}\%u_{i_0}u_{i_1} \cdots u_{i_k}\#\&.$$

Now,  $\beta' \in C_\theta(\alpha')$  and  $\beta' \neq \alpha'$ . Thus, as  $\alpha', \beta' \in L(G)$ ,  $L(G)$  is not  $\theta$ -conjugate-free.

Now, we prove the other part, i.e., if  $L(G)$  is not  $\theta$ -conjugate-free, then the PCP instance  $(U, V)$  has a solution. Assume that  $L(G)$  is not  $\theta$ -conjugate-free, i.e., there exists  $w = xy \in L(G)$  such that for  $y \in T^+$ ,  $\theta(y)x \in L(G)$  and  $w \neq \theta(y)x$ . Now, we have two cases:

- Consider  $xy$  is in the form of  $\alpha$ . Then, for some integer  $m$ ,  
 $xy = (\%\$)^{m+1}\%u_{i_0}u_{i_1} \cdots u_{i_m}\#\&$ . Since,  $\{\#, \%, \$, \&\} \cap \Sigma = \emptyset$ ,  $y \in T^+$  and  $\theta(a) \in \Sigma$  for all  $a \in \Sigma$ ,  $\theta(y)x$  will be in the form of  $\beta$ . Then, there are two possible decompositions as follows:
  - (1)  $x = (\%\$)^{m+1}\%$  and  $y = u_{i_0}u_{i_1} \cdots u_{i_m}\#\&$ . Now,  $\theta(y)x$  is in form  $\beta$  and  $(\%\$)^{m+1}\%$  is a suffix of  $\theta(y)x$  for some integer  $m+1$ . Then, as  $\theta(u_{i_0}u_{i_1} \cdots u_{i_m}\#\&) = \theta(\&)\theta(\#)\theta(v_{i_m}) \cdots \theta(v_{i_1})\theta(v_{i_0})$ , the PCP instance  $(U, V)$  has a solution.
  - (2)  $x = (\%\$)^{m'}\%$  and  $y = (\%\$)^{m''}u_{i_0}u_{i_1} \cdots u_{i_m}\#\&$  where  $m+1 = m'+m''$  and  $m'' \geq 1$ . Since,  $\theta(y)x$  is in form  $\beta$  and  $\theta(a) \in \Sigma$  for all  $a \in \Sigma$ ,  $\theta(\%) = \%$  and  $\theta(\$) = \$$ . Then,  $(\%\$)^{m+1}\%$  is a suffix of  $\theta(y)x$  for some integer  $m+1$ . Thus, as  $\theta(u_{i_0}u_{i_1} \cdots u_{i_m}\#\&) = \theta(\&)\theta(\#)\theta(v_{i_m}) \cdots \theta(v_{i_1})\theta(v_{i_0})$ , the PCP instance  $(U, V)$  has a solution.
- Consider  $xy$  is in the form of  $\beta$ . Then, for some integer  $r$ ,  
 $xy = \theta(\&)\theta(\#)\theta(v_{i_r}) \cdots \theta(v_{i_1})\theta(v_{i_0})(\%\$)^{r+1}\%$ . Since,  $\{\#, \%, \$, \&\} \cap \Sigma = \emptyset$ ,  $y \in T^+$  and  $\theta(a) \in \Sigma$  for all  $a \in \Sigma$ ,  $\theta(y)x$  will be in the form of  $\alpha$ . Then, there are three possible decompositions as follows:
  - (1)  $x = \theta(\&)\theta(\#)$  and  $y = \theta(v_{i_r}) \cdots \theta(v_{i_1})\theta(v_{i_0})(\%\$)^{r+1}\%$ .
  - (2)  $x = \theta(\&)$  and  $y = \theta(\#)\theta(v_{i_r}) \cdots \theta(v_{i_1})\theta(v_{i_0})(\%\$)^{r+1}\%$ .
  - (3)  $x = \lambda$  and  $y = \theta(\&)\theta(\#)\theta(v_{i_r}) \cdots \theta(v_{i_1})\theta(v_{i_0})(\%\$)^{r+1}\%$ .
Since,  $\theta(y)x$  is in form  $\alpha$  and  $\theta(a) \in \Sigma$  for all  $a \in \Sigma$ , we have,  $\theta(\%) = \%$ ,  $\theta(\$) = \$$  and  $(\%\$)^{r+1}\%$  is a prefix of  $\theta(y)x$  for some integer  $r+1$ . Thus, as  $\theta(\theta(v_{i_r}) \cdots \theta(v_{i_1})\theta(v_{i_0})(\%\$)^{r+1}\%) = (\%\$)^{r+1}\%u_{i_0}u_{i_1} \cdots u_{i_r}$ , the PCP instance  $(U, V)$  has a solution.

Therefore, as PCP is undecidable [19], it is also undecidable whether or not  $L$  is  $\theta$ -conjugate-free when  $L$  is context-free.  $\square$

## 7. CONCLUSIONS

In this paper, we have discussed the properties of the  $\theta$ -conjugates, being motivated by encoding information into DNA molecules. First we have characterized the elements of  $\theta$ -conjugates. Then, we have characterized  $u$  and  $v$  such that  $C_\theta(u) = C_\theta(v)$ . Also, we have investigated the closure property of certain family of languages under  $\theta$ -conjugate operation. We have discussed the iterated  $\theta$ -conjugates of a word and languages. Then, we have studied the concept of  $\theta$ -conjugate-free language and some decidability problems for  $\theta$ -conjugate-freeness for different language classes. For regular language  $L$  with no  $\theta$ -palindrome, it is decidable whether or not  $L$  is  $\theta$ -conjugate-free. But for general regular language  $L$ , the result is still unknown. Also, it would be interesting if we find maximum number of primitive,  $\theta$ -primitive, bordered,  $\theta$ -bordered and squared words in the set of all Watson-Crick conjugates of a word.

## REFERENCES

- [1] A. Amir and Y. Aumann and Gad M. Landau and M. Lewenstein and N. Lewenstein. Pattern Matching with Swaps. *Journal of Algorithms*, 37(2): 247-266, 2000.
- [2] F. Blanchet-Sadri and D.K. Luhmann. Conjugacy on partial words. *Theoretical Computer Science*, 289(1): 297-312, 2002.
- [3] A. Brandstädt. Closure properties of certain families of formal languages with respect to a generalization of cyclic closure. *RAIRO. Informatique théorique*, 15(3): 233-252, 1981.
- [4] Elena Czeizler and Lila Kari and Shinnosuke Seki. On a special class of primitive words. *Theoretical Computer Science*, 411(3): 617-630, 2010.
- [5] J. Endrullis and J. Shallit, and T. Smith. Undecidability and Finite Automata. *Developments in Language Theory, Lecture Notes in Computer Science*, 10396: 160-172, Springer, 2017.
- [6] Chuan Guo and Jeffrey Shallit, and Arseny M Shur. On the combinatorics of palindromes and antipalindromes. *arXiv preprint arXiv:1503.09112*, 2015.
- [7] László Hegedüs and Benedek Nagy. On periodic properties of circular words. *Discrete Mathematics*, 339(3): 1189-1197, 2016.
- [8] G. Jirásková and A. Okhotin. State complexity of cyclic shift. *RAIRO-Theoretical Informatics and Applications*, 42(2): 335-360, 2008.
- [9] Lila Kari and Kalpana Mahalingam. Watson-Crick Conjugate and Commutative Words. *DNA Computing, Lecture Notes in Computer Science*, 4848: 273-283, Springer, 2008.
- [10] Lila Kari and Kalpana Mahalingam. Watson-Crick palindromes in DNA computing. *Natural Computing*, 9(2): 297-316, 2010.
- [11] O. Lipsky, B. Porat, E. Porat, B. Riva Shalom, and A. Tzur. String matching with up to  $k$  swaps and mismatches. *Information and Computation*, 208(9): 1020-1030, 2010.
- [12] O. Lipsky, B. Porat, E. Porat, B. Riva Shalom, and A. Tzur. Approximate String Matching with Swap and Mismatch. *Algorithms and Computation, Lecture Notes in Computer Science*, 4835: 869-880, Springer, 2007.
- [13] M. Lothaire. *Combinatorics on words*. Cambridge University Press, 1997.
- [14] R. C. Lyndon and M.P. Schützenberger. The equation  $a^M = b^N c^P$  in a free group. *Michigan Mathematical Journal*, 9: 289-298, 1962.

- [15] Kalpana Mahalingam and Anuran Maity, and Palak Pandoh. Counting (Watson-Crick) palindromes in Watson-Crick conjugates. *Information and Computation*, 285: 104863, 2022.
- [16] Kalpana Mahalingam and Anuran Maity, and Palak Pandoh, and Rama Raghavan. Block reversal on finite words. *Theoretical Computer Science*, 894: 135-151, 2021.
- [17] A. Maslov. Estimates of the number of states of finite automata. *Doklady Akademii Nauk SSSR*, 194(6): 1266-1268, 1970.
- [18] T. Oshiba. Closure property of the family of context-free languages under the cyclic shift operation. *Electronics & communications in Japan*, 55(4): 119-122, 1972.
- [19] Emil L Post. A variant of a recursively unsolvable problem. *Bulletin of the American Mathematical Society*, 52(4): 264-268, 1946
- [20] G. Rozenberg and A. Salomaa(Eds.). Handbook of Formal Languages. Volume 1. *Springer*, 1997.
- [21] Jeffrey Shallit. A second course in formal languages and automata theory. *Cambridge University Press*, 2008.
- [22] H. J. Shyr. Free Monoids and Languages. *Hon Min Book Company*, 2001.

DEPARTMENT OF MATHEMATICS,, INDIAN INSTITUTE OF TECHNOLOGY MADRAS, CHENNAI, 600036, INDIA

*Email address:* [kmahalingam@iitm.ac.in](mailto:kmahalingam@iitm.ac.in), [anuran.maity@gmail.com](mailto:anuran.maity@gmail.com)