

TOEPLITZ SEPARABILITY, ENTANGLEMENT, AND COMPLETE POSITIVITY USING OPERATOR SYSTEM DUALITY

DOUGLAS FARENICK AND MICHELLE MCBURNEY

ABSTRACT. A new proof is presented of a theorem of L. Gurvits, which states that the cone of positive block-Toeplitz matrices with matrix entries has no entangled elements. We also show that in the cone of positive Toeplitz matrices with Toeplitz entries, entangled elements exist in all dimensions. The proof of the Gurvits separation theorem is achieved by making use of the structure of the operator system dual of the operator system $C(S^1)^{(n)}$ of $n \times n$ Toeplitz matrices over the complex field, and by determining precisely the structure of the generators of the extremal rays of the positive cones of the operator systems $C(S^1)^{(n)} \otimes_{\min} \mathcal{B}(\mathcal{H})$ and $C(S^1)_{(n)} \otimes_{\min} \mathcal{B}(\mathcal{H})$, where \mathcal{H} is an arbitrary Hilbert space and $C(S^1)_{(n)}$ is the operator system dual of $C(S^1)^{(n)}$. Our approach also has the advantage of providing some new information concerning positive Toeplitz matrices whose entries are from the type I factor $\mathcal{B}(\mathcal{H})$. In particular, we prove that normal positive linear maps ψ on $\mathcal{B}(\mathcal{H})$ are partially completely positive in the sense that $\psi^{(n)}(x)$ is positive whenever x is a positive $n \times n$ Toeplitz matrix with entries from $\mathcal{B}(\mathcal{H})$. We also establish a certain factorisation theorem for positive Toeplitz matrices (of operators), and explain how the factorisation theorem can be used to show an equivalence between the Gurvits approach to separation and an earlier approach of T. Ando to universality.

1. INTRODUCTION

If \otimes_{σ} is an operator system tensor product of operator systems \mathcal{R} and \mathcal{S} , then an element x in the positive cone of $\mathcal{R} \otimes_{\sigma} \mathcal{S}$ is said to be \otimes_{σ} -separable if

$$x = \sum_{j=1}^m r_j \otimes s_j,$$

for some positive $r_j \in \mathcal{R}$ and positive $s_j \in \mathcal{S}$. If a positive x is not \otimes_{σ} -separable, then it is said to be \otimes_{σ} -entangled. When the operator system tensor product \otimes_{σ} is the spatial one \otimes_{\min} , then we simply used the terms “separable” and “entangled” to describe the two possible scenarios for positive elements of $\mathcal{R} \otimes_{\min} \mathcal{S}$.

Following [5, 6, 10, 15], the operator system of $n \times n$ Toeplitz matrices over the complex numbers is denoted by $C(S^1)^{(n)}$. The canonical linear basis for $C(S^1)^{(n)}$

2020 *Mathematics Subject Classification.* 46L07, 47L05, 47L07.
Supported in part by the NSERC Discovery Grant program.

is the one given by $\{r_{-n+1}, \dots, r_{-1}, r_0, r_1, \dots, r_{n-1}\}$, whereby

$$r_k = \begin{cases} s^k & : \text{ if } k \geq 0 \\ (s^*)^k & : \text{ if } k < 0 \end{cases} \quad \text{and} \quad s = \begin{bmatrix} 0 & & & & & \\ 1 & 0 & & & & \\ & 1 & 0 & & & \\ & & \ddots & \ddots & & \\ & & & \ddots & \ddots & \\ & & & & 1 & 0 \end{bmatrix}.$$

The identity matrix r_0 serves as the Archimedean order unit for the operator system $C(S^1)^{(n)}$.

If \mathcal{S} is any complex vector space, then an element $x = \sum_{\ell=-n+1}^{n-1} r_\ell \otimes \tau_\ell$ of the algebraic tensor product $C(S^1)^{(n)} \otimes \mathcal{S}$ can be canonically identified with the (block Toeplitz) matrix

$$(1.1) \quad x = \begin{bmatrix} \tau_0 & \tau_{-1} & \tau_{-2} & \dots & \tau_{-n+2} & \tau_{-n+1} \\ \tau_1 & \tau_0 & \tau_{-1} & \tau_{-2} & \dots & \tau_{-n+2} \\ \tau_2 & \tau_1 & \tau_0 & \tau_{-1} & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \tau_{-2} \\ \tau_{n-2} & & \ddots & \ddots & \ddots & \tau_{-1} \\ \tau_{n-1} & \tau_{n-2} & \dots & \tau_2 & \tau_1 & \tau_0 \end{bmatrix}.$$

Our first interest in the present paper is with the case where $\mathcal{S} = \mathcal{M}_p(\mathbb{C})$, the C^* -algebra of $p \times p$ complex matrices. In this regard, the following theorem was communicated in [8, 9].

Theorem 1.1 (Gurvits). *Every positive element of $C(S^1)^{(n)} \otimes_{\min} \mathcal{M}_p(\mathbb{C})$ is separable.*

One of the main goals of the present paper is to provide an alternative approach to proving Theorem 1.1, which we present in Section §3 by making use of recent work in [5, 6] on operator system duality and by considering the structure of the extremal rays of the positive cone of $C(S^1)^{(n)} \otimes_{\min} \mathcal{M}_p(\mathbb{C})$. The advantage of this approach is that it is somewhat more general in that we consider Toeplitz matrices whose entries are bounded linear operators on Hilbert spaces of arbitrary dimension. A secondary advantage of this approach is that it applies to other operator systems—in particular, to the operator system dual of $C(S^1)^{(n)}$.

The operator system $C(S^1)^{(n)}$ and its operator system dual, $C(S^1)_{(n)}$, are receiving recent attention because of their importance to spectral truncation in non-commutative geometry [5, 10, 15]. It is, therefore, interesting to determine whether there is an analogue of Theorem 1.1 when $C(S^1)^{(n)}$ is replaced by its operator system dual $C(S^1)_{(n)}$. There is not; in fact, we shall show in Section §4 that there are extremal rays of the positive cone of $C(S^1)_{(n)} \otimes_{\min} \mathcal{M}_n(\mathbb{C})$ whose generators are entangled.

Throughout this paper, \mathbb{C}^p is considered as a p -dimensional Hilbert space, where the inner product on \mathbb{C}^p is the standard inner product. Hilbert spaces of arbitrary (finite or infinite) dimension are denoted by \mathcal{H} , and $\mathcal{B}(\mathcal{H})$ denotes the algebra of all bounded linear operators $x : \mathcal{H} \rightarrow \mathcal{H}$. In the case where $\mathcal{H} = \mathbb{C}^p$, then $\mathcal{B}(\mathcal{H}) = \mathcal{M}_p(\mathbb{C})$.

2. DUALITY, PURITY, AND UNIVERSALITY

Recall from [4] that if \mathcal{R} is an operator system and \mathcal{R}^d denotes its dual space, then \mathcal{R}^d is a matrix-ordered $*$ -vector space in which a matrix $\Phi = [\varphi_{ij}]_{i,j=1}^p$ of linear functionals $\varphi_{ij} : \mathcal{R} \rightarrow \mathbb{C}$ is considered positive whenever the linear map $r \mapsto [\varphi_{ij}(r)]_{i,j=1}^p$ is a completely positive linear map $\hat{\Phi} : \mathcal{R} \rightarrow \mathcal{M}_p(\mathbb{C})$. Furthermore, if $\phi : \mathcal{R} \rightarrow \mathcal{S}$ is a linear map of operator systems and $\phi^d : \mathcal{S}^d \rightarrow \mathcal{R}^d$ denotes the adjoint transformation as linear mapping of matrix-ordered $*$ -vector spaces, then ϕ is positive if and only if ϕ^d is positive. Likewise, $\phi \otimes \text{id}_{\mathcal{M}_p(\mathbb{C})}$ is positive if and only if $\phi^d \otimes \text{id}_{\mathcal{M}_p(\mathbb{C})}$ is positive, for $p \in \mathbb{N}$. Importantly, if \mathcal{R} has finite dimension, then the matrix-ordered space \mathcal{R}^d is in fact an operator system, and any faithful state on \mathcal{R} serves as an Archimedean order unit for the matrix ordering of \mathcal{R}^d .

For any operator system \mathcal{R} , the cone of positive $p \times p$ matrices over \mathcal{R} is denoted by $\mathcal{M}_p(\mathcal{R})_+$. With respect to the operator system tensor product \otimes_{\min} , we have

$$\mathcal{M}_p(\mathcal{R})_+ = (\mathcal{R} \otimes_{\min} \mathcal{M}_p(\mathcal{R}))_+.$$

Because $\mathcal{M}_p(\mathcal{R})_+$ is a cone, it admits extremal rays. (Recall that a face \mathcal{F} in a proper convex cone \mathcal{C} is an extremal ray if there exists an element $\phi \in \mathcal{C}$, called a generator of the extremal ray, such that $\mathcal{F} = \{\alpha\phi \mid t \in \mathbb{R}, \alpha \geq 0\}$.)

We shall adopt the term “pure” for elements in convex cones that generate extremal rays.

Definition 2.1. An element ϕ of a convex cone \mathcal{C} is *pure* if the equation $\vartheta + \omega = \phi$, for some $\vartheta, \omega \in \mathcal{C}$, holds only if there exists a scalar $s \in [0, 1]$ such that $\vartheta = s\phi$ and $\omega = (1 - s)\phi$.

The purpose in identifying pure elements of proper convex cones \mathcal{C} in finite-dimensional vector spaces is that they generate, through arbitrary linear combinations of pure elements using positive real coefficients, all elements of \mathcal{C} . The cones \mathcal{C} that shall be of interest here are the cones $\mathcal{M}_p(\mathcal{R})_+$ and $\mathcal{CP}(\mathcal{R}, \mathcal{S})$, the latter being the cone of completely positive linear maps of the operator system \mathcal{R} into the operator system \mathcal{S} .

The duality mentioned earlier amounts to the following positivity criterion for a matrix $\Phi \in \mathcal{M}_p(\mathcal{R}^d)$ of linear functionals on \mathcal{R} :

$$(2.1) \quad \Phi \in \mathcal{M}_p(\mathcal{R}^d)_+ \text{ if and only if } \hat{\Phi} \in \mathcal{CP}(\mathcal{R}, \mathcal{M}_p(\mathbb{C})).$$

Furthermore, when \mathcal{R} has finite dimension, the positivity condition (2.1) admits a more general form whereby the matrix algebra $\mathcal{M}_p(\mathbb{C})$ is replaced by an arbitrary operator system \mathcal{S} [11, Section 4], [12, Lemma 8.5]. Specifically, if

$$t = \sum_{j=1}^m x_j \otimes y_j \in \mathcal{R} \otimes_{\min} \mathcal{S},$$

and if $\hat{t} : \mathcal{R}^d \rightarrow \mathcal{S}$ is the linear map defined by

$$\hat{t}(\varphi) = \sum_{j=1}^m \varphi(x_j) y_j, \text{ for } \varphi \in \mathcal{R}^d,$$

then $t \in (\mathcal{R} \otimes_{\min} \mathcal{S})_+$ if and only if \hat{t} is completely positive. Thus, from $(\mathcal{R}^d)^d = \mathcal{R}$, we have

$$(2.2) \quad \Phi \in (\mathcal{R}^d \otimes_{\min} \mathcal{S})_+ \text{ if and only if } \hat{\Phi} \in \mathcal{CP}(\mathcal{R}, \mathcal{S}).$$

Returning now to the study of positive Toeplitz matrices, let $C(S^1)$ denote the unital abelian C^* -algebra of all continuous functions $f : S^1 \rightarrow \mathbb{C}$, where $S^1 \subset \mathbb{C}$ is the unit circle. For each $n \in \mathbb{N}$, $C(S^1)_{(n)}$ shall denote the vector space of those $f \in C(S^1)$ for which the Fourier coefficients $\hat{f}(k)$ of f satisfy $\hat{f}(k) = 0$ for every $k \in \mathbb{Z}$ such that $|k| \geq n$. Hence, every $f \in C(S^1)_{(n)}$ is given by

$$f(z) = \sum_{k=-n+1}^{n-1} \alpha_k z^k,$$

as a function of $z \in S^1$, where each $\alpha_k = \hat{f}(k) = \frac{1}{2\pi} \int_0^{2\pi} f(e^{i\theta}) e^{-ik\theta} d\theta$. The vector space $C(S^1)_{(n)}$ is an operator system via the matrix ordering that arises from the identification of $\mathcal{M}_p(C(S^1)_{(n)})$, the space of $p \times p$ matrices with entries from $C(S^1)_{(n)}$, with the space of continuous functions $F : S^1 \rightarrow \mathcal{M}_p(\mathbb{C})$, and where the Archimedean order unit is the canonical one (namely, the constant function $\chi_0 : S^1 \rightarrow \mathbb{C}$ given by $\chi_0(z) = 1$, for $z \in S^1$). (The operator system of all $n \times n$ Toeplitz matrices over \mathbb{C} , which we have been denoting by $C(S^1)^{(n)}$, has the identity matrix in $\mathcal{M}_n(\mathbb{C})$ as the canonical Archimedean order unit for $C(S^1)^{(n)}$.)

The following theorem was obtained in [5, 6].

Theorem 2.2 (Toeplitz Duality). *The linear map $\phi : C(S^1)^{(n)} \rightarrow (C(S^1)_{(n)})^d$ that sends a Toeplitz matrix $t = [\tau_{k-\ell}]_{k,\ell=0}^{n-1} \in \mathcal{M}_n(\mathbb{C})$ to the linear functional $\varphi_t : C(S^1)_{(n)} \rightarrow \mathbb{C}$ defined by*

$$(2.3) \quad \varphi_t(f) = \sum_{k=-n+1}^{n-1} \tau_{-k} \hat{f}(k),$$

for $f \in C(S^1)_{(n)}$, is a unital complete order isomorphism. That is, $C(S^1)^{(n)}$ is the operator system dual of $C(S^1)_{(n)}$.

Definition 2.3. The universal positive $n \times n$ Toeplitz matrix is the Toeplitz-matrix-valued function $T_n : S^1 \rightarrow \mathcal{M}_n(\mathbb{C})$ given by

$$(2.4) \quad T_n(z) = \begin{bmatrix} 1 & z^{-1} & z^{-2} & \dots & z^{-n+2} & z^{-n+1} \\ z & 1 & z^{-1} & z^{-2} & \dots & z^{-n+2} \\ z^2 & z & 1 & z^{-1} & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & \ddots & z^{-2} \\ z^{n-2} & & \ddots & \ddots & \ddots & z^{-1} \\ z^{n-1} & z^{n-2} & \dots & z^2 & z & 1 \end{bmatrix}.$$

Because the matrix $T_n(z)$ defined in (2.4) admits a factorisation of the form $T_n(z) = \gamma_n(z)\gamma_n(z)^*$, where

$$\gamma_n(z) = \begin{bmatrix} 1 \\ z \\ z^2 \\ \vdots \\ z^{n-1} \end{bmatrix},$$

the matrix-valued function $T_n : S^1 \rightarrow \mathcal{M}_n(\mathbb{C})$ maps S^1 into the rank-1 positive $n \times n$ Toeplitz matrices. The term “universal” is used because, for every operator system \mathcal{S} and positive Toeplitz matrix $x = [\tau_{k-j}]_{k,j=1}^n \in C(S^1)^{(n)} \otimes_{\min} \mathcal{S}$, there exists a completely positive linear map $\phi : C(S^1)_{(n)} \rightarrow \mathcal{S}$ for which $\tau_\ell = \phi(z^\ell)$, for every ℓ [1], [6, Theorem 7.10].

We shall show in Section §4 that T_n is an entangled element of the positive cone of $C(S^1)_{(n)} \otimes_{\min} \mathcal{M}_n(\mathbb{C})$, for every $n \geq 2$, in contrast to the dual situation of $C(S^1)^{(n)} \otimes_{\min} \mathcal{M}_n(\mathbb{C})$, which has no entangled positive elements.

3. TOEPLITZ SEPARABILITY

The primary purpose of this section is to give a proof of the separation theorem, Theorem 1.1, by determining the structure of the pure elements of the cone $(C(S^1)^{(n)} \otimes_{\min} \mathcal{B}(\mathcal{H}))_+$ for Hilbert spaces \mathcal{H} of finite or infinite dimension. Thus, this approach addresses a more general context than that considered by Gurvits in [8, 9]. In Section §4, the same methodology is used in determining the structure of the pure elements of the cone $(C(S^1)_{(n)} \otimes_{\min} \mathcal{B}(\mathcal{H}))_+$, replacing $C(S^1)^{(n)}$ by its operator system dual $C(S^1)_{(n)}$.

The following observation, inspired by [5, Corollary 2.20], is key to our approach.

Lemma 3.1. *If \mathcal{S} is any operator system, then an element $\Phi \in (C(S^1)^{(n)} \otimes_{\min} \mathcal{S})_+$ is pure if and only if $\hat{\Phi}$ is a pure completely positive linear map $C(S^1)_{(n)} \rightarrow \mathcal{S}$.*

Proof. Assume $\Phi \in (C(S^1)^{(n)} \otimes_{\min} \mathcal{S})_+$ is pure. Thus, if $\Phi = \sum_{\ell=-n+1}^{n-1} r_\ell \otimes \tau_\ell$, then

$$\hat{\Phi} = \sum_{\ell=-n+1}^{n-1} \varphi_{r_\ell} \tau_\ell, \text{ where the action of } \hat{\Phi} \text{ on } C(S^1)_{(n)} \text{ is given by}$$

$$\hat{\Phi}(f) = \sum_{\ell=-n+1}^{n-1} \hat{f}(-\ell) \tau_\ell,$$

for every $f \in C(S^1)_{(n)}$.

Suppose that there exists completely positive maps of $\theta, \psi : C(S^1)_{(n)} \rightarrow \mathcal{S}$ such that $\hat{\Phi} = \theta + \psi$. Let $\theta(z^k) = t_k$ and $\psi(z^k) = s_k$, so that

$$\theta(f) = \theta \left(\sum_{k=-n+1}^{n-1} \hat{f}(k) z^k \right) = \sum_{k=-n+1}^{n-1} \hat{f}(k) \theta(z^k) = \sum_{k=-n+1}^{n-1} \hat{f}(k) t_k$$

and, similarly,

$$\psi(f) = \sum_{k=-n+1}^{n-1} \hat{f}(k) s_k,$$

for every $f \in C(S^1)_{(n)}$. We obtain, therefore, Toeplitz matrices

$$T = \begin{bmatrix} t_0 & t_1 & t_2 & \cdots & t_{n-1} \\ t_{-1} & t_0 & t_1 & & \\ \vdots & & \ddots & & t_1 \\ t_{-n+1} & \cdots & \cdots & t_{-1} & t_0 \end{bmatrix} \text{ and } S = \begin{bmatrix} s_0 & s_1 & s_2 & \cdots & s_{n-1} \\ s_{-1} & s_0 & s_1 & & \\ \vdots & & \ddots & & s_1 \\ s_{-n+1} & \cdots & \cdots & s_{-1} & s_0 \end{bmatrix}.$$

For each k ,

$$\tau_{-k} = \hat{\Phi}(z^k) = \vartheta(z^k) + \psi(z^k) = t_k + s_k,$$

and so $\tau_{-k} = t_k + s_k = a_{-k} + b_{-k}$, where $a_k = t_{-k}$ and $b_k = s_{-k}$. Thus, $\Phi = A + B$. If A and B were positive elements of $C(S^1)^{(n)} \otimes_{\min} \mathcal{S}$, then by the purity of Φ we would have $A = \alpha\Phi$ and $B = (1 - \alpha)\Phi$ for some $\alpha \in [0, 1]$, implying that $\theta = \alpha\hat{\Phi}$ and $\psi = (1 - \alpha)\hat{\Phi}$ and thereby proving the purity of $\hat{\Phi}$. To confirm $A, B \in (C(S^1)^{(n)} \otimes_{\min} \mathcal{S})_+$, we need only show that \hat{A} and \hat{B} are completely positive. However, this is true by the way in which A and B are defined leads $\hat{A} = \theta$ and $\hat{B} = \psi$.

The proof of the converse is similar and it, thus, omitted. \square

Theorem 3.2. *For every Hilbert space \mathcal{H} , a matrix $T \in (C(S^1)^{(n)} \otimes_{\min} \mathcal{B}(\mathcal{H}))_+$ is pure if and only if there exists a complex number $\lambda \in S^1$, a rank-1 projection $Q \in \mathcal{B}(\mathcal{H})$, and a scalar $\alpha \geq 0$ such that*

$$T = T_n(\lambda^{-1}) \otimes (\alpha Q).$$

Proof. Suppose that $T = [\tau_{k-j}]_{k,j=1}^n \in (C(S^1)^{(n)} \otimes_{\min} \mathcal{B}(\mathcal{H}))_+$ is pure, and express T in its canonical tensor representation with respect to the canonical linear basis of $C(S^1)^{(n)}$:

$$T = \sum_{\ell=-n+1}^{n-1} r_\ell \otimes \tau_\ell.$$

By the Duality Theorem, Theorem 2.2, each matrix r_ℓ induces a linear functional φ_{r_ℓ} on $C(S^1)_{(n)}$ via the action

$$\varphi_{r_\ell}(f) = \hat{f}(-\ell),$$

for each $f \in C(S^1)_{(n)}$. The positivity condition (2.2) and Proposition 3.1 indicate that $\hat{T} : C(S^1)_{(n)} \rightarrow \mathcal{B}(\mathcal{H})$ is a pure completely positive linear map, where the action of \hat{T} on $C(S^1)_{(n)}$ is given by

$$\hat{T}(f) = \sum_{\ell=-n+1}^{n-1} \varphi_{r_\ell}(f) \tau_\ell = \sum_{\ell=-n+1}^{n-1} \hat{f}(-\ell) \tau_\ell.$$

The operator system $C(S^1)_{(n)}$ is an operator subsystem of the unital C^* -algebra $C(S^1)$, and so the pure completely positive linear map \hat{T} has an extension to a pure completely positive linear map on $C(S^1)$ [2, p. 180], which we denote again by \hat{T} .

The irreducible representations of the commutative C^* -algebra $C(S^1)$ are point evaluations. Because pure completely positive maps of C^* -algebras into $\mathcal{B}(\mathcal{H})$ have Stinespring dilations [14] in which the dilating homomorphisms are irreducible representations [2, Corollary 1.4.3], there is an element $\lambda \in S^1$ and a linear map $w : \mathcal{H} \rightarrow \mathbb{C}$ such that

$$\hat{T}(g) = w^* g(\lambda) w = g(\lambda) w^* w, \text{ for every } g \in C(S^1).$$

Because $w^* w$ is of rank-1, there exists a scalar $\alpha \geq 0$ and a rank-1 projection $Q \in \mathcal{B}(\mathcal{H})$ such that $w^* w = \alpha Q$. Hence,

$$\hat{T}(g) = \alpha g(\lambda) Q, \text{ for every } g \in C(S^1).$$

If $k \in \{-n+1, \dots, -1, 0, 1, \dots, n-1\}$, then the function $z \mapsto z^k$ is an element of $C(S^1)_{(n)}$ and

$$\hat{T}(z^k) = \alpha \lambda^k Q.$$

Thus,

$$\alpha \lambda^k Q = \hat{T}(z^k) = \sum_{\ell=-n+1}^{n-1} \varphi_{r_\ell}(z^k) \tau_\ell = \tau_{-k}.$$

As the equation above holds for every choice of k , we deduce $\tau_\ell = \lambda^{-\ell}(\alpha Q)$ for every ℓ , which implies that

$$T = \sum_{\ell=-n+1}^{n-1} r_\ell \otimes \tau_\ell = \sum_{\ell=-n+1}^{n-1} \lambda^{-\ell} r_\ell \otimes (\alpha Q) = T_n(\lambda^{-1}) \otimes (\alpha Q).$$

Hence, the pure element T has the form $T_n(\lambda^{-1}) \otimes (\alpha Q)$, for some $\lambda \in S^1$, $\alpha \geq 0$, and rank-1 projection Q .

Conversely, consider suppose that $T = T_n(\lambda^{-1}) \otimes (\alpha Q)$, for some $\lambda \in S^1$, $\alpha \geq 0$, and rank-1 projection $Q \in \mathcal{B}(\mathcal{H})$. Then $\hat{T}_n : C(S^1)^{(n)} \rightarrow \mathcal{B}(\mathcal{H})$ is given by $\hat{T}_n(f) = \alpha f(\lambda) Q$. Considered as a map on $C(S^1)$, \hat{T}_n has the form $\hat{T}_n(g) = w^* g(\lambda) w$, where $w : \mathcal{H} \rightarrow \mathbb{C}$ is a linear map such that $w^* w = \alpha Q$. As such a map is pure in the cone $\mathcal{CP}(C(S^1), \mathcal{B}(\mathcal{H}))$ [2, Corollary 1.4.3], it is also pure in the cone $\mathcal{CP}(C(S^1)_{(n)}, \mathcal{B}(\mathcal{H}))$. Hence, as the purity of \hat{T} implies the purity of T , this completes the proof. \square

Corollary 3.3 (Separation Theorem). *Every element of $(C(S^1)^{(n)} \otimes_{\min} \mathcal{M}_p(\mathbb{C}))_+$ is separable. That is, a matrix $T \in C(S^1)^{(n)} \otimes_{\min} \mathcal{M}_p(\mathbb{C})$ is positive if and only if there are $\lambda_1, \dots, \lambda_m \in S^1$ and positive $b_1, \dots, b_m \in \mathcal{M}_p(\mathbb{C})$ such that*

$$T = \sum_{j=1}^m T_n(\lambda_j) \otimes b_j.$$

Proof. As the operator system $C(S^1)^{(n)}$ and the Hilbert space $\mathcal{H} = \mathbb{C}^p$ have finite dimension, the convex cone $(C(S^1)^{(n)} \otimes_{\min} \mathcal{M}_p(\mathbb{C}))_+$ is closed. Hence, by [13, Theorem 18.5], each element T of $(C(S^1)^{(n)} \otimes_{\min} \mathcal{M}_p(\mathbb{C}))_+$ has the form

$$T = \sum_{j=1}^m T_j,$$

for some pure elements T_j of $(C(S^1)^{(n)} \otimes_{\min} \mathcal{M}_p(\mathbb{C}))_+$. Because each T_j is separable, by Theorem 3.2, so is T . \square

The following approximation result, which was proved in [6], is noted here because it is a direct consequence of the separation theorem.

Proposition 3.4 (Approximate Separation). *If \mathcal{A} is a unital nuclear C^* -algebra, and if $x \in (C(S^1)^{(n)} \otimes_{\min} \mathcal{A})_+$, then for every $\varepsilon > 0$ there exists a separable $y \in (C(S^1)^{(n)} \otimes_{\min} \mathcal{A})_+$ such that $\|x - y\| < \varepsilon$.*

4. TOEPLITZ ENTANGLEMENT

4.1. Entanglement in block-Toeplitz matrices with Toeplitz blocks. Theorem 1.1 asserts that every positive matrix in $(C(S^1)^{(n)} \otimes_{\min} \mathcal{M}_p(\mathbb{C}))_+$ is separable. It is also known (see [6]) that the replacement of $\mathcal{M}_p(\mathbb{C})$ by some other operator system \mathcal{S} could lead to entangled elements in $(C(S^1)^{(n)} \otimes_{\min} \mathcal{S})_+$. One such example occurs with $\mathcal{S} = C(S^1)_{(n)}$, the operator system dual of $C(S^1)^{(n)}$. However, it is also of interest to know whether $(C(S^1)^{(n)} \otimes_{\min} C(S^1)^{(p)})_+$ admits entangled elements. (This is the positive cone of the operator system of $n \times n$ Toeplitz matrices whose entries are $p \times p$ Toeplitz matrices, which is an operator system of interest in certain areas of applied mathematics.) We explain below why entangled matrices are present in the cone $(C(S^1)^{(n)} \otimes_{\min} C(S^1)^{(p)})_+$.

First, let us recall that the maximal operator system tensor product, \otimes_{\max} , on operator system \mathcal{R} and \mathcal{S} is the operator system structure on the algebraic tensor product $\mathcal{R} \otimes \mathcal{S}$ obtained by declaring a matrix $x \in \mathcal{M}_p(\mathcal{R} \otimes \mathcal{S})$ to be positive if, for each $\varepsilon > 0$, there are $n, p \in \mathbb{N}$, $a \in \mathcal{M}_n(\mathcal{R})_+$, $b \in \mathcal{M}_p(\mathcal{S})_+$, and a linear map $\delta : \mathbb{C}^q \rightarrow \mathbb{C}^n \otimes \mathbb{C}^p$ such that

$$\varepsilon(e_{\mathcal{R}} \otimes e_{\mathcal{S}}) + x = \delta^*(a \otimes b)\delta.$$

(Here, $e_{\mathcal{R}}$ and $e_{\mathcal{S}}$ denote the Archimedean order units of \mathcal{R} and \mathcal{S} .) As sets, the inclusion $(\mathcal{R} \otimes_{\max} \mathcal{S})_+ \subseteq (\mathcal{R} \otimes_{\min} \mathcal{S})_+$ always holds; however, it is typically more difficult for a matrix over $\mathcal{R} \otimes \mathcal{S}$ to be “max positive” than “min positive.”

Proposition 4.1. *There exist entangled elements in $(C(S^1)^{(n)} \otimes_{\min} C(S^1)^{(p)})_+$, for every $n, p \geq 2$.*

Proof. Suppose, on the contrary, that every $x \in (C(S^1)^{(n)} \otimes_{\min} C(S^1)^{(p)})_+$ is sep-

arable. Then $x = \sum_{j=1}^k a_j \otimes b_j$, for some positive Toeplitz matrices $a_j \in C(S^1)^{(n)}$, $b_j \in C(S^1)^{(p)}$. By definition of the maximal operator system tensor product, each $a_j \otimes b_j$ is an element of the positive cone $(C(S^1)^{(n)} \otimes_{\max} C(S^1)^{(p)})_+$. Hence, x also belongs to $(C(S^1)^{(n)} \otimes_{\max} C(S^1)^{(p)})_+$, implying that

$$\left(C(S^1)^{(n)} \otimes_{\min} C(S^1)^{(p)}\right)_+ \subseteq \left(C(S^1)^{(n)} \otimes_{\max} C(S^1)^{(p)}\right)_+.$$

Because the reverse inclusion always holds, we deduce that the operator systems $C(S^1)^{(n)} \otimes_{\min} C(S^1)^{(p)}$ and $C(S^1)^{(n)} \otimes_{\max} C(S^1)^{(p)}$ coincide. However, for $n, p \geq 2$, these operator systems are in fact distinct [6, Corollary 6.6]. Hence, the positive cone $(C(S^1)^{(n)} \otimes_{\min} C(S^1)^{(p)})_+$ must admit entangled elements. \square

4.2. Toeplitz duality admits entanglement. We now consider the situation whereby the operator system $C(S^1)^{(n)}$ is replaced by its dual $C(S^1)_{(n)}$ when tensoring with $\mathcal{B}(\mathcal{H})$.

The operator system $C(S^1)_{(n)}$ has a canonical linear basis consisting of functions $\chi_k : S^1 \rightarrow \mathbb{C}$ defined by $\chi_k(z) = z^k$, for $k = -n + 1, \dots, n - 1$, where χ_0 is the Archimedean order unit for $C(S^1)_{(n)}$. Thus, an arbitrary element $F \in$

$C(S^1)_{(n)} \otimes_{\min} \mathcal{S}$ has the form

$$(4.1) \quad F = \sum_{\ell=-n+1}^{n-1} \chi_\ell \otimes s_\ell,$$

for some $s_\ell \in \mathcal{S}$. Alternatively, the operator system $C(S^1)_{(n)} \otimes_{\min} \mathcal{S}$ coincides with the operator system of all continuous functions $F : S^1 \rightarrow \mathcal{S}$ of the form

$$(4.2) \quad F(z) = \sum_{\ell=-n+1}^{n-1} z^\ell s_\ell, \text{ for } z \in S^1.$$

The latter viewpoint is particularly attractive for $\mathcal{S} = \mathcal{M}_p(\mathbb{C})$, for in such cases F arises as a $p \times p$ matrix of trigonometric polynomials.

Although (4.1) and (4.2) give rise to equivalent forms of positivity, for the clarity of the proofs below, it is useful to maintain a formal distinction between the two. To this end, recall that the linear functional φ_{χ_ℓ} on $C(S^1)^{(n)}$ induced by $\chi_\ell \in C(S^1)_{(n)}$ has the property that

$$\varphi_{\chi_\ell}(r_k) = \left\{ \begin{array}{ll} 1 & : \text{ if } k = -\ell \\ 0 & : \text{ if } k \neq -\ell \end{array} \right\}.$$

Lemma 4.2. *If \mathcal{S} is any operator system, then an element $F \in (C(S^1)_{(n)} \otimes_{\min} \mathcal{S})_+$ is pure if and only if $\hat{F} : C(S^1)^{(n)} \rightarrow \mathcal{S}$ is a pure completely positive linear map.*

Proof. The proof is similar to the proof of Lemma 3.1, where the role of r_ℓ in Lemma 3.1 is played by χ_ℓ here. \square

The following result describes the structure of pure elements in the positive cone of $C(S^1)_{(n)} \otimes_{\min} \mathcal{B}(\mathcal{H})$.

Proposition 4.3. *Consider $F \in (C(S^1)_{(n)} \otimes_{\min} \mathcal{B}(\mathcal{H}))_+$ in the function form (4.2). If F is pure, then there exists an operator $w : \mathcal{H} \rightarrow \mathbb{C}^n$ such that*

$$F(z) = w^* T_n(z^{-1}) w,$$

for every $z \in S^1$.

Proof. Because $F = \sum_{\ell=-n+1}^{n-1} \chi_\ell \otimes a_\ell$ is positive and pure, the linear map $\hat{F} : C(S^1)^{(n)} \rightarrow \mathcal{S}$ given by $\hat{F} = \sum_{\ell=-n+1}^{n-1} \varphi_{\chi_\ell} a_\ell$ is completely positive. By the Arveson pure extension theorem [2, p. 180], \hat{F} extends to a pure completely positive linear map of $\mathcal{M}_n(\mathbb{C})$ into $\mathcal{B}(\mathcal{H})$, which we continue to denote by \hat{F} . Since all irreducible representations of the C^* -algebra $\mathcal{M}_n(\mathbb{C})$ are unitarily equivalent to the identity representation, the pure completely positive linear map \hat{F} necessarily has the form $\hat{F}(x) = w^* x w$, for all $x \in \mathcal{M}_n(\mathbb{C})$, for some linear operator $w : \mathcal{H} \rightarrow \mathbb{C}^n$ [2, Corollary 1.4.3]. Thus, for each $k \in \{-n+1, \dots, -1, 0, 1, \dots, n-1\}$,

$$\hat{F}(r_k) = w^* r_k w = \sum_{\ell=-n+1}^{n-1} \varphi_{\chi_\ell}(r_k) a_\ell = a_{-k}.$$

Hence, considering the function form (4.2) of F , we deduce that

$$F(z) = \sum_{\ell=-n+1}^{n-1} z^\ell (w^* r_{-\ell} w) = w^* \left(\sum_{\ell=-n+1}^{n-1} z^\ell r_{-\ell} \right) w = w^* T_n(z^{-1}) w,$$

for all $z \in S^1$. \square

Because of finite dimensionality, the cone $(C(S^1)_{(n)} \otimes \mathcal{M}_p(\mathbb{C}))_+$ is closed. Hence, we have the following consequence:

Corollary 4.4. *A matrix $F \in C(S^1)_{(n)} \otimes_{\min} \mathcal{M}_p(\mathbb{C})$ is positive if and only if there are linear maps $w_j : \mathbb{C}^p \rightarrow \mathbb{C}^n$, for $j = 1, \dots, m$, such that*

$$F(z) = \sum_{j=1}^m w_j^* T_n(z) w_j,$$

for $z \in S^1$.

Our main result in this section is the following one: the universal Toeplitz matrices are entangled.

Theorem 4.5. *For every $n \geq 2$, the universal Toeplitz matrix T_n is pure and entangled in $(C(S^1)_{(n)} \otimes_{\min} \mathcal{M}_n(\mathbb{C}))_+$.*

Proof. Suppose, on the contrary, that T_n is separable. Thus, using the form (4.2), there are nonzero $f_1, \dots, f_m \in (C(S^1)_{(n)})_+$ and nonzero $b_1, \dots, b_m \in \mathcal{M}_n(\mathbb{C})_+$ such that $T_n(z) = \sum_{j=1}^m f_j(z) b_j$, for all $z \in S^1$. Let

$$\mathcal{Z} = \{z_0 \in S^1 \mid f_j(z_0) = 0, \text{ for some } j = 1, \dots, m\}.$$

Because $g_j(z) = z^{n-1} f_j(z)$ is a complex polynomial of degree $2n-2$, there are, for each j , at most finitely many $z \in S^1$ for which $f_j(z) = 0$. Thus, the set \mathcal{Z} is finite.

Select $z \in S^1 \setminus \mathcal{Z}$. Because $T_n(z)$ is rank 1 and positive, there exist orthonormal vectors $\xi_1^z, \dots, \xi_{n-1}^z \in \mathbb{C}^n$ such that $T_n(z) \xi_k^z = 0$ for all $k = 1, \dots, n-1$. Thus,

$$0 = \langle T_n(z) \xi_k^z, \xi_k^z \rangle = \sum_{j=1}^m f_j(z) \langle b_j \xi_k^z, \xi_k^z \rangle.$$

Since $f_j(z) > 0$, for each j , it must be that $\langle b_j \xi_k^z, \xi_k^z \rangle = 0$ and, hence, $b_j \xi_j^z = 0$ for each k and j . Thus, $\ker T_n(z) \subseteq \ker b_j$ for all j , and so $\text{ran } T_n(z) \supseteq \text{ran } b_j$. Therefore, as $b_j \neq 0$ and $T_n(z)$ has rank 1, we have the equality $\text{ran } T_n(z) = \text{ran } b_j$ for each j , and such is true for all $z \in S^1 \setminus \mathcal{Z}$. Hence,

$$\text{ran } b_j = \text{ran } T_n(z) = \left\{ \delta \begin{bmatrix} 1 \\ z^2 \\ \vdots \\ z^{n-1} \end{bmatrix} \mid \delta \in \mathbb{C} \right\},$$

for all $z \in S^1 \setminus \mathcal{Z}$ and all j . Because $S^1 \setminus \mathcal{Z}$ is infinite, there exist $z \in S^1$ such that $z, -z \notin \mathcal{Z}$. With such as z ,

$$\begin{bmatrix} 1 \\ z \\ z^2 \\ z^3 \\ \vdots \\ z^{n-1} \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 1 \\ -z \\ z^2 \\ -z^3 \\ \vdots \\ (-z)^{n-1} \end{bmatrix}$$

are linearly independent and belong to the range of each b_j . Hence, $\dim(\text{ran } b_j) \geq 2$, in contradiction to $\dim(\text{ran } b_j) = 1$. Therefore, it cannot be that T_n is separable.

To show that T_n is pure, we need only show that the completely positive linear map $\hat{T}_n : C(S^1)^{(n)} \rightarrow \mathcal{M}_n(\mathbb{C})$ is pure (by Lemma 4.2). For each k , $\hat{T}_n(r_k) = r_{-k} = u^* r_k u$, where $u \in \mathcal{M}_n(\mathbb{C})$ is the unitary matrix that implements the transpose map on the Toeplitz operator system. Hence, \hat{T}_n is the restriction to $C(S^1)^{(n)}$ of the irreducible representation $\pi : \mathcal{M}_n(\mathbb{C}) \rightarrow \mathcal{M}_n(\mathbb{C})$ given by $\pi(x) = u^* x u$. Because $C(S^1)^{(n)}$ generates the C*-algebra $\mathcal{M}_n(\mathbb{C})$, and because π is a boundary representation for $C(S^1)^{(n)}$ [3, Theorem 2.1.1 and Remark 2 on p. 288], the restriction \hat{T}_n of π to $C(S^1)^{(n)}$ is necessarily pure [7, Proposition 2.12]. \square

5. TOEPLITZ COMPLETE POSITIVITY

The innovation introduced by Stinespring in his classic paper [14] was to focus on those positive linear maps $\psi : \mathcal{A} \rightarrow \mathcal{B}(\mathcal{K})$, where \mathcal{A} is a unital C*-algebra and \mathcal{K} is a Hilbert space, for which $\psi^{(n)} = \text{id}_{\mathcal{M}_n(\mathbb{C})} \otimes \psi$ is a positive linear map of $\mathcal{M}_n(\mathbb{C}) \otimes_{\min} \mathcal{A}$ into $\mathcal{M}_n(\mathbb{C}) \otimes_{\min} \mathcal{B}(\mathcal{K})$ for every $n \in \mathbb{N}$. With the analogy encapsulated by the definition below, the role of $\mathcal{M}_n(\mathbb{C})$ is taken by $C(S^1)^{(n)}$.

Definition 5.1. If \mathcal{R} is an operator system, then a linear map $\psi : \mathcal{R} \rightarrow \mathcal{B}(\mathcal{K})$, for some Hilbert space \mathcal{K} , is *Toeplitz completely positive* if, for every $n \in \mathbb{N}$ and matrix

$$\begin{bmatrix} \tau_0 & \tau_{-1} & \cdots & \tau_{-n+1} \\ \tau_1 & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \tau_{-1} \\ \tau_{n-1} & \cdots & \tau_1 & \tau_0 \end{bmatrix} \in \mathcal{M}_n(\mathcal{R})_+,$$

the matrix

$$\begin{bmatrix} \psi(\tau_0) & \psi(\tau_{-1}) & \cdots & \psi(\tau_{-n+1}) \\ \psi(\tau_1) & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \psi(\tau_{-1}) \\ \psi(\tau_{n-1}) & \cdots & \psi(\tau_1) & \psi(\tau_0) \end{bmatrix}$$

is a positive operator on $\bigoplus_1^n \mathcal{K}$.

Theorem 1.1 was used in [6] to establish the following result.

Theorem 5.2. *Every positive linear map on a unital nuclear C*-algebra is Toeplitz completely positive.*

Our aim here is to prove the following similar result.

Theorem 5.3. *If \mathcal{H} is a Hilbert space of infinite dimension, then every normal positive linear map on $\mathcal{B}(\mathcal{H})$ is Toeplitz completely positive.*

Proof. For each $n \in \mathbb{N}$, consider $C(S^1)^{(n)} \otimes_{\min} \mathcal{B}(\mathcal{H})$ in the topology inherited from the ultraweak topology of the von Neumann algebra $\mathcal{M}_n(\mathbb{C}) \otimes \mathcal{B}(\mathcal{H})$. Thus, if $\psi : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{K})$ is a normal positive linear map, then $\psi^{(n)} = \text{id}_{\mathcal{M}_n(\mathbb{C})} \otimes \psi$ is continuous with respect to the ultraweak topology of $\mathcal{M}_n(\mathbb{C}) \otimes \mathcal{B}(\mathcal{H})$.

By Theorem 1.1, the extremal rays of $(C(S^1)^{(n)} \otimes_{\min} \mathcal{B}(\mathcal{H}))_+$ are generated by elements of the form $T_n(\lambda^{-1}) \otimes (\alpha Q)$, for some $\lambda \in S^1$, $\alpha \geq 0$, and rank-1 projection Q . As the set of all finite sums of elements of the form αQ coincides with all finite-rank positive linear operators on $\mathcal{B}(\mathcal{H})$, and because this set is ultraweakly dense in $\mathcal{B}(\mathcal{H})_+$, the set

$$\left\{ \sum_{j=1}^m T_n(\lambda_j^{-1}) \otimes (\alpha_j Q_j) \mid m \in \mathbb{N}, \lambda_j \in S^1, \alpha_j \geq 0, Q_j \text{ is a rank-1 projection} \right\},$$

is ultraweakly dense in $(C(S^1)^{(n)} \otimes_{\min} \mathcal{B}(\mathcal{H}))_+$. Furthermore, the action of $\psi^{(n)}$ on a pure element $T_n(\lambda^{-1}) \otimes (\alpha Q)$ is given by

$$\psi^{(n)} [T_n(\lambda^{-1}) \otimes (\alpha Q)] = T_n(\lambda^{-1}) \otimes \alpha \psi(Q),$$

which is positive in $\mathcal{M}_n(\mathbb{C}) \otimes \mathcal{B}(\mathcal{K})$. Hence, by ultraweak continuity, $\psi^{(n)}$ is positive on $C(S^1)^{(n)} \otimes_{\min} \mathcal{B}(\mathcal{H})$. \square

6. APPENDIX: FACTORISATION AND UNIVERSALITY ARE EQUIVALENT

Our purpose in this paper has been to underscore the role and use of Toeplitz duality in results pertaining to separability and entanglement. Nevertheless, it useful to make note of the relationship between Ando's universality theorem [1] and Gurvits' separation theorem [8, 9], the latter depending on a certain factorisation of Toeplitz matrices that Gurvits established through some elegant algebraic arguments. How these two results intersect is described by Theorem 6.1 below.

Before turning to Theorem 6.1, let us first observe that the universal Toeplitz matrix valued function $T_n : S^1 \rightarrow \mathcal{M}_n(\mathbb{C})$ can be evaluated, via continuous functional calculus, at any unitary operator u , thereby producing a positive $n \times n$ Toeplitz matrix $T_n(u)$ whose entries are integral powers u^ℓ of u .

Theorem 6.1 (Factorisation Theorem). *If $T = [\tau_{k-j}]_{k,j=1}^n \in C(S^1)^{(n)} \otimes_{\min} \mathcal{B}(\mathcal{H})$ is a positive Toeplitz matrix, then there are a Hilbert space \mathcal{K} , a unitary operator $u \in \mathcal{B}(\mathcal{K})$, and a bounded linear operator $w : \mathcal{K} \rightarrow \mathcal{H}$ such that*

$$(6.1) \quad T = (1_n \otimes w)^* T_n(u) (1_n \otimes w),$$

where 1_n denotes the $n \times n$ identity matrix. If, moreover, the unitary operator u has finite spectrum, then T is separable in the positive cone of $C(S^1)^{(n)} \otimes_{\min} \mathcal{B}(\mathcal{H})$.

Proof. By Ando's theorem [1], the positivity of T implies that there is a completely positive linear map $\phi : C(S^1) \rightarrow \mathcal{B}(\mathcal{H})$ such that $\phi(\chi_\ell) = \tau_\ell$, for each $\ell = -n + 1, \dots, n-1$, and where $\chi_\ell(z) = z^\ell$. By Stinespring's theorem [14], there are a Hilbert space \mathcal{K} , a unital $*$ -representation $\pi : C(S^1) \rightarrow \mathcal{B}(\mathcal{K})$ of $C(S^1)$, and a bounded linear operator $w : \mathcal{K} \rightarrow \mathcal{H}$ such that $\phi(f) = w^* \pi(f) w$, for every $f \in C(S^1)$. Let $u \in \mathcal{B}(\mathcal{K})$ be the unitary operator given by $u = \pi(\chi_1)$. Thus,

$$\tau_\ell = \phi(\chi_\ell) = w^* \pi(\chi_\ell) w = w^* \phi(\chi_1)^\ell w^* = w^* u^\ell w,$$

for each ℓ , and so

$$\begin{aligned} T &= \sum_{\ell=-n+1}^{n-1} r_\ell \otimes w^* u^\ell w \\ &= (1_n \otimes w)^* \left(\sum_{\ell=-n+1}^{n-1} r_\ell \otimes u^\ell \right) (1_n \otimes w) \\ &= (1_n \otimes w)^* T_n(u) (1_n \otimes w), \end{aligned}$$

which is precisely the sought-for form (6.1).

Suppose now that $u \in \mathcal{B}(\mathcal{K})$ has finite spectrum. By the spectral theorem, there are $\lambda_1, \dots, \lambda_m \in S^1$ and pairwise-orthogonal projections $q_1, \dots, q_m \in \mathcal{B}(\mathcal{K})$ such that $u^\ell = \sum_{j=1}^m \lambda_j^\ell q_j$, for all $\ell \in \mathbb{Z}$. Therefore,

$$\begin{aligned} T_n(u) &= \sum_{\ell=-n+1}^{n-1} r_\ell \otimes \left(\sum_j \lambda_j^\ell q_j \right) \\ &= \sum_{j=1}^m \sum_{\ell=-n+1}^{n-1} \lambda_j^\ell r_\ell \otimes q_j \\ &= \sum_{j=1}^m T_n(\lambda_j^{-1}) \otimes q_j. \end{aligned}$$

Hence,

$$T = (1_n \otimes w)^* T_n(u) (1_n \otimes w) = \sum_{j=1}^m T_n(\lambda_j^{-1}) \otimes w^* q_j w,$$

which is a separable element of the positive cone of $C(S^1)^{(n)} \otimes_{\min} \mathcal{B}(\mathcal{H})$. \square

The proof of Theorem 6.1 shows that the factorisation given in (6.1) of a positive Toeplitz matrix T is a consequence of Ando's universality theorem. Conversely, if $T = [\tau_{k-j}]_{i,j=1}^n$ is a Toeplitz matrix of operators on \mathcal{H} whose (k, j) -entries are given by $\tau_{k-j} = w^* u^{k-j} w$, for some unitary $u \in \mathcal{B}(\mathcal{K})$ and operator $w : \mathcal{K} \rightarrow \mathcal{H}$, then T is positive and there is a completely positive linear map $\phi : C(S^1)_{(n)} \rightarrow \mathcal{B}(\mathcal{H})$ such that $\phi(\chi_\ell) = \tau_\ell$, for each $\ell = -n+1, \dots, n-1$. The completely positive linear map ϕ in question arises from the fact that $C(S^1)$ is the universal C^* -algebra generated by a unitary operator, and so there is a unital $*$ -homomorphism $\varrho : C(S^1) \rightarrow \mathcal{B}(\mathcal{K})$ such that $\varrho(z^\ell) = u^\ell$, for every $\ell \in \mathbb{Z}$; thus, ϕ is given by $\phi(z^\ell) = w^* \varrho(z^\ell) w$, for $\ell = -n+1, \dots, n-1$. Hence, Ando's universality theorem is a consequence of the factorisation theorem (Theorem 6.1).

Separation depends on the finiteness of spectrum. The approach of Gurvits in [8, 9] to his separation theorem (Theorem 1.1) is to show that a positive Toeplitz matrix $T \in C(S^1)^{(n)} \otimes_{\min} \mathcal{M}_p(\mathbb{C})$ admits a factorisation of the form $(1_n \otimes w)^* T_n(u) (1_n \otimes w)$ in which the unitary u has finite spectrum.

REFERENCES

- [1] T. Andô, *Truncated moment problems for operators*, Acta Sci. Math. (Szeged) **31** (1970), 319–334. MR 290157
- [2] William Arveson, *Subalgebras of C^* -algebras*, Acta Math. **123** (1969), 141–224. MR MR0253059 (40 #6274)

- [3] ———, *Subalgebras of C^* -algebras. II*, Acta Math. **128** (1972), no. 3-4, 271–308. MR MR0394232 (52 #15035)
- [4] Man Duen Choi and Edward G. Effros, *Injectivity and operator spaces*, J. Functional Analysis **24** (1977), no. 2, 156–209. MR 0430809 (55 #3814)
- [5] Alain Connes and Walter D. van Suijlekom, *Spectral truncations in noncommutative geometry and operator systems*, Comm. Math. Phys. **383** (2021), no. 3, 2021–2067. MR 4244265
- [6] Douglas Farenick, *The operator system of Toeplitz matrices*, Trans. Amer. Math. Soc. Ser. B **8** (2021), 999–1023. MR 4340832
- [7] Douglas Farenick and Ryan Tessier, *Purity of the embeddings of operator systems into their C^* - and injective envelopes*, Pacific J. Math. **317** (2022), no. 2, 317–338.
- [8] Leonid Gurvits, LANL Unclassified Technical Report (2001), LAUR-01-2030.
- [9] Leonid Gurvits and Howard Burnam, *Largest separable balls around the maximally mixed bipartite quantum states*, Phys. Rev. A **66** (2002), 062311.
- [10] E. Hekkelman, *Truncated geometry on the circle*, Lett. Math. Phys. **112** (2022), no. 2, Paper No. 20, 19. MR 4392480
- [11] Ali Kavruk, Vern I. Paulsen, Ivan G. Todorov, and Mark Tomforde, *Tensor products of operator systems*, J. Funct. Anal. **261** (2011), no. 2, 267–299. MR 2793115
- [12] Ali S. Kavruk, Vern I. Paulsen, Ivan G. Todorov, and Mark Tomforde, *Quotients, exactness, and nuclearity in the operator system category*, Adv. Math. **235** (2013), 321–360. MR 3010061
- [13] R. Tyrrell Rockafellar, *Convex analysis*, Princeton Mathematical Series, No. 28, Princeton University Press, Princeton, N.J., 1970. MR 0274683
- [14] W. Forrest Stinespring, *Positive functions on C^* -algebras*, Proc. Amer. Math. Soc. **6** (1955), 211–216. MR 69403
- [15] Walter D. van Suijlekom, *Gromov-Hausdorff convergence of state spaces for spectral truncations*, J. Geom. Phys. **162** (2021), Paper No. 104075, 11. MR 4199398

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF REGINA, REGINA, SASKATCHEWAN S4S 0A2, CANADA

Email address: douglas.farenick@uregina.ca

Email address: mmm034@uregina.ca