

Expanded-clique graphs and the domination problem*

Mitre C. Dourado^{†1}, Rodolfo A. Oliveira^{‡2}, Vitor Ponciano^{§1},
Alessandra B. Queiróz^{¶1}, and Rômulo L. O. Silva^{||1}

¹Instituto de Computação, Universidade Federal do Rio de Janeiro,
Rio de Janeiro, Brazil

²Instituto do Noroeste Fluminense, Universidade Federal Fluminense, Brazil

August 9, 2022

Abstract

Given a graph G and a function $f : V(G) \rightarrow \mathbb{N}$ such that $f(v_i) \geq d(v_i)$ for every $v_i \in V(G)$, where $d(v_i)$ is the degree of v_i , the expanded-clique graph H of (G, f) exists and is defined as follows: for every $v_i \in V(G)$, there is a set $V_i \subseteq V(H)$ with $f(v_i)$ vertices forming a clique, and $d_G(v_i)$ vertices of V_i have a neighbor outside V_i , one in each V_j such that $v_j \in N(v_i)$. In this work, we present two characterizations of the expanded-clique graphs. One of them leads to a linear-time algorithm of recognition. We show that given an expanded-graph H , the domination number of H plus the 2-independence number of the root G of H is equal to $|V(G)|$. We also show that the domination problem is NP-complete for some subclasses of the expanded-clique graphs.

1 Introduction

We consider finite, simple and undirected graphs. The degree, the open and the closed neighborhoods of a vertex v are denoted by $d(v)$, $N(v)$ and $N[v]$, respectively. In this text, whenever we refer a graph by G , we will denote its vertex by $\{v_1, \dots, v_n\}$. Given a function $f : V(G) \rightarrow \mathbb{N}$ such that $f(v_i) \geq d(v_i)$ for every $v_i \in V(G)$, the *expanded-clique graph* H of (G, f) exists and is defined as follows: for every $v_i \in V(G)$, there is a set $V_i \subseteq V(H)$ with $f(v_i)$ vertices forming a clique. For every $v_j \in N(v_i)$, there is a vertex $v_{i,j} \in V_i$ and the edge $v_{i,j}v_{j,i}$ in $E(H)$. If $f(v_i) - d(v_i) = k_i > 0$, then V_i has k_i simplicial vertices, which are denoted by $v'_{i,1}, \dots, v'_{i,k_i}$. In

*Partially supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico, Brazil, Grant numbers 305404/2020-2, FAPERJ ...

[†]mitre@ic.ufrj.br

[‡]rodolfooliveira@id.uff.br

[§]vtponciano@ufrj.br

[¶]alessandrabarbosa@ufrj.br

^{||}romulolo@ufpa.br

this case, G is the *root of H* under the f -clique-expansion operation. The set V_i will be referred to as the *expanded clique (associated with v_i)*. Note that for every $v \in V_i$, $d(v) \in \{|V_i| - 1, |V_i|\}$. We say that H is a *expanded-clique graph* when H is the expanded-clique graph for some graph G and function f . If $f(v_i) = k$ for some $k \in \mathbb{N}$, we can say that H is a *k -expanded-clique graph*.

In this work, we are interested in the complexity aspects of the recognition problem of the expanded-clique graphs and of the domination problem on this class. We remark that the well-studied inflated graphs [4, 8, 10] form a subclass of the clique-expanded graphs, since a graph H is *inflated* if it is the expanded-graph of a pair (G, f) satisfying $f(v_i) = d(v_i)$.

Recall that $D \subseteq V(G)$ is a *dominating set* of a graph G if every vertex of $V(G) \setminus D$ has a neighbor in D . The domination problem is a classic problem in graphs having many relevant applications, such as in communication and computer networks [20] (especially playing an important role as the backbone of the wireless design [3, 16]), influence analysis in social medias [1], and topics regarding Internet of Robotic Things [17]. It can be stated as:

DOMINATING SET

Input: A graph G and a positive integer ℓ .

Question: Is there a dominating set $S \subseteq V(G)$ so that $|S| \leq \ell$?

Since this problem and some variations are NP-complete [2, 18], techniques based on heuristic methods [14], approximation algorithms [15], and FPT-algorithms [9] have been used to deal somehow with it.

The text is organized as follows. In Section 2, we show that the subdivided-line graph [7] and the Sierpiński graphs [12] are proper subclasses of expanded-clique graphs. We also show that the expanded-clique graphs form a proper subclass of the line graph of bipartite graphs [6]. we present two characterizations of the expanded-clique graphs. One of them leads to a linear-time algorithm of recognition. Section 3 deals with the domination problem. We show that this problem is NP-complete for expanded-clique graphs restricted to planar line graphs of bipartite graphs with a maximum degree 3 and for cubic line graphs of bipartite graphs. For $\Delta(G) \leq k$, we show that this problem is NP-complete for any fixed $k \geq 3$. We also show that given an expanded-graph H , the domination number of H plus the 2-independence number of the root G of H is equal to $|V(G)|$. As a consequence of this result, we derive lower and upper bounds for the dominating number of expanded-clique graphs, which lead to the fact that a dominating set of H can be easily found within the ratio $(1 + \frac{1}{\Delta}(G))$ of the minimum.

We conclude this section by presenting useful notation. Let G be a graph. The minimum and maximum degrees of G are denoted by $\delta(G)$ and $\Delta(G)$, respectively. We say that G is a *cubic graph* if G is 3-regular. We write K_n for the complete graph with n vertices. A graph having at least one plane representation without crossing edges is called *planar graph*. A subgraph of G *induced* by V' is denoted by $G[V']$. A vertex $v \in V$ is a *simplicial vertex* if $N_G(v)$ induces a clique. For the case of $V' \subseteq V$, denote the *closed neighborhood of V' in G* as $N_G[V'] = \{v \in N_G[v'] : \text{for all } v' \in V'\}$, and *open neighborhood of V' in G* as $N_G(V') = N_G[V'] \setminus V'$. We write $K_{p,q}$ for the complete bipartite graph, where stable sets

have respectively the size p and q , and all possible edges among them is in the graph. A *claw* is a $K_{1,3}$ graph. A *diamond* is an induced cycle C_4 with one diagonal. An *odd-hole* is any induced cycle C_q where q is an odd number greater than 4. A *butterfly graph*, also called hourglass graph, is a graph with 5 vertices and 6 edges. It can be constructed by joining 2 copies of the C_3 with a common vertex.

2 Characterization and recognition

The *subdivision* of a graph F , $S(F)$, is the replacement of every edge $uv \in E(F)$ for a new vertex x_{uv} and edges $x_{uv}u$ and $x_{uv}v$. The *line graph* of F , written $L(F)$, is the graph whose vertex set is $E(F)$ and in which two distinct vertices uv and xy are adjacent if and only if they are adjacent in F , i.e., $\{u, v\} \cap \{x, y\} \neq \emptyset$. We say that $L(S(F))$ is a *subdivided-line graph* [7]. See an example in Figure 1. The subdivided-line graphs are also known in the literature as *inflated graphs* [8].




Figure 1: An example of a line graph of subdivision.

The Sierpiński graphs were introduced by Klavžar and Milutinović as a generalization of the graph of the Tower of Hanoi problem [11]. Given integers $p \geq 1$ and $q \geq 1$, the *Sierpiński graph* $S(p, q)$ has a vertex for each p -tuple that can be formed from $\{1, \dots, q\}$ and, for two distinct vertices $u = (u_1, u_2, \dots, u_p)$ and $w = (w_1, w_2, \dots, w_p)$, $uw \in E(S(p, q))$ if and only if there exists an $h \in \{1, \dots, p\}$ such that

1. $u_t = w_t$ for $t \in \{1, \dots, h - 1\}$;
2. $u_h \neq w_h$;
3. $u_t = w_h$ and $w_t = u_h$ for $t \in \{h + 1, p\}$.

In Figure 2, we show examples of $S(1, 3)$, $S(2, 3)$, and $S(3, 3)$.

Observe that a subdivided-line graph has no simplicial vertices, whereas the number of simplicial vertices of a Sierpiński graph is q , which means that they are disjoint graph classes. In the next result, we show that both are proper subclasses of the expanded-clique graphs.

Proposition 2.1. *If H is a subdivided-line graph or a Sierpiński graph, then H is expanded-clique.*

Proof. For both cases, we need to find a root G and its corresponding f . Let H be a subdivided-line graph. Since $H = L(S(G))$, we can take G as the root and $f(v_i) = d(v_i)$ for every $v_i \in V(G)$.




Figure 2: Examples of Sierpiński graphs.

Now, let H be a Sierpiński graph $S(p, q)$. For $p = 1$, we have $V(G) = \{v_1\}$ and $f(v_1) = q$. For $p > 1$, let $G = S(p-1, q)$ and $f(v_i) = q$ for every $v_i \in V(G)$. The expanded-clique graph of (G, f) has $q|V(G)|$ vertices and this is the number of vertices of $S(p, q)$ because for any $(p-1)$ -tuple that can be formed with q elements, we can form q tuples by adding one coordinate. The definition of Sierpiński graph and the definition of expanded-clique graphs imply that these q vertices form a clique. Denote such clique by V_i . Let $v_i \in V(S(p-1, q))$. We can write $v_i = (u_1, \dots, u_{p-1})$. Note that $d(v_i) \in \{q-1, q\}$. We know that every vertex of $S(p, q)$ and every vertex of a q -expanded-clique graph has also degree $q-1$ or q . If $d(v_i) = q$, then since $p \geq 2$, the definition implies that there are two coordinates of v_i that are different. Then, for every $v_j \in N(v_i)$, there is a vertex in V_i and a vertex in V_j that are adjacent in $S(p, q)$. If $d(v_i) = q-1$, then all coordinates of v_i are equal. Then for every $v_j \in N(v_i)$, there is a vertex in V_i and a vertex in V_j that are adjacent in $S(p, q)$. Since these edges coincide with the edges of the definition of expanded-clique graphs, we conclude that $H = S(p, q)$ is an expanded-clique graph. \square

Before showing that the line graphs of bipartite graphs form a superclass of the expanded-clique graphs, we recall an useful result.

Proposition 2.2. [6] *A graph is a line graph of bipartite graph if and only if it is (claw, diamond, odd-hole)-free.*

Theorem 2.3. *If H is expanded-clique, then H is a line graph of bipartite graph.*

proof. Let H be an expanded-clique graph. By Proposition 2.2, it suffices to show that H is (claw, diamond, odd-hole)-free. By the definition of expanded-clique graph, for every $v \in V(H)$, either $N_H(v)$ is a clique or there is $u \in N_H(v)$ such that $N_H(v) \setminus \{u\}$ is a clique and u has no neighbors in $N_H(v)$. Since a claw and a diamond have vertices not satisfying this property, we conclude that H is (claw, diamond)-free. Now, let $C_q = u_0 u_1 \dots u_q$ be a cycle of H where $u_0 = u_q$ and $q \geq 3$, and let G be a root of H . The definition of expanded-clique graph implies that for every $i \in \{1, \dots, q-1\}$, either $u_{i-1}, u_i \in V_j$ or $u_i, u_{i+1} \in V_j$, but there are no three vertices of C_q contained in a same V_j . Therefore, q is even, which means that H is odd-hole-free. \square

For $k \geq 3$, a sequence of vertices u_1, \dots, u_k of a graph F is a *chain* if $u_i u_{i+1} \in E(F)$ for $i \in \{1, \dots, k-1\}$, $d(u_i) = 2$ for $i \in \{2, \dots, k-1\}$ and is maximal with these properties. We

say that a chain is *bad* if k is odd, $d(u_1) \geq 3$ and $d(u_k) \geq 3$. If a chain is not bad, then it is *good*. We say that a vertex $v \in V(F)$ is *1-simplicial* if there is $u \in N(v)$ such that $N(v) \setminus \{u\}$ is a clique and u has no neighbors in $N(v)$.

Theorem 2.4. *A connected graph H that is not a cycle is expanded-clique if and only if every chain of H is good and every vertex of $V(H)$ is either simplicial or 1-simplicial.*

Proof. We begin by considering the case where H is a path. Since, in this case, every vertex of H is simplicial or 1-simplicial and H does not have bad chains, we have to show a pair (G, f) such that G is the root of H under the f -expanded-clique operation. It is easy to see that for $k \geq 2$, $P_{\frac{k}{2}}$ is a root for P_k if k is even, while $P_{\lceil \frac{k}{2} \rceil}$ is a root for P_k if k is odd. From now on, we can assume that H is neither a cycle nor a path.

For the necessity, consider that H is the expanded-clique of a pair (G, f) . First, suppose by contradiction that u_1, \dots, u_k is a bad chain of H . Since H is neither a cycle nor a path, we can assume that the neighbor w of u_1 has degree at least 3. Therefore, w and u_1 belong to different expanded cliques of H . Furthermore, the expanded clique containing u_1 also contains u_2 and no more vertices. Then, $\{v_i, v_{i+1}\}$ is an expanded clique for every $i < k$ such that i is odd. If k is odd, then $d(v_k) = 2$ and v_k does not belong to any expanded clique, which is not possible. Therefore, any chain of H is good.

Now, suppose by contradiction that there is $v \in V(H)$ such that $N(v)$ is not simplicial neither 1-simplicial. Note that $d(v) \geq 3$. Then, there are $u, w \in N(v)$ such that $uw \notin E(H)$ and let $x \in N_H(v) \setminus \{u, w\}$. If $u, w \in N_H(x)$, then H has a diamond, which is not possible by Proposition 2.2 and Theorem 2.3. Then, without loss of generality, we can assume that $ux \notin E(H)$. If $xw \notin E(H)$, then H has a claw, which is also not possible. Then, the subgraph of H induced by $N(v)$ has exactly two connected components C_1 and C_2 each one being a complete graph such that $|V(C_i)| \geq 2$ for $i \in \{1, 2\}$. Without loss of generality, we can assume that the expanded clique containing v is contained in $V(C_1)$. Now, we reach a contradiction because v has at least 2 neighbors outside C_1 while the definition of expanded-clique graph guarantees that each vertex has at most one neighbor outside the expanded clique containing it.

For the sufficiency, consider that every chain of H is good and for every $v \in V(H)$, either v is a simplicial or an 1-simplicial vertex. We will construct a graph G and a function f such that H is the expanded-clique graph of (G, f) . In order to do this, we define $S(v)$ for every $v \in V(H)$ as follows.

First, consider the vertices v with degree at least 3 in any order. If v is a simplicial vertex, then set $S(v) = N(v)$; otherwise, let $u \in N(v)$ such that $N(v) \setminus \{u\}$ is a clique and u has no neighbors in $N(v)$ and set $S(v) = N(v) \setminus \{u\}$. For the vertices v with degree 2, we choose first the ones having a neighbor w such that $S(w)$ has already been defined. Denote by u the neighbor of v different of w . Then, define $S(v) = \{v, u\}$ and $S(u) = \{v, u\}$. Finally, for the pendant vertices v , define $S(v) = \{v\}$. The assumptions on H guarantees that $S(v)$ will be defined for every vertex $v \in V(H)$.

Now, consider any ordering v_1, \dots, v_p of $V(H)$. Then, for i from 1 to p , add to G a vertex

v'_i if $S(v_i) \neq S(v_j)$ for every $j \in [i - 1]$ and set $f(v'_i) = |S(v_i)|$. Finally, add to G the edge $v'_i v'_j$ if there is an edge joining some vertex of $S(v_i)$ to some vertex of $S(v_j)$. Noting that $S(v_i)$ is a clique for every $v_i \in V(H)$ and that every vertex of $S(v_i)$ has at most one neighbor outside $S(v_i)$, we conclude that H is the expanded-clique graph of (G, f) . \square

Proposition 2.5. *If H is an expanded-clique graph, then H is C_4 -free.*

Proof. Let H be the expanded-clique graph of a pair (G, f) . Suppose by contradiction that H contains an induced C_4 formed by the vertices u_1, u_2, u_3 and u_4 and the edges $u_1 u_2, u_2 u_3, u_3 u_4$ and $u_4 u_1$. By the definition, the expanded cliques of H form a partition of $V(H)$. Consider the expanded clique K_1 containing u_1 . We also know that u_1 has at most one neighbor outside K_1 . Without loss of generality, we can assume that $u_4 \in K_1$ and that u_2 belongs to an expanded clique K_2 different of K_1 . The same reasoning implies that $u_3 \in K_2$. With these facts, we conclude that there are two edges in G joining the vertices $v_i, v_j \in V(G)$ associated with K_1 and K_2 , which contradicts the assumption that G is a simple graph. \square

The proof of Theorem 2.4 implies that one can recognize whether a graph is expanded-clique in linear time, since the cycles are covered by Propositions 2.5 and 2.2 and Theorem 2.3.

Corollary 2.6. *One can recognize an expanded-clique graph in $O(n + m)$ steps.*

Next, we present another characterization of expanded-clique graphs.

Corollary 2.7. *A graph is expanded-clique if and only if it is (bad chain, butterfly, claw, C_4 , diamond, odd-hole)-free.*

Proof. Let H be an expanded-clique graph. Due to Theorem 2.3, we know that H is a line graph of a bipartite graph. Then, Proposition 2.2 implies that H is (claw, diamond, odd-hole)-free. By Theorem 2.4, H is bad chain free. Since a butterfly has a vertex that is neither simplicial nor 1-simplicial, Theorem 2.4 also implies that H is butterfly free. Since Proposition 2.5 guarantees that H is C_4 free, H is (bad chain, butterfly, claw, C_4 , diamond, odd-hole)-free.

Conversely, let H be a graph that is (bad chain, butterfly, claw, C_4 , diamond, odd-hole)-free. If H is a cycle C_k , then k is even greater than 4. It is clear that $C_{2k'}$ for $k' \geq 3$ is an expanded-clique graph. Then, consider that H is not a cycle. Assume by contradiction that H is not an expanded-clique graph. As H does not contain no bad chains, by Theorem 2.4, H contains some vertex v that is neither simplicial nor 1-simplicial. Thus, we can choose vertices u_1, u_2, u_3 in $N(v)$ such that $u_1 u_2 \notin E(H)$. If $\{u_1, u_2, u_3\}$ is an independent set, we have a claw and it is a contradiction. If there is only one edge among the vertices of $\{u_1, u_2, u_3\}$, then v has a fourth neighbor that may create a claw, a diamond or a butterfly, which is also not possible. If there edges $u_1 u_3$ and $u_2 u_3$ exist, we have a contradiction since they yield a diamond. \square

3 The domination problem

In this section, we consider the DOMINATING SET problem on expanded-clique graphs. Given a graph G , denote by $\gamma(G)$ the size of a minimum dominating set of G . The edge version of the

domination problem can be stated as follows.

EDGE DOMINATING SET

Input: A graph G and a positive integer ℓ .

Question: Is there a set $E' \subseteq E(G)$ so that $|E'| \leq \ell$ and every edge of $E(G) \setminus E'$ is adjacent to some edge of E' ?

It is known that the EDGE DOMINATING SET problem is NP-complete for bipartite graphs with maximum degree 3 [19], which means that DOMINATING SET is NP-complete for graphs with maximum degree 4 that are line graphs of bipartite graphs. It is also known that the DOMINATING SET problem is NP-complete for planar graphs with maximum degree 3 [5]. Up to our best knowledge, for no proper subclass of these two classes, the DOMINATING SET problem is known to be NP-complete. Due to Theorem 2.3, the next result presents a proper subclass of the intersection of these two classes for which the DOMINATING SET problem is NP-complete.

Theorem 3.1. *DOMINATING SET is NP-complete for 3-expanded-clique graphs even for planar graphs with maximum degree 3.*

proof. Since DOMINATING SET belongs to NP for general graphs, that condition holds for our particular case. For the reduction, we use the DOMINATING SET problem restricted to planar graphs with maximum degree 3 since it is NP-complete [5]. Let $\langle G, \ell \rangle$ be an instance of DOMINATING SET where G is planar with maximum degree 3 and let G' be the 3-expanded-clique of G . We construct an instance $\langle H, \ell' \rangle$ of DOMINATING SET where H is a 3-expanded-clique graph of G . Set $\ell' = 2(|V(G)| - \ell) + 3\ell = 2|V(G)| + \ell$. See Figure 3 for an example.

Note that H is also planar with a maximum degree 3. For $u \in V(G)$, denote the expanded-clique of G' associated with u by $\{u_1, u_2, u_3\}$, and for $i \in [3]$ and $u_i \in V(G')$, denote the expanded-clique of H associated with u_i by $\{u_{i,1}, u_{i,2}, u_{i,3}\}$. For $u \in V(G)$, denote by H_u the subgraph of H induced by $\{u_{i,j} : i \in [3] \text{ and } j \in [3]\}$. Figure 4.

Claim 1. *For every $u \in V(G)$, it holds $\gamma(H_u) = 3$.*

Proof of Claim 1. We know that $\gamma(H_u) \geq 3$ because $\Delta(H_u) = 3$ and $|V(H_u)| = 9$. On the other hand, note that one vertex of each K_3 is enough for a dominating set, and, therefore, $\gamma(H_u) = 3$. See Figure 4-(ii). ■

Claim 2. *If D_H is a dominating set of H , then $|V(H_u) \cap D_H| \geq 2$. Furthermore, if $|V(H_u) \cap D_H| = 2$, then $V(H_u) \setminus N_H[D_H]$ has only one vertex which is dominated by some vertex in H_v whose $|V(H_v) \cap D_H| = 3$.*

Proof of Claim 2. Note that the vertices of H_u with degree 3 do not have neighbors in H which are outside H_u . Since there is no vertices adjacent to all such vertices, we conclude that $|V(H_u) \cap D_H| \geq 2$. Now, consider that $V(H_u) \cap D_H = \{x, y\}$. Observe that if x would be a vertex of H_u with degree 2, then D_H would not be a dominating set of H . Therefore, we can assume that x, y are the vertices depicted in Figure 4-(iii) and $V(H_u) \setminus N_H[D_H]$ has only one




Figure 3: Graph resulting from polynomial transformation.




Figure 4: Vertex domination in H_u .

vertex which is dominated by some vertex in H_v . It remains to show that $|V(H_v) \cap D_H| = 3$. But this is consequence of the fact that the 3 vertices of H_v having neighbors in H outside H_v form a dominating set of H_v . ■

Now, we shall prove that G has a dominating set with at most ℓ vertices if and only if H has a dominating set with at most $2|V(G)| + \ell$ vertices.

(\Rightarrow) Consider that D is a dominating set of G with $|D| \leq \ell$. Starting with D_H empty, for each $v \in D$, add to D_H the vertices of H_v with degree 2. Now, for each $u \in V(G) \setminus D$, let $v \in D$ such that $uv \in E(G)$. Denote by x the vertex of H_u having a neighbor in H_v . Then, add to D_H the two vertices of degree 3 in H_u having a common neighbor with x . Observe that $|D_H| = 2(|V(G)| - |D|) + 3|D| = 2|V(G)| + |D| \leq 2|V(G)| + \ell$ and, furthermore, D_H is a

dominating set of H .

(\Leftarrow) Now, let D_H be a dominating set of H such that $|D_H| \leq 2|V(G)| + \ell$. By Claim 2, we know that $|V(H_u) \cap D_H| \geq 2$ for every $u \in V(G)$. Therefore, the number of vertices $v \in V(G)$ such that $|V(H_v) \cap D_H| \leq 3$ is at most ℓ . Claim 2 also says that if $|V(H_u) \cap D_H| = 2$ for $u \in V(G)$, then H_u has a simplicial dominated by some vertex in H_v where $|V(H_v) \cap D_H| \geq 3$. Therefore, should we assume $D \subseteq V(G)$ composed by vertices v such that $|V(H_v) \cap D_H| \geq 3$ we have a dominating set D with size at most ℓ . \square

In [19], we have that the EDGE DOMINATING SET is NP-complete for bipartite graphs with the maximum degree 3, thereby implying the NP-completeness for line graphs of bipartite graphs with the degree at most 3. Despite that result, we combine Theorems 2.3 and 3.1 for obtaining:

Corollary 3.2. *DOMINATING SET is NP-complete for planar line graphs of bipartite graphs with a maximum degree 3.*

Furthermore, the domination problem is NP-complete even for cubic graphs [13], and the reduction used in Theorem 3.1 can be used to show the NP-completeness for cubic expanded-clique graphs. Thus, we can conclude:

Corollary 3.3. *DOMINATING SET is NP-complete even for cubic line graphs of bipartite graphs.*

Proposition 3.4. *Let H be the expanded-clique graph of a graph G and function f . If S is a dominating set of H , then there is a dominating set S' of H such that $|S'| \leq |S|$ and $|S' \cap V_i| \leq 1$ for every $v_i \in V(G)$.*

Proof. Let S be a dominating set of H . Apply the following process. If $|S \cap V_i| \leq 1$ for every $v_i \in V(G)$, then set $S' = S$ and stop. Otherwise, there is $v_i \in V(G)$ such that $|S \cap V_i| \geq 2$. Let $v_{i,j} \in S \cap V_i$. If $|S \cap V_j| \geq 1$, then $S \setminus \{v_{i,j}\}$ is also a dominating set of H . Then, redefine S as $S \setminus \{v_{i,j}\}$ and repeat. If $|S \cap V_j| = 0$, then $(S \setminus \{v_{i,j}\}) \cup \{v_{j,i}\}$ is also a dominating set of H . Then, redefine S as $(S \setminus \{v_{i,j}\}) \cup \{v_{j,i}\}$ and repeat. Observe that this process eventually finishes with a set S' such that $|S' \cap V_i| \leq 1$ for every $v_i \in V(G)$. \square

A *2-independent set* in a graph G is a subset I of the vertices such that the distance between any two vertices of I in G is at least three. We denote by $\alpha_2(G, f)$ the maximum cardinality of a set S such that S is a 2-independent set of G and every $v \in S$ satisfies $f(v) = d_G(v)$.

Theorem 3.5. *Let H be the expanded-clique graph of a pair (G, f) . Then, $\gamma(H) + \alpha_2(G, f) = |V(G)|$.*

Proof. Let S be a dominating set of H . By Proposition 3.4, there is a dominating set S' of H such that $|S'| \leq |S|$ and $|S' \cap V_i| \leq 1$ for every $v_i \in V(G)$. It is clear that $|S'| \leq n$. Note that for any $v_i \in V(G)$, if $f(v_i) > d_G(v_i)$, then $|S' \cap V_i| = 1$. Therefore, if $|S' \cap V_i| = 0$, then $f(v_i) = d_G(v_i)$. Observe that if $f(v_i) = d_G(v_i)$, $S' \cap V_i = \emptyset$ and $v_i v_j \in E(G)$, then vertex $v_{j,i}$ belongs to S . Hence, the vertices v_i with $|S' \cap V_i| = 0$ form a 2-independent set T in G

with cardinality $|T| = |V(G)| - |S'|$ containing only vertices v_i such that $f(v_i) = d_G(v_i)$. Since $|S'| \leq |S|$, we conclude that $\alpha_2(G, f) \geq |V(G)| - \gamma(H)$.

Now, let T be a 2-independent set of G containing only vertices v_i such that $f(v_i) = d_G(v_i)$ and set $S = \emptyset$. If $v_i \in T$, then for every $v_j \in N_G(v_i)$, add $v_{j,i}$ to S . If v_j is not in T neither has a neighbor in T , then choose a vertex of V_j and add it to S . Since no vertex of G has more than one neighbor in T , we have that S is a dominating set of H with $|V(G)| - |T|$ vertices. Therefore, $\gamma(H) \leq |V(G)| - \alpha_2(G, f)$. \square

Corollary 3.6. *If H is the expanded-clique graph of a pair (G, f) where $f(v) > d_G(v)$ for every $v \in V(G)$, then $\gamma(H) = |V(G)|$.*

Corollary 3.6 implies that DOMINATING SET is easy for expanded-clique graphs if $f(v_i) = k$ for every $v_i \in V(G)$ for any $k > \Delta(G)$. Hence, it is interesting to think about $k = \Delta = \Delta(G)$. Another consequence of Theorem 3.5 restricted to Δ -expanded-clique graphs is that $\gamma(H) \geq \frac{|V(G)|\Delta}{\Delta+1}$.

Corollary 3.7. *If H is the Δ -expanded-clique graph of G , then $\gamma(H) \geq \frac{|V(G)|\Delta}{\Delta+1}$.*

Proof. Let S be a minimum dominating set of H . By Proposition 3.4, we know that $|S \cap V_j| \leq 1$ for every $v_j \in V(G)$, which means that $|S| \leq |V(G)|$.

Consider that all vertices of G are initially unmarked and repeat the following process until that all vertices be marked. If there is an unmarked $v_i \in V(G)$ with $d_G(v_i) = \Delta(G)$. If $|S \cap V_i| = 1$, then mark v_i and repeat. If $S \cap V_i = \emptyset$ and every vertex of $N[v_i]$ is unmarked, then for every $v_j \in N_G(v_i)$, we know that $v_{j,i}$ belongs to S . Then, mark the vertices of $N[v_i]$ and repeat. If $S \cap V_i = \emptyset$ and there is some $v_j \in N(v_i)$ that is marked. Observe $v_{j,i} \notin S$, which means that some vertex of V_i belongs to S . Then, mark v_i and repeat. If there is an unmarked vertex v_i such that $d_G(v_i) < \Delta(G)$. Observe that there is a vertex of V_i in S . Then, mark v_i and repeat.

Since $V(G)$ is finite, the above process eventually ends. Observe that in each step i , if the number of vertices that are marked is denoted by x_i and, among these vertices, the number of vertices that belong to S is x'_i , it holds that $\frac{x'_i}{x_i} \geq \frac{\Delta}{\Delta+1}$. Since each vertex is marked exactly once, we have that $\gamma(H) \geq \frac{|V(G)|\Delta}{\Delta+1}$. \square

A consequence of Corollaries 3.6 and 3.7 is that an approximation of dominating set of k -expanded-clique graphs within $(1 + \frac{1}{\Delta})$ -approximation by simply selecting one vertex of each expanded-clique.

We conclude by showing that DOMINATING SET can be solved in linear time for $k = 2$ since.

Proposition 3.8. *The following hold for a 2-expanded-clique graph H with a connected root G .*

- *If G is a path with two or three vertices, then $\gamma(H) = k$;*
- *If G is a path with $q \geq 4$ vertices or G is a cycle with q vertices, then $\gamma(H) = \lceil \frac{q}{3} \rceil$.*

Proof. For G being a path with two or three vertices, we have $\gamma(H) = 1$ for $k = 1$ because H is a P_2 as well, or $\gamma(H) = 2$ since H is a P_4 or P_6 when $k = 2$, and so $\gamma(H) = k$. The next ones handled together, taking G a path (cycle) with $q \geq 4$, it implies H is path (cycle) with $2q$ vertices. Since one vertex dominates two others, we can remove a P_3 from G and obtain a new graph G_1 so that G' is a connected subgraph of G_1 . Applying repeatedly the previous process, let it be r times, until the resulting graph G_r has no more than 2 vertices, then $\gamma(H) = r = \lceil \frac{q}{3} \rceil$ if G_r is empty or $\gamma(H) = r + 1 = \lceil \frac{q}{3} \rceil$ otherwise. Thus, it finishes the proof. \square

References

- [1] F. N. Abu-Khzam and K. Lamaa, *Efficient heuristic algorithms for positive-influence dominating set in social networks*, IEEE INFOCOM 2018 - IEEE Conference on Computer Communications Workshops (INFOCOM WKSHPS), 2018, pp. 610–615.
- [2] Derek G. Corneil and Lorna K. Stewart, *Dominating sets in perfect graphs*, Discrete Mathematics **86** (1990), no. 1, 145 – 164.
- [3] Ding-Zhu Du and Peng-Jun Wan, *Connected dominating set: Theory and applications*, vol. 77, Springer Optimization and Its Applications, no. 1, Springer Publishing Company, Incorporated, 2013.
- [4] O. Favaron, *Irredundance in inflated graphs*, J. Graph Theory **28** (1998), 97–104.
- [5] Michael R. Garey and David S. Johnson, *Computer and intractability: A guide to the np-completeness*, W. H. Freeman & Co., USA, 1979.
- [6] F. Harary and C. Holzmann, *Line graphs of bipartite graphs*, Rev. Soc. Mat. Chile **1** (1974), 19–22.
- [7] Toru Hasunuma, *Structural properties of subdivided-line graphs*, Combinatorial Algorithms (Berlin, Heidelberg) (Thierry Lecroq and Laurent Mouchard, eds.), Springer Berlin Heidelberg, 2013, pp. 216–229.
- [8] Michael A. Henning and Adel P. Kazemi, *Total domination in inflated graphs*, Discrete Applied Mathematics **160** (2012), no. 1, 164–169.
- [9] Jisu Jeong, Sigve Hortemo Sæther, and Jan Arne Telle, *Maximum matching width: New characterizations and a fast algorithm for dominating set*, Discrete Applied Mathematics **248** (2018), 114 – 124, Seventh Workshop on Graph Classes, Optimization, and Width Parameters, Aussois, France, October 2015.
- [10] Liying Kang, Moo Young Sohn, and T.C.E. Cheng, *Paired-domination in inflated graphs*, Theoretical Computer Science **320** (2004), no. 2, 485–494.
- [11] S. Klavžar and U. Milutinović, *Graphs $s(n, k)$ and a variant of the tower of Hanoi problem*, Czechoslovak Mathematical Journal **47** (1997), 95–104.

- [12] Sandi Klavžar, Uroš Milutinović, and Ciril Petr, *1-perfect codes in Sierpiński graphs*, Bulletin of the Australian Mathematical Society **66** (2002), no. 3, 369–384.
- [13] Jan Kratochvíl, *Regular codes in regular graphs are difficult*, Discrete Mathematics **133** (1994), no. 1, 191 – 205.
- [14] Ruizhi Li, Shuli Hu, Jian Gao, Yupeng Zhou, Yiyuan Wang, and Minghao Yin, *Grasp for connected dominating set problems*, Neural Computing and Applications **28** (2018), 1059 – 1067.
- [15] Subhrangsu Mandal and Arobinda Gupta, *Approximation algorithms for permanent dominating set problem on dynamic networks*, Distributed Computing and Internet Technology (Cham) (Atul Negi, Raj Bhatnagar, and Laxmi Parida, eds.), Springer International Publishing, 2018, pp. 265–279.
- [16] T. Pino, S. Choudhury, and F. Al-Turjman, *Dominating set algorithms for wireless sensor networks survivability*, IEEE Access **6** (2018), 17527–17532.
- [17] Cristanel Razafimandimby, Valeria Loscrí, Anna Maria Vegni, and Abderrahim Benslimane, *Neuro-dominating set scheme for a fast and efficient robot deployment in internet of robotic things*, Ad Hoc Networks **86** (2019), 36 – 45.
- [18] Oliver Schaudt and Rainer Schrader, *The complexity of connected dominating sets and total dominating sets with specified induced subgraphs*, Information Processing Letters **112** (2012), no. 24, 953 – 957.
- [19] Mihalis Yannakakis and Fanica Gavril, *Edge dominating sets in graphs*, SIAM J. Appl. Math. **38** (1980), 364–372.
- [20] D. Zhao, G. Xiao, Z. Wang, L. Wang, and L. Xu, *Minimum dominating set of multiplex networks: Definition, application, and identification*, IEEE Transactions on Systems, Man, and Cybernetics: Systems (2020), 1–15.