

Quantum Field Theory based Quantum Information: Measurements and Correlations

Charis Anastopoulos ^(a) *, Bei-Lok Hu ^(b) †, and Konstantina Savvidou ^(a) ‡

^(a) Department of Physics, University of Patras, 26500 Greece

^(b) Maryland Center for Fundamental Physics and Joint Quantum Institute,
University of Maryland, College Park, Maryland 20742-4111 U.S.A.

August 9, 2022

Abstract

This is the first in a series of papers aiming to develop a relativistic quantum information theory in terms of unequal-time correlation functions in quantum field theory. In this work, we highlight two formalisms which together can provide a useful theoretical platform suitable for further developments: 1) Quantum field measurements using the Quantum Temporal Probabilities (QTP) method; 2) Closed-Time-Path (CTP) formalism for causal time evolutions. QTP incorporates the detector into the quantum description, while emphasising that the records of measurement are macroscopic and can be expressed in terms of classical spacetime coordinates. We first present a new, elementary derivation of the QTP formulas for the probabilities of n measurement events. We then demonstrate the relation of QTP with the Closed-Time-Path formalism, by writing an explicit formula that relates the associated generating functionals. We exploit the path integral representation of the CTP formalism, in order to express the measured probabilities in terms of path integrals. After this, we provide some simple applications of the QTP formalism. In particular, we show how Unruh-DeWitt detector models and Glauber's photodetection theory appear as limiting cases. Finally, with quantum correlation being the pivotal notion in relativistic quantum information and measurements, we highlight the role played by the CTP two-particle irreducible effective action which enables one to tap into the resources of non-equilibrium quantum field theory for our stated purpose.

*anastop@upatras.gr

†blhu@umd.edu

‡ksavvidou@upatras.gr

1 Introduction

Quantum information theory (QIT) is a quantum extension of classical information theory. The primary motive force behind explosive developments of QIT in the last quarter of a century is entanglement, the uniquely quantum feature which provides the new and powerful resource for quantum computing, quantum metrology, quantum communication and more.

While potential applications of quantum information technologies have drawn close attention of government and private research and development, the theoretical foundation of QIT is surprisingly lacking in a major portion. Our understanding of QIT is far lagging behind the fully developed quantum theory of nature, namely, quantum field theory (QFT), which has proven its validity and worth in the full range of physical sciences from particle-nuclear physics to atomic, optical and condensed matter physics, from quarks to black holes and the early universe. So far, quantum information theory has been largely developed in the context of non-relativistic quantum mechanics, a small corner of full QFT. It is ostensibly inadequate when basic relativistic effects – both special and general – such as causality and covariance, need be accounted for.

1.1 QITs not based on QFT are incomplete

QFT is a quantum theory that also includes axioms about the effect of spacetime structure on the properties of quantum systems, especially regarding the causal propagation of signals. In contrast, current quantum information theories do not incorporate the latter axioms. Their notion of causality, based on the sequence of successive operations on a quantum system, lacks a direct spacetime representation. As a result, current QITs cannot make crucial relativistic distinctions, for example, between timelike and spacelike correlations, it does not describe real-time signal propagation, and it ignores relativistic constraints on permissible measurements. A genuinely relativistic QIT must overcome such limitations [1].

Furthermore, experiments that study causal information transfer or gravitational interaction in multi-partite quantum systems require a QFT treatment of interactions for consistency. A non-QFT description may severely misrepresent the theoretical modeling of the system or the physical interpretation of the results. This point is crucial for tests of foundational issues of quantum theory invoking quantum information concepts such as entanglement and decoherence or methods such as signaling and communication. This is especially so for quantum information experiments in space [2, 3] and for experiments designed to explore quantum effects from gravity [4–7].

The introduction of the key concepts of spacetime covariance and causality in QIT forces us to address problems that originate from the foundations of QFT. We draw an incomplete list of noteworthy issues below.

Quantum States. It has long been known that the quantum state is genuinely different when recorded from different Lorentz frames in set-ups that involve more than two quan-

tum measurements [8–10]. There is no problem with the theory’s physical predictions that are expressed in terms of (multi-time) probabilities [11]. However, the usual notions of quantum information (entropy, entanglement) are defined through the quantum state, and as such, they are ambiguous in relativistic measurement set-ups.

Local Operations. It is a challenging problem to formalize the notion of a localized quantum system in QFT. There are powerful theorems demonstrating that even allowing for unsharp localization in a spatial region leads to faster than light signals [12, 13]. Hence, expressing the crucial quantum informational notion of a local operation in terms of spatial localization can lead to conflicts with relativistic causality.

Projective Measurements. There are strong arguments that ideal (i.e., projective) measurements in QFT are incompatible with causality [14], essentially because they change the quantum state over a full Cauchy surface. However, existing quantum information notions (even the very notion of a qubit) presuppose maximal extraction of information through ideal measurements.

We contend that the development of consistent relativistic QIT requires a measurement theory that respects causality and locality while being expressed in terms of quantum fields¹. Furthermore, this measurement theory ought to be practical, i.e., it should lead to non-trivial predictions for setups that are accessible now or in the near future. For other reasons why a QFT measurement theory is needed, see Ref. [19].

1.2 Past work on QFT measurements

The earliest discussion of QFT measurements, to our knowledge, was by Landau and Peierls [20], who derived an inequality for the localization of particles. Bohr and Rosenfeld criticized some of their assumptions [21], and proved the crucial result that the measurement of field properties requires a test particle of macroscopic scale, in the sense that its charge Q must be much larger than the electron charge e .

Arguably the first explicit model for QFT measurement was Glauber’s photodetection theory [22, 23]. The theory was developed as a quantum generalization of the classical theory of coherence for the EM field. It expresses unnormalized probabilities for photon detection in terms of the electric field operators $\mathbf{E}(x)$ and the field state $|\psi\rangle$. The joint probability density $P(x_1, x_2, \dots, x_n)$ for the detection of one photon at each of the spacetime points x_1, x_2, \dots, x_n is given by

$$P(x_1, x_2, \dots, x_n) = \langle \psi | E^{(-)}(x_1) E^{(-)}(x_2) \dots E^{(-)}(x_n) E^{(+)}(x_n) \dots E^{(+)}(x_2) E^{(+)}(x_1) | \psi \rangle, \quad (1)$$

where $E^{(+)}$ is the positive frequency component and $E^{(-)}$ the negative frequency component of the projected field vector field $\mathbf{n} \cdot \mathbf{E}(x)$. The joint detection probability of photons at different moments of time is essential for the definition of higher order coherences of

¹The explicit formulation in terms of QFT is the point of divergence of our approach from past works on formulating a relativistic QIT [15–18].

the electromagnetic field, and for describing phenomena like the Hanbury-Brown-Twiss effect, photon bunching and anti-bunching [24].

Glauber’s theory has been immensely successful in quantum optics. However, its applicability is limited to measurements of photons. Another problem is that the field splitting into positive and negative frequency components is non-local: it follows from the so-called Rotating Wave Approximation for the interaction of the field to the detector. The latter misrepresents the retarded propagation of the electromagnetic field, hence, the model may face problems with causality in set-ups that involve propagation along distances.

A very common class of models employed for QFT measurements are the *Unruh-DeWitt* detectors [25, 26]. They first appeared in the study of the Unruh effect, in order to clarify the physical properties of the field that are experienced by accelerated observers. In an Unruh-DeWitt detector, the quantum field is coupled to a point-like system that moves along a pre-determined spacetime trajectory $x(\tau)$, where τ is the trajectory proper-time.

While a first-principles derivation of the detector’s response for non-inertial motion is still missing, Unruh-DeWitt detector models are the simplest models for describing QFT measurements [27], and they have found several applications—for a sampling of the latter, see Ref. [28]. Unruh-DeWitt detectors are limited in that detector degrees of freedom are not described by a QFT. As a result, they may lead to non-causal signals in systems that involve more than one detector—see, for example, the discussions in Refs. [29–31].

Measurement models have also been constructed in the context of algebraic QFT [32–36], primarily in order to address issues of causality and locality. For example, Ref. [35] considers a system and a probe, both described by a QFT. The two field systems are independent and they interact within a bounded spacetime region. Their interaction can be described by an S matrix, thus leading to correlations between observables on the system and records on the probe. One can define probabilities for the latter using measurement theory for some operators that are well defined on the probe Hilbert space. However, this program has not quite yet been developed into a practical tool capable of concrete physical predictions.

1.3 Our approach

In this work, we describe QFT measurements using the Quantum Temporal Probabilities method [37–39]. The original motivation of this method was to provide a general framework for temporally extended quantum observables [40–42], hence, the name. The key idea is to distinguish between the time parameter of Schrödinger equation from the time variable associated to particle detection [43, 44]. The latter are then treated as macroscopic quasi-classical variables of the detector. Here, we use the word ‘quasi-classical’ as in the decoherent histories approach to quantum theory [45–50]. It refers to coarse-grained quantum variables that satisfy appropriate decoherence conditions, and

they approximately satisfy classical evolution equations [49, 50]. Hence, although the detector is described in microscopic scales by quantum theory, its macroscopic records are expressed in terms of classical spacetime coordinates. Furthermore, the detector is also described in terms of quantum fields. Its interactions with the detector respect causality, because the interaction Hamiltonian is a local functional of quantum fields.

Glauber’s detection theory and Unruh-DeWitt detector models emerge from QTP as limiting cases that characterize specific regimes: Glauber’s theory appears at the limit of very small characteristic timescales for the detector, while Unruh-DeWitt models appear at the limit of very short length-scales. In comparison to the algebraic QFT approaches to measurements, QTP has a more nuanced description of the detectors, which allows it to develop into a practical tool to be employed in a large class of problems.

A key insight from QTP is that probabilities for measurements are defined in terms on specific unequal-time correlation functions of the quantum field. This is particularly important, as such correlation functions are a staple of QFT. Powerful methods have been developed for their calculation and the analysis of their properties. The specific correlation functions relevant to QTP are not the usual ones of S-matrix theory (in-out formalism), but they appear in the Closed-Time-Path (CTP) (Schwinger-Keldysh or ‘in-in’) formalism [51–55]. The CTP formalism improves over the in-out formalism, in that it allows for real and causal equations of motion, and it has found many applications in early universe cosmology [56, 57], nuclear-particle process [55, 58–60] and condensed matter physics [61, 62].

The connection that we establish between the two formalisms allows us to translate between the concepts of quantum measurement theory and of quantum field theory. The former is based on operational notions like Positive-Operator-Valued measures (POVMs) and effects, while the latter employs functional methods like path integrals in order to express its predictions in a manifestly covariant language. In this work, we show, for example, how to express POVMs for particle detection in terms of path-integrals.

A key point of our analysis is the central role played by the unequal-time correlation functions. In the QTP description, they contain all information about measured probabilities. The detection probability for N events depends on an $2N$ -unequal time correlation function. This has the following implication. Two-event measurements typically characterize bipartite systems, and as such the corresponding probabilities are related to entanglement. Hence, field four-point functions contain all information that pertains to bipartite entanglement. A relation at the level of non-relativistic fields was shown in [39].

Working at the level of the correlation functions brings us closer to the main ideas of non-equilibrium QFT, where irreversibility is implemented by the truncation of the hierarchy of field correlation functions, and the slaving of higher correlation functions to the two-point function. Again, there is a natural relation between QTP and non-equilibrium formalisms that are based on CTP. We argue that the QTP probabilities function as a registrar of information for the quantum field keeping track of how much information resides in which, and how it flows from one order to another through the

dynamics.

We believe that the scheme presented here has good potential to systemize quantum information in QFT, and to identify the parts of this information that is relevant to the field's statistical, stochastic and thermodynamic behavior. Hence, this formalism could provide a concrete method for defining quantum information in QFT via the correlation hierarchy, as has been proposed in [63]. Such a definition would be very different from definitions of information in standard QIT that is based on the properties of the single-time quantum state.

1.4 Our results

Our results are the following.

First, we present a new derivation of the QTP formulas for the probabilities of n measurement events. The derivation is intuitive and pedagogical, as it only requires elementary perturbation theory. It reproduces the results of the more rigorous QTP analysis to leading order in perturbation theory with respect to the system-apparatus coupling. This suffices for most applications. Then we analyze the mathematical structure of the probability formulas. In particular, we show how they provide an explicit connection between the concepts of quantum measurement theory and QFT concepts, namely, unequal-time field correlation functions.

Second, we demonstrate the relation of QTP with the Closed-Time-Path formalism, by deriving an explicit formula that relates the associated generating functionals. The Closed-Time-Path formalism has a natural path integral formulation, which allows us to express the QTP probabilities in terms of path integrals. Conversely, QTP provides a theory of observables appropriate to the in-in formalism.

Third, we provide some demonstrative applications of the formalism. We briefly review the construction of time-of-arrival probabilities for relativistic particles. Then we show how the commonly employed models of Glauber's photodetection and Unruh-DeWitt detectors appear as limiting cases of the QTP formalism. This implies, in particular, that there exists regimes in which QTP provides different predictions from Glauber's theory, and these differences may be experimentally distinguishable.

Fourth, we explore the links between QTP and non-equilibrium QFT. For field states with a large number of particles, QTP probabilities provide natural definitions for Boltzmann-type observables. They also allow us to explore how information flows between the different levels of the Schwinger-Dyson hierarchy of correlation functions.

This paper is structured as follows. In Sec. 2, we present the main ideas of the QTP method, and we derive the probability formula for n -detections. In Sec. 3, we derive the connection with the CTP formalism, and also derive a path integral expression for QTP probabilities. In Sec. 3, we present some applications of the formalism, and in Sec. 4, we explore the links to non-equilibrium QFT.

2 Probabilities for QFT measurements

In this section, we present the QTP approach to quantum field measurements. First, we explain in more detail the need for a QFT measurement theory. Then, we derive the QTP probability formulas and we analyze their structure. The key feature of the probability formula is that probability densities pertaining to n measurement events are linear functionals of specific $2n$ unequal-time correlation functions.

2.1 The need of a QFT measurement theory

Most QFT predictions involve set-ups with a single state preparation and a single-detection event, which can be described in terms of the S-matrix. For example, S-matrix amplitudes determine scattering cross-sections; S-matrix poles determine the spectrum of composite particles and decay rates. This gives the general impression that there is no further need for an elaborate measurement theory.

However, this over-simplified view is deceptive. There are at least two cases where the S-matrix analysis is insufficient. First, in quantum optics, we need joint probabilities of detection in order to describe phenomena that involve high-order coherences of the EM field, like photon bunching and anti-bunching [24]. A first-principles calculation of joint probabilities requires a non-trivial description of quantum measurements. In non-relativistic physics, such a calculation involves the use of the state-update rule, i.e., quantum state reduction. In QFT, such a rule is missing, even from axiomatizations—see Ref. [1]. In practice, joint probabilities are expressed in terms of photodetection models, like Glauber’s, whose derivation is rather heuristic. However, planned experiments in deep space [2,3] that involve measurement of EM field correlations will arguably require a first-principles analysis of joint probabilities in order to take into account the relative motion of detectors and delayed propagation at long distances.

The second case where S matrix theory does not suffice is when the expectation values of physical quantities at finite moments of time are called for, rather than just the asymptotic amplitudes offered in the ‘in-out’ S-matrix theory. A notable example is the construction of dynamical equations for the evolution of spacetime metric in the early universe and in black holes from the solutions to the semiclassical Einstein equation [64, 65]; these equations require the evaluation of the expectation values of the stress-energy and its fluctuations. Wherever causal time developments are needed, instead of transition between asymptotic states, this ‘in-in’ formulation of QFT, as in the CTP formalism, is indispensable. This is certainly true for the description of non-equilibrium processes such as quantum transport in quantum many-body systems. Powerful functional techniques have been developed to deal with such problems. Indeed, in this paper we demonstrate a connection between those techniques and our account of QFT measurements.

A crucial challenge in incorporating quantum informational concepts in QFT is the lack of a common mathematical language. QIT focuses on the Hilbert space aspects of quantum theory, for example, the geometry of the space of quantum states, the

characterization of entanglement and the identification of physically permissible state transformations. It relies crucially on the way quantum theory describes the extraction of information through measurements. In contrast, QFT is commonly described through functional methods that lead to the evaluation of unequal-time field correlation functions. The latter contain all information about the system, and they can be easily manipulated to derive quantities like perturbative scattering amplitudes, expectations of stress-energy tensors and single-particle Wigner functions.

The mathematical difference is not only technical, as it pertains to fundamental notions like locality and causality. In QIT, these issues are usually expressed through the Local Operations and Classical Communication (LOCC) paradigm [66]: The Hilbert space of any informational system is split up as a tensor product $\otimes_i \mathcal{H}_i$, where \mathcal{H}_i is the Hilbert space of the i -th subsystem. A local operation on the i -th subsystem is a set of completely positive maps $\mathcal{C}^{(i)}(a)$ on states of \mathcal{H}_i , such that $\sum_a \mathcal{C}^{(i)}(a) = \hat{I}$, where a are the measurement outcomes of the operation. Then, causality is implemented through the concept of classical communication. An operation $\mathcal{C}^{(i)}(a)$ on a subsystem i may depend on the outcome b of an operation $\mathcal{D}^{(j)}(b)$, if the outcome can be communicated to i through a classical channel, prior to the operation $\mathcal{C}^{(i)}(a)$. Hence, QIT carries the causal structure of classical communication. It is usually not concerned with real-time quantum signal propagation between disconnected subsystems.

In contrast, QFTs focus on spacetime propagation of information, as expressed in the theory's dynamics and spacetime symmetries. These properties are nicely captured by the unequal-time correlation functions, which can be chosen to be manifestly spacetime covariant. Causality is implemented by the requirement that specific correlation functions vanish if their arguments are spacelike separated².

2.2 The Quantum Temporal Probabilities Approach: key points

The key features of the QTP approach to measurements on quantum fields are the following. First, the apparatus is fully incorporated into the quantum description and it is also treated via QFT. The interaction between the measured system and the apparatus local and causal, in the sense that it is governed by an interaction Hamiltonian that is a local functional of quantum fields.

The measurement apparatus is a macroscopic system that exhibits classical behavior. We describe this behavior through the decoherent histories approach to emergent classicality [46, 49]. We assume that the pointer variable is a highly coarse-grained observable, so that histories for measurement outcomes satisfy appropriate decoherence conditions.

QTP treats measurements events as localized in space and in time, as is the case for

²To be precise, causality is incorporated in the *cluster decomposition* property of the S-matrix [67]. Cluster decomposition is a property of a specific hierarchy of correlation functions $G_n(X_1, X_2, \dots, X_n)$, where $n = 0, 1, 2, \dots$. It asserts that $G_{n+m}(X_1, \dots, X_n, X'_1, \dots, X'_m) = G_n(X_1, X_2, \dots, X_n)G_m(X'_1, X'_2, \dots, X'_m)$, if the cluster of points X_1, \dots, X_n is spacelike separated from the cluster X'_1, \dots, X'_m .

all physical measurement outcomes. For example, an elementary solid state detector has a specific location in a lab and it records a particle at a specific moment of time that is determined with finite accuracy. In principle, both position and time can be random variables. For example, when directing a single particle towards an array of detectors, both the elementary detector that records the particle (i.e., the location of the detection record) and the time of recording vary from one run of the experiments to the other. Hence, the predictions of the theory must be expressed in terms of probability densities

$$P(x_1, q_1; x_2, q_2, \dots, x_n, q_n), \quad (2)$$

for multiple detection events. Here, x_i are spacetime points, q_i stand for any other observable that has been recorded and P is a probability density with respect to both x_i and q_i .

Since there is no self-adjoint operator for time, there are no ideal (i.e., projective) measurements for time. It follows that none of the probabilities densities (2) corresponds to an ideal measurement, not even ones that involve single-measurement event. The probabilities (2) are defined through Positive Operator Valued Measures (POVMs).

2.3 Detection probability for a single detector

Consider a QFT on Minkowski spacetime M , described by a Hilbert space \mathcal{F} , which carries a unitary representation of the Poincaré group. We denote the fields in the Heisenberg picture (with respect to the Hamiltonian in \mathcal{F} by $\hat{\phi}_r(x)$, where r runs over spacetime and internal indices.

An apparatus is a physical system described by a Hilbert space \mathcal{K} . The apparatus follows a world tube $\mathcal{W} = \mathbb{R} \times U$ in Minkowski spacetime, for some $U \subset \mathbb{R}^3$. We assume that the size of the apparatus is finite, but still much larger than the scale of microscopic dynamics. We will elaborate on the properties of the apparatus later.

We assume a coupling between the system and the apparatus with support only in a small spacetime region around a point x . The finite spacetime extent of the coupling is arguably unphysical, but it serves to mimic the effect of a detection record localized at x . To leading order in perturbation theory, the resulting probabilities coincide with those obtained from a more detailed modeling of the measurement act through localized POVMs for the time of detection [39]. Working in the interaction picture, we express the coupling term as

$$\hat{V}_x = \int F_x(y) \hat{C}_a(y) \otimes \hat{J}^a(y), \quad (3)$$

where $\hat{C}_a(x)$ is a composite operator on \mathcal{F} that is local with respect to the field $\hat{\phi}_r(x)$ and a runs over spacetime and internal indices. The current operators $\hat{J}^a(x)$ are defined on \mathcal{K} .

The switching functions $F_x(y)$ are dimensionless, they depend on the motion of the apparatus and they vanish outside the apparatus world tube and at times that the

interaction is switched-off. The total spacetime volume associated to a switching function is $v = \int dY F_x^2(y)$.

The S-matrix associated to Eq. (3) is $\hat{S}_x = \mathcal{T} \exp[-i \int d^4y F_x(y) \hat{C}_a(y) \otimes \hat{J}^a(y)]$, where \mathcal{T} stands for time ordering. To leading order in the interaction,

$$\hat{S}_x = \hat{I} - i\hat{V}_x. \quad (4)$$

Let the initial state of the system be $|\psi\rangle \in \mathcal{F}$ and the initial state of the apparatus be $|\Omega\rangle$. A particle record appears if the detector transitions from $|\Omega\rangle$ to its complementary subspace \mathcal{K}' . Furthermore, on \mathcal{K}' , we measure a property of the particle through a pointer observable q . The latter is described by a family of positive operators $\hat{\Pi}(q)$, such that $\sum_q \hat{\Pi}(q) = \hat{I} - |\Omega\rangle\langle\Omega|$. The pointer observable is typically very coarse, and we assume that it is stationary under spacetime translations by the self-dynamics of the detector, so that the record is preserved after the end of the measurement.

Then, we evaluate the probability

$$\text{Prob}(x, q) = \langle\psi, \Omega | \hat{S}_x^\dagger [\hat{I} \otimes \hat{\Pi}(q)] \hat{S}_x |\psi, \Omega\rangle \quad (5)$$

that the detector is excited and records a value q to leading order in perturbation theory

$$\text{Prob}(x, q) = \int d^4y_1 d^4y_2 F_x(y_1) F_x(y_2) G_{ab}(y_1, y_2) \langle\Omega | \hat{J}^a(y_1) \hat{\Pi}(q) \hat{J}^b(y_2) | \Omega\rangle, \quad (6)$$

where

$$G_{ab}(x, x') = \langle\psi | \hat{C}_a(x) \hat{C}_b(x') | \psi\rangle, \quad (7)$$

is a correlation function for the quantum field.

The probability $\text{Prob}(x, q)$ of Eq. (6) is not a density with respect to x ; x appears as a parameter of the switching function. In classical probability theory, we could define an unnormalized probability density $W(x, q)$ with respect to x by dividing $\text{Prob}(x, q)$ with the effective spacetime volume v ,

$$W(x, q) = v^{-1} \text{Prob}(x, q). \quad (8)$$

To further proceed in our analysis, we have to make specific assumptions about the detector. First, we assume that the detector carries a representation of the spacetime translation group with generators \hat{p}^μ . We also take $|\Omega\rangle$ to be approximately translation invariant³. This is a reasonable assumption for a macroscopic detector, with size much larger than the atomic scale.

³In general, we assume that the detector Hilbert space also carries a representation of the Poincaré group. However, the state $|\Omega\rangle$ is not the Poincaré invariant vacuum. An actual detector involves a macroscopically large number of fermions—to be precise, it is described by the subspace of the fermionic fields for leptons and baryons that corresponds to macroscopically large, constant, values of the leptonic and baryonic quantum numbers—and it defines a preferred reference system at which its center of momentum has zero three-momentum.

Let x_0 be a reference point along the world-tube of the detector. Then,

$$\hat{J}^a(y) = e^{-i\hat{p}\cdot(y-x_0)} \hat{J}^a(x_0) e^{i\hat{p}\cdot(y-x_0)}, \quad (9)$$

and we can write $\langle \Omega | \hat{J}^a(y_1) \hat{\Pi}(q) \hat{J}^b(y_2) | \Omega \rangle = R^{ab}(y_2 - y_1, q)$, where

$$R^{ab}(x, q) := \langle \Omega | \hat{J}^a(x_0) \sqrt{\hat{\Pi}(q)} e^{-i\hat{p}\cdot x} \sqrt{\hat{\Pi}(q)} \hat{J}^b(x_0) | \Omega \rangle \quad (10)$$

We will refer to $R^{ab}(y, q)$ as the *detector kernel*.

The simplest choice for the switching functions F_x are Gaussians,

$$F_x(y) = \exp\left[-\frac{1}{2}D(x, y)\right], \quad (11)$$

where D a Euclidean distance function on Minkowski spacetime. Such functions are defined in terms of a dimensionless Euclidean metric $g_{\mu\nu}^{(E)}$ on Minkowski spacetime. A simple and physically relevant metric is determined by the timelike vector field u_μ normal along the world-tube, by

$$g_{\mu\nu} = \frac{1}{\delta_t^2} u_\mu u_\nu + \frac{1}{\delta_x^2} (u_\mu u_\nu + \eta_{\mu\nu}). \quad (12)$$

where δ_t is the temporal accuracy and δ_x is the special accuracy of the detector. Both are assumed to be macroscopic scales, in order to make contact with the main assumption of the QTP method, namely the existence of macroscopic records of observation [39].

The Gaussian switching functions satisfy a useful identity

$$f(x)f(x') = f^2\left(\frac{x+x'}{2}\right) \sqrt{f(x-x')}. \quad (13)$$

The volume v of the spacetime region in which the field-apparatus interaction is switched on is $v = \pi^2 \delta_t \delta_x^3$. We note that the function $\sigma(x) := \frac{1}{v} f^2(x)$ is a normalized probability density on M . Then, we write

$$W(x, q) = \int d^4y \sigma(x-x') P(x', q), \quad (14)$$

where

$$P(x, q) = \int d^4\xi \sqrt{f(y)} R^{ab}(y, q) G_{ab}\left(x - \frac{1}{2}y, x + \frac{1}{2}y\right), \quad (15)$$

The probability distribution $W(x, q)$ is the convolution of $P(x, q)$ with the probability density $\sigma(x)$ that incorporates the accuracy of our measurements. If $P(x, q)$ is non-negative and the scale of variation in x is much larger than both δ_t and δ_x , we can treat $P(x, q)$ as a finer-grained version of $W(x, q)$ and employ this as our probability density for detection.

The kernel $R^{ab}(x, q)$ is typically characterized by some correlation length-scale ℓ and some correlation time-scale τ , such that $R^{ab}(x, q) \simeq 0$ if $|t(\xi)| \gg \tau$ or $|\mathbf{x}(\xi)| \gg \ell$. Both scales ℓ and τ are microscopic and characterize the constituents of the apparatus and their dynamics. If $\ell \ll \delta_x$ and $\tau \ll \delta_t$, then $R^{ab}(x, q)\sqrt{f}(x) \simeq R^{ab}(x, q)$ and the QTP probability formula simplifies to

$$P(x, q) = \int d^4\xi R^{ab}(y, q) G_{ab}(x - \frac{1}{2}y, x + \frac{1}{2}y). \quad (16)$$

The probability densities (16) are not normalized to unity. In general, the total probability of detection $P_{det} = \sum_q \int_{\mathcal{W}} d^4x P(q, x)$ must be a small number, for perturbation theory to be applicable. There is always a probability $P(\emptyset) = 1 - P_{det}$ of no detection. We normalize probabilities by dividing $P(x, q)/P_{det}$, i.e., by conditioning the probability densities $P(q, x)$ with respect to the existence of a detection record.

2.4 Remarks

1. The definition (8) of a spacetime density with respect to time is not rigorous for quantum probabilities, because it involves the combination of probabilities defined with respect to different experimental set-ups, i.e., different switching functions for the Hamiltonians. There are numerous theorems asserting that such combinations may not be acceptable in quantum theory. Nonetheless, Eq. (8) can be derived as a genuine probability density in the context of the QTP method [39, 68], as long as we restrict to the leading order of perturbation theory.
2. Probabilities are defined here using the Born rule for the pointer variable. The more rigorous derivation employs the decoherent histories probability assignment which incorporates the Born rule.
3. In QTP, the interaction is present at all times, as it should be in any first-principles derivation. The smearing functions $F_x(y)$ are not interpreted in terms of a switching-on of the interaction, but they describe the *sampling* of the spacetime point. Hence, the spacetime volume \hat{v} is a measure of *coarse-graining*, i.e., of the inaccuracy in the determination of the spacetime point. This point is important for a proper derivation, because probabilities can only be defined for histories that satisfy a decoherence condition, for which coarse-graining is a prerequisite. In principle, a preferred value of v that corresponds to the coarse-graining scale at which probabilities are well-defined is determined from first-principles—see [37] for explicit calculations in simple models. This means that not all sampling functions $F_x(y)$ are acceptable: their support cannot be made arbitrarily small. Such constraints cannot be seen in the derivation that we presented here.
4. QTP leads to different predictions from the method presented here at higher orders of perturbation theory for the system-apparatus interaction. However, in many set-ups only the lowest order terms can be said to provide a meaningful signal, i.e., a correlation between observables in the measured field and pointer variables in the apparatus. Higher

order interaction terms often degrade such correlations, and for this reason they can be conceptualized as noise. This is the case, for example, in Glauber's photodetection theory.

5. In general, the field-apparatus couplings are fixed by the standard model of particle physics. Even in idealized models for detectors, we have to specify the physical process through which detection takes place. This information is encoded in the composite operators $\hat{C}(x)$. Consider, for example, the case of a QFT with a single scalar field $\hat{\phi}(x)$. The choice $\hat{C}(x) = \hat{\phi}(x)$ corresponds to detection of particles through absorption. The choice $\hat{C}(x) =: \hat{\phi}^2(x) :$ corresponds to particle detection through scattering, through terms that involve one creation and one annihilation operator. This observation generalizes to fields of arbitrary spin.

2.5 Detection probability for multiple detectors

We proceed to the derivation of a probability formula for the case that the field interacts with n detectors. Each detector is described by a Hilbert space \mathcal{K}_i , $i = 1, 2, \dots, n$. The total Hilbert space of the system is $\mathcal{F} \otimes \mathcal{K}_1 \otimes \mathcal{K}_2 \otimes \dots \otimes \mathcal{K}_n$. The coupling operator of the i -th detector is

$$\hat{V}_x^{(i)} = \int F_x(y) \hat{C}_a^{(i)}(y) \otimes \hat{I} \otimes \dots \otimes \hat{J}_{(i)}^a(y) \otimes \dots \otimes \hat{I}, \quad (17)$$

where $\hat{J}_{(i)}^a(y)$ is a current operator for the i -th detector and $\hat{C}_a^{(i)}$ is the associated composite operator.

We denote the S-matrix for the i -th detector by $\hat{S}_x^{(i)}$. Again, to leading order in the interaction $\hat{S}_x^{(i)} = \hat{I} - i\hat{V}_x^{(i)}$. We also denote the initial state of the i -th detector by $|\Omega_i\rangle$, and the measurement operators as $\hat{\Pi}^{(i)}(q_i)$.

The key point is that the S matrix for the total interaction of the field is defined by time ordering with respect to the spacetime points x_1, x_2, \dots, x_n associated to the detectors,

$$\hat{S}_{x_1, x_2, \dots, x_n} = \mathcal{T} \left[\hat{S}_{x_1}^{(1)} \hat{S}_{x_2}^{(2)} \dots \hat{S}_{x_n}^{(n)} \right]; \quad (18)$$

here, \mathcal{T} stands for time ordering.

To leading order in perturbation theory, the probability density for n measurement events is

$$W_n(x_1, q_1; x_2, q_2; \dots; x_n, q_n) = \int d^4x'_1 \dots d^4x'_n \sigma^{(1)}(x_1 - x'_1) \dots \sigma^{(n)}(x_n - x'_n) P(x'_1, q_1; x'_2, q_2; \dots; x'_n, q_n) \quad (19)$$

where

$$P_n(x_1, q_1; x_2, q_2; \dots; x_n, q_n) = \int d^4y_1 \dots d^4y_n \sqrt{f^{(1)}(y_1)} \dots \sqrt{f^{(n)}(y_n)} R_{(1)}^{a_1 b_1}(y_1, q_1) \dots \times \dots R_{(n)}^{a_n b_n}(y_n, q_n) G_{a_1 \dots a_n, b_1 \dots b_n}(x_1 - \frac{1}{2}y_1, \dots, x_n - \frac{1}{2}y_n; x_1 + \frac{1}{2}y_1, \dots, x_n - \frac{1}{2}y_n). \quad (20)$$

Here $R^{(i)}(x, q)$ is the measurement kernel for the i -th detector. The field correlation function $G_{a_1 \dots a_n, b_1 \dots b_n}(x_1, \dots, x_n; x'_1, \dots, x'_n)$ is given by

$$G_{a_1 \dots a_n, b_1 \dots b_n}(x_1, \dots, x_n; x'_1, \dots, x'_n) = \langle \psi | \mathcal{T}^* [\hat{C}_{b_1}^{(1)}(x'_1) \dots \hat{C}_{b_n}^{(n)}(x'_n)] \times \mathcal{T} [\hat{C}_{a_n}^{(n)}(x_n) \dots \hat{C}_{a_1}^{(1)}(x_1)] | \psi \rangle \quad (21)$$

where \mathcal{T}^* stands for reverse time ordering.

Again we note that in the appropriate regime, the probability becomes independent of the switching functions, and equal to

$$P_n(x_1, q_1; x_2, q_2; \dots; x_n, q_n) = \int d^4 y_1 \dots d^4 y_n R_{(1)}^{a_1 b_1}(y_1, q_1) \dots R_{(n)}^{a_n b_n}(y_n, q_n) \times G_{a_1 \dots a_n, b_1 \dots b_n}(x_1 - \frac{1}{2}y_1, \dots, x_n - \frac{1}{2}y_n; x_1 + \frac{1}{2}y_1, \dots, x_n + \frac{1}{2}y_n) \quad (22)$$

2.6 The detector kernel

The contribution of each detector to the probability is determined by the detector kernel $R^{ab}(x, q)$, defined by Eq. (10). The detector kernel coincides with the matrix elements of a unitary operator,

$$R^{ab}(x, q) = \langle a, q | e^{i\hat{p} \cdot (x-x_0)} | b, q \rangle, \quad (23)$$

where $|a, q\rangle = \sqrt{\hat{\Pi}(q)} \hat{J}^a(x_0) |\Omega\rangle$. If $\hat{\Pi}$ is a projector, then vectors $|a, q\rangle$ with different values of q are orthogonal: $\langle a, q | b, q' \rangle = 0$ for $q \neq q'$.

The Fourier transform of the detector kernel

$$\tilde{R}^{ab}(\xi, q) = \int d^4 x e^{-i\xi \cdot x} R^{ab}(x, q), \quad (24)$$

is given by

$$\tilde{R}^{ab}(\xi, q) = (2\pi)^4 e^{i\xi \cdot x_0} \langle a, q | \hat{E}_\xi | b, q \rangle, \quad (25)$$

where $\hat{E}_\xi = \delta^4(\hat{p} - \xi)$ is the projector onto the subspace with four-momentum ξ^μ , $\langle a, q | \hat{E}_\xi | a, q \rangle \geq 0$. The momentum four vector associated to the detector is timelike and the associated energy p^0 is positive. This means that $\tilde{R}^{ab}(p, q) = 0$ for spacelike p , or if $p^0 < 0$.

For a scalar composite operator $\hat{C}(x)$, we can drop the indices a, b, \dots from the detector kernel and write simply $R(x, q)$. If we record no other observable, we simply write $R(x) := \langle \omega | e^{i\hat{p} \cdot (x-x_0)} | \omega \rangle$, where $|\omega\rangle = \hat{J}^a(x_0) |\Omega\rangle$. Hence, the detector kernel is determined by the energy-momentum distribution of the ‘threshold’ state ω .

2.7 Index notation for probabilities

It is convenient to express the probability densities (22) using an abstract notation. We use small Greek indices $\alpha, \beta, \gamma \dots$ for the pairs (x, a) where x is a spacetime point and a the internal index for the composite operators \hat{C}_a . All indices in a time-ordered product are upper, and all indices in an anti-time-ordered product are lower. Hence, we write the correlation functions (21) as

$$G_{\beta_1 \beta_2 \dots \beta_n}^{\alpha_1 \alpha_2 \dots \alpha_n}$$

We denote by z the pairs (x, q) , where x is a spacetime point and q any other recorded observable or the event \emptyset of no detection. If we denote by Γ the set of possible values for q , then z takes values on the set $Z := M \times \Gamma \cup \{\emptyset\}$. We will write the kernel

$$\sigma[x - \frac{1}{2}(y + y')] \sqrt{f}(y - y') R^{ab}(y - y', q)$$

as $R_\alpha^\beta(z)$ where α stands for (y, a) , β for (y', b) and z for (x, q) . We will use the same symbol for the approximate expression $\delta[x - \frac{1}{2}(y + y')] R^{ab}(y - y', q)$.

Then, the probability formula (22) becomes

$$P_n(z_1, z_2, \dots, z_n) = G_{\beta_1 \beta_2 \dots \beta_n}^{\alpha_1 \alpha_2 \dots \alpha_n} {}^{(1)}R_{\alpha_1}^{\beta_1}(z_1) {}^{(2)}R_{\alpha_2}^{\beta_2}(z_2) \dots {}^{(n)}R_{\alpha_n}^{\beta_n}(z_n). \quad (26)$$

There are two ways of viewing Eq. (26). On one hand, it corresponds to a POVM defined on the Hilbert space \mathcal{F} of the quantum field, with alternatives in the set Z^n . On the other hand, we can consider a vector space \mathcal{S} of smearing functions f_a to the composite operators $\hat{C}^a(x)$, so that $\hat{C}(f) = \int d^4x \hat{C}^a(x) f_a(x)$. The detector kernel is an element of $\mathcal{S} \otimes \mathcal{S}^*$, where \mathcal{S}^* stands for the dual vector space of \mathcal{S} . Then, Eq. (26) defines a positive linear functional on $(\mathcal{S} \otimes \mathcal{S}^*)^n$ that corresponds to the correlation functions $G_{\beta_1 \beta_2 \dots \beta_n}^{\alpha_1 \alpha_2 \dots \alpha_n}$.

3 Relation of the QTP approach to the Closed-Time-Path formalism

The probability density (20) for n measurement events is a linear functional of the $2n$ -point unequal-time correlation function (21). This correlation function has n time-ordered arguments and n arguments in reversed time order. It does not appear in the usual S-matrix description of QFT; the correlation functions in the S-matrix description involve only time-ordered arguments. Rather, the correlation function (21) appears in the Schwinger-Keldysh or Closed-Time-Path (CTP) formalism of QFT.

In what follows, we will review the motivation and function of the CTP formalism, and then we will show how it relates to the QTP probability assignment.

3.1 S-matrix vs CTP correlation functions

By the Dyson formula, the S-matrix for any QFT involves Feynman (i.e., time-ordered) correlation functions of local composite operators $\hat{C}(x)$ of the fields. A common way to generate the Feynman Green functions of a quantum field is from the variational derivatives of a generating functional $Z[J] = \langle 0 | \hat{U}[J] | 0 \rangle$, where

$$\hat{U}[J] = \mathcal{T} \exp \left[i \int d^4 X J^a(X) \hat{C}_a(X) \right]. \quad (27)$$

The generating functional admits the path integral representation

$$Z[J] = \int D\phi e^{i\{S[\Phi] + \int d^4 x J(x) C(x)\}}, \quad (28)$$

where $S[\phi]$ is the classical action functional and $C(x)$ is the classical counterpart of the composite operator $\hat{C}(x)$.

It is convenient to work with the generating functional $W[J] = -i \ln Z[J]$. In perturbation theory, the coefficients of the Taylor expansion of W with respect to J are the connected Feynman graphs. In particular, we may introduce the ‘background’ or ‘mean’ field

$$C_a(x) = \frac{\delta W[J]}{\delta J^a(x)} \quad (29)$$

A more efficient representation of the Green functions is afforded by the Legendre transformation of W

$$\Gamma[C] = W[J] - \int d^4 x J^a(x) C_a(x) \quad (30)$$

since the Taylor expansion of Γ involves only one-particle irreducible Feynman graphs. It is tempting to consider the inverse of (29) as the equation of motion for the background field

$$\frac{\delta \Gamma}{\delta C_a(x)} = -J^a(x). \quad (31)$$

However, this equation is generally complex. This property follows from the fact that the ‘background’ field is not a true expectation value, but rather a matrix element of the field between *in* and *out* states [69].

We can get around this difficulty by adopting the Closed-Time-Path formalism of Schwinger [51] and Keldysh [52]. To this end, we couple the field to two different external sources $J^a(x)$ and $\bar{J}^a(x)$, and we define the CTP generating functional

$$Z_{CTP}[J, \bar{J}] = \langle \psi_0 | \hat{U}^\dagger[\bar{J}] \hat{U}[J] | \psi_0 \rangle, \quad (32)$$

By definition, $Z[J, J] = 1$ and $Z^*[J, \bar{J}] = Z[\bar{J}, J]$. The state $|\psi_0\rangle$ is defined in the distant part, i.e., prior to any time at which $J(x)$ has support—it is an *in* state. Note that a large class of initial states, relevant to particle experiments, can be reconstructed

by the action of source terms of the form (27). Hence, for a large class of problems, the generating function Z_{CTP} for the QFT vacuum $|0\rangle$ contains all physically relevant information.

The CTP generating functional describes correlation functions with n time-ordered and m anti-time-ordered entries,

$$G_{a_1\dots a_m, b_1\dots b_n}^{n,m}(x_1, \dots, x_n; x'_1, \dots, x'_n) = \langle \psi_0 | \mathcal{T}^* [\hat{C}_{b_1}^{(1)}(x'_1) \dots \hat{C}_{b_m}^{(n)}(x'_m)] \times \mathcal{T} [\hat{C}_{a_n}^{(i)}(x_n) \dots \hat{C}_{a_1}^{(1)}(x_1)] | \psi_0 \rangle \quad (33)$$

as functional derivatives of $Z_{CTP}[J, \bar{J}]$,

$$G_{a_1\dots a_m, b_1\dots b_n}^{n,m}(x_1, \dots, x_n; x'_1, \dots, x'_n) = i^{n-m} \left(\frac{\partial^{n+m} Z_{CTP}}{\delta J_1^a(x_1) \dots \delta J^{a_n}(x_n) \delta \bar{J}_1^b(x'_1) \dots \delta \bar{J}^{b_m}(x'_n)} \right)_{J=\bar{J}=0}.$$

For a vacuum initial state, the generating functional has a path integral expression

$$Z_{CTP}[J, \bar{J}] = \int D\phi D\bar{\phi} e^{i\{S[\phi] - S[\bar{\phi}] + \int d^4x [J^a(x)C_a(x) - \bar{J}(x)\bar{C}_a(x)]\}}, \quad (34)$$

where \bar{C} is defined a functional of $\bar{\phi}$. Thus we are led to formally doubling the degrees of freedom and integrating over pairs of histories $(\phi, \bar{\phi})$ that vanish in the asymptotic past and future.

We can actually define two background fields $C_a(x)$ and $\bar{C}_a(x)$ by taking variations with respect to either source,

$$C_a(x) = \frac{\delta W}{\delta J^a(x)} \quad \bar{C}_a(x) = -\frac{\delta W}{\delta \bar{J}^a(x)} \quad (35)$$

where $W[J, \bar{J}] = -i \ln Z_{CTP}[J, \bar{J}]$. The physical situation corresponds to the case when both sources agree. The closed time-path effective action is the double Legendre transform of W

$$\Gamma[C, \bar{C}] = W[J, \bar{J}] - \int d^4x J^a(x)C_a(x) + \int d^4x \bar{J}^a(x)\bar{C}_a(x). \quad (36)$$

The equations of motion now form a coupled system

$$\frac{\delta \Gamma}{\delta C_a(x)} = J_a(x) \quad \frac{\delta \Gamma}{\delta \bar{C}_a(x)} = \bar{J}_a(x) \quad (37)$$

They are identical to each other when $J = \bar{J}$ is the physical external source, and they admit a solution where $C = \bar{C}$ is the physical mean field. Of course, both background fields must be identified only after the variational derivative has been computed [56]. As opposed to the in-out formulation, the equations of motion (37) are always real and causal whereas in cases where the in-out effective action becomes complex, the in-in effective action leads to dissipative evolution. This underscores the fact that the doubling

of degrees of freedom is essential, since no action functional of the physical mean field alone could possibly lead to the right equations of motion.

The CTP effective action formalism can be used to treat fully non-equilibrium dynamics of open quantum systems, going far beyond the traditional linear response theory which are confined to near-equilibrium conditions. In this context, we are often interested in calculating mean-field expectation values $\langle \hat{C}_a(x) \rangle$ that may describe the evolution of classicalized macroscopic field observables.

Observe that quite generally the CTP effective action obeys $\Gamma[C, C] = 0$ and $\Gamma[C, \bar{C}] = -\Gamma[\bar{C}, C]^*$. Therefore, if we separate the CTP effective action in its real and imaginary parts, we find that the former is an odd function of C_a and \bar{C}_a , while the latter is even. Let $C_a^{(0)} = \bar{C}_a^{(0)}$ be an extremum of the CTP effective action, which corresponds to a solution of the mean field equations of motion with zero sources. We take linearized perturbations $\Delta C_a = C_a - C_a^{(0)}$ and $\Delta \bar{C}_a = \bar{C}_a - C_a^{(0)}$. Then, the CTP effective action must take the form

$$\begin{aligned} \Gamma[\Delta C, \Delta \bar{C}] = \frac{1}{2} \int d^d x d^d x' \{ & -[\Delta C_a(x)](x) \mathcal{D}^{ab}(x, x') \{\Delta C_b\}(x') \\ & + i[\Delta C_a](x) N^{ab}(x, x') [\Delta C_b](x') \} \end{aligned} \quad (38)$$

where $[\Delta C_a] = (\Delta C_a - \Delta \bar{C}_a)$, $\{\Delta C_a\} = (\Delta C_a + \Delta \bar{C}_a)$, and \mathcal{D}^{ab} and N^{ab} are two non-local kernels. A seeming redundancy appears because N^{ab} does not contribute to the mean field equations of motion. The latter take the simple

$$\int d^d x' \mathcal{D}^{ab}(x, x') \Delta C_b(x') = J^a(x), \quad (39)$$

where $J^a(x)$ is an external source. The kernel N^{ab} contains information about the *departure* of the actual evolution from the mean-field equation (39). This deviation may be accounted for by coupling the fluctuations to a *stochastic* external source [70–73], rather than a deterministic source like $J^a(x)$ in Eq. (39). This means that the effective classical equations of motion are of the form

$$\int d^d x' \mathcal{D}^{ab}(x, x') \Delta C_b(x') = J^a(x) + \xi^a(x), \quad (40)$$

where $\xi^a(x)$ has vanishing expectation—thus justifying Eq. (39)—but non-zero correlator $\langle \xi(x)^a \xi^b(x') \rangle = N^{ab}(x, x')$. This is why N is referred to as a the *noise kernel*. Hence, CTP allows us to explore how higher-order correlation functions are manifested as noise in the mean-field (classical) equations of motion.

3.2 Measurements and the CTP generating functional

The relation between the QTP description of measurements and the CTP formalism is more transparent, if we use the index notation of Sec. 2.7. For consistency, the sources

J have a lower Greek index, and the sources \bar{J} have an upper Greek index. Then, we write the CTP generating functional as

$$Z_{CTP}[J, \bar{J}] = \sum_{n,m=0}^{\infty} \frac{i^{m-n}}{n!m!} G_{\beta_1 \dots \beta_m}^{\alpha_1 \dots \alpha_n} J_{\alpha_1} \dots J_{\alpha_n} \bar{J}^{\beta_1} \dots \bar{J}^{\beta_m}, \quad (41)$$

or, conversely,

$$G_{\beta_1 \dots \beta_m}^{\alpha_1 \dots \alpha_n} = i^{n-m} \left(\frac{\partial^{n+m} Z_{CTP}[J, \bar{J}]}{\partial J_{\alpha_1} \dots \partial J_{\alpha_n} \partial \bar{J}^{\beta_1} \dots \partial \bar{J}^{\beta_m}} \right)_{J=\bar{J}=0}. \quad (42)$$

The probability densities (26) involve *balanced* correlation functions, i.e., correlation functions with an equal number of upper and lower indices. We can construct a generating functional that contains only such functions. The key observation is that such correlations contribute to the sum only through products of the form $J_{\alpha} \bar{J}^{\beta}$. Hence, the natural source for a diagonal generating functional Z_{CTP}^d that only involves balanced correlation functions is a ‘tensor’ L_{α}^{β} . We define

$$Z_{CTP}^d[L] = \sum_{n=0}^{\infty} \frac{1}{n!} G_{\beta_1 \dots \beta_m}^{\alpha_1 \dots \alpha_n} L_{\alpha_1}^{\beta_1} \dots L_{\alpha_n}^{\beta_n}. \quad (43)$$

Suppose now that we consider only measurements of a single type, i.e., all detector kernels $R_{\alpha}^{\beta}(z)$ are identical. Then, we can define a moment-generating functional for all probability densities (26), in terms of sources $j(z)$,

$$Z_{QTP}[j] = \sum_{n=0}^{\infty} \sum_{z_1, z_2, \dots, z_n} \frac{1}{n!} P_n(z_1, z_2, \dots, z_n) j(z_1) \dots j(z_n). \quad (44)$$

It is straightforward to show that

$$Z_{QTP}[j] = Z_{CTP}^d[R \cdot j], \quad (45)$$

where $(R \cdot j)_A^B = \sum_z R_A^B(z) j(z)$.

Eq. (45) is a fundamental relation for quantum measurements in QFT, as it relates the moment generating functional for a hierarchy of measured probability densities to the generating functional of unequal-time correlation functions

It is straightforward to write a path integral expression for $Z_{CTP}^d[L]$ for a vacuum initial state

$$Z_{CTP}^d[L] = \int D\phi D\bar{\phi} e^{iS[\phi] - iS[\bar{\phi}] + \int d^4x d^4x' C_a(x) \bar{C}_b(x') L^{ab}(x, x')}. \quad (46)$$

To obtain a simple path integral expression for a broader class of states, we recall that many field initial states can be obtained from the action of an external source $\zeta(x)$

on the vacuum, i.e., they are of the form $|\psi_0\rangle = \hat{U}[\zeta]|0\rangle$, where now we write $\hat{U}[\zeta] = \mathcal{T} \exp \left[i \int d^4 X \zeta^k(X) \hat{A}_k(X) \right]$ in terms of composite operators $\hat{A}_k(x)$ that differ, in general from $\hat{C}_a(x)$.

To see this, consider the case of a free scalar field $\hat{\phi}(x)$. Choosing $\hat{A}(x) = \hat{\phi}(x)$, $\hat{U}[\zeta]$ is a Weyl operator, and $|\psi_0\rangle$ is a field coherent state. For single-event measurements, and for any state with a fixed number N of particles, the probability density (16) depends only on the one-particle reduced density matrix $\rho_1(\mathbf{k}, \mathbf{k}') = N^{-1} \langle \psi_0 | \hat{a}_{\mathbf{k}}^\dagger \hat{a}_{\mathbf{k}'} | \psi_0 \rangle$, where $\hat{a}_{\mathbf{k}}$ and $\hat{a}_{\mathbf{k}}^\dagger$ are the creation and annihilation operators of the field. It is straightforward to show that one can reproduce any pure ρ_1 with an appropriate choice of $|\zeta(x)\rangle$. For $\hat{A}(x) =: \hat{\phi}(x)^2$, $\hat{U}[\zeta]$ is a product of a Bogoliubov transformation with a Weyl operator, and, hence, $|\psi_0\rangle$ is a squeezed state. We can reproduce a large class of two-particle reduced density matrices $\rho_2(\mathbf{k}_1, \mathbf{k}_2; \mathbf{k}'_1, \mathbf{k}'_2) = N^{-2} \langle \psi_0 | \hat{a}_{\mathbf{k}_1}^\dagger \hat{a}_{\mathbf{k}_2}^\dagger \hat{a}_{\mathbf{k}'_2} \hat{a}_{\mathbf{k}'_1} | \psi_0 \rangle$, including entangled ones, with appropriate choices of ζ . This

Hence, for a quantum state that is obtained from an external source ζ_k , we write the path integral expression

$$Z_{CTP}^d[f, L] = \int D\phi D\bar{\phi} e^{iS[\phi] - iS[\bar{\phi}] + i \int d^4 x \zeta^k(x) [A_k(x) - \bar{A}_k(x)] + \int d^4 x d^4 x' C_a(x) \bar{C}_b(x') L^{ab}(x, x')}, \quad (47)$$

where we must assume that the spacetime support of the kernel L^{ab} is later than the support of ζ (state preparation is prior to measurement). By Eq. (45)

$$Z_{QTP}[f, j] = \int D\phi D\bar{\phi} e^{iS[\phi] - iS[\bar{\phi}] + i \int d^4 x \zeta^k(x) [A_k(x) - \bar{A}_k(x)]} \times e^{\sum_z \int d^4 x d^4 x' C_a(x) \bar{C}_b(x') R^{ab}(x, x'; z) j(z)} \quad (48)$$

The probability densities for n measurement events are obtained from functional variation of $Z_{QTP}[\zeta, j]$ with respect to j at $j = 0$. For example, the single-event probability density $\hat{P}_1(x, z)$ of Eq. (16) is given by the path integral

$$P_1(x, q) = \int D\phi D\bar{\phi} \left(\int d^4 y C_a(x + \frac{1}{2}y) \bar{C}_b(x - \frac{1}{2}y) K^{ab}(y, q) \right) \times e^{iS[\phi] - iS[\bar{\phi}] + i \int d^4 x \zeta^k(x) [A_k(x) - \bar{A}_k(x)]}.$$

Expressions such as the above provide an explicit link between concepts of quantum measurement theory like POVMs and the practical and highly successful functional language of QFT. We believe that this link is essential for a local and covariant definition of quantum informational notions in QFT.

4 Applications

Next, we present three immediate applications of the QTP formalism. First, we review how the QTP formalism leads to the construction of time of arrival probabilities for

relativistic particles. Second, we derive Unruh-DeWitt detectors from the QTP probability assignment at a particular limit. Third, we identify the regime in which Glauber's photodetection theory applies.

4.1 Time-of-arrival probabilities

The first application of the QTP formalism is the construction of probabilities for the time-of-arrival. The simplest time-of-arrival measurement involves a particle prepared on an initial state $|\psi_0\rangle$ that is localized around $x = 0$ and has positive mean momentum. The issue is to construct the probability $P(L, t)dt$ that the particle is detected at $x = L$ at some moment between t and $t + \delta t$. The problem is that no unique time-of-arrival probability exists [74, 75]. The lack of a self-adjoint operator for time [76] means that we cannot employ Born's rule in order to obtain an unambiguous answer.

A time-of-arrival probability distribution can be constructed from Eq. (16), in absence of any additional observable q . The simplest case, corresponding to a free scalar field $\hat{\phi}(X)$ of mass m , was analysed in Ref. [39]. For a scalar composite operator $\hat{C}(X)$, we obtain a probability density for detection

$$P(x) = \int d^4\xi R(\xi)G(x - \frac{1}{2}\xi, x + \frac{1}{2}\xi). \quad (49)$$

In a time-of-arrival measurement, a detector is placed at a macroscopic distance L from the particle source. If L is much larger than the size of the detector, only particles with momenta along the axis that connects the source to the detector are recorded. Hence, the problem is reduced to two spacetime dimensions. We set the spacetime coordinate $x = (t, L)$. Then, Eq. (49) becomes

$$P(t, L) = \int \frac{dpdp'}{2\pi} \frac{\rho(p, p')}{2\sqrt{\epsilon_p\epsilon_{p'}}} \tilde{R}\left(\frac{p+p'}{2}, \frac{\epsilon_p+\epsilon_{p'}}{2}\right) e^{i(p-p')L - i(\epsilon_p - \epsilon_{p'})t}, \quad (50)$$

where $\epsilon_p = \sqrt{p^2 + m^2}$, and $\tilde{R}(E, p)$ is the Fourier transform of $R(x) = R(t, L)$.

We normalize Eq. (50) by treating L as a parameter and t as a random variable. The procedure is described in Ref. [39]. The end result is a probability distribution

$$P(t, L) = \int \frac{dpdp'}{2\pi} \tilde{\rho}(p, p') \sqrt{v_p v_{p'}} S(p, p') e^{i(p-p')L - i(\epsilon_p - \epsilon_{p'})t}, \quad (51)$$

where $v_p = p/\epsilon_p$ is the particle velocity. $S(p, p')$ are the matrix elements $\langle p|\hat{S}|p'\rangle$ of the *localization operator* \hat{S} , defined by

$$\langle p|\hat{S}|p'\rangle := \frac{\tilde{R}\left(\frac{p+p'}{2}, \frac{\epsilon_p+\epsilon_{p'}}{2}\right)}{\sqrt{\tilde{R}(p, \epsilon_p)\tilde{R}(p', \epsilon_{p'})}}. \quad (52)$$

By definition, $\langle p|\hat{S}|p'\rangle \geq 0$ and $S(p, p) = 1$. Its name originates from the fact that \hat{S} describes the localization of an elementary measurement event.

For a pure state initial state $|\psi\rangle$, Eq. (50) becomes

$$P(t, L) = \langle \psi | \hat{U}^\dagger(t, L) \sqrt{|\hat{v}|} \hat{S} \sqrt{|\hat{v}|} \hat{U}(t, L) | \psi \rangle, \quad (53)$$

where $\hat{U}(t, L)$ is the spacetime-translation operator $\hat{U}(t, L) = e^{i\hat{p}L - i\hat{H}t}$ and $\hat{v} = \hat{p}\hat{H}^{-1}$ is the velocity operator. Eq. (53) defines a positive probability distribution if and only if \hat{S} is a positive operator. It is always satisfied if $\ln \tilde{R}(p, \epsilon_p)$ is a convex function of p . Then, the Cauchy-Schwarz inequality applies,

$$\langle p|\hat{S}|p'\rangle \leq \sqrt{\langle p|\hat{S}|p\rangle \langle p'|\hat{S}|p'\rangle} = 1. \quad (54)$$

Maximal localization is achieved when Eq. (54) is saturated, i.e., for $\langle p|\hat{S}|p'\rangle = 1$. This means that $\hat{S} = \delta(\hat{x})$, where $\hat{x} = i\frac{\partial}{\partial p}$. The probability density (53) for \hat{S} corresponding to maximal localization was first identified by León [77], in terms of the eigenvectors of a non-self-adjoint time-of-arrival operator. In the non-relativistic limit, it reduces to Kijowski's probability distribution [78].

4.2 The limit of a pointlike detector

In a pointlike detector, the detector's world-tube shrinks to a single timelike curve $x_0^\mu(\tau)$, parameterized by its proper time τ . We implement this limit, by expressing the smearing function as

$$F_x(y) = \int d\tau f(\tau - s) \delta^4[y - x_0(\tau)], \quad (55)$$

where $f(\tau)$ is a smearing function peaked around 0, and s is defined by the condition $x_0(s) = x$.

We choose a Gaussian switching function $f(s) = \exp\left[-\frac{s^2}{2\delta_t^2}\right]$. Then, we obtain a probability density for detection $W(q, \tau) = \int ds \sigma(\tau - s) P(q, s)$, where $\sigma(s) = (\sqrt{\pi}\delta_t)^{-1} f^2(s)$, and

$$P(q, \tau) = \int ds \sqrt{f(s)} G_{ab} \left[x_0\left(\tau - \frac{1}{2}s\right), x_0\left(\tau + \frac{1}{2}s\right) \right] \langle \Omega | \hat{J}^a \left[x_0\left(\tau - \frac{1}{2}s\right) \right] \Pi(q) \hat{J}^b \left[x_0\left(\tau + \frac{1}{2}s\right) \right] | \Omega \rangle. \quad (56)$$

The Unruh-DeWitt detection models [25, 26] are a special case of pointlike detectors, in which

$$\hat{J}^a[x_0(\tau)] = e^{i\hat{h}\tau} \hat{J}^a[x_0(0)] e^{-i\hat{h}\tau}, \quad (57)$$

in terms of a Hamiltonian \hat{h} that is defined with respect to the rest frame of the detector. Such Hamiltonians are well defined if the four-velocity $\dot{x}_0^\mu(\tau)$ is a Killing vector of Minkowski spacetime, because then time evolution with respect to τ corresponds to the time translation of a Lorentzian time coordinate. It is not obvious that such a Hamiltonian exists for non-inertial trajectories. In a general QFT, Hamiltonians that generate time translations along non inertial time coordinates do not exist [79, 80]. However, it may be the case that they can be defined as appropriate limits when the spatial size of the system goes to zero, the vanishing of the system's size canceling the ultraviolet divergences that characterize such Hamiltonians. To the best of our knowledge such an analysis has not been carried out. In this sense, the defining assumption for Unruh-DeWitt detectors—for non-inertial trajectories—remains unjustified from first principles.

Assuming the Unruh-DeWitt condition (57), and taking $|\Omega\rangle$ as the lowest eigenstate of \hat{h} , we find

$$P(q, \tau) = \int ds \sqrt{f(s)} G_{ab} [x_0(\tau - \frac{1}{2}s), x_0(\tau + \frac{1}{2}s)] \langle \Omega | \hat{\mu}^a \Pi(q) e^{i\hat{h}s} \hat{\mu}^b | \Omega \rangle, \quad (58)$$

where we wrote $\hat{\mu}^a = \hat{J}^a[x_0(0)]$. If we identify q with the energy ϵ in the rest frame of the detector, we find that

$$P(\epsilon, \tau) = \int ds \sqrt{f(s)} e^{i\epsilon s} G_{ab} [x_0(\tau - \frac{1}{2}s), x_0(\tau + \frac{1}{2}s)] \langle \Omega | \hat{\mu}^a | \epsilon \rangle \langle \epsilon | \hat{\mu}^b | \Omega \rangle. \quad (59)$$

For a scalar coupling operator, where we drop the indices a, b and so on, and for energies such that $\epsilon \delta_t \gg 1$, we write

$$P(\epsilon, \tau) = |\langle \Omega | \hat{\mu} | \epsilon \rangle|^2 \int ds e^{i\epsilon s} G_{ab} [x_0(\tau - \frac{1}{2}s), x_0(\tau + \frac{1}{2}s)], \quad (60)$$

i.e., we obtain the standard formula for the response of the Unruh-DeWitt detector.

Hence, the probabilities of the Unruh-DeWitt detectors for inertial path can be obtained from the QTP probability formula—that is derived from full QFT—at the limit where the world-tube of the detector shrinks to a single timelike curve.

4.3 Derivation of Glauber's theory as a limit

Consider a detector in inertial motion, i.e., the timelike normal of its world-tube is a time-translation vector field in Minkowski spacetime. For simplicity, we consider a quantum scalar field $\hat{\phi}(x)$, with a composite operators $\hat{C}(x) = \hat{\phi}(x)$.

It is convenient to employ coherent states $|z\rangle$ of the scalar field. They are defined as eigenvalues of the annihilation operator

$$\hat{a}_k |z\rangle = z(k) |z\rangle, \quad (61)$$

in terms of a square-integrable complex-valued function $z(k)$; here $k = (\mathbf{k}, \omega_{\mathbf{k}})$ is an on-shell four-vector. We employ the P representation for the initial state

$$\hat{\rho} = \int Dz f_P(z) |z\rangle\langle z|, \quad (62)$$

where $f_P(z)$ is a functional known as the P -symbol of quantum state $\hat{\rho}$ [81]. The functional integration in Eq. (62) is defined in terms of the Gaussian measure for the Bargmann representation of the quantum field [82].

For simplicity, we assume that no magnitude q is recorded. The probability density Eq. (16) is expressed as

$$P(x) = \int Dz f_P(z) P_z(x), \quad (63)$$

where $P_z(x)$ is the probability density Eq. (16) evaluated for a coherent state. We find that P_z is a sum of three contributions

$$P_z(x) = P_z^{(0)}(x) + P_z^{(1)}(X) + P_z^{(2)}(x), \quad (64)$$

where

$$\begin{aligned} P_z^{(0)}(X) &= \int \frac{d^3k}{(2\pi)^3} \frac{\tilde{R}(k)}{2\omega_{\mathbf{k}}} \\ P_z^{(1)}(x) &= \int \frac{d^3k}{(2\pi)^3} \frac{d^3k'}{\sqrt{2\omega_{\mathbf{k}}}} \frac{d^3k'}{(2\pi)^3 \sqrt{2\omega'_{\mathbf{k}}}} z(k) z^*(k') e^{i(k-k') \cdot x} \tilde{R}\left(\frac{k+k'}{2}\right) \\ P_z^{(2)}(x) &= 2\text{Re} \int \frac{d^3k}{(2\pi)^3 \sqrt{2\omega_{\mathbf{k}}}} \frac{d^3k'}{(2\pi)^3 \sqrt{2\omega'_{\mathbf{k}}}} z(k) z(k') e^{i(k+k') \cdot x} \tilde{R}\left(\frac{k'-k}{2}\right) \end{aligned} \quad (65)$$

The meaning of the three terms in Eq. (64) is the following.

- The term $P_z^{(0)}$ is state-independent. It corresponds to a background noise of ‘false alarms’ in the detector and its contribution to the detection probability is negligible.
- The term $P_z^{(1)}$ is generated by the terms in the density matrix that corresponds to a fixed number of particles, i.e., by the restriction of the density matrix in the N particle subspaces.
- The term $P_z^{(2)}$ involves contributions from terms with different numbers of photons.

The second term in Eq. (64) involves a double momentum integral over $e^{-i(\omega_k - \omega_{k'})\tau}$ (co-rotating terms), while the third term involves a double integral over $e^{-i(\omega_k + \omega_{k'})\tau}$ (counter-rotating terms). The dominant contributions to the double integral arise from

values of k and k' that correspond to a slowly varying phase. This implies that co-rotating terms typically dominate over counter-rotating ones. It is therefore meaningful to invoke the Rotating Wave Approximation (RWA) and drop the contribution of the counter rotating terms.

The RWA is a defining approximation of Glauber's theory of photo-detection. The domain of validity of the RWA in photodetection has been a matter of some debate. Some authors have shown that photodetection probabilities that are obtained with the RWA appear to violate causality at short times and they proposed corrections to Glauber's theory [83,84]. Others indicate that a different form of the RWA in photodetection theory can guarantee causality [85]. A fundamental problem of the RWA is that it corresponds to a Hamiltonian that has no lower bound, and thus it is thermodynamically unacceptable [86]. Regarding the degree of accuracy of the RWA and related approximations, see, Ref. [87] and references therein.

To examine the validity of the RWA, we consider a coherent state that corresponds to a pulse of mean wave-number \mathbf{k}_0 localized around $\mathbf{x} = 0$ at $t = 0$. We consider massless particles (i.e., scalar photons). We choose a Gaussian

$$z(k) = (2\pi)^3 z_0 (2\pi\Delta^2)^{-3/2} e^{-\frac{(\mathbf{k}-\mathbf{k}_0)^2}{2\Delta^2}}, \quad (66)$$

where z_0 is a constant complex valued vector and Δ is the spectral width of the pulse.

First, we evaluate the term $P_z^{(1)}(\tau, \mathbf{Q})$ in Eq. (64). We employ a saddle point approximation. We expand $\omega_k = \omega_{k_0} + \mathbf{v}_{k_0} \cdot (\mathbf{k} - \mathbf{k}_0)$, where $\mathbf{v}_k = \mathbf{k}/|\mathbf{k}|$ is the three-velocity vector for a photon. Then, we assume that \tilde{R} varies slowly with k , so that the contribution of the degradation function is a constant $\tilde{R}(k_0)$, we find

$$P_z^{(1)}(t, \mathbf{x}) = \frac{1}{2} |z_0|^2 \tilde{R}(k_0) e^{-\Delta^2(\mathbf{x} - \mathbf{v}_{k_0} t)^2}. \quad (67)$$

We also evaluate the term $P_z^{(2)}$ in the saddle-point approximation. We find

$$P_z^{(2)}(t, \mathbf{x}) \sim \text{Re} [z_0^2 e^{ik_0 \cdot x}]. \quad (68)$$

The key point here is that the oscillatory terms in Eq. (68) must be averaged over a spatial region of size δ_x^3 and for times of order δ_t . For Gaussian smearing functions, this averaging leads to suppression factors of order $e^{-\delta^2 |\mathbf{k}_0|^2 e^{-\sigma^2 \omega_{k_0}^2}}$. Thus, the contribution of the term $P_z^{(2)}$ to the probability density (64) is suppressed, and the RWA is justified. We expect that the suppression of counter-rotating terms due to spatial and temporal coarse graining holds for general initial states that represent almost monochromatic wave packets.

The RWA is equivalent with the expression of the detection probability as

$$P(x) = \int d^4 y R(y) G_{RWA}[x - \frac{1}{2}y, x + \frac{1}{2}y], \quad (69)$$

in terms of the correlation function

$$G_{RWA}(x, x') = 2\langle\Psi|\hat{\phi}^{(-)}(x')\hat{\phi}^{(+)}(x)|\Psi\rangle, \quad (70)$$

where $\hat{\phi}^{(+)}$ and $\hat{\phi}^{(-)}$ denote the positive and negative frequency part of $\hat{\phi}$.

Suppose now that the spacetime spread of $R(x)$ is much smaller than the spatial and temporal variation of the field. For massless particles, this is a stronger form of the dipole approximation that is often employed in quantum optics: we assume that the typical wavelengths and frequencies in the initial state are much larger than the size of the detector and the associated time spread of detection. Then, we can approximate $R(x)$ with a delta function, to obtain

$$P(x) \sim G_{RWA}[x, x]. \quad (71)$$

This expression coincided with the standard formula of Glauber's photo-detection theory.

For a single measurement event we recover Glauber's probabilities by assuming (i) the RWA and (ii) $R(y) \sim \delta(y)$. As we showed, for a large class of initial states, the RWA is not an independent assumption, as it arises due to the spatial and temporal coarse-graining of the detector for a large class of initial states. Thus, the crucial assumption of Glauber's theory is that the characteristic scales of the detector kernel are much smaller than any time / length parameter that characterize the initial state of the field. A more detailed analysis, with an emphasis on the divergence from Glauber's theory, will be undertaken elsewhere.

5 Links to non-equilibrium QFT

In Sec. 2, we saw that the QTP probabilities are linear functional of balanced correlation functions. The measurements do not probe unbalanced correlation functions. Since the latter include $\langle\hat{C}_a(x)\rangle$, QTP probabilities cannot access mean field information. This limitation is not fundamental. Remember that the operator $\hat{C}(x)$ appears in the interaction term with the apparatus. This restriction means that we cannot use couplings of the form $\int d^4x \hat{C}_a(x) \otimes \hat{J}^a(X)$, in order to directly measure the operator $\hat{C}_a(x)$. At least such measurements are not possible with weak field-apparatus coupling where perturbation theory is applicable⁴.

Suppose, for example that \hat{C} coincides with the field operator $\hat{\phi}$ —we drop the index a for simplicity. Then, single-detector probabilities record only local information about particles. Let the field is in a state with a macroscopically large number of particles, so that it can be viewed as a thermodynamic system. Then the single-detector probability essentially coincides with a particle-number density function. If we also measure the recorded particle's momentum k , the QTP probability density $P(x, k)$ is an operationally

⁴It is possible to measure $\hat{C}_a(x)$ for non-perturbative couplings [91], but the resulting probabilities are very different from those of QTP, and they are closer to those of von Neumann measurements.

defined version of Boltzmann’s distribution function. By Eq. (16), $P(x, k)$ is a linear functional of the correlation function $G(x, x') = \langle \hat{\phi}(x) \hat{\phi}(x') \rangle$, which is usually taken to satisfy the Baym-Kadanoff equations.

From the above analysis, it follows that Boltzmann’s thermodynamic entropy

$$S_B(\Sigma) = - \int_{\Sigma} d^3x d^3k P(x, k) \ln P(x, k) \quad (72)$$

where Σ is a Cauchy surface is a Shannon-type entropy for single-detection measurements. This means that quantum informational quantities, defined through measurements, have a direct application to non-equilibrium QFT. Furthermore, n -detector QTP probabilities probe higher-order correlation functions of the quantum field, thus allowing an analysis that is not accessible by traditional methods.

Such an analysis is beyond the scope of this paper. In this section, we will only describe some structural similarities and connections between the QTP analysis and methods of non-equilibrium QFT. A detailed analysis will be taken up in other publications.

5.1 Stochastic correlation dynamics from two-particle irreducible effective action

As shown in Sec. 3.2, the generating functional of QTP correlation functions is defined in terms of non-local source terms $L^{ab}(x, x')$. It is structurally similar to the two-particle irreducible effective action (2PIEA) [55, 94] that has found many applications in non-equilibrium QFT—see, for example, [58–60].

We proceed to a brief review of the 2PIEA formalism following [95, 96], emphasizing how it can be used to define evolution equations with noise from higher-order correlation functions.

For ease of notation, we use a version of DeWitt condensed notation where capital indices A correspond to both the spacetime dependence and the branch of the CTP field (forward or backward in time, ϕ or $\bar{\phi}$). Hence, we will be writing ϕ_A , C_A , and so on. Hence, the action in the CTP generating functional will be $S[\phi_A] = S[\phi_a] - S[\bar{\phi}_a]$. In the two-particle irreducible representation the (two-point) correlation function stands as an independent variable apart from the mean field. Thus there is a separate source K^{AB} driving $C_A C_B$ over the usual $J^A C_A$ term in the one-particle irreducible representation—see the similarity to Eq. (48).

From the generating functional

$$Z[K^{AB}] = e^{iW[K^{AB}]} = \int D\phi_A e^{i\{S + \frac{1}{2}K^{AB}C_A C_B\}} \quad (73)$$

we have

$$G^{AB} = \langle \hat{C}_A \hat{C}_B \rangle = 2 \left. \frac{\delta W}{\delta K^{AB}} \right|_{K=0} \quad (74)$$

and

$$\frac{\delta^2 W}{\delta K^{AB} \delta K^{CD}} \Big|_{K=0} = \frac{i}{4} \left\{ \langle \hat{C}_A \hat{C}_B \hat{C}_C \hat{C}_D \rangle - \langle \hat{C}_A \hat{C}_B \rangle \langle \hat{C}_C \hat{C}_D \rangle \right\}. \quad (75)$$

Suppose that we want to express the effective dynamics of G_{AB} in a closed form, but to go beyond the Baym-Kadanoff equations, by taking into account noise from higher order correlations. For an non-equilibrium system, we seek a formulation in terms of a new object \mathbf{G}_{AB} , namely, a stochastic correlation function whose expectation value over the noise average gives the usual two point functions while its fluctuations reproduce the quantum fluctuations in the binary products of field operators. The simplest assumption is to take \mathbf{G}_{AB} as a Gaussian process that is defined formally by the relationships

$$\langle \mathbf{G}_{AB} \rangle = \langle \hat{C}_A \hat{C}_B \rangle; \quad \langle \mathbf{G}_{AB} \mathbf{G}_{CD} \rangle = \langle \hat{C}_A \hat{C}_B \hat{C}_C \hat{C}_D \rangle \quad (76)$$

The Legendre transform of W is the 2PIEA,

$$\Gamma_{2PI} [G_{AB}] = W [K^{AB}] - \frac{1}{2} K^{AB} G_{AB}; \quad K^{AB} = -2 \frac{\delta \Gamma}{\delta G_{AB}} \quad (77)$$

The Schwinger-Dyson equation for the propagators is simply, $\frac{\delta \Gamma_{2PI}}{\delta G_{AB}} = 0$. When including the stochastic source \mathbf{G}_{AB} , it becomes

$$\frac{\delta \Gamma_{2PI}}{\delta \mathbf{G}_{AB}} = -\frac{1}{2} \kappa^{AB} \quad (78)$$

where κ_{ab} is a stochastic nonlocal Gaussian source defined by

$$\langle \kappa^{AB} \rangle = 0; \quad \langle \kappa^{AB} \kappa^{CD} \rangle = 4i \left[\frac{\delta^2 \Gamma_{2PI}}{\delta G_{AB} \delta G_{CD}} \right]^\dagger \quad (79)$$

It is well established that the noiseless Eq. (78) ($\kappa = 0$) can be used as a basis for the derivation of transport equations in the near equilibrium limit. Indeed, for a $\lambda\phi^4$ type theory, the resulting equation is simply the Boltzmann equation for a distribution function f defined from the Wigner transform of G^{ab} . The full stochastic equation (78) leads, in the same limit, to a Boltzmann - Langevin equation [95], thus providing the microscopic basis for this equation in a manifestly relativistic quantum field theory.

5.2 Correlation Histories

The two-particle irreducible formalism can be extended to an n-particle irreducible formalism, for any n . There is an effective action Γ_{nPI} for each n , from which all effective actions for $n' < n$ can be derived. Taking $n \rightarrow \infty$, we obtain a master effective action. The functional variation of the master effective action yields the hierarchy of Schwinger-Dyson equations [96].

To obtain effective closed dynamics for the correlations at order n , we must truncate the Schwinger-Dyson hierarchy upon this order. Truncation renders the master effective

action complex. Its imaginary part arises from correlation functions of order higher than n , the fluctuations of which Calzetta and Hu define as *correlation noises* [95] at order n . For example, the noise κ^{AB} in Eq. (78) is the correlation noise of order two.

Calzetta and Hu defined the notion of *correlation histories* [97], in analogy to the decoherent histories program. A fine-grained correlation history corresponds to the full Schwinger-Dyson hierarchy of correlation functions. When we truncate the hierarchy at finite order n , we treat only correlation functions of order n as independent. Higher-order correlations are ignored or slaved to the lowest-order ones. A truncated hierarchy defines a *coarse-grained* correlation history. For example, mean field theory studies *coarse-grained* correlation histories at order $n = 1$; the Baym-Kadanoff equation, or Boltzmann equation and their stochastic generalizations refer to *coarse-grained* correlation histories of order $n = 2$.

The key point is that the truncation of the master effective action always leads to dissipation and noise for the coarse-grained histories. Any truncated theory is an effective field theory in the correlation hierarchy formulation. This effective field theory carries only partial information. The missing information resides in the correlation noise, and manifests as dissipation in the subsystem dynamics. The higher-order correlations are analogous to an environment in the theory of open quantum systems [98]. This noise may lead to decoherence of correlation histories [97], i.e., to the classicalization of the effective description.

The QTP approach demonstrates that the different levels of correlation histories can be accessed by the measurement of n -detector joint probabilities. Eq. (26) assigns to each initial state $|\psi\rangle$ of the field a hierarchy of joint probability distributions $P_n(z_1, z_2, \dots, z_n)$.

In classical probability theory, a hierarchy of correlation functions defines a classical stochastic process, if it satisfies the Kolmogorov additivity condition,

$$P_{n-1}(z_1, \dots, z_{n-1}) = \int dz_n P_n(z_1, z_2, \dots, z_n) \quad (80)$$

Quantum probability distributions for sequential measurements do not satisfy this condition [99]. Hence, the violation of Eq. (80) is a genuine signature of quantum dynamics; it cannot be reproduced by classical physics, including classical stochastic processes. It is rather different from the Leggett-Garg inequalities [100] that also refer to the behavior of quantum multi-time probabilities. Some authors refer to the violation of Eq. (80) as "temporal non-locality" [101]. In contrast, if measurements on a quantum field approximately satisfy Eq. (80), then the measurement outcomes can be simulated by a stochastic process with n -time probabilities given by the probability distributions (80). Then the generating functional Z_{QTP} corresponds to a stochastic process, i.e., it is obtained as the functional Laplace transform of a classical stochastic probability measure.

Hence, the hierarchy $P_n(z_1, z_2, \dots, z_n)$ provides a natural and unambiguous classicality criterion⁵. Given the relation between QTP probabilities and QFT correlation

⁵For a different classicality criterion, based on the decoherent histories approach, that applies to the special case of measurements of the field operator, see [102].

functions, this criterion can be used to probe the information content of different levels for correlation histories. For example, the validity of Eq. (80) is necessary for deriving deterministic or classical stochastic dynamics for $P_1(z)$, i.e., for the validity of the Boltzmann or the Boltzmann-Langevin equation. The failure of (80) means that the four-point correlation functions are too ‘quantum’ to allow effective classical stochastic dynamics for the two-point correlation function.

Conversely, the failure of Eq. (80) can be used to provide a measure of irreducibly quantum information at the level $n = 2$. An example of such a measure is

$$S_Q = \int dz_2 \left| \int dz_1 P_2(z_1, z_2) - P_1(z_2) \right|. \quad (81)$$

A second informational quantity is the correlation of the probability distribution, i.e., a measure of the deviation of $P_2(z_1, z_2)$ from $P_1(z_1)P_1(z_2)$ ⁶. This information is typically quantified by the correlation entropy

$$S_C = \int dz_1 dz_2 P_2(z_1, z_2) \ln \frac{P_2(z_1, z_2)}{P_1(z_1)P_1(z_2)}. \quad (82)$$

In general, the correlation entropy will contain information for both quantum correlations (if $S_Q \neq 0$) and classical stochastic ones. Indeed, S_C may not have the usual properties of correlation entropy if $S_Q \neq 0$, and other measures that will distinguish will be more convenient. The third relevant informational quantity in Boltzmann’s entropy (72), defined in terms of $P_1(z_1)$. These three quantities are the most important for describing the flow of information at the level of the 2PIEA.

Hence, the QTP hierarchy functions as a registrar of information of the quantum system, keeping track of how much information resides in what order, and how it flows from one order to another through the dynamics. There is a good potential for this scheme to systemize quantum information in QFT⁷: keeping track of the contents and the flow of information and measuring the degree of coherence in a quantum system.

6 Conclusions

In this work, we presented the QTP formalism for measurements in quantum fields, and we explored its relations to the Closed-Time-Path description of QFT. These relations

⁶The factorization assumption of the joint probability for two events is equivalent to Boltzmann’s Stosszahlansatz in the derivation of Boltzmann’s equation through the Bogoliubov–Born–Green–Kirkwood–Yvon hierarchy [103]. The joint probability density P_2 is then slaved to P_1

⁷Of course, several issues remain to be addressed. The probability distributions P_n depend on the measurement apparatus through the detector kernels. We would like to avoid strong apparatus dependence when deriving informational quantities. We may have to consider ideal, maximal information, apparatuses (like the ones appearing in the study of the time-of-arrival), or look for quantities that can be minimized with respect to all physical detector kernels.

provide a direct translation between the operational language of measurement theory (POVMs, effects, and so on) to the manifestly covariant description of QFT through functional methods. This is an essential step towards formulating a general theory of relativistic quantum information.

Our work highlights the central role of the hierarchy of unequal-time correlation functions. On one hand, POVMs for measurement are linear functional of such correlation functions, on the other hand, these correlation functions are crucial for the implementation of perturbation theory, but also give real causal dynamics in the CTP approach. When working at the level of non-equilibrium QFT, correlation functions are employed in order to define both observables and effective irreversible dynamics.

Equally important is the fact that correlation functions are covariant and causal objects. For this reason, we contend that a relativistic QIT that respects both causality and spacetime symmetry must define all informational quantities in terms of such correlation functions. This contrasts the standard approach of QIT that is based on properties of single-time quantum states, like von Neumann entropy or entanglement.

Building a sound theoretical foundation for relativistic QIT is not only important for theoretical completeness. It is also needed for describing quantum experiments in space [2,3] that will explore the effects of non-inertial motion (acceleration, rotation) and gravity on quantum resources like entanglement. Corrections to standard photodetection models, like the ones predicted by QTP, may be measurable in experiments that involve long separations or large relative velocities between detectors.

Acknowledgements

Research was supported by grant JSF-19-07-0001 from the Julian Schwinger Foundation.

References

- [1] C. Anastopoulos and N. Savvidou, *Quantum Information in Relativity: The Challenge of QFT Measurements*, Entropy 24, 4 (2022).
- [2] D. Rideout et al, *Fundamental Quantum Optics Experiments Conceivable with Satellites – Reaching Relativistic Distances and Velocities*, Class. Quantum Grav. 29, 224011 (2012).
- [3] M. Mohageg et al, *The Deep Space Quantum Link: Prospective Fundamental Physics Experiments using Long-Baseline Quantum Optics*, arXiv:2111.15591.
- [4] C. Anastopoulos and B. L. Hu, *Probing a Gravitational Cat State*, Class. Quant. Grav. 32, 165022 (2015).
- [5] S. Bose, A. Mazumdar, G. W. Morley, H. Ulbricht, M. Toro, M. Paternostro, A. A. Geraci, P. F. Barker, M. S. Kim, and G. Milburn, *A Spin Entanglement Witness for Quantum Gravity*, Phys. Rev. Lett. 119, 240401 (2017).

- [6] C. Marletto and V. Vedral, *Gravitationally Induced Entanglement between Two Massive Particles is Sufficient Evidence of Quantum Effects in Gravity*, Phys. Rev. Lett. 119, 240402 (2017).
- [7] C. Anastopoulos and B. L. Hu, *Quantum Superposition of Two Gravitational Cat States*, Class. Quant. Grav. 37, 235012 (2020).
- [8] I. Bloch, *Some Relativistic Oddities in the Quantum Theory of Observation*, Phys. Rev. 156, 137 (1967).
- [9] Y. Aharonov and D. Z. Albert, *Can we make sense out of the measurement process in relativistic quantum mechanics?*, Phys. Rev. D4, 359 (1981).
- [10] A. Peres and D. Terno, *Quantum Information and Relativity Theory*, Rev. Mod. Phys. 76, 93 (2004).
- [11] R. M F. Houtappel, H. van Dam, and E. P. Wigner, *The Conceptual Basis and Use of the Geometric Invariance Principles*, Rev. Mod. Phys. 37, 595 (1965).
- [12] D. Malament, *In Defense of Dogma: Why There Cannot Be A Relativistic Quantum Mechanics of (Localizable) Particles*, in "Perspectives on Quantum Reality", ed. R. Clifton (Kluwer Academic, Dordrecht 1996).
- [13] G. C. Hegerfeldt, *Instantaneous Spreading and Einstein Causality in Quantum Theory*, Annalen der Physik 7, 716 (1998).
- [14] R. Sorkin, *Impossible Measurements on Quantum Fields*, in "Directions in General Relativity", eds. B. L. Hu and T. A. Jacobson (Cambridge University Press, 1993).
- [15] A. Peres, *Classical interventions in quantum systems. II. Relativistic invariance*, Phys. Rev. A61, 022117 (2000).
- [16] S. Popescu, and L. Vaidman, *Causality constraints on nonlocal quantum measurements*, Phys. Rev. A49, 4331 (1994).
- [17] L. Vaidman, *Instantaneous Measurement of Nonlocal Variables*, Phys. Rev. Lett. 90, 010402 (2003).
- [18] D. Beckman, D. Gottesman, M. A. Nielsen, J. Preskill, *Causal and localizable quantum operations*, Phys. Rev. A64, 052309 (2001).
- [19] D. Grimmer, *The Pragmatic QFT Measurement Problem and the need for a Heisenberg-like Cut in QFT*, arXiv:2205.09608.
- [20] L. Landau and R. Peierls, *Erweiterung des Unbestimmtheitsprinzips für die Relativistische Quantentheorie*, Zeit. Phys. 69, 56 (1931).
- [21] N. Bohr and L. Rosenfeld, *On the Question of the Measurability of Electromagnetic Field Quantities*, Mat.-fys. Medd. Dan. Vid. Selsk.12, (1933).
- [22] R. J. Glauber, *The Quantum Theory of Optical Coherence*, Phys. Rev. 130, 2529 (1963).
- [23] R. J. Glauber, *Coherent and Incoherent States of the Radiation Field*, Phys. Rev. 131, 2766 (1963).
- [24] M. O. Scully and M. S. Zubairy, *Quantum Optics* (Cambridge University Press, 2012).
- [25] W. G. Unruh, *Notes on Black Hole Evaporation*, Phys. Rev. D14, 870 (1976).
- [26] B. S. DeWitt, *Quantum Gravity: the New Synthesis* in General Relativity: An Einstein Centenary Survey, ed. by S. W. Hawking and W. Israel (Cambridge University Press, Cambridge, 1979), p. 680.

- [27] J. Polo-Gómez, L. J. Garay, L. J. and E. Martín-Martínez, *A detector-based measurement theory for quantum field theory*, Phys. Rev. D 105, 065003 (2022).
- [28] B. L. Hu, S-Y Lin, J. Louko, *Relativistic Quantum Information in Detectors–Field Interactions*, Class. Quantum Grav. 29, 224005 (2012).
- [29] M. Cliche and A.Kempf, *The relativistic quantum channel of communication through field quanta*, Phys. Rev. A 81, 012330 (2010).
- [30] E. Martin-Martinez, *Causality issues of particle detector models in QFT and Quantum Optics*, Phys. Rev. D 92, 104019 (2015).
- [31] J. de Ramon, M. Papageorgiou and E. Martin-Martinez, *Relativistic causality in particle detector models: Faster-than-light signalling and "Impossible measurements"*, Phys. Rev. D 103, 085002 (2021).
- [32] K. E. Hellwig and K. Kraus, *Formal description of measurements in local quantum field theory*, Phys. Rev. D 1, 566 (1970).
- [33] K. Okamura and M. Ozawa, *Measurement theory in local quantum physics*, J. Math. Phys. 57, 015209 (2015).
- [34] S. Doplicher, *The measurement process in local quantum physics and the EPR paradox*, Commun. Math. Phys. 357, 407 (2018).
- [35] C. J. Fewster and R. Verch, *Quantum fields and local measurements*, Comm. Math. Phys. 378, 851 (2020).
- [36] H. Bostelmann, C. J. Fewster and M. H. Ruep, *Impossible measurements require impossible apparatus*, Phys. Rev. D103, 025017 (2021).
- [37] C. Anastopoulos and N. Savvidou, *Time-of-Arrival Probabilities for General Particle Detectors*, Phys. Rev. A86, 012111 (2012).
- [38] C. Anastopoulos and N. Savvidou, *Time-of-Arrival Correlations*, Phys. Rev. A95, 032105 (2017).
- [39] C. Anastopoulos and N. Savvidou, *Time of arrival and Localization of Relativistic Particles*, J. Math. Phys. 60, 0323301 (2019).
- [40] C. Anastopoulos and N. Savvidou, *Time-of-arrival Probabilities and Quantum Measurements*, J. Math. Phys. 47, 122106 (2006).
- [41] C. Anastopoulos and N. Savvidou, *Time-of-arrival probabilities and quantum measurements. II. Application to tunneling times*, J. Math. Phys. 49, 022101 (2008).
- [42] C. Anastopoulos, *Time-of-arrival probabilities and quantum measurements. III. Decay of unstable states*, J. Math. Phys. 49, 022103 (2008).
- [43] K. Savvidou, *The Action Operator for Continuous-time Histories* J. Math. Phys. 40, 5657 (1999); *Continuous Time in Consistent Histories*, gr-qc/9912076.
- [44] N. Savvidou, *Space-time Symmetries in Histories Canonical Gravity*, in "Approaches to Quantum Gravity", edited by D. Oriti (Cambridge University Press, Cambridge 2009).
- [45] R. B. Griffiths, *Consistent Quantum Theory* (Cambridge University Press, Cambridge 2003).
- [46] R. Omnés, *The Interpretation of Quantum Mechanics*, (Princeton University Press, Princeton 1994).

- [47] R. Omnés, *Understanding Quantum Mechanics* (Princeton University Press, Princeton 1999).
- [48] M. Gell-Mann and J. B. Hartle, *Quantum Mechanics in the Light of Quantum Cosmology*, in ‘Complexity, Entropy, and the Physics of Information’, ed. by W. Zurek, (Addison Wesley, Reading 1990);
- [49] M. Gell-Mann and J. B. Hartle, *Classical Equations for Quantum Systems*, Phys. Rev. D47, 3345 (1993).
- [50] J.B. Hartle, *Spacetime Quantum Mechanics and the Quantum Mechanics of Spacetime* in ‘Gravitation and Quantizations’, in the Proceedings of the 1992 Les Houches Summer School, ed. by B. Julia and J. Zinn-Justin, Les Houches Summer School Proceedings, Vol. LVII, (North Holland, Amsterdam, 1995); [gr-qc/9304006].
- [51] J. S. Schwinger, *Brownian Motion of a Quantum Oscillator*, J. Math. Phys. 2, 407 (1961).
- [52] L. V. Keldysh, *Diagram Technique for Nonequilibrium Processes*, Zh. Eksp. Teor. Fiz. 47, 1515 (1964).
- [53] G. Zhou, Z. Su, B. Hao and L. Yu, *Equilibrium and nonequilibrium formalisms made unified*, Phys. Rep. 118, 1 (1985).
- [54] B. S. DeWitt, *Effective action for expectation values*, in ”Quantum Concepts in Space and Time”, ed. R. Penrose and C. J. Isham (Clarendon Press, Oxford, 1986)
- [55] E. Calzetta and B. L. Hu, *Nonequilibrium quantum fields: Closed-time-path effective action, Wigner function, and Boltzmann equation*, Phys. Rev. D37, 2878 (1988).
- [56] E. Calzetta and B. L. Hu, *Closed-time-path functional formalism in curved spacetime*, Phys. Rev. D35, 495 (1987).
- [57] S. Weinberg, *Quantum contributions to cosmological correlations* Phys. Rev. D 72, 043514 (2005).
- [58] E. A. Calzetta and B. L. Hu, *Nonequilibrium quantum field theory* (Cambridge University Press, 2008).
- [59] J. Berges, *Introduction to nonequilibrium quantum field theory*, AIP Conf. Proc. 739, 3 (2004).
- [60] J. Berges, *Nonequilibrium Quantum Fields: From Cold Atoms to Cosmology*, Lecture Notes of the Les Houches Summer School: Strongly Interacting Quantum Systems out of Equilibrium (Oxford University Press, Oxford, 2016).
- [61] A. Kamenev, *Field Theory of Non-Equilibrium Systems* (Cambridge University Press, Cambridge, 2011).
- [62] J. Rammer, *Quantum Field Theory of Non-equilibrium States* (Cambridge University Press, Cambridge, 2009)
- [63] B. L. Hu, *Correlation Dynamics of Quantum Fields and Black Hole Information Paradox*, talk at the International School of Astro-fundamental Physics, Sept. 1995. Proceedings edited by N. Sanchez and Zichichi (Kluwer Publishers, Dordrecht, 1996).
- [64] E. Calzetta and B. L. Hu, ‘*Noise and Fluctuations in Semiclassical Gravity*’, Phys. Rev. D49, 6636 (1994).
- [65] B. L. Hu and A. Matacz, *Backreaction in Semiclassical Gravity: the Einstein-Langevin Equation*, Phys. Rev. 51 (1995).

- [66] E. Chitambar, D. Leung, L. Mancinska, M. Ozols, and A. Winter, *Everything You Always Wanted to Know About LOCC (But Were Afraid to Ask)*, Commun. Math. Phys., 328, 303 (2014).
- [67] S. Weinberg, *The Quantum Theory of Fields: I. Foundations* (Cambridge University Press, Cambridge, 1996).
- [68] C. Anastopoulos and N. Savvidou, *Measurements on relativistic quantum fields: I. Probability assignment*, arXiv:1509.01837.
- [69] J. Hartle and B. L. Hu, *Quantum effects in the early universe. II. Effective action for scalar fields in homogeneous cosmologies with small anisotropy*, Phys. Rev. D20, 1772 (1979).
- [70] R. Feynman and F. Vernon, *The Theory of a general quantum system interacting with a linear dissipative system*, Ann. Phys. (NY) **24**, 118 (1963).
- [71] R. Feynman and A. Hibbs, *Quantum Mechanics and Path Integrals*, (McGraw - Hill, New York, 1965).
- [72] A. O. Caldeira and A. J. Leggett, *Path integral approach to quantum Brownian motion*, Physica **121A**, 587 (1983).
- [73] B. L. Hu, J. P. Paz and Y. Zhang, *Quantum Brownian motion in a general environment: Exact master equation with nonlocal dissipation and colored noise*, Phys. Rev. **D45**, 2843 (1992).
- [74] J. C. Muga and J. R. Leavens, *Arrival time in quantum mechanics*, Phys. Rep. 338, 353 (2000).
- [75] J. C. Muga, R. S. Mayato, and I. L. Equisquiza, *Time in Quantum Mechanics, vol 1* (Springer 2008); J. G. Muga, A Ruschhaupt and A. Del Campo, *Time in Quantum Mechanics, vol 2* (Springer 2010).
- [76] W. Pauli, *The Principles of Quantum Mechanics*, in *Encyclopedia of Physics*, edited by S. Flugge, Vol. 5/1 (Springer, Berlin, 1958).
- [77] J. León, *Time-of-arrival formalism for the relativistic particle*, J. Phys A: Math. Gen. 30, 4791 (1997).
- [78] J. Kijowski, *On the time operator in quantum mechanics and the Heisenberg uncertainty relation for energy and time*, Rep. Math. Phys. 6, 361 (1974).
- [79] A. D. Helfer, *The stress - energy operator*, Class. Quantum Grav. 13, L129 (1996).
- [80] C. G. Torre and M. Varadarajan, *Functional Evolution of Free Quantum Fields*, Class. Quantum. Grav. 16, 2651 (1999).
- [81] JR Klauder and B Skagerstam, *Coherent States: Applications in Physics and Mathematical Physics* (World Scientific, Singapore 1985).
- [82] F. A. Berezin, *The Method of Second Quantization* (Academic Press, 1966).
- [83] V.P. Bykov and V.I. Tatarskii, *Causality violation in the Glauber theory of photodetection*, Phys. Lett. A 136, 77 (1989).
- [84] M. Fleischhauer, *Quantum-theory of photodetection without the rotating wave approximation* J. Phys. A: Math. Gen. 31, 453 (1998).
- [85] P. W. Milonni, D. F. V. James, and H. Fearn, *Photodetection and causality in quantum optics*, Phys. Rev. A 52, 1525 (1995).

- [86] G. W. Ford and R. F. O'Connell, *The rotating wave approximation (RWA) of quantum optics: serious defect*, Physica A243, 377 (1997).
- [87] C. Fleming, N. I. Cummings, C. Anastopoulos and B. L. Hu, *The Rotating-Wave Approximation: Consistency and Applicability from an Open Quantum System Analysis*, J. Phys. A: Math. Theor. 43, 405304 (2010).
- [88] C. Anastopoulos and N. Savvidou, *Quantum Temporal Probabilities in Tunneling Systems*, Ann. Phys. 336, 281 (2013).
- [89] C. Anastopoulos and N. Savvidou, *Path of a Tunneling Particle*, Phys Rev. A95 , 052120 (2017).
- [90] C. Anastopoulos and N. Savvidou, *Coherences of Accelerated Detectors and the Local Character of the Unruh Effect*, J. Math. Phys. 53, 012107 (2012).
- [91] M. Papageorgiou and C. Anastopoulos, *Field observables recorded by Unruh-Dewitt detectors* (in preparation).
- [92] G. Baym and L. P. Kadanoff, *Conservation Laws and Correlation Functions*, Phys. Rev. 124, 287 (1961).
- [93] G. Baym and L. P. Kadanoff, *Quantum Statistical Mechanics*, (Benjamin, New York 1962).
- [94] S. A. Ramsey, and B. L. Hu, B.L., *O(N) quantum fields in curved spacetime*, Phys. Rev. D56, 661 (1997).
- [95] E. Calzetta and B. L. Hu, *Stochastic dynamics of correlations in quantum field theory: From the Schwinger-Dyson to Boltzmann-Langevin equation*, Phys. Rev. D61, 025012 (1999).
- [96] E. Calzetta and B. L. Hu, "Correlations, Decoherence, Dissipation and Noise in Quantum Field Theory", in *Heat Kernel Techniques and Quantum Gravity*, ed. S. A. Fulling (Texas A& M Press, College Station 1995).
- [97] E. Calzetta and B. L. Hu, "Decoherence of Correlation Histories" in *Directions in General Relativity, Vol II: Brill Festschrift*, eds B. L. Hu and T. A. Jacobson (Cambridge University Press, Cambridge, 1993).
- [98] C. Anastopoulos, *Coarse grainings and irreversibility in quantum field theory*, Phys. Rev. D56, 1009 (1997).
- [99] C. Anastopoulos, *Classical versus quantum probability in sequential measurements*, Found. Phys. 36, 1601 (2006).
- [100] A. J. Leggett and A. Garg, *Quantum mechanics versus macroscopic realism: Is the flux there when nobody looks?*, Phys. Rev. Lett. 54, 857 (1985).
- [101] J. Kofler and Č. Brukner, *Condition for macroscopic realism beyond the Leggett-Garg inequalities*, Phys. Rev. A 87, 052115 (2013).
- [102] C. Anastopoulos, *Quantum Correlation Functions and the Classical Limit*, C. Anastopoulos, Phys. Rev. D63, 125024 (2001).
- [103] R. Balescu, *Equilibrium and Nonequilibrium Statistical Mechanics* (John Wiley, New York, 1975).